

The impact of the rice production system (irrigated vs lowland) on root-associated microbiome from farmer's fields in western Burkina Faso

Mariam Barro^{1,2,3}, Issa Wonni², Marie Simonin^{1,4}, Abalo Itolou Kassankogno², Agnieszka Klonowska^{1,2}, Lionel Moulin¹, Gilles Béna¹, Irénée Somda³, Caroline Brunel^{1,5,§}, Charlotte Tollenaere^{1,*,§}

¹PHIM Plant Health Institute, Univ Montpellier, IRD, CIRAD, INRAE, Institut Agro, Montpellier, 34394, France

²INERA, Institut de l'Environnement et de Recherches Agricoles du Burkina Faso, Laboratoire de Phytopathologie, Bobo-Dioulasso, 01, Burkina Faso

³Univ Nazi Boni, Institut du Développement rural, Laboratoire des Systèmes naturels, Agrosystèmes et Ingénierie de l'Environnement (SyNAIE), Bobo-Dioulasso, 01, Burkina Faso

⁴Present Address: Univ Angers, Institut Agro, INRAE, IRHS, SFR QUASAV, F-49000 Angers, France

⁵Present Address: CIRAD, UPR HortSys, F-34398, Montpellier, F-49000 Angers, France

*Corresponding author: PHIM (Plant Health Institute of Montpellier). IRD (Institut de Recherche pour le Développement) - 911, avenue Agropolis - BP64501 - 34394 Montpellier cedex 05 - France. E-mail: charlotte.tollenaere@ird.fr

One sentence summary: Irrigated rice and rainfed lowlands harbor contrasted root-associated microbiomes in small farmer's fields from Burkina Faso, with contrasted richness, and peculiar indicator taxa in each system.

Editor: Angela Sessitsch

[§]equal contribution.

ABSTRACT

Due to their potential applications for food safety, there is a growing interest in rice root-associated microbial communities, but some systems remain understudied. Here, we compare the assemblage of root-associated microbiota in rice sampled in 19 small farmer's fields from irrigated and rainfed lowlands in Burkina Faso, using an amplicon metabarcoding approach of the 16S rRNA gene (prokaryotes, three plant samples per field) and ITS (fungi, one sample per field). In addition to the expected structure by root compartments (root vs rhizosphere) and geographical zones, we showed that the rice production system is a major driver of microbiome structure. In irrigated systems, we found a higher diversity of prokaryotic communities from the rhizosphere and more complex co-occurrence networks, compared to rainfed lowlands, while fungal communities exhibited an opposite pattern (higher richness in rainfed lowlands). Core taxa were different between the two systems, and indicator species were identified: mostly within *Bacillaceae* in rainfed lowlands, and within *Burkholderiaceae* and *Moraxellaceae* in irrigated areas. Finally, a higher abundance in rainfed lowlands was found for mycorrhizal fungi (both compartments) and rhizobia (rhizosphere only). Our results highlight deep microbiome differences induced by contrasted rice production systems that should consequently be considered for microbial engineering applications.

Keywords: irrigated rice, rainfed lowlands; root associated microorganisms; metabarcoding, *Oryza sativa*, rice production system, West Africa

INTRODUCTION

Soil and rhizosphere host megadiverse and dynamic communities of microorganisms that are crucial to the plants they associate with. Their role is particularly recognized for crops as the below-ground microbiota supply plants with nutrients and provide protection against pathogens (Singh et al. 2020, Chialva et al. 2021). Recent research suggests that root-associated microbes can improve plant tolerance to environmental stressors (Chialva et al. 2020), and modify phenology (Lu et al. 2018) and morphological traits (Kumar et al. 2018). Cultivated plants and their associated microbial communities are thus increasingly studied jointly, as holobionts (a concept reviewed by Vandenkoornhuysen et al. in 2015), because a deeper understanding of their interaction might help to develop microbial engineering applications for modern sustainable agricultural systems (Chialva et al. 2021). While much progress has been made, and despite the growing interest in crop microbiome, the mechanisms that control root-associated microbiome assembly (i.e. structure, composition, and dynamics) remain difficult to disentangle (Brunel et al. 2020).

Rice is the most important food crop in the world, grown in variable climatic conditions and representing the staple food of more than half of the world's population, mostly in Asia, Africa, and Latin America (Pandey et al. 2010). The major species cultivated worldwide is *Oryza sativa* L. (known as 'Asian rice'). This is also true in Africa, where a second rice species was domesticated (*O. glaberrima*, referred to as 'African rice'), but is much less grown because of lower yield (Linares 2002). Given its importance for food security, and the impact of microbiota on plant productivity, metabarcoding approaches are more and more used to describe the microbiome of *O. sativa* (Kim and Lee 2020), particularly its root-associated microbial communities (reviewed by Ding et al. 2019). Rice has the particularity to be cultivated in flooded paddy soils over most growth stages, so that the rhizosphere is located in an oxic-anoxic interface (Ding et al. 2019). Compared to other crops, rice root microbial communities are particularly enriched in *Deltaproteobacteria*, *Euryarchaeota*, *Chytridiomycota* (Ding et al. 2019). Communities' structuring is driven both by the host plant (in terms of root compartment/microhabitats,

Received: March 25, 2022. Revised: July 4, 2022. Accepted: July 20, 2022

© The Author(s) 2022. Published by Oxford University Press on behalf of FEMS. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com

plant genotype) and its environment (geographical zone, bioclimate, soil properties, agricultural practices; Ding et al. 2019). The host genotype has been shown to structure rice associated microbial communities (Edwards et al. 2015, Alonso et al. 2020), but its influence is generally weak compared to environmental factors (Edwards et al. 2018, Guo et al. 2021). For instance, Edwards et al. (2015) evidenced important effects of pedoclimate and agricultural practices. Other environmental factors, such as drought stress (Santos-Medellín et al. 2017), water management (Chialva et al. 2020), phosphorus (Long and Yao 2020), were also shown to affect rice root-associated microbiota. However, the rice microbiome has been poorly explored in the African context in spite of (i) the importance to document the soil diversity as shaped by cultural practices, and (ii) the growing importance of rice in Africa (Soullier et al. 2020). An exception is the recent work by Kanasugi et al. (2020) that evidenced an effect of the region in structuring of rice microbiome described in six tropic savanna regions in Ghana. More generally, there is a lack of knowledge concerning crop-associated microbiota in the African continent, that results in a biased view of the microbial world associated with crops due to an unbalanced worldwide sampling repartition (Brunel et al. 2020, Hughes et al. 2021).

Rice is grown around the world in three dominant growing systems (Rao et al. 2017). First, irrigated lowlands, with full water control, produce 75% of the global rice production. Second, rainfed lowlands (including flood prone), represents around 19% of the world's rice production. Finally, rainfed upland rice, only possible under high rainfall, results in 4% of the global total rice production. In Burkina Faso, irrigated rice represents small areas (costly infrastructures representing less than 30% of harvested areas; CountrySTAT 2020), but produces more than half of the production, because of a relatively high productivity, up to 4 to 7 t/ha per year (MAHRH 2011). Because of lower yields (for example, only 2 t/ha per year was estimated in Dano, Serpantié et al. 2019), rainfed lowlands that cover a wider area (ca. 70%), represent only 42% of the production (MAHRH 2011). The full water control is the typical feature that distinguishes irrigated areas (IR) and rainfed lowlands (RL). In addition, different agricultural practices characterize each system in West Africa (see Nonvide et al. 2018 in Benin). In western Burkina Faso, these contrasted practices have been documented with three main differences: (i) the possibility to grow rice twice a year is restricted to irrigated areas, (ii) mineral fertilization is more frequent in irrigated areas, and (iii) direct sowing is performed only in rainfed lowlands (Barro et al. 2021a). The effect of these two contrasted production systems on rice microbiota have never been investigated.

This study aims at describing rice root-associated microbial communities in farmer's fields from western Burkina Faso. More specifically, we investigate whether rice roots from two contrasted rice growing systems (irrigated and rainfed lowland areas) host different microbial communities. Our study focused on prokaryote communities, investigated through 16S rRNA sequencing of three samples per analyzed field, and we also characterized fungal communities by ITS sequencing, with a lower sample size (one sample per field, see methods). We collected rice root and rhizosphere samples in farmer's fields from three geographical zones, each consisting of an irrigated rice-growing area and neighboring rainfed lowlands. A previous study performed in the same study fields documented more intensive agricultural practices in irrigated areas, compared to rainfed lowlands (e.g. mineral fertilization, two rice cropping per year; Barro et al. 2021a). Considering the effect of intensification on belowground biodiversity and microbial network complexity (Banerjee et al. 2019, Tamburini et al.

2020), we hypothesize an effect of the rice growing system on root-associated microbial communities. If true, this may have consequences on plant health (Wei et al. 2015), particularly in this system where rice diseases were shown to circulate at higher levels in irrigated areas compared to rainfed lowlands (Barro et al. 2021a).

MATERIAL AND METHODS

Study sites in western Burkina Faso

The study sites are located in three geographic zones in western Burkina Faso, with maximum distance between each zone about 90 kilometers (Fig. 1A). Each zone comprises one irrigated area and the neighboring rainfed lowland, with maximum distance between the two rice growing systems of each zone being 7 kilometers (Fig. 1B). The climate consequently do not differ between rice growing systems within each zone, but average precipitation during the rice growing season (July to early December) differ between the three geographical zones (WorldClim 2 data; Fick and Hijmans 2017; see Fig. S1). The three geographical zones were not studied for themselves, but instead considered as replicates to investigate how rice growing systems affect microbial communities in general. These six sites were studied from 2016 to 2019, with the characterization of agricultural practices and the follow-up of major rice diseases symptoms (Barro et al. 2021a; see further details on the methodology and raw data at: <https://doi.org/10.23708/8FDWIE>).

Rice root sampling

Within the six sites, we investigated a total of 19 fields, with three fields per site, except in Bama, the largest irrigated perimeter studied, where four fields were sampled (Fig. 1B). Rice genotyping data are available for these 19 fields, with as much as 18 different rice genotypes (only two fields with the same rice genotype), illustrating the high rice genetic diversity in the area (see Barro et al. 2021b).

Each field studied corresponds to a square of approximately 25 m on a side. Root and rhizosphere sampling was performed at rice maturation, between October, 16th and December 3rd 2018. We chose this developmental stage based on a previous study showing that the rice microbiome composition evolves during the growing season until a 'mature' microbiome at the flowering stage (Edwards et al. 2018). Within each field, we sampled three plants located on the square diagonal, with at least 5 meters distance. These three plants per field were also inspected visually and specific symptoms of the major rice diseases were reported (as in Barro et al. 2021a). Sampling was performed with gloves and scissors (ethanol sanitized between two samples) and involved nodal, bases and seminal roots (all sampled together). Roots were roughly shaken to remove non-adherent soil, and placed in 50 ml sterile tubes containing sterile Phosphate Buffered Saline (PBS) solution for a rapid (15 s) rinse and then stored in another 50 ml sterile PBS-containing tube. We placed the tubes in a cooler and then at 4°C when back to the laboratory, on the same day.

Soil physicochemical properties: data acquisition and analysis

The three geographical zones studied (Fig. 1A) are characterized by Lixisols (soils with subsurface accumulation of low activity clays and high base saturation) according to the Harmonized World Soil Database (HWSD) map (FAO/IIASA/ISRIC/ISSCAS/JRC 2012).




Figure 1. Location of the study sites and soil physico-chemical properties. (A) Location of the study sites in western Burkina Faso on the soil Harmonized World Soil Database (HWSD) map of West Africa (<https://webarchive.iiasa.ac.at/Research/LUC/External-World-soil-database/HTML/>). The three geographical zones studied are in Lixisols (LX: soils with subsurface accumulation of low activity clays and high base saturation). (B) Location of the field studied within each geographical zones. Soil texture, are indicated for each of the 19 studied fields, with pie charts representing relative proportions of sand (in yellow), silt (in green), and clay (in blue). (C) Soil chemical properties estimated in each geographical zones, with colors representing the rice growing system (irrigated areas in blue and rainfed lowlands in red). Each point corresponds to one field studied. Averages over the 19 studied fields are indicated for each parameter, as well as the results of statistical tests for the geographical zone effect.

Soil sampling was performed on the same day as rice root sampling, and in three locations nearby sampled plants, using a 10 cm depth auger. Back from the field, sampled soil was dried in the shade at ambient temperature and stored until analysis.

INERA/GRN service performed the analyses of soil samples according to a standardized methodology. Briefly, soil physical properties were assessed by soil particle size distribution following Bouyoucos (1962), pH was estimated according to AFNOR (1981), total organic carbon with the Walkley and Black (1934)'s method, total concentrations of nitrogen with Kjeldhal method Hillebrand et al. (1953), and finally phosphorus and potassium content as described respectively in Novozansky et al. (1983) and Walinga et al. (1989). Then, cation exchange capacity (CEC), a measure of fertility, nutrient retention capacity, and the capacity to protect groundwater from cation contamination, was estimated, as well as sorptive bioaccessibility extraction (SBE), that relates to the environmental mobility, partitioning and toxicity of soil pollutants, following Metson (1956). Soil data are publicly available on the IRD Dataverse: <https://doi.org/10.23708/LZ8A5B>.

Rice root conditioning, DNA extraction, and sequencing

Less than 24 h after sampling, root samples (rice roots including rhizosphere) stored at 4°C were processed. In order to separate the different root compartments, the tubes were vortexed vigorously one minute and then, roots were removed from the PBS solution using sterile forceps. The remaining PBS solution was considered as the 'rhizosphere' compartment. Roots were then surface-sterilized with 70% alcohol (30 s), 1% bleach (30 s) and finally rinsed three times in sterile water. We considered these surface-sterilized roots as the 'root' compartment, which comprises both endosphere microorganisms as well as persistent DNA from the rhizoplane. DNA extraction from the rhizosphere and root samples were performed on the same day as the process of compartment separation.

For DNA extractions, 0.25 g of root samples (crushed in liquid nitrogen beforehand) and 0.25 g of rhizosphere were extracted using the DNeasy PowerSoil Kit (Qiagen, Hilden, Germany), following manufacturer's recommendations. DNA quality and quantity was verified using a NanoDrop ND-1000 spectrophotometer. PCR amplification, library, and MiSeq Illumina sequencing were performed by Macrogen (Seoul, South Korea) using primers 341F (16S V3F, 5'-CCTACGGGNGGCWGCAG-3') and 785R (16S V4R, 5'-GACTACHVGGGTATCTAATCC-3') to amplify the V3-V4 regions of the 16S rRNA gene (Thijs et al. 2017), and using primers ITS1f: CTTGGTCATTTAGAGGAAGTAA and ITS2: GCTGCGTTCTTCATC-GATGC to amplify the internal transcribed spacer 1 (ITS1) region (Op De Beeck et al. 2014). We specifically focused on the factors structuring prokaryotic communities (16S rRNA gene sequencing), but were also interested to see whether similar tendencies hold for fungal communities (ITS sequencing). We consequently chose the following approach: sequencing was performed for each sampled plant (19 fields * 3 plants = 57 samples per compartment) for 16S rRNA gene sequencing and for a composite sample (three plant samples were pooled to result in one sample per field, so that the total number of samples per compartment is 19) for the ITS marker. Negative controls (three for 16S rRNA gene and one for ITS) were sequenced to remove potential contaminants.

Sequence data are retrievable from NCBI (National Center for Biotechnology Information) under the Bioproject ID: PR-JNA763095.

Bioinformatic analyses of obtained sequences

All bioinformatics and statistical analyses were performed in R software v 3.6.3 (R core Team 2018) and the package *ggplot2* (Wickham 2016) was used for the visualization.

Raw sequences were processed using a custom script from the *dada2* pipeline, which is designed to resolve exact biological sequences (ASVs for Amplicon Sequence Variants) from Illumina sequence data without sequence clustering (Callahan et al. 2016). Raw sequences were first demultiplexed by comparing index reads with a key. Primers and adapters were screened and removed using a custom script with *cutadapt* (Martin 2011). Paired sequences were trimmed. Sequences were dereplicated, and the unique sequence pairs were denoised using the *dada* function. Next, paired-end sequences were merged, and chimeras were removed. Taxonomy assignments were determined against the SILVA SSU r138 (Quast et al. 2013) and the UNITE 2021 (Abarenkov et al. 2021) taxonomic databases for 16S rRNA gene and ITS, respectively using the *idtaxa* function from the *decipher* R package (Wright 2016). Mitochondria and chloroplast sequences were then removed, and contaminants were identified and removed using negative controls and the *decontam* package (Davis et al. 2018). Rarefaction curves were drawn for each sample, using the *rarecurve* function of the *vegan* R package (Oksanen et al. 2007), and the rarefaction plateau was reached for all samples (Fig. S2). Two filtering steps were performed, first, to account for differences in sequencing depths, samples were rarefied to 4236 and 7828, for 16S rRNA gene and ITS, respectively removing 1092 prokaryote and 350 fungal ASVs. Then, ASVs not seen more than once in at least 2% of the samples were removed. We then obtained 2116969 (8260 ASVs) final sequences for 16S rRNA gene and 172719 (566 ASVs) sequences for ITS. A figure showing the phyla relative abundances was constructed and presented according to the rice-growing system, the geographic zone and the root compartment.

Indices of α -diversity (observed richness and Shannon diversity index) were calculated using the *estimate_richness* function from the *phyloseq* package (McMurdie and Holmes 2013). Members of the core microbiota were identified for 16S rRNA gene and ITS communities (including both rhizosphere and roots compartments) in each rice grown system using the prevalence threshold of 60% of samples. This threshold allows identifying the ecological core microbiome in the soil heterogeneous matrix considering the degree of accuracy generated by the processing into exact sequences (ASV) while ensuring not to miss lower abundance taxa that have fallen below the threshold of detection in one or more samples, but are of functional and ecological importance (Neu et al. 2021).

For 16S rRNA gene dataset only, we inferred co-occurrence networks using the *SpiecEasi* pipeline (Kurtz et al. 2015), independently for each rice growing system: rainfed lowland and irrigated areas. Networks were computed for ASVs present in more than 15% of the samples to meet the data requirement to draw correlations. They were calculated using the method 'mb' and setting the *lambda.min.ratio* to 1e-3 and *nlambda* to 50. We identified hub taxa, i.e. the potential keystone of the microbial network belonging to the most connected ASVs, based on their node parameters (method adapted from Berry and Widder 2014): a low betweenness centrality (lower quantile, < 0.9), and a high closeness centrality (higher quantile, > 0.75), transitivity (higher quantile, > 0.25) and degree (higher quantile, > 0.75). The node and network parameters were determined using the R package *igraph* (Csardi and Nepusz 2006) and *qgraph* (Epskamp et al. 2012). Complete networks were further described by calculating the number of nodes and

hubs, the network mean degree, mean closeness and betweenness centralities, the total number of edges, and the positive to negative edges ratio. The taxonomic affiliation of all ASVs was refined using nucleotide basic local alignment search tool (BLASTn) analyses on NCBI nr database. We screened the table of blast best hits of all 16S rRNA gene and ITS ASVs (present in the rarefied but unfiltered dataset) in order to search the genus or species names of a number of pathogen species listed a priori (Table S1), based on the reference book *Compendium of Rice Diseases* (Cartwright et al. 2018). We also searched within blast best hits for ITS ASVs assigned to the *Glomeromycetes* class, as well as 16S rRNA gene and ITS ASV assigned to a species name's including 'rhizob' to identify rhizobia, and including 'oryz', as this may correspond to a specific interaction (pathogen or beneficial) with rice.

Statistical analyses

We first analyzed soil physico-chemical parameters. To this purpose, PERMANOVA were performed on soil physical properties (texture, i.e. relative amount of sand, silt, and clay) on the one hand, and soil chemical properties (seven variables: pH, total carbon, total nitrogen, total phosphorus, total potassium, SBE, CEC) on the other hand. For both models, we included as explanatory factors the 'geographical zone' and 'rice-growing system', as well as their interaction, using *adonis2* function from the *vegan* R package, with 999 permutations. Posthoc tests were done using the *pairwiseAdonis* function (<https://github.com/pmartinezarbizu/pairwiseAdonis>). In addition, we performed linear models (*lm* function) on each soil variable independently, after having tested for normality (function *shapiro.test*) function, and transformed the variable (using the package *bestNormalize*, Peterson 2021) if required. The linear models included as explanatory variables: the rice growing system, the geographical zone, and their interaction (the last one being removed from the final model if revealed non-significant). Correction for multiple testing (10 soil parameters) was performed using the FDR (false discovery rate) method, using the function *p.adjust* (option method 'BH') from the package *stats* (R core Team 2018).

For microbial root and rhizosphere communities, respectively, PERMANOVA based on a Bray–Curtis dissimilarity matrix were used to test for significant effects geographical zones, rice growing systems and their interaction on β -diversity. If significant, differences were further tested using the *pairwise.adonis* function of the *pairwiseAdonis* package (version 0.4). The graphical representation of β -diversity was based on Non-metric Multi-Dimensional Scaling (NMDS, *metaMDS* function). The effect of edaphic variables (i.e. pH, total organic carbon, total phosphorus, total nitrogen, total potassium, CEC, and SBE) in structuring β -diversity was tested using the *envfit* function (9999 permutations) in R package *vegan*. Structuring soil properties were thus fitted onto the ordination space.

We tested for an effect of the rice growing system, the geographical zone and their interaction on obtained indices of alpha diversity, using linear models for Shannon diversity indices, and generalized linear models (Poisson distribution) for observed richness. Models used for ITS data set did not include random effect (function *lm* for Shannon index and *glm* for observed richness). On the other hand, as three samples of different plants but same rice fields were included in the 16S data set, we built generalized linear mixed models (GLMMs) for prokaryote kingdom to include the particular field as random factor. The package *lme4* (Bates et al. 2015) was used for this purpose (function *lmer* for Shannon index and *glmer* for observed richness). These models were run indepen-

dently for each compartment (rhizosphere on the one hand, and root associated on the other) and were analyzed by Type III ANOVA with the package *car* (Fox and Weisberg 2019).

Then, we identified particular soil taxa that were associated with rainfed lowlands or irrigated systems using indicator analyses with the function *multipatt* implemented in the *indicspecies* package (De Cáceres, Legendre and Moretti 2010). The algorithm determines both fidelity and consistency to a system.

Finally, we analyzed the repartition of ASVs belonging to rhizobia, and assigned to the class *Glomeromycetes* (arbuscular mycorrhizal fungi, AMFs). The summed abundance of all ASVs which best blast hit is within the two groups were modeled with Poisson distribution, and analyzed with Type III ANOVA with the package *car*. Explanatory variables included the rice growing system, the geographical zone, and their interaction. For ITS data (*Glomeromycetes*), GLMs were used with no random factor. For 16S rRNA gene data (rhizobia), GLMMs were built, including the particular field as a random factor.

R code used to perform the analyses and generate the figures are available upon request.

RESULTS

Structure of rice soil properties and rice root associated microbial communities


PERMANOVA analysis performed on the Bray–Curtis distance matrix of soil characteristics to describe the overall soil properties, highlighted no significant influence of the rice growing system, but a differentiation according to the geographical zone, for both physical ($F = 6.420$, $r^2 = 0.448$, $P = 0.006$, Fig. 1B) and chemical ($F = 4.121$, $r^2 = 0.346$, $P = 0.026$) soil parameters. More precisely, we found no effect of the rice growing system, but a significant effect of the geographical zone on the clay and sand contents, as well as total Phosphorus, total Potassium, SBE, and CEC (Table S2). Posthoc tests revealed that the geographic zones that differed statistically were the same for the six above-mentioned variables, namely, Bazon and Karfiguela zones (Fig. 1C).

To determine whether the geographical zones (Karfiguela, Bama, or Bazon), the rice-growing systems (irrigated vs. rainfed lowlands), or their interactions, structured root-associated or rhizosphere microbial communities, we performed PERMANOVA analysis on the Bray–Curtis distance matrix of 16S rRNA gene and ITS ASVs, respectively. As different communities' structures were revealed between root and rhizosphere for both prokaryotes ($F = 6.863$, $r^2 = 0.052$, $P < 0.001$) and fungi ($F = 3.753$, $r^2 = 0.080$, $P < 0.001$), we further subsetted 16S and ITS data sets to better observe the relative influence of the rice-growing systems and the geographical zones in shaping root and rhizosphere communities separately (Table 1, Fig. 2).

In the rhizosphere, prokaryotic communities were structured by the rice growing system, the geographical zone and the interaction between rice growing system and zone (Table 1). Posthoc tests revealed that all pairs of sites (i.e. irrigated and lowland from the same geographical zone) were significantly different, but interestingly revealed no significant difference between communities originating from irrigated systems, whereas all communities from rainfed lowland sites exhibited distinct structures (Table S3). Fungal communities of the rhizosphere were mostly structured by the rice growing system (Table 1). The geographical zone and the interaction between rice growing system and the geographical zone were also driving the rhizosphere fungal microbiome. The lower number of samples did not allow to detect, if any, statistically sig-

Table 1. Results of PERMANOVA analysis performed independently for rhizosphere compartment and root compartment, for 16S and ITS microbiome data.

| | | Prokaryotes | | | | | 16S | | | | |
|-------------|--------------------|-------------|----------|-------|-------|--------|-----|----------|-------|-------|--------|
| | | Df | SumOfSqs | R2 | F | Pr(>F) | Df | SumOfSqs | R2 | F | Pr(>F) |
| Rhizosphere | Rice system | 1 | 1.890 | 0.079 | 5.096 | 0.001 | 1 | 0,687 | 0,115 | 2,452 | 0,001 |
| | Zone | 2 | 1.646 | 0.069 | 2.218 | 0.001 | 2 | 0,872 | 0,146 | 1,555 | 0,007 |
| | Rice system * Zone | 2 | 1.393 | 0.058 | 1.877 | 0.001 | 2 | 0,748 | 0,126 | 1,335 | 0,036 |
| | Residual | 51 | 18.919 | 0.793 | NA | NA | 13 | 3,644 | 0,612 | NA | NA |
| | Total | 56 | 23.848 | 1.000 | NA | NA | 18 | 5,951 | 1,000 | NA | NA |
| Root | Rice system | 1 | 1.614 | 0.079 | 5.155 | 0.001 | 1 | 0,846 | 0,111 | 2,289 | 0,002 |
| | Zone | 2 | 1.407 | 0.069 | 2.247 | 0.001 | 2 | 0,891 | 0,117 | 1,206 | 0,123 |
| | Rice system * Zone | 2 | 1.535 | 0.075 | 2.451 | 0.002 | 2 | 1,076 | 0,141 | 1,457 | 0,013 |
| | Residual | 51 | 15.966 | 0.778 | NA | NA | 13 | 4,803 | 0,631 | NA | NA |
| | Total | 56 | 20.522 | 1.000 | NA | NA | 18 | 7,616 | 1,000 | NA | NA |

**Figure 2.** NMDS ordination showing the three factors identified as drivers of the structuration of rice root microbial communities: the color of points represent the rice growing system (irrigated vs. rainfed lowland), while the shape shows the geographical zone (Banzon, Karfiguela and Bama). On the left side are presented the analyses based on 16S rRNA gene reflecting Prokaryote communities, where one point corresponds to one plant. On the right side are shown the analyses based on ITS reflecting fungal communities, where one point corresponds to one field. The root compartment is presented on the upper side of the figure, while the rhizosphere data appear on the bottom side. Only the soil physicochemical parameter that revealed as having a significant effect (see Supplementary Table S4) are represented with arrows: cation exchange capacity (CEC), Sorptive Bioaccessibility Extraction (SBE), total concentrations of phosphorus (Total P) and total concentrations of nitrogen (Total N).

nificant differences between sites in communities' structures for ITS (Table S3).

As observed for rhizosphere communities, the root-associated prokaryotic communities were mainly shaped by the rice growing system, the interaction between rice growing system and the geographical zone and the geographical zone (Table 1). For all pairs of sites (within the same geographical zone), irrigated areas and rainfed lowlands were significantly different. No significant difference was detected between communities originating from irrigated systems, whereas in two out of three comparisons, communities from rainfed lowland sites exhibited distinct struc-

tures (Table S3). Root-associated fungal communities were also mostly influenced by the rice growing system and by the interaction between rice growing system and the geographical zone (Table 1). The effect of the geographical zone (three zones) on root-associated fungal communities was not evidenced. As for rhizosphere, posthoc tests on root-associated fungal communities were all non-significant (Table S3).

The influence of soil chemical parameters on microbial community structure is reported in Fig. 2 as arrows and in Table S4. We noticed that the prokaryotic communities of both rhizosphere and roots were affected by the same three parameters: SBE ($r^2 = 0.482$,

$P < 0.001$ for rhizosphere, and $r^2 = 0.175$, $P = 0.006$ for roots), CEC ($r^2 = 0.314$, $P < 0.001$ for rhizosphere, and $r^2 = 0.204$, $P = 0.003$ for roots) and total phosphorus ($r^2 = 0.132$, $P = 0.023$ for rhizosphere, and $r^2 = 0.179$, $P = 0.004$ for roots). For fungi, although various parameters were marginally significant in each compartment (Table S4), we only detected a significant effect of total nitrogen on rhizosphere communities ($r^2 = 0.320$, $P = 0.043$).

Composition of rice root microbiomes and comparison of alpha-diversity

While 16S rRNA gene data were assigned at the genus level for 64% of ASVs, only 34% of ITS ASVs could be assigned (see assignments at the phylum level in Fig. S4). Assignations at the phylum level were obtained for all (100%) 16S ASVs, but only for 62% for ITS ASVs (see Fig. S3). Assigned prokaryotic taxa represent 17 phyla, most abundant ones being Proteobacteria, Firmicutes, Mixococcota and Acidobacteriota. For ITS, seven phyla were found, with the most abundant ones being Ascomycota followed by Basidiomycota.

We tested the effect of the rice growing system on the diversity indices (alpha-diversity). No effect of the rice growing system could be evidenced on the root-associated compartment, although the interaction between rice growing system and geographical zone significantly affected observed richness for both prokaryotes and fungi (Table 2). In addition, the rice growing system had a significant effect on the prokaryote diversity of the rhizosphere (Table 2), with a higher Shannon diversity index in irrigated areas (5.03 ± 0.13), compared to rainfed lowlands (4.39 ± 0.12) and a higher observed richness (275.6 ± 26.4) in irrigated areas compared to rainfed lowlands (143.7 ± 17.4) (Fig. 3). An opposite pattern was found for fungal communities of the rhizosphere with higher observed richness in rainfed lowlands (75.78 ± 4.34), compared to irrigated areas (63.80 ± 4.51) (Table 2 and Fig. 3).

Core microbiome and co-occurrence networks in the two rice-growing systems

ASVs belonging to the core microbiome of lowland vs irrigated rice were respectively identified with a prevalence threshold set to 60%. For the 16S rRNA gene, we identified 26 core ASVs associated with the irrigated systems, and two core ASVs in lowlands (Fig. 4). Among the core taxa in irrigated areas, the vast majority of phylotypes (25/26) belonged to the Burkholderiaceae family, with 24 assigned to *Ralstonia pickettii* and one to *Paraburkholderia kururiensis*. One of the core ASVs is common to both irrigated area and rainfed lowlands systems. Its best blast hit corresponds to *Bradyrhizobium tropiciagri* (Bradyrhizobiaceae) with a 99.5% sequence similarity. Another core phylotype in rainfed lowlands is assigned to the same species with 99.3% sequence similarity.

For ITS, we identified 5 core ASVs in the irrigated systems, compared to 11 core ASVs associated with the lowlands, 4 of them being common to both rice growing systems (Fig. 4).

Then, we compared the prokaryotic co-occurrence networks in each rice growing system respectively (Fig. 5). We identified 15 hub ASVs in the irrigated systems and 20 in rainfed lowlands. We found a higher edge number in irrigated compared to rainfed lowlands: 1720 positive and 269 negative resulting in 2029 total edges in irrigated areas, while only 1163 positive and 85 negative resulting in 1248 total edges were found in rainfed lowlands. Finally, the network computed from irrigated areas had higher connectivity compared to the one from rainfed lowlands (9.8 vs 7.9 node mean degrees, respectively). None of the identified hub taxa were also core in any of the two rice growing systems.

Table 2. Results of the statistical tests performed on 16S and ITS alpha diversity indices (Shannon diversity index and observed richness) for rhizosphere and root compartments.

| | Prokaryotes 16S | | | | | | Fungi ITS | | | | | | |
|-------------|-------------------------|-------|-------|-------------------|-------|--------------|-------------------------|--------|---------|-------------------|----|--------|--------------|
| | Shannon diversity index | | | Observed richness | | | Shannon diversity index | | | Observed richness | | | |
| | Df | Chisq | P | Df | Chisq | P | Df | Sum Sq | F value | P | Df | Chisq | P |
| Rhizosphere | Rice system | 1 | 5.776 | 0.016 | 1 | 6.979 | 0.008 | 0.000 | 0.004 | 0.949 | 1 | 6.826 | 0.009 |
| | Zone | 2 | 5.087 | 0.078 | 2 | 5.510 | 0.064 | 0.014 | 0.070 | 0.932 | 2 | 0.056 | 0.973 |
| Root | Rice system * Zone | | ns | | ns | | ns | | ns | | 2 | 6.088 | 0.048 |
| | Rice system | 1 | 0.554 | 0.456 | 1 | 0.003 | 0.958 | 0.077 | 0.158 | 0.461 | 1 | 0.334 | 0.563 |
| | Zone | 2 | 1.084 | 0.586 | 2 | 7.422 | 0.024 | 0.505 | 0.522 | 0.604 | 2 | 32.476 | 0.001 |
| | Rice system * Zone | | ns | | 7.769 | 0.021 | ns | | ns | | 2 | 13.440 | 0.001 |

The interaction between rice growing system and geographical zone was first included in the model, but then removed if non significant (indicated as 'ns' in the table). Mixed models (using the functions: `lmer` for Shannon index, and `glmer` with poisson family for observed richness) were used for 16S data, analyzed at the plant level, in order to include the field sampled as random effect, while no random effect were used for ITS data (functions: `lm` for Shannon index, and `glm` with poisson family for observed richness).





Figure 3. Comparison of root-associated microbiota α -diversity in contrasted rice-growing systems: irrigated (in blue) vs rainfed lowland (in red). Observed richness and Shannon indices are reported for each sample (i.e. one plant for 16S and one field for ITS), as violin diagram for each rice growing system. The left side of the figure presents the results obtained for 16S microbiome, while the right side shows the results obtained for ITS analysis. On top are shown the results of the rhizosphere compartment and on the bottom are the results obtained for the root associated compartment. The shape of the points shows the geographical zone (Banzon, Karfiguela, and Bama)

(A) Prokaryotes (16S)

Irrigated

- *Paraburkholderia kururiensis*
- *Ralstonia pickettii*



(B) Fungi (ITS)

Irrigated

- *Pulveroboletus sinensis*



Both

- *Epicoccum polychromum*
- *Nigrospora magnoliae*
- *Staurosphaeria rhamnocola*

Rainfed Lowland

- *Coniochaeta rosae*
- *Hypomyces semicircularis*
- *Paraphaeosphaeria michotii*
- *Pseudothielavia arxii*
- *Symptodiella acicola*
- *Tristartiperidium microsporium*
- *Vermiculariopsiella eucalypti*

Figure 4. Venn diagram representing the core sequence variants for each rice growing system: irrigated vs rainfed lowlands. (A) For prokaryotes (B) For fungi.

Only one ASVs was identified as a hub in both irrigated and rainfed lowland systems, assigned to *Anaeromyxobacter dehalogenans* (Anaeromyxobacteraceae). Hub taxa in irrigated areas (15 ASVs) were assigned to 15 different species from 12 families, while hub taxa in rainfed lowland (20 ASVs) only corresponded to 17 species from 11 families (Fig. 5).

Indicator taxa of the two rice growing systems

For 16S rRNA gene data, we found 128 indicator taxa in irrigated areas, including ASVs from 24 bacterial families, most of them assigned to *Acinetobacter*, *Ralstonia*, *Comamonas*, and *Clostridium* (Table 3). On the other hand, only 63 were identified in rainfed lowlands, most of them within the Bacillaceae family, including ASVs assigned to *Exiguobacterium* and *Priestia* (Table 3). The ASV assigned to *Paraburkholderia kururiensis* (Burkholderiaceae) revealed

as indicator in irrigated areas (Table 3) was also a core taxa in irrigated areas. Also, among the 21 indicator ASVs in irrigated areas assigned to *Ralstonia pickettii* (Burkholderiaceae), 16 were also core in irrigated areas. In addition, four indicator ASVs in irrigated areas were also hubs in this system: one assigned to *Pseudogulbenkiania subflava* (Chromobacteriaceae), one to *Clostridium intestinale* (Clostridiaceae), another to *Enterobacter cloacae* (Enterobacteriaceae), and finally one corresponding to *Pseudomonas flavescens* (Pseudomonadaceae). One ASVs assigned to *Priestia flexa* (Bacillaceae) and one assigned to *Methylosinus trichosporium* (Methylocystaceae) were hubs in rainfed lowlands.

For ITS data, we found 16 indicator taxa in irrigated areas, and 27 in rainfed lowlands (Table 3). Indicator taxa in irrigated areas were assigned to four classes: *Agaricomycetes*, *Microbotryomycetes*, and *Sordariomycetes*. Indicator taxa in rainfed lowlands were assigned to five classes: *Chytridiomycetes*, *Dothideomycetes*, *Sac-*



Figure 5. Network properties for each of the co-occurrence networks obtained from 16S rRNA data: **(A)** Samples from irrigated perimeters, and **(B)** samples from rainfed lowlands. Graphical representation is shown, as well as a table showing the assignation of hub ASV in each rice growing system.

charomycetes, Sordariomycetes, and Ustilaginomycetes. One ITS ASV identified as indicator taxa in irrigated, with best hit *Pulverobolus sinensis* (Agaricomycetes), was also core in this rice growing system, and two indicator taxa in rainfed lowlands were also core in this system: one assigned to *Paraphaeosphaeria michotii* (Dothideomycetes) and the other to *Coniochaeta rosae* (Sordariomycetes).

Putative pathogen or phytobeneficial taxa

First, responses for an 'oryz' query within assignment and blast, found matching records only in the 16S rRNA gene data set. A total of 200 ASVs included 'oryz' in their names, from 21 different genera, none of these species corresponded to pathogens from Table S6. Among them, putative beneficial taxa were found, particularly

the following: *Azospirillum oryzae*, *Novosphingobium oryzae*, *Paenibacillus oryzae*, and *Rhizobium oryzae*, *R. rhizoryzae*, and *R. straminoryzae*.

Next, we made a subset of the 16S rRNA gene data set for ASVs assigned to rhizobia. This corresponds to a total of 96 ASV among which one was indicator species for rainfed lowlands (Table 3), two are core in rainfed lowlands with one also core in irrigated (Fig. 4). Summed abundances of rhizobia of the rhizosphere are affected by the rice growing system (Table 4), with higher abundances in rainfed lowlands (Fig. 6). In roots, summed abundances of rhizobia were affected by the geographic zone, as well as the interaction between rice growing system and geographic zone (Table 4, Fig. 6).

The same kind of analysis was then performed for the *Glomeromycetes* class (total of 14 ASVs) in ITS data set. AMF summed abundance was affected by the rice growing system, the geograph-

Table 3. List of species assignment and number of sequence variants (ASVs) identified as indicator taxa in irrigated and rainfed lowland environment. The species in bold were also found as potential hub or core taxa.

| Kingdom | Family (Prokaryotes)/Class (Fungi) | Irrigated | Rainfed Lowlands | |
|-----------------------|---|---|--|--|
| Prokaryotes | Acidobacteriaceae | | <i>Occallatibacter savannae</i> (2) | |
| | Anaerolineaceae | <i>Ornatilinea apprima</i> (2) | <i>Ornatilinea apprima</i> (1) | |
| | Anaeromyxobacteraceae | <i>Anaeromyxobacter dehalogenans</i> (2) | | |
| | Azospirillaceae | <i>Azospirillum soli</i> (1) | | |
| | Bacillaceae | | <i>Bacillus zanthoxyli</i> (3) | <i>Bacillus zanthoxyli</i> (2) |
| | | | <i>Neobacillus cucumis</i> (2) | <i>Neobacillus cucumis</i> (1) |
| | | | <i>Priestia flexa</i> (1) | <i>Priestia flexa</i> (4) |
| | | | <i>Exiguobacterium acetylicum</i> (4) | <i>Exiguobacterium acetylicum</i> (25) |
| | | | | <i>Exiguobacterium indicum</i> (19) |
| | Bradyrhizobiaceae | <i>Bradyrhizobium oligotrophicum</i> (4) | | |
| | Bryobacteraceae | <i>Paludibaculum fermentans</i> (2) | | |
| | Burkholderiaceae | | <i>Burkholderia vietnamiensis</i> (1) | <i>Burkholderia vietnamiensis</i> (1) |
| | | | <i>Paraburkholderia kururiensis</i> (1) | |
| | | <i>Ralstonia pickettii</i> (21) | | |
| | Caulobacteraceae | | <i>Caulobacter hibisci</i> (1) | |
| | | | <i>Phenylobacterium panacis</i> (1) | |
| | Chitinophagaceae | | | <i>Flavisolibacter ginsenosidimitans</i> (1) |
| | Chromobacteriaceae | <i>Pseudogulbenkiania subflava</i> (1) | | |
| | Clostridiaceae | | <i>Clostridium beijerinckii</i> (8) | |
| | | | <i>Clostridium huakuii</i> (5) | |
| | | | <i>Clostridium intestinale</i> (1) | |
| | | | <i>Albitalea terrae</i> (1) | |
| | Comamonadaceae | <i>Comamonas testosteroni</i> (9) | | |
| | Enterobacteriaceae | | <i>Enterobacter cloacae</i> (1) | |
| | | | <i>Enterobacter mori</i> (1) | |
| | | | <i>Klebsiella variicola</i> (1) | |
| | Erythrobacteraceae | <i>Altericroceibacterium xinjiangense</i> (1) | | |
| | Gallionellaceae | | | <i>Sideroxydans lithotrophicus</i> (1) |
| | Geobacteraceae | <i>Geobacter luticola</i> (1) | | |
| | Kineosporiaceae | | <i>Angustibacter speluncae</i> (4) | |
| | | | <i>Kineococcus glutinatus</i> (1) | |
| | Methylobacteriaceae | <i>Methylobacterium radiotolerans</i> (1) | | |
| | Methylocystaceae | | <i>Methylocystis parvus</i> (2) | <i>Methylocystis parvus</i> (1) |
| | | <i>Methylosinus trichosporium</i> (1) | <i>Methylosinus trichosporium</i> (1) | |
| Moraxellaceae | | <i>Acinetobacter brisouii</i> (1) | | |
| | | <i>Acinetobacter modestus</i> (2) | | |
| | | <i>Acinetobacter pittii</i> (1) | | |
| | | <i>Acinetobacter seifertii</i> (1) | | |
| | | <i>Acinetobacter soli</i> (20) | <i>Acinetobacter soli</i> (1) | |
| | | <i>Pseudomonas citronellolis</i> (2) | <i>Pseudomonas paralactis</i> (2) | |
| Pseudomonadaceae | | <i>Pseudomonas flavescens</i> (1) | | |
| | | <i>Pseudomonas glareae</i> (1) | | |
| | | <i>Pseudomonas paralactis</i> (2) | | |
| | | <i>Pseudomonas weihenstephanensis</i> (2) | | |
| | | <i>Rhizobium etli</i> (1) | | |
| | | <i>Dyella marensis</i> (2) | | |
| | | <i>Shewanella xiamenensis</i> (5) | | |
| Sphingomonadaceae | <i>Sphingomonas limnosedimicola</i> (1) | | | |
| Sterolibacteriaceae | <i>Sulfurisoma sedimicola</i> (1) | | | |
| Thermosporotrichaceae | | | <i>Thermosporothrix hazakensis</i> (1) | |
| Fungi | Number of ASVs | 128 | 63 | |
| | Agaricomycetes | <i>Crepidotus albolanatus</i> (1) | | |
| | | <i>Marasmiellus celebanticus</i> (1) | | |
| | | <i>Pseudosperma notodryinum</i> (1) | | |
| | | <i>Pulveroboletus sinensis</i> (1) | | |
| Chytridiomycetes | | | <i>Clydaea vesicula</i> (1) | |
| | | | <i>Protrudomyces lateralis</i> (1) | |

Table 3. Continued

| Kingdom | Family (Prokaryotes)/Class (Fungi) | Irrigated | Rainfed Lowlands |
|---------|------------------------------------|--|---|
| | Dothideomycetes | <i>Epicoccum polychromum</i> (1) <i>Muyocopron laterale</i> (1) <i>Poaceascoma filiforme</i> (1) | <i>Camarosporidiella melnikii</i> (1) <i>Cladosporium chasmanthicola</i> (1) <i>Dendryphion europaeum</i> (1) <i>Epicoccum polychromum</i> (1) <i>Gordonomyces mucovaginatius</i> (1) <i>Helminthosporium erythrinicola</i> (1) <i>Paraphaeosphaeria michotii</i> (1) <i>Stemphylium botryosum</i> (1) |
| | Microbotryomycetes | <i>Rhodosporidiobolus colostri</i> (1) | <i>[Candida] boidinii</i> (1) |
| | Saccharomycetes | | <i>Nakaseomyces delphensis</i> (2) |
| | Sordariomycetes | <i>Diatrypella vulgaris</i> (1) <i>Neurospora lineolata</i> (1) <i>Ophiostoma pityokteinis</i> (2) <i>Vermiculariopsiella eucalypti</i> (3) | <i>Coniochaeta rosae</i> (1) <i>Cordana bisbyi</i> (1) <i>Cordana ellipsoidea</i> (1) <i>Diatrypella atlantica</i> (1) <i>Neurospora lineolata</i> (5) <i>Ophiostoma pityokteinis</i> (1) <i>Tristatiperidium microsporium</i> (1) <i>Vermiculariopsiella eucalypti</i> (1) <i>Xylotumulus gibbispurus</i> (1) <i>Violaceomyces palustris</i> (1) |
| | Ustilaginomycetes | | |
| | NA | <i>Asteromyces cruciatus</i> (1) | |
| | Number of ASVs | 16 | 27 |

Table 4. Results of the statistical tests performed on summed abundance of Rhizobia on the one hand, and Glomeromycota on the other hand, for both rhizosphere and root compartment.

| | | Rhizobia | | | Glomeromycota | | |
|-------------|--------------------|----------|--------|--------------|---------------|----------|--------------|
| | | Df | Chisq | P | Df | Chisq | P |
| Rhizosphere | Rice system | 1 | 20.043 | 0.000 | 1 | 26.159 | 0.000 |
| | Zone | 2 | 1.684 | 0.431 | 2 | 32.543 | 0.000 |
| | Rice system * Zone | | ns | | 2 | 282.909 | 0.000 |
| Root | Rice system | 1 | 1.090 | 0.297 | 1 | 106.760 | 0.000 |
| | Zone | 2 | 13.001 | 0.002 | 2 | 1999.660 | 0.000 |
| | Rice system * Zone | 2 | 6.069 | 0.048 | 2 | 20.710 | 0.000 |

The interaction between rice growing system and geographical zone was first included in the model, but then removed if non significant (indicated as 'ns' in the table). Mixed models (using the functions: glmer with poisson family) were used for 16S data, analyzed at the plant level, in order to include the field sampled as random effect, while no random effect were used for ITS data (functions: glm with poisson family).

ical zone and their interaction, both for rhizosphere and root compartments (Table 4). Higher abundances of *Glomeromycetes* are observed in rainfed lowlands, compared to irrigated areas (Fig. 6).


We then screened the list of all assigned ASVs for a set of pathogen species defined *a priori* (see the list in the Table S1). For prokaryotes (16S rRNA gene data), a number of ASVs corresponded to the genera of pathogens, but only *Burkholderia glumae* (two ASVs), *Acidovorax avenae* (four ASVs), and *Dickeya chrysanthemi* (six ASVs) were identified at the species level. These 12 ASVs identified at the species level were however only found in one sample. The *Xanthomonas* genus was found, but with no assignment to *X. oryzae* (instead, assigned to *X. theicola*, which is phylogenetically closed to the rice associated *X. sontii*, Bansal et al. 2020), although two sampled plants had specific symptoms of bacterial leaf streak (caused by *X. oryzae* pathovar *oryzicola*). A similar situation (assignment at the genus level) was observed for the genera *Pseudomonas*, *Pantoea*, and *Sphingomonas*. The same analysis of putative pathogens for ITS revealed the presence of the following ten genera: *Alternaria*, *Bipolaris*, *Ceratobasidium*, *Curvularia*, *Fusarium*, *Helminthosporium*, *Microdochium*, *Rhizoctonia*, *Sarocladium*, *Bipo-*

laris spp. causes brown spot symptoms, which were frequently observed (33 out of 57 sampled plants, i.e. 57.9%). On the other hand, one sampled plants presented rice leaf blast symptoms, but no *Pyricularia oryzae* was found in this ITS dataset from rhizosphere and root samples.

DISCUSSION

This study aimed at describing the rice root-associated microbiome by comparing contrasted rice growing systems in farmer's fields in Burkina Faso. We found that the rice growing system was a structuring factor for rice root-associated microbiomes, with an effect on alpha diversity in the rhizosphere, higher in irrigated areas for bacteria, while rainfed lowlands harbor higher fungal richness. In addition, we identified a number of phylotypes with potential key roles (hub, core, indicators) in the two contrasted systems, as well as putative phytobeneficial and pathogen species. Although the results on fungi (ITS region) must be taken with caution due to a smaller sample size and the poor representation of obtained sequences in available taxonomic databases, this study

(A) Rhizobia summed abundance



(B) Glomeromycota summed abundance




Figure 6. Spatial repartition of summed abundance of potentially phyto-beneficial families in the two rice root microbiome compartments (rhizosphere and roots), and in each rice growing system (irrigated perimeters vs rainfed lowlands). (A) Summed abundance for 14 ITS ASVs assigned to the class *Glomeromycetes*; (B) Summed abundance for 96 ITS ASVs assigned to rhizobia. The shape of the point corresponds to each geographic zone.

shed light on some drivers of assemblage of rice root associated microbial communities in a sparsely documented African system.

Structure and diversity of the root-associated microbiomes is affected by the rice growing system

Although Edwards et al. (2018) showed that the root-associated microbiome of distant field sites converge in similarity during the growing season, our study performed at the maturity stage of rice still evidenced some drivers of rice root-associated microbial communities structure. First, as for most rice microbiome studies, we found an effect of the compartment/micro-habitat (Edwards et al. 2015; Santos-Medellin et al. 2017; Guo et al. 2021; Kawasaki et al. 2021), and the geographical zone (Edwards et al. 2015; Kanasugi et al. 2020) on the beta-diversity of rice root-associated microbiome. In addition, our study shows that the contrasted rice-growing systems, namely irrigated perimeters vs rainfed lowlands, harbor contrasted rice root-associated microbial communities, both for prokaryotic and fungal communities, and for rhizosphere and root compartment. We notice that the soil physicochemical properties weakly differ between irrigated areas and rainfed lowland, the soil composition was instead mostly affected by the geographical zones. Consequently, we evidence a structuring effect of the rice-growing system that was only slightly related to contrasted soil physicochemical properties. Our results are in line with a previous study comparing microbiomes from two contrasted water management conditions (upland vs. lowland rice) in controlled settings (a field experiment in northern Italy), which showed differentiation in microbial communities, particularly for root microbiome, and to a lesser extent in soil samples (Chialva et al. 2020).

In our study, some of the soil physicochemical parameters affected rice root-associated microbial communities. In particular,

CEC and SBE, that reflect soil exchange capacity and bioaccessibility, were the most important soil parameters for the structure of both rhizosphere and root prokaryotic communities. These parameters are not commonly measured in other studies of the root-associated microbiome, and our results argue for including them in soil chemical characterization, to investigate whether their impact in microbiome structure is general or not. In addition, phosphorus content significantly structured the prokaryotic communities, both in rhizosphere and roots. Such an effect of phosphorus is known for the rice root associated microbiome (Long and Yao 2020). On the other hand, the soil chemical parameter evidenced in this study to structure fungal rhizosphere communities was the total nitrogen (N). This is in accordance with a study by Chen et al. (2019) showing that nitrogen input drives changes in the microbial root-associated community structure in wheat. Moreover, Wang and Huang (2021) showed the effect of optimized N application on fungal community structure from paddy soils. Kanasugi et al. (2020) also evidenced an effect of soil nitrate on rice root fungal communities in Ghana.

We found a higher taxonomic diversity in irrigated areas, compared to rainfed lowlands, for prokaryotic communities of the rhizosphere, while the opposite (higher diversity in rainfed lowlands compared to irrigated areas) was found for fungal communities. Chialva et al. (2020)'s found that the 16S rRNA gene diversity was similar in lowland and upland rice, but they found a significantly higher ITS diversity in lowland rice compared to upland. It is important to note that our sampling was performed directly in farmer's fields, with no control of management practices, while most results, including those of Chialva et al. (2020), were obtained in field trials. Indeed, in our study, various factors, such as rice cultivars, fertilization regime and rotation, exhibit large variability. However, levels of rice genetic diversity were shown to be compa-

rable within each site, both in irrigated areas and rainfed lowlands (Barro et al. 2021b), therefore the effect of rice growing system on alpha-diversity of microbiomes could not be attributed to result from difference in terms of genetic diversity of the host plant.

Considering the irrigated areas as systems with more intensive agricultural practices, compared to rainfed lowland, we expected to observe higher diversity of microorganisms in rainfed systems, as it was the case for fungal communities, but not the opposite one obtained for bacterial communities. Indeed, agricultural intensification was shown to reduce microbial network complexity and the abundance of keystone taxa in roots (Banerjee et al. 2019). In addition, the fertilization regime is known to have strong impact on root-associated microbiota (Ding et al. 2019; Xiong et al. 2021). Various studies showed that organic fertilization enhances microbial diversity (Liu et al. 2020). For example, recommended fertilization preserved belowground microbial populations, compared to the fertilization mostly used (conventional fertilization) that depressed bacterial diversity, in experiments performed in China (Ullah et al. 2020). We considered the irrigated areas as more intensified systems, compared to rainfed lowland, particularly because only irrigated areas allow growing rice twice a year, and because only rainfed lowland sites presented fields with no mineral fertilization at all (Barro et al. 2021a). We noticed, however, that organic fertilization remained rare, and its frequency was not drastically affected by the rice growing system. Finally, transplantation was always performed in irrigated areas, while direct sowing was the most common practice in rainfed lowlands.

On the other hand, paddy soils studied in western Burkina Faso (all over the six sites) are particularly poor if compared for example to a study of more than 8000 soils in Hunan Province (Duan et al. 2020). The studies previously cited evidencing fertilization effects, were performed in soils with higher carbon and nitrogen contents (see for example Ullah et al. 2020). The effect of fertilization on microbial diversity may actually depend on various aspects, including the soil type. Notably, a positive relationship was found between rice fertilization and soil bacterial richness and diversity in a 19-years inorganic fertilization assay in a reddish paddy soil in southern China (Huang et al. 2019); while Wang and Huang (2021) showed an effect of the fertilization on paddy soils microbial community composition but no effect on the diversity. The poverty of soils in the sampled area may be involved in the complexity of fertilization-diversity relationships observed in this study.

Our results showed that the prokaryotic communities in the rice rhizosphere and roots from the three irrigated sites do not differ significantly from each other. On the other hand, the same analysis revealed significant differentiation between the three rainfed lowland study sites (in all three cases for rhizosphere and two out of three comparisons in roots). Also, we found very few core phylotypes in rainfed lowland, with only two core ASVs for 16S rRNA gene, which reinforces the above-mentioned observation. These results are likely driven by a higher heterogeneity between rainfed sites, in terms of water control, agricultural practices or rice genotypes. Indeed, in irrigated rice, the farmer has the potential to control irrigation water during the whole growing season. On the other hand, irrigation in rainfed lowland is dependent on precipitation that differs between the three geographical zones sampled within the rice growing season. In addition, we showed a high heterogeneity of agricultural practices in rainfed lowlands: for example, legume rotation was common only in the rainfed lowland of Bama zone, and organic fertilization was especially frequent in the rainfed lowland of Karfiguela zone (Barro et al. 2021a). Finally, in terms of rice genetics, a high rice ge-

netic differentiation was found between the rainfed lowland site of Karfiguela zone and the five other sites: a distinct genetic group *O. sativa* Aus, and other distinct landraces were found in this peculiar site, compared to the five others where only *O. sativa indica* was grown (Barro et al. 2021b). Lower sampling size for ITS may explain the absence of significant differences obtained between pairs of sites. Alternatively, the pattern may be different for fungal diversity, as suggested by the higher number of core taxa in rainfed lowlands compared to irrigated areas.

Key taxa reveal different ecological functioning between irrigated systems and lowlands

The core microbiome is hypothesized to represent the most ecologically and functionally important microbial associates of that environment under the conditions sampled (Neu et al. 2021), so that describing the cores and determining whether it is shared between irrigated systems and rainfed lowlands is of major interest. While four fungal taxa were found to be cores in both systems, only one bacterial core taxa was shared between the two rice growing systems: assigned to *Bradyrhizobium tropiciagri*, a nitrogen-fixing symbiont isolated from tropical forage legumes (Delamuta et al. 2015). Some legumes (crops in rotation, or wild plants such as *Aeschynomene* or *Stylosanthes*) may share the lowland soils with rice, and bacteria could interact with different plant host. Indeed, several strains of *Bradyrhizobium spp.* isolated from rice roots were described to be capable of fixing N_2 (Chaintreuil et al. 2000, Ding et al. 2019). In addition, the core taxa in irrigated areas likely includes *Paraburkholderia kururiensis*, a bacterium with potential phyto-beneficial properties (bioremediation, biofertilization, and biocontrol of pathogens; Dias et al. 2019). Various ASVs identified as core in irrigated areas were assigned to *Ralstonia pickettii*, an ubiquitous Betaproteobacteria found in water and soil, capable to thrive in low nutrient (oligotrophic) conditions (Ryan et al. 2007), considered as potential biocontrol agent against its congeneric pathogen *R. solanacearum* (Wei et al. 2013), and also, noteworthy, described as human emerging pathogen causing nosocomial infections (Ryan et al. 2006).

Hub taxa are potential keystones, i.e. taxa that drive community composition and function irrespective of their abundance (Banerjee et al. 2018). Only one ASVs was identified as hub of the prokaryote co-occurrence networks both in irrigated areas and rainfed lowland site. It is assigned to *Anaeromyxobacter dehalogenans* (Anaeromyxobacteraceae), a versatile soil bacterium capable of denitrification in the presence of iron (Onley et al. 2018). On the other hand, although various ASVs indicator in either irrigated areas or rainfed lowlands were assigned to the same species, we also found many bacterial taxa identified as hubs that differ between irrigated areas and rainfed lowlands (even at the family level), reflecting a highly contrasted structuring and functioning of bacterial communities in the two rice growing systems.

To identify specific ASVs that characterized microbiome variations between rice growing systems, we further analyzed our data for the presence of candidate indicator taxa. Most indicator taxa were found in irrigated systems for prokaryotes (128 vs. only 63 in rainfed lowlands) while the opposite was found for fungi (27 in rainfed lowlands vs only 16 in irrigated areas). For prokaryotes, five taxa identified as indicator species in irrigated areas were also core or hub: the previously mentioned *Paraburkholderia kururiensis* and *Ralstonia pickettii* as well as *Pseudogulbenkiania subflava*, *Clostridium intestinale*, *Acinetobacter soli*, and *Pseudomonas flavescens*. In rainfed lowlands, it was the case for *Priestia flexa* and *Methylosinus trichosporium*. In particular, we notice that *Acinetobac-*

ter soli was identified as a potent phosphorus solubilizer in rice and consequently promising for plant growth promotion (Rasul et al. 2019). For fungi, we identified as both indicator and core taxa: *Pulveroboletus sinensis* in irrigated areas, as well as both *Coniochaeta rosae* and *Paraphaeosphaeria michotii* in rainfed lowlands. Two ASVs assigned to Chytridiomycetes were identified as indicator species in rainfed lowland systems. These aquatic fungi (Barr 2001), found in the rhizosphere compartment in our study, are known as particularly abundant in microbial communities associated with rice roots, compared to other crops (Ding et al. 2019) and were preferentially associated to lowland conditions compared to upland (Chialva et al. 2020).

We identified a few potentially beneficial taxa that could be investigated further. In particular, rhizobia abundance in the rhizosphere was higher in rainfed lowlands compared to irrigated areas. This was expected considering that rainfed lowlands correspond to lower fertilizer input, and, in some cases legume rotations between rice cycles (frequent at least in the rainfed lowland of Bama zone, see above). Rhizobia populations are released from nodules when legume plant dies, and rhizobia are rice-colonizing bacteria (Chi et al. 2005). We also showed that AMF abundance is affected by the rice growing system (in both rhizosphere and roots). This is congruent with other studies reporting AMF colonization depending on farming regimes. For example, the rice roots cultivated in the conventional agrosystem (N and P fertilization and pesticides) or under permanent flooding showed no AMF colonization, while the rice plants grown with organic conditions showed typical mycorrhization patterns (Lumini et al. 2011).

Various pathogen species were suspected from the sequence variants identified in this study. In particular, the 16S rRNA gene dataset contained ASVs assigned to *Burkholderia glumae*, *Acidovorax avenae* and *Dickeya chrysanthemi*. All three remained rare, with each of these sequences only found in one sample. The presence of *B. glumae* was described in Burkina Faso (but with no molecular data; Ouedraogo et al. 2004), and targeted detection performed in two sites failed to detect *B. glumae* and *A. avenae* (Bangratz et al. 2020). For fungal pathogens, we found ASVs assigned to ten genera comprising rice pathogens, including *Bipolaris*, *Curvularia*, *Fusarium*, and *Rhizoctonia*. Some of these putative fungal pathogens are frequent, particularly one, whose best blast hit is *Curvularia chonburiensis*, identified as core taxa in both irrigated perimeters and rainfed lowlands. Various *Curvularia* species are known to be pathogenic in rice (Gao et al. 2012, Majeed et al. 2015), and their widespread repartition evidenced here argues for more plant pathology work on the interactions between *Curvularia* and rice.

The literature shows that higher microbiome diversity may be associated with a lower infection rate (see for example Rutten et al. 2021). For rice in western Burkina Faso, when compared to irrigated perimeters, we found that rainfed lowlands harbor higher rhizosphere fungal richness and lower prevalence of major rice diseases, particularly bacterial leaf streak and the fungal rice blast disease, based on the observation of foliar symptoms (Barro et al. 2021a). On the other hand, for prokaryote communities, the pattern is actually opposite, with a higher diversity in irrigated areas compared to rainfed lowlands. The potential relationship between the diversity of root-associated microbiome and diseases may actually be complex, requiring other studies, including under controlled conditions. More generally, scientific interest in the relationship between root-associated microbiota and plant diseases is growing (Vannier et al. 2019, Trivedi et al. 2020). These aspects constitute a promising avenue of research in this system as well, especially considering reciprocal effects evidenced in rice from experimental studies (Spence et al. 2014, Tian et al. 2021).

Perspectives

We are only at the beginning of understanding the complexity of rice root microbial communities, especially for rice cultivation in Africa. The originality, but also a limitation of our study, lies in the fact that the samples were collected in farmer fields, and it globally compares the two contrasting rice production systems that differ in various management practices, so that it could not tease apart the specific effect of each individual factor (water management, cultivar, fertilization, etc). More investigations are now required to decipher each structuring factor at a smaller scale: in particular between fields within each site, where the rice cultivar and specific agricultural practices are likely to play a significant role (Delitte et al. 2021).

Describing rice microbiota through metabarcoding is a first mandatory step that needs to be combined with culturomics for a greater accuracy and a deeper description, in particular for such systems where some taxa are poorly described in taxonomic databases. Experimental work in an integrative approach is also required to move on towards microbiota management methodologies. Such microbiota-based strategies could contribute to improving rice health and productivity (Sessitsch and Mitter 2015), while preserving human health. They are consequently an important component of the toolbox of science-based strategies to achieve zero-hunger in Africa.

SUPPLEMENTARY DATA

Supplementary data are available at [FEMSEC](#) online.

ACKNOWLEDGMENTS

This work was performed thanks to the facilities of the 'International joint Laboratory LMI PathoBios: Observatory of plant pathogens in West Africa: biodiversity and biosafety' (www.pathobios.com; twitter.com/PathoBios), and conducted within E-SPACE project (www6.inrae.fr/e-space/), led by Claire Neema. We are very grateful to Abdoul Kader Guigma, Yacouba Kone, Moumouni Traoré, and Arzouma Diasso for their contributions to the fieldwork in Burkina Faso. We thank the rice farmers from Badala, Bama, Senzon, Banzon, Tengrela, and Karfiguela for their kind collaboration. We thank Corinne Vacher and Alain Sarniguet for constructive discussions at earlier stage of the project. The manuscript benefited from helpful comments by Philippe Roumagnac, who also contributed to the initial design of the study. Finally, we dedicate this manuscript to the late Christian Vernière for his kind feedback at different stages of this project.

FUNDING

This work was publicly funded through ANR (the French National Research Agency) under 'Investissements d'avenir' programme with the reference ANR-10-LABX-001-01 Labex Agro (E-Space and Ri-PaBIOME projects), coordinated by Agropolis Fondation under the frame of I-SITE MUSE (ANR-16-IDEX-006) and by the CGIAR Research Program on Rice Agri-food Systems (RICE).

Conflict of interest statement: None declared.

REFERENCES

Abarenkov K, Zirk A, Piirmann T et al. UNITE general FASTA release for Fungi 2. 2021; Version 10.05.2021. UNITE Community.

- AFNOR. Détermination du pH (Association Française de Normalisation). 1981; NF ISO 10390. 339-348. AFNOR qualité des sols. Paris.
- Alonso P, Blondin L, Gladieux P et al. Heterogeneity of the rice microbial community of the chinese centuries-old honghe hani rice terraces system. *Environ Microbiol* 2020;**22**:3429–45.
- Banerjee S, Schlaeppi K, van der Heijden MGA. Keystone taxa as drivers of microbiome structure and functioning. *Nat Rev Microbiol* 2018;**16**:567–76.
- Banerjee S, Walder F, Büchi L et al. Agricultural intensification reduces microbial network complexity and the abundance of keystone taxa in roots. *ISME J* 2019;**13**:1722–36.
- Bangratz M, Wonni I, Kini K et al. Design of a new multiplex PCR assay for rice pathogenic bacteria detection and its application to infer disease incidence and detect co-infection in rice fields in burkina faso. *PLoS One* 2020;**15**:e0232115.
- Bansal K, Kumar S, Patil Prabhu B et al. Phylogenomic insights into diversity and evolution of nonpathogenic xanthomonas strains associated with citrus. *mSphere* 2020;**5**:e00087–00020.
- Barr DJS. Chytridiomycota. Systematics and Evolution: Part A. In: McLaughlin DJ, McLaughlin EG, Lemke PA, (eds), Berlin, Heidelberg: Springer Berlin Heidelberg, 2001, 93–112.
- Barro M, Kassankogno AI, Wonni I et al. Spatiotemporal survey of multiple rice diseases in irrigated areas compared to rainfed lowlands in the western Burkina faso. *Plant Dis* 2021a;**105**:3889–99.
- Barro M, Konate KA, Wonni I et al., Assessment of genetic diversity of rice in registered cultivars and farmers' fields in Burkina faso. *Crops* 2021b;**1**:129–40.
- Bates D, Mächler M, Bolker B et al. Fitting linear mixed-effects models using lme4. *J Statist Software* 2015;**67**:1–48.
- Berry D, Widder S. Deciphering microbial interactions and detecting keystone species with co-occurrence networks. *Front Microbiol* 2014;**5**:219.
- Bouyoucos GJ. Hydrometer method improved for making particle size analysis of soils. *Agron J* 1962;**54**:464–5.
- Brunel C, Pouteau R, Dawson W et al. Towards unraveling macroecological patterns in rhizosphere microbiomes. *Trends Plant Sci* 2020;**25**:1017–29.
- Callahan BJ, McMurdie PJ, Rosen MJ et al. DADA2: high-resolution sample inference from illumina amplicon data. *Nat Methods* 2016;**13**:581–3.
- Cartwright RD, Groth DE, Wamishe YA et al. *Compendium of Rice Diseases and Pests, Second Edition*. 2018; The American Phytopathological Society.
- Chaintreuil C, Giraud E, Prin Y et al. Photosynthetic bradyrhizobia are natural endophytes of the african wild rice *Oryza breviligulata*. *Appl Environ Microbiol* 2000;**66**:5437–47.
- Chen S, Waghmode TR, Sun R et al. Root-associated microbiomes of wheat under the combined effect of plant development and nitrogen fertilization. *Microbiome* 2019;**7**:136.
- Chi F, Shen SH, Cheng HP et al. Ascending migration of endophytic rhizobia, from roots to leaves, inside rice plants and assessment of benefits to rice growth physiology. *Appl Environ Microbiol* 2005;**71**:7271–8.
- Chialva M, Ghignone S, Cozzi P et al. Water management and phenology influence the root-associated rice field microbiota. *FEMS Microbiol Ecol* 2020;**96**:faa146.
- Chialva M, Lanfranco L, Bonfante P. The plant microbiota: composition, functions, and engineering. *Curr Opin Biotechnol* 2021;**73**:135–142.
- CountrySTAT. Annuaire statistique agricole 1993-2021: Superficie des céréales. (Faso CB, ed.) 2020.
- Csardi G, Nepusz T. The igraph software package for complex network research. 2006;**1695**
- Davis NM, Proctor D, Holmes SP et al. Simple statistical identification and removal of contaminant sequences in marker-gene and metagenomics data. *Microbiome* 2018;**6**:226. doi:doi.org/10.1186/s40168-018-0605-2
- De Cáceres M, Legendre P, Moretti M. Improving indicator species analysis by combining groups of sites. *Oikos* 2010;**119**:1674–84.
- Delamuta JRM, Ribeiro RA, Ormeño-Orrillo E et al. *Bradyrhizobium tropiciagri* sp. nov. and *bradyrhizobium embrapense* sp. nov., nitrogen-fixing symbionts of tropical forage legumes. *Int J Syst Evol Microbiol* 2015;**65**:4424–33.
- Delitte M, Caulier S, Bragard C et al. Plant microbiota beyond farming practices: a review. *Front Sustain Food Syst* 2021;**5**:624203.
- Dias GM, de Sousa Pires A, Grilo VS et al. Comparative genomics of *Paraburkholderia kururiensis* and its potential in bioremediation, biofertilization, and biocontrol of plant pathogens. *Microbiology-Open* 2019;**8**:e00801.
- Ding L-J, Cui H-L, Nie S-A et al. Microbiomes inhabiting rice roots and rhizosphere. *FEMS Microbiol Ecol* 2019;**95**:fz040.
- Duan L, Li Z, Xie H et al. Large-scale spatial variability of eight soil chemical properties within paddy fields. *Catena* 2020;**188**:104350.
- Edwards J, Johnson C, Santos-Medellin C et al. Structure, variation, and assembly of the root-associated microbiomes of rice. *Proc Natl Acad Sci* 2015;**112**:E911–20.
- Edwards JA, Santos-Medellin CM, Liechty ZS et al. Compositional shifts in root-associated bacterial and archaeal microbiota track the plant life cycle in field-grown rice. *PLoS Biol* 2018;**16**:e2003862.
- Epskamp S, Cramer AOJ, Waldorp LJ et al. qgraph: network visualizations of relationships in psychometric data. *J Stat Soft* 2012;**48**:1–18.
- FAO/IIASA/ISRIC/ISSCAS/JRC. *Harmonized World Soil Database (version 1.2)*. FAO and IIASA: Rome, Italy and Laxenburg, Austria. 2012.
- Fick SE, Hijmans RJ. WorldClim 2: new 1 km spatial resolution climate surfaces for global land areas. *Int J Climatol* 2017;**37**:4302–15.
- Fox J, Weisberg S. *An R Companion to Applied Regression*. Sage, Thousand Oaks CA. 2019.
- Gao BD, Huang W, Xia H. A New Rice Disease, Black Sheath Spot, Caused by *Curvularia fallax* in China. *Plant Dis* 2012;**96**:1224.
- Guo J, Ling N, Li Y et al. Seed-borne, endospheric and rhizospheric core microbiota as predictors of plant functional traits across rice cultivars are dominated by deterministic processes. *New Phytol* 2021;**230**:2047–60.
- Hillebrand WF, Lundel GEF, Bright HA et al. *Applied inorganic analysis*. 2nd edition. New York, USA. 1953.
- Huang Q, Wang J, Wang C et al. The 19-years inorganic fertilization increased bacterial diversity and altered bacterial community composition and potential functions in a paddy soil. *Applied Soil Ecology* 2019;**144**:60–7.
- Hughes AC, Orr MC, Ma K et al. Sampling biases shape our view of the natural world. *Ecography* 2021;**44**:1259–69.
- Kanasugi M, Sarkodee-Addo E, Ansong Omari R et al. Exploring rice root microbiome; the variation, specialization and interaction of bacteria and fungi in six tropic savanna regions in ghana. *Sustainability* 2020;**12**:5835.
- Kawasaki A, Dennis PG, Forstner C et al. The microbiomes on the roots of wheat (*Triticum aestivum* L.) and rice (*Oryza sativa* L.) exhibit significant differences in structure between root types and along root axes. *Funct Plant Biol*. 2021;**48**:871–88.
- Kim H, Lee Y-H. The rice microbiome: a model platform for crop holobio. *Phytobiomes Journal* 2020;**4**:5–18.

- Kurtz ZD, Müller CL, Miraldi ER et al. Sparse and compositionally robust inference of microbial ecological networks. *PLoS Comput Biol* 2015;**11**:e1004226.
- Linares OF. African rice (*Oryza glaberrima*): history and future potential. *Proc Natl Acad Sci* 2002;**99**:16360.
- Liu Z, Guo Q, Feng Z et al. Long-term organic fertilization improves the productivity of kiwifruit (*Actinidia chinensis* planch.) through increasing rhizosphere microbial diversity and network complexity. *Appl Soil Ecol* 2020;**147**:103426.
- Long X-E, Yao H. Phosphorus input alters the assembly of rice (*Oryza sativa* L.) root-associated communities. *Microb Ecol* 2020;**79**:357–66.
- Lu T, Ke M, Lavoie M et al. Rhizosphere microorganisms can influence the timing of plant flowering. *Microbiome* 2018;**6**:231.
- Lumini E, Vallino M, Alguacil MM et al. Different farming and water regimes in Italian rice fields affect arbuscular mycorrhizal fungal soil communities. *Ecol Appl* 2011;**21**:1696–707.
- MAHRH. Burkina Faso: National Rice Development Strategy (NRDS). 2011.
- Majeed RA, Shahid AA, Ashfaq M et al. First report of *Curvularia lunata* causing brown leaf spots of rice in Punjab, Pakistan. *Plant Dis* 2015;**100**:219.
- Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMB Net J* 2011;**17**:10–2.
- McMurdie PJ, Holmes S. phyloseq: An R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One* 2013;**8**:e61217.
- Metson AJ. Methods of chemical analysis for soil survey samples. *N Z Soil Bur Bull* 1956;**12**.
- Neu AT, Allen EE, Roy K. Defining and quantifying the core microbiome: challenges and prospects. *Proc Natl Acad Sci* 2021;**118**:e2104429118.
- Nonvide GMA, Sarpong DB, Kwadzo GTM et al. Farmers' perceptions of irrigation and constraints on rice production in Benin: a stakeholder-consultation approach. *Int J Water Resour Dev* 2018;**34**:1001–21.
- Novozansky IV, Houba JG, Vaneck R et al. A novel digestion technique for multi-element analysis. *Commun Soil Sci Plant Anal* **14**, 1983:239–49.
- Oksanen J, Kindt R, Legendre P et al. The vegan package. Community Ecol Package 2007. <https://cran.r-project.org/web/packages/vegan/>.
- Onley JR, Ahsan S, Sanford RA et al. Denitrification by *Anaeromyxobacter dehalogenans*, a common soil bacterium lacking the nitrite reductase genes nirS and nirK. *Appl Environ Microbiol* 2018;**84**:e01985–17.
- Op De Beeck M, Lievens B, Busschaert P et al. Comparison and validation of some ITS primer pairs useful for fungal metabarcoding studies. *PLoS One* 2014;**9**:e97629.
- Ouedraogo S, Somda I, Borro F et al. Detection and characterization of plant pathogenic bacteria transmitted by seed rice in Burkina Faso. *Agronomie Africaine* 2004;**6**:9–17.
- Pandey S, Byerlee D, Dawe D. Rice in the global economy: strategic research and policy issues for food security. Los Banos (Philippines). 2010.
- Peterson RA. Finding optimal normalizing transformations via best-Normalize. *R J* 2021;**13**:310–29. doi: 10.32614/RJ-2021-041.
- Quast C, Pruesse E, Yilmaz P et al. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res* 2013;**41**:D590–6.
- R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing. 2018; Vienna, Austria. URL <https://www.R-project.org/>.
- Rao A, Wani S, Ramesha M et al. Chapter 8 rice production systems. *Rice Production Worldwide*, In: Chauhan B, Jabran K, Mahajan G, (eds). Springer International Publishing AG 2017.
- Rasul M, Yasmin S, Suleman M et al. Glucose dehydrogenase gene containing phosphobacteria for biofortification of phosphorus with growth promotion of rice. *Microbiol Res* 2019;**223–225**: 1–12.
- Rutten G, Hönig L, Schwaß R et al. More diverse tree communities promote foliar fungal pathogen diversity, but decrease infestation rates per tree species, in a subtropical biodiversity experiment. *J Ecol* 2021;**109**:2068–80.
- Ryan MP, Pembroke JT, Adley CC. *Ralstonia pickettii* in environmental biotechnology: potential and applications. *J Appl Microbiol* 2007;**103**:754–64.
- Ryan MP, Pembroke JT, Adley CC. *Ralstonia pickettii*: a persistent Gram-negative nosocomial infectious organism. *J Hosp Infect* 2006;**62**:278–84.
- Santos-Medellín C, Edwards J, Liechty Z et al. Drought stress results in a compartment-specific restructuring of the rice root-associated microbiomes. *Mbio* 2017;**8**:e00764–00717.
- Senthil Kumar CM, Jacob TK, Devasahayam S et al. Multifarious plant growth promotion by an entomopathogenic fungus *Lecanicillium psalliotae*. *Microbiol Res* 2018;**207**:153–60.
- Serpantié G, Dorée A, Fusillier J-L et al. Nouveaux risques dans les bas-fonds des terroirs soudanais. Une étude de cas au Burkina Faso. *Cah Agri* 2019;**28**:19.
- Sessitsch A, Mitter B. 21st century agriculture: integration of plant microbiomes for improved crop production and food security. *Microb Biotechnol* 2015;**8**:32–3.
- Singh BK, Trivedi P, Egedi E et al. Crop microbiome and sustainable agriculture. *Nat Rev Microbiol* 2020;**18**:601–2.
- Soullier G, Demont M, Arouna A et al. The state of rice value chain upgrading in West Africa. *Global Food Security-Agriculture Policy Econom Environ* 2020;**25**:100365.
- Spence C, Alff E, Johnson C et al. Natural rice rhizospheric microbes suppress rice blast infections. *BMC Plant Biol* 2014;**14**:130.
- Tamburini G, Bommarco R, Wanger TC et al. Agricultural diversification promotes multiple ecosystem services without compromising yield. *Sci Adv* 2020;**6**:eaba1715.
- Thijs S, Op De Beeck M, Beckers B et al. Comparative evaluation of four bacteria-specific primer pairs for 16S rRNA gene surveys. *Front Microbiol* 2017;**8**:494. doi:10.3389/fmicb.2017.00494
- Tian D, Chen Z, Lin Y et al. The interaction between rice genotype and *Magnaporthe oryzae* regulates the assembly of rice root-associated microbiota. *Rice* 2021;**14**:40.
- Trivedi P, Leach JE, Tringe SG et al. Plant-microbiome interactions: from community assembly to plant health. *Nat Rev Microbiol* 2020;**18**:607–21.
- Ullah S, He P, Ai C et al. How do soil bacterial diversity and community composition respond under recommended and conventional nitrogen fertilization regimes? *Microorganisms* 2020;**8**:1193.
- Vandenkoornhuysen P, Quaiser A, Duhamel M et al. The importance of the microbiome of the plant holobiont. *New Phytol* 2015;**206**:1196–206.
- Vannier N, Agler M, Hacquard S. Microbiota-mediated disease resistance in plants. *PLoS Pathog* 2019;**15**:e1007740.
- Walinga J, Van Vark W, Houba VJ et al. Plant analysis procedures. Wageningen, University A, *Soil and Plant Analysis*, (7:Wageningen: Landbouwniversiteit, 1989;197–200. eds), 1986–1995.
- Walkley A, Black IA. An examination method of the detjareff and a proposed modification of the chromic acid titration method. *Soil Sci* 1934;**37**:29–38.

- Wang L, Huang D. Soil microbial community composition in a paddy field with different fertilization managements. *Can J Microbiol* 2021;**67**:864–74.
- Wei Z, Huang J, Tan S et al. The congeneric strain *Ralstonia pickettii* QL-A6 of *Ralstonia solanacearum* as an effective biocontrol agent for bacterial wilt of tomato. *Biol Control* 2013;**65**: 278–85.
- Wei Z, Yang T, Friman V-P et al. Trophic network architecture of root-associated bacterial communities determines pathogen invasion and plant health. *Nat Commun* 2015;**6**:8413.
- Wickham H. ggplot2: Elegant Graphics for Data Analysis. 2016; ISBN 978-3-319-24277-4. <https://ggplot2.tidyverse.org>
- Wright ES. Using DECIPHER v2.0 to analyze big biological sequence data in R. *The R Journal* 2016;**8**:352–9.
- Xiong QQ, Hu JL, Wei HY et al. Relationship between plant roots, rhizosphere microorganisms, and nitrogen and its special focus on rice. *Agriculture* 2021;**11**:234.