

available at www.sciencedirect.comjournal homepage: www.elsevier.com/locate/mycres

Forest stand age and the occurrence of chanterelle (*Cantharellus*) species in Oregon's central Cascade Mountains

Susie M. DUNHAM^{a,*}, Thomas E. O'DELL^b, Randy MOLINA^c

^aOregon State University, Department of Botany and Plant Pathology, 2082 Cordley Hall, Corvallis, OR 97331, USA

^bUSDI National Park Service, Northern Colorado Plateau Inventory and Monitoring Network, 2282 SW Resource Blvd., Moab, UT 84532, USA

^cUSDA Forest Service, Pacific Northwest Research Station, 620 SW Main, Suite 400, Portland, OR, USA

ARTICLE INFO

Article history:

Received 14 March 2006

Received in revised form

31 July 2006

Accepted 11 September 2006

Published online 22 November 2006

Corresponding Editor: John Dighton

Keywords:

Cantharellus formosus

Cantharellus subalbidus

Cantharellus cascadenis

Habitat

Pacific Northwest

ABSTRACT

We describe watershed-scale habitat associations of three *Cantharellus* species with respect to stand age. During the 1998 autumn fruiting season we collected chanterelle sporocarps from 18 forest stands in and adjacent to the H.J. Andrews experimental forest in the central Cascade Mountains of Oregon. Sampled stands represented two age categories: old growth (~350+ y) and 40–60-y-old second growth naturally regenerated from clear-cut harvest. Old growth and second growth stands were spatially paired to reduce the chance of spurious habitat relationships caused by unmeasured correlated variables. We found stand age to be a good predictor of the distribution of *C. subalbidus* and *C. formosus*, but only marginally useful for predicting the occurrence of *C. cascadenis*. The odds that a randomly located chanterelle sporocarp will be *C. subalbidus*, compared to other chanterelles, are 3–23.5 times higher in old growth than in second growth. Alternatively, there is only a 4–38% chance that a randomly located sporocarp will be *C. formosus* in old growth. *C. cascadenis* was found to be uncommon throughout the study area and showed no significant habitat associations. The abundance of *C. cascadenis* increased substantially with decreasing elevation indicating that landscape features other than stand age may be more useful in predicting its occurrence.

Published by Elsevier Ltd on behalf of The British Mycological Society.

Introduction

Ectomycorrhizal (EM) fungi play a prominent role in forest community dynamics because they form vital mutualisms with forest trees (Smith & Read 1997). Although we have gained important new knowledge on EM fungal communities from sporocarp and root tip studies (Horton & Bruns 2001), our knowledge of actual habitat requirements for individual fungal species remains incomplete (Molina et al. 2001; Dreisbach et al. 2002). This lack of information makes it difficult to include fungi in land management plans where habitat

attributes are maintained to ensure species persistence (USDA & USDI 1994).

The intimate link between EM fungi and their host associates (Molina et al. 1992; Smith & Read 1997), makes plant community composition an important factor for predicting EM species occurrence at regional scales. Within individual watersheds, forest stand age and various disturbance regimes influence the composition of EM communities (Molina et al. 1992; O'Dell et al. 1992; Vogt et al. 1992; Waters et al. 1997; Baar et al. 1999; Colgan et al. 1999; Taylor & Bruns 1999; Krana-better et al. 2005). For example, fire is a primary disturbance

* Corresponding author.

E-mail address: susie.dunham@oregonstate.edu

0953-7562/\$ – see front matter Published by Elsevier Ltd on behalf of The British Mycological Society.

doi:10.1016/j.mycres.2006.09.007

mechanism in Pacific Northwest (PNW) forests (Morrison & Swanson 1990; Agee 1993) and several studies demonstrate its impact on EM communities (Horton et al. 1998; Jonsson et al. 1999; Dahlberg 2002; Smith et al. 2005). Similarly, disturbance associated with forest harvest can affect the local persistence of EM species (Pilz & Perry 1984; Amaranthus & Perry 1987; Jones et al. 2003). In Sweden, where 95 % of forests are used for commercial purposes (Berg et al. 1994), negative impacts of forest management practices are implicated in the high proportion of threatened macrofungi in spruce forests (Rydin et al. 1997; Arnolds 2001).

In the PNW, timber harvest over the last 50 y has significantly reduced the area and fragmented the distribution of old growth forests (Spies & Franklin 1988). As a result, forests previously dominated by old growth Douglas fir (*Pseudotsuga menziesii*) and western hemlock (*Tsuga heterophylla*) have been converted to a mosaic of mature and old growth remnants in a matrix of younger, managed forest (Franklin & Forman 1987; Morrison 1988). Several ecological characteristics that distinguish old growth (~200–1000 y of age) forests from young forests include high photosynthetic productivity that is largely invested in respiration and maintenance rather than new growth, and high nutrient retention. These distinctive functional characteristics are correlated with a complex set of structural characteristics (diverse tree age and size, multi-layered canopies, and abundant dead wood) that also differentiate old growth from young forests (Franklin & Spies 1991). As a result old growth forests support a taxonomically diverse set of organisms, including many fungi (Smith et al. 2002), that may be restricted to this increasingly rare habitat (Lehmkuhl & Ruggiero 1991; Thomas et al. 1993).

Among species with suspected affinities to old growth habitats are some chanterelles in the genus *Cantharellus*, which are popular edible EM fungi both commercially and recreationally harvested from wild populations in the western USA (Danell 1999; Pilz et al. 2003; Pilz et al. 2006). Timber harvests have been reduced on Federal lands over the last decade and concurrently the demand for non-timber forest products has increased. As a result, chanterelles (primarily *C. formosus*; Redhead et al. 1997) have become one of the most widely collected edible mushrooms taken from PNW forests (Pilz & Molina 2002). Because chanterelle species are both economically and ecologically important in the PNW region, forest managers desire information about habitat preferences of each species.

During the autumn of 1997, we conducted genetic studies to determine the size of chanterelle genets (individual mycelial networks) and to assess whether variation in genet size is related to forest disturbance (Dunham et al. 2003a). The species under study were not evenly distributed across the sampled disturbance treatments. A recently described species, *C. cascadenis* (Dunham et al. 2003b), along with *C. subalbidus* primarily occurred on old growth (OG) plots, whereas *C. formosus* dominated second growth (SG) plots. In this study we expanded sampling of OG and SG stands to the watershed scale to characterize the relative abundances and habitat associations of the Pacific golden chanterelle (*C. formosus*), white chanterelle (*C. subalbidus*), and a recently described species (*C. cascadenis*), in old growth and second growth forests.

Methods

Study areas

The H. J. Andrews Experimental Forest (HJA; lat. 44°15'N, long. 122°10'W) is a 6400 ha watershed that has been administered as a long-term ecological research site on the Willamette National Forest since 1948 (Franklin et al. 1990). Elevations within the HJA range from 420–1630 m and the area is typical of the central portion of the western Cascade range in Oregon (Franklin et al. 1990). About 90 % of the annual precipitation falls from October to April with the wettest period in December and driest conditions occurring in July.

Within the HJA, seven OG stands (350+ y old) with no history of recent fire and seven SG stands naturally regenerated from clear cuts (1950–1960) were selected based on their accessibility and spatial arrangement. Two additional pairs of OG and SG stands were selected in the Mill Creek area that lies along the southeast boundary of the HJA. To reduce the chance of spurious habitat associations caused by unmeasured habitat variables (slope, aspect, elevation, etc.) each OG stand was spatially paired with a nearby SG stand. Stands selected for this study ranged from 420–1100 m in elevation, had north, south, southwest, or southeast aspects, and ranged from 20–35 % in average slope. To avoid the confounding affects of additional EM host species (e.g., Pacific silver fir, *Abies amabilis*), we selected only stands dominated by Douglas fir and western hemlock. All stands fall within the western hemlock community type patterns of Franklin (1979) with understory vascular plants including *Polystichum munitum*/*Oxalis oregana* in wet sites, *Berberis nervosa*/*Rhododendron macrophyllum* in mesic sites, *Berberis nervosa*/*Gautheria shallon* in drier sites, and *Linnaea borealis* in cooler, drier sites. The post cutting history of Mill Creek SG stands are not well documented, but the forest zone (Franklin & Dyrness 1973) and habitat associations (Franklin 1979) are similar to the HJA.

Field sampling methods

Collecting began two weeks after the first substantial autumn rain in October of 1998 and ended when the first winter snows covered the area in December. Two individuals searched in each stand for a total of 1.5 h once in October and once in November and data from these two time periods were pooled for analysis. Within each stand, random sampling was initiated by walking to a tree for which the ground at the base was not visible from the road used to access the site. Randomly compass bearings selected at this initial point and at each chanterelle sporocarp collection location were used to direct a continuous series of 150–200 m transects. If a random compass bearing directed a collector out of the stand either the reverse azimuth was followed or a completely new random bearing was selected. Second growth stands ranged from 10–20 ha in total area and the area searched within larger old growth stands roughly approximated this range. When chanterelle sporocarps were located within visual range, a minimum of 5 m (approximate genet size, Dunham et al. 2003a) was kept between any two collections to avoid re-sampling genets. Each collection location was flagged to

avoid re-sampling genets encountered by the two collectors at different times within a sampling effort or during the November collection period. Genetic samples were taken from each sporocarp collected and the species identification confirmed using molecular genetic techniques and vouchers collected during previous studies (Dunham et al. 2003a, b).

Molecular methods

DNA extraction and PCR amplifications of the ITS region were carried out using standard methods described in Dunham et al. (2003a, b). According to the manufacturer's recommendations (GIBCO BRL, Grand Island, NY), unpurified PCR products amplified with ITS1-F (Gardes & Bruns 1993) and ITS4 (White et al. 1990) were digested separately with two restriction enzymes (*Hinf*I and *Hae*III) known to differentiate the three species of interest (Dunham et al. 2003b).

Statistical analysis

Two-dimensional Chi-square analysis was used to determine if the frequency of chanterelle occurrence in the two stand age categories (i.e., OG & SG) was dependent on species identity. The habitat responses of individual species were determined using paired t-test analysis of abundance data with individual stands as replicates. To better characterize the strength of chanterelle habitat association, binomial logistic regression (Allison 1999) was used to test if, relative to two other species, the odds of occurrence for each species is equal in OG and SG. The LOGISTIC procedure available in SAS v.9.1 (SAS Institute Inc.; Allison 1999) was used to estimate intercept and slope parameters for three binary logit models with each equation corresponding to a comparison between the species of interest and all other collections. For example, we used a single binary analysis that included *Cantharellus subalbidus* versus *C. formosus* and *C. cascadenis* combined as alternate response levels to model the relative odds of randomly encountering each species in OG habitat. Drop-in deviance F-tests were used to test the significance of regression slopes that were used to calculate the relative odds of the three chanterelle species occurring in OG stands. Stand location was included as an additional categorical predictive variable in initial regression models to test whether model predictions were equal across the study area.

Results

Distribution and abundance of chanterelle species

A total of 596 chanterelle sporocarps were collected across the 18 OG and SG stands during the autumn fruiting seasons of 1998 (Table 1). Using RFLP analysis we detected only the three genotypes expected from the three species of interest, *Cantharellus formosus*, *C. subalbidus*, and *C. cascadenis*. The overall Chi-square analysis showed that stand age was strongly associated with the identity of species found in stands ($\chi^2 = 144.85$, $df = 2$, $P < 0.0001$). *C. formosus* and *C. subalbidus* occurred in stands across the entire study area with *C. formosus*

significantly associated with SG stands ($t = -2.82$, $df = 8$, $P = 0.023$; Table 1) and infrequent or not detected in OG stands. The reverse was true for *C. subalbidus*, which was significantly associated with OG stands ($t = 4.35$, $df = 8$, $P = 0.003$; Table 1) and infrequent or not detected in SG stands. The total number of *C. cascadenis* collections was lower compared with *C. formosus* and *C. subalbidus* but still adequate for analyses. The distribution of *C. cascadenis* collections was patchy across the study area and its occurrence within stands not associated with stand age ($t = -0.05$, $df = 8$, $P = 0.96$; Table 1).

Stand location had no significant influence in the binary logit models. In other words, the location of paired OG and SG stands did not affect the response (species). However, *C. cascadenis* was not found in four SG and two OG stands and 26 of 34 *C. cascadenis* SG collections were taken from a single SG stand in the Mill Creek area (Table 1). This stand has the lowest elevation (426 m) of all the stands sampled. An OG stand, also in the Mill Creek area, with the second lowest elevation (490 m) had the second highest abundance of *C. cascadenis* with 12 collections. All other stands sampled were 600 m or higher in elevation and contained eight or fewer *C. cascadenis* collections (Table 1, Fig 1). It appears that elevation may be associated with the distribution of *C. cascadenis*, but because elevation was not a factor of interest in the design of this study, it was not sampled well enough to include it as an explanatory variable in logistic regression analyses. To better understand the distribution of this species in stands above 600 m, and independently to examine the influence of stands located below 600 m, we repeated all analyses both with and without stands with mean elevations below 600 m. This did not change the overall results of the Chi-square and paired t-test analyses, but it did dramatically decrease the P-value associated with the *C. cascadenis* comparison ($t = 1.99$, $df = 6$, $P = 0.09$; Table 1) indicating that stand age may weakly influence the distribution of this species at higher elevations.

Strength of habitat associations

Intercept and slope parameters for the logistic regression analyses were estimated for both the data sets including all stands and excluding stands below 600 m (Table 2). When all the observations are considered, *Cantharellus subalbidus* was more likely to occur in OG stands relative to *C. formosus* and *C. cascadenis* ($P = 0.0006$). Conversely, *C. formosus* was more likely to occur in SG relative to *C. subalbidus* and *C. cascadenis* ($P = 0.002$). The likelihood that a randomly located sporocarp will be *C. subalbidus* (versus *C. cascadenis* or *C. formosus*) is 3–23.5 (95 % CI, $P = 0.0006$) times higher when that sporocarp is found in OG than in SG (Table 2).

When stands with mean elevations below 600 m are deleted from the analysis the strong habitat associations of *C. subalbidus* (OG) and *C. formosus* (SG) remain statistically significant with only minor shifts in the 95 % confidence intervals for the odds ratios (Table 2). The comparisons between *C. cascadenis* and the other two species are similar to the t-test results with non-significant results in the overall analysis that move toward a marginal OG association when only stands above 600 m are analysed ($P = 0.98$ and 0.18 , respectively).

Table 1 – Abundances (sporocarp number) of the three chanterelle species pooled across 18 stands

Age	<i>Cantharellus cascadensis</i> (n=67; 16)	Mean Difference ^a (95 % CI)	<i>Cantharellus formosus</i> (n=281; 227)	Mean Difference ^a (95 % CI)	<i>Cantharellus subalbidus</i> (n=248; 217)	Mean Difference ^a (95 % CI)
Old growth (all data)	33 (11.3 %)	–0.1	68 (23.4 %)	–15.0	190 (65.3 %)	14.7
Second growth (all data)	34 (11.1 %)	(–5.6, 5.4)	213 (69.8 %)	(–27.3, –2.7)	58 (19.1 %)	(6.9, 22.5)
Old growth (600 m+)	13 (5.4 %)	1.4	62 (26.0 %)	–13.3	164 (68.6 %)	15.9
Second growth (600 m+)	3 (1.0 %)	(–0.3, 3.2)	165 (75.0 %)	(–26.1, –0.5)	53 (24 %)	(5.5, 26.2)

In the three abundance columns, the values represent the abundance for that species followed by the percentage of the total sporocarps found at that age class and elevation (shown in parentheses). The first two rows show results for the entire data set and the second two rows (600 m+) show data from stands above 600 m in elevation.

a The mean of the differences in abundance (OG–SG) across all stand pairs and the associated 95 % confidence interval around this mean difference.

Discussion

Long-term species persistence is dependent on both the size and geographic distributions of populations (Rabinowitz 1981; Rey Benayas et al. 1999). For EM fungi, these population characteristics are dependent on the status of small and large-scale landscape habitat characteristics. Here we describe the habitat associations of three chanterelle species with respect to stand age. At the watershed scale where stands of 10–20 ha are the sampling units, stand age is a good predictor of the distribution of *Cantharellus subalbidus* and *C. formosus*, but is only marginally important for *C. cascadensis*. Other patterns not captured by this study may explain the distribution of *C. cascadensis* better. Our sampling scale was too coarse to tease out the biological processes that control the presence of species within stands, but clearly stand age and possibly elevation are correlated with the spatial arrangement of finer scale habitat variables important to chanterelle species. A logical extension of this work would be to study the fine scale soil and host tree characteristics that control the distribution of each species within stands (e.g., Bergemann & Largent 2000; Trappe 2004).

The SG stands that we selected are the youngest where abundant chanterelle fruiting is common (S. Dunham, pers. obs.) and the OG stands are some of the oldest and least disturbed sites on the HJA. PNW forest ecosystems are complex and the treatment categories used here (OG and SG) encompass multiple ecological characteristics. We can speculate on potential causal factors, but more research is needed to characterize species distributions in the broad range of intermediate age classes.

Cantharellus subalbidus

Our results demonstrate that the probability of locating *Cantharellus subalbidus* fruit bodies increases significantly in OG habitats. On the HJA, OG surrounds all sampled SG stands creating a habitat mosaic that would maximize the chances of *C. subalbidus* colonizing SG stands as they matured. The reduced odds of finding *C. subalbidus* in SG indicate that they did not persist following logging and that biological, physical or chemical characteristics of the soil or competition from other

newly establishing species are limiting its re-establishment in young stands. Alternatively, *C. subalbidus* may be able to establish mycelia in young stands, but subsequently experience suppressed fruiting until OG characteristics begin to develop. Characteristic features of OG forests include the abundance of coarse woody debris in all stages of decay, uneven aged trees, gaps and uneven soil surface caused by large uprooted trees (Spies et al. 1988; Spies & Franklin 1989; 1991).




Fig 1 – Number of *Cantharellus cascadensis* collections plotted against the mean elevation of the paired old growth and second growth stands where collections were made. Open circles represent old growth collections and closed circles represent second growth collections. At two elevations equal abundances are represented by overlapping filled and open circles.

Table 2 – Parameter estimates for the relationship between species type (response) and stand age using three binary logit models

Estimate	<i>Cantharellus cascadenis</i> All data; above 600 m	<i>Cantharellus subalbidus</i> All data; above 600 m	<i>Cantharellus formosus</i> All data; above 600 m
Intercept	–2.06; –3.51	–1.45; –1.32	0.84; 1.15
Standard error	0.62; 0.69	0.40; 0.41	0.38; 0.37
95 % CI ^a	–3.53, –1.01; –5.24, –2.38	–2.31, –0.72; –2.19, –0.58	0.13, 1.63; 0.46, 1.93
Stand age	0.019; 1.09	2.08; 2.01	–2.03; –2.24
Standard error	0.88; 0.82	0.52; 0.54	0.57; 0.52
95 % CI	–1.80, 1.82; –0.42, 2.97	1.10, 3.16; 0.98, 3.13	–3.19, –0.96; –3.32, –1.25
P-value ^b	$F_{1,16}=0.00$; $F_{1,14}=1.97$ $p = 0.98$; $p = 0.18$	$F_{1,16}=18.26$; $F_{1,14}=15.49$ $p = 0.0006$; $p = 0.002$	$F_{1,16}=14.59$; $F_{1,14}=21.10$ $p = 0.002$; $p = 0.0004$
Odds ratio	1.02; 2.97	8.00; 7.46	0.13; 0.11
95 % CI	0.17, 6.17; 0.66, 19.49	3.00, 23.57; 2.66, 22.87	0.04, 0.38; 0.04, 0.29

In each column the second set of values are parameter estimates from analyses where stands below 600 m in elevation were not included.
a Maximum likelihood ratio estimates.
b From drop in deviance likelihood ratio tests.

The importance of coarse woody debris in predicting the occurrence of threatened macrofungi in late successional conifer forests in Europe has been demonstrated (Berg *et al.* 1994; Rydin *et al.* 1997). Wide confidence intervals for the odds ratio characterizing the association between *C. subalbidus* and OG may result from the retention of coarse woody debris in SG stands allowing for patchy development of pre-harvest soil conditions required by this species.

The PNW region has been an important supplier to the world timber market (Waddell *et al.* 1989). Clear-cut harvesting and subsequent even-aged forest management has been practised on private lands for well over a century (Robbins 1988) and on federal lands since the 1940s (Harris 1984). During the past few decades, rotation lengths on public lands have generally been in the range of 80–100 y, while private landowners generally cut on rotation lengths of 50 y (Spies *et al.* 1994). This intense forest harvest activity has resulted in profound alterations in the age structure of the forest environment since extensive logging began (Wallin *et al.* 1996). Estimates of the percentage of old growth forests >200 y old remaining from pre-logging time periods range from 13.1 % (Spies & Franklin 1988; Booth 1991) to 17.5 % (Haynes 1986). Depending on the minimum stand age at which the OG characteristics necessary for the establishment and fruiting of *C. subalbidus* develop, the conversion of PNW forests from OG to SG may have significantly reduced the distribution of *C. subalbidus* in the western Cascades.

This assertion depends on how representative the HJA is of the western Cascades as a whole. The HJA is representative of the winter climate of the northern Cascades and of the PNW in general (Greenland 1994) and has been included in an enormous amount of basic and applied research used in the development of forest management plans for the region (Cissel *et al.* 1999). If these results can be extended to other areas, *C. subalbidus* has likely been largely excluded from private lands with short cutting rotations and possibly also from large portions of public lands if 80–100 y rotation cycles are not long enough for it to establish and maintain viable populations. *C. subalbidus* was also listed under the survey and manage guidelines of the Northwest Forest Plan (USDA & USDI 1994) that were intended to protect rare, little-known

species associated with late-successional (OG) forests. Our results confirm the strong association of *C. subalbidus* with late-successional forests, thus verifying original assumptions made by experts who developed the plan. Future research on this species should be directed towards identifying factors that limit its distribution and determining if appropriate conditions for its establishment and persistence develop as SG stands continue to mature beyond 50 y.

Cantharellus formosus

Analyses presented here indicate that *Cantharellus formosus* is strongly associated with 40–60-y-old SG habitats so the large-scale harvesting disturbance that has taken place throughout the PNW probably has increased the distribution of this species. Given the proximity of our sampled SG stands to OG, it is likely that *C. formosus* is adapted to soils altered by disturbance or to physiological differences associated with younger host trees. Buried coarse woody debris has been identified as a factor important to the ecology of *C. formosus* (Largent & Sime 1994; Norvell *et al.* 1996) but in our OG soils this was not a limiting resource. Conversely, quantitative work completed by Bergemann & Largent (2000) demonstrated that above-ground coarse woody debris is not a significant predictor of *C. formosus* occurrence. Instead, the percent cover of bare humus unoccupied by moss, moderate duff depth, and low exchangeable acidity, were found to be useful variables for predicting *C. formosus* fruiting (Bergemann & Largent 2000) on small study plots.

Disturbances associated with forest harvest usually result in a loss of soil spatial heterogeneity (Miller & Lodge 1997) that can alter many soil characteristics important to EM fungi (Harvey *et al.* 1980a, b; Pilz & Perry 1984; Amaranthus & Perry 1987). After such a major disturbance, stress tolerance can allow some fungal species to replace others that are less tolerant (Lodge & Cantrell 1995). Fungi that are favoured by disturbance generally exhibit effective dispersal, rapid nutrient uptake, and rapid mycelial extension for capturing nutrients and water (Pugh & Boddy 1988). The age at which stands regenerating from clear-cut harvest are colonized by *C. formosus* is unclear because the fungus may establish the EM

symbiosis with young trees early on but delay fruiting until the stand is somewhat recovered from the initial disturbance. Future research on this species should be directed at exploring conditions favourable to both host tree colonization and the onset of fruiting to determine if there is a significant time lag between the two events.

Cantharellus cascadenis

Uneven abundances and the patchy distribution of *Cantharellus cascadenis* across the study area make characterization of its habitat associations difficult given our study design. However, the spatial pairing of stands has allowed us to detect a confounding variable possibly associated with the distribution of *C. cascadenis*. It is possible that environmental variables correlated with stand age have some weak control over the distribution of *C. cascadenis* but this relationship is not constant over the range of elevation we sampled as the size of *C. cascadenis* populations increased substantially at elevations below 600 m. It was clear from this sampling that *C. cascadenis* is uncommon in the study area relative to other chanterelles and much survey work still is required to fully understand its distribution in Oregon. This analysis indicates that future survey work should focus on lower elevation habitats.

Acknowledgements

We thank Jason Dunham, Lisa Ganio, Manuela Huso, Dave Pilz and Julia Jones for advice on experimental design and data analysis and for comments on previous versions of this manuscript, Teresa Lebel and Mike Taft for assistance in the field, and Dylan Nelson for laboratory assistance. S.D. thanks Joey Spatafora for support during the preparation of this manuscript. The constructive comments of two anonymous reviewers greatly improved the clarity of this manuscript. This research was funded by cooperative agreement PNW 97-5090-2-CA between the USDA Forest Service and Oregon State University.

REFERENCES

- Agee JK, 1993. *Fire Ecology of Pacific Northwest Forests*. Island Press, Washington, DC.
- Allison PD, 1999. *Logistic Regression using the SAS System: Theory and Application*. SAS Institute, Cary, NC.
- Amaranthus MP, Perry DA, 1987. Effect of soil transfer on ectomycorrhiza formation and survival and growth of conifer seedlings in disturbed forest sites. *Canadian Journal of Forest Research* **17**: 944–950.
- Arnolds E, 2001. The future of fungi in Europe: threats, conservation and management. In: Moore D, Nauta MM, Evans SE, Rotheroe M (eds), *Fungal Conservation: Issues and Solutions*. Cambridge University Press, Cambridge, pp. 64–80.
- Baar J, Horton TR, Kretzer AM, Bruns TD, 1999. Mycorrhizal colonization of *Pinus muricata* from resistant propagules after a stand-replacing wildfire. *New Phytologist* **143**: 409–418.
- Berg Å, Ehnström B, Gustafsson L, Hallingbäck T, Jonsell M, Weslien J, 1994. Threatened plant, animal, and fungus species in Swedish forests: distributions and habitat associations. *Conservation Biology* **8**: 718–731.
- Bergemann SE, Largent DL, 2000. The site specific variables that correlate with the distribution of the Pacific Golden Chanterelle, *Cantharellus formosus*. *Forest Ecology and Management* **130**: 99–107.
- Booth DE, 1991. Estimating prelogging old-growth in the Pacific Northwest. *Journal of Forestry* **89**: 25–29.
- Cissel JH, Swanson FJ, Weisberg PJ, 1999. Landscape management using historical fire regimes: Blue River, Oregon. *Ecological Applications* **9**: 1217–1231.
- Colgan W, Carey AB, Trappe JM, Molina R, Thysell D, 1999. Diversity and productivity of hypogeous fungal sporocarps in a variably thinned Douglas fir forest. *Canadian Journal of Forest Research* **29**: 1259–1268.
- Dahlberg A, 2002. Effects of fire on ectomycorrhizal fungi in fenoscandian boreal forests. *Silva Fennica* **36**: 69–80.
- Danell E, 1999. *Cantharellus*. In: Cairney JWG, Chambers SM (eds), *Ectomycorrhizal Fungi: Key Genera in Profile*. Springer-Verlag, Heidelberg, pp. 253–267.
- Dreisbach TA, Smith JE, Molina R, 2002. Challenges of modeling fungal habitat: when and where do you find chanterelles? In: Scott JM, Heglund PJ, Morrison M, Raphael M, Haufler J, Wall B (eds), *Predicting Species Occurrence: Issues of Scale and Accuracy*. Island Press, Covello, CA, pp. 475–481.
- Dunham SM, Kretzer A, Pfrender ME, 2003a. Characterization of the extent and distribution of Pacific golden chanterelle (*Cantharellus formosus*) individuals using co-dominant microsatellite loci. *Molecular Ecology* **12**: 1607–1618.
- Dunham SM, O'Dell TE, Molina R, 2003b. Analysis of nrDNA sequences and microsatellite allele frequencies reveals a cryptic chanterelle species *Cantharellus cascadenis* sp. nov. from the American Pacific Northwest. *Mycological Research* **107**: 1163–1177.
- Franklin JF, 1979. Vegetation of the Douglas fir region. In: Heilman PE, Anderson HW, Baumgartner DM (eds), *Forest Soils of the Douglas Fir Region*. Washington State University, Cooperative Extension, Pullman, pp. 93–112.
- Franklin JF, Bledsoe CS, Callahan JT, 1990. Contributions of the long-term ecological research program. *Bioscience* **40**: 509–523.
- Franklin JF, Dyrness CT, 1973. *Natural Vegetation of Oregon and Washington* USDA Forest Service General Technical Report PNW-8. Pacific Northwest Forest and Range Exp. Station, Portland, Oregon.
- Franklin JF, Forman RTT, 1987. Creating landscape patterns by forest cutting: ecological consequences and principles. *Landscape Ecology* **1**: 518.
- Franklin JF, Spies TA, 1991. Composition, function, and structure of old-growth Douglas fir forests. In: Ruggiero LF, Aubry KB, Huff MH (eds), *Wildlife and Vegetation of Unmanaged Douglas Fir Forests* USDA Forest Service General Technical Report PNW-GTR-285, pp. 71–92.
- Gardes M, Bruns TD, 1993. ITS primers with enhanced specificity for Basidiomycetes: applications to the identification of mycorrhizae and rusts. *Molecular Ecology* **2**: 113–118.
- Greenland D, 1994. The Pacific Northwest regional context of the climate of the H.J. Andrews Experimental Forest. *Northwest Science* **69**: 81–96.
- Harris LD, 1984. *The Fragmented Forest*. University of Chicago Press, Chicago.
- Harvey AE, Jurgensen MF, Larsen MJ, 1980a. Clearcut harvesting and ectomycorrhizae: survival of activity on residual roots and influence on bordering forests stance in western Montana. *Canadian Journal of Forest Research* **10**: 300–303.
- Harvey AE, Larsen MJ, Jurgensen MF, 1980b. Early effects of partial cutting on the number and distribution of active ectomycorrhizae in steep, rocky soils of Douglas fir/larch forests in

- western Montana. *Canadian Journal of Forest Research* 10: 436–440.
- Haynes RW, 1986. *Inventory and Value of Old-growth in the Douglas fir region* USDA Forest Service Research Note PNW-437.
- Horton TR, Bruns TD, 2001. The molecular revolution in ectomycorrhizal ecology: peeking into the black box. *Molecular Ecology* 10: 1855–1871.
- Horton TR, Cazares E, Bruns TD, 1998. Ectomycorrhizal vesicular-arbuscular and dark septate fungal colonization of bishop pine (*Pinus muricata*) seedlings in the first five months of growth after wildfire. *Mycorrhiza* 8: 11–18.
- Jones MD, Durall DM, Cairney JWJ, 2003. Ectomycorrhizal fungal communities in young forest stands regenerating after clear-cut logging. *New Phytologist* 157: 399–422.
- Jonsson L, Dahlberg A, Nilsson M-C, Zackrisson O, Kårén O, 1999. Ectomycorrhizal fungal communities in late-successional Swedish boreal forests, and their composition following wildfire. *Molecular Ecology* 8: 205–215.
- Kranabetter JM, Friesen J, Gamiet S, Kroeger P, 2005. Ectomycorrhizal mushroom distribution by stand age in western hemlock – lodgepole pine forests of northwestern British Columbia. *Canadian Journal of Forest Research* 35: 1527–1539.
- Largent DL, Sime AD, 1994. A preliminary report on the phenology, sporulation, and lifespan of *Cantharellus cibarius* and *Boletus edulis* basidiomes at Patrick's Point State Park. In: Adams DH, Rios JE, Stoere AJ (eds), *Proceedings of the Symposium of the 43rd Annual Meeting of the California Forest Pest Council*. 16–17 November 1994, Rancho Cordova, CA.
- Lehmkuhl JF, Ruggiero LF, 1991. Forest fragmentation in the Pacific Northwest and its potential effects on wildlife. In: Ruggiero LF, Aubry AB, Huff MH (eds), *Wildlife and Vegetation of Unmanaged Douglas Fir Forests*. USDA Forest Service General Technical Report PNW-GTR-285, pp. 35–69.
- Lodge DJ, Cantrell S, 1995. Fungal communities in wet tropical forests: variation in time and space. *Canadian Journal of Botany* 73: S1391–S1398.
- Miller RM, Lodge DJ, 1997. Fungal responses to disturbance: agriculture and forestry. In: Wicklow DT, Söderström B (eds), *The Mycota Vol. 4. Environmental and Microbial Relationships*. Springer-Verlag, Berlin, pp. 65–84.
- Molina R, Massicotte H, Trappe JM, 1992. Specificity phenomena in mycorrhizal symbioses: community-ecological consequences and practical implications. In: Allen M (ed), *Mycorrhizal Functioning, and Integrative Plant–Fungal Process*. Chapman and Hall, New York, pp. 357–423.
- Molina R, Pilz D, Smith J, Dunham S, Dreisbach T, O'Dell T, Castellano M, 2001. Conservation and management of forest fungi in the Pacific Northwestern United States: an integrated ecosystem approach. In: Moore D, Nauta MM, Evans SE, Rotheroe M (eds), *Fungal Conservation: Issues and Solutions*. Cambridge University Press, Cambridge, pp. 19–63.
- Morrison PM, 1988. *Old Growth in the Pacific Northwest: A Status Report*. The Wilderness Society, Washington, DC.
- Morrison P, Swanson FJ, 1990. *Fire History and Pattern in a Cascade Range Landscape*. Pacific Northwest Research Station, Portland Oregon, USA, U.S. Forest Service General Technical Report PNW-GTR-254.
- Norvell L, Roger J, Lindgren J, Kopecky F, 1996. Oregon *Cantharellus* study project 1986–96. In: Pilz D, Molina R (eds), *Managing Forest Ecosystems to Conserve Fungus Diversity and Sustain Wild Mushroom Harvests*. General Technical Report PNW-GTR-371. US Department of Agriculture, Forest Service, Pacific Northwest Research Station, Portland, OR, pp. 69–72.
- O'Dell TE, Luoma DL, Molina RJ, 1992. Ectomycorrhizal fungal communities in young, managed, and old-growth Douglas fir stands. *Northwest Environmental Journal* 8: 166–168.
- Pilz D, Molina R, 2002. Commercial harvests of edible mushrooms from the forests of the Pacific Northwest United States: issues, management, and monitoring for sustainability. *Forest Ecology and Management* 155: 3–16.
- Pilz D, Molina R, Mayo J, 2006. Effects of thinning young forests on chanterelle mushroom production. *Journal of Forestry* 104: 9–14.
- Pilz D, Norvell L, Danell E, Molina R, 2003. *Ecology and Management of Commercially Harvested Chanterelle Mushrooms*. General Technical Report PNW-GTR-576. USDA Forestry Service, Pacific Northwest Research Station, Portland, OR.
- Pilz D, Perry DA, 1984. Impact of clearcutting and slash burning on ectomycorrhizal associations of Douglas fir seedlings. *Canadian Journal of Forest Research* 14: 94–100.
- Pugh GJF, Boddy L, 1988. A view of disturbance and life strategies in fungi. *Proceedings of the Royal Society of Edinburgh* 94B: 3–11.
- Rabinowitz D, 1981. Seven forms of rarity. In: Synge H (ed), *The Biological Aspects of Rare Plant Conservation*. John Wiley, Chichester, pp. 205–207.
- Redhead SA, Norvell LL, Danell E, 1997. *Cantharellus formosus* and the Pacific Golden Chanterelle harvest in Western North America. *Mycotaxon* 65: 285–322.
- Rey Benayas JM, Scheiner SM, García Sánchez-Colomer M, Levassor C, 1999. Commonness and rarity: theory and application of a new model to Mediterranean montane grasslands. *Conservation Ecology* 3: 5.
- Robbins WG, 1988. *Hard Times in Paradise, Coos Bay, Oregon 1850–1986*. University of Washington Press, Seattle, WA.
- Rydin H, Martin D, Hallingbäck T, 1997. Biological characteristics, habitat associations, and distribution of macrofungi in Sweden. *Conservation Biology* 11: 628–640.
- Smith JE, McKay D, Brenner G, McIvers J, Spatafora JW, 2005. Early impacts of forest restoration treatments on the ectomycorrhizal fungal community and fine root biomass in a mixed conifer forest. *Journal of Applied Ecology* 42: 526–535.
- Smith JE, Molina R, Huso MMP, Luoma D, McKay D, Castellano MA, Lebel T, Valachovic Y, 2002. Species richness, abundance, and composition of hypogeous and epigeous ectomycorrhizal fungal sporocarps in young, rotationage, and old-growth stands of Douglas fir (*Pseudotsuga menziesii*) in the Cascade Range of Oregon, USA. *Canadian Journal of Botany* 80: 186–204.
- Smith SE, Read DJ, 1997. *Mycorrhizal Symbiosis*, 2nd edn. Academic Press, San Diego and London.
- Spies TA, Franklin JF, 1988. Old-growth and forest dynamics in the Douglas fir region of western Oregon and Washington. *Natural Areas Journal* 8: 190–201.
- Spies TA, Franklin JF, 1989. Gap characteristics and vegetation response in coniferous forests of the Pacific Northwest. *Ecology* 70: 543–545.
- Spies TA, Franklin JF, 1991. The structure of natural young, mature, and old-growth Douglas fir forests in Oregon and Washington. In: Ruggiero LF, Aubry KB, Carey AB, Huff MH (eds), *Wildlife and Vegetation of Unmanaged Douglas Fir Forests*, pp. 91–109. USDA Forest Service General Technical Report PNW-GTR-285.
- Spies TA, Franklin JF, Thomas TB, 1988. Coarse woody debris in Douglas fir forests of western Oregon and Washington. *Ecology* 69: 1689–1702.
- Spies TA, Ripple WJ, Bradshaw GA, 1994. Dynamics and pattern of a managed coniferous forest landscape. *Ecological Applications* 4: 555–568.
- Taylor DL, Bruns TD, 1999. Community structure of ectomycorrhizal fungi in a *Pinus muricata* forest: minimal overlap between the mature forest and resistant propagule communities. *Molecular Ecology* 8: 1837–1850.
- Thomas JW, Raphael MG, Anthony RG, Forsman ED, Gunderson AG, Holthausen RS, Marcot BG, Reeves GH, Sedell JR, Solis DM, 1993. *Viability Assessments and Management*

- Considerations for Species Associated with Late-successional and Old-Growth Forests of the Pacific Northwest*. USDA Forest Service, Portland, Oregon.
- Trappe MJ, 2004. Habitat and host associations of *Craterellus tubaeformis* in northwestern Oregon. *Mycologia* **96**: 498–509.
- USDA (U.S. Department of Agriculture) Forest Service, USDI (U.S. Department of Interior) Bureau of Land Management, 1994. *Record of Decision on Management of Habitat for Late-successional and Old-growth Forest Related Species within the Range of the Northern Spotted Owl (Northwest Forest Plan)*. U.S. Department of Agriculture, Forest Service and U.S. Department of Interior, Bureau of Land Management, Portland, OR.
- Vogt KA, Bloomfield J, Ammirati JF, Ammirati SR, 1992. Sporocarp production by *Basidiomycetes*, with emphasis on forest ecosystems. In: Carroll GC, Wicklow DT (eds), *The Fungal Community: Its Organization and Role in the Ecosystem*, 2nd edn. Marcel Dekker, New York, pp. 563–581.
- Waddell KL, Oswald DD, Powell DS, 1989. *Forest Statistics of the United States, 1987*. Pacific Northwest Forest and Range Experiment Station, U.S. Department of Agriculture Forest Service, Resource Bulletin PNW-RB-168.
- Wallin DO, Swanson FJ, Marks B, Cissel JH, Kertis J, 1996. Comparison of managed and pre-settlement landscape dynamics in forests of the Pacific Northwest, USA. *Forest Ecology and Management* **85**: 291–309.
- Waters JR, McKelvey KS, Luoma DL, Zabel CJ, 1997. Truffle production in old-growth and mature fir stands in northeastern California. *Forest Ecology and Management* **96**: 155–166.
- White TJ, Bruns TD, Lee SB, Taylor JW, 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis N, Gelfand D, Sninsky J, White T (eds), *PCR Protocols: A Guide to Methods and Applications*. Academic Press, San Diego, pp. 315–322.