

***Vesicomycicola trifurcatus*, a new genus and species of commensal polychaete (Annelida: Polychaeta: Nautiliniellidae) found in deep-sea clams from the Blake Ridge cold seep**

Jennifer Dreyer, Tomoyuki Miura, and Cindy Lee Van Dover

(JD, CLVD) The College of William and Mary, Department of Biology, Millington Hall, Williamsburg, Virginia 23187, U.S.A., email: jcdrey@wm.edu; clvand@wm.edu

(TM) Department of Biological Production and Environmental Science, Faculty of Agriculture, Miyazaki University, 1-1 Gakuen-Kibanadai-Nishi, Miyazaki 889-2192, Japan, email: miura@cc.miyazaki-u.ac.jp

Abstract.—A new genus and species of deep-sea polychaete belonging to the family Nautiliniellidae is described from the Blake Ridge cold seep off the coast of South Carolina at a depth of 2155 m. This species is commensal within the mantle cavity of ~60% of the vesicomimid clams collected at the seep site. *Vesicomycicola trifurcatus* is distinguished from previously described nautiliniellid genera and species by the presence of two pairs of tentacular cirri and up to seven trifurcate hooked chaetae on the posterior parapodia. The new species resembles *Theyomytilidicola tridentatus* in having trifurcate hooks, but the arrangement and number of chaetae differs. Only two types of chaetae are present in *V. trifurcatus*: four to seven stout, simple hooks anteriorly to mid-body, and up to seven trifurcate hooks posteriorly. In contrast, there are three types of chaetae in *I. tridentatus*: up to five stout hooks per parapodium, each with a minute projection on cutting edge of the main fang, 10–20 simple, slender tridentate chaetae, and numerous minute mucronate chaetae. A key to species of Nautiliniellidae is included.

The Nautiliniellidae is a small group of deep-sea polychaetes that live in the mantle cavity of a clam or mussel host. Nautiliniellids have been collected from chemosynthetically based deep-sea habitats, including cold seeps and hydrothermal vents. Since nautiliniellids were first reported by Miura & Laubier (1989), 10 genera and 14 species have been described (Table 1). Two undescribed species have also been reported, one from a cold seep at Barbados Trench (4960 m; Olu et al. 1996) and one off the Pacific coast of Mexico (3221 m; Olu, pers. comm.).

An additional genus, *Santelma*, has been assigned to the family Nautiliniellidae (Blake 1993, Glasby 1993), but its affiliation with the Nautiliniellidae remains questionable. The only known species, *Santelma*

miraseta (Fauchald, 1972), was first placed in the family Pilargidae and the genus *Pilargis*. Blake (1993) redescribed the species and assigned it to *Santelma*, a new nautiliniellid genus, based on chaetal similarities. Unlike nautiliniellids, *S. miraseta* has extruded neuroaciculae, a median antenna (or its trace), and it lacks neuropodial hooks and parapodial cirri. Based on these features, *S. miraseta* fits better within the original family Pilargidae (Salazar-Vallejo, pers. comm.). We follow the precedent of Miura & Hashimoto (1996) and exclude *S. miraseta* from the Nautiliniellidae.

Nautiliniellids have reduced and simplified body structures that are associated with a commensal or parasitic life. These modifications include a less developed anterior region, the presence of only simple hooked

Table 1.—Family Nautiliniellidae: list of genera and species, host bivalve genus and family, collection depth of type specimen, location where type specimen was collected and author reference.

Genus and species	Host bivalve Genus (Family)	Depth (m)	Location	Reference
1 <i>Flascarpia alvinae</i>	Unknown	3303	Florida Escarpment	Blake (1993)
2 <i>Itheyomytilidicola tridentatus</i>	<i>Bathymodiolus</i> (Mytilidae)	1395	Okinawa Trough	Miura & Hashimoto (1996)
3 <i>Laubierus mucronatus</i>	(Mytilidae)	3243	Florida Escarpment	Blake (1993)
4 <i>Miura spinosa</i>	Unknown	565	Santa Maria Basin	Blake (1993)
5 <i>Mytilidiphila enseiensis</i>	near <i>Adula</i> (Mytilidae)	625	Okinawa Trough	Miura & Hashimoto (1993)
6 <i>Mytilidiphila okinawaensis</i>	<i>Bathymodiolus</i> (Mytilidae)	701	Okinawa Trough	Miura & Hashimoto (1993)
7 <i>Natsushima bifurcata</i>	<i>Solemya</i> (Solemyidae)	1114	Sagami Bay	Miura & Laubier (1990)
8 <i>Natsushima graciliceps</i>	<i>Solemya</i> (Solemyidae)	98	Kagoshima Bay	Miura & Hashimoto (1996)
9 <i>Nautiliniella cabypotogenicola</i>	<i>Calyptogena</i> (Vesicomomyidae)	5650	Japan Trench	Miura & Laubier (1989)
10 <i>Petrecca thyasira</i>	<i>Thyasira</i> (Thyasiridae)	3700	Laurentian Fan	Blake (1990)
11 <i>Shinkai longipedata</i>	<i>Calyptogena</i> (Vesicomomyidae)	1400	Okinawa Trough	Miura & Ohra (1991)
12 <i>Shinkai sagamiensis</i>	<i>Calyptogena</i> (Vesicomomyidae)	1170	Sagami Bay	Miura & Laubier (1990)
13 <i>Shinkai semitlonga</i>	<i>Calyptogena</i> (Vesicomomyidae)	625	Okinawa Trough	Miura & Hashimoto (1996)
14 <i>Thyasiridicola branchiatus</i>	<i>Conchocèle</i> (Thyasiridae)	1160	Sagami Bay	Miura & Hashimoto (1996)
15 <i>Vesicomomyicola trifurcatus</i>	<i>Vesicomomya</i> (Vesicomomyidae)	2155	Blake Ridge Diapir	Present study

chaetae modified to grasp host tissue, and the absence of anal cirri on the pygidium. Diagnostic characters of the family include the number of prostomial appendages, number of tentacular cirri, and chaetal morphology and number. These characters are specific to each genus but are useful for species identifications since seven of the ten nautiliniellid genera are monospecific. Based on these morphological characters, we determined that the specimens collected from the Blake Ridge cold seep belong to a new genus and species described herein.

Material and Methods

Biological samples were collected at the Blake Ridge Diapir site (ODP Site 996; 32°30'N, 76°11'W; 2155 m) on 25 to 28 Sep 2001, using the DSV *Alvin*. A description of the study site can be found in Van Dover et al. (2003). Although geological and chemical properties of this site have been explored during the past decade, the *Alvin* 2001 samples represent the first collections of megafauna and macrofauna from this area.

Host clams were collected using a suction sampler. The clams were identified as a new genus and species in the Family Vesicomomyidae, based on morphological characters, molecular differences in comparison to described species, and geographic and bathymetric location (E. Kryolora, pers. comm.). Clams were dissected and nautiliniellids were removed and placed into either 10% buffered formalin or 3% glutaraldehyde and 0.1 M phosphate buffer with 0.25 M sucrose (pH 7.4). After 24 hours, formalin-fixed nautiliniellids were rinsed and stored in 70% ethanol.

Photographs of the external morphology were taken with a compound light microscope (LM) and a scanning electron microscope (SEM). Specimens for LM were mounted in glycerol and ethanol and observed with a Zeiss Axioskop 2 binocular compound microscope. Specimens for SEM were dehydrated through a graded series of

ethanol, terminating with 100% ethanol. Samples were then critical-point dried, gold sputter coated (20 nm thick), and observed with an Amray SEM 1810. Images were captured using a Spot camera (Diagnostic Instruments) or a DP11 digital camera (Olympus). Line illustrations were prepared using a camera lucida attached to a Wild Heerbrugg compound microscope.

Systematics

Family Nautiliniellidae Miura & Laubier,
1989

Vesicomycicola, new genus

Type species.—*Vesicomycicola trifurcatus*, new species, by present designation.

Diagnosis.—Body with strong dorsal arch, ventrally flattened. Prostomium with one pair of palps, without eyes. Tentacular segment fused with prostomium, with dorsal and ventral cirri, neuroacicula, and neuropodial hooked chaetae. Parapodia sub-biramous, with dorsal and ventral cirri. Notopodia and neuropodia each with one embedded acicula. Chaetae absent on notopodia. Two types of chaetae present on neuropodia: simple hooked chaetae on anterior segments (some with single subapical tooth present on anterior to mid-body segments), and tricurcate hooked chaetae on posterior segments. Pygidium cylindrical, without anal cirri.

Gender.—Masculine.

Etymology.—The generic name is derived from the name of the host vesicomycid clams these polychaetes inhabit.

Vesicomycicola trifurcatus, new species (Figs. 1–4)

Type material.—Holotype (ODP Site 996; 32°30'N, 76°11'W; 2155 m, 28 Sep 2001, Alvin Dive 3712; USNM 1016220) and five paratypes (USNM 1016221) from same dive and date were deposited in the collections of the National Museum of Natural History, Smithsonian Institution, Washington, District of Columbia. An ad-

ditional five paratypes, each from the same dive and date, were deposited in the Museum National d'Histoire Naturelle, Paris (MNHN POLY TYPE 1405) and the National Science Museum, Tokyo (NSMT—Pol P 458).

Additional material.—Voucher specimens were retained in the collection of CLVD in the Department of Biology at The College of William and Mary.

Description.—Holotype female, ovigerous, measuring 8.4 mm long, 1.3 mm wide, including parapodia, with 37 segments. Paratypes ranging from 4.4–12.7 mm long, 0.8–1.6 mm wide, including parapodia, and with 28–41 segments. Body flattened ventrally, arched dorsally. Some live specimens with green pigment in parapodia, others with pale pink color; preserved specimens in alcohol pink to white in color. Some preserved females pale green; internal oocytes evident through transparent parapodial epidermis. Preserved holotype and paratypes curled (Fig. 1A).

Prostomium rounded, with palps (Fig. 2A–C). Eyes absent. Tentacular segment fused with prostomium, with one pair of dorsal and ventral cirri, neuroacicula, and neuropodial hooked chaetae (Fig. 2C). Foregut with well-developed muscular region (Fig. 2A–C). Pygidium rounded, without anal cirri (Fig. 2D).

Parapodia subbiramous, with dorsal and ventral cirri. Dorsal cirri with inflated base and tapering tip, twice as long as ventral cirri. Notopodia with single embedded acicula, lacking chaetae (Fig. 3A). Neuropodia with a single bent acicula and hooked chaetae (Fig. 3B).

Neuropodial hooks of two types. Anterior neuropodia with simple stout hooks with recurved tips, four to seven on each parapodium (Fig. 4A, B), some anterior to mid-body chaetae with single small apical tooth near tip, appearing slightly bifid (Fig. 4C). Posterior neuropodia with thinner, simple hooks with trifurcate tips, up to seven per neuropodium (Fig. 4D, E).

Etymology.—The specific name comes




Fig. 1. *Vesicomycicola trifurcatus* new genus, new species. Scanning electron micrograph (SEM) of whole body.

from *tri-* = three times, + *furcatus* = forked, in reference to the trifurcate chaetae present on the posterior segments.

Biology.—The mantle cavities of ~60% of the Blake Ridge clams sampled contained one to five nautiliniellid polychaetes. Carbon and nitrogen stable isotope compositions of worm and clam tissues were consistent with a parasitic life-style for the worm, but the sulfur isotope composition of the worms was so distinct from that of the clams that an alternative diet must be inferred (Van Dover et al. 2003). Van Dover et al. (2003) proposed a feeding strategy whereby ciliary activity of the clam gills moves sufficient volumes of seawater to allow the polychaetes to collect and consume suspended organic particles either from gill mucus or from a worm-generated mucus net.

Discussion

Vesicomycicola trifurcatus resembles species in the genera *Nautiliniella*, *Natushi-*

ma, *Shinkai*, and *Thyasiridicola*, based on shared characters of the tentacular segment, which in these four genera includes dorsal and ventral cirri and neurochaetae (with the exception of the genus *Thyasiridicola*, which lacks neurochaetae). The genus *Vesicomycicola* differs from these four genera in the number and morphology of the neuropodial chaetae.

Vesicomycicola trifurcatus resembles *Iheyomytilidicola tridentatus* Miura & Hashimoto, 1996 based on the trifurcate chaetal morphology, but the arrangement and number of chaetae on the parapodia differs. There are only two types of chaetae present in *V. trifurcatus*: stout, simple hooks (four to seven; sometimes bifid) on the anterior to mid-body parapodia, and trifurcate hooks (up to seven) on the posterior parapodia. In contrast, there are three types of chaetae in *I. tridentatus*: stout hooks (up to five), each with a minute projection on the cutting edge of the main fang; simple,




Fig. 2. *Vesicomycicola trifurcatus* new genus, new species. A. Light micrograph (LM) of anterior end, dorsal view. B. Drawing of anterior end, dorsal view. C. Drawing of anterior end, ventral view. D. LM of pygidium, dorsal view.

slender tridentate chaetae (10–20); and numerous minute chaetae with mucronate tips (Miura & Hashimoto 1996).

Based on its unique set of morphological characters, we consider *V. trifurcatus* to be a new genus and species. A key to nautiliniellid species is provided to aid in identification; most species are location and host specific.

The terminology and interpretation of prostomial appendages in this family is the subject of some debate (Blake 1993, Miura & Hashimoto 1996), suggesting the need for a re-evaluation and revision of this family and its genera once a consistent diagnosis of prostomial appendages can be applied.

Color dimorphism was a distinctive character of live *V. trifurcatus*, but on preservation the color variation was lost. Polychaetes with green parapodia in new collections (2003) were all gravid females. In other nautiliniellid species, color dimorphism corresponds to sexual dimorphism (Miura & Hashimoto 1996, Miura 1998). We have yet to confirm that the pale colored specimens are males. With the discovery of each new species in the Nautiliniellidae, we learn more about the ecology of these worms and their relationship with their host bivalves; we still know little about the internal anatomy, reproductive biology and larval characteristics, or the trophic ecology of this polychaete family.




Fig. 3. *Vesicomycicola trifurcatus* new genus, new species. A. Drawing of mid-body parapodium with embedded aciculum and dorsal cirrus; lateral view. B. Drawing of mid-body neuropodium and ventral cirrus; ventral view.

Key to the species of Nautiliniellidae

- 1a. Prostomial appendages (palps or antennae) absent *Miura spinosa* Blake, 1993
- 1b. One or two pairs of prostomial appendages present 2
- 2a. Tentacular segment with only one pair of cirri 3
- 2b. Tentacular segment with one pair of dorsal and ventral cirri 7
- 3a. Tentacular segment with or without neurochaetae; all neuropodial hooks slender 4
- 3b. Tentacular segment without neurochaetae; some neuropodial hooks stout 5
- 4a. Neurochaetae ≥ 20 (up to 35) per parapodium; neurochaetae with inflated, subdistal stems and slightly curved, pointed distal ends *Mytilidiphila enseisensis* Miura & Hashimoto, 1993
- 4b. Neurochaetae ≤ 20 per parapodium; neurochaetae with rounded tips and slightly curved, distal ends *Mytilidiphila okinawaensis* Miura & Hashimoto, 1993
- 5a. Only one type of neurochaeta present: large, stout hooks 6
- 5b. Two types of neurochaetae present: One to two large, stout hooks and 15–20 small, mucronate tipped chaetae (in crow's of 2) *Laubierus mucronatus* Blake, 1993
- 5c. Three types of neurochaetae present: \leq five stout hooks with minute projection on cutting fang, 10–20 tridentate chaetae, and numerous minute, slender chaetae with single mucronate spin *Theyomytilidicola tridentatus* Miura & Hashimoto, 1996
- 6a. Maximum of one to two stout hooks per parapodium *Petrecca thyasira* Blake, 1990
- 6b. Maximum of seven to eight stout hooks per parapodium *Flascarpia alvinae* Blake, 1993
- 7a. One type of neurochaetae present ... 8
- 7b. Two types of neurochaetae present .. 9
- 8a. One large, stout hook per parapodium *Nautiliniella calyptogenicola* Miura & Laubier, 1989
- 8b. Maximum of four stout hooks per parapodium, and branchiae-like notopodial projections present *Thyasiridicola branchiatus* Miura & Hashimoto, 1996
- 8c. Number of anterior stout hooks variable (2–25) and notopodial branchiae-like projections absent 10




Fig. 4. *Vesicomycicola trifurcatus* new genus, new species. A. SEM of simple anterior neuropodial hooks. B. LM of anterior neuropodial hooks. C. SEM of mid-body chaetae with small apical tooth near tip, appearing slightly bifid. D, E. SEM of posterior trifurcate chaetae.

- 9a. Neurochaetae with two to three stout hooks and numerous bifurcate simple chaetae 11
- 9b. Neurochaetae with four to seven stout hooks per parapodium present anteriorly and five to seven trifurcate hooks posteriorly . . . *Vesicomycicola trifurcatus*, new genus and species
- 10a. Notopodia in middle regions especially elongate; middle to posterior neuropodia with a single hook with strongly curved distal fang . . . *Shinkai longipedata* Miura & Ohta, 1991
- 10b. Notopodia not elongate in any regions; middle to posterior neuropodia with a single hook, strongly curved on distal end with knob on tip *Shinkai sagamiensis* Miura & Laubier, 1990
- 10c. Notopodia in middle regions slightly

elongated; middle to posterior neuropodia with \geq five, slightly curved hooks *Shinkai semilonga*
Miura & Hashimoto, 1996

11a. Short, conical notopodia on middle segments *Natsushima bifurcata*
Miura & Laubier, 1990

11b. Elongate notopodia on middle segments *Natsushima graciliceps*
Miura & Hashimoto, 1996

Acknowledgments

We thank Captain Silva, the crew of R/V *Atlantis*, Expedition Leader Dudley Foster, the pilots and technicians of DSV *Alvin*, and members of the science party for their assistance at sea, and Karine Olu and Daniel Desbruyeres for loaning us nautiliniellid specimens. We are grateful to Joe Scott, Jewel Thomas and Megan Ward for help with illustration preparations and layout and Dr. Norman Fashing for use of his camera lucida. The manuscript benefited from reviews of Brigitte Hilbig, Stephen Gardiner and one anonymous reviewer. This research was supported by National Oceanic & Atmospheric Administration's National Undersea Research Program (University of North Carolina NC-Wilmington National Undersea Research Center) and Ocean Exploration Program. The Carol Woody Internship Program (College of William and Mary) and the Lerner Gray Memorial Fund of the American Museum of Natural History provided support to JD for collaboration with TM in Japan.

Literature Cited

Blake, J. A. 1990. A new genus and species of Polychaeta commensal with a deep-sea thyasirid clam.—Proceedings of the Biological Society of Washington 103:681–686.
———. 1993. New genera and species of deep-sea polychaetes of the Family Nautiliniellidae from the Gulf of Mexico and the Eastern Pacific.—Proceedings of the Biological Society of Washington 106:147–157.

Dall, W. H. 1886. Reports on the results of dredging, under the supervision of Alexander Agassiz, in the Gulf of Mexico, and in the Caribbean Sea, 1877–79, by the U.S. coast survey steamer Blake XXIV. Report on the Mollusca. Part 1. Brachiopoda and Pelecypoda.—Bulletin of the Museum of Comparative Zoology at Harvard University 12:171–318.
Fauchald, K. 1972. Benthic polychaetous annelids from deep water off western Mexico and adjacent areas in the eastern Pacific Ocean.—Allan Hancock Monographs in Marine Biology No. 7: 1–575.
Glasby, C. J. 1993. Family revision and cladistic analysis of the Nereidoidea (Polychaeta: Phyllozoidea).—Invertebrate Taxonomy 7:1551–1573.
Miura, T. 1998. Annelida Polychaeta (in part). Pp. 70–75 in D. Desbruyeres & M. Segonzac, eds. Handbook of deep-sea hydrothermal vent fauna. French Research Institute for the Exploitation of the Sea (IFREMER), France, pp. 279.
———, & J. Hashimoto. 1993. *Mytilidiphila*, a new genus of nautiliniellid polychaete living in the mantle cavity of deep-sea mytilid bivalves collected from the Okinawa Trough.—Zoological Science 10:169–174.
———, & ———. 1996. Nautiliniellid polychaetes living in the mantle cavity of bivalve molluscs from cold seeps and hydrothermal vents around Japan.—Publications from the Seto Marine Laboratory 37(316):257–274.
———, & L. Laubier. 1989. *Nautilina calyptogenicola*, a new genus and species of parasitic polychaete on a vesicomid bivalve from the Japan Trench, representative of a new Family Nautiliniidae.—Zoological Science 6:387–390.
———, & ———. 1990. Nautiliniellid polychaetes collected from the Hatsushima cold-seep Site in Sagami Bay, with descriptions of the new genera and species.—Zoological Science 7:319–325.
———, & S. Ohta. 1991. Two polychaete species from the deep-sea hydrothermal vent in the middle Okinawa Trough.—Zoological Science 8:83–387.
Olu, K., M. Sibuet, F. Harmegnies, J.-P. Foucher, & A. Fiala-Medioni. 1996. Spatial distribution of diverse cold-seep communities living on various diapiric structures of the southern Barbados prism.—Progress in Oceanography 38:347–376.
Van Dover, C. L. et al. 2003. Blake Ridge methane seeps: characterization of a soft-sediment, chemosynthetically based ecosystem.—Deep-Sea Research I 50:281–300.