



# Atypical, high-diversity assemblages of foraminifera in a mangrove estuary in northern Brazil

Nisan Sariaslan and Martin R. Langer

Institute für Geowissenschaften, Paläontologie, Universität Bonn, Bonn 53115, Germany

**Correspondence:** Martin R. Langer (martin.langer@uni-bonn.de)

Received: 1 March 2021 – Discussion started: 29 March 2021

Revised: 31 May 2021 – Accepted: 1 June 2021 – Published: 7 July 2021

**Abstract.** Mangrove forests are extremely productive ecosystems, are sources and sinks of organic carbon, and provide essential services both to the marine environment and people. We have studied the composition and species richness of modern benthic foraminiferal assemblages from mangrove swamps along the Mamanguape River in Paraíba, northern Brazil. Sampling points for foraminifera were selected to acquire information on the composition of foraminiferal assemblages from dense mangrove stands collected along a river transect. Almost 100 species of benthic foraminifera were identified within the shallow mangrove habitats. The large number of identified mangrove taxa is the highest recorded so far for true mangrove habitats. The high species richness rivals shallow-water assemblages recorded from nearby offshore and reef environments and indicates that a particularly large number of species is capable of growing and flourishing under multiple stressor conditions. Numerical analysis of the faunal assemblages shows that specific taxa, which were previously known to be uncommon in mangrove environments, are abundant in the Mamanguape River estuary. The atypical foraminiferal fauna found in the Mamanguape River estuary resembles shallow-water offshore assemblages, is characterized by high percent abundances of perforate and miliolid taxa, and contains only very few of the otherwise typical and numerically abundant agglutinated mangrove taxa. The unusual structure of the assemblages recorded provides insight into what combination of environmental variables controls their composition and novel perspectives to reconstruct past mangrove environments. Distribution, diversity, and species-specific analysis will provide guidance on the use of Brazilian mangrove foraminifera as indicators for the strength of tidal activity, pollution, and anoxia in coastal waters and sea-level reconstructions.

## 1 Introduction

Mangrove ecosystems around the world play an important role in protecting biodiversity, preserving shorelines, and regulating carbon cycling. They respond actively to coastal processes and sediment input and are considered one of the best geological indicators for the detection of modifications in coastal zone dynamics (Cunha-Lignon et al., 2009). Brazil is home to approximately 15 % of the world's total mangal forest areas but the release of effluents and untreated wastewater pose threats to ecosystems and marine biota. Nearly 6700 km (90 %) of Brazil's 7400 km of coastline hosts mangrove forests. The Mamanguape River estuary is the second largest estuary in the northeastern state of Paraíba (Brazil) covering a mangrove swamp area of more than 57 km<sup>2</sup> (Bezerra et al., 2012).

Mangrove ecosystems are subject to a suite of disturbances that vary in their intrinsic nature (e.g., geological, physical, chemical, and biological) in time and space. Inhabiting the interface between land and sea at low latitudes, these ecosystems occupy a harsh environment and are subject to daily tidal, temperature, and salinity variations as well as varying degrees of anoxia. Mangrove forests and their inhabitants are therefore rather robust and highly tolerant to life in their saline environments within warm, subtropical, and tropical seascapes (Alongi, 2008). In addition to the natural fluctuations, anthropogenic activities also affect these ecosystems. The estuaries of the northeastern ecoregion are among the most affected by the human occupation processes in Brazil and need actions that guide an integrated management to maintain ecosystem sustainability. Cities close to the Mamanguape River estuary have a total of nearly 40 000 inhabitants (Instituto Brasileiro de Geografia e Estatística, 2019) and extensive sugarcane and shrimp aquaculture fields place

environmental pressure on the mangrove ecosystem. The synergistic effects of multiple stressors may cause broad-scale changes in estuarine and coastal ecosystems impacting the abundance, species richness, and distribution of functionally important taxa. The interactions of multiple stressors are likely to increase as climate change and anthropogenic pressures will alter the delivery of freshwater and associated nutrients and pollutants to estuarine and coastal ecosystems (Scavia et al., 2002; Paerl et al., 2006; Gillanders et al., 2011; Schiedek et al., 2008). Understanding the response of benthic communities to key stressors is vital for managing mangrove environments and the first step towards setting ecologically relevant limits.

Foraminifera in tropical mangrove environments of South America and the islands nearby have been studied since the late 1940s in Trinidad (Cushman and Brönnimann, 1948a, b; Wilson et al., 2008), the Gulf of Paria (Todd and Brönnimann, 1957), Bahia (Zaninetti et al., 1979; Hiltermann et al., 1981; Eichler et al., 2015; Laut et al., 2016), Rio de Janeiro (Brönnimann et al., 1981; Debenay et al., 2001; Barbosa et al., 2005; Laut et al., 2016; Martins et al., 2016; Gasparini and Vilela, 2017; Belart et al., 2019), Colombia (Boltovskoy and Hincapié de Martínez, 1983), Puerto Rico (Culver, 1990), French Guiana (Debenay et al., 2002, 2004), Santa Catarina (Laut et al., 2016), Sao Paulo (Eichler et al., 2007; Passos et al., 2017; Eichler et al., 2019), and Rio Grande do Sul (Laut et al., 2016; Damasio et al., 2020; Semensatto et al., 2009). Most foraminifera assemblages studied from true mangrove settings were previously reported to be dominated by agglutinated species in a low-diversity ensemble (e.g., Boltovskoy, 1984; Culver, 1990; Debenay, 1990; Murray, 1991; Brönnimann et al., 1992 and references therein). However, we were persuaded to perform an in-depth investigation of our samples upon observing highly diverse and calcareous-dominated assemblages in our preliminary results. The objectives of the present study were (i) to provide detailed documentation on the structure and species richness of foraminiferal assemblages in mangroves of the Mamanguape River estuary of northern Brazil, (ii) to understand the driving forces contributing to the unusual composition and high diversity of these assemblages, and (iii) to discuss implications for interpreting the fossil record of foraminiferal mangrove assemblages.

## 2 Regional setting


The Mamanguape River estuary system is located on the coast of Paraíba State in northeastern Brazil and bound by latitudes 6°43'02" S to 6°51'54" S and longitudes 35°07'46" W to 34°54'04" W (Fig. 1). It is situated in an incised river valley, formed along a graben structure and oriented perpendicular to the coastline (Bezerra et al., 2001). The Mamanguape River estuary belongs to the Northeastern Marine Ecoregion (NEME; Spalding et al., 2007), covers an

area of ~ 658 km<sup>2</sup>, and is characterized by different rainfall regimes with varying rates of precipitation and duration during the wet and dry periods. As a result, a pronounced environmental variability among NEME estuaries is observed, where the highest reported pH (9.5) was recorded in the Mamanguape River (see Table 3.1 in Rafaela et al., 2018).


Radiocarbon dates collected from core sample material provide evidence that the inundation of the estuary occurred over the last 6000 years and more abruptly within the last 1000 years (Alvez, 2015). The dissolved oxygen levels in the estuary are controlled by a balance between the deoxygenation caused by the intense vegetation and tidal cycles and the oxygenation brought by bioturbating organisms. It is likely that hypoxia is a common condition of intertidal Mamanguape mangrove environments due to the mineralization of a large amount of organic matter produced by mangrove trees, which is responsible for a high consumption of oxygen by bacteria (Alongi et al., 2004) but also due to the exchange of porewater between sediments and the water column, known as “tidal pumping” (Li et al., 2009; Gleeson et al., 2013; Call et al., 2015; Leopold et al., 2017). According to Nordi et al. (2009), the tidal cycle is semi-diurnal on the northeastern coast of Brazil, producing two floods and two ebbs per day with a tidal range of approximately 2.8 m (Paludo and Klonowski, 1999). The estuary is heavily bioturbated by macrofaunal organisms such as oligochaetes, polychaetes, gastropods, bivalves, and shipworms. In addition, crabs and fish are diverse and abundant and are part of the vibrant ecosystem found in the estuary (Leonel et al., 2002; Nascimento et al., 2016; Van der Linden et al., 2017). Thus, sediments of the Mamanguape River estuary are heterogeneous in terms of their oxygen content; they rapidly become anoxic below the sediment surface, but bioturbation results in localized oxygenation of sediments around macrofaunal burrows (see also Langer et al., 1990).

## 3 Material and methods

Sediment samples were collected from the top 2 cm in September 2006 along the Mamanguape River estuary (Fig. 1). This time of the year falls within the dry season (August to November; Debenay et al., 2004) and the year 2006 is considered a “very dry” year and the driest between 2002 and 2006 (Santos et al., 2015). The sampling sites selected are located along the main and tributary mangrove channels and are scattered along a 2 km traverse upstream (Fig. 2). Samples were taken during a post-conference field trip at FORAMS 2006, the largest international symposium on foraminifera promoting the exchange of foraminiferal material under permission of the national science community. Sampling was conducted within dense mangrove stands and with distance from the channels (> 10 m) to avoid potentially allochthonous species transported by currents along the channel. Sampling points are located directly around the



**Figure 1.** Map of the Mamanguape River estuary with sample site locations (modified from Dolbeth et al., 2016).



**Figure 2.** Fisher  $\alpha$  diversity index graph for foraminiferal samples from the Mamanguape River estuary in comparison to total assemblages reported in previous mangrove studies (data from Brönnimann et al., 1981 (Guaratiba, Brazil); Rouvillois, 1982 (Casamance, Senegal); Lipps and Langer, 1999 (Jellyfish Lake, Palau); Debenay et al., 2002, 2004 (Kaw River, French Guiana); Langer and Lipps, 2003 (Madang, Papua New Guinea); Langer and Lipps, 2006 (Moorea, French Polynesia); Wilson et al., 2008 (Trinidad); Fajemila et al., 2015 (Moorea, French Polynesia); Langer et al., 2016 (Akanda, Gabon); Fajemila and Langer, 2017 (Sao Tome and Príncipe); Eichler, 2018 (Sao Paulo, Brazil); and Semensatto et al., 2009 (Trapandé Bay, Brazil)).

roots within the mangrove trees and the sediments collected are composed of organic-rich mud and silt.

The samples were washed over 63  $\mu$ m sieves and dried at room temperature; a total of  $\sim$ 1352 foraminifera specimens (death assemblage) were picked from these samples

(Table 1). To standardize the number of individuals recorded in individual samples and to avoid potential pitfalls (Patterson and Fishbein, 1989; Murray, 2006), picking was limited to a weight of 0.05 g of washed and dried sediment. All specimens were identified to species level, illustrated by scanning electron microscopy (SEM), and arranged into plates using Adobe Illustrator CS5 (Figs. 3–9). The total number of individuals was then counted for each species (Table 1) and their abundance and distribution patterns were analyzed.

As a measure of diversity, species richness was determined for each sample and is illustrated by the Fisher  $\alpha$  diversity index (Fig. 2, Table 2; Fisher et al., 1943; Murray, 1973). To this end, the total number of individuals has been plotted against the total number of species via the PAST software (Hammer et al., 2001) to compare the foraminiferal assemblage diversity and species richness with respect to the sampled locations. In addition, the Shannon diversity index ( $H$ ) was calculated to characterize species diversity in foraminiferal communities. The Shannon diversity index accounts for both abundance and evenness of the species present. The proportion of species  $i$  relative to the total number of species ( $p_i$ ) is calculated and then multiplied by the natural logarithm of this proportion ( $\ln p_i$ ). The resulting product is summed across species and multiplied by  $-1$ :

$$H = - \sum_{j=1}^s p_j \ln(p_j). \tag{1}$$

Species richness and Fisher  $\alpha$  diversity values were then compared to other mangrove studies from around the world to place our study in perspective. Lastly, the foraminiferal fauna was documented on plates (Figs. 3–9). All specimens are deposited in the micropaleontological collection at the Institute of Geosciences, University of Bonn (Germany).

**Table 1.** Alphabetical list, abundance of all species of foraminifera identified, and total abundance of specimens, genera, and species in the samples from the Mamanguape River estuary.

Species	Sample 1	Sample 2	Sample 3	Sample 4
<i>Abditodentrix rhomboidalis</i>	2	2	2	4
<i>Adelosina milletti</i>	3	4	2	3
<i>Ammonia advena</i>	2	3	3	1
<i>Ammonia veneta</i>	23	11	11	20
<i>Amphistegina lessonii</i>	1	1		
<i>Amphistegina radiata</i>	1			
<i>Angulogerina</i> cf. <i>A. occidentalis</i>	2	7	5	7
<i>Anomalinulla glabrata</i>	1	2		
<i>Anomalinulla</i> sp. 1	25	17	18	17
<i>Arenoparella mexicana</i>				1
<i>Articulina alticostata</i>				1
<i>Astrononion gallowayi</i>	2	1		
<i>Bigenerina</i> sp. 1	1	2	2	1
<i>Bolivina brevior</i>	2		1	
<i>Bolivina densipunctata</i>	4	3		
<i>Bolivina ordinaria</i>	9	9	2	2
<i>Bolivina striatula</i>	4	5	1	7
<i>Bolivina</i> sp. 1		2		
<i>Bolivina variabilis</i>	1	1		7
<i>Bolivina</i> cf. <i>B. variabilis</i>	6	6	13	10
<i>Buliminella elegantissima</i>	1		1	
<i>Cassidelina</i> sp. 1	2	2	1	2
<i>Cornuspira involvens</i>	5	5	3	1
<i>Cornuspira planorbis</i>			1	1
<i>Criboelphidium mirum</i>	2	6	12	5
<i>Cushmanina bricei</i>			1	2
<i>Discorbis williamsoni</i>	1			1
<i>Discorbitina pustulata</i>		1	1	2
<i>Edentostomina</i> sp. 1	2	3	2	3
<i>Elphidium sagrum</i>		2	3	3
<i>Elongobula parallela</i>	4	10	9	7
<i>Eoponidella pulchella</i>	4	1		
<i>Fischerina</i> sp. 1	1			
<i>Fissurina</i> sp. 2	2	1	2	1
<i>Fissurina bispinata</i>			1	
<i>Fissurina colomboensis</i>	1	2	2	3
<i>Fissurina semimarginata</i>		2	1	2
<i>Fissurina</i> ? sp. 1	1			
<i>Glabratella carinata</i>	2	4	2	3
<i>Glabratella mirabilis</i>	2	1	1	2
<i>Globocassidulina crassa</i>	2	2	5	4
<i>Globocassidulina rossensis</i>	2	4	3	
<i>Hauerina atlantica</i>				1
<i>Inaequalina</i> sp. 1				1
<i>Laevipeneroplis bradyi</i>				1
<i>Lagena tenuis</i>				1
<i>Loxostomina costulata</i>	5	1	1	
<i>Miliolinella webbiana</i>		1	2	3
<i>Miliolinella</i> sp. 1	1		1	1
<i>Mychostomina revertens</i>				1
<i>Neoconorbina radiatogranulata</i>		1	1	2
<i>Neoconorbina</i> sp. 1	3	3	4	4
<i>Neoconorbina terquemi</i>		5	1	1
<i>Nonionoides grateloupii</i>	8	5	2	4

Table 1. Continued.


Species	Sample 1	Sample 2	Sample 3	Sample 4
<i>Orbitina carinata</i>	12	25	30	22
<i>Pararotalia cananeaensis</i>	81	76	75	71
<i>Procerolagena oceanica</i>				1
<i>Pseudolachlanella eburnea</i>	1	2	2	3
<i>Pseudolachlanella bermudezi</i>	9	6	6	12
<i>Pseudotriloculina</i> sp. 1	4	2	3	2
<i>Pseudotriloculina</i> sp. 2	2	2	1	2
<i>Pseudotriloculina</i> sp. 3	2	1	2	3
<i>Quinqueloculina</i> cf. <i>Q. boschiana</i>	5	5	2	3
<i>Quinqueloculina</i> cf. <i>Q. carinatastriata</i>	3	3	3	3
<i>Quinqueloculina cuvierina</i>	10	8	16	15
<i>Quinqueloculina moynensis</i>	19	22	15	17
<i>Quinqueloculina poeyana</i>	1			
<i>Quinqueloculina</i> cf. <i>Q. rebecca</i>	1	1		
<i>Quinqueloculina samoensis</i>	3	5	4	4
<i>Quinqueloculina tantabiddyensis</i>	1	1		
<i>Quinqueloculina</i> sp. 1	2		1	2
<i>Quinqueloculina quinquecarinata</i>	1		1	1
<i>Quinqueloculina</i> sp. 2			3	2
<i>Quinqueloculina</i> sp. 3	2	2	3	3
<i>Quinqueloculina</i> sp. 4			1	
<i>Quinqueloculina</i> cf. <i>Q. compta</i>			1	2
<i>Quinqueloculina</i> cf. <i>Q. zhengi</i>	1			
<i>Rosalina</i> sp. 1		2	2	1
<i>Rosalina</i> sp. 2	15	14	8	11
<i>Rosalina bradyi</i>		7		1
<i>Rotaliammina trumbulli</i>	2		1	
<i>Rotorbis auberii</i>	4	5		3
<i>Sagrina pulchella</i>		1	3	3
<i>Sigmoilinita costata</i>	2		2	1
<i>Sigmavirgulina tortuosa</i>	1	3	6	2
<i>Svratkina acuta</i>	2	1	1	2
<i>Siphonina reticulata</i>	4	3	4	5
<i>Spirillina grosseperforata</i>			1	1
<i>Textularia</i> sp. 1		1	2	1
<i>Textularia</i> sp. 2		2	1	1
<i>Textularia</i> cf. <i>T. semialata</i>		1		
<i>Trochammina inflata</i>	1		1	
<i>Wiesnerella auriculata</i>	4	5	5	
Total number of specimens	335	346	328	343
Number of genera	40	38	38	44
Number of species	65	66	66	70

## 4 Results

### 4.1 Composition of foraminiferal assemblages

A total of 1352 benthic foraminifera specimens belonging to 93 species and 53 genera, including hyaline-perforate, porcelaneous, and agglutinated taxa, were recovered from the samples collected in the Mamanguape estuary (Fig. 1 and Table 1). The foraminiferal tests obtained were found to be particularly well preserved, with even the most fragile forms

in good condition, lacking signs of dissolution, abrasion, or breakage. Many specimens were found to contain pyrite framboids inside their tests, a feature considered to be indicative of anoxic conditions in the surrounding sediment (Fig. 9, figs. 22–23). Hyaline-perforate foraminifera comprise 34 genera (64.2% of the total assemblage) and 53 species (57% of the total assemblage), porcelaneous types account for 14 genera (26.4%) and 33 species (35%), and agglutinated taxa are represented by 5 genera (9.4%) and 7 species (8%). Percent abundances of wall structural types (agglu-




**Figure 3.** (1–2) *Trochammina inflata* (Montagu); (3–4) *Rotaliammina trumbulli* Seiglie; (5–6) *Arenoparrella mexicana* (Kornfeld); (7) *Bigenerina* sp. 1; (8–9) *Textularia* sp. 1; (10) *Textularia* cf. *T. semialata* Cushman; (11) *Textularia* sp. 2; (12) *Mychostomina revertens* (Rhum- bler); (13–14) *Spirillina grosseperforata* Zheng; (15–16) *Cornuspira involvens* Reuss; (17) *Cornuspira planorbis* Schultz; (18–19) *Fische- rina* sp. 1; (20–21) *Wiesnerella auriculata* (Egger); (22–23) *Edentostomina* sp. 1; (24) *Adelosina milletti* (Wiesner); (25) *Articulina alticostata* Cushman; (26–27) *Pseudotriloculina* sp. 1; (28–29) *Pseudotriloculina* sp. 2; (30–33) *Pseudotriloculina* sp. 3. The scale bar is 100 and 50  $\mu\text{m}$  for figs. (8)–(14), (17)–(19), (21)–(24), and (26)–(29).

tinated, hyaline-perforate, and porcelaneous) do not show any significant variation among the sampling sites. Hyaline-perforate foraminifera range between 72 % and 75 %, porcelaneous foraminifera constitute 23 % to 27 %, and agglutinated species contribute only 1 % or 2 % to the total assemblage at each site (Table 2). The total number of benthic foraminifera species increases slightly from sample site 1 near the mouth of the estuary (65 species) towards the innermost sample site 4, where 70 species were recorded.

#### 4.2 Species richness and diversity

Species richness values vary between 65–70 among the sample sites (Table 1). The highest value (70 species) was recorded at site 4 and the lowest at site 1 (65 species). Fisher  $\alpha$  values range between 24.04–26.60 (Table 2 and Fig. 3), with the highest values at site 4 and the lowest at site 1. As a general trend, species richness and Fisher  $\alpha$  values were found to increase from the sampling sites closer to the ocean




**Figure 4.** (1–3) *Quinqueloculina* cf. *Q. compta* Cushman; (4–6) *Quinqueloculina cuvierina* (d’Orbigny); (7) *Quinqueloculina* cf. *Q. zhengi* Parker; (8–10) *Quinqueloculina poeyana* d’Orbigny; (11–12) *Pseudolachlanella eburnea* (d’Orbigny); (13–14) *Pseudolachlanella bermudezi* (Acosta); (15–16) *Quinqueloculina moynensis* Collins; (17) *Quinqueloculina quinquecarinata* Collins; (18–21) *Quinqueloculina* cf. *Q. rebecca* Vella; (22–25) *Sigmolinita costata* Schlumberger; (26–27) *Quinqueloculina samoensis* Cushman; (28) *Quinqueloculina tantabid-dyensis* Parker; (29) *Quinqueloculina* sp. 2; (30–32) *Quinqueloculina* sp. 1; (33–35) *Quinqueloculina* sp. 4. The scale bar is 100 and 50  $\mu\text{m}$  for figs. (11)–(12), (16), (21), (24), (26), (29), (30)–(31), and (33).

towards the inner parts of the estuary with the highest values at site 4 (Table 1 and Fig. 2). Dominance, evenness and the Shannon diversity index values recorded revealed only minor variability and were found to be comparatively uniform across all sample sites analyzed (Table 2).

#### 4.3 Distribution patterns

Samples were found to have a comparatively uniform composition without much difference in species richness, species

diversity, or community structure. The assemblages analyzed from each sampling site show a striking dominance of *Pararotalia cananeaensis* (20.7%–24.2% of the total assemblages). The second most abundant taxon is *Orbitina carinata*, a hyaline-perforate species with the highest percent abundances distant from the mouth of the estuary. *Ammonia veneta* and *Anomalina* sp. 1 are also abundant in all our samples, constituting 3.2%–7.5% and 4.9%–7.5% of the total assemblages, respectively. The porcela-




**Figure 5.** (1–3) *Quinqueloculina* cf. *Q. carinatastriata* Wiesner; (4–6) *Quinqueloculina* sp. 3; (7) *Quinqueloculina* cf. *Q. bosciana* d’Orbigny; (8–9) *Miliolinella webbiana* d’Orbigny; (10) *Miliolinella* sp. 1; (11) *Hauerina atlantica* Cushman; (12–13) *Inaequalina* sp. 1; (14) *Laevipeneroplis bradyi* Cushman; (15) *Procerolagena oceanica* (Albani); (16–17) *Cushmanina bricei* McCulloch; (18–19) *Fissurina colomboensis* McCulloch; (20) *Fissurina bispinata* Ujjié; (21) *Fissurina* sp. 2; (22–23) *Fissurina semimarginata* Reuss; (24–25) *Fissurina?* sp. 1; (26–28) *Bolivina densipunctata* Sellier de Civrieux; (29–32) *Bolivina ordinaria* Phleger and Parker. The scale bar is 100 and 50  $\mu\text{m}$  for figs. (2), (6)–(7), (17), (20), (22)–(25), (26), and (28)–(32).

neous species *Quinqueloculina moynensis* constitutes between 4.57%–6.36% of the total assemblage at individual sites. Species of the genus *Quinqueloculina* are represented in the assemblages by a highly diverse group and constitute 15 species. Among the 93 species recorded, a total of 40 species were present at all four sampling sites including the five most abundant species (*P. cananeaensis*, *A. veneta*, *Anomalinulla* sp. 1, *O. carinata*, and *Q. moynensis*).

Occurrence records of a few species display preferences for particular sites in the estuary. Among these, *Angulogerrina* cf. *A. occidentalis*, *Criboelphidium mirum*, and *Elongobula parallela* are particularly abundant at site 2, 3, and 4, which correspond to the inner reaches of the estuary. Species recorded exclusively at sites located near the mouth region of the estuary (sites 1 and 2) include *Anomalinulla glabrata*, *Bolivina densipunctata*, *Bolivina variabilis*, *Eoponidella pulchella*, *Fischerina* sp. 1, *Fissurina?* sp. 1, *Quin-*






**Figure 6.** (1) *Bolivina ordinaria* Phleger and Parker; (2–3) *Bolivina striatula* Cushman; (4–5) *Bolivina variabilis* Williamson; (6–8) *Bolivina* cf. *B. variabilis* (Williamson); (9) *Bolivina* sp. 1; (10–11) *Abditodentrix rhomboidalis* (Millett); (12–13) *Bolivina brevior* Cushman; (14–15) *Globocassidulina rossensis* Kennett; (16–17) *Globocassidulina crassa* (d’Orbigny); (18) *Lagena tenuis* Bornemann; (19) *Loxostomina costulata* (Cushman); (20–21) *Sagrina pulchella* d’Orbigny; (22–23) *Cassidelina* sp. 1; (24) *Buliminella elegantissima* (d’Orbigny); (25–26) *Angulogerina* cf. *A. occidentalis* (Cushman); (27–28) *Sigmavirgulina tortuosa* (Brady); (29–30) *Siphonina reticulata* (Czjzek); (31–33) *Discorbitina pustulata* (Heron-Allen and Earland). The scale bar is 100 and 50  $\mu\text{m}$  for figs. (1), (4)–(5), (12)–(17), (21)–(23), (25)–(26), and (28)–(33).

*queloculina poeyana*, *Q.* cf. *Q. rebecca*, and *Q. tantabid-yensis*. Amphisteginid foraminifera, although present in low abundances, are also restricted to sample site 1 and 2.

## 5 Discussion

Analyses of foraminiferal assemblages from the Maman-guape mangrove estuary revealed highly diverse, particularly species-rich, and structurally complex biota of benthic taxa.

With a total of 93 taxa, the species richness exceeds previous species counts and was found to be the highest among all true mangrove environments studied so far (see below and Fig. 2). In addition, our samples yielded unusual assemblages heavily dominated by calcareous taxa, lacking agglutinated specimens to a great degree, as reflected by the dominance of hyaline-perforate taxa that contribute between 72.3%–75.1% to the total population. Porcelaneous miliolids, representatives of shallow marine environments, constitute the




**Figure 7.** (1–3) *Orbitina carinata* Sellier de Civrieux; (4–6) *Rotorbis auberii* (d’Orbigny); (7–9) *Neoconorbina* sp. 1; (10–11) *Neoconorbina radiatogranulata* Parker; (12) *Neoconorbina terquemi* (Rzehak); (13–15) *Rosalina bradyi* (Cushman); (16) *Rosalina* sp. 1; (17–19) *Rosalina* sp. 2; (20–23) *Glabratella carinata* Seiglie and Bermúdez; (24–27) *Glabratella mirabilis* Seiglie and Bermúdez; (28–29) *Glabratella carinata* Seiglie and Bermúdez. The scale bar is 100 and 50  $\mu\text{m}$  for figs. (3)–(5), (7), (10)–(12), (14)–(15), (17), and (20)–(29).

second most abundant group with an abundance of 23.8%–26.5%. Agglutinated species, which are typically dominant in mangrove environments (Murray, 1991), contribute only 1.7%–2.1% to the total assemblages (Table 2). Species richness values recorded range between 65 and 70 at individual sites, whereas the average species richness was found to be generally much lower in other studies (Fig. 2; Debénay, 1990, 2001; Murray, 2006 and references therein). The atypical composition and highly diverse Mamanguape assemblages raise the question of which driving forces are decisive for the development of such unusual mangrove biota.

Because such assemblages rather resemble foraminiferal faunas from shallow coastal settings and are difficult to relate to mangrove habitats, the potential implications for the interpretation of the fossil record are considered.


As outlined above and illustrated in the Fisher  $\alpha$  diagram (Fig. 2), the Mamanguape foraminiferal assemblages were found to deviate from usual foraminiferal mangrove biota in species richness, in the composition of wall-structural types, the presence of abundant hyaline-perforate and porcelainous miliolid taxa, and in particularly low abundances of agglutinated species. While Fisher  $\alpha$  value recordings



**Figure 8.** (1) *Glabrattella mirabilis* Seiglie and Bermúdez; (2–4) *Elongobula parallela* (Cushman and Parker); (5–6) *Eoponidella pulchella* (Parker); (7–8) *Amphistegina lessonii* (d’Orbigny); (9) *Amphistegina radiata* (Fichtel and Moll); (10–12) *Nonionides grateloupii* (d’Orbigny); (13–14) *Syratkina acuta* (Sidebottom); (15) *Anomalinulla glabrata* (Cushman); (16–20) *Anomalinulla* sp. 1.; (21–30) *Pararotalia cananeaensis* Debenay, Duleba, Bonetti De Melo e Souza & Eichler. The scale bar is 100 and 50  $\mu\text{m}$  for figs. (1), (3)–(4), (6)–(9), (13)–(14), (18)–(20), (24), and (27)–(30).

from previous mangrove studies ranged between 0.3 and 11, the Mamanguape faunas ranged between 24.4 and 26.6 (Fig. 2). The higher numbers are mainly due to the presence of both porcelaneous-miliolid and hyaline-perforate taxa and are more similar to values commonly encountered in coastal nearshore environments at the seaward end of river estuaries or in tropical lagoon or back-reef settings where foraminiferal diversity is much higher than in mangroves (Saunders, 1958; Halicz et al., 1984; Debenay et al., 2001; Langer and Lipps, 2003; Thissen and Langer, 2017; Langer et al., 2013; Fajemila et al., 2020a).

The composition of our benthic foraminifera assemblages also contrasts with the ones found in previous studies on total assemblages of mangrove foraminifera from the southern Atlantic, where benthic foraminifera assemblages are exclusively dominated by agglutinated and hyaline-perforate species and mostly lack porcelaneous taxa (Fig. 2; Murray, 1991, 2006). Agglutinated foraminifera within the Mamanguape mangroves contribute a mere 1% to 2% to the total assemblages, while other studies showed abundances that generally range above 40% (Culver, 1990; Brönniman et al., 1992; Debenay and Guillou, 2002; Murray, 2006).



**Figure 9.** (1–3) *Ammonia advena* Cushman; (4–10) *Ammonia veneta* (Schultze); (11–14) *Criboelphidium mirum* Langer and Schmidt-Sinns; (15–16) *Elphidium sagrum* (d’Orbigny); (17–19) *Discorbis willamsoni* Chapman and Parr; (20–21) *Astrononion gallowayi* Loeblich and Tappan; (22–23) Pyrite framboids found in our specimens, (24) *Globigerinoides* sp. 1; (25–26) *Globigerinoides* sp. 2. The scale bar is 100 and 50  $\mu\text{m}$  for figs. (6), (13), (15)–(16), and (18)–(20), 25  $\mu\text{m}$  for figs. (21) and (26), and 10  $\mu\text{m}$  for fig. (23).

Deviations from “typical” mangrove assemblages are also marked by the dominance and numerical abundance of *Pararotalia cananeiaensis* (Debenay et al., 2001), a taxon of a nearshore, shallow-water genus that is commonly not expected in large numbers in mangrove habitats (Geslin et al., 2002; Debenay et al., 2001). Within the Mamanguape mangroves, percent abundances of *P. cananeiaensis* were found to range between 20.1 % and 24.2 %, making it the most abundant taxon of all species. Similar high abundances were recently reported by Damasio et al. (2020) from the Santos estuary (southeast Brazil), where the species was reported

from the bay and along the main channel but absent from the low-saline northernmost mangrove habitats. Eichler et al. (1995) reported the species from the Bay of Trapandé (Brazil) and attributed its inshore occurrence to the marine influence. Similarly, Murray et al. (1982) attributed the upstream presence of small coastal species to be indicative of an upstream transport of sediment of marine origin. Debenay et al. (2001) studied in detail the occurrence and transportation of *P. cananeiaensis* along Brazilian coasts under variable hydrodynamic conditions. They associated the presence of *P. cananeiaensis* in estuary channels with the intensity of

**Table 2.** Sample site information including geographical position, salinity (Medeiros et al., 2018), number of species, Fisher  $\alpha$ , Shannon's  $H$ , dominance (SID), evenness ( $e^H/S$ ), and percent abundances of the agglutinated, miliolid, and perforate specimens as well as the five most abundant species.

	Sample 1	Sample 2	Sample 3	Sample 4
Longitude	6°46'59.61" S	6°46'57.84" S	6°47'57.89" S	6°46'55.93" S
Latitude	34°56'40.83" W	34°57'54.95" W	34°57'49.94" W	34°58'55.04" W
Salinity ‰	32 ± 3.9	30.3 ± 3.9	28.3 ± 3.0	28.7 ± 3.1
Fisher $\alpha$	24.04	24.19	24.89	26.6
Shannon H	3.324	3.439	3.345	3.484
Dominance D	0.08149	0.0696	0.07689	0.06473
Evenness ( $e^H/S$ )	0.4271	0.4721	0.4297	0.4656
Agglutinated %	1.19	1.73	2.13	1.17
Porcelaneous %	24.78	23.12	23.78	26.53
Hyaline-perforate %	74.02	75.14	73.17	72.30
<i>Ammonia veneta</i> %	7.46	3.18	3.35	5.83
<i>Anomalinulla</i> sp. 1 %	7.46	4.91	5.49	4.96
<i>Orbitina carinata</i>	3.58	7.22	13.16	6.41
<i>Pararotalia cananeaensis</i> %	24.18	21.97	22.87	20.70
<i>Quinqueloculina moynensis</i> %	5.67	6.36	4.57	4.96

marine intrusion into paralic systems and considered the upstream presence more likely to be a function of transport (allochthonous) than of in situ growth (see also Burone et al., 2006).

Seven different lines of evidence, however, strongly argue for an in situ deposition of assemblages and autochthonous presence of *P. cananeaensis*, at least within the Mamanguape estuary. (i) Numerical abundances of *P. cananeaensis*, Shannon's  $H$ , evenness, Fisher  $\alpha$ , and dominance index values do not vary substantially along the sampled transect and are largely uniform across the sample sites. Evidence for decreasing percent abundances upstream or features of successive filtering through the dense tributary mangrove channels is thus lacking. (ii) The Mamanguape River estuary is partially dammed by a natural sand barrier (Fig. 1) and acts as a hindrance in terms of water exchange and transport of coastal species deep into the estuary. (iii) Except for two individuals, planktic foraminifera are absent, thus arguing against a large-scale transport of open-ocean taxa upstream. (iv) The preservation of the taxa recovered ranges from good to excellent and includes both fragile, robust, smaller, and larger species. Features of abrasion, transport, or rolling over large distances were not observed. (v) The sample sites investigated all revealed a particularly high species richness of benthic foraminifera. (vi) The lack of both peneroplid and soritid foraminifera, two groups that are widely present in shallow-water nearshore biofacies in the region (Weinmann, 2009; Disaró et al., 2014), argues against a transport of species from nearshore to estuarine mangrove habitats. (vii) Pyrite framboids were frequently observed within the chambers in the majority of species, reflecting the presence of anoxic conditions at the site of deposition (see Fig. 9, figs. 22–23). Pyrite formation is tightly intertwined with the presence of organic

matter (Berner, 1970, 1984; Thiel et al., 2019), suggesting that (a) foraminifera cell cytoplasm was present during deposition; (b) autolytic, bacterial, or prey-mediated protoplasm decay in foraminifera has been largely prevented (Murray and Bowser, 2000); and (c) foraminiferal assemblages are largely autochthonous. The time required for pyrite framboids to form the largest crystals, like the ones found in our assemblages, was previously estimated to be around 35 d (Rickard, 2019).

If a large-scale upstream transport of coastal species would indeed have taken place, a successive filtering and numerical reduction effect of coastal species through the dense mangrove stands would be expected. However, this has not been recorded in the assemblages. Instead, the number of benthic species was found to increase towards the inner parts of the estuary and the assemblages are therefore considered to be largely autochthonous.

Marine conditions supporting such diverse assemblages are manifested in high salinity values in our sampling area during both dry and wet seasons and are interpreted as the result of a longer residence time of marine waters in the estuary (Fig. 2; Dolbeth et al., 2016; Medeiros et al., 2018). Prevailing marine conditions within the dense Mamanguape mangrove estuary were also considered a major factor contributing to fish diversity in previous studies (MacDonald et al., 2008; Blaber, 2013).

The results obtained revealed unprecedentedly high species richness values and assemblages dominated by calcareous species instead of agglutinated forms and represent an atypical example for a mangrove benthic foraminiferal fauna when compared to previous studies. While these findings may seem puzzling at first sight, they are considered to mainly result from high salinity values in the upstream

areas of the Mamanguape River estuary (Table 2). The resemblance to shallow-water nearshore foraminiferal communities is both indicated by the presence of a large number of porcelaneous miliolid and hyaline-perforate species and in the abundance of *P. cananeiaensis*. In addition, it should be noted that the low number of agglutinated specimens and species found in our assemblages is not an artifact of preservation, as samples were carefully treated and the material was picked immediately after transportation to the laboratory.

Salinity has long been recognized as a major driving force governing the species richness, composition of wall structural types, and diversity of foraminiferal assemblages (Murray, 1991; Debenay, 1990). A distinct separation between agglutinated and calcareous taxa along a pronounced salinity gradient has recently been reported from Lagos Lagoon, where mangrove and low salinity environments are dominated by agglutinated foraminifera (Fajemila et al., 2020b). While the majority of mangrove assemblages from along the Atlantic coast of South America were reported to be dominated by agglutinated taxa, the Mamanguape mangrove estuary was found to contain abundant calcareous wall types and to be almost devoid of agglutinated taxa. Among the environmental framework conditions promoting such atypical mangrove assemblages of foraminifera, salinity stands out as a prominent control factor. Salinity recordings along the sampling traverse revealed values ranging between  $\sim 25\%$  and  $35\%$ , indicating mostly euhaline to brackish water conditions. The prevalence of marine conditions within the studied area of the Mamanguape estuary is supported by comparatively long residence times of marine waters, high tidal amplitudes, and semi-diurnal tidal flushing of the estuarine environment. The resemblance of the atypical mangrove assemblages to shallow-water nearshore biota may thus be attributed to the intensity of marine conditions, where calcareous foraminifera dominate over agglutinated taxa (Boltovskoy, 1954; Todd and Brönnimann, 1957; Boltovskoy and Hincapié de Martínez, 1983; Scott et al., 1990; Debenay, 2001; Debenay et al., 2002, 2004; Moreno et al., 2005; Eichler et al., 2007; Camacho et al., 2015; Shaw et al., 2016; Passos et al., 2017). The penetration of calcareous species into the Mamanguape estuary has also been reported from the neighboring Paraíba estuary (Debenay et al., 2002). However, unlike the Mamanguape estuary, the Paraíba estuary receives freshwater discharge from more rivers and invasions of marine foraminifera only occur during the dry season and completely disappear during the rainy season (Debenay et al., 2002).

Independent of whether the foraminiferal assemblages are allochthonous or autochthonous, the atypical Mamanguape mangrove faunas raise questions concerning implications for the interpretations of the fossil record. Unlike previous studies on foraminifera from South American and many other mangrove environments around the world, where specialized agglutinated taxa dominated mangrove environments, excep-

tionally species-rich and diverse assemblages of foraminifera prevail within the dense mangroves at Mamanguape. Both Debenay et al. (2001, 2002) and Woodroffe et al. (2005) demonstrated that a range of environmental factors jointly govern both the composition, distribution, and preservation of foraminiferal biota in mangrove environments. Besides salinity, these include the elevation, grain size, organic content, and taphonomic processes related to low pH excursions resulting in the removal of agglutinated and calcareous taxa. Post-mortem disaggregation and taphonomic loss may thus bias the fossil record and constitute a serious constraint regarding paleoenvironmental reconstructions.

Mangrove ecosystems have existed at least since the Late Cretaceous and fossil evidence used for paleoecological reconstructions is mostly based on organic remains (fruit; flowers; wood or leaves; or microfossils, particularly pollen; Ellison et al., 1999) and sedimentological features (Augustinus, 1995). Sedimentation of typical fine-grained mangrove deposits results from the reduction in current velocity where tree trunks, prop roots and pneumatophores exercise a filter function and result in typical mangrove mud sediments. Within the Mamanguape estuary, the accumulation and vertical accretion of mangrove mud is a mixture of river-born clastic sediments, organic material produced by mangrove trees and associated fauna, and suspension-rich ocean floating debris penetrating mangrove swamps. In high precipitation areas and under low salinity conditions, the production of organic material is generally promoted but reduces pH conditions within the sediment and limits carbonate availability and thus reproduction of calcareous foraminifera. On the other hand, agglutinated tests are known to resist low pH conditions and dissolution while they are removed through oxidation–reduction reactions (Ellison and Nichols, 1976; Scott and Medioli, 1978; Boltovskoy, 1984; Goldstein, 1988 and the references therein; Thomas and Varekamp, 1991; Debenay et al., 2002, 2004). However, in the absence of considerable rainfall and during the warm periods, penetration of coastal waters into the estuarine system occurs. Based on the intensity of the marine influence and tides, calcareous foraminifera start to dominate and get preserved in the otherwise brackish or freshwater realm (Boltovskoy, 1954; Todd and Brönnimann, 1957; Scott et al., 1990; Debenay, 2001; Debenay et al., 2002, 2004; Moreno et al., 2005; Eichler et al., 2007; Camacho et al., 2015; Shaw et al., 2016; Passos et al., 2017; Woodroffe et al., 2005). Normally, salinity tends to be higher in the downstream areas in estuaries, yet the upstream areas of the Mamanguape are characterized by high salinity (Table 2), indicative of a consistent influence of oceanic waters penetrating even further upstream of the sampling sites.

## 6 Conclusions

Mangrove environments of the Mamanguape River estuary are home to diverse assemblages of benthic foraminifera including 33 porcelaneous, 53 hyaline-perforate, and 7 agglutinated species and represent an extraordinary ecosystem characterized by atypical, highly diverse, and species-rich benthic foraminiferal biota. The atypical population structure features prominent components of hyaline-perforate and miliolid-porcelaneous benthic foraminifera and are unprecedented in previous mangrove studies. The assemblages resemble coastal nearshore biota that are traditionally not classified as mangrove foraminiferal faunas and are here considered to be the result of highly saline ocean waters consistently penetrating deep into the estuary and promoting the presence of compositionally diverse and species-rich biota. When preserved in the fossil record, such assemblages are not readily related to mangrove ecosystems and would most likely be interpreted as a coastal, nearshore, shallow-water environment. Our findings thus have implications for inferring environmental conditions of past mangrove ecosystems. Salinity was found to be the main controlling factor structuring benthic foraminiferal assemblages, where both the comparatively long residence time of marine waters and the low freshwater inflow support prevailing euhaline to brackish water conditions and the presence of calcareous-dominated, species-rich foraminiferal assemblages.

*Code availability.* The PAST software of Hammer et al. (2001) is available at: <https://www.nhm.uio.no/english/research/infrastructure/past/>.

*Data availability.* All data used in the present study are provided in Tables 1 and 2.

*Author contributions.* MRL conceptualized the study and collected the material. NS and MRL analyzed the data and prepared and wrote the manuscript.

*Competing interests.* The authors declare that they have no conflict of interest.

*Disclaimer.* Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

*Acknowledgements.* We are grateful to the reviewers and the editor for their constructive comments and suggestions. We also thank Stephanie Pietsch for assistance with the collection of the material and Georg Oleschinski for the SEM pictures of the framboidal

pyrite. The analysis and study of the material was supported by a grant from the German Federal Foreign Office and the University of Bonn.

*Financial support.* This open-access publication was funded by the University of Bonn.

*Review statement.* This paper was edited by Hiroshi Kitazato and reviewed by Johann Hohenegger and one anonymous referee.

## References

- Alongi, D.: Mangrove forests: Resilience, protection from tsunamis, and responses to global climate change, *Estuar. Coast. Shelf S.*, 76, 1–13, <https://doi.org/10.1016/j.ecss.2007.08.024>, 2008.
- Alongi, D., Wattayakorn, G., Boyle, S., Tirendi, F., Payn, C., and Dixon, P.: Influence of roots and climate on mineral and trace element storage and flux in tropical mangrove soils, *Biogeochemistry*, 69, 105–123, <https://doi.org/10.1023/B:BIOG.0000031043.06245.af>, 2004.
- Alves, F. C.: Análise morfoestrutural das bacias dos rios e Mamanguape (PB) com base em MDE-SRTM e imagem PALSAR/ALOS-1, MSc Diss., Inst. Nac. Pes. Esp., INPE, 5 151 pp., available at: <http://mtc-m16d.sid.inpe.br/col/sid.inpe.br/mtc-m19/2015/02.26.13.32/doc/publicacao.pdf> (last access: 29 June 2021), 2015.
- Augustinus, P. G. E. F.: Geomorphology and Sedimentology of Mangroves: *Geom. Sediment, Est.*, 53, 333–357, 1995.
- Barbosa, C. F., Scott D. B., Seoane J. C. S., and Turcq, B. J.: Foraminiferal zonation as base lines for Quaternary sea-level fluctuations in south-southeast Brazilian mangroves and marshes, *J. Foramin. Res.*, 35, 22–43, 2005.
- Belart, P., Renan, H., Raposo, D., Martins, V., Frontalini, F., Figueiredo, M., Lorini, M. L., and Laut, L.: Seasonal Dynamics of Benthic Foraminiferal Biocoenosis in the Tropical Saquarema Lagoonal System (Brazil), *Estuar. Coast.*, 42, 1–20, 2019.
- Berner, R. A.: Sedimentary pyrite formation, *Am. J. Sci.*, 268, 1–23, 1970.
- Berner, R. A.: Sedimentary pyrite formation – An update, *Geochim. Cosmochim. Ac.*, 48, 605–615, 1984.
- Bezerra, D., Nascimento, D., Ferreira, E., Rocha, P., and Mourão, J.: Influence of tides and winds on fishing techniques and strategies in the Mamanguape River Estuary, Paraíba State, NE Brazil, *Anais da Academia Brasileira de Ciências*, 84, 775–88, [10.1590/S0001-37652012005000046](https://doi.org/10.1590/S0001-37652012005000046), 2012.
- Bezerra, F. H., Amaro, V. E., Vita-Finze, C., and Saady, A.: Pliocene-Quaternary fault control of sedimentation and coastal plain morphology in NE Brazil, *J. S. Am. Earth Sci.*, 14, 61–75, [https://doi.org/10.1016/S0895-9811\(01\)00009-8](https://doi.org/10.1016/S0895-9811(01)00009-8), 2001.
- Blaber, S. J. M.: Fishes and fisheries in tropical estuaries: the last 10 years, *Estuar Coast Shelf S.*, 135, 57–65, 2013.
- Boltovskoy, E.: Foraminíferos del Golfo San Jorge. *Rev. Inst. Nac. Investig. Cien. Nat. Mus. Argen. Cien. Nat. “Bernardino Rivadavia”*, *Cien. Geol.*, III, Buenos Aires, 3, 79–246, 1954.
- Boltovskoy, E.: Foraminifera of mangrove swamps, *Physis A*, 42, 1–9, 1984.

- Boltovskoy, E. and Hincapié de Martínez, S.: Foraminiferos del manglar de Tesca, Cartagena, Colombia. *Rev. Esp. Micropal.*, 15, 205–220, 1983.
- Brönnimann, P., Dias-Brito, D., and Moura, J., A.: Foraminiferos da Facies Mangue Da Planície de Mare de Guaratiba, Rio de Janeiro, Brasil, *Anais do Congresso Latino-Americano de Paleontologia*, Porto Alegre, 2, 877–891, 1981.
- Brönnimann, P., Whittaker, J. E., and Zaninetti, L.: Brackish water foraminifera from mangrove sediments of southwest Viti Levu, Fiji Islands, Southwest Pacific. *Rev. Paléobiol.*, 11, 13–65, 1992.
- Burone, L., Venturini, N., Sprechmann, P., Valente, P., and Muniz, P.: Foraminiferal responses to polluted sediments in the Montevideo coastal zone, Uruguay. *Mar. Pollut. Bull.*, 52, 61–73, 2006.
- Call, M., Maher, D., Ruiz-Halpern, S., Mangion, P., Erler, D., Oakes, J., Rosentreter, J., Murray, R., and Eyre, B.: Spatial and temporal variability of carbon dioxide and methane fluxes over semi-diurnal and spring–neap–spring timescales in a mangrove creek. *Geochim. Cosmochim. Ac.*, 150, 211–225, <https://doi.org/10.1016/j.gca.2014.11.023>, 2015.
- Camacho, S., Moura, D., Connor, S., David, S., and Boski, T.: Ecological zonation of benthic foraminifera in the lower Guadiana Estuary (southeastern Portugal). *Mar. Micropaleontol.*, 114, 1–18, <https://doi.org/10.1016/j.marmicro.2014.10.004>, 2015.
- Cunha-Lignon, M., Coelho-Jr., C., Almeida, R., Menghini, R., Correa, F., Schaeffer-Novelli, Y., Cintrón-Molero, G., and Dahdouh-Guebas F.: Mangrove Forests and Sedimentary Processes on the South Coast of São Paulo State (Brazil). *J. Coast. Res., Spec. Iss.*, 56, 405–409, 2009.
- Cushman, J. A. and Brönnimann, P.: Some new genera and species of foraminifera from brackish water of Trinidad. *Contr. Lab. Foramin. Res.*, 24, 15–21, 1948a.
- Cushman, J. A. and Brönnimann, P.: Additional new species of arenaceous Foraminifera from shallow waters of Trinidad. *Contr. Lab. Foramin. Res.*, 24, 37–42, 1948b.
- Culver, S.: Benthic Foraminifera of Puerto Rican Mangrove-Lagoon Systems: Potential for Paleoenvironmental Interpretations. *Palaios*, 5, 34–51, <https://doi.org/10.2307/3514995>, 1990.
- Damasio, B., Timoszczuk, C., Kim, B. S. M., Sousa, S., Bicego, M., Siegle, E., and Figueira, R.: Impacts of hydrodynamics and pollutants on foraminiferal fauna distribution in the Santos Estuary (SE Brazil). *J. Sediment. Environ.*, 5, 61–86, <https://doi.org/10.1007/s43217-020-00003-w>, 2020.
- Debenay, J. P.: Recent foraminiferal assemblages and their distribution relative to environmental stress in the paralic environments of West Africa (Cape Timiris to Ebrie Lagoon). *J. Foramin. Res.*, 20, 267–282, 1990.
- Debenay, J. P.: Foraminiferal assemblages in a hypersaline lagoon, Araruama (RJ) Brazil. *J. Foramin. Res.*, 31, 133–151, 2001.
- Debenay, J. P. and Guillou, J. J.: Ecological transitions indicated by foraminiferal assemblages in paralic environments. *Estuaries*, 25, 1107–1120, 2002.
- Debenay, J. P., Duleba, W., Bonetti, C., De, S., Souza, M., and Eichler, B.: *Pararotalia cananeiaensis* n. sp.: Indicator of marine influence and water circulation in Brazilian coastal and paralic environments. *J. Foramin. Res.*, 31, 152–163, 2001.
- Debenay, J. P., Guiral, D., and Parra, M.: Ecological Factors Acting on the Microfauna in Mangrove Swamps. The Case of Foraminiferal Assemblages in French Guiana. *Estuar. Coast. Shelf S.*, 55, 509–533, <https://doi.org/10.1006/ecss.2001.0906>, 2002.
- Debenay, J. P., Guiral, D., and Parra, M.: Behaviour and taphonomic loss in foraminiferal assemblages of mangrove swamps of French Guiana. *Mar. Geol.*, 208, 295–296, 2004.
- Disaró, S. T., Totah, V. I., Watanabe, S., Ribas, E. R., and Pupo, D. V.: Biodiversidade Marinha da Bacia Potiguar: Foraminifera. *Mus. Nacional*, Rio de Janeiro, 1–218, 2014.
- Dolbeth, M., Vendel, A. L., Baeta, A., Pessanha, A., and Patrício, J.: Exploring ecosystem functioning in two Brazilian estuaries with fish richness, traits and food webs. *Mar. Ecol.-Prog. Ser.*, 560, 41–55, <https://doi.org/10.3354/meps11895>, 2016.
- Eichler, P.: Foraminiferal zonation from a subtropical mangrove in Bertioga Channel (São Paulo, SP, Brazil). *Reg. Stud. Mar. Sci.*, 25, 100460, <https://doi.org/10.1016/j.rsma.2018.100460>, 2018.
- Eichler, B. B., Debenay, J. P., Bonetti, C., and Duleba, W.: Distribution of benthic foraminifera in the southwestern zone of the estuarine-lagoonal system of Iguapé-Cananéia (Brazil). *Bol. Inst. Oceanogr.*, 43, 1–17, <https://doi.org/10.1590/S0373-55241995000100001>, 1995.
- Eichler, P. P. B., Eichler, B. B., De Miranda, L. B., and Rodrigues, A. R.: Modern foraminiferal facies in a subtropical estuarine Channel, Bertioga, São Paulo, Brazil. *J. Foramin. Res.*, 37, 234–247, 2007.
- Eichler, P., Rodrigues, A., Pereira, E., Eichler, B., Kahn, A., and Vital, H.: Foraminifera as Environmental Condition Indicators in Todos os Santos Bay (Bahia, Brazil). *Open J. Ecol.*, 5, 326–342, <https://doi.org/10.4236/oje.2015.57027>, 2015.
- Eichler, P., Gomes, C., Costa, R., Corrêa, A., Gomes, M., and Vital, H.: Foraminifera Distribution in Itapitangui River Mangrove, Cananeia (Sao Paulo, Brazil) and its Environmental Relations. *Modern Approaches in Oceanography and Petrochemical Sciences*, 3, 219–224, 2019.
- Ellison, A., Farnsworth, E., and Merkt, R. E.: Origins of mangrove ecosystems and the mangrove biodiversity anomaly. *Global Ecol. Biogeogr.*, 8, 95–115, 1999.
- Ellison, R. L. and Nichols, M. M.: Modern and Holocene foraminifera in the Chesapeake Bay region, in: *First International Symposium on Benthic Foraminifera of the Continental Margins*, edited by: Schafer, C. T. and Pelletier, B. R., 14 Part A – Ecology and Biology: *Marit. Sed., Spec. Publ.*, 1, 131–151, Halifax, Nova Scotia, Canada, 1976.
- Fajemila, O. T. and Langer, M. R.: Spatial distribution and biogeographic significance of foraminiferal assemblages from São Tomé and Príncipe, Gulf of Guinea, West Africa. *Neues Jahrb. Geol. Paläontol.*, 285, 337–360, <https://doi.org/10.1127/njgpa/2017/0686>, 2017.
- Fajemila, O. T., Langer, M. R., and Lipps, J. H.: Spatial Patterns in the Distribution, Diversity and Abundance of Benthic Foraminifera around Moorea (Society Archipelago, French Polynesia). *PLoS One*, 10, e0145752, <https://doi.org/10.1371/journal.pone.0145752>, 2015.
- Fajemila, O. T., Langer, M. R., and Lipps, J. H.: Atlas of Shallow-Water Tropical Benthic Foraminifera from Moorea (Society Islands, French Polynesia). *Cushman Found. Foramin. Res., Spec. Publ.*, 48, 1–107, 2020a.
- Fajemila, O. T., Sariaslan, N., and Langer, M. R.: Spatial distribution of benthic foraminifera in the Lagos Lagoon (Nigeria): Trac-



- ing the impact of environmental perturbations, *PLoS One*, 15, e0243481, <https://doi.org/10.1371/journal.pone.0243481>, 2020b.
- Fisher R. A., Corbet A. S., and Williams, C. B.: The relationship between the number of species and the number of individuals in a random sample of animal populations, *J. Anim. Ecol.*, 12, 42–58, 1943.
- Gasparini, S. and Vilela, C.: Paleoenvironmental evolution based on benthic foraminifera biofacies of the Paraíba do Sul Deltaic Complex, eastern Brazil, *J. S. Am. Earth Sci.*, 80, 291–303, <https://doi.org/10.1016/j.jsames.2017.09.026>, 2017.
- Geslin, E., Debenay, J. P., Duleba, W., and Bonetti, C.: Morphological abnormalities of foraminiferal tests in Brazilian environments: comparison between polluted and non-polluted areas, *Mar. Micropaleontol.*, 45, 151–168, 2002.
- Gillanders, B. M., Elsdon, T. S., Halliday, I. A., Jenkins, G. P., Robins, J. B., and Valesini, F. J.: Potential effects of climate change on Australian estuaries and fish-utilising estuaries: a review, *Mar. Freshwater Res.*, 62, 1115–1131, <https://doi.org/10.1071/MF11047>, 2011.
- Goldstein, S. T.: Foraminifera of relict salt marsh deposits, St. Catherines Island, Georgia: Taphonomic implications *Palaios*, 3, 327–334, 1988.
- Gleeson, J., Santos, I. R., Maher, D. T., and Golsby-Smith, L.: Groundwater-surface water exchange in a mangrove tidal creek: evidence from natural geochemical tracers and implications for nutrient budgets, *Mar. Chem.*, 156, 27–37, 2013.
- Hammer, Ø., Harper, D. A. T., and Ryan, P. D.: PAST: Paleontological statistics software package for education and data analysis, *Palaeontol. Electron.*, 4, 1–9, 2001 (software available at: <https://www.nhm.uio.no/english/research/infrastructure/past/>, last access: 30 June 2021).
- Halicz, E., Noy, N., and Reiss, Z.: Foraminifera from Shura Arwashie mangrove (Sinai), in: *Hydrobiology of the mangal*, edited by: Por, F. D. and Dor, I., Dr. W. Junk Publ., The Hague, 145–149, 1984.
- Hiltermann, H., Brönnimann, P., and Zaninetti, L.: Neue Biozönosen in den Sedimenten der Mangrove bei Acupe, Bahia, Brasilien, *Notes Lab. Pal. Univ. Geneve*, 8, 1–6, 1981.
- Instituto Brasileiro de Geografia e Estatística: Population Estimates for the cities Mamanguape, Rio Tinto and Marcação, available at: <https://www.ibge.gov.br/en/cities-and-states.html?view=municipio> (last access: 1 February 2021), 2019.
- Langer, M. and Lipps, J.: Foraminiferal distribution and diversity, Madang Reef and Lagoon, Papua New Guinea, *Coral Reefs*, 22, 143–154, <https://doi.org/10.1007/s00338-003-0298-1>, 2003.
- Langer, M., Thissen, J., Makled, W., and Weinmann, A.: The Foraminifera from the Bazaruto Archipelago (Mozambique), *Neues Jahrb. Geol. Paläontol.*, 267, 155–170, <https://doi.org/10.1127/0077-7749/2013/0302>, 2013.
- Langer, M. R. and Lipps, J. H.: Assembly and persistence of foraminifera in introduced mangroves on Moorea, French Polynesia, *Micropaleontology*, 52, 343–355, 2006.
- Langer, M. R., Hottinger, L., and Huber, B.: Functional Morphology in Low-Diverse Benthic Foraminiferal Assemblages from Tidal-Flats of the North Sea, *Senck. Marit.*, 20, 81–99, 1990.
- Langer, M. R., Fajemila, O. T., and Mannl, S.: Assemblages of recent intertidal mangrove foraminifera from the Akanda National Park, Gabon: sea level proxies preserved in faunal assemblages, *Neues Jahrb. Geol. Paläontol.*, 281, 327–338, 2016.
- Laut, L. L. M., Martins, V., da Silva, F. S., Crapez, M. A. C., Fontana, L. F., Carvalhal-Gomes, S. B. V., and Souza, R. C. C. L.: Foraminifera, thecamoebians, and bacterial activity in polluted intertropical and subtropical Brazilian estuarine systems, *J. Coast. Res.*, 32, 56–69, 2016.
- Leonel, R. M. V., Lopes, S. G. B. C., and Aversari, M.: Distribution of wood-boring bivalves in the Mamanguape river estuary, Paraíba, Brazil, *J. Mar. Biol. Assoc.*, 82, 1039–1040, 2002.
- Leopold, A., Marchand, C., Deborde, J., and Allenbach, M.: Water Biogeochemistry of a Mangrove-Dominated Estuary Under a Semi-Arid Climate (New Caledonia), *Estuar. Coast.*, 40, 773–791, <https://doi.org/10.1007/s12237-016-0179-9>, 2017.
- Li X., Burnett W., and Chanton, J.: Submarine Ground Water Discharge Driven by Tidal Pumping in a Heterogeneous Aquifer, *Ground Water*, 47, 558–568, <https://doi.org/10.1111/j.1745-6584.2009.00563.x>, 2009.
- Lipps, J. and Langer, M.: Benthic Foraminifera from the Meromictic Mecherchar Jellyfish Lake, Palau (Western Pacific), *Micropaleontology*, 45, 278–284, <https://doi.org/10.2307/1486137>, 1999.
- MacDonald, J. A., Glover, T., and Weis, J. S.: The impact of mangrove prop-root epibionts on juvenile reef fishes: a field experiment using artificial roots and epifauna, *Estuar. Coast.*, 31, 981, <https://doi.org/10.1007/s12237-008-9083-2>, 2008.
- Martins, V., Laut, L., Frontalini, F., Sequeira, C., Rodrigues, R., Fonseca, M., Bergamaschi, S., Pereira, E., Delavy, F., Figueiredo Jr., A., Miranda, P., Terroso, D., Pena, A., Laut, V., Figueira, R., and Rocha, F.: Controlling Factors on the Abundance, Diversity and Size of Living Benthic Foraminifera in the NE Sector of Guanabara Bay (Brazil), *J. Sediment. Environ.*, 1, 393–410, <https://doi.org/10.12957/jse.2016.26872>, 2016.
- Medeiros, A. P. M., Xavier, J. H. D. A., Da Silva, M. B., Aires-Souza, L., and Rosa, I. M. D. L.: Distribution patterns of the fish assemblage in the Mamanguape River Estuary, North-eastern Brazil, *Mar. Biol. Res.*, 14, 524–536, 2018.
- Moreno, J., Fatela, F., Andrade, C., Cascalho, J., Moreno, F., and Drago, T.: Living foraminiferal assemblages from the Minho and Coura estuaries (Northern Portugal): a stressful environment, *Thalassas*, 21, 17–28, 2005.
- Murray, J.: Ecology and applications of benthic foraminifera, Cambridge University Press, 426 pp., 2006.
- Murray, J. J. and Bowser, S. S.: Mortality, protoplasm decay rate, and reliability of staining techniques to recognize “living” foraminifera: a review, *J. Foramin. Res.*, 30, 66–70, 2000.
- Murray, J. W.: *Distribution and Ecology of Living Benthic Foraminiferids*, Heinemann Educational Books, London, 274 pp., 1973.
- Murray, J. W.: *Ecology and Palaeoecology of Benthic Foraminifera*, Longman, Harlow, 397 pp., 1991.
- Murray, J. W., Sturrock, S., and Weston J.: Suspended load transport of foraminiferal tests in a tide- and wave-swept sea, *J. Foramin. Res.*, 12, 51–85, 1982.
- Nascimento, D., Alves, A. G., Alves, R., Barboza, R., Diele, K., and Mourão, J.: An examination of the techniques used to capture mangrove crabs, *Ucides cordatus*, in the Mamanguape River estuary, northeastern Brazil, with implications for management, *Ocean Coast Manage.*, 130, 50–57, <https://doi.org/10.1016/j.ocecoaman.2016.05.010>, 2016.
- Nordi, N., Nishida, A. K., and Alves, R. R.: Effectiveness of two gathering techniques for *Ucides cordatus* in Northeast Brazil: im-

- plications for the sustainability of mangrove ecosystems, *Hum. Ecol.*, 37, 121–127, <https://doi.org/10.1007/s10745-009-9214-9>, 2009.
- Patterson, R. T. and Fishbein, E.: Re-examination of the statistical methods used to determine the number of point counts needed for micropaleontological quantitative research, *J. Paleontol.*, 63, 245–248, 1989.
- Paerl, H., Valdes, L., Peierls, B., Adolf, J., and Harding, L.: Anthropogenic and climate influences on the eutrophication of large estuarine ecosystems, *Limnol. Oceanogr.*, 51, 448–462, [https://doi.org/10.4319/lo.2006.51.1\\_part\\_2.0448](https://doi.org/10.4319/lo.2006.51.1_part_2.0448), 2006.
- Paludo, D. and Klonowski, V. S.: Barra de Mamanguape e PB: Estudo do impacto do uso de madeira de manguezal pela população extrativista e da possibilidade de reflorestamento e manejo dos recursos madeireiros, Conselho Nacional da Reserva da Biosfera da Mata Atlântica, Sao Paulo, 1999.
- Passos, C., Kukimodo, I., and Semensatto Junior, D.: Foraminiferal assemblages along the intertidal zone of Itapanhaú River, Bertioga (Brazil), *J. S. Am. Earth Sci.*, 79, 297–306, <https://doi.org/10.1016/j.jsames.2017.08.017>, 2017.
- Rafaela, M., Filho, J., Rocha-Barreira, C., Matthews-Cascon, H., Santos, E., David, H., and Matos, A.: Benthic Estuarine Assemblages of the Northeastern Brazil Marine Ecoregion: A Benthic Perspective, in: *Brazilian Estuaries*, edited by: Lana, P. da C. and Bernardino, A. F., Springer, Berlin, Heidelberg, Germany, 75–94, [https://doi.org/10.1007/978-3-319-77779-5\\_3](https://doi.org/10.1007/978-3-319-77779-5_3), 2018.
- Rickard, D.: How long does it take a pyrite framboid to form?, *Earth Planet. Sc. Lett.*, 513, 64–68, <https://doi.org/10.1016/j.epsl.2019.02.019>, 2019.
- Rouvilleis, A.: Comparaison entre la faune de foraminifères de la surface des mangroves actuelles du Sénégal de Bassid (Sine Saloum), d'Oussouye et de Balingor (Casamance) et celle d'un sondage à Bango (Delta du Fleuve Sénégal), *Cah. Micropal.*, 2, 149–154, 1982.
- Santos, E., Araújo, L., and Marcelino, A.: Análise climática da Bacia Hidrográfica do Rio Mamanguape, *Rev. Brasil. Engen. Agrí. Amb.*, 19, 9–14, <https://doi.org/10.1590/1807-1929/agriambi.v19n1p9-14>, 2015.
- Saunders, J. B.: Recent foraminifera of mangrove swamps and river estuaries and their fossil counterparts in Trinidad, *Micropaleontology*, 4, 79–92, 1958.
- Scavia, D., Field, J., Boesch, D., Buddemeier, R., Burkett, V., Cayan, D., Fogarty, M., Harwell, M., Howarth, R., Mason, C., Reed, D., Royer, T., Sallenger, A., and Titus, J.: Climate Change Impacts on U.S. Coastal and Marine Ecosystems, *Estuar. Coast.*, 25, 149–164, <https://doi.org/10.1007/BF02691304>, 2002.
- Schiedek, D., Sundelin, B., Readman, J., and Macdonald, R.: Interactions between climate change and contaminants, *Marine Pollution Bulletin*, 54, 1845–1856, <https://doi.org/10.1016/j.marpolbul.2007.09.020>, 2008.
- Scott, D. and Medioli, F.: Vertical zonation of marsh foraminifera as accurate indicators of former sea-levels, *Nature*, 272, 528–531, <https://doi.org/10.1038/272528a0>, 1978.
- Scott, D. B., Schnack, E. J., Espinosa, L., and Barbosa, C. F.: Recent Marsh Foraminifera from the East Coast of South America: Comparison to the Northern Hemisphere, *Paleoecology, Biostratigraphy, Paleooceanography and Taxonomy of Agglutinated Foraminifera*, 717–737, Kluwer Academic Publishers, Dordrecht, the Netherlands, 1990.
- Semensatto Jr., D. L., Funo, R. H. F., Dias-Brito, D., and Coelho Jr., C.: Foraminiferal ecological zonation along a Brazilian mangrove transect: Diversity, morphotypes and the influence of subaerial exposure time, *Rev. Micropaléontol.*, 52, 67–74, <https://doi.org/10.1016/j.revmic.2008.06.004>, 2009.
- Shaw, T. A., Kirby, J. R., Holgate, S., Tutman, P., and Plater, A. J.: Contemporary salt-marsh foraminiferal distribution from the adriatic coast of croatia and its potential for sea-level studies, *J. Foramin. Res.*, 46, 314–332, <https://doi.org/10.2113/gsjfr.46.3.314>, 2016.
- Spalding, M., Fox, H., Allen, G., Davidson, N., Ferdaña, Z., Finlayson, M., Halpern, B., Jorge, M., Lombana, A. Lourie, S., Martin, K., McManus, E., Molnar, J., Recchia, C., and Robertson, J.: Marine Ecoregions of the World: A Bioregionalization of Coastal and Shelf Areas, *BioScience*, 57, 573–583, <https://doi.org/10.1641/B570707>, 2007.
- Thiel, J., Byrne, J. M., Kappler, A., Schink, S., and Pester, M.: Pyrite formation from FeS and H<sub>2</sub>S is mediated through microbial redox activity, *P. Natl. Acad. Sci. USA*, 116, 6897–6902, <https://doi.org/10.1073/pnas.1814412116>, 2019.
- Thissen, J. M. and Langer, M. R.: Spatial Patterns and Structural Composition of Foraminiferal Assemblages from the Zanzibar Archipelago (Tanzania), *Paleontographica, Abt. A: Palaeozoology – Stratigraphy*, 308, 1–67, 2017.
- Thomas, E. and Varekamp J. C.: Palaeo-environmental analyses of marsh sequences (Clinton, Connecticut): evidence for punctuated rise in relative sea level during the latest Holocene, *J. Coast. Res.*, 11, Spec. Iss., 125–158, 1991.
- Todd, R. and Brönnimann, P.: Recent foraminifera and thecamoebina from the eastern Gulf of Paria, Trinidad, Cushman Foundation for Foraminiferal Research, 3, 1–43, 1957.
- Van der Linden, P., Marchini, A., Smith, C., Dolbeth, M., Simone, L., Marques, J., Joseline, M., Medeiros, C., and Patrício, J.: Functional changes in polychaete and mollusc communities in two tropical estuaries, *Estuar. Coast. Shelf S.*, 187, 62–73, <https://doi.org/10.1016/j.ecss.2016.12.019>, 2017.
- Weinmann, A. E.: Benthic shallow-water foraminifera from Fernando de Noronha and Abrolhos, Brazil: A comparative study between two archipelagos and biogeographic analysis of larger foraminifera within the tropical western Atlantic Ocean, Master thesis, University of Bonn, 172 pp., 2009.
- Wilson, B., Miller, K., Thomas, A. L., Cooke, N., and Ramsingh, R.: Foraminifera in the Mangal at the Caroni Swamp, Trinidad: Diversity, Population Structure and Relation to Sea Level, *J. Foramin. Res.*, 38, 127–136, 2008.
- Woodroffe, S. A., Horton, B. P., and Larcombe, P.: Contemporary intertidal foraminifera distributions of Cleveland Bay, Central Great Barrier Reef shelf, Australia: implications for sea-level reconstructions, *J. Foramin. Res.*, 35, 259–270, 2005.
- Zaninetti, L., Brönnimann, P., Dias-Brito, D. A., Mitsuru Casaletti, P., Koutsoukos, E., and Silveira, S.: Distribution écologique des Foraminifères dans la Mangrove d'Acupe, Etat de Bahia, Brésil, Notes du Laboratoire de Paleontologie de l'Université de Genève, 4, 1–17, 1979.