

Opisthobranchia Of Japan (II)

Baba, Kikutaro

The Amakusa Marine Biological Laboratory, Tomioka, Kumamoto-ken.

<https://doi.org/10.5109/22586>

出版情報：九州大学大学院農学研究院紀要. 5 (7), pp.289-344, 1937-11. Kyushu Imperial University
バージョン：
権利関係：

OPISTHOBRANCHIA OF JAPAN (II)¹⁾

Kikutarō BABA

Suborder Nudibranchia

Tribe 1. Holohepatica (= Doridacea)

Family Hexabanchidae

Genus *Hexabanchus* EHRENBERG, 1831

Hexabanchus EHRENBERG, Symb. phys., 1831.

Type: *Hexabanchus praetextus* EHRENBERG.

34. *Hexabanchus marginatus* (QUOY & GAIMARD, 1832)

Family Euphuridae (= Polyceridae)

Subfamily Euphurinae

Genus *Euphurus* RAFINESQUE, 1815

Euphurus RAFINESQUE, Analyse Nature, 1815, p. 142.

Triopa JOHNSTON, Ann. Nat. Hist., vol. 1, 1838, p. 123.

Type: *Doris clavigera* O. F. MÜLLER.

35. *Euphurus ornatus* nov. sp.

(Pl. 2, fig. 8)

Distribution in Japan: Tateyama and Seto (Kii).

The living animal is limaciform, measuring about 10 mm in length. The back is smooth save for the presence of about a dozen small blunt tubercles in rather regular rows. It is marked off from

¹⁾ Contributions from the Zoological Laboratory, Kyūshū Imperial University, No. 103. Papers from the Amakusa Marine Biological Laboratory, No. 62.

sides by a series of 7 processes consisting of an ovoid body with a short stalk. The perfoliate rhinophores are retractile within low sheaths with an entire margin. The non-retractile branchiae, 3-4 in number, are simply pinnate, and lie closely in front of the anus. The margin of the head veil bears 4-5 processes similar in shape to those along the edge of the body but having small granules towards the top. The oral tentacles are canaliculiform. The anterior edge of the foot is abruptly truncated.

The general body colour is yellowish white beautifully ornamented with tiny orange spots. The rhinophores are dull yellow tipped with vermilion.

Localities: Tateyama (May 1925; 1 sp.) and Seto (Apr. 1937; 2 sps., coll. by Mr. F. HIRO).

This species is especially characteristic in the shape of the processes along the edge of the head and back. It has also an unmistakable coloration.

Genus *Polycera* CUVIER, 1817

Polycera CUVIER, Règne Animal, tom. 2, 1817, p. 389.

Type: *Doris quadrilineata* O. F. MÜLLER.

36. *Polycera fujitai* nov. sp.

(Text-fig. 1)


Distribution in Japan: Tomioka (Amakusa).

The animal in life is limaciform and measures up to 45 mm in length. The rhinophore consists of a long stalk and a small perfoliate clavus, and is without a sheath. The branchiae, 7 in number, are tripinnate and arranged in little more than a semi-circle, about half way back. The hinder part of the circle is occupied by the anus. The frontal veil is slightly expanded, and bears 6-10 simple digitiform processes. The back margin is distinct, having a row of small subconical papillae. Similar ones are also distributed on the back, sides and tail crest. The genital orifice lies on the right side a short distance behind the head. The oral tentacles are lobiform. The antero-lateral corners of the foot are slightly produced into obtuse points.

The ground colour of the body is translucent yellowish white, but it is almost everywhere obscured by black speckles and also by irregular streaks of yellowish brown and dark. The stalk of

the rhinophore is dark, the clavus black and the tip white. The dermal papillae tipped with light yellow. The sole uniformly yellowish white.

The jaw-plates are well-developed and of a horn-yellow colour. They are approximately triangular in outline, and the anterior edge



Text-fig. 1.—*Polycera fujitai*. A. Dorsal view of entire animal ($\times 1.5$); B. A half-row of radula ($\times 70$), a. 1st lateral tooth b. 2nd tooth; C. A pair of jaw-plates, in frontal lateral view ($\times 10$).

is continued out into wing-like flange which stands laterally recurved. The radula is of a characteristic black-brown, consisting of about 10-13 rows, 4-5 in each half-row, and may be formulated as $10 (-13) \times 4.5.0.4-5$. The 1st lateral tooth is roughly hamate with a blunt spine about half way down. The 2nd lateral tooth is larger and is simply hamate. The succeeding teeth, 2-3 in number, are much reduced to become scale-like plates.

Locality: Tomioka (Jan. 1933; fairly common on *Zostera*).

The present species resembles a Vancouver Island Nudibranch, *Polycera zosterae* O'DONOGHUE, in the habitat and external form, but is distinguished from that form by having a different coloration, much more numerous branchiae and fewer lateral teeth. It is here named *P. fujitai* in honour of Dr. Tsunenobu FUJITA.

Genus *Gymnodoris* STIMPSON, 1855*Gymnodoris* STIMPSON, Proc. Acad. Nat. Sci. Philadelphia, 1855, p. 379.*Trevelyana* KELAART, Ann. Mag. Nat. Hist., ser. 3, vol. 1, 1858, p. 257.Type: *Gymnodoris maculata* STIMPSON.*Synopsis of the Japanese species:*

- A. Branchiae in a transverse crescentic row *striata*.
 B. Branchiae in a circular or a horseshoe row.
1. Innermost lateral tooth larger than the succeeding teeth.
 - a. Body uniformly orange-yellow *inornata*.
 - b. Body yellow with orange spots, sometimes white with yellow or orange spots *citrina*.
 2. Innermost lateral tooth smaller than the succeeding teeth.
 - a. Body white with orange spots *alba*.
 - b. Body yellowish with orange-yellow spots and streaks *okinawae*.

37. *Gymnodoris maculata* STIMPSON, 185538. *Gymnodoris citrina* (BERGH, 1877)

(Pl. 1, fig. 9)

By examining many specimens of *Gymnodoris japonica* (BABA) I have found that this is identical with *G. citrina* (BERGH), the 2nd lateral tooth of the radula showing no constant specific difference. In some specimens the tooth in question has a small basal denticle, but in others it is entirely smooth.

39. *Gymnodoris alba* (BERGH, 1877)40. *Gymnodoris inornata* (BERGH, 1880)

(Pl. 1, fig. 8)

41. *Gymnodoris striata* (ELIOT, 1908)42. *Gymnodoris okinawae* BABA, 1936Genus *Nembrotha* BERGH, 1877*Nembrotha* BERGH, Malac. Unters., Hft. 11, 1877, p. 450.Type: *Nembrotha nigerrima* BERGH.43. *Nembrotha luteolineata* BABA, 1936Genus *Caloplocamus* BERGH, 1879*Euplocamus* PHILIPPI, Enum. Moll. Siciliae, 1, 1836, p. 103.

Kaloplocamus BERGH, Verh. k. k. zool.-bot. Gesell. Wien, Bd. 29, 1879, p. 623, foot-note.
Caloplocamus THIELE, Handb. syst. Weicht., Th. 2, 1931, p. 425.

Type: *Euplocamus croceus* PHILIPPI.

44. *Caloplocamus ramosus* (CANTRAINE, 1835)¹⁾

Genus *Plocamophorus* RÜPPELL & LEUCKART, 1828

Plocamopherus RÜPPELL & LEUCKART, Wirbellose Thiere, 1828, p. 17.

Type: *Plocamopherus ocellatus* RÜPPELL & LEUCKART.

- a. First lateral tooth bicuspid, the succeeding inner teeth simply hamate. Body yellow or orange-yellow thickly covered with chocolate mottles ... *imperialis*.
 b. Inner lateral teeth roughly bicuspid. Body yellowish shaded with chocolate, and almost everywhere scattered with deep chocolate and chrome-yellow spots
 *tilesii*.

45. *Plocamophorus imperialis* ANGAS, 1864

Plocamophorus imperialis ANGAS, Journ. Conchyl., Paris, tom. 12, 1864, pp. 59-60, pl. 5, fig. 7.—Vaucluse (Australia).

Plocamopherus imperialis ELIOT, Journ. Coll. Sci. Imp. Univ. Tôkyô, vol. 35, art. 1, 1913, p. 30.—Japan.

Locality: Seto, Kii (Mar. 1937; 2 sps.).

46. *Plocamophorus tilesii* BERGH, 1877

47. *Plocamophorus* sp. HIRASÉ, 1927

Plocamophorus sp. HIRASÉ, Moluskoj, 1927, p. 1472, fig. 2832.—Misaki.

Genus *Kalinga* ALDER & HANCOCK, 1864

Kalinga ALDER & HANCOCK, Trans. Zool. Soc. London, vol. 5, pt. 3, 1864, pp. 134-135.

Type: *Kalinga ornata* ALDER & HANCOCK.

48. *Kalinga ornata* ALDER & HANCOCK, 1864

Subfamily Acanthodoridinae

Genus *Acanthodoris* GRAY, 1850

Acanthodoris GRAY, Figs. Mollusc. Anim., vol. 4, 1850, p. 103.

Type: *Doris pilosa* ABILDGAARD.

¹⁾ Synonym: *C. croceus* (PHILIPPI), see O'DONOGHUE (1929, p. 775).

Synopsis of the Japanese species :

- a. Back closely covered with conical papillae... .. *pilosa*.
 b. Back smooth *uchidai*.

49. *Acanthodoris pilosa* (ABILDGAARD, 1789)

50. *Acanthodoris uchidai* BABA, 1935

Subfamily Okeniinae (=Goniodoridinae)

Genus *Goniodoris* FORBES & GOODSIR, 1839

Goniodoris FORBES & GOODSIR, Athenaeum, no. 618, 1839, p. 647.

Type: *Doris nodosa* MONTAGU.

Synopsis of the Japanese species :

- a. Back with a median longitudinal crest *castanea*.
 b. Mid-dorsal crest replaced by 3 large obtuse tubercles *glabra* nov. sp.

51. *Goniodoris castanea* ALDER & HANCOCK, 1845

(Pl. 1, fig. 5)

52. *Goniodoris glabra* nov. sp.

(Pl. 1, fig. 6; text-fig. 2)

Distribution in Japan: Tomioka (Amakusa).

The body is limaciform and elongated, measuring about 25 mm in length. The back is marked off from the rest of the body by a



Text-fig. 2.—*Goniodoris glabra*.
 A half-row of radula ($\times 150$).

pallial flange standing recurved all around excepting at the posterior median notch. The mid-dorsal crest is replaced in this species by 3 large obtuse tubercles disposed longitudinally between rhinophores and branchiae. The rhinophore is non-retractile and has a perfoliate clavus. The tail has a mid-dorsal crest. The back and sides are covered with sparse almost indistinct tubercles. The head is produced laterally into a pair of oral tentacles. The branchiae consist of 3 non-retractile plumes arranged separately around the anus.

The general body colour is a dark chocolate sparsely spotted

with yellow, and there is a yellow cross band a short distance behind the rhinophores.

The cuticular labial disk is without marked armatures. The radula formula is $20 \times 1.1.0.1.1$. The inner lateral tooth is large, consisting of a roughly oblong base and a smooth hook without denticulations. The outer lateral tooth is simply scale-like.

Locality: Tomioka (Mar. 1934; 1 sp.).

This species is quite distinctive and differs from other members of the genus in the external form with 3 branchiae and 3 mid-dorsal tubercles, and in the inner tooth of the radula which has no denticulation. Also the coloration of this species is unmistakable.

53. *Goniodoris* sp. BABA, 1935

Genus *Okenia* MENKE, 1830

Okenia MENKE, Synops. method. Mollusc., ed. 2, 1830, p. 10.

Type: *Idalia elegans* LEUCKART.

With 2 subgenera: *Okenia* s. s. and *Idaliella*.

54. *Okenia (Idaliella) barnardi* nov. sp.

(Pl. 2, fig. 6; text-fig. 3)


Distribution in Japan: Tomioka (Amakusa).

The living animal is small, limaciform and about 10 mm in length. The back is marked off from the rest of the body by a low pallial ridge which bears 16 slender clavate papillae, 8 on each side, the posterior ones being bifurcated. The rhinophores are slender and non-retractile, and the greater part of their length is occupied by a perfoliate clavus. The non-retractile branchiae, 7 in number, are simply pinnate and arranged around the anus. Over the mouth is an oral veil produced laterally into a pair of tentacular expansions. The back and sides are nearly smooth; the tail is produced behind; the foot is very large and expanded. The genital orifice lies on the right side, well behind the level of the rhinophores.

The ground colour of the body is yellowish white. Inside the margin of the back there is a dark chocolate band, inclined to a brownish colour towards the centre. A similar band surrounds

the sides of the body. The back and sides are spotted with opaque white. The rhinophores and branchial plumes are of a chocolate colour.

The labial armature takes the form of a semicircle. It consists of a large number of closely-set hooks, each bearing 5-7 irregular



Text-fig. 3.—*Okenia barnardi*. A. Labial armature ($\times 90$); B. Elements of the same ($\times 500$); C. A row of radula ($\times 220$), a. inner lateral tooth, b. outer tooth.

denticles at the tip. Radula formula $22 \times 1.1.0.1.1$. The inner lateral tooth is large and consists of a hook at right angles to the base; this hook bears a series of denticles about 15 in number. The outer lateral tooth is decidedly smaller and simply scale-like.

Locality: Tomioka (May 1933; 1 sp.).

The present species is referred to the subgenus *Idaliella* by the absence of papillae at the centre of the back, but differs from all the other species so far recorded in external features, in colours and in the shape of the labial armature. I propose to call it *Okenia barnardi*, after my friend Dr. K. H. BARNARD of the South African Museum, who assisted me in identifying the species.

Genus *Trapania* PRUVOT-FOL, 1931

Drepania LAFONT, Journ. Conchyl., Paris, tom. 22, 1874, p. 369 (non *Drepania* HÜBNER, 1816, for Lepidoptera).

Trapania PRUVOT-FOL, Bull. Mus. Hist. Nat. Paris, ser. 2, tom. 3, no. 3, 1931 (June), p. 309; PRUVOT-FOL, ditto, no. 8, 1931, p. 747.

Drepanida MAGFARLAND, Nautilus, vol. 45, no. 1, 1931 (July), p. 31.

Type: *Drepania fusca* LAFONT.

55. *Trapania japonica* (BABA, 1935)¹⁾

Family Vayssiereidae (=Okadaidae)

Genus *Okadaia* BABA, 1930

56. *Okadaia elegans* BABA, 1930

Family Dorididae

Subfamily Glossodoridinae

Genus *Glossodoris* EHRENBERG, 1831

Glossodoris EHRENBERG, Symb. phys., 1831.

Chromodoris ALDER & HANCOCK, Monogr. Brit. Nudib., pt. 7, 1855, appendix, p. 17.

Type: *Glossodoris xantholeuca* EHRENBERG.

Synopsis of the Japanese species:

- a. First lateral tooth tricuspid, succeeding teeth bicuspid... .. *festiva*.
- b. Lateral teeth very numerous (162.0.162), each with two cusps *thalassopora*.
- c. First lateral tooth denticulate on both sides, succeeding teeth with an external denticulate margin *pallescens, alderi, sibogae, aurcopurpurea, lineolata, ditonota, pantharcla*.

57. *Glossodoris lineolata* (VAN HASSELT, 1824)

Doris lineolata VAN HASSELT, Allgem. Konst en Letter-Bode, 1824, p. 22.—Java.

Locality: Hachijō-shima (1 sp., coll. by. Mr. T. FUJITA).

58. *Glossodoris festiva* (ADAMS, 1861)

59. *Glossodoris alderi* (COLLINGWOOD, 1881)²⁾

60. *Glossodoris pallescens* (BERGH, 1875)

61. *Glossodoris thalassopora* (BERGH, 1879)

Chromodoris thalassopora BERGH, Journ. Mus. Godeffroy, Hft. 14, 1879, pp. 19-21, pl. 4, figs. 35-40; pl. 5, figs. 1-2.—'M. japonicum'.

¹⁾ Synonym: *Drepania japonica*, see BABA (1935, p. 336).

²⁾ Synonym: *G. reticulata* (PEASE), see BABA (1933, p. 169).

62. *Glossodoris pantharella* (BERGH, 1879)

Chromodoris pantharella BERGH, Journ. Mus. Godeffroy, Hft. 14, 1879, p. 3.—Formosa Strait; BERGH, Malakoz. Blätt., Bd. 1, 1879, pp. 94-98, pl. 3, figs. 12-25.—Formosa Strait.

63. *Glossodoris aureopurpurea* (COLLINGWOOD, 1881)64. *Glossodoris clitonota* (BERGH, 1905)

Chromodoris clitonota BERGH, Siboga-Exped., 1905, p. 160, pl. 5, fig. 16.—Java; RISBEC, Faune Colon. Franç., tom. 2, 1928, pp. 156-158; pl. 6, fig. 9, text-fig. 45.—New Caledonia.

Glossodoris clitonota TAKI, Venus, vol. 3, no. 4, 1932, pp. 215-216, fig. 7.—Seto.

65. *Glossodoris sibogae* (BERGH, 1905)

Chromodoris sibogae BERGH, Siboga-Exped., 1905, p. 157, pl. 16, figs. 38-39.—Timor; ELIOT, Journ. Coll. Sci. Imp. Univ. Tôkyô, vol. 35, art. 1, 1913, pp. 27-28, pl. 2, fig. 7.—Misaki; HIRASÉ, Moluskoj, 1927, p. 1471, fig. 2830.—Misaki.

66. *Glossodoris* sp. FUJITA, 1893

Chromodoris sp. FUJITA, Dôbutsugaku Zasshi, vol. 5, no. 55, 1893, pp. 163-164, fig. 3.—Misaki.

Genus *Noumea* RISBEC, 1928

Noumea RISBEC, Faune Colon. Franç., tom. 2, 1928, p. 165.

Type: *Noumea romeri* RISBEC.

67. *Noumea nivalis* nov. sp.


(Text-fig. 4)

Distribution in Japan: Seto (Kii), Zushi and Kominato.

The body is typically glossodoridiform, ranging from 8 to 15 mm in length. The branchiae consist of 7-9, small, simply pinnate plumes. No knob-like bodies are present on the postero-ventral margin of the mantle.

The ground colour of the body is snow-white. The back is ornamented with almost inconspicuous colours as follows: (1) a yellow, often discontinuous line, running all around the pallial margin; (2) a few, tiny, orange-yellow spots irregularly scattered over the centre. The upper half of the rhinophore is orange-yellow, and the basal half white. The branchial plumes are of a uniform white, sometimes tipped with yellow. The foot is without markings.

The elements of the labial armature are closely-set tiny hooks, each with a simple or bifid tip. The radula formula is about $30 \times 25.0.25$. The rachis is narrow, without a thickening. The 1st lateral tooth is large, consisting of an exceedingly broad base and a pointed blade. This blade bears a single denticle on the inside and 2-3 denticles on the outside. The succeeding teeth are smaller and typically hamate, each with a series of 3-5 denticles. The outermost teeth are also hamate, the denticles being confined to the tip of the hook.



Text-fig. 4.—*Noumea nivalis*. A. Entire animal ($\times 5$); B. Elements of labial armature ($\times 500$); C. A half-row of radula ($\times 500$), a. 1st lateral tooth, b. succeeding teeth, c. outermost tooth.

Localities: Seto (Mar. 1937; 1 sp.), Zushi (May 1931; 1 sp.) and Kominato (June 1933; 2 sps.).

The present species resembles the New Caledonian members of *Noumea* in the 1st lateral tooth consisting of a large broad base, but differs in colours and in details of the radula from all of them.

Genus *Cadlina* BERGH, 1879

Cadlina BERGH, Proc. Acad. Nat. Sci. Philadelphia, 1879, p. 170 (114).

Type: *Doris repanda* ALDER & HANCOCK.

Synopsis of the Japanese species:

- a. Tubercles on back granular, closely set... .. *japonica*.
 b. Tubercles on back bluntly conical, sparsely set *sagamiensis*.

68. *Cadlina japonica* BABA, 1937

69. *Cadlina sagamiensis* BABA, 1937

70. *Cadlina* sp. ELIOT, 1913

Cadlina (or *Tyrinna*) sp. ELIOT, Journ. Coll. Sci. Imp. Univ. Tôkyô, vol. 35, art. 1, 1913, p. 21.—Ago Bay.

71. *Cadlina* sp. BABA, 1935Genus *Casella* H. & A. ADAMS, 1854*Casella* H. & A. ADAMS, Genera rec. Moll., vol. 2, 1854, p. 57.Type: *Casella gouldii* H. & A. ADAMS.72. *Casella atromarginata* (CUVIER, 1804)*Doris atromarginata* CUVIER, Ann. Mus. Hist. Nat., tom. 4, 1804, p. 473, pl. 2, fig. 6.—
'Mer des Indes'.*Casella atromarginata* TAKI, Venus, vol. 3, no. 4, 1932, pp. 214-215, fig. 6.—Seto.Genus *Ceratosoma* ADAMS & REEVE, 1850*Ceratosoma* ADAMS & REEVE, Samarang, 1850, pp. 67-68.Type: *Ceratosoma cornigerum* ADAMS & REEVE.73. *Ceratosoma cornigerum* ADAMS & REEVE, 1850

Subfamily Actinocyclusinae

Genus *Actinocyclus* EHRENBERG, 1831*Actinocyclus* EHRENBERG, Symb. phys., 1831.Type: *Actinocyclus verrucosus* EHRENBERG.74. *Actinocyclus japonicus* (ELIOT, 1913)

(Pl. 1, fig. 14)

Subfamily Thorunninae

Genus *Rostanga* BERGH, 1879*Rostanga* BERGH, Arch. f. Naturgesch., Jahrg. 45, Bd. 1, 1879, p. 353.Type: *Doris coccinea* ALDER & HANCOCK.75. *Rostanga arbutus* (ANGAS, 1864)

Subfamily Doridinae

Genus *Echinodoris* BERGH, 1874*Echinodoris* BERGH, Journ. Mus. Godeffroy, Hft. 6, 1874, p. 109.Type: *Doris eolida* QUOY & GAIMARD.76. *Echinodoris armata* ELIOT, 1913*Echinodoris armata* ELIOT, Journ. Coll. Sci. Imp. Univ. Tôkyô, vol. 35, art. 1, 1913,
pp. 7-8.—Japan.

Genus *Ctenodoris* ELIOT, 1907

Ctenodoris ELIOT, Proc. Malac. Soc. London, vol. 7, 1907, p. 338.

Ctenodoris is especially characteristic in the transverse arrangement of the simply pinnate branchiae. These latter are retractile within a cavity, whose foremost lip is valve-like. The back is granulate. The oral tentacles are lobiform. No labial armatures. Radula formula $\infty.0.\infty$. The lateral teeth are all hamate and smooth. The hermaphrodite gland spreads over the liver.

Type: *Staurodoris pecten* ELIOT.

In the external configuration, *Ctenodoris* closely resembles *Guyonia* RISBEC, 1928, but the latter is said to have digitate oral tentacles, a couple of jaw-plates and a massive hermaphrodite gland separated from the liver.


77. *Ctenodoris aurantiaca* (ELIOT, 1913)

(Text-fig. 5)

Doris (*Ctenodoris*) *aurantiaca* ELIOT, Journ. Coll. Sci. Imp. Univ. Tôkyô, vol 35, art. 1, 1913, pp. 5-7, pl. 1, fig. 1.—Misaki.

Distribution in Japan: Tomioka (Amakusa) and Misaki.

The body is typically doridiform and has the shape of an elongated ellipse with the mantle extending beyond the foot all around. It measures 12-26 mm in length. The back is closely covered with tiny obtuse granules. The branchial plumes, 14-16 in number, are simply pinnate and are arranged in a transverse row. The branchial cavity is not circular but is crescentic with the horns directed anteriorly and with the foremost valve-like lip extending behind. The oral tentacles are lobiform and contiguous at the bases. The anterior edge of the foot is abruptly truncated and bilabiate.



Text-fig. 5.—*Ctenodoris aurantiaca*. A. Entire animal in dorsal and ventral views ($\times 2$); B. A half-row of radula ($\times 500$), a. innermost lateral teeth, b. outermost tooth.

The colour of the entire animal is yellow.

The labial cuticle is without armatures. The radula formula is about $45 \times 40-60.0.40-60$. The lateral teeth are all simply hamate and smooth. The hermaphrodite gland covers the liver.

Locality: Tomioka (June 1935-36; 3 sps. on a yellow sponge).

Subfamily Archidoridinae

Genus *Trippa* BERGH, 1877

Trippa BERGH, Jahrb. Deutsch. Malakoz. Gesell., Bd. 4, 1877, p. 63; BERGH, Malac. Unters., Hft. 12, 1877, p. 543.

Type: *Trippa ornata* BERGH.

78. *Trippa intecta* (KELAART, 1858)

Genus *Petelodoris* BERGH, 1881

Petelodoris BERGH, Verh. k. k. zool.-bot. Gesell. Wien, Bd. 31, 1881, pp. 11 (227)-12 (228).

Type: *Petelodoris triphylla* BERGH.

79. *Petelodoris triphylla* BERGH, 1881

Petelodoris triphylla BERGH, Verh. k. k. zool.-bot. Gesell. Wien, Bd. 31, 1881, pp. 12 (228)-14 (230), pl. 7, figs. 4-15.—Enoshima.

Genus *Geitodoris* BERGH, 1892

Geitodoris BERGH, Zool. Jahrb., Syst., Bd. 6, 1892, p. 130.

Type: *Doris complanata* VERRILL.

Synopsis of the Japanese species:

- a. Back grayish yellow mottled with chocolate brown. A half-row of radula composed of 3 types of teeth: inner teeth simply hamate, middle teeth hamate with a serrulated edge, outer teeth thin and crowded together ... *ohshimai*.
- b. Back yellow. A half-row of radula composed of 2 types of teeth: inner teeth simply hamate, outer teeth thin and crowded together ... *lutea* nov. sp.

80. *Geitodoris ohshimai* BABA, 1936


81. *Geitodoris lutea* nov. sp.

(Text-fig. 6)

Distribution in Japan: Tomioka (Amakusa).

The animal is doridiform and somewhat flattened, and the mantle projects beyond the foot all around. It is about 32 mm

long. The back is closely covered with tiny granules. The rhinophore sheath has a granulated margin. The branchial plumes are in a circle around the anal papilla, and can be withdrawn into a cavity with a granulated margin. They number 3 on each side, with the last ones bifurcated. The oral tentacles are digitate. The foot is elongate-oval, the bilabiate front end is rounded off and the hinder end comes to a blunt point.



Text-fig. 6.—*Geitodoris lutea*. A. Entire animal in dorsal view ($\times 1.3$); B. Elements of labial armature ($\times 380$); C. A half-row of radula ($\times 250$), a. innermost lateral teeth, b. outermost teeth.

The general colour of the back is yellow with a dark shade towards the centre. The tips of all the granules, and the rhinophore clavi are dark brown. The under side of the body is yellow.

The labial armature is an incomplete ring, consisting of irregular rodlets closely packed together. The radula contains about 17 rows of teeth, each half-row consisting of inner and outer teeth. The inner teeth, 22-26 in number, are simply hamate without denticles, and gradually increase in size outward. The outer teeth, about 20 in number, are spatular and closely gathered, the general appearance being thus brush-like. The hermaphrodite gland spreads over the liver.

Locality: Near Tomioka (June 1935; 2 sps.).

This species is closely allied to *Geitodoris capensis* BERGH in colours and radula type, but differs in having closely-set granules on the back.


Subfamily Discodoridinae

Genus *Peltodoris* BERGH, 1880*Peltodoris* BERGH, Mittlg. Zool. Stat. Neapel, Hft. 2, 1880, pp. 223-224.Type: *Peltodoris atromaculata* BERGH.82. *Peltodoris mauritiana* BERGH, 1889Genus *Thordisa* BERGH, 1877*Thordisa* BERGH, Malac. Unters., Hft. 12, 1877, p. 540.Type: *Thordisa maculigera* BERGH.83. *Thordisa amakusana* nov. sp.

(Pl. 1, fig. 12; text-fig. 7)

Distribution in Japan: Tomioka (Amakusa).

The animal, about 30 mm in length, is doridiform and flattened, and has the shape of an elongated ellipse with the ends equally rounded. The mantle is soft (spicules not well developed) and wide, and the whole upper surface is closely covered with minute



Text-fig. 7.—*Thordisa amakusana*. A half-row of radula ($\times 250$), a. innermost lateral teeth, b. outermost teeth.

villous papillae. The rhinophore sheath has a smooth margin. The branchial plumes, about 12 in number, are all simply pinnate and arranged in an almost complete circle around the anus. The margin of the branchial cavity is slightly elevated and is guarded by villous papillae. The oral tentacles are digitiform. The foot is elongate-oval; in front it is rounded off and bilabiate with a deep notch in the middle of the upper lip.

The back is orange-yellow with a dark shade over the viscera and with minute chocolate spots on the whole surface. The rhinophore clavi are chocolate and the branchiae pale orange-

yellow; the under surface of the body is yellowish; the brim of the foot bears a series of chocolate spots.

The cuticular labial disk is without armatures. The radula formula is $20 \times 22-25.0.22-25$. The majority of the lateral teeth are simply hamate, but the outermost 3-4 are much reduced with split tips.

Locality: Tomioka (Mar. 1935; 1 sp.).

The present species is especially characteristic in the coloration, simply pinnate branchiae and small radula formula. In colours it recalls *Thordisa hilaris* BERGH, but differs mainly in the configuration of the branchiae.

Genus *Discodoris* BERGH, 1877

Discodoris BERGH, Jahrb. Deutsch. Malakoz. Gesell., Bd. 4, 1877, p. 61.

Type: *Discodoris boholiensis* BERGH.

Synopsis of the Japanese species:

1. Body usually large (L. 40-80 mm). Radula large (40-60 laterals on each side).
 - a. Brown, back with large dark mottles, the darkest (3-4) ones on sides of back *concinna*.
 - b. Yellowish brown, back with indistinct brown shades. Also back often with reticulate figures due to peculiar arrangement of various granules *pardalis*.
2. Body usually small (L. 15-25 mm). Radula small (14-30 laterals on each side).
 - a. Branchiae 6. Back pale chocolate with dark mottles *yaeyamensis*.
 - b. Branchiae 9. Ashy yellow with a few dark flecks on back *pallida*.

84. *Discodoris concinna* (ALDER & HANCOCK, 1864)

85. *Discodoris pardalis* (ALDER & HANCOCK, 1864)

86. *Discodoris yaeyamensis* BABA, 1936

87. *Discodoris pallida* nov. sp.


(Pl. I, fig. 11; text-fig. 8)

Distribution in Japan: Tomioka (Amakusa).

The animal in life is soft and flattened, and measures about 15 mm in length. The back appears almost smooth, but is actually covered everywhere with villous papillae. The rhinophore sheath has a villous margin. The tripinnate branchial plumes, about 9 in number, are small and lie in a complete circle around the anus.

The margin of the branchial cavity is guarded by villous papillae. The oral tentacles are digitiform. The front edge of the foot is bilabiate with a deep median notch in the upper lip. Posteriorly the foot passes to a bluntly pointed tail.

The general colour of the body is an ashy yellow, becoming darker towards the median line. A small number of dark brown flecks are irregularly scattered over the back, the majority of them



Text-fig. 8.—*Discodoris pallida*. A. A half-row of radula ($\times 240$), a. innermost lateral teeth, b. outermost teeth; B. A pair of labial armatures ($\times 80$).

being in the mid-dorsal region. The tips of all the papillae are dark brown. The rhinophores and branchiae are of the same colour as the mantle. The under side of the body is uniformly ashy yellow.

The cuticular labial disk is armed with a pair of roughly oblong plates, each consisting of numerous tiny rods closely packed together. The radula contains 15 rows of teeth, each half-row being composed of 14-17 lateral teeth. The 1st lateral tooth is small and simply hamate. The succeeding teeth gradually increase in size outward, and decrease again to the edge. The outermost tooth is usually simple, sometimes split at the tip.

Locality: Tomioka (Oct. 1935; 1 sp.).

The present species closely resembles the Californian form *Discodoris heathi* MACFARLAND in colours, but differs from it in the type of the radula. In the latter species the radula is broad

with the formula $20 \times 36-42.0.36-42$, each half-row falling in two groups of different types.

Subfamily Homoeodoridinae

Genus *Homoeodoris* BERGH, 1881

Homoeodoris BERGH, Verh. k. k. zool.-bot. Gesell. Wien, Bd. 31, 1881, pp. 6 (222)-7 (223).

Homoeodoris THIELE, Handb. syst. Weicht., Th. 2, 1931, p. 438.

Type: *Homoeodoris japonica* BERGH.

88. *Homoeodoris japonica* BERGH, 1881

Subfamily Asteronotinae

Genus *Halgerda* BERGH, 1880

Halgerda BERGH, Verh. k. k. zool.-bot. Gesell. Wien, Bd. 30, 1880, p. 38 (190).

Type: *Halgerda formosa* BERGH.

Synopsis of the Japanese species:

- a. Back with prominent subconical tubercles interlinked by low ridges ... *graphica*.
 b. Back covered with fine villous papillae *japonica*.

89. *Halgerda graphica* BASEDOW & HEDLEY, 1905

90. *Halgerda japonica* ELIOT, 1913

Genus *Asteronotus* EHRENBERG, 1831

Asteronotus EHRENBERG, Symb. phys., 1831.

Type: *Asteronotus hemprichii* EHRENBERG.

91. *Asteronotus cespitosus* (VAN HASSELT, 1824)

Subfamily Arginae (=Platydoridinae)

Genus *Argus* BOHADSCH, 1761

Argus BOHADSCH, Anim. mar., 1761, p. 65.

Platydoris BERGH, Jahrb. Deutsch. Malakoz. Gesell., Bd. 4, 1877, p. 73.

Type: *Argus argo* BOHADSCH.

Synopsis of the Japanese species:

- A. Body large (L. 60-90 mm). Radula large (90-150 laterals on each side).
 1. Back with cloudy chocolate figures, under side orange-yellow with oval chocolate markings *speciosus*.

2. Back finely dotted with chocolate *tabulatus*.
 3. Back with wavy chocolate lines.
 a. Crimson blotches on back and under side... .. *cruentus*.
 b. No crimson blotches *striatus*.
 B. Body small (L. 25 mm). Radula small (50 laterals on each side). Yellowish, with chocolate mottles on back, mantle and branchial cavity margined with yellow *esakii*.

92. *Argus cruentus* (QUOY & GAIMARD, 1832)

93. *Argus striatus* (KELAART, 1858)

Doris striata KELAART, Journ. Asiatic Soc. Colombo, vol. 3, pt. 1, 1858.

Platydoris striata ELIOT, Journ. Coll. Sci. Imp. Univ. Tôkyô, vol. 35, art. 1, 1913, pp. 18-20.—Kataura.

94. *Argus speciosus* (ABRAHAM, 1877)

95. *Argus tabulatus* (ABRAHAM, 1877)

96. *Argus esakii* BABA, 1936

Genus *Doriopsis* PEASE, 1860

Doriopsis PEASE, Proc. Zool. Soc. London, 1860, p. 32.

Guyonia RISBEC, Faune Colon. Franç., tom. 2, 1928, p. 102.

Type: *Doriopsis granulosa* PEASE.

97. *Doriopsis viridis* PEASE, 1861

Doriopsis viridis PEASE, Proc. Zool. Soc. London, 1861, pp. 244-245.—Tahiti.

Doris pecten COLLINGWOOD, Trans. Linn. Soc. London, Zool., vol. 2, 1881, p. 126, pl. 9, figs. 1-5.—Bush Island (Keelung).

Subfamily Dendrodoridinae (=Doridopsinae)

Genus *Dendrodoris* EHRENBERG, 1831

Dendrodoris EHRENBERG, Symb. phys., 1831.

Doriopsis BERGH, Journ. Mus. Godeffroy, Hft. 8, 1875, pp. 82-87.

Doridopsis ALDER & HANCOCK, Trans. Zool. Soc. London, vol. 5, pt. 3, 1864, pp. 124-126.

Type: *Dendrodoris lugubris* EHRENBERG.

With two subgenera: *Dendrodoris* s. s. and *Doriopsilla*.

Synopsis of the Japanese species:

A. Body very soft (*Dendrodoris* s. s.).

1. Back entirely smooth.

a. Colours variable, usually black *nigra*.

- b. Yellow or orange yellow, the back with black irregular mottles
... .. *rubra* var. *nigromaculata*.
- c. Orange-yellow, the back with small, black, well defined mottles
... .. *guttata*.
- 2. Back sparsely granulate. Body exceedingly elongated. Dirty yellow, mottled
with dark brown *elongata*.
- 3. Back with warty compound protuberances.
 - a. Back dirty yellow with a large dark brown network ... *tuberculosa* var.
 - b. Back yellowish white with 3 longitudinal rows of dark brown areas, each
bearing blue spots *gemmacea*.
- B. Body spiculate, rather rigid (*Doriopsilla*). Orange-yellow, the back with a whitish
network *miniata*.

98. *Dendrodoris (Dendrodoris) tuberculosa*
(QUOY & GAIMARD, 1832), variety

99. *Dendrodoris (Dendrodoris) nigra* (STIMPSON, 1855)¹⁾

100. *Dendrodoris (Dendrodoris) gemmacea*
(ALDER & HANCOCK, 1864)
(Pl. I, fig. 4)

101. *Dendrodoris (Dendrodoris) rubra* var. *nigromaculata*
(ELIOT, 1913)

102. *Dendrodoris (Dendrodoris) guttata* (ODHNER, 1917)
(Pl. I, fig. 13)

103. *Dendrodoris (Dendrodoris) elongata* BABA, 1936

104. *Dendrodoris (Doriopsilla) miniata* (ALDER & HANCOCK, 1864)
(Pl. I, fig. 1)

Family Phyllididae

Genus *Phyllidia* CUVIER, 1796

Phyllidia CUVIER, Bull. Sci. Soc. Philom. Paris, 1796, p. 105.

Type: *Phyllidia varicosa* LAMARCK.

With 4 subgenera: *Phyllidia* s. s., *Phyllidiella*, *Phyllidiopsis*,
and *Ceratophyllidia*.

Synopsis of the Japanese species:

- a. Warts on back fused into (3) longitudinal and (many) short marginal ridges
... .. *varicosa*.

¹⁾ Synonym: *D. nigra* variety, see BABA (1935), p. 348.

- b. Warts simple, or composed of 2-4 tubercles *pustulosa*.
 c. Warts highly compound, often consisting of 5-16 tubercles *nobilis*.
 d. Warts mostly solitary but those in the mid-dorsal line fused to form a ridge
 *japonica* nom. nov.

105. *Phyllidia* (*Phyllidia*) *varicosa* LAMARCK, 1801

106. *Phyllidia* (*Phyllidiella*) *pustulosa* CUVIER, 1804

107. *Phyllidia* (*Phyllidiella*) *nobilis* (BERGH, 1869)

108. *Phyllidia* (*Phyllidiella*) *japonica* nom. nov.

Phyllidia tuberculata BABA, Venus, vol. 2, no. 3, 1930, pp. 117-118, pl. 4, fig. 1.—Tateyama Bay.

In 1930 *Phyllidia tuberculata* was proposed as a new species. But as the specific name *tuberculata* is already preoccupied by *P. tuberculata* RISBEC¹⁾, I must rename the present species *P. japonica* nom. nov.

Tribe. 2. Cladohepatica (=Aeolidiacea)

Family Duvauceliidae

Genus *Duvaucelia* RISSO, 1826

Duvaucelia RISSO, Hist. nat. Europe mérid., 4, 1826, p. 38.

Type: *Duvaucelia gracilis* RISSO.

With 5 subgenera: *Duvaucelia* s. s., *Candellista*, *Microlophus*, *Tritonidoxa* and *Sphaerostoma*.

Synopsis of the Japanese species:

- a. Back and sides smooth, vermilion-tinted *exsulans*.
 b. Back with traces of tubercles, sides smooth, whitish reticulation on back
 *reticulata*.

109. *Duvaucelia* (*Duvaucelia*) *reticulata* (BERGH, 1881)

Tritonia reticulata BERGH, Verh. k. k. zool.-bot. Gesell. Wien, Bd. 31, 1881, pp. 23 (239)-34 (250), pl. 8, figs. 7-20; pl. 9, figs. 1-12; pl. 10, figs. 1-10.—'M. japonicum'.

110. *Duvaucelia* (*Duvaucelia*) *exsulans* (BERGH, 1894)

(Text-fig. 9)


Tritonia exsulans BERGH, Bull. Mus. Comp. Zool. Harvard, vol. 25, no. 10, 1894, pp. 150-152, pl. 3, figs. 11-12; pl. 4, fig. 6.—Off California; O'Donoghue, Trans.

¹⁾ RISBEC (1928), p. 59.

Roy. Canad. Inst., vol. 13, pt. 1, 1921, pp. 152-154, pl. 7, figs. 4-6.—near Departure Bay.

Distribution in Japan: Sakhalin.

The animal is about 75 mm long. The frontal veil is expanded with a median notch, and bears on each side a series of 6-7 small conical papillae and on the ventral side a spoon-shaped, grooved tentacle at the lateral edge. The rhinophore sheath is elevated with an upper crenulate margin. The back is sharply marked off from sides by dorso-lateral ridges, each bearing 15-16 branchial tufts. The anal papilla lies about half way back on the right side of the body between the branchial tufts, 6 and 7; the genital



Text-fig. 9.—*Duvaucelia exsulans*. A. Entire animal in lateral view ($\times 0.8$); B. Masticatory edge of jaw-plate ($\times 90$); C. Jaw-plate ($\times 2$); D. Mouth parts ($\times 1.3$); E. A half-row of radula ($\times 70$), a. central tooth, b. 1st lateral tooth, c. 2nd tooth, d. 40th tooth, e. outermost teeth.

orifice is situated a short distance in front of the anus, beneath the 4th tuft. The nephroproct opens immediately above the anus. The back and sides appear to be quite smooth. The anterior end of the foot is abruptly rounded and bilabiate.

According to Mr. URITA the living animal was vermilion-tinted with whitish branchial tufts.

The masticatory process of the jaw-plate has several rows of imbricating scales. The radula formula is $65 \times 75-95.1.1.1.75-95$. The central tooth is broad and tricuspid. The 1st lateral tooth is clumsy with a small blunt spine, and is followed by a series of hamate lateral teeth reaching a maximum near the middle of the row. The liver is a large mass covered partially by the hermaphrodite gland. There are no stomachal plates.

Locality: Sakhalin (1 sp., coll. by Mr. T. URITA).

Genus *Tritoniopsilla* PRUVOT-FOL, 1933

Tritoniopsilla PRUVOT-FOL, Mém. Inst. Egypte, tom. 21, 1933, p. 108.

Type: *Tritoniopsis brucei* ELIOT.

111. *Tritoniopsilla tetraquetra* (PALLAS, 1788)¹⁾

Limax tetraquetra PALLAS, Nova Acta Petropolit., vol. 2, 1788, pp. 237, 239, pl. 5, fig. 22.
—'Mari Kurilas'.

Tritonia tetraquetra BERGH, Proc. Acad. Nat. Sci. Philadelphia, 1879, pp. 154 (98)-161 (105), pl. 3, figs. 13-16; pl. 4, figs. 5-12; pl. 5, figs. 1-2.—'Insulae Kurilae, Aleutianae'.

Genus *Marionia* VAYSSIÈRE, 1877

Marionia VAYSSIÈRE, Ann. Mag. Nat. Hist., ser. 4, vol. 20, 1877, pp. 367-368.

Type: *Marionia berghi* VAYSSIÈRE.

Synopsis of the Japanese species:

- a. Back without distinct polygons but with warts. Radula formula $38 \times 50.1.1.1.50$; masticatory process of jaw-plate with 4-5 rows of denticles *granularis*.
- b. Back with polygons and warts. Radula formula $52-70 \times 125-130.1.1.1.125-130$; masticatory process covered partially with irregular rudimentary armatures
... .. *pustulosa odhneri* nov. subsp.

112. *Marionia granularis* ODHNER, 1936

Marionia granularis ODHNER, Mém. Mus. Roy. Hist. Nat. Belgique, ser. 2, fasc. 3, 1936, p. 1090, pl., fig. 6; text-fig. 21.—Formosa Channel (20° 20' N., 121° 30' E.).


¹⁾ See ODHNER (1936), p. 1080.

113. *Marionia pustulosa odhneri* nov. subsp.

(Text-figs. 10-11)

Distribution in Japan: Momotori near Toba.

Several living specimens ranging from 40 to 80 mm in length. The frontal veil is two-lobed with a slight median notch, and bears on each side 8-10 small conical papillae and at its postero-lateral edge a grooved spoon-shaped tentacle. These papillae have each 1-3 accessory knobs mainly on the ventral side. The margin of the rhinophore sheath is crenulate. From 10 to 12 large branchial tufts are present on dorso-lateral ridges, each with a short stalk. The anal papilla lies on the right of the body between the branchial tufts 4 and 5, or 5 and 6; the genital orifice is situated beneath the 3rd or 4th tuft. The position of the nephroproct




Text-fig. 10.—*Marionia pustulosa odhneri*. A. Entire animal in lateral view ($\times 1$); B. Ventral view of head-veil; C. Polygons and warts of back; D-E. Right side of body to show local relations among anus, nephroproct and genital orifice, a. nephroproct, b. anus, c. genital orifice.

differs in different individuals. In some specimens this opening is found above and in front of, and in others just in front of, the anus. The whole upper surface of the body (back, sides, frontal veil,

branchial stalks and rhinophore sheaths) is closely covered with granular tubercles which are apt to fuse into compound warts on the back and sides. Dorsally the warts and tubercles are interlined with the so-called indistinct polygons.

The back and frontal veil are olive green covered with a fine purple reticulation. In addition there is a pale broad median band from the head to the tip of the tail, sending off similar lateral bands to the bases of the branchial tufts. The sides of the body and the foot are olive green.

The jaw-plate has a smooth cutting edge. The surface of the



Text-fig. 11.—*Marionia pustulosa odhneri*. A. Jaw-plate ($\times 3$); B. Masticatory process ($\times 30$); C. Scales on masticatory process ($\times 300$); D. General aspect of digestive organs, a. pharyngeal bulb, b. oesophagus, c. intestine, d. posterior liver, e. stomach, f. anterior liver; E. A half-row of radula ($\times 70$), a. central tooth, b. 1st lateral tooth, c. succeeding teeth, d. outermost teeth; F. Stomachal plates ($\times 10$).

masticatory process appears to be almost even, but a closer examination reveals that it is partially covered with irregular, almost rudimentary armatures, the outermost being spiny and the rest scale-like. The radulae in two specimens dissected contain 52-70 rows of teeth, each half-row consisting of exceedingly numerous (125-130) lateral teeth. The central tooth consist of an approximately rectangular plate bearing 3 large cusps, of which the median is the largest. The 1st lateral tooth differs in form from the succeeding teeth which are all simply hamate without denticles. The stomach is armed with a girdle of about 28 horn-yellow plates. The liver consists of a small anterior and a large posterior lobe.

Locality: Momotori near Toba (July 1931; fairly common).

The present form agrees quite well with the Queensland species, *Marionia pustulosa* ODHNER, in all the essential characters (general external aspects, type of jaws and radula), but differs slightly in the possession of well-developed stomachal plates, having rhinophore sheaths with a crenulate edge and smaller dorsal polygons. It is recorded here as a subspecies (or variety?) of the above-said species, under the name *M. pustulosa odhneri* nov. subsp.

Genus *Marioniopsis* ODHNER, 1934

Marioniopsis ODHNER, "Terra Nova" Exped., Zool., vol. 7, no. 5, 1934, p. 286.

Type: *Tritonia cyanobranchiata* RÜPPELL & LEUCKART.

114. *Marioniopsis ? babai* ODHNER, 1936

Duvaucelia irrorata BABA, Annot. Zool. Japon., vol. 14, no. 2, 1933, pp. 274-275, fig. 1.
—Tomioka.

Marioniopsis babai ODHNER, Mém. Mus. Roy. Hist. Nat. Belgique, ser. 2, fasc. 3, 1936, p. 1085, 1087.

The main characters are those previously given, to which may be added: on each latero-ventral edge of the frontal veil is borne a grooved spoon-like tentacle; the genital orifice lies on the right body side below the 2nd, and the anus below the 4th branchial tuft; the inner wall of the stomach is armed with a girdle of whitish chitinous plates.

As is suggested by Dr. ODHNER, a further study on jaws and liver system is needed before the present form is satisfactorily placed within the genus *Marioniopsis*.

Family Arminidae (=Pleurophyllidiidae)

Subfamily Dermatobranchinae

Genus *Dermatobranchus* VAN HASSELT, 1824*Dermatobranchus* VAN HASSELT, Allgem. Konst en Letter-Bode, 1824, p. 37.*Pleuroleura* BERGH, Malac. Unters., Hft. 6, 1874, p. 277.Type: *Dermatobranchus striatus* VAN HASSELT.115. *Dermatobranchus striatus* VAN HASSELT, 1824


(Pl. 2, fig. 1; text-fig. 12)

Dermatobranchus striatus VAN HASSELT, Allgem. Konst en Letter-Bode, 1824, pp. 37-38.

—Java; BERGH, Notes Leyden Mus., vol. 9, 1887, pp. 311-312, pl. 6, figs. 1-2.

Pleuroleura striata BERGH, Zool. Jahrb., Syst., Bd. 3, 1888, p. 362; ELIOT, Journ. Coll. Sci. Imp. Univ. Tôkyô, vol. 35, art. 1, 1913, pp. 41-42.—Misaki.*Distribution in Japan*: Tomioka (Amakusa), Seto (Kii), Misaki and Tateyama.

The animal measures 20-30 mm in length. The head is expanded in a smooth semicircular veil, behind which is a sort of neck bearing a pair of approximating rhinophores. It continues



Text-fig. 12.—*Dermatobranchus striatus*. A. Jaw-plate ($\times 50$); B. A half-row of radula ($\times 470$), a. central tooth, b. 1st lateral tooth, c. 2nd tooth, d. outermost teeth.

back into a tongue-shaped mantle without line of demarcation. The rhinophore is a non-retractile cylindrical structure with a longitudinally perfoliate clavus. Dorsally the mantle is marked with (about 18) low longitudinal ridges, while on the lower extreme margin there is a series of cnidophores. The sides of

the body are quite smooth and are free from branchial ridges. The genital orifice lies on the right body side immediately behind the head-veil. The anus opens a short distance behind the genital orifice. The foot is lanceolate, often extending beyond the mantle all around. The front end is broad and continued out laterally into slight obtuse expansions. A median longitudinal furrow runs down on the posterior two-thirds of the pedal sole.

The ground-colour of the body is a fleshy white. The back is marked with fine, dark, continuous or discontinuous lines running along the grooves among the longitudinal ridges; with irregularly scattered black dots defined by pale chocolate ocelli; with a dark semilunar band across the level of the anus. The upper half of the rhinophore is vivid orange, and the lower half white. The head-veil, sides, and foot are of a uniform fleshy white without markings.

The jaw-plate has a produced, horn-like masticatory process, and the edge is armed with several rows of spiny denticles. The radula formulae in two specimens are $30 \times 15.1.1.1.15$, and $34 \times 10-13.1.1.1.10-13$. The central tooth is wide with a series (13-17) of denticles on either side of a large median cusp. The 1st lateral tooth is like a half of the central tooth, bearing 15-17 denticles. The succeeding teeth are all simply hamate without denticles.

Localities: Tomioka (Feb. 1935; 2 sps.), Seto (Mar. 1937; 1 sp.) and Tateyama (May 1929; several sps.).

The present species is distinguished from the other members of *Dermatobranchus* mainly by the folded mantle, peculiar coloration and type of the radula.

Subfamily Armininae

Genus *Armina* RAFINESQUE, 1814

Armina RAFINESQUE, Précis découv. somiol., 1814, p. 30.

Pleurophyllidia MECKEL, MECKELS Deutsch. Arch. Physiol., Bd. 8, 1823, p. 197.

Type: *Armina tigrina* RAFINESQUE.

With 3 subgenera, viz., *Armina* s. s., *Linguella* and *Camarga*.

Synopsis of the Japanese species:

- I. Head-veil-separated from mantle by a clear boundary; rhinophores close together (*Armina* s. s.).

- A. Head-veil papillated.
 - a. Papillae few (11-14); a single side-lamella on each side ... *comta*.
 - b. Papillae many (50-80); side-lamellae 4-7 on each side ... *papillata*.
- B. Head-veil smooth.
 - 1. Side-lamellae few, in longitudinal rows.
 - a. Number of lateral teeth moderate (40-50); 1st and succeeding inner teeth denticulate; side-lamellae 3-5 on each side ... *japonica*.
 - b. Number of lateral teeth exceedingly large (231); 1st lateral tooth denticulate, succeeding teeth smooth; side-lamellae 8 on each side ... *similis*.
 - 2. Side-lamellae many (about 50), in oblique rows... *taeniolata*.
- II. No clear boundary between head-veil and mantle; rhinophores widely separated (*Linguella*).
 - A. Number of lateral teeth exceedingly large (120-140) ... *lairra*.
 - B. Number of lateral teeth moderate (22-50).
 - 1. Back smooth ... *fallax*.
 - 2. Back granulated.
 - a. Mantle-flange with two colossal appendages ... *variolosa*.
 - b. Mantle-flange without appendages... *babai*.

116. *Armina (Armina) taeniolata* (BERGH, 1866)

(Text-fig. 13)


Pleurophyllidia taeniolata BERGH, Naturh. Tidsskr., ser. 3, vol. 4, 1866, pp. 42-46, pl. 5, figs. 1-22; pl. 6, figs. 1-9.—Mozambique; ELIOT, Journ. Conch., London, vol. 11, no. 8, 1905, pp. 239-240, pl. 5, fig. 1.—Maskat; ELIOT, Proc. Zool. Soc. London, 1906, pt. 2, pp. 679-680.—India.

Distribution in Japan: Kii and Tateyama.

The animal, preserved, measures about 60 mm in length. The head veil is smooth above. It is marked off from the mantle by a clear boundary. The rhinophores lie close together as usual. Dorsally the mantle is marked with (about 30-35) longitudinal ridges. The anterior gills consist of delicate longitudinal folds, some of which are continuous behind with the side-lamellae. These latter are in the form of many (about 50) short ridges arranged in oblique rows and extending from the anterior gill-folds to the posterior end of the mantle. The genital orifice opens on the right side below the anterior gill-folds. The anus is situated about half way back on the same side. The nephroproct lies about midway between genital orifice and anus. The foot has a median ventral furrow.

According to a coloured sketch by Mr. OKAMOTO, the head-shield is black bordered with yellow; the upper side of the mantle

is black ornamented with many whitish longitudinal lines (ridges) and a yellow border; the under side of the mantle and the side-lamellae are brown; the foot is purplish black.



Text-fig. 13.—*Armina taeniolata*. A. Lateral view of entire animal ($\times 1$); B. Jaw-plate ($\times 6$); C. A half-row of radula ($\times 90$), a. central tooth, b. 1st lateral tooth, c. outermost teeth.

The masticatory edge of the jaw-plate is covered with several rows of scale-like armatures. The radula formula for the Kii specimen is about $35 \times 55.1.1.1.55$. The central tooth is broad with about 9 denticles on each side of a large median cusp. The 1st lateral tooth has a rectangular base, which gives off an inner claw-like spine and bears a few (1-2) denticles (or sometimes these are missing altogether). The succeeding teeth excepting several (20) outermost are all hamate with a few (5-8), indistinct denticles.

The radula formula for the Tateyama specimen is about $55 \times 45.1.1.1.45$. The central tooth is broad with 5-6 denticles on each side of a median spine. The 1st lateral tooth differs in form from the succeeding teeth. All the lateral teeth are without denticles.

Localities: Kii (May 1933; 1 sp. coll. by Mr. K. OKAMOTO) and Tateyama (1 sp.).

The above description agrees well with those given by BERGH and ELIOT for *Pleurophyllidia taeniolata*. This species appears to be especially characteristic in the oblique arrangement of the side-lamellae and in the type of the radula.

117. *Armina (Armina) comta* (BERGH, 1880)

118. *Armina (Armina) japonica* (ELIOT, 1913)

119. *Armina (Armina) similis* (ELIOT, 1913)

Pleurophyllidia similis ELIOT, Journ. Coll. Sci. Imp. Univ. Tôkyô, vol. 35, art. 1, 1913, pp. 39-40.—Misaki.

120. *Armina (Armina) papillata* (BABA, 1933)

121. *Armina (Linguella) iaira* (BERGH, 1865)

Sancara iaira BERGH, Vidensk. Meddel. Naturh. Foren., (1864), 1865, pp. 178-194, pl. 3; BERGH, Naturh. Tidsskr., ser. 3, vol. 4, 1866, pp. 64-68.—Japan.

Linguella iaira BERGH, Malac. Unters., Hft. 6, 1874, p. 268, pl. 34, figs. 25-26.

122. *Armina (Linguella) fallax* (BERGH, 1880)

Linguella fallax BERGH, Verh. k. k. zool.-bot. Gesell. Wien, Bd. 30, 1880, pp. 25 (177)-28 (180), pl. 2, figs. 13-15; pl. 3, fig. 13; pl. 4, figs. 4-7.—Enoshima.

123. *Armina (Linguella) variolosa* (BERGH, 1904)

Linguella varicosa BERGH, SEMPER'S Reisen, Bd. 9, Th. 6, Lief. 1, 1904, pp. 21-24.—Tonsang Harbour (China); ELIOT, Journ. Coll. Sci. Imp. Univ. Tôkyô, vol. 35, art. 1, 1913, pp. 40-41.—Minatomura.

124. *Armina (Linguella) babai* (TCHANG, 1936)

Linguella babai TCHANG, Contrib. Inst. Zool. Nat. Acad. Peiping, vol. 2, no. 2, 1936, pp. 100-120, 146-147, figs. 47-55; pl. 2, figs. 7-10; pl. 3, fig. 12; pl. 15.—Tsingtao.

Locality: Momotori near Toba (July 1931; 3 sps.).

Family Dironidae

Genus *Dirona* COCKERELL & ELIOT, 1905

Dirona COCKERELL & ELIOT, Journ. Malac., vol. 12, no. 3, 1905, pp. 45-46.

Type: *Dirona picta* COCKERELL & ELIOT.

125. *Dirona albolineata* MACFARLAND, 1912

Family Bornellidae

Genus *Bornella* GRAY, 1850*Bornella* GRAY, Figs. Mollusc. Anim., vol. 4, 1850, p. 107.Type: *Bornella adamsii* GRAY.126. *Bornella digitata* ADAMS & REEVE, 1850Genus *Pseudobornella* BABA, 1932*Pseudobornella* BABA, Annot. Zool. Japon., vol. 13, no. 4, 1932, pp. 369-370.Type: *Pseudobornella orientalis* BABA.127. *Pseudobornella orientalis* BABA, 1932

Family Scyllaeidae


Genus *Scyllaea* LINNÉ, 1758*Scyllaea* LINNÉ, Syst. Nat., ed. 10, 1758, p. 656.Type: *Scyllaea pelagica* LINNÉ.128. *Scyllaea bicolor* BERGH, 1880*Scyllaea bicolor* BERGH, Verh. k. k. zool.-bot. Gesell. Wien, Bd. 30, 1880, pp. 15 (167)-20 (172), pl. 1, figs. 12-17; pl. 3, figs. 3-6.—Enoshima; ELIOT, Journ. Coll. Sci. Imp. Univ. Tôkyô, vol. 35, art. 1, 1913, p. 34.—Misaki.Genus *Notobryon* ODHNER, 1936*Notobryon* ODHNER, Mém. Mus. Roy. Hist. Nat. Belgique, ser. 2, fasc. 3, 1936, p. 1099.Type: *Notobryon wardi* ODHNER.129. *Notobryon wardi* ODHNER, 1936

(Pl. 2, fig. 3; text-fig. 14)

Notobryon wardi ODHNER, Mém. Mus. Roy. Hist. Nat. Belgique, ser. 2, fasc. 3, 1936, pp. 1099-1103, pl., figs. 1-3; text-figs. 31 b, 32-38.—Queensland.*Distribution in Japan*: Tomioka (Amakusa) and Seto (Kii).

The body is elongated and limaciform, ending posteriorly in a blunt tail. It measures about 60 mm in the living state. The head is small, but is expanded ventrally in a semicircular veil with a wavy margin. The clavus of the rhinophore is small and conical,

and bears oblique folds. It is retractile within a stout elevated sheath which terminates distally in a slightly crenulate margin. There is a longitudinal crest or flange along the posterior margin of the sheath. The dorso-lateral margins of the body are produced



Text-fig. 14.—*Notobryon wardi*. A. General aspect of digestive organs ($\times 2$), a. pharyngeal bulb, b. right anterior liver-lobe, c. intestine, d. hermaphrodite gland, e. posterior liver-lobe, f. left anterior liver-lobe; B. Jaw-plate ($\times 10$); C. Scales on masticatory process ($\times 400$); D. A half-row of radula ($\times 140$), a. 1st lateral tooth, b. 2nd tooth, c. 10th tooth, d. outermost teeth.

into two pairs of large foliaceous lobes. The edges of these lobes are indistinctly crenulate, and the tips are rolled to form a sort of short upstanding siphons. The branchial tufts, 4 on each dorso-lateral lobe and one at the hinder end of the mantle, are long cylindrical processes which give off branches, themselves dividing into smaller ramose twigs. The back is smooth. The neck has

on each side a small conical papilla. The sides are fairly smooth though there is a row of (10-12) conical papillae of various sizes extending from the neck to the tail. The tail has a mid-dorsal crest. The foot is narrow but is capable of expansion. It is rounded in front and continued into a blunt tail posteriorly. The anus opens on the right side between the dorso-lateral lobes. The genital orifice lies on the same side a short distance behind the level of the rhinophore sheath.

The general ground-colour of the body is a beautiful saffron-yellow with deeper dots and mottles. An olive colour predominates on the rhinophore sheaths, dorso-lateral margins, tail crest, and dorso-lateral lobes. Dorsally there is a bold opaque marking at the base of the dorso-lateral lobe. The papillae on the neck and sides are opaque white. The rhinophore clavi light saffron-yellow; the anal papilla deep red; the branchial plumes colourless; the foot saffron-yellow.

The jaw-plates are shell-shaped and of a horn-yellow colour. The masticatory edge is produced dorsally into a wide flange with a recurved border. This flange is covered everywhere with fine close-fit scales. The radula formulae in two specimens are $13 \times 14-20.0.14-20$, and $15 \times 14-21.0.14-21$, respectively. The older end of the radula is black-tinted while the newer end is colourless. In both radulae the rachis is wide and bare, without a central tooth. The 1st lateral tooth is the smallest and is sometimes degraded. Typically it takes the form of an oblong plate which has a short claw-like spine and an outer row of (3-4) denticles. The next lateral tooth has (8-12) inner and (about 5) outer denticles. The succeeding teeth present the same general form, getting larger towards the middle of the half-row. The denticles are decidedly smaller on the inner margin than on the outer. At the extreme end of the row the teeth decrease considerably in size. The stomach has a girdle of black chitinous plates, each being broadly triangular in form with a mammiliform apex. The liver, deep blue black in colour, is divided into two anterior and one posterior lobes, the last of them being the largest. The hermaphrodite gland consists of two large grayish lobes.

Locality: Tomioka (May 1936; 2 sps.) and Seto (Mar. 1937; 1 sp.).

I have been able to identify the present specimens as *Notobryon*

wardi. They agree with that species in the general body form and in the type of the radula, liver and hermaphrodite gland.

Genus *Crosslandia* ELIOT, 1902

Crosslandia ELIOT, Proc. Zool. Soc. London, 1902, pt. 2, pp. 64-68.

Type: *Crosslandia viridis* ELIOT.

130. *Crosslandia viridis* ELIOT, 1902

(Pl. 1, fig. 7; text-fig. 15)

Crosslandia viridis ELIOT, Proc. Zool. Soc. London, 1902, pt. 2, pp. 64-68, pl. 5, figs. 1-2, 4-8; text-figs. 2-4.—Zanzibar.

Crosslandia fusca ELIOT, ditto, p. 68, pl. 5, fig. 3.—Zanzibar; ELIOT, Journ. Linn. Soc. London, Zool., vol. 31, 1908, pp. 90-94, figs. a-c.—Red Sea.

Crosslandia orientalis THIELE, Tiefsee-Exped., 1925, p. 287, pl. 33, fig. 20.—near Zanzibar.


Distribution in Japan: Tomioka (Amakusa).

The living animal measures about 30 mm in length. It is produced into a long neck and tail, and laterally into two foliaceous lobes, one on each side. The lobe itself has a posterior notch down the tip, showing a sign of subdivision into two halves, the anterior being a little larger than the posterior. The head is small with a triangular veil which has ventrally a slit-like mouth. The rhinophore is small, conical and perfoliated. It is entirely retractile within an elevated sheath whose upper margin is wavy. There is a longitudinal crest along the posterior margin of the sheath. Small conical papillae are borne on the dorso-lateral margins and on the sides of the body. The branchial tufts, about 40 in number, are small and ramose and are irregularly scattered over the inner surface of the foliaceous lobes and on the back between and behind these lobes. The tail has a mid-dorsal crest. The genital orifice lies on the right side immediately behind the rhinophore sheath. The anus is situated on the same side just beneath the foliaceous lobe. The foot is small with an abruptly rounded anterior edge.

The upper side of the body is of a translucent yellowish brown sparsely maculated with white, and is ornamented with additional blue markings. Of these latter, one is on the rhinophore sheath, two on the mid-dorsal line, about 15 along the margin of the foliaceous lobe, and 3-4 on the side of the body.

The jaw-plates are shallow shell-shaped. At the dorsal edge it

is produced into a wide masticatory process with a recurved border. This process is covered as usual with fine, closely approximated scales. The radula formula is $18 \times 16 \cdot 22 \cdot 1 \cdot 16 \cdot 22$. The central tooth is small and consists of an oblong base bearing a



Text-fig. 15.—*Crosslandia viridis*. A. Mouth-parts; B. General aspect of digestive organs ($\times 4$), a. oesophagus, b. intestine, c. hermaphrodite glands, d. liver-lobes; C. A half-row of radula ($\times 200$), a. central tooth, b. innermost lateral teeth, c. outermost teeth; D. Jaw-plate ($\times 18$); E. Scales on masticatory process ($\times 470$).

series of 4-6 denticles on both sides of a small median cusp. The 1st lateral tooth is plate-like with 3-5 denticles on both sides of a small median cusp. The succeeding teeth do not differ much in shape from the 1st tooth, but get larger as they pass outward. At the extreme end of the row the teeth decrease in size, becoming roughly hamate with irregular flanking denticles.

The stomach is armed with a girdle of chitinous plates which are colourless and triangular in shape. The liver is in two large masses, anterior and posterior. The hermaphrodite glands consist of 3 spherical masses.

Locality: Tomioka (Feb. 1935; 1 sp.).

In 1902 ELIOT created two species, *Crosslandia viridis* and *C. fusca*, which, however, were subsequently regarded by him (1908) as varieties of a single species. In 1929 O'DONOGHUE accepted *viridis* as the type of the genus *Crosslandia* and *fusca* as a synonym. *C. viridis* ELIOT is known to be subject to a vast range of colour variation, from brown to green, and to this species, in particular to its brown form, is referable the present Japanese specimen.

Family Phylliroidae

Genus *Phylliroe* PÉRON & LESUEUR, 1810

Phylliroe PÉRON & LESUEUR, Ann. Mus. Hist. Nat., tom. 15, 1810, p. 65.

Type: *Phylliroe bucephala* PÉRON & LESUEUR.

131. *Phylliroe bucephala* PÉRON & LESUEUR, 1810

Phylliroe bucephala PÉRON & LESUEUR, Ann. Mus. Hist. Nat., tom. 15, 1810, p. 65, pl. 2, figs. 1-3; ELIOT, Journ. Coll. Sci. Imp. Univ. Tôkyô, vol. 35, art. 1, 1913, p. 34.

—Bonin Islands.

Genus *Cephalopyge* HANEL, 1905

Cephalopyge HANEL, Zool. Jahrb., Syst., Bd. 21, 1905, pp. 451-462.

Type: *Phylliroe trematoides* CHUN.

132. *Cephalopyge orientalis* BABA, 1933

Family Fimbriidae (=Tethyidae, Tethymelibidae)

Genus *Melibe* RANG, 1829

Melibe RANG, Manuel Mollusq., 1829, p. 129.

Type: *Milibe rosea* RANG.

Synopsis of the Japanese species:

- A. Foot very much expanded, hood large, body enormous (250-500 mm long)
 - *japonica*.
- B. Foot very narrow, hood of moderate size, body not large.
 - 1. Cirri of hood-margin few, in 1-3 rows; body small, pale yellow with sparsely set brownish mottles *pilosa*.
 - 2. Cirri of hood-margin many, in 5-10 rows.
 - a. Lip reddish brown, body reddish brown with dark shades and mottles *pilosa*.
 - b. Lip not reddish brown, body yellowish brown... .. *vexillifera*.

133. *Melibe pilosa* PEASE, 1860

134. *Melibe papillosa* (DE FILIPPI, 1867)

135. *Melibe vexillifera* BERGH, 1880

Melibe vexillifera BERGH, Verh. k. k. zool.-bot. Gesell. Wien, Bd. 30, 1880, pp. 10 (162)-13 (165), pl. 2, figs. 1-11; pl. 3, figs. 1-2.—Enoshima.

Melibe vexillifera? ELIOT, Journ. Coll. Sci. Imp. Univ. Tôkyô, vol. 35, art. 1, 1913, p. 37, pl. 2, fig. 11; HIRASÉ, Moluskoj, 1927, p. 1475, fig. 2837.—Misaki.

136. *Melibe japonica* ELIOT, 1913

Family Dotonidae

Genus *Doto* OKEN, 1815

Doto OKEN, Lehrb. Naturg., Th. 3, Zool., pt. 1, 1815, pp. 10, 278.

Type: *Doris maculata* MONTAGU.

137. *Doto japonica* ODHNER, 1936

Doto japonica ODHNER, Mém. Mus. Roy. Hist. Nat. Belgique, ser. 2, fasc. 3, 1936, pp. 1121-1122, pl., figs. 11-16; text-fig. 47.—Moroiso near Misaki.

Family Madrellidae

Genus *Madrella* ALDER & HANCOCK, 1864

Madrella ALDER & HANCOCK, Trans. Zool. Soc. London, vol. 5, pt. 3, 1864, pp. 141-142.

Type: *Madrella ferruginosa* ALDER & HANCOCK.

138. *Madrella sanguinea* (ANGAS, 1864)

(Pl. 1, fig. 2)

Family Flabellinidae

Genus *Coryphella* GRAY, 1850

Coryphella GRAY, Figs. Mollusc. Anim., vol. 4, 1850, p. 109.

Type: *Eolis rufibranchialis* JOHNSTON.

Synopsis of the Japanese species:

- a. Branchial papillae many, in (30) oblique rows on each side *athadona*.
 b. Branchial papillae very few, 6-7 groups on each side, each group containing from 1 to 2 of them *ornata*.

139. *Coryphella alderi* ADAMS, 1861

Coryphella alderi ADAMS, Ann. Mag. Nat. Hist., ser. 3, vol. 8, 1861, p. 140.—Off Fuku-yama, Tsugaru Strait.

140. *Coryphella athadona* BERGH, 1875

141. *Coryphella ornata* RISBEC, 1928

Family Tergipedidae

Genus *Cuthona* ALDER & HANCOCK, 1855

Cuthona ALDER & HANCOCK, Monogr. Brit. Nudib., pt. 7, 1855, appendix, p. 22.

Type: *Eolis nana* ALDER & HANCOCK.

Cuthona, with *Cratena*, *Amphorina* (of some authors), *Hervia* and *Rizzolia* as synonyms, may possibly be classified as follows:

Subgenus *Cuthona* s. s. (+ *Cratena*, *Amphorina* spp. of some authors). Antero-lateral corners of foot rounded, or slightly angulated.

Subgenus *Hervia* BERGH, 1871. (+ *Rizzolia*). Antero-lateral corners of foot tentaculiform.

Synopsis of the Japanese species belonging to the subgenus Cuthona:

- a. Branchial rows many (15-19); teeth numerous (70-82); each comb-like with 8-10 denticles in front, median spine indistinct. Body orange-yellow, branchial papillae orange-yellow with a dark blue vein and a white tip *bicolor*.
- b. Branchial rows 8-9; teeth few (15), each horseshoe-shaped with a prominent median cusp and 5-6 flanking denticles. Back, rhinophores and oral tentacles black-spotted, branchial veins pale green *yatsui*.

142. *Cuthona (Cuthona) bicolor* BERGH, 1904

143. *Cuthona (Cuthona) sp.* ELIOT, 1913

Amphorina? sp. ELIOT, Journ. Coll. Sci. Imp. Univ. Tôkyô, vol. 35, art. 1, 1913, p. 43.
—Misaki

144. *Cuthona (Cuthona) yatsui* (BABA, 1930)

(Pl. 2, fig. 2)

145. *Cuthona (Cuthona) sp.* BABA, 1935

Synopsis of the Japanese species belonging to the subgenus Hervia:

1. Branchial papillae in many (9-17) oblique rows.
 - a. Genital orifice below the 4th right branchial row. Two chocolate lines on each side of body *quadrilineata*.
 - b. Genital orifice below the 2nd right branchial row. Body orange-yellow, branchial papillae yellow with a cobalt-blue vein *ornata* nov. sp.
2. Branchial papillae in (4-8) groups and on horseshoe-shaped bases, the 1st group being widely separated from the rest.
 - a. Genital orifice below the 2nd right branchial row. Branchial groups from

- 2nd to last moderately close together. Body yellowish, branchial veins reddish brown, yellowish brown, deep orange yellow or chrome yellow
 *ceylonica*.
- b. Genital orifice below the 2nd right branchial row. Branchial groups from 3rd to last very close together and moderately separated from the 2nd. Body white, branchial veins reddish brown *enosimensis*.
- c. Genital orifice below and between the 1st and 2nd right branchial groups. Tail very long *japonica* nom. nov.
- d. Genital orifice below the 1st right branchial row. Back and sides with blue and vermilion lines *emurai* nov. sp.

146. ? *Cuthona (Hervia) rosea* (BERGH, 1888)

Hervia rosea? ELIOT, Journ. Coll. Sci. Imp. Univ. Tôkyô, vol. 35, art. 1, 1913, pp. 42-43.—Misaki.

147. *Cuthona (Hervia) ceylonica* (FARRAN, 1905)148. *Cuthona (Hervia) enosimensis* (BABA, 1930)

In my previous paper (BABA 1930, pp. 118-119) the branchial papillae have been described to be in 3 groups on each side. The reexamination of the original specimen reveals that the apparent 3rd group consists of 2-3 small, closely approximating groups, each lying upon a horseshoe-shaped base. In all, therefore, the branchial groups are 4 on the left and 5 on the right side.

149. *Cuthona (Hervia) quadrilineata* (BABA, 1930)150. *Cuthona (Hervia) quadrilineata* (BABA, 1930), variety151. *Cuthona (Hervia) japonica* nom. nov.

Rizzolia modesta BERGH, Verh. k. k. zool.-bot. Gesell. Wien, Bd. 30, 1880, p. 4 (156)-8 (160), pl. I, figs. 1-11.—Enoshima; ? ELIOT, Proc. Malac. Soc. London, vol. 6, pt. 4, 1905, pp. 230-231.—Inland Sea.

Rizzolia modesta of BERGH appears to be a distinct species of the genus *Cuthona*. I propose to call it as above because the name *modesta* is preoccupied (*Hervia modesta* BERGH, 1874).


152. *Cuthona (Hervia) emurai* nov. sp.

(Text-fig. 16)

Distribution in Japan: Niigata (Japan Sea).

Several specimens accompanied by a coloured sketch have

been sent for identification by Prof. EMURA of Niigata Kôtô-Gakkô. The body in life ranges from 10 to 35 mm in length. The antero-lateral corners of the head are continued out into two strong pointed oral tentacles. The rhinophores are shorter, being constricted along the whole length but not perfoliated. The branchial papillae are fusiform, very deciduous and arranged superficially in 3 tufts, but actually in 5-6 groups on each side of the body. Each group lies upon a roughly horseshoe-shaped base opening outward. The 1st group commences postero-lateral to the rhinophore, contains over 50 papillae, and is widely separated from the 2nd. The genital opening lies immediately below the 1st group on the right side. The 2nd group, with nearly 50 papillae, is as large as the 1st,



Text-fig. 16.—*Cuthona emurai*. A. Entire animal ($\times 2$); B. Tooth in dorsal view ($\times 60$); C. The same in lateral view; D. Jaw-plate ($\times 10$). The original sketch of the animal has been made from life by Prof. EMURA.

and on the right side it surrounds the anal papilla. The succeeding groups are close together, and decrease in size as they pass backwards. In the last group the papillae are crowded irregularly so as to cover the posterior extremity of the back. The foot

is bilabiate with a notch in the middle of the upper lip, and is continued out laterally into two well-marked tentaculiform processes.

The ground-colour of the body is of a pale (fleshy) yellow. Along the mid-dorsal region there run two bluish bilateral lines which pass forward and run right up the rhinophores and oral tentacles; posteriorly they converge to the tip of the tail. A broken mid-dorsal vermilion line runs about half way down from the head. The sides of the body are each marked with two lines running parallel with each other, the upper bluish and the lower shorter and vermilion. The branchial papillae are chocolate-coloured with usually a white vein and a vermilion marking immediately below the whitish tip, sometimes a white broken vein running up to the tip. The antero-lateral tentaculiform processes of the foot are each marked with a bluish line.

The cutting edge of the jaw-plate is continued out in a strong pointed masticatory process bearing a series of (55) denticles. The radula is uniseriate with about 20 teeth and is formulated as $20 \times 0.1.0$. Each tooth is of a characteristic horseshoe-shape. The apex is marked by a strong median spine flanked by 1-2 small denticles near the base; ventrally and distally the spine bears a series of (6-8) almost indistinct denticles. On each side of the median spine there runs a row of sharp denticles, the latter varying from 6 to 7 in number.

Locality: Niigata (Apr. 1934; fairly ~~common~~).

This is a new species characteristic especially of colours, arrangement of the branchiae, and position of the genital orifice.


153. *Cuthona (Hervia) ornata* nov. sp.

(Pl. 2, fig. 4; text-fig. 17)

Distribution in Japan: Tomioka (Amakusa).

The body in life ranges from 8 to 12 mm in length. The antero-lateral corners of the head are continued out into two slender oral tentacles. The rhinophores are a little longer, and are simply cylindrico-conical without perfoliations. The margin of the back bears a small number of branchial papillae in 9 transverse rows, the largest being on the inside and the smallest outside. They are small, short and roughly fusiform, and are about 2 in the 1st, 4 in the 2nd, 5 in the 3rd and 4th, 4 in the 5th and 6th, 3 in the 7th and 8th, and 1 in the 9th row. The end of the body extends

in a long tail behind the level of the last rows. The anal papilla lies between the 4th and 5th rows on the right side. The genital orifice is situated below the 2nd row on the same side. The foot



Text-fig. 17.—*Cuthona ornata*. A. Tooth ($\times 230$); B. Jaw-plate ($\times 70$).

is fairly narrow; the front edge is rounded off and its corners are produced outward into two horn-like processes.

The general colour of the head, back and sides is orange-yellow and this also is true of the basal half of the rhinophores and oral tentacles. The branchial papillae are yellow with a cobalt-blue vein which terminates distally in an opaque yellow marking. The upper half of the rhinophores and oral tentacles are yellow.

The jaw-plates are shell-shaped with a short masticatory process. The cutting edge probably bears a series of denticles which are now worn jagged. The uniseriate radula consists of 55 teeth and is formulated as $55 \times 0.1.0$. The tooth is roughly horseshoe-shaped. In front it is marked by a fairly stout median spine rising far back, on each side of which runs a row of 5-7 sharp denticles, some of the denticles being smaller than others.

Locality: Tomioka (Mar. 1935; 2 sps.).

This species resembles *Cuthona (Hervia) sibogae* (BERGH) in the body form (and probably in the radula consisting of numerous teeth), but cannot be safely identified with that species by having a different coloration and a fairly stout median spine of the teeth.

Family Fionidae

Genus *Fiona* FORBES & HANLEY, 1851

Fiona FORBES & HANLEY, Brit. Moll., pts. 41-42, vol. 3 (contents, p. 10, note), 1851.

Type: *Oithona nobilis* FORBES & HANLEY.

154. *Fiona pinnata* (ESCHSCHOLTZ, 1831)

Eolidia pinnata ESCHSCHOLTZ, Zool. Atlas, Hft. 4, 1831, p. 14, pl. 19, fig. 1.—Sitka (Alaska).

Fiona pinnata BERGH, Verh. k. k. zool.-bot. Gesell. Wien, Bd. 23, 1873, pp. 609-610.—Japan.

Fiona marina ELIOT, Journ. Coll. Sci. Imp. Univ. Tôkyô, vol. 35, art. 1, 1913, pp. 44-45.—Japan.

Family Aeolidiidae

Subfamily Favorininae

Genus *Favorinus* GRAY, 1850

Favorinus GRAY, Figs. Mollusc. Anim., vol. 4, 1850, p. 109.

Type: *Eolis alba* ALDER & HANCOCK.


155. *Favorinus pacificus* nov. sp.

(Pl. 2, fig. 7; text-fig. 18)

Distribution in Japan: Tomioka (Amakusa).

The body in life measures about 10 mm in length. The anterolateral corners of the head are continued out into two exceedingly slender oral tentacles. The rhinophores are a little shorter and cylindrico-conical, with indistinct constrictions along the whole length. The margin of the back bears a small number of fusiform branchial papillae in 7 oblique rows, the largest being on the inside and the smallest outside; they are about 3 in the 1st, 2nd, 3rd and 4th rows, and 2 in the succeeding rows. The foot is fairly narrow, passing off to a point behind. The front edge is rounded off and its corners are produced outward into tentaculiform processes.

The general colour of the body is a translucent white and this also is true of the oral tentacles, tail and foot. The rhinophores are chocolate-tinted, except at the distal third which is colourless.



Text-fig. 18.—*Favorinus pacificus*. A. Lateral view of teeth ($\times 220$); B. Masticatory process of jaw-plate ($\times 90$); C. Masticatory edge ($\times 300$).

The branchial papillae are dark yellow almost up to the tip which terminates in a white cap, with a purple marking below.

The jaw-plates have a produced masticatory process. The whole length of the cutting edge is armed with a series of closely set spiny denticles, at the base of which are several (2-3) rows of small additional denticles. The uniseriate radula contains about 21 teeth and is formulated as $21 \times 0.1.0$. The tooth consists of a V-shaped base bearing in front a large median spine. The latter is quite smooth without any trace of denticle.

Locality: Tomioka (Feb. 1935; 1 sp.).

This species to a certain extent in its general appearance and type of radula recalls *Favorinus albus* (ALDER & HANCOCK). But it has a different arrangement of branchial papillae and also a somewhat different coloration, and I propose to put it on record as a new species with the name *F. pacificus*.

Genus *Pteraeolidia* BERGH, 1875

Pteraeolidia BERGH, Verh. k. k. zool.-bot. Gesell. Wien, Bd. 25, 1875, p. 652.

Type: *Flabellina semperi* BERGH.

156. *Pteraeolidia semperi* (BERGH, 1870)

Subfamily Phyllodesmiinae

Genus *Phyllodesmium* EHRENBERG, 1831*Phyllodesmium* EHRENBERG, Symb. phys., 1831.Type: *Phyllodesmium hyalinum* EHRENBERG.157. *Phyllodesmium hyalinum* EHRENBERG, 1831

(Pl. 2, fig. 5)

Subfamily Aeolidiinae

Genus *Aeolidia* CUVIER, 1798*Aeolidia* CUVIER, Tabl. élém. Hist. Nat., 1798, p. 388.Type: *Limax papillosus* LINNÉ.158. *Aeolidia papillosa* (LINNÉ, 1761)Genus *Baeolidia* BERGH, 1888*Baeolidia* BERGH, Malac. Unters., Hft. 16, pt. 1, 1888, pp. 777-778.Type: *Baeolidia moebii* BERGH.*Synopsis of the Japanese species:*

- a. Body large (40-70 mm), branchial groups many (10-15), tooth very broad
 *major amakusana* nov. subsp.
- b. Body small (12 mm), branchial groups 5, tooth not broad *japonica*.

159. *Baeolidia major amakusana* nov. subsp.¹⁾

(Pl. 1, fig. 10)

Baeolidia major BARA, Annot. Zool. Japon., vol. 14, no. 1, 1933, p. 178.—Tomioka.

As our specimen previously referred to *Baeolidia major* is different from the type of ELIOT in the position of the anus, it seems better to call the former *B. major amakusana* nov. subsp. Two more specimens from Tomioka (length 35 and 70 mm respectively) have been examined. The rhinophore is closely studded with small round granules in irregular rows. The branchial papillae are in about 12-15 groups, each lying upon a horseshoe-shaped base. The back is yellowish brown spotted with yellow,

1) *Baeolidia major* at p. 201 (Opisthobranchia of Japan, I), is altered to *B. m. amakusana*.

paler on the pericardium. The branchial papillae have dark brown liver diverticula shining through the integument which is spotted with yellow.

160. *Baeolidia japonica* BABA, 1933

A few remarks may be added to the previous description. The branchial papillae are arranged on both sides in about 5 groups, the first two lying upon horseshoe-shaped bases and the remainder upon simple oblique bases. The genital orifice lies immediately below the 1st group and the anus in the centre of the 2nd group on the right side. The body is dark brown, covered with white spots which are most conspicuous on the back and head.

Genus *Eolidina* QUATREFAGES, 1843

Eolidina QUATREFAGES, Ann. Sci. Nat., Zool., ser. 2, tom. 19, 1843, p. 276.

Aeolidiella BERGH, Vidensk. Meddel. Naturh. Foren., (1866) 1867, p. 99.

Type: *Eolidina paradoxum* QUATREFAGES.

With 3 subgenera, viz. *Eolidina* s. s. (*Aeolidiella*), *Spurilla* and *Berghia*.

Synopsis of the Japanese species:

- a. Body stout, rhinophores with 2-3 annular swellings, antero-lateral corners of foot rounded *japonica*.
- b. Body small, rhinophores smooth, antero-lateral corners of foot slightly angulated *takanosimensis*.

161. *Eolidina (Eolidina) japonica* (ELIOT, 1913)

162. *Eolidina (Eolidina) takanosimensis* (BABA, 1930)

Subfamily Glaucinae

Genus *Glaucus* FORSTER, 1777

Glaucus FORSTER, Voy. Resolution, 1, 1777, p. 49.

Type: *Glaucus atlanticus* FORSTER.

163. *Glaucus marinus* (DUPONT, 1763)

Hirudo marina DUPONT, Phil. Trans. Roy. Soc., 1763, p. 57.

Glaucus lineatus ELIOT, Journ. Coll. Sci. Imp. Univ. Tōkyō, vol. 35, art. 1, 1913, pp. 45-46.—Sagami Sea; HIRASÉ, Moluskoj, 1927, p. 1475, fig. 2838.—Kii, Sagami.

SUMMARY

This paper gives a comprehensive list of the species of Nudi-branchia from Japanese waters. The following eleven are recognized as new species: *Euphurus ornatus*, *Polycera fujitai*, *Goniodoris glabra*, *Okenia barnardi*, *Noumea nivalis*, *Geitodoris lutea*, *Thordisa amakusana*, *Discodoris pallida*, *Cuthona emurai*, *C. ornata*, *Favorinus pacificus*.

The following two are regarded as new subspecies: *Marionia pustulosa odhneri*, *Baeolidia major amakusana*.

Two new specific names have been proposed for the previously noted forms:—*Phyllidia japonica*, *Cuthona japonica*.

CORRECTION OF ERRORS IN "OPISTHOBRANCHIA OF JAPAN (I)"

- P. 196, line 21, for "The Foundation for the Promotion of Scientific and Industrial Research of Japan," read "The Japan Society for the Promotion of Scientific Research."
- P. 211, line 4, for "*Tethys japonica* (CLESSIN, 1899)," read "*T. sieboldiana* (CLESSIN, 1899)." See CLESSIN (1899), p. 54.
- P. 222, line 27, for "colour," read "ground-colour."
- P. 223, line 5, for "descending," read "ascending;" line 6, for "ascending," read "descending."
- P. 226, line 14, for "ascending," read "descending;" line 15, for "descending," read "ascending."
- P. 234, line 41, for "Si, T.," read "Tchang Si."

POSTSCRIPT

Pleurobranchaea japonica has been recorded by the late Prof. J. THIELE from Kôbe, Japan (Gastrop. d. Deutsch. Tiefsee-Exped., Th. 2, 1925, p. 283, pl. 33, fig. 8). The description is very short, but it appears probable that this is identical with the common Japanese species, *P. novaezealandiae* CHEESEMAN (see "Opisthobranchia of Japan," I, p. 229).

BIBLIOGRAPHY

- ALDER, J. & A. HANCOCK 1845 Notice of a new genus and several new species of nudibranchiate mollusca. Ann. Mag. Nat. Hist., ser. 1, vol. 16.
- 1855 A monograph of the British nudibranchiate mollusca, with figures of all the species. pt. 7, appendix.
- ALLAN, J. K. 1932 A new genus and species of sea-slug, and two new species of sea-hares from Australia. Rec. Austral. Mus., vol. 18, no. 6.
- 1932 Australian nudibranchs. Austr. Zoologist, vol. 7, pt. 2.
- BABA, K. 1937 Opisthobranchia of Japan (I). Journ. Dept. Agric. Kyûshû Imp. Univ., vol. 5, no. 4.
- 1937 Contribution to the knowledge of a nudibranch, *Okadaia elegans* BABA. Japan. Journ. Zool., vol. 7, no. 2.
- 1937 Note on the colour variation of a dorid, *Rostanga arbutus* (ANGAS, 1864). Venus, vol. 7, no. 1.
- 1937 Two new species of the nudibranchiate genus *Cadlina* from Sagami Bay, Japan. Venus, vol. 7, no. 2.
- 1937 Record of a nudibranch, *Gymnodoris striata* (ELIOT), from Amakusa, Japan. Zool. Mag. (Japan), vol. 49, no. 6.
- BERGH, R. 1874 Malacologische Untersuchungen, Hft. 6.
- 1874 Neue Nacktschnecken der Südsee. 2. Journ. Mus. Godeffroy, Hft. 6.
- 1875 Neue Nacktschnecken der Südsee. 3. Journ. Mus. Godeffroy, Hft. 8.
- 1877 Kritische Untersuchungen der EHRENBURG'schen Doriden. Jahrb. Deutsch. Malakoz. Gesell., Bd. 4.
- 1878 Malacologische Untersuchungen, Hft. 13.
- 1879 Gattungen nordischer Doriden. Arch. f. Naturgesch., Jahrg. 45, Bd. 1.
- 1880 Die Gattung *Gymnodoris* FORBES. Malakoz. Blätt., Bd. 2.
- 1888 Beiträge zur Kenntniss der Aeolidiaden. 9. Verh. k. k. zool.-bot. Gesell. Wien, Bd. 38.
- 1888 Die Pleuroleuriden, eine Familie der nudibranchiaten Gaströpoden. Zool. Jahrb., Syst., Bd. 3.
- 1892 Die cryptobranchiaten Dorididen. Zool. Jahrb., Syst., Bd. 6.
- 1894 Die Opisthobranchien. Bull. Mus. Comp. Zool. Harvard, vol. 25, no. 10.
- COCKERELL, T. D. A. & C. ELIOT 1905 Notes on a collection of Californian nudibranchs. Journ. Malac., vol. 12, no. 3.
- ELIOT, C. 1902 On some nudibranchs from Zanzibar. Proc. Zool. Soc. London, pt. 2.
- 1904 On some nudibranchs from East Africa and Zanzibar. pt. 4. Dorididae Cryptobranchiatae, 2. Proc. Zool. Soc. London, pt. 1.
- 1905 Nudibranchs from the Indo-Pacific. 1. Notes on a collection dredged near Karachi and Maskat. Journ. Conch., London, vol. 11, no. 8.
- HANEL, E. 1905 *Cephalopyge trematoides* (CHUN). Eine neue Mollusken-Gattung. Zool. Jahrb., Syst., Bd. 21.
- HIRASÉ, S. 1937 On a new *Berthella* from Japan. Proc. Malac. Soc. London, vol. 22, pt. 4.
- KELAART, E. F. 1858 Description of a new Ceylonese nudibranch. Ann. Mag. Nat. Hist., ser. 3, vol. 1.
- MACFARLAND, F. M. 1931 *Drepanida*, new name for *Drepania* LAFONT, preoccupied. Nautilus, vol. 45, no. 1.

- PRUVOT-FOL, A. 1930 Du genre *Dendrodoris* EHRENBERG et de ses rapports avec le genre *Doriopsis* PEASE et avec quelques autres. Note sur la taxonomie des nudibranches. Bull. Mus. Hist. Nat. Paris, ser. 2, tom. 2, no. 3.
- 1931 Notes de systématique sur les opisthobranches (suite). Bull. Mus. Hist. Nat. Paris, ser. 2, tom. 3, no. 8.
- VAYSSIÈRE, A. 1877 On a new genus of the family Tritoniidae. Ann. Mag. Nat. Hist., ser. 4, vol. 20.
- 1879 Description du *Marionia berghii*. Journ. Conchyl., Paris, tom. 27.
- 1901, 1919 Recherches zoologiques et anatomiques sur les mollusques opisthobranches du Golfe de Marseille. Ann. Mus. Hist. Nat. Marseille, toms. 6, 17.

INDEX OF THE GENERA AND SPECIES

(I) = Opisthobranchia of Japan I

(II) = Opisthobranchia of Japan II

- Acanthodoris* (II) 293
pilosa (I) 199, (II) 294
uchidai (II) 294
- Aclesia freeri* (I) 218
- Actinocyclus* (II) 300
japonicus (I) 202, (II) 300
- Aeolidia* (II) 335
papillosa (I) 199, (II) 335
- Aeolidiella* (II) 336
- Aglaja* (I) 204
cyanea (I) 201, 205
gigliolii (I) 197, 202, 204
- Aglaja* (I) 204
cyanea var. *vittata* (I) 205
- Albania formosa* (I) 197
- Amphorina* (II) 328
- Aplysia* (I) 207
atromarginata (I) 208
euchlora (I) 197, 208
fimbriata (I) 196
japonica (I) 197, 211
laevigata (I) 196
marginata (I) 197, 208
marmorea (I) 197, 208
nigrocincta (I) 208
parvula (I) 208
sibogae (I) 211
- Argus* (II) 307
cruentus (I) 201, (II) 308
esakii (II) 308
speciosus (I) 198, 201, (II) 308
striatus (I) 201, (II) 308
tabulatus (I) 198, 201, (II) 308
- Armina* (II) 317
babai (II) 320
comta (I) 198, 202, (II) 320
fallax (I) 198, 202, (II) 320
iaira (I) 197, 202, (II) 320
japonica (I) 202, (II) 320
papillata (II) 320
similis (I) 202, (II) 320
taeniolata (I) 201, (II) 318
variolosa (II) 320
- Asteronotus* (II) 307
cespitosus (I) 201, (II) 307
- Baeolidia* (II) 335
japonica (II) 336
major (I) 201, (II) 335
major amakusana (II) 335
- Berthella* (I) 227
borneensis (I) 227
gotoi (I) 227
plumula delicata (I) 201, 227
- Bornella* (II) 321
digitata (I) 201, (II) 321
- Bulla hirundinina* (I) 206
- Bursatella leachii* var. *freeri* (I) 218
leachii var. *hirasei* (I) 218
- Cadlina* (II) 299
japonica (II) 299
sagamiensis (II) 299
- Calliopaea dendritica* (I) 223
- Caloplocamus* (II) 292, 293
croceus (II) 293
ramosus (I) 197, 198, 200, (II) 293
- Casella* (II) 300
atromarginata (I) 201, (II) 300
- Cephalopyge* (II) 326
orientalis (I) 200, (II) 326
- Ceratossoma* (II) 300
cornigerum (I) 198, 201, (II) 300
- Chelidonura* (I) 206
hirundinina (I) 201, 206
- Chromodoris* (II) 297
alderi (I) 197
aureopurpurea (I) 198
clitonota (II) 298
marenzelleri (I) 198
pantharella (I) 197
petechialis (I) 198
sibogae (II) 298
thalassopora (I) 197, (II) 297
- Coryphella* (II) 327
alderi (I) 197, 202, (II) 327

- athadona (I) 197, 199, (II) 328
 ornata (I) 201, (II) 328
Cratena (II) 328
Crosslandia (II) 324
 fusca (II) 324
 orientalis (II) 324
 viridis (I) 201, (II) 324
Cryptophthalmus (I) 202
 smaragdinus (I) 201, 202
Ctenodoris (II) 301
 aurantiaca (I) 202, (II) 301
Cuthona (II) 328
 bicolor (I) 202, (II) 328
 ceylonica (I) 198, 201, (II) 329
 emurai (I) 199, (II) 329
 enosimensis (II) 329
 japonica (II) 329
 ornata (II) 331
 quadrilineata (II) 329
 quadrilineata var. (II) 329
 rosea (II) 329
 yatsui (II) 328
Cyerce (I) 221
 nigricans (I) 201, 222

Dendrodoris (II) 308
 elongata (II) 309
 gemmacea (I) 198, 201, (II) 309
 guttata (I) 201, (II) 309
 miniata (II) 309
 nigra (I) 196, 197, 198, 201, (II) 309
 nigra var. (II) 309
 rubra var. *nigromaculata* (I) 202,
 (II) 309
 tuberculosa (I) 201
 tuberculosa var. (II) 309
Dermatobranchus (II) 316
 striatus (I) 201, (II) 316
Dirona (II) 320
 albolineata (I) 199, (II) 321
Discodoris (II) 305
 concinna (I) 201, (II) 305
 pallida (II) 305
 pardalis (I) 201, (II) 305
 yaeyamensis (II) 305
Dolabella (I) 221
 scapula (I) 201, 221

Dolabrifera (I) 215
 dolabrifera (I) 201, 216
 tahitensis (I) 216
Doridium (I) 204
 cyaneum var. *vittatum* (I) 205
Doridopsis (II) 308
 gemmacea (I) 198
 indacus (I) 197
Doriopsis (II) 308
 nigra var. *coerulea* (I) 198
 viridis (I) 197, 201, (II) 308
Doriprismatica festiva (I) 197
Doris areolata (I) 196
 atromarginata (II) 300
 aurantiaca (II) 301
 latens (I) 196
 lineolata (II) 297
 nigra (I) 196
 olivacea (I) 196
 pecten (I) 197, (II) 308
 rogersii (I) 196
 striata (II) 308
Doto (II) 327
 japonica (I) 202, (II) 327
Drepania (II) 297
 japonica (II) 297
Drepanida (II) 297
Duvaucelia (II) 310
 exsulans (I) 199, (II) 310
 irrorata (II) 315
 reticulata (I) 198, 202, (II) 310
 tetraquetra (I) 197, 199

Echinodoris (II) 300
 armata (I) 202, (II) 300
Elysia (I) 224
 grandifolia (I) 201, 225
 japonica (I) 202, 225
 viridis (I) 200, 225
 yaeyamana (I) 226
Eolidia pinnata (II) 333
Eolidina (II) 336
 japonica (I) 202, (II) 336
 takanosimensis (II) 336
Euphurus (II) 289
 ornatus (II) 289
Euplocamus (II) 292
 japonicus (I) 197, 198

- Favorinus** (II) 333
 pacificus (II) 333
Fiona (II) 333
 marina (II) 333
 pinnata (I) 197, 200, (II) 333
Geitodoris (II) 302
 lutea (II) 302
 ohshimai (II) 302
Glaucus (II) 336
 lineatus (II) 336
 marinus (I) 200, (II) 336
Glossodoris (II) 297
 alderi (I) 197, 198, 201, (II) 297
 aureopurpurea (I) 198, (II) 298
 clitonota (I) 201, (II) 298
 festiva (I) 197, 198, 202, (II) 297
 lineolata (I) 201, (II) 297
 pallescens (I) 198, 201, (II) 297
 pantharella (I) 197, 202, (II) 298
 reticulata (II) 297
 sibogae (I) 201, (II) 298
 thalassopora (I) 197, 202, (II) 297
Goniodoris (II) 294
 castanea (I) 200, 201, (II) 294
 glabra (II) 294
Guyonia (II) 308
Gymnodoris (II) 292
 alba (I) 201, (II) 292
 citrina (I) 201, (II) 292
 inornata (I) 198, 201, (II) 292
 japonica (II) 292
 maculata (I) 196, 202, (II) 292
 okinawae (II) 292
 striata (II) 292
Halgerda (II) 307
 graphica (I) 201, (II) 307
 japonica (I) 202, (II) 307
Hermaea (I) 223
 dendritica (I) 200, 223
Hervia (II) 328
Hexabranchnus (II) 289
 lacer (I) 197
 marginatus (I) 201, (II) 289
Hirudo marina (II) 336
Homoeodoris (II) 307
 japonica (I) 198, 202, (II) 307
Homoiodoris (II) 307
Kalinga (II) 293
 ornata (I) 201, (II) 293
Kaloplocamus (II) 293
Laplysia (I) 207
Limax tetraquetra (II) 312
Linguella babai (II) 320
 fallax (I) 198, (II) 320
 iaira (II) 320
 variolosa (II) 320
Madrella (II) 327
 sanguinea (I) 201, (II) 327
Marionia (II) 312
 granularis (II) 312
 pustulosa odhneri (II) 313
Marioniopsis (II) 315
 babai (II) 315
Melibe (II) 326
 japonica (I) 202, (II) 327
 papillosa (I) 197, 198, (II) 327
 pilosa (I) 198, 201, (II) 327
 vexillifera (I) 198, 202, (II) 327
Nembrotha (II) 292
 luteolineata (II) 292
Notarchus (I) 218
 leachii var. *freeri* (I) 201, 218
 lineolatus (I) 196
 longicaudus (I) 196, 201, 221
 stimpsoni (I) 221
Notobryon (II) 321
 wardi (I) 201, (II) 321
Noumea (II) 298
 nivalis (II) 298
Okadaia (I) 197, (II) 297
 elegans (I) 199, (II) 297
Okenia (II) 295
 barnardi (II) 295
Oscanius sp. (I) 227
Pelta (I) 202
Peltodoris (II) 304
 mauritiana (I) 199, 201, (II) 304
Petalifera (I) 216
 punctulata (I) 197, 126

- Petelodoris* (II) 302
 triphylla (I) 198, 202, (II) 302
Phyllaplysia punctulata (I) 197, 216
Phyllidia (II) 309
 japonica (II) 310
 nobilis (I) 201, (II) 310
 pustulosa (I) 201, (II) 310
 tuberculata (II) 310
 varicosa (I) 201, (II) 310
Phylliroe (II) 326
 bucephala (I) 200, (II) 326
Phyllodesmium (II) 335
 hyalinum (I) 201, (II) 335
Placobranchus (I) 224
 guttatus (I) 196
 ocellatus (I) 196, 201, 224
Platydoris (I) 307
 striata (II) 308
Pleurobranchaea (I) 229
 japonica (II) 337
 novaezealandiae (I) 229
 novaezealandiae var. *granulosa*
 (I) 229
Pleurobranchus (I) 229
 delicatus (I) 227
Pleuroleura (II) 316
 striata (II) 316
Pleurophyllidia (II) 317
 comta (I) 198
 similis (II) 320
 taeniolata (II) 318
Plocamopherus (II) 293
Plocamophorus (II) 293
 imperialis (I) 201, (II) 293
 tilesii (I) 197, 198, 202, (II) 293
Polycera (II) 290
 fujitai (II) 290
Pseudobornella (II) 321
 orientalis (II) 321
Pteraeolidia (II) 334
 semperi (I) 201, (II) 335

Rizzolia (II) 328
 modesta (I) 198, (II) 329

Rostanga (II) 300
 arbutus (I) 201, (II) 300
Runcina (I) 202
 elioti (I) 202

Sancara iaira (I) 197, (II) 320
Scyllaea (II) 321
 bicolor (I) 198, (II) 321
Stiliger (I) 222
 akkeshiensis (I) 223
 berghi (I) 222
Stylocheilus longicaudus (I) 221

Tethys (I) 207
 dactylomela (I) 196, 201, 208
 euchlora (I) 197, 202, 208
 hirasei (I) 211
 japonica (I) 197, 211
 kurodai (I) 213
 laevigata (I) 196, 208
 marginata (I) 197, 208
 marmorea (I) 197
 nigrocincta (I) 208
 norfolkensis (I) 208
 parvula (I) 199, 201, 208
 sibogae (I) 201, 211
Thordisa (II) 304
 amakusana (II) 304
Trapania (II) 296
 japonica (II) 297
Trevelyana (II) 292
 felis (I) 197
 inornata (I) 198
Triopa (II) 289
Trippa (II) 302
 intecta (I) 201, (II) 302
Tritonia exsulans (II) 310
 reticulata (I) 198, (II) 310
 tetraquetra (I) 197, (II) 312
Tritoniopsilla (II) 312
 tetraquetra (II) 312

Vayssierea (I) 197

EXPLANATION OF PLATES

Plate 1.

- Fig. 1. *Dendrodoris miniata* (ALDER & HANCOCK). × 1.
Fig. 2. *Madrella sanguinea* (ANGAS). × 3.
Fig. 3. *Discodoris pardalis* (ALDER & HANCOCK). × 1.
Fig. 4. *Dendrodoris gemmacea* (ALDER & HANCOCK). × 1.
Fig. 5. *Goniodoris castanea* ALDER & HANCOCK. × 1.
Fig. 6. *Goniodoris glabra* nov. sp. × 2.
Fig. 7. *Crosslandia viridis* ELIOT. × 2.
Fig. 8. *Gymnodoris inornata* (BERGH). × 2.
Fig. 9. *Gymnodoris citrina* (BERGH). × 6. (drawn from a Tomioka specimen with a rather unusual coloration).
Fig. 10. *Baeolidia major amakusana* nov. subsp. × 1.
Fig. 11. *Discodoris pallida* nov. sp. × 3.
Fig. 12. *Thordisa amakusana* nov. sp. × 1.3.
Fig. 13. *Dendrodoris guttata* (ODHNER). × 1.
Fig. 14. *Actinocyclus japonicus* (ELIOT). × 1.

Plate 2.

- Fig. 1. *Dermatobranchus striatus* VAN HASSELT. × 4.
Fig. 2. *Cuthona yatsui* (BABA). × 5.
Fig. 3. *Notobryon wardi* ODHNER. × 1.
Fig. 4. *Cuthona ornata* nov. sp. × 8.
Fig. 5. *Phylloidesmium hyalinum* EHRENBERG. × 2.
Fig. 6. *Okenia barnardi* nov. sp. × 6.
Fig. 7. *Favorinus pacificus* nov. sp. × 8.
Fig. 8. *Euphurus ornatus* nov. sp. × 7.



