



## **DOCUMENTING OTITIS MEDIA (H65-H67)**

- Include the following:
  - Laterality (e.g., left right or bilateral)
  - Tympanic membrane rupture, if present
  - Secondary causes such as tobacco smoke
  
- Inquire and collect information to assist in identifying:
  - Type as serous, sanguinous, suppurative, nonsuppurative, allergic or mucoid
  - Infectious agent such as strp, staph, Scarlett fever, influenza measles, mumps
  - Temporal factors as acute, subacute, chronic or recurrent
  
- Use additional code(s) to identify the following:
  - Exposure to environmental tobacco smoke
  - Exposure to tobacco smoke in the perinatal period
  - History of tobacco use
  - Occupational exposure to environmental tobacco smoke
  - Tobacco dependence
  - Tobacco use

### **OTITIS MEDIA EXAMPLES**

ICD-10 Codes	ICD-10 Descriptions
B05.3	Measles complicated by otitis media
H65.01	Acute serous otitis media, right ear
H65.113	Acute and subacute allergic otitis media (mucoid) sanguinous) (serous), bilateral
H65.194	Other acute non suppurative otitis media, recurrent, right ear
H65.32	Chronic mucoid otitis media, left ear
H66.012	Acute suppurative otitis media with spontaneous rupture of ear drum, left ear
H66.13	Chronic tubotympanic suppurative otitis media, bilateral
H66.22	Chronic atticoantral suppurative otitis media, left ear

J11.83

Influenza due to unidentified influenza virus with otitis media