

New records of *Phascolosoma turnerae* (Sipuncula: Phascolosomatidae) from the Balearic Basin, Mediterranean Sea

J.I. SAIZ¹, J.E. CARTES², V. MAMOURIDIS², A. MECO² AND M.A. PANCUCCI-PAPADOPOULOU³

¹Department of Zoology and Cell Biology, UPV/EHU, PO Box 644, 48080 Bilbao, Spain, ²Institut de Ciències del Mar (CSIC), Passeig Marítim de la Barceloneta 37-49, 08003 Barcelona, Spain, ³Hellenic Centre for Marine Research, PO Box 712, 19013 Anavissos, Greece

Specimens of the deep-water sipunculan worm Phascolosoma (Phascolosoma) turnerae were recently collected from the western part of the Mediterranean Sea. This species is characterized by hooks showing a peculiar anterior stout and long projection at their base. A key to all the Phascolosoma species found in the Mediterranean Sea is included.

Keywords: *Phascolosoma turnerae*, new records, western Mediterranean, sipunculan worm

Submitted 6 August 2013; accepted 24 January 2014

INTRODUCTION

The genus *Phascolosoma* Leuckart, 1828, is one of the most species-rich genera within the phylum Sipuncula (Murina, 1984; Gibbs & Cutler, 1987; Cutler, 1994), with the majority of its species showing a preference for the warmer shallow waters of the world's oceans (Murina, 1975). However, a few phascolosomatid species are rare in their geographical distribution (Cutler, 1994), since they show preferences for deep cold waters instead.

Deep-sea communities in the Mediterranean are poorly known (Bazairi *et al.*, 2010). Recent studies conducted over the slope of the Balearic Basin (western Mediterranean) revealed the presence of deep-sea phascolosomatids (Cartes *et al.*, 2009). They were identified as *Phascolosoma turnerae* Rice, 1985 in a taxonomic checklist of the sipunculan fauna for the Mediterranean Sea (Coll *et al.*, 2010). The new phascolosomatids inhabited muddy bottoms in the Balearic Basin. From this area, a total of 7 other species of sipunculans have been collected (Cartes *et al.*, 2009; Mamouridis *et al.*, 2011; Tecchio *et al.*, 2013) by using different bottom trawls and box-corers at depths between 427 and 2265 m. This relatively high diversity of sipunculans was related to their ability in exploiting particulate organic matter of different quality (more or less degraded), arriving at bathyal depths, as inferred by the quite different stable isotope $\delta^{15}\text{N}$ found in *Aspidosiphon muelleri* Diesing, 1851 and *Sipunculus norvegicus* Danielssen, 1869 (respectively 5.39‰ and 9.57‰, Fanelli *et al.*, 2011), two dominant sipunculan species inhabiting the investigated area (Cartes *et al.*, 2009).

After a detailed anatomical study of the phascolosomatid specimens collected, we observed anatomical features unknown to any previously recorded Mediterranean species (Pancucci-Papadopoulou *et al.*, 1999) of the genus

Phascolosoma Leuckart, 1828. Thus, the aims of the present study are: (1) to report this new faunistic finding for the Mediterranean Sea; and (2) to select reliable characters for the distinction of closely related species. The ecological information of those collected specimens is also discussed.

MATERIALS AND METHODS

Specimens of deep-sea phascolosomatids were collected during trawling operations of the RV 'García del Cid' in the north-western part of the Mediterranean Sea. Station names, coordinates, depths, sampling dates and number of specimens are listed as follows:

—Station BIOM3-OTSB4; 41°07.130''–41°07.841''N 2°22.497''–2°27.977''E; 1039–1103 m; 01/07/2007; 2 specimens.
—Station PROMETEO3 M-28; 41°07'60''N 02°52'16''E; 1500 m; 13 May 2009; 1 specimen.

Immediately after sampling, all samples were preserved in 4% borax-buffered formaldehyde, prepared using seawater. Once in the laboratory, the material was studied using standard dissecting techniques and both binocular and compound microscopes. The species identification was based mainly on the works of Cutler (1994) and Pancucci-Papadopoulou *et al.* (1999). Voucher material was deposited at the Institut de Ciències del Mar (CSIC).

RESULTS

SYSTEMATICS

Phascolosoma (Phascolosoma) turnerae Rice, 1985
(Figure 1A–F)

Trunk 23–44 mm long and 9–12 mm wide, light brown in colour (Figure 1A). Skin opaque to translucent. Papillae

Corresponding author:

J.I. Saiz

Email: ji.saiz@ehu.es




Fig. 1. *Phascolosoma (Phascolosoma) turnerae*: (A) external view; (B) papillae in the anterior part of the trunk; (C) papillae in the middle part of the trunk; (D) papillae at the trunk end; (E) scanning electron micrograph of a hook ring; (F) detail of a hook under the light microscope. Red arrow indicates the characteristic long basilateral extension of the hook. Scale bars: A, 5 mm; B and D, 0.5 mm; C, 0.3 mm; F, 20 μ m.

distributed over the entire trunk, larger and more densely packed at both ends (Figure 1B, D). Introvert 25–42 mm long with some pigmented bands. About 28 digitiform tentacles unpigmented. About 90 hook rings, many incomplete. Under the microscope hooks appear slightly curved without a secondary tooth (Figure 1E). Height 50–75 μ m. At the posterior base of the hooks there are long toes and a basal wart. A long clear streak with some expansion is noted. No triangles, but with a characteristic long basilateral extension (Figure 1F). Tubuliform papillae about 10 μ m high are placed between the hook rings. Internally, the longitudinal musculature has about 30–34 anastomosing bands. Circular musculature also split in fine bands. Four retractors the ventral arising from 6–8 bands, whereas the dorsal arise from 7–8 bands. The distance from the ventral insertions to the trunk end is 5–10 mm. Gut organized in about 15 spirals. Spindle muscle present and attached posteriorly. Two nephridia, 11 mm long with the posterior

2 mm free. Nephridiopores open a little anterior to the anus. Gonads observed at the base of the ventral retractors.

DISCUSSION

A total of 5 species of *Phascolosoma* have been reported from the Mediterranean Sea at coastal waters varying from 0 to 230 m deep (Pancucci-Papadopoulou *et al.*, 1999). The collection of deep-water phascolosomatids in the mainland slope of the Balearic Basin was remarkable, since we were unable to relate them to any representative of the genus *Phascolosoma* for the Mediterranean Sea. According to Cutler (1994), the nature of the hooks appears to be species-specific within this sipunculid genus and a global key for the identification of all valid species was constructed by using basically the detailed study of the hooks under the light microscope. In this way, the

presence of characteristic hooks with anterior prong-like extensions related our specimens to only the two species exhibiting this character: *Phascolosoma turnerae* Rice, 1985; and *P. saprophagicum* Gibbs, 1987. Gibbs (1987) differentiated the two species by: (i) the height of the hooks ($> 45\text{--}80\ \mu\text{m}$ in *P. turnerae* vs $20\text{--}25\ \mu\text{m}$ in *P. saprophagicum*); and (ii) the external shape of the hooks (sharply-pointed in *P. turnerae* vs bluntly-rounded in *P. saprophagicum*). Since our specimens showed sharply bent hooks, $50\text{--}75\ \mu\text{m}$ tall, we have identified them with *P. turnerae*.

The faunistic finding is also interesting from an ecological approach. The species was described originally by Rice (1985) inhabiting submerged wood at bathyal depths in the Strait of Florida and the Gulf of Mexico. The large abundance of sipunculans boring the wood persuaded this author to note a biological association between this sipunculan species and the wood. Similarly, our BIOM₃-OTSB₄ specimens were collected (after 4.5 ha of trawling) together with a large piece ($\sim 25\ \text{kg}$) of a tree trunk, some tree leaves (from purple beech, *Fagus sylvatica*) and *Posidonia oceanica* remains, suggesting this ground located near Besós Canyon could be a depocentre area for terrestrial-shelf material. These terrestrial inputs are patchily distributed over the slope (authors, unpublished data), although they are more frequent in the mainland than in the insular slope of the Balearic Basin, with higher levels of total organic matter at $1000\text{--}1100\ \text{m}$ (Fanelli *et al.*, 2013). $\delta^{13}\text{C}$ stable isotope signals on sediments close to the BIOM₃-OTSB₄ haul (between -23.6‰ – -24.8‰ Fanelli *et al.*, 2013) also suggests an enrichment of terrigenous sediments there.

Almost simultaneously Edmonds (1985) described a new species, *Phascolosoma kapalum*, from deep bottoms off Australia. In this paper, there was no indication of wood occupation by the sipunculans. Later on, Edmonds (1992) compared specimens of *P. kapalum* with *P. turnerae* and decided that the two taxa were conspecific, the first name being a junior synonym of the last one. The finding of an extra specimen from a piece of rotting wood off the Great Barrier Reef (Australia) confirmed the character of a deep-sea wood dweller. However, in the review of the subgenus *Phascolosoma* achieved by Cutler & Cutler (1990), they identified further specimens from bathyal depths in the Gulf of Mexico, but this time the association was noted with tubeworm aggregations near a cold water seep. Other more ecological papers (Olu *et al.*, 1996, 2010; MacAvoy *et al.* 2005; Cordes *et al.*, 2006, 2007) reported the species also in mussel beds and sediments at the base of the tubeworm aggregations from different cold seep sites around the Gulf of Mexico, but also offshore the Orinoco River delta in South America (at $1950\text{--}2080\ \text{m}$ deep in sediment samples).

Another rare phascolosomatid, *P. saprophagicum* Gibbs, 1987, was collected from bathyal depths close to Chatham Island (New Zealand). Several specimens were obtained from the crevices and between the attached mussels on a large piece of decaying bone corresponding to a whale skull. Both *P. saprophagicum* and *P. turnerae* are the only representatives of the genus colonizing bizarre ephemeral habitats in deeper waters. Moreover, both species share this unique basal projection at the anterior base of the hook. Whereas we can suspect the existence of some genetic affinity between the two species, only DNA has been sequenced by now from *P. turnerae* (Kawauchi *et al.*, 2012). Consequently, *P. turnerae* and *P. saprophagicum*, remain separate valid entities, as was proposed in the last monograph about the phylum (Cutler, 1994).

Recently, a comprehensive and detailed study on the reproductive biology and life-history of *P. turnerae* was published (Rice *et al.*, 2012), since the authors were able to maintain several specimens in captivity for long periods of time. Interestingly, other kinds of habitats were reported for the species, such as authigenic carbonate rocks and bundles of plant fibres, which inspired the authors to design artificial collectors of deep-sea phascolosomatids. The species was additionally found in bathyal depths close to the Bahamas and Barbados, showing by now a disjunct geographical distribution of two remote areas: the western Atlantic and south-western Pacific Oceans. To explain this gap, Rice *et al.* (2012) suggested the possibility of a wider distribution than currently known for *P. turnerae*. This hypothesis is today more plausible, with the discovery of *P. turnerae* in bathyal depths of the Mediterranean Sea (Bienhold *et al.*, 2013; our findings). In this way, these locations within the Mediterranean Sea represent an intermediate spot between the two remote areas where the species was previously collected. The presence of *P. turnerae* in the Mediterranean Sea has also implications in a potential connection of disjunct metapopulations of *P. turnerae* from both sides of the Atlantic Ocean. Dispersal trajectories of planktonic larvae were simulated in a recent study of this wood-dwelling species (Young *et al.*, 2012). Its larvae were capable of reaching the mid-Atlantic off Newfoundland, a distance of more than $3000\ \text{km}$, during a long drifting period. Consequently, the possibility of a genetic exchange from west to east in the North Atlantic Drift current is even more probable, specimens of *P. turnerae* having been collected in the Mediterranean Sea.

Finally, we were unable to note a direct association of *P. turnerae* with sunken wood in the western part of the Mediterranean Sea. However, Bienhold *et al.* (2013) did recently by deploying wood colonization experiments offshore the River Nile delta at a depth of $1690\ \text{m}$ (eastern Mediterranean Sea). In our case, *P. turnerae* was the only sipunculan species collected in both hauls, while other sipunculans are relatively abundant from other sampling sites (Cartes *et al.*, 2009; Mamouridis *et al.*, 2011; Tecchio *et al.*, 2013) over the investigated slope. In general, surface deposit-feeder invertebrates were poorly represented in the first reported haul (only 1 holothurian, *Mesothuria intestinalis*), while the rest of invertebrates were—based on stable isotope analyses in the same area (Fanelli *et al.*, 2011, in 2013)—both filter feeders (1 bivalve *Abra longicallus*; 1 holothurian *Ypsilothuria bitentaculata*) and carnivores (1 seastar *Ceramaster grenadensis*). This suggests that inputs of fresh organic matter, as found deeper (e.g. over $1600\ \text{m}$, Cartes *et al.*, 2009), are low within the trawled area, indirectly reinforcing the idea that *P. turnerae* would have as food source ephemeral material derived from terrestrial inputs, such as remains of wood.

KEY TO THE GENUS
PHASCOLOSOMA LEUCKART
1828, FROM THE MEDITERRANEAN
SEA [MODIFIED FROM PANCUCCI-
PAPADOPOULOU *ET AL.* (1999) TO
ACCOMMODATE THE NEW RECORD]

- 1a. Hooks under the light microscope exhibiting an anterior stout and long projection at the base of their convex side *P. turnerae*

- 1b. Hooks without an anterior projection at the base of their convex side 2
- 2a. More than 50 complete and incomplete rings of hooks over the full everted introvert 3
- 2b. Less than 50 complete and incomplete rings of hooks ..4
- 3a. Hooks with a posterior crescent area, many >75 µm tall; preanal papillae are smooth cones; pigment bands on introvert *P. stephensoni*
- 3b. Hooks with a granular triangle at their anterior base, but without crescents, most <75 µm tall; no pigment bands on introvert *P. granulatum*
- 4a. Large rounded hump on concave side of the hook; preanal papillae smooth, posteriorly directed, cone shaped *P. perlucens*
- 4b. Concave side of hook smooth or with small tooth 5
- 5a. Hooks with distinct triangle; narrow band of red cone-shaped preanal papillae *P. scolops*
- 5b. Hook triangle indistinct or absent; preanal papillae not distinct from dome-shaped trunk papillae
P. agassizii agassizii.

ACKNOWLEDGEMENTS

This study was funded by the Spanish Ministry of Science projects ANTROMARE (J.E.C. and V.M. ref. CTM2009-12214-Co2-01/MAR) and PROMETEO (A.M. ref. CTM2007-66316-Co2/MAR). The authors thank all the participants of both scientific cruises. Special thanks to Dr Th. Kanellopoulos for scanning electronic microscope shooting at the HCMR premises in Anavyssos.

REFERENCES

- Bazairi H., Ben Haj S., Boero F., Cebrian D., De Juan S., Limam A., Leonart J., Torchia G. and Rais C. (2010) *The Mediterranean Sea biodiversity: state of the ecosystems, pressures, impacts and future priorities*. Tunis: UNEP-MAP RAC/SPA, 100 pp.
- Bienhold C., Pop Ristova P., Wenzhöfer F., Dittmar T. and Boetius A. (2013) How deep-sea wood falls sustain chemosynthetic life. *PLoS ONE* 8, e53590. doi:10.1371/journal.pone.0053590.
- Cartes J., Maynou F., Fanelli E., Romano C., Mamouridis V. and Papiol V. (2009) The distribution of megabenthic, invertebrate epifauna in the Balearic Basin (western Mediterranean) between 400 and 2300 m: environmental gradients influencing assemblages composition and biomass trends. *Journal of Sea Research* 61, 244–257.
- Coll M., Piroddi C., Steenbeek J., Kaschner K., Lasram F.B.R., Aguzzi J., Ballesteros E., Bianchi C.N., Corbera J., Dailianis T., Danovaro R., Estrada M., Froggia C., Galil B.S., Gasol J.M., Gertwagen R., Gil J., Guilhaumon F., Kesner-Reyes K., Kitsos M.-S., Koukouras A., Lampadariou N., Laxamana E., Lotze H.K., Martin D., Mouillot D., Oro D., Raicevich S., Rius-Barile J., Saiz-Salinas J.L., San Vicente C., Somot S., Templado J., Turon X., Vafidis D., Villanueva R. and Voultsiadou E. (2010) The biodiversity of the Mediterranean Sea: estimates, patterns, and threats. *PLoS ONE* 5, e11842. doi:10.1371/journal.pone.0011842.
- Cordes E.E., Bergquist D.C., Predmore B.L., Jones C., Deines P., Telesnicki G. and Fisher C.R. (2006) Alternate unstable states: convergent paths of succession in hydrocarbon-seep tubeworm-associated communities. *Journal of Experimental Marine Biology and Ecology* 339, 159–176.
- Cordes E.E., Carney S.L., Hourdez S., Carney R.S., Brooks J.M. and Fisher C.R. (2007) Cold seeps of the deep Gulf of Mexico: community structure and biogeographic comparisons to Atlantic equatorial belt seep communities. *Deep-Sea Research I* 54, 637–653.
- Cutler N.J. and Cutler E.B. (1990) A revision of the subgenus *Phascolosoma* (Sipuncula: *Phascolosoma*). *Proceedings of the Biological Society of Washington* 103, 691–730.
- Cutler E.B. (1994) *The Sipuncula: their systematics, biology, and evolution*. Ithaca, NY: Cornell University Press, 453 pp.
- Edmonds S.J. (1985) A new species of *Phascolosoma* (Sipuncula) from Australia. *Transactions of the Royal Society of South Australia* 109, 43–44.
- Edmonds S.J. (1992) A note on *Phascolosoma turnerae* (Sipuncula). *Transactions of the Royal Society of South Australia* 116, 151.
- Fanelli E., Papiol V., Cartes J.E., Rumolo P., Brunet C. and Sprovieri M. (2011) Food web structure of the megabenthic, invertebrate epifauna on the Catalan slope (NW Mediterranean): evidence from $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analysis. *Deep-Sea Research I* 58, 98–109.
- Fanelli E., Papiol V., Cartes J.E., Rumolo P. and López-Pérez C. (2013) Trophic webs of deep-sea megafauna on mainland and insular slopes of the NW Mediterranean: a comparison by stable isotope analysis. *Marine Ecology Progress Series* 490, 199–221.
- Gibbs P.E. (1987) A new species of *Phascolosoma* (Sipuncula) associated with a decaying whale's skull trawled at 880 m depth in the south-west Pacific. *New Zealand Journal of Zoology* 14, 135–137.
- Gibbs P.E. and Cutler E.B. (1987) A classification of the phylum Sipuncula. *Bulletin of the British Museum of Natural History, Zoology* 52, 43–58.
- Kawauchi G.Y., Sharma P.P. and Giribet G. (2012) Sipunculan phylogeny based on six genes, with a new classification and the descriptions of two new families. *Zoologica Scripta* 41, 186–210.
- MacAvoy S.E., Fisher C.R., Carney R.S. and Macko S.A. (2005) Nutritional associations among fauna at hydrocarbon seep communities in the Gulf of Mexico. *Marine Ecology Progress Series* 292, 51–60.
- Mamouridis V., Cartes J.E., Parra S., Fanelli E. and Saiz Salinas J.I. (2011) A first temporal analysis on the dynamics of deep-sea macrofauna: influence of environmental variability off Catalonia coasts (western Mediterranean). *Deep-Sea Research I* 58, 323–337.
- Murina V.V. (1975) The geographical distribution of marine worms of the phylum Sipuncula of the World Ocean. In Rice M.E. and Todorović M. (eds) *Proceedings of the International Symposium on the Biology of Sipuncula and Echiura Volume 1*. Belgrade, Serbia: Naučno Delo, pp. 3–18.
- Murina G.-V.V. (1984) Ecology of Sipuncula. *Marine Ecology Progress Series* 17, 1–7.
- Olu K., Sibuet M., Harmegnies F., Foucher J.-P. and Fiala-Medioni A. (1996) Spatial distribution of diverse cold seep communities living on various diapiric structures of the southern Barbados prism. *Progress in Oceanography* 38, 347–376.
- Olu K., Cordes E.E., Fisher C.R., Brooks J.M., Sibuet M. and Desbruyères D. (2010) Biogeography and potential exchanges among the Atlantic Equatorial Belt cold-seep faunas. *PLoS ONE* 5, e11967. doi: 10.1371/journal.pone.0011967.
- Pancucci-Papadopoulou M.A., Murina G.V.V. and Zenetos A. (1999) *The phylum Sipuncula in the Mediterranean Sea. Monographs on Marine Sciences*. Athens: National Center for Marine Research, 109 pp.
- Rice M.E. (1985) Description of a wood dwelling sipunculan, *Phascolosoma turnerae*, new species. *Proceedings of the Biological Society of Washington* 98, 54–60.
- Rice M.E., Reichardt H.G., Piraino J. and Young C.M. (2012) Reproduction, development, growth, and the length of larval life of *Phascolosoma turnerae*, a wood-dwelling deep-sea sipunculan. *Invertebrate Biology* 131, 204–215.

Tecchio S., Ramírez-Llodra E., Aguzzi J., Sanchez-Vidal A., Flexas M., Sardà F. and Company J.B. (2013) Seasonal fluctuations of deep megabenthos: finding evidence of standing stock accumulation in a flux-rich continental slope. *Progress in Oceanography* 118, 188–198.

and

Young C.M., He R., Emlet R.B., Li Y., Qian H., Arellano S.M., Van Gaest A., Bennett K.C., Wolf M., Smart T.I. and Rice M.E. (2012) Dispersal of deep-sea larvae from the Intra-American Seas:

simulations of trajectories using ocean models. *Integrative and Comparative Biology* 52, 483–496.

Correspondence should be addressed to:

J.I. Saiz
Department of Zoology and Cell Biology
University of the Basque Country/EHU
PO Box 644, 48080 Bilbao, Spain
email: ji.saiz@ehu.es