

SCIENTIFIC PUBLICATIONS OF THE AMERICAN MUSEUM OF NATURAL HISTORY

AMERICAN MUSEUM NOVITATES

BULLETIN OF THE AMERICAN MUSEUM OF NATURAL HISTORY

ANTHROPOLOGICAL PAPERS OF THE AMERICAN MUSEUM OF NATURAL HISTORY

PUBLICATIONS COMMITTEE

ROBERT S. VOSS, CHAIR

BOARD OF EDITORS

JIN MENG, PALEONTOLOGY

LORENZO PRENDINI, INVERTEBRATE ZOOLOGY

ROBERT S. VOSS, VERTEBRATE ZOOLOGY

PETER M. WHITELEY, ANTHROPOLOGY

MANAGING EDITOR

MARY KNIGHT

Submission procedures can be found at <http://research.amnh.org/scipubs>

All issues of *Novitates* and *Bulletin* are available on the web (<http://digilibRARY.amnh.org/dspace>). Order printed copies on the web from:

<http://shop.amnh.org/a701/shop-by-category/books/scientific-publications.html>

or via standard mail from:

American Museum of Natural History—Scientific Publications
Central Park West at 79th Street
New York, NY 10024

⑧ This paper meets the requirements of ANSI/NISO Z39.48-1992 (permanence of paper).

ON THE COVER: UPPER IMAGE: THE SHRIMP *ALPHEUS MICROSTYLUS* (BATE, 1888) (ALPHEIDAE) INFESTED VENTRALLY BY THE ABDOMINAL BOPYRID PARASITE *EOPHRIXUS CAUDATUS*, N. SP. LOWER IMAGE: *E. CAUDATUS*, N. SP., MALE (LEFT) AND FEMALE (RIGHT).


AN ET AL.: REVIEW OF BOPYRIDS PARASITIC ON CARIDEAN SHRIMP

AMNH BULLETIN 399

2015

A REVIEW OF BOPYRIDS (CRUSTACEA: ISOPODA: BOPYRIDAE) PARASITIC ON CARIDEAN SHRIMPS (CRUSTACEA: DECAPODA: CARIDEA) FROM CHINA

JIANMEI AN, CHRISTOPHER B. BOYKO, AND XINZHENG LI



BULLETIN OF THE AMERICAN MUSEUM OF NATURAL HISTORY

A REVIEW OF BOPYRIDS (CRUSTACEA:
ISOPODA: BOPYRIDAE) PARASITIC ON
CARIDEAN SHRIMPS (CRUSTACEA: DECAPODA:
CARIDEA) FROM CHINA

JIANMEI AN

School of Life Science, Shanxi Normal University, Linfen, China

CHRISTOPHER B. BOYKO

*Department of Biology, Dowling College, Oakdale, NY; and Division of
Invertebrate Zoology, American Museum of Natural History*

XINZHENG LI

Institute of Oceanology, Chinese Academy of Sciences, Qingdao, China

BULLETIN OF THE AMERICAN MUSEUM OF NATURAL HISTORY

Number 399, 85 pp., 23 figures, 2 tables

Issued December 18, 2015

CONTENTS

Abstract	4
Introduction	4
Summary List of New Taxa, Taxonomic Decisions, and Chinese Records	13
Materials and Methods	14
Systematic Account	14
<i>Crustacea</i> Brünnich, 1772	14
<i>Isopoda</i> Latreille, 1817	14
<i>Cymothoida</i> Wägele, 1989	14
<i>Bopyridae</i> Rafinesque, 1815	14
<i>Argeiinae</i> Markham, 1977	14
Key to Genera of Argeiinae	15
<i>Argeia</i> Dana, 1852	15
<i>Argeia pugettensis</i> Dana, 1852	15
<i>Argeia</i> sp.	19
<i>Stegoalpheon</i> Chopra, 1923	20
<i>Stegoalpheon kempfi</i> Chopra, 1923	20
<i>Pseudioninae</i> Codoreanu, 1967	22
<i>Paranikione</i> , n. gen.	23
<i>Paranikione sibogae</i> , n. sp.	23
<i>Paranikione distorta</i> , n. sp.	25
<i>Pseudione</i> Kossmann, 1881	27
<i>Pseudione tattersalli</i> Nierstrasz and Brender à Brandis, 1923	27
<i>Bopyrininae</i> Rafinesque, 1815	28
Key to Genera of Bopyrininae	28
<i>Bathygyge</i> Hansen, 1897	29
<i>Bathygyge grandis</i> Hansen, 1897	29
<i>Bopyrella</i> Bonnier, 1900	31
<i>Bopyrella articulata</i> , n. sp.	32
<i>Bopyrella tanytelson</i> Markham, 1982	34
Key to Species of <i>Bopyrella</i>	34
<i>Bopyrinella</i> Nierstrasz and Brender à Brandis, 1925	34
<i>Bopyrinella albida</i> Shiino, 1958	35
<i>Bopyrinella parameces</i> , n. sp.	35
Key to Species of <i>Bopyrinella</i> Nierstrasz and Brender à Brandis, 1925	37
<i>Bopyrione</i> Bourdon and Markham, 1980	37
<i>Bopyrione longicapitata</i> Markham, 1982	38
<i>Bopyrione multifeminae</i> , n. sp.	38
<i>Bopyrione toloensis</i> Markham, 1982	41
Key to Species of <i>Bopyrione</i> Bourdon and Markham, 1980	41
<i>Bopyroides</i> Stimpson, 1864	41
<i>Bopyroides hippolytes</i> (Kröyer, 1838)	41
<i>Bopyroides shiinoi</i> Rybakov and Andeev, 1991	43
Key to Species of <i>Bopyroides</i> Stimpson, 1864	45
<i>Capitetragonia</i> Pearse, 1953	45
<i>Capitetragonia elliptica</i> (Markham, 1992), n. comb.	46
<i>Discorsobopyrus</i> Boyko, 2004	46
<i>Discorsobopyrus stebbingi</i> (Nierstrasz and Brender à Brandis, 1923)	46
<i>Litobopyrus</i> Markham, 1982	47
<i>Litobopyrus longicaudatus</i> Markham, 1982	47

<i>Parabopyrella</i> Markham, 1982	47
<i>Parabopyrella choprai</i> (Nierstrasz and Brender à Brandis, 1929)	48
<i>Parabopyrella cuspidata</i> , n. sp.	48
<i>Parabopyrella distincta</i> (Nierstrasz and Brender à Brandis).....	50
<i>Parabopyrella elongata</i> (Shiino, 1949)	52
<i>Parabopyrella hodgarti</i> (Chopra, 1923).....	54
<i>Parabopyrella indica</i> (Chopra, 1923)	56
<i>Parabopyrella perplexa</i> Markham, 1990.....	57
<i>Parabopyrella symmetros</i> , n. sp.....	59
Key to Species of <i>Parabopyrella</i> Markham, 1985	59
<i>Probopyriscus</i> Markham, 1982	62
<i>Probopyriscus novempalensis</i> Markham, 1982	62
<i>Septembopyrina</i> , n. gen.	62
<i>Septembopyrina tozeumaophila</i> , n. sp.....	63
<i>Hemiarthrinae</i> Markham, 1972	65
<i>Anchiarthrus</i> Markham, 1992	65
<i>Anchiarthrus derelictus</i> Markham, 1992	65
<i>Apophrixus</i> Nierstrasz and Brender à Brandis, 1931	65
<i>Apophrixus constrictus</i> Markham, 1982.....	65
<i>Dicopleon</i> Markham, 1972	66
<i>Dicopleon morator</i> Markham, 1982.....	66
<i>Eophrixus</i> Caroli, 1930	66
<i>Eophrixus brevicauda</i> (Chopra, 1923), n. comb.	67
<i>Eophrixus caudatus</i> , n. sp.....	67
<i>Eophrixus shoiji</i> Shiino, 1941	70
Key to Species of <i>Eophrixus</i> Caroli, 1930	70
<i>Hyperphrixus</i> Nierstrasz and Brender à Brandis, 1931	71
<i>Hyperphrixus filiformis</i> (Chopra, 1923), n. comb.	71
<i>Sigyn</i> , n. gen.	71
<i>Sigyn branchialis</i> , n. sp.	72
Acknowledgments	75
References	75

ABSTRACT

A review of 37 bopyrid isopods known to infest 43 species of caridean shrimps in China is presented, based on literature records and new material. Synonyms are provided for all species, with descriptions and detailed illustrations given for species represented by material examined herein. Bopyrids recorded as new species or new Chinese records belong to the subfamilies Argeiinae (2 or 3 species, depending on the identity of *Argeia* sp.: *Argeia pugettensis* Dana, 1852, *Argeia* sp., *Stegoalpheon kempfi* Chopra, 1923), Pseudioninae (2 species: *Paranikione sibogae*, n. gen. n. sp., and *Paranikione distorta*, n. sp.), Bopyrinae (13 species: *Bathygyge grandis* Hansen, 1897, *Bopyrella articulata*, n. sp., *Bopyrinella parameces*, n. sp., *Bopyrione multifeminae*, n. sp., *Bopyroides hippolytes* (Kröyer, 1838), *Bopyroides shiinoi* Rybakov and Andeev, 1991, *Parabopyrella cuspidata*, n. sp., *Parabopyrella distincta* (Nierstrasz and Brender à Brandis, 1923), *Parabopyrella elongata* (Shiino, 1949), *Parabopyrella hodgarti* (Chopra, 1923), *Parabopyrella perplexa* Markham, 1990, *Parabopyrella symmetros*, n. sp., and *Septembopyrina tozeumaophila*, n. gen., n. sp.), and Hemiarthrinae (2 species: *Eophrixus caudatus*, n. sp., and *Sigyn branchialis*, n. gen., n. sp.), collected from the South China Sea (7 species), East China Sea (3 species), Yellow Sea (4 species), and East and South China seas (2 species). Three new genera and 10 new species are described, with an additional four species newly recorded from Chinese waters. The genus *Argeia* is reviewed and a new genus is erected for *Argeia lowisi* Chopra, 1923, and *Argeia nierstraszi* Shiino, 1958. *Capitetragonia* Pearse, 1953, is recognized as the senior synonym of *Probopyria* Markham, 1985. The genus *Parabopyriscus* and species *Parabopyriscus stellatus* are deemed to be available from Markham, 1982, not Markham 1985. *Hemiarthrus filiformis* Chopra, 1923, is transferred to *Hyperphrixus* Nierstrasz and Brender à Brandis, 1931. The species composition of *Eophrixus* Caroli, 1930, is reviewed, the type species designation is clarified, and two species of *Hypophryxus* Shiino, 1934, are transferred into this genus. A key to all genera of Argeiinae and Bopyrinae and keys to species of seven bopyrine genera are provided, as is a list (with hosts) of the 82 bopyrid and ionid species currently known to infest Chinese noncaridean decapods.

INTRODUCTION

Bopyridae Rafinesque, 1815, is the largest and best-defined epicaridean family, species of which infest their decapod hosts primarily bran-chially or abdominally. Markham (1985a) and Martin and Davis (2001) considered Bopyridae as divided into 10 subfamilies. Subsequently, Bopyrophryxinae Condreanu, 1965, was synonymized with Pseudioninae Codreanu, 1967 (Bourdon and Boyko, 2005). Boyko et al. (2013) used 18S rDNA to explore the phylogeny of epicarideans and their results showed that Ioninae H. Milne Edwards, 1840, should be restricted to the type genus and removed from Bopyridae; the remaining species formerly in Ioninae were placed into a newly described Keponinae Boyko et al., 2013. Additionally Boyko et al. (2013) showed that Entophilinae Richardson, 1903, did not belong to Bopyridae but rather was a family within Cryptoniscoidea. Therefore, Bopyridae currently includes eight subfamilies, four of which (Argeiinae Markham, 1977, Bopyrinae Rafinesque, 1815, Hemiarthrinae Markham, 1972, and Pseudioninae

Codreanu, 1967) are found parasitizing caridean shrimp, with all but the latter subfamily found exclusively on carideans.

Studies on bopyrids from China, except for Hong Kong and Taiwan, were conducted primarily from 2006 onward. Prior to 2006, Nierstrasz and Brender à Brandis (1929, 1930, 1931) reported six bopyrids from Chinese waters, while Markham (1982, 1990, 1992a) and Boyko (2004) recorded 54 species. An et al. (e.g., 2006a, b, 2007a, 2007b, 2008, 2009a, 2009b, 2010, 2011, 2012a, 2012b, 2012c, 2012d, 2013a, 2013b, 2013c, 2013d, 2013e, 2014a, 2014b), Duan et al., (2008), and Williams and An (2009) reported 84 species in seven subfamilies and 46 genera. Currently, there are 119 bopyrid species known from Chinese waters (caridean parasites listed in the text plus the noncaridean parasites in table 1).

The materials reported on herein were included in an unpublished dissertation by one of us (An, 2006, not a published work according to the criteria of the ICZN). We reexamined all the specimens and have

TABLE 1

All Species of Bopyridae and Ionidae, Exclusive of Those Known from Caridean Hosts, from China
Abbreviations of subfamilies of Bopyridae and Ionidae: A, Athelginae; I, Ionidae; K, Keponinae; O, Orbioninae; P, Pseudioninae.

Subfamily	Bopyrid species	Hosts in China	Chinese localities	Other localities	Host(s) elsewhere
P	<i>Albunione yoda</i> Markham and Boyko, 2003	<i>Albunea groeningi</i> Boyko, 2002 (Anomura: Albuneidae)	Taiwan	—	—
K	<i>Allokepon hendersoni</i> (Giard and Bonnier, 1888)	<i>Charybdis bimaculata</i> (Miers, 1886) (Brachyura: Portunidae)	East China Sea; Yellow Sea	India	<i>Charybdis callianassa</i> (Herbst, 1789) (Brachyura: Portunidae)
K	<i>Allokepon longicauda</i> Duan et al., 2008	<i>Portunus pulchrirastatus</i> (Gordon, 1931) (Brachyura: Portunidae)	Beibu Gulf	—	—
K	<i>Allokepon monodi</i> (Boudon, 1967a)	<i>Portunus pelagicus</i> (Linnaeus, 1758); <i>Portunus trituberclatus</i> (Miers, 1876); <i>Thalamita</i> sp. (Brachyura: Portunidae)	Hainan Province; Zhejiang Province	Senegal	<i>Stenorhynchus seticornis</i> (Herbst, 1788) (Brachyura: Inachidae)
K	<i>Allokepon sinensis</i> (Danforth, 1971b)	<i>Lissocarcinus orbicularis</i> Dana, 1852 (Brachyura: Portunidae)	Hong Kong	Thailand; Philippines	<i>Portunus pelagicus</i> (Linnaeus, 1758) (Brachyura: Portunidae)
P	<i>Allorhinmorphus</i> <i>tuberculatus</i> An et al., 2012d	<i>Lissoporella quadrilobata</i> (Miers, 1884) (originally as <i>Porcellana</i> sp.) (Anomura: Porcellanidae)	East China Sea; Xisha (Paracel Islands)	—	—
K	<i>Apoceton digitatum</i> Stock, 1960	<i>Senocia vittata</i> (Stimpson, 1858) (Brachyura: Leucosidae)	South China Sea	Singapore	<i>Leucosia craniolarts</i> (Linnaeus, 1758) (Brachyura: Leucosidae)
K	<i>Apoceton leucosiae</i> An et al., 2006b	<i>Leucosia anatum</i> (Herbst, 1783) (Brachyura: Leucosidae)	South China Sea	—	—
K	<i>Apoceton pulcher</i> Nierstrasz and Brender à Brandis, 1930	<i>Pyrhila pisum</i> (De Haan, 1841); <i>Pyrhila</i> <i>carinata</i> (Bell, 1855); <i>Lyphira</i> <i>heterograna</i> (Ortmann, 1892); “ <i>Phlyra</i> ” spp. (Brachyura: Leucosidae)	Liaoning Province; Shandong Province; Jiangsu Province; Bohai Bay; Hebei Province; Zhejiang Province	Japan; Korea	<i>Pyrhila pisum</i> (De Haan, 1841) (Brachyura: Leucosidae)
P	<i>Aporobopyrus</i> <i>enosteoidis</i> (Markham, 1982)	<i>Enosteoides ornatus</i> (Stimpson, 1858) (Anomura: Porcellanidae)	Hong Kong	—	—
P	<i>Aporobopyrus</i> <i>megacephalon</i> (Nierstrasz and Brender à Brandis, 1929)	<i>Pachychelus pectinifarpus</i> Stimpson, 1858 (Anomura: Porcellanidae)	Hong Kong	Gulf of Thailand	porcellanid sp. (Anomura: Porcellanidae)

TABLE 1
(Continued)

Subfamily	Bopyrid species	Hosts in China	Chinese localities	Other localities	Host(s) elsewhere
P	<i>Aporobopyrus retrosa</i> Richardson, 1910	<i>Munida compressa</i> Baba, 1988; <i>Munida japonica</i> Stimpson, 1858 (Anomura: Munididae)	Taiwan	Philippines	<i>Munida kuboi</i> Yanagita, 1943; <i>Munida philippinensis</i> Macpherson and Baba, 1993 (Anomura: Munididae)
P	<i>Asymmetrone globifera</i> An et al., 2010	<i>Dardanus hessii</i> (Miers, 1884) (Anomura: Diogenidae)	Beibu Gulf; Xisha (Paracel Islands)	—	—
A	<i>Athelges takanoshimensis</i> Ishii, 1914	<i>Pagurodolfinia doederleini</i> (Doflein, 1902); <i>Pagurus aff. filholi</i> (de Man, 1887); <i>Pagurus pectinatus</i> (Stimpson, 1858); <i>Pagurus trigonocheirus</i> (Stimpson, 1858) (Anomura: Paguridae); <i>Clibanarius</i> sp.; <i>Diogenes edwardsii</i> (De Haan, 1849) (Anomura: Diogenidae)	Yellow Sea; Taiwan; Hong Kong	Japan	<i>Pagurus filholi</i> (de Man, 1887); <i>Pagurus japonicus</i> (Stimpson, 1858); <i>Pagurus pectinatus</i> (Stimpson, 1858) (Anomura: Paguridae)
P	<i>Bonyissa pyriforma</i> (Shiino, 1928)	<i>Clibanarius binaculans</i> (De Haan, 1849); <i>Diogenes edwardsii</i> (De Haan, 1849) (Anomura: Diogenidae)	Hong Kong	Japan	<i>Diogenes edwardsii</i> (De Haan, 1849) (Anomura: Diogenidae)
K	<i>Cancricepon choproae</i> (Nierstrasz and Brender à Brandis, 1925)	<i>Scalopidina spinosipes</i> Stimpson, 1858	Beibu Gulf	North Carolina to Florida, Mississippi; Mexico; Bermuda; Bahamas; <i>herbstii</i> H. Milne Edwards, 1834; <i>Panoplax depressa</i> Stimpson, 1871; <i>Rhithropanopeus harrisi</i> (Gould, 1841) (Brachyura: Panopeidae); <i>Garthiope barbadensis</i> (Rathbun, 1921); <i>Paraliomera dispar</i> (Stimpson, 1871) (Brachyura: Xanthidae)	—
K	<i>Cancricepon multiuberosum</i> An et al., 2012c	<i>Lionera laevis</i> (A. Milne-Edwards, 1873) (Brachyura: Xanthidae)	Xisha (Paracel Islands)	—	—
K	<i>Dactylolepon barbuladigitus</i> An et al., 2007b	<i>Ligare rubromaculata</i> (De Haan, 1835) (Brachyura: Xanthidae)	South China Sea	—	—
K	<i>Dactylolepon caribaeus</i> Markham, 1975	<i>Toru trituberculatus</i> (Sakai, 1961) (Brachyura: Leucosiidae)	South China Sea	SE coast of Dominican Republic	<i>Iliacantha liodactylus</i> Rathbun, 1898; <i>Iliacantha subglobosa</i> Stimpson, 1871 (Brachyura: Leucosiidae)

TABLE 1
(Continued)

Subfamily	Bopyrid species	Hosts in China	Chinese localities	Other localities	Host(s) elsewhere
K	<i>Dactylokepon holthuisi</i> Bourdon, 1967b	<i>Eduarcus martensi</i> (Pfeiffer, 1881) (Achelata: Scyllaridae)	South China Sea	Akaba Gulf	<i>Eduarcus lewiniolphi</i> (Holthuis, 1967) (Achelata: Scyllaridae)
K	<i>Dactylokepon richardsonae</i> Siebbing, 1910b	<i>Portunus argentatus</i> (A. Milne-Edwards, 1867) (Brachyura: Portunidae)	Guangdong Province	Seychelles; Thailand; Moluccas	<i>Portunus tuberculatus</i> (A. Milne-Edwards, 1861) (Brachyura: Portunidae); <i>Trapezia cymodoce</i> (Herbst, 1801) (Brachyura: Trapeziidae)
K	<i>Dactylokepon semipennatus</i> Bourdon, 1983	<i>Lupoclytus rotundatus</i> Adams and White, 1849; <i>Portunus gladiator</i> Fabricius, 1798 (Brachyura: Portunidae)	Guangdong Province; Beibu Gulf	Seleman Bay	<i>Lentilhammus latinanus</i> (Gordon, 1934) (Brachyura: Pilumnidae)
O	<i>Epipenaeon fissurae</i> Kensley, 1974	<i>Parapenaeus lanceolatus</i> Kubo, 1949; <i>Parapenaeus longipes</i> Alcock, 1905 (Dendrobranchiata: Penaeidae)	South China Sea	South Africa; Bay of Bengal	<i>Parapenaeus fissurus</i> Bate, 1881; <i>Parapenaeus longipes</i> Alcock, 1905 (Dendrobranchiata: Penaeidae)
O	<i>Epipenaeon grande</i> Nierstrasz and Brender à Brandis, 1931	<i>Penaeus monodon</i> Fabricius, 1798; <i>Penaeus penicillatus</i> (Alcock, 1905) (Dendrobranchiata: Penaeidae)	Guangdong Province; Fujian Province South China Sea	—	—
O	<i>Epipenaeon ingens</i> Nobili, 1906	<i>Penaeus chinensis</i> (Osbeck, 1765); <i>Penaeus penicillatus</i> (Alcock, 1905); <i>Penaeus semisulcatus</i> De Haan, 1844 (Dendrobranchiata: Penaeidae)	Guangdong Province; Fujian Province; Hong Kong	Indo-West Pacific	<i>Penaeus</i> spp. (Dendrobranchiata: Penaeidae)
O	<i>Epipenaeon latifrons</i> Bourdon, 1979b	<i>Penaeus penicillatus</i> (Alcock, 1905); <i>Penaeus semisulcatus</i> De Haan, 1844 (Dendrobranchiata: Penaeidae)	Guangdong Province; Fujian Province; Hainan Province	Australia	<i>Penaeus semisulcatus</i> De Haan, 1844 (Dendrobranchiata: Penaeidae)
P	<i>Gigantione hainanensis</i> An et al., 2009b	<i>Atergatis floridus</i> (Linnaeus, 1767); <i>Atergatis</i> spp. (Brachyura: Xanthidae)	Hainan Province	—	—
P	<i>Gigantione ishigakiensis</i> Shiino, 1941	<i>Liagore rubromaculata</i> (De Haan, 1835) (Brachyura: Xanthidae)	South China Sea	Japan	<i>Carpilius convexus</i> (Forskal, 1775) (Brachyura: Carpiliidae)
P	<i>Gigantione rhombos</i> An et al., 2009b	<i>Eucrate alcocki</i> Serène, 1973; <i>Eucrate</i> sp.; <i>Heteroplax dentata</i> Simpson, 1858 (Brachyura: Euryplacidae)	South China Sea; Beibu Gulf	—	—
P	<i>Gigantione tau</i> An et al., 2009b	<i>Carcinopax longimana</i> (De Haan, 1833) (Brachyura: Gonoplatidae)	East China Sea	—	—

TABLE 1
(Continued)

Subfamily	Bopyrid species	Hosts in China	Chinese localities	Other localities	Host(s) elsewhere
P	<i>Gyge fujianensis</i> An et al., 2009a	<i>Upogebia major</i> (De Haan, 1841) (Gebiidae: Upogebiidae)	Fujian Province	—	—
P	<i>Gyge ovalis</i> (Shinno, 1939)	<i>Austinogebia eutulis</i> (Ngoc-Ho and Chan, 1992); <i>Austinogebia wahsiemwei</i> (Yu, 1931); <i>Upogebia major</i> (De Haan, 1841) (Gebiidae: Upogebiidae)	Shandong Province; Taiwan	Japan Korea	<i>Upogebia major</i> (De Haan, 1841) (Gebiidae: Upogebiidae)
K	<i>Hypocopon globosus</i> Markham, 1922	Pinnotheridae sp. (Brachyura: Pinnotheridae)	Hong Kong	—	—
I	<i>Ione cornuta</i> Bate, 1864	<i>Nihonotrypaea harmandi</i> (Bouvier, 1901); <i>Nihonotrypaea japonica</i> (Ortmann, 1891); <i>Nihonotrypaea petalura</i> (Stimpson, 1860) (Axidea: Callianassidae); <i>Upogebia major</i> (De Haan, 1841) (Gebiidae: Upogebiidae)	Shandong Province	British Columbia to California; Japan; Korea	<i>Neotrypaea californiensis</i> (Dana, 1854); <i>Neotrypaea gigas</i> (Dana, 1852); <i>Nihonotrypaea japonica</i> (Ortmann, 1891) (Axidea: Callianassidae)
I	<i>Ione tainwanensis</i> Markham, 1995	<i>Callianixa</i> sp. (Axidea: Callianixidae)	Taiwan	—	—
K	<i>Leidya ucae</i> Pearse, 1930	<i>Uca forcipata</i> (Adams and White, 1849) (Brachyura: Ocypodidae)	Fujian Province	—	—
K	<i>Lobocepon grapsi</i> Nobili, 1905	<i>Grapsus grapsus</i> (Linnaeus, 1758) (Brachyura: Grapsidae)	Taiwan	—	—
K	<i>Megacepon chopardi</i> George, 1947	<i>Perisesarma malpoense</i> (Soh, 1978) (Brachyura: Sesarmidae).	Hong Kong	India; Japan; Thailand	<i>Muradium tetragonum</i> (Fabricius, 1798); <i>Sesarma dehaani</i> (H. Milne Edwards, 1853); <i>Episesarma mederi</i> (H. Milne Edwards, 1853) (Brachyura: Sesarmidae)
K	<i>Megacepon disparatum</i> An et al., 2012a	<i>Metaplagia longipes</i> Stimpson, 1858; <i>Metaplagia</i> sp. (Brachyura: Varniidae)	Hainan Province; Zhejiang Province	—	—
K	<i>Megacepon goetici</i> (Shinno, 1934)	<i>Gaetice depressus</i> (De Haan, 1833) (Brachyura: Varunidae)	Hong Kong	Japan	<i>Gaetice depressus</i> (De Haan, 1833) (Brachyura: Varunidae)
K	<i>Megacepon sexarmae</i> (Pearse, 1930)	<i>Sesarma dehaani</i> H. Milne Edwards, 1853 (Brachyura: Sesarmidae)	Fujian Province	—	—
K	<i>Megacepon sheni</i> An et al., 2012a	<i>Metaplagia sheni</i> Gordon, 1930 (Brachyura: Varunidae)	Hainan Province	—	—
K	<i>Metacepon pleopodata</i> Bourdon and Stock, 1979	<i>Ceratoplax ciliata</i> Stimpson, 1858 (Brachyura: Pilumnidae)	Hong Kong	Amboina (Indonesia)	<i>Heteroplumhus setosus</i> (A. Milne-Edwards, 1873) (Brachyura: Pilumnidae)

TABLE 1
(Continued)

Subfamily	Bopyrid species	Hosts in China	Chinese localities	Other localities	Host(s) elsewhere
O	<i>Minicopenaon crosnieri</i> (Bourdon, 1979a)	<i>Metapenaeopsis philippi</i> (Bate, 1881), <i>Metapenaeopsis provocatoria longirostris</i> Crosier, 1987	Zhejiang Province; Taiwan	Madagascar	<i>Metapenaeopsis andamanensis</i> (Wood-Mason, 1891) (Dendrobranchiata: Penaedae)
O	<i>Minicopenaon intermedium</i> Bourdon, 1981	<i>Metapenaeopsis conger</i> (Wood-Mason, 1891), <i>Metapenaeopsis philippi</i> (Bate, 1881) (Dendrobranchiata: Penaedae)	East China Sea	Philippines	<i>Metapenaeopsis</i> sp. (Dendrobranchiata: Penaedae)
O	<i>Minicopenaon liuriavui</i> An et al., 2013c	<i>Metapenaeopsis conger</i> (Wood-Mason, 1891), <i>Metapenaeopsis philippi</i> (Bate, 1881) (Dendrobranchiata: Penaedae)	East China Sea South China Sea	—	—
A	<i>Minimathelges minutus</i> (Markham, 1992a)	<i>Paguridae</i> sp. (Anomura: Paguridae)	Hong Kong	—	—
K	<i>Onkokepon articulatus</i> An et al., 2006a	<i>Euclosiana unidentata</i> (De Haan, 1841), <i>Leucosia longibrachia</i> Shen and Chen, 1978 (Brachyura: Leucosiidae)	Beibu Gulf	—	—
K	<i>Onkokepon beihuensis</i> An et al., 2006a	<i>Leucosia longibrachia</i> Shen and Chen, 1978 (Brachyura: Leucosiidae)	Beibu Gulf	—	—
O	<i>Orbione halipori</i> Nierszasz and Brender à Brandis, 1923	<i>Metapenaeus ensis</i> (De Haan, 1844), <i>Metapenaeus joyneri</i> (Miers, 1880), <i>Metapenaeus monoceros</i> (Fabricius, 1798) (Dendrobranchiata: Penaedae)	Guangdong Province; Beibu Gulf; Jiaozhou Bay; Hong Kong	Indo-West Pacific	<i>Aristacomorpha foliacea</i> (Risso, 1827) (Dendrobranchiata: Aristidae), <i>Hadopenaeus lucasii</i> (Spence Bate, 1881), <i>Haliporoides sibogae sibogae</i> (de Man, 1907), <i>H. sibogae madagascariensis</i> Crozier, 1978, <i>H. triarthrus triarthrus</i> Stebbing, 1914 (Dendrobranchiata: Solenoceridae), <i>Metapenaeus ensis</i> (De Haan, 1844), <i>M. joyneri joyneri</i> (Miers, 1880), <i>M. monoceros</i> (Fabricius, 1798) (Dendrobranchiata: Penaedae)
O	<i>Orbione penei</i> Bonnier, 1900	“ <i>Penaeus</i> ” sp. (Dendrobranchiata: Penaedae)	Hong Kong	—	—
P	<i>Orthione griffenis</i> Markham, 2004	<i>Austinogebia wuhsiense</i> (Yu, 1931) (Gebidae: Upogebiidae)	Shandong Province	Oregon, USA; Canada; Japan;	<i>Austinogebia marutensis</i> (Sakai, 1986), <i>Upogebia issaeffi</i> (Balss, 1913), <i>Upogebia pugettensis</i> (Dana, 1852) (Gebidae: Upogebiidae)

TABLE I
(Continued)

Subfamily	Bopyrid species	Hosts in China	Chinese localities	Other localities	Host(s) elsewhere
P	<i>Ovionella obovata</i> (Shiino, 1958)	<i>Munida japonica</i> Stimpson, 1858 (Anomura: Munididae)	Taiwan	Japan	<i>Munida japonica</i> Stimpson, 1858 (Anomura: Munididae)
P	<i>Pagurion arrosor</i> An et al., 2013d	<i>Dardanus arrosor</i> (Herbst, 1796) (Anomura: Diogenidae)	South China Sea	—	—
P	<i>Pagurion tuberculata</i> Shiino, 1933	<i>Dardanus aspersus</i> (Berthold, 1846) (Anomura: Diogenidae)	South China Sea	Japan; Philippines; Australia	<i>Calcinus gaimardi</i> (H. Milne Edwards, 1848), <i>C. minutus</i> (Buitendijk, 1937), <i>Dardanus langopodes</i> (Forsskål, 1775), <i>D. scutellatus</i> (H. Milne Edwards, 1848) (Anomura: Diogenidae)
P	<i>Parapagurion calcincola</i> Shiino, 1933	<i>Pagurus aff. hedleyi</i> (Grant and McCulloch, 1906), <i>Pagurus kulkarnii</i> Sankoli, 1962 (Anomura: Paguridae)	Hong Kong	Japan; Gulf of Thailand	<i>Calcinus elegans</i> (H. Milne Edwards, 1836), <i>Paguristes</i> sp. (Anomura: Diogenidae)
O	<i>Parapenaeon expansa</i> Bourdon, 1979a	<i>Penaeus penicillatus</i> Alcock, 1905, <i>Penaeus japonicus</i> Bate, 1888 (Dendrobranchiata: Penaeidae)	Guangdong Province Fujian Province; South China Sea	Madagascar; Australia; Pakistan; New Caledonia; Indonesia; Seychelles	<i>Metapenaeopsis faonizii</i> (Ramadan, 1938), <i>Metapenaeopsis gailliardi</i> Crossner, 1991, <i>Metapenaeopsis mogrenensis</i> consobrina (Nobili, 1904), <i>Metapenaeopsis sinica</i> Liu and Zhong, 1988, <i>Penaeus indicus</i> H. Milne Edwards, 1837, <i>Penaeus longisyllus</i> Kubo, 1943, <i>Penaeus marginatus</i> Randall, 1840, <i>Penaeus merguensis</i> de Man, 1888, <i>Penaeus pfebejus</i> Hess, 1865 (Dendrobranchiata: Penaeidae)
O	<i>Parapenaeon japonica</i> (Thielemann, 1910)	<i>Metapenaeopsis barbata</i> (De Haan, 1844), <i>Metapenaeopsis dura</i> Kubo, 1949, <i>Metapenaeopsis lamellata</i> (De Haan, 1844), <i>Metapenaeopsis mogrensis</i> Rathbun, 1902, <i>Metapenaeopsis novaguineae</i> (Haswell, 1879), <i>Metapenaeopsis toloensis</i> Hall, 1962, <i>Metapenaeopsis stridulans</i> (Alcock, 1905), <i>Penaeus japonicus</i> Bate, 1888 (Dendrobranchiata: Penaeidae)	Beibu Gulf; East China Sea; South China Sea; Hong Kong	Indo-West Pacific	<i>Metapenaeopsis acclivis</i> (Rathbun, 1902), <i>Metapenaeopsis barbata</i> (De Haan, 1844), <i>Metapenaeopsis lamellata</i> (De Haan, 1844), <i>Metapenaeopsis sinica</i> Liu and Zhong, 1988, <i>Metapenaeopsis velutina</i> (Dana, 1852), <i>Metapenaeus monoceros</i> (Fabricius, 1798), <i>Melicerus latisulcatus</i> (Kishinouye, 1896), <i>Mierspenaeopsis sculptilis</i> (Heller, 1862), <i>Penaeus japonicus</i> Bate, 1888, <i>Penaeus longistylus</i> Kubo, 1943, <i>Penaeus semiculatus</i> De Haan,

TABLE 1
(Continued)

Subfamily	Bopyrid species	Hosts in China	Chinese localities	Other localities	Host(s) elsewhere
O	<i>Parapenaeon tertium</i> Nierstrasz and Brender à Brandis, 1932	<i>Metapenaeopsis provocatoria longirostris</i> Crosier, 1987 (Dendrobranchiata: Penaeidae)	Taiwan	Indonesia	1844, <i>Trachysalambria fulva</i> (Dall, 1857) (Dendrobranchiata: Penaeidae) <i>Parapenaeus longipes</i> Alcock, 1905 (Dendrobranchiata: Penaeidae)
O	<i>Parapenaeonella</i> <i>distincta</i> Shiino, 1949b	<i>Metapenaeus affinis</i> (H. Milne-Edwards, 1837), <i>Trachysalambria curvirostris</i> (Stimpson, 1860) (Dendrobranchiata: Penaeidae)	Guangdong Province; Shandong Province	Japan Philippines	<i>Trachysalambria curvirostris</i> (Stimpson, 1860), Unidentified Penaeidae sp. (Dendrobranchiata: Penaeidae)
O	<i>Parapenaeonella</i> <i>minutopoda</i> An et al., 2013b	<i>Atypopeneaus compressipes</i> (Henderson, 1893) (Dendrobranchiata: Penaeidae)	East China Sea; South China Sea	—	—
P	<i>Parasymmetrione</i> <i>tuberculinea</i> An et al., 2010	<i>Clibanarius corallinus</i> (H. Milne Edwards, 1848) (Anomura: Diogenidae)	Xisha (Paracel Islands)	—	—
P	<i>Parasymmetriorbione</i> <i>bicauda</i> An et al., 2013a	<i>Solenocera alicarnaria</i> Kubo, 1949, <i>Solenocera comata</i> Stebbing, 1915 (Dendrobranchiata: Solenoceridae)	Beibu Gulf	—	—
A	<i>Parathelges</i> <i>enoshimensis</i> Shiino, 1933	<i>Spiriopagurus</i> sp. (Anomura: Paguridae)	South China Sea	Japan Korea	<i>Pagurus filholi</i> (de Man, 1887), <i>Pagurus</i> <i>minutus</i> Hess, 1865, <i>Pagurus</i> sp. (Anomura: Paguridae)
P	<i>Parione pachychelii</i> Shiino, 1950	<i>Pachycheloides stevensi</i> Stimpson, 1858 (Anomura: Porcellanidae)	Hong Kong	Japan	<i>Pachycheloides stevensi</i> Stimpson, 1858 (Anomura: Porcellanidae)
P	<i>Parionella decidens</i> Nierstrasz and Brender à Brandis, 1929	<i>Porcellanidae</i> sp. (Anomura: Porcellanidae)	South China Sea	—	—
P	<i>Parionella notexochia</i> Bourdon, 1972b	<i>Eumunida balssi</i> Gordon, 1930 (Anomura: Eumunididae)	East China Sea (32°26'N, 128°37'E)	East China Sea	—
P	<i>Parionina pacifica</i> Nierstrasz and Brender à Brandis, 1929	<i>Galathea</i> sp. (Anomura: Galatheidae)	East China Sea	—	—

TABLE 1
(Continued)

Subfamily	Bopyrid species	Hosts in China	Chinese localities	Other localities	Host(s) elsewhere
P	<i>Parioninella astridae</i> Nierszrasz and Brenner à Brandis, 1930	<i>Pachycheles sculptus</i> (H. Milne Edwards, 1857). <i>Pachycheles</i> sp. (Anomura: Porcellanidae)	Hong Kong	Indonesia; Philippines	<i>Pachycheles sculptus</i> (H. Milne Edwards, 1857), (Anomura: Porcellanidae)
K	<i>Procepon liuruijui</i> An et al., 2009a	<i>Austinogebia wuhsiemensi</i> (Yu, 1931) (Gebiidae: Upogebiidae)	Guangdong Province	—	—
P	<i>Progebiophilus</i> <i>elongatus</i> An et al., 2009a	<i>Nihonotrypaea japonica</i> (Ortmann, 1891) (Axiiidae: Callianassidae)	Shandong Province	—	—
P	<i>Progebiophilus</i> <i>sinicus</i> Markham, 1982	<i>Upogebia major</i> (De Haan, 1841); <i>Upogebia</i> sp. (Gebiidae: Upogebiidae)	Fujian Province; Hong Kong	—	—
P	<i>Pseudione longicauda</i> Shiino, 1937b	<i>Nihonotrypaea japonica</i> (Ortmann, 1891), <i>Nihonotrypaea petalura</i> (Stimpson, 1860) (Axiiidae: Callianassidae)	Shandong Province; Hong Kong	Japan	<i>Callianassa subterranea</i> (Montagu, 1808) (Axiiidae: Callianassidae)
P	<i>Pseudionella</i> <i>spiropaguri</i> An et al., 2013d	<i>Spiropagurus profundorum</i> Alcock, 1905, <i>Spiropagurus spiriger</i> (De Haan, 1849) (Anomura: Paguridae)	South China Sea	—	—
A	<i>Pseudostegias</i> <i>dulcicum</i> Markham, 1982	<i>Diogenes</i> aff. <i>edwardsii</i> (De Haan, 1849) (Anomura: Diogenidae)	Hong Kong	Thailand	<i>Clibanarius merguiensis</i> de Man, 1888 (Anomura: Diogenidae)
A	<i>Pseudostegias</i> <i>seroensis</i> Shiino, 1933	<i>Calcinus laevimanus</i> (Randall, 1840), <i>Clibanarius bernaculatus</i> (De Haan, 1849), <i>Clibanarius ransonni</i> Forest, 1953, <i>Clibanarius striolatus</i> Dana, 1852 (Anomura: Diogenidae)	Hainan Province; Taiwan; Hong Kong	Japan; Thailand	<i>Clibanarius bernaculatus</i> (De Haan, 1849), <i>Clibanarius padavensis</i> de Man, 1888 (Anomura: Diogenidae)
K	<i>Rhopalione sinensis</i> Markham, 1990	<i>Arcotheres sinensis</i> (Shen, 1932), <i>Pinnotheres arrinae</i> Sakai, 1939 (Brachyura: Pinnotheridae)	Fujian Province; Hong Kong	—	—
K	<i>Scyracepion distincta</i> An et al., 2012c	<i>Jonas distinctus</i> (De Haan, 1835) (Brachyura: Corystidae)	South China Sea	—	—
K	<i>Tylokepon biturus</i> An, 2009	<i>Menaethius monoceros</i> (Latrelle, 1825) (Brachyura: Epialtidae)	Hainan Province	—	—
K	<i>Tylokepon bonnieri</i> Siebbing, 1904	<i>Hyastenus diacanthus</i> (De Haan, 1839) (Brachyura: Epialtidae), <i>Enoplolambrus validus</i> (De Haan, 1837) (Brachyura: Parthenopidae)	Beibu Gulf; South China Sea	India	<i>Tylocarcinus styx</i> (Herbst, 1803) (Brachyura: Epialtidae)

TABLE I
(Continued)

Subfamily	Bopyrid species	Hosts in China	Chinese localities	Other localities	Host(s) elsewhere
K	<i>Tylokepon naxiae</i> (Bonnier, 1900)	<i>Hyastenus diacanthus</i> (De Haan, 1839) (Brachyura: Epialtidae)	Hong Kong	—	—
P	<i>Upogebione bidigitatus</i> An et al., 2009a	<i>Upogebia carcinicanda</i> (Stimpson, 1860), <i>Austinogetbia wuhsiensis</i> (Yu, 1931) (Gebidae: Upogebiidae)	Guangdong Province; Shandong Province	—	—

concluded that they represent 20 species in 12 genera belonging to four subfamilies. The names for all species as given in An (2006) are included in the synonymy lists, but with the notation that several names used as new in that dissertation are treated here in synonymy and are not available names. All these species are described (or redescribed) with figures of males and females presented. A key to all genera of Argeinae and Bopyrinae and keys to species of seven bopyrine genera are provided, as is a list (with hosts) of the 82 bopyrid and ionid species currently known to infest Chinese noncaridean decapods.

In addition to the new Chinese species and records, several taxonomic decisions are made in this paper. The genus *Argeia* is reviewed and a new genus is erected for *Argeia lowisi* Chopra, 1923, and *Argeia nierstraszi* Shiino, 1958. *Capitetragnonia* Pearse, 1953, is recognized as the senior synonym of *Probopyria* Markham, 1985. The genus *Parabopyrus* Markham and species *Parabopyrus stellatus* Markham are deemed to be available from Markham, 1982, not Markham 1985. *Hemiarthrus filiformis* Chopra, 1923, is transferred to *Hyperphrixus* Nierstrasz and Brender à Brandis, 1931. The species composition of *Eophrixus* Caroli, 1930, is reviewed, the type species designation is clarified, and two species of *Hypophryxus* Shiino, 1934, are transferred into this genus.

SUMMARY LIST OF NEW TAXA, TAXONOMIC DECISIONS, AND CHINESE RECORDS

Argeinae

Stegoargeia, n. gen.

Stegoargeia lowisi (Chopra, 1923), n. comb.

Stegoargeia nierstraszi (Shiino, 1958), n. comb.

Pseudioninae

Paranikione sibogae, n. gen., n. sp.

Paranikione distorta, n. sp.

Bopyrinae

Bopyrella articulata, n. sp.

Bopyrinella parameces, n. sp.

Bopyrione multifeminae, n. sp.

Bopyroides shiinoi Rybakov and Andeev, 1991
(new Chinese record)

Capitetragnonia Pearse, 1953 (= *Probopyria* Markham, 1985), new synonymy

- Capitetragonia alphei* (Richardson, 1900), n. comb.
Capitetragonia elliptica (Markham, 1992), n. comb.
Parabopyrella cuspidata, n. sp.
Parabopyrella distincta (Nierstrasz and Brender à Brandis, 1923) (new Chinese record)
Parabopyrella elongata (Shiino, 1949) (new Chinese record)
Parabopyrella hodgarti (Chopra, 1923) (new Chinese record)
Parabopyrella symmetros, n. sp.
Septembopyrina tozeumaophila, n. gen., n. sp.

Hemiarthrinae

- Eophrixus brevicaudata* (Chopra, 1923), n. comb.
Eophrixus caudatus, n. sp.
Eophrixus leptochelae (Pillai, 1966), n. comb.
Eophrixus pikei (Bruce, 1968), n. comb.
Sigyn branchialis, n. gen., n. sp.

MATERIALS AND METHODS

Materials for this study originated from the China Comprehensive Oceanographic Survey of the 1950s and 1989–1991. All materials examined are deposited in the Institute of Oceanology, Chinese Academy of Sciences, Qingdao, China (IOCAS), except those specimens studied at the Smithsonian Institution, Washington, D.C. (USNM). IOCAS collection abbreviations are: CIECR, CIEAL, CIEPR, CIEGL, CIEOG, CIEHI, where C = Crustacea, I = Isopoda, E = Epicaridea and the last two letters = host family as follows: AL = Alpheidae, CR = Crangonidae, GL = Glyphocrangonidae, HI = Hippolytidae, OG = Ogyrididae, PR = Processidae. Animals were viewed and drawn using a Zeiss Stemi SV Apo. Males for scanning electron microscope study were fixed in 2.5% glutaraldehyde in 0.2 M Millonig's phosphate buffer at pH 7.4 for 1.5 hr and postfixed in 1% osmium tetroxide in 0.2 M Millonig's buffer for 1 hr. The specimens were then dehydrated through a graded series of ethanol, followed by critical point drying. After sputter coating with colloidal gold, the specimens were examined with a KYKY2800B scanning electron microscope. Synonyms do not include all records of species from sources such as textbooks, government documents, and most

publications where species are mentioned without locality or host data. Bibliographic references are provided for all bopyrid taxa, including those listed in the tables.

SYSTEMATIC ACCOUNT

- Crustacea Brünnich, 1772
 Isopoda Latreille, 1817
 Cymothoida Wägele, 1989
 Bopyridae Rafinesque, 1815
 Argeiinae Markham, 1977

DIAGNOSIS (modified from Markham, 1977): Female: Body rounded, ranging from subtriangular to nearly circular; weakly distorted, up to 20°; all body regions and usually all segments distinct; head much broader than long, oval to fusiform, barbula with two or three tapered lobes on each side, otherwise smooth; maxilliped with articulated palp or tuft of setae in palp area; pereopods small; brood pouch usually wide open; pleon of six pleomeres, first five bearing prominent lateral plates, pleomere 6 with uniramous uropods subequal in size and shape to lateral plates; pleopods uni- or biramous, usually five pairs (occasionally with only first three or four pairs), decreasing in size posteriorly, tubercular or as small oval flaps.

Male: All body regions and pereomeres distinct; head small, broader than long, markedly narrower than pereon; pleon fused, anteriorly as broad as final pereomere, semi-circular to subtriangular, no indications of segmentation; no pleonal appendages.

REMARKS: Some of Markham's (1977) characters used to justify the erection of Argeiinae, such as the presence of lateral plates and uniramous pleopods on females, are not found in all the genera currently placed in this subfamily (see above diagnosis). As a consequence, the monophyly of this subfamily has been questioned by earlier authors (e.g., Adkison et al., 1982; Bourdon and Bruce, 1983). Adkison et al. (1982) considered *Parargeia* Hansen, 1897, to be closer to Pseudioninae and the other genera nearer to Bopyrinae, but the molecular phylogeny of Boyko et al. (2013) showed *Argeia* being the sister taxon to Hemiarthrinae. A molecular analysis of species in all the argeiine genera is needed to

address questions of Argeinae monophyly and validity.

KEY TO GENERA OF ARGEINAE, BASED ON MATURE FEMALES

- 1a. Pleon gradually tapering from pereon ... 2
- 1b. Pleon with pleomere 1 markedly narrower than pereomere 7 ... *Eragia* Markam, 1977
- 2a. Pleomeres fused *Gareia* Bourdon and Bruce, 1983
- 2b. Pleomeres distinct 3
- 3a. Coxal plates present on at least pereomeres 1 to 4, or those pereomeres divided laterally; pereopods reduced; body ovoid, ovate or subtriangular 4
- 3b. Coxal plates absent, no pereomeres prominently divided laterally; pereopods prominent; body nearly circular *Argeiopsis* Kensley, 1974
- 4a. Body broader than long 5
- 4b. Body longer than broad 6
- 5a. Dorsally directed ovate lateral plates present *Stegoalpheon* Chopra, 1923
- 5b. Dorsally directed lateral plates lacking *Stegoargeia*, n. gen. (see below)
- 6a. Brood pouch almost completely enclosed; lateral plates extending farther to sides than width of pereon ... *Parargeia* Hansen, 1897
- 6b. Brood pouch open; lateral plates not extending farther to sides than width of pereon 7
- 7a. Length of lateral plates and uropods not exceeding width of first pleomere *Argeia* Dana, 1852
- 7b. Length of final lateral plates and uropods exceeding width of first pleomere ... *Bopyrosa* Nierstrasz and Brender à Brandis, 1923

Argeia Dana, 1852

DIAGNOSIS (modified from Markham, 1977):

Female: Body ovoid, about 3/4 as wide as long, slightly distorted, all body regions and segments distinct; head subrectangular, wider than long; maxilliped palp setose (sometimes articulated); barbula with two or three short projections on each side; pereomeres either bearing coxal plates or prominently divided laterally; oostegites surrounding but not covering marsupium, oostegite 1 with large rounded posterolateral lobe, not tapered; pleon of six pleomeres, first five produced into prominent oval to lanceolate lateral

plates; pleopods uniramous, exopodites often reduced to knobs on ventral surface, especially posteriorly; uropods uniramous, of same size and shape as lateral plates.

Male: Body approximately three times as long as wide; head usually much narrower than pereon; pereomeres distinctly separated, subequal in width; pleon triangular, ending in rounded point.

TYPE SPECIES: *Argeia pugettensis* Dana, 1852, by monotypy.

OTHER SPECIES: *Argeia atlantica* Markham, 1977.

REMARKS: *Argeia lowisi* Chopra, 1923, and *A. nierstraszi* Shiino, 1958, do not belong in *Argeia*. Females of both *A. pugettensis* and *A. atlantica* have an oostegite 1 with a rounded (not tapered) posterior lobe, the first 5 pleomeres with lateral plates and a pair of uniramous pleopods on pleomeres 1–5. However, females of *A. lowisi* and *A. nierstraszi* both have an elongate and tapered oostegite 1 posterior lobe (similar to that seen in *Stegoalpheon kempfi* Chopra, 1923), no lateral plates or biramous pleopods. Although the oostegite 1 is very similar in *A. lowisi*, *A. nierstraszi*, and *S. kempfi*, the presence of five or six pairs of uniramous lateral plates and five pairs of biramous pleopods in *S. kempfi* precludes the inclusion of *A. lowisi* and *A. nierstraszi* in that genus. Therefore, we erect *Stegoargeia*, n. gen. (type species *Argeia lowisi* Chopra, 1923), for the type species as well as *A. nierstraszi*. The hosts for both *Argeia* species are crangonids, while those of *Stegoalpheon* and *Stegoargeia*, n. gen., are alpheids. The males of all three genera are very similar and suggest a close relationship between the genera.

Argeia pugettensis Dana, 1852

Figure 1

Argeia pugettensis Dana, 1852: 804–805 [Puget Sound, Washington, infesting *Metacrangon munita* (Dana, 1852)]; Dana, 1855: pl. 53, fig. 7a–d; Stimpson, 1857: 511 [Puget Sound, Washington, infesting *M. munita*]; Stebbing, 1893: 336, 337; Calman, 1898: 281; Richardson, 1899a: 868; Richardson, 1899b: 336–337; Bonnier, 1900: 48, 61, 171, 221, 327–328, fig. 55; Richardson, 1900: 308; Gerstaecker, 1901: 185; Richardson, 1904a: 45, 60–64, figs. 35–40 [Bering

Sea to San Diego Bay, California, infesting *Argis alaskensis* (Kingsley, 1883), *Argis crassa* (Rathbun, 1899), *Argis dentata* (Rathbun, 1902), *Argis lar* (Owen, 1839), *Argis ovifer* (Rathbun, 1902), *Crangon alaskensis* Lockington, 1877, *Crangon alba* Holmes, 1900, *Crangon dalli* Rathbun, 1902, *Crangon franciscorum angustimanus* Rathbun, 1902, *Crangon nigromaculata* Lockington, 1877, *Crangon propinquus* Stimpson, 1860, *M. munita*, *Neocrangon communis* Rathbun, 1899]; Richardson, 1904b: 858; Richardson, 1905a: 220 [British Columbia and Alaska, infesting *Argis dentata* (Rathbun, 1902), *C. alaskensis*, *N. communis*]; Richardson, 1905b: 544–550, figs 586–597 [same material as Richardson, 1904a]; Richardson, 1909: 122 [Japan and Korea, infesting *Argis* sp.]; Chopra, 1923: 476–478, 480; Nierstrasz and Brender à Brandis, 1923: 87; Fee, 1927: 16, 40 [British Columbia, infesting *Lissocrangon stylirostris* (Holmes, 1900), *M. munita*]; Nierstrasz and Brender à Brandis, 1929: 16 [California, infesting *Crangon* sp.]; Fraser, 1932: 64; Hiraiwa, 1933: 53; Shiino, 1933: 249, 277–279, fig. 11 [Japan, infesting unknown host]; Gurjanova, 1936a: 219–221, fig. 138 [Kamchatka Peninsula and Bering Sea, infesting *Argis* spp., *Crangon* spp.]; Gurjanova, 1936b: 258; Shiino, 1937a: 299 [Japan, infesting *Crangon affinis* De Haan, 1849]; Ricketts and Calvin, 1939: 185, 281; Baer, 1946: 65; Hatch, 1947: 164, 224, pl. 9, figs. 110–112; Ricketts and Calvin, 1948: 185, 320; Baer, 1951: 69–70; Ricketts and Calvin, 1952: 239, 438; Shiino, 1952: 34; Menzies and Miller, 1954: 141, 153, fig. 65c, d; Shiino, 1958: 56, pl. 3, fig. 8 [Japan, infesting *A. lar*, *C. affinis*, *Neocrangon sagamiensis* (Balss, 1913)]; Oguro, 1961: 43–47, figs. 1–3, 5–7 [Japan, infesting *C. affinis*]; Danforth, 1963: 35, 52, 54, 60, 61, 63, 64, 68, pl. 2, fig. 1, pl. 4, figs. 4–6; Sindermann and Rosenfield, 1967: 351; Bourdon, 1968: 358; George and Strömborg, 1968: 253 [Washington, infesting *A. dentata*, *C. alaskensis*, *Crangon franciscorum* Stimpson, 1856, *Eualus suckleyi* (Stimpson, 1854), *Mesocrangon munitella* (Walker, 1898), *M. munita*, *N. communis*]; Ricketts and Calvin, 1968: 286, 488; Schultz, 1969: 323–324, figs. 516–517;

Danforth, 1970: 41, 42, 56–57, 143, 150, fig. 5a–c; Sindermann, 1970: 171; Strömborg, 1971: 28; Kozloff, 1974: 148; Miller, 1975: 285, 287, 305, pl. 64, fig. 18; Markham, 1977: 112–114 [Washington and California, infesting *C. alaskensis*, *C. franciscorum*, *Neocrangon resima* (Rathbun, 1902)]; Rudy and Rudy, 1979: 122, 124, 126; Beck, 1980: 135; Bourdon et al., 1981: 502; Jay, 1985: i–vii, 1–63 [California, infesting *C. franciscorum*]; Ricketts et al., 1985: 328, 533; Markham, 1986: 156, 161; Bourdon, 1987: 342–343, fig. 17.23; Strathmann, 1987: 436; Kim and Kwon, 1988: 199, 201, 207–210, fig. 6 [Korea, infesting *A. lar*, *C. affinis*, possibly *Crangon hakodatei* Rathbun, 1902]; Jay, 1989: 68–76 [California, infesting *C. franciscorum*]; Sindermann, 1990: 196; Brusca and Wilson, 1991: 148, fig. 2a; Feeney, 1995: 6; Trilles, 1999: 285, 292, 334, fig. 8.13C; Brusca et al., 2001: 20; Shields, 2001: 313; An, 2006: 61–63, fig. 26 [China, infesting *C. affinis*, *Crangon cassiope* de Man, 1906, *Crangon crangon* (Linnaeus, 1758)]; Espinosa-Pérez and Hendrickx, 2006: 237; Chaplin-Ebanks and Curran, 2007: 73; Marin Jarrin, 2007: x, 54, 90–92, 96, 100, 118 [Oregon, infesting *Lissocrangon stylirostris* (Holmes, 1900)]; An et al., 2008: 225–226 [China, infesting *C. affinis*, *C. cassiope*, *C. crangon*]; Liu, 2008: Marin Jarrin and Shanks, 2008: 691; 613–620 [Oregon, infesting *L. stylirostris*]; Romero-Rodríguez and Román-Contreras, 2008: 1207; Wicksten, 2008: 143; Espinosa-Pérez et al., 2009: 229; Williams and An, 2009: 122; Dumbauld et al., 2011: 337; Han and Li, 2010: 228; An, 2011: 130–133, figs. 5.1–5.3 [China, infesting *C. affinis*, *C. cassiope*, *C. crangon*]; Marin Jarrin and Shanks, 2011: 237; Wicksten, 2012: 106; Williams and Boyko, 2012: 7; Penha-Lopes et al., 2013: 35; Rudy and Rudy, 2013: 208, 210, 272; Sherman and Curran, 2013: 1336; Romero-Rodríguez and Román-Contreras, 2014: 470; Seo et al., 2014: 413–417, fig. 2 [Korea, infesting *A. lar*]; Cericola and Williams, 2015: 239.

Argeia pauperata Stimpson, 1857: 511 [San Francisco, California, infesting *C. franciscorum*]; Stebbing, 1893: 415; Calman, 1898: 281; Bonnier, 1900: 61, 171, 221, 328, 381; Gerstaecker, 1901: 185, 266;

- Richardson, 1905b: 551; Chopra, 1923: 476–478; Nierstrasz and Brender à Brandis, 1923: 87; Gifford, 1934: 1–20, pls. 1–5; [California, infesting *C. franciscorum*]; Danforth, 1963: 20, 28, 34; 68, 75, pl. 1, fig. 4, pl. 2, fig. 2, pl. 4, figs. 1–3; Schultz, 1969: 323; Danforth, 1970: 41, 56, 150, fig. 4d–f; Strömborg, 1971: 28; Miller, 1975: 287, 305; Nelson et al., 1986: 121–124.
- Argeja* [sic] *pugettensis* Cornalia and Panceri, 1861: 86, 16.
- ?*Bopyrus* sp. Bate, 1888: 485 [Japan, infesting *C. affinis*].
- Argeia* sp. Calman, 1898: 261, 281 [Washington, USA, infesting *C. affinis*].
- Argeia depauperata* [sic] Richardson, 1899a: 868; Richardson, 1899b: 336, 337; Richardson, 1900: 308.
- Argeia* (?) [spp.] Bonnier, 1900: 221.
- Argeia Calmani* Bonnier, 1900: 171, 221, 329, 381 [Washington, infesting *C. affinis*] (nomen nudum).
- Argeia Pingi* Yü, 1935: 52–53 [Hopei and Shantung Provinces, China, infesting *C. cassiope*].
- Argaeia* [sic] *pugettensis*—MacGinitie and MacGinitie, 1949: 266 [west coast of United States, infesting *C. alaskensis*]; MacGinitie and MacGinitie, 1968: 265–266.

MATERIAL EXAMINED: Infesting *Crangon affinis* De Haan, 1849. 2♀♀, ♂ (CIECR 007601), South Yellow Sea, Stn. 76, 123°30' E, 27°20' N, 45 m, 18 April 1975; 2♀♀, 2♂♂ (CIECR020801), South Yellow Sea, Stn. 208, 127°00' E, 30°30' N, 46 m, 12 July 1968; ♀, ♂ (CIECR104941), Yellow Sea, Stn.10494, 122°00' E, 35°00' N, 52 m, 13 June 2004; 2♀♀, 2♂♂ (CIECR007602), South Yellow Sea, Stn.76, 123°30' E, 27°20' N, 45 m, 10 June 1975; ♀, ♂ (CIECR030601), Yellow Sea, Stn. 306, 126°00' E, 29°30' N, 60 m, 13 November 1957; ♀ (CIECR350801), Yellow Sea, Stn. 3058, 120°00' E, 35°00' N, 81 m, 12 September 1992; ♀, ♂ (CIECR626301); South China Sea, Stn. 6263, 107°30' E, 19°00' N, 77 m, 8 March 1962; ♀, ♂ (CIECR008101), South Yellow Sea, Stn. 81, 122°00' E, 27°10' N, 31 m, 8 June 1975; ♀, ♂ (CIECR005101), South Yellow Sea, Stn. 51, 123°20' E, 28°30' N, 65 m, 18 April, 1975; ♀, ♂ (CIECR000101), South Yellow Sea, Stn.1, 121°00' E, 26°30' N, 75 m, 29 April 1976; ♀, ♂ (CIECR590501),

Shazikou, Qingdao, Shandong Province, 16 May 1959, coll. Fenxuan Zhang; 2♀♀, 2♂♂ (CIECR580901), South Yellow Sea, Stn. 81, 122°00' E, 27°10' N, 31 m, 20 September 1958; 2♀♀, 2♂♂ (CIECR303801), Yellow Sea, Stn. 3038, 123°45' E, 36°00' N, 71.5 m, 10 November 1958.

Infesting *Crangon crangon* (Linnaeus, 1758). 2♀♀, 2♂♂ (CIECR530401), Yantai Fish Market, Shandong Province, 20 April 1953; 2♀♀, 2♂♂ (CIECR570101), Shazikou, Qingdao, Shandong Province, 20 January 1957, coll. Shaowu Wang, Xiuxi Zhang and Zhengang Fan; ♀, ♂ (CIECR550401), Yantai, Shandong Province, 5 April 1955, coll. Engze Yang; ♀, ♂ (CIECR500301), Beidai He, Hebei Province, 5 March 1950.

Infesting *Crangon cassiope* de Man, 1906. 2♀♀, 2♂♂ (CIECR510401), 2♀♀, 2♂♂ (CIECR510402), Yantai, Shandong Province, 6 April 1951.

DESCRIPTION: (CIECR007601): Length 8.12 mm, pereon length 4.58 mm, maximal width 6.2 mm, head length 0.94 mm, head width 1.87 mm, distorted 36° (fig. 1A, B). All segments distinct. Head with narrow frontal lamina, anterior edge of head deeply bilobate (fig. 1A), eyes lacking. Antennule of three articles, antenna of five articles, terminally setose. Maxilliped (fig. 1C, D) with setose palp, plectron sharp, anterior article two times larger than posterior article. Barbula with two pairs of smooth falcate lateral projections on each side (fig. 1E).

Pereon broadest across third pereomere (fig. 1A). Coxal plates very reduced, only visible on first three pereomeres of long side. Dorsolateral bosses on first three pereomeres of longer side, and first four pereomeres on shorter side. Tergal projections on longer side extended, tapering distally to sharp tip. Brood pouch open (fig. 1B). Oostegite 1 (fig. 1F, G) with nearly smooth internal ridge, posterolateral point lacking. Posterior pereopods larger than anterior pairs, ischium with enlarged tubercular lobe, dactyli blunt (fig. 1H, I). Pleon of six pleomeres, first five pleomeres with lateral plates and uniramous pleopods. Sixth pleomere without lateral plates, with uniramous uropods (fig. 1B).

DESCRIPTION: Male (CIECR007601): Length 2.26 mm, maximal width across pereomere 4, 0.71 mm, head width 0.43 mm,




Fig. 1. *Argeia pugettensis* Dana, 1852, female (CIECR007601) (A–I): A. Dorsal view. B. Ventral view. C. Right maxilliped, external view. D. Palp of right maxilliped. E. Right side of barbula. F. Right oostegite 1, external view. G. Right oostegite 1, internal view. H. Right pereopod 7. I. Right pereopod 1. Female (CIECR530401). J. Dorsal view. Male (CIECR007601) (K–N): K. Dorsal view. L. Ventral view. M. Antenna and antennule. N. Pereopod 7. Male (CIECR510401). O. Ventral view of pleon. Scale: 1 mm (A–C, E–L, O), 0.5 mm (D, M, N).

head length 0.20 mm, pleonal length 0.55 mm. (fig. 1K, L). Head elliptical, separated from first pereomere (fig. 1K). Black eyes near posterior edge of head (fig. 1K). Antennule of three articles, antenna of four articles, terminally setose (fig. 1M). Pereomeres

distinct, subequal in width, midventral projections lacking (fig. 1L). Dactyli of first pereopod larger others (fig. 1N). Pleon fused into single ovate piece, pleopods and uropods lacking; anal cone usually lacking (occasionally present) (fig. 1O).

VARIATIONS: The head of some females is so deeply bilobed it looks like a pair of spheres united. One female specimen (CIECR530401) with an extremely bulging head, very extended, terminally sharp tergal projections (fig. 1J) and the surface of the lateral plates are crenulated. One male (CIECR510401) with anal cone on the terminal end of the pleon (fig. 1O).

HOSTS AND LOCALITIES: Infesting *Argis alasensis* (Kingsley, 1883), *Argis crassa* (Rathbun, 1899), *Argis dentata* (Rathbun, 1902), *Argis lar* (Owen, 1839), *Argis ovifer* (Rathbun, 1902), *Crangon affinis* De Haan, 1849, *Crangon alaskensis* Lockington, 1877, *Crangon alba* Holmes, 1900, *Crangon cassiope* de Man, 1906, *Crangon crangon* (Linnaeus, 1758), *Crangon dalli* Rathbun, 1902, *Crangon franciscorum* Stimpson, 1856, *Crangon franciscorum angustimanus* Rathbun, 1902, *Crangon hakodatei* Rathbun, 1902, *Crangon nigromaculata* Lockington, 1877, *Crangon propinquus* Stimpson, 1860, *Eualus suckleyi* (Stimpson, 1854), *Lissocrangon stylirostris* (Holmes, 1900), *Mesocrangon munitella* (Walker, 1898), *Metacrangon munita* (Dana, 1852), *Neocrangon communis* Rathbun, 1899, *Neocrangon resima* (Rathbun, 1902), *Neocrangon sagamiensis* (Balss, 1913) (Crangoniidae), Bering Sea to San Diego, Japan, Korea, China.

REMARKS: This species has an extremely wide range, occurring in both the northeast and northwest Pacific on at least 23 crangonid hosts. Specimens from Japan, Korea, and China have been well described, but northwestern Pacific specimens are much more poorly known. Richardson (e.g., 1905b) described the species in some detail, but a fuller redescription, especially of characters now known to be informative at the species level, is needed. An example of a possible difference between northwestern and northeastern Pacific specimens, based on literature review, is that the northwestern specimens appear to have moderate to pronounced bilobation of the head, while northeastern specimens have a nearly flat head in dorsal view (see, e.g., Gifford, 1934). Additionally, given the wide geographic range of the species, a molecular analysis of specimens from both sides of the Pacific should be undertaken to test the possibility of cryptic species.




Fig. 2. *Argeia* sp., female (CIECR979401) (A–B): A. Dorsal view. B. Right maxilliped, external view. Scale: 1 mm (A); 0.5 mm (B).

There are three hosts of this species in China and one of these, *Crangon cassiope*, is the type host for *Argeia pingi* Yü, 1935, which was only very briefly described, not illustrated, and never reported subsequent to its original description. Markham (1977) thought *A. pingi* might be a synonym of *A. pugettensis*. Because this host has now been recollected and is infested with *Argeia* specimens that do not differ from *A. pugettensis* as currently defined, *A. pingi* is here considered a synonym of *A. pugettensis*. However, if future molecular and/or morphological analyses show any differences warranting recognition of the northeastern and northwestern populations as distinct species, *A. pingi* would likely be the valid name for northwestern Pacific specimens, assuming that all northwestern Pacific specimens are conspecific. However, the type specimens of *A. pingi* appear to be lost and a neotype would be required.

Argeia sp.
Figure 2

Argeia atlantica: An, 2006: 63–64; An et al., 2008: 226; An, 2011: 133–134, fig. 5–4 (not *Argeia atlantica* Markham, 1977).

MATERIAL EXAMINED: Infesting *Crangon affinis* De Haan, 1849, ♀ (CIECR979401), ♂ (CIECR979402), Yellow Sea, Stn. 9794, 123°30'E, 35°30'N, 9 January 2004.

DESCRIPTION: Female (CIECR979401): Length 10.04 mm, pereon length 4.46 mm, maximal width 7.92 mm, head length 0.92 mm, head width 2.31 mm (fig. 2A).

All segments distinct except median regions of pleomeres 1 and 2. Anterior edge of head deeply bilobate, frontal lamina extending beyond head (fig. 2A); eyes lacking. Antennule and antenna of three and five articles, respectively, terminally setose. Maxilliped (fig. 2B) with setose palp, sharp plectron. Barbula with two pairs of simple falcate lateral projections on each side.

Pereon broadest across third pereomere (fig. 2A). Coxal plates, dorsolateral bosses and tergal projections enlarged on first four pereomeres. Tergal projections of pereomeres 3 and 4 acute and posteriorly directed. Brood pouch open. Oostegite 1 with nearly smooth internal ridge, without posterolateral point, posterior margin with setae. Posterior pereopods larger than anterior pairs, ischia without tubercles.

Pleon of six pleomeres, first five bearing lateral plates with serrated margins and uniramous smooth pleopods. Sixth pleomere without lateral plates, with uniramous uropods.

DESCRIPTION: Male (CIECR979402): Length 2.0 mm, maximal width (across pereomere 4) 0.65 mm, head width 0.41 mm, head length 0.16 mm, pleonal length 0.62 mm. Head elliptical, separated from first pereomere; eyes present. Antennule and antennae of three and four articles, respectively. Pereomere segments distinct, subequal in width, midventral projections lacking. Pereopods subequal in size and shape. Pleon fused into single ovate piece, pleopods and uropods lacking.

REMARKS: The female specimen differs from typical *A. pugettensis* in that it has the ischium of each pereopod without any large tubercular swelling and the lateral plates of the pleon have serrated margins. The male is not obviously different from those of *A. pugettensis*. Earlier records of these specimens (An, 2006, 2011; An et al., 2008) identified the specimens as *A. atlantica*. Aside from the geographical distance between China and the western Atlantic, the Chinese specimens do not match *A. atlantica* as females of that species do not have serrated lateral plates and the head is flat (the head of

the Chinese specimen, like those of *A. pugettensis*, is bilobed). Because this female lacks swellings on the ischia and has serrated lateral plates, it may represent an undescribed species. However, as it might also be an aberrant specimen of *A. pugettensis*, more material is needed for study.

Stegoalpheon Chopra, 1923

DIAGNOSIS (modified from Markham, 1977): Female: body ovate, wider than long, slightly asymmetrical; head separated from pereon, large frontal lamina. Pereomeres distinct, all except first subequal in length, first shorter, sixth segment not shorter than others, coxal plates on segments 1–4; dorsolateral bosses indistinct, marsupium open. Oostegite 1 with prominent posterolateral falcate points extending far beyond sides. Pleomere of 6 segments, distinct dorsally and laterally; segments 1–5 (sometimes 6) with dorsally directed lateral plates and biramous pleopods ventrally located and extending to sides of pleon. Uropods uniramous, similar to pleopods in shape.

Male: Head distinct from pereon, eyes large. Pereomeres well defined. Pleon with all segments fused, pleopods and uropods absent.

TYPE SPECIES: *Stegoalpheon kempfi* Chopra, 1923, by original designation.

OTHER SPECIES: None.

REMARKS: Chopra (1923) originally described *Stegoalpheon kempfi* as having five pairs of triramous pleopods. However, Shiino (1951) and Pillai (1966) convincingly showed that the dorsally directed lobes were lateral plates and that the pleopods were biramous. Markham (1977) stated that the type species has five pairs of uniramous lateral plates and five pairs of uniramous pleopods, but this is incorrect. There is clearly some variability in the number of lateral plates, as Chopra (1923) indicated five pairs, which agrees with the present material, while both Shiino (1951) and Pillai (1966) described specimens with six pairs.

Stegoalpheon kempfi Chopra, 1923 Figure 3

Stegoalpheon kempfi Chopra, 1923: 462, 464–467, text fig. 8, pl. 13, fig. 1–7 [India,




Fig. 3. *Stegoalpheon kempfi* Chopra, 1923, female (CIEAL880701) (A–H): **A.** Dorsal view. **B.** Ventral view. **C.** Left antenna and antennule. **D.** Right maxilliped, external view. **E.** Barbula. **F.** Right oostegite 1, external view. **G.** Right oostegite 1, internal view. **H.** Right pereopod 2. Male (CIEAL880701) (I–K): **I.** Dorsal view. **J.** Ventral view. **K.** Antenna and antennule. Scale: 1 mm (A, B, D–G,); 0.5 mm (H–J), 0.25 mm (C, K).

infesting *Alpheus* cf. *lobidens* (De Haan, 1849)]; Shiino, 1951: 26–29, figs. 1–2 [Japan, infesting *Alpheus rapax* Fabricius, 1798]; Pillai, 1954: 20; Pillai, 1966: 188–190, figs. 12–20 [India, infesting *Alpheus* sp.]; Shiino, 1958: 61 [Japan, infesting *Alpheus rapax* Fabricius, 1798]; Rao and Ramaprasad, 1964: 588–589, figs. 1–2 [India, infesting *Alpheus malabaricus* (Fabricius, 1775)]; Shiino, 1972: 8; Kannupandi, 1976: 87–93, figs. 1–3; Markham, 1977: 109–111, 119–120; Adkison et al., 1982: 337; Kensley, 2001: 226; An, 2006: 65–65, fig. 27; An et al., 2008: 225–226, fig. 2 [China, infesting *Alpheus* sp.]; Liu, 2008: 693; Williams and Boyko, 2010: 150; An, 2011: 134–136, figs. 5–5–7 [same material as An et al., 2008].

Stegoalpheon choproai Pillai, 1954: 20 [India, infesting *Alpheus paludicola* Kemp, 1915]; Rao and Ramaprasad, 1964: 588; Markham, 1977: 109, 111, 120; Kensley, 2001: 226; Williams and Boyko, 2010: 150.

Stegoalpheen [sic] *kempi*—Rao and Ramaprasad, 1964: 588 (figure caption).

MATERIAL EXAMINED: Infesting *Alpheus* sp., 1 ♀, 1 ♂ (CIEAL880701), Niuchelun Jiao of Nansha, 116°10'E, 9°36'N, 23 July 1988; 1 ♀, 1 ♂ (CIEAL940901), Banyue Jiao of Nansha, 116°16'E, 8°52'N, 29 September 1994.

DESCRIPTION: Female (CIEAL880701): Length 4.28 mm, maximal width across pereomere 3 4.43 mm, head length 0.88 mm, head width 1.29 mm, almost symmetrical (fig. 3A, B).

Body oval, length 3/4 width. All segments distinct. Head pentagonal, frontal lamina obscure, but anterior margin of the head extended (fig. 3A). Black eyes on lateral corner of anterior margin of head. Antennule of three articles, antennae of four articles, terminally setose (fig. 3C). Maxilliped (fig. 3D) without palp, but tuft of setae indicating palp location. Plectron extend and sharp. Barbulae (fig. 3E) with two pairs of falcate lateral projections on each side, one median small tubercle present.

Pereon broadest across third pereomere (fig. 3A). Coxal plates reduced, dorsolateral bosses on first five pereomeres. Tergal projections of pereomeres 2–4 distinct. Brood pouch completely open (fig. 3B). Oostegite 1 with

smooth internal ridge, posterolateral point rounded laterally (fig. 3F, G). Pereopods short, with blunt dactyli (fig. 3H). Pleon of six pleomeres, first five pleomeres with small, dorsally directed, tubercular lateral plates. Five pairs of biramous ovate pleopods, those on longer side of body much larger than those on shorter side (fig. 3B). Sixth pleomeres with uniramous uropods, similar in shape to pleopods.

DESCRIPTION: Male (CIEAL880701): Length 2.03 mm, maximal width (across pereomere 4) 0.95 mm, head width 0.52 mm, head length 0.34 mm, pleonal length 0.35 mm (fig. 3I, J). Head pentagonal, separated from first pereomere (fig. 3I). Irregular black eyes near postolateral corner of the head (fig. 3I). Antennule of three articles, antenna of seven articles, much longer than antennule, antennulae and antenna terminally setose (fig. 3K). Pereon much wider than head, segments distinct (fig. 3I), all subequal in width. Pereomeres 6 and 7 with midventral projections (fig. 3J). First four pereopods larger than last three, dactyli of first two larger than others (fig. 3J). Pleon fused into single piece, without any pleopods or uropods (fig. 3J).

HOSTS AND LOCALITIES: Infesting *Alpheus* cf. *lobidens* (De Haan, 1849), *Alpheus malabaricus* (Fabricius, 1775), *Alpheus paludicola* Kemp, 1915, *Alpheus rapax* (Fabricius, 1798), *Alpheus* sp. (Alpheidae), India, Japan, China.

REMARKS: As pointed out by Williams and Boyko (2010), *Stegoalpheon choproai* was synonymized with *S. kempi* by Pillai (1966), despite later (Markham, 1977; Kensley, 2001) citations of the species as valid. The present females conform well with Chopra's (1923) holotype although with much smaller dorsally directed lateral plates; however, the present male possesses midventral projections on the last two pereomeres and a wider pereon (allotype male without any midventral projections and a pereon subequal in width compared to the head).

Pseudioninae Codreanu, 1967

DIAGNOSIS: Female: All body segments distinct, head usually fused with frontal lamina. Pereon with coxal plates, dorsolateral bosses,

and tergal projections. Pleopods usually biramous, uropods present.

Male: Head separated from first pereomere, or fused on posterior edge. All pereomeres distinct, often with midventral tubercles. Pleon with one (fused) or six distinct pleomeres, usually with tubercular or flaplike pleopods on first five pleomeres. Uropods sometimes present.

REMARKS: Pseudioninae is largest and most heterogeneous subfamily in Bopyridae, includes 48 genera and 236 species, and may not be monophyletic (Williams and Boyko, 2012). Some species possess putatively primitive characters, such as males having lateral plates on the pleon and well-developed pleopods. All species are branchial parasites and most species infest anomurans, axiideans, and gebiideans, with relatively few being found on carideans and brachyurans.

Paranikione, n. gen.

DIAGNOSIS: Female: Body asymmetrical. Head wider than long, with well-developed frontal lamina. Barbula with two pairs of smooth lateral projections. All pereomeres with coxal plates, first four with dorsolateral bosses. Brood pouch open; internal ridge of oostegite 1 smooth. Pleon with six segments, five pairs of pleopods and tuberculate biramous uropods.

Male: Body segments distinct, pereomeres subequal in width, pleon with five pairs of tuberculate pleopods, sixth pleomere with posterolateral uropods.

TYPE SPECIES: *Paranikione sibogae*, n. sp., by original designation.

OTHER SPECIES: *Paranikione distorta*, n. sp.

ETYMOLOGY: *Paranikione* refers to the close relationship of the new genus to *Nikione* Kensley, 1974. The genus name is feminine.

REMARKS: Females of both species of *Paranikione*, n. gen., have a well-developed frontal lamina, coxal plates on all pereomeres and pleomeres, and biramous pleopods and uropods. These two species are allied to the monotypic genera *Nikione* and *Urobopyrus* Richardson, 1904, which also infest processid shrimp. The new genus is most closely related to *Nikione*, but differs in that females of *Nikione*

have only five pleomeres while those of *Paranikione*, n. gen., have six distinct pleomeres. The key characteristic of *Urobopyrus* is that the female has no lateral plates on the pleomeres, but species of *Paranikione*, n. gen., have well-developed lateral plates on all pleomeres.

Paranikione sibogae, n. sp.

Figure 4

Urobopyrus processae: An, 2006: 69–70, fig. 29 (not *Urobopyrus processae* Richardson, 1904).

MATERIAL EXAMINED: Infesting *Nikoides sibogae* de Man, 1918. Holotype ♀ (CIEPR 615401) and allotype ♂ (CIEPR 615402), South China Sea, Stn. 6154, 110°45'E, 19°15'N, 38 m, 10 July 1959, coll. Fengshan Xu. Paratypes: 1 ♀, 1 ♂ (CIEPR 603001), South China Sea, Stn. 6030, 115°30'E, 21°30'N, 115 m, 13 April 1960, coll. Jingzuo Qu.

OTHER MATERIAL: 1 ♀, 2 ♂♂ (CIEPR 607701), South China Sea, Stn. 6077, 113°00'E, 21°00'N, 54 m, 21 April 1959, coll. Fuzeng Sun; 1 ♀, 1 ♂ (CIEPR 615901), South China Sea, Stn. 6159, 110°30'E, 18°45'N, 31 m, 6 May 1960, coll. Shoupeng Shen.

DESCRIPTION: Holotype female (CIEPR 615401): Length 4.39 mm, maximal width 4.54 mm, head length 0.69 mm, head width 1.32 mm, sinistral body shape, distorted 23° (fig. 4A, B). Body wider than long, left side longer than right, all segments distinct. Head elliptical with bilobate anterior margin, well-developed frontal lamina extending beyond head. Eyes absent (fig. 4A). Antennule of three articles, antenna of five articles, setae not observed. Maxilliped rectangular, with stout and short palp bearing three stout setae distally (fig. 4D). Barbula with two pairs of smooth falcate lateral projections on each side, median pair of small triangular projections near center (fig. 4D).

Pereon broadest across third pereomere. First four pereomeres with dorsolateral bosses, all pereomeres with coxal plates. Coxal plates of left side larger than right (fig. 4A). Brood pouch widely open (fig. 4B). Oostegite 1 (fig. 4E, F) with smooth internal




Fig. 4. *Paranikione sibogae* n. gen., n. sp., holotype female (CIEPR615401) (A–F): **A.** Dorsal view. **B.** Ventral view. **C.** Left maxilliped, external view. **D.** Right side of barbula. **E.** Right oostegite 1, external view. **F.** Right oostegite 1, internal view. Allotype male (CIEPR615402) (G–J): **G.** Dorsal view. **H.** Ventral view. **I.** Pereopod 1. **J.** Pereopod 7. Scale: 1 mm (A, B); 0.5 mm (C–F); 0.25 mm (G–J).

ridge, posterolateral point round, laterally directed, mesiolateral margin setose. Posterior pereopods larger than anterior pairs, carpi and meri smooth, bases of all pereopods produced into smooth lobes. Pleon of six pleomeres, lateral plates well developed. First five pleomeres with biramous pleopods and lateral plates, sixth pleomeres with biramous uropods. Surface of pleopods and uropods tubercular (fig. 4B).

DESCRIPTION: Allotype male (CIEPR 615402): Length 1.50 mm, maximal width across pleomere 4 0.44 mm, head width 0.28 mm, head length 0.16 mm. All pereomeres distinctly segmented (fig. 4G, H). Head semicircular (fig. 4G), dark eyes medially (fig. 4G). Antennule of three articles, with setae on distal article, antenna of five articles, terminally setose (fig. 4H). All pereomeres subequal in width, midventral projections lacking (fig. 4H). First two pereopods each with slightly larger dactylus and longer ischium (fig. 4I, J) than other pereopods. Pleon of six segments, midventral projections lacking, five pleomeres with small tubercular pleopods. Sixth pleomere (pleotelson) with medial anal cone and pair of distolateral uropods with setose terminal margins (fig. 4H).

VARIATION: The paratype female agrees with the holotype in most characters, but is weakly tubercular on the surface of the head and sinistral at 39°. Two immature males (CIEPR607701) are attached to the pleomere of this female. Another female (CIEPR615901) has a white body, as opposed to the typical preserved tan coloration, but this may be an artifact of preservation.

ETYMOLOGY: The specific name, *sibogae*, refers to the host name and honors the Dutch *Siboga* Expedition to Indonesia (March 1899–February 1900) on which bopyrid researcher Hugo Frederik Nierstrasz (1872–1937) served as zoologist. The species is feminine because ships names are considered female.

HOST AND LOCALITY: Infesting *Nikoides sibogae* de Man, 1918 (Processidae), South China Sea, 38–54 m.

REMARKS: See the generic diagnosis for discussion of the differences between this species and those in *Nikione* and *Urobopyrus* and Remarks under *Paranikione distorta*, n. sp.,

for the differences between the two species in the new genus.

***Paranikione distorta*, n. sp.**

Figure 5

Urobopyrus bicornis An, 2006: 70–71, fig. 30 (unavailable name).

MATERIAL EXAMINED: Infesting *Hayashidonus japonica* (De Haan, 1844). Holotype ♀ (CIEPR010801), allotype ♂ (CIEPR010802), eastern Yellow Sea, Stn. 0108, 122°45'E, 31°15'N, 94 m, 15 August 1964, coll. unknown. Paratypes: 1 ♀, 1 ♂ (CIEPR601201), Shantou, Guangdong Province, Stn. 6012, 117°00'E, 22°30'N, 47.3 m, 23 December 1959, coll. Yu.

OTHER MATERIAL: 1 ♀, 1 ♂ (CIEPR625401), Beibu Gulf, Stn. 6254, 107°30'E, 18°30'N, 63.9 m, 10 July 1960, coll. Zhengang Fan.

DESCRIPTION: Holotype female (CIEPR 010801): Length 5.15 mm, maximal width 4.96 mm, head length 0.82 mm, frontal lamina 0.22 mm, head width 1.37 mm, pleon length 1.38 mm, sinistral 67° (fig. 5A, B).

Head triangular, frontal lamina extending beyond sides of head and produced into ear-like lamellae at anterolateral corners. Eyes absent (fig. 5A). Antennule of three articles, antenna of four articles, both terminally setose (fig. 5C). Maxilliped of two segments with short stout nonarticulated palp (fig. 5D), terminal end of palp and adjacent margin fringed with setae (fig. 5E). Barbula with two pairs of smooth falcate lateral projections on each side (fig. 5F).

Pereon broadest across third pereomere (fig. 5A), boundary between pereomeres 2–4 obscure, weakly fused. All pereomeres with coxal plates, those on left side much larger than right. Dorsolateral bosses on first four pereomeres. Brood pouch widely open (fig. 5B). Oostegite 1 visible from ventral view (fig. 5B), two articles subequal in length, smooth internal ridge and nearly straight posterior edge (fig. 5G, H). Pereopods smaller posteriorly (fig. 5I, J), each with five articles, carpi and meri fused, dactyli small. Pleon of six pleomeres, medially fused and sharply distorted sinistrally. First five pleomeres with biramous pleopods and lateral plates. Lateral plates of left side larger than right. Sixth




Fig. 5. *Paranikione distorta* n. sp., holotype female (CIEPR010801) (A–J): A. Dorsal view. B. Ventral view. C. Right antenna and antennule. D. Right maxilliped, external view. E. Palp of maxilliped. F. Left side of barbula. G. Left oostegite 1, external view. H. Left oostegite 1, internal view. I. Left pereopod 4. J. Left pereopod 6. Allotype male (CIEPR010802) (K, L, N, O): K. Dorsal view. L. Ventral view. M. Male (CIEPR010802) dorsal view. N. Left antenna and antennule. O. Left pereopod 5. Scale: 2 mm (A, B); 0.5 mm (D, F–H, K–M,); 0.25 mm (C, E, I, J, N, O).

pleomere with lobate biramous uropods. Ventral surface of pleopods thickly tubercular (fig. 5B), but uropods nearly smooth and lanceolate.

DESCRIPTION: Allotype male (CIEPR010 802): Length 2.15 mm, maximal width (across pleon 4) 0.38 mm, head width 0.31 mm, pleotelson length 0.83 mm. All pereomeres distinctly segmented (fig. 5K, L). Head semicircular (fig. 2K); dark eyes in postero-lateral corners (fig. 5K). Antennule of three articles, antenna of four articles, terminal articles setose (fig. 5N). Pereomeres subequal in width, midventral projections lacking (fig. 5L). All pereopods with six articles and subequal in size (fig. 5O). Pleon of six segments, first five pleomeres with large tubercular pleopods. Sixth pleomere with minute medial anal cone and long posteriorly directed tapering uropods (fig. 5L), distal tips of uropods with setae.

VARIATION: The paratype female (CIEPR60 1201) agrees with the holotype, but the antennae can be seen in dorsal view, and the median part of the pereon bulges outward. The paratype male (CIEPR601201) has a curved posterior margin of the head and its uropods are longer than those of the allotype (fig. 2M). The other female (CIEPR625401) is much larger (total length 6.09 mm) and has a wider frontal lamina and longer pleopods than the types.

ETYMOLOGY: The specific name, *distorta*, refers to the sharply sinistrally distorted pleon of the female.

HOST AND LOCALITIES: Infesting *Hayashidonus japonica* (De Haan, 1844) (Processidae). Yellow Sea, Beibu Gulf, and Guangdong Province, China, 47–94 m.

REMARKS: *Paranikione distorta*, n. sp., has a well-developed frontal lamina, pleon of six segments, lateral plates of the pleon lamellar, and is clearly congeneric with *Paranikione sibogae*, n. sp. The new species differ from *P. sibogae*, n. sp., in having (1) median part of female pleon fused, (2), female lacking acute median projections on the barbula, (3) female with sharply distorted pleon, (4) pereopods of the female with carpi and meri fused, (5) male with large tubercular pleopods and (6) male with long posteriorly directed uropods.

Pseudione Kossmann, 1881

DIAGNOSIS (after Markham, 1985a): Female: Body oval or pyriform; all segments distinct. Frontal lamina moderately developed. Coxal plates reduced; oostegite 1 with rounded posterolateral point. Six pleomeres, first five produced into moderately to greatly developed lateral plates; pleopods biramous, lanceolate; uropods usually uniramous, lanceolate.

Male: Body smoothly tapered anteriorly and posteriorly from midpoint (pleon occasionally abruptly wider or narrower than final pereomere); all body regions distinct. Pereopods subequal, anterior pereopods not markedly larger than others. Pleon 1/5 to 1/3 of total body length, of six pleomeres (specimens of some species with pleomeres variously fused and pleons proportionately shorter); pleopods uniramous, tuberculiform to flaplike; no uropods, posterior border of terminal pleomere slightly to greatly produced into posterolateral points.

TYPE SPECIES: *Pseudione callianassae* Kossmann, 1881, by monotypy.

SPECIES HOSTED BY CARIDEANS: *Pseudione affinis* (G. O. Sars, 1883), *P. ampla* Markham, 1988, *P. clevai* Boyko, 2004, *P. cognata* Markham, 1985, *P. elongata africana* Kensley, 1968, *P. elongata elongata* (Hansen, 1897), *P. indica* Chopra, 1930, *P. magna* Shiino, 1951, *P. parviramus* Adkison, 1988, *P. pontocari* Page, 1985, *P. stylopoda* Boyko, 2004,

REMARKS: The remainder of the 53 species and subspecies of *Pseudione* are parasites of anomurans (26 species), axiideans (10 species), nephropids (3 species), a brachyuran (1 species) and one from an unknown host. The genus is likely paraphyletic, but the type species is very poorly known.

Pseudione tattersalli Nierstrasz and Brender à Brandis, 1923

Pseudione tattersalli Nierstrasz and Brender à Brandis, 1923: 72, 74–76, pl. 5, fig. 8, pl. 9, fig. 36 [Indonesia, infesting unknown host]; Nierstrasz and Brender à Brandis, 1931: 167; Danforth, 1963: 10; Danforth, 1970: 3; Danforth, 1971a: 101; Bourdon, 1972a: 108–110, fig. 3 [Indonesia, infesting *Plesionika ensis* (A. Milne Edwards, 1881)];

Adkison, 1988: 577–578; Høeg and Rybakov, 1992: 604; Markham, 1985a: 14; Boyko, 2004: 678, fig. 1 [Taiwan, infesting *Plesionika fimbriata* Chace, 1985].

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Plesionika ensis* (A. Milne Edwards, 1881) and *Plesionika fimbriata* Chace, 1985 (Pandalidae), Taiwan, Indonesia.

REMARKS: Nierstrasz and Brender à Brandis (1923) did not know the identity of the type host; this was later determined by Bourdon (1972a).

Bopyrinae Rafinesque, 1815

DIAGNOSIS: Female: Head usually fused with first pereomere, frontal lamina absent. Pereomeres often distinct, coxal plates and dorsolateral bosses reduced. Brood pouch often partly or widely open. Pleomeres distinct or fused at least on one side, without lateral plates or uropods. Pleopods smooth, biramous or uniramous.

Male: Head often fused with first pereomere, other pereomeres distinct. Pleon usually fused into single piece, pleopods and uropods often reduced.

REMARKS: Bopyrinae contains 27 genera and 122 species (Boyko, 2014; herein). All are branchial parasites of caridean shrimps.

KEY TO THE 27 GENERA OF
BOPYRINAЕ, BASED ON FEMALE
CHARACTERS (MODIFIED FROM
MARKHAM, 1985a)

- 1a. Pleon sharply rotated, with posterior edge pointing forward . . . *Bathygyge* Hansen, 1897
- 1b. Pleon extending straight back or nearly so . . . 2
- 2a. Uropods present 3
- 2b. Uropods absent 5
- 3a. Biramous uropods, with long, thin rami *Urobopyrus* Richardson, 1904
- 3b. Uniramous uropods, long and wide 4
- 4a. Maxilliped without palp. *Litobopyrus* Markham, 1982
- 4b. Maxilliped with setose palp *Parapleurocrypta* Chopra, 1923
- 5a. Pleopods reduced 6
- 5b. Pleopods well developed 7

- 6a. Pleopods of one side reduced *Palaemonellione* Markham, 1989
- 6b. All pleopods reduced *Ogyridione* Markham, 1988
- 7a. Biramous pleopods 8
- 7b. At least some uniramous pleopods 14
- 8a. Head distinctly extended into anterolateral horns *Capitetragonia* Pearse, 1953
(= *Probopyria* Markham, 1985)
- 8b. Head not anterolaterally extended 9
- 9a. Head and pereon separate 10
- 9b. Head and pereon fused at least medially 13
- 10a. Body subcordate (heart shaped), barbula with two pairs of projections *Probopyrus* Giard and Bonnier, 1888
- 10b. Body not cordate, barbula with single pair of projections 11
- 11a. Oostegite 1 without posterolateral point *Parabopyriscus* Markham, 1982
- 11b. Oostegite 1 with posterolateral point 12
- 12a. Pleon of six distinct pleomeres well separated laterally *Probopyriscus* Markham, 1982
- 12b. Pleon of six pleomeres separated on long side only *Probopyrinella*
Nierstrasz and Brender à Brandis, 1929
- 13a. Pleomeres of at least one side fused *Bopyrella* Bonnier, 1900
- 13b. Pleomeres distinct on both sides *Parabopyrella* Markham, 1985
- 14a. First three pleopods biramous, fourth uniramous *Allobopyrus* Bourdon, 1983
- 14b. All pleopods uniramous 15
- 15a. Five pairs of pleopods 16
- 15b. Fewer than five pairs of pleopods 23
- 16a. Five pairs of oostegites 17
- 16b. Seven pairs of oostegites *Septembopyrina*, n. gen.
- 17a. Coxal plates absent *Probopyrione* Bourdon, 1983
- 17b. Coxal plates present 18
- 18a. Coxal plates on pereomeres 1–4 19
- 18b. Coxal plate on pereomeres 2–4 22
- 19a. Pleopods rudimentary, represented by ill-defined tubercles *Bopyroides* Stimpson, 1864
- 19b. Pleopods flaplike, not rudimentary 20
- 20a. Body ovate *Discorsobopyrus* Boyko, 2004
- 20b. Body elongate 21
- 21a. First pair oostegites symmetrical *Bopyrus* Latreille, 1802
- 21b. First pair oostegites asymmetrical *Bopyrinella* Nierstrasz and Brender à Brandis, 1925
- 22a. Pleomeres fused *Parabopyrus* Shiino, 1934

- 22b. Pleomeres distinct *Probynia*
 Bourdon and Bruce, 1983
- 23a. Pleomeres laterally extended, digitate
 *Bopyrinina* Shiino, 1933
- 23b. Pleomeres rounded laterally 24
- 24a. Pleomeres fused on one side
 *Bopyrina* Kossmann, 1881
- 24b. Pleomeres distinct on both sides 25
- 25a. Oostegite 1 of left and right side asymmetrical *Schizobopyrina* Markham, 1985
- 25b. Oostegite 1 of left and right side symmetrical 26
- 26a. Head fused with first pereomere at least medially *Synsynella* Hay, 1917
- 26b. Head separate from first pereomere
 *Bopyrione* Bourdon and Markham, 1980

Bathygyge Hansen, 1897

DIAGNOSIS (after Markham, 1985a): Female: Body subcircular. Coxal plates large; oostegites about half covering marsupium. Small pleon sharply rotated, of six distinct separated pleomeres, lacking lateral plates, pleopods and uropods biramous.

Male: All pereomeres distinct. Propodus of each pereopod produced into ventrodistal projection receiving tip of dactylus. Pleon fused, ovate, bulbous, about 1/4 total body length, lacking appendages and most traces of segmentation.

TYPE SPECIES: *Bathygyge grandis* Hansen, 1897, by monotypy.

OTHER SPECIES: None.

Bathygyge grandis Hansen, 1897

Figure 6

Bopyrus Faxon, 1895: 140 [material described by Hansen, 1897].

Bathygyge grandis Hansen, 1897: 122–124, pl. 6, fig. 2 [off Nayarit, Mexico, infesting *Glyphocrangon spinulosa* Faxon, 1893]; Richardson, 1899a: 869; Richardson, 1899b: 338; Richard, 1900: 71; Bonnier, 1900: 8, 48, 221, 291–292, 381, text fig. 53; Richardson, 1905b: xx, xlvi, 537–539, 720, fig. 581; Stebbing, 1908: 38, 57–59, 94, pl. 7 (33) [South Africa, infesting *Glyphocrangon sculpta* (Smith, 1882)]; Stebbing, 1910a: 436, pl. 7 [South Africa, infesting *G. sculpta*]; Nierstrasz and Brender à Brandis, 1923: 86; Barnard, 1940: 494; Danforth,

1963: 33, 37, 91, 92, pl. 5, figs. 1–2; Sadoğlu, 1969: 197; Schultz, 1969: 312, fig. 496; Danforth, 1970: 9, 43, 57–58, 149, fig. 5d, e; Holthuis, 1971: 285; Wenner, 1978: 1058 [Middle Atlantic Bight, infesting *G. sculpta* and *Glyphocrangon longirostris* (Smith, 1882)]; Bourdon, 1979c: 510; Markham, 1979: 771–772; Markham 1985a: 19–20, 131 [Virginia, infesting *G. sculpta* and *G. longirostris*]; Markham, 1986: 155–156, fig. 4B; Markham, 1988: 57; Kaufmann et al., 1989: 1882 [Magellan Rise, NE Pacific, infesting *Glyphocrangon vicaria* Faxon, 1896 (the only *Glyphocrangon* listed)]; Salazar-Vallejo and Leija-Tristán, 1990: 429; Leija-Tristán and Salazar-Vallejo, 1991: 1; Markham, 1992b: 3; Espinosa-Pérez and Hendrickx, 2001: 50; Román-Contreras and Soto, 2002: 379; An, 2006: 67–68, fig. 28; An et al., 2007a: 1002–1003, fig. 1 [East China Sea, infesting *G. megalophthalma* de Man, 1918]; Liu, 2008: 691; Román-Contreras, 2008: 93; An, 2011: i, iv, 138–140, figs. 1–1, 5–8–5–10 [East China Sea, infesting *G. megalophthalma*].

? *Bathygyge* sp. Bourdon, 1967c: 857 [Canary Islands, infesting *Glyphocrangon* sp., parasite misidentified as *Gigantione bouvieri* Bonnier, 1900, by Richardson; hyperparasitized by *Rolandiscus serratus* (Bourdon, 1967) (Cabiropidae)]; Lemos de Castro, 1970: 2; Restivo, 1971: 71; Restivo, 1975: 153; Bourdon, 1979c: 510 [Azores, infesting *G. longirostris*]; Bourdon et al., 1981: 498; Rybakov, 1990: 415.

? “bopyrid parasites” Holthuis, 1971: 339 [Nigeria, infesting *G. longirostris*].

? *Munidion* sp. Wicksten, 1979: 222 [San Clemente Basin, California, infesting *G. vicaria*]; Wicksten, 2008: 164 [not *Munidion* sp.].

? “branchial bopyrid” Chace, 1984: 11 [West of Halmahera, Indonesia, infesting *Glyphocrangon faxoni* de Man, 1918], 20 [Philippines, infesting *Glyphocrangon pugnax* de Man, 1918].

Bathygege [sic] *grandis* Campos and Rosa de Campos, 1989: 33.

? “bopyrid isopod” Moore et al., 2003: 368 [Bear Seamount, 39°55'N, 67°30'W; probably infesting *G. sculpta* (as that is the only *Glyphocrangon* listed)]; Ahyong, 2006: 68




Fig. 6. *Bathygyge grandis* Hansen, 1897, female (CIEGL002801) (A–G); A. Dorsal view. B. Ventral view. C. Left maxilliped, external view. D. Left side of barbula. E. Left oostegite 1, external view. F. Left oostegite 1, internal view. G. Right pereopod 1. Male (CIEGL002801) (H–M); H. Dorsal view. I. Ventral view. J. Antennae and antennule. K. Pereopod 1. L. Pereopod 7. Scale: 2 mm (A, B); 1 mm (D–I); 0.5 mm (C, K–M); 0.25 mm (J).

[Norfolk Ridge, Tasman Sea, infesting *Glyphocrangon dimorpha* Komai, 2004]. “bopyrid isopod” Han and Li, 2007: 550 [East China Sea, infesting *Glyphocrangon megalophthalma* de Man, 1918, material examined herein].

MATERIAL EXAMINED: Infesting *Glyphocrangon megalophthalma* de Man, 1918, 1♀, 1♂, 3 cryptoniscus larvae (CIEGL002801), East China Sea, Stn. 28, 126°00'E, 26°10'N, 2000–2150 m, 10 June 1978.

DESCRIPTION: Female: Length 12.19 mm, maximal width across pereomere 3, 11.74 mm, head length 2.38 mm, head width 3.74 mm, pereon length 6.33 mm, distorted approximately 66° (fig. 6A, B).

Body ovate, all pereomeres distinct. Head ovate, frontal lamina well developed, extended, and covering anterior part of head (fig. 6A). Eyes and antennae indistinct. Maxilliped (fig. 6C) with triangular palp, fringed with thick setae. Plectron long and sharp. Barbula (fig. 6D) with two pairs of long falcate lateral projections on each side and one small tubercle near median.

Coxal plates of pereon very long, those of first two pereomeres covering head. Rounded dorsolateral bosses on first four pereomeres. Tergal projections of pereomeres 2–4 distinct (fig. 6A). Brood pouch closed (fig. 6B). Oostegite 1 with two articles, first article rounded with smooth internal ridge, second article with posterolateral point, posterior edge with setae (fig. 6E, F). Pereopods larger posteriorly, subequal in shape (fig. 6H). Pleon short, fused in median portion and abruptly distorted. Pleon of five pleomeres, first four pleomeres with smooth biramous pleopods. Fifth pleomere with uniramous uropods.

Male: Length 5.91 mm, maximal width, across pereomere 4, 1.81 mm, head width 1.00 mm, head length 0.47 mm, pereonal length 3.80 mm. Head elliptical, separated from first pereomere (fig. 6H). Round yellow eyes near posterior edge of head (fig. 6H). Antennule of 3 articles, antenna of five articles, all terminally setose (fig. 6J). Pereon segments distinct (fig. 6H), subequal in width. Pereopods subequal in size. Dactyli of first four pereomeres much larger than others (fig. 6K, L). Carpi of last two peropods long,

other pereopods with rounded carpi (fig. 6I, K, L). First four pereopods larger than last three, dactyli of first two larger than others (fig. 6L). Pleon fused into single piece, one proximal tubercle on medioventral surface of pleon; pleopods and uropods lacking (fig. 6I).

HOSTS AND LOCALITIES: Infesting *Glyphocrangon longirostris* (Smith, 1882), *G. megalophthalma* de Man, 1918, *G. sculpta* (Smith, 1882), *G. spinulosa* (Faxon, 1893) (Crangonidae), off Nayarit, Mexico, Virginia, Middle Atlantic Bight, South Africa, East China Sea. Possibly also on *G. dimorpha* Komai, 2004, *G. saxonii* de Man, 1918, *G. pugnax* de Man, 1918, and *G. vicaria* Faxon, 1896, from Tasman Sea, Indonesia, Philippines and California.

REMARKS: This species is in need of reexamination, as is it unclear whether it is or is not a single species with a circumglobal distribution. Unfortunately, the type female was severely damaged even before it was described and no additional eastern Pacific specimens have been collected. Markham (1985a) deferred to Adkison (cited in Markham, 1985a), who indicated he was going to redescribe the species and discuss its systematic position, but that study was never published. Markham (1985a) thought this species was closest to *Pseudione affinis* (G. O. Sars, 1883), and that it probably should be reassigned to its own subfamily. The present female has three cryptoniscus larvae (fig. 6M) attached to the surface of the oostegites.

Bopyrella Bonnier, 1900

DIAGNOSIS: Female: Body outline elliptical, asymmetrical. Head fused completely or at least medially with first pereomere. Maxilliped with setose palp. Coxal plates very reduced; dorsolateral bosses usually on first four pereomeres. Barbula with two pairs of projections on each side. Brood pouch widely open. All pereomeres fused, segments indicated laterally or dorsally. Four or five biramous flaplike pleopods. Uropods absent.

Male: Head and first pereomere usually fused, juncture indicated by lateral indentations. Pleon much wider than pereon,

especially first pleomere. Pleopods and uropods lacking.

TYPE SPECIES: *Bopyrella thomsoni* Bonnier, 1900, by original designation.

OTHER SPECIES: *Bopyrella articulata*, n. sp., *B. calmani* (Richardson, 1905), *B. harmopleon* Bowman, 1956, *B. malensis* Bourdon, 1980a, *B. moluccensis* Bourdon, 1983, *B. tanytelson* Markham, 1985.

REMARKS: Bonnier (1900) erected this genus for *Bopyrella thomsoni* infesting *Alpheus strenuus* Dana, 1852, from "îles des Amis" (= Tonga). Besides the type species, Bonnier (1900) questionably included three other species in *Bopyrella*: *Bopyrus palaemonis* Risso, 1816 (nomen dubium), *Bopyrina nitescens* Giard and Bonnier, 1890 (nomen nudum), and *Bopyrus alphei* Giard and Bonnier, 1890 (nomen nudum; = *Bopyrus alphei* Richardson, 1900 = *Capitetragonia alphei* (Richardson, 1900), n. comb., herein). Bourdon (1980a) reviewed *Bopyrella* and divided it into three groups according to the degree of fusion of the female pleon. Markham (1985a) concluded that the degree of fusion of the female pleon was important at the generic level and transferred 30 species in *Bopyrella* to other genera (e.g., *Probopyria* [= *Capitetragonia*], *Parabopyrella*, *Probopyrinella*, and *Synsynella*). He retained only five species in *Bopyrella*, all parasitizing alpheid hosts: *B. thomsoni*, *B. malensis* Bourdon, 1980a, from the Maldives, *B. macginitiei* Shiino, 1964, from California, *B. harmopleon* Bowman, 1956, from Venezuela, and *B. calmani* (Richardson, 1905), from California. Sassaman et al. (1984) regarded *B. macginitiei* as a synonym of *B. calmani*. Bourdon (1983) described *B. moluccensis* as a new species from the Moluccas and Markham (1985b) described *B. tanytelson* from Thailand. Therefore, the genus *Bopyrella* currently includes six well-described species (Boyko, 2014) plus the poorly known *B. palaemonis*. An eighth species is described herein.

Bopyrella articulata, n. sp.

Figure 7

Bopyrella glabra An, 2006: 72–73, fig. 31
(unavailable name).

MATERIAL EXAMINED: Infesting *Alpheus hoplocheles* Coutière, 1897. Holotype ♀ (CIEAL570603), allotype ♂ (CIEAL570604), Dazhou Island, Hainan Province, 111°20'E, 18°40'N, 11 June 1957. Paratypes: 1 ♀ (CIEAL570605), 1 ♂ (CIEAL570606), Dazhou Island, Hainan Province, 111°20'E, 18°40'N, 11 June 1957.

DESCRIPTION: Holotype female (CIEAL570603): Length 11.24 mm, maximal width 7.06 mm, head length 2.57 mm, head width 2.8 mm, dextral 29° (fig. 7A, B).

Head subquadrate, fused with pereomere 1 medially, but lateral boundary visible. Frontal lamina lacking, small eyes in lateral corners (fig. 7A). Antennule of three articles, basal article much larger than second article, terminal article with setae. Antenna of four articles, nonsetose (fig. 7C). Maxilliped with large three-segmented palp (fig. 7D), palp and adjacent margin with setae (fig. 7E). Barbulula with two pairs of falcate lateral projections on each side (fig. 7F).

Pereon segments distinct, broadest across third pereomere (fig. 7A). First four pereomeres with coxal plates and dorsolateral bosses slightly larger on longer side (fig. 7A). Brood pouch completely open, oostegite 1 visible in ventral view (fig. 7B). Oostegite 1 (fig. 7G, H) with irregular anterior margin, internal ridge bearing many small projections, postero-lateral point blunt laterally, curved on posterior edge. Pereopods subequal in size and structure (fig. 7H), dactyli blunt. Pleon lateral margin completely fused, but four obscure segments radiate around median of fused pleon. Pleon with five flaplike biramous pleopods, endopodite of pleopod 1 much longer than others (fig. 7B); uropods lacking (fig. 7B).

DESCRIPTION: Allotype male (CIEAL570604): Length 3.25 mm, maximal width (across pleon 1) 1.49 mm, head width 0.75 mm, head length 0.42 mm. All pereon segments distinct, pleon fused medially (fig. 7I, J). Head ovate with curved posterior edge (fig. 7I). Eyes mediolateral (fig. 7I). Antennae and antennules of three articles each, not visibly setose (fig. 7K). Pereomeres almost equal in width, each with small midventral projection (fig. 7J). All pereopods with six articles, dactyli of first three pereopods much larger than others (fig. 7J, L). Pleon of six segments, dorsomedian region weakly fused, but distinct in ventral view.




Fig. 7. *Bopyrella articulata* n. sp., holotype female (CIEAL570603) (A–H): A. Dorsal view. B. Ventral view. C. Left antenna and antennule. D. Left maxilliped, external view. E. Palp of maxilliped. F. Left side of barbula. G. Left oostegite 1, external view. H. Left oostegite 1, internal view. Allotype male (CIEAL570604) (I–L): I. Dorsal view. J. Ventral view. K. Right antenna and antennule. L. Left pereopod 4. Scale: 2 mm (A, B); 1 mm (G, H); 0.5 mm (D–F, I, J); 0.25 mm (C, K, L).

Pleomere 1 widest, with midventral tubercle; pleopods and uropods lacking (fig. 7J).

VARIATION: Paratype female (CIEAL 570605) immature, total length 7.14 mm, widest pereomere 4.68 mm, almost symmetrical. Paratype male (CIEAL 570606) mature, similar to allotype.

ETYMOLOGY: The specific name, *articulata*, refers to the maxilliped of the female with its three-segmented palp.

HOST AND LOCALITY: Infesting *Alpheus hoplochelus* Coutière, 1897 (Alpheidae), Hainan Province, China.

REMARKS: *Bopyrella articulata* has the female head fused with the first pereomere, a triarticulated maxilliped palp, a fused pleon, and four pairs of biramous pleopods. The first pleomere of the male is much wider than the other pereomeres and pleomeres. The new species is most related to *B. thomsoni*, which has both a similar (complete) fusion of the female pleomeres as well as a similar male pleon shape, but (1) the maxilliped of the new species has a triarticulated palp (*B. thomsoni* female with nonarticulated palp) and (2) male of *B. articulata* with midventral tubercles on all pereomeres and pleomere 1 (lacking in *B. thomsoni*). The new species and *B. thomsoni* are distinguished from the other five species of *Bopyrella* by having the pleon completely fused, without any lateral indentations indicating segmentation on either side.

Bopyrella tanytelson Markham, 1982

Bopyrella tanytelson Markham, 1985b: 20–23, fig. 10, table 1 [Thailand, infesting *Alpheus* sp.]; Markham, 1990: 559–560 (Hong Kong, infesting *Synalpheus streptodactylus* Coutière, 1905); Markham, 1992a: 298, table 1; Kensley, 2001: 223; Li, 2003: 139, 154, 157; Liu, 2008: 691.

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Synalpheus streptodactylus* Coutière, 1905, *Alpheus* sp. (Alpheidae), Hong Kong, Thailand.

REMARKS: As noted by Markham, the mentions by Miya (1972) and Monod (1976) of bopyrids on *Synalpheus streptodactylus* from Japan and New Caledonia, respectively, could be *Bopyrella tanytelson*, but no specimens

from these collections have ever been critically examined.

KEY TO THE SEVEN SPECIES OF *BOPYRELLA* BONNIER, 1900, BASED ON FEMALES

- | | | |
|-----|--|--------------------------------------|
| 1a. | Four pairs of pleopods | 2 |
| 1b. | Five pairs of pleopods | 3 |
| 2a. | Pereomeres distinct, median part of pleon fused, but pleomeres indicated by lateral indentations | <i>B. moluccensis</i> Bourdon, 1983 |
| 2b. | Pereon fused medially, pleon fused but with lateral indentations on long side | <i>B. malensis</i> Bourdon, 1980a |
| 3a. | Pleon completely fused, without lateral indentations | 4 |
| 3b. | Pleon incompletely fused, lateral indentations on at least one side | 5 |
| 4a. | Maxilliped with nonarticulated palp | <i>B. thomsoni</i> Bonnier, 1900 |
| 4b. | Maxilliped with triarticulated palp | <i>B. articulata</i> , n. sp. |
| 5a. | Maxilliped palp articulated | <i>B. harmopleon</i> Bowman, 1956 |
| 5b. | Maxilliped palp nonarticulated | 6 |
| 6a. | Pleomeres 1–4 distinct, last two pleomeres fused | <i>B. calmani</i> (Richardson, 1905) |
| 6b. | Pleomeres indistinct, indicated only by lateral indentations on long side | <i>B. tanytelson</i> Markham, 1985 |

Bopyrinella Nierstrasz and Brender à Brandis, 1925

DIAGNOSIS: Female Body long and moderately distorted. Head fused with first pereomere at least medially. Maxilliped without palp, or with setose palp. Narrow coxal plates and obscure dorsolateral bosses on long side. Five or seven pairs of oostegites, only oostegite 1 well developed, right and left oostegites not symmetrical, other oostegites reduced. Pleon of six segments laterally distinct but fused dorsally and sometimes ventrally. Five pairs of flaplike uniramous pleopods. Uropods absent.

Male: Body much longer than wide. Head fused with first pereomere on posterior margin. Pereomeres distinct on dorsal and lateral sides, midventral tubercles lacking. Pleon of six segments, fused at least on ventral surface. Pleopods reduced, no uropods.

TYPE SPECIES: *Bopyrinella antillensis* Nierstrasz and Brender à Brandis, 1925 (= *Bopyrina thorii* Richardson, 1904), by monotypy.

OTHER SPECIES: *Bopyrinella albida* Shiino, 1958, *B. nipponica* Shiino, 1936, *B. parameces*, n. sp., *B. stricticauda* Monod, 1933.

REMARKS: Nierstrasz and Brender à Brandis (1925) erected *Bopyrinella* with the type species *B. antillensis* parasitizing *Thor floridanus* Kingsley, 1878, from Curaçao. Markham (1985a) regarded *B. antillensis* as a junior synonym of *Bopyrina thorii* Richardson, 1904, known from the same host in Florida; he then removed *B. thorii* from *Bopyrina*, making the correct name for the type species *Bopyrinella thorii*. Monod (1933) described “*Bopyrinella* (?) *stricticauda*” from an unidentified alpheid in the Gulf of Suez, but while Markham (1985a) thought it did not belong to *Bopyrinella*, he did not specify what characters indicated that it was not in the correct genus nor did he state where it should be placed. Based on the description and illustrations of Monod (1933), the species may be correctly placed in *Bopyrinella*, but some characters (particularly those of the mouthparts) are not well described. Shiino (1936) described *Bopyrinella antillensis* var. *nipponica* infesting *Spirontocaris rectirostris* (= *Heptacarpus rectirostris* (Stimpson, 1860)) from Japan and Markham (1985a) raised it to a full species as *B. nipponica*. Shiino (1958) described *B. albida* infesting *Athanass kominatoensis* Kubo, 1942 (= *Arete indicus* Coutière, 1903), from Japan. Therefore, there are currently four species in *Bopyrinella*, two on hosts in Hippolytidae and two from hosts in Alpheidae. A new species from a new host species and family, Ogyrididae, is described herein.

Bopyrinella albida Shiino, 1958

Bopyrinella albida Shiino, 1958: 62–64, fig. 17 [Japan, infesting *Arete indicus* Coutière, 1903]; Bourdon, 1968: 170; Shiino, 1972: 8; Markham, 1985a: 81; Markham, 1985b: 3, 25–28, 62, fig. 12, table 1 [Thailand, infesting *Athanass dimorphus* Ortmann, 1894]; Markham, 1990: 559 [Hong Kong, infesting *Arete dorsalis* Stimpson, 1860]; Bruce, 1990: 585; Nakashima, 1995: 12, table 3 [Japan, infesting *A. indicus*]; Kensley, 2001: 223; Li, 2003: 139, 154, 157; Liu, 2008: 691.

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Arete indicus* Coutière, 1903, *Athanass dimorphus* Ortmann, 1894, *Arete dorsalis* Stimpson, 1860 (Alpheidae), Japan, Thailand, Hong Kong.

REMARKS: The type host was originally cited as *Athanass kominatoensis* Kubo, 1942, which is treated as a doubtful synonym of *Arete indicus* by De Grave and Fransen (2011). *Bopyrinella albida* is one of only three species of bopyrines known to possess seven pairs of oostegites.

Bopyrinella parameces, n. sp.

Figure 8

Bopyrinella parameces An, 2006: 76–77, fig. 33 (unavailable name).

MATERIAL EXAMINED: Infesting *Ogyrides orientalis* (Stimpson, 1860). Holotype ♀ (CIEOG501201), allotype ♂ (CIEOG501202), Huiquan Bay, Qingdao, Shandong Province, 120°20'E, 36°00'N, 10 December 1950.

DESCRIPTION: Holotype female (CIEOG 501201): Length 5.66 mm, maximal width 2.41 mm, head length 0.67 mm, head width 0.69 mm, pereon length 2.15 mm, pleon length 2.88 mm (fig. 8A, B).

Body width about 1/3 length, pereon length subequal to pleon length, flattened, calceiform, slightly distorted. Head oval, without frontal lamina, fused with first pereomere medially, eyes lacking (fig. 8A). Antennule reduced to single article, antenna lacking (fig. 8B). Maxilliped without palp (fig. 8C), spur blunt and short. Barbula (fig. 8D) with two pairs of smooth projections on each side, flattened near center.

Pereomeres distinct, broadest across third and fourth pereomeres (fig. 8A). Narrow coxal plates on left side of all pereomeres, right side only on first pereomere. Dorsolateral bosses on first four pereomeres of left side, and only first pereomere of right side. Brood pouch widely open (fig. 8B), oostegite 1 almost symmetrical except for posterolateral point (fig. 8B, E, F), small and depressed anterior article, large posterior article. Internal ridge smooth, posterolateral point rounded. Other four pairs of oostegites small, triangular (fig. 8B). All pereopods subequal in size and structure (fig. 8G), carpi and meri smooth,




Fig. 8. *Bopyrinella parameces* n. sp., holotype female (CIEOG501201) (A–G): A. Dorsal view. B. Ventral view. C. Right maxilliped, external view. D. Right side of barbula. E. Right oostegite 1, external view. F. Right oostegite 1, internal view. G. Right pereopod 2. Allotype male (CIEOG501202) (H, I): H. Dorsal view. I. Ventral view. Scale: 2 mm (A, B); 1 mm (E, F); 0.5 mm (C, D, G–I).

bases of all pereopods stout. Pleon of six pleomeres, elongated, first pleomere identifiable from dorsal suture, other pleomeres detected only from lateral indentations. Five pairs of uniramous flaplike pleopods, posteriorly becoming smaller (fig. 8B). Sixth pleomere indented posteromedially, uropods lacking (fig. 8B).

DESCRIPTION: Allotype male (CIEOG501 202): Length 0.87 mm, maximal width (across pereon 3) 0.29 mm, head width 0.18 mm, head length 0.14 mm. All pereon segments distinct, pleomere distinct in dorsal view (fig. 8H, I). Head elliptical, fused with first pereomere (fig. 8H), eyes absent (fig. 8H). Antennule of three articles, antenna of four articles, terminally nonsetose (fig. 8I). Pereomeres subequal in width, midventral projections lacking (fig. 8H). All pereopods small, posteriorly becoming slightly larger (fig. 8I). Pleon sharply narrow, pleomeres distinct only on dorsal surface, no pleopods or uropods (fig. 8I).

ETYMOLOGY: The specific name, *parameces*, refers to the female's long elliptical, calceiform, outline.

HOST AND LOCALITY: Infesting *Ogyrides orientalis* (Stimpson, 1860) (Ogyrididae), Shandong Province, China.

REMARKS: The present specimens agree with three of the other species of *Bopyrinella* in having (1) female with relatively long body outline, (2) only oostegite 1 well developed and different in shape on left and right sides, (3) brood pouch completely open, (4) five pairs of flaplike uniramous pleopods and lacking uropods. The fourth species, *B. albida*, has rudimentary sixth and seventh oostegites, a character state that was surprisingly not emphasized by Shiino (1958) or Markham (1985b) as this was the first recorded species in Bopyrinae bearing seven pairs of oostegites. Previously reported hosts of *Bopyrinella* species are from Hippolytidae and Alpheidae, but the present host is in Ogyrididae. All three families, however, belong to Alpheoidea. The only other bopyrid known to infest a species in Ogyrididae is *Ogyridione caroliniana* Markham, 1988, which is not at all similar to the new species. *Bopyrinella parameces*, n. sp., is most related to *Bopyrinella albida* Shiino, 1958, but can be distinguished from it by (1) the new species lacks a frontal lamina (*B.*

albida with large frontal lamina), (2) the body of the new species is only slightly distorted (*B. albida* with body highly distorted), (3) the oostegite 1 of the new species differ only slightly from each other in shape (*B. albida* with very different oostegite 1), (4) the new species has only five pairs of oostegites (*B. albida* with seven pairs), (5) the new species with only one pair of antennae (antennule) (*B. albida* with antennule and antennae), (6) male of the new species without pleopods (*B. albida* with tubercular pleopods), and (7) the new species lacks eyes (*B. albida* with eye spots).

KEY TO FOUR SPECIES OF *BOPYRINELLA* NIERSTRASZ AND BRENDER À BRANDIS, 1925, BASED ON FEMALES (*BOPYRINELLA* *STRICTICAUDA* MONOD, 1933, NOT INCLUDED IN KEY)

- | | | |
|-----|---|------------------------------------|
| 1a. | Four pairs of pleopods | <i>B. thori</i> (Richardson, 1904) |
| 1b. | Five pairs of pleopods | 2 |
| 2a. | Maxilliped with setose palp | <i>B. nipponica</i> Shiino, 1936 |
| 2b. | Maxilliped without palp | 3 |
| 3a. | Head with frontal lamina, seven pairs of oostegites | <i>B. albida</i> Shiino, 1958 |
| 3b. | Head without frontal lamina, five pairs of oostegites | <i>B. parameces</i> , n. sp. |

Bopyrione Bourdon and Markham, 1980

DIAGNOSIS: Female: Body outline ovate, slightly distorted. Head separated from first pereomere, but deeply embedded into pereon. Maxilliped without palp. Barbula with two pairs of falcate lateral projections on each side. Brood pouch medially open or closed, oostegite 1 with long posterolateral point and smooth internal ridge. Pleon of five or six obscure pleomeres, lateral margins almost entire. Four or five pairs of uniramous flap-like pleopods, lateral plates and uropods lacking.

Male: Body longer than wide, pereomeres distinct, pleomeres fused. Most or all pereomeres with midventral tubercles, pleopods and uropods lacking.

TYPE SPECIES: *Bopyrione synalphei* Bourdon and Markham, 1980, by original designation.

OTHER SPECIES: *Bopyrione longicapitata* Markham, 1982, *B. multifeminae*, n. sp., *B. toloensis* Markham, 1982, *B. woodmasoni* (Chopra, 1923).

REMARKS: Bourdon and Markham (1980) erected *Bopyrione* for *B. synalphei* infesting *Synalpheus goodei* Coutière, 1909, *S. bousfieldi* Chace, 1972, and *S. pectiniger* Coutière, 1907, plus three other unidentified *Synalpheus* sp. (sp. near *S. goodei*, sp. near *S. longicarpus* (Herrick, 1891) and *Synalpheus* sp.) from the western Atlantic Ocean (Gulf of Mexico off Florida, Haiti, and Curaçao). *Bopyroides woodmasoni* Chopra, 1923, infesting *Synalpheus* sp. from India, was transferred to *Bopyrione* by Bourdon and Markham (1980). Markham (1982) later described two new species from Hong Kong: *B. longicapitata*, infesting an *Alpheus* sp. questionably identified as *Alpheus lobidens* De Haan, 1849, and *B. toloensis* infesting an *Alpheus* sp. questionably identified as *Alpheus microstylus* (Bate, 1888). All hosts of the four previously known *Bopyrione* species are in Alpheidae.

Bopyrione longicapitata Markham, 1982

Bopyrione longicapitata Markham, 1982: 348–351, figs. 14, 15 [Hong Kong, infesting *Alpheus lobidens?* De Haan, 1849]; Markham, 1985a: 59; Markham, 1992a: 285 [Hong Kong, infesting *Alpheus bisincisus* De Haan, 1849]; Li, 2003: 140, 157; Liu, 2008: 691.

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Alpheus lobidens?* De Haan, 1849, and *Alpheus bisincisus* De Haan, 1849 (Alpheidae), Hong Kong.

REMARKS: The type host was only provisionally identified as *A. lobidens* in Markham (1982).

Bopyrione multifeminae, n. sp.

Figures 9, 10

Bopyrione multifemina An, 2006: 87–88, fig. 41, pl. 5 (unavailable name).

MATERIAL EXAMINED: Infesting *Alpheus* sp., holotype ♀ (CIEAL991101), allotype ♂ (CIEAL991103), Jiaozhou Bay, Stn. 1, 120°21'E, 36°10'N, 6 m, 10 November 1999,

coll. Zhang. Paratypes: 1 ♀ (CIEAL991102), same locality and same branchial chamber as holotype; 3♀♀, host infested in both branchial chambers (CIEAL981101), 1 ♂ (CIEAL981102), Jiaozhou Bay, Stn. 8, 120°14'E, 36°02'N, 20 m, 6 November 1998, coll. Zhang; 2 ♀♀, 1 ♂ host infested in both branchial chambers (CIEAL111941), southern Yellow Sea, Stn. 11194, 120°30'E, 34°30'N, 17 m, 14 June 2004, coll. Hongfa Wang; 1 ♀, 1 ♂ from right branchial chamber (CIEAL111942), southern Yellow Sea, Stn. 11194, 120°30'E, 34°30'N, 17 m, 14 June 2004, coll. Hongfa Wang.

DESCRIPTION: Holotype female (CIEAL991101): Length 5.01 mm, maximal width 3.12 mm, head length 1.56 mm, head width 1.38 mm (fig. 9A, B).

Head triangular, posteriorly acute, distinctly separated from first pereomere. Frontal lamina and eyes lacking (fig. 9A). Antennule of three articles, antenna of four articles, both nonsetose (fig. 9C). Maxilliped triangular, without palp, spur sharp (fig. 9D). Barbula with two pairs of acute falcate lateral projections on each side, smooth near median (fig. 9E).

Pereon broadest across third pereomere (fig. 9A). All pereomeres without coxal plates or dorsolateral bosses. Brood pouch partly open (fig. 9B). Oostegite 1 (fig. 9F, G) without median groove, internal ridge smooth. First article of oostegite 1 much wider than second article. Posterolateral point curved posteriorly. Pereopods subequal in size and shape (fig. 9H), bases of all pereopods produced into small lobes. Pleon of six pleomeres, margins entire, first five bearing flaplike uniramous pleopods, posteriorly smaller, margins undulating (fig. 9B, I). Uropods lacking (fig. 9B).

DESCRIPTION: Allotype male (CIEAL991103): Length 1.02 mm, maximal width (across pereon 1) 0.40 mm, head width 0.32 mm, pleonal length 0.24 mm. All dorsal surfaces setose (fig. 10A), pereon segments distinct, pleon fused into single piece (fig. 9J, 10A). Head elliptical, posterior edge curved (fig. 9J); black eye pigment present on posterolateral corners (fig. 9J). One pair of antennae (? antennules) of three articles, setose (fig. 10B). Pereomeres almost subequal in width, first pereomere widest (fig. 9J), midventral




Fig. 9. *Bopyrione multifeminae* n. sp., holotype female (CIEAL991101) (A–I): **A.** Dorsal view. **B.** Ventral view. **C.** Right antenna and antennule. **D.** Right maxilliped, external view. **E.** Right side of barbula. **F.** Right oostegite 1, external view. **G.** Right oostegite 1, internal view. **H.** Right pereopod 2. **I.** Right pleopod 3. Allotype male (CIEAL991103) (**J**): **J.** Dorsal view. Scale: 1 mm (A, B); 0.25 mm (C–J).

projections (fig. 10B) lacking. All pereopods with platelike scales on ventral surface (fig. 10C), first three pereopods largest, all carpi and meri nearly fused (fig. 10D, E). Pleon fused into single piece, only first pleomeres visible with dorsal indentations, others completely fused; pleopods and uropods lacking (fig. 9J).

VARIATION: All paratype females agree with the holotype except one immature female

(CIEAL111941) that has a proportionally longer body and the brood pouch completely closed.

ETYMOLOGY: The specific name, *multifeminae*, refers to the occupation of the same branchial chamber of the host by two females.

HOST(S) AND LOCALITIES: Infesting *Alpheus* sp. (likely more than one species) (Alpheidae), Shandong Province and southern Yellow Sea, China, 6–20 m.




Fig. 10. *Bopyrione multifeminae* n. sp. SEM image of male (CIEAL981102) (A–E): A. Ventral view of the male. B. Right antenna and antennule. C. Right pereopod 1. D. Left pereopods 1–3. E. Left pereopod 5. Scale: 100 µm (A, D, E); 10 µm (B, C).

REMARKS: *Bopyrione multifeminae*, n. sp., shares with the other four species in *Bopyrione*: female with head deeply embedded into the pereon, maxilliped without palp, barbula with two pairs of falcate lateral projections on each side, uniramous pleopods; lateral plates and uropods lacking. The new species is most closely related to *B. longicapitata*, but (1) the female of *B. multifeminae*, n. sp., lacks eyes and a frontal lamina (*B. longicapitata* female with eyes and frontal lamina), (2) *B. multifeminae*, n. sp., female without any coxal plates or dorsolateral bosses (*B. longicapitata* female with reduced coxal plates), (3) *B. multifeminae*, n. sp., female pleopods small and flaplike, not reaching the median of the pleon (*B. longicapitata* female with larger flaplike pleopods, extending to the median of the pleon), (4) *B. multifeminae*, n. sp., male without midventral tubercles (*B. longicapitata* male with midventral tubercles on all pereomeres). Males with a single pair of antennae occur in both *B. multifeminae*, n. sp., and *B. toloensis*. It is typical for bopyrids to occur with one female and one male in any occupied branchial chamber, occasionally with more than one male accompanying a female, but more than one collection of *B. multifeminae*, n. sp., was made with the unusual condition of two females and one male in the same branchial chamber.

Bopyrione toloensis Markham, 1982

Bopyrione toloensis Markham, 1982: 351–354, figs. 16–17 [Hong Kong, infesting *Alpheus microstylus?* (Bate, 1888)]; Markham, 1985a: 59; Morton, 2003: 37; Li, 2003: 140, 157; Liu, 2008: 691.

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Alpheus microstylus?* (Bate, 1888) (Alpheidae), Hong Kong.

REMARKS: The host was only provisionally identified as *A. microstylus* in Markham (1982).

KEY TO THE FIVE SPECIES OF *BOPYRIONE* BOURDON AND MARKHAM, 1980

- | | |
|---|---|
| 1a. Head triangular, pleopods large, flaplike | 2 |
| 1b. Head subovate, pleopods small, tubercular | 4 |

- | | |
|--|---|
| 2a. Anterior margin of head entire | 3 |
| 2b. Anterior margin of head crenulate | <i>B. toloensis</i> Markham, 1982 |
| 3a. Eyes present, brood pouch closed | <i>B. longicapitata</i> Markham, 1982 |
| 3b. Eyes lacking, brood pouch medially open | <i>B. multifeminae</i> , n. sp. |
| 4a. First four pereomeres with dorsolateral bosses | <i>B. woodmasoni</i> (Chopra, 1923) |
| 4b. All pereomeres without dorsolateral bosses | <i>B. synalphei</i> Bourdon and Markham, 1980 |

Bopyroides Stimpson, 1864

DIAGNOSIS: Female: Body subovate, moderately distorted. Head separate from pereon. Distinct frontal lamina, palp, coxal plates, dorsolateral bosses. Posterolateral point of first oostigite not directed backward. Pleopods uniramous or absent, uropods lacking.

Male: Body elongate, without midventral tubercles. Pleon completely fused, without pleopods or uropods.

TYPE SPECIES: *Bopyroides acutimarginatus* Stimpson, 1864 (= *Bopyrus hippolytes* Kröyer, 1838), by original designation.

OTHER SPECIES: *Bopyroides cluthae* (Scott, 1902), *B. shiinoi* Rybakov and Avdeev, 1991.

REMARKS: Stimpson (1864) erected this genus for *Bopyroides acutimarginatus* from Puget Sound. Bourdon (1968) reviewed the genus and considered *B. acutimarginatus* as a synonym of *Bopyrus hippolytes*, and also synonymized *B. furcata* Norman, 1905, and *Pleurocrypta patiencei* Scott, 1902, with *B. cluthae*. Markham (1985a) regarded *B. sarsi* Bonnier, 1900, as identical with *B. hippolytes*. Rybakov and Avdeev (1991) described *B. shiinoi* from the northwestern Pacific.

Bopyroides hippolytes (Kröyer, 1838)

Figure 11

Bopyrus hippolytes Kröyer, 1838a: 306–310, 318, pl. 4, fig. 22 [type locality Greenland; infesting *Lebbeus polaris*]; Kröyer, 1838b: 78–82, 90, pl. 4, fig. 22.

Bopyroides hippolytes—Markham, 1985a: 52–53 (extensive synonymy); Bourdon, 1987: 342; Kim and Kwon, 1988, 212–213, fig. 8 [Korea, infesting *Pandalus borealis*]; Rybakov, 1990: 409 [Sea of Okhotsk, infesting *P. borealis*, host for *Bourdonia tridentata* Rybakov, 1990 (Cabiropidae)]; Rybakov




Fig. 11. *Bopyroides hippolytes* (Kröyer, 1838), reference female (CIEHI401001) (A–H): **A**. Dorsal view. **B**. Ventral view. **C**. Left maxilliped, external view. **D**. Left side of barbula. **E**. Left oostegite 1, external view. **F**. Left oostegite 1, internal view. **G**. Left pereopod 1. **H**. Left pereopod 7. Reference male (CIEHI401002) (I–L): **I**. Dorsal view. **J**. Ventral view. **K**. Right antenna and antennule. **L**. Left pereopod 1. Scale: 1 mm (A, B); 0.5 mm (C, E–H); 0.25 mm (D, I–L).

and Avdeev, 1991: 167–168, fig. 1 [comparison with *B. shiinoi*]; Saito et al., 2000: 39–40 (list); Boyko, 2004: 692–693, fig. 10 [Taiwan, infesting *Lebbeus* cf. *spinirostris* (Kobyakova, 1936)]; An, 2006: 78–79, fig. 34.

MATERIAL EXAMINED: Infesting *Birulia kishinouyei*, 1 ♀ (CIEHI401001), 1 ♂ (CIEHI401002), Yellow Sea, Stn. 4010, 123°30'E, 33°30'N, 60 m, 6 December 1959, coll. Jie-shan Xu and Mu Chen.

DESCRIPTION: Female (CIEHI401001): Length 8.38 mm, maximal width 7.30 mm, head length 1.92 mm, head width 2.17 mm, pleon length 2.60 mm, distorted 19°. All body segments distinct (fig. 7A, B). Head oval, frontal lamina extending beyond head, anterior edge shallowly bilobate, posterior edge curved, eyes absent (fig. 7A). Maxilliped (fig. 7C) with short setose palp and obscure spur. Barbula (fig. 7D) with two pairs of long falcate lateral projections on each side, outer one more slender.

Pereon broadest across third pereomere (fig. 11A). Coxal plates rudimentary on both sides. Brood pouch completely open (fig. 11B). Oostegite 1 (fig. 11E, F) with two equally long articles, nearly smooth internal ridge bearing one large and two or three small projections, posterolateral point directed laterally. Pereopods larger posteriorly (fig. 11G, H), with blunt and small dactyli. Pleon of six pleomeres, lateral plates lacking. Four pairs of small tubercular uniramous pleopods, uropods lacking.

DESCRIPTION: Male (CIEHI401002): Length 2.95 mm, maximal width (across pereon 4) 0.83 mm, head width 0.54 mm, pleonal length 0.79 mm. All pereomeres distinctly segmented, pleon fused (fig. 11I, J). Head a flattened ellipse, posterior edge curved (fig. 11I), black eyes in posterolateral corners (fig. 11I). Antennule of two articles, antenna of four articles, terminally setose (fig. 11K). Pereomeres subequal in width, lacking midventral projections (fig. 11J). Pereopods of subequal size and structure (fig. 11L). Pleon completely fused, without pleopods or uropods, obscure segment demarcations visible in ventral view (fig. 11J).

HOSTS AND LOCALITIES: *Birulia kishinouyei* (Yokoya, 1930), *Eualus fabricii* (Kröyer, 1841), *E. gaimardi* (H. Milne Edwards,

1837), *E. pusiolus* (Kröyer, 1841), *E. suckleyi* (Stimpson, 1864), *Heptacarpus brevirostris* (Dana, 1852), *H. herdmani* (Walker, 1898), *Hippolyte varians* Leach, 1814, *Lebbeus groenlandicus* (Fabricius, 1775), *Lebbeus polaris* (Sabine, 1824), *L. cf. spinirostris* (Kobyakova, 1936), *Spirontocaris arcuata* Rathbun, 1902, *S. holmesi* Holthuis, 1947, *S. lamellicornis* (Dana, 1852), *S. lilljeborgii* (Danielssen, 1859), *S. murdochii* Rathbun, 1902, *S. phippsii* (Kröyer, 1841), *S. spinus* (Sowerby, 1805) (Hippolytidae) and *Pandalopsis aleutica* Rathbun, 1902, *P. dispar* Rathbun, 1902, *Pandalus borealis* Kröyer, 1838, *P. goniurus* Stimpson, 1860, *P. jordani* Rathbun, 1902, *P. montagui* Leach, 1814 (Pandalidae), from Greenland, Iceland, Ireland, England, Denmark, Norway, Arctic Ocean, White Sea, Barents Sea, Sea of Okhotsk, Bering Sea, Japan, Korea, Yellow Sea, Taiwan, Kodiak, Alaska to Puget Sound, Washington, Ellsmere Island, Baffin Bay, Gulf of St. Lawrence, Nova Scotia, Maine to Cape Cod Bay, Massachusetts, 60 m (Yellow Sea), 506–680 m (Taiwan) (most other depths not recorded).

REMARKS: *Bopyroides hippolytes* has been described many times in detail (see Markham, 1985a, for a more complete synonymy and discussion). The synonymy list provided here includes only select references published after Markham (1985a). All hosts are in Hippolytidae and Pandalidae. Boyko (2004) recorded this species from Chinese waters infesting *Lebbeus* cf. *spinirostris* (Kobyakova) off Taiwan. The present specimens conform well to previous descriptions of this species, but the host and locality are new records. *Birulia kishinouyei* has not previously been reported as hosting any bopyrid.

Bopyroides shiinoi Rybakov and Andeev, 1991
Figure 12

Bopyroides hippolytes: Shiino, 1937a: 293–296, fig. 1 [Japan, infesting *Spirontocaris alcimede* de Man, 1906 (= *Heptacarpus geniculatus* (Stimpson, 1860)), *S. mororani* Rathbun, 1902 (= *S. ochotensis* (Brandt, 1851)), *S. pandaloides* Stimpson (= *Heptacarpus pandaloides* (Stimpson, 1860))] (not *Bopyroides hippolytes* Kröyer, 1838).




Fig. 12. *Bopyroides shiinoi* Rybakov and Andeev, 1991, reference female (CIEHI560901) (A-G): **A.** Dorsal view. **B.** Ventral view. **C.** Left maxilliped, external view. **D.** Left side of barbula. **E.** Left oostegite 1, external view. **F.** Left oostegite 1, internal view. **G.** Left pereopod 1. Reference male (CIEHI560902) (H, I): **H.** Dorsal view. **I.** Ventral view. Scale: 1 mm (A, B); 0.25 mm (C, D, G); 0.5 mm (E, F, H, I); 0.25 mm (G).

Bopyroides shiinoi Rybakov and Andeev, 1991: 169–171 [Peter the Great Bay, Sea of Japan, infesting *Heptacarpus camtschaticus* (Stimpson, 1860), *H. geniculata longirostris* (Kobjakova) (= *H. longirostris* (Kobjakova, 1936)), and *H. grebnitzkii* Rathbun, 1902]; Boyko, 2004: 692–693 (mention, no new records); An, 2006: 79–80, fig. 35.

MATERIAL EXAMINED: Infesting *Hippolyte* sp. (Hippolytidae), 1 ♀ (CIEHI560901), 1 ♂ (CIEHI560902), Sea Island, Dalian City, Jiangsu Province, 123°15'E, 39°00'N, 27 September 1956.

DESCRIPTION: Female (CIEHI560901): Length 4.86 mm, maximal width 3.60 mm, head length 0.84 mm, head width 1.28 mm, sinistrally distorted 36° (fig. 12A, B).

Head ovate, anterior edge bilobate, frontal lamina present, eyes absent (fig. 12A). Maxilliped (fig. 12C) with setose, articulated palp, inner edge fringes with setae. Barbula (fig. 12D) with two pairs of sharp, elongate falcate lateral projections on each side.

Pereon broadest across third pereomere (fig. 12A). Narrow rudimentary coxal plates on first four pereomeres of both sides. Lateral margin of pereomeres 5–7 indented (fig. 12A). Brood pouch completely open (fig. 12B). Oostegite 1 (fig. 12E, F) with two equally long articles, internal ridge almost smooth, posterolateral point blunt and directed laterally; fifth oostegite longest. Pereopods with subequal size and structure (fig. 12G), blunt dactyli, bases with carinae. Pleon of five pleomeres, fused medially. Pleopods and uropods lacking.

DESCRIPTION: Male (CIEHI560902): Length 1.87 mm, maximal width (across pereon 5) 0.58 mm, head width 0.34 mm, pleonal length 0.51 mm. All pereon segments distinct, pleomeres fused (fig. 12H, I). Head a flattened ellipse (fig. 12H), eyes in posterolateral corners (fig. 12H). Antennule of three articles, antenna of five articles, terminally setose, extending beyond margins of head (fig. 12I). Pereomeres subequal in width, lacking midventral projections (fig. 12I). Pereopods with subequal size and structure (fig. 12I). Pleon completely fused, pleopods and uropods lacking (fig. 12I).

HOSTS AND LOCALITIES: Infesting *Heptacarpus geniculatus* (Stimpson, 1860), *Heptacarpus*

pandaloides (Stimpson, 1860), *Hippolyte* sp., *Spirontocaris ochotensis* (Brandt, 1851) (Hippolytidae), Japan and Jiangsu Province, China.

REMARKS: Shiino (1937a) recorded *Bopyroides hippolytes* infesting three *Spirontocaris* spp. from Japan, but this material was considered as belonging to *B. shiinoi* by Rybakov and Avdeev (1991), which was described from a more northerly locality in the Sea of Japan and from different hippolytid hosts. Rybakov and Avdeev (1991) noted that the most obvious difference between *B. hippolytes* and *B. shiinoi* is in the fusion of the terminal pleomeres. The present female does not have tubercular pleopods but does have fusion of the last two pleomeres, suggesting that it is conspecific with *B. shiinoi*. The host of the present specimens must be new, as no species of *Hippolyte* was previously known as a host for *B. shiinoi*, but this host specimen is not identifiable beyond the genus level.

KEY TO THE THREE SPECIES OF *BOPYROIDES* STIMPSON, 1864

- 1a. Pleon of five pleomeres (last two pleomeres fused) . . . *B. shiinoi* Rybakov and Avdeev, 1991
- 1b. Pleon of six pleomeres 2
- 2a. Sixth pleomere extended and produced into two divergent rami . . . *B. cluthae* (Scott, 1902)
- 2b. Sixth pleomere not extended, terminally smooth, round . . . *B. hippolytes* (Kröyer, 1838)

Capitetragonia Pearse, 1953

DIAGNOSIS (modified from Markham, 1985a): Female: All body segments distinct. Head extended and produced anterolaterally; barbula with two pairs of lateral projections, both longer than wide; maxilliped lacking palp. Five or seven oostegites; oostegite 1 approximately as wide as long, extended into posterolateral projection at least 1/5 of total length; coxal plates absent. Pleon of five or six pleomeres divided by deep notches along both sides; four pairs of flaplike biramous pleopods; no uropods.

Male: Body less than three times as long as broad, sides subparallel. Head, pereomeres, and pleon distinct. No midventral tubercles. Pleon fused, margins undulate, wider than adjacent pereomeres, much shorter than

wide; five pairs of uniramous, flap like pleopods; no uropods.

TYPE SPECIES: *Capitetragnonia asperotibialis* Pearse, 1953 (= *Bopyrus alphei* Richardson, 1900).

OTHER SPECIES: *Capitetragnonia elliptica* (Markham, 1992), n. comb.

REMARKS: Markham (1985a) erected *Probopyria* for the single species *Bopyrus alphei* Richardson, 1900. He examined the syntypes of *Capitetragnonia asperotibialis* Pearse, 1953 (USNM 93720, see Pearse, 1953), and concluded they were conspecific with *B. alphei*. One of us (CBB) has also examined these syntypes and we concur with Markham's (1985a) conclusion as to this synonymy. However, because the type species of *Capitetragnonia* and *Probopyria* are synonyms of each other, the correct name for the genus must be *Capitetragnonia*, while the correct name for the type species is *Capitetragnonia alphei* (Richardson, 1900).

Capitetragnonia elliptica (Markham, 1992),
n. comb.

Probopyria elliptica Markham, 1992a: 283–285, figs. 6–7 [Hong Kong, infesting *Alpheus malleodigitus* (Bate, 1888)]; Li, 2003: 140, 153, 158.

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Alpheus malleodigitus* (Bate, 1888) (Alpheidae), Hong Kong.

REMARKS: Markham (1992a) provisionally placed this species in *Probopyria* because the female has seven pairs of oostegites, known only in two other species of bopyrines (see Remarks under *Septembopyrina*, n. gen.), while the type species has the more usual five pairs. The male is much more similar, although that of *P. elliptica* has one fewer pleomere.

Discorsobopyrus Boyko, 2004

DIAGNOSIS (after Boyko, 2004): Female: body ovate, one side of pereon slightly longer than other; head triangular, weakly produced with narrow frontal lamina. Antennae and antennules reduced to single segment each. Maxilliped with stout, distally rounded spur;

palm lacking. Oostegite 1 proximal lobe ovate, distal lobe subtriangular, internal ridge smooth. Pereon composed of seven pereomeres, broadest across third pereomere. Coxal plates well developed on both sides, all elongate. Dorsolateral bosses well developed on some pereomeres, indistinct on others. Tergal projections lacking. Oostegites nearly completely enclosing marsupium. Basis of all pereopods bearing pronounced rounded medial boss; propodus with cup for insertion of dactylus. Pleon with five pleomeres plus pleotelson; first through fifth pleomeres with uniramous elongate pleopods and uniramous, short subquadrate lateral plates (some indistinct); edges and surfaces of all lateral plates smooth; pleopods smooth, uropods lacking.

Male: Head ovate, fused with first segment of pereon. Body shape elongate; fourth pereomere broadest; first three directed slightly anterolaterally, fourth through sixth laterally directed, seventh directed slightly posterolaterally. Posterior pereopods slightly larger. Pleon with one segment; faint lateral indication of segmentation and tapered pleotelson tip. Midventral tubercles on second to seventh pereomeres, pleopods and uropods lacking.

TYPE SPECIES: *Bopyrus stebbingi* Nierstrasz and Brender à Brandis, 1923, by original designation.

OTHER SPECIES: None.

Discorsobopyrus stebbingi (Nierstrasz and Brender à Brandis, 1923)

Bopyrus stebbingi Nierstrasz and Brender à Brandis, 1923: 97–98, pl. 7, fig. 23a–c [Indonesia, infesting unknown host]; Chopra, 1923: 518, 541–542; Bourdon, 1968: 372.

Discorsobopyrus stebbingi: Boyko, 2004: 694–697, figs. 11–13 [Taiwan, Indonesia, infesting *Heterocarpus sibogae* de Man, 1917]; Markham, 2010: 159.

? “unidentified bopyrid” Li and Chan, 2014: 135 [Philippines, infesting *H. sibogae*].

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Heterocarpus sibogae* de Man, 1917 (Pandalidae), Philippines, Indonesia, and Taiwan.

REMARKS: The unidentified bopyrid on a specimen of *H. sibogae* collected from the

Philippines and cited by Li and Chan (2014) may belong to this species or possibly to *Pseudione magna* Shiino, 1951 (see Markham, 2010); no other bopyrid is known from this host species.

Litobopyrus Markham, 1982

DIAGNOSIS (after Markham, 1982): Female: Body distorted; head distinct from pereon; frontal lamina obscure; maxilliped with tuft of setae, palp lacking; barbula of two moderately long projections on each side; oostegite 1 lacking internal ornamentation and postero-lateral point; lateral plates elongate; five pairs of biramous pleopods with endopodites reduced to sessile knobs; uropods broad and long.

Male: Greater than three times as long as broad; head medially fused with first pereomere; pereopods subequal, with all articles distinct; pleomeres fused, lacking appendages.

TYPE SPECIES: *Litobopyrus longicaudatus* Markham, 1982, by original designation.

OTHER SPECIES: None.

Litobopyrus longicaudatus Markham, 1982

Litobopyrus longicaudatus Markham, 1982: 346–348, fig. 15 [Hong Kong, infesting *Athanas* sp.]; Morton, 2003: 37; Li, 2003: 140, 153, 157.

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Athanas* sp. (Alpheidae), Hong Kong.

REMARKS: The taxonomic position of *L. longicatudatus* is unclear, as the female characters are more like those of the bopyrine *Urobopyrus* Richardson, 1904, while the male characters more resemble those of the pseudionine *Pleurocrypta* Hesse, 1865.

Parabopyrella Markham, 1982

DIAGNOSIS: Female: Head fused with first pereomere except at anterolateral corners; maxilliped with setose palp; barbula with one or two pairs of lateral projections on each side; oostegite 1 with unadorned or sparsely digitate internal ridge; dorsolateral bosses usually on first four pereomeres; pleon of six segments indicated by lateral indentations; pleotelson rounded or produced into posterolateral points;

five pairs of flaplike biramous pleopods; uropods absent.

Male: Head usually fused with first pereomere; eyes present or absent; pereopods subequal in size and shape; pleon variously fused; first pleomere as broad as seventh pereomere or much broader; pleopods tuberculate if present; uropods usually absent.

TYPE SPECIES: *Bopyrella mortenseni* Nierstrasz and Brender à Brandis, 1929, by original designation.

OTHER SPECIES: *P. angulosa* (Bourdon, 1980a), *P. angusta* (Shiino, 1936), *P. australiensis* (Bourdon, 1980a), *P. barnardi* (Nierstrasz and Brender à Brandis, 1931), *P. bonnieri* (Nierstrasz and Brender à Brandis, 1923), *P. choprai* (Nierstrasz and Brender à Brandis, 1929), *P. crenulata* (Shiino, 1939), *P. cuspidata*, n. sp., *P. delagoae* (Bourdon, 1982), *P. distincta* (Nierstrasz and Brender à Brandis, 1923), *P. elongata* (Shiino, 1949), *P. essingtoni* (Bourdon and Bruce 1983), *P. hoggarti* (Chopra, 1923), *P. incisa* (Chopra, 1923), *P. indica* (Chopra, 1923), *P. intermedia* (Nierstrasz and Brender à Brandis, 1923), *P. lata* (Nierstrasz and Brender à Brandis, 1929), *P. megatelson* (Nierstrasz and Brender à Brandis, 1929), *P. nierstraszi* (Chopra, 1930), *P. pacifica* (Shiino, 1933), *P. perplexa* Markham, 1990, *P. richardsonae* (Nierstrasz and Brender à Brandis, 1929), *P. saronae* (Bourdon and Bruce, 1979), *P. setoensis* (Shiino, 1939), *P. symmetros*, n. sp., *P. tanyensis* (Bourdon, 1979), and *P. thomasi* (Nierstrasz and Brender à Brandis, 1929).

REMARKS: Although Markham (1985a) purported to erect this genus and restricted the definition of *Bopyrella*, transferring four western Atlantic species and 20 Indo-West Pacific species to *Parabopyrella*, in fact the genus name is available from Markham (1982) where it was cited as “in press” but with a type species, *Bopyrella mortenseni*, designated (the same species was “designated” as the type species in Markham, 1985a). Only one other taxon, *Bopyrella deformans indica* Chopra, 1923, was listed as included in the genus when it was introduced by Markham (1982). Markham (1985a) erroneously referred to the 1982 usage of the name as a nomen nudum.

Parabopyrella can be distinguished from *Bopyrella* in that females of *Parabopyrella*

have lateral indentations that indicate the pleomeres on both sides while the pleomeres of females in *Bopyrella* are completely fused. Previously, this genus included 26 species, all known from hosts in Alpheidae and Hippolytidae. Two new species are described herein.

According to the shape of the distal margin of the pleotelson of the female, species of *Parabopyrella* can be divided into three groups (after Bourdon, 1980a). “A” group has the pleotelson tip entire, truncate, or convex and produced into a point in the median and includes six species (*P. angulosa*, *P. chopræ*, *P. crenulata*, *P. delagoae*, *P. perplexa*, *P. symmetros*, n. sp.). “B” group has the pleotelson tip entire, but convex and produced into a rounded distal region and includes eight species (*P. angusta*, *P. essingtoni*, *P. mortensenii*, *P. nierstraszi*, *P. richardsonae*, *P. saronae*, *P. setoensis*, *P. tanyensis*). “C” group has the pleotelson incised medially, with divergent posterolateral lobes. Within the “C” group, there are four types: “C1” group with the medial incision shallow and the distal ends of the lobes quadrate, and includes five species (*P. australiensis*, *P. barnardi*, *P. cuspidata*, n. sp., *P. elongata*, *P. pacifica*), “C2” group with the medial incision shallow, but distal ends of the lobes rounded, and includes three species (*P. distincta*, *P. intermedia*, *P. thomasi*), “C3” group with the medial incision deep and the distal ends of lobes pointed and long, and includes two species (*P. lata*, *P. megatelson*), “C4” group with the medial incision shallow but produced into two small points, and includes four species (*P. bonnieri*, *P. hodgarti*, *P. incisa*, *P. indica*).

Parabopyrella chopræ (Nierstrasz and Brender à Brandis, 1929)

Bopyrella chopræ Nierstrasz and Brender à Brandis, 1929: 29–30, figs. 32–33 [Nicobar Islands, infesting *Alpheus lottini* Guérin-Méneville, 1838]; Taiwan, infesting *Lysmata vittata* (Stimpson, 1860) or *Synalpheus* sp. (see below)]; Chopra, 1930: 134, 138; Shiino, 1933: 282–283.

Bopyrella chopræ: Bourdon, 1980a: 187, 213–215, fig. 13.

Parabopyrella chopræ: Kensley and Chan, 2001: 481; Boyko, 2004: 677; Boyko, 2006: 43; Liu, 2008: 692.

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Alpheus lottini* Guérin-Méneville, 1838) (Alpheidae) and *Lysmata vittata* (Stimpson, 1860) (Hippolytidae) or *Synalpheus* sp. (Alpheidae), Nicobar Islands and Taiwan.

REMARKS: As pointed out by Boyko (2006), there is some doubt as to the host in Taiwan as given by Nierstrasz and Brender à Brandis (1929), but both hippolytid and alpheid shrimp are known as hosts of *Parabopyrella* species, so neither can be ruled out (although the fact that the host from the Nicobar Islands was an alpheid suggests that the *Synalpheus* sp. is more likely). Bourdon (1980a) changed the spelling of the specific name without comment, presumably because it is generally agreed that Chopra was a man. Although it is possible that Nierstrasz and Brender à Brandis (1929) treated the name as if it were a first-declension Latin noun, which are usually but not always feminine, and used the appropriate genitive, this seems unlikely. On the second page of their paper, they remarked “Die späteren Forscher, wie Miss Richardson, Chopra, und auch wir selbst, haben das Prinzip Bonnier's übernommen aus Armut, weil es kein besseres gibt.” The lack of an honorific for Chopra, as was provided for Richardson, suggests that Nierstrasz and Brender à Brandis thought Chopra was a male but does not explain why the –ae ending was used.

Parabopyrella cuspidata, n. sp.
Figure 13

Parabopyrella cuspidatus An, 2006: 85–86, fig. 40 (unavailable name).

MATERIAL EXAMINED: Infesting *Alpheus digitatus* De Haan, 1844, holotype ♀ (CIEAL 560301), allotype ♂ (CIEAL 560302), Haimen, Jiangsu Province, 121°15'E, 31°54'N, 12–14 March 1956, coll. Ruiyu Liu and Zhen-gang Fan.

DESCRIPTION: Holotype female (CIEAL 560301): Length 9.24 mm, maximal width 5.53 mm, head length 2.06 mm, head width 2.1 mm, distorted 22° (fig. 13A, B).

Head fused with first pereomere except in anterolateral corners, obscure posterior




Fig. 13. *Parabopyrella cuspidata*, n. sp., holotype female (CIEAL560301) (A–I): A. Dorsal view. B. Ventral view. C. Left maxilliped, external view. D. Palp of maxilliped, internal view. E. Left side of barbula. F. Left oostegite 1, external view. G. Left oostegite 1, internal view. H. Right pereopod 2. I. Right pereopod 6. Allotype male (CIEAL560302) (J–M): J. Dorsal view. K. Ventral view. L. Right antennae and antennule. M. Left pereopod 7. Scale: 2 mm (A, B); 0.5 mm (C, E–G); 0.25 mm (D, H–K); 0.15 mm (L, M).

margin sharply curved and anterior margin of head irregular. Small black eyes near posterior margin of head (fig. 13A). Antennule of three articles, antenna of five articles. Maxilliped articulated with triangular palp laterally fringed with setae (fig. 13C, D). Barbula with two pairs of short falcate lateral projections on each side (fig. 13E).

Pereon broadest across third pereomere (fig. 13A). Pereomeres 2–5 medially fused. Coxal plates well developed on left side of all pereomeres and on right side of first four pereomeres. Round dorsolateral bosses on first four pereomeres. Brood pouch widely open (fig. 13B). Oostegite 1 (fig. 13F, G) with two articles, internal ridge bearing small simple projections, posterolateral point directed straight backward, distal portion of posterior edge fringed with setae. Posterior pereopods slightly larger (fig. 13H, I), carpi and meri smooth, dactyli short, bases of all pereopods produced into lobes. Pleon of six pleomeres, segments indicated by dorsal sutures and deep lateral indentations (fig. 13A), first five bearing biramous flaplike pleopods (fig. 13B), sixth pleomere incised medially, uropods lacking (fig. 13B).

DESCRIPTION: Allotype male (CIEAL560302): Length 2.20 mm, maximal width across pereomere 4 0.82 mm, head width 0.47 mm, head length 0.28 mm. All segments distinct (fig. 13J, K). Head elliptical with curved posterior edge (fig. 13J), large black eyes near posterior margin of head (fig. 13J). Antennule of three articles with setae on distal article, antenna of one article, terminally smooth (fig. 13L). Pereomeres widest across fourth pereomere, lacking midventral projections (fig. 13K). All pereopods of similar size and structure, dactyli sharp but short (fig. 13M). Pleon of six distinct segments, without midventral projections, pleopods or uropods (fig. 13K). Sixth pleomere produced into two extend subacute lobes with medioventral anal cone (fig. 13K).

ETYMOLOGY: The specific name, *cuspidata*, refers to the head of the female having a sharply curved posterior margin and an irregular anterior margin.

HOST AND LOCALITY: Infesting *Alpheus digitalis* De Haan, 1844 (Alpheidae), Jiangsu Province, China.

REMARKS: The new species belongs to the “C1” group with a medially incised pleotelson and can be distinguished from the other four species of the “C1” group by characters of oostegite 1, maxilliped, and the posterior margin of the head. The male of the new species differs from those of all other *Parabopyrella* species by the characters given below in (4). *Parabopyrella cuspidata*, n. sp., is most closely related to *P. barnardi*, but (1) the female of the new species bears a pleon with dorsal sutures and deep indentations on both sides (the female of *P. barnardi* has the pleon fused dorsally and with only the left side deeply indented), (2) the maxilliped of the new species has a triangular palp (*P. barnardi* with cylindrical palp), (3) the posterolateral point of oostegite 1 of the new species is sharp and backwardly directed (that of *P. barnardi* round and laterally directed), (4) the male of the new species has distinct pleomeres and laterally separated lobes on the pleotelson (male of *P. barnardi* with fused pleomeres and rounded pleotelson).

Parabopyrella distincta (Nierstrasz and

Brender à Brandis, 1923)

Figure 14

Bopyrella distincta Nierstrasz and Brender à Brandis, 1923: 97, pl. 6, fig. 22 a–d [type locality: Pulau Kaniungan-Ketjil, Indonesia; infesting *Synalpheus amboinæ* (Zehntner, 1984) = *Synalpheus stimpsonii* (de Man, 1888); Chopra, 1927: 121–122, figs. 1–2 [Tamban, Tamil Nadu, India; infesting unidentified alpheid]; Shiino, 1936: 159; Shiino, 1939: 93; Shiino, 1949a: 49; Bourdon, 1980a: 205–208, fig. 9 [redescription of holotype].

Parabopyrella distincta: Markham, 1985a: 67 [transfer to *Parabopyrella*]. —An, 2006: 82–83, fig. 37.

MATERIAL EXAMINED: Infesting *Alpheus* sp., 1 ♀ (CIEAL550001), 1 ♂ (CIEAL550002), 1955, Hainan Province, Qionghai Province.

DESCRIPTION: Female (CIEAL550001): Length 7.52 mm, maximal width 5.53 mm,




Fig. 14. *Parabopyrella distincta* (Nierstrasz and Brender à Brandis, 1923), reference female (CIEAL550001) (A–G): **A.** Dorsal view. **B.** Ventral view. **C.** Left maxilliped, external view. **D.** Left side of barbula. **E.** Left oostegite 1, external view. **F.** Left oostegite 1, internal view. **G.** Right pereopod 7. Reference male (CIEAL550002) (H–J): **H.** Dorsal view. **I.** Ventral view. **J.** Right antennae and antennule. Scale: 2 mm (A, B); 1 mm (E, F); 0.5 mm (C, D, H, I, G); 0.25 mm (J).

head width 2.19 mm, distorted 31° (fig. 14A, B).

Head fused with first pereomere, with anterolateral corner notch. Frontal lamina and eyes lacking (fig. 14A). Maxilliped (fig. 14C) with unarticulated setose palp. Barbula (fig. 14D) with two pairs of smooth falcate lateral projections on each side and shallow median indentation in center.

Pereon broadest across third pereomere (fig. 14A). Coxal plates rudimentary on short side, but well developed on first three pereomeres on long side. First three pereomeres of left side and only first pereomere of right side with dorsolateral bosses. Brood pouch completely open (fig. 14B). Oostegite 1 (fig. 14E, F) with two equally long articles, internal ridge bearing simple small projections, posterolateral point sharply directed laterally. Fifth oostegite much longer than other oostegites (fig. 14B). Pereopods of similar size, basis of each pereopod with carina (fig. 14G). Pleon of six pleomeres, bearing five pairs of biramous flaplike pleopods. Pleotelson medially incised, but rami distally round, uropods lacking.

DESCRIPTION: Male (CIEAL550002): Length 2.11 mm, maximal width (across pleon 1) 0.82 mm, head width 0.41 mm, pleonal length 0.26 mm. All pereomeres distinct (fig. 14H, I). Head ovate, anterior margin rounded, posterior margin curved (fig. 14H), black eyes near posterior margin (fig. 14H). Antenna of one article, antennule of three articles, both terminally setose (fig. 14J). Pereomeres subequal in width, lacking midventral projections (fig. 14I). Pereopods subequal in size, but dactyli of first five pairs larger than others, carpi of first pereopod rounded, others of elongate columnar form (fig. 14I). First three pleomeres distinct, with midventral tubercles. Last two pleomeres fused into single piece, but segments indicated by lateral indentations, without uropods (fig. 14H, I).

HOSTS AND LOCALITIES: Infesting *Alpheus* sp. (possibly more than one species), *Synalpheus stimpsonii* (de Man, 1888) (Alpheidae), Hainan Province, Qionghai Province, China and Pulau, Indonesia, 11 m.

REMARKS: Nierstrasz and Brender à Brandis (1923) described *Bopyrella distincta* based on a single female collected by the Siboga-Expedition. Chopra (1927) reported on a pair of

specimens from southeast India and described the male for the first time. Bourdon (1980a) redescribed and figured the female holotype specimen, incorrectly citing the host as *Synalpheus ambonicae* (sic). Markham (1985a) reviewed *Bopyrella*, and transferred this species to *Parabopyrella*. The present specimens are therefore only the third female and second male to be reported. The female conforms well to the holotype, but differ in some minor characters, such as the pleon of the present female not fused medially, and the posterolateral point of oostegite 1 more elongate and directed laterally.

Parabopyrella elongata (Shiino, 1949)

Figure 15

Bopyrella elongata Shiino, 1949a: 45–50, fig. 1 a–c. [Japan; infesting “*Crangon bis-incisus* (De Haan)” (= *Alpheus bisincisus* De Haan, 1849]. —Bourdon, 1980a: 194–196, fig. 4, tab. 1 [Queensland, Australia; infesting *Alpheus* sp.]

Parabopyrella elongata: Markham, 1985a: 67 [transfer to *Parabopyrella*]. —An, 2006: 83–84, fig. 38.

MATERIAL EXAMINED: Infesting *Alpheus* sp., 1 ♀ (CIEAL851001a), 1 ♂ (CIEAL851001b), Xiamen City, 118°15'E, 24°25'N, 15 October 1985.

DESCRIPTION: Female (CIEAL851001a): Length 7.03 mm, maximal width 4.49 mm, head length 1.46 mm, head width 1.63 mm, pereon length 2.84 mm, distorted 21° (fig. 15A, B). Head subquadrate, fused with first pereomere except in anterolateral corners, with thin frontal lamina. Black eyes near anterolateral corners (fig. 15A). Antennule of two articles, antenna of three articles, distally setose. Maxilliped (fig. 15C) with short stout setose palp. Barbula (fig. 15D) with two pairs of short smooth lateral projections on each side, small round tubercle near median.

Pereon segments distinct, broadest across third pereomere (fig. 15A). Coxal plates and dorsolateral bosses on first four pereomeres. Brood pouch widely open (fig. 15B). Oostegite 1 (fig. 15E, F) with nearly smooth internal ridge, posterolateral point sharp, directed posteriorly. Pereopods larger posteriorly (fig. 15G), bases of all pereopods with small




Fig. 15. *Parabopyrella elongata* (Shiino, 1949), reference female (CIEAL851001a) (A–G): **A.** Dorsal view. **B.** Ventral view. **C.** Right maxilliped, external view. **D.** Right side of barbula. **E.** Right oostegite 1, external view. **F.** Right oostegite 1, internal view. **G.** Right pereopod 7. Reference male (CIEAL851001b) (H, I): **H.** Dorsal view. **I.** Ventral view. Scale: 2 mm (A, B); 1 mm (E, F, H, I); 0.5 mm (C, D); 0.35 mm (G).

carina. Pleon of six pleomeres, segments indistinct but indicated by dorsal sutures and lateral deep indentations. First five pleomeres with flaplike biramous pleopods and weakly developed lateral plates; uropods absent.

DESCRIPTION: Male (CIEAL851001b): Length 1.63 mm, maximal width (across pereomere 7) 0.65 mm, pleonal length 0.41 mm (fig. 15H, I). Head ovate, completely fused with first pereomere (fig. 15I), without eyes (fig. 15H). Antenna obscure. Pereomeres 2–7 distinct, with similar pereopods (fig. 15I). Pereomeres subequal in width, lacking midventral projections (fig. 15I). Pleon fused into single piece, without lateral indentations, ventral suture visible and indicative of pleomeres, five tubercular pleopods (fig. 15I), uropods lacking.

HOSTS AND LOCALITY: Infesting *Alpheus bisincisus* De Haan, 1849, *Alpheus* sp. (probably more than one species) (Alpheidae), Japan, Queensland, Australia, and Fujian Province, China.

REMARKS: Shiino (1949a) described *Bopyrella elongata* infesting *Alpheus bisincisus* from Seto, Japan, and Bourdon (1980a) reported it infesting *Alpheus* sp. at Queensland; Markham (1985a) transferred it to *Parabopyrella*. The present specimens conform well to the redescription of Bourdon (1980a), but the male differs from both previously reported males of *P. elongata*. Shiino's (1949a) and Bourdon's (1980a) males have fused pleon segments but with the pleomeres indicated by lateral indentations. The present male has completely smooth lateral margins and has the pleomeres indicated by ventral sutures. Although the male differs in this respect from the other specimens, the female is very similar to the previously reported females and this material is therefore identified as *P. elongata*.

Parabopyrella hodgarti (Chopra, 1923)
Figure 16

Bopyrella hodgarti Chopra, 1923: 416, 469, 473–475, fig. 10; pl. 14, figs. 7–12 [type locality Vizagapatam, Madras Presidency (Bay of Bengal) (= Visakhapatnam, Andhra Pradesh, India); infesting *Alpheus crassimanus* Heller, 1865 (= *A. lobidens* De Haan, 1849)]. —Shiino, 1949a: 47 (mention). —Barnard, 1955: 79, fig. 37b, c (in part; Inhambane, Mozambique; infesting

A. crassimanus (= *A. lobidens*). —Bourdon, 1968: 407–408 (mention). —Kensley, 1974: 261 (off Natal, South Africa; infesting *Alpheus* sp.). —Bourdon, 1980a: 190–191 (mention).

Parabopyrella barnardi australiensis: An, 2006: 81–82, fig. 36 (not *Parabopyrella australiensis* (Bourdon, 1980a)).
not *Bopyrella hodgarti*: Barnard, 1955: 79 (in part; Mozambique; infesting *Hippolyte* sp.).

MATERIAL EXAMINED: Infesting *Alpheus japonicus* Miers, 1879, 1 ♀ (CIEAL570601a), 1 ♂ (CIEAL570601b), Jiaozhou Bay, Qingdao, Shandong Province, 120°15'E, 36°15'N, 5.5 m, 18 June 1957; 1 ♀, 1 ♂ (CIEAL570602), Jiaozhou Bay, Qingdao, Shandong Province, 120°15'E, 36°15'N, 5.5 m, 18 June 1957.

Infesting *Synalpheus streptodactylus* Couxière, 1905, 1 ♀ (CIEAL760801) East China Sea, Stn. V-3, 123°00'E, 26°30'N, 105 m, 27 August 1976, coll. Zhican Tang and Zhongyan Qi.

DESCRIPTION: Female (CIEAL570601a): Length 6.86 mm, maximal width 4.69 mm, head length 1.17 mm, head width 1.49 mm (fig. 16A, B). Head an inverted trapezoid, without frontal lamina; eyes absent (fig. 16A). Antennule and antenna absent. Maxilliped (fig. 16C) with short straight and stout palp, long setae terminally. Barbula (fig. 16D) with two pairs of smooth falcate lateral projections on each side and shallow indentation near center.

Pereon broadest across third pereomere (fig. 16A). Boundary between head and first pereomere only visible laterally. Coxal plates small and only on first four pereomeres of both sides, those on left side slightly larger than those on right side. Brood pouch widely open, oostegite 1 and median part of barbula visible in ventral view (fig. 16B); oostegite 5 longest. Internal ridge of oostegite 1 (fig. 16E, F) bearing six to eight small digitate projections, posterolateral point rounded posteriorly. Pereopods larger posteriorly (fig. 16G). Pleon of six pleomeres, but sutures obscure, wide lateral indentations indicating pleomeres. Five pairs of triangular flaplike biramous pleopods, surfaces smooth, uropods lacking.

DESCRIPTION: Male (CIEAL570601b): Length 1.64 mm, maximal width (across




Fig. 16. *Parabopyrella hodgarti* (Chopra, 1923), reference female (CIEAL570601a) (A–G): **A**. Dorsal view. **B**. Ventral view. **C**. Left maxilliped, external view. **D**. Left side of barbula. **E**. Left oostegite 1, external view. **F**. Left oostegite 1, internal view. **G**. Left pereopod 1. Reference male (CIEAL570601b) (H, I): **H**. Dorsal view. **I**. Ventral view. Scale: 2 mm (A, B); 0.5 mm (C); 1 mm (D–F); 0.35 mm (G); 0.45 mm (H, I).

pleomere 1) 0.54 mm, head width 0.27 mm, head and pleomere length 1.25 mm. (fig. 16H, I). Head ovate, posterior edge curved, but boundary between head and first pereomere obscure (fig. 16H), eyes present mediolaterally (fig. 16H). Antennule of three articles, antenna of five articles, terminally smooth (fig. 16I). Pereomeres subequal in width, lacking midventral projections (fig. 16I). Pereopods smaller posteriorly, first three pereopods with largest dactyli, last two pereopods with smallest dactyli (fig. 16I). Pleon fused, segments indicated by strong lateral indentations, first pleomere widest with pair of large round tubercular pleopods, all other pleopods and uropods absent (fig. 16I).

HOSTS AND LOCALITIES: Infesting *Alpheus japonicus* Miers, 1879, *A. lobidens* De Haan, 1849, *A.* sp. indet., *Synalpheus streptodactylus* Coutière, 1905 (Alpheidae), Andhra Pradesh, India, Mozambique, South Africa, Shandong Province, China, East China Sea, 5.5–105 m.

REMARKS: This species was previously known from the western Pacific in the Bay of Bengal, Mozambique, and South Africa (Chopra, 1923; Barnard, 1955; Kensley, 1974). Chopra (1923) erroneously stated that a “single specimen” was collected but described both the male and female; both are syntypes because Chopra failed to designate a holotype.

This is first record from China, and both hosts are first recorded as bearing parasitic isopods. The present specimens conform well to the female syntype of Chopra (1923), but the male differs from the male syntype in that (1) the present males have only the first pleomere distinct and other pleomeres indicated by lateral indentations; but the syntype male has the first three pleomeres distinct with the others fused without any lateral indentations (the Mozambique male of Barnard (1955) is somewhat intermediate in this character with weak indications of segmentation on the first four pleomeres); (2) the present male has one pair of pleopods, but the syntype male lacks all pleopods (character state in Mozambique male unknown); (3) the present male has only the first pleomere wider than the pereon, but the syntype male has the first three pleomeres wider than the pereon (the Mozambique male is somewhat intermediate with pleomeres 2 and 3 wider than

in the present specimen but not as wide as in the male syntype).

Based on the host and the morphological differences indicated above, it is very likely that the specimens reported by Barnard (1955) from *Hippolyte* sp. are distinct from *P. hodgarti*, but, as he did not describe or illustrate them, their identity is unknown.

Parabopyrella indica (Chopra, 1923)

“a subspecies of Hay’s *B. deformans*” Chopra, 1922: 70.

Bopyrella deformans var. *indica* Chopra, 1923: 416, 420

Bopyrella deformans indica Chopra, 1923: 468–473, text fig. 9, pl. 14, fig. 1–6 [Pakistan and India, infesting *Synapheus tumidomanus tumidomanus* (Paul’son, 1875); NE of Sri Lanka, infesting *Synapheus nilandensis* Coutière, 1905]; Chopra, 1927: 119–121 [Gulf of Manaar, infesting alpheid sp.]; Chopra, 1930: 133; Monod, 1933: 236; Qazi, 1959: 55, 60; Savant, 1973: 331; Kruczynski and Menzies, 1977: 551, 556; Kensley, 2001: 222.

Synsynella deformans var. *indica*: Nierstrasz and Brender à Brandis, 1929: 4, 38, fig. 48 [Hong Kong, infesting *Synalpheus* sp.]; Monod, 1933: 227–232, figs. 50–51 [Red Sea, infesting alpheid sp.]; Shiino, 1949a: 49; Kruczynski and Menzies, 1977: 552.

“Chopra’s ‘sub-species’” Kruczynski and Menzies, 1977: 556.

Bopyrella indica: Bourdon, 1979a: 501–503, fig. 21 [Madagascar, infesting *Synalpheus* sp.]; Bourdon, 1980a: 187, 208–210, fig. 10 [Hong Kong, infesting *Synalpheus* sp.; Indonesia, infesting *Synalpheus* sp.; New Caledonia, infesting *Synapheus neomeris* (de Man, 1897); Queensland, infesting *Synalpheus* sp.]; Bourdon, 1980b: 243; Kazmi and Bourdon, 1997: 59.

Bopyrella distincta [sic]: Bourdon, 1980a: 210 (not *Bopyrella distincta* Nierstrasz and Brender à Brandis, 1923).

Parabopyrella indica: Markham, 1982: 345; Markham, 1992a: 298; Kensley, 2001: 224; Kazmi et al., 2002: 61, fig. 18 [Arabian Sea, infesting *Synapheus tumidomanus tumidomanus* (Paul’son, 1875)]; Poore et al., 2002: 119; Li, 2003: 140, 158; Liu, 2008: 693; Markham, 2010: 152, 162.

[*Parabopyrella*] *indica*: Kazmi and Bourdon, 1997: 59.

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Synapheus neomeris* (de Man, 1897), *Synapheus nilandensis* Coutière, 1905, *Synapheus tumidomanus tumidomanus* (Paul'son, 1875), *Synalpheus* sp(p). (Alpheidae), Pakistan, India, NE of Sri Lanka, Gulf of Manaar, Hong Kong, Red Sea, Arabian Sea, Madagascar, New Caledonia.

REMARKS: Chopra (1923) variously referred to this taxon as a new subspecies or variety of *Bopyrella deformans* (Hay, 1917) (the type species of *Synsynella* Hay, 1917), but the name has always been used at the subspecific level.

Parabopyrella perplexa Markham, 1990
Figure 17

Parabopyrella perplexa Markham, 1990: 556–559, fig. 2 [Hong Kong, China; infesting “unidentified (caridean?) shrimp”]. —Morton, 2003: 43 (list). —Li, 2003: 140 (list). *Ovobopyrus dentatus* An, 2006: 89–90, fig. 42 (unavailable name).

MATERIAL EXAMINED: Infesting *Alpheus digitalis* De Haan, 1844, 1 ♀ (CIEAL603301), 2 ♂♂ (CIEAL603302), South China Sea, Stn. 6033, 115°00'E, 22°30'N, 11 December 1959, coll. Jinzhao Zhu. 1 ♀, 1 ♂ (CIEAL920301), Sanya, Hainan Province, 109°30'E, 18°20'N, 23 March 1992.

DESCRIPTION: Female (CIEAL603301): Length 5.93 mm, maximal width 4.19 mm, head length 1.11 mm, head width 1.54 mm, distorted 26° (fig. 17A, B).

Head an inverted trapezoid, without frontal lamina, fused with first pereomere on posterior margin; eyes absent (fig. 17A). Antennule of two articles, antenna of four articles, both nonsetose (fig. 17C). Maxilliped triangular with straight palp, fringed with setae on inner margin (fig. 17D, E). Barbula with two pairs of falcate lateral projections on each side and one pair of rounded projections near median (fig. 17F).

Pereomeres distinct; pereon broadest across third pereomere (fig. 17A). Coxal plates and

dorsolateral bosses on first three pereomeres on both sides, but larger on longer side. Brood pouch completely open (fig. 17B), oostegite 1 visible in ventral view, second to fourth oostegite small, fifth oostegite longer than others. Oostegite 1 (fig. 17G, H) with two articles nearly equal in length, internal ridge smooth, posterolateral point directed backward, posterior margin setose. Bases of first four pereopods smaller than those of other three pereopods (fig. 17B). Pleon of six pleomeres, first two pleomeres indicated by dorsal sutures and lateral notches, other pleomeres only distinguished by deep lateral notches. Male inserted in median of five pairs of flaplike biramous pleopods (fig. 17B); uropods absent (fig. 17B).

DESCRIPTION: Male (CIEAL603302): Length 1.77 mm, maximal width (across pereomere 3) 0.65 mm, head width 0.47 mm, head length 0.16 mm, pleonal length 0.42 mm (fig. 17I, J). Head elliptical, fusion with first pereomere indicated by lateral notch between head and first pereomere (fig. 17I); eyes absent (fig. 17I). Antennule of three articles, antenna lacking, nonsetose (fig. 17J). Pereomeres subequal in width, without midventral projections (fig. 17J). Dactyli of first three pereopods larger than those of last four pereopods (fig. 17J). Pleon fused and ovate, segments indicated by lateral indentations. Three midventral projections on median of pleon, pleopods and uropods lacking, two clusters of setae on terminal margin of pleon (fig. 17J).

HOSTS AND LOCALITIES: Infesting *Alpheus digitalis* De Haan, 1844 (Alpheidae), and unidentified (caridean?) shrimp, Hong Kong and South China Sea.

REMARKS: Markham (1990) described this species from Hong Kong, but with the host and male both unknown, so the present records are the first with identified hosts and with males. The present females conform well to the holotype female, with only some minor differences: the lateral plates and dorsolateral bosses of the present specimens are on the first three pereomeres but on the first four pereomeres of the holotype; the posterior margin of oostegite 1 of the present specimens is fringed with setae, but that of the holotype is smooth.




Fig. 17. *Parabopyrella perplexa* Markham, 1990, reference female (CIEAL603301) (A–H): A. Dorsal view. B. Ventral view. C. Left antennae and antennule. D. Right maxilliped, external view. E. Palp of right maxilliped. F. Right side of barbula. G. Right oostegite 1, external view. H. Right oostegite 1, internal view. Reference male (CIEAL603302) (I, J): I. Dorsal view. J. Ventral view. Scale: 2 mm (A, B); 0.5 mm (C, I, J); 1 mm (D, F–H); 0.55 mm (E).

Parabopyrella symmetros, n. sp.

Figure 18

Parabopyrella mortenseni: An, 2006: 84–85, fig. 39 (not *Parabopyrella mortenseni* Nierstrasz and Brender à Brandis, 1929)).

MATERIAL EXAMINED: Infesting *Alpheus microstylus* (Bate, 1888), holotype ♀ (CIEAL 800505), allotype ♂ (CIEAL800506), Xisha, Shanhua Islands, 111°40'E, 16°45'N, 9–12 May 1980. Paratypes: 3 ♀♀ (CIEAL800507), 3 ♂♂ (CIEAL800508), same data as for holotype.

DESCRIPTION: Holotype female (CIEAL 800505): Length 5.92 mm, maximal width 3.81 mm, head length 1.48 mm, head width 1.55 mm, almost symmetrical (fig. 18A, B). Head an inverted trapezoid, lateral margins distinct, posterior margin fused with first per-ecomore; black eyes near anterolateral corners (fig. 18A). Antennule of three articles, terminally setose, visible in dorsal view, basal article stout. Antenna of two articles, terminally setose. Maxilliped (fig. 18C) with stout palp, terminally truncate, setose. Barbula (fig. 18D) with two pairs of falcate lateral projections on each side, medially flattened.

Pereon broadest across third pereomere (fig. 18A). Coxal plates and dorsolateral bosses on first four pereomeres. Brood pouch completely open (fig. 18B), oostegite 1 and barbula visible in ventral view. Oostegite 1 (fig. 18E, F) with two articles, anterior margin of first article concave, internal ridge bearing two or three small projections, posterolateral point directed laterally. Pereopods of similar size (fig. 18G), with slender bases and small dactyli. Pleon of six pleomeres, first pleomere distinct, second to fourth pleomeres on right side distinct, but those on left side obscure and only indicated by lateral indentations. Fifth and sixth pleomeres fused and indicated by lateral notches. Pleotelson fan shaped, terminal margin entire. First five pleomeres with flaplike biramous pleopods; uropods absent.

DESCRIPTION: Allotype male (CIEAL800 506): Length 1.95 mm, maximal width across pereomere 3 0.70 mm, head width 0.38 mm, head length 0.17 mm, pleonal length 0.39 mm (fig. 18H, I). Head elliptical, fused with first pereomere, but with lateral notch between head and pereon (fig. 18H); black

eyes mediolaterally (fig. 18H). Antennule of two articles, antenna of three articles, lacking terminal setae (fig. 18I). Pereomeres subequal in width, lacking midventral projections (fig. 18I). Pereopods of similar size, but first two with slightly larger dactyli, last four pereopods with longer carpi than first three (fig. 18I). Pleon with first three segments distinct, last three segments fused but indicated by weak lateral indentations. First three pleomeres with midventral tubercles. Pleopods and uropods lacking (fig. 18I).

ETYMOLOGY: The specific name *symmetros* is feminine and refers to the female having an almost symmetrical body shape.

HOSTS AND LOCALITY: Infesting *Alpheus microstylius* (Bate, 1888) (Alpheidae), Xisha (Paracel Islands), China.

REMARKS: *Parabopyrella symmetros*, n. sp., belongs to the "A" group with the female pleotelson posterior margin entire, not incised or indented. The new species can be distinguished from the other five species of "A" group by the symmetrical female having the first, second, and third pleomeres on the right side distinct, and the male with the first three pleomeres distinct. This species is most closely related to *P. perplexa*, but *P. perplexa* females have an asymmetrical body, lack eyes, have the posterolateral point of oostegite 1 acute, and the palp of the maxilliped triangular. *Alpheus microstylus* is also known to be infested by *Bopyrione toloensis* Markham, 1982.

KEY TO THE 28 SPECIES OF
PARABOPYRELLA MARKHAM, 1985,
BASED ON MALE AND FEMALE
CHARACTERS

- 1a. Posterior edge of female pleotelson entire 2
 - 1b. Posterior edge of female pleotelson not entire 15
 - 2a. Posterior edge of female pleotelson truncate or very weakly convex 3
 - 2b. Posterior edge of female pleotelson strongly convex 4
 - 3a. Body symmetrical *P. symmetros*, n. sp.
 - 3b. Body asymmetrical
..... *P. perplexa* Markham, 1990
 - 4a. Posterior edge of female pleotelson rounded 8
 - 4b. Posterior edge of female pleotelson not rounded 5




Fig. 18. *Parabopyrella symmetros* n. sp., holotype female (CIEAL800505) (A–G): A. Dorsal view. B. Ventral view. C. Right maxilliped, external view. D. Right side of barbula. E. Right oostegite 1, external view. F. Right oostegite 1, internal view. G. Right pereopod 7. Allotype male (CIEAL800506) (H, I): H. Dorsal view. I. Ventral view. Scale: 2 mm (A, B); 1 mm (C, D); 1.5 mm (E, F); 0.55 mm (G); 0.5 mm (H, I).

- 5a. Anterior margin of female head crenulated *P. crenulata* (Shiino, 1939)

5b. Anterior margin of female head smooth .. 6

6a. Dorsolateral bosses reduced on both sides of female pereon *P. choprail* (Nierstrasz and Brender à Brandis, 1929)

6b. Dorsolateral bosses on at least one side of female's first four pereomeres 7

7a. Distal margin of female pleotelson acute *P. delagoae* (Bourdon, 1982)

7b. Distal margin of female pleotelson blunt *P. angulosa* (Bourdon, 1980a)

8a. Coxal plates and dorsolateral bosses of female absent ... *P. setoensis* (Shiino, 1939)

8b. Coxal plates and dorsolateral bosses of female present 9

9a. Lateral margin of pleon of male deeply indented 10

9b. Lateral margin of pleon of male slightly indented 12

10a. Median of female pleon fused, segments obscure *P. angusta* (Shiino, 1963)

10b. Median of female pleon not fused, only last two pleomeres fused 11

11a. Male with five pairs of pleopods. *P. nierstrazi* (Chopra, 1930)

11b. Male with four pairs of pleopods *P. tanyensis* (Bourdon, 1979)

12a. Female barbula with one pair of lateral projections *P. saronae* (Bourdon and Bruce, 1979)

12b. Female barbula with two pairs of lateral projections 13

13a. First oostegite with long posterior portion ... *P. essingtoni* (Bourdon and Bruce 1983)

13b. Oostegite 1 with short posterior portion .. 14

14a. Pleon of male much broader than pereon *P. richardsonae* (Nierstrasz and Brender à Brandis, 1929)

14b. Pereon of male equal to or broader than pleon *P. mortensenii* (Nierstrasz and Brender à Brandis, 1929)

15a. Distal edge of female pleon deeply separated 16

15b. Distal edge of female pleon slightly indented 17

16a. Pereon of female much wider than pleon *P. lata* (Nierstrasz and Brender à Brandis, 1929)

16b. Pereon of female equal to or slightly wider than than pleon *P. megatelson* (Nierstrasz and Brender à Brandis, 1929)

17a. Divergent pleotelson of female terminally rounded 18

17b. Divergent pleotelson of female not terminally rounded 20

18a. Male with slender pleon, all segments fused, without lateral indentations *P. thomasi* (Nierstrasz and Brender à Brandis, 1929)

18b. Male with broad pleon, lateral indentations indicating segments 19

19a. Barbula of female with one pair of lateral projections *P. intermedia* (Nierstrasz and Brender à Brandis, 1923)

19b. Barbula of female with two pairs of lateral projections *P. distincta* (Nierstrasz and Brender à Brandis, 1923)

20a. Pleotelson of female divergent and terminally quadrate 21

20b. Pleotelson of female slightly divergent, terminus with two points 25

21a. Male with uropods *P. pacifica* (Shiino, 1933)

21b. Male without uropods 22

22a. Male pleomeres not completely fused ... 23

22b. Male pleon fused, segments indicated only by lateral indentations 24

23a. Male pleon segments indicated by dorsal obscure sutures and deep lateral indentations *P. barnardi* (Nierstrasz and Brender à Brandis, 1931)

23b. Male pleon segments without any fusion, distinct *P. cuspidata*, n. sp.

24a. Female with eyes *P. elongata* (Shiino, 1949)

24b. Female without eyes *P. australiensis* (Bourdon, 1980a)

25a. Pleomeres of male fused, with very weak lateral indentations 26

25b. Pleomeres of male fused only medially ... 27

26a. Male first two pleomeres with midventral tubercles *P. bonnieri* (Nierstrasz and Brender à Brandis, 1923)

26b. Male first pleomere only with midventral tubercle *P. incisa* (Chopra, 1923)

27a. Male first two pleomeres much wider than pleon *P. hodgarti* (Chopra, 1923)

27b. Male first pleomere only wider than pleon *P. indica* (Chopra, 1923)

Probopyriscus Markham, 1982

DIAGNOSIS (modified from Markham, 1982): Female: Body approximately half as long as

wide, slightly distorted; head distinct and extended anteriorly, with long frontal lamina; maxilliped palp extended and articulated with large spur; barbula with single blunt projection on each side with trace of second projection mesially; all pereomeres distinct with coxal plates covering margins; oostegites reduced and exposing brood pouch, oostegite 1 slightly pointed and internally unadorned; pleon of six distinct pleomeres well separated laterally; five pairs of flaplike biramous pleopods; no uropods.

Male: Head distinctly separated laterally from first pleomere; all pereomeres distinct and separated by lateral notches; pereopods small, all equally developed, with meri and carpi fused; pleon of six pleomeres indistinctly separated dorsally but deeply separated laterally; five pairs of buttonlike pleopods; no uropods.

TYPE SPECIES: *Probopyriscus novempalensis* Markham, 1982, by original designation.

OTHER SPECIES: None.

REMARKS: Markham (1982) compared the characters of the type and sole species of *Probopyriscus* with those of several other bopyrines, including the western Atlantic *Parabopyriscus stellatus* and he indicated that the diagnosis of the latter genus and species was in press; that paper was eventually published as Markham (1985a). By introducing *Parabopyriscus stellatus* as a genus and species combination with detailed morphological characters that clearly differentiated these taxa from all others, Markham (1982, not 1985a) made these names available (ICZN, 1999: Article 13) and, as there was only a single species stated to be in *Parabopyriscus*, *P. stellatus* is the type species by monotypy.

Probopyriscus novempalensis Markham, 1982

Probopyriscus novempalensis Markham, 1982: 293–296, fig. 1 [Hong Kong, infesting *Alpheus* sp.]; Li, 2003: 140, 158.

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Alpheus* sp. (Alpheidae), Hong Kong.

REMARKS: This species is known only from the holotype and allotype pair.

Septembopyrina, n. gen.

DIAGNOSIS: Female: Body slightly distorted, pleomeres distinct. Head fused with first pereomere medially and with well-developed frontal lamina, boundary with head obscure. Maxilliped without palp, barbula with pair of projections. Dorsolateral bosses only on first pereomere. Seven pairs of oostegites, brood pouch widely open. Oostegite 1 of left side and right side slightly different in shape, internal ridges smooth. Seventh oostegite smallest. Pleon of five pleomeres, final one very broad. Five pairs of uniramous flaplike pleopods, uropods lacking.

Male: Body width about 1/3 length. Head fused with first pereomere, lateral indentations present. No midventral tubercles. Pereopods smaller posteriorly. Pleon segments distinct ventrally but fused dorsally.

ETYMOLOGY: The prefix "septem" (for the seven oostegites of the female) plus the generic name *Bopyrina* are selected to emphasize the close relationship to *Bopyrina*; gender feminine.

TYPE SPECIES: *Septembopyrina tozeumao-phila*, n. sp., by original designation.

OTHER SPECIES: None.

REMARKS: The new genus is related to *Schizobopyrina* Markham, 1985, and *Bopyrina* in that all have unequal formation of oostegite 1, head somewhat fused with the first pereomere, uniramous pleopods and lack uropods. Markham (1985a) separated *Schizobopyrina* from *Bopyrina* and transferred 10 species from *Bopyrina* to *Schizobopyrina*. He showed that *Schizobopyrina* females have a maxilliped palp, elongate oostegites 2–5, and at least lateral separation of the pleomeres, but *Bopyrina* females lack a maxilliped palp, have tiny oostegites 2–5, and pleomere fusion on the shorter side. This new genus is most closely related to *Bopyrina*, but *Septembopyrina*, n. gen., has the female pleomeres distinct on both sides and seven pairs of oostegites. The new genus also can be distinguished from *Schizobopyrina* by the weakly asymmetrical body (strongly asymmetrical in *Schizobopyrina*), five pairs of uniramous pleopods (four pairs in *Schizobopyrina*), and maxilliped without palp (with palp in *Schizobopyrina*). The presence of seven pairs of oostegites is very uncommon and considered a primitive

character state in bopyrids (Boyko and Williams, 2010); the only prior records from any bopyrines are from *Bopyrinella albida* and *Capitetragnia elliptica*, but the type species of *Septembopyrina*, n. gen., does not otherwise resemble these two species except in this one character.

***Septembopyrina tozeumaophila*, n. sp.**

Figure 19

Parabopyrina tozeuma An, 2006: 74–75, fig. 32 (unavailable name).

MATERIAL EXAMINED: Infesting *Tozeuma lanceolatum* Stimpson, 1860, holotype ♀ (CIEHI790502a), allotype ♂ (CIEHI790502b), Beibu Gulf, Stn. 7905, 108°30'E, 18°30'N, 29 m, 6 January 1962, coll. Fuzeng Sun. Paratypes: 1 ♀, 2 ♂ (CIEHI620301), South China Sea, Stn. 6203, 109°00'E, 18°15'N, 23.8 m, 16 July 1959, coll. Fengshan Xu. Other material: 2 ♀♀, 1 ♂ (CIEHI790501), Beibu Gulf, Stn. 7905, 108°30'E, 18°30'N, 26 m, 16 August 1962, coll. Fuzeng Sun; 2 ♀♀ (CIEHI405401), East China Sea, Stn. 4054, 123°30'E, 29°30'N, 69 m, 8 December 1959, coll. Yongliang Wang and Jinzhou Zhu; 1 ♀ (CIEHI590201), East China Sea, Stn. 4054, 123°30'E, 29°30'N, 86 m, 1 February 1959, coll., Jieshan Xu; 2♂♂ (CIEHI622201), Beibu Gulf, Stn. 6222, 108°30'E, 18°45'N, 23 m, 26 January 1959, coll. Zhengang Fan; 1 ♀ (CIEHI606601), South China Sea, Stn. 6066, 113°30'E, 20°30'N, 88 m, 9 April 1960, coll. Zhican Tang; 2 ♀♀ (CIEHI405402), East China Sea, Stn. 4054, 123°30'E, 29°30'N, 75 m, 7 April 1959, coll. Yulin Liao and Mingshou Li.

DESCRIPTION: Holotype female (CIEHI790502a): Length 4.41 mm, maximal width 2.94 mm, head length 0.76 mm, head width 1.24 mm, slightly distorted (fig. 19A, B).

Head an inverted trapezoid, lateral margin distinct, posterior margin fused and embedded in first pereomere. Head with large frontal lamina, boundary between head and frontal lamina obscure; without eyes (fig. 19A). Antennule and antenna of two and three articles, respectively, terminally without setae (fig. 19C). Maxilliped (fig. 19D) without palp, first article cube shaped, short and with blunt spur. Barbula (fig. 19E) with pair of

falcate lateral projections on each side and pair of triangular tubercles medially.

Pereon broadest across third and fourth pereomeres (fig. 19A). Dorsolateral bosses only on first pereomeres (fig. 19A). Brood pouch widely open (fig. 19B), with seven pairs of oostegites. Oostegite 1 of left side and right side asymmetrical: oostegite 1 of left side (fig. 19F, G) with broad posterolateral point, of right side (fig. 15H, I) with triangular posterolateral point. Oostegite 1 with smooth internal ridge; oostegites 2, 3 larger than others, oostegite 7 smallest. Pereopods of similar size and structure (fig. 19J, K). Pleon of five pleomeres, first four narrow with obscure sutures. Fifth pleomeres broad and terminally extended into two small tubercles. Five pairs of uniramous flaplike pleopods, uropods lacking (fig. 19B).

DESCRIPTION: Allotype male (CIEHI790502b): Length 1.50 mm, maximal width (across pereomere 2) 0.45 mm, head width 0.31 mm, head length 0.17 mm, pereon length 0.95 mm (fig. 19L, M). Head elliptical, fused with first pereomere but separated by lateral notch between head and pereon (fig. 19L). Black eyes in lateral corners (fig. 19L). Antennule of two articles, antenna of three articles, terminally nonsetose. Pereomeres distinct, lacking midventral projections (fig. 19M). Pereopods with similar structure, but posteriorly smaller (fig. 19M). Pleon with first three segments distinct, last three segments fused but indicated by lateral indentations and ventral sutures. Pleopods and uropods lacking (fig. 19I).

VARIATION: One female from CIEHI790501 is different from the other female specimen in the lot as well as all other material as it has very reduced oostegites.

ETYMOLOGY: The specific name *tozeumaophila* refers to the generic name of the host shrimp.

HOST AND LOCALITIES: Infesting *Tozeuma lanceolatum* Stimpson, 1860 (Hippolytidae), Beibu Gulf, South China Sea, East China Sea, 23–88 m.

REMARKS: The present specimens represent a new genus and species related to *Bopyrina* and *Schizobopyrina*. *Tozeuma lanceolatum* is first recorded here as a host for parasitic isopods. Other species of bopyrids known to




Fig. 19. *Septembopyrina tozeumaophila* n. sp., holotype female (CIEHI790502a) (A–K): A. Dorsal view. B. Ventral view. C. Right antenna and antennule. D. Right maxilliped, external view. E. Right side of barbula. F. Right oostegite 1, external view. G. Right oostegite 1, internal view. H. Left oostegite 1, external view. I. Left oostegite 1, internal view. J. Right pereopod 1. K. Right pereopod 7. Allotype male (CIEHI790502b) (L, M): L. Dorsal view. M. Ventral view. Scale: 2 mm (A, B); 0.25 mm (C); 0.5 mm (D–K); 0.6 mm (L, M).

parasitize *Tozeuma* hosts are *Parabopyrella thomasi* (Nierstrasz and Brender à Brandis, 1929) (infesting *T. carolinense* Kingsley, 1878, in St. Thomas) and *Schizobopyrina lobata* (Bourdon and Bruce, 1983) (infesting *Tozeuma* sp. in Queensland, Australia).

Hemiarthrinae Markham, 1972

DIAGNOSIS: Female: Body highly distorted. Head deeply embedded into pereon. Pereomeres only distinct on short side. Brood pouch closed, covering entire ventral and lateral side of body. Seven pairs of pereopods usually present on short side. Pleomeres with variable degree of fusion, first four pairs of pleopods with lateral plates. Uropods present.

Male: Pereon segments distinct, without midventral tubercles. Pleon fused, without any pleopods or uropods.

REMARKS: Most of the 55 species, in 27 genera, placed in Hemiarthrinae are ventral abdominal parasites of caridean shrimps (although a few species infest caridean hosts dorsoventrally or in the region of the mouth-parts). Previously, only five species from Chinese waters were reported, all from Hong Kong (Markham, 1982; 1990).

Anchiarthrus Markham, 1992

DIAGNOSIS (modified from Markham, 1992a): Female: Body distorted less than 90°; head lacking antennal groove; barbula with single projection on each side. First and second pereopods of long side large, others small; first and second pereopods on short side absent others small and clustered near pleon. Oostegite 1 dimorphic, smaller one much longer than wide. Pleonal separation indistinct with three pleomeres indicated; three pairs of lanceolate lateral plates, progressively smaller posteriorly; three pairs of biramous pleopods, progressively larger posteriorly.

Male: Body slightly more than twice as long as broad. Head distinct from pereon; antennae prominently extended beyond margin of head. Last pereomere fused with pleon, remnants of pleomeres indicated laterally.

TYPE SPECIES: *Anchiarthrus derelictus* Markham, 1992, by original designation.

OTHER SPECIES: None.

Anchiarthrus derelictus Markham, 1992

Anisarthrus (?), sp. nov. Markham, 1990: 555, 564.

Anchiarthrus derelictus Markham, 1992a: 296–297, fig. 17.

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Arete dorsalis* Stimpson, 1860 (Alpheidae), Hong Kong, 1 m.

REMARKS: As noted by Markham (1990, 1992a), *Anchiarthrus derelictus* is most similar to *Anisarthrus pelseneeri* Giard, 1907 (the type species of *Anisarthrus* Giard, 1907; see Giard, 1907), but differs from it in the female having two large pereopods, fewer than seven pereopods on the shorter side of the body, and three distinguishable pleomeres.

Apophrixus Nierstrasz and Brender à Brandis, 1931

DIAGNOSIS: (modified from Nierstrasz and Brender à Brandis, 1931, Markham, 1982): Female: Body weakly asymmetrical, one side more swollen and longer than other. Head large. Dorsal side with four distinct pereomeres; pleomeres fused. Seven pereopods on short side of body, 6 or 7 on long side; pereopods 4–6 smaller than others. Pleon with four pairs of well-developed lateral plates and four pairs of uniramous pleopods; uropods uniramous.

Males: Pleomeres fused into single large segment, no pleopods or uropods.

TYPE SPECIES: *Apophrixus philippinensis* Nierstrasz and Brender à Brandis, 1931, by monotypy.

OTHER SPECIES: *Apophrixus constrictus* Markham, 1982.

Apophrixus constrictus Markham, 1982

Apophrixus constrictus Markham, 1982: 378–381, fig. 30–31.

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Alpheus* sp. (Alpheidae), Hong Kong.

Dicopleon Markham, 1972

DIAGNOSIS (modified from Markham, 1972): Female: Body weakly asymmetrical; head deeply embedded in pleon. First two pereopods on long side well developed, third pereopod present but much smaller, other pereopods lacking. Pleon with four pairs of well developed lateral plates and four pairs of pleopods, first three biramous; fifth pleomere divided into two large terminal lobes, uropods lacking.

Male: Head and pleon medially fused with pereon; eyes present, antennae elongate. Pleon ending in sharp double point.

TYPE SPECIES: *Dicopleon periclimenis* Markham, 1972, by original designation.

OTHER SPECIES: *Dicopleon bifidus* (Bourdon, 1967), *D. morator* Markham, 1982, *D. processae* Markham, 1980.

Dicopleon morator Markham, 1982

Dicopleon morator Markham, 1982: 381–384, figs. 32–33 [Hong Kong, infesting *Periclimenis commensalis* Borradaile, 1915]; Bruce, 1982: 237 (species erroneously cited as described in "Markham, 1981" (lapsus for 1982)); Li, 2003: 153, 157.

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Periclimenis commensalis* Borradaile, 1915) (Palaemonidae), Hong Kong.

REMARKS: Although the mention of the species by Bruce (1982) occurred on an earlier page in the same volume in which Markham (1982) described the species, it is technically not a nomen nudum as Bruce (1982) was published at the same time as Markham (1982) and there is no page priority rule in the International Code of Zoological Nomenclature.

Eophrixus Caroli, 1930

DIAGNOSIS: Female: Body distorted with swollen brood pouch. All seven pairs of pereopods well developed and of same size. Pleon of five segments, first four pleomeres with lobate lateral plates and biramous pleopods. Endopodite of pleopods often smaller than exopodites; uropods usually small.

Male: Longer than wide, head often fused with first pereomere. Pleon fused into single piece, without pleopods.

TYPE SPECIES: *Phrixus (Eophrixus) lysmatae* Caroli, 1930, by subsequent designation of Markham, 1982.

OTHER SPECIES: *Eophrixus adriaticus* (Nierstrasz and Brender à Brandis, 1931), *E. brevicauda* (Chopra, 1923) n. comb., *E. caudatus*, n. sp., *E. enchocephalus* Caroli, 1930, *E. kuboi* (Shiino, 1939), *E. laevimanus* Caroli, 1930, *E. leptochelae* (Pillai, 1966) n. comb., *E. nigrocinctus* (Chopra, 1923), *E. pikei* (Bruce, 1968), n. comb., *E. shoji* Shiino, 1941, *E. subcaudalis* (Hay, 1917).

REMARKS: Caroli (1930) established *Eophrixus*, containing three species, as a subgenus of "Phrixus" (= *Hemiarthrus* Giard and Bonnier, 1887), but only one species, *Phrixus (Eophrixus) lysmatae* Caroli, 1930, was well described, leading Markham (1985a) to conclude that the other two species, *Phrixus (Eophrixus) enchocephalus* Caroli, 1930, and *Phrixus (Eophrixus) laevimanus* Caroli, 1930, were nomina nuda. However, Caroli (1930) mentioned morphological characters (mostly color) and provided photographs (of poor quality) for both these species therefore, although both are poorly described and not included in the key below, the names are available under ICZN Article 12.2.7 (applying to names published before 1931). The earliest author to mention a type species for the genus appears to be Markham (1982), but he indicated that *Phrixus (Eophrixus) lysmatae* was the type species by monotypy, which is incorrect. Markham (1982) actually selected *Phrixus (Eophrixus) lysmatae* as the type species of *Eophrixus* by subsequent designation. Nierstrasz and Brender à Brandis (1931) raised *Eophrixus* to generic status and Caroli (1949) synonymized *Hypophryxus* Shiino, 1934, with *Eophrixus*.

Markham (1992a) purportedly removed *Hypophryxus* from synonymy with *Eophrixus*, but a reading of that paper shows that he was, in fact, purporting to remove from synonymy *Hyperphrixus*, a genus that has never been synonymized with any other. Markham (1992a) appeared to transfer *Hemiarthrus filiformis* Chopra, 1923, to *Hypophryxus* (e.g., pp. 277, 291 and fig. 15, 16 figure captions

on pp. 292–293) but also on (pp. 292–293) gave the new combination as *Hyperphryxus* [sic] *filiformis* and erroneously referred to *Hyperphryxus* [sic] Shiino, 1934, when *Hyperphryxus* was a genus erected by Nierstrasz and Brender à Brandis, 1931. He also cited the type species of *Hyperphryxus* as *Hyperphryxus* [sic] *yusakiensis* Shiino, 1934, when it is *Hyperphryxus tattersalli* Nierstrasz and Brender à Brandis, 1931. Apparently, Markham (1992a) became confused by the similarity of the genus names *Hyperphryxus* and *Hypophryxus* and confounded their characters and type species. Based on the key characters of *H. filiformis*, including having fewer than seven pereopods on one side of the body, this species belongs to *Hyperphryxus*, not *Hypophryxus*. The correct name for the species is therefore *Hyperphryxus filiformis* (Chopra, 1923) (new combination). Markham's (1992a) statement about removing *Hyperphryxus* [sic] from synonymy with *Eophrixus* is likewise incorrect as *Hyperphryxus* was never synonymized with *Eophrixus* by Caroli (1949) or any other author. *Hypophryxus* was synonymized with *Eophrixus* by Caroli (1949) bases on the fact that the type species (*H. yusakiensis*) has all the characters of *Eophrixus* and thus we concur that the two genera should be considered synonymous. Pillai (1966) and Bruce (1968) each described a species in *Hypophryxus*, both unaware of Caroli's (1949) synonymy of that genus with *Eophrixus*. The result of this is that *Eophrixus* contains 13 species: the seven included by Markham (1985a), the two poorly described species of Caroli (1930), *Hemiarthrus brevicauda* Chopra, 1923, *Hypophryxus leptochelae* Pillai, 1966, *Hypophryxus pikei* Bruce, 1968, and the new species described herein.

Eophrixus brevicauda (Chopra, 1923),
n. comb.

? “Parasite” de Man, 1913: 263.
Hemiarthrus brevicauda Chopra, 1923: 416,
 419, 430, 431, 439–440, pl. 11, figs. 8–9. —
 Monod, 1933: 244.
Phrixus (Paraphrixus) brevicauda: Caroli,
 1930: 259.
Paraphrixus brevicauda: Nierstrasz and Bren-
 der a Brandis, 1931: 205. —Markham,
 1992a: table 1.

? *Paraphrixus brevicauda*: Markham, 1990:
 563–564.

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Synalpheus hastilicrassus* Coutière, 1905, *S. tumidomanus tumidomanus* (Paulson, 1875) (Alpheidae), Indonesia (?), Andaman Islands, Hong Kong, 2–9 fathoms (= 3.7–16.5 m).

REMARKS: Markham (1985a) transferred the type species of *Paraphrixus*, *Phryxus subcaudalis* Hay, 1917, to *Eophrixus*, synonymizing the two genera. However, Markham (1990, 1992a) continued to use the combination *Paraphrixus brevicauda* while admitting (1990) that “*Paraphrixus* is no longer a valid name, but revision of the genus is not appropriate here.” It is clear that, based on current data, *Hemiarthrus brevicauda* must be placed in *Eophrixus*, as this is the genus with which it shares the most characters and, as a synonymized genus, it cannot be used as a valid name. Rediscovery of *E. brevicauda* is desirable in order to determine whether it really belongs in *Eophrixus*, as Chopra's (1923) description is incomplete and the type specimens are inaccessible.

Eophrixus caudatus, n. sp.
 Figures 20, 21A, B

Eophrixus shojii An, 2006: 172–173, fig. 88
 (not *Eophrixus shojii* Shiino, 1941).

MATERIAL EXAMINED: Infesting *Alpheus microstythus* (Bate, 1888) (fig. 17A, B), holotype ♀ (CIEAL800501), allotype ♂ (CIEAL 800502): Xisha Coral Island, 16°32' N, 111°36'E, 9–21 May 1980.

DESCRIPTION: Holotype female (CIEAL 800501): Length 6.11 mm, including lateral plates, maximal width 4.43 mm, including brood pouch, head length 0.56 mm, head width 1.56 mm (figs. 20A, 21A, B).

Body strongly asymmetrical, with swollen pereon. Head inserted into pereon, with bilobate anterior and posterior margins, without eyes (fig. 20A). Antennule of three articles, antenna of four articles, both setose, with stout bases (fig. 20C). Maxilliped rectangular, without palp, plectron blunt, anterior article 3 times larger than posterior article (fig. 20D).




Fig. 20. *Eophrixus caudatus*, n. sp., holotype female (CIEAL800501) (A–G): A. Dorsal view. B. Ventral view of pleon. C. Left antenna and antennule. D. Left maxilliped. E. Barbula of right side. F. Right oostegite 1, external view. G. Right oostegite 1, internal view. H. Left pereopod 5. I. Left pereopod 6. Allotype male (CIEAL800502) (J–N): J. Dorsal view. K. Ventral view. L. Right antenna and antennule. M. Ventral view of pleon. N. Right pereopod 7. Scale: 1 mm (A); 0.6 mm (B, E–G, J, K); 0.45 mm (D); 0.2 mm (C, N); 0.15 mm (H, I, L, M).

Barbula with two pairs of simple falcate lateral projections on each side (fig. 20E).

Left side of body somewhat fused, but right side distinctly segmented with pereopods

closely crowded together. Dorsolateral bosses and coxal plates absent. Swollen brood pouch completely closed (fig. 20A). Oostegite 1 (fig. 20F, G) with smooth internal ridge,




Fig. 21. Hemiarthrine bopyrids and hosts. **A.** Host of *Eophrixus caudatus*, n. sp. **B.** Female, male and eggs of *E. caudatus*. **C.** Host of *Sigyn branchialis*, n. gen., n. sp. (right branchial chamber bulge indicating bopyrid, arrows indicating rhizocephalans). **D.** Dorsal view of female of *S. branchialis*, n. gen., n. sp. with attached male.

posterolateral point rounded and directed laterally. Seven pereopods crowded together on short side, but first two pereopods on long side near head, third pereopod inserted in brood pouch, last four pereopods crowded between brood pouch and lateral plates. All pereopods slender, with long meri, dactyli longer in posterior pereopods (fig. 20H, I). Pleon with four pairs of lateral plates and biramous pleopods, segmentation indistinct. All lateral plates and exopodites of pleopods lobate, endopodites well developed (fig. 20B). Terminal pleomere with distinct uniramous uropods (fig. 20B).

DESCRIPTION: Allotype male (CIEAL8 00502): Length 3.53 mm, maximal width (across pleon 4) 0.93 mm, head width 0.67 mm, head length 0.33 mm (fig. 20J, K). Head semicircular (fig. 20J), fused with first pereomere; small black eyes posterolaterally (fig. 20J). Antennule of three articles, almost smooth, antenna of six articles, terminal two articles setose (fig. 20L). All pereomeres subequal in width, lacking midventral projections (fig. 20K). All pereopods of same size and structure, carpi and meri terminally setose, dactyli blunt (fig. 20N). Pleon fused into single piece, without any pleopods; lateral margins straight, without any notches or incisions. Posterior margin of pleon with pair of fingerlike uropodal rami, anal cone in median (fig. 20M).

ETYMOLOGY: The specific name, *caudatus*, refers to the female bearing a pair of distinct, forked, fingerlike uropods.

HOST AND LOCALITY: Infesting *Alpheus microstylus* (Bate, 1888) (Alpheidae), Xisha (Paracel Islands), China.

REMARKS: The female of the new species has a swollen brood pouch, seven pairs of pereopods on both sides, and the male has a fused pleon, indicating that the present specimens belong to *Eophrixus*. The new species can be distinguished from the other 10 well-described species by its unique forked fingerlike uropods. *Eophrixus caudatus*, n. sp., is most closely related to *E. shoji* Shiino, 1941, but the new species differs from *E. shoji* as follows: (1) the first lateral plates of the new species are almost equal in size to the others, whereas the first lateral plates of *E. shoji* are much smaller than the others; (2) the female of the new species has prominent forked

uropods while the uropods of *E. shoji* are small and globular; (3) the male of the new species lacks any indentation or notch on the pleon margin, but the male of *E. shoji* shows a distinct series of indentations indicating the pleomeres.

Eophrixus shoji Shiino, 1941

Eophrixus shoji Shiino, 1941: 155–156, figs. 1–2 [Japan, infesting *Alpheus japonicus* Miers, 1879]; Shiino, 1958: 71 [Japan, infesting *Alpheus rapax* Fabricius, 1798]; Danforth, 1963: 8; Markham, 1972: 48; Markham, 1982: 275, figs. 28–29 [Hong Kong, infesting *Alpheus lobidens?* De Haan, 1849]; Page, 1985: 185, 205–208, fig. 15 [New Zealand, infesting *Alpheus socialis* Heller, 1862]; Kim and Kwon, 1988: 199, 218–219, fig. 11 [Korea, infesting *A. japonicus*]; Markham, 1990: 555, 563 [Hong Kong, infesting *Alpheus brevirostris* (Olivier, 1811)]; Markham, 1991: 289, 294, fig. 3 [Thailand, infesting *Alpheus* sp.]; Markham, 1992a: 298; Kensley, 2001: 223; Li, 2003: 140, 154, 157; Liu, 2008: 692.

Anisarthrus shoji: Codreanu and Codreanu, 1956: 119; Codreanu, 1961: 138.

Eophryxus [sic] *shoji*: Shiino, 1972: 9.

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Alpheus brevirostris* (Olivier, 1811), *A. japonicus* Miers, 1879, *A. lobidens?* De Haan, 1849, *A. rapax* Fabricius, 1798, *A. socialis* Heller, 1862, *Alpheus* sp. (Alpheidae), Japan, Korea, Hong Kong, Thailand, New Zealand.

KEY TO 11 SPECIES OF *EOPHRIXUS* CAROLI, 1930, BASED ON MALE AND FEMALE CHARACTERS (*E. ENCHOPHYLLUS* AND *E. LAEVIMANUS* NOT INCLUDED IN THE KEY)

- | | | |
|-----|--|---|
| 1a. | Female without uropods | 2 |
| 1b. | Female with uropods | 3 |
| 2a. | Five pleomeres, exopods of pleopods well developed . . . <i>E. pikei</i> (Bruce, 1968), n. comb. | |
| 2b. | Four pleomeres, exopods of pleopods weakly developed . . . <i>E. kuboi</i> (Shiino, 1939) | |
| 3a. | Pereomeres I and II with serrate margins . . . <i>E. leptochelae</i> (Pillai, 1966), n. comb. | |
| 3b. | Pereomeres I and II with smooth margins | 4 |

- 4a. Uropods as slender rami 5
 4b. Uropods large, foliaceous 6
 5a. Lateral plates much larger than associated pleopods 7
 5b. Lateral plates nearly same size as associated pleopods *E. caudatus*, n. sp.
 6a. Male pleotelson with mediodistal bifurcation *E. subcaudalis* (Hay, 1917)
 6b. Male pleotelson elongate, entire *E. yusakiensis* (Shiino, 1934)
 7a. Lateral plates of pleon globular *E. brevicauda* (Chopra, 1923)
 7b. Lateral plates of pleon plate shaped 8
 8a. First pair of lateral plates much smaller than others *E. shoji* Shiino, 1941
 8b. Lateral plates subequal in size 9
 9a. Pereon with distinct pigment *E. nigrocinctus* (Chopra, 1923)
 9b. Pereon without pigment 10
 10a. Pleon with single small globular lobe terminally *E. lysmatae* Caroli, 1930
 10b. Pleon with a pair of globular uropods terminally *E. adriaticus* (Nierstrasz and Brender à Brandis, 1931)

Hyperphrixus Nierstrasz and Brender à Brandis, 1931

DIAGNOSIS (after Markham, 1985a): Female: Body axis distorted more than 90°, outline subcircular. Head deeply set into pereon, not bisecting any pereomeres; seven pereopods on concave side, clustered near head or evenly spaced between head and pleon; one or two pereopods on convex side. Five pleomeres, first four bearing well-developed, lanceolate, uniramous lateral plates and biramous pleopods; uropods lacking.

Male: Body more than three times as long as broad. Head and pereon separated. Sides of pereon subparallel; all pereomeres distinct but not deeply separated laterally. Pleon swollen, without traces of segmentation or appendages.

TYPE SPECIES: *Hyperphrixus tattersalli* Nierstrasz and Brender à Brandis, 1931, by monotypy.

OTHER SPECIES: *Hyperphrixus castrensis* Markham, 1985a; *Hyperphrixus filiformis* (Chopra, 1923)

Hyperphrixus filiformis (Chopra, 1923), n. comb.

Hemiarthrus filiformis Chopra, 1923: 416, 428, 431, 435–438, text fig. 2, pl. 9, figs. 4–6 [Andaman Islands, infesting *Alpheus paracyone* Coutière, 1905]; Chopra, 1930: 118; Monod, 1933: 244.

Hemiarthrus filiformis var. *attenuata* Chopra, 1923: 419, 428, 430, 431, 438–439, pl. 11, fig. 7 [Andaman Islands, infesting *Alpheus paracyone* Coutière, 1905]; Chopra, 1930: 118–119; Monod, 1933: 244.

Phrixus (Paraphrixus) filiformis—Caroli, 1930: 259, 265.

Paraphrixus filiformis: Nierstrasz and Brender à Brandis, 1931: 205.

Hypophryxus filiformis: Markham, 1992a: 293–295, fig. 15–16 [Hong Kong, infesting *Alpheus malleodigitus* (Bate, 1888)]; Kensley, 2001: 224; Li, 2003: 140, 158.

Eophrixus filiformis: Kensley, 2001: 223.

MATERIAL EXAMINED: None.

HOSTS AND LOCALITIES: Infesting *Alpheus paracyone* Coutière, 1905, and *Alpheus malleodigitus* (Bate, 1888) (Alpheidae), Andaman Islands and Hong Kong.

REMARKS: Markham (1992a) synonymized *Hemiarthrus filiformis* var. *attenuata* with the nominal species, but erroneously placed it into *Hypophryxus* (see Remarks under *Eophrixus* above).

Sigyn, n. gen.

DIAGNOSIS: Female: Head highly distorted, embedded in pereon. Seven pereopods on both sides, but crowded together on short side. First two pereopods of long side near head, third on brood pouch located opposite head, other four pereopods of long side crowded together near pleon. Pleon of five segments, first four with lateral plates and uniramous pleopods. Pleon produced into a pedunculated globe, surface covered with small tubercles.

Male: Head fused with first pereomere, eyes present. Pereomeres distinct. Pleon without segmentation or appendages.

ETYMOLOGY: The name *Sigyn* (“victorious girlfriend” in Old Norse) calls attention to the closely related genus *Loki* Markham,

1972; Sigyn is the wife of Loki in Norse mythology; gender feminine.

***Sigyn branchialis*, n. sp.**

Figures 21C, D, 22, 23

Eophrixus branchialis An, 2006: 174, fig. 89
(unavailable name).

MATERIAL EXAMINED: Infesting *Alpheus digitalis* De Haan, 1844, holotype ♀ (CIEAL 920301), Sanya, 18°14'N, 109°30'E, 23 March 1992. Allotype ♂ (CIEAL920301b), same locality as holotype. Female bopyrid parasiting right branchial chamber of host (fig. 21C, D).

DESCRIPTION: Holotype female (CIEAL 920301a): Length 4.67 mm (including brood pouch), head length 0.67 mm, head width 0.67 mm. pereon length 0.78 mm (fig. 22A, B). Head length equal to width, eyes absent (fig. 22A). Head with frontal lamina. Only a single pair of antennae of five articles present (fig. 22C). Maxilliped with much larger anterior article, without palp, plectron highly blunted and not visible as distinct structure (fig. 22D). Barbula (fig. 22E) not located at exact bottom of maxilliped, inclines left, with seven to eight acute lobes on margin. First pereomere concealed by head, only visible lateral to head. Second to seventh pereomeres distinct only on short side (fig. 22A). Brood pouch large and completely closed (fig. 22B). Oostegite 1 (fig. 22F–I) with distinct asymmetry, left side much larger than right side. Oostegite 1 of left side (fig. 22F, G) with nearly smooth internal ridge, posterolateral point long and sharp, directed laterally. Oostegite of right side (fig. 22H, I) with smooth internal ridge, posterolateral point blunt and posteriorly directed. First two pereopods of long side near head, third pereopod attached to ventral surface of brood pouch, opposite head (fig. 22B). Fourth to seventh pleopods crowded near pleon (fig. 22B). All pereopods of short side arranged in a line (fig. 22A). Pereopods of same size and structure, with long bases and blunt dactyli (fig. 22J, K).

Pleon of five pleomeres, first four pleomeres bearing uniramous flaplike pleopods and lateral plates. Fifth pleomere a bulging sphere, surface covered with tubercles (fig. 22L).

DESCRIPTION: Allotype male (CIEA920302): Length 1.64 mm, maximal width (across

pereomere 4) 0.52 mm, head length 0.16 mm, head width 0.36 mm, pleon length 0.44 mm. All pereon segments distinct (fig. 22M, 23A). Allotype attached at pereon/pleon boundary of holotype. Body elongate, sides nearly parallel except for rounded head and posterior abdomen (fig. 22M). Head subovate, wider than long, fused with pereomere 1. Small dark eyes near posterolateral corners (fig. 22M). Antennule of three articles, terminally setose (fig. 23C). Antenna of five articles, much longer than antennule, terminal two articles setose (fig. 22B). Pereomeres subequal in width, with truncate and setose margins (fig. 23D). All pereopods with similar structure and proportions, carpi and meri fused (fig. 23D, E). Flap-shaped scales with serrated edges on surface of ischia, meri, and carpi (fig. 23D, F). Propodi with many flap-shaped scales embedded in dactyli (fig. 23E), scale surfaces not smooth but covered with micro-scales (fig. 23G). Pereon without midventral projections, but with many setae on ventral surface of pereon (fig. 23H). Pleon fused into single piece, without pleopods or uropods, terminally setose (fig. 23J).

ETYMOLOGY: The specific name, *branchialis*, refers to the type specimens found in the branchial chamber of the host.

HOST AND LOCALITY: Infesting *Alpheus digitalis* De Haan, 1844 (Alpheidae), Hainan Province, China.

REMARKS: The present specimens parasitize a caridean shrimp host, the female has a swollen brood pouch formed by the oostegites from one side of the body, asymmetrical oostegite 1, a highly distorted body, and the male has a fused pleon; therefore this species belongs to Hemiarthrinae. However, no genera in this subfamily described to date are known to infest the branchial chamber of their host shrimps. Most hemiarthrine species are ventral abdominal parasites, but a few are found parasitizing other parts of the shrimp hosts, such as the dorsal pleon (*Filophryxus dorsalis* Bruce, 1972b), or even embedded in the mouthparts (*Orophryxus shiinoi*, Bruce, 1972a). The closest position of a hemiarthrine species on a host to that found with *Sigyn branchialis*, n. gen., n. sp., is seen with *Metaphrixus intutus* Bruce, 1966, which is positioned over the host branchial chamber and is attached underneath a raised flange




Fig. 22. *Sigyn branchialis* n. gen., n. sp., holotype female (CIEAL920301a) (A–L): A. Dorsal view. B. Ventral view. C. Right antenna. D. Right maxilliped. E. Barbula and maxilliped. F. Left oostegite 1, external view. G. Left oostegite 1, internal view. H. Right oostegite 1, external view. I. Right oostegite 1, internal view. J. Left pereopod 4. K. Left pereopod 7. L. Ventral view of the pleon. Allotype male (CIEAL920301b) (M): M. Dorsal view. Scale: 1 mm (A, B); 0.15 mm (C, D); 0.35 mm (E); 0.25 mm (F, G, K); 0.20 mm (H–J); 0.4 mm (L); 0.3 mm (M).

derived from the first abdominal pleuron of the host. The branchial mode of parasitism is considered to be the ancestral condition for bopyrids (see Boyko et al., 2013) and *Sigyn branchialis*, n. gen., n. sp., with its branchial position and full complement of pereopods on both sides of the female body may represent the most primitive hemiarthrine known to date.

The present specimens differ sufficiently from other hemiarthrine genera and deserve generic status. The new genus is most closely

related to *Loki* Markham, 1972, *Eophrixus* Caroli, 1930, *Anisarthrus* Giard, 1907, and *Cataphryxus* Shiino, 1936. *Sigyn*, n. gen., can be distinguished from those four genera by its mode of parasitism within the branchial chamber, female with uniramous pleopods and globular terminal pleomere, and male with fused pleon lacking any lateral notches or appendages. The differences are summarized in table 2. Additionally, the fourth pleopod of the female has a tubercle that may indicate an endopodite (fig. 22L).




Fig. 23. *Sigyn branchialis*, n. gen., n. sp., SEM image of male (CIEAL920301b) (A–J): A. Ventral view. B. Left antenna. C. Right antennule. D. First right three pereopods. E. Right pereopods 6, 7. F. Carpi and meri of left pereopod 2. G. Propodus of left pereopod 2. H. Ventral surface of pereopod 3. I. Terminal edge of pleon. Scale: 100 µm (A); 10 µm (B–F, H, I); 1 µm (G).

There are two saclike organisms parasitizing the abdomen of the host shrimp (fig. 21C). The more anterior specimen is a shriveled sac with a distinct mantle opening,

indicating that it is a rhizocephalan. The posterior specimen also appears to be a rhizocephalan, is ovate and filled with eggs, but there is no mantle pore. It is unclear whether

TABLE 2
Comparison of *Sigyn* n. gen. with Four Other Closely Related Hemiarthrine Genera

	Characters	<i>Sigyn</i>	<i>Loki</i>	<i>Eophrixus</i>	<i>Cataphryxus</i>	<i>Anisarthrus</i>
Female	Parasite mode	Branchial	Abdominal	Abdominal	Abdominal	Abdominal
	Lateral plates	4 pairs	3 pairs	4 pairs	4 pairs	4 pairs
	Pleopods	Uniramous	Uniramous	Biramous	Biramous	Uniramous
	Number of pleopods	7	6 (absent on pleomere 4)	7	7	7
Male	Uropods	Absent	Fork-shaped	Globular	Absent	Absent
	Eyes	Small	Absent	Small	Absent	Large
	Head and pereon	Fused	Fused	Fused	Separated	Fused
	Pereomeres	Distinct	Fused medially	Distinct	Distinct	Distinct
	Pereon and pleon	Separated	Fused	Separated	Separated	Separated
	Pleomeres	Fused	Fused	Defined laterally	Fused	Fused
	Pleotelson posterior margin	Smooth	Anal tube	Anal tube	Smooth	Anal tube

these represent two species or different stages of development of female externae of the same species. The only rhizocephalans known from alpheids are members of *Thylacoplethus* Coutière, 1902 (Cirripedia: Rhizocephala: Akentrogonida), but neither of these belongs to that genus.

ACKNOWLEDGMENTS

The first and second authors contributed equally to this publication. This study was supported by the National Natural Science Youth Foundation (no. 31101614) and National Natural Science Foundation (no. 31471970). We are indebted to all collectors from IOCAS in 1950s and 1989–1991. Thanks to Masayuki Osawa (Shimane University) for identification of *Allorbimorphus tuberculatus*.

REFERENCES

- Adkison, D.L. 1988. *Pseudione parviramus* and *Aporobopyrus collardi*, two new species of Bopyridae (Isopoda: Epicaridea) from the Gulf of Mexico. Proceedings of the Biological Society of Washington 101: 576–584.
- Adkison, D.L., R.W. Heard, and G.T. Clark. 1982. Description of the male and notes on the female of *Argeiopsis inhacae* (Crustacea: Isopoda: Bopyridae). Proceedings of the Biological Society of Washington 95: 334–337.
- Ahyong, S.T. 2006. [Crustacea: Galatheidae, Polychelidae, and Glyphocrangonidae (squatlobsters, blind, deep-sea lobsters and deepwater shrimps)]. In Explore lost worlds of the deep NORFANZ Voyage (chapter 3.1, results, taxonomy): 48–83. Final Report to the Department of Environments and Heritage (Australia's National Oceans Office). Online resource (<http://www.environment.gov.au/apps/coasts/discovery/publications/norfanz-voyage-report.html>).
- An, J.-M. 2006. Study on the taxonomy and zoogeography of the family Bopyridae (Crustacea: Isopoda) in the China Seas [sic]. Ph.D. dissertation, Institute of Oceanology of Chinese Academy of Sciences, vii + 225 pp.
- An, J. 2009. A review of bopyrid isopods infesting crabs from China. Integrative and Comparative Biology 49: 95–105.
- An, J. 2011. Taxonomy of parasitic isopods from China. Beijing: Science Press.
- An, J., H. Yu, and X.-Z. Li. 2006a. Description of a new genus and two new species of Ioninae (Isopoda: Epicaridea: Bopyridae) parasites of Leucosiidae (Decapoda: Brachyura) from Beibu Gulf. Zootaxa 1186: 57–68.
- An, J., H. Yu, and X.-Z. Li. 2006b. A review of the genus *Apocepon* Nierstrasz and Brender à Brandis, 1930 (Isopoda: Epicaridea: Bopyridae) from Chinese waters, with description of a new species and a new record species. Zootaxa 1199: 1–17.
- An, J., H. Yu, and X.-Z. Li. 2007a. One new record genus of Bopyridae (Crustacea, Isopoda, Epicaridea) from China. Acta Zootaxonomica Sinica 32: 1002–1003.
- An, J., H. Yu, and J.D. Williams. 2007b. Four new records and a new species of *Dactylokepon* Stebbing, 1910 (Epicaridea: Bopyridae: Ioninae) from Chinese waters. Journal of Natural History 41: 2063–2079.
- An, J., H. Yu, and X.-Z. Li. 2008. Subfamily Argeiinae (Crustacea, Isopoda, Epicaridea) from China. Acta Zootaxonomica Sinica 33: 225–226.
- An, J., J.D. Williams, and H. Yu. 2009a. The Bopyridae (Crustacea: Isopoda) parasitic on thalassinideans

- (Crustacea: Decapoda) from China. Proceedings of the Biological Society of Washington 122: 225–246.
- An, J., H. Yu, and J.C. Markham. 2009b. First record of the genus *Gigantione* (Epicaridea: Bopyridae: Pseudioninae) from Chinese waters, with descriptions of three new species. Journal of Natural History 43: 335–353.
- An, J., J.C. Markham, and H. Yu. 2010. Description of two new species and a new genus of bopyrid isopod parasites (Bopyridae: Pseudioninae) of hermit crabs from China. Journal of Natural History 44: 2065–2073.
- An, J., J.D. Williams, and H. Yu. 2011. Three abdominal parasitic isopods (Isopoda: Epicaridea: Bopyridae: Athelginae) on hermit crabs from China and Hong Kong. Journal of Natural History 45: 2901–2913.
- An, J., C.B. Boyko, and X.-Z. Li. 2012a. Two new species of the genus *Megacepon* George, 1947 (Crustacea: Isopoda: Bopyridae) infesting Varunidae (Crustacea: Brachyura: Grapoidea) from China. Journal of Natural History 46: 131–143.
- An, J., H. Yu, and C.B. Boyko. 2012b. A review of the genus *Scyracepon* Tattersall, 1905 (Crustacea: Isopoda: Bopyridae), with description of a new species from China. Journal of Natural History 46: 2889–2895.
- An, J., H. Yu, and J.D. Williams. 2012c. A new species of *Cancricepon* Giard and Bonnier, 1887 (Isopoda: Bopyridae) from China. Systematic Parasitology 83: 243–248.
- An, J., X.-Q. Zhang, and X.-Z. Li. 2012d. A new species of Bopyridae (Crustacea, Isopoda, Epicaridea) infesting *Porcellana* sp. Acta Zootaxonomica Sinica 37: 722–725.
- An, J., C.B. Boyko, and X.-Z. Li. 2013a. A new genus and species of parasitic isopod (Isopoda: Bopyridae: Orbioninae) from Chinese waters, with a new locality and host record for *Orbione halipori* Nierstrasz and Brender à Brandis, 1923. Systematic Parasitology 84: 149–156.
- An, J., C.B. Boyko, and X.-Z. Li. 2013b. A review of the genus *Parapenaeonella* (Isopoda: Bopyridae: Orbioninae), with description of a new species from Chinese waters and synonymy of *Parapenaeonella lamellata* Bourdon, 1979. Journal of Natural History 47: 2633–2648.
- An, J., C.B. Boyko, and X.-Z. Li. 2013c. A new species and two new records of *Minicopenaon* Bourdon, 1981 (Crustacea: Isopoda: Bopyridae) from China. Systematic Parasitology 85: 255–262.
- An, J., X. Li, and J.C. Markham. 2013d. Three isopod parasites (Bopyridae: Pseudioninae), including two new species, of hermit crabs from the South China Sea. Raffles Bulletin of Zoology 61: 561–569.
- An, J., C.B. Boyko, and X.-Z. Li. 2014a. Review of the parasitic genus *Epipenaeon* Nobili, 1906 (Crustacea: Isopoda: Bopyridae), with new records and redescription of four species from China. Journal of Natural History 48: 2027–2048.
- An, J., X. Niu, J.C. Markham, and W. Jiang. 2014b. A review of the genus *Rhopalione* Pérez, 1920 (Crustacea: Isopoda: Bopyridae), with first record of the male and parasitic mode of *R. sinensis* Markham, 1990 from China. Chinese Journal of Oceanology and Limnology. 32: 1–4.
- Baer, J.G. 1946. Le parasitisme. Lausanne: F. Rouge et Cie, 232 pp.
- Baer, J.G. 1951. Ecology of animal parasites. Urbana: University of Illinois Press, x + 224 pp.
- Barnard, K.H. 1940. Contributions to the crustacean fauna of South Africa. XII. Further additions to the Tanaidacea, Isopoda, and Amphipoda, together with keys for the identification of the hitherto recorded marine and fresh-water species. Annals of the South African Museum 32: 381–543.
- Barnard, K.H. 1955. Additions to the fauna-list of South African Crustacea and Pycnogonida. Annals of the South African Museum 43: 1–107.
- Bate, C.S. 1864. Characters of new species of crustaceans discovered by J.K. Lord on the coast of Vancouver Island. Proceedings of the Zoological Society of London 1864: 661–668.
- Bate, C.S. 1888. Report on the Crustacea Macrura dredged by H.M.S. *Challenger* during the years 1873–1876. Reports on the Scientific results of the voyage of H.M.S. *Challenger* during the years 1873–76. Zoology 24: i–xc, 1–918, 927–942, pls. 1–148.
- Beck, J.T. 1980. Life history relationships between the bopyrid isopod *Probopyrus pandalicola* and one of its freshwater shrimp hosts *Palaemonetes paludosus*. American Midland Naturalist 104: 135–154.
- Bonnier, J. 1900. Contribution à l'étude des épicarides. Les Bopyridae. Travaux de la Station Zoologique de Wimereux 8: 1–476, pls. 1–41.
- Bourdon, R. 1967a. Sur trois nouveaux Bopyridae du Sénégal. Bulletin de l'Institut Fondamental d'Afrique Noire, sér. A 29: 107–122.
- Bourdon, R. 1967b. Sur deux nouveaux épicarides (Isopoda) parasites de crustacés décapodes. Zoologische Mededelingen 42: 167–174.
- Bourdon, R. 1967c. Sur quelques nouvelles espèces de Cabiropsidae (Isopoda Epicaridea). Bulletin du Muséum National d'Histoire Naturelle (2nd série) 38: 846–868.
- Bourdon, R. 1968. Les Bopyridae des mers européennes. Mémoires du Muséum National d'Histoire Naturelle (Série A) Zoologie 50: 77–424.
- Bourdon, R. 1972a. Epicaridea de Java, Ile Maurice et Afrique du Sud (Crustacea, Isopoda). Steenstrupia 2:105–119.
- Bourdon, R. 1972b. Sur quelques Bopyridae (Crustacea, Isopoda) parasites de galathéides. Bulletin du Muséum National d'Histoire Naturelle (3e sér.) 66: 817–838.
- Bourdon, R. 1979a. Épicarides de Madagascar. II. Bulletin du Muséum National d'Histoire Naturelle (4e sér.) 1 (A): 471–506.

- Bourdon, R. 1979b. Sur la taxonomie et l'éthologie de quelques orbioninés (Isopoda Epicaridea). Internationale Revue der Gesammte Hydrobiologie 64: 425–435.
- Bourdon, R. 1979c. Bopyridae de la campagne Biaçores (Isopoda Epicaridea). Bulletin du Muséum National d'Histoire Naturelle 4e sér. 1 (A): 507–512.
- Bourdon, R. 1980a. Les espèces du genre *Bopyrella* J. Bonnier (Crustacea, Isopoda, Bopyridae). Bulletin du Muséum National d'Histoire Naturelle (4e sér.) 2 (A): 185–236.
- Bourdon, R. 1980b. *Aporobopyrus dollfusi* n. sp. (Crustacea, Epicaridea, Bopyridae) parasite de porcellanes de la Mer Rouge. Bulletin du Muséum National d'Histoire Naturelle (4e sér.) 2 (A): 237–244.
- Bourdon, R. 1981. Crustacés isopodes. I. Bopyridae parasites des Pénéides. Résultats des Campagnes MUSORSTOM. I. Philippines (18–28 Mars 1976), vol. 1 (10). Mémoires d'ORSTOM 91: 237–260.
- Bourdon, R. 1982. Sur deux nouveaux isopodes épicarides du genre *Bopyrella* Bonnier. Internationale Revue der Gesammte Hydrobiologie 67: 749–756.
- Bourdon, R. 1983. Expédition Rumphius II (1975). Crustacés parasites, commensaux, etc. (Th. Monod éd.). VIII. Crustacés isopodes (3e partie: épicarides Bopyridae). Bulletin du Muséum National d'Histoire Naturelle (4e sér.) 5 (A): 845–869.
- Bourdon, R. 1987. Suborder Epicaridea. In E.N. Kozloff (editor), Marine invertebrates of the Pacific Northwest: 341–344. Seattle: University of Washington Press.
- Bourdon, R., and C.B. Boyko. 2005. Redescription of *Bopyrophryxus branchiabdominalis* Codreanu, 1965 (Crustacea: Isopoda: Bopyridae) with a reappraisal of the subfamily Bopyrophryxinae Codreau, 1965. Proceedings of the Biological Society of Washington 118: 108–116.
- Bourdon, R., and A.J. Bruce. 1979. *Bopyrella saronae* sp. nov., a new bopyrid parasite (Isopoda) of the shrimp *Saron marmoratus* (Olivier). Crustaceana 37: 191–197.
- Bourdon, R., and A.J. Bruce. 1983. Six bopyrid shrimp parasites (Isopoda, Epicaridea) new to the Australian fauna. Crustaceana 45: 96–106.
- Bourdon, R. and J.C. Markham. 1980. A new genus and species of bopyrid isopod infesting alpheid shrimps of the genus *Synalpheus* in the western Atlantic Ocean. Zoologische Mededelingen 55: 221–230.
- Bourdon, R., and J.H. Stock. 1979. On some Indo-West Pacific Bopyridae (Isopoda, Epicaridea) in the collections of the Zoölogisch Museum, Amsterdam. Beaufortia 28: 205–218.
- Bourdon, R., J.-L. d'Hondt, and A. Veillet. 1981. Note préliminaire sur les microsètes et les «fentes céphaliques» chez les bopyriens (crustacés épicarides). Bulletin de la Société Zoologique de France 105: 495–504.
- Bowman, T. E. 1956. Una especie nueva de *Bopyrella* (Crustacea: Isopoda) de los Roques, Venezuela. Novedades Científicas Contribuciones Occasionales del Museo de Historia Natural La Salle, Serie Zoológica 19: 1–4.
- Boyko, C.B. 2004. The Bopyridae (Crustacea: Isopoda) of Taiwan. Zoological Studies 43: 677–703.
- Boyko, C.B. 2006. A new shrimp host for *Parabopyrella lata* (Nierstrasz and Brender à Brandis, 1929) (Crustacea: Isopoda: Bopyridae) from the Florida gulf coast: a novel host/parasite relationship or a case of mistaken identity? Gulf of Mexico Science 24: 41–44.
- Boyko, C.B. 2014. Bopyrinae Rafinesque, 1815. In M. Schotte et al. (editors), World marine, freshwater and terrestrial isopod crustaceans database. World Register of Marine Species. Online resource (<http://www.marinespecies.org/aphia.php?p=taxdetails&id=589329> on 2014-04-07).
- Boyko, C.B., and J.D. Williams. 2010. A new genus and species of primitive bopyrid (Isopoda, Bopyridae) parasitizing hermit crabs (Anomura) from deep waters in the eastern Atlantic and Japan. In C.H.J. M. Fransen, S. De Grave, and P.K.L. Ng (editors), Studies on Malacostraca, Lipke Bijdeley Holthuis memorial vol. (Crustacean Monographs 14): 145–157. Leiden: Brill.
- Boyko, C.B., J. Moss, J.D. Williams, and J.D. Shields. 2013. A molecular phylogeny of Bopyroidea and Cryptoniscoidea (Crustacea: Isopoda). Systematics and Biodiversity 11: 495–506.
- Bruce, A.J. 1966. An interesting new phrytid (Epicaridea) parasitic on a pontoniid shrimp. Annals and Magazine of Natural History, 13th ser. 8: 385–390.
- Bruce, A.J. 1972a. *Orophryxus shiinoi* gen nov., sp. nov., an unusual phrytid (Crustacea, Isopoda, Bopyridae) parasitic upon a pontoniid shrimp from Zanzibar. Parasitology 64: 445–450.
- Bruce, A.J. 1972b. *Filophryxus dorsalis* gen. nov., sp. nov., an unusual bopyrid parasite from eastern Australia. Parasitology 65: 351–358.
- Bruce, A.J. 1968. A new species of *Hypophryxus* (Isopoda, Bopyridae) from an Indo-Pacific pontoniid shrimp. Crustaceana 14: 13–20.
- Bruce, A.J. 1982. The pontoniine shrimp fauna of Hong Kong. In B. Morton and C.K. Tseng (editors), Proceedings of the First International Marine Biological Workshop: the marine flora and fauna of Hong Kong and southern China, Hong Kong, 1980: 233–284. Hong Kong: Hong Kong University Press.
- Bruce, A.J. 1990. Redescriptions of five Hong Kong carideans first described by William Stimpson, 1860. In B. Morton (editor), Proceedings of the Second International Marine Biological Workshop: the marine flora and fauna of Hong Kong and Southern

- China, Hong Kong, 1986: 569–610. Hong Kong: Hong Kong University Press.
- Brusca, R.C., and G.D.F. Wilson. 1991. A phylogenetic analysis of the Isopoda with some classificatory recommendations. *Memoirs of the Queensland Museum* 31: 143–204.
- Brusca, R.C., V.R. Coelho, and S. Taiti. 2001. A guide to the coastal isopods of California. Online resource (http://tolweb.org/notes/?note_id=3004).
- Calman, W.T. 1898. On a collection of Crustacea from Puget Sound. *Annals of the New York Academy of Sciences* 11: 259–292, pls. 31–34.
- Campos, E., and A.R. De Campos. 1989. Epicarideos de Baja California: distribución y notas ecológicas de *Probopyrus pandalicola* (Packard, 1879) en el Pacífico oriental. *Revista de Biología Tropical* 37: 29–36.
- Caroli, E. 1930. Notizia di tre specie nuove ed una poco nota di bopiridi addominali, parassiti di caridei del golfo di Napoli (contributo alla conoscenza del genere *Phrixus* Rathke). *Bollettino della Società Naturalisti in Napoli* 41: 258–269, pl. 9.
- Caroli, E. 1949. Breve rassegna dei bopiridi addominali parassiti di caridei. *Pubblicazioni della Stazione Zoologica di Napoli* 21: 232–235.
- Cericola, M.J., and J.D. Williams. 2015. Prevalence, reproduction and morphology of the parasitic isopod *Athelges takanoshimensis* Ishii, 1914 (Isopoda: Bopyridae) from Hong Kong hermit crabs. *Marine Biology Research* 11: 236–252.
- Chace, F.A., Jr. 1984. The caridean shrimps (Crustacea: Decapoda) of the Albatross Philippine Expedition, 1907–1910, part 2: families Glyphocrangonidae and Crangonidae. *Smithsonian Contributions to Zoology* 397: i–iv, 1–63.
- Chaplin-Ebanks, S.A., and M.C. Curran. 2007. Prevalence of the bopyrid isopod *Probopyrus pandalicola* in the grass shrimp, *Palaemonetes pugio*, in four tidal creeks on the South Carolina–Georgia coast. *Journal of Parasitology* 93: 73–77.
- Chopra, B. 1922. Preliminary note on Isopoda of the family Bopyridae parasitic on Indian Decapoda Macrura. *Journal and Proceedings of the Asiatic Society of Bengal*, New Series 18: 69–71.
- Chopra, B. 1923. Bopyrid isopods parasitic on Indian Decapoda Macrura. *Records of the Indian Museum* 25: 411–550, pls. 11–21.
- Chopra, B. 1927. Bopyrid isopods. *Bulletin of the Madras Government Museum*, New Series, Natural History Section 1: 119–122.
- Chopra, B. 1930. Further notes on bopyrid isopods parasitic on Indian Decapoda Macrura. *Records of the Indian Museum* 32: 113–147, pls. 4–6.
- Codreanu, R. 1961. Crustacei paraziti cu afinitati indo-pacifice în Marea Neagră. *Hidrobiologia* 3: 133–146. [in Romanian]
- Codreanu, R. 1967. Clasificarea evolutiva a bopirienilor, isopode parazite ale crustaceelor decapode si importanta lor biologica generala. *Studii si Cercetari de Biologie Seria Zoologie* 19: 203–211.
- Codreanu, M., and R. Codreanu. 1956. Sur l'*Anisarthrus pelseneeri*, épicaride parasite abdominal de la crevette *Athanas nitescens*; sa présence dans la Mer Noire et la dispersion du genre *Anisarthrus*. *Bulletin Biologique de la France et de la Belgique* 90: 111–121.
- Cornalia, E., and P. Panceri. 1861. Osservazioni zoologiche ed anatomiche sopra un nuovo genere di isopodi sedentarii (*Gyge branchialis*). *Memorie delle Reale Accademia delle Scienze di Torino* (ser. 2) 19: 85–118, pls. 1–2.
- Coutière, H. 1902. Sur un nouveau de rhizocéphale, parasite des Alpheidae. *Comptes Rendus Hebdomadiers des Séances de l'Académie de Sciences* 134: 913–915.
- Dana J.D. 1852. Crustacea, part I. *United States Exploring Expedition during the years 1838–42 under the command of Ch. Wilkes U.S.N.* 13: i–viii + 1–685. Philadelphia: C. Sherman.
- Dana, J.D. 1855. *Atlas Crustacea. United States Exploring Expedition, during the years 1838, 1839, 1840, 1841, 1842. Under the command of Charles Wilkes, U.S.N.*. Philadelphia: C. Sherman.
- Danforth, C.G. 1963. Bopyridian (Crustacea, Isopoda) parasites found in the eastern Pacific of the United States. Ph.D. dissertation, Oregon State University, Corvallis.
- Danforth, C.G. 1970. Epicaridea (Crustacea: Isopoda) of North America. Ann Arbor, MI: University Microfilms, ii + 1–191, pls. 1–48.
- Danforth, C.G. 1971a. Two bopyrids (Isopoda) from New Guinea. *Bulletin of the Southern California Academy of Sciences* 70: 99–102.
- Danforth, C.G. 1971b. New bopyrids (Isopoda) from the Indian and Pacific Oceans. *Micronesica* 7: 163–177.
- De Grave, S., and C.H.J.M. Fransen. 2011. Cari-deorum catalogus: the Recent species of the dendrobranchiate, stenopodidean, procarididean and caridean shrimps (Crustacea: Decapoda). *Zoologische Mededelingen* 85: 195–589.
- de Man, J.G. 1913. The Decapoda of the Siboga Expedition. Part II. Family Alpheidae. *Siboga Expeditie Monographie* 39al: 113–465.
- Duan J., J. An, and H. Yu. 2008. A new species and two new record species of genus *Allokepon* Markham, 1982 (Isopoda: Epicaridea: Bopyridae) from China. *Zootaxa* 1682: 62–68.
- Dumbauld, B., J. Chapman, M. Torchin, and A. Kuris. 2011. Is the collapse of mud shrimp (*Upogebia pugettensis*) populations along the Pacific Coast of North America caused by outbreaks of a previously unknown bopyrid isopod parasite (*Orthione griffensis*)? *Estuaries and Coasts* 34: 336–350.
- Espinosa-Pérez, M.D.C., and M.E. Hendrickx. 2001. Checklist of isopods (Crustacea: Peracarida: Isopoda) from the eastern tropical Pacific. *Belgian Journal of Zoology* 131: 43–55.

- Espinosa-Pérez, M.D.C., and M.E. Hendrickx. 2006. A comparative analysis of biodiversity and distribution of shallow-water marine isopods (Crustacea: Isopoda) from polar and temperate waters in the East Pacific. *Belgian Journal of Zoology* 136: 219–247.
- Espinosa-Pérez, M.D. C., M.E. Hendrickx, and J.J. Morrone. 2009. Identification of generalized tracks for the species of Isopoda (Peracarida) from the eastern Pacific. *Journal of Crustacean Biology* 29: 224–231.
- Faxon, W. 1895. Reports on an exploration off the west coasts of Mexico, Central and South America, and off the Galapagos Islands, in charge of Alexander Agassiz, by the U.S. Fish Commission Steamer "Albatross" during 1891, Lieut.-Commander Z.L. Tanner, U.S.N., commanding. XV. The stalk-eyed Crustacea. *Memoirs of the Museum of Comparative Zoölogy at Harvard College* 18: 1–292, pls. A–K, 1–56, 1 map.
- Fee, A.R. 1927. The Isopoda of Departure Bay and vicinity, with descriptions of new species, variations and colour notes. *Contributions to Canadian Biology and Fisheries (New Series)* 3: 13–46, 1 pl.
- Feeney, T.D. 1995. Physical controls on the distribution of contaminants on Sturgeon Bank, Fraser River Delta, British Columbia. Master of Science thesis, University of British Columbia, Vancouver.
- Fraser, C.M. 1932. A comparison of the marine fauna of the Nanaimo region with that of the San Juan Archipelago. *Transactions of the Royal Society of Canada (3rd series)* 26: 49–70.
- George, M.J., and J.-O. Strömberg. 1968. Some new species and new records of marine isopods from San Juan Archipelago, Washington, U.S.A. *Crustaceana* 14: 225–254.
- George, P.C. 1947. *Megacepon choprai*, gen. et sp. nov., a bopyrid isopod from the gill chamber of *Sesarma tetragonum* (Fabr.). *Records of the Indian Museum* 44: 385–390.
- Gerstaecker, A. 1901. Die Klassen und Ordnungen der Arthropoden wissenschaftlich dargestellt in Wort und Bild. Band 5, Abtheilung 2. Crustacea (Zweite Hälfte: Malacostraca). Leipzig: C.F. Winter'sche Verlagshandlung, i–viii + 1–1319, pls. 1–128.
- Giard, A. 1907. Sur l'*Anisarthrus pelseneeri* (nov. gen. et nov. sp.) bopyrien parasite d'*Athanas nitescens* Leach et sur la synonymie du genre *Hemiarthrus*. *Comptes Rendus Hebdomadaires des Séances et Mémoires de la Société de Biologie* 63: 321–324.
- Giard, A., and J. Bonnier. 1887. Contributions à l'étude des bopyriens. Travaux de l'Institut Zoologique de Lille et du Laboratoire de Zoologie Maritime de Wimereux 5: 1–272, pls. 1–10.
- Giard, A., and J. Bonnier. 1888. Sur quelques espèces nouvelles de céponiens. *Comptes Rendus Hebdomadaires des Séances de l'Académie de Sciences* 107: 44–47.
- Giard, A., and J. Bonnier. 1890. Prodrome d'une monographie des épicarides du Golfe de Naples. *Bulletin Scientifique de la France et de la Belgique* 22: 367–391.
- Gifford, J. 1934. The life history of *Argeia pauperata* From *Crago franciscorum*. Master's thesis, Stanford University, Stanford, 21 pp. 2 tables, 1 chart, 5 pls.
- Gurjanova, E., 1936a. Faune de l'URSS. Crustaces. Vol. 7, no. 3, isopodes des mers orientales. Institut Zoologique de l'Academie des Sciences de l'URSS 6, vii + 278 pp.; figs. 1–150. [in Russian with German summary]
- Gurjanova, E. 1936b. Beiträge zur Kenntnis der Isopodenfauna des Pazifischen Ozeans. IV. Neue Isopodenarten aus dem Japanischen und Beringmeer. *Zoologischer Anzeiger* 114: 250–265.
- Han, Q., and X. Li. 2007. An account of the *Glyphocrangon* shrimps (Decapoda, Caridea) in the collections of the Institute of Oceanology, Chinese Academy of Sciences. *Crustaceana* 80: 545–553.
- Han, Q.-X., and X.-Z. Li. 2010. Records of the Cranionidae (Crustacea, Decapoda, Caridea) species from the Yellow Sea and Bohai Sea. *Acta Zootaxonomica Sinica* 35: 227–239.
- Hansen, H.J. 1897. Reports on the dredging operations off the west coast of Central America to the Galapagos, to the west coast of Mexico, and in the Gulf of California, in charge of Alexander Agassiz, carried on by the U.S. Fish Commission Steamer "Albatross," during 1891, Lieut. Commander Z.L. Tanner, U.S.N., commanding. XXII. The Isopoda. *Bulletin of the Museum of Comparative Zoölogy at Harvard College* 31: 95–129, pls. 1–6, 1 map.
- Hatch, M.H. 1947. The Chelifera and Isopoda of Washington and adjacent regions. *University of Washington Publications in Biology* 10: 155–274.
- Hay, W. P. 1917. A new genus and three new species of parasitic isopod crustaceans. *Proceedings of the United States National Museum* 51: 569–574, pls. 98–100.
- Hesse, C. E. 1865. Recherches sur les crustacés rares ou nouveaux des côtes de France (complément du troisième article). *Annales des Sciences Naturelles, Zoologie*, sér. 5 3: 221–242, pl. 4.
- Hiraiwa, Y.K. 1933. Studies on a bopyrid, *Epipenaeon japonica* Thielemann. I. Morphological studies in both sexes. *Journal of Science of the Hiroshima University (Ser. B.) Div. 1 (Zoology)* 2 (4): 49–70, pl. 1.
- Høeg, J.T., and A.V. Rybakov. 1992. Revision of the Rhizocephala Akentrogonida (Cirripedia), with a list of all the species and a key to the identification of families. *Journal of Crustacean Biology* 12: 600–609.
- Holthuis, L.B. 1971. Biological results of the University of Miami Deep-Sea Expeditions. 75. The Atlantic shrimps of the deep-sea genus *Glyphocrangon* A. Milne Edwards, 1881. *Bulletin of Marine Science* 21: 267–373.

- International Commission on Zoological Nomenclature (ICZN). 1999. International code of zoological nomenclature. 4th ed. London: International Trust for Zoological Nomenclature.
- Ishii, S. 1914. On a new epicaridean isopod (*Athelges takanoshimensis* sp. nov.) from *Eupagurus samuelis* Stimp. *Annotationes Zoologicae Japonenses* 8: 519–530, pl. 7.
- Jay, C.V. 1985. Life history relationships between the parasitic isopod *Argeia pugettensis* (Epicaridea: Bopyridae) and its host shrimp *Crangon franciscorum* (Caridea: Crangonidae). M.A. thesis, Humboldt State University, Arcata, California.
- Jay, C.V. 1989. Prevalence, size and fecundity of the parasitic isopod *Argeia pugettensis* on its host shrimp *Crangon francisorum* (sic). *American Midland Naturalist* 121: 68–77.
- Kannupandi, T. 1976. A study on the cuticle of *Stegoalpheon kempfi* Chopra (Isopoda: Crustacea) in relation to its parasitic mode of life. *Journal of Experimental Marine Biology and Ecology* 25: 87–94.
- Kaufmann, R.S., W.W. Wakefield, and A. Genin. 1989. Distribution of epibenthic mega fauna and lebensspuren on two central North Pacific seamounts. *Deep Sea Research* 36: 1863–1896.
- Kazmi, Q. B., and R. Bourdon. 1997. A new bopyrid isopod *Progebiophilus assisi* on the mud shrimp *Upogebia (U.) assisi* Barnard (Thalassinoidea) from Pakistan. *Pakistan Journal of Marine Sciences* 61/62: 59–67.
- Kazmi, Q. B., M. Schotte, and F. Yousuf. 2002. An illustrated key to the Malacostraca (Crustacea) of the northern Arabian Sea, part V: Isopoda. *Pakistan Journal of Marine Sciences* 2: 47–116.
- Kensley, B. 1968. *Pseudione elongata africana*, a new subspecies of bopyrid isopod from the west coast of the Cape Peninsula, South Africa. *Crustaceana* 15: 188–192.
- Kensley, B. 1974. Bopyrid Isopoda from southern Africa. *Crustaceana* 26: 259–266.
- Kensley, B. 2001. Biogeography of the marine Isopoda of the Indian Ocean, with a check-list of species and records. In B. Kensley and R.C. Brusca (editors), *Crustacean Issues* 13. Isopod systematics and evolution: 205–264. Rotterdam: Balkema.
- Kensley, B., and T.-Y. Chan. 2001. Two species of deep-sea flabelliferan isopods from Taiwan (Crustacea: Peracarida: Aegidae, Anuropidae). *Journal of Natural History* 35: 481–496.
- Kim, H.S., and D.-H. Kwon. 1988. Bopyrid isopods parasitic on decapod crustaceans in Korea. *Korean Journal of Systematic Zoology Special Issue* 2: 199–221.
- Kossmann, R. 1881. Studien über Bopyriden. *Zeitschrift für Wissenschaftliche Zoologie* 35: 652–665, pls. 32–35.
- Kozloff, E.N. 1974. Keys to the marine invertebrates of Puget Sound, the Strait of Georgia, and the San Juan Archipelago, and adjacent regions. Seattle: University of Washington Press, x + 226 pp.
- Kröyer, H. 1838a. Grönlands Amfipoder. Det Kongelige Danske Videnskabernes Selskabs Naturvidenskabelige og Mathematiske Afhandlinger, series 4 (7): 229–326, pls. 1–4.
- Kröyer, H. 1838b. Grönlands Amfipoder. Copenhagen: Bianco Lunos. [Separately paginated reprinting of Kröyer, 1838a.]
- Kruczynski, W.L., and R.J. Menzies. 1977. Taxonomic status of *Synsynella* Hay and *Bopyro* Pearse (Isopoda: Bopyridae). *Proceedings of the Biological Society of Washington* 89: 551–558.
- Latrelle, P. A. 1802. *Histoire Naturelle Generale et Particulière, des Crustacés et des Insectes*. F. Dufart, Paris.
- Leija-Tristán, A., and S.I. Salazar-Vallejo. 1991. Parasitismo de *Progebiophilus bruscae* (Isopoda: Bopyridae) sobre el camarón *Upogebia dawsoni* (Thalassinoidea: Upogebiidae), en Baja California Sur, México. *Revista de Biología Tropical* 39: 1–5.
- Lemos de Castro, A. 1970. Crustáceos isópodos epicarídeos do Brasil. V. Duas espécies novas de hiperparasitas pertencentes ao gênero *Cabirops* Kossmann (Isopoda, Cabiropsidae). *Boletim do Museu Nacional (Nova Série) Zoologia* 277: 1–7, pls. 1–4.
- Li, L. 2003. Hong Kong's isopods. In B. Morton (editor), *Perspectives on marine environmental change in Hong Kong and southern China, 1977–2001. Proceedings of an International Workshop Reunion Conference, Hong Kong 21–26 October 2001*: 137–166. Hong Kong: Hong Kong University Press.
- Li, X., and T.-Y. Chan. 2014. Pandalid shrimps (Crustacea, Decapoda, Caridea) collected from the Philippines PANGLAO 2005 deep-sea expedition. *Mémoires du Muséum National d'Histoire Naturelle* 204: 129–154.
- Liu, R.E. (editor). 2008. *Checklist of marine biota of China seas*. Beijing: Science Press, 1267 pp.
- MacGinitie, G.E., and N. MacGinitie. 1949. *Natural history of marine animals*. New York: McGraw-Hill, xii + 473 pp.; 282 figs.
- MacGinitie, G.E., and N. MacGinitie. 1968. *Natural history of marine animals*. 2nd ed. New York: McGraw-Hill, xii + 523 pp.; 286 figs.
- Marin Jarrin, J.R. 2007. The ecology of surf zone fauna of dissipative sandy beaches in southern Oregon, U.S.A. Master of Science thesis, University of Oregon, Eugene.
- Marin Jarrin, J.R., and A.L. Shanks. 2008. Ecology of a population of *Lissocrangon stylirostris* (Caridea: Crangonidae), with notes on the occurrence and biology of its parasite, *Argeia pugettensis* (Isopoda: Bopyridae). *Journal of Crustacean Biology* 28: 613–621.
- Marin Jarrin, J.R., and A.L. Shanks. 2011. Spatio-temporal dynamics of the surf-zone faunal assemblages at a southern Oregon sandy beach. *Marine Ecology* 32: 232–242.

- Markham, J.C. 1972. Two new genera of western Atlantic abdominally parasitizing Bopyridae (Isopoda, Epicaridea), with a proposed new name for their subfamily. *Crustaceana Supplement* 3: 39–56.
- Markham, J.C. 1975. New records of two species of parasitic isopods of the bopyrid subfamily Ioninae in the western Atlantic. *Crustaceana* 29: 55–67.
- Markham, J.C. 1977. Description of a new western Atlantic species of *Argeia* Dana with a proposed new subfamily for this and related genera (Crustacea, Isopoda, Bopyridae). *Zoologische Mededelingen* 52: 107–123.
- Markham, J.C. 1979. A new genus and species of bopyrid isopod infesting the crangonid shrimp *Pontophilus abyssi* Smith in deep water of the northwestern Atlantic Ocean. *Proceedings of the Biological Society of Washington* 92: 769–772.
- Markham, J.C. 1980. A new species of *Dicropleon* and new records of three other species of bopyrid isopods from the Indian Ocean. *Bulletin of Marine Science* 30: 623–633.
- Markham, J.C. 1982. Bopyrid isopods parasitic on decapod crustaceans in Hong Kong and Southern China. In B. Morton and C.K. Tseng (editors), *Proceedings of the First International Marine Biological Workshop: the marine flora and fauna of Hong Kong and southern China*, Hong Kong, 1980: 325–391. Hong Kong: Hong Kong University Press.
- Markham, J.C. 1985a. A review of the bopyrid isopods infesting caridean shrimps in the northwestern Atlantic Ocean, with special reference to those collected during the Hourglass Cruises in the Gulf of Mexico. *Memoirs of the Hourglass Cruises* 8: 1–156.
- Markham, J.C. 1985b. Additions to the bopyrid isopod fauna of Thailand. *Zoologische Verhandelingen* 224: 1–63.
- Markham, J.C. 1986. Evolution and zoogeography of the Isopoda Bopyridae, parasites of Crustacea Decapoda. In R.H. Gore and K.L. Heck (editors), *Crustacean Issues 4. Crustacean biogeography*: 143–164. Rotterdam: A.A. Balkema.
- Markham, J.C. 1988. Descriptions and revisions of some species of Isopoda Bopyridae of the north western Atlantic Ocean. *Zoologische Verhandelingen* 246: 1–63.
- Markham, J.C. 1989. Three species of Isopoda Bopyridae new to the fauna of the Philippines. The Beagle, Records of the Northern Territory Museum of Arts and Sciences 6: 141–148.
- Markham, J.C. 1990. Further notes on the Isopoda Bopyridae of Hong Kong. In B. Morton (editor), *Proceedings of the Second International Marine Biological Workshop: the marine flora and fauna of Hong Kong and Southern China*, Hong Kong, 1986: 555–568. Hong Kong: Hong Kong University Press.
- Markham, J.C. 1991. Redescriptions and new records of Isopoda Bopyridae (Crustacea) from Thailand. *Raffles Bulletin of Zoology* 39: 289–297.
- Markham, J.C. 1992a. Second list of additions to the Isopoda Bopyridae of Hong Kong. In B. Morton (editor), *The marine flora and fauna of Hong Kong and southern China III. Proceedings of the Fourth International Marine Biological Workshop: the marine flora and fauna of Hong Kong and southern China*, Hong Kong, 11–29 April 1989: 277–302. Hong Kong: Hong Kong University Press.
- Markham, J.C. 1992b. The Isopoda Bopyridae of the eastern Pacific – missing or just hiding? *Proceedings of the San Diego Society of Natural History* 17: 1–4.
- Markham, J.C. 1995. Two new species of Isopoda Bopyridae (Crustacea) infesting thalassinideans in the western Pacific. *Raffles Bulletin of Zoology* 43: 83–89.
- Markham, J.C. 2004. New species and records of Bopyridae (Crustacea: Isopoda) infesting species of the genus *Upogebia* (Crustacea: Decapoda: Upogebiidae): the genera *Orthione* Markham, 1988, and *Gyge* Cornalia and Panceri, 1861. *Proceedings of the Biological Society of Washington* 117: 186–198.
- Markham, J.C. 2010. The isopod parasites (Crustacea: Isopoda: Bopyridae) of decapod Crustacea of Queensland, Australia, with descriptions of three new species. In P.J.F. Davie and J.A. Phillips (editors), *Proceedings of the 13th International Marine Biological Workshop, Marine Fauna and Flora of Moreton Bay, Queensland. Memoirs of the Queensland Museum—Nature* 54: 151–197.
- Markham, J.C., and C.B. Boyko. 2003. A new species of *Albunione* Markham & Boyko, 1999 (Crustacea: Isopoda: Bopyridae: Pseudioninae) from Taiwan. *American Museum Novitates* 3410: 1–7.
- Martin, J.W., and G.E. Davis. 2001. An updated classification of the recent Crustacea. *Natural History Museum of Los Angeles County Science Series* 39: 1–124.
- Menzies, R.J., and M.A. Miller. 1954. Key to the Chelifera and the suborders of the Isopoda. In S.F. Light (editor), *Intertidal invertebrates of the central California Coast*: 138–159. Berkeley: University of California Press.
- Miller, M.A. 1975. Phylum Arthropoda: Crustacea, Tanaidacea and Isopoda. In R.I. Smith and J.T. Carlton (editors), *Light's manual: intertidal invertebrates of the central California coast*: 277–312. Berkeley: University of California Press, xviii + 716 pp.
- Miya, Y. 1972. The Alpheidae (Crustacea, Decapoda) of Japan and its adjacent waters. Part I. Publications of the Amakusa Marine Biological Laboratory 3: 23–101.
- Monod, T. 1933. Mission Robert-Ph. Dollfus en Égypte. Tanaidacea et Isopoda. Mémoires Présentés à l'Institut d'Égypte 21: 161–264.
- Monod, T. 1976. Sur une nouvelle collection de Crustacés Decapodes de Nouméa (Nouvelle Calédonie). *Cahiers du Pacifique* 19: 133–152.
- Moore, J.A., et al. 2003. Biodiversity of Bear Seamount, New England Seamount Chain: results of exploratory trawling. *Journal of Northwestern Atlantic Fishery Science* 31: 363–372.

- Morton, B. 2003. Hong Kong's international malacological wetland and marine biological workshops (1977–1998): changing local attitudes towards marine conservation. In B. Morton (editor), Perspectives on marine environmental change in Hong Kong and southern China, 1977–2001. Proceedings of an International Workshop Reunion Conference, Hong Kong 21–26 October 2001: 31–71. Hong Kong: Hong Kong University Press.
- Nakashima, Y. 1995. Can small male shrimps achieve copulation in the presence of larger ones? *Journal of Ethology* 13: 9–16.
- Nelson, S.G., M.A. Simmons, and A.W. Knight. 1986. The energy burden of the bopyrid parasite *Argeia pauperata* (Crustacea, Isopoda) on the grass shrimp *Crangon franciscorum* (Crustacea, Crangonidae). *Comparative Biochemistry and Physiology* 83A: 121–124.
- Nierstrasz, H.F., and G.A. Brender à Brandis. 1923. Die Isopoden der Siboga-Expedition. II. Isopoda Genuina. I. Epicaridea. *Siboga-Expeditie* 32b: 57–121, pls. 4–9.
- Nierstrasz, H.F., and G.A. Brender à Brandis. 1925. Bijdragen tot de kennis der fauna van Curaçao. Resultaten eener reis van Dr. C.J. van der Horst in 1920. Epicaridea. *Bijdragen tot de Dierkunde* 24: 1–8, pl. 1.
- Nierstrasz, H.F., and G.A. Brender à Brandis. 1929. Papers from Dr. Th. Mortensen's Pacific Expedition 1914–16. XLVIII. Epicaridea. I. Videnskabelige Meddelelser fra den Dansk Naturhistoriske Forening i København 87: 1–44.
- Nierstrasz, H.F., and G.A. Brender à Brandis. 1930. Résultats scientifiques du voyage aux Indes orientales néerlandaises de LL. AA. RR. le Prince et la Princesse Léopold de Belgique. Isopoda Epicaridea. Mémoires du Musée Royal d'Histoire Naturelle de Belgique, Hors Série 3: 12–17.
- Nierstrasz, H.F., and G.A. Brender à Brandis. 1931. Papers from Dr. Th. Mortensen's Pacific Expedition 1914–16. LVII. Epicaridea II. Videnskabelige Meddelelser fra den Dansk Naturhistoriske Forening i København 91: 147–226, pl. 1.
- Nierstrasz, H.F., and G.A. Brender à Brandis. 1932. Alte und neue Epicaridea. *Zoologischer Anzeiger* 101: 90–100.
- Nobili, G. 1905. Decapodi e isopodi della Nuova Guinea Tedesca raccolti dal Sign. L. Biró. *Annales Historico-Naturales Musei Nationalis Hungarici* 3: 480–507, pls. 12–13.
- Nobili, G. 1906. Nuovi bopiridi. Atti della Real Accademia della Scienze di Torino 41: 1098–1113, pl. 1.
- Norman, A. M. 1905. Museum normanianum, or a catalogue of the Invertebrata of the Arctic and North Atlantic temperate ocean and Palaearctic Region, which are contained in the collection of the Rev. Canon A. M. Norman, M. A., D. C. L., LL. D., F. R. S., F. L. S., &c. III. Crustacea Second Edition. Durham: Thos. Cadocleugh & Son.
- Oguro, C. 1961. On the neurosecretory system of two parasitic isopods, *Argeia pugettensis* Dana and *Athelges japonicus* Shiino. *Annotationes Zoologicae Japonenses* 34: 43–48.
- Page, R.D.M. 1985. Review of the New Zealand Bopyridae (Crustacea: Isopoda: Epicaridea). *New Zealand Journal of Zoology* 12: 185–212.
- Pearse, A.S. 1930. Parasites of Fukien crabs. *Proceedings of the Natural History Society of Fukien Christian University* 3: 10–18.
- Pearse, A.S. 1953. Parasitic crustaceans from Alligator Harbor, Florida. *Quarterly Journal of the Florida Academy of Sciences* 15: 187–243.
- Penha-Lopes, G., J.F. Marques, M.C. Leal, A.F. Carvalho, and J. Paula. 2013. Population structure and reproduction of *Pseudione elongata africana* (Bopyridae, Isopoda). *Western Indian Ocean Journal of Marine Science* 11: 27–39.
- Pillai, N.K. 1954. A preliminary note on the Tanaidacea and Isopoda of Travancore. *Bulletin of the Central Research Institute, University of Travancore* (series C, Natural Sciences) 3: 1–21.
- Pillai, N.K. 1966. Isopod parasites of south Indian crustaceans. *Crustaceana* 10: 183–191.
- Poore, G.C.B., J.C. Markham, and H.M. Lew Tan 2002. Superfamily: Bopyroidea. In W.W.K. Houston and P.L. Beesley (editors), *Zoological catalogue of Australia*, vol. 19.2A. Crustacea: Malacostraca: Syncarida, Peracarida: Isopoda, Tanaidacea, Mictacea, Thermosbaenacea, Spelaeogriphacea: 111–137. Melbourne: CSIRO Publishing.
- Qazi, M.H. 1959. Some bopyrid isopods of West Pakistan. *Scientist (Scientific Society of Pakistan)* 3: 55–62, 2 unnumbered pls.
- Rafinesque, C. S. 1815. Analyse de la nature ou tableau de l'univers et des corps organisés. Palerme.
- Rao, K.S.P.B., and T.N.C. Ramaprasad. 1964. A note on *Stegoalpheon kempfi* Chopra. *Current Science* 33: 588–589.
- Restivo, F. 1971. Su di una nuova specie di *Cabiroops* parassita di *Pseudione*. *Pubblicazioni della Stazione Zoologica di Napoli* 39: 70–86.
- Restivo, F. 1975. Nuovi dati su *Paracabiroops* (n.d. *Cabiroops*) *marsupialis* Caroli, parassita di *Gyge branchialis*. *Pubblicazione della Stazione Zoologica di Napoli* 39: 150–168.
- Richard, J. 1900. Essai sur les crustacés considérés dans leurs rapports avec l'hygiène, la médecine et la parasitologie. Lille: Bigot Frères.
- Richardson, H. 1899a. Key to the isopods of the Pacific coast of North America, with descriptions of 22 new species. *Proceedings of the United States National Museum* 21: 815–869.
- Richardson, H. 1899b. Key to the isopods of the Pacific coast of North America, with descriptions of 22 new

- species. Annals and Magazine of Natural History, ser. 7, 4: 321–338.
- Richardson, H. 1900. Synopses of North-American Invertebrates. VIII. The Isopoda. — Part II. Asel-lota, Oniscoidea, Epicaridea. American Naturalist 34: 295–309.
- Richardson, H. 1904a. Contributions to the natural history of the Isopoda. Proceedings of the United States National Museum 27: 1–89.
- Richardson, H. 1904b. A reply to certain criticisms of Prof. Alfred Giard respecting the bopyrids. Comptes Rendus Hebdomadaires des Séances et Mémoires de la Société de Biologie 56: 856–858.
- Richardson, H. 1905a. Isopods from the Alaska Salmon Investigation. Bulletin of the Bureau of Fisheries 24: 209–221.
- Richardson, H. 1905b. A monograph on the isopods of North America. Bulletin of the United States National Museum 54: i–lili, 1–727.
- Richardson, H. 1909. Isopods collected in the northwest Pacific by the U.S. Bureau of Fisheries Steamer “Albatross” in 1906. Proceedings of the United States National Museum 37: 75–129.
- Richardson, H. 1910. Marine isopods collected in the Philippines by the U.S. Fisheries steamer *Albatross* in 1907–8. Bureau of Fisheries Document 736: 1–44.
- Ricketts, E.F., and J. Calvin. 1939. Between Pacific tides. Stanford, CA: Stanford University Press, xxii + 320 pp.; 112 figs.
- Ricketts, E.F., and J. Calvin. 1948. Between Pacific tides. Revised ed. Stanford, CA: Stanford University Press, xxvii + 365 pp.; 129 figs; frontispiece.
- Ricketts, E.F., and J. Calvin. 1952. Between Pacific tides. 3rd ed. Stanford, CA: Stanford University Press, xiii + 501 pp.; 132 figs.; 46 plates; 2 unnumbered maps; 9 unnumbered plates.
- Ricketts, E.F., and J. Calvin, 1968. Between Pacific tides. 4th ed, revised by Joel W. Hedgpeth. Stanford, CA: Stanford University Press, xiv + 614 pp.
- Ricketts, E.F., J. Calvin and J.W. Hedgpeth. 1985. Between Pacific tides. 5th ed., revised by David W. Phillips. Stanford: Stanford University Press, vii + 652 pp.
- Risso, A. 1816. Histoire Naturelle des Crustacés des Environs de Nice. Librairie Grecque-Latine-Allemande, Paris.
- Román-Contreras, R. 2008. Chapter 5. Estudios y registros de isópodos epicarídeos en México: 1897–2005. In F. Álvarez Noguera and G.A. Rodríguez Almaraz (editors), Crustáceos de México: Estado Actual de su Conocimiento: 81–114. Monterrey, Mexico: Biblioteca Universitaria Raúl Rangel Frías.
- Román-Contreras, R., and L.A. Soto. 2002. A new deep-water genus and species of a branchial bopyrid infesting the galatheid crab *Munidopsis erinaceus* from the southwestern Gulf of Mexico. Journal of Crustacean Biology 27: 370–379.
- Romero-Rodríguez, J., and R. Román-Contreras. 2008. Aspects of the reproduction of *Bopyrinella thorii* (Richardson, 1904) (Isopoda, Bopyridae), a branchial parasite of *Thor floridanus* Kingsley, 1878 (Decapoda, Hippolytidae) in Bahía de la Ascensión, Mexican Caribbean. Crustaceana 81: 1201–1210.
- Romero-Rodríguez, J., and R. Román-Contreras. 2014. Relationships of the branchial parasite *Bopyrinella thorii* (Isopoda, Bopyridae) and its host *Thor floridanus* (Decapoda, Hippolytidae). Crustaceana 87: 463–475.
- Rudy, J., P., and L.H. Rudy. 1979. Oregon Estuarine invertebrates: an illustrated guide to the common and important invertebrate animals. Portland, OR: U.S. Fish and Wildlife Service.
- Rudy, P., and L. Rudy. 2013. Oregon estuarine invertebrates. 2nd ed. University of Oregon, Eugene. Available online (<https://library.uoregon.edu/scilib/oimb/oci>).
- Rybakov, A.V. 1990. *Bourdonia tridentata* gen. n., sp. n. (Isopoda: Cabriopsidae) a hyperparasite of *Bopyroides hippolytes* Kroyer [sic] from the shrimp *Pandalus borealis*. Parazitologija 24: 408–416.
- Rybakov, A.V., and V.V. Avdeev. 1991. Two *Bopyroides* species (Isopoda, Bopyridae) from the far-eastern shrimps. Parazitologija 25: 167–172.
- Saito, N., G. Itani, and N. Nunomura. 2000. A preliminary checklist of isopod crustaceans of Japan. Bulletin of the Toyama Science Museum 23: 11–107.
- Sadoglu, P. 1969. Variations in eye degeneration and pigment in some parasitic isopods during their life cycle. Pubblicazione della Stazione Zoologica di Napoli 37: 173–209.
- Salazar-Vallejo, S.I., and A. Leija-Tristán. 1990. *Progebiophilus bruscai* n. sp., a new bopyrid isopod parasitic on the mud shrimp, *Upogebia dawsoni* Williams (Thalassinoidea), from the Gulf of California. Cahiers de Biologie Marine 30: 423–432.
- Sars, G. O. 1883. Oversigt af Norges Crustaceer med foreløbige Bemaerkninger over de nye eller mindre bekjendte Arter. I. (Podophthalmata — Cumacea — Isopoda — Amphipoda). Forhandlinger i Videnskabs-Selskabet i Christiania for 1882: 1–124, pls. 1–6.
- Sassaman, C., G.A. Schultz, and R. Garthwaite. 1984. Host, synonymy, and parasitic incidence of *Bopyrella calmani* (Richardson) from central California (Isopoda: Epicaridea: Bopyridae). Proceedings of the Biological Society of Washington 97: 645–654.
- Savant, K.B. 1973. A note on the first record of isopod parasite *Bopyrella hogarti* [sic] (Chopra) on west coast of India. Current Science 42: 330–331.
- Schultz, G.A. 1969. How to know the marine isopod crustaceans. Dubuque, IA: Wm. C. Brown Company.
- Scott, T. 1902. Notes on Scottish Crustacea. Annals and Magazine of Natural History, ser. 7 10: 1–5, pl. 1.
- Seo, Y.S., C.-W. Oh, and A. Kim. 2014. Effect of the bopyrid isopod *Argeia puggettensis* on the growth and reproduction of the Kuro shrimp *Argis lar*.

- Korean Journal of Fisheries and Aquatic Science 47: 413–418.
- Sherman, M. B., and M.C. Curran. 2013. The effect of the bopyrid isopod *Probopyrus pandalicola* (Packard, 1879) (Isopoda, Bopyridae) on the survival time of the daggerblade grass shrimp *Palaemonetes pugio* Holthuis, 1949 (Decapoda, Palaemonidae) during starvation at two different temperatures. *Crustaceana* 86: 1328–1342.
- Shields, J.D. 2001. CD-ROM Review [of] Goater, Timothy M. 1999, *An Exploration of Marine Parasitic Crustacea*. *Journal of Crustacean Biology* 21: 313–314.
- Shiino, S.M. 1933. Bopyrids from Tanabe Bay. Memoirs of the College of Science, Kyoto Imperial University (ser. B) 8: 249–300.
- Shiino, S.M. 1934. Bopyrids from Tanabe Bay II. Memoirs of the College of Science, Kyoto Imperial University (ser. B) 9: 257–287.
- Shiino, S.M. 1936. Bopyrids from Tanabe Bay III. Memoirs of the College of Science, Kyoto Imperial University (ser. B) 11: 157–174.
- Shiino, S.M. 1937a. Some additions to the bopyrid fauna of Japan. *Annotationes Zoologicae Japonenses* 16: 293–300.
- Shiino, S.M. 1937b. Bopyrids from Tanabe Bay, IV. Memoirs of the College of Science, Kyoto Imperial University (ser. B) 12: 479–493.
- Shiino, S.M. 1939. Bopyrids from Kyūsyū and Ryūkyū. *Records of Oceanographic Works in Japan* 10: 79–99.
- Shiino, S.M. 1941. Further notes on bopyrids from Kyūsyū and Ryūkyū. *Annotationes Zoologicae Japonenses* 20: 154–158.
- Shiino, S.M. 1949a. On two species of the bopyrid genus, *Bopyrella*, found in Japan. *Bulletin of the Biogeographical Society of Japan* 14: 45–50.
- Shiino, S.M. 1949b. On two new genera of Bopyridae found in Japan. *Bulletin of the Biogeographical Society of Japan* 14: 57–63.
- Shiino, S.M. 1950. Notes on some new bopyrids from Japan. *Journal of Mie Medical College* 1: 151–167.
- Shiino, S.M. 1951. Some bopyrid parasites found on the decapod crustaceans from the waters along Mie Prefecture. *Report of Faculty of Fisheries, Prefectural University of Mie* 1: 26–40.
- Shiino, S.M. 1952. Phylogeny of the family Bopyridae. *Annual Report of the Prefectural University of Mie*, section 2, Natural Science 1: 33–56.
- Shiino, S.M. 1958. Note on the bopyrid fauna of Japan. *Report of the Faculty of Fisheries, Prefectural University of Mie* 3: 27–74, pl. 3.
- Shiino, S. M. 1964. On three bopyrid isopods from California. *Report of the Faculty of Fisheries, Prefectural University of Mie* 5: 19–25.
- Shiino, S.M. 1972. [The Epicaridea (list of species) from Japan]. *Kansai Shizenkagaku* 24: 7–10. [in Japanese]
- Sindermann, C.J. 1970. Principal diseases of marine fish and shellfish. New York: Academic Press, x + 369 pp.
- Sindermann, C.J. 1990. Principal diseases of marine fish and shellfish, 2nd ed. Vol. 2. London: Academic Press, xii + 516 pp.
- Sindermann, C.J., and A. Rosenfield. 1967. Principal diseases of commercially important marine bivalve Mollusca and Crustacea. United States Department of the Interior, Fish and Wildlife Service, Bureau of Commercial Fisheries, *Fishery Bulletin* 66: 335–385, figs. 1–9.
- Stebbing, T.R.R. 1893. A history of Crustacea, Recent Malacostraca. London: Kegan Paul, Trench, Trübner and Co.
- Stebbing, T.R.R. 1904. South African Crustacea. Part II. *Marine Investigations in South Africa* 2: 1–92, pls. 5–16.
- Stebbing, T.R.R. 1908. South African Crustacea. Part IV. *Annals of the South African Museum* 6: 1–96, pls. 28–40.
- Stebbing, T.R.R. 1910a. General catalogue of South African Crustacea (part 5 of S.A. Crustacea, for the Marine Investigations in South Africa). *Annals of the South African Museum* 6: 281–593, pls. 15–22.
- Stebbing, T.R.R. 1910b. No. VI.—Isopoda from the Indian Ocean and British East Africa. *Transactions of the Linnean Society of London*, 2nd ser., *Zoology* 14: 83–122, pls. 5–11.
- Stimpson, W. 1857. On the Crustacea and Echinodermata of the Pacific shores of North America. *Boston Journal of Natural History* 6: 444–532.
- Stimpson, W. 1864. Descriptions of new species of marine Invertebrata from Puget Sound, collected by the naturalists of the North-West Boundary Commission, A.H. Campbell, Esq., Commissioner. *Proceedings of the Academy of Natural Sciences of Philadelphia* 1864: 153–161.
- Stock, J.H. 1960. Notes on Epicaridea. *Crustaceana* 1: 28–33, pl. 1.
- Strathmann, M.F. 1987. Phylum or Subphylum Crustacea class Malacostraca order Decapoda, Caridea. In M.F. Strathmann (editor), *Reproduction and development of marine invertebrates of the northern Pacific Coast. Data and methods for the study of eggs, embryos, and larvae*: 432–440. Seattle: University of Washington Press.
- Strömberg, J.-O. 1971. Contribution to the embryology of bopyrid isopods with special reference to *Bopyroides*, *Hemiarthrus* and *Peseudione* (sic) (Isopoda, Epicaridea). *Sarsia* 47: 1–47.
- Thielemann, M. 1910. Beiträge zur Naturgeschichte Ostasiens. Herausgegeben von Dr. F. Doflein. Beiträge zur Kenntnis der Isopodenfauna Ostasiens. Abhandlungen der mathematisch-physischen Klasse der Königlichen Bayerischen Akademie von Wissenschaften Supplementarischer Band 2 (Abhandlung 3): 1–109, pls. 1–2.

- Trilles, J.-P. 1999. Ordre des isopodes sous-ordre des épicerides (Epicaridea Latreille, 1825). In J. Forest (editor), *Traité de Zoologie. Anatomie, Systématique, Biologie* (Pierre-P. Grassé). Tome VII, Fascicule III A, Crustacés Péracarides. Mémoires de l'Institut Oceanographique 19: 279–352.
- Wenner, E.L. 1978. Comparative biology of four species of glyphocrangonid and crangonid shrimp from the continental slope of the Middle Atlantic Bight. Canadian Journal of Zoology 56: 1052–1065.
- Wicksten, M.K. 1979. New records of the species of *Glypocrangon* in the northeastern Pacific Ocean (Caridea: Glyphocrangonidae). Proceedings of the Biological Society of Washington 92: 217–224.
- Wicksten, M.K. 2008. Decapod Crustacea of the Californian and Oregonian zoogeographic provinces. Online resource (<http://repositories.cdlib.org/sio/lib/26/>). [This is the original publication; this document has been revised several times as an e-document with changes in page numbers, e.g., p. 164 in the 2008 version is p. 168 in the 2009 version]
- Wicksten, M.K. 2012. Decapod Crustacea of the Californian and Oregonian zoogeographic provinces. Zootaxa 3371: 1–307.
- Williams J.D. and J. An. 2009. The cryptogenic parasitic isopod *Orthione griffenis* Markham, 2004 from the east and west Pacific. Integrative and Comparative Biology 49: 1–13.
- Williams, J.D., and C.B. Boyko. 2010. First description of the male of *Eragia profunda* Markham, 1994 (Crustacea: Isopoda: Bopyridae), with comments on relationships in the subfamily Argeiinae. Proceedings of the Biological Society of Washington 123: 149–152.
- Williams, J.D., and C.B. Boyko. 2012. The global diversity of parasitic isopods associated with crustacean hosts (Isopoda: Bopyroidea and Cryptoniscoidea). PLoS ONE 7: e35350.
- Yü, S.C. 1935. Notes on a bopyrid isopod parasitic on *Crago crago cassiope* de Man. Scientific Conference Nanning, Kwangsi Province, Aug. 11–15, 1935: 52–53.