


Supplemental Online Material for  
S. M. Kidwell and T. A. Rothfus, 2010. The living, the dead, and the expected dead: little  
bias of proportional abundances in bivalve death assemblages from variation in lifespan.  
*Paleobiology* 36(4)


Submitted April 27, 2010

This supplement consists of 7 supplemental figures and 2 supplemental tables.  
Supplemental Table 1 = information on datasets of living and dead abundances, gathered  
from other authors  
Supplemental Table 2 = raw published data on bivalve maximum lifespans (global  
compilation)


SUPPLEMENTAL FIGURE 1. Histogram (A) and cumulative percentage plot (B) of the frequency of species lifespans in the abundance database, in increments of one year. Calculations are based on all 413 living species occurrences in the abundance database, but only results for lifespans up to 30 years are displayed (comprise 97% of occurrences; maximum lifespan ranges up to 75 years). In A, the white bar at 9 years denotes species occurrences (33 total) that were assigned this lifespan as a default (no documented lifespan for that species or for any species in its genus or family). These occurrences are not included in the calculations for B.




SUPPLEMENTAL FIGURE 2. Additional simulations of expected death assemblages (exp DA) for the various hypothetical living assemblages (hypoth LA) in Text-figure 1, using an approximately 40-fold variation in lifespan (the set 2, 5, 8, 35, and 75 years), which reflects the minimum, median, and maximum lifespans in the dataset that shows the largest spread in lifespan among the 30 datasets analyzed (San Juan Channel shell gravel; see Supplemental Table 1). Larger values of  $|LB|$  are generated than when lifespans vary by 10-fold (Text-Figure 1), but switching in rank abundance between the LA and the expected DA still occurs largely among the most abundant species in the LA and the effects of LB are moderated as the evenness of the LA decreases.





**SUPPLEMENTAL FIGURE 3.** Additional simulations of expected death assemblages (exp DA) for the various hypothetical living assemblages (hypoth LA) in Text-figure 1, using an approximately 80-fold variation in lifespan (the set 1, 3, 8, 35, and 75 years), which reflects the minimum, first mode, median, and maximum lifespans in the entire abundance database (see Supplemental Figure 1). Only slightly larger values of  $|LB|$  are generated than when lifespans vary by 40-fold (Supplemental Figure 2). Switching in rank abundance between the LA and the expected DA still occurs largely among the most abundant species in the LA and the effects of LB are moderated as the evenness of the LA decreases.




SUPPLEMENTAL FIGURE 4. A. Frequency distribution of observed discordance (OD) and lifespan bias (LB) using the same data as plotted in Text-figure 2 (all 413 living species occurrences). Note expansion in scale to increments of  $|0.01|$  around zero. B. Cumulative proportional frequency of OD and LB.




SUPPLEMENTAL FIGURE 5. Results of Text-figure 5 parsed according to degree of human activities in the study area. A. Species occurrences from pristine areas. B. Species occurrences from areas subject to human activities.



SUPPLEMENTAL FIGURE 6. A. Among the 85 unique bivalve species (out of 205 unique species in the abundance database) for which we have data from the literature, body size tends to increase with lifespan (trend line is approximate). Body sizes range from 2.5 to 181 mm, where size is the geometric mean of width and height of the largest reported specimens in the global literature. The lifespans of these species range from 2 to 53 years, using data from our lifespan database. Linear regression of  $\ln$  body size and  $\ln$  lifespan, coefficient = 0.52,  $p = 0.001$ , adjusted  $r^2 = 0.11$ . A rank-order test of the same data yields Spearman rho = 0.29,  $p = 0.007$ . B. OD is not correlated significantly with body size among the 228 species occurrences for which we have body size data (Spearman rank-order test rho = 0.005).



SUPPLEMENTAL FIGURE 7. Log-log version of the plot of living proportional abundance and lifespan in Text-Figure 7, showing lack of correlation among the 413 living species occurrences. Black icons = most abundant species in a dataset; superposed white asterisks denote dataset is from a pristine study area.



SUPPLEMENTAL TABLE 1. Summary features of the 30 datasets of living and dead bivalve species abundances used to build the abundance database, drawn from live-dead studies conducted by other authors in subtidal habitats of coastal embayments or the open continental shelf. Datasets are listed by degree of suspected modification from anthropogenic eutrophication (AE) and bottom trawling (BT; see Kidwell 2007 for sources of information on human impacts). AE0 = no extra input of nutrients; AE0.5 = possible input from watershed; AE1 = definite input, e.g. from agriculture, coastal industry or urban areas; AE1.5 =definite input and possibly severe; AE2 = severe or near point source. BT0 = no commercial exploitation of demersal fin- and shellfish or only artisanal methods with minimal damage to habitat and negligible bycatch; BT0.5 = bottom trawling might be active but uncertain; BT1 = commercial fishery using heavy gear; BT2 = especially intense commercial trawling of the seafloor, e.g. every year or more than once a year. For studies where the original author sampled the living assemblage repeatedly, we used the census that yielded the largest number of living individuals (e.g., “LA 4” indicates this was the fourth census in the series). These 30 datasets constitute all available live-dead molluscan datasets containing at least 100 living bivalve individuals after pooling of all available raw samples from the habitat, with each habitat constituting a lithologically homogeneous area within the study. Many well-known live-dead studies that have been included in other meta-analyses (e.g., Kidwell 2001, 2007) are not included here because their bivalve LA did not reach this minimal sample size in a single sampling of the habitat. S denotes number of species; those species constituting at least 0.001 of the LA were included in the analysis, and generate the analytical S of the dataset (contrasted with the raw S of that dataset). N denotes number of bivalve individuals in the raw LA. PIE = evenness in distribution of individuals among species, based on Hurlbert 1971. Minimum and maximum lifespans encountered in the analytical LA and in the raw LA. The quality of information (“knowledge”) of lifespans for a dataset is expressed as the proportion of species whose lifespan is estimated on the basis of published data for that species as opposed to data on congeneric or confamilial species.

Study area and seafloor type	Lead author	Setting	AE score	BT score	Mesh size (mm)	Latitude (°N)	LA analytic S (only species constituti ng ≥0.001 of raw sampled LA)	LA raw S (includes species rarer than 0.001)	LA raw N				DA raw N			
									LA raw PIE	DA raw S	DA raw N	DA raw PIE	LA raw N	DA raw N	DA raw PIE	
<b>Pristine areas, processed using coarse-mesh &gt;1 mm:</b>																
Laguna Madre shelly sand	Smith	Embay	0	0	5	26.5	16	16	1647	0.47	12	735	0.30			
Lower Laguna Madre sand	Smith	Embay	0	0	5	26.5	19	19	175	0.89	18	366	0.82			
Mugu Lagoon sand (LA census 5)	Peterson	Embay	0	0	2	34	13	13	1925	0.56	29	3786	0.67			
San Juan Strait shell gravel	Kowalewski	Open	0	0	2.3	48.5	23	23	827	0.82	31	4669	0.84			

Benthic Habitat Data Summary													
Location		Site Type		Depth (m)		Area (km²)		Mean Depth (m)		Mean Area (km²)		Mean Density (ind/m²)	
Region	Site Name	Sampling Method	Setting	Min	Max	Mean	SD	Min	Max	Mean	SD	Min	Max
Yucatan backreef gravelly	Ekdale	Embay	0	0	3	21	21	21	378	0.79	77	12328	0.90
Yucatan lagoon sandy mud	Ekdale	Embay	0	0	3	21	23	23	981	0.86	62	29710	0.89
Yucatan Strait shell gravel	Ekdale	Open	0	0	3	21	12	21	5087	0.16	75	68052	0.65
<b>Pristine areas, processed using fine-mesh ≤1 mm:</b>													
Mannin Bay algal bank	Bosence	Embay	0	0	0.5	53	14	14	242	0.76	21	895	0.86
Mannin Bay clean algal gravel	Bosence	Embay	0	0	0.5	53	17	20	2129	0.21	35	3678	0.81
Mannin Bay fine sand	Bosence	Embay	0	0	0.5	53	13	13	243	0.57	36	1248	0.87
Mannin Bay muddy algal gravel	Bosence	Embay	0	0	0.5	53	22	22	339	0.84	26	1950	0.70
<b>Human-modified areas, all mesh sizes:</b>													
Tijuana Slough sand (LA census 5)	Peterson Garcia-Cubas	Embay	0.5	0	2	32.5	13	13	571	0.58	22	691	0.78
Laguna Chica mud	Embay	1	0	1.5	20	8	8	404	0.25	15	1006	0.72	
Tomales Bay mud	Johnson	Embay	0.5	0.5	1.5	38	11	11	271	0.57	24	10156	0.05
Tomales Bay muddy sand	Johnson	Embay	0.5	0.5	1.5	38	11	11	162	0.77	16	66	0.88
Tomales Bay sand	Johnson	Embay	0.5	0.5	1.5	38	10	10	176	0.69	15	54	0.83
Chesapeake Bay eelgrass	Jackson	Embay	1	0.5	1	37.5	10	10	396	0.73	12	568	0.85
West Bay muddy sand	White	Embay	1	0.5	1	29.5	16	16	419	0.73	33	1992	0.68
West Bay sand	White	Embay	1	0.5	1	29.5	10	10	260	0.64	24	1074	0.33
West Bay sandy mud	White	Embay	1	0.5	1	29.5	14	14	350	0.81	30	833	0.75
Corpus Christi shelf sand	White	Open	0	2	0.5	27.5	11	15	3112	0.10	57	34357	0.55
Corpus Christi shelf sandy mud (LA census 4)	Staff	Open	0	2	1	27.5	20	20	988	0.17	47	7986	0.91
Laguna Pueblo Viejo mud	Reguero	Embay	2	0	1.5	22	7	7	122	0.63	10	3168	0.61
Galveston shelf shelly muddy sand	White	Open	1	2	0.5	29.5	14	14	125	0.85	66	31541	0.78
Plymouth Sound shell gravel (LA census 1)	Carthew	Open	1	2	2	50	10	11	4818	0.06	44	14817	0.37
Rhodes shelf silty sand (LA census Summer)	Zenetos	Open	2	1	0.5	36	37	37	175	0.91	25	167	0.92
<b>Small datasets (S &lt;7), all areas and mesh sizes:</b>													
Laguna Camaronera shell gravel	Reguero	Embay	1	0	1.5	19	4	4	139	0.12	18	1460	0.61
Laguna Pueblo Viejo muddy sand	Reguero	Embay	2	0	1.5	22	4	4	98	0.48	8	4587	0.60
Laguna Alvarado sand	Reguero	Embay	2	0	1.5	19	4	4	350	0.57	9	4278	0.46
Laguna Alvarado shell gravel	Reguero	Embay	2	0	1.5	19	6	6	1238	0.72	9	45013	0.32

Total	30 datasets												
Median													

413  
living  
species  
occurren  
ces  
13

[384  
dead-  
only]  
286661  
1971  
0.732

**Pooling of multiple samplings of LA:**

Mugu Lagoon sand (pool 7 samplings of LA over 24 mo, 1 sampling of DA)	Peterson	Embay	0	0	2	34	15	21	9463	0.46	29	3786	0.67
Tijuana Slough sand (pool 7 LAs over 24 mo, 1 of DA)	Peterson	Embay	0.5	0	2	32.5	18	19	2981	0.64	22	691	0.78
Corpus Christi shelf sandy mud (pool 9 LAs over 14 mo, 1 final DA)	Staff	Open	0	2	1	27.5	36	48	3801	0.73	42	7325	0.83
Plymouth Bay shell gravel (pool 4 LAs over 7 mo, 1 DA)	Carthew	Open	1	2	2	50	14	17	8396	0.16	53	37965	0.37
Rhodes shelf silty sand (pool 4 LAs over 9 mo)	Zenetos	Open	2	1	0.5	36	55	55	462	0.90	25	167	0.92

Study area and seafloor type	Min, max lifespan (years), analytic LA	Min, max lifespan (years), raw LA	Median lifespan (years), raw LA	Proportion of lifespans known for species	Proportion of lifespans known for genus	top-ranked species in LA & its maximum lifespan (y)
<b>Pristine areas, processed using coarse-mesh &gt;1 mm:</b>						
Laguna Madre shelly sand	3, 19	2, 19	8	0.16	0.42	Anomalocardia cuneimeris 5
Lower Laguna Madre sand	2, 19	2, 19	6	0.19	0.33	Chione cancellata 6
Mugu Lagoon sand (LA census 5)	4, 53	2, 53	10	0.20	0.23	Cryptomya californica 21
San Juan Strait shell gravel	2, 75	2, 75	16	0.31	0.29	Modiolus modiolus 75
Yucatan backreef gravelly	2, 38	2, 41	5	0.09	0.21	Transennella cubaniana 14
Yucatan lagoon sandy mud	2, 18	2, 41	5	0.06	0.26	Carditamera floridana 18
Yucatan Strait shell gravel	2, 18	2, 44	5	0.11	0.23	Ervilia nitens 4

**Pristine areas, processed using fine-mesh****≤1 mm:**

Mannin Bay algal bank	2, 14	1, 27	7	0.61	0.04	<i>Musculus discors</i> 6
Mannin Bay clean algal gravel	2, 29	1, 27	10	0.41	0.14	<i>Goodallia triangularis</i> 29
Mannin Bay fine sand	2, 19	1, 27	7	0.51	0.11	<i>Thyasira flexuosa</i> 7
Mannin Bay muddy algal gravel	2, 19	1, 28	9	0.53	0.10	<i>Kurtiella bidentata</i> 7

**Human-modified areas, all mesh sizes:**

Tijuana Slough sand (LA census 5)	3, 32	2, 32	8.5	0.18	0.32	<i>Nuttallia nuttallii</i> 32
Laguna Chica mud	2, 17	2, 18	6	0.27	0.40	<i>Cyrenoida floridana</i> 11
Tomales Bay mud	1, 21	1, 25	9	0.38	0.31	<i>Lyonsia californica</i> 9
Tomales Bay muddy sand	1, 17	1, 25	10	0.39	0.17	<i>Lyonsia californica</i> 9
Tomales Bay sand	1, 17	1, 53	10.5	0.28	0.28	<i>Cadella nuculoides</i> 17
Chesapeake Bay eelgrass	3, 46	3, 46	8.5	0.29	0.24	<i>Amygdalum papyrium</i> 6
West Bay muddy sand	2, 19	2, 19	9	0.22	0.28	<i>Mulinia lateralis</i> 3
West Bay sand	3, 19	3, 19	7	0.19	0.27	<i>Mulinia lateralis</i> 3
West Bay sandy mud	3, 15	2, 15	7	0.22	0.16	<i>Mulinia lateralis</i> 3
Corpus Christi shelf sand	2, 14	1, 18	6	0.16	0.26	<i>Abra aequalis</i> 3
Corpus Christi shelf sandy mud (LA census 4)	2, 40	1, 46	6	0.22	0.24	<i>Abra aequalis</i> 3
Laguna Pueblo Viejo mud	3, 17	3, 18	6	0.40	0.30	<i>Cyrenoida floridana</i> 11
Galveston shelf shelly muddy sand	2, 17	1, 23	8	0.19	0.24	<i>Sacella concentrica</i> 9
Plymouth Sound shell gravel (LA census 1)	2, 15	2, 40	9	0.34	0.18	<i>Spisula elliptica</i> 10
Rhodes shelf silty sand (LA census Summer)	2, 29	1, 45	9	0.23	0.25	<i>Nuculana fragilis</i> 10

**Small datasets (S <7), all areas and mesh sizes:**

Laguna Camaronera shell gravel	3, 17	2, 18	6	0.22	0.22	<i>Mulinia lateralis</i> 3
Laguna Pueblo Viejo muddy sand	3, 17	3, 18	6	0.40	0.30	<i>Cyrenoida floridana</i> 11
Laguna Alvarado sand	3, 10	3, 15	6	0.33	0.56	<i>Hormomya exustus</i> 3
Laguna Alvarado shell gravel	3, 15	3, 15	6	0.33	0.50	<i>Polymesoda caroliniana</i> 6

Total

Median (2, 18.5) 2, 26 7 0.25 0.25

**Pooling of multiple samplings of LA:**

Mugu Lagoon sand (pool 7 samplings of LA over 24 mo, 1 sampling of DA)	3, 53	2, 53	10	0.29	0.38	Cryptomya californica
Tijuana Slough sand (pool 7 LAs over 24 mo, 1 of DA)	3, 53	2, 53	9	0.26	0.42	Nuttallia nuttallii
Corpus Christi shelf sandy mud (pool 9 LAs over 14 mo, 1 final DA)	2, 46	11, 46	6	0.23	0.33	Abra aequalis
Plymouth Bay shell gravel (pool 4 LAs over 7 mo, 1 DA)	2, 20	2, 40	9	0.37	0.17	Parvicardium scabrum
Rhodes shelf silty sand (pool 4 LAs over 9 mo)	2, 38	1, 45	9	0.27	0.25	Nuculana fragilis

**REFERENCES (sources of raw abundance data) :**

- BOSENCE, D.W.J. 1979, Live and dead faunas from coralline algal gravels, Co. Galway, Eire: Palaeontology, v. 22, p. 449-478. [raw data provided by authors but archived at British Museum]
- CARTHEW, R. and D. BOSENCE. 1986: Community preservation in Recent shell-gravels, English Channel. Palaeontology 29, 243-268. [raw data provided by authors but archived at British Museum]
- EKDALE, A. A. 1972. Ecology and paleoecology of marine invertebrate communities in calcareous substrates, northeast Quintana Roo, Mexico [M.S. thesis]: Houston, Rice University, 159 p.
- EKDALE, A. A. 1977, Quantitative paleoecological aspects of modern marine mollusk distribution, northeast Yucatan coast, Mexico. In: Frost, S. H., M. P. Weiss and J. B. Saunders, eds., American Association of Petroleum Geologists, Studies in Geology 4: 195-207.

- GARCIA-CUBAS, A., M. REGUERO, and R. ELIZARRARAS. 1992. Moluscos del sistema lagunar Chica-Grande, Veracruz, Mexico: sistematica y ecologia: Anales del Instituto de Ciencias del Mar y Limnologia, Universidad nacional Autonoma de Mexico 19(1): 1-121.
- JACKSON, J.B.C.. 1968. Neontological and paleontological study of the autecology and synecology of the molluscan fauna of Fleets Bay, Virginia [M.S. thesis]: Washington, D.C., George Washington University, 111p.
- JOHNSON, R. G. 1965. Pelecypod death assemblages in Tomales Bay, California. Journal of Paleontology 39: 80-85. [raw data archived at California Academy of Sciences]
- KOWALEWSKI, M., M. CARROLL, L. CASAZZA, N. GUPTA, B. HANNISDAL, et al. 2003. Quantitative fidelity of brachiopod-mollusk assemblages from modern subtidal environments of San Juan Islands, USA. Journal of Taphonomy 1: 43-65. [authors provided raw data on counts of living individuals, all identifiable dead specimens (includes small fragments that are not unique individuals), and whole dead shells (unique individuals; each whole bivalve shell is assumed here to represent a unique bivalve individual). We used whole dead shells only as the death assemblage]
- PETERSON, C.H. 1976. Relative abundance of living and dead molluscs in two California lagoons: *Lethaia*, v. 9, p. 137-148.
- PETERSON, C.H. 1972. Species diversity, disturbance and time in the bivalve communities of some California lagoons. PhD dissertation, University of California Santa Barbara, 230 p.
- REGEUERO, M. and A. GARCIA-CUBAS. 1989. Moluscos de la Laguna de Alvarado, Veracruz: sistematica y ecologia. Anales del Instituto de Ciencias del Mar y Limnologia, Universidad nacional Autonoma de Mexico 16(2): 279-306.
- REGEUERO, M. and A. GARCIA-CUBAS. 1991. Moluscos de la Laguna Camaronera, Veracruz, Mexico: sistematica y ecologia. Anales del Instituto de Ciencias del Mar y Limnologia, Universidad nacional Autonoma de Mexico 18(1): 1-23.
- REGEUERO, M. and A. GARCIA-CUBAS. 1993. Moluscos de la Laguna Pueblo Viejo, Veracruz, Mexico: sistematica y ecologia. Anales del Instituto de Ciencias del Mar y Limnologia, Universidad nacional Autonoma de Mexico 20(1): 77-104.

- SMITH, E.J. 1985. Paleoecologic aspects of modern macroinvertebrate communities of southern Laguna Madre, Texas [M.S. thesis]: Nacogdoches, Texas, Stephen F. Austin State University, 79 p.
- STAFF, G.M. and E.N. POWELL. 1999: Onshore-offshore trends in community structural attributes: death assemblages from the shallow continental shelf of Texas. *Continental Shelf Research* 19, 717-756. [raw data provided by authors]
- WHITE, W. A., T. R. CALNAN, R. A. MORTON, R. S. KIMBLE, T. G. LITTLETON, J.H. McGOWEN and H.S. NANCE. 1983. Submerged lands of Texas, Corpus Christi area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands. Austin, Texas: Bureau of Economic Geology, University of Texas at Austin. [raw data archived at Univ Texas Austin Library]
- WHITE, W. A., T. R. CALNAN, R. A. MORTON, R. S. KIMBLE, T. G. LITTLETON, J.H. McGOWEN, H.S. NANCE, and K.E. SCHMEDES. 1985. Submerged lands of Texas, Galveston-Houston area: sediments, geochemistry, benthic macroinvertebrates, and associated wetlands. Austin, Texas: Bureau of Economic Geology, University of Texas at Austin. [raw data archived at Univ Texas Austin Library]
- ZENETOS, A. and J. J. VAN AARTSEN. 1994 (1995). The deep sea molluscan fauna of the S.E. Aegean Sea and its relation to the neighbouring fauna. *Bollettino Malacologico* 30: p. 253-268. [raw data provided by authors]

SUPPLEMENTAL TABLE 2. Database of published maximum lifespans of marine and estuarine bivalve species, in years, used to estimate the maximum lifespans of species in 30 live-dead datasets (Supplemental Table 1). Full references at end of this file. Lifespan data were gathered from (1) fully referenced secondary compilations (Comfort 1957; Robertson 1979; Powell and Cummins 1985; Powell and Stanton 1985; Heller 1990; Callender and Powell 1997; websites “MarLIN BIOTIC” and “MSAP”), (2) primary studies of multiple species in a region (Zaika 1973; Petersen 1958 and 1977; Zolotarev 1980 and 1982; Maximovich 1984; Le Pennec and Beninger 2000; Banqueiro Cardenas and Aranda 2003 (latter provides instantaneous mortality rates rather than lifespans)), and (3) approximately 100 primary studies of single species located by electronic search of the *Zoological Record* using the keywords “Bivalvia”, “lifespan”, and “longevity” (September 2008).

**Note 1. Taxonomic consistency.** In compiling lifespan data, we encountered inconsistency among published reports in the genus assignments of some species because the lifespan literature draws on papers published from the 1920s to the 2000’s, during which time bivalve systematics underwent considerable revision at the genus level. For example, many large Linnean genera such as *Nuculana*, *Arca*, *Pecten*, *Ostrea*, *Cardium*, *Tellina*, *Venus*, and *Corbula* have been subdivided and most subgenera recognized by the Bivalve Treatise on Invertebrate Paleontology have been elevated to full genus rank (see the global compendia for pectinids of Djikstra (2004) and for cardiids by ter Poorten (2005), and, for all other groups, the regional compendia of CLEMAM (western Europe), Malacolog 4.1.0 (western Atlantic), Coan et al. (2000; eastern Pacific), and OBIS (Indo-Pacific)). For internal consistency, we used these taxonomic authorities to update the genus assignments of species so that, for example, all published lifespan estimates for the common European cockle appear as *Cerastoderma edule* and not also as *Cardium edule*. In cases where taxonomic authorities differed, we deferred to the 2008 version of an ongoing effort to reconcile these and other sources on a global scale by Jablonski, Roy, and Valentine (described in Krug et al. 2008, 2009). We used the same procedure to make consistent all bivalve species names in the 30 live-dead datasets (Supplemental Table 1) so that the lifespan and abundance databases could be merged. In all cases we assumed that the original author had correctly identified the species and family of the specimens in his/her analysis. Our updating procedure thus decreased the number of unique species recognized in either the lifespan or the abundance database (because entities like *Cerastoderma edule* and *Cardium edule* were merged). This procedure did not change the number of families recognized, but increased the number of recognized genera (in the lifespan database, an increase from ~100 to ~150 genera, largely owing to elevation of subgenera to genus rank). Genus names in quotes are assignments of species in the lifespan literature that could not be verified.

**Note 2. Availability of multiple estimates of lifespan.** We were able to locate more than one estimate of maximum lifespan for 42% of bivalve species encountered in the lifespan literature, for 36% of (updated) genera, and for 67% of families. Multiple estimates arise from analysis of geographically distinct populations of a species and the use of different methods of age determination, which include growth lines, size-frequency distributions, growth rates, and mark-recapture experiments. Multiple entries in the table below for a single species by a single author indicate that the author examined multiple populations or applied more than one method to a single population. Like stratigraphic and geographic ranges, the maximum lifespan of a species will tend to be underestimated: the maximum lifespan achieved within a given stock can be

depressed by an array of environmental conditions and is also sensitive to sampling intensity (see Beukema 1989; Powell et al 2001; and discussion in main text). Thus, when multiple estimates of maximum lifespan were available for a species, we used the largest of all reported values as its global maximum lifespan.

Note 3. Estimating the maximum lifespans of species occurring in the abundance database. The global maximum lifespan of a species (see Note 2) was applied as the relevant lifespan to all occurrences of that species in the abundance database. 32 % of all species occurrences in the abundance database could be assigned a lifespan with this species-level resolution. When a published report was not available for a species, we used the median of maximum lifespans of other species in that same genus (rounding up to full years; each species contributed only one maximum lifespan value to the calculation for the genus). Such genus-based estimates were the finest estimate possible for 26 % of all species occurrences in the abundance database. When a published report was not available for either a species or its genus, we used the median of other genera in the same family (each genus contributed only one maximum lifespan value to the calculation for the family). Family-level estimates were assigned to 37% of species occurrences in the abundance database.

Note 4. Clade-level differences in lifespan. Among the families where we had lifespan estimates for at least 2 genera and at least 5 species, some families contain mostly short-lived species (maximum lifespan <8 years; e.g., Lucinidae, Donacidae, Semelidae, Montacutidae, Corbulidae, Corbiculidae, Teredinidae, perhaps Dreisseneidae) whereas others contain mostly long-lived species (>9 years; Carditidae, Cultellidae, Myidae, Hiatellidae, perhaps Arcidae) and yet others contain mostly intermediate-lifespan species (8-9 years; Nuculidae, Nuculanidae). Some very large families are best described as heterogeneous in maximum lifespan (e.g., Mactridae, Mytilidae, Pectinidae, Ostreidae, Tellinidae).

Family	Genus (updated from the primary source as needed)	Species	Lifespan (yrs)	Primary source of information (secondary source, if used)
Anomiidae	Pododesmus	squamula	6	Maximovich, N.V. 1984.
Arcidae	Anadara	tuberculosa	13	Baqueiro Cardenas and Aranda, 2003
Arcidae	Arca	boucardi	14	Zolotarev, 1980
Arcidae	Arca	boucardi	13	Zolotarev, 1980
Arcidae	Arca	boucardi	20	Zolotarev, 1980 (Heller, 1990)
Arcidae	Arca	noae	16	Peharda et al, 2002.
Arcidae	Mabellarca	notabilis	4	Jiminez et al, 2004.
Arcidae	Scapharca	broughtoni	46	Zolotarev, 1980 (Heller, 1990)
Arcidae	Senilia	senilis	9	Okera, 1976 (Powell & Stanton, 1985)
Arcidae	Senilia	senilis	8	Zolotarev 1982; Table 1
Arcidae	Tegillarca	granosa	4	de Goeij et al, 2003.
Arcticidae	Arctica	islandica	192	Abele et al, 2008.
Arcticidae	Arctica	islandica	400	Abele et al, 2008.
Arcticidae	Arctica	islandica	100	Duineveld et al., 2002
Arcticidae	Arctica	islandica	70	Jones, 1980 (Powell and Stanton, 1985)

Arcticidae	Arctica	islandica	220	Jones, 1983 (Heller, 1990)
Arcticidae	Arctica	islandica	14	Petersen, G. H., 1958.
Arcticidae	Arctica	islandica	100	Thorarinsdottir, 1999 (MarLIN BIOTIC)
Astartidae	Astarte	sulcata	45	Weber et al, 2001.
Astartidae	Tridonta	arctica	7	Selin, NI, 2007.
Astartidae	Tridonta	borealis	8	Selin, NI, 2007.
Astartidae	Tridonta	elliptica	10	Erlenkeuser et al. 1975 ( Trutschler, K and C Samtleben, 1988.)
Astartidae	Tridonta	elliptica	2	Jaeckel 1952 (Trutschler, K and C Samtleben, 1988.)
Astartidae	Tridonta	elliptica	20	Trutschler, K and C Samtleben, 1988.
Astartidae	Tridonta	elliptica	6	Maximovich, N.V. 1984.
Astartidae	Tridonta	montagui	20	Petersen, G. H., 1958.
Astartidae	Tridonta	montagui	3	Maximovich, N.V. 1984.
Astartidae	Tridonta	borealis	6	Maximovich, N.V. 1984.
Axinopsidae	Axinopsida	sericata	4	Zaika, VE, 1973.
Cardiidae	Acanthocardia	echinata	15	Petersen, G. H., 1958.
Cardiidae	Cerastoderma	edule	6	Beukema 1989
Cardiidae	Cerastoderma	edule	7	Boyden 1972 (Powell & Stanton, 1985)
Cardiidae	Cerastoderma	edule	7	Cerrato, 1980 (Heller, 1990)
Cardiidae	Cerastoderma	edule	14	Cole, 1956 (Comfort, 1957)
Cardiidae	Cerastoderma	edule	11	Evans and Tallmark, 1977 (Powell and Stanton, 1985)
Cardiidae	Cerastoderma	edule	10	Fretter and Graham 1964 (MarLIN BIOTIC)
Cardiidae	Cerastoderma	edule	7	Hibbert 1976 (Roberston 1979)
Cardiidae	Cerastoderma	edule	3	Muus, BJ, 1967.
Cardiidae	Cerastoderma	edule	5	Stephen, 1931 (Comfort, 1957)
Cardiidae	Cerastoderma	edule	6	Warwick and Price, 1975 (Robertson, 1979)
Cardiidae	Cerastoderma	edule	5	Wolff and deWolf, 1977 (Robertson, 1979)
Cardiidae	Cerastoderma	edule	5	Zaika, VE, 1973.
Cardiidae	Cerastoderma	edule	10	Zolotarev 1982; Table 1
Cardiidae	Cerastoderma	glaucum	7	Boyden 1972 (Powell & Stanton, 1985)
Cardiidae	Cerastoderma	glaucum	5	MarLIN BIOTIC
Cardiidae	Cerastoderma	glaucum	9	Zolotarev 1982; Table 1
Cardiidae	Cerastoderma	glaucum	4	Muus, BJ, 1967. [lamarcki =jr syn of glaucum]
Cardiidae	Ciliatocardium	ciliatum	25	Petersen, 1978 (Heller, 1990)
Cardiidae	Ciliatocardium	ciliatum	33	Zolotarev 1982; Table 1
Cardiidae	Clinocardium	corbis	10	Cerrato, 1980 (Heller, 1990)
Cardiidae	Clinocardium	corbis	7	Fraser, 1931 (Comfort, 1957)
Cardiidae	Clinocardium	corbis	16	Taylor, 1960 (Powell & Cummins, 1985)
Cardiidae	Clinocardium	corbis	16	Weymouth & Thompson, 1930 (Comfort, 1957)
Cardiidae	Clinocardium	nuttallii	18	Hoening, 1983 (Powell & Cummins, 1985)
Cardiidae	Clinocardium	nuttallii	14	Zolotarev, 1980 (Heller, 1990)
Cardiidae	Keenocardium	californiense	11	Zolotarev, 1980 (Heller, 1990)
Cardiidae	Keenocardium	californiense	7	Zolotarev, 1980 (Heller, 1990)

Cardiidae	Parvicardium	exiguum	1	Lastra, M, A Sanchez, and J Mora, 1993.
Cardiidae	Parvicardium	exiguum	3	Petersen, G. H., 1958.
Cardiidae	Serripes	groenlandicus	22	Petersen, 1978 (Heller, 1990)
Cardiidae	Serripes	groenlandicus	33	Zolotarev 1982; Table 1
Carditidae	Carditamera	affinis	18	Riascos et al, 2008.
Carditidae	Cyclocardia	crebricostata	58	Zolotarev, 1980 (Heller, 1990)
Corbiculidae	Corbicula	fluminea	7	Hall 1984 The Nautilus
Corbiculidae	Corbicula	fluminea	3	Hornbach, DJ, 1992.
Corbiculidae	Corbicula	fluminea	5	Mouthon, J, 2001.
Corbiculidae	Corbicula	largillieri	3	Ituarte, CF, 1984.
Corbiculidae	Polymesoda (Geloina)	erosa	6	Morton, B, 1988.
Corbulidae	Aniscorbula	venusta	8	Zolotarev, 1980 (Heller, 1990)
Corbulidae	Corbula	trigona	2	Maslin & Bouvet 1986 (Callender and Powell, 1997)
Corbulidae	Corbula	trigona	2	Maslin & Bouvet, 1986
Corbulidae	Corbula	vicaria	4	Rainer, 1982 (Powell & Cummins, 1985)
Corbulidae	Potamocorbula	amurensis	5	Zolotarev, 1980 (Heller, 1990) ( <i>ustulata</i> is a syn)
Corbulidae	Varicorbula	gibba	5	Holmes, SP and N Miler 2006.
Corbulidae	Varicorbula	gibba	2	Hrs-Brenko, M., 2006.
Corbulidae	Varicorbula	gibba	6	Janesen, 1990 (MarLIN BIOTIC)
Corbulidae	Varicorbula	gibba	4	Petersen, G. H., 1977.
Corbulidae	Varicorbula	gibba	2	Power et al, 2006.
Cultellidae	Ensis	macha	14	Urban, 1996
Cultellidae	Ensis	siliqua	12	Comfort, 1964 (Heller, 1990)
Cultellidae	Siliqua	alta	22	Zolotarev, 1980
Cultellidae	Siliqua	alta	24	Zolotarev, 1980
Cultellidae	Siliqua	patula	17	Cerrato, 1980 (Heller, 1990)
Cultellidae	Siliqua	patula	25	Weymouth & McMillin, 1931 (Comfort, 1957)
Cyrenoididae	Cyrenoida	floridana	11	Veira et al, 2006.
Donacidae	"Donax"	hanleyanus	3	Ansell, 1983 (Heller, 1990)
Donacidae	"Donax"	hanleyanus	3	Penchaszadeh & Olivier, 1975 (Powell & Cummins, 1985)
Donacidae	"Donax"	spiculum	1	Ansell et al, 1972 (Powell & Cummins, 1985)
Donacidae	"Donax"	tumida	1	Loesch, 1957 (Powell & Cummins, 1985)
Donacidae	Chion	denticulatus	2	Wade, 1968 (Powell & Cummins, 1985)
Donacidae	Chion	incarnatus	3	Ansell et al, 1972 (Powell & Cummins, 1985)
Donacidae	Chion	variabilis	1	Ansell, 1983 (Heller, 1990)
Donacidae	Chion	variabilis	1	Jones, et al, 2004.
Donacidae	Chion	variabilis	1	Loesch, 1957 (Powell & Cummins, 1985)
Donacidae	Cuneus	semistriatus	1	Ansell, 1983 (Heller, 1990)
Donacidae	Cuneus	venustus	1	Ansell, 1983 (Heller, 1990)
Donacidae	Cuneus	vittatus	7	Ansell, 1983 (Heller, 1990)
Donacidae	Cuneus	vittatus	10	Petersen, G. H., 1977.
Donacidae	Donax	gouldi	3	Coe, 1955 (Powell & Cummins, 1985)
Donacidae	Donax	gouldi	2	Fitch, JE, 1952.

Donacidae	Grammatodonax	sordidus	2	McLachlan, A, 1979.
Donacidae	Latona	cuneatus	2	Nayar, 1955 (Comfort, 1957)
Donacidae	Latona	serra	1	Ansell, 1983 (Heller, 1990)
Donacidae	Latona	serra	5	Laudien, J, 2002.
Donacidae	Latona	serra	5	McLachlan & Hanekom, 1979 (Powell & Cummins, 1985)
Donacidae	Serrula	trunculus	3	Ansell, 1983 (Heller, 1990)
Donacidae	Serrula	trunculus	5	Bayed, A, 1998.
Donacidae	Serrula	trunculus	4	Manca Zeichen et al, 2002.
Dreisseneidae	Dreissena	polymorpha	4	Chase, ME, and RC Bailey, 1999.
Dreisseneidae	Dreissena	polymorpha	6	Kornobis, 1977 (Callender and Powell, 1997)
Dreisseneidae	Dreissena	polymorpha	7	Stanezykowska, 1977 (Callender and Powell, 1997)
Dreisseneidae	Dreissena	polymorpha	19	Verween et al, 2006.
Dreisseneidae	Dreissena	polymorpha	4	Verween et al, 2006.
Dreisseneidae	Dreissena	polymorpha	2	Verween et al, 2006.
Dreisseneidae	Mytilopsis	leucophaeata	6	Verween et al, 2006.
Dreisseneidae	Mytilopsis	sallei	1	Morton, B, 1981 (Heller, 1990)
Dreisseneidae	Mytilopsis	sallei	2	Morton, B, 1989.
Dreisseneidae	Mytilopsis	sallei	2	Power et al, 2006.
Dreisseneidae	Dreissena	polymorpha	6	Zaika, VE, 1973.
Glycymeridae	Glycymeris	glycymeris	17	Menesguen and Dreves, 1987 (Callender and Powell, 1997)
Glycymeridae	Glycymeris	yessoensis	64	Zolotarev, 1980 (Heller, 1990)
Hiatellidae	Hiatella	arctica	3	Lees, DC, 2005 (unpub) Assessment of Bivalve Recovery, Prince William Sound
Hiatellidae	Hiatella	arctica	6	Zolotarev 1982; Table 1
Hiatellidae	Hiatella	arctica	6	Maximovich, N.V. 1984.
Hiatellidae	Hiatella	byssifera	15	Petersen, GH, 1978 (Heller, 1990)
Hiatellidae	Panopea	abbreviata	40	Morsan, E and NF Ciocco, 2004.
Hiatellidae	Panopea	abrupta	160	Bureau et al. 2002 (cit by Strom et al, 2005)
Hiatellidae	Panopea	generosa	15	Fitch, JE, 1952.
Hiatellidae	Panopea	generosa	120	Jones, 1983 (Heller, 1990)
Hiatellidae	Panopea	generosa	100	Shaul & Goodwin, 1982 (Powell & Cummins, 1985)
Hiatellidae	Panopea	zelandica	34	Gribben, PE, and RG Creese, 2005.
Hiatellidae	Panopea	zelandica	85	Gribben, PE, and RG Creese, 2005.
Lasaeidae (or Galeommatidae)	Lasaea	rubra	4	McGrath & O'Foighil, 1986 (Heller, 1990)
Laternulidae	Laternula	elliptica	36	Philipp et al, 2005.
Laternulidae	Laternula	elliptica	13	Ralph & Maxwell, 1977 (Powell & Cummins, 1985)
Lucinidae	Anodontia	omissa	1	de Goeij et al, 2003.
Lucinidae	Anodontia	omissa	2	Rainer, 1982 (Powell & Cummins, 1985)
Lucinidae	Codakia	orbicularis	3	Le Pennec, M, and PG Beninger, 2000.
Lucinidae	Divaricella	irpex	1	de Goeij et al, 2003.
Lucinidae	Loripes	lacteus	5	Velosa et al, 2007.
Lucinidae	Loripes lacteus	lucinalis	3	Le Pennec, M, and PG Beninger, 2000.
Lucinidae	Pillucina	neglecta	2	Mukai, 1974 (Robertson, 1979)

Mactridae	Mactra	corallina	6	Petersen, G. H., 1977.
Mactridae	Mactra	incongrua	2	Tanaka and Kikuchi, 1971 (Robertson, 1979)
Mactridae	Mactra	sulcataria	4	Kim et al. 1983 (Callender and Powell 1997)
Mactridae	Mactra	sulcataria	12	Zolotarev, 1980 (Heller, 1990)
Mactridae	Mactromeris	polynyma	16	Feder et al, 1979 (Powell & Cummins, 1985)
Mactridae	Mactromeris	polynyma	25	Hughes, SE, and N Bourne, 1981.
Mactridae	Mactromeris polynyma	voyi	52	Zolotarev, 1980 (Heller, 1990)
Mactridae	Mulinia	lateralis	3	Cerrato, 1980 (Heller, 1990)
Mactridae	Pseudocardium	sachalinensis	12	Sasaki, 1981 (Powell & Cummins, 1985)
Mactridae	Pseudocardium	sachalinensis	55	Zolotarev, 1980 (Heller, 1990)
Mactridae	Rangia	cuneata	5	Coen, LD, and RE Grizzle, 2007.
Mactridae	Rangia	cuneata	10	Wolfe & Petteway, 1968 (Powell & Cummins, 1985)
Mactridae	Spisula	elliptica	10	Petersen, G. H., 1977.
Mactridae	Spisula	elliptica	1	Zaika, VE, 1973.
Mactridae	Spisula	ovalis	7	Menesguen and Dreves, 1987 (Callender and Powell, 1997)
Mactridae	Spisula	solida	10	Fahy et al, 2003 (MarLIN BIOTIC) (easily confused with elliptica)
Mactridae	Spisula	ravenili or solidissima similis	4	Walker, RL and PB Heffernan, 1994. (southern surfclam; ravenili is syn)
Mactridae	Spisula	ravenili or solidissima similis	6	Walker, RL and PB Heffernan, 1994. (southern surfclam; ravenili is syn)
Mactridae	Spisula	solidissima	10	Cerrato, RM, and DL Keith, 1992.
Mactridae	Spisula	solidissima	22	Cerrato, RM, and DL Keith, 1992.
Mactridae	Spisula	solidissima	25	Jones 1980 (Powell and Stanton 1985)
Mactridae	Spisula	solidissima	31	Jones et al, 1978 (Powell & Cummins, 1985)
Mactridae	Spisula	solidissima	37	Walker, RL and PB Heffernan, 1994. (Atlantic surfclam, northern)
Mactridae	Spisula	subtruncata	1	Bodoy, A, 1980.
Mactridae	Spisula	subtruncata	4	Deval, MC and D Gokturk, 2008.
Mactridae	Spisula	subtruncata	6	Petersen, G. H., 1977.
Mactridae	Tresus	capax	16	Hancock et al, 1979 in Emmett et al. 1991 (MSAP)
Mactridae	Tresus	capax	5	Hancock, et al, 1979. (Rudy & Rudy, 1979)
Mactridae	Tresus	capax	16	Wendell et al, 1976; Bourne & Smith, 1972 (Powell & Cummins, 1985)
Mactridae	Tresus	nuttalli	17	Frey 1971, Wolotira et al. 1989 in Emmett et al. 1991 (MSAP)
Malletiidae	Tindaria	callistiformis	100	Turekian et al, 1975 (Heller, 1990)
Mesodesmatidae	Mesodesma	arctatum	24	Brethes et al, 1986.
Mesodesmatidae	Mesodesma	mactroides	7	Fiori, S and O Defeo, 2006.
Mesodesmatidae	Paphies	ventricosum	9	Comfort, 1957 (Heller, 1990)
Montacutidae	Kurtiella	bidentata		Kunitzer, A, 1989
Montacutidae	Kurtiella	bidentata		MarLIN BIOTIC
Montacutidae	Kurtiella	bidentata		O'Brien, K, and BJ Keegan, 2006.
Montacutidae	Kurtiella	bidentata	7	Ockelmann & Muus, 1978 (Heller, 1990)
Montacutidae	Kurtiella	bidentata	4	O'Foighil et al, 1984.

Montacutidae	Kurtiella	planulata	4	Franz, 1973 (Powell & Stanton 1985)
Montacutidae	Kurtiella	planulata		Santos, SL and JL Simon, 1980.
Montacutidae	Mysella	cuneata	6	Gage, 1968 (Heller, 1990)
Montacutidae	Rochefortia	tumida	7	Lees, DC, 2005 (unpub) Assessment of Bivalve Recovery, Prince William Sound
Montacutidae	Tellimya	ferruginosa	3	Petersen, G. H., 1977.
Myidae	Arenomya	arenaria	2	Burke & Mann, 1974 (Robertson, 1979)
Myidae	Arenomya	arenaria	12	Evans and Tallmark, 1977 (Powell and Stanton 1985)
Myidae	Arenomya	arenaria	28	Jones, 1983 (Heller, 1990)
Myidae	Arenomya	arenaria	28	MacDonald and Thomas 1980 in Emmett et al. 1991; Brousseau 1978, Brousseau and Baglivo 1987 in Emmett et al. 1991 (MSAP)
Myidae	Arenomya	arenaria	8	Maximovich, N.V. 1984.
Myidae	Arenomya	arenaria	8	Newcoombe, 1935; 1936 (Comfort, 1957)
Myidae	Arenomya	arenaria	13	Philipp et al, 2005.
Myidae	Arenomya	arenaria	28	Strasser, M., 1999. (MarLIN BIOTIC)
Myidae	Arenomya	arenaria	17	Warwick and Price, 1975 (Powell & Stanton, 1985)
Myidae	Arenomya	japonica	42	Zolotarev, 1980
Myidae	Mya	priapus	15	Zolotarev, 1980 (Heller, 1990)
Myidae	Mya	truncata	40	Amaro et al, 2003
Myidae	Mya	truncata	40	Duineveld et al.
Myidae	Mya	truncata	18	Petersen, 1978 (Heller, 1990)
Myidae	Panomya	priapus	15	Zolotarev, 1980
Myidae	Sphenia	sincera	4	Hanks, RW, and DB Packer, 1985.
Mytilidae	"Mytilus"	variabilis	5	Sewell, 1924 (Comfort, 1957)
Mytilidae	Adula	californiensis	5	Cerrato, 1980 (Heller, 1990)
Mytilidae	Adula	californiensis	3	Coe & Fox, 1942 (Comfort, 1957)
Mytilidae	Arcuatula	arcuatula	2	George & Nair, 1974 (Powell & Cummins, 1985)
Mytilidae	Bathymodiolus	thermophila	19	Rhoads et al, 1981 (Heller, 1990)
Mytilidae	Brachidontes	variabilis	3	Wilson & Hodgkin, 1967 (Powell & Cummins, 1985)
Mytilidae	Brachidontes	variabilis	3	Morton, B, 1988.
Mytilidae	Crenomytilus	grayanus	150	Jones, 1983 (Heller, 1990)
Mytilidae	Crenomytilus	grayanus	88	Selin, 1980 (Powell & Stanton, 1985)
Mytilidae	Crenomytilus	grayanus	150	Selin, NI, 2004.
Mytilidae	Crenomytilus	grayanus	108	Zolotarev, 1980
Mytilidae	Crenomytilus	grayanus	95	Zolotarev, 1980
Mytilidae	Crenomytilus	grayanus	150	Zolotarev, 1980
Mytilidae	Geukensia	demissa	3	Baqueiro Cardenas, E, and DA Aranda, 2003.
Mytilidae	Geukensia	demissa	15	Franz, DR, 2001.
Mytilidae	Geukensia	demissa	23	Lutz & Castagna, 1980 (Heller, 1990)
Mytilidae	Geukensia	demissus	8	Zaika, VE, 1973.
Mytilidae	Limnoperna	pulex	2	Wilson & Hodgkin, 1967 (Powell & Cummins, 1985)
Mytilidae	Lithophaga	lithophaga	54	Galinou-Mitsoudi, S and AI Sinis, 1995.
Mytilidae	Modiolus	barbatus	14	Peherda et al, 2007.

Mytilidae	Modiolus	modiolus	48	Anwar et al, 1990.
Mytilidae	Modiolus	modiolus	38	Anwar et al, 1990.
Mytilidae	Modiolus	modiolus	75	Anwar et al, 1990. (MarLIN BIOTIC)
Mytilidae	Modiolus	modiolus	35	Comely, 1978 (Heller, 1990)
Mytilidae	Modiolus	modiolus	36	Zolotarev 1982; Table 1
Mytilidae	Modiolus	modiolus	61	Zolotarev, 1980 (Heller, 1990)
Mytilidae	Musculista	senhausius	2	Tanaka and Kikuchi, 1971 (Robertson, 1979)
Mytilidae	Musculista	senhousia	2	Crooks, JA, 1996.
Mytilidae	Musculista	senhousia	2	Power et al, 2006.
Mytilidae	Musculus	discors	6	Maximovich, N.V. 1984.
Mytilidae	Musculus	laevigatus	3	Maximovich, N.V. 1984.
Mytilidae	Musculus	niger	9	Petersen, G. H., 1977.
Mytilidae	Mytella	strigata	4	Baqueiro Cardenas, E, and DA Aranda, 2003.
Mytilidae	Mytella	strigata	18	Baqueiro Cardenas, E, and DA Aranda, 2003.
Mytilidae	Mytilaster	lineatus	3	Zaika, VE, 1973.
Mytilidae	Mytilisepta	keenae	15	Zolotarev, 1980 (Heller, 1990)
Mytilidae	Mytilus	coruscus	39	Zolotarev, 1980 (Heller, 1990)
Mytilidae	Mytilus	edulis	8	Maximovich, N.V. 1984.
Mytilidae	Mytilus	edulis	10	Milne and Dunnert, 1972 (Robertson, 1979)
Mytilidae	Mytilus	edulis	24	Seed 1974 in Emmett et al. 1991 (MSAP)
Mytilidae	Mytilus	edulis	24	Theisen, 1973 (Powell & Cummins, 1985)
Mytilidae	Mytilus	edulis	25	Thiesen, 1973 (MarLIN BIOTIC)
Mytilidae	Mytilus	edulis	10	Williamson, 1908 (Comfort, 1957)
Mytilidae	Mytilus	edulis	17	Zolotarev 1982; Table 1
Mytilidae	Mytilus	edulis	15	Zolotarev, 1980 (Heller, 1990)
Mytilidae	Mytilus	galloprovincialis	9	Zaika, VE, 1973.
Mytilidae	Mytilus	galloprovincialis	3	Abada-Boudjema, Y-M, and JC Dauvin, 1995.
Mytilidae	Mytilus	galloprovincialis	12	Zaika, VE, 1973.
Mytilidae	Perna	perna	2	Abada-Boudjema, Y-M, and JC Dauvin, 1995.
Mytilidae	Perna	viridis	3	Lee, 1985 (Heller, 1980) plus Power et al, 2004.
Nuculanidae	Jupiteria	minuta	7	Ansell & Parulekar, 1978 (Heller, 1990)
Nuculanidae	Jupiteria	minuta	8	Ansell et al, 1978 (Powell & Stanton 1985)
Nuculanidae	Jupiteria	minuta	6	Petersen, G. H., 1977.
Nuculanidae	Nuculana	pernula	5	Ansell et al, 1978 (Powell & Stanton, 1985)
Nuculanidae	Nuculana	pernula	10	Hutchings and Haedrich, 1984 (Callender and Powell, 1997)
Nuculanidae	Nuculana	pernula	9	Zolotarev 1982; Table 1
Nuculanidae	Nuculana	pernula	9	Zolotarev, 1980
Nuculidae	Ennucula	tenuis	13	Harvey, R and J.D. Gage, 1995.
Nuculidae	Ennucula	tenuis	6	Petersen, G. H., 1977.
Nuculidae	Lamellinucula	sulcata	17	Comfort, 1964 (Heller, 1990)
Nuculidae	Nucula	nitidosa	5	Rachor, E, 1976.
Nuculidae	Nucula	nitidosa	12	Wilson, JG 1992. Plus others (MarLIN BIOTIC)

Nuculidae	Nucula	nitidosa	10	Wilson, JG 1992. Plus others (MarLIN BIOTIC)
Nuculidae	Nucula	nitidosa	7	Wilson, JG 1992. Plus others (MarLIN BIOTIC)
Nuculidae	Nucula	nucleus	10	Chardy et al, 1984 (Callender & Powell, 1997)
Nuculidae	Nucula	nucleus	12	Comfort, 1964 (Heller, 1990)
Nuculidae	Nucula	paulula	2	Mukai, 1974 (Robertson, 1979)
Nuculidae	Nucula	turgida	11	Allen, 1952 (Comfort, 1957)
Nuculidae	Nucula	turgida	6	Davis, JP and JG Wilson, 1982 [1983].
Nuculidae	Nucula	turgida	8	Petersen, G. H., 1977.
Nuculidae	Nucula proxima	annulata	8	Cerrato, 1980 (Heller, 1990)
Nuculidae	Truncacila	insignis	9	Zolotarev, 1980 (Heller, 1990)
Ostreidae	Crassostrea	madrasensis	4	Somasekar et al, 1982 (Powell & Cummins, 1985)
Ostreidae	Crassostrea	virginiaca	6	Grave, 1933 (Comfort, 1957)
Ostreidae	Crassostrea	virginica	6	Comfort, 1957 (Heller, 1990)
Ostreidae	Crassostrea	virginica	2	Dame, 1976 (Roberston, 1979)
Ostreidae	Ostrea	edulis	20	Christensen & Dance, 1980 (Heller, 1990)
Ostreidae	Ostrea	edulis	12	Orton & Amirthalingam, 1930 (Comfort, 1957)
Ostreidae	Ostrea	edulis	20	Rodhouse, 1978 (Powell & Stanton, 1985)
Ostreidae	Ostrea	edulis	15	Yonge, CM 1960 plus others (MarLIN BIOTIC)
Ostreidae	Ostrea	lutaria	10	Allen, 1979 (Callender and Powell, 1997)
Ostreidae	Ostreola	lurida	na	MSAP
Pandoridae	Clidiophora	gouldi	3	Sanders, 1956 (Roberston, 1979)
Pandoridae	Pandora	pulchella	11	Zolotarev, 1980 (Heller, 1990)
Pectinidae	Adamusium	colbecki	40	Berkman, PA, et al, 2004.
Pectinidae	Adamusium	colbecki	18	Heilmayer et al, 2004.
Pectinidae	Adamusium	colbecki	10	Ralph & Maxwell, 1977 (Heller, 1990)
Pectinidae	Adamusium	colbecki	7	Ralph & Maxwell, 1977 (Powell & Cummins, 1985)
Pectinidae	Adamusium	colbecki	12	Stockton, 1984 (Callender and Powell, 1997)
Pectinidae	Adamusium	colbecki	7	Zolotarev 1982; Table 1
Pectinidae	Aequipecten	opercularis	8	Taylor & Venn, 1978 (Powell & Cummins, 1985)
Pectinidae	Aequipecten	opercularis	6	Williams & Dredge, 1981 (Heller, 1990)
Pectinidae	Amusium	balloti	4	Heald, 1978 (Powell & Cummins, 1985)
Pectinidae	Amusium	balloti	3	Heald, DI and N Caputi, 1981.
Pectinidae	Amusium	japonicum	4	Williams & Dredge, 1981 (Heller, 1990)
Pectinidae	Amusium	pleuronectes	2	Del Norte 1988 (Callender and Powell 1997)
Pectinidae	Argopecten	circularis	5	Baqueiro Cardenas, E, and DA Aranda, 2003.
Pectinidae	Argopecten	circularis	3	Baqueiro Cardenas, E, and DA Aranda, 2003.
Pectinidae	Argopecten	gibbus	2	Williams & Dredge, 1981 (Powell & Cummins, 1985)
Pectinidae	Argopecten	irradians	2	Gutsell, 1930 (Comfort, 1957)
Pectinidae	Argopecten	irradians	1	Sastray, 1979 (Heller, 1990)

Pectinidae	Argopecten	irradians	2	Williams & Dredge, 1981 (Powell & Cummins, 1985)
Pectinidae	Argopecten	irradians irradians	2	Brielj, VM and MK Krause, 1992.
Pectinidae	Argopecten	japonicum	2	Williams & Dredge, 1981 (Powell & Cummins, 1985)
Pectinidae	Chlamys	albidus	8	Zolotarev, 1980 (Heller, 1990)
Pectinidae	Chlamys	islandica	24	Brielj, VM and MK Krause, 1992.
Pectinidae	Chlamys	islandica	23	Vahl, 1981 (Powell & Cummins, 1985)
Pectinidae	Hinnites	giganteum	25	Haderlie and Abbott in Morris et al. 1980 (MSAP)
Pectinidae	Mimachlamys	varia	7	Conan & Shafee, 1978 (Powell & Cummins, 1985)
Pectinidae	Mizuhopecten	yessoensis	8	Bazykalova, 1934 (Comfort, 1957)
Pectinidae	Mizuhopecten	yessoensis	12	Ventilla, 1982 (Heller, 1990)
Pectinidae	Mizuhopecten	yessoensis	9	Zolotarev 1982; Table 1
Pectinidae	Mizuhopecten	yessoensis	13	Zolotarev, 1980
Pectinidae	Mizuhopecten	yessoensis	16	Zolotarev, 1980
Pectinidae	Mizuhopecten	yessoensis	9	Zolotarev, 1980
Pectinidae	Patinopecten	caurinus	NA	Elfstrom et al, 2005.
Pectinidae	Patinopecten	caurinus	15	Haynes & Hitz, 1971 (Powell & Cummins, 1985)
Pectinidae	Pecten	maximus	12	Zolotarev 1982; Table 1
Pectinidae	Pecten	maximus	20	Cerrato, 1980 (Heller, 1990)
Pectinidae	Pecten	maximus	22	Minchin 2003, Mason 1983 (MarLIN BIOTIC)
Pectinidae	Pecten	maximus	18	Tang, 1941 (Comfort, 1957)
Pectinidae	Pecten	meridionalis	16	Zolotarev 1982; Table 1
Pectinidae	Pecten	meridionalis	11	Fairbridge, 1953 (Comfort, 1957)
Pectinidae	Placopecten	magellanicus	12	Williams & Dredge, 1981 (Heller, 1990)
Pectinidae	Placopecten	magellanicus	12	Posgay, 1979 (Callender and Powell, 1997)
Pectinidae	Placopecten	magellanicus	12	Stevenson & Dickie (Powell & Cummins, 1985)
Pectinidae	Placopecten	magellanicus	12	Williams & Dredge, 1981 (Heller, 1990)
Pectinidae	Swiftopecten	swifti	15	Zolotarev, 1980 (Heller, 1990)
Periplomatidae	Bontaea	praetenua	5	Petersen, G. H., 1977.
Petricolidae	Petricolaria	pholadiformis	10	Duval, 1963
Pharidae	Ensis	arcuatus	18	MarLIN BIOTIC
Pharidae	Ensis	ensis	20	MarLIN BIOTIC
Pharidae	Ensis	ensis	5	Petersen, G. H., 1977.
Pharidae	Ensis	siliqua	18	MarLIN BIOTIC
Pharidae	Ensis	spp	20	MarLIN BIOTIC
Pharidae	Phaxas	pellucidus	3	Petersen, G. H., 1977.
Pharidae	Siliqua	patula	17	Fitch, JE, 1952.
Pharidae	Siliqua	patula	7	Lassuy and Simons, 1989. (MSAP)
Pharidae	Siliqua	patula	19	Zolotarev 1982; Table 1
Pharidae	Siliqua	pulchella	2	de Goeij et al, 2003.
Pholadidae	Barnea	candida	12	Pinn et al, 2005.
Pholadidae	Barnea	parva	12	Pinn et al, 2005.
Pholadidae	Penitella	penita	21	Evans, 1968 (Powell & Cummins, 1985)
Pholadidae	Penitella	penita	6	Rudy & Rudy, 1979.

Pholadidae	Pholas	dactylus	14	MarLIN BIOTIC
Pholadidae	Pholas	dactylus	12	Pinn et al, 2005.
Pholadidae	Zirfaea gabbi	pilsbryi	8	Evans, J W. 1970 (Rudy & Rudy, 1979.)
Pinnidae	Atrina	fragilis	10	MarLIN BIOTIC
Pinnidae	Atrina	maura	6	Baqueiro Cardenas, E, and DA Aranda, 2003.
Pinnidae	Atrina	rigida	4	Baqueiro Cardenas, E, and DA Aranda, 2003.
Pinnidae	Pinna	atropurpurea	12	Butler & Brewster, 1979 (Heller, 1990)
Pinnidae	Pinna	nobilis	18	Butler et al., 1993 (MarLIN BIOTIC)
Pinnidae	Pinna	nobilis	28	Garcia-march, JR and A Marquez-Aliaga, 2007.
Pinnidae	Pinna	nobilis	15	Katsanevakis 2005 Endangered species research
Psammobiidae	"Gari"	kazunensis	14	Zolotarev, 1980 (Heller, 1990)
Psammobiidae	Gobraeus	solida	14	Urban & Campos, 1994
Psammobiidae	Nuttallia	ezonis	40	Zolotarev, 1980
Psammobiidae	Nuttallia	ezonis	25	Zolotarev, 1980
Psammobiidae	Nuttallia	olivacea	24	Zolotarev, 1980
Psammobiidae	Nuttallia	olivacea	20	Zolotarev, 1980
Psammobiidae	Psammobia	fervens	8	Petersen, G. H., 1977.
Pteriidae	Meleagrina or Pteria	vulgaris	7	Comfort, 1957 (Heller, 1990)
Pteriidae	Meleagrina or Pteria	vulgaris	7	Herdman, 1904 (Comfort, 1957)
Pteriidae	Meleagrina or Pteria	vulgaris	8	Narayanan & Michael, 1968 (Powell & Cummins, 1985)
Pteriidae	Pinctada	imbricata	3	Marcano et al, 2005.
Pteriidae	Pinctada	imbricata	5	Verginelli, R and A Prieto, 1991.
Pteriidae	Pinctada	martensi	8	Narayanan & Michael, 1968 (Powell & Cummins, 1985)
Scrobiculariidae	Scrobicularia	plana	5	Coelho et al, 2006.
Scrobiculariidae	Scrobicularia	plana	18	Green, 1957 (Comfort, 1957)
Scrobiculariidae	Scrobicularia	plana	5	Hughes, 1970 (Robertson, 1979)
Semelidae	Abra	nitida	2	Buchanan & Warwick, 1974 (Robertston, 1979)
Semelidae	Abra	nitida	3	Josefson, 1982 (Callender and Powell, 1997)
Semelidae	Abra	nitida	3	Petersen, G. H., 1977.
Semelidae	Abra	nitida	4	Velosa et al, 2007.
Semelidae	Abra	nitida	9	Velosa et al, 2007.
Semelidae	Abra	ovata	4	Zaika, 1973 (Heller, 1990)
Semelidae	Abra	prismatic	1	Dauvin, 1986b (Callender and Powell, 1997)
Semelidae	Abra	prismatic	5	Petersen, G. H., 1977.
Semelidae	Abra	tenuis	2	Bachelet, G, 1989. (Dekker & Beukema, 1993)
Semelidae	Abra	tenuis	2.5	Dekker, R, and JJ Beukema, 1993.
Semelidae	Abra	tenuis	2	Gibbs, PE, 1984 (Dekker & Beukema, 1993)
Semelidae	Amphidesma	ventricosum	9	Rapson, 1952 (Comfort, 1957 & Powell & Cummins 1985)
Semelidae	Cumingia	tellinoides	4	Grave, 1933 (Comfort, 1957)
Semelidae	Endopleura	lubrica	2	Mukai, 1974 (Robertson, 1979)
Semelidae	Endopleura	lubrica	1	Tanaka and Kikuchi, 1971 (Robertson, 1979)
Semelidae	Semele	solida	13	Urban & Campos, 1994

Semelidae	Syndosmya	alba	2	Dauvin 1986a (Callender and Powell 1997)
Semelidae	Syndosmya	alba	2	Dauvin and Gentil, 1989. (MarLIN BIOTIC)
Semelidae	Syndosmya	alba	4	Petersen, G. H., 1977.
Semelidae	Syndosmya	alba	2	Lastra, M, J Palacio, and J Mora, 1993.
Semelidae	Syndosmya	ovata	4	Zaika, VE, 1973.
Semelidae	Theora	fragilis	2	Rainer, 1982 (Powell & Cummins, 1985)
Solecurtidae	Mesopleura	diviscus	5	Fraser 1967 (Powell & Stanton, 1985)
Solecurtidae	Tagelus	dombeii	15	Urban, 1996
Solemyidae	Solemya	reidi	3	Le Pennec, M, and PG Beninger, 2000.
Solenidae	Ensisolen	krustensterni	12	Zolotarev, 1980 (Heller, 1990)
Solenidae	Ensisolen	rosaceus	17	Fitch, JE, 1952.
Solenidae	Ensisolen	sicaru	17	Fitch, JE, 1952.
Solenidae	Solen	corneus	5	McLachlan, 1974 (Powell & Cummins, 1985)
Tellinidae	Angulus	tenuis	5	Faure, G, 1969.
Tellinidae	Angulus	tenuis	8	MacIntyre, 1970 (Dekker & Beukema, 1999)
Tellinidae	Angulus	tenuis	4	Petersen, G. H., 1977.
Tellinidae	Angulus	tenuis	5	Stephen, 1931 (Comfort, 1957)
Tellinidae	Angulus	tenuis	7	Trevallion, 1971 (Powell & Stanton, 1985)
Tellinidae	Cadella	lubrica	17	Zolotarev, 1980
Tellinidae	Cadella	lubrica	15	Zolotarev, 1980
Tellinidae	Eurytellina	alternata	3	Moore & Lopez, 1970 (Powell & Cummins, 1985)
Tellinidae	Fabulina	fabula	7	Petersen, G. H., 1977.
Tellinidae	Fabulina	fabula	5	Salzwedel, 1979 (MarLIN BIOTIC)
Tellinidae	Heteromacoma	inquinata	5	Lees, DC, 2005 (unpub) Assessment of Bivalve Recovery, Prince William Sound
Tellinidae	Macoma	balthica	5	Lees, DC, 2005 (unpub) Assessment of Bivalve Recovery, Prince William Sound
Tellinidae	Macoma	balthica	2	Burke & Mann, 1974 (Robertson, 1979)
Tellinidae	Macoma	balthica	6	Chambers and Milne, 1975 (Robertson, 1979)
Tellinidae	Macoma	balthica	9	Evans and Tallmark, 1977 (Powell and Stanton, 1985)
Tellinidae	Macoma	balthica	10	Gilbert, 1973 (MarLIN BIOTIC)
Tellinidae	Macoma	balthica	30	Gilbert, MA, 1973.
Tellinidae	Macoma	balthica	6	Maximovich, N.V. 1984.
Tellinidae	Macoma	balthica	10	Salzwedel, 1979 (MarLIN BIOTIC)
Tellinidae	Macoma	balthica	7	Warwick and Price, 1975 (Powell & Stanton, 1985)
Tellinidae	Macoma	balthica	10	Wolff and deWolf, 1977 (Robertson, 1979)
Tellinidae	Macoma	balthica	18	Zolotarev, 1980 (Heller, 1990)
Tellinidae	Macoma	calcarea	17	Petersen, 1978 (Heller, 1990)
Tellinidae	Macoma	calcarea	5	Petersen, G. H., 1977.
Tellinidae	Macoma	contabulata	15	Zolotarev, 1980 (Heller, 1990)
Tellinidae	Macoma	litoralis	5	McLachlan, 1974 (Powell & Cummins, 1985)

Tellinidae	Macoma	middendorffii	24	Zolotarev, 1980 (Heller, 1990)
Tellinidae	Macomona	deltoidalis	4	Rainer, 1982 (Powell & Cummins, 1985)
Tellinidae	Macomona deltoidalis	deltoides	4	Robertson, in press (Robertson, 1979) [not found]
Tellinidae	Megangulus	venulosa	31	Zolotarev, 1980
Tellinidae	Megangulus	venulosa	29	Zolotarev, 1980
Tellinidae	Megangulus	venulosa	26	Zolotarev, 1980
Tellinidae	Megangulus	zyonoensis	61	Zolotarev, 1980 (Heller, 1990)
Tellinidae	Merisca	martinicensis	2	Penzias, 1969 (Powell and Stanton, 1985)
Tellinidae	Merisca	piratica	2	de Goeij et al, 2003.
Tellinidae	Serratina	capsoides	3	de Goeij et al, 2003.
Teredinidae	Bankia	gouldi	1	Hoagland, 1986 (Heller, 1990)
Teredinidae	Bankia	setacea	2	Hill, C. L. and C. A. Kofoid, eds. 1927 (Rudy & Rudy, 1979.)
Teredinidae	Teredo	bartschi	1	Hoagland, 1986 (Heller, 1990)
Teredinidae	Teredo	navalis	2	Grave, 1928 (Comfort, 1957)
Teredinidae	Teredo	navalis	1	Hoagland, 1986 (Heller, 1990)
Thraciidae	Thracia	papyracea	4	Petersen, G. H., 1977.
Thraciidae	Trigonothracia	jinxigae	1	Morton, B, 1995.
Thyasiridae	Thyasira	flexuosa	3	Lopez-Jamar et al, 1987 (Callender and Powell, 1997)
Thyasiridae	Thyasira	flexuosa	1	Lopez-Jamar et al, 1987 (Heller, 1990)
Thyasiridae	Thyasira	flexuosa	7	Petersen, G. H., 1958.
Thyasiridae	Thyasira	flexuosa	3	Velosa et al, 2007.
Thyasiridae	Thyasira	gouldi	10	Blacknell and Ansell, 1975 (MarLIN BIOTIC)
Tridacnidae	Tridacna	gigas	60	Watanabe et al, 2004.
Tridacnidae	Tridacna	gigas	200	Watanabe et al, 2004.
Turtoniidae	Turtonia	minuta	2	Tokmakova et al, 2006.
Ungulinidae	Felaniella	usta	9	Zolotarev, 1980 (Heller, 1990)
Veneridae	Anomalocardia	squamosa	5	Morton, 1978 (Powell & Stanton, 1985)
Veneridae	Callista	chione	40	Forster, 1981 (Powell & Cummins, 1985)
Veneridae	Callista	chione	16	Leontarakis, PK, and C.A. Richardson, 2005.
Veneridae	Callista	chione	17	Metaxatos, A, 2004.
Veneridae	Callista	chione	18	Metaxatos, A, 2004.
Veneridae	Callista	chione	30	Metaxatos, A, 2004.
Veneridae	Callithaca	adamsi	29	Zolotarev, 1980 (Heller, 1990)
Veneridae	Callithaca	tenerrima	10	Fitch, JE, 1952.
Veneridae	Chamelea	gallina	9	Boltacheva & Mazlumyan, 2003
Veneridae	Chamelea	gallina	9	Boltacheva, NA and SA Mazlumyan, 2003.
Veneridae	Chamelea	gallina	8	Cerrato, 1980 (Heller, 1990)
Veneridae	Chamelea	gallina	25	Duineveld et al., 2002
Veneridae	Chamelea	gallina	5	Gaspar et al, 2004.
Veneridae	Chamelea	striatula	10	Guillou, J and PG Sauriau, 1985.
Veneridae	Chamelea	striatula	8	Petersen, G. H., 1958.
Veneridae	Chione	cancellata	5	Baqueiro Cardenas, E, and DA Aranda, 2003.
Veneridae	Chione	cancellata	6	Moore and Lopez, 1969 (Robertson, 1979)

Veneridae	<i>Chione</i>	<i>undatella</i>	2	Baqueiro Cardenas, E, and DA Aranda, 2003.
Veneridae	<i>Chione</i>	<i>undatella</i>	4	Baqueiro Cardenas, E, and DA Aranda, 2003.
Veneridae	<i>Chionista</i>	<i>cortezi</i>	8	Schone et al, 2002.
Veneridae	<i>Chionista</i>	<i>fluctifraga</i>	16	Schone et al, 2002.
Veneridae	<i>Dosinella</i>	<i>angulosa</i>	26	Zolotarev, 1980 (Heller, 1990)
Veneridae	<i>Dosinella</i>	<i>hepatica</i>	9	Hanekom, N, 1986.
Veneridae	<i>Dosinella</i>	<i>hepatica</i>	6	McLachlan, 1974 (Powell & Cummins, 1985)
Veneridae	<i>Dosinia</i>	<i>elegans</i>	3	Moore & Lopez, 1970 (Powell & Cummins, 1985)
Veneridae	<i>Dosinia</i>	<i>lupinus</i>	8	Petersen, G. H., 1958.
Veneridae	<i>Dosinia</i>	<i>lupinus</i>	15	Tunberg, B, 1983a. <i>Sarsia</i> 68: 33-40
Veneridae	<i>Dosinia</i>	<i>ponderosa</i>	6	Baqueiro Cardenas, E, and DA Aranda, 2003.
Veneridae	<i>Eucallista</i>	<i>purpurata</i>	30	Morosan and Orensan 2004
Veneridae	<i>Gemma</i>	<i>gemma</i>	4	Bradley & Cooke, 1959 (Powell & Cummins, 1985)
Veneridae	<i>Gemma</i>	<i>gemma</i>	1	Sellmer, 1967 (Heller, 1990)
Veneridae	<i>Liocyma</i>	<i>fluctuosa</i>	21	Zolotarev, 1980
Veneridae	<i>Marcia</i>	<i>japonica</i>	11	Zolotarev, 1980
Veneridae	<i>Marcia</i>	<i>japonica</i>	27	Zolotarev, 1980 (Heller, 1990)
Veneridae	<i>Marcia</i>	<i>japonica</i>	25	Zolotarev, 1980 (Heller, 1990)
Veneridae	<i>Marcia</i>	<i>opima</i>	3	Mane, 1974 (Powell & Cummins, 1985)
Veneridae	<i>Megapitaria</i>	<i>aurantiaca</i>	7	Baqueiro Cardenas, E, and DA Aranda, 2003.
Veneridae	<i>Megapitaria</i>	<i>squalida</i>	3	Baqueiro Cardenas, E, and DA Aranda, 2003.
Veneridae	<i>Megapitaria</i>	<i>squalida</i>	6	Baqueiro Cardenas, E, and DA Aranda, 2003.
Veneridae	<i>Mercenaria</i>	<i>campechiensis</i>	5	Baqueiro Cardenas, E, and DA Aranda, 2003.
Veneridae	<i>Mercenaria</i>	<i>mercenaria</i>	17	Belden, 1912 (Comfort, 1957)
Veneridae	<i>Mercenaria</i>	<i>mercenaria</i>	15	Cerrato, 1980 (Heller, 1990)
Veneridae	<i>Mercenaria</i>	<i>mercenaria</i>	9	Hibbert, 1976 (Roberston, 1979)
Veneridae	<i>Mercenaria</i>	<i>mercenaria</i>	40	Hopkins, 1930 (Comfort, 1957)
Veneridae	<i>Mercenaria</i>	<i>mercenaria</i>	9	Kennish, 1980 (Heller, 1990)
Veneridae	<i>Mercenaria</i>	<i>mercenaria</i>	40	Loesch and Haven, 1973 (Powell & Stanton, 1985)
Veneridae	<i>Mercenaria</i>	<i>mercenaria</i>	46	Peterson, CH, 1986.
Veneridae	<i>Mercenaria</i>	<i>stimpsoni</i>	40	Zolotarev, 1980 (Heller, 1990)
Veneridae	<i>Notochione</i>	<i>jedoensis</i>	15	Zolotarev, 1980 (Heller, 1990)
Veneridae	<i>Nutricola</i>	<i>lordi</i>	2	Zaika, VE, 1973.
Veneridae	<i>Nutricola</i>	<i>tantilla</i>	1	Rudy & Rudy, 1979.
Veneridae	<i>Nutricola</i>	<i>tantilla</i>	1	Kabat, AR, 1985.
Veneridae	<i>Pachydesma</i>	<i>stultorum</i>	14	Cerrato, 1980 (Heller, 1990)
Veneridae	<i>Pachydesma</i>	<i>stultorum</i>	12	Coe, 1947 (Comfort, 1957)
Veneridae	<i>Pachydesma</i>	<i>stultorum</i>	35	Fitch, JE, 1952.
Veneridae	<i>Pachydesma</i>	<i>stultorum</i>	20	Weymouth, 1923 (Comfort, 1957)
Veneridae	<i>Paphia</i>	<i>euglypta</i>	14	Zolotarev, 1980 (Heller, 1990)
Veneridae	<i>Paphia</i>	<i>laterisulca</i>	3	Mane & Nagabhushanam, 1979 (Powell & Cummins, 1985)
Veneridae	<i>Pectunculus</i>	<i>exoleta</i>	12	Tunberg, B, 1983b. <i>Sarsia</i> 68: 41-45.

Veneridae	Pectunculus	exoleta	9	Petersen, G. H., 1958.
Veneridae	Pectunculus	exoleta	7	Tunberg, 1984b (Callender and Powell, 1997)
Veneridae	Pectunculus	exoleta		Tunberg, B, 1983a. Sarsia 68: 33-41
Veneridae	Petunculus	exoleta	7	Comfort, 1964 (Heller, 1990)
Veneridae	Phacosoma	japonicum	10	Tanabe, K, 1988.
Veneridae	Protothaca	antiqua	12	Urban, 1996
Veneridae	Protothaca	laciniata	10	Fitch, JE, 1952.
Veneridae	Protothaca	staminea	10	Lees, DC, 2005 (unpub) Assessment of Bivalve Recovery, Prince William Sound
Veneridae	Protothaca	staminea	16	Chew and Ma 1987, Fraser and Smith 1928; Abbott 1974; Chew and Ma 1987 all in Emmett et al. 1991 (MSAP)
Veneridae	Protothaca	staminea	10	Fitch, JE, 1952.
Veneridae	Protothaca	staminea	8	Harrington, RJ 1985 dissertation (Takesue, RK and A. van Geen, 2004.)
Veneridae	Protothaca	staminea	13	Paul et al, 1976 (Powell & Cummins, 1985)
Veneridae	Protothaca	staminea	10	Schmidt, RR. and JE. Warme, 1969.
Veneridae	Protothaca	staminea	16	Zolotarev 1982; Table 1
Veneridae	Protothaca	thaca	17	Urban & Campos, 1994
Veneridae	Retrotapes	exalbida	70	Lomovasky et al, 2002
Veneridae	Ruditapes	japonica	3	Ohba, 1959 (Powell and Stanton, 1985)
Veneridae	Ruditapes	japonica	10	Frey 1971 in Emmett et al. 1991 (MSAP) (syn w V. philippinarum)
Veneridae	Ruditapes	philippinarum	2	Yap, 1977 (Powell & Cummins, 1985)
Veneridae	Saxidomus	brevisiphonata	63	Zolotarev, 1980
Veneridae	Saxidomus	brevisiphonata	76	Zolotarev, 1980 (Heller, 1990)
Veneridae	Saxidomus	gigantea	11	Zolotarev 1982; Table 1
Veneridae	Saxidomus gigantea	giganteus	20	Fraser, CM, 1929. (Rudy & Rudy, 1979.)
Veneridae	Saxidomus gigantea	giganteus	20	MSAP
Veneridae	Tapes	aurea	5	Hibbert, 1976 (Roberston, 1979)
Veneridae	Tawera	gayi	15	Lomovasky et al, 2005
Veneridae	Tawera	mawsoni	15	Lomovasky et al, 2005
Veneridae	Timoclea	micra	1.5	Mukai, 1974 (Robertson, 1979)
Veneridae	Timoclea	micra	1	Tanaka and Kikuchi, 1971 (Robertson, 1979)
Veneridae	Tivela	mactroides	2	Arriehe, D, and A Prieto, 2006.
Veneridae	Venerupis	pullastra	9	Cerrato, 1980 (Heller, 1990)
Veneridae	Venerupis	pullastra	8	Quayle, 1952 (Comfort, 1957)
Veneridae	Venerupis	pullastra	10	Johannessen, 1973 (Roberston, 1979)
Veneridae	Venerupis	rhomboides	10	Menesguen and Dreves, 1987 (Callender and Powell, 1997)
Veneridae	Venerupis	senegensis	10	MarLIN BIOTIC = same as V. pullastra, V. saxatilis, V. corrugata
Vesicomyidae	Calyptogena	magnifica	11	Jones, 1983 (Heller, 1990)
Vesicomyidae	Calyptogena	magnifica	50	Lutz et al, 1988 (Callender and Powell, 1997)
Vesicomyidae	Calyptogena	magnifica	37	Rao and Roux, 1984 (Callender and Powell, 1997)

Vesicomyidae	Calyptogena	magnifica	37	Roux et al, 1985 (Callender and Powell, 1997)
Yoldiidae	Cnesterium	notabilis	17	Nakaoka, M and S Matsui, 1994.
Yoldiidae	Megayoldia	thraciaeformis	15	Hutchings and Haedrich, 1984 (Callender and Powell, 1997)
Yoldiidae	Yoldia	limatula	4	Lewis et al, 1982 (Powell & Cummins, 1985)
Yoldiidae	Yoldia	limatula	4	Nakaoka, M and S Matsui, 1994.
Yoldiidae	Yoldia	limatula	3	Sanders, 1956 (Roberston, 1979)

## REFERENCES

### **Secondary sources (all are fully referenced compilations of the literature):**

- Callender, WR and EN Powell, 1997. Autochthonous death assemblages from chemoautotrophic communities at petroleum seeps: palaeoproduction, energy flow, and implications for the fossil record. *Historical Biology* 12: 165-198.
- Comfort, A, 1957. The duration of life in molluscs. *Proc Malacological Soc London* 32: 219-240.
- Heller, J, 1990. Longevity in molluscs. *Malacologia* 31: 259-295.
- MarLIN BIOTIC (Marine Life Information Network, Biological Traits Information Catalogue). [www.marlin.ac.uk/biotic/](http://www.marlin.ac.uk/biotic/), accessed September 2008.
- MSAP (Marine Species with Aquaculture Potential off the coast of Oregon and Pacific Northwest), Species Profiles. <http://hmsc.oregonstate.edu/projects/msap/PS/masterlist/index.html>, accessed September 2008.
- Powell, EN and H Cummins, 1985. Are molluscan maximum life spans determined by long-term cycles in benthic communities? *Oecologia* 67: 177-182.
- Powell, EN and RJ Stanton, Jr, 1985. Estimating biomass and energy flow of molluscs in palaeo-communities. *Palaeontology* 28: 1-34.
- Robertson, AI, 1979. The relationship between annual production:biomass ratios and lifespans for marine macrobenthos. *Oecologia* 38: 193-202.

### **Primary sources (secondary source of information, if primary not examined directly):**

- Abada-Boudjema, Y-M, and JC Dauvin, 1995. Recruitment and life span of two natural mussel populations *Perna perna* (Linnaeus) and *Mytilis galloprovincialis* (Lamarck) from the Algeria coast. *Journal of Molluscan Studies* 61: 467-481.
- Abele, D, J Strahl, T Brey, and EER Philipp, 2008. Imperceptible senescence: ageing in the ocean quahog *Arctica islandica*. *Free Radical Research* 42: 474-480.
- Allen, 1952 (Comfort, 1957)
- Allen, 1979 (Callender and Powell 1997)
- Amaro, T, G Duineveld, M Bergman, and R Witbaard, 2003. Growth variations in the bivalve *Mya truncata*: a tool to trace changes in the Frisian Front macrofauna (southern North Sea)? *Helgoland Marine Research* 57: 132–138.
- Ansell & Parulekar, 1978 (Heller, 1990)
- Ansell et al, 1972 (Powell & Cummins, 1985)
- Ansell et al, 1978 (Powell & Stanton 1985)
- Ansell, 1983 (Heller, 1990)
- Anwar, NA, CA Richardson, and R Seed, 1990. Age determination, growth rate and populations structure of the horse mussel *Modiolus modiolus*. *Journal of the Marine Biological Association of the U. K.* 70: 441-457.
- Arriehe, D, and A Prieto, 2006. Population parameters of the trigonal tivela *Tivela mactroides* (Bivalvia: Veneridae) from Caicara Beach, Anzoategui, Venezuela. *Ciencias Marinas* 32: 285-296.
- Bachelet, G 1989. Recruitment in *Abra tenuis* (Montagu)(Bivalvia, Semelidae), a species with direct development and a protracted meiobenthic phase. In JS Ryland and PA Tyler,

- Reproduction , genetics and distributions of marine organisms. Fredensborg, Olsen & Olsen
- Baqueiro Cardenas, E and DA Aranda, 2003. Patroes en la biologia poblacional de moluscos de importancia comercial en Mexico. Revista de Biología Tropical 51 (suppl 4): 97-107 (Mem. 30th Reunion).
- Bayed, A, 1998. Growth variability of *Donax trunculus* in the coasts of Morocco. Cahiers Options Mediterraneennes 35: 11-23.
- Bazykalova, 1934 (Comfort, 1957)
- Belden, 1912 (Comfort, 1957)
- Berkman, PA, R Cattaneo-Vietti, M Chiantore, and C Howard-Williams, 2004. Polar emergence and the influence of increased sea-ice extent on the Cenozoic biogeography of pectinid molluscs in Antarctic coastal areas. Deep-Sea Research II 51: 1839-1855.
- Beukema JJ (1989) Bias in estimates of maximum life span, with an example of the edible cockle, *Cerastoderma edule*. Neth. J. Zool. 39: 79-85.
- Blacknell, W. M. and Ansell, A. D., 1975. Features of the reproductive cycle of an arctic bivalve from a Scottish sea loch. Marine Ecology, Pubblicazioni Della Stazione Zoologica Di Napoli I, 39, Suppl., 26-52.
- Bodoy, A, 1980. Croissance et variations de la composition biochimique du bivalve *Spisula subtruncata* (Da Costa) dans le golfe de Marseille. Thethys 9(4): 345-354.
- Boltacheva, NA and SA Mazlumyan, 2003. The growth and longevity of *Chamelea gallina* (Mollusca, Veneridae) in the Black Sea. Vestnik Zoologii 37(3): 71-74.
- Bourne and Smith, 1972 (Powell and Cummins, 1985)
- Boyden 1972 (Powell and Stanton, 1985)
- Bradley and Cooke, 1959 (Powell and Cummins, 1985)
- Brethes, JC, G Desrosiers, and G Fortin, 1986. Croissance et production due bivalve *Mesodesma arctatum* (Conrad) sur la cote nord du golf du Saint-Laurent. Canadian Journal of Zoology 64: 1914-1919.
- Brielj, VM and MK Krause, 1992. Resource allocation and population genetics of the bay scallop, *Argopecten irradians irradians*: effects of age and allozyme heterozygosity on reproductive output. Marine Biology 113: 253-261.
- Buchanan and Warwick, 1974 (Robertston, 1979)
- Burke and Mann, 1974 (Robertson, 1979)
- Butler and Brewster, 1979 (Heller, 1990)
- Butler, A.J., Vicente, N. and de Gaulejac, B., 1993. Ecology of the pteroid bivalves *Pinna bicolor* Gmelin and *Pinna nobilis* Linnaeus. Marine Life, 3, 37-45.
- Cerrato, 1980 (Heller, 1990)
- Cerrato, RM, and DL Keith, 1992. Age structure, growth, and morphometric variations in the Atlantic surf clam, *Spisula solidissima*, from estuarine and inshore waters. Marine Biology (Berlin) 114: 581-593.
- Chambers and Milne 1975 (Robertson 1979)
- Chardy et al. 1984 (Callender and Powell 1997)
- Chase, ME, and RC Bailey, 1999. The ecology of the zebra mussel (*Dreissena polymorpha*) in the Lower Great Lakes of North America: 1. Population dynamics and growth. Journal of Great Lakes Research 25: 107-121.
- Christensen and Dance, 1980 (Heller, 1990)
- Coe and Fox, 1942 (Comfort, 1957)

- Coe, 1947 (Comfort, 1957)
- Coe, 1955 (Powell and Cummins, 1985)
- Coelho, JP, M Rosa, E Pereira, A Duarte, and MA Pardal, 2006. Pattern and annual rates of *Scrobicularia plana* mercury bioaccumulation in a human induced mercury gradient (Ria de Aveiro, Portugal). *Estuarine Coastal and shelf science* 69: 629-635.
- Coen, LD, and RE Grizzle, 2007. The importance of habitat created by molluscan shellfish to managed species along the Atlantic coast of the United States. *Atlantic States Marine Fisheries Commission, Habitat Management Series* 8
- Cole, 1956 (Comfort, 1957)
- Comely, 1978 (Heller, 1990)
- Comfort, 1964 (Heller, 1990)
- Conan and Shafee, 1978 (Powell and Cummins, 1985)
- Crooks, JA, 1996. The population ecology of an exotic mussel, *Musculista senhousia*, in a southern California Bay. *Estuaries* 19: 45-50. [San Diego]
- Dame 1976 (Roberston, 1979)
- Dauvin, 1986a (Callender and Powell, 1997)
- Dauvin, 1986b (Callender and Powell, 1997)
- Dauvin, J-C. and Gentil, F., 1989. Long-term changes in populations of subtidal bivalves (*Abra alba* and *Abra prismatica*) from the Bay of Morlaix (Western English Channel). *Marine Biology*, 103, 63-73.
- Davis, JP and JG Wilson, 1982 [1983]. The population structure and ecology of *Nucula turgida* (Leckenby and Marshall) in Dublin Bay. *Progress in Underwater Science* 8: 53-60.
- de Goeij, P, M Lavaleye, GB Pearson and T. Piersma, 2003. Seasonal changes in the macrozoobenthos of a tropical mudflat. NIOZ Report 2003-4 [Australia]
- Dekker, R, and JJ Beukema, 1993. Dynamics and growth of a bivalve, *Abra tenuis*, at the northern edge of its distribution. *Journal of the Marine Biological Association of the U. K.* 73: 497-511
- Del Norte, 1988 (Callender and Powell, 1997)
- Deval, MC and D Gokturk, 2008. Population structure and dynamics of the cut trough shell *Spisula subtruncata* (da Costa) in the Sea of Marmara, Turkey. *Fisheries Research* (Amsterdam) 89: 241-247.
- Duineveld, G., M. Bergman, R. Witbaard and T. Amaro. 2002. Population dynamics of large bivalves in offshore southern North Sea (BIVALFF). Annual Report of the Royal Nederlands Instituut voor Onderzoek der Zee, p. 14-15.
- Duval, D.M., 1963. The biology of *Petricola pholadiformis* Lamarck (Lamellibranchiata: Petricolidae). *Proceedings of the Malacological Society*. 35: 89-100.
- Elfstrom, CM, CT Smith, KC Jones, and JE Seeb, 2005. Characterization of 16 polymorphic microsatellite loci in weathervane scallop *Patinopecten caurinus*. *Molecular Ecology Notes* 5: 514-516.
- Evans and Tallmark, 1977. (Powell and Stanton, 1985)
- Evans, 1968. (Powell and Cummins, 1985)
- Evans. J W. 1970. Palaeontological implications of a biological study of rock-boring clams (Family Pholadidae) In Trace fossils. ed. T.P.Crimes and J. Harper.
- Fahy, E, J. Carroll, M. O'Toole, and J. Hickey, 2003. A preliminary account of fisheries for the surf clam *Spisula solidissima* (L) (Mactacea) in Ireland. *Fisheries Bulletin (Dublin)* Volume: 21 Pages: i-iv, 1-27

- Fairbridge, 1953. (Comfort, 1957)
- Faure, G, 1969. Ecologie et croissance de *Tellina tenuis* Da Costa sur les cotes de la Charente-maritime. *Tethys* 1: 383-393.
- Feder et al, 1979. (Powell and Cummins, 1985)
- Fiori, S and O Defeo, 2006. Biogeographic patterns in life-history traits of the yellow clam, *Mesodesma mactroides*, in sandy baches of South America. *Journal of Coastal Research* 22: 872-880.
- Fitch, JE, 1952. Common marine bivalves of California. *Fish Bulletin No 90* (Calif. State Dept. Fish and Game)
- Forster, 1981 (Powell and Cummins, 1985)
- Franz, 1973 (Powell and Stanton, 1985)
- Franz, DR, 2001. Recruitment, survivorship, and age structure of a New York ribbed mussel population (*Geukensia dimissa*) in relation to shore level: a nine year study. *Estuaries* 24: 319-327.
- Fraser 1967 (Powell and Stanton, 1985)
- Fraser, 1931 (Comfort, 1957)
- Fraser, C. McLean, 1929. The spawning and free swimming larval periods of *Saxidomus* and *Paphia*. *Transactions of the Royal Society of Canada* (4)23:195-98.
- Fretter, V. and Graham, A., 1964. Reproduction. In *Physiology of Mollusca*, vol. 1, (ed. K.M. Wilbur and C.M. Yonge). New York: Academic Press.
- Gage, 1968 (Heller, 1990)
- Galinou-Mitsoudi, S and AI Sinis, 1995. Age and growth of *Lithophaga lithophaga* (Linnaeus, 1758) (Bivalvia: Mytilidae), based on annual growth lines in the shell. *Journal of Molluscan Studies* 61: 435-453.
- Garcia-March, JR and A Marquez-Aliaga, 2007. *Pinna nobilis* L, 1758 age determination by internal shell register. *Marine Biology* 151: 1077-1085.
- Gaspar, MG, AM Pereira, P Vasconcelos, and CC. Monteiro, 2004. Age and growth of *Chamelea gallina* from the Algarve Coast (southern Portugal): influence of seawater temperature and gametogenic cycle on growth rate. *Journal of Molluscan Studies* 70: 371 - 377.
- George and Nair, 1974 (Powell and Cummins, 1985)
- Gibbs, PE 1984. The population cycle of the bivalve *Abra tenuis* and its mode of reproduction. *Journal of the Marine Biological Association of the U. K.* 64: 791-800.
- Gilbert, M.A., 1973. Growth rate, longevity and maximum size of *Macoma balthica* (L.). *Biological Bulletin of the Marine Laboratory, Woods Hole*, 145, 119-126.
- Grave, 1928 (Comfort, 1957)
- Grave, 1933 (Comfort, 1957)
- Green, 1957 (Comfort, 1957)
- Gribben, PE, and RG Creese, 2005. Age, growth, and mortality of the New Zealand geoduck clam, *Panopea zelandica* (Bivalvia: Hiatellidae) in two North Island populations. *Bulletin of Marine Science* 77: 119-135.
- Guillou, J and PG Sauriau, 1985. Some observations on the biology and the ecology of a *Venus striatula* population in the Bay of Douarnenez, Brittany. *Journal of the Marine Biological Association of the U.K.* 65: 889-900.
- Gutsell, 1930 (Comfort, 1957)

- Hall, J.J., 1984. Production of immature *Corbicula fluminea* (Bivalvia:Corbiculidae), in Lake Norman, North Carolina. *The Nautilus* 98(4):153-159.
- Hancock, D. R., T. F. Gaumer, G. B. Willeke, G. P. Robart, and J. Flynn. 1979. Subtidal clam populations: Distribution, abundance, and ecology. Oregon State University Sea Grant Program. ORESU-T-79-002.
- Hanekom, N, 1986. Growth and somatic production estimates of *Dosinia hepatica* (Lamarck) (Mollusca: Bivalvia) in the Swartkops Estuary, South Africa. *South African Journal of Zoology* 21: 325-330.
- Hanks, RW, ad DB Packer, 1985. A new species of *Sphenia* (Bivalvia: Myidae) from the Gulf of Maine. *Veliger* 27: 320-330.
- Harrington, RJ 1985 dissertation, cited by Takesue, RK and A. van Geen, 2004. Mg/Ca, Sr/Ca, and stable isotopes in modern and Holocene *Protothaca staminea* shells from a northern California coastal upwelling region. *Geochimica et Cosmochimica Acta* 68: 3845
- Harvey, R. and J.D. Gage, 1995. Reproduction and recruitment of *Nuculoma tenuis* (Bivalvia: Nuculoida) from Loch Etive, Scotland. *Journal of Molluscan Studies* 1995 61: 409-419
- Haynes and Hitz, 1971 (Powell and Cummins, 1985)
- Heald, 1978 (Powell and Cummins, 1985)
- Heald, DI and N Caputi, 1981. Some aspects of growth, recruitment and reproduction in the southern saucer scallop, *Amusium balloti* (Bernardi, 1861) in Shark Bay, Western Australia. *Western Australian Marine Research Laboratories Fisheries Research Bulletin* No 25: 1-31.
- Heilmayer O, Brey T, Chiantore M, Cattaneo-Vietti R, Arntz WE (2003) Age and productivity of the Antarctic scallop, *Adamussium colbecki*, in Terra Nova Bay (Ross Sea, Antarctica). *Journal of Experimental Marine Biology and Ecology* 288:239–256
- Herdman, 1904 (Comfort, 1957)
- Hibbert 1976 (Roberston 1979)
- Hill, C. L. and C. A. Kofoid, eds. 1927. Marine Borers and their Relation to Marine Construction on the Pacific Coast. Final Report, San Francisco Bay Marine Piling Committee, San Francisco. 357 pp. C. A. Kofoid and R. C. Miller: Biological Section, pp. 188-343.
- Hoagland, 1986 (Heller, 1990)
- Hoenig, 1983 (Powell and Cummins, 1985)
- Holmes, SP and N Miler 2006. Aspects of the ecology and population genetics of the bivalve *Corbula gibba*. *Mar Ecol Progr Ser* 315: 129-140.
- Hopkins, 1930 (Comfort, 1957)
- Hornbach, DJ, 1992. Life history traits of a riverine population of the Asian clam *Corbicula fluminea*. *Amer Midland Nat* 127: 248-257. [Charlottesville, VA stream]
- Hrs-Brenko, M., 2006. The basket shell, *Corbula gibba* Olivi, 1792 (Bivalve Mollusks) as a species resistant to environmental disturbances: a review. *Acta Adriatica* 47(1): 49-64.
- Hughes, 1970 (Robertson, 1979)
- Hughes, SE, and N Bourne, 1981. Stock assessment and life history of a newly discovered Alaska surf clam (*Spisula polynyma*) resource in the southeastern Bering Sea. *Canad J Fisheries and Aquatic Sciences* 38: 1173-1181.
- Hutchings and Haedrich, 1984 (Callender and Powell, 1997)
- Ituarte, CF, 1984. Aspectos biológicos de las poblaciones de *Corbicula largillierti* Philippi (Mollusca, Pelecypoda) en el Rio de la Plata. *Revista del Museo de La Plata, Sección Zoológica* 13 (143): 231-247. [Argentina]

- Jensen, J., 1990. Increased abundance and growth of the suspension feeding bivalve *Corbula gibba* in a shallow part of the eutrophic Limfjord Denmark. Netherlands Journal of Sea Research, 27, 101-108.
- Jiminez, M, A Prieto, L Riz, B Marin, and T Allen, 2004. Distribucion de tallas, crecimiento y mortalidad de *Anadara notabilis* (Bivalvia: Arcidae) en la Bahia de Mochima, Estado Sucre, Venezuela. Boletin del Instituto Oceanografico de Venezuela Universidad de Oriente 43 (1-2): 41-47.
- Johannessen, 1973 (Roberston, 1979)
- Jones et al, 1978 (Powell and Cummins, 1985)
- Jones, 1980 (Powell and Stanton, 1985)
- Jones, 1983 (Heller, 1990)
- Jones, DS, IR Quitmyer, and CFT Andrus, 2004. Seasonal shell growth and longevity in *Donax variabilis* from northwestern Florida: evidence from oxygen isotopes. Journal of Shellfisheries Research 23: 707-714
- Josefson, 1982 (Callender and Powell, 1997)
- Kabat, AR, 1985. The allometry of brooding in *Transennella tantilla* (Gould) (Mollusca: Bivalvia). Journal of Experimental marine Biology Ecology 91: 271-279.
- Katsanevakis, S, 2005. Population ecology of the endangered fan mussel *Pinna nobilis* in a marine lake. Endangered Species Research 1;1-9.
- Kennish, 1980 (Heller, 1990)
- Kim et al, 1983 (Callender and Powell, 1997)
- Kornobis, 1977 (Callender and Powell, 1997)
- Korringa, P., 1952. Recent advances in oyster biology. Quarterly Review of Biology, 27, 266-308 and 339-365.
- Kunitzer, A, 1989 Factors affecting the population dynamics of *Amphiura filiformis* (Echinodermata: Ophiuroidae) and *Mysella bidentata* (Bivalvia: Galeommatacea) in the North Sea, 395-406. In JS Ryland and PA Tyler, Reproduction , genetics and distributions of marine organisms. Fredensborg, Olsen and Olsen.
- Lassuy, D. R., and D. Simons. 1989. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Pacific Northwest) --Pacific razor clam. U. S. Fish and Wildlife Service Biological Report 82(11.89). U. S. Army Corps of Engineers, TR-EL-824, 16 pp.
- Lastra, M, A Sanchez, and J Mora, 1993. Population dynamics and secondary production of *Parvicardium exiguum* (GMELIN, 1790) in Santander Bay (N of Spain). Journal of Molluscan Studies 59: 73-81.
- Lastra, M, J Palacio, and J Mora, 1993. Populations dynamics and secondary production of *Abra alba* (wood)(Bivalvia) in the Santander Bay, northern Spain. Sarsia 78: 35-42.
- Laudien, J, 2002. Population dynamics and ecology of the surf clam *Donax serra* (Vialvia, Donacidae) inhabiting beaches of the Benguela upwelling system. Ber. Polarforsch. Meeresforsch. 432. [[cites Laudien et al. 2001]]
- Le Pennec, M, and PG Beninger, 2000. Reproductive characteristics and strategies of reducing-system bivalves. Comparative Biochemistry and Physiology Part A 126: 1-16.
- Lee, 1985 (Heller, 1980)
- Lees, Dennis C., 2005 (unpub proposal). Assessment of Bivalve Recovery on Treated Mixed-Soft Beaches in Prince William Sound [370.pdf; Assessment of Bivalve Recovery, Prince William Sound; Project Number: 02574-BAA]

- Leontarakis, PK, and C.A. Richardson, 2005. Growth of the smooth clam, *Callista chione* (Linnaeus, 1758) (Bivalvia: Veneridae) from the Thracian Sea, northeastern Mediterranean. Journal of Molluscan Studies 71: 189 - 192.
- Lewis et al, 1982 (Powell and Cummins, 1985)
- Loesch and Haven, 1973 (Powell and Stanton, 1985)
- Loesch, 1957 (Powell and Cummins, 1985)
- Lomovasky, B.J., Brey, T., Morriconi, E. and Calvo, J. 2002. Growth and production of the venerid bivalve *Eurhomalea exalbida* in the Beagle Channel, Tierra del Fuego. Journal of Sea Research 48: 209-216.
- Lomovasky, BJ, T. Brey, and E. Morriconi, 2005. Population dynamics of the venerid bivalve *Tawera gayi* (Hupé, 1854) in the Ushuaia Bay, Beagle Channel. Journal of Applied Ichthyology 21 (1): 64 – 69.
- Lopez-Jamar et al, 1987 (Callender and Powell 1997) (Heller, 1990)
- Lutz and Castagna, 1980 (Heller, 1990)
- Lutz et al, 1988 (Callender and Powell 1997)
- MacIntyre 1970 (cited by Dekker and Beukema 1999)
- Manca Zeichen, M, S Agnesi, A Mariani, A Maccaroni, and GD Ardizzone, 2002. Biology and populations dynamics of *Donax trunculus* L. (Bivalvia: Donacidae) in the South Adriatic coast (Italy). Estuarine Coastal and Shelf Science 54: 971-982.
- Mane and Nagabhushanam, 1979 (Powell and Cummins, 1985)
- Mane, 1974 (Powell and Cummins, 1985)
- Marcano, JS, A Priet, A Larez, JJ Alio and H Sanabria, 2005. Growth and nortality of *Pinctada imbricata* (Mollusca: Peteridae) in Guamachito, Araya Peninsula, Sucre State, Venezuela. Ciencias Marinas 31: 387-397.
- Maslin and Bouvet ,1986 (Callender and Powell, 1997)
- Mason, J., 1983. Scallop and queen fisheries in the British Isles. Farnham: Fishing News Books
- Maximovich, N.V., 1984. Peculiarities of the ecology of abundant bivalve molluscs from the Chupa Inlet. Malacological Review 17 (1-2): 125-126
- McGrath and O'Foighil, 1986 (Heller, 1990)
- McLachlan and Hanekom, 1979 (Powell and Cummins, 1985)
- McLachlan, 1974 (Powell and Cummins, 1985)
- McLachlan, A, 1979. Growth and production of *Donax sordidus* Hanley on an open sandy beach in Algoa Bay. South African Journal of Zoology 14: 61-66.
- Menesguen and Dreves, 1987 (Callender and Powell, 1997)
- Metaxatos, A, 2004. Population dynamics of the venerid bivalve *Callista chione* (L.) in a coastal area of the eastern Mediterranean. Journal of Sea Research 52: 293-305.
- Milne and Dunnert, 1972 (Robertson, 1979)
- Moore and Lopez, 1970 (Powell and Cummins, 1985)
- Moore and Lopez, 1969 (Robertson, 1979)
- Morosan, EM and JML Orensanz, 2004. Age struture and growth in an unusual population of purple clams, *Amiantis purpuratus* (Lamarck, 1818) (Bivalvia: Veneridae), from Argentine Patagonia. Journ Shellfish Research 23: 73-80.
- Morsan, E and NF Ciocco, 2004. Age and growth model for the southern geoduck, *Panopea abbreviata*, off Puerto Lobos (Patagonia, Argentina). Fisheries Research (Amsterdam) 69: 343-348.
- Morton, B, 1978 (Powell and Stanton, 1985)

- Morton, B, 1981 (Heller, 1990)
- Morton, B, 1988. The population dynamics and reproductive cycle of *Brachidontes variabilis* (Bivalvia: Mytilidae) in a Hong Kong mangrove. *Malacological Review* 21: 109=117.
- Morton, B, 1988. The population structure and age of *Polymesoda (Geloina) erosa* (Bivalvia: Corbiculacea) from a Hong Kong mangrove. *Asian Marine Biology* 5: 107-113.
- Morton, B, 1989. Life-history characteristics and sexual strategy of *Mytilopsis sallei* (Bivalvia: Dreissenacea), introduced into Hong Kong. *Journal of Zoology (London)* 219: 469-485.
- Morton, B, 1995. The ecology and functional morphology of *Trigonothracia jinxingae* (Bivalvia: Anomalodesmata: Thraciodea) from Xiamen, China. *Journal of Zoology (London)* 237: 445-468.
- Mouthon, J, 2001. Life cycle and population dynamics of the Asian clam *Corbicula fluminea* (Bivalvia: Corbiculidae) in the Saone River at Lyon (France). *Hydrobiologia* 452: 109-119.
- Mukai, 1974 (Robertson, 1979)
- Muus, BJ, 1967. The fauna of Danish estuaries and lagoons. *Meddelelser fra Danmarks Fiskeri- og Havundersogelser NS* 5(1): 1-316
- Nakaoka, M and S Matsui, 1994. Annual variation in the growth rate of *Yoldia notabilis* (Bivalvia: Nuculanidae) in Otsuchi Bay, northwestern Japan, analyzed using shell microgrowth patterns. *Marine Biology* 119: 397-404.
- Narasimham, KA, 1988. Fishery and the population dynamics of the blood clam, *Anadara granosa* (Linnaeus) in the Kakinada Bay. *CMFRI Bulletin* 42: 130-134.
- Narayanan and Michael, 1968 (Powell and Cummins, 1985)
- Nayar, 1955 (Comfort, 1957)
- Newcoombe, 1935; 1936 (Comfort, 1957)
- O'Brien, K, and BJ Keegan, 2006. Age-related reproductive biology of the bivalve *Mysella bidentata* (Montagu) (Bivalvia: Galeommatacea) in Kinsale Harbour (south coast of Ireland). *Irish Naturalists' Journal* 28: 284-299.
- Ockelmann and Muus, 1978 (Heller, 1990)
- O'Foighil, D, D McGrath, ME Conneely, BF Keegan and M Costelloe, 1984. Population dynamics and reproduction of *Mysella bidentata* (Bivalvia: Galeommatacea) in Galway Bay, Irish west coast. *Marine Biology* 81: 283-291.
- Ohba, 1959 (Powell and Stanton, 1985)
- Okera, 1976 (Powell and Stanton, 1985)
- Orton and Amirthalingam, 1930 (Comfort, 1957)
- Paul et al, 1976 (Powell and Cummins, 1985)
- Peharda, M, CA Richardson, I Mladineo, S Sestanovic, Z Popovic, J Bolotin, and N Vrgoc, 2007. Age, growth and population structure of *Modiolus barbatus* from the Adriatic. *Marine Biology* 151: 629-638.
- Peharda, M, CA Richardson, V Onofri, A Bratos, and M Crnceanic, 2002. Age and growth of the bivalve *Arca noae* L. in the Croatian Adriatic Sea. *Journal of Molluscan Studies* 68: 307-310.
- Penchaszadeh and Olivier, 1975 (Powell and Cummins, 1985)
- Penzias, 1969 (Powell and Stanton, 1985)
- Peterson, CH, 1986. Quantitative allometry of gamete production by *Mercenaria mercenaria* into old age. *Marine Ecology Progress Series* 29: 93-97.

- Petersen, G. H., 1958. Notes on the growth and biology of the different *Cardium* species in Danish brackish water areas. Meddelelser fra Danmarks Fiskeri-og Havundersøgelser NS2(22):1-31 (cited by Muus, BJ 1967)
- Petersen, G. H., 1977. The density, biomass and origin of the bivalves of the central North Sea. Meddelelser fra Danmarks Fiskeri-og Havundersøgler, New Series Vol. 7: 221-273.
- Philipp, E, T. Brey, H-O Portner, and D Abele, 2005. Chronological and physiological ageing in a polar and a temperate mud clam. Mechanisms of Ageing and Development 126: 598-609.
- Pinn, EH, CA Richardson, RC Thompson, and SJ Hawkins, 2005. Burrow morphology, biometry, age and growth of piddocks (Mollusca: Bivalvia: Pholadidae) on the south coast of England. Marine Biology (Berlin) 147: 943-953.
- Posgay, 1979 (Callender and Powell, 1997)
- Power, AJ, J Nunez, M Mitchell and RL Walker 2004. Reproductive pattern of the blood ark, *Anadara ovalis*, from the northeast coast of Florida. Journal of Shellfisheries Research 23(1): 173-178.
- Power, AJ, M Mitchell, R Walker, M Posey, T Alphin, and C Belcher, 2006. Baseline Port Surveys for Introduced Molluscan, Crustacean, and Polychaete species in the South Atlantic Bight. NOAA's National Sea Grant Aquatic Nuisance Species Program, Project R/HAB-15, Grant NA06RG0029.  
[http://www.marsci.uga.edu/gaseagrant/pdf/Port\\_Survey.pdf](http://www.marsci.uga.edu/gaseagrant/pdf/Port_Survey.pdf)
- Quayle, 1952 (Comfort, 1957)
- Rachor, E, 1976. Structure, dynamics and productivity of a population of *Nucula nitidosa* (Bivalvia: Protobranchia). Berichte der Deutschen Wissenschaftlichen Komission fur Meeresforschung 24: 296-331.
- Rainer, 1982 (Powell and Cummins, 1985)
- Ralph and Maxwell, 1977 (Powell and Cummins, 1985)
- Rao and Roux, 1984 (Callender and Powell, 1997)
- Rapson, 1952 (Comfort, 1957 and Powell and Cummins, 1985)
- Rhoads et al, 1981 (Heller, 1990)
- Riascos, JM, O Heilmayer, and J Laudien, 2008. Population dynamics of the tropical bivalve *Cardita affinis* from Malaga Bay, Colombian Pacific related to La Nina 1999-2000. Helgoland Marine Research 62 (Supplement 1): S63-S71.
- Robertson, in press (Robertson, 1979) [not found]
- Rodhouse, 1978 (Powell and Stanton, 1985)
- Roux et al, 1985 (Callender and Powell, 1997)
- Rudy, P., and L.H. Rudy. 1979. Oregon estuarine invertebrates: an illustrated guide to the common and important invertebrate animals. U.S. Fish and Wildlife Service Biological Service Program. FWS/OBS-79/111:166-168.
- Salzwedel, H., 1979. Reproduction, growth, mortality and variations in abundance and biomass of *Tellina fabula* (Bivalvia) in the German Bight in 1975/1976. Veroffentlichungen des Instituts fur Meeresforschung in Bremerhaven, 18, 111-202.
- Sanders, 1956 (Roberston, 1979)
- Santos, SL and JL Simon, 1980. Response of soft-bottom benthos to annual catastrophic disturbance in a south Florida estuary. Marine Ecology Progress Series 3: 347-355.
- Sasaki, 1981 (Powell and Cummins, 1985)
- Sastray, 1979 (Heller, 1990)

- Schmidt, RR and JE Warme, 1969. Population characteristics of *Protothaca staminea* (Conrad) from Magu Lagoon, California. *Veliger* 12(2):193-9.
- Schöne, BR, DH Goodwin, KW Flessa, DL Dettman, and P Roopnarine, 2002. Sclerochronology and growth of the bivalve mollusks *Chione (Chionista) fluctifraga* and *C. (Chionista) cortezi* in the northern Gulf of California, Mexico. *Veliger* 45: 45-54.
- Selin, NI, 2004. Life span of marine bivalve mollusks: ecological aspects. Abstracts of the conference Mollusks of the northeastern Asia and Northern Pacific: Biodiversity, Ecology, Biogeography and Faunal History. Vladivostok, Russia, p 141-143.
- Selin, NI, 2007. Shell form, growth and life span of *Astarte arctica* and *A. borealis* (Mollusca: Bivalvia) from the subtidal zone of northwestern Sakhalin Island. *Biologiya Morya* (Vladivostok) 33: 278-283.
- Sellmer, 1967 (Heller, 1990)
- Sewell, 1924 (Comfort, 1957)
- Shaul and Goodwin, 1982 (Powell and Cummins, 1985)
- Somasekar et al, 1982 (Powell and Cummins, 1985)
- Stanezykowska, 1977 (Callender and Powell, 1997)
- Stephen, 1931 (Comfort, 1957)
- Stevenson and Dickie, 1954 (Powell and Cummins, 1985)
- Stockton, 1984 (Callender and Powell, 1997)
- Strasser, M., 1999. *Mya arenaria* - an ancient invader of the North Sea coast. *Helgoländer Meeresuntersuchungen*, 52, 309-324.
- Strom, A, RC Francis, NJ Mantua, EL Miles, and DL Peterson, 2005. Preserving low-frequency climate signals in growth records of geoduck clams (*Panopea abrupta*). *Palaeogeography Palaeoclimatology Palaeoecology* 228: 167-178.
- Takesue, RK and A. van Geen, 2004. Mg/Ca, St/Ca, and stable isotopes in modern and Holocene *Protothaca staminea* shells from a northern California coastal upwelling region, *Geochimica et Cosmochimica Acta* 68(19): 3845-3861.
- Tanabe, K, 1988. Age and growth rate determinations of an intertidal bivalve, *Phacosoma japonicum* using internal shell increments. *Lethaia* 21: 231-241.
- Tanaka and Kikuchi, 1971 (Robertson 1979)
- Tang, 1941 (Comfort, 1957)
- Taylor and Venn, 1978 (Powell and Cummins, 1985)
- Taylor, 1960 (Powell and Cummins, 1985)
- Theisen, 1973 (Powell and Cummins, 1985)
- Thiesen, B.F., 1973. The growth of *Mytilus edulis* L. (Bivalvia) from Disko and Thule district, Greenland. *Ophelia*, 12, 59-77.
- Thorarinsdottir, G.G., 1999. Lifespan of two long-lived bivalves *Arctica islandica* and *Panopea generosa*. Phuket Marine Biological Center Special Publication, no. 9, pp. 41 -46.
- Tokmakova, NP, NA Galimulina, and AP Anisimov, 2006. Morphofunctional characterization and ploidy levels of cells in the bivalve digestive gland, with special reference to somatic polyploidy. *Russian Journal of Marine Biology* 32(4): 229-235.
- Trevallion, 1971 (Powell and Stanton, 1985)
- Trutschler, K and C Samtleben, 1988. Shell growth of *Astarte elliptica* (Bivalvia) from Kiel Bay (western Baltic Sea). *Mar Ecol Progr Ser* 42: 155-162.
- Tunberg, 1984b (Callender and Powell, 1997)
- Tunberg, B, 1983. Growth of *Dosinia exoleta* (L) (Bivalvia) in Raunefjorden, western Norway. *Sarsia* 68: 41-45.

- Tunberg, B, 1983. Population structure, size distribution, and shell growth of *Dosinia lupinus* (L) (Bivalvia) in Ravnefjorden, western Norway, with biometrical comparison to *Dosinia exoleta* (L). *Sarsia* 68: 33-40
- Turekian et al, 1975 (Heller, 1990)
- Urban, H-J, and B Campos, 1994. Population dynamics of the bivalves *Gari solida*, *Semele solida* and *Protothaca thaca* from a small bay in Chile at 36°S\*. *Marine Ecology Progress Series* 115: 93-102.
- Urban, H-J, 1996. Population dynamics of the bivalves *Venus antiqua*, *Tagelus dombeii*, and *Ensis macha* from Chile at 36°S. *Journal of Shellfisheries Research* 15: 719-727.
- Vahl, 1981 (Powell and Cummins, 1985)
- Velosa, V, J Moreira, and JS Toncoso, 2007. Annual dynamics of bivalve populations in muddy bottoms of the Ensenada de Baiona (Galicia, NW Iberian Peninsula). *Iberus* 25: 19-28.
- Ventilla, 1982 (Heller, 1990)
- Verginelli, R and A Prieto, 1991. Produccion secundaria de *Pinctada imbricata* (Roding, 1798) (Pteriidae) en una poblacion del Golfo de Cariaco, Venezuela. *Acta Cientifica Venezolana* 42: 138-144.
- Verween, A, M Vincx, and S Degraer, 2006. Growth patterns of *Mytilopsis leucophaeata*, an invasive biofouling bivalve in Europe. *Biofouling* 22: 221-231.
- Vieira, NKM, NL Poff, DM Carlisle, SR Moulton II, ML Koski, and BC Kondratieff, 2006, A database of lotic invertebrate traits for North America: U.S. Geological Survey Data Series 187, <http://pubs.water.usgs.gov/ds187>.
- Wade, 1968 (Powell and Cummins, 1985)
- Walker, RL and PB Heffernan, 1994. Age, growth rate, and size of the southern surfclam, *Spisula solidissima similis* (Say, 1822). *Journ Shellfish Res* 13: 433-441.
- Warwick and Price, 1975 (Robertson, 1979)
- Watanabe, T, A Suzuki, H Kawahata, H Kan, and S Ogawa, 2004. A 60-year isotopic record from a mid-Holocene fossil giant clam (*Tridacna gigas*) in the Ryukyu Islands: physiological and paleoclimatic implications. *Palaeogeogr Palaeoclimatol Palaeoecol* 212: 3
- Weber, A, R Witbaard, and S van Steenpaal, 2001. Patterns of growth and undetectable growth lines of *Astarte sulcata* (Bivalvia) in the Faroe-Shetland Channel. *Senckenbergiana maritima* 31: 235-244.
- Wendell et al, 1976 1972 (Powell and Cummins, 1985)
- Weymouth and McMillin, 1931 (Comfort, 1957)
- Weymouth and Thompson, 1930 (Comfort, 1957)
- Weymouth, 1923 (Comfort, 1957)
- Williams and Dredge, 1981 (Powell and Cummins, 1985)
- Williamson, 1908 (Comfort, 1957)
- Wilson and Hodgkin, 1967 (Powell and Cummins, 1985)
- Wilson, J.G., 1992. Age specific energetics of reproduction in *Nucula turgida* (Leckenby and Marshall) a bivalve with lecithotrophic larval development. *Invertebrate Reproduction and Development*, 22, 275-280.
- Wolfe and Petteway, 1968 (Powell and Cummins, 1985)
- Wolff and deWolf, 1977 (Robertson 1979)
- Yap, 1977 (Powell and Cummins, 1985)
- Yonge, C.M., 1960. Oysters. London: Collins

- Zaika, VE, 1973. (transl. A. Mercado). Specific Production of Aquatic Invertebrates. J Wiley, New York, 154 p.
- Zolotarev, VN, 1980. The life span of bivalves from the Sea of Japan and Sea of Okhotsk. Soviet Journal of Marine Biology 6(6): 301-308.
- Zolotarev, VN, 1982. Assessment of life span in bivalve populations. *Zhurn. obsch. biol.* 43(2): 249–260.

**Sources for genus updating:**

- DIJKSTRA, H.H. 2004. Classification of Pectinoidea (Propeamussiidae and Pectinidae)eb publication <http://www.scallop.nl/>
- TER POORTEN, J. J. 2005. Outline of a Systematic Index -- Recent CARDIIDAE (Lamarck, 1809). Visaya 2005: 1-13.
- CLEMAM (western Europe),  
Malacolog 4.1.0 (western Atlantic),
- COAN, E. V., P. V. SCOTT and F. R. BERNARD. 2000. Bivalve Seashells of Western North America. Santa Barbara Museum of Natural History Monographs No. 2, 766 p.
- OBIS (Indo-Pacific)
- KRUG, A. Z., D. JABLONSKI and J. W. VALENTINE, 2008. Species-genus ratios reflect a global history of diversification and range expansion in marine bivalves. Proceedings of the Royal Society of London B 275: 1117-1123.
- KRUG, A. Z., D. JABLONSKI, J. W. VALENTINE and K. ROY. 2009. Generation of Earth's first-order biodiversity pattern. Astrobiology 9: 113-124.