

## Molecular Systematics of the Freshwater Prawn Genus *Macrobrachium* Bate, 1868 (Crustacea: Decapoda: Palaemonidae) Inferred from mtDNA Sequences, with Emphasis on East Asian Species

Min-Yun Liu<sup>1</sup>, Yi-Xiong Cai<sup>2</sup>, and Chyng-Shyan Tzeng<sup>1,\*</sup>

<sup>1</sup>Department of Life Sciences, National Tsing Hua University, 101 Kang-Fu Road, Sec. 2, Hsinchu 300, Taiwan

<sup>2</sup>Biodiversity Centre, National Parks Board, 1 Cluny Road, Singapore, 259569, Republic of Singapore

(Accepted June 12, 2006)

**Min-Yun Liu, Yi-Xiong Cai, and Chyng-Shyan Tzeng (2007)** Molecular systematics of the freshwater prawn genus *Macrobrachium* Bate, 1868 (Crustacea: Decapoda: Palaemonidae) inferred from mtDNA sequences, with emphasis on East Asian species. *Zoological Studies* 46(3): 272-289. Based on the mitochondrial DNA fragment of the large subunit (16S) ribosomal RNA gene, the monophyletic phylogeny of the genus *Macrobrachium*, including both land-locked and euryhaline species, was supported. There was, however, poor support for the internal structure. Results suggest that the evolution of this group was marked by rapid radiation. The 2 hypotheses that the prawns originated from marine ancestors and subsequently migrated towards fresh water in more than 1 wave of migration and that the abbreviated larval development of land-locked species represents adaptive convergence are both supported. It appears that most of the morphological characters commonly used to determine the taxonomy of genus evolved independently during the invasion of inland waters. Based on a fragment of 16S and a fragment of the cytochrome oxidase subunit I (COI) gene, several species groups, including species from different geographic regions and endemic groups were recognized from various continents (or regions). The presence of 5 cryptic species was indicated, and the identities of 4 supposedly undescribed species are confirmed. Localized speciation events in East Asia can be correlated with morphological similarities, notably members of the *M. asperulum* species group. Morphological characters tend to be conserved within species but quite variable between geographically distant populations, making species identification difficult. The present molecular results combined with morphological datasets can be used to help reorganize the taxonomy of various species groups. <http://zoolstud.sinica.edu.tw/Journals/46.3/272.pdf>

**Key words:** Molecular systematics, Phylogeny, Land-locked species, *Macrobrachium*, Cryptic species.

Freshwater prawns of the genus *Macrobrachium* Bate, 1868 (Crustacea: Palaemonidae) are a highly diverse group of decapod crustaceans thought to have originated from marine ancestors, some of which subsequently migrated towards fresh water in more than 1 wave; hence its members are known to inhabit the entire range of habitats from purely marine areas to inland hill streams and impounded water bodies (Tiwari 1955, Shokita 1979, Jalihal et al. 1993). To date, approximately 210 species are recognized (Short 2004); and there are numerous yet undescribed cryptic species (Chace and Bruce 1993,

Wowor and Choy 2001, Cai and Ng 2002, Cai et al. 2004, Short 2004, D. Wowor, pers. comm., Y. Cai, pers. obs.).

The genus *Macrobrachium* can be ecologically separated into 2 groups: most species are widely distributed and require a certain saline concentration (i.e., 10‰-35‰) to complete their larval development, as euryhaline species; others are land-locked species, with limited distributions and complete their entire life cycle in fresh water (Holthuis 1950, Johnson 1973, Shokita 1979). As *Macrobrachium* migrated towards fresh water, the prawns gradually evolved several adaptive fea-

\*To whom correspondence and reprint requests should be addressed. Tel: 886-3-5742765. Fax: 886-3-5742765. E-mail: [cstzeng@life.nthu.edu.tw](mailto:cstzeng@life.nthu.edu.tw)

tures. One of them is the abbreviated larval development achieved by reducing both the number of larval stages and the duration of the larval period (Shokita 1979, Jalihal et al. 1993). The abbreviated development of larvae in land-locked species was suggested to be a result of selective pressures for becoming established in freshwater environments (Shokita 1979, Magalhães and Walker 1988), and is a convergent phenomenon overriding phylogenetic relationships even above the generic level (Magalhães and Walker 1988). On the contrary, Pereira and Garcia (1995) suggested that since primitive palaemonids, like *Troglocubanus*, *Palaemonetes*, and *Pseudopalaemon*, possess abbreviated development, it can possibly be considered a primitive trait, and the abbreviated development took place early in the origin of the family Palaemonidae, rather than being a recent process.

Because of the degree of conservation of morphological characteristics, much debate has surrounded the systematic relationships of many species within this group, and its taxonomy and phylogenetic inferences have until recently been exclusively based on comparisons of external morphological characters (Holthuis 1950 1952, Johnson 1973, Pereira 1997). Some species groups were proposed based on morphological similarities, mainly of the rostrum and 2nd pereopod (Holthuis 1950, Johnson 1973). The phylogenetic significance of these groupings remains to be tested. Apart from their taxonomy, the phylogenetic affinities among *Macrobrachium* species are poorly understood. Pereira (1997) carried out the first phylogenetic study based on morphological characters on the family Palaemonidae. In recent years, Murphy and Austin (2002 2003 2004 2005) published a series of results for the phylogeny of *Macrobrachium* species based on the mitochondrial (mt)DNA fragment of the large subunit (16S) ribosomal (r)RNA gene marker. Their studies produced some interesting results and led to the generic clarification of some local species.

East Asia has a large landmass that resides within both the tropics and subtropics and has a long history of overland connections with tropical areas possessing rich species pools (Hamilton 1983, Guo et al. 1998). This region exhibits high species diversity of plant taxa (Guo et al. 1998, Qian and Ricklefs 2000), and its freshwater fish fauna also represents one of the richest ichthyofaunas in the Sino-Indian region (Koltelat 1989, Banareescu 1990). Climatic changes and geographic heterogeneity have played major roles in the diversification and speciation of East Asia's

biota, and it is a superior continental model for studying increases in regional diversity through allopatric speciation (Qian and Ricklefs 2000).

Previous studies of the genus *Macrobrachium* in East Asia comprised regional species surveys or zoogeographic distributions, including in China (Yu 1936, Dai 1984, Liu et al. 1990, Cai and Dai, 1999), Taiwan (Hwang and Yu 1982 1983, Shy and Yu 1998), and Japan (Hayashi 2000a b c). In total, about 37 species of *Macrobrachium* are found in China (Li et al. 2003), Taiwan (Shy and Yu 1998), and Japan (Hayashi 2000a b c), and there is a relatively high level of endemism. Some population studies of *M. nipponense* were based on allozyme variations and reproductive traits (see Mashiko and Numachi 2000 for review). Shokita (1979) inferred the speciation and origin of land-locked species based on prawns from the Ryukyu Archipelago, southern Japan, and discussed the biogeography of the genus *Macrobrachium* with special reference to larval dispersal (Shokita 1985). However, little work has been carried out on the phylogenetic relationships among *Macrobrachium* species of East Asia.

For this study utilizing the mtDNA 16S rRNA marker, we attempted to investigate the phylogeny and evolution of land-locked species of the genus *Macrobrachium*, based on species from the Indo-West Pacific region and by using sequences available from GenBank. Then, we focused on species distributed in East Asia, including China, Taiwan, and Japan, including the Ryukyus. We used a combination of the 16S rRNA fragment and a fragment of the cytochrome oxidase subunit I (COI) gene to elucidate the phylogenetic relationships of East Asian species, to test if speciation patterns of endemic species in the region resulted from multiple lineages or from a single event, and to reveal any cryptic species that are difficult to distinguish using more-traditional techniques (Kowltun 2000, Hendrixon and Bond 2005, Ellis et al. 2006).

## MATERIALS AND METHODS

### Collection of materials

In total, 238 specimens, representing 34 putative and 4 undescribed species of *Macrobrachium*, were collected from East and Southeast Asia for the sequence analysis. Moreover, for 15 species, multiple individuals were sequenced from geographically distant populations to assess the monophyly of the putative species (Sites and

Marshall 2003, Peters et al. 2005). One to 8 specimens were analyzed per locality (Table 1). Specimens used in the present study (Table 1) were caught in the wild and preserved in 75%-95% ethanol. Five species of 3 closely related genera in the same family (the Palaemonidae), namely *Exopalaemon modestus*, *E. orientis*, *Palaemonetes sinensis*, *P. atrinubes*, and *Palaemon siuenus*, together with an atyid shrimp, *Caridina pseudodenticulata*, were included as the outgroup. Additional mtDNA 16S sequences available from GenBank were included in this analysis (Table 2), to encompass a total of 62 *Macrobrachium* species.

### DNA extraction, PCR amplification, and sequencing

Total genomic DNA was extracted from the abdominal muscle by proteinase K/sodium dodecylsulfate (SDS) dissolution, phenol-chloroform extraction, and ethanol precipitation according to standard procedures (Sambrook et al. 1989).

Fragments of 2 mitochondrial genes, the 16S rRNA and COI genes, were amplified from total genomic (g)DNA by a polymerase chain reaction (PCR) using the conserved primers 1471 (5'-CCT GTTTANCAAAAACAT-3') and 1472 (5'-AGATA GAAACCAACCTGG-3') (Crandall and Fitzpatrick 1996) for 16S rRNA, and COI-a (5'-AGTATAAGCGTCTGGGTAGTC-3') and COI-f (5'-CCTGCAGGAGGA GGAGACCC-3') (Palumbi and Benzie 1991) for the COI gene. These primers of 16S and COI did not work well for some species, so a new primer pair, 1471B (5'-CCTGTTTAN CAAAACATGTCTG-3') and 1472B (5'-AGATA GAAACCAACCTGGCTCAC-3'), was modified for the 16S rRNA gene, and a new COI-fR (5'-CGTCGTGGTATGCCDTTARWCCTA-3') primer replaced the COI-a primer.

The amplification (50  $\mu$ l) of 16S rRNA used 1 mM of each primer, 0.2 mM of each dNTP, 1 unit of *Taq* polymerase (Promega, Madison, WI, USA), template DNA (50-100 ng), and 1X amplification buffer containing 1.5 mM MgCl<sub>2</sub> and either the

**Table 1.** Species of *Macrobrachium* and the outgroup included in this study. Sample localities, the number of samples examined, distribution, life cycle, and GenBank accession nos. are provided

Species	Sample locality	Locality code	Number examined	Distribution	Life cycle	Accession nos.	
						16S rRNA	COI
<i>M. anhuiense</i>	Anhui Prov., China	CN	1	East Asia	Land-locked	DQ194909	AB235240
<i>M. asperulum</i>	Fujian Prov., China	CN	6	East Asia	Land-locked	DQ194908	AB235241
	Ilan Co., Taiwan	N-TW <sup>a</sup>	7			DQ194906	AB235242
	Pingtung Co., Taiwan	S-TW <sup>b</sup>	5			DQ194907	AB235243
<i>M. australe</i>	Hualien Co., Taiwan	TW	6	Indo-West Pacific	Euryhaline	DQ194904	AB235245
	Cebu Is., the Philippines	PH	1			DQ194905	AB235244
<i>M. edentatum</i>	Sichuan Prov., China	CN	3	Indo-West Pacific	Land-locked	DQ194912	AB235247
<i>M. equidens</i>	Ilan Co., Taiwan	TW	4	Indo-West Pacific	Euryhaline	DQ194918	AB235250
	Bohol, the Philippines	PH	3			DQ194916	AB235248
	Seletor Is., Singapore	SG	5			DQ194917	AB235249
<i>M. esculentum</i>	Hualien Co., Taiwan	TW	6	Indo-West Pacific	Euryhaline	DQ194913	AB235252
	Cebu Is., Philippines	PH	1			DQ194914	AB235251
<i>M. formosense</i>	Yamaguchi Pref., Japan	JP	2	East Asia	Euryhaline	DQ194920	AB235253
	Okinawa I., Ryukyus, Japan	RK	3			DQ194922	AB235255
	Hainan Prov., China	CN	2			DQ194921	AB235254
	Hualien Co., Taiwan	TW	5			DQ194919	AB235256
<i>M. fukiense</i>	Fujian Prov., China	CN	2	East Asia	Land-locked	DQ194923	AB235257
<i>M. gracilirostre</i>	Hualien Co., Taiwan	TW	5	Indo-West Pacific	Euryhaline	DQ194924	AB235258
<i>M. grandimanus</i>	Okinawa I., Ryukyu, Japan	RK	2	Indo-West Pacific	Euryhaline	DQ194926	AB235260
	Hawaii Is., USA	HW	3			DQ194925	AB235259
<i>M. hainanense</i>	Guangdong Prov., China	CN	3	East Asia	Euryhaline	DQ194927	AB235261
	Hainan Prov., China	CN	2				
<i>M. cf. horstii</i>	Hualien Co., Taiwan	TW	9	Indo-West Pacific	Euryhaline	DQ194928	AB235291
<i>M. idae</i>	Khanom, Thailand	TH	2	Indo-West Pacific	Euryhaline	DQ194930	AB235262
<i>M. inflatum</i>	Anhui Prov., China	CN	3	East Asia	Land-locked	DQ194931	AB235263

**Table 1.** (Cont.)

Species	Sample locality	Locality code	Number examined	Distribution	Life cycle	Accession nos.	
						16S rRNA	COI
<i>M. japonicum</i>	Yamaguchi Pref., Japan	JP	2	East Asia	Euryhaline	DQ194934	AB235264
	Okinawa I., Ryukyu, Japan	RK	4			DQ194935	AB235265
	Hualien Co., Taiwan	TW	8			DQ194933	AB235266
<i>M. jaroense</i>	Hualien Co., Taiwan	TW	6	Indo-West Pacific	Euryhaline	DQ194932	AB235267
<i>M. lar</i>	Iriomote I., Ryukyu, Japan	RK	2	Indo-West Pacific	Euryhaline	DQ194941	AB235269
	Hualien Co., Taiwan	TW	8			DQ194939	AB235270
	Cebu Is., the Philippines	PH	2			DQ194940	AB235268
<i>M. lanatum</i>	Myanmar (Burma)	IN	3	Indo-West Pacific	Euryhaline	DQ194911	AB235246
<i>M. latidactylus</i>	Hainan Prov., China	CN	1	Indo-West Pacific	Euryhaline	DQ194943	AB235271
	Hualien Co., Taiwan	TW	8			DQ194942	AB235275
	Bohol Is., the Philippines	PH	2			DQ194945	AB235273
	Tioman Is., Malaysia	MY	4			DQ194944	AB235272
	Dodaga Halmahera, Thailand	TH	1			DQ194946	AB235274
<i>M. latimanus</i>	Okinawa I., Ryukyu, Japan	RK	1	Indo-West Pacific	Euryhaline	DQ194938	AB235277
	Hualien Co., Taiwan	TW	3			DQ194936	AB235278
	Cebu Is., the Philippines	PH	3			DQ194937	AB235276
<i>M. lepidactyloides</i>	Hualien Co., Taiwan	TW	7	Indo-West Pacific	Euryhaline	DQ194929	AB235279
<i>M. maculatum</i>	Anhui Prov., China	CN	4	Indo-West Pacific	Land-locked	DQ194910	AB235280
<i>M. malayanum</i>	Gunonghedang, Malaysia	MY	2	Indo-West Pacific	Land-locked	DQ194947	AB235281
<i>M. mammillodactylus</i>	Bohol Is., the Philippines	PH	1	Indo-West Pacific	Euryhaline	DQ194915	AB235282
<i>M. meridionalis</i>	Hainan Prov., China	CN	6	Indo-West Pacific	Euryhaline	DQ194948	AB235283
	Tiomam I., Malaysia	MY	1			DQ194949	AB235284
<i>M. naso</i>	Yaungwhe, Myanmar	BM	3	Indo-West Pacific	Euryhaline	DQ194950	AB235285
<i>M. neglectum</i>	Langkawi I., Malaysia	MY	1	Indo-West Pacific	Euryhaline	DQ194953	AB235286
<i>M. nipponense</i>	Guangxi Prov., China	CN	5	Indo-West Pacific	Land-locked	DQ194952	AB235287
	Hualien Co., Taiwan	TW	6			DQ194951	AB235288
<i>M. pinguis</i>	Fujian Prov., China	CN	2	East Asia	Land-locked	DQ194958	AB235289
<i>M. placidulum</i>	Leyte, the Philippines	PH	4	Indo-West Pacific	Euryhaline	DQ194956	AB235290
<i>M. placidum</i>	Leyte, the Philippines	PH	3	Indo-West Pacific	Euryhaline	DQ194957	AB235292
<i>M. platycheles</i>	Nee Soon, Singapore	SG	3	Indo-West Pacific	Land-locked	DQ194955	AB235294
	Johor, Malaysia	MY	2			DQ194954	AB235293
<i>M. rosenbergii</i>	Kaohsiung Co., Taiwan	TW	2	Indo-West Pacific	Euryhaline	DQ194959	AB235295
<i>M. shokitai</i>	Iriomote I., Ryukyu, Japan	RK	5	East Asia	Land-locked	DQ194961	AB235296
<i>M. yui</i>	Yunnan Prov., China	CN	2	South East Asia	Land-locked	DQ194960	AB235297
<i>M. sp.1</i>	Hualien Co., Taiwan	TW	1	East Asia	?	DQ194962	AB235298
<i>M. sp.2</i>	Guangxi Prov., China	CN	5	East Asia	?	DQ194963	AB235299
<i>M. sp.3</i>	Jiangxi Prov., China	CN	1	East Asia	?	DQ194964	AB235300
<i>M. sp.4</i>	Siem Reap, Cambodia	KH	1	South East Asia	?	DQ194965	AB235301
<i>M. sp.5</i>	Hualien Co., Taiwan	TW	8	East Asia	?	DQ194966	AB235302
<i>M. sp.6</i>	Bohol, the Philippines	PH	3	Indo-West Pacific	?	DQ194967	AB235303
<i>M. sp.7</i>	Okinawa I., Ryukyu, Japan	RK	1	East Asia	?	DQ194968	AB235304
<i>M. sp.8</i>	Okinawa I., Ryukyu, Japan	RK	1	East Asia	?	DQ194969	AB235305
Outgroup							
<i>Exopalaemon modestus</i>	Jiangxi Prov., China	CN	2	East Asia	Land-locked	DQ194971	AB235307
<i>Exopalaemon orientis</i>	Taipei Co., Taiwan	TW	3	East Asia	Euryhaline	DQ194972	AB235306
<i>Palaemonetes sinensis</i>	Jiangxi Prov., China	CN	1	East Asia	Land-locked	DQ194970	-
<i>Caridina pseudodenticulata</i>	Hsinchu Co., Taiwan	TW	3	East Asia	Land-locked	DQ194973	AB235308

<sup>a</sup>Illan County is in northeastern Taiwan. <sup>b</sup>Pingtung County is in southern Taiwan.

primer pair 1471+1472 or 1471B+1472B. The amplification conditions involved an initial cycle of denaturation at 94°C for 5 min, and then 35 cycles of denaturation at 94°C for 1 min, annealing at 48-55°C for 1 min, and extension at 72°C for 1 min, followed by a final extension at 72°C for 5 min. For the COI gene, the amplification used either primer pair COI-f+COI-a or COI-f+COI-fR. A similar profile to that of 16S rRNA was employed except that the annealing temperature ranged from 45 to 50°C. The size and quality of the PCR products were visualized on 1.5% agarose gels.

Prior to sequencing, the PCR products were purified using a gel purification kit according to the manufacturer's instructions (QIAGEN, Valencia, CA, USA). In order to control the sequence accuracy and resolve any ambiguous bases, sequencing was carried out in both directions using the same primer pairs for PCR by cycle sequencing using the ABI PRISM Dye-Terminator Sequencing Kit (Applied Biosystems, Foster City, CA, USA) and electrophoresis on an Applied Biosystems Automated Sequencer (model 377 or 3100).

### Data analysis

Forward and reverse sequences for an individual were edited using SeqMan (DNASTAR, LaserGene, Madison, WI, USA). All sequences were aligned using MegAlign (DNASTAR, LaserGene), with checks and adjustments made by eye using BioEdit v5.0.9 (Hall 1999). Exploratory data analysis of the sequences was performed using MEGA vers. 2.1 (Kumar et al. 2001) and DnaSP 4.00 (Rozas et al. 2003). Pairwise sequence comparisons provided an assessment of levels of saturation by plotting the number of transitions and transversions against the uncorrected proportional distances (p-distances) for each pair of unique sequences of *Macrobrachium* species (Morrison et al. 2004). Inter- and intraspecific genetic distances were calculated using the Kimura (1980) 2-parameter model with the pairwise deletion option in the MEGA program.

For the 16S sequences, the *Macrobrachium* species collected in this study, including 4 undescribed species, 7 species in the outgroup (Table 1), and additional sequences available from GenBank (Table 2), were analyzed to determine the molecular systematics of the genus. To elucidate the phylogenetic relationships of East Asian species, we analyzed the COI sequences and the combination of 16S and COI sequences only for

species collected in this study (Table 1); species containing only 16S sequences from GenBank (Table 2) were not included in this analysis. Sequences of both the 16S and COI genes were combined (Table 1) as both genes are in effect linked, and this is an appropriate way of dealing with random topological differences that are attributable to sampling error (Hipp et al. 2004). In order to test for the consistency of the phylogenetic signals in the data, phylogenetic relationships were inferred using 4 different analytical approaches. Maximum-parsimony (MP) (Camin and Sokal 1965) analysis was conducted assuming equal weightings for all characters, and results were compared when gaps were treated either as missing data or as a 5th character state. Neighbor-joining (NJ) (Saitou and Nei 1987) and maximum-likelihood (ML) (Felsenstein 1981) analyses were carried out using appropriate DNA substitution models calculated with ModelTest vers. 3.5 (Posada and Crandall 1998). In the MP analysis, the heuristic search option with tree-bisection-reconnection (TBR) branch swapping and 100 stepwise random additions of taxa was used. In the NJ and MP analyses, levels of branch support were assessed using bootstrap resampling (Felsenstein 1985) with 1000 replicates to evaluate the reliability of the inferred topologies. Bootstrap resamplings were run with the "fast" stepwise addition algorithm and 100 replicates for ML, because of the large number of taxa involved and the computational time requirements. The NJ, MP, and ML analyses were carried out with PAUP\* vers. 4.0b10 (Swofford 2000). Bayesian analyses (BI) were performed with MRBAYES vers. 3.0 (Ronquist and Huelsenbeck 2003) using the models selected by MrModeltest (Nylander 2004). Markov chain Monte Carlo (MCMC) chains were run for  $1 \times 10^6$  generations, and trees were saved each 100 generations (with the 1st 1000 trees being discarded as "burn-in"). In BI, posterior probabilities are true probabilities of clades, and those with values of 95% or greater were deemed to be significantly supported.

## RESULTS

### Sequence characteristics and variations

The 16S rRNA sequence amplified by the 1471B+1472B primer pair varied from 524 to 533 bp in length (Table 1). Additional sequences obtained from GenBank (Table 2) were shorter;

final truncated lengths for the multiple alignments were 442 bp, including 20 sites with gaps, 191 variable sites, with 160 parsimoniously informative sites. The numbers of transitions outnumbered transversions in all comparisons by a factor of approximately 1.7. The COI sequences of species collected in this study (Table 1) and amplified by

the COI-f+COI-fR primer pair contained 608 bp, including 276 sites which were variable and 258 which were parsimoniously informative. No stop codon was revealed when the COI sequences were translated into amino acids. The numbers of transitions outnumbered transversions by an average of 2.2. Base frequencies in both mtDNA

**Table 2.** Additional 16S sequences of *Macrobrachium* and the outgroup used in this study obtained from GenBank

Species	Sample locality	Locality code	Distribution	Life cycle	Accession no.
<i>M. acanthochirus</i>	Mexico	MX	South/Central America	Euryhaline	AY377837 <sup>c</sup>
<i>M. acanthurus</i>	Mexico	MX	South/Central America	Euryhaline	AY282780 <sup>b</sup>
<i>M. aemulum</i>	Australia	AU	Indo-West Pacific	Euryhaline	AY282769 <sup>b</sup>
<i>M. auratum</i>	Australia	AU	Indo-West Pacific	Euryhaline	AY282775 <sup>b</sup>
<i>M. australiense</i>	Australia	AU	Indo-West Pacific	Land-locked	AY282764 <sup>b</sup>
<i>M. brasiliense</i>	Brazil	BR	South/Central America	Land-locked	AY377839 <sup>b</sup>
<i>M. bullatum</i>	Australia	AU	Indo-West Pacific	Land-locked	AY282778 <sup>b</sup>
<i>M. carcinus</i>	Puerto Rico	PR	South/Central America	Euryhaline	AY282779 <sup>b</sup>
<i>M. crenulatum</i>	Puerto Rico	PR	South/Central America	Euryhaline	AY377840 <sup>c</sup>
<i>M. equidens</i>	Australia	AU	Indo-West Pacific	Euryhaline	AY282773 <sup>b</sup>
<i>M. gangeticum</i>	India	IN	Indo-West Pacific	Euryhaline	AY730054 <sup>d</sup>
<i>M. hainanense</i>	Hong Kong	HK	Indo-West Pacific	Euryhaline	AY377841 <sup>c</sup>
<i>M. handschini</i>	Australia	AU	Indo-West Pacific	Land-locked	AY282781 <sup>b</sup>
<i>M. heterochirus</i>	Puerto Rico	PR	South/Central America	Euryhaline	AY377842 <sup>c</sup>
<i>M. idea</i>	Australia	AU	Indo-West Pacific	Euryhaline	AY282777 <sup>b</sup>
<i>M. intermedium</i>	Australia	AU	Indo-West Pacific	Marine <sup>e</sup>	AF439515 <sup>a</sup>
<i>M. koombooloomba</i>	Australia	AU	Indo-West Pacific	Land-locked	AY282767 <sup>b</sup>
<i>M. lamarrei</i>	India	IN	Indo-West Pacific	Land-locked	AY730051 <sup>d</sup>
<i>M. lar</i>	Australia	AU	Indo-West Pacific	Euryhaline	AY282766 <sup>b</sup>
	Indonesia	ID			AY377843 <sup>c</sup>
<i>M. latimanus</i>	Australia	AU	Indo-West Pacific	Euryhaline	AY282765 <sup>b</sup>
<i>M. latidactylus</i>	Australia	AU	Indo-West Pacific	Euryhaline	AY282770 <sup>b</sup>
<i>M. malcolmsonii</i>	India	IN	Indo-West Pacific	Euryhaline	AY730050 <sup>d</sup>
<i>M. malayanum</i>	Singapore	SG	Indo-West Pacific	Land-locked	AY377844 <sup>c</sup>
<i>M. mammillodactylus</i>	Indonesia	ID	Indo-West Pacific	Euryhaline	AY377845 <sup>c</sup>
	Australia	AU			AY282776 <sup>b</sup>
<i>M. neglectum</i>	Indonesia	ID	Indo-West Pacific	Euryhaline	AY377846 <sup>c</sup>
<i>M. novaehollandiae</i>	Australia	AU	Indo-West Pacific	Euryhaline	AY282772 <sup>b</sup>
<i>M. olfersii</i>	Brazil	BR	South/Central America	Euryhaline	AY377848 <sup>c</sup>
	Mexico	MX	South/Central America		AY377849 <sup>c</sup>
<i>M. platycheles</i>	Singapore	SG	Indo-West Pacific	Land-locked	AY377850 <sup>c</sup>
<i>M. potiuna</i>	Brazil	BR	South/Central America	Land-locked	AY377851 <sup>c</sup>
<i>M. rosenbergii</i>	Australia	AU	Indo-West Pacific	Euryhaline	AY282774 <sup>b</sup>
<i>M. sankollii</i>	India	IN	Indo-West Pacific	Land-Locked	AY730052 <sup>d</sup>
<i>M. scabriculum</i>	India	IN	Indo-West Pacific	Euryhaline	AY730055 <sup>d</sup>
<i>M. tolmerum</i>	Australia	AU	Indo-West Pacific	Euryhaline	AY282768 <sup>b</sup>
<i>M. trompii</i>	Singapore	SG	Indo-West Pacific	Land-locked	AY377852 <sup>c</sup>
<i>M. zariqueyi</i>	Annobon	AN	West Africa	Euryhaline	AY377847 <sup>c</sup>
<i>Palaemon serenus</i>	Australia	AU	Indo-West Pacific	Euryhaline	AF439518 <sup>a</sup>
<i>Palaemonetes atrinubes</i>	Australia	AU	Indo-West Pacific	Euryhaline	AF439520 <sup>a</sup>
<i>Palaemonetes australis</i>	Australia	AU	Indo-West Pacific	Euryhaline	AF439517 <sup>a</sup>

<sup>a</sup>Murphy and Austin 2003. <sup>b</sup>Murphy and Austin 2004. <sup>c</sup>Murphy and Austin 2005. <sup>d</sup>available in GenBank. <sup>e</sup>the only known species of the genus that spends its entire life in the sea (Holthuis 1952).

genes showed an AT bias (with G+C contents of 35.8% for 16S and 40.7% for COI). The combined data set of 16S+COI multiple alignments was 1142 bp in length including 20 sites with gaps, 472 variable sites, and 410 parsimoniously informative sites. All mtDNA sequences determined in this study were deposited in the GenBank/DDBJ databases under accession numbers DQ194904-DQ194973 and AB235240-AB235308 (Table 1).


In the gene-saturation analyses, substitutions of transitions and transversions for 16S were approximately linear in distribution with a positive slope for the regression ( $R^2 = 0.714$  and  $0.660$  for transitions and transversions, respectively, data not shown), indicating that 16S rRNA is not saturated. For the COI data, transitions and transversions were plotted by separate codon position, and a saturation tendency was shown for transitions in the 3rd position, not for the 1st or 2nd positions, which appeared to reach a plateau at p-distances above 20% (Fig. 1). Sequence divergence estimates among the *Macrobrachium* species ranged 0.47%-22.44% for 16S and 0.16%-25.54% for COI; for conspecific individuals from the same locality, they ranged 0.00%-0.11% for 16S and 0.00%-3.70% for COI. Between populations (or individuals) from different localities of conspecifics, they ranged 0.00%-3.20% for 16S and 0.00%-12.63% for COI, (the most-distant populations of *M. grandimanus* being from the Ryukyus and Hawaii in this study). Four undescribed species were confirmed to be genetically distinct from other species, with interspecific divergences of 3.5%-19.9% for 16S and 9.48%-27.67% for COI. These will, hereafter, be referred to as *Macrobrachium* sp.1, *M. sp.2*, *M. sp.3*, and *M. sp.4*. We also found some discordant cases in which intraspecific divergences (4.9%-9.2% for 16S and 13.23%-17.24% for COI) greatly exceeded the usual ranges in this study, and might therefore reflect interspecific differences. Such discordant situations were found in samples of *M. latidactylus*, *M. latimanus*, *M. jaroense*, *M. placidum*, and *M. equidens* taken from multiple localities. According to these results, we believe such specimens may represent "cryptic species" (sensu Gusmão et al. 2000, Knowlton 2000, Hendrixon and Bond 2005, Ellis et al. 2006). These species were thus respectively referred to as *M. sp.5*, *M. sp.6*, *M. sp.7*, and *M. sp.8* for *M. latidactylus*, *M. latimanus*, *M. jaroense*, and *M. placidum* (Figs. 2, 3). *Macrobrachium equidens* is discussed below. In contrast, the divergence between 2 morphologically distinct species, *M. formosense* and *M. hainanense* (0.01%-0.02% for

16S and 0.16%-0.66% for COI), was as close as that of the intraspecific level. This was also found for *M. cf. horstii* and *M. placidulum* (0.37% for 16S and 1.33% for COI) (Table 3).

### Phylogenetic analyses

Based on results from Modeltest, the best-fit models in the NJ and ML analyses were as follows: 16S, HKY+I+G (Hasegawa et al. 1985), with a correction for the among-site rate variation estimation (G) of 0.6420 and a proportion of invariable sites (I) of 0.4499; COI: GTR+I+G (general time reversible model; Rodriguez et al. 1990), with a correction for G of 0.4023 and I of 0.4704; and for the combined dataset: TrN+I+G (Tamura-Nei model; Tamura and Nei 1993) with a correction for G of 0.7019 and I of 0.5354. For BI, the best-fit models selected by the Akaike information criterion (AIC) in MrModeltest were as follows: 16S: GTR+I+G, with a correction for G of 0.5897 and I of 0.4106; COI: GTR+I+G, with a correction for G of 0.4133 and I of 0.4770; and for the combined dataset: GTR+I+G with a correction for G of 0.6176 and I of 0.4675.

For 16S, 103 haplotypes for *Macrobrachium* species, including 4 undescribed species, 7 species of the outgroup (Table 1), and sequences available from GenBank (Table 2), were analyzed. A 50% majority consensus tree was obtained from the MP analyses. All 4 phylogenetic analyses generated similar tree topologies when gaps were treated as a 5th character (Fig. 2). The major differences between the 4 analyses lay in the levels of support provided for the various clades. Tree topologies supported the monophyly of the genus



**Fig. 1.** Saturation test of cytochrome oxidase subunit I (COI), relationships of uncorrected p-distances between pairs of taxa, and the number of transitional changes at the 3rd codon position. A saturation tendency was shown to reach a plateau at p-distances above 20%.

*Macrobrachium* with high bootstrap values in the NJ, MP, and BI analyses. The outgroup genera of *Palaemonetes*, *Palaemon*, and *Exopalaemon* formed a monophyletic group with high bootstrap support, and were paraphyletic to *Macrobrachium*. *Exopalaemon modestus* and *E. orientis* formed a sister taxon pair with high bootstrap support in all analyses. *Palaemonetes atrinubes* and *P. sinensis* formed a sister taxon pair with weak bootstrap support, *Palaemon serenus* and *M. intermedium* formed a sister taxon pair with high bootstrap support. The species *M. intermedium*, located outside the *Macrobrachium* clade, was more-closely related to *Palaemon serenus* than to any other species (Fig. 2).

The deeper internal nodes were generally unresolved in the 16S analysis (Fig. 2). Some species groups, with good support for many terminal clades, were revealed giving some support or revealing some incongruence with earlier classifications. In addition to the Central/South American and Central/South American/West African species clades as reported by Murphy and Austin (2005), another 4 monophyletic groups were revealed (Fig. 2). The India group (INDIA) contains *Macrobrachium rosenbergii*, 2 Indian euryhaline species, *M. gangeticum* and *M. malcolmsonii*, and 2 Indian land-locked species, *M. lamarrei* and *M. sankollii*. The Indo-West Pacific group (IWP) includes several widely distributed euryhaline species, namely *M. cf. horstii*, *M. placidulum*, *M. placidum*, and 1 cryptic species, *M. sp.8*, which is morphologically very

close to *M. placidum*. The next 2 distinct groups are endemic to East Asia; East Asia group I (EA I) is the land-locked *M. asperulum* species group, containing *M. asperulum*, *M. anhuiense*, *M. pinguis*, *M. shokitai*, and *M. maculatum*. East Asia group II (EA II) contains the euryhaline species *M. formosense* and *M. hainanense* and the land-locked species, *M. nipponense*, *M. inflatum*, and an undescribed species, *M. sp.4* (Fig. 2).

Multiple samples from distant geographic populations of putative species were grouped into species-specific monophyletic groups with low to high bootstrap support (i.e., bootstrap values of 51-100), with the exception of 2 species (Fig. 2). The *M. equidens* species group, suggested by Johnson (1973), including *M. equidens*, *M. idae*, *M. mammillodactylus*, and *M. novaehollandiae*, did not form a monophyletic group in our study (Fig. 2). Moreover, specimens of *M. equidens* from 4 different localities, those of Taiwan, the Philippines, and Australia, formed a lineage and were separated, with distinct genetic distances (16.7%-17.4%), from specimens from Singapore, the type locality of *M. equidens*. Such disagreement was more evident in other species. Two populations of *M. hainanense* from China (Guangdong Prov. and Hainan I., Table 1) had an inter-population distance of 0.08%. When we compared our *M. hainanense* (*M. hainanense* (ML)) sequences with *M. hainanense* (HK) (accession no.: AY377841), collected from Hong Kong (Murphy and Austin 2005), there were inconsistencies with a significant dis-

**Table 3.** Genetic distance matrix of the large subunit (16S) (lower left) and cytochrome oxidase subunit I (COI) (upper right) among species of 4 species groups. SpG 1-4 are species groups 1-4, as revealed in figure 3


16S \ COI	SpG 1								SpG 2			
	<i>M. asperulum</i> <sup>a</sup>	<i>M. anhuiense</i>	<i>M. asperulum</i> <sup>b</sup>	<i>M. pinguis</i>	<i>M. asperulum</i> <sup>c</sup>	<i>M. shokitai</i>	<i>M. maculatum</i>	<i>M. sp. 2</i>	<i>M. sp. 3</i>	<i>M. edentatum</i>	<i>M. sp. 4</i>	<i>M. hainanense</i>
SpG 1												
<i>M. asperulum</i> <sup>a</sup>		0.0252	0.0534	0.0537	0.0555	0.0922	0.1522	0.1559	0.1382	0.1248	0.1713	0.1737
<i>M. anhuiense</i>	0.0090		0.0588	0.0553	0.0413	0.0998	0.1498	0.1581	0.1463	0.1248	0.1757	0.1848
<i>M. asperulum</i> <sup>b</sup>	0.0261	0.0284		0.0408	0.0532	0.0998	0.1521	0.1537	0.1526	0.1245	0.1562	0.1651
<i>M. pinguis</i>	0.0236	0.0235	0.0164		0.0393	0.1059	0.1518	0.1622	0.1485	0.1206	0.1711	0.1891
<i>M. asperulum</i> <sup>c</sup>	0.0283	0.0257	0.0256	0.0205		0.0694	0.1359	0.1255	0.1718	0.1311	0.1711	0.1822
<i>M. shokitai</i>	0.0406	0.0381	0.0332	0.0405	0.0405		0.1539	0.1454	0.1741	0.1557	0.1774	0.1842
<i>M. maculatum</i>	0.0555	0.0529	0.0579	0.0630	0.0631	0.0554		0.0948	0.1446	0.1565	0.1718	0.1851
<i>M. sp. 2</i>	0.0657	0.0631	0.0630	0.0605	0.0631	0.0604	0.0358		0.1549	0.1457	0.1588	0.1918
<i>M. sp. 3</i>	0.0734	0.0707	0.0784	0.0783	0.0759	0.0757	0.0710	0.0789		0.1404	0.1902	0.2115
<i>M. edentatum</i>	0.0657	0.0607	0.0656	0.0707	0.0682	0.0504	0.0711	0.0736	0.0761		0.1793	0.1982
SpG 2												
<i>M. sp. 4</i>	0.1086	0.1116	0.0999	0.1053	0.1140	0.0998	0.1001	0.1139	0.1191	0.1062		0.1302




Table 3. (Cont.)

16S \ COI	SpG 1						Sp G 2					
	<i>M. asperulum</i> <sup>a</sup>	<i>M. anhuiense</i>	<i>M. asperulum</i> <sup>b</sup>	<i>M. pinguis</i>	<i>M. asperulum</i> <sup>c</sup>	<i>M. shokitai</i>	<i>M. maculatum</i>	<i>M. sp. 2</i>	<i>M. sp. 3</i>	<i>M. edentatum</i>	<i>M. sp. 4</i>	<i>M. hainanense</i>
<i>M. hainanense</i>	0.0769	0.0741	0.0687	0.0712	0.0794	0.0633	0.0714	0.0713	0.0791	0.0766	0.0791	
<i>M. formosense</i>	0.0769	0.0741	0.0687	0.0712	0.0794	0.0633	0.0714	0.0713	0.0792	0.0766	0.0792	0.0021
<i>M. nipponense</i>	0.0716	0.0688	0.0687	0.0712	0.0741	0.0582	0.0662	0.0713	0.0737	0.0713	0.0791	0.0141
SpG 3												
<i>M. inflatum</i>	0.0689	0.0662	0.0661	0.0685	0.0714	0.0608	0.0636	0.0687	0.0712	0.0687	0.0763	0.0117
<i>M. lanatum</i>	0.0975	0.0946	0.0945	0.0998	0.0892	0.0916	0.0949	0.1002	0.0945	0.0893	0.1221	0.0981
<i>M. latidactylus</i>	0.1021	0.1019	0.0886	0.0965	0.1074	0.1046	0.1049	0.1158	0.1209	0.1158	0.1438	0.1104
SpG 4												
<i>M. sp. 5</i>	0.1184	0.1210	0.1184	0.1239	0.1239	0.1264	0.1270	0.1297	0.1406	0.1268	0.1583	0.1356
<i>M. esculentum</i>	0.0895	0.0893	0.0838	0.0839	0.0839	0.0837	0.0815	0.0814	0.0919	0.0814	0.1109	0.0954
<i>M. cf. horstii</i>	0.1082	0.1053	0.1135	0.1078	0.1053	0.1025	0.1161	0.1274	0.1299	0.0888	0.1327	0.1197
<i>M. placidulum</i>	0.1111	0.1081	0.1107	0.1105	0.0998	0.0970	0.1133	0.1188	0.1299	0.0942	0.1299	0.1112
<i>M. placidum</i>	0.1216	0.1243	0.1242	0.1297	0.1158	0.1186	0.1243	0.1302	0.1499	0.1104	0.1495	0.1304
<i>M. sp. 8</i>	0.1051	0.1049	0.1048	0.1049	0.0940	0.0915	0.0942	0.1105	0.1102	0.0993	0.1294	0.1191
<i>M. lepidactyloides</i>	0.1294	0.1293	0.1406	0.1376	0.1433	0.1290	0.1495	0.1464	0.1521	0.1321	0.1726	0.1212
Sp G 2												
16S \ COI	<i>M. formosense</i>	<i>M. nipponense</i>	<i>M. inflatum</i>	<i>M. lanatum</i>	<i>M. latidactylus</i>	<i>M. sp. 5</i>	<i>M. esculentum</i>	<i>M. cf. horstii</i>	<i>M. placidulum</i>	<i>M. placidum</i>	<i>M. sp. 8</i>	<i>M. lepidactyloides</i>
SpG 1												
<i>M. asperulum</i> <sup>a</sup>	0.1759	0.1651	0.1715	0.1662	0.2165	0.2115	0.1635	0.2171	0.2193	0.2151	0.1916	0.2102
<i>M. anhuiense</i>	0.1870	0.1759	0.1825	0.1579	0.2146	0.2120	0.1638	0.2077	0.2241	0.2174	0.1938	0.2102
<i>M. asperulum</i> <sup>b</sup>	0.1671	0.1612	0.1675	0.1468	0.2094	0.2096	0.1530	0.2081	0.1943	0.1943	0.1656	0.2102
<i>M. pinguis</i>	0.1914	0.1670	0.1823	0.1638	0.2234	0.2043	0.1787	0.2144	0.2075	0.2172	0.1868	0.2288
<i>M. asperulum</i> <sup>c</sup>	0.1842	0.1820	0.1864	0.1701	0.1863	0.2108	0.1633	0.2003	0.2137	0.2163	0.1928	0.2051
<i>M. shokitai</i>	0.1864	0.1798	0.1754	0.1638	0.1934	0.2094	0.1651	0.2277	0.2373	0.2281	0.1996	0.2279
<i>M. maculatum</i>	0.1872	0.1633	0.1676	0.1871	0.2115	0.1932	0.1809	0.2212	0.2224	0.2344	0.2038	0.2027
<i>M. sp. 2</i>	0.1941	0.1764	0.1809	0.2083	0.2197	0.2099	0.1832	0.2115	0.2229	0.2162	0.2057	0.2069
<i>M. sp. 3</i>	0.2138	0.1822	0.1733	0.2066	0.2372	0.2201	0.1848	0.2284	0.2366	0.2663	0.1980	0.2563
<i>M. edentatum</i>	0.2005	0.1757	0.1823	0.1943	0.2049	0.1914	0.1673	0.1834	0.1961	0.2267	0.1914	0.2011
SpG 2												
<i>M. sp. 4</i>	0.1323	0.1137	0.1117	0.1787	0.1964	0.1611	0.1642	0.1754	0.1947	0.1909	0.1506	0.1809
<i>M. hainanense</i>	0.0016	0.0591	0.0591	0.1722	0.1943	0.1742	0.1597	0.1923	0.2029	0.2176	0.1901	0.2155
<i>M. formosense</i>		0.0610	0.0573	0.1711	0.1966	0.1720	0.1618	0.1945	0.2052	0.2201	0.1922	0.2179
<i>M. nipponense</i>	0.0141		0.0271	0.1678	0.2059	0.1832	0.1610	0.1991	0.2011	0.2083	0.1813	0.2274
SpG 3												
<i>M. inflatum</i>	0.0117	0.0070		0.1678	0.2012	0.1787	0.1601	0.1945	0.2032	0.2106	0.1904	0.2274
<i>M. lanatum</i>	0.0981	0.0898	0.0898		0.1801	0.1664	0.1463	0.1821	0.2043	0.1883	0.2008	0.1999
<i>M. latidactylus</i>	0.1104	0.1076	0.1132	0.0863		0.1772	0.1861	0.1914	0.2141	0.1864	0.1866	0.1999
SpG 4												
<i>M. sp. 5</i>	0.1356	0.1242	0.1299	0.0941	0.0482		0.1702	0.1755	0.2087	0.2003	0.2165	0.2057
<i>M. esculentum</i>	0.0954	0.0872	0.0927	0.0611	0.0839	0.0891		0.1731	0.1889	0.2045	0.1881	0.1835
<i>M. cf. horstii</i>	0.1197	0.1084	0.1056	0.1167	0.1413	0.1442	0.1084		0.0133	0.1293	0.1943	0.1777
<i>M. placidulum</i>	0.1112	0.1056	0.1028	0.1139	0.1384	0.1413	0.0946	0.0037		0.1426	0.1811	0.2045
<i>M. placidum</i>	0.1304	0.1191	0.1161	0.1249	0.1384	0.1355	0.1193	0.0636	0.0584		0.1989	0.1926
<i>M. sp. 8</i>	0.1191	0.1024	0.1051	0.0808	0.0992	0.1074	0.0732	0.0791	0.0790	0.0868		0.2033
<i>M. lepidactyloides</i>	0.1212	0.1211	0.1239	0.1409	0.1571	0.1720	0.1382	0.1186	0.1158	0.1297	0.1132	

<sup>a</sup>Samples collected from northern Taiwan. <sup>b</sup>samples collected from southern Taiwan. <sup>c</sup>samples collected from China.



**Fig. 2.** Phylogenetic relationships based on large subunit (16S) sequences represented by a maximum parsimony (MP) 50% majority tree. Numbers on the branches indicate bootstrap values for the Neighbor-joining (NJ), MP, and maximum-likelihood (ML) analyses, and Bayesian analysis (BI). LL, land-locked species; EA, East Asia; IWP, Indo-West Pacific; S/C AM, South/Central America; WA, West Africa. EA I and EA II, 2 species groups found in East Asia. Abbreviations in brackets after the species names are the locality codes, which are given only when multiple samples of species were collected from different localities to assess the monophyly of the putative species. Locality codes are explained in tables 1 and 2. An asterisk (\*) indicates a bootstrap value of < 50.



**Fig. 3.** Phylogenetic relationships based on combined large subunit (16S) and cytochrome oxidase subunit I (COI) sequences represented by a maximum-parsimony (MP) consensus tree. Numbers above the branches indicate bootstrap values for the Neighbor-joining (NJ) and MP analyses, while numbers below the branches indicate maximum-likelihood (ML) bootstrap values and Bayesian analysis (BI). LL, land-locked species; SpG1-4, species group 1-4. Abbreviations in brackets after the species names are the locality codes, which are given only when multiple samples of species were collected from different localities to assess the monophyly of the putative species. Locality codes are explained in table 1. An asterisk (\*) indicates a bootstrap value of < 50.

tance of 7.6%. The *M. hainanense* (HK) sequence was placed in a different clade with land-locked species and was closely correlated to *M. maculatum* with a divergence of 3.8%, wherein our *M. hainanense* sequences formed a clade with the euryhaline species group and was closely correlated with *M. formosense*.

Species containing only 16S rRNA sequences in previous studies or only available in GenBank (Table 2) were not included in constructing the COI phylogenetic tree (COI tree not shown). Most of the species groups revealed above (Fig. 2), including the 4 undescribed *M. spp*1-4, 4 cryptic species, *M. spp*.5-8, and the incongruence of *M. equidens*. The phylogenetic tree of COI sequences revealed similar topologies as constructed by the phylogenetic analyses of the 16S sequences (Fig. 2). The monophyly of the *Macrobrachium* species clade could not be confirmed using COI data, as outgroup species of *Exopalaemon modestus* and *E. orientis* nested in one of the polyphyletic clades of *Macrobrachium* species taxa. The non-monophyletic structure of *Macrobrachium* species may be attributed to saturation of the 3rd codon of the COI gene (Fig. 1).

### Analysis of the combined dataset

We used a combination of 16S and COI fragments of species collected in this study (Table 1) to elucidate the phylogenetic relationships of East Asian species. Although the deeper internal nodes were generally unresolved and the relationships among the species were not well-resolved by the combined sequences, certain phylogenetic relationships of the species complexes were shown to be well supported (Fig. 3). For example, the morphologically similar land-locked species endemic to East Asia, including the *M. asperulum* species group (*M. anhuiense*, *M. pinguis*, and *M. shokitai*) and other land-locked species of *M. edentatum* and *M. maculatum* and 2 undescribed species of *M. sp*.2 and *M. sp*.3 formed a monophyletic group (species group 1; SpG 1). The morphologically dissimilar land-locked species, *M. fukiense*, and Southeast Asian species, *M. malayanum*, *M. platycheles*, and *M. yui*, were not included. *Macrobrachium nipponense*, *M. inflatum*, *M. formosense*, *M. hainanense*, and 1 undescribed species, *M. sp*.4, formed a clade as a species group (species group 2; SpG 2). Another species group (species group 3; SpG 3) with a similar morphology of an unequal 2nd periopod, containing *M. esculentum*, *M. lanatum*, *M. latidactylus*, and cryp-

tic species *M. sp*.5, formed a monophyletic group. Species *M. lepidactyloides* formed a clade with the morphologically similar species of *M. cf. horstii*, *M. placidulum*, *M. placidum*, and cryptic species *M. sp*.8 forming the monophyletic species group 4 (SpG 4).

The incongruence of the non-monophyly of *M. equidens* in the 16S (Fig. 2), COI (tree not shown) and 16S+COI (Fig. 3) analyses implies the existence of a 5th cryptic species.

## DISCUSSION

### Phylogenetic relationships

Although we have included more species distributed in East Asia and the Indo-West Pacific region than Murphy and Austin (2005) did, the phylogeny of *Macrobrachium* species based on mtDNA 16S rRNA still showed poorly resolved “starburst” relationships, which lack internal structure with short internal branch lengths and longer tips among species of *Macrobrachium* (Fig. 2). Such phylogenetic relationships may have been caused by a weakness of the marker when it reaches saturation or by a lack of power of the data to resolve relationships among taxa (Albertson et al. 1999). 16S rRNA, used in this study, is not saturated (data not shown). Alternatively, the unresolved phylogeny detected herein may also be explained by rapid radiation, as suggested by earlier studies of *Sebastes* rockfishes (Johns and Avise 1998), fairy shrimp (Daniels et al. 2004), Caribbean sponge-dwelling snapping shrimp (Morrison et al. 2004), squat lobsters (Machordoma and Macpherson 2004), and freshwater crayfish (Shull et al. 2005). In all of those cases, resolution and/or support for the nodes in question were poor, suggesting a real phenomenon resulting from rapid radiation, rather than simply a paucity of appropriate data. When we focused on prawn species distributed in East Asia, the combined dataset of 16S and COI sequences revealed the same pattern of phylogenetic relationships (Fig. 3), whereas relationships among many of the terminal taxa were moderately to well supported which indicates several important features.

The monophyly of the genus *Macrobrachium* with the outgroup was supported by the phylogenetic analyses (Figs. 2, 3); *M. intermedium* was excluded from the genus *Macrobrachium* in earlier studies (Pereira 1997, Murphy and Austin 2002 2003, Short 2004). Multiple origins of

*Macrobrachium* fauna on various continents (or regions), like Central and South America, East Asia, the Indo-West Pacific, and India (Fig. 2), are suggested, supporting the results of Murphy and Austin (2005). The planktonic larval stage with salinity tolerance may play an important role in its long-distance dispersal and may have contributed to the widespread distribution of species (Shokita 1985, Jalihal et al. 1993, de Bruyn et al. 2004), such as those of *M. latidactylus*, *M. latimanus*, *M. grandimanus*, *M. rosenbergii*, and *M. lar*. Minor genetic differences (0.00%-3.20% for 16S and 0.00%-12.63% for COI) among conspecific, widespread euryhaline populations over broad geographic areas (Tables 1, 2, Figs. 2, 3), suggest that gene flow has been continually ongoing, and that they tend to have low levels of differentiation compared to species with a non-planktonic larval phase (Cameron 1986).

None of the land-locked species, including *Exopalaemon modestus* and *Palaemonetes sinensis*, formed a monophyletic group, or was located in the basal position, suggesting that they did not diverge from a single marine ancestor, but likely originated from marine ancestors and subsequently moved towards fresh water in multiple waves of migration (Figs. 2, 3) (Tiwari 1955, Jalihal et al. 1993). An abbreviated larval development pattern (ovigerous female with large eggs) has been suggested as being a process which resulted from selective pressures caused by attempts to become established in freshwater environments, and is a result of adaptive convergence (Shokita 1979, Magalhães and Walker 1988). This pattern is parallel to that of another freshwater shrimp in the family Atyidae (Magalhães and Walker 1988) and also in Jamaican's *Sesarma* crabs (Schubart et al. 1998).

Jayachandran's (2001) suggestion that the genus *Macrobrachium* could be grouped into 2 categories based on morphological characters of the 2nd pereopod is not supported by our findings. Our results (Figs. 2, 3) demonstrated that grouping species according to the most-common morphological characters used in the taxonomy of the genus *Macrobrachium* (e.g., the 2nd pereopod or the rostrum) does not form a monophyletic group, and these characters do not always have phylogenetic value. Morphological characters (such as the unequal 2nd pereopod, big robust claws, spine, etc.) likely independently developed during the invasion of inland waters.

## Taxonomic implications

Based on mtDNA sequences, 11 of 15 species we obtained from geographically distant populations formed monophyletic lineages; 4 undescribed species (*M. spp.1-4*) were identified, and 4 cryptic species (*M. spp.5-8*), respectively grouped with *M. latidactylus*, *M. latimanus*, *M. jaroense*, and *M. placidulum*, were inferred according to the phylogenetic reconstructions and sequence divergence levels. The incongruence of the non-monophyly of *M. equidens* in the 16S, COI, and 16S+COI analyses (Figs. 2, 3, COI tree not shown) implies the existence of a 5th cryptic species.

Some misidentifications or invalid species were also revealed. The species *M. hainanense* (accession no.: AY377841), collected from Hong Kong by Murphy and Austin (2005) was inconsistent with our specimens from 2 Chinese populations (sample localities of Guangdong Prov. and Hainan I., Table 1) with a significant genetic distance (7.6% in 16S) and different position in the phylogenetic tree (Fig. 2). Holthuis (1950) commented that *M. hainanense* was so closely related to *M. formosense* that it should perhaps only be considered a subspecies. According to the divergence and phylogenetic tree (Fig. 2), *M. hainanense* used by Murphy and Austin (2005) is most probably a "misidentified species". Examination of Hong Kong specimens by the second author showed that there is an undescribed species of the *Macrobrachium asperurum* species group in Hong Kong. The subadult specimens of this species could easily be confused with those of *M. hainanense*, which also occurs in Hong Kong.

*Macrobrachium anhuiense* and *M. pinguis* formed sister pairs to *M. asperurum* distributed in northern and southern Taiwan (Table 1), respectively, with high bootstrap support, and showed intraspecific levels of genetic divergences (0.90%-1.64% for 16S and 2.52%-4.08% for COI) (Table 3) in the 16S and combined dataset analyses (Fig. 2, 3). Liu et al. (1990) suggested that *M. pinguis* is an invalid species described based on undeveloped males, and was synonymous with *M. asperurum*. *Macrobrachium anhuiense* was also excluded from the species list of Chinese palaemonoid fauna (Li et al. 2003). The present DNA data support both taxonomic actions. Neither species, based on morphological and phylogenetic evidence, could be separated from *M. asperurum*, and both should be treated as invalid species.

*Macrobrachium shokitai* has a restricted distribution on Iriomote I. of the southern Ryukyus, Japan and is closely related morphologically and ontogenetically to *M. asperulum*. When these 2 species were crossed, their hybrid was found to be sterile. Shokita (1979) considered *M. shokitai* to be an offshoot of *M. asperulum*.

When the combined dataset was analyzed, a species group containing *M. cf. horstii*, *M. placidulum*, *M. placidum*, and *M. lepidactyloides* formed a monophyletic group, species group 4 (SpG 4) (Fig. 3). Molecular studies did not support the close relationship of *M. lepidactyloides* and *M. placidum* (with genetic distances of 12.97% for 16S and 19.26% for COI) (Table 3) as indicated by morphological similarities. This contradicts a suggestion by Chace and Bruce (1993) that *M. lepidactyloides* may be synonymous with *M. placidum* based on morphology. The species *M. cf. horstii* (cf. Shy and Yu 1998) was closely allied to *M. placidulum* with an intraspecific level of genetic divergence (0.37% for 16S and 1.33% for COI) (Table 3), and it is also morphologically similar to the latter species; thus, it should be regarded as conspecific with *M. placidulum*.

The 4 cryptic species, (*M. spp.*5-8, and the incongruence of the non-monophyly of *M. equidens* (Figs. 2, 3) imply the existence of a 5th cryptic species, which showed minor morphological differences of “intraspecific” variations. However, they were genetically very distinct from other populations (localities) with values of interspecific divergences (4.9%-9.2% for 16S and 13.2%-17.2% for COI) and were closely allied with high bootstrap values. This also suggests that the use of traditional morphological characters alone is insufficient to accurately diagnose natural species groups of *Macrobrachium* (Holthuis 1950 1952, Johnson 1973).

In our study, the intraspecific 16S sequence divergence estimates between populations (or individuals) from different localities ranged 0.0%-3.2%. The significant divergence of 16S (5.1%-6.2%) between eastern and western *M. rosenbergii* clades along Huxley's line (de Bruyn et al. 2004) is far beyond the ranges of intraspecific divergence determined in the present study when compared with *M. lar*, *M. latidactylus*, *M. mammillodactylus*, *M. latimanus*, and *M. grandimanus* which are also distributed across Huxley's line. Genetic evidence in the present study supports the suggestion of previous studies (Johnson 1973, Lindenfelser 1984, Wowor and Ng 2001, de Bruyn et al. 2004) that *M. rosenbergii* may actually represent 2 distinct taxa: eastern and western forms.

*Macrobrachium* is a notoriously difficult genus taxonomically, as the morphological plasticity of taxonomically important traits (e.g., the rostrum and/or the 2nd pereopod) change so much and so gradually during their growth (Holthuis 1950) and are influenced by environmental parameters (Dimmock et al. 2004). Morphologically similar species are often quite genetically distinct. However, this might not be reflected in the phylogenetic relationships, as shown in the *M. equidens* species group (Johnson 1973) (Figs. 2, 3), suggesting that conservative systematic traditions or morphological stasis may be involved (Knowlton 2000). Most genetic analyses of species boundaries in marine crustaceans (see Knowlton 2000 for review) and freshwater macroinvertebrates (Baker et al. 2004, Shih et al. 2004 2005 2006) confirm or reveal the existence of cryptic species that are difficult to distinguish using traditional techniques. Some cryptic species are distinguished by surprisingly large genetic differences (Knowlton 2000, Hendrixon and Bond 2005, Ellis et al. 2006). This problem highlights a number of features of species group in these analyses. First, despite the diversity of species of this genus, they are all relatively conservative in general appearance, and taxonomic mistakes are easily made. Second, it shows the considerable value of having multiple samplings within each taxonomic group of interest, so that possible errors can be detected (Sites and Marshall 2003, Peters et al. 2005). It also suggests that using a single example of an individual or population to represent a species in the overall analysis is not justified and should be treated with caution. It is necessary to reevaluate the practical species concept based on such a multiple-sample analysis (Figs. 2, 3). The cryptic species detected here (Figs. 2, 3) suggest that the use of molecular techniques will be a significant help in delimiting species and understanding their relationships (Knowlton 2000, Hendrixon and Bond 2005, Ellis et al. 2006).

Some of the species groups, including species spanning different geographic regions, could be suggested from Central and South America, India, the Indo-West Pacific, and East Asia (Fig. 2). *Macrobrachium rosenbergii*, the widely distributed euryhaline species (De Man 1879, Johnson 1973) with an extended type of larval development and regarded as probably fairly “ancient” in nature by Johnson (1973), was not placed at the basal position of the *Macrobrachium* species clade. However, it formed a species group, as suggested by Johnson (1973), together

with 2 Indian euryhaline species, *M. gangeticum* and *M. malcolmsonii*, and 2 land-locked species endemic to India, *M. lamarrei* and *M. sankollii* (Fig. 2). Species groups containing such a mixture of species with different life cycles suggests that the species group may have evolved from a single ancestral lineage. Such a lineage was also found in species group 2 (SpG 2) containing *M. nipponense*, *M. inflatum*, *M. formosense*, *M. hainanense*, and 1 undescribed species *M. sp.4* (Figs. 2, 3), and species group 3 (SpG 3), another euryhaline species group with an asymmetrical 2nd pereopod, containing *M. lanatum*, *M. esculentum*, *M. latidactylus*, and cryptic species *M. sp.5*, (Fig. 3). This euryhaline species group contains a land-locked species endemic to Australia, *M. handschini*, which formed a sister pair with *M. esculentum* (specimens collected from Taiwan and the Philippines) in the 16S analysis (Fig. 2).

An endemic speciation event in East Asia was suggested in the land-locked species group, including the *M. asperulum* species group (species group 1) (Fig. 3). This land-locked species group did not form a monophyletic group with the land-locked species endemic to Southeast Asia (*M. yui*, *M. malayanum*, and *M. platycheles*), which implies a single lineage. Among them, *M. asperulum* is the most widely distributed species, being known from southern Siberia to southeastern China (Holthuis 1950). The other land-locked species have restricted distribution ranges. For freshwater-dependent prawns, factors responsible for dispersal generally involve land continuity and therefore river confluences (e.g., during sea-level lowering in glacial maxima), as well as river capture in headwaters (Banareescu 1990). The best explanation is that it represents fragmentation of a widespread ancestral taxon (vicariance) through allopatric speciation (Qian and Ricklefs 2000), rather than a dispersal phenomenon from a more-restricted "center of origin" (Wiley 1988).

In addition to the ancient speciation by radiation, there is evidence for ongoing freshwater invasion and recent speciation in East Asia. The processes of freshwater invasion and penetration of cool temperate areas (at high latitudes) may be represented by *M. nipponense*. *Macrobrachium nipponense* is distributed along the coastline from northern Southeast Asia north to East Asia and Japan (Liu et al. 1990, Cai and Dai, 1999). Its narrow salinity tolerance, and relatively shorter zoeal period may have limited the dispersal distance to a dispersal pattern from estuary to estuary, apparently not a transoceanic dispersal in which larval

stages of euryhaline species can disperse across the ocean by ocean currents as can *M. gradimanus* (Shokita 1985) and *M. rosenbergii* (de Bruyn et al. 2004). Some *M. nipponense* populations, on the basis of allozyme variation and reproductive traits, have apparently split into freshwater and estuarine populations in the same river in Japan (see Mashiko and Numachi 2000 for review), while populations in China and Taiwan are now land-locked species (Liu et al. 1990, Cai and Dai, 1999). This probably represents different steps in the process of inland water invasion. The land-locked populations of *M. nipponense* in China and Taiwan represent an advanced state of freshwater invasion, while populations of *M. nipponense* in Japan may represent both the advanced state (the land-locked populations inhabiting fresh water) and a euryhaline state. *Macrobrachium hainanense* and *M. formosense* are closely related species morphologically; Holthuis (1950) commented that *M. hainanense* should perhaps only be considered a subspecies. These 2 species exhibited the lowest interspecific DNA divergence (Table 3) and are closely related (Figs. 2, 3). Based on the fact that *M. formosense* has a more-restricted distribution than *M. hainanense* (Li et al. 2003); we believe that *M. formosense* may represent a newly derived species in recent geological times.

This study was based on only 2 mitochondrial markers. Further studies, particularly using other mitochondrial or nuclear sequence data and including more species, are obviously required to further investigate the evolutionary history of this genus. New data on the biology and ecology of these species and their habitats, and updated knowledge of the paleogeographical history will help clarify the possibility of microhabitat and behavioral specialization giving rise to radiation by *Macrobrachium*.

**Acknowledgments:** We thank Ken-Ichi Hayashi, Tohru Naruse, Daisy Wowor, and Te-Yu Liao for specimen collection. Tzi-Yuan Wang helped with data analyses. Thanks are given to Peter K.L. Ng for helpful discussion and the Zoological Reference Collection (ZRC) of the Raffles Museum of Biodiversity Research, National University of Singapore (NUS) for providing specimens for morphological study. We particularly thank Hsi-Te Shih for specimen collection, data analyses, and discussions. Two anonymous reviewers who greatly improved this manuscript are also acknowledged.

## REFERENCES

- Albertson RC, JA Markert, PD Danley, TD Kocher. 1999. Phylogeny of a rapidly evolving clade: the cichlid fishes of Lake Malawi, East Africa. *Proc. Natl. Acad. Sci. USA*. **96**: 5107-5110.
- Baker AM, JM Hughes, JC Dean, SE Bunn. 2004. Mitochondrial DNA reveals phylogenetic structuring and cryptic diversity in Australian freshwater macroinvertebrate assemblages. *Mar. Freshwater Res.* **55**: 629-640.
- Banarescu P. 1991. Distribution and dispersal of freshwater animals in North America and Eurasia. *In: Zoogeography of fresh waters*. Vol. 2. Wiesbaden, Germany: AULA-Verlag, pp. 512-1091.
- Cai Y, A Dai. 1999. Freshwater shrimps (Crustacea: Decapoda: Caridea) from the Xishuangbanna region of Yunnan Province, southern China. *Hydrobiologia* **400**: 211-241.
- Cai Y, P Naiyanetr, PKL Ng. 2004. The freshwater prawns of the genus *Macrobrachium* Bate, 1868, of Thailand (Crustacea: Decapoda: Palaemonidae). *J. Nat. Hist.* **38**: 581-649.
- Cai Y, PKL Ng. 2002. The freshwater palaemonid prawns (Crustacea: Decapoda: Caridea) of Myanmar. *Hydrobiologia* **487**: 59-83.
- Cameron RA. 1986. Introduction to the invertebrate larval biology workshop: a brief background. *Bull. Mar. Sci.* **39**: 145-161.
- Camin JH, RR Sokal. 1965. A method for deducing branching sequences in phylogeny. *Evolution* **19**: 311-326.
- Chace FA, AJ Bruce. 1993. The Caridean shrimps (Crustacea: Decapoda) of the Albatross Philippine expedition, 1907-1910, Part 6: Superfamily Palaemonoidea. Washington, DC: Smithsonian Institution Press, 152 pp.
- Crandall KA, JFJ Fitzpatrick. 1996. Crayfish systematics: using a combination of procedures to estimate phylogeny. *Syst. Biol.* **45**: 1-26.
- Dai AY. 1984. A preliminary study on the freshwater prawn genus *Macrobrachium* of China (Decapoda: Caridea). *Acta Zool. Sin.* **9**: 244-252. (in Chinese with English abstract)
- Daniels SR, M Hamer, C Rogers. 2004. Molecular evidence suggests an ancient radiation for the fairy shrimp genus *Streptocephalus* (Branchiopoda: Anostraca). *Biol. J. Linn. Soc.* **82**: 313-327.
- de Bruyn M, JA Wilson, PB Mather. 2004. Huxley's line demarcates extensive genetic divergence between eastern and western forms of the giant freshwater prawn, *Macrobrachium rosenbergii*. *Mol. Phylogenet. Evol.* **30**: 251-257.
- De Man JG. 1879. On some species of the genus *Palaemon* Fabr. with descriptions of two new forms. *Notes Leyden Mus.* **1**: 165-184.
- Dimmock A, I Williamson, PB Mather. 2004. The influence of environment on the morphology of *Macrobrachium australiense* (Decapoda: Palaemonidae). *Aquacult. Int.* **12**: 435-456.
- Ellis JS, ME Knight, C Carvell, D Goulson. 2006. Cryptic species identification: a simple diagnostic tool for discriminating between two problematic bumblebee species. *Mol. Ecol. Notes* **6**: 540-542.
- Felsenstein J. 1981. Evolutionary trees from DNA sequences: a maximum likelihood approach. *J. Mol. Evol.* **17**: 368-376.
- Felsenstein J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**: 783-791.
- Guo Q, RE Ricklefs, ML Cody. 1998. Vascular plant diversity in eastern Asia and North America: historical and ecological explanations. *Bot. J. Linn. Soc.* **128**: 123-136.
- Gusmão J, C Lazoski, AM Sole-Cava. 2000. A new species of *Penaeus* (Crustacea: Penaeidae) revealed by allozyme and cytochrome oxidase I analyses. *Mar. Biol.* **137**: 435-446.
- Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl. Acids. Symp. Ser.* **41**: 95-98.
- Hamilton W. 1983. Cretaceous and Cenozoic history of the northern continents. *Ann. Mo. Bot. Gard.* **70**: 440-458.
- Hasegawa M, H Kishino, T Yano. 1985. Dating of the human-ape splitting by a molecular clock of mitochondrial DNA. *J. Mol. Evol.* **21**: 160-174.
- Hayashi KI. 2000a. Prawns, shrimp and lobsters from Japan (112). Family Palaemonidae, subfamily Palaemoninae-genus *Macrobrachium*. *Aquabiology* **22**: 240-245. (in Japanese)
- Hayashi KI. 2000b. Prawns, shrimp and lobsters from Japan (113). Family Palaemonidae, subfamily Palaemoninae-genus *Macrobrachium*. *Aquabiology* **22**: 360-363. (in Japanese)
- Hayashi KI. 2000c. Prawns, shrimp and lobsters from Japan (114). Family Palaemonidae, subfamily Palaemoninae-genus *Macrobrachium*. *Aquabiology* **22**: 468-472. (in Japanese)
- Hendrixson BE, JE Bond. 2005. Testing species boundaries in the *Antrodiaetus unicolor* complex (Araneae: Mygalomorphae: Antrodiaetidae): "paraphyly" and cryptic diversity. *Mol. Phylogenet. Evol.* **36**: 405-416.
- Hipp AL, JC Hall, KJ Sytsma. 2004. Congruence versus phylogenetic accuracy: revisiting the incongruence length difference test. *Syst. Biol.* **53**: 81-89.
- Holthuis LB. 1950. The Decapoda of the Siboga Expedition. Part X. The Palaemonidae collected by the Siboga and Snellius expeditions, with remarks on other species, Part I: Subfamily Palaemoninae. *Siboga-Exped. Leiden* **39a**: 1-268.
- Holthuis LB. 1952. A general revision of the Palaemonidae (Crustacea, Decapoda, Natantia) of the Americas. II. The subfamily Palaemoninae. *Occas. Pap. Allan Hancock Found.* **12**: 1-396.
- Hwang JJ, HP Yu. 1982. Studies on the freshwater shrimps of the genus *Macrobrachium* (Crustacea, Decapoda, Palaemonidae) from Taiwan. *J. Taiwan Mus.* **25**: 157-179. (in Chinese with English abstract)
- Hwang JJ, HP Yu. 1983. Key to the freshwater shrimps of the genus *Macrobrachium* from Taiwan with their habitat and distribution. *China Fish.* **326**: 14-18. (in Chinese with English abstract)
- Jalihal DR, KN Sankolli, S Shenoy. 1993. Evolution of larval developmental patterns and the process of freshwaterization in the prawn genus *Macrobrachium* Bate, 1868 (Decapoda, Palaemonidae). *Crustaceana* **65**: 365-376.
- Jayachandran KV. 2001. Palaemonid prawns: biodiversity, taxonomy, biology and management. Enfield, NH: Science Publishers Press.
- Johns GC, JC Avise. 1998. Tests for ancient species flocks based on molecular phylogenetic appraisals of *Sebastes* rockfishes and other marine fishes. *Evolution* **52**: 1135-1146.
- Johnson DS. 1973. Notes on some species of the genus



- Macrobrachium*. J. Singapore Nat. Acad. Sci. **3**: 273-291.
- Knowlton N. 2000. Molecular genetic analyses of species boundaries in the sea. *Hydrobiologia* **420**: 73-90.
- Kotelat M. 1989. Zoogeography of the fishes from Indochinese inland waters with an annotated check-list. *Bull. Zool. Mus. Univ. Amsterdam* **12**: 1-54.
- Kumar S, K Tamura, IB Jakobsen, M Nei. 2001. MEGA2: Molecular Evolutionary Genetics Analysis Software. Tempe, AZ: Arizona State Univ.
- Li X, R Liu, X Liang. 2003. Study on the zoogeography of Chinese Palaemonoid fauna. *In* Transactions of Chinese Crustacean Society no. 4. Beijing: Science Press, pp. 123-145. (in Chinese with English abstract)
- Lindenfelser ME. 1984. Morphometric and allozymic congruence: evolution in the prawn *Macrobrachium rosenbergii* (Decapoda: Palaemonidae). *Syst. Zool.* **33**: 195-204.
- Liu JY, XQ Liang, SL Yan. 1990. A study of the Palaemoninae (Crustacea: Decapoda) from China I. *Macrobrachium, Leander and Leandrites*. *In*: Transactions of the Chinese Crustacean Society Vol. 2. Beijing: Science Press, pp. 102-134. (in Chinese with English abstract)
- Machordoma A, E Macpherson. 2004. Rapid radiation and cryptic speciation in squat lobsters of the genus *Munida* (Crustacea, Decapoda) and related genera in the South West Pacific: molecular and morphological evidence. *Mol. Phylogenet. Evol.* **33**: 259-279.
- Magalhães C, I Walker. 1988. Larval development and ecological distribution of central Amazonian palaemonid shrimps (Decapoda, Caridea). *Crustaceana* **55**: 279-292.
- Mashiko K, KI Numachi. 2000. Derivation of populations with different-sized eggs in the palaemonid prawn *Macrobrachium nipponense*. *J. Crustacean Biol.* **20**: 118-127.
- Morrison CL, R Ríos, JE Duffy. 2004. Phylogenetic evidence for an ancient rapid radiation of Caribbean sponge-dwelling snapping shrimps (*Synalpheus*). *Mol. Phylogenet. Evol.* **30**: 563-581.
- Murphy NP, CM Austin. 2002. A preliminary study of 16S rRNA sequence variation in Australian *Macrobrachium* shrimps (Palaemonidae: Decapoda) reveals inconsistencies in their current classification. *Invertebr. Syst.* **16**: 697-701.
- Murphy NP, CM Austin. 2003. Molecular taxonomy and phylogenetics of some species of Australian palaemonid shrimps. *J. Crustacean Biol.* **23**: 169-177.
- Murphy NP, CM Austin. 2004. Multiple origins of the endemic Australia *Macrobrachium* (Decapoda: Palaemonidae) based on 16S rRNA mitochondrial sequences. *Aust. J. Zool.* **52**: 549-559.
- Murphy NP, CM Austin. 2005. Phylogenetic relationships of the globally distributed freshwater prawn genus *Macrobrachium* (Crustacea: Decapoda: Palaemonidae): biogeography, taxonomy and the convergent evolution of abbreviated larval development. *Zool. Scr.* **34**: 187-197.
- Nylander JAA 2004. MrModeltest v2. Program distributed by the author. Uppsala, Sweden: Evolutionary Biology Centre, Uppsala Univ.
- Palumbi SR, J Benzie. 1991. Large mitochondrial DNA differences between morphologically similar penaeid shrimp. *J. Mol. Mar. Biol. Biotech.* **1**: 27-34.
- Pereira G. 1997. A cladistic analysis of the freshwater shrimps of the family Palaemonidae (Crustacea, Decapoda, Caridea). *Acta Biol. Venez.* **17**: 1-69.
- Pereira G, A Garcia. 1995. Larval development of *Macrobrachium reyesi* Pereira (Decapoda: Palaemonidae), with a discussion on the origin of abbreviated development in Palaemonids. *J. Crustacean Biol.* **15**: 117-133.
- Peters JL, KG McCracken, YN Zhuravlev, Y Lua, RE Wilson, KP Johnson, KE Omland. 2005. Phylogenetics of wigeons and allies (Anatidae: *Anas*): the importance of sampling multiple loci and multiple individuals. *Mol. Phylogenet. Evol.* **35**: 209-224.
- Posada D, KA Crandall. 1998. Modeltest: testing the model of DNA substitution. *Bioinformatics.* **14**: 817-818.
- Qian H, RE Ricklefs. 2000. Large-scale processes and Asian bias in species diversity of temperate plants. *Science* **407**: 180-182.
- Rodriguez R, JL Oliver, A Marin, JR Medina. 1990. The general stochastic model of nucleotide substitution. *J. Theor. Biol.* **142**: 485-501.
- Ronquist F, JP Huelsenbeck. 2003. MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**: 1572-1574.
- Rozas J, JC Sanchez-DelBarrio, X Messeguer, R Rozas. 2003. DNASP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics* **19**: 2496-2497.
- Saitou N, M Nei. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**: 406-425.
- Sambrook J, EF Fritsch, T Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. New York: Cold Spring Harbor Laboratory Press.
- Schubart CD, R Diesel, SB Hedges. 1998. Rapid evolution to terrestrial life in Jamaican crabs. *Nature* **393**: 363-365.
- Shih HT, GX Chen, LM Wang. 2005. A new species of freshwater crab (Decapoda: Brachyura: Potamidae) from Dongyin Island, Matsu, Taiwan, defined by morphological and molecular characters, with notes on its biogeography. *J. Nat. Hist.* **39**: 2901-2911.
- Shih HT, HC Hung, CD Schubart, CA Chen, HW Chang. 2006. Intraspecific genetic diversity of the endemic freshwater crab *Candidiopotamon rathbunae* (Decapoda, Brachyura, Potamidae) reflects five million years of geological history of Taiwan. *J. Biogeogr.* **33**: 980-989.
- Shih HT, PKL Ng, HW Chang. 2004. Systematics of the genus *Geothelphusa* (Crustacea, Decapoda, Brachyura, Potamidae) from southern Taiwan: a molecular appraisal. *Zool. Stud.* **43**: 561-570.
- Shokita S. 1979. The distribution and speciation of the inland water shrimps and prawns from the Ryukyu Islands-II. *Bull. Tokai Reg. Fish. Res. Lab.* **28**: 193-278. (in Japanese with English abstract)
- Shokita S. 1985. Larval development of the Palaemonid prawn, *Macrobrachium grandimanus* (Randall), reared in the laboratory, with special reference to larval dispersal. *Zool. Sci.* **2**: 785-803.
- Short JW. 2004. A revision of Australian river prawns, *Macrobrachium* (Crustacea: Decapoda: Palaemonidae). *Hydrobiologia* **525**: 1-100.
- Shull HC, M Perez-Losada, D Blair, K Sewell, EA Sinclair, S Lawler, M Ponniah, KA Crandall. 2005. Phylogeny and biogeography of the freshwater crayfish *Euastacus* (Decapoda: Parastacidae) based on nuclear and mitochondrial DNA. *Mol. Phylogenet. Evol.* **37**: 249-263.
- Shy JY, HP Yu. 1998. *Freshwater shrimps of Taiwan*. Kaohsiung, Taiwan: National Museum of Marine Biology and Aquarium Press. (in Chinese)
- Sites JW, J Marshall. 2003. Delimiting species: a Renaissance issue in systematic biology. *Trends Ecol. Evol.* **18**: 462-

- 470.
- Swofford DL. 2000. PAUP\*: Phylogenetic Analysis Using Parsimony (\* and other methods). Vers. 4. Sunderland, MA: Sinauer Associates Press.
- Tamura K, M Nei. 1993. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Mol. Biol. Evol.* **10**: 512-526.
- Tiwari KK. 1955. Distribution of Indo-Burmese freshwater prawns of the genus *Palaemon* Fabr., and its bearing on the Satpura hypothesis. *Bull. Natl. Inst. Sci. India.* **7**: 230-239.
- Wiley E. 1988. Vicariance biogeography. *Ann. Rev. Ecol. Syst.* **19**: 513-542.
- Wowor D, SC Choy. 2001. The freshwater prawns of the genus *Macrobrachium* Bate, 1868 (Crustacea: Decapoda: Palaemonidae) from Brunei Darussalam. *Raffles Buff. Zool.* **49**: 269-289.
- Wowor D, Ng PKL. 2001. Identity of the giant prawn, *Macrobrachium rosenbergii* (De Man, 1879) (Crustacea: Decapoda: Caridea: Palaemonidae). *In*: Proceedings of the Fifth International Crustacean Congress, Melbourne, Australia, 9-13 July 2001. Australia: The Crustacean Society.
- Yu SC. 1936. Notes on new freshwater prawns of the genus *Palaemon* from Yunnan. *Bull. Fan Mem. Inst. Biol. (Zool.)* **6**: 305-314.