

Chan, Tin-Yam & Hsiang-Ping Yu
1986

CRUSTACEA LIBRARY
SMITHSONIAN INSTITUTION
RETURN TO VENTS

ASIAN MARINE BIOLOGY (1986). 3: pp 89-99

THE DEEP-SEA SHRIMPS OF THE FAMILY OPLOPHORIDAE (CRUSTACEA: DECAPODA) FROM TAIWAN*

Tin-Yam Chan and Hsiang-Ping Yu

Department of Aquaculture; and Graduate School of Fisheries; National Taiwan College of Marine Science and Technology, Keelung, 200 Taiwan, Republic of China.

Abstract

This paper reports upon four species of deep-sea oplophorid shrimps collected off north-east and southern Taiwan at depths of about 100–450 m. All are new to the marine fauna list of Taiwan. *Oplophorus gracilirostris* and *Oplophorus typus* are found to have the ability to lock the laterally extended scaphocerites. A key is given for their identification and their colours are fully illustrated. Sexual differences are also discussed.

Introduction

Most oplophorids are well-known bathypelagic shrimps that have the ability to luminesce and exhibit vertical migration. Interestingly, a number of specimens belonging to this family have been obtained from the coastal shrimp fishery and are used by local fishermen as part of the natural food in the fish and prawn culture industry. In the present paper, four oplophorid species are described from our collections, namely: *Oplophorus typus* H. Milne Edwards, 1837, *Oplophorus gracilirostris* A. Milne Edwards, 1881, *Oplophorus spinicauda* A. Milne Edwards, 1883 and *Systellaspis pellucida* (Filhol, 1885). All have the same local name "spiny shrimp" and are new to the fauna list of Taiwan. The first two species have the ability to lock their laterally extended scaphocerite by unique structures on the scaphocerite and the basicerite. This report includes descriptions of the external characteristics and the sexual differences of these shrimps, together with keys and colour illustrations. Restricted synonymies are also provided.

Materials and Methods

All the specimens were obtained from local fish markets at fishing harbours and were said to be caught at about 100–450 m depth off north-east and southern Taiwan. Reference specimens are deposited in the Fisheries Department of the National Taiwan College of Marine Science and Technology. The stated measurement is body length which was measured from the postorbital margin to the posterior margin of the telson with the specimen outstretched.

Key to the genera of Oplophoridae from Taiwan:

- A. Exopods of all thoracic appendages conventionally slender; outer margin of scaphocerite smooth; abdominal somite VI never dorsally carinate *Systellaspis* Bate, 1888
- B. Exopods of maxilliped III and pereopod I foliaceous; outer margin of scaphocerite usually spinous (all Taiwan species with outer margin of scaphocerite spinous)
. *Oplophorus* H. Milne Edwards, 1837

* Contribution No. 975-0409-B019-01 from the National Science Council, Republic of China.

Genus *Systellaspis* Bate, 1888
Systellaspis pellucida (Filhol, 1885)
(Plate IA; Figure 1)

Acanthephyra pellucida -- Filhol, 1885: 144, 162.

Systellaspis affinis -- Des Man, 1920: 43; Chace, 1936: 29; Calman, 1939: 190; Dennell, 1940: 345; Chace, 1947: 39, fig. 3; Crosnier and Forest, 1968: 1133.

? not *Systellaspis lanceocaudata* Bate, 1888 -- Bals, 1925: 243, figs. 12, 13.

Systellaspis pellucida -- Crosnier and Forest, 1973: 92, figs. 26c, 27c.

• 'Material Examined': Ta-Chi, I-Lan County;

Chan leg.: 14 October 1984; 1 ♂ 54 mm, 4 ovigerous ♀♀ 57–62 mm. 31 December 1984; 1 ovigerous ♀ 59 mm, 1 ♀ 60 mm. 26 March 1985; 4 ovigerous ♀♀ 54–56 mm, 25 ♀♀ 52–57 mm. 8 May 1985; 1 ♂ 52 mm, 5 ovigerous ♀♀ 50–53 mm, 5 ♀♀ 48–52 mm. 14 July 1985; 3 ♂♂ 41–44 mm, 4 ♀♀ 47–50 mm. Tong-Kang, Ping-Tong County; Chan leg.: 31 October 1984; 2 ovigerous ♀♀ 56 and 62 mm. 2 December 1984; 2 ovigerous ♀♀ 56 and 62 mm. 2 December 1984; 3 ♀♀ 51–58 mm. Su-Ao, I-Lan County; Chan leg.: 3 May 1985; 2 ovigerous ♀♀ 51 and 52 mm, 2 ♀♀ 52 and 53 mm.

Body slender and shell flexible. Rostrum




Fig. 1. *Systellaspis pellucida*.
A, Pleopod II of male (X 12.5);
B, Abdominal pleura I and II of male (X 12.5);
C, Abdominal pleura I and II of female (X 12.5).




Plate I. A, Lateral view of a 53 mm ovigerous female of *Systellaspis pellucida*, showing partly scarlet and partly white eggs; B, Lateral view of a 47 mm ovigerous female of *Oplophorus spinicauda*, showing near hatching white eggs; C, Lateral view of a 56 mm ovigerous female of *Oplophorus gracilirostris*, showing scarlet eggs; D, Lateral view of a 39 mm female of *Oplophorus typus*.

armed dorsally with 4–6 (usually 6) teeth and ventrally with 4–6 (usually 5) teeth. Usually 3–4 post-rostral teeth present. Rostrum generally slightly longer than scaphocerite. No lateral carinae on carapace. Exopods of all pereopods slender. Abdominal tergites III, IV and V produced into short but acute spine posteriorly. Hind margins of abdominal tergites IV and V not serrated laterally. Dorsal midline of abdominal tergite VI and anterior telson rounded or flattened, not sulcate. Telson more or less as long as uropods (slightly shorter in some specimens). Eggs few, ranging from 6–18 per ovigerous female, large and beanshaped, about 2×3 mm.

- **Colour:** Generally transparent but covered with scattered scarlet spots. Eye brown-red. Internal organs inside carapace and eggs scarlet. Carapace with violet-blue longitudinal bars (photophores) near ventral margin. Thoracic appendages, bases of pleopods and posterior margins of posterior three abdominal segments vermilion. Photophores of deep purple colour.
- **Distribution:** World wide tropical seas: Philippines, Malay Archipelago, West Africa, Bahamas, West Indies, China Sea and Zanzibar.
- **Remarks:** Detailed morphological figures have been given by Crosnier and Forest (1973). The distribution and structure of the photophores have been extensively studied by Calman (1939) and Dennell (1940). The male specimens investigated in this study possess an appendix masculina on the endopod of the pleopod II and both appendix interna and appendix masculina are translucent and slender. The sex of the species can be somewhat determined from the outline of the abdominal pleuron I, where the lower margin is concave in the male but is nearly straight in the female. Moreover, in males, a median blunt protuberance is present on the abdominal sternites I and II, the pleopods of the abdominal segment III are very close to each other and the coxae of the anterior three pleopods are relatively short, about half the length of the basis. However, in females only a

rudimentary median elevation is visible on the abdominal sternites I and II, the pleopods of the abdominal segment III are widely separated and the coxae of the anterior three pleopods are long, markedly longer than half the length of the basis. *S. pellucida* is relatively larger than other local oplophorids and more numerous off north-eastern Taiwan.

Genus *Oplophorus* H. Milne Edwards, 1837

Key to the species of Oplophorus from Taiwan:

1. Abdominal somites II, III and IV produced into overhanging spine posteriorly
..... *Oplophorus spinicauda*
A Milne Edwards, 1883
- Abdominal somites III, IV and V produced into overhanging spine posteriorly 2
2. Rostrum provided with teeth near tip of scaphocerite; posterior spine of abdominal somite V shorter than IV
..... *Oplophorus gracilirostris*
A. Milne Edwards, 1881
- Rostrum devoid of teeth near tip of scaphocerite; posterior spine of abdominal somite V longer than IV
..... *Oplophorus typus*
H. Milne Edwards, 1837

Oplophorus spinicauda A. Milne Edwards, 1883
(Plate IB; Figure 2)

Oplophorus spinicauda -- A. Milne Edwards, 1883: pl. 30; Chace, 1936: 30; Chace, 1940: 184, fig. 54; Chace, 1947: 42; Hayashi and Miyake, 1969: 68; Crosnier and Forest, 1973: 24; Hanamura, 1979: 171; Miyake, 1982: 174.

Oplophorus foliaceus -- Rathbun, 1906: 922, pl. 20 fig. 8.

Hoplophorous spinicauda -- De Man, 1920: 42.

Hoplophorous foliaceus -- Balss, 1925: 249.

- **Material Examined:** Tong-Kang, Ping-Tong County: 27 January 1983; 1 ♀ 43.5 mm. Yu leg. 31 October 1984; 4 ovigerous ♀♀ 43–47 mm. 2 December 1984; 1 ♂ 32 mm, 17 ovigerous ♂♂ 42–54 mm. 26 March 1985; 1




Fig. 2. *Oplophorus spinicauda*.

- A, Scaphocerite, latero-ventral view (X 12.5);
 B, Lateral view of scaphocerite and basicerite (X 12.5);

♂ 26 mm, 2 ovigerous ♀♀ 44 and 45 mm. 28 July 1985; 3 ovigerous ♀♀ 45–47 mm. Chan leg.

Body slender and shell flexible. Rostrum always with 8 teeth above and 7 teeth below, extending slightly beyond scaphocerite. Post-rostral carina usually provided with 4 teeth. Outer margin of scaphocerite armed with 4–5 large acute spines. Exopods of all pereopods foliaceous. Conspicuous overhanging median posterior spine present on abdominal tergites II, III and IV. Telson more or less longer than uropods. Eggs large, few (about 10 in one ovigerous female) and bean-shaped, about 2×2.5 mm.

- **Colour:** Similar to *S. pellucida* but with scattered red dots and no violet longitudinal bars (photophores) on ventral carapace. Lateral surfaces of abdomen with transparent, yellow-green, patches and tips of uropods reddish. Eggs scarlet, becoming white and with blackish eye spot when close to time of hatching.
- **Distribution:** World wide tropical seas: Belize, Morocco, northern Madagascar, southern India, Hawaiian Islands, Florida, West Indies, Philippines, E. Indian Ocean and Japan.
- **Remarks:** As pointed out by Chace (1940) and Hayashi and Miyake (1969), there is no appendix masculina on the endopod of the pleopod II in males. The sex of the species can, however, be determined by the shape of

the lower margin of the abdominal pleuron I, which is concave and rough in males whereas it is horizontally straight and smooth in females. No median protuberance is present on the abdominal sternites I and II of either sex but the appearance of the coxae of the anterior three pleopods are similar to those of *S. pellucida*. This species closely resembles *S. pellucida* and is very different from other *Oplophorus* species in the general body shape, soft integument, non-rigid exopods of the pereopods, shape of the rostrum and telson, and the outline of the abdominal pleuron I in each sex. Moreover, the appearance of the appendix interna is also thin and slender. Additional to lacking the ability to lock the scaphocerite as in other *Oplophorus* species reported upon in this paper, it appears rather unsuitable to place this species in the genus *Oplophorus* only on account of the characteristic foliaceous exopods of the pereopods, as this character is somewhat vague. This species is mainly found off southern Taiwan and is smaller than other local oplophorids.

Oplophorus gracilirostris A. Milne
 Edwards, 1881

(Plates IC and IIA, B; Figures 3, 4)

Oplophorus gracilirostris -- A. Milne Edwards, 1881: 6; Rathbun, 1906: 921; Yokoya, 1933: 29, fig. 12; Chace, 1936: 30; Chace, 1947: 44, figs. 4–7; George and Rao, 1966: 329; Kensley, 1972: 38, fig. 17F; Hanamura, 1979: 171; Miyake, 1982: 174.

Oplophorus longirostris -- Bate, 1888: 765, pl. 127 fig. 2.

Hoplophorus Smithii -- Wood-Mason and Alcock, 1891: 194.

Hoplophorus gracilirostris -- Alcock, 1901: 73; Kemp and Sewell, 1912: 20; De Man, 1920: 42.

? not *Hoplophorus typus* (H. Milne Edwards, 1837) -- Blass, 1925: 248, figs. 21–23 (in part); Calman, 1939: 188 (in part).