

Joel W. Martin


419

a catalogue &
bibliography to the

**CRABS (BRACHYURA)
OF THE
GULF OF MEXICO**

LAWRENCE W. POWERS

contributions in marine science—supplement to vol. 20



9119

Joel W. Martin

CONTRIBUTIONS IN MARINE SCIENCE
Supplement to Volume 20
December 1977

A Catalogue and Bibliography
TO THE
CRABS (BRACHYURA) OF THE GULF OF MEXICO

LAWRENCE W. POWERS¹

PORT ARANSAS MARINE LABORATORY
University of Texas Marine Science Institute
Port Aransas, Texas 78373

December 1977

¹ Present address: Biology Department, City College (CUNY), New York, New York 10031.
University of Texas Marine Science Institute Contribution No. 215.

Editor: DONALD E. WOHLSCHLAG
Editorial Assistant: RUTH GRUNDY

EDITOR'S NOTE

We are offering, "A Catalogue and Bibliography to the Crabs (Brachyura) of the Gulf of Mexico," By Lawrence W. Powers as a supplement to Volume 20 of our Contributions in Marine Science. We hope that this type of monographic work will stimulate further syntheses.

Additional copies of this Catalogue and Bibliography may be obtained by writing:

The Librarian
Port Aransas Marine Laboratory
The University of Texas Marine Science Institute
Port Aransas, Texas 78373

Price: \$8.00 per copy

ACKNOWLEDGMENTS

I wish to express my appreciation for those who reviewed the preliminary draft of this manuscript. The following offered valuable suggestions for its improvement: Lawrence G. Abele, Fenner A. Chace, Jr., Darryl L. Felder, John S. Garth, and Austin B. Williams. Samuel R. Haley and Warren Pulich reviewed the revised introduction. I assume full responsibility for remaining errors and omissions.

I would like to dedicate this effort to Mrs. Ruth Grundy, librarian of the Marine Science Laboratory at Port Aransas, for her initial encouragement of this project, continued enthusiasm during compilation, patience with the many missed deadlines, and her labors to help see it through to completion. I also appreciate the help and support of others at the Marine Science Laboratory, especially Donald Wohlschlag, William Brogden, Carl Oppenheimer, Sarah Lowrie, Phyllis Picarazzi, and Mary Ellen Tomberlein. Dinah Bowman prepared the cover illustration.

TABLE OF CONTENTS

	Page
Acknowledgments	3
Table of Contents	5
INTRODUCTION	7
CLASSIFICATION	8
BIOGEOGRAPHY	10
DISCUSSION	13
SPECIES ENTRIES	19
BRACHYURA	19
Section DROMIACEA	19
Superfamily DROMIOIDEA	19
Family DROMIIDAE	19
Family PROSOPIDAE	21
Subfamily HOMOLODROMIINAE	21
Superfamily HOMOLOIDEA	21
Family HOMOLIDAE	21
Family LATREILLIDAE	23
Section OXYSTOMATA	23
Superfamily RANINOIDEA	23
Family RANINIDAE	23
Superfamily DORIPPOIDEA	26
Family DORIPPIDAE	26
Superfamily CALAPPOIDEA	29
Family CALAPPIDAE	29
Subfamily CALAPPINAE	29
Subfamily MATUTINAE	32
Family LEUCOSIIDAE	35
Subfamily EBALIINAE	35
Subfamily LEUCOSIINAE	36
Subfamily PHILYRINAE	38
Section HAPALOCARCINIDEA	40
Superfamily HAPALOCARCINOIDEA	40
Family HAPALOCARCINIDAE	40
Section OXYRHYNCHA	41
Family MAJIDAE	41
Subfamily ACANTHONYCHINAE	41
Subfamily INACHINAE	42
Subfamily MAJINAE	49
Subfamily MITHRACINAE	50
Subfamily OPHTHALMIINAE	60
Subfamily PISINAE	62
Family PARTHENOPIDAE	67
Subfamily PARTHENOPINAE	67

Section CANCRIDEA	71
Superfamily CANCROIDEA	71
Family ATELECYCLIDAE	71
Subfamily ATELECYCLINAE	71
Family CANCRINAE	71
Subfamily CANCRINAE	71
Section BRACHYRHYNCHA	72
Superfamily PORTUNOIDEA	72
Family PORTUNIDAE	72
Subfamily POLYBIINAE	72
Subfamily PORTUNINAE	74
Superfamily XANTHOIDEA	86
Family POTAMIDAE	86
Family XANTHIDAE	87
Family GERYONIDAE	110
Family GONEPLACIDAE	111
Family PALICIDAE	117
Family PINNOTHERIDAE	119
Subfamily PINNOTHERINAE	119
Subfamily PINNOTHERELIINAE	125
Family GRAPSIDAE	128
Subfamily GRAPSINAE	128
Subfamily PLAGUSIINAE	131
Subfamily SESARMINAE	132
Subfamily VARUNINAE	136
Family GECARCINIDAE	138
Superfamily OCYPODOIDEA	140
Family OCYPODIDAE	140
Subfamily OCYPODINAE	140
BIBLIOGRAPHY	150
INDEX	184

INTRODUCTION

The purpose of this catalogue and bibliography is to provide a convenient reference source for information on the brachyuran (true) crabs of the Gulf of Mexico. Although it is intended primarily for zoologists and others familiar with taxonomic resources, the catalogue includes some informal features in order to increase its usefulness to students and persons without systematic experience. The catalogue is a compilation of recent literature and the taxonomic organization of the species entries represents the author's interpretation of revisions by carcinological authorities. As a reference source, it is intended to be a compromise between the systematic works with formal synonymies and descriptions of species and various bibliographies that could be generated by the use of computerized key word entries.

The major part of the catalogue consists of a checklist of species from the Gulf of Mexico (referred throughout the text as the Gulf) that was compiled from the published literature. Unpublished materials, such as theses, manuscripts, and uncirculated reports, were included sparingly. Doctoral dissertations were cited in cases where the material was not published elsewhere or when the topic was considered essential for inclusion in this work. It was not possible to examine and include every known reference and some sources were inadvertently missed; omissions and errors should be brought to the attention of the author. A complete checklist of the Brachyura of the Gulf has not been published since Rathbun's multi-volume work, "The Crabs of America" (1918, 1925, 1930, 1937). Since that time, numerous local and regional surveys have yielded new findings on the distribution and biology of many species, new species have been described, and revisions in the systematic organization of several groups have appeared. For most non-specialists, much of this documentary literature is inaccessible and the taxonomic revisions may be difficult to evaluate. The present work should serve as an introduction to the literature for a species and it should also indicate the amount of data and type of research available for a species or group of crabs.

References to the literature are divided into two sections. Those of value in identification and characterization of a species are listed under the name and source of original description. These citations include synonyms and other invalid names; they occasionally emphasize a name or part of a description that is not synonymous (e.g., Williams, 1974a, p. 731, figs. 4, 18a (not 18b), 20c, etc.). Other references are listed under "Remarks": they provide information on the natural history, ecology, development, physiology, behavior, pathology, or commercial fishery of a species. For some commercially-important species, the available literature is large and widely scattered. The citations for these species are listed with a *minimum* of descriptive explanation and the number of omitted references will be proportionately higher than for those species with only a few known references.

The species entries consist of the current valid name, the published original reference of the species description, common names (if any), taxonomic and

descriptive references, geographical and bathymetric distributions, habitats, and bibliography with annotations and comments. Species are arranged alphabetically within genera and subgenera, as are genera within subfamilies.

Geographical ranges are based on published sources. If the identification of a species or locality is doubtful, the locality is preceded by a question mark. Place names are listed from north to south and east to west; eastern Atlantic localities follow western Atlantic; eastern Pacific and Indo-Pacific localities are cited last. The presence of a species is not assumed in the absence of collection records from large gaps in otherwise continuous distributions. For example, many species are listed for the Carolinas and the east coast of Florida, without indicating presence in Georgian waters.

Bathymetric ranges are presented in meters and in fathoms, in order to facilitate comparisons with new data as they appear in the literature. Depths are omitted for semiterrestrial and terrestrial crabs.

Habitat descriptions include substrate types, terrain features (land crabs), associations with other organisms (commensals, parasites), type of water (marine, brackish), and general biotope (estuarine, marsh, rocky intertidal, offshore benthic, etc.). Terrestrial habitats are usually described in greater detail than aquatic habitats.

Remarks include literature citations on all aspects of the biology of the species. In many cases, the available literature may consist only of one or a few published descriptions or citations of the species. Regional lists include faunal surveys in the Gulf of Mexico area. These lists may include data on the habitats, seasonal occurrence, and biology of the crabs, or they may list only the species names. Species identifications were usually verified by a specialist; most of the lists were published. Other remarks include discussions of taxonomic revisions, identification problems, comparisons with other related species, records of ovigerous females, and ecological notes.

CLASSIFICATION

A review of attempts to classify decapod crustaceans, including arrangements of brachyuran families, was presented by Glaessner (1969) in *Treatise on Invertebrate Paleontology*. The scheme adopted for the *Treatise*, a compromise of paleontological and zoological considerations, is used in the present catalogue. Table I presents an outline of the extant families of Brachyura as recognized by Glaessner (1969) and modified for the present work.

The family Hapalocarcinidae is not included in Glaessner (1969) and the Palicidae is listed within the Brachyrhyncha as Superfamily Uncertain. Fenner Chace, in a review of a preliminary manuscript of this catalogue, suggested the placement of the hapalocarcinids as a separate section, although Verrill (1908, p. 426) proposed the inclusion of these crabs as "a peculiar superfamily group," apparently among the Oxystomata. Chace also suggested the inclusion of the Palicidae among the Xanthoidea as a matter of convenience. The freshwater family of crabs, Potamidae, has been revised by Bott (1955b) and separated into two families, the Pseudothelphusidae and the Trichodactylidae. The Prosopidae

TABLE I
Classification of Brachyuran Families

INFRAORDER BRACHYURA

SECTION DROMIACEA de Haan, 1833

- Superfamily DROMIOIDEA de Haan, 1833
- Family PROSOPIDAE von Meyer, 1860
- Family DYNOMENIDAE Ortmann, 1892
- Family DROMIIDAE de Haan, 1833
- Superfamily HOMOLOIDEA White, 1847
- Family HOMOLIDAE White, 1847

SECTION OXYSTOMATA H. Milne Edwards, 1834

- Superfamily DORIPPOIDEA de Haan, 1841
- Family DORIPPIDAE de Haan, 1841
- Superfamily CALAPPOIDEA de Haan, 1833
- Family CALAPPIDAE de Haan, 1833
- Family LEUCOSIIDAE Samouelle, 1819
- Superfamily RANINOIDEA de Haan, 1841
- Family RANINIDAE de Haan, 1841

SECTION HAPALOCARCINIDEA Calman, 1900

- Superfamily HAPALOCARCINOIDEA Calman, 1900
- Family HAPALOCARCINIDAE Calman, 1900

SECTION OXYRHYNCHA Latreille, 1803

- Family MAJIDAE Samouelle, 1819
- Family PARTHENOPIDAE Macleay, 1838

SECTION BRACHYRHYNCHA Borradaile, 1907

- Superfamily PORTUNOIDEA Rafinesque, 1815
- Family PORTUNIDAE Rafinesque, 1815
- Superfamily XANTHOIDEA Dana, 1851
- Family XANTHIDAE Dana, 1851
- Family GERYONIDAE Colosi, 1924
- Family GONEPLACIDAE Macleay, 1838
- Family POTAMIDAE Ortmann, 1896
- Family PINNOTHERIDAE de Haan, 1833
- Family GRAPSIDAE Macleay, 1838
- Family GECARCINIDAE Macleay, 1838
- Family PALICIDAE Bouvier, 1898
- Superfamily OCYPODOIDEA Rafinesque, 1815
- Family OCYPODIDAE Rafinesque, 1815
- Family RETROPLUMIDAE Gill, 1894

SECTION CANCRIDEA Latreille, 1803

- Superfamily CANCROIDEA Latreille, 1803
 - Family CANCRIDAE Latreille, 1803
 - Family CORYSTIDAE Samouelle, 1819
 - Family ATELECYCLIDAE Ortmann, 1893
-

includes the subfamily Homolodromiinae Alcock, 1899, usually treated as a family in other systematic classifications. The Latreillidae was recognized as a separate family by Rathbun (1937), whereas Balss (1957) included the genus *Latreillia* Roux in the Homolidae. Glaessner (1969) tentatively adopted the arrangement of Balss, but left the question of placement open; the Latreillidae are presented as a separate family in the present work. The families Dynomenidae and Retroplumidae do not contain species represented in the Gulf of Mexico.

The arrangement of species and genera within their respective higher taxa may differ according to the current stability of nomenclature within the group. Within the Xanthidae, subfamilies are not listed because the arrangement of genera is incomplete, pending further revisions by Guinot and others. The genus *Uca* of the Ocypodidae has received considerable attention from carcinologists, notably Bott (1973) and Crane (1975). Both of these workers have proposed subgenera to accommodate findings of morphological, biogeographical, and behavioral similarities among species groups. Because the changes are too recent to permit evaluation by other specialists, the species of *Uca* are listed in the present text alphabetically; controversial forms, proposed as full species by some workers and as subspecies by others, are treated as species for the present. It is much easier to combine separate sets of references and records than to attempt to separate them at a later date. Generally, this plan has been followed throughout the systematic presentation; when a taxon can be treated in two or more ways, the simplest arrangement was selected.

A discussion of brachyuran evolution is beyond the scope of this catalogue and the reader is referred to papers by Bourne (1922), Gordon (1963), Stevcic (1971a, 1971b) and the monograph, *Treatise on Invertebrate Paleontology*, which includes the review by Glaessner (1969). This latter reference also compares and discusses the various classification schemes that have been proposed to account for presumed affinities of living and fossil specimens.

BIOGEOGRAPHY

The species catalogue recognizes 352 species of 158 genera, belonging to 22 families. Enumerations of taxa are somewhat arbitrary, depending on the amount of splitting or lumping of various groups. In addition to the 352 species recorded from the Gulf of Mexico, two subspecies and two varietal forms are listed as distinct entries, three species of uncertain Gulf distribution are included (recorded from Cuban waters, coast unspecified), and one species from outside of the Gulf is listed because of nomenclatural relationships with a Gulf species. Forty-one of the 352 species are presently regarded as endemic to the Gulf; the distribution of species and genera by family are listed in Table II.

The greatest number of endemic species occurred among the Pinnotheridae, a family composed mainly of crabs commensal with other organisms. Specificity for hosts and other restrictions imposed by the biology of commensal associates may account for the high ratio of endemism in this family. The largest families, in terms of total numbers of species, contain relatively few species confined to the

TABLE II
Endemic and Total Species and Genera of Brachyura

Family	Endemic species	Species	Total Genera
Dromiidae	0	5	3
Prosopidae	0	2	2
Homolidae	1	4	3
Latreillidae	0	1	1
Raninidae	1	7	4
Dorippidae	6	16	7
Calappidae	0	13	5
Leucosiidae	0	15	8
Hapalocarcinidae	0	1	1
Majidae	4	77	33
Parthenopidae	0	13	8
Atelecyclidae	0	1	1
Cancriidae	0	1	1
Portunidae	2	27	8
Xanthidae	3	69	32
Geryonidae	0	1	1
Goneplacidae	5	19	15
Palicidae	1	9	1
Pinnotheridae	12	31	7
Grapsidae	1	21	12
Gecarcinidae	0	4	2
Ocypodidae	5	15	3
Total	41	352	158

Gulf. The Majidae, Xanthidae, Portunidae, and Grapsidae have endemic percentages of less than 10% of total Gulf species, whereas the Dorippidae, Goneplacidae, and Ocypodidae contain endemic ratios of one-fourth or greater.

The geographical distribution of Gulf species within other areas is compared in Table III. The degree of affinity with the Caribbean fauna is indicated by the large percentage of Gulf crabs also occurring in the Greater Antilles (63.1%), Lesser Antilles (55.7%), and north coast of South America (34.7%). Within these regions, however, there exist considerable differences in collection effort and subsequent taxonomic study. Thus, the number of apparent species recorded from a particular island or section of continental coastline may be a better measure of the number of marine science institutes present than actual species diversity or richness.

Eastern Florida, the Bahamas, and Bermuda contain successively fewer Gulf species, whereas the combined Virginian and Carolinian Provinces of the Atlantic coast contain 44.3% of the total Gulf crabs. The number of Gulf species present in the eastern Atlantic comprises only 7.7% of the Gulf total and one-third of these belong to the ubiquitous Grapsidae. Only 16 Gulf species (4.5%) are also recorded from the eastern Pacific; seven species have been collected from other parts of the Indo-Pacific region.

TABLE III
Distribution of Gulf Crabs in Other Areas

Family	Total species	GA	Caribbean			WC	Western Atlantic				Other	
			LA	SA			EF	BA	BE	NC	EA	EP
Dromiidae	5	4	3	3	0	2	2	2	4	0	0	
Prosopidae	2	2	2	0	0	1	0	0	0	0	0	
Homolidae	4	3	2	0	1	1	1	0	2	2	0	
Iatreillidae	1	1	0	0	0	0	0	0	1	1	0	
Raninidae	7	5	1	1	2	1	1	0	3	1	2	
Dorippidae	16	12	5	0	0	2	0	0	5	1	1	
Calappidae	13	8	8	4	4	5	3	4	10	2	2	
Leucosiidae	15	10	10	5	0	5	3	0	9	0	0	
Hapalocarcinidae	1	0	1	0	0	0	0	1	0	0	0	
Majidae	77	55	53	26	21	28	25	8	37	1	1	
Parthenopidae	13	8	8	4	0	5	4	1	9	0	0	
Atelecyclidae	1	0	1	0	0	0	0	0	0	0	0	
Cancriidae	1	0	0	0	0	1	0	1	1	0	0	
Portunidae	27	20	17	16	13	16	11	13	16	6	1	
Xanthidae	69	46	49	34	20	20	31	15	27	4	2	
Geryonidae	1	1	0	0	0	1	0	0	1	0	0	
Goneplacidae	19	8	5	1	0	2	2	2	5	0	0	
Palicidae	9	2	4	2	0	3	1	0	4	0	0	
Pinnotheridae	31	10	3	2	1	5	0	0	13	0	0	
Grapsidae	21	15	13	13	8	6	11	9	6	9	6	
Gecarcinidae	4	4	4	4	4	3	3	2	0	0	1	
Ocypodidae	15	8	7	7	6	8	6	1	3	0	0	
Total	352	222	196	122	80	115	104	59	156	27	16	

GA = Greater Antilles; LA = Lesser Antilles; SA = north coast of South America; WC = western Caribbean, including east coast of Central America; EF = east coast of Florida; BA = Bahamas; BE = Bermuda; NC = U.S. Atlantic coast, from New England to Georgia; EA = eastern Atlantic, from Europe to South Africa and mid-South Atlantic islands; EP = eastern Pacific, from California to Chile, including Galapagos.

The distribution of species within the Gulf of Mexico is presented in Table IV, with the Gulf regions diagrammed in Figure 1. Again, some areas have been sampled often (e.g., west and northwest coasts of Florida) and other areas have received only sporadic attention (east coast of Mexico). More species occur on the limestone-based continental shelf of the eastern Gulf than on the sand and mud substrates of the western Gulf. The Gulf regions in Figure 1 correspond to the geophysical features described by Antoine (1972) as Gulf provinces. The dominant sediment types are shown in Figure 2 and reef patches are indicated in Figure 3. Most Gulf species are probably West Indian in origin or affinity (see discussion below), where the shallow water habitats are similar to Gulf regions 1, 2, and 6. The north coast of Cuba (region 7) also contains a relatively high species abundance, probably associated with the greater degree of habitat diversity and wide depth range in that part of the Gulf. A detailed analysis of species distribution patterns, by families, will not be attempted in the present contribution. Table IV summarizes the species recorded to date from each of

TABLE IV
Species Distribution Within the Gulf of Mexico

Family	Total species	Gulf Region (Figure 1)						
		1	2	3	4	5	6	7
Dromiidae	5	5	2	3	0	0	2	2
Prosopidae	2	1	0	0	0	0	0	2
Homolidae	4	2	0	0	1	0	0	3
Latreillidae	1	1	0	0	0	0	0	1
Raninidae	7	5	1	2	2	2	0	4
Dorippidae	16	8	2	2	1	1	1	11
Calappidae	13	11	5	5	3	3	4	5
Leucosiidae	15	13	8	3	1	1	1	3
Hapalocarcinidae	1	1	0	0	0	0	0	0
Majidae	77	61	38	20	7	1	15	31
Parthenopidae	13	10	6	4	0	0	3	4
Atelecyclidae	1	0	1	1	1	1	0	0
Cancriidae	1	1	0	0	0	0	0	0
Portunidae	27	23	11	13	6	6	11	16
Xanthidae	69	61	28	26	4	0	7	35
Geryonidae	1	1	1	1	1	0	0	1
Goneplacidae	19	12	5	8	2	0	8	8
Palicidae	9	5	5	2	1	0	3	4
Pinnotheridae	31	16	12	10	1	0	2	6
Grapsidae	21	16	3	8	6	1	3	11
Gecarcinidae	4	2	0	2	3	3	3	3
Ocypodidae	15	6	7	8	7	3	6	7
Total	352	261	145	118	47	22	69	161

these regions. An analysis of breeding times and collection sites of ovigerous females, combined with data on circulation patterns of currents, larval maturation periods, and tolerances to environmental parameters (temperature, salinity, oxygen content, etc.), will be necessary to establish a comprehensive biogeography of Gulf crabs.

DISCUSSION

The distribution of decapod crustaceans in the western Atlantic has been the subject of several discussions in recent years. Hedgpeth (1953) summarized much of what was known about the geology and zoogeography of the Gulf of Mexico. He emphasized the role of Pleistocene sea level changes and their effects on continuous distributions of marine organisms in the Carolinian Province, a region encompassing shores and adjacent waters from Texas to Cape Hatteras. After the Florida peninsula emerged, isolation of the Gulf and Atlantic populations occurred for many species not able to circumvent the tropical waters of South Florida. The distributions of *Ovalipes* and *Uca* were cited by Abele (1970) as examples. For this reason the range citations in the present contribution attempt to distinguish eastern, southern, and western Florida records.




FIG. 1. Regions of the Gulf of Mexico, based on geological assemblages, depth, and geo-physical characteristics. After Antoine (1972); numbers refer to Table IV and do not correspond with province numbers used by Antoine in his Figure 1-1.

Williams (1965) computed an Atlantic-Gulf disjunct ratio of about 10%, based on studies of Carolina decapods. Only 3 (1.4%) of the species presented in his Table 1 were considered endemic to the Carolinas, further evidence of the ubiquitous nature of the northern fauna. Abele (1970) emphasized the northern and southern affinities of the northeastern Gulf decapods, which he believed was better characterized as a heterogenous assemblage rather than a typical fauna of a province he considered to be ill-defined. In his analysis, 91 species (36.5%) of decapods were similar to the typical Carolinian fauna, but many species previously thought to have a disjunct distribution have been more recently reported from southern Florida (Tabb and Manning, 1961; Rouse, 1970). Abele listed 33 species (13.4% of the northeast decapod total) as endemic and 41 species (16.4%) as tropical in origin, confined to the Caribbean and Gulf by higher temperature requirements. The remaining 84 species in his study were the ubiquitous fauna that ranged from the Carolinas to the tropics.

Studies of terrestrial and freshwater crabs can provide further insights into possible sources of origin. Chace and Hobbs (1969) listed 57 species of crabs from the Caribbean area, none of which had apparent origins in North America alone (exclusive of Mexico). They concluded that the majority of endemics in the




FIG. 2. Areas of the Gulf of Mexico based on dominant sediment type. After Lynch (1954). Coral reefs and patches are shown in Figure 3.

- A = sand, includes other narrow stippled areas adjacent to continental coastline.
- B = sand-mud.
- C = mud.
- D = blue mud.
- E = calcareous mud.
- F = limestone with thin veneer of detrital sediments.
- G = *Globigerina* ooze.
- H = pteropod ooze.

Greater Antilles had originated in Central America or southern Mexico and a few had arrived from South America. The directions of prevalent surface currents (Figure 4) indicate the one-way flow of potential larval populations out of the Caribbean into the Gulf. Although the patterns of loop currents within the Gulf change seasonally, successive recruitment of Caribbean fauna to various parts of the Gulf are possible throughout the year. Various sources (Williams, 1965 summarizes many of the records) indicate that many Caribbean crab species are ovigerous throughout the year; thus Gulf recruitment to a particular shore segment would be a function of current pattern and speed, seasonal temperature, and length of larval life. Laboratory studies of larval development (summarized by Garth, 1965b) indicate that warmer temperatures prolong maturation time. Larval populations thus carried into the Gulf could be diverted into one of several directions. The westward entering currents sweep the shores




FIG. 3. Location of major coral reefs and other features of the Gulf of Mexico, derived from several sources, including Lynch (1954). Dashed line represents the 100 fathom (600 feet) isobath.

- 1—Florida Keys
- 2—Dry Tortugas
- 3—Apalachicola Bay and Cape San Blas
- 4—Mississippi River Delta
- 5—East and West Flower Garden Banks, off Texas
- 6—Seven and One-Half Fathom Reef, off Padre Island
- 7—Pink Shrimp Grounds in Campeche Bay
- 8—Alacrán Reef on Campeche Bank

of Yucatan, eastern Mexico, and eventually, converge with other currents off Texas. Central entering currents flow northwest, into region 3 (Figure 1). The most complex patterns occur as the eastward currents diverge and converge in seasonal patterns of loops. Much of the resultant flow sweeps through the Straits of Florida to emerge into the Atlantic as the warm, northward moving Gulf Stream. Other loops circulate up the west coast of Florida, reaching as far as the Mississippi delta in the winter (Figure 4B). Although these large scale current diagrams do not indicate the complexity of water movements in the Gulf region, nor do they allow one to evaluate the spatial and temporal variations in local currents that might be used for the migration of larval crabs, Figure 4 (after Leipper, 1954) does indicate the successive relationship between Caribbean, Gulf, and western Atlantic faunal provinces, respectively. Undoubtedly, many western




FIG. 4. Surface currents of the Gulf of Mexico (after Leipper, 1954).
4A. Current patterns in June.
4B. Current patterns in December.

Atlantic faunal affinities with West Indian elements occur as a result of direct recruitment, bypassing the Gulf of Mexico.

Specific genera that have been recently investigated and summarized with respect to zoogeography include *Sesarma* (Abele, 1973), *Ovalipes* (Williams, 1976), *Callinectes* (Williams, 1974a), and *Uca* (Crane, 1975). Abele (1973) divided the six Florida species of *Sesarma* into those with Carolinian and West Indian affinities. Although Hedgpeth (1953) had emphasized salinity and temperature tolerances as reasons for the absence of *Sesarma cinereum* and *S. reticulatum* from southern Florida, Abele (1973) believes that competitive exclusion by *S. ricordi* may account for the absence of the first and possibly the second disjunct species. Interspecific competition is an unknown factor in the distribution of virtually every brachyuran species, especially the aquatic crabs. An exception is the genus *Uca*, where intensive studies by several authorities over the past half century have produced the largest amount of biological data on any one crab genus. Crane (1975) summarized the zoogeographical and evolutionary information on *Uca* distribution. She remarked on the relatively depauperate Atlantic fauna of this genus, compared to the rich and diverse Pacific assemblage, also known for many other groups of invertebrates. The differences are attributed to cooling during the Oligocene to Pleistocene, which was far more severe in the Atlantic than in the Pacific, resulting in the extinction of many sensitive tropical species. The isolation of the Caribbean at the end of the Pliocene, due to the emergence of Panama, resulted in the Atlantic assemblage, each species of which has a Pacific analogue. Only *U. subcylindrica*, a poorly-known Gulf endemic, has no eastern Pacific counterpart. Other Gulf species of *Uca*, five of which are endemic, may be in the process of rapid speciation. The most recently-described species, *Uca panacea* (Novak and Salmon, 1974), may have diverged from *U. pugilator* behaviorally and ecologically with only minimal morphological change. Many authorities are reluctant to consider this form a separate species without further evidence of isolation or distinction. The same applies to *U. virens* and *U. longisignalis* (see entries in systematic section), two forms similar to *U. pugnax* of the Atlantic coast. An analysis of courtship displays and habitat requirements reveals the divergence of the species not detected from preserved specimens alone. This evidence is substantiated by preliminary comparisons of isozyme patterns for Atlantic and Gulf populations of *U. panacea*, *U. pugilator*, *U. pugnax*, *U. virens*, and *U. longisignalis* (Selander, Johnson and Avise, 1971; Selander, pers. comm.). Unfortunately, this glimpse of speciation in progress is a rare exception to the usual type of information available. Most accounts of zoogeography depend on collection records derived from preserved specimens. Misidentifications, erroneous locality labels, and gaps in collection efforts are familiar difficulties.

This contribution is offered as a preliminary step toward compiling the necessary information for a synthesis on the distribution and evolution of crabs in the Gulf region. It should facilitate the collection of data on particular groups of crabs and the comparison of specific lines of evidence relative to ecological, behavioral, physiological, or developmental aspects. It can also indicate the rela-

tive lack of information on the majority of species that have been collected and formally described, but in which observations on living animals are minimal or lacking. Hopefully, it will serve to stimulate increased efforts at erasing these deficiencies in our knowledge of some of the most fascinating animals man can hope to meet.

SPECIES ENTRIES

Infraorder BRACHYURA Latreille, 1803

SECTION DROMIACEA de Haan, 1833

Superfamily DROMIOIDEA de Haan, 1833

Family DROMIIDAE de Haan, 1833

Dromia Weber, 1795

Dromia erythropus (George Edwards, 1771) (Cat. Anim. Catesby's Nat. Hist. Carolinas, with Linnaean Names)

Rathbun, 1933, p. 107, fig. 105; Rathbun, 1937, p. 31, text-fig. 11, pl. 6, figs. 1, 2, pl. 8, figs. 1, 2; Felder, 1973a, p. 44, pl. 6, fig. 2.

Range: Bermuda; Bahamas; Florida Keys and Dry Tortugas; off Louisiana and Texas; north coast of Cuba; Jamaica; Haiti; Puerto Rico; St. Thomas to Barbados; Netherlands Antilles; Pernambuco to São Paulo, Brazil.

Depth: shallow water to 360 m (to 197 fm), most common at depths of less than 46 m (25 fm).

Habitat: on hard substrates (coral, shell, near rocks); dorsal carapace is always covered with sponges or compound ascidians.

Remarks: Felder (1973) lists collection localities of Seven and One-Half Fathom Reef off Texas and a sublittoral prominence about 90 miles south of Pecan Island, in the South Atlantic, but a recent (1977) personal communication from Dr. Chace indicates that the St. Helena specimens are still not positively identified; however, they are not *D. erythropus*. Coelho and Ramos (1972) list this species from Brazil. Hazlett (1971) has examined the antennule chemosensitivity of this species.

Dromidia Stimpson, 1858

Dromidia antillensis Stimpson, 1858 (Ann. Lyc. Nat. Hist. New York 7: 71)

Hay & Shore, 1918, p. 417, pl. 31, fig. 5; Rathbun, 1933, p. 108, fig. 106; Rathbun, 1937, p. 33, text-fig. 12, pl. 7, figs. 1-3; Chace, 1940, p. 6; Williams, 1965, p. 143, fig. 118; Felder, 1973a, p. 44, pl. 6, figs. 1, 3.

Range: Bermuda; North Carolina; Bahamas; southeast Florida; Florida Keys, Straits, and Dry Tortugas; west coast of Florida to Texas; northwest and north coasts of Yucatan; off west and south coasts of Cuba; Jamaica; Haiti; Puerto Rico; Virgin Islands; Grenada; off Bahia to Espirito Santo, Brazil.

Depth: shore to 331 m (to 170 fm).

Habitat: from hard bottoms (shell, rock, or coral); usually carries a sponge or compound ascidian over the dorsal carapace. Uncommon to rare in many Gulf areas, but Hildebrand (1955) reported this crab as common on the pink shrimp grounds of Campeche, Mexico, in 6 to 16 fm (11 to 29 m) of water.

Remarks: Rathbun (1937) reported ovigerous females in winter, spring, and summer from Florida and the West Indies, and she also listed specimens with infestations of bopyrid parasites. Williams (1965) noted crabs that carried zoanthoid polyps; specimens from Alligator Harbor in northwest Florida carried the ascidian, *Eudistoma capsulatum* (Wass, 1955). Regional lists include Florida (Dragovich and Kelly, 1964; Abele, 1970; Menzel, 1971), Mississippi (Franks *et al.*, 1972), Texas (Hildebrand, 1955; Parker, 1959; Leary, 1967), and Mexico (Hildebrand, 1955). Felder (1973a) indicates that this species is common on Seven and One-Half Fathom Reef, off Texas; Chace (1956) listed this species from several Gulf stations of the R/V *Oregon*. Listed from French Guiana by Guinot-Dumortier (1959) and from Brazil by Coelho and Ramos (1972). Larval development under laboratory conditions was studied by Rice & Provenzano (1966).

Hypoconcha Guerin, 1854

Hypoconcha arcuata Stimpson, 1858 (Proc. Acad. Nat. Sci. Philadelphia 1858: 226)

Hay & Shore, 1918, p. 418, pl. 31, fig. 3 (not 2); Rathbun, 1937, p. 47, pl. 11, figs. 1-4; Williams, 1965, p. 144, fig. 119.

Range: North Carolina to southern Florida; Dry Tortugas; west coast of Florida; St. Thomas, Virgin Islands; Surinam to Espirito Santo, Brazil.

Depth: 2 to 40 m (1 to 22 fm).

Habitat: sand and shell substrates. Williams (1965) notes that this species is always found with a lamellibranch mollusc shell, usually a clam, which it carries over its back by its claws and fourth and fifth walking legs.

Remarks: Listed from Florida by Wass (1955), Dragovich and Kelly (1964), Abele (1970), and Menzel (1971). Kircher (1970) studied larval development under laboratory conditions.

Hypoconcha sabulosa (Herbst, 1799) (Vers. Natur. Krabben u. Krebse, vol. 3, p. 57)

Hay & Shore, 1918, p. 418, pl. 31, fig. 2 (not 3); Rathbun, 1937, p. 44, pl. 8, figs. 3-4, pl. 9, figs. 1-5; Williams, 1965, p. 145, fig. 120; Felder, 1973a, p. 44, pl. 6, fig. 5.

Range: off North Carolina; Florida Keys and Dry Tortugas; off Texas; Jamaica; Guianas; Moranhão to Bahia, Brazil.

Depth: 1 to 90 m (to 49 fm).

Habitat: from sand, shell, and coral bottoms; Williams (1965) states that the habits of this crab are similar to those of *H. arcuata*, but that it is a much rarer species.

Remarks: Listed from Florida by Hulings (1961) and Abele (1970) and from Brazil by Coelho and Ramos (1972).

Hypoconcha spinosissima Rathbun, 1933 (Proc. Biol. Soc. Washington 46: 185)

Rathbun, 1937, p. 46, text-fig. 14, pl. 10, figs. 1-2; Felder, 1973a, p. 44, pl. 6, fig. 4.

Range: off North Carolina; Dry Tortugas; west coast of Florida; ? Texas; off north coast of Yucatan; Jamaica.

Depth: 26 to 110 m (14 to 60 fm).

Habitat: broken coral or shell, sand bottoms.

Remarks: Although listed by Leary (1967) for Texas, I know of no published collection records to verify its presence in the northwestern Gulf. Felder (1973a) includes this species, based on the listing by Leary.

Family PROSOPIDAE von Meyer, 1860

Subfamily HOMOLODROMIINAE Alcock, 1899

(Glaessner, 1969, corrected the family name Prosoponidae of von Meyer, 1860 and listed three subfamilies: Prosopinae, Pithonotinae, and Homolodromiinae. Only the latter subfamily is extant and many authors treat this group as a family.)

Dicranodromia A. Milne Edwards, 1880

Dicranodromia ovata A. Milne Edwards, 1880 (Bull. Mus. Comp. Zool. 8: 32)

Rathbun, 1937, p. 60, text-fig. 15, pl. 13, figs. 3-4; Chace, 1940, p. 7; Pequegnat, 1970, p. 173.

Range: east and west coasts of Florida; Florida Keys and Straits; off north coast of Cuba; northwest Caribbean Sea; Guadeloupe; Barbados.

Depth: 128 to 895 m (70 to 490 fm).

Habitat: no data available.

Remarks: Rathbun (1937) reported ovigerous females from Florida in June and Chace (1940) reported ovigerous females from Cuba in early May.

Homolodromia A. Milne Edwards, 1880

Homolodromia paradoxa A. Milne Edwards, 1880 (Bull. Mus. Comp. Zool. 8: 33)

Rathbun, 1937, p. 58, pl. 13, figs. 1-2, pl. 14, figs. 1-4; Chace, 1940, p. 7.

Range: north coast of Cuba; off Nevis, Leeward Islands

Depth: 651 to 896 m (356 to 490 fm), possibly to 1106 m (605 fm).

Habitat: no data available.

Superfamily HOMOLOIDEA White, 1848

Family HOMOLIDAE White, 1847

Homola Leach, 1815

Homola barbata (Fabricius, 1793) (Entomol. system., vol. 2, p. 460)

As *Thelxiope barbata*—Rathbun, 1937, p. 63, text-fig. 16, pl. 15, figs. 1–2; Chace, 1940, p. 8; Barnard, 1950; p. 338, fig. 65d–e.

As *Homola barbata*—Williams, 1965, p. 146, fig. 121.

Range: Massachusetts to southern Florida; Florida Keys and Dry Tortugas; north coast of Cuba; off east coast of Yucatan (Caribbean); in eastern Atlantic, off Naples, Portugal, Azores, Madeira Islands; off South Africa.

Depth: 55 to 682 m (30 to 373 fm).

Habitat: sand, shell, and coral substrates; occasionally on mud bottoms.

Remarks: The genus names *Thelxiope* and *Homola* have been interchanged by various authors up to 1958, when the latter name was adopted for genus, family, and superfamily designation. Gordon (1950) described reproductive structures and the evolution of this genus among the Dromiacea. Rice (1964) and Rice and Provenzano (1970) studied larval development. Ovigerous females occur in June–July off North Carolina and Florida (Williams, 1965) and in October off Delaware (Rathbun, 1937). Hartnoll (1970, 1971) noted swimming behavior.

Homola vigil A. Milne Edwards, 1880 (Bull. Mus. Comp. Zool. 8: 33)

As *Thelxiope vigil*—Rathbun, 1937, p. 66, pl. 16, figs. 1–3; Chace, 1940, p. 9.

Range: off Georgia; north and south coasts of Cuba; Guadeloupe; Martinique.

Depth: 309 to 804 m (169 to 440 fm).

Habitat: sand, broken shell, and coral bottoms.

Remarks: Chace (1940) reported ovigerous females from Cuba in April and May.

Homologenus A. Milne Edwards, 1888

Homologenus rostratus (A. Milne Edwards, 1880) (Bull. Mus. Comp. Zool. 8: 34)

Rathbun, 1937, p. 70, text-fig. 17, pl. 17, figs. 1–3; Chace, 1940, p. 9; Pequegnat, 1970, p. 174, fig. 6–1.

Range: Bahamas; east coast of Mexico in southwest Gulf of Mexico; north and south coasts of Cuba; Virgin Islands; near Aves Island (Lesser Antilles); Azores; off Morocco.

Depth: 600 to 1600 m (330 to 875 fm).

Habitat: fine sand, mud, and ooze substrates.

Remarks: Ovigerous females were reported from Cuba in March and May (Chace, 1940) and from the Windward Passage in April (Rathbun, 1937).

Hypsophrys Wood-Mason, 1891

Hypsophrys noar Williams, 1974 (Proc. Biol. Soc. Washington 87: 485)

Williams, 1974b, p. 485, figs. 1–12.

Range: southwest of Dry Tortugas, in Florida Straits.

Depth: 732 m (400 fm).

Remarks: Williams (1974b) compares this species, known only from the type specimen, with the other two species of *Hypsophrys*. The male holotype carried a number of small barnacles, *Poecilasma inaequilaterale*, on the abdominal setae and left cheliped.

Family LATREILLIIDAE Alcock, 1899

(This family, including the genera *Latreillia* Roux and *Latreillopsis* Henderson, was recognized by Rathbun (1937), but these genera were included in the Homolidae by Balss (1957). Glaessner (1969) leaves the position of these genera undetermined.)

Latreillia Roux, 1830

Latreillia elegans P. Roux, 1830 (Crust. Méditerranée et de son littoral, pl. 22, 1828 (1830))

Hay & Shore, 1918, p. 419, pl. 31, fig. 4; Rathbun, 1937, p. 73, text-fig. 18, pl. 20, pl. 21, figs. 1-8; Chace, 1940, p. 10; Williams, McCloskey & Gray, 1968, p. 42, fig. 1.

Range: Massachusetts to North Carolina; off South Florida; Florida Keys; north coast of Cuba; eastern North Atlantic Ocean; Mediterranean Sea; off Natal, South Africa.

Depth: 46 to 366 m (25 to 200 fm).

Habitat: sand, shell, and coral bottoms; from soft mud; off sponges.

Remarks: Rathbun (1937) reported ovigerous females from Massachusetts in August and from Florida in February.

SECTION OXYSTOMATA H. Milne Edwards, 1834

Superfamily RANINOIDEA de Haan, 1841

Family RANINIDAE de Haan, 1841

(This family is treated as a "subtribe" *Gymnopleura* of the "tribe" *Brachyura* by Bourne (1922). Balss (1957) and Glaessner (1969) include this group as family and superfamily within the Oxystomata. The systematic position of the raninids depends, in part, on the relative degree of specialization and primitiveness assigned to the morphological characters of these aberrant crabs.)

Lyreidus de Haan, 1841

Lyreidus bairdii Smith, 1881 (Proc. U.S. Nat. Mus. 3: 420)

Rathbun, 1937, p. 23, pl. 5, figs. 5-6; Chace, 1940, p. 6; Pequegnat, 1970, p. 180.

Range: Massachusetts; Dry Tortugas; west coast of Florida; off Louisiana, Texas, and Mexico; north coast of Cuba; north of Puerto Rico.

Depth: 119 to 823 m (65 to 450 fm).

Habitat: soft mud substrates.

Remarks: Chace (1940) reported a specimen with rhizocephalan parasites from off the north coast of Cuba. Listed from several stations of the R/V *Oregon* by Chace (1956) in the Gulf of Mexico. Pequegnat (1970) considered this species to be the most common raninid in the Gulf and he provides some data on densities at different depths along the continental slope.

Ranilia H. Milne Edwards, 1837

Ranilia constricta (A. Milne Edwards, 1880) (Bull. Mus. Comp. Zool. 8: 35)

Rathbun, 1937, p. 20, pl. 4, fig. 5, pl. 5, figs. 1, 2; Gomes Corrêa, 1970, p. 2; Pequegnat, 1970, p. 180.

Range: Florida Straits; off north coast of Cuba and southeast Gulf of Mexico; Rio de Janeiro, Brazil; eastern Atlantic, Ascension Island; Senegal to Congo.

Depth: off shallow reef (Rathbun, 1937); 183 to 336 m (100 to 200 fm) in Gulf of Mexico (Pequegnat, 1970); original type is from 86 m (47 fm), off Florida.

Habitat: coral reefs; hard bottoms in deep water.

Remarks: Listed from Brazil by Coelho and Ramos (1972). Hartnoll (1971) cites an observation by Darwin on a species of *Ranilia* in the southern Atlantic in which swimming was noted, but the extent to which raninids are able to swim is not known.

Ranilia muricata H. Milne Edwards, 1837 (Hist. nat. Crust., vol. 2, p. 196)

Hay & Shore, 1918, p. 420, pl. 31, fig. 1; Rathbun, 1937, p. 18, pl. 3, figs. 3-6, pl. 4, figs. 1-4; Williams, 1965, p. 142, fig. 117.

Range: North Carolina; Bahamas; Florida Straits; southern to northwestern Florida; Swan Island (Caribbean).

Depth: 9 to 102 m (5 to 56 fm).

Habitat: offshore, on sandy and broken shell substrates.

Remarks: Listed from Florida by Wass (1955), Abele (1970) and Menzel (1971). Rathbun (1937) listed ovigerous females from North Carolina in September. This species has been recovered from fish stomachs in North Carolina offshore waters (Williams, 1965).

Raninoides H. Milne Edwards, 1837

Raninoides lamarcki A. Milne Edwards & Bouvier, 1923 (Mem. Mus. Comp. Zool. 47: 299)

Rathbun, 1937, p. 13, text-fig. 8, pl. 1, figs. 3, 4; Chace, 1940, p. 5.

Range: north of Cuba; north of Puerto Rico; off Colon, Panama (Caribbean).

Depth: 46 to 366 m (25 to 200 fm).

Habitat: no data available.

Remarks: Chace (1940) notes an error in plate 2 of Rathbun, 1937: figure 3 is a chela of *R. lamarcki* and not *R. fossor*, as the label indicates. The error was due to a label interchange on figures a and b of Milne Edwards and Bouvier's (1923) original drawing. Manning (1975) has subsequently indicated that *R.*

fossor is a synonym of an Indo-West Pacific species, *Notosceles chimmonis* Bourne.

Raninoides loevis (Latreille, 1825) (Encycl. méth., Hist. nat., vol. 10, p. 268)

Rathbun, 1937, p. 8, text-fig 3, pl. 1, figs 1, 2; Guinot-Dumortier, 1959, p. 246 fig. 2a-c; Gomes Corrêa, 1970, p. 9.

Range: Florida Keys and Dry Tortugas; southwest coast of Florida; Campeche Bay, off Tabasco, Mexico; Barbados; Colombia (Caribbean); Guianas to Bahia, Brazil; Pacific coasts of Panama and Colombia.

Depth: 18 to 196 m (10 to 107 fm).

Habitat: bottom types include ooze, mud, shelly mud, coral, and broken shell.

Remarks: Listed from the R/V *Oregon* collections in the Gulf of Mexico by Chace (1956). Guinot-Dumortier (1959) and Knight (1968) compared this species with *R. benedicti* Rathbun. Listed from Brazil by Coelho and Ramos (1972).

Raninoides louisianensis Rathbun, 1933 (Proc. Biol. Soc. Washington 46: 186)

Common Name: Frog Crab

Rathbun, 1937, p. 12, text-figs. 6, 7, pl. 1, figs. 5, 6; Pequegnat, 1970, p. 181; Felder, 1973a, p. 38, pl. 4, fig. 6.

Range: Gulf of Mexico, from the Mississippi Delta to Campeche Banks.

Depth: 55 to 400 m (30 to 220 fm). Collection records of the R/V *Alaminos* showed an extension of this species on the middle and outer continental shelf to only 115 fm and Pequegnat (1970) believes that the R/V *Oregon* records at 200 and 220 fm may be due to trawl contamination from earlier, shallower stations.

Habitat: muddy and fine sand-mud bottoms.

Remarks: Listed from Texas by Leary (1967) and from the Gulf of Mexico by Chace (1956). Oviparous females were taken in February, June, July, and October by the R/V *Alaminos* (Pequegnat, 1970). Franks *et al.* (1972) reported salinity and temperature ranges of collections off Mississippi.

Symethis Weber, 1795

Symethis variolosa (Fabricius, 1793) (Entomol. System. emend. et aucta, vol. 2, p. 476)

Rathbun, 1937, p. 26, text-fig. 10, pl. 5, figs. 7, 8; Gomes Corrêa, 1970, p. 10.

Range: North Carolina; southeast Florida; Florida Keys and Dry Tortugas; north of Puerto Rico; Fernando do Noronha to Bahia, Brazil; Pacific coast of Panama.

Depth: 18 to 110 m (10 to 60 fm).

Habitat: sand, mud bottoms; on calcareous algae; under stones.

Remarks: Cerase-Vivas and Gray (1966) extended the previously known range of this species to North Carolina. Listed from Brazil by Coelho and Ramos (1972) and by Fausto Filho (1974).

Superfamily DORIPPOIDEA de Haan, 1841

Family DORIPPIDAE de Hann, 1841

Clythrocerus A. Milne Edwards & Bouvier, 1899

Clythrocerus nitidus (A. Milne Edwards, 1880) (Bull. Mus. Comp. Zool. 8: 24)

Rathbun, 1937, p. 109, text-figs. 26, 27, pl. 33, figs. 1–2.

Range: South Carolina; southeast and northwest Florida; Florida Keys; Grenada.

Depth: 12 to 479 m (6.5 to 262 fm).

Habitat: rock, coral, sand, shell, and gravel bottoms.

Remarks: Listed by Wass (1955) at 25 fm off Cape San Blas, Florida. Rathbun (1937) reported ovigerous females from Florida in February, March, and late June.

Clythrocerus stimpsoni Rathbun, 1937 (Bull. U.S. Nat. Mus. 166: 121)

Rathbun, 1937, pl. 121, text-fig. 32, pl. 34, figs. 5, 6.

Range: west coast of Florida.

Depth: 183 m (100 fm).

Remarks: Known only from the single female type specimen.

Corycodus A. Milne Edwards, 1880

Corycodus bullatus A. Milne Edwards, 1880 (Bull. Mus. Comp. Zool. 8: 23)

Rathbun, 1937, p. 103, pl. 29, figs. 1–4, pl. 30, fig. 1, pl. 31, fig. 1.

Range: off north coast of Cuba.

Depth: 320 to 457 m (175 to 250 fm).

Cyclodorippe A. Milne Edwards, 1880

Cyclodorippe antennaria A. Milne Edwards, 1880 (Bull. Mus. Comp. Zool. 8: 25)

Rathbun, 1937, p. 104, text-fig. 24, pl. 32, figs. 1, 2; Pequegnat, 1970, p. 177.

Range: west coast of Cuba; north coast of Yucatan (Gulf); north coast of Cuba; Puerto Rico; Lesser Antilles, from Dominica to Grenada.

Depth: 91 to 653 m (50 to 357 fm).

Habitat: primarily hard bottoms (coral, sand, and shell).

Cyclodorippe bouvieri Rathbun, 1934 (Smithsonian Misc. Coll. 91: 1)

Rathbun, 1937, p. 106, pl. 32, figs. 3–4, pl. 81, figs. 1–2.

Range: north coast of Cuba; northeast of Puerto Rico.

Depth: 274 to 549 m (150 to 300 fm).

Remarks: Rathbun (1937) lists an ovigerous female from Puerto Rico, taken in March at 150 fm.

Cyclodorippe ornata Chace, 1940 (Torreia 3: 19)

Chace, 1940, p. 19, figs. 7, 8.

Range: off north coast of Cuba.

Depth: 375 to 439 m (205 to 240 fm).

Remarks: Chace (1940) reported an ovigerous female from off Cuba, taken in May at 240 fm.

Cyonomus A. Milne Edwards, 1880

Cyonomus caecus Chace, 1940 (Torreia 3: 12)

Chace, 1940, p. 12, figs. 1-2.

Range: north coast of Cuba.

Depth: 841 m (460 fm).

Cyonomus cubensis Chace, 1940 (Torreia 3: 16)

Chace, 1940, p. 16, figs. 5-6.

Range: north coast of Cuba.

Depth: 475 to 1006 m (260 to 550 fm).

Remarks: Chace (1940) reported ovigerous females from off Cuba in March and May. He feels that this species is a possible link between the genera *Cyonomus* and *Cymopolus* and it may be eventually elevated to generic status when studied further.

Cyonomus quadratus A. Milne Edwards, 1880 (Bull. Mus. Comp. Zool. 8: 26)

Rathbun, 1933, p. 106, fig. 104; Rathbun, 1937, p. 98, text-fig. 23, pl. 30, fig. 3, pl. 31, fig. 3.

Range: northwest of Dry Tortugas; west, north and south coasts of Cuba; Puerto Rico; Lesser Antilles, from St. Croix to Grenada.

Depth: 185 to 929 m (101 to 508 fm).

Habitat: soft bottoms, fine sand, mud, and ooze.

Cyonomus rostratus Chace, 1940 (Torreia 3: 14)

Chace, 1940, p. 14, figs. 3, 4.

Range: north coast of Cuba.

Depth: 658 m (360 fm).

Cymopolus A. Milne Edwards, 1880

(This genus should not be confused with *Cymopolia* Roux, a synonym of *Palicus* Phillipi, of the family Palicidae)

Cymopolus agassizi A. Milne Edwards & Bouvier, 1899 (Bull. Mus. Hist. Nat., Paris 5: 385)

Rathbun, 1937, p. 100, pl. 30, fig. 2, pl. 31, fig. 2.

Range: Florida Straits; Florida Keys; Puerto Rico.

Depth: 128 to 549 m (70 to 300 fm).

Habitat: sand and rocky bottoms.

Remarks: Rathbun (1937) lists ovigerous females from off Florida in February and March.

Ethusa Roux, 1828

Ethusa mascarone americana A. Milne Edwards, 1880 (Bull. Mus. Comp. Zool. 8: 30)

Rathbun, 1933, p. 105, fig. 102; Rathbun, 1937, p. 78, pl. 22, fig. 2, pl. 23, fig. 2; Williams, McCloskey & Gray, 1968, p. 43, fig. 2.

Range: North Carolina; Florida Keys and Dry Tortugas: west coast of Florida; Puerto Rico; Virgin Islands; Maranhão to Bahia, Brazil; Gulf of California and the Pacific coast of Panama.

Depth: shallow water to 82 m (to 45 fm).

Habitat: rock, coral, coarse shell, and sand substrates; from surfaces of algae, bryozoans, and seaweeds.

Remarks: Abele (1970) remarks on the need for revision of this species and its subspecies, noting the variation in growth of critical characters, also cited by Finnegan (1931) and Garth (1966). Williams, McCloskey and Gray (1968) observed a specimen in the laboratory, taken from a reef off North Carolina, that clasped objects over its carapace, in the manner of dromiid crabs and similar to that reported for *E. lata* in the Pacific (Garth, 1946). Listed from Brazil by Rodrigues da Costa (1968a) and Coelho and Ramos (1972).

Ethusa microphthalmma Smith, 1881 (Proc. U.S. Nat. Mus. 3: 418)

Rathbun, 1937, p. 82, pl. 22, fig. 3, pl. 23, fig. 3; Chace, 1940, p. 10; Pequegnat, 1970, p. 175.

Range: Massachusetts to North Carolina; Dry Tortugas; off west coast of Florida; off Mississippi; off east coast of Mexico and Tabasco (Gulf); north and south coasts of Cuba; northeast Caribbean Sea.

Depth: 110 to 752 m (60 to 411 fm).

Habitat: soft bottoms, fine sand, mud, and mud with shell.

Remarks: Listed from the Gulf of Mexico by Chace (1956) and from Texas by Leary (1967), but not by Felder (1973a) who didn't include deep sea species. Rathbun (1937) reported ovigerous females from off Dry Tortugas in July.

Ethusa tenuipes Rathbun, 1897 (Proc. Biol. Soc. Washington 11: 110)

Rathbun, 1937, p. 87, pl. 24, fig. 3, pl. 25, fig. 3; Chace, 1940, p. 11; Williams, McCloskey & Gray, 1968, p. 44.

Range: off North Carolina; east coast of Florida; Florida Keys and Dry Tortugas; off Alabama; north and south coasts of Cuba.

Depth: 46 to 402 m (25 to 220 fm).

Habitat: sand and coral bottoms.

Remarks: Rathbun (1937) reported ovigerous females from North Carolina in July, from off Cuba in April, and from off Dry Tortugas in July.

Ethusa truncata A. Milne Edwards & Bouvier, 1899 (Bull. Mus. Hist. Nat., Paris 5: 384)

Rathbun, 1937, p. 85, pl. 28, figs. 1-2.

Range: off west coast of Florida; off Mississippi Delta and Louisiana; north-west of Trinidad.

Depth: 133 to 219 m (73 to 119 fm).

Ethusina Smith, 1884

Ethusina abyssicola Smith, 1884 (Rept. U.S. Comm. Fish Fisher. 1882, p. 349 (1884))

Rathbun, 1937, p. 91, text-fig. 21, pl. 26, fig. 1, pl. 27, fig. 1; Pequegnat, 1970, p. 175, fig. 6-2.

Range: Massachusetts to North Carolina; in deep waters of northwest, north-east, and southwest quadrants of Gulf of Mexico; off Cape Frio, Brazil; off west coast of Spain.

Depth: 860 to 4061 m (470 to 2220 fm).

Habitat: very soft bottoms, muds and oozes.

Remarks: Pequegnat (1970) presents evidence for two modes of bathymetric distribution, one at 860 to 1399 m (470 to 765 fm) with specimens resembling typical *E. abyssicola*, and a deeper mode at 2551 to 4061 m (1395 to 2220 fm) with specimens that approach *E. faxonii* in size and shape of exorbital teeth. Further studies may reveal two distinct species. Pequegnat (1970) also reports that an ovigerous female with a few advanced embryos was collected at 765 fm in mid-November.

Superfamily CALAPPOIDEA de Haan, 1833

Family CALAPPIDAE de Haan, 1833

Subfamily CALAPPINAE de Haan, 1833

Acanthocarpus Stimpson, 1871

Acanthocarpus alexandri Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 153)

Rathbun, 1937, p. 221, pl. 69, figs. 1-2; Chace, 1940, p. 26; Williams, 1965, p. 156, fig. 137; Pequegnat, 1970, p. 177, fig. 6-3.

Range: Massachusetts; North Carolina to south Florida; Florida Keys and Dry Tortugas; west and northwest Florida; Mississippi; Texas; east coast of Mexico; north coast of Cuba; Puerto Rico to the Grenadines; off Brazil.

Depth: 68 to 476 m (37 to 260 fm).

Habitat: primarily soft bottoms, fine sand, mud and ooze.

Remarks: Chase (1940) and Pequegnat (1970) note that the carapace of this species is broader than long, contrary to the description of Rathbun (1937). Chace (1956) listed this species from the Gulf of Mexico collections of the R/V

Oregon and Pequegnat (1970) reported this crab as the most abundant deep water calappid from the *Alaminos* collections. Rathbun (1937) listed ovigerous females from off Florida in June and Pequegnat (1970) listed the same from the deep Gulf in early August to mid-November. Listed from Brazil by Coelho and Ramos (1972).

Acanthocarpus bispinosus A. Milne Edwards, 1880 (Bull. Mus. Comp. Zool. 8: 19)

Rathbun, 1937, p. 224, pl. 68, figs. 1-3.

Range: off west and northwest coasts of Florida; Dry Tortugas; Grenadines, Windward Islands.

Depth: 201 to 360 m (110 to 197 fm).

Habitat: mud-shell, coral, and clay-mud bottoms.

Remarks: The depth and location records off Florida for the R/V *Oregon* stations (Chace, 1956) may be confused. Stations 1007 and 1010 are positioned well within the 100 fm isobath, yet the depths for these stations are listed at 180 and 225 fm, respectively.

Calappa Weber, 1795

Calappa angusta A. Milne Edwards, 1880 (Bull. Mus. Comp. Zool. 8: 18)

Hay & Shore, 1918, p. 421, pl. 31, fig. 7; Rathbun, 1937, p. 210, pl. 64, figs. 1-6; Williams, 1965, p. 154, fig. 134; Pequegnat, 1970, p. 177.

Range: North Carolina; southeast Florida; Florida Keys and Dry Tortugas; west coast of Florida and mid-eastern Gulf of Mexico; off east coast of Mexico; off North coast of Yucatan; St. Thomas to Grenada; Barbados.

Depth: 13 to 274 m (7 to 150 fm).

Habitat: coral, sand, broken shell, and gravel substrates.

Remarks: Rathbun (1937) reported ovigerous females from southern Florida in March. Chace (1956) recorded this species from the eastern Gulf of Mexico and Williams (1965) stated that this crab is more abundant in the Gulf Stream than in adjacent inshore waters. Shoup (1968) described shell opening behavior by this species.

Calappa flammea (Herbst, 1794) (Versuch Naturgesch. Krabben u. Krebse, vol. 2, p. 161)

Common Names: Flame-streaked Box Crab; Shame-faced Crab

Hay & Shore, 1918, p. 421, pl. 31, fig. 8; Rathbun, 1933, p. 103; Rathbun, 1937, p. 198 (part), pl. 59, figs. 1-2, pl. 60, fig. 1; Reed, 1941, p. 44; Holthuis, 1958, p. 148, figs. 28-35; Williams, 1965, p. 152, figs. 130-131; Felder, 1973a, p. 43, pl. 5, fig. 11.

Range: Massachusetts to south Florida (see Remarks); Bermuda; Bahamas; Florida Straits and Keys; Dry Tortugas; west and northwest Florida; all inshore Gulf of Mexico areas from Florida to Yucatan, Mexico.

Depth: shore to 73 m (to 40 fm), rarely to 229 m (125 fm).

Habitat: hard bottoms, primarily sand.

Remarks: The breeding range for this crab extends only to North Carolina, but larvae as far north as New England may occasionally survive a mild winter to provide temporary range extensions (Hay and Shore, 1918; Holthuis, 1958). Some of the larval stages are figured by Lebour (1944). This species was revised by Holthuis (1958), who found at least two species that had been previously combined under this name. One of these, *C. ocellata*, also occurs in the Gulf of Mexico, thus previous records, particularly from off west Florida, may refer to either or both species. Listed from the R/V *Oregon* collections in the Gulf by Chace (1956). Cheliped modifications associated with shell opening of molluscs were described by Shoup (1968).

Calappa gallus (Herbst, 1803) (Versuch Naturgesch. Krabben u. Krebse, vol. 3, pt. 3, pp. 18 and 46)

Common Name: Yellow Box Crab

Rathbun, 1933, p. 103; Rathbun, 1937, p. 214, pl. 65, figs. 2-3; Barnard, 1950, p. 350, fig. 66e-i; Sakai, 1965, p. 55, pl. 21, fig. 3; Fausto Filho, 1967, p. 48, fig. 4, pl. VI, figs. 7-8; Sakai, 1976b, p. 131, pl. 39, fig. 2.

Range: Bermuda; Bahamas; Florida Keys and Dry Tortugas; northwest Cuba; Jamaica; Puerto Rico; St. Croix to Barbados; off Campeche snapper banks (Gulf of Mexico); Panama (Carib.) to Venezuela; Netherlands Antilles; Ceará to Bahia, Brazil; St. Helena Island (So. Atlantic); off western Africa, from Senegal to Angola; South Africa; Red Sea; Reunion and Seychelles, in Persian Gulf; off India and Maldives; Philippines; Formosa; Japan; Marshall Inlands; Samoa; Hawaiian Islands.

Depth: low tide mark to 218 m (to 119 fm).

Habitat: hard substrates; on reef flats; coral, sand, shell and rock bottoms.

Remarks: Barnard (1950) and Sakai (1965) provide some earlier references for the African and Asian areas, respectively. Coelho and Ramos (1972) and Fausto Filho (1974) list Brazilian records for this species.

Calappa ocellata Holthuis, 1958 (Stud. Fauna Curaçao 8: 158)

Rathbun, 1901, p. 84 (part), pl. 2; Verrill, 1908, p. 420 (part), pl. 25, fig. 1; Rathbun, 1937, p. 198 (part), not pl. 59 or 60; Holthuis, 1958, p. 158, figs. 36-40; Williams, 1965, p. 153, figs. 132-133.

As *C. ocellata*—Fausto Filho, 1967, p. 42, fig. I, pl. 1, figs. 1-2.

Range: Bermuda; off North Carolina and in Beaufort Harbor (rare); Florida Keys and Dry Tortugas; Jamaica; Hispaniola; Puerto Rico; Virgin Islands; Caribbean coasts of Panama and Colombia; Netherlands Antilles; Rocas to Pernambuco, Brazil.

Depth: shallow water to 52 m (to 28 fm).

Remarks: Plates 59 and 60 of Rathbun (1937) are of *C. flammea*, not *C. ocellata*. Other records of *C. flammea* may also include specimens of this species, especially reports prior to the revision by Holthuis (1958). Shoup (1968) described shell opening behavior by this species and other *Calappa*. Coelho and Ramos (1972) list this crab from Brazil.

Calappa springeri Rathbun, 1931.

This species proved to be identical to *C. sulcata* Rathbun, 1898 when examined by Holthuis (1958) and so this name is a junior synonym. Rathbun's earlier description was based on a juvenile of the species and her later description, as *C. springeri*, was based on an adult form. Many of the earlier Gulf of Mexico faunal surveys list *C. springeri* and these can all be referred to *Calappa sulcata*.

Calappa sulcata Rathbun, 1898 (Bull. Lab. Nat. Hist. State Univ. Iowa 4: 289)
Common Names: Yellow Box Crab; Shame-faced Crab; Parrot Crab.

As *C. springeri*—Rathbun, 1937, p. 205, pl. 60, fig. 1, pl. 61, figs. 1–2.

As *C. sulcata*—Rathbun, 1933, p. 103, fig. 99; Rathbun, 1937, p. 211, pl. 64, figs. 7–8, pl. 65, fig. 1; Holthuis, 1958, p. 179, figs. 51–54; Williams, 1965, p. 155, figs. 135–136; Fausto Filho, 1967, p. 46, fig. 3, pl. II, figs. 5–6; Felder, 1973a, p. 42, pl. 5, fig. 10.

Range: North Carolina; Chesapeake Bight; Dry Tortugas; Alabama to south Texas; Tabasco, Mexico; Puerto Rico; Venezuela to Surinam; Amapá to Sergipe, Brazil.

Depth: shore to 183 m (to 100 fm).

Habitat: sand and sand-mud bottoms.

Remarks: Listed from the Gulf of Mexico (some under *C. springeri*) by Chace (1956) and Fotheringham and Brunenmeister (1975), from Mississippi (Franks *et al.*, 1972), and off Texas (Gunter, 1950; Hildebrand, 1954; Leary, 1967). Hildebrand (1954) reported ovigerous females from Texas in May and that claws of this crab were occasionally served at restaurants in Port Aransas. A report of mollusc shell-opening by this and other *Calappa* species (Shoup, 1968) provides one of the few behavioral studies. Listed from Chesapeake Bight by Musick and McEachran (1972) and from Brazil by Coelho and Ramos (1972).

Cycloes de Haan, 1837***Cycloes bairdii*** Stimpson, 1860 (Ann. Lyc. Nat. Hist. New York 7: 237)

Rathbun, 1933, p. 101, fig. 98; Rathbun, 1937, p. 225, pl. 69, figs. 3–4; Garth, 1946, p. 362, pl. 62, figs. 7–8; Williams, McCloskey & Gray, 1968, p. 49, fig. 6.

Range: Bermuda; Bahamas; North Carolina; southeast Florida; Florida Keys and Dry Tortugas; Florida Straits; west coast of Florida; Cuba; Puerto Rico; St. Thomas to Barbados; Old Providence Island (Carib.); in the Pacific, from west coast of Mexico to Ecuador; Rocas to Espirito Santo, Brazil.

Depth: 3 to 229 m (1.5 to 125 fm).

Habitat: sand, rock, coral, and shell bottoms; buries in sand.

Remarks: Guinot-Dumortier & Dumortier (1961) described a stridulatory apparatus in this species and within the genus. Juvenile forms of this crab from Brazil were described by Rodrigues da Costa (1968b) and this species is also listed from Brazil by Coelho and Ramos (1972) and by Fausto Filho (1974).

Subfamily MATUTINAE Macleay, 1838

(Several genera of this subfamily, including *Hepatus* and *Osachila*, were

examined by Guinot (1966) and aligned with *Aethra* of the Parthenopidae, along with *Actaeomorpha* of the Leucosiidae. She proposed a new subfamily, Aethrinae, to contain these genera, pending further studies. The status of these changes is still in doubt, as is the status of the genus *Matuta* Weber, which is not represented in the Gulf of Mexico. Until such studies are available, *Hepatus* and *Osachila* are included within the Matutinae and they are listed with the other calappids, while recognizing their probable relationship to the Parthenopidae.)

Hepatus Latreille, 1802

Hepatus epheliticus (Johansson, in Linnaeus, 1763) (*Amoenitates academicae*, etc., vol. 6, p. 414)

Common Names: Calico Crab; Leopard Crab; Dolly Varden Crab

Hay & Shore, 1918, p. 422, pl. 37, fig. 1; Rathbun, 1937, p. 238, pl. 70, figs. 3-4, pl. 71, figs. 1-4; Williams, 1965, p. 158, fig. 140; Felder, 1973a, p. 43, pl. 5, fig. 14.

Range: Chesapeake Bay to south Florida; Florida Keys and Dry Tortugas; west coast of Florida to south Texas; Campeche Banks, off Yucatan; Cuba; Jamaica; Dominican Republic.

Depth: near shore to 46 m (to 25 fm).

Habitat: sand, sand-shell, and mud-sand substrates; found inside passes, channels, and harbors, but more common in shallow, open marine waters. Buries in substrate, probably nocturnal. Frequently collected with sea anemones attached to dorsal carapace.

Remarks: *Calliactis tricolor* is the most common anemone found on this crab, usually a single anemone located in the middle of the anterior margin, where the exhalent current of the crab induces a current over the anemone's basal disc (Carlgren and Hedgpeth, 1952). Larval development of this crab was studied by Costlow and Bookhout (1962). Gray (1957) measured the total gill area. Considerable variation exists in the dorsal carapace color patterns: some are spotted and others are marked with horizontal bands, including continuous gradations between these forms. Ovigerous females are not often collected, but have been reported off Texas in July (Hildebrand, 1954). Regional lists include Florida (Wass, 1955; Dragovich and Kelly, 1964; Abele, 1970; Menzel, 1971), Mississippi (Richmond, 1962; Franks *et al.*, 1972), Louisiana (Behre, 1950; Hoese and Valentine, 1972), Texas (Gunter, 1950; Hedgpeth, 1953; Hildebrand, 1954; Parker, 1959; Breuer, 1962; Leary, 1967), Campeche (Hildebrand, 1955), and offshore waters of the Gulf (Chace, 1956). Fotheringham and Brunenmeister (1975) summarize some of the natural history of this crab. Guinot (1966) reviews the taxonomic status of this and related genera.

Hepatus princeps (Herbst, 1794).

This name was determined to be a junior synonym of *H. pudibundus* (Herbst, 1785) in a revision by Holthuis (1959, p. 167). Rathbun (1937) and earlier regional surveys use the junior name and these are referred to *H. pudibundus*.

Hepatus pudibundus (Herbst, 1785) (Versuch Naturgesch. Krabben u. Krebse, vol. 1, p. 199)

As *H. princeps*—Rathbun, 1933, p. 104, fig. 101; Rathbun, 1937, p. 235, pl. 70, figs. 1–2.

As *H. pudibundus*—Holthuis, 1959, p. 167, figs. 36–37, 38a–b; Williams, 1965, p. 157, figs. 138–139; Guinot, 1966, p. 755, figs. 9, 18, 30; Fausto Filho, 1967, p. 50, fig. 5, pl. II, figs. 9–10; Felder, 1973a, p. 43, pl. 5, fig. 13.

Range: off Georgia; off Louisiana and Texas; north and south coasts of Cuba; Jamaica; Haiti; Puerto Rico; St. Thomas to Guadeloupe; Panama (Carib.); Surinam; Bahia to Santa Catarina, Brazil; off Guinea, western Africa; Cape of Good Hope, South Africa.

Depth: shore to 49 m (to 27 fm).

Habitat: very shallow waters with sand-mud and shell-mud bottoms. Often carries anemones and barnacles on carapace.

Remarks: The systematic status of this species and the genus have been reviewed by Holthuis (1959) and by Guinot (1966). Holthuis reported ovigerous females from Surinam in April. Behre (1950) collected this crab at Grand Isle, Louisiana, only once or twice, from among oyster beds. Leary (1967) lists this species from Texas and Felder (1973a) confirms this with a record of his own. Nomura and Fausto Filho (1966) reported biometric data on this crab from Brazil and additional Brazilian records were provided by Rodrigues da Costa (1968a) and Coelho and Ramos (1972).

Osachila Stimpson, 1871

(See remarks under Matutinae with regard to this genus and *Hepatus*.)

Osachila antillensis Rathbun, 1898 (Bull. Lab. Nat. Hist. State Univ. Iowa 4: 290)

Rathbun, 1933, p. 104, fig. 100; Rathbun, 1937, p. 251, pl. 77, fig. 2.

Range: north coast of Cuba; St. Croix, Virgin Islands; Montserrat; Dominica; Barbados; Grenada.

Depth: 123 to 304 m (67 to 164 fm).

Habitat: sand, shell, coral, and rock substrates; on sand-mud bottoms.

Osachila semilevis Rathbun, 1916 (Proc. U.S. Nat. Mus. 50: 652)

Hay & Shore, 1918, p. 422, pl. 31, fig. 9; Rathbun, 1937, p. 251, pl. 77, fig. 1; Williams, 1965, p. 159, fig. 142.

Range: North Carolina; Dry Tortugas; northwest Florida.

Depth: 2 to 91 m (1 to 50 fm).

Habitat: sand, shell, and rocky bottoms.

Remarks: Williams (1965) reports that this species has been collected north of Florida only twice, both times in the Beaufort, North Carolina area. He also reports ovigerous females from Florida in July. Listed from Florida by Wass (1955) and Abele (1970), but not by Menzel (1971).

Osachila tuberosa Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 154)

Hay & Shore, 1918, p. 423, pl. 31, fig. 10; Rathbun, 1937, p. 250, pl. 77, fig. 3;

Williams, 1965, p. 159, fig. 141; Guinot, 1966, p. 748, figs. 3, 15, 17, 27, 31, 34, 35; Pequegnat, 1970, p. 178.

Range: North Carolina (rare); Florida Keys and Dry Tortugas; south and west coasts of Florida; eastern portions of the Gulf of Mexico.

Depth: 66 to 183 m (36 to 100 fm), but more common at the shallower end of its bathymetric range.

Remarks: Listed from the Gulf of Mexico by Chace (1956) for the R/V *Oregon* collections.

Family LEUCOSIIDAE Samouelle, 1819

Subfamily EBALIINAE Stimpson, 1871

Ebalia Leach, 1817

Ebalia cariosa (Stimpson, 1860) (Ann. Lyc. Nat. Hist. New York 7: 238)

Rathbun, 1937, p. 125, pl. 35, figs. 6-7; Williams, 1965, p. 147, fig. 122.

Range: North Carolina; southeast Florida; Florida Keys and Dry Tortugas; west and northwest coasts of Florida; Jamaica; northeastern South America to São Paulo, Brazil.

Depth: low tide mark to 131 m (to 72 fm).

Habitat: from coral, rock, sand, and broken shell substrates.

Remarks: Florida listings include Wass (1955), Tabb & Manning (1961), Rouse (1970), Abele (1970), Lyons *et al.* (1971), and Menzel (1971). Williams (1965) reports ovigerous females from North Carolina throughout the summer and he notes the death-feigning behavior of this crab when it is captured. The crab resembles pebbles and shell pieces with which it is associated. Listed from Brazil by Coelho and Ramos (1972).

Ebalia stimpsoni A. Milne Edwards, 1880 (Bull. Mus. Comp. Zool. 8: 22)

Rathbun, 1933, p. 100, fig. 96; Rathbun, 1937, p. 124, text-fig. 33, pl. 35, figs. 1-3, pl. 37, figs. 1-3; Williams, McCloskey & Gray, 1968, p. 46.

Range: off North Carolina; Bahamas; Florida Keys and Dry Tortugas; west coast of Florida; Puerto Rico; Barbados; Maranhão to Bahia, Brazil.

Depth: 7 to 183 m (4 to 100 fm).

Habitat: sand, shell, and coral bottoms.

Remarks: Listed from Florida by Hulings (1961) and Abele (1970). Williams, McCloskey and Gray (1968) reported an ovigerous female from off North Carolina in May. Listed from Brazil by Rodrigues da Costa (1968a) and by Coelho and Ramos (1972).

Lithadia Bell, 1855

Lithadia cadaverosa Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 159)

Rathbun, 1937, p. 137, pl. 38, figs. 3-6.

Range: Bahamas; northwest coast of Florida and northeast portion of Gulf.

Depth: 46 to 62 m (25 to 34 fm).

Habitat: sand, broken shell, and gravel substrates.

Speloeophorus A. Milne Edwards, 1865

Speloeophorus elevatus Rathbun, 1898 (Bull. Lab. Nat. Hist. State Univ. Iowa 4: 290)

Rathbun, 1937, p. 145, pl. 39, figs. 7-9.

Range: Florida Keys; Jamaica; off Cape St. Roque and from Maranhão to Alagoas, Brazil.

Depth: 1 to 83 m (1 to 45 fm).

Habitat: broken shell substrate.

Remarks: Listed from Brazil by Coelho (1971a) and Coelho and Ramos (1972).

Speloeophorus nodosus (Bell, 1855) (Trans. Linn. Soc. London 21: 307)

Hay & Shore, 1918, p. 425, pl. 32, fig. 5; Rathbun, 1933, p. 99, fig. 95; Rathbun, 1937, p. 142, pl. 40, figs. 1-5; Williams, 1965, p. 148, figs. 123-124.

Range: North Carolina; Florida Keys and Dry Tortugas; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Guadeloupe.

Depth: 3 to 18 m (1.5 to 10 fm).

Habitat: coral reefs and coral-sand bottoms.

Remarks: Williams (1965) states that this species is rare in the northern part of its range and he cites a report by Pearse and Williams (1951) of its occurrence on a reef off North Carolina. This crab readily feigns death when captured.

Speloeophorus pontifer (Stimpson, 1871) (Ann. Lyc. Nat. Hist. New York 10: 115)

Hay & Shore, 1918, p. 425, pl. 32, fig. 5; Rathbun, 1933, p. 100; Rathbun, 1937, p. 144, pl. 39, figs. 1-3; Williams, 1965, p. 149, figs. 125-126.

Range: off North Carolina; Florida Keys and Dry Tortugas; west coast of Florida; Cuba; Puerto Rico; Barbados.

Depth: low tide mark to 229 m (to 125 fm).

Habitat: coral, sand, and shell bottoms.

Uhlías Stimpson, 1871

Uhlías limbatus Stimpson, 1871 (Ann. Lyc. Nat. Hist. New York 10: 118)

Rathbun, 1937, p. 150, pl. 36, figs. 3-5.

Range: west of Key West, Florida; north coast of Cuba; Jamaica; Haiti; St. Thomas, Virgin Islands.

Depth: 4 to 64 m (2 to 35 fm).

Habitat: shell and grassy substrates.

Callidactylus Stimpson, 1871***Callidactylus asper*** Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 158)

Rathbun, 1937, p. 193, pl. 58, figs. 1-3; Williams, McCloskey & Gray, 1968, p. 48, fig. 5.

Range: off North Carolina; southeast Florida; Florida Keys and Dry Tortugas; off southwest Florida; east coast of Haiti; Maranhão to Alagoas, Brazil.

Depth: 27 to 91 m (15 to 50 fm).

Habitat: sand substrates; off *Lithothamnium* reef.Remarks: Listed from Brazil by Coelho (1971a, 1971b) and Coelho and Ramos (1972). See remarks for *Iliacantha liodactylus*.***Iliacantha*** Stimpson, 1871***Iliacantha intermedia*** Miers, 1886 (Voyage of H.M.S. *Challenger*, Brachyura, vol. 17, p. 302.

Hay & Shore, 1918, p. 424, pl. 32, fig. 3; Rathbun, 1937, p. 186, pl. 54, figs. 1-2; Williams, 1965, p. 151, fig. 129.

Range: North Carolina; Florida Keys and Dry Tortugas; northwest coast of Florida; St. Thomas, Virgin Islands; off Venezuela; from Maranhão to Bahia, Brazil.

Depth: 10 to 329 m (5.5 to 180 fm).

Habitat: sand, gravel, coral, and broken shell bottoms.

Remarks: Williams (1965) comments on the close resemblance between juveniles of this species and those of *Persephona mediterranea*. Listed from Brazil by Coelho (1971a) and Coelho and Ramos (1972).***Iliacantha liodactylus*** Rathbun, 1898 (Bull. Lab. Nat. Hist. State Univ. Iowa 4: 291)

Rathbun, 1937, p. 186, text-fig. 41, pl. 55, figs. 1-2; Felder, 1973a, p. 39, pl. 5, fig. 2.

Range: west coast of Florida; ? Haiti; Puerto Rico; St. John, Virgin Islands; Trinidad; Alagoas to Bahia, Brazil.

Depth: 9 to 130 m (5 to 71 fm).

Habitat: mud substrates.

Remarks: Recorded by Chace (1956) from the R/V *Oregon* cruise in the Gulf of Mexico. The locality records cited for this species and for *Callidactylus asper* by Rathbun (1937) are not in agreement with the latitudes and longitudes listed. Both species are listed from the east coast of Haiti, as recorded by the Johnson-Smithsonian Expedition of 1933, but the coordinates as listed would place *C. asper* east of Puerto Rico and place *I. liodactylus* on the east coast of the Dominican Republic. Leary (1967) lists *I. liodactylus* from Texas and Felder (1973a) repeats Leary's listing, but I know of no published collection records to verify the presence of this species in the northwestern Gulf. Listed from Brazil by Coelho (1971b) and Coelho and Ramos (1972).

Iliacantha sparsa Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 156)

Rathbun, 1937, p. 190, pl. 56, figs. 1–2.

Range: northwest of the Dry Tortugas; off north and southeast coasts of Puerto Rico; Barbados; Maranhão to Bahia, Brazil.

Depth: 23 to 73 m (13 to 40 fm).

Habitat: gravel, shell, and coral bottoms; on calcareous algae.

Remarks: Only male specimens were listed by Rathbun (1937), one of which was heavily infested with parasites. Listed from Brazil by Coelho (1971a) and Coelho and Ramos (1972).

Iliacantha subglobosa Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 155)

Hay & Shore, 1918, p. 424, pl. 32, fig. 2; Rathbun, 1937, p. 185, pl. 53, figs. 1–2; Williams, 1965, p. 150, fig. 128; Pequegnat, 1970, p. 179.

Range: North Carolina; southeast Florida; Florida Keys and Dry Tortugas; northwest Florida; north coast of Cuba; Jamaica; Lesser Antilles, from Montserrat to Barbados; from Amapá to Alagoas, Brazil.

Depth: 27 to 393 m (15 to 215 fm).

Habitat: coral, sand, and rock bottoms.

Remarks: Listed by Chace (1956) from the Gulf of Mexico. Williams (1965) reported ovigerous females from the Gulf in June. Brazilian records include Coelho (1971a, 1971b) and Coelho and Ramos (1972).

Subfamily PHILYRINAE Rathbun, 1937

Myropsis Stimpson, 1871

Myropsis quinquespinosa Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 157)

Rathbun, 1937, p. 164, pl. 46, figs. 1–3; Chace, 1940, p. 24; Williams, McCloskey & Gray, 1968, p. 46, fig. 4; Pequegnat, 1970, p. 179; Felder, 1973a, p. 39, pl. 5, fig. 5.

Range: Massachusetts; North Carolina; Bahamas; southeast Florida; Florida Keys and Dry Tortugas; west coast of Florida; all portions of Gulf of Mexico; off Alabama, Texas, and Campeche Bank in Mexico; north and south coasts of Cuba; Jamaica; Puerto Rico; Lesser Antilles, from Martinique to Grenada; Barbados; Venezuela.

Depth: 91 to 329 m (50 to 185 fm), rare report to 1045 m (572 fm).

Habitat: commonly on mud bottoms, also on sand and shell substrates.

Remarks: Rathbun (1937) reported ovigerous females from Florida in May and July and Williams, McCloskey and Gray (1968) reported an ovigerous female from off North Carolina in July, where they believe this species is restricted to a depth range of 120 to 160 m. Chace (1956) recorded this crab from 50 fm off Sabine, Texas and it is listed from Texas by Leary (1967). Pequegnat (1970) notes that the depth record of 572 fm may be excessive.

Persephona* Leach, 1817**Persephona aquilonaris* Rathbun, 1933.**

This name was given to the western Atlantic species, including the Gulf of Mexico form, originally to indicate a subspecies of *P. punctata*. A revision of the genus by Guinot-Dumortier (1959) indicated that the two forms were distinct species and she extended the range through the Caribbean to South America, where it co-exists with *P. punctata*. Rathbun (1937) had restricted *P. punctata aquilonaris* to North America. At the same time as her revision, Guinot-Dumortier (1959) indicated that the species illustrated by Herbst, *P. mediterranea*, may be the correct name of *P. aquilonaris*. Rathbun (1937, p. 153) listed *P. mediterranea* as a synonym of *P. punctata*, but she indicated that the type locality (Mediterranean Sea) was probably incorrect. Abele (1970) reviews the synonymy and notes that L. B. Holthuis verified the identity and synonymy of *P. mediterranea* and *P. aquilonaris*. Abele (1970) also notes that the figure in Rathbun (1933, p. 99, fig. 94) for *P. punctata* from Puerto Rico is a figure of *P. mediterranea*.

***Persephona crinita* Rathbun, 1931 (J. Washington Acad. Sci. 21: 128)**

Rathbun, 1937, p. 163, pl. 43, figs. 2-3, pl. 44, figs. 1-3; Felder, 1973a, p. 39, pl. 5, fig. 3.

Range: northwest Florida to Texas; Trinidad; Ilha São Sebastião, Brazil.

Depth: 5.5 to 91 m (3 to 50 fm).

Habitat: mud and mud-sand bottoms; more common in open marine waters than is *P. mediterranea*.

Remarks: Hildebrand (1954) collected an ovigerous female from off Texas in June. Wass (1955) stated that this species is rarer than *P. aquilonaris* (= *P. mediterranea*) off northwest Florida and Menzel (1971) listed this crab as uncommon on sand bottoms in the same area. Franks *et al.* (1972) extended the known depth range to 50 fm, from collections off Mississippi. Listed from the Gulf collections of the R/V *Oregon* by Chace (1956).

***Persephona mediterranea* (Herbst, 1794) (Versuch Naturgesch. Krabben u. Krebse, vol. 2, p. 150)**

Common Name: Purse Crab

As *P. punctata*—Hay & Shore, 1918, p. 423, pl. 32, fig. 9 (part); Rathbun, 1933, p. 99 (part), fig. 94.

As *P. punctata aquilonaris*—Rathbun, 1937, p. 154, pl. 42, figs. 6-7; Williams, 1965, p. 150, fig. 127.

As *P. aquilonaris*—Guinot-Dumortier, 1959, p. 429, figs. 7, 9; Felder, 1973a, p. 42, pl. 5, fig. 4.

As *P. mediterranea*—Abele, 1970, p. 62.

Range: New Jersey to south Florida; Dry Tortugas; west coast of Florida to south Texas; off Campeche, Mexico; Guadeloupe, Lesser Antilles; French Guiana; Santa Catarina, Brazil.

Depth: 4 to 55 m (2 to 30 fm).

Habitat: sand, shell, and coral bottoms; inshore waters, in passes; in very shallow water just below low tide mark; from mud-shell substrates in north-western Gulf of Mexico.

Remarks: For explanation of nomenclatural revisions, see *P. aquilonaris*. Williams (1965) reported abundant colonies of this crab, with ovigerous females present throughout the spring and summer in North Carolina waters. Regional listings include the Gulf (Chace, 1956; Fotheringham and Brunenmeister, 1975), Florida (Wass, 1955; Tabb and Manning, 1961; Dragovich and Kelly, 1964), Louisiana (Behre, 1950), and Texas (Hedgpeth, 1953; Hildebrand, 1954; Leary, 1967).

***Persephona punctata aquilonaris* Rathbun, 1933.**

Revised by Guinot-Dumortier (1959) to a separate species and later determined to be a junior synonym of *P. mediterranea* (Herbst). See *P. aquilonaris* for nomenclatural discussion.

SECTION HAPALOCARCINIDEA Verrill, 1908

Superfamily HAPALOCARCINOIDEA (Verrill, 1908)

Family HAPALOCARCINIDAE Calman, 1900

(The taxonomic status of these unique crabs is uncertain. Verrill (1908, p. 426) proposed the designation "Hapalocarcinidea" for "a peculiar superfamily," apparently including them among the Oxystomata. Glaesner (1969) does not include this group in his classification schemes. Chace (personal comm.) has suggested the retention of "Hapalocarcinidea" for a section name and the appropriate ending for the superfamily name, both of which are credited to Verrill. The status and relationships of the hapalocarcinids will undoubtedly change with further study.)

***Cryptochirus* Heller, 1861**

***Cryptochirus corallicola* (Verrill, 1908)** (Trans. Connecticut Acad. Arts Sci. 13:427)

Rathbun, 1937, p. 262 (part), text-fig. 47, pl. 78, figs. 5-7.

Range: Florida Straits; Dry Tortugas; Bermuda; Dominica; Maranhão to Bahia, Brazil.

Depth: 0 to 75 m (0 to 41 fm).

Habitat: in ovoid cavities in the upper surfaces of corals, such as *Meandra areolata* and *Meandrina* sp.

Remarks: Serene (1966) reviews the taxonomy and geographical distribution of hapalocarcinids. Rathbun (1937) commented on the relationship between the crabs and their living coral habitat: "the opening of the den is usually semi-circular or lunate, commonly oblique to surface of coral. The downturned, rough,

and dirt-covered front of the crab serves as an operculum, closing the aperture. Full grown crabs are probably unable to leave their dens." The western African record listed by Rathbun (1937) was referred to *Troglocarcinus balssi* by Monod (1956). Listed from Brazil (as *Troglocarcinus corallicola*) by Coelho (1971a) and Coelho and Ramos (1972). Fausto Filho (1974) provides some habitat notes and extended the known depth to 75 m, based on Brazilian specimens collected by Coelho.

SECTION OXYRHYNCHA Latreille, 1803

Family MAJIDAE Samouelle, 1819

Subfamily ACANTHONYCHINAE Stimpson, 1870

Acanthonyx Latreille, 1825

Acanthonyx petiverii H. Milne Edwards, 1834 (Hist. Nat. Crust., vol. 1, p. 343)

Rathbun, 1925, p. 142, text-fig. 52, pl. 44, pl. 222, figs. 1–6; Rathbun, 1933, p. 13, fig. 11; Garth, 1958, p. 225, pl. O, fig. 3, pl. 25, fig. 2; Felder, 1973a, p. 53, fig. 14.

Range: Bahamas; southeast and northwest Florida; Texas; Cuba; Jamaica; Puerto Rico; Virgin Islands; Netherlands Antilles; Panama (Carib.) to Rio de Janeiro, Brazil; along the Pacific coast, from Baja California to Caldera, Chile; Galapagos Islands.

Depth: shore to 29 m (16 fm).

Habitat: in tide pools of rocky, surf-beaten shores; algal-covered surfaces and on seaweeds; sandy shores; coral flats.

Remarks: Leary (1967) lists this crab from Texas and Felder (1973a) found this crab on rock jetties in southern Texas. Chace (1966) figured four types of carapace variation in this species. Abele (1970) also collected this crab, a single female, from seaweed on jetties in northwest Florida. Listed from Brazil by Coelho (1971a, 1971c) and Coelho and Ramos (1972).

Epialtus H. Milne Edwards, 1834

Epialtus bituberculatus H. Milne Edwards, 1834 (Hist. Nat. Crust., vol. 1, p. 345)

Rathbun, 1925, p. 148, text-figs. 53a, 54, pl. 45, figs. 3–4; Rathbun, 1933, p. 14, fig. 15; Garth, 1958, p. 228.

Range: east coast of Florida; Key West, Florida; Puerto Rico; Panama (Carib.) to Colombia; Ceará to Pernambuco, Brazil.

Depth: shallow waters.

Habitat: like others of this genus, on hard surfaces and in tide pools, feeding among seaweed and other plant growth.

Remarks: Although this species has been listed from Chile and from southern California, Garth (1958) recognizes an exclusively Atlantic range. Listed from Brazil by Coelho (1971c) and Coelho and Ramos (1972).

Epialtus dilatatus A. Milne Edwards, 1878 (Crust. Rég. Mex., p. 140)

Rathbun, 1925, p. 153, text-fig. 53j, pl. 45, fig. 2; Rathbun, 1933, p. 15, fig. 14; Williams, 1965, p. 249, figs. 228, 233D.

Range: North Carolina; Bahamas; south Florida; Dry Tortugas; Isla Mujeres, Yucatan (Carib.); Puerto Rico; St. Thomas, Virgin Islands.

Depth: 5 to 22 m (2.5 to 12 fm).

Habitat: sand, coral, and broken shell substrates.

Remarks: Yang (1968) describes the development of this crab to the first adult stage under laboratory conditions. The taxonomic status of the elongated form is still unclear and is described separately below.

Epialtus dilatatus forma ***elongata*** Rathbun, 1923 (Proc. Biol. Soc. Washington 36: 72)

Rathbun, 1925, p. 154, fig. 53k, pl. 48.

Range: Florida Keys; south, west, and northwest coasts of Florida.

Depth: 2 to 14 m (1 to 7.5 fm).

Habitat: on sandy bottoms with grass; in patches of *Sargassum*.

Remarks: Abele (1970) notes the considerable variation in this species and genus and he expresses reservations about the distinct status of this form. Florida listings include Wass (1955), Abele (1970), Menzel (1971), and Lyons *et al.*, (1971). Wass (1955) collected this crab from *Sargassum*, off Ochlockonee Bay, Florida. Rathbun (1925) reported ovigerous females from Florida in January and February; other records include ovigerous females from Florida in March and July.

Epialtus longirostris Stimpson, 1860 (Ann. Lyc. Nat. Hist. New York 7: 199)

Rathbun, 1925, p. 151, text-figs. 53g, 56.

Range: Key West and west coast of Florida; Cuba; Jamaica; St. Thomas, Virgin Islands; northeast Brazil.

Depth: shallow water, 3 to 5 m (2 to 3 fm) off Cuba; 19 to 54 m (10 to 30 fm) off Brazil.

Remarks: Listed from Brazil by Coelho (1971c).

Mocosoa Stimpson, 1871

Mocosoa crebripunctata Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 128)

Rathbun, 1925, p. 159, text-fig. 59, pl. 49, figs. 3-4.

Range: Florida Straits; off Cape San Blas, northwest Florida; Maranhão to Espírito Santo, Brazil.

Depth: 27 to 131 m (15 to 72 fm).

Remarks: Listed from Brazil by Coelho (1971a) and by Coelho and Ramos (1972).

Aepinus* Rathbun, 1897**Aepinus septemspinosus* (A. Milne Edwards, 1879) (Crust. Rég. Mex., p. 185)**

Rathbun, 1925, p. 92, text-figs. 28–29, pl. 32, figs. 3–4, pl. 219, figs. 1–3; Rathbun, 1933, p. 10, fig. 7.

Range: Bahamas; Florida Straits; Dry Tortugas; west and northwest coasts of Florida; Puerto Rico; St. Thomas, Virgin Islands; Pará to Bahia, Brazil.

Depth: 13 to 85 m (7 to 47 fm).

Habitat: on hard surfaces, mainly coral; from calcareous algae and rock.

Remarks: Listed from off northwest Florida by Wass (1955). Coelho (1971c) extended the known depth range and commented on the ecology of this crab in Brazil. Also listed from Brazil by Coelho and Ramos (1972) and Fausto Filho (1974).

Anasimus* A. Milne Edwards, 1880**Anasimus latus* Rathbun, 1894 (Proc. U.S. Nat. Mus. 17: 58)**

Rathbun, 1925, p. 65, pl. 214; Guinot-Dumortier, 1960, p. 177, fig. 18a–b; Williams, 1965, p. 240, figs. 217, 223F; Felder, 1973a, p. 49, pl. 7, fig. 5.

Range: North Carolina to south Florida; Florida Keys; northwest Florida to off coast of Tabasco, Mexico; west of Trinidad; Guianas; Amapá, Brazil.

Depth: 48 to 161 m (26 to 88 fm).

Habitat: mud, sand, broken shell, and coral bottoms.

Remarks: Rathbun (1925) compares differences in the morphology of adults and juveniles. Reports of ovigerous females were summarized by Williams (1965); females apparently carry eggs throughout the year in the Gulf of Mexico. Holthuis (1959) describes this species from off Surinam, where it is ovigerous from April to August. Sandifer and van Engel (1972) studied larval development under laboratory conditions. Regional lists include Florida (Wass, 1955), Mississippi (Franks *et al.*, 1972), Texas (Hildebrand, 1954; Leary, 1967), Mexico (Hildebrand, 1954), and the Gulf of Mexico collections of the R/V *Oregon* (Chace, 1956). Listed from Brazil by Coelho (1971c) and Coelho and Ramos (1972).

Anomalothir* Miers, 1879**Anomalothir frontalis* (A. Milne Edwards, 1879) (Crust. Rég. Mex., p. 189)**

Rathbun, 1925, p. 25, pl. 8, fig. 1, pl. 9, fig. 1, pl. 207; Chace, 1940, p. 56.

Range: north coast of Cuba; Montserrat to Barbados.

Depth: 133 to 421 m (73 to 230 fm).

Habitat: sand, coral, and broken shell bottoms.

Remarks: Rathbun (1925) compares this species with *A. furcillatus*.

***Anomalothir furcillatus* (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 125)**

Rathbun, 1925, p. 24, text-fig. 6, pl. 8, fig. 2, pl. 9, fig. 2, pl. 206; Rathbun, 1933, p. 6, fig. 2; Chace, 1940, p. 55; Williams, 1965, p. 236, figs. 212, 223A.

Range: North Carolina; Florida Straits and Keys; Dry Tortugas; off northwest

Florida; north coast of Cuba; north coast of Yucatan; Jamaica; St. Croix, Virgin Island to Grenada.

Depth: 55 to 686 m (30 to 375 fm), more commonly at depths of greater than 183 m (100 fm).

Habitat: mud, sand, shell, stone, and coral bottoms.

Remarks: Rathbun (1925) listed ovigerous females from the northeast quadrant of the Gulf in February and March.

Arachnopsis Stimpson, 1871

Arachnopsis flipes Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 121) ,

Rathbun, 1925, p. 89, text-figs. 26-27, pl. 32, figs. 1-2, pl. 219, figs. 4-5; Williams, McCloskey & Gray, 1968, p. 58.

Range: off North Carolina; Florida Straits; Dry Tortugas; west coast of Florida; off Dominica; Barbados; Ceará to Rio Grande do Norte, Brazil.

Depth: 27 to 238 m (15 to 130 fm).

Habit: sand, shell, and coral bottoms.

Remarks: Listed by Wass (1955) from northwest Florida and by Chace (1956) from R/V *Oregon* collections in the Gulf of Mexico. Recorded from Brazil by Coelho (1971a, 1971b) and Coelho and Ramos (1972).

Batrachonotus Stimpson, 1871

Batrachonotus fragosus Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 122)

Rathbun, 1925, p. 123, text-fig. 48, pl. 39, figs. 1-4; Rathbun, 1933, p. 13, fig. 10; Williams, 1965, p. 238, figs. 214, 223C.

Range: North Carolina; south, west and northwest coasts of Florida; Dry Tortugas; north coast of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Curaçao, Netherlands Antilles; Rio de Janeiro, Brazil.

Depth: shore to 137 m (to 75 fm).

Habitat: sand, shell and coral bottoms; rarely on mud.

Remarks: Listed from northwest Florida by Wass (1955), Hulings (1961), and Abele (1970). Williams (1965) reported ovigerous females from Dry Tortugas in June.

Collodes Stimpson, 1860

Collodes armatus Rathbun, 1898 (Bull. Lab. Nat. Hist. State Univ. Iowa 4: 252)

Rathbun, 1925, p. 122, pl. 217, fig. 6.

Range: off Havana, Cuba.

Remarks: known only from the type specimen, collected in 1893.

Collodes leptocheles Rathbun, 1894 (Proc. U.S. Nat. Mus. 17: 53)

Rathbun, 1925, p. 117, text-fig. 42, pl. 38, figs. 5-6; Pequegnat, 1970, p. 181; Felder, 1973a, p. 49, pl. 7, fig. 3.

Range: all quadrants of Gulf of Mexico except southeast, off coasts of Florida, Alabama, and Texas; off Vera Cruz, Mexico.

Depth: 124 to 384 m (68 to 210 fm).

Habitat: gray mud, broken shell substrates.

Remarks: Chace (1956) reported the first record of this species from off Texas, taken at 150 fm, and Leary (1967) also lists this crab from Texas. Pequegnat (1970) reported two ovigerous females, taken in August from 111 fm.

Collodes trispinosus Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 120)

Rathbun, 1925, p. 107, text-fig. 32, pl. 36, figs. 5-6; Williams, 1965, p. 239, figs. 215, 223D.

Range: North Carolina to south Florida; Florida Straits and Keys; Dry Tortugas; west and northwest coasts of Florida.

Depth: 7 to 150 m (4 to 82 fm).

Habitat: sand, broken shell, and gravel substrates.

Remarks: Listed from northwest Florida by Wass (1955). Williams (1965) lists ovigerous females from North Carolina in October and from Florida in July.

Euprognatha Stimpson, 1871

Euprognatha gracilipes A. Milne Edwards, 1878 (Crust. Rég. Mex., p. 184)

Rathbun, 1925, p. 101, pl. 34, figs. 3-4; Rathbun, 1933, p. 11.

Range: Florida Keys; north coast of Yucatan; off north coast of Cuba; Puerto Rico; St. Croix, Virgin Islands; Barbados; Amapá to São Paulo, Brazil.

Depth: 72 to 368 m (39 to 201 fm).

Habitat: sand, shell, coral, and rocky bottoms.

Remarks: Rathbun (1925) comments on variation within this species and the possibility of two different subspecies within the described population. Listed from Brazil by Coelho (1971c) and Coelho and Ramos (1972).

Euprognatha rastellifera acuta A. Milne Edwards, 1880 (Crust. Rég. Mex., p. 348)

Rathbun, 1925, p. 96, pl. 34, figs. 1-2; Rathbun, 1933, p. 11, fig. 8; Chace, 1940, p. 57.

Range: Massachusetts; North and South Carolina; Florida Keys; north coast of Cuba; Puerto Rico; St. Kitts; Martinique.

Depth: 102 to 708 m (56 to 387 fm).

Habitat: sand, coral, and shell substrates.

Remarks: There is considerable geographical overlap among the three forms of this species, the typical form, *E. r. acuta*, and *E. r. marthae*. The subspecies *acuta* is distributed over the entire species range, although it is more common in the southern portion, whereas *marthae* is listed by Rathbun (1925) as restricted to north of southern Florida, along the Atlantic coast. Williams (1965) notes the need for clarification of these forms and their distributions.

Inachoides H. Milne Edwards & Lucas, 1843

Inachoides forceps A. Milne Edwards, 1879 (Crust. Rég. Mex., p. 199)

As *I. laevis*—Rathbun, 1925, p. 61 (part), text-fig. 17, not pl. 22, figs. 3-6; Rathbun, 1933, p. 8, fig. 4.

As *I. forceps*—Garth, 1958, p. 101; Williams, McCloskey & Gray, 1968, p. 60, fig. 14.

Range: North Carolina; west and northwest coasts of Florida; Jamaica; Puerto Rico; Guadeloupe; St. Thomas, Virgin Islands; Guianas to Santa Catarina, Brazil.

Depth: shallow water to 38 m (to 21 fm).

Habitat: sand, gravel, and coral substrates.

Remarks: Garth (1958) separated the Pacific and Atlantic populations, previously combined as *I. laevis*, into two distinct species, retaining *laevis* for the original Pacific form designated by Stimpson. Williams, McCloskey and Gray (1968) cite pl. 22 of Rathbun (1925) as a synonymy, but this is of a specimen from the Pacific coast of Panama and it should be referred to *I. laevis*. Listed from northwest Florida by Wass (1955) and Menzel (1971) and from Brazil by Coelho (1971c) and Coelho and Ramos (1972).

Inachoides laevis Stimpson, 1860.

Restricted to Pacific records by Garth (1958); all Atlantic records are referred to *I. forceps* A. Milne Edwards.

Metoporphaphis Stimpson, 1860

Metoporphaphis calcarata (Say, 1818) (J. Acad. Nat. Sci. Philadelphia 1: 455)

Hay & Shore, 1918, p. 454, pl. 37, fig. 5; Rathbun, 1925, p. 21, text-fig. 5, pls. 6-7; Williams, 1965, p. 243, figs. 221, 223J; Felder, 1973a, p. 48, pl. 7, fig. 2.

Range: North Carolina to south Florida; Florida Keys; west coast of Florida to south Texas; Rio de Janeiro, Brazil.

Depth: shallow water to 90 m (to 49 fm).

Habitat: primarily hard surfaces, common on sand; rock, coral and grassy areas; among hydroids at North Carolina (Williams, 1965) and oyster beds at Grand Isle, Louisiana (Behre, 1950).

Remarks: Regional lists include Florida (Wass, 1955; Tabb and Manning, 1961; Dragovich and Kelly, 1964; Abele, 1970; Lyons *et al.*, 1971; Menzel, 1971), Mississippi (Richmond, 1968), Louisiana (Behre, 1950), and Texas (Breuer, 1962). Oviparous females are known from South Carolina in August (Williams, 1965) and from Florida in March and August (Wass, 1955). Abele (1970) noted sexual dimorphism in the shape of the chelae. Wass (1955) commented on the ability of this crab to remain suspended in water by the rhythmic waving of the elongate legs, which are covered with fine setae on the distal portion. Listed from Brazil by Coelho and Ramos (1972).

Podocheila Stimpson, 1860

Podocheila curvirostris (A. Milne Edwards, 1879) (Crust. Rég. Mex., p. 196)

Rathbun, 1925, p. 50, pls. 19, 210; Chace, 1940, p. 56.

Range: Florida Straits; north coast of Cuba; Caribbean coast of Yucatan; Montserrat; Barbados; Grenadines.

Depth: 133 to 384 m (73 to 210 fm).

Habitat: coral, sand, and broken shell bottoms.

Remarks: Rathbun (1925) reported an ovigerous female from off Cuba in May.

Podochela gracilipes Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 126)

Hay & Shore, 1918, p. 454, pl. 37, fig. 6; Rathbun, 1925, p. 47, text-fig. 12, pl. 17; Williams, 1965, p. 243, figs. 220, 223I.

Range: North and South Carolina; Florida Keys and Straits; Dry Tortugas; west coast of Florida to off Alabama; north coast of Yucatan; Barbados; Colombia (Carib.); Guianas to Santa Catarina, Brazil.

Depth: 6 to 220 m (3 to 120 fm).

Habitat: sand, gravel, broken shell, rocky, and coral bottoms.

Remarks: Ovigerous females are known from North Carolina in December (Williams, 1965) and from Florida in March (Rathbun, 1925). Listed from shallow waters in Brazil by Coelho and Ramos (1972).

Podochela lamelligera (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 126)

Rathbun, 1925, p. 52, pl. 20, figs. 1-2.

Range: southeast Florida; off Key West, Florida; off northwest Florida.

Depth: 38 to 110 m (21 to 60 fm).

Habitat: sand, coral, and rocky substrates.

Remarks: Rathbun (1925) reports an ovigerous female from off Cape Florida in March. Listed by Chace (1956) from the Gulf of Mexico collections of the R/V *Oregon*.

Podochela macrodera Stimpson, 1860 (Ann. Lyc. Nat. Hist. New York 7: 196)

Rathbun, 1925, p. 44, text-fig. 11, pl. 16; Rathbun, 1933, p. 8.

Range: Bahamas; Florida Keys; west coast of Florida; off Caribbean coast of Yucatan; Cuba; Puerto Rico; Virgin Islands; Guadalupe; Curaçao, Netherlands Antilles; Brazil.

Depth: shallow water to 53 m (to 29 fm), rare to 91 m (50 fm).

Habitat: coral, sand, and gravel bottoms; from sponges; among rocks.

Remarks: Rathbun (1925) listed ovigerous females from off Florida in February and from Cuba in June. Coelho (1971c) listed this crab from depths of 20 to 53 m off Brazil.

Podochela riisei Stimpson, 1860 (Ann. Lyc. Nat. Hist. New York 7: 196)

Hay & Shore, 1918, p. 453, pl. 37, fig. 9; Rathbun, 1925, p. 33, text-fig. 9, pl. 11, figs. 1-2, pl. 208, fig. 2; Rathbun, 1933, p. 7, fig. 3; Chace, 1940, p. 56; Williams, 1965, p. 241, figs. 218, 223G.

Range: North Carolina; Bermuda; Bahamas; Florida Keys and Dry Tortugas; west and northwest coasts of Florida; Campeche, off Yucatan; Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Rio de Janeiro and south of Pernambuco, Brazil.

Depth: shallow water to 90 m (to 49 fm).

Habitat: coral, sand, shell, rock, and gravel substrates; on alga beds, grasses on sand bottoms; among *Sargassum* (Wass, 1955).

Remarks: Records of ovigerous females include the west coast of Florida in March, April, June and September (Lyons *et al.*, 1971) and from November through February (Rathbun, 1925). These crabs are often covered with a number of other organisms, including actinians, sponges, and rhizocephalan barnacles (Rathbun, 1925), and bryozoans, ascidians, and a red algae, *Calathamnion byssoideum*, as reported by Wass (1955). Regional lists include Florida (Wass, 1955; Tabb and Manning, 1961; Dragovich and Kelly, 1964; Abele, 1970; Lyons *et al.*, 1971; Menzel, 1971), Campeche (Hildebrand, 1955), and the Gulf of Mexico (Chace, 1956). Coelho (1971c) listed it from depths of 24 to 90 m, off Brazil.

Podochela sidneyi Rathbun, 1924 (Proc. U.S. Nat. Mus. 64: 1)

Rathbun, 1925, p. 39, pls. 12–13; Williams, 1965, p. 242, figs. 219, 223H; Felder, 1973a, p. 49, pl. 7, fig. 4.

Range: North Carolina; Dry Tortugas; west coast of Florida to central Texas coast; off north coast of Yucatan; northwest coast of Cuba.

Depth: shallow water to 187 m (to 102 fm).

Habitat: mud, broken shell, sand, coral, and rock bottoms; on alga-covered surfaces.

Remarks: Hildebrand (1954) collected specimens off Louisiana and Texas with small *Styela* (ascidians) attached to the legs, and some were covered with a dense hydroid growth. Listed from Florida by Wass (1955), Abele (1970), and Menzel (1971) and from Texas by Hildebrand (1954) and Leary (1967).

Pyromaia Stimpson, 1871

Pyromaia arachna Rathbun, 1924 (Proc. U.S. Nat. Mus. 64: 1)

Rathbun, 1925, p. 131, pls. 42–43; Pequegnat, 1970, p. 182.

Range: off South Carolina; off west coast of Florida to off east coast of Mexico; throughout all quadrants of the Gulf of Mexico.

Depth: 183 to 384 m (100 to 210 fm).

Habitat: off mud, mud-sand, and mud-shell bottoms.

Remarks: Rathbun (1925) reports ovigerous females off Florida in March. Listed from the Gulf collections of the R/V *Oregon* by Chace (1956) and from Texas by Leary (1967).

Pyromaia cuspidata Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 110)

Hay & Shore, 1918, p. 455, pl. 38, fig. 4; Rathbun, 1925, p. 129, text-fig. 49, pl. 41; Chace, 1940, p. 57; Williams, 1965, p. 240, figs. 216, 223E; Pequegnat, 1970, p. 181.

Range: North Carolina; south Florida; Florida Keys and Straits; Dry Tortugas; west coast of Florida; off Caribbean coast of Yucatan; north and south coasts of Cuba.

Depth: 27 to 549 m (15 to 300 fm).

Habitat: mud, sand, rock, coral, and pebble substrates.

Remarks: Chace (1940) noted that this crab autotomizes its legs readily. Williams (1965) reports ovigerous females from off Florida in February and July.

Stenorhynchus Lamarck, 1818

Stenorhynchus seticornis (Herbst, 1788) (Versuch Naturgesch. Krabben u. Krebse, vol. 1, p. 229)

Common Names: Arrow Crab; Araña del Mar

Hay & Shore, 1918, p. 455, pl. 37, fig. 8; Rathbun, 1925, p. 13, text-fig. 3, pls. 2-3; Rathbun, 1933, p. 6, fig. 1; Chace, 1940, p. 55; Williams, 1965, p. 244, figs. 222, 223K; Felder, 1973a, p. 48, pl. 7, fig. 1; Pequegnat & Ray, 1974, p. 237, figs. 11-12; Yang, 1976, p. 157.

Range: North Carolina to south Florida; Bermuda; Bahamas; Florida Keys and Straits; Dry Tortugas; west coast of Florida to south Texas; north and east coasts of Yucatan; north and south coasts of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands to Dominica; Netherlands Antilles; Colombia; Maranhão to Santa Catarina, Brazil.

Depth: near surface to 1489 m (814 fm).

Habitat: rock, coral, sand, sand-shell, and pebble bottoms; from sponges; off wharf pilings and rock jetties.

Remarks: The eastern Atlantic records cited by Rathbun (1925) are referred to *S. lanceolatus* (Brullé) by Yang (1967) and Barr (1975). Yang (1976) provides evidence that *S. seticornis* contains two species, one in shallow water, the other a deep water form. Williams (1965) states that this crab is more commonly collected at depths of 100 fm or less. Various authors use a variation of the genus name (e.g., *Stenorynchus*, in Williams, 1965). Yang (1967) describes larval stages. Ovigerous females occur off Texas in May and June (Hildebrand, 1954) and throughout the spring and summer over most of its range. Regional lists include Florida (Wass, 1955; Menzel, 1971), Mississippi (Franks *et al.*, 1972), Texas (Hildebrand, 1954; Leary, 1967), and the mid-ocean Gulf (Chace, 1956). Hartnoll (1965a) provided some notes on the biology and growth of Jamaican populations and Barr (1971, 1975) studied its biology in the Virgin Islands. Its occurrence in Brazil was noted by Coelho (1971c) and Coelho and Ramos (1972). Agonistic behavior of this crab is discussed by Schone (1968) and antennule chemosensitivity was tested by Hazlett (1971). Herrnkind, Stanton and Conklin (1976) described commensal relationships with an anemone in Florida.

Subfamily MAJINAE Samouelle, 1819

Temnonotus A. Milne Edwards, 1875

Temnonotus granulosus A. Milne Edwards, 1875 (Crust. Rég. Mex., p. 83)

As *T. simplex*—Rathbun, 1925, p. 342, pl. 249, figs. 10-12.

As *T. granulosus*—Rathbun, 1925, p. 341, pl. 249, figs. 7-9; Chace, 1940, p. 65, fig. 22.

Range: north coast of Cuba; Barbados.

Depth: 183 to 478 m (100 to 260 fm).

Habitat: coral, broken shell bottoms.

Remarks: Rathbun (1925) commented on the likelihood that the female type of *T. granulatus* was conspecific with the males of *T. simplex*, the only known specimens of these two species. Chace (1940) synonymized the two names on the basis of a male *T. granulatus* taken off Cuba. He also noted that distinctions in individual specimens are attributable to age variation as well as sexual dimorphism.

Temnonotus simplex A. Milne Edwards, 1875

Synonym of *T. granulatus* A. Milne Edwards, as revised by Chace (1940).

Subfamily MITHRACINAE Balss, 1929 (sensu Garth, 1958).

(This subfamily consists of Mithracinae Balss, 1929 plus Macrocoelominae Balss, 1929, derived from the Majinae Periceroida of Alcock, 1895. Glaessner treats the genera of this group under the subfamily Majinae Samouelle, 1819.)

Coelocerus A. Milne Edwards, 1875

Coelocerus spinosus A. Milne Edwards, 1875 (Crust. Rég. Mex., p. 85)

Rathbun, 1925, p. 446, text-fig. 130, pl. 263, pl. 264, figs. 1-2; Felder, 1973a, p. 49, pl. 7, fig. 6.

Range: west coast of Florida to off Alabama.

Depth: 24 to 64 m (13 to 35 fm).

Habitat: rock, coral, and sand bottoms.

Remarks: Garth (1958) states that this genus may be a link between *Libinia* and *Neodoclea* of the Pisinae and *Stenocionops* and *Macrocoeloma* of the Mithracinae.

Hemus A. Milne Edwards, 1875

Hemus cristulipes A. Milne Edwards, 1875 (Crust. Rég. Mex., p. 88)

Rathbun, 1925, p. 345, text-fig. 110, pl. 124, fig. 1, pl. 248, figs. 9-15; Rathbun, 1933, p. 21, fig. 23.

Range: northwest coast of Florida; north coast of Yucatan; Puerto Rico; Curaçao, Netherlands Antilles, Maranhão to Pernambuco, Brazil.

Depth 15 to 69 m (8 to 38 fm).

Habitat: sand, rock, and coral bottoms; in horn sponges and in the coral *Porites porites*.

Remarks: Listed by Wass (1955) and Menzel (1971) from northwest Florida. Brazilian listings include Coelho (1971c), Coelho and Ramos (1972), and Fausto Filho (1974).

Macrocoeloma Miers, 1879

Macrocoeloma camptocerum (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 112)

Hay & Shore, 1918, p. 457, pl. 38, fig. 12; Rathbun, 1925, p. 469, pl. 174, fig. 4, pl. 270, fig. 2; Williams, 1965, p. 264, figs. 244, 245K.

Range: North Carolina; southeast Florida; Florida Keys; south Florida to northwest Florida.

Depth: 4 to 35 m (2 to 19 fm).

Habitat: rock, sand, coral, and broken shell substrates; often from grassy areas.

Remarks: Rathbun (1925) reported specimens covered with various sponges, hydroids, bryozoans, ascidians, and infestations of rhizocephalid barnacles. Oviparous females were listed from North Carolina in August and from Florida in January to March. Wass (1955) and Menzel (1971) list this species from northwest Florida and Lyons *et al.* (1971) state that it is ovigerous in April, June, and October at Crystal River, Florida.

Macrocoeloma diplacanthum (Stimpson, 1860) (Ann. Lyc. Nat. Hist. New York 7: 183)

Rathbun, 1925, p. 478, pl. 169, fig. 1, pl. 269, figs. 1-3; Rathbun, 1933, p. 36.

Range: Bahamas; Key West, Florida; Cuba; Jamaica; Puerto Rico; Virgin Islands; Guadeloupe; Curaçao, Netherlands Antilles; Old Providence Island (Carib.).

Depth: 5 to 24 m (3 to 13 fm).

Habitat: off shallow reefs; sandy substrates.

Remarks: Rathbun (1925) listed specimens infected with rhizocephalid barnacles and others that were encrusted with algae.

Macrocoeloma eutheca (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 112)

Rathbun, 1925, p. 484, text-fig. 137, pl. 170, fig. 1, pl. 171, fig. 1; Rathbun, 1933, p. 37.

Range: Bahamas; southeast Florida; Florida Keys; north coast of Cuba; St. Croix, Virgin Islands; Barbados; Caribbean coast of Panama; Maranhão to Bahia, Brazil.

Depth: 30 to 214 m (16 to 117 fm).

Habitat: rock, broken shell, and coral substrates.

Remarks: Chace (1956) listed this crab from the R/V *Oregon* collections in the Gulf of Mexico. Listings from Brazil include Coelho (1971a, 1971c) and Coelho and Ramos (1972).

Macrocoeloma intermedium Rathbun, 1901 (Bull. U.S. Fish. Comm. 20: 75)

Rathbun, 1925, p. 486, text-fig. 138, pl. 170, fig. 2, pl. 171, fig. 2.

Range: north coast of Cuba; Dominica; Caribbean coast of Panama.

Depth: 62 to 298 m (34 to 163 fm).

Habitat: coral and broken shell bottoms.

Macrocoeloma laevigatum (Stimpson, 1860) (Ann. Lyc. Nat. Hist. New York 7: 181)

Rathbun, 1925, p. 483, text-fig. 136, pl. 169, figs. 2-3; Rathbun, 1933, p. 36.

Range: Florida Keys; north coast of Cuba; Jamaica; St. Thomas, Virgin Islands; Guadeloupe; Piauí to Alagoas, Brazil.

Depth: shore to 31 m (to 17 fm).

Habitat: rock and sand bottoms, often weedy.

Remarks: Listed from Brazil by Coelho (1971a, 1971c) and Coelho and Ramos (1972).

Macrocoeloma septemspinusum (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 113)

Rathbun, 1925, p. 477, pl. 173, figs. 2-3.

Range: South Carolina; Bahamas; Florida Keys; northeast quadrant of Gulf; Ceará to Rio Grande do Norte, Brazil.

Depth: shallow water to 145 m (to 79 fm), rarely to 212 m (116 fm).

Habitat: sand, broken shell, and coral substrates; on calcareous algae.

Remarks: Listed by Chace (1956) from the Gulf of Mexico. Brazilian records include Coelho (1971a, 1971c) and Coelho and Ramos (1972).

Macrocoeloma subparallelum (Stimpson, 1860) (Ann. Lyc. Nat. Hist. New York 7: 182)

Rathbun, 1925, p. 480, pl. 172; Rathbun, 1933, p. 36.

Range: north coast of Cuba; Jamaica; Haiti; Puerto Rico; St. Thomas, Virgin Islands; Guadeloupe; Barbados; Old Providence Island (Carib.); Rio Grande do Norte to Pernambuco, Brazil.

Depth: shallow water to 22 m (to 12 fm).

Habitat: on coral reefs; in tide pools; on bottoms of sand, grasses and weeds.

Remarks: Brazilian records include notes on its occurrence in the littoral zone (Coelho, 1971c) and listings by Coelho and Ramos (1972) and Fausto Filho (1974).

Macrocoeloma trispinosum trispinosum (Latreille, 1825) (Encyc. meth., Hist. Nat., vol 10, p. 142)

Common Names: Grass Crab; Sponge Crab; Decorator Crab

Hay & Shore, 1918, p. 457, pl. 38, fig. 11; Rathbun, 1925, p. 466, text-fig. 132, pl. 166, fig. 1, pl. 167; Rathbun, 1933, p. 35, fig. 31; Williams, 1965, p. 263, fig. 243; Felder, 1973a, p. 53, pl. 7, fig. 9.

Range: North Carolina; Bermuda; south Florida to northwest Florida; off Louisiana and Texas; Gulf and Caribbean coasts of Yucatan, Mexico; Cuba; Jamaica; Puerto Rico; St. Thomas to St. Lucia; Curaçao, Netherlands Antilles; Piauí to Bahia, Brazil.

Depth: shallow water to 82 m (to 45 fm).

Habitat: sand, rock and shell bottoms; among submerged mangrove roots; from wharf pilings; from floating masses of *Sargassum*.

Remarks: Rathbun (1925) listed three varieties of this species, two of which are listed here as subspecies (the typical form and *M. t. nodipes*), and a third, considered an intermediate form. Behre (1950) listed an unspecified form of *Macrocoeloma* from Louisiana and Menzel (1971) and Lyons *et al.* (1971) record this crab as uncommon in Florida. Ovipigerous female records were summarized by Williams (1965). Hartnoll (1965a) described the biology and growth of this crab in Jamaica. Many specimens are encrusted with sponges, which are

attached to the hairs of the carapace and legs, providing the basis for two of its common names. Records from Brazil include Coelho (1971a, 1971c), Coelho and Ramos (1972) and Fausto Filho (1974).

Macrocoeloma trispinosum nodipes (Desbonne, 1867) (Crust. de la Guadeloupe, p. 15)

Rathbun, 1925, p. 468, pl. 166, fig. 2, pl. 168, fig. 2; Rathbun, 1933, p. 36; Williams, 1965, p. 264.

Range: North Carolina; Bermuda; south to northwest coasts of Florida; Florida Keys and Dry Tortugas; Cuba; Puerto Rico; Antigua; Brazil.

Depth: shore to 48 m (to 26 fm).

Habitat: sand, shell, rock, and coral bottoms; grassy areas.

Remarks: Rathbun (1925) lists several specimens that were covered with sponges and one crab that was encrusted with worm tubes. She reported ovigerous females from Florida in December and from Cuba in June. Listed from Florida by Wass (1955), Abele (1970), and Menzel (1971), and from the Gulf collections of the R/V *Oregon* by Chace (1956). Abele (1970) treats this form as a separate species and notes the need for revision of this genus.

Macrocoeloma trispinosum variety Rathbun, 1925 (Bull. U.S. Nat. Mus. 129: 468)

Rathbun, 1925, p. 468, pl. 168, fig. 1; Rathbun, 1933, p. 36.

Range: North Carolina; Bahamas; southeast and west coasts of Florida; Florida Keys and Dry Tortugas; Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Curaçao, Netherlands Antilles.

Depth: low tide mark to 51 m (to 28 fm).

Habitat: sand, shell, rock, and coral bottoms.

Remarks: Williams (1965) followed the practice of Rathbun (1925) in treating this variety as an unnamed but distinct form, an intermediate linking the typical subspecies and *M. t. nodipes*. If the latter is raised to the status of a full species (as in Abele, 1970), then this variety will require reevaluation as a subspecies or species.

Microphrys H. Milne Edwards, 1851

Microphrys antillensis Rathbun, 1920 (Proc. Washington Acad. Sci. 33: 24)

As *M. platysoma*—Hay & Shore, 1918, p. 459, pl. 38, fig. 9.

As *M. antillensis*—Rathbun, 1925, p. 498, text-fig. 141, pl. 176, figs. 3, 4; Rathbun, 1933, p. 38; Williams, 1965, p. 260, figs. 240, 245G.

Range: North Carolina; Bimini; west coast of Florida; north coast of Cuba; Jamaica; Puerto Rico; Brazil.

Depth: 4 to 38 m (2 to 21 fm).

Habitat: mud, coral, sand, shell, and weed bottoms.

Remarks: Ovigerous females were reported from North Carolina in September (Rathbun, 1925) and from Florida in June and from Bimini in November (Williams, 1965). Listed from Brazil by Coelho (1971c) and Coelho and Ramos (1972).

Microphrys bicornutus (Latreille, 1825) (Encyc. meth., Hist. Nat. Insectes, vol. 10, p. 141)

Common Name: Dirty Decorator Crab

Hay & Shore, 1918, p. 459, pl. 38, fig. 10; Rathbun, 1925, p. 489, text-fig. 139, pl. 175; Rathbun, 1933, p. 37, fig. 32; Williams, 1965, p. 259, figs. 239, 245F.

Range: North Carolina to south Florida; Bermuda; Bahamas; Florida Keys and Dry Tortugas; west and northwest coasts of Florida; north coast of Cuba; Jamaica; Puerto Rico; St. Thomas to Barbados; Old Providence Island in the Caribbean Sea; Caribbean coast of Panama to Venezuela; Curaçao; Trinidad; Island of Santa Anna to Santa Catarina, Brazil.

Depth: shallow water to 30 m (to 16.5 fm).

Habitat: common on coral reefs; on a variety of substrates, including rock, shell, sand, and mud; among grass, mangrove roots, and on sponges; often covered with anemones, algae, sponges, etc.

Remarks: Rathbun (1925) notes that Latreille's original type locality, "Nouvelle Hollande," is an error. Ovipigerous females occur from March to August in the Caribbean and from November to January in the West Indies, Brazil, and Venezuela (Williams, 1965). Wass (1955) and Menzel (1971) list this species as rare in northwest Florida. Listed from Brazil by Coelho (1971a), Coelho and Ramos (1972), and Fausto Filho (1974). Hartnoll (1965a) commented on the biology and growth of this crab in Jamaica and Coelho (1971c) provided ecological notes on Brazilian specimens. Zoeal stages have been described by Lebour (1944) and by Hartnoll (1964b). Hazlett (1972a, 1972b) and Hazlett and Estabrook (1974) analyzed agonistic behavior and Hazlett (1971) tested the antennular chemosensitivity of this species. Williams (1965) lists records of copepods and a tapeworm from this crab.

Microphrys platysoma (Rathbun, 1901), not (Stimpson, 1860).

Specimens from Puerto Rico were designated by Rathbun (1920) as *M. antillensis*, to which all of the Atlantic specimens are referred, separating them from the Pacific form of Stimpson.

Mithrax Desmarest, 1823

(Although Latreille, 1817 had been traditionally recognized as the author of this genus, Garth (1958) questioned the validity of Latreille's citation and attributes the first valid citation to Desmarest.)

Subgenus ***Mithraculus*** White, 1847

Mithrax (Mithraculus) cinctimanus (Stimpson, 1860) (Amer. J. Sci. 29: 132)

Rathbun, 1925, p. 438, pl. 158; Rathbun, 1933, p. 32.

Range: Bahamas; southeast Florida; Florida Keys and Dry Tortugas; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; St. Martin; Antigua; Curaçao.

Depth: shallow water.

Habitat: on coral reefs; rocky bottoms; inside sponges.

Mithrax (Mithraculus) coryphe (Herbst, 1801) (Natur. Krabben u. Krebse, vol. 3, pt. 2, p. 8)

Rathbun, 1925, p. 426, pl. 153; Rathbun, 1933, p. 31.

Range: Bahamas; southeast Florida; Florida Keys; north coast of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands to Barbados; Caribbean coast of Panama to Colombia; Curaçao; Trinidad; Ceará and Fernando de Noronha to São Paulo, Brazil.

Depth: shallow water to 55 m (to 30 fm).

Habitat: in cavities of corals, rocks, and sponges; on sand, broken shell, grass, and mud bottoms.

Remarks: Listed from Brazil by Coelho and Ramos (1972).

Mithrax (Mithraculus) forceps (A. Milne Edwards, 1875) (Crust. Rég. Mex., p. 109)

Hay & Shore, 1918, p. 457, pl. 38, fig. 1; Rathbun, 1925, p. 431, pl. 156; Rathbun, 1933, p. 32; Chace, 1940, p. 67; Williams, 1965, p. 258, figs. 238, 245E; Pequegnat & Ray, 1974, p. 236, figs. 1-4.

Range: Bermuda; North Carolina to south Florida; Florida Keys and Dry Tortugas; west and northwest coasts of Florida; West Flower Garden Bank, off Texas; south coast of Cuba; Puerto Rico; St. Thomas, Virgin Islands; Barbados; Old Providence Island (Carib.); Netherlands Antilles; Venezuela; Trinidad; Ceará to Rio de Janeiro, Brazil.

Depth: intertidal to 90 m (to 49 fm).

Habitat: under stones and dead coral; in crevices along rocky shores and reefs; in sponges; on sand, shell, coral, rock, and grass bottoms.

Remarks: Ovigerous females have been reported from Florida in November to February, from the Gulf of Mexico in February, and from the southern Caribbean in April, mid-summer, September, and November (Williams, 1965). Listed from northwest Florida by Wass (1955), Abele (1970), and Menzel (1971). Chace (1956) recorded this crab from the Gulf of Mexico collections of the R/V *Oregon* and Pequegnat and Ray (1974) state that this species is one of the most common brachyurans on the West Flower Garden reefs, often taken on sponges. Williams (1965) reports this crab from the sponge, *Stematumenia strobilinia*, off North Carolina. Lebour (1944) described some of the larval stages. Threat behavior was described and illustrated by Schone (1968). Listed from Brazil by Coelho (1971a, 1971c), Coelho and Ramos (1972) and Fausto Filho (1974).

Mithrax (Mithraculus) ruber (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 118)

Rathbun, 1925, p. 432, pl. 157; Rathbun, 1933, p. 32.

Range: north coast of Cuba; Puerto Rico; St. Thomas, Virgin Islands to Barbados; Curaçao.

Depth: shallow water to 46 m (to 25 fm), rare to 154 m (84 fm).

Habitat: sand, shell, grass, coral, and mud substrates; off coral reefs; in sponges.

Mithrax (Mithraculus) sculptus (Lamarck, 1818) (Hist. Anim. sans Vert., vol. 5, p. 242)

Rathbun, 1925, p. 422, text-figs. 125–126, pl. 152; Rathbun, 1933, p. 31.

Range: Bahamas; southeast Florida; Florida Keys and Dry Tortugas; north coast of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Antigua; Barbados; Swan Island and Old Providence Island (Carib.); Belize; Curaçao; Rio Grande do Norte to Bahia, Brazil.

Depth: shallow water to 55 m (to 30 fm).

Habitat: abundant on coral reefs; under stones at low tide; on sand, shell, grass, and mud bottoms.

Remarks: Hartnoll (1965a) provides data on the biology and growth of this crab in Jamaica. Listed from Brazil by Coelho and Ramos (1972) and by Fausto Filho (1974).

Subgenus ***Mithrax*** Desmarest, 1823

Mithrax (Mithrax) acuticornis Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 116)

Rathbun, 1925, p. 388, pl. 136, figs. 1–2, pl. 257, fig. 1; Rathbun, 1933, p. 29, fig. 28; Felder, 1973a, p. 52, pl. 7, fig. 10.

Range: southeast coast of Florida; Florida Keys and Straits; Dry Tortugas; west and northwest coasts of Florida; off Texas; north and east coasts of Yucatan, Mexico; Puerto Rico; Santa Cruz to Grenadines in the Lesser Antilles; Amapá to Espirito Santo, Brazil.

Depth: 11 to 103 m (6 to 56 fm).

Habitat: sand, mud, broken shell, rock, and coral substrates.

Remarks: Rathbun (1925) compared this species with the young of *M. cornutus* and *M. spinosissimus*, with which it can be confused. Felder (1973a) provided the first northwestern Gulf of Mexico records, based on specimens collected off Galveston and Port Mansfield, Texas. Listed from Brazil by Coelho (1971a, 1971c) and Coelho and Ramos (1972).

Mithrax (Mithrax) cornutus Saussure, 1857 (Rev. Mag. Zool., ser. 2, 9: 501)

Common Names: Coral Crab; Red Spider Crab

Rathbun, 1925, p. 386, pl. 137, figs. 3–4, pl. 256.

Range: Bermuda; Florida Straits; north coast of Cuba; between Jamaica and Haiti; Dominica; Martinique; off Bahia, Brazil.

Depth: shallow water to 1077 m (589 fm).

Habitat: coral, sand, and broken shell bottoms.

Remarks: Listed by Chace (1956) from the Gulf of Mexico.

Mithrax (Mithrax) hispidus (Herbst, 1790) (Natur. Krabben u. Krebse, vol. 1, p. 245 (not p. 247)).

Common Name: Coral Crab

Rathbun, 1925, p. 406, text-fig. 124, pls. 145–146, pl. 147, fig. 3; Rathbun, 1933, p. 30; Williams, 1965, p. 256, figs. 236, 245C; Pequegnat & Ray, 1974, p. 236, figs. 5–10.

Range: Delaware Bay to south Florida; Bermuda; Bahamas; Florida Keys and Dry Tortugas; West Flower Garden Bank, off Texas; Jamaica; Curaçao; Pará to São Paulo, Brazil.

Depth: shallow water to 65 m (36 fm).

Habitat: sand, shell, and stone bottoms; on coral reefs; inside sponges; occasionally on the sea grass, *Halodule*.

Remarks: Plentiful on the coral reefs of the West Flower Garden Bank (Pequegnat and Ray, 1974). Listed from Brazil by Coelho (1971a, 1971c) and by Coelho and Ramos (1972).

Mithrax (Mithrax) holderi Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 117)

Rathbun, 1925, p. 392, pl. 138, figs. 1–2, pl. 257, fig. 2; Rathbun, 1933, p. 29.

Range: Florida Keys and Dry Tortugas; north and south coasts of Cuba; Jamaica; Puerto Rico; Virgin Islands.

Depth: intertidal to 38 m (to 21 fm).

Habitat: coral bottoms.

Mithrax (Mithrax) pilosus Rathbun, 1892 (Proc. U.S. Nat. Mus. 15: 262)

Rathbun, 1925, p. 394, pl. 138, fig. 3, pl. 258; Rathbun, 1933, p. 29.

Range: Bahamas; Florida Keys and Dry Tortugas; Vera Cruz, Mexico; Cuba; Puerto Rico; St. Thomas, Virgin Islands to Barbados; Caribbean coast of Panama; Venezuela.

Depth: data not available.

Habitat: rare on stony bottoms; off reefs.

Mithrax (Mithrax) pleuracanthus Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 116)

As *M. depressus*—Hay & Shore, 1918, p. 458, pl. 38, fig. 2.

As *M. pleuracanthus*—Hay & Shore, 1918, p. 458, pl. 38, fig. 3; Rathbun, 1925, p. 411, pl. 150; Rathbun, 1933, p. 31; Williams, 1965, p. 257, figs. 237, 245D.

Range: Bermuda; North and South Carolina; southeast Florida; Florida Keys and Dry Tortugas; Bahamas; west and northwest coasts of Florida; north coast of Yucatan; north coast of Cuba; Puerto Rico; St. Thomas, Virgin Islands; St. Martin; Old Providence Island (Carib.); Curaçao; Venezuela.

Depth: shallow water to 51 m (to 28 fm).

Habitat: common on rocky, gravel, and broken shell substrates; occasionally on sand and mud bottoms; in the sponge *Stematumenia strobilinia* at Tortugas (Pearse, 1934). Often encrusted with bryozoans, serpulid worms, etc. (after Williams, 1965).

Remarks: Listed as common in northwest Florida by Wass (1955), Abele (1970), and Menzel (1971). Williams (1965) notes that this species is often associated with *Mithrax forceps* on the banks off North Carolina in April, from St. Thomas in July, and from Venezuela in September.

Mithrax (Mithrax) spinosissimus (Lamarck, 1818) (Hist. Nat. Anim. sans Vert., vol. 5, p. 241)

Common Names: Cangrejo de la Santa Virgen; Cabouca; Lazy Crab

Rathbun, 1925, p. 383, pl. 135; Rathbun, 1933, p. 29; Chace, 1940, p. 67; Williams, 1965, p. 254, figs. 234, 245A.

Range: North or South (?) Carolina; Florida Keys and Dry Tortugas; north and south coasts of Cuba; Jamaica; Haiti; Virgin Islands; Guadeloupe. Type locality of "Ile-de-France" is erroneous.

Depth: shallow water to 179 m (to 98 fm).

Habitat: among rocks and on coral sand bottoms; crab is often encrusted with stalked barnacles, bryozoans, serpulid worms, red foraminiferans, etc.

Remarks: Oviparous females were reported from Cuba in May and June by Rathbun (1925). Hazlett and Rittschof (1975) reported on spatial patterns of activity. Brownell, Provenzano and Martinez (1977) reported on attempts to commercially culture this crab in Venezuela.

Mithrax (Mithrax) verrucosus H. Milne Edwards, 1832 (Mag. Zool., vol. 2, class 7, pl. 4)

Rathbun, 1925, p. 400, pl. 144; Rathbun, 1933, p. 30; Williams, 1965, p. 255, figs. 235, 245B.

Range: South Carolina; Bahamas; southeast Florida; Florida Keys and Dry Tortugas; north coast of Cuba; Swan Islands (Carib.) Jamaica; Hispaniola; Puerto Rico; St. Thomas, Virgin Islands; Guadeloupe; Curaçao; Martinique; Rocas and Fernando do Noronha, Brazil.

Depth: shallow water, near shore.

Habitat: among rocks; hides in crevices and holes; often found near madrepores; has nocturnal habits.

Remarks: Found in *Porites porites* at Curaçao. Pearse (1932b) determined the freezing point of hemolymph from this crab at Dry Tortugas. Listed from Brazil by Coelho (1971a, 1971c), Coelho and Ramos (1972) and by Fausto Filho (1974).

Stenocionops Desmarest, 1823

Stenocionops furcata furcata (Oliver, 1791) (Encyc. méth., Hist. Nat., Insects, vol. 6, p. 174)

Common Names: Decorator Crab; Macca Crab

Rathbun, 1925, p. 449, text-fig. 131, pls. 160–161; Rathbun, 1933, p. 33, fig. 30; Guinot-Dumortier, 1960, p. 180, fig. 21a–b.

Range: Georgia; Florida (location unspecified); ? Gulf of Mexico; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Dominica; Barbados; French Guiana; Paraíba to Rio de Janeiro, Brazil; South Africa.

Depth: shallow water to 64 m (to 35 fm).

Habitat: mud, sand, coral, rock, and shell bottoms; on wharf pilings.

Remarks: The inclusion of this form in the Gulf fauna is doubtful. Although Leary (1967) and Felder (1973a) report this form from Texas, both reports are

based on a listing by Hildebrand (1954) in which the subspecies identification may have been in error. Felder (1973a) indicates that more recent collections in the northwestern Gulf are of crabs similar to *S. furcata coelata*, and other records confirm the presence of this latter form in the region. Neither of the two specimens cited by Rathbun (1925, p. 542) are determined for the Gulf. Guinot-Dumortier (1960) remarked that the male pleopod of this species is very similar to the pleopods of three Pacific species figured by Garth (1958). Listed from Brazil by Coelho and Ramos (1972).

Stenocionops furcata coelata (A. Milne Edwards, 1878) (Bull. Soc. Philom., ser. 7, 2:224)

Hay & Shore, 1918, p. 460, pl. 39, fig. 3; Rathbun, 1925, p. 450, pl. 164; Rathbun, 1933, p. 34; Williams, 1965, p. 261, figs. 241, 245H; Felder, 1973a, p. 53, pl. 7, fig. 12.

Range: North Carolina to south Florida; Florida Keys and Dry Tortugas; west and northwest coasts of Florida; Alabama to Texas; north and east coasts of Yucatan; north coast of Cuba; Puerto Rico; St. Lucia; Barbados.

Depth: shallow water to 110 m (to 60 fm), rarely to 508 m (278 fm).

Habitat: sand, shell, and coral bottoms, usually of coarse material; on shelly reefs off North Carolina.

Remarks: Early Gulf records distinguishing this subspecies from the typical form may be confused (Felder, 1973a). A single specimen taken off St. Joseph Island, Texas (Hildebrand, 1954) is probably *coelata*; it was covered with algae, hydroids, bryozoans, three small *Ostrea*, and three large *Calliactis tricolor*. Listed from the Gulf of Mexico (Chace, 1956) and from northwest Florida (Wass, 1955; Hulings, 1961; Abele, 1970). Oviparous females occur in Florida from March to August (Williams, 1965).

Stenocionops spinimana (Rathbun, 1892) (Proc. U.S. Nat. Mus. 15: 240)

Hay & Shore, 1918, p. 460, pl. 39, fig. 2; Rathbun, 1925, p. 457, pl. 267; Williams, 1965, p. 262, figs. 242, 245I; Pequegnat, 1970, p. 182.

Range: North and South Carolina; Florida Keys; west coast of Florida to off Mississippi.

Depth: 37 to 227 m (20 to 124 fm).

Habitat: mud, sand, shell, coral, and rock bottoms.

Remarks: Williams (1965) noted the age-related variability in the morphology of this and related species. Rathbun (1925) listed oviparous females from South Carolina in December. Listed by Chace (1956) from the Gulf of Mexico and by Franks *et al.* (1972) off Mississippi at 37 to 91 m depth.

Stenocionops spinosissima (Saussure, 1857) (Rev. Mag. Zool., ser. 2, 9: 501)

Rathbun, 1925, p. 455, pl. 165, fig. 2, pl. 264, figs. 3-4, pl. 265; Chace, 1940, p. 67; Williams, McCloskey & Gray, 1968, p. 62; Pequegnat, 1970, p. 182; Felder, 1973a, p. 52, pl. 7, fig. 11.

Range: North Carolina; south and southwest Florida; off Texas and east coast of Mexico; north coast of Cuba; Haiti; Guadeloupe; Dominica; Rio de Janeiro and Fernando de Noronha, Brazil.

Depth: 46 to 480 m (25 to 260 fm); center of depth distribution in Gulf is at 110 to 183 m (60 to 100 fm) (Pequegnat, 1970).

Habitat: mud and sand bottoms.

Remarks: Hildebrand (1954) found this crab to be common off the Texas coast at depths of 25 to 37 fm (46 to 68 m) and he reported ovigerous females in February and April. Listed from Texas by Leary (1967) and from the Gulf of Mexico by Chace (1956). Recorded from Brazil by Coelho and Ramos (1972) and by Fausto Filho (1974).

***Teleophrys* Stimpson, 1860**

***Teleophrys ornatus* Rathbun, 1901 (Bull. U.S. Fish. Comm. 20: 65)**

Rathbun, 1925, p. 444, text-fig. 129, pl. 159, figs. 3-4, pl. 262, figs. 8-9; Rathbun, 1933, p. 33, fig. 29.

Range: off northeast Yucatan (Gulf); Puerto Rico; St. Croix; Fernando de Noronha, Brazil.

Depth: 7 to 44 m (4 to 24 fm).

Habitat: rock and coral bottoms.

Remarks: Listed from Brazil by Coelho and Ramos (1972) and Fausto Filho (1974).

***Thoe* Bell, 1836**

***Thoe puella* Stimpson, 1860 (Ann. Lyc. Nat. Hist. New York 7: 178)**

Rathbun, 1925, p. 348, text-figs. 111-112, pl. 125, figs. 1-2; Rathbun, 1933, p. 21, fig. 24.

Range: Florida Keys and Dry Tortugas; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Guadeloupe; Curaçao.

Depth: Shallow water.

Habitat: in and on coral reefs.

Subfamily OPHTHALMIINAE Balss, 1929

***Picroceroides* Miers, 1886**

***Picroceroides tubularis* Miers, 1886 (*Challenger* Rept., Zool. 17: 77)**

Rathbun, 1925, p. 354, text-fig. 115, pl. 126, pl. 254, figs. 2-5.

Range: Bahamas; southeast Florida; north and south coasts of Cuba; between Jamaica and Haiti; St. Thomas, Virgin Islands; Maranhão to Espirito Santo, Brazil.

Depth: shallow water to 110 m (to 60 fm).

Habitat: coral and broken shell bottoms.

Remarks: Listed from Brazil by Coelho (1971c), Coelho and Ramos (1972), and Fausto Filho (1974).

Pitho Bell, 1835***Pitho aculeata*** (Gibbes, 1850) (Proc. Amer. Assoc. Adv. Sci. 3: 171)

Rathbun, 1925, p. 357, text-fig. 116c, pl. 127, pl. 251, fig. 1; Rathbun, 1933, p. 23, fig. 26.

Range: Bahamas; Florida Keys and Dry Tortugas; west coast of Florida; north coast of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Guadeloupe; Old Providence Island (Carib.); Netherlands Antilles.

Depth: Shallow water.

Habitat: sand, shell, coral, grass, and mud bottoms; among algae in lagoons; on *Sargassum*; on banks at low tide.

Pitho anisodon (von Martens, 1872) (Arch. f. Naturg. 38: 83)

Rathbun, 1925, p. 368, text-figs. 116b, 117d, 118, pl. 131, pl. 251, fig. 2; Rathbun, 1933, p. 24.

Range: Bahamas; south, west, and northwest coasts of Florida; Florida Keys; north coast of Cuba; Jamaica; Puerto Rico; Guadeloupe; Curaçao, Netherlands Antilles.

Depth: shallow water to 22 m (to 12 fm).

Habitat: rock, sand, mud, coral, and grass bottoms.

Remarks: Reported from Florida by Wass (1955), Tabb and Manning (1961), Dragovich and Kelly (1964), Abele (1970), and Menzel (1971). Tabb and Manning (1961) noted the presence of ovigerous females in Florida Bay when salinities were fully marine and in Tampa Bay in March. Abele (1970) remarked on the variability of lateral and orbital spines with size of crab.

Pitho laevigata (A. Milne Edwards, 1875) (Crust. Rég. Mex., p. 116)

Rathbun, 1925, p. 372, pl. 132, figs. 3-4, pl. 133, fig. 3, pl. 250, figs. 11-13.

Range: west and northwest coasts of Florida; Antilles, loc. unsp.; Colombia; Trinidad.

Depth: shallow water, listed from 10.5 m (about 6 fm).

Habitat: coral, rock, and grass bottoms.

Remarks: Listed from northwest Florida by Wass (1955). Rathbun (1925) described a variety of this crab based on a female collected from the west coast of Florida.

Pitho lherminieri (Schramm, 1867) (Crust. Guadeloupe, Desbonne & Schramm, p. 20)

Hay & Shore, 1918, p. 459, pl. 38, fig. 8; Rathbun, 1925, p. 362, text-fig. 116a, 117b, pl. 128, figs. 1-2, pl. 129, figs. 1-2, pl. 252, fig. 2; Rathbun, 1933, p. 24; Williams, 1965, p. 246, figs. 224, 233A.

Range: North and South Carolina; Bahamas; Florida Keys; west and northwest coasts of Florida; Vera Cruz, Mexico; north coast of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands to Martinique; Old Providence Island (Carib.); Curaçao; Cape St. Roque to São Paulo, Brazil.

Depth: shallow water to 51 m (to 28 fm), rarely to 220 m (120 fm).

Habitat: coral, sand, shell, rock, and grass bottoms; rarely on mud.

Remarks: Listed from northwest Florida by Wass (1955); recorded from Brazil by Coelho (1971c), Coelho and Ramos (1972) and Fausto Filho (1974). Ovipigerous females have been collected from the Bahamas and Florida in May to November and from Brazil in December (Williams, 1965).

Pitho mirabilis (Herbst, 1794) (Naturg. Krabben u. Krebse, vol. 2, p. 152)

Rathbun, 1925, p. 366, text-figs. 116d, 117c, pl. 128, fig. 3, pl. 129, fig. 3, pl. 253, fig. 1; Rathbun, 1933, p. 24.

Range: Bahamas; Florida Keys; Puerto Rico; Guadeloupe.

Depth: shallow water.

Habitat: rock and coral bottoms.

Remarks: Only part of the original species description by Herbst is valid (Rathbun, 1925).

Tyche Bell, 1835

Tyche emarginata White, 1847 (Ann. Mag. Nat. Hist. 20: 206)

Hay & Shore, 1918, p. 461, pl. 39, fig. 4; Rathbun, 1925, p. 508, pl. 272, pl. 273, figs. 7-12; Garth, 1946, p. 406, text-fig. 1; Williams, 1965, p. 247, figs. 225, 226, 233B.

Range: North Carolina; Bahamas; Florida Keys and Dry Tortugas; west and northwest coasts of Florida; St. Thomas, Virgin Islands; Guadeloupe; Cape St. Roque, Brazil.

Depth: shallow water to 37 m (to 20 fm).

Habitat: shell and rock bottoms.

Remarks: Garth (1946) compared this species with its Pacific analogue, *T. lamellifrons*. Listed from Florida by Wass (1955) and Lyons *et al.* (1971) and from the Gulf collections of the R/V *Oregon* by Chace (1956).

Subfamily PISINAE Dana, 1852

Chorinus Latreille, 1825

Chorinus heros (Herbst, 1790) (Naturg. Krabben u. Krebse, vol. 1, pl. 18, fig. 2)

Rathbun, 1925, p. 305, text-fig. 101, pl. 107, pl. 246, figs. 3-5; Rathbun, 1933, p. 20, fig. 21.

Range: Bermuda; Florida Keys and Dry Tortugas; Cuba; Caribbean coast of Yucatan, Mexico; Jamaica; Hispaniola; Puerto Rico; St. Croix; Barbados; Ceará to Bahia, Brazil.

Depth: shallow water to 48 m (to 26 fm).

Habitat: sand, shell, rock, and coral bottoms.

Remarks: Listed from the Gulf of Mexico by Chace (1956) and from Brazil by Coelho (1971c) and Coelho and Ramos (1972).

Holoplites Rathbun, 1894**Holoplites armata** (A. Milne Edwards, 1880) (Crust. Rég. Mex., p. 348)

Rathbun, 1925, p. 307, text-fig. 102, pl. 108, pl. 245, figs. 6-8; Chace, 1940, p. 64; Bullis & Thompson, 1965, p. 12.

Range: north coast of Cuba; St. Vincent; Grenadines; Barbados; Pará, Brazil.

Depth: 161 to 798 m (88 to 387 fm).

Habitat: coral, shell, and rock bottoms.

Remarks: Listed from Brazil by Coelho and Ramos (1972).

Libinia Leach, 1815**Libinia dubia** H. Milne Edwards, 1834 (Hist. Nat. Crust., vol. 1, p. 300)

Hay & Shore, 1918, p. 456, pl. 38, fig. 5; Rathbun, 1925, p. 313, text-figs. 105-106, pls. 114-115, 122, fig. 1; Williams, 1965, p. 252, figs. 232, 233G; Felder, 1973a, p. 52, pl. 7, fig. 8.

Range: Cape Cod, Massachusetts to south Florida; Bahamas; Florida Keys; west Florida to south Texas; Cuba; off Gabon, western Africa.

Depth: shallow water to 46 m (to 25 fm).

Habitat: primarily on mud and mud-sand substrates; also on sand, gravel, and shell bottoms; often near shore and occasionally in tide pools. Juveniles are usually covered with hydroids, ascidians, sponges, etc., but older adults are almost always clean.

Remarks: Gulf reports of this species may be confused with those for *L. emarginata*, especially records of juveniles for which identification is difficult and further complicated by the variety of attached organisms. Regional lists include Florida (Wass, 1955; Tabb and Manning, 1961; Dragovich and Kelly, 1964; Lyons *et al.*, 1971; Menzel, 1971), Mississippi (Richmond, 1962; Christmas and Langley, 1973), Louisiana (Behre, 1950; Hoese and Valentine, 1972), and Texas (Leary, 1967). Fotheringham and Brunenmeister (1975) describe some aspects of natural history for this crab in the Gulf.

Tabb and Manning (1961) found that larger adults were prevalent in the sponge-algae areas of Florida Bay, whereas smaller individuals were more common in the *Thalassia* beds. Wass (1955) states that this species is more common than *L. emarginata* in the shallower harbor and bay waters, but that the reverse is true in the more marine outer waters of northwest Florida. This crab occurred in salinity ranges of 20.4 to 39 ppt in Texas bays after a drought (Hoese, 1960) and down to 12 ppt at Crystal River, Florida (Lyons *et al.*, 1971). In Tampa Bay, juveniles are common on *Gracilaria* beds and this species is the most common spider crab of the area (Dragovich and Kelly, 1964). Oviparous females have been reported in Florida from January to July (Dragovich and Kelly, 1964; Lyons *et al.*, 1971).

Several authors have commented on the presence of young *L. dubia* in or on the cabbagehead jellyfish, *Stomolophus meleagris*. Corrington (1927) and Gutsell (1928) record this relationship in Carolina waters and Williams (1965) states that small crabs have been found in the genital pits and subumbrellar space.

Jachowski (1963) noted a similar relationship with *Aurelia aurita*. Wass (1955) noted that one female crab was covered with 93 barnacles (*Balanus*) and Pearse (1952) reported copepods in the gills and on the carapace of this crab in Texas waters.

Development of this species in the laboratory was described by Sandifer and van Engel (1971). Ayres (1938) compared relationships between habitat and oxygen consumption and Gray (1957) correlated habitat and gill surface area in this species. Other physiological studies include tolerance to desiccation and salinity changes (Pearse, 1929) and the physiological activity and neurosecretions of the pericardial organs (Berlind and Cooke, 1970).

Libinia emarginata Leach, 1815 (Zool. Misc., 2: 130)

Common Name: Common Spider Crab

Hay & Shore, 1918, p. 456, pl. 38, fig. 6; Rathbun, 1925, p. 311, text-figs. 103-104, pls. 110-113; Williams, 1965, p. 252, figs. 231, 233H; Felder, 1973a, p. 52, pl. 7, fig. 7.

Range: Nova Scotia to south Florida; Florida Keys; west coast of Florida to Mexico. Pacific coast records are probably erroneous (Garth, 1958).

Depth: shore to 49 m (to 27 fm), rarely to 124 m (68 fm).

Habitat: on all types of substrates, but most common on mud and mud-sand in shallow waters.

Remarks: Gulf records of this species and *L. dubia* may be confused due to similarity of the juvenile stages. Wass (1955) compares the key morphological features of the two species. Listed from Florida (Wass, 1955; Menzel, 1971), Mississippi (Richmond, 1962; Franks *et al.*, 1972), Louisiana (Behre, 1950; Hoese and Valentine, 1972), Texas (Gunter, 1950; Hildebrand, 1954; Parker, 1959; Leary, 1967), Mexico (Hildebrand, 1954), and the Gulf of Mexico (Chace, 1956).

Hildebrand (1954) states that this species is the most common large spider crab in the western Gulf of Mexico, reversing the numerical dominance relationship with *L. dubia* that is found in the eastern Gulf. Like *L. dubia*, young *L. emarginata* are often associated with the cabbagehead jellyfish, *Stomolophus meleagris*. Ovigerous females are present in the western Gulf in February and are common in July. Hoese and Valentine (1972) collected a crab from the Chandeleur Islands that was covered with the bryozoan, *Bugula*. Musick and McEachran (1972) reported a depth range of 18 to 51 m for this crab in Chesapeake Bight. Aldrich (1976) reported predation by this crab on the starfish, *Asterias*. Forward (1977) described shadow responses of the larval stages.

Other studies include: measurement of gill surface area in relation to habitat (Gray, 1957), biochemical adaptations to temperature variation (Vernberg and Vernberg, 1968), osmoregulation (Gilles, 1970), physiology of molting (Skinner and Graham, 1972), sterol synthesis in larval stages (Whitney, 1969), vitellogenesis (Hinsch and Cone, 1969), sperm structure (Hinsch, 1973), growth biometrics (Aldrich, 1974), reproductive physiology (Hinsch, 1970), neural fine structure (Skobe and Nunnemacher, 1970), antennule chemosensitivity (Haz-

lett, 1971), pericardial organ neurosecretion (Berlind and Cook, 1970), and behavior related to copulation and reproduction (Hinsch, 1968).

Libinia erinacea (A. Milne Edwards, 1879) (Crust. Rég. Mex., p. 202)

Rathbun, 1925, p. 321, pl. 109.

Range: Florida Keys; southeast to northwest Florida; north coast of Cuba.

Depth: 4 to 68 m (2 to 37 fm).

Habitat: sand-mud and rock bottoms; in patches of moss.

Remarks: The status of this species is confused due to a paucity of mature specimens. Rathbun (1925) remarked on the close resemblance of this species to *L. dubia*, but she also listed differences between the two forms. Tabb and Manning (1961) also questioned the status of this crab as a distinct species and Abele (1970) compared the types of the two species and concluded that further examination of more mature specimens will be required. Abele (1970) also noted that the type specimen of *L. erinacea* is in the Museum of Comparative Zoology (Harvard) and not in the Paris Museum, as indicated by Rathbun (1925). Listed as uncommon at Apalachee Bay, Florida by Menzel (1971).

Libinia rhomboidea Streets, 1870 (Proc. Acad. Nat. Sci. Philadelphia, p. 106)

Rathbun, 1925, p. 323, pls. 116-117, pl. 246, figs. 1-3.

Range: west and north coasts of Cuba; off Merida, Yucatan (Gulf coast).

Depth and Habitat: no data available.

Remarks: Rathbun (1925) believed that the type locality of "East Indies" was probably an error. She also compared this species with *L. dubia* and noted variation in size of the spines.

Nibilia A. Milne Edwards, 1878

Nibilia antilocapra (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 110)

Rathbun, 1925, p. 290, text-fig. 97, pls. 102-103, 239; Williams, 1965, p. 251, figs. 230, 233F.

Range: North Carolina; southeast Florida; off Mobile, Alabama; Gulf of Campeche, Mexico; St. Vincent; Barbados.

Depth: 71 to 256 m (39 to 140 fm).

Habitat: mud, sand, broken shell, rock, and coral bottoms.

Remarks: Ovigerous females were collected at St. Vincent in February and at Barbados in March (Rathbun, 1925). Recorded by Chace (1956) from the Gulf of Mexico, some distance west of Dry Tortugas.

Pelia Bell, 1836

Pelia mutica (Gibbes, 1850) (Proc. Amer. Assoc. Adv. Sci. 3: 171)

Hay & Shore, 1918, p. 455, pl. 38, fig. 7; Rathbun, 1925, p. 278, text-fig. 94, pl. 98, figs. 2-3; Rathbun, 1933, p. 18, fig. 19; Williams, 1965, p. 250, figs. 229, 233E; Felder, 1973a, p. 53, pl. 7, fig. 13.

Range: Massachusetts to south Florida; Florida Keys; west coast of Florida to

south Texas; north coast of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands.

Depth: shallow water to 51 m (to 28 fm).

Habitat: shell or rock rubble; mud, sand, gravel, and coral substrates; among hydroids, ascidians, and sponges on wharf pilings; from *Chaetopterus* tubes.

Remarks: Ovigerous females have been collected from Florida in February to July, through the summer in the Carolinas, and from Massachusetts in July. Regional lists include Florida (Wass, 1955; Tabb and Manning, 1961; Menzel, 1971), Louisiana (Behre, 1950; Hoese and Valentine, 1972), and Texas (Felder, 1973a). Tabb and Manning (1961) reported that all of the crabs collected in Florida Bay were covered with algae, sponges, or hydroids; Wass (1955) noted that this crab is common in clumps of the ascidian, *Styela*. Gray (1961) reported the crab in tubes of the annelid, *Chaetopterus*, in North Carolina. Hartnoll (1965a) studied growth and other aspects of biology of this species in Jamaica.

Rochinia A. Milne Edwards, 1875

Rochinia crassa (A. Milne Edwards, 1879) (Crust. Rég. Mex., p. 203)

Rathbun, 1925, p. 210, text-figs. 83-84, pls. 68-69, 226; Chace, 1940, p. 62; Williams, McCloskey & Gray, 1968, p. 60; Pequegnat, 1970, p. 183.

Range: Massachusetts to south Florida; Florida Straits; off Alabama and Texas; east coast of Mexico; north coast of Cuba.

Depth: 128 to 860 m (70 to 470 fm).

Habitat: mud, sand, and coral oozes.

Remarks: Rathbun (1925) reported ovigerous females from off South Carolina in December. Musick and McEachran (1972) listed this crab from 194 m depth in Chesapeake Bight. Listed by Chace (1956) from the R/V *Oregon* collections in the Gulf of Mexico.

Rochinia hystrix (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 124)

Rathbun, 1925, p. 214, pls. 70-71; Rathbun, 1933, p. 17, fig. 17; Chace, 1940, p. 62.

Range: Florida Keys and Dry Tortugas; north coast of Cuba; off northwest Florida and Mississippi; Puerto Rico; Martinique to Barbados.

Depth: 150 to 708 m (82 to 387 fm).

Habitat: sand, coral, and rocky substrates.

Remarks: Rathbun (1925) listed ovigerous females from off Key West in February. Chace (1940) commented on sample numbers from off Cuba and Chace (1956) listed this species from the Gulf collections of the R/V *Oregon*.

Rochinia tanneri (Smith, 1883) (Proc. U.S. Nat. Mus. 6: 4)

Rathbun, 1925, p. 216, pl. 227, fig. 1; Williams, McCloskey & Gray, 1968, p. 60, fig. 15.

Range: Massachusetts to North Carolina; southeast Florida; off Key West and Florida Straits.

Depth: 128 to 708 m (70 to 387 fm).

Habitat: sand and shell bottoms.

Rochinia umbonata (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 115)

Rathbun, 1925, p. 222, text-fig. 85, pl. 72, pl. 73, fig. 1; Chace, 1940, p. 63; Williams, McCloskey & Gray, 1968, fig. 16; Pequegnat, 1970, p. 183.

Range: North Carolina to south Florida; Florida Straits, off Key West; off Mississippi; St. Vincent, Windward Islands.

Depth: 161 to 900 m (88 to 492 fm).

Habitat: sand, gravel, coral, shell, and foraminiferan substrates.

Remarks: Listed by Chace (1956) from the Gulf of Mexico.

Family PARTHENOPIDAE Macleay, 1838

Subfamily PARTHENOPINAE Macleay, 1838

Cryptopodia H. Milne Edwards, 1834***Cryptopodia concava*** Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 137)

Rathbun, 1925, p. 553, text-fig. 151, pl. 202, figs. 3-4, pl. 282, figs. 6-11; Rathbun, 1933, p. 42, fig. 37; Williams, McCloskey & Gray, 1968, p. 64.

Range: off North Carolina; Bahamas; Florida Keys and Dry Tortugas; west coast of Florida; St. Thomas, Virgin Islands; Ceará to Bahia, Brazil.

Depth: 7 to 62 m (4 to 34 fm).

Habitat: mud, sand, shell, and coral bottoms.

Remarks: Listed from Brazil by Coelho and Ramos (1972).

Heterocrypta Stimpson, 1871***Heterocrypta granulata*** (Gibbes, 1850) (Proc. Amer. Assoc. Adv. Sci. 3: 173)

Common Name: Pentagon Crab

Hay & Shore, 1918, p. 464, pl. 39, fig. 9; Rathbun, 1925, p. 555, text-fig. 152, pl. 203, figs. 1-2, pl. 282, figs. 1-3; Rathbun, 1933, p. 43, fig. 38; Williams, 1965, p. 270, figs. 251, 252E; Felder, 1973a, p. 45, pl. 6, fig. 6.

Range: Massachusetts to Georgia; Florida Keys and Straits; west coast of Florida to south Texas; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Ceará to Bahia, Brazil.

Depth: 4 to 137 m (2 to 75 fm).

Habitat: mud, sand, gravel, shell, rock, and coral bottoms; this crab is difficult to detect on pebble and shell substrates, where its form and coloration provide excellent camouflage.

Remarks: Williams (1965) reports ovigerous females throughout the summer off North Carolina. Regional lists include Florida (Wass, 1955; Tabb and Manning, 1961; Dragovich and Kelly, 1964; Rouse, 1970; Menzel, 1971), Louisiana (Behre, 1950) and Texas (Hedgpeth, 1953; Parker, 1959; Leary, 1967). Listed from Brazil by Coelho and Ramos (1972).

Leiolambrus A. Milne Edwards, 1878

Leiolambrus nitidus Rathbun, 1901 (Bull. U.S. Fish. Comm. 20: 80)

Rathbun, 1925, p. 545, pl. 199, pl. 281, fig. 1; Rathbun, 1933, p. 41, fig. 35; Guinot-Dumortier, 1960, p. 182, figs. 23a-b, 26; Felder, 1973a, p. 45, pl. 6, fig. 7.

Range: Gulf of Mexico, from off Alabama to south Texas; Jamaica; Puerto Rico; French Guiana.

Depth: 7 to 73 m (4 to 40 fm).

Habitat: mud, mud-sand, and mud-shell bottoms.

Remarks: Regional lists include Alabama (Chace, 1956), Louisiana (Dawson, 1966), and Texas (Hildebrand, 1954; Leary, 1967). Hildebrand (1954) reported ovigerous females from off Texas in June. Listed from French Guiana by Guinot-Dumortier (1960).

Mesorhoa Stimpson, 1871

Mesorhoa sexspinosa Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 136)

Rathbun, 1925, p. 547, text-fig. 150, pl. 200; Rathbun, 1933, p. 42, fig. 36; Williams, McCloskey & Gray, 1968, p. 64, fig. 17.

Range: North Carolina; Florida Keys and Dry Tortugas; northwest coast of Florida; Puerto Rico; Virgin Islands.

Depth: 7 to 49 m (4 to 27 fm).

Habitat: sand and shell substrates.

Remarks: Rathbun (1925) reported an ovigerous female from off northwest Florida in January.

Parthenope Weber, 1795

Parthenope (Parthenope) agonus (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 131)

As *P. agona*—Hay & Shore, 1918, p. 462, pl. 39, fig. 5; Williams, 1965, p. 266, figs. 246, 252A; Pequegnat, 1970, p. 183.

As *P. agonus*—Rathbun, 1925, p. 513, text-fig. 146, pls. 178-179, pl. 275, figs. 1-3; Rathbun, 1933, p. 39.

Range: North Carolina; Florida Straits; Dry Tortugas; west coast of Florida; Puerto Rico; Trinidad; off Guianas.

Depth: 46 to 391 m (25 to 214 fm).

Habitat: sand, broken shell, gravel, coral, and mud bottoms.

Remarks: Ovigerous females were reported from northwest Florida in March (Rathbun, 1925) and from the Guianas in September (Williams, 1965). Chace (1956) lists this crab from the eastern Gulf of Mexico.

Subgenus *Platylambrus* Stimpson, 1871

Parthenope (Platylambrus) fraterculus (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 130)

Rathbun, 1925, p. 525, pls. 186-187, pl. 190, fig. 2; Williams, 1965, p. 269, figs. 249, 252D.

Range: North Carolina; southeast Florida; Florida Keys and Straits; Dry Tortugas; west and northwest coasts of Florida; northeast coast of Yucatan; Barbados; mouth of Amazon River, Brazil; Surinam.

Depth: 7 to 201 m (4 to 110 fm).

Habitat: sand, shell, gravel, rock, and coral bottoms.

Remarks: Williams (1965) listed ovigerous females from south Florida in May and from northwestern Florida in August. Listed by Chace (1956) from the northwestern Gulf of Mexico and by Holthuis (1959) from Surinam.

Parthenope (Platylambrus) pourtalesii (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 129)

Hay & Shore, 1918, p. 462, pl. 39, fig. 6; Rathbun, 1925, p. 521, pls. 182–183, 276; Rathbun, 1933, p. 39, fig. 33; Chace, 1940, p. 53; Williams, 1965, p. 268, figs. 248, 252C; Pequegnat, 1970, p. 183; Felder, 1973a, p. 48, pl. 6, fig. 9.

Range: New Jersey to south Florida; Florida Keys and Straits; Dry Tortugas; north coast of Yucatan; off north coast of Cuba; Grenada.

Depth: 18 to 348 m (10 to 190 fm).

Habitat: primarily mud and sand-mud bottoms; also on sand, shell, and gravel substrates. Pequegnat (1970) states that Gulf populations are probably centered at the middle of the continental shelf.

Remarks: Ovigerous females are known from North Carolina in December (Williams, 1965) and from the southeast Gulf in July (Pequegnat, 1970). Listed by Chace (1956) from off Yucatan. Leary (1967) includes this species on a Texas list and this is repeated by Felder (1973a), but I know of no actual records to confirm collection of this crab in the northwestern Gulf.

Parthenope (Platylambrus) serrata (H. Milne Edwards, 1834) (Hist. Nat. Crust., vol. 1, p. 357)

As *Platylambrus serratus*—Hay & Shore, 1918, p. 463, pl. 39, fig. 7.

As *Parthenope serrata*—Rathbun, 1925, p. 516, pls. 180–181, pl. 275, figs. 7–10; Rathbun, 1933, p. 39; Williams, 1965, p. 267, figs. 247, 252B; Türkay, 1968, p. 251; Felder, 1973a, p. 45, pl. 6, fig. 8.

Range: North Carolina; Bermuda; Bahamas; southeast and south Florida; Florida Keys and Dry Tortugas; west Florida to off Texas; Bay of Campeche, off Yucatan; north coast of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Curaçao; Surinam; Bahia, Brazil.

Depth: shallow water to 110 m (to 60 fm).

Habitat: mainly on mud and mud-sand bottoms; also from sand, shell, gravel and coral substrates.

Remarks: Gore (1977) reviews this species and concludes that two distinct species have been confused under this name. Ovigerous females are listed from North Carolina in June, from Florida in summer, from Cuba in October, and from Surinam in May to June (Williams, 1965). Hildebrand (1955) reported this crab present on the pink shrimp grounds at Campeche. Wass (1955) and Menzel (1971) indicated that this crab was rare off northwestern Florida; Chace (1956) listed a single record from off Mississippi for the collections of the R/V *Oregon*. Listed from Texas by Leary (1967), but Felder (1973a) cites only a

single collection record from the northwestern Gulf, a specimen taken from a snapper stomach off south Texas. Listed from Surinam by Holthuis (1959).

***Solenolambrus* Stimpson, 1871**

***Solenolambrus decemspinus* Rathbun, 1894** (Proc. U.S. Nat. Mus. 17: 84)

Rathbun, 1925, p. 540, pl. 194, figs. 1–2; Rathbun, 1933, p. 41.

Range: off northwest Florida; Puerto Rico.

Depth: 82 to 110 m (45 to 60 fm).

Habitat: sand and sand-mud bottoms.

***Solenolambrus tenellus* Stimpson, 1871** (Bull. Mus. Comp. Zool. 2: 134)

Hay & Shore, 1918, p. 463, pl. 39, fig. 8; Rathbun, 1925, p. 541, pl. 194, figs. 3–4, pl. 279, figs. 5–9; Rathbun, 1933, p. 41; Williams, 1965, p. 270, fig. 250.

Range: North Carolina; southeast Florida; Bahamas; Florida Keys; west coast of Florida; Barbados.

Depth: 55 to 210 m (30 to 115 fm).

Habitat: sand, broken coral, and rocky bottoms.

Remarks: Ovigerous females are known from Florida in May-June and from Barbados in May (Rathbun, 1925).

***Solenolambrus typicus* Stimpson, 1871** (Bull. Mus. Comp. Zool. 2: 133)

Rathbun, 1925, p. 537, text-fig. 148, pls. 192–193, pl. 279, figs. 1–4; Rathbun, 1933, p. 40, fig. 34; Chace, 1940, p. 53; Williams, McCloskey & Gray, 1968, p. 63; Pequegnat, 1970, p. 184.

Range: North Carolina; Bahamas; southeast Florida; Florida Keys and Straits; Dry Tortugas; off Texas; north and south coasts of Cuba.

Depth: 91 to 618 m (50 to 338 fm).

Habitat: sand, broken shell, and coral substrates.

Remarks: Location of the R/V *Alaminos* record is confused. Table 6–9 of Pequegnat (1970, p. 201) lists this species from the southwestern Gulf and a synopsis of distribution (p. 184) includes a range from the Bahamas to northwest Florida, but the actual collecting station of record (64-A-10-13) is located off south Texas.

***Thyrolambrus* Rathbun, 1894**

***Thyrolambrus astroides* Rathbun, 1894** (Proc. U.S. Nat. Mus. 17: 83)

Rathbun, 1925, p. 532, text-fig. 147, pl. 196, pl. 280, figs. 5–6.

Range: off north coast of Cuba; Mauritius and Andaman Sea, in the Indo-Pacific region.

Depth: 123 to 366 m (67 to 200 fm) off Cuba; 66 m (36 fm) in Andaman Sea.

Habitat: coral bottoms.

Tutankhamen Rathbun, 1925***Tutankhamen cristatipes*** (A. Milne Edwards, 1880) (Crust. Rég. Mex., p. 352)

Rathbun, 1925, p. 530, pl. 277, figs. 3-5.

Range: Pourtales Plateau, Florida Straits; St. Vincent, Windward Islands.

Depth: 227 to 366 m (124 to 200 fm).

SECTION CANCRIDEA Latreille, 1803

Superfamily CANCROIDEA Latreille, 1803

Family ATELECYCLIDAE Ortmann, 1893

Subfamily ATELECYCLINAE Ortmann, 1893

Trachycarcinus Faxon, 1893***Trachycarcinus spinulifer*** Rathbun, 1898.Determined by Pequegnat (1970) to be a synonym of *Trichopeltarion nobile* A. Milne Edwards, to which all previous records are referred.**Trichopeltarion** A. Milne Edwards, 1880***Trichopeltarion nobile*** A. Milne Edwards, 1880 (Bull. Mus. Comp. Zool. 8: 20)As *Trachycarcinus spinulifer*—Rathbun, 1930, p. 166, text-figs. 26-27, pls. 70-71.As *Trichopeltarion nobile*—Rathbun, 1930, p. 168, pl. 73; Pequegnat, 1970, p. 184, figs. 6-4, 6-5.

Range: off northwest Florida and Mississippi; east coast of Mexico; Bay of Campeche; off St. Lucia.

Depth: 274 to 752 m (150 to 411 fm).

Habitat: mud bottoms.

Remarks: Pequegnat (1970) synonymized the two species on the basis of fresh material collected by the R/V *Alaminos* in the Gulf of Mexico. The largest of 9 females collected was ovigerous, from 195 fm depth, in September. Listed by Chace (1956), as *T. spinulifer*, from off Mississippi.

Family CANCRIDAE Latreille, 1803.

Subfamily CANCRINAE Latreille, 1803

Cancer Linnaeus, 1758***Cancer borealis*** Stimpson, 1859 (Ann. Lyc. Nat. Hist. New York 7: 50)

Common Names: Jonah Crab; Northern Crab

Hay & Shore, 1918, p. 434, pl. 35, fig. 2; Rathbun, 1930, p. 182, text-figs. 30-31;

Chace, 1940, p. 38; Williams, 1965, p. 175, fig. 156.

Range: Nova Scotia to east coast of Florida; Bermuda; south coast of Florida; south of Dry Tortugas, in Florida Straits.

Depth: intertidal to 796 m (to 435 fm).

Habitat: immature and sub-adult stages occur intertidally among rocks and in shallower harbors and bays; larger specimens are found over a wide range of depths on sand, shell, gravel, mud, and ooze substrates.

Remarks: Ovigerous females were listed from Florida in June (Williams, 1965). This species has been collected in the western portions of the Florida Straits and can be considered to inhabit the Gulf of Mexico. *Cancer irroratus* Say, 1817 has been collected in the Florida Straits slightly west of 80° W, but is more typically restricted to the Atlantic coast and is not included in the Gulf fauna at the present time (based on pers. comm. of F. A. Chace).

SECTION BRACHYRHYNCHA Borradaile, 1907

(This group of crabs includes over 3000 species, world-wide. Older systematic literature traditionally divided the Brachyryncha into two subgroups, based on the general shape of the carapace: the Cyclometopa (round-fronted) or "cancroid" crabs and the Catametopa (square-fronted) or "grapsoid" crabs. The first group, as treated by Rathbun (1930), consisted of the families Portunidae, Xanthidae, Potamidae, Atelecyclidae, and Cancridae. These last two families are treated by some authors and in the present list within a separate section, the Cancridea, following the format of Glaessner (1969). Sakai (1965) groups the Goneplacidae with the other cancroid families in his key to the Brachyryncha, whereas Rathbun (1918) treated the Goneplacidae as a grapsoid family. This family has long been recognized as containing genera that link certain genera of the Xanthidae and Grapsidae, thus blurring many of the sharp distinctions between grapsoid and cancroid characteristics. The position of the Palicidae is treated as "uncertain" by Glaessner (1969), but it is included here with the Xanthoidea as a matter of convenience, suggested by Chace (pers. comm.).)

Superfamily PORTUNOIDEA Rafinesque, 1815.

Family PORTUNIDAE Rafinesque, 1815

Subfamily POLYBIINAE Ortmann, 1893

(*Ovalipes* is listed as a genus of the Macropipinae in Glaessner (1969), but an addendum, p. R627, indicates the synonymy with Polybiinae.)

Bathynectes Stimpson, 1871

Bathynectes superba (Costa, 1853) (Fauna Regno Napoli, Addiz. Decapodi Brachyuri, p. 19)

Milne Edwards & Bouvier, 1923, p. 311; Rathbun, 1930, p. 28, pls. 9-10; Chace, 1940, p. 30; Monod, 1956, p. 183, figs. 210-212; Williams, McCloskey & Gray, 1968, p. 50; Christiansen, 1969, p. 70, fig. 28, map 22.

Range: Massachusetts to south Florida; Florida Keys and Straits; west of Dry Tortugas and mid-eastern Gulf of Mexico; off Alabama and Mississippi; north-east of Yucatan; north and south coasts of Cuba; in the eastern Atlantic—Norway; north of Scotland; west coast of France; Cape Verde Islands; Mediterranean and Black Seas.

Depth: 100 to 1455 m (55 to 769 fm).

Habitat: on sand, gravel, shell and coral bottoms; rocky areas.

Remarks: Most of the Gulf of Mexico records are from Chace (1956). Musick and McEachran (1972) collected this crab at depths of 159 to 274 m in Chesapeake Bight. Roberts (1969) studied larval development and epizoites of this species were described by Lewis (1976).

Benthochascon Alcock & Anderson, 1899

Benthochascon schmitti Rathbun, 1931 (J. Washington Acad. Sci. 21: 125)

Rathbun, 1931a, p. 125, pls. 1-2; Pequegnat, 1970, p. 187, fig. 6-6.

Range: off Dry Tortugas; deep waters off Mississippi to Texas; off southern Gulf coast of Mexico; recently found off New England.

Depth: 201 to 510 m (110 to 279 fm).

Habitat: mud, silt-clay substrates.

Remarks: This species was first reported from the Dry Tortugas by Rathbun (1931a), based on a specimen collected by W. L. Schmitt. Schmitt (1931) provided some note on this crab. Reported by Chace (1956) from the Gulf collections of the R/V *Oregon*. Pequegnat (1970) collected numerous specimens from various parts of the Gulf of Mexico and Wigley and Messersmith (1976) collected a single male from 252 m depth off southern New England.

Ovalipes Rathbun, 1898

(This genus has undergone a number of recent revisions, resulting in considerable confusion to non-specialists and complicating the comparison of field records in the non-systematic literature. Prior to revision by Williams (1962), *Ovalipes ocellatus* was considered to be represented in the Atlantic by the typical form and in the Gulf of Mexico by a subspecies, *O. o. guadulpensis*. Many of the earlier surveys and checklists reported *O. ocellatus* from the Gulf, but without specifying subspecies or authority. Felder (1973a) listed both forms for the northwestern Gulf, but with reservations, noting the nomenclatural confusion in the records. Türkay (1971) had discovered, in the meantime, that the original type of Saussure's *guadulpensis* belonged to the genus *Macropipus* and not *Ovalipes* and that it was named for a locality in the Azores rather than in the Caribbean. The next available name for the western Atlantic form thus became *Ovalipes ocellatus floridanus*, as used by Hay and Shore (1918) and recognized by Türkay (1971). Two forms of this crab were recognized by Williams (1962, 1965) and by Stephenson and Rees (1968), *Form a* from the Carolinian Province off the southeastern United States and *Form b* from the Gulf of Mexico. Williams (1976) finally

presented evidence that the two forms of *O. o. floridanus* should be regarded as separate species, but only one, *O. floridanus*, inhabits the Gulf of Mexico. See Williams (1976) for a nomenclatural history of these species.)

Ovalipes floridanus Hay & Shore, 1918 (Bull. U.S. Bur. Fish. 35: 427)

Common Names: Lady Crab; Sand Crab

As *O. ocellatus floridanus*—Hay & Shore, 1918, p. 427, pl. 32, fig. 8; Türkay, 1971, p. 139, fig. 3.

As *O. ocellatus gadulpenis*—Rathbun, 1930, p. 23, pl. 4 (part, the Pensacola, Alabama, and Gulf of Mexico specimens).

As *O. gadulpenis (Form b)*—Stephenson & Rees, 1968, p. 243, pls. 37D, 40F, 41E, 42K, fig. 1K.

As *O. gadulpenis*—Williams, 1965, p. 161 (part, Gulf specimens only); Felder, 1973a, p. 54, pl. 8, fig. 2.

As *O. floridanus*—Williams, 1976, p. 206, fig. 1a–d.

Range: southwest Florida to south Texas.

Depth: near surface and shallow water to 31 m (to 17 fm).

Habitat: mainly sandy substrates; also on coral and broken shell bottoms. In coastal lagoons, channels, and bays in southern and central Texas.

Remarks: Williams (1976) describes and compares the two species of *Ovalipes* previously considered to be *O. gadulpenis*. The Carolinian form was named *O. stephensoni*, Form a of Stephenson and Rees (1968). The type locality for *O. floridanus* is Pensacola, Florida. Williams (1976) lists ovigerous females in February from Florida. Abele (1970) noted that juveniles were common on shallow grass flats in northwest Florida, while adults were more often found offshore on sandy bottoms. Regional lists include, as *O. gadulpenis* or *O. ocellatus gadulpenis*, Florida (Wass, 1955, Abele, 1970; Menzel, 1971), Mississippi (Richmond, 1962; Franks *et al.*, 1972; Christmas and Langley, 1973), Louisiana (Behre, 1950; Hildebrand, 1954), Texas (Gunter, 1950; Leary, 1967), and the northwestern Gulf (Fotheringham and Brunenmeister, 1975). Caine (1974) provided a detailed description of feeding behavior, ecology, and burrowing activities. Abele (1970) commented on behavior of the animal in sand substrates. Notes on the related species are provided in Williams (1965).

Ovalipes gadulpenis and ***Ovalipes ocellatus gadulpenis*** (Saussure, 1858).

Both names are invalid and all Gulf of Mexico reports should be referred to *Ovalipes floridanus* Hay and Shore, 1918. See Williams (1976) for a review of the nomenclatural history of this species.

Subfamily PORTUNINAE Rafinesque, 1815

Arenaeus Dana, 1851

Arenaeus cribrarius (Lamarck, 1818) (His. Nat. Anim. sans Vert., vol 5, p. 259)

Common Name: Speckled Crab

Hay & Shore, 1918, p. 434, pl. 34, fig. 3; Rathbun, 1930, p. 134, pl. 58, figs. 2–3,

pls. 59-60; Rathbun, 1933, p. 50; Williams, 1965, p. 173, fig. 153; Felder, 1973a, p. 55, pl. 8, fig. 4.

Range: Massachusetts to North Carolina; Bermuda; east coast of Florida; Florida Keys and Dry Tortugas; west coast of Florida to Tabasco, Mexico; Jamaica; Puerto Rico; Dominica to St. Lucia; Belize to Colombia; Curaçao; Ceará to Santa Catarina, Brazil.

Depth: shore to 68 m (to 37 fm).

Habitat: in surf zone of sandy beaches; an adroit swimmer, it is seldom washed ashore and it is rarely found in estuaries and back lagoons.

Remarks: Williams (1965) lists ovigerous females from Florida in August. Regional lists include Florida (Wass, 1955; Abele, 1970; Menzel, 1971), Mississippi (Richmond, 1962), off the Mississippi Delta (Chace, 1956), Louisiana (Behre, 1950; Hoese and Valentine, 1972), Texas (Gunter, 1950, Hildebrand, 1954), and the northwestern Gulf of Mexico (Fotheringham and Brunenmeister, 1975). Listed from Brazil by Coelho and Ramos (1972).

Callinectes Stimpson, 1860

(This commercially important genus has been revised and reviewed by Williams (1974a), including details of reproductive morphology, notes on larval development, fossil records, a discussion of zoogeographic affinities, and keys for the identification of the 14 species known worldwide. In the Gulf region, as well as elsewhere, confusion has existed as to the identity of the lesser or smaller blue crab species, cited locally as *Callinectes danae* and *C. ornatus*. A prior revision by Williams (1966) established a new species, *C. similis*, for the Gulf populations and many of the east coast specimens, limiting the former two species to the southeastern Gulf and beyond. The present list includes eight species in the Gulf of Mexico, five of which occur only in the eastern or southern margins of the region. Only *C. rathbunae* is endemic to the Gulf; *C. sapidus* and *C. similis* range widely along the U.S. Gulf and Atlantic coasts and beyond (see Figures 24-27 in Williams, 1974a).)

Callinectes bocourti A. Milne Edwards, 1879 (Crust. Rég. Mex., p. 226)

Rathbun, 1930, p. 128, text-figs. 15g, 16e, 17h, 18f, pl. 55; Rathbun, 1933, p. 49; Holthuis, 1959, p. 201, text-fig. 47, pl. 5, fig. 2; Chace & Hobbs, 1969, p. 127, text-figs. 35, 37a; Williams, 1974a, p. 767, figs. 12, 18j, 20m, 22j, 27.

Range: southeast Florida; Mississippi (rare); Jamaica; Hispaniola; Puerto Rico; Dominica; Virgin Islands to Barbados; Trinidad; Netherlands Antilles; Belize to Panama; Colombia to Santa Catarina, Brazil.

Depth: shallow water, near shore and inshore waters.

Habitat: in shallow, brackish waters of estuaries and river mouths. Tolerates salinities as low as 5 ppt; females usually move to saltier waters after mating. Williams (1974a) notes that this species is often associated with *C. sapidus*, but appears to be more tolerant than the latter to stagnant and polluted waters. Collected from a mud bottom in 3 m of water in Biloxi Bay, Mississippi (Perry, 1973). See Chace and Hobbs (1969) for extensive habitat notes in Dominica.

Remarks: The first reports of this species in North America were from Florida (Provenzano, 1961), from Mississippi, the only Gulf records (Perry, 1973), and a second Florida finding (Gore and Grizzle, 1974). Norse (1972) noted habitat preferences in Jamaica. Listed from Brazil by Coelho (1971a) and Coelho and Ramos (1972).

Callinectes danae Smith, 1869 (Trans. Connecticut Acad. Sci. 2: 7)

Rathbun, 1930, p. 118 (part)), text-figs. 15d, 16d, 17b, 18d, pl. 51; Rathbun, 1933, p. 49; Chace, 1940, p. 33; Chace & Hobbs, 1969, p. 130, fig. 37b; Holthuis, 1959, p. 201; Williams, 1966, p. 86, figs. 2A-D, 4C-D; Jones, 1968, p. 187; Williams, 1974a, p. 746, figs. 7, 18e, 20e-f, 22e, 24.

Range: Bermuda; southeast Florida; Dry Tortugas; north coast of Cuba; Caribbean coast of Yucatan; Jamaica; Hispaniola; Puerto Rico; St. Croix to Barbados; Trinidad; Netherlands Antilles; Belize to Panama; Colombia to Santa Catarina, Brazil.

Depth: shore to 75 m (to 41 fm).

Habitat: from nearly fresh to full sea water, possibly in hypersaline lagoons; Occurs among mangroves and in muddy estuaries in Brazil; Off beaches and in open water.

Remarks: Williams (1974a) believes that earlier records of this species from Chile are erroneous. Records of this species along the U.S. Gulf coast should be referred to *C. similis*, as per the revision by Williams (1966, 1974a). Records from east Florida indicated this crab's presence on the ocean side of islands in Biscayne Bay and its absence from the Florida Keys (Park, 1969). Norse (1972) noted habitat preferences in Jamaica. Listed from Brazil by Coelho and Ramos (1972). Morrison and Morrison (1952) studied hemolymph coagulation in Bermuda specimens.

Callinectes exasperatus (Gerstaecker 1856) (Arch. f. Naturg. 22: 129)

Rathbun, 1930, p. 130, text-figs. 15f, 16f, 17e, 18e, pl. 56; Contreras, 1930, p. 236, fig. 7; Rathbun, 1933, p. 49; Chace, 1940, p. 44; Chace & Hobbs, 1969, p. 131, fig. 37c; Felder, 1973a, p. 58, pl. 8, fig. 8; Williams, 1974a, p. 757, figs. 9, 18g, 20i, 22g, 26.

Range: Bermuda; Bahamas; southeast Florida; Florida Keys and Dry Tortugas; Vera Cruz to Yucatan, Mexico; north and south coasts of Cuba; Jamaica; Hispaniola; Puerto Rico; St. Croix to Grenadines; Trinidad; Netherlands Antilles; Caribbean coast of Yucatan to Panama; Venezuela and Isla de Margarita; Rio Grande do Norte to Santa Catarina, Brazil.

Depth: shore to 7.5 m (to 4 fm).

Habitat: estuarine and marine waters, possibly fresh water; around river mouths and in mangrove areas.

Remarks: Williams (1974a) comments on a lack of specimens from the Guianas and northern Brazil and that a locality record from Chile is erroneous. Leary (1967) listed this species from Texas on the basis of a single specimen from near Port Aransas, tentatively identified by Gordon Gunter, but not available for confirmation by Williams. Leary (1967) stated that the frontal teeth pattern was very similar to *C. danae* (= *C. similis*), thus the Gulf record for this

species is questionable. Felder (1973a) includes this species in his key for the northwestern Gulf on the basis of the same specimen. Reports of this crab in southern Florida are provided by Futch (1965) and Park (1969). Norse (1972) noted habitat preferences in Jamaica. Listed from Brazil by Coelho and Ramos (1972).

Callinectes marginatus (A. Milne Edwards, 1861) (Arch. Mus. Hist. Nat., Paris 10: 318)

Rathbun, 1930, p. 123, figs. 15e, 16d, 17d, 18c, pl. 53; Contreras, 1930, p. 235, fig. 6; Rathbun, 1933, p. 49; Chace & Hobbs, 1969, p. 131, fig. 37d; Felder, 1973a, p. 59, pl. 8, fig. 9; Williams, 1974a, p. 722, figs. 3, 18b (not 18a), 20a, 22b, 27.

Range: North Carolina; Bermuda; Bahamas; southeast Florida; Florida Keys and Dry Tortugas; southwest Florida; Louisiana (rare); Vera Cruz to Yucatan, Mexico; north and south coasts of Cuba; Jamaica; Hispaniola; Puerto Rico; St. Croix to Grenadines; Tobago and Trinidad; Netherlands Antilles; Yucatan to Panama; Colombia to São Paulo, Brazil; in eastern Atlantic—Senegal to Angola; Mauritania; Cape Verde Islands.

Depth: intertidal to 15 m, rarely to 25 m (to 14 fm).

Habitat: shallow water on sand and mud flats; edges of mangrove swamps; grass flats, oyster bars, and rocky pools; often in brackish water, rarely in open marine water.

Remarks: The records from North Carolina and Louisiana are extralimital occurrences. Behre (1950) listed this crab as a common component of the *Sargassum* community in Louisiana waters, but Felder (1973a) commented that this species is by no means common and that Behre probably confused it with *Portunus sayi*. Williams (1974a) reports spawning records throughout the range of this crab, from December to July. Norse (1972) noted habitat preferences in Jamaica. Listed from Brazil by Coelho and Ramos (1972) and from West Africa by Forest and Guinot (1966).

Callinectes ornatus Ordway, 1863 (J. Boston Soc. Nat. Hist. 7: 571)

Rathbun, 1930, p. 114 (part), text-figs. 15b, 16a, 17a, 18b, pl. 50; Contreras, 1930, p. 232 (part), fig. 4; Rathbun, 1933, p. 48, fig. 40; Chace, 1940, p. 33; Chace & Hobbs, 1969, p. 132, fig. 37e; Holthuis, 1959, p. 200; Williams, 1965, p. 172 (part); Williams, 1966, p. 84, figs. 1A–B, 4A–B; Williams, 1974a, p. 739, figs. 6, 18d, 20d, 22d, 25.

Range: Bermuda; Bahamas; North Carolina to south Florida; Florida Keys and Dry Tortugas; west Florida to Tampa Bay; Gulf coast of Yucatan; north and south coasts of Cuba; Jamaica; Hispaniola; Puerto Rico; St. Croix to Barbados; Trinidad; Netherlands Antilles; Caribbean coast of Yucatan to Belize; Colombia to Santa Catarina, Brazil.

Depth: shore to 75 m (41 fm).

Habitat: sand, mud, shell substrates; off sponges; near river mouths and bays; in fresh waters, but more common in waters of moderate salinities (Williams, 1974a).

Remarks: As with many portunids, juveniles of this species may be difficult to distinguish from others, especially those of *C. danae* and *C. similis*. Records

of *C. ornatus* in New Jersey, Louisiana, and Texas (Rathbun, 1930; Leary, 1967) should be referred to *C. similis*, based on the revision and restriction of this species by Williams (1966). Brues (1927) discussed the ecology of this crab and Norse (1972) noted habitat preferences for the species in Jamaica. Listed from Brazil by Coelho (1971a) and Coelho and Ramos (1972).

Callinectes rathbunae Contreras, 1930 (An. Inst. Biol. Univ. Nac. Auton., Mex. 1: 238)

Contreras, 1930, p. 238, figs. 9–10; Felder, 1973a, p. 58, pl. 8, fig. 10; Williams, 1974a, p. 772, figs. 13, 19a, 20n, 22k, 27.

Range: eastern coast of Mexico, from mouth of Rio Grande to southern Vera Cruz.

Depth: very shallow waters.

Habitat: estuarine waters of ditches, lagoons, and river mouths; in shallow coastal bays and a broad range of salinities.

Remarks: This species is apparently localized to the central Mexican coast, with occasional specimens as far north as the Rio Grande. Listed by Leary (1967) for Texas, based on a single male specimen found at the mouth of the Rio Grande River (H. H. Hildebrand, collector).

Callinectes sapidus Rathbun, 1896 (Proc. U.S. Nat. Mus. 18: 352)

Common Names: Blue Crab; Common Edible Crab

Hay & Shore, 1918, p. 432, pl. 35, fig. 1; Rathbun, 1930, p. 99, text-figs. 15a, 16c, 17c, 18a, 19, pl. 47; Holthuis, 1961, p. 50, pl. 1, fig. 2, pl. 2, fig. 2; Williams, 1965, p. 168, fig. 151; Chace & Hobbs, 1969, p. 133, figs. 36, 37f; Christiansen, 1969, p. 72, fig. 29; Felder, 1973a, p. 55, pl. 8, fig. 7; Williams, 1974a, p. 778, figs. 1, 16, 17, 19d, 21, 23b–c, 26.

As *C. sapidus acutidens*—Rathbun, 1930, p. 111, text-fig. 15c, pl. 48; Contreras, 1930, p. 228, fig. 1; Rathbun, 1933, p. 48.

Range: Bermuda; Bahamas; Nova Scotia to south Florida; Florida Keys and Dry Tortugas; south Florida to Yucatan, along entire Gulf coast of the United States and Mexico; north and south coasts of Cuba; Jamaica; Haiti; Puerto Rico; Virgin Islands; Dominica; Trinidad and Tobago; Netherlands Antilles; Yucatan to Guatamala; Nicaragua to Panama; Venezuela; central Brazil to northern Argentina; in the eastern Atlantic—Denmark; Netherlands and adjacent North Sea; southwest France; Golfo di Genova; northern Adriatic Sea; Aegean Sea; western part of Black Sea; eastern Mediterranean Sea; Japan (Sakai, 1976a). Williams (1974a) reviews the distribution of this species and cites pertinent literature on the ship transport of crab larvae and their introduction into the Old World during recent times.

Depth: shore (intertidal) to 90 m (49 fm), more commonly to 35 m (19 fm).

Habitat: occurs in a wide range of salinities, from freshwater to hypersaline; along coasts in shallow water off ocean beaches (mainly females), in bays, estuaries, lagoons, ponds, ditches; well upstream in larger rivers (mainly males). Copeland and Bechtel (1974) listed an optimum salinity range of 0 to 27 ppt and optimum temperature range of 10 to 35°C, but extremes of 117 ppt (Hildebrand, 1957) and 45°C. (personal observations) under natural conditions are

tolerated for short periods. Mahood *et al.*, (1970) showed that temperature and salinity tolerances were interdependent. Blue crabs are also able to tolerate low oxygen conditions and they are occasionally found in very polluted and anoxic waters. Along the Texas coast, blue crabs in shallow ponds on sand flats are exposed to temperatures in excess of 40°C. during midday. These crabs have been observed to leave the water for *Salicornia* patches around the pools, where they rely on aerial respiration for 2 to 4 hours in the cooler (30 to 35°C.), humidity-saturated environment (personal observations). Blue crabs can travel some distance overland at night and during wet periods, again relying on aerial respiration.

Remarks: The vast literature, much of it non-technical or of a commercial nature, precludes anything approaching a complete bibliography on this species. The original type for this species was obtained from the eastern coast of the United States, a variant of the form that is more typical throughout its range. The "typical" form, most often encountered from Florida southward, was considered a subspecies by Rathbun (1930) and many others, *C. sapidus acutidens*, so-named because of the surface features and pronounced spines, teeth, and prominent ridges. Because the type-based form from farther north was the basis for comparison, confusion existed for some years over the designation of a type to replace the original, which had been lost. Williams (1974a) discusses these variations and the nomenclatural history of this crab, agreeing with Chace and Hobbs (1969) that a variety of extreme forms exist and that they could be considered separate species if they were considered in isolation from each other. However, these forms are intergraded and form a continuum, without morphological, bathymetrical, or geographical discontinuity, thus all the forms of *Callinectes sapidus* are considered, at present, to represent a single species in the process of local speciation which is still morphologically incomplete.

Recent literature compilations on this species were provided by Cronin *et al.* (1957) and by Tagatz and Hall (1971). Gulf regional lists include Florida (Wass, 1955; Tabb and Manning, 1961; Dragovich and Kelly, 1964; Rouse, 1970; Menzel, 1971; Lyons *et al.*, 1971), Mississippi (Richmond, 1962; Franks *et al.*, 1972; Christmas and Langley, 1973), Louisiana (Behre, 1950; Darnell, 1959; Hoese and Valentine, 1972), Texas (Gunter, 1950; Hedgpeth, 1950; Hoese, 1960; Copeland, 1965; Leary, 1967; More, 1969; Copeland and Bechtel, 1974), Mexico (Contreras, 1930; Hildebrand, 1957), and north coast of Cuba (Chace, 1940).

Hay (1905) and Churchill (1919) provided comprehensive life history studies. Other information on ecology includes: habitat relationships in Texas (Hedgpeth, 1953; Simmons, 1957; Breuer, 1962; Fotheringham and Brunenmeister, 1975; Trent, Pullen and Proctor, 1976), megalops ecology in Maryland (Cargo, 1960), habitat notes in Mississippi (Franks *et al.*, 1972; Christmas and Langley, 1973), effects of environmental variables on juveniles (Holland, Aldrich and Strawn, 1971), larval ecology in Virginia (Sandifer, 1973), seasonal population changes in Chesapeake Bight (Musick and McEachran, 1972), field observations of freshwater populations (Gunter, 1938), temperature and thermal tolerance (Tagatz, 1969a), and habitats in Jamaica (Norse, 1972).

Behavioral studies include: agonistic displays (Jachowski, 1974), larval shadow responses (Forward, 1977), analyses of swimming behavior (Spirito, 1972), predation on oyster spats (Lunz, 1947), quahogs (Carriker, 1951; Haven and Andrews, 1957), and gastropods (Hamilton, 1976), predator avoidance (Gunter, 1954), sex recognition (Teytaud, 1971), and climbing behavior on vegetation (Abbott, 1967).

Developmental studies include: larval rearing in the laboratory (Costlow and Bookhout, 1959, 1960a; Rust and Carlson, 1960; Davis, 1965), feeding of larvae (Sulkin and Epifanio, 1975), variability in morphology of larvae (Tyler and Cargo, 1963; Costlow, 1965), hormonal control of development (Costlow, 1963), zoeal growth and survival (Sandoz and Rogers, 1944), effects of pesticides on larval development (Bookhout and Costlow, 1976), salinity-temperature effects on larvae (Costlow, 1967; Rosenberg and Costlow, 1976), growth of juveniles (Gray and Newcombe, 1938; Tagatz, 1969b; Holland, Aldrich and Strawn, 1971), and development of cheliped laterality (Hamilton, Nishimoto and Halusky, 1976).

Physiological studies include: effects of salinity on growth at terminal molt (Haefner and Shuster, 1964), ionic and osmotic regulation (Gifford, 1962a; Tan and van Engel, 1966; Mantel, 1967; Copeland and Fitzjarrell, 1968; Ballard and Abbott, 1969; Tagatz, 1971; Gerard and Gilles, 1972; Mangum and Amende, 1972; Lynch *et al.*, 1973; Towles *et al.*, 1976; Mangum *et al.*, 1976; Mangum and Towle, 1977), respiration and respiratory pigments (Ayres, 1938; Gray, 1957; Horn and Kerr, 1963, 1969; Bonaventura *et al.*, 1974; Engel and Eggert, 1974; Mangum and Weiland, 1975; Weiland and Mangum, 1975; Lewis and Haefner, 1976; Laird and Haefner, 1976), hemolymph volume (Gleeson and Zubkoff, 1977), responses of megalops to pressure (Naylor and Isaac, 1973), gross anatomy and fine structure (Cochran, 1935; Cronin, 1947; Pyle and Cronin, 1950; Jahromi and Govind, 1976), ultrastructure of sperm (G. C. Brown, 1966), partial albinism (Sims and Joyce, 1966), neuroendocrinology (Costlow, 1963; Payen *et al.*, 1971; Andrews, Copeland and Fingerman, 1971; Skinner and Graham, 1972; Andrews, 1973; Ludolph, Paganalli and Mote, 1973), biochemical adaptations (Vernberg and Vernberg, 1968; Robert and Gray, 1972), neurobiology (Mendelson, 1963, 1966; Skobe and Nunnemacher, 1970; Hazlett, 1971; White and Spirito, 1973; Maynard and Dando, 1974; Reingold, 1975; Steinacker, 1975), fluoride metabolism (Moore, 1971), effects of DDT (Sheridan, 1975), homolymph coagulation (Morrison and Morrison, 1952), and temperature effects on growth and metabolism (Tagatz, 1969a; Leffler, 1972).

Studies on parasitology and pathology include: parasitic barnacles (Humes, 1941a; Reinhard, 1950a, 1950b; Adkins, 1972a), external barnacles (Williams and Porter, 1964), effects of nemertean on reproduction (Hopkins, 1947), infections of nemertean in gills (Pearse, 1949), amoebic infections (Sprague and Beckett, 1966; Sawyer, 1969; Sprague, Beckett and Sawyer, 1969; Pauley, Newman and Gould, 1975), effects of amoeba infections on hemocyte values (Sawyer, Cox and Higginbottom, 1970), gas-bubble disease (Johnson, 1976), microsporid infections (Sprague, 1965, 1966), bacterial infections (Krantz, Colwell and

Lovelace, 1969; Cook and Lofton, 1973), fungal infections (Couch, 1942; Newcombe and Rogers, 1947; Rogers-Talbert, 1948; Bland *et al.*, 1976) and trematode infections (Overstreet and Perry, 1972).

Commercial fishery and mariculture reports include: Texas fishery (Daugherty, 1952a; More, 1969), Louisiana fishery (Jaworski, 1970; Adkins, 1972b), Mississippi fishery (Perret, 1967; Perry, 1975), Chesapeake Bay fishery (Truitt, 1939; van Engel, 1958; Miller, Sulkin and Lippson, 1975), methods for handling soft crabs (Haefner and Garten, 1974), migration in bays and estuaries (Fiedler, 1930), and mariculture (Rust and Carlson, 1960; Sulkin and Epifanio, 1975). The preceding does not include unpublished reports of state Fish and Game Commissions, which contain statistics on commercial landings and are further summarized by publications of the U.S. Fish and Wildlife Service.

Callinectes similis Williams, 1966 (Tulane Stud. Zool. 13: 87)

Common Name: Lesser Blue Crab

As *C. danae*—Rathbun, 1930, p. 118 (part).

As *C. ornatus*—Hay & Shore, 1918, p. 433, pl. 34, fig. 2; Contreras, 1930, p. 231 (part), ? fig. 4; Rathbun, 1930, p. 114 (part); Williams, 1965, p. 172, fig. 152.

As *C. similis*—Williams, 1966, p. 87, figs. 3, 4E–F; Felder, 1973a, p. 58, pl. 8, fig. 1; Williams, 1974a, p. 731, figs. 4, 18a (not 18b), 20c, 22a, 24.

Range: Delaware Bay to southern Florida; Florida Keys; northwest Florida to Campeche, Yucatan, including Gulf coast of United States and Mexico.

Depth: shore to 92 m (50 fm).

Habitat: in ocean waters, near shore on sand and mud bottoms, often in association with populations of *C. sapidus*; in bays and estuaries, rarely below salinities of 15 ppt (Hoese, 1960 lists a range of 4.7 to 45 ppt), most common at 25 to 37 ppt; temperature ranges of 13 to 29°C. in the Gulf, slightly higher in Texas bays.

Remarks: With the exception of southwest Florida, all Gulf coast records of *C. danae* and *C. ornatus* for the United States and Mexico should be referred to *C. similis*. Regional lists, including records under *danae* and *ornatus*, include Florida (Wass, 1955; Tagatz, 1967; Abele, 1970), Mississippi (Richmond, 1962; Franks *et al.*, 1972; Christmas and Langley, 1973), Louisiana (Behre, 1950; Dawson, 1966), Texas (Gunter, 1950; Daugherty, 1952b; Hildebrand, 1954; Simmons, 1957; Hoese, 1960; Breuer, 1962; Leary, 1967; Hoese *et al.*, 1968), and Campeche (Hildebrand, 1955). Nocturnal swimming at the surface was noted by Franks *et al.* (1972).

Cronius Stimpson, 1860

Cronius ruber (Lamarck, 1818) (Hist. Nat. Anim. sans Vert., vol. 5, p. 260)

Rathbun, 1930, p. 139, pls. 62–63; Rathbun, 1933, p. 51; Garth, 1965a, p. 15; Williams, 1965, p. 174, fig. 154; Felder, 1973a, p. 55, pl. 8, fig. 3.

Range: South Carolina to south Florida; Dry Tortugas; off Texas; off Campeche, Mexico; Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands to Dominica; Caribbean coast of Panama; Amapá to Santa Catarina, Brazil; in

eastern Atlantic—from Cape Verde Island and Senegal to Angola; in eastern Pacific—from Lower California (Mexico) to Peru; Galapagos Islands; Clipperton Island.

Depth: shallow water to 105 m (to 57 fm).

Habitat: sandy bottoms; from areas of reefs, rocks, and shell rubble.

Remarks: Listed from Texas by Leary (1967) and confirmed by Felder (1973a). Garth and Stephenson (1966) commented on Pacific distribution and African collections were examined by Forest and Guinot (1966). Listed from Brazil to a depth of 105 m by Coelho and Ramos (1972).

Cronius tumidulus (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 149)

Rathbun, 1930, p. 142, pl. 64; Rathbun, 1933, p. 51, fig. 43.

Range: Bermuda; Bahamas; Florida Keys and Dry Tortugas; west coast of Florida; north and south coasts of Cuba; Jamaica; Puerto Rico; Virgin Islands; Netherlands Antilles; Old Providence Island (Carib.); Ceará to Bahia, Brazil.

Depth: 5 to 73 m (3 to 40 fm).

Habitat: coral, sand, and rock bottoms; grass, seaweed, and *Sargassum*.

Remarks: Listed from Brazil by Coelho and Ramos, 1972.

Lupella Rathbun, 1897

Lupella forceps (Fabricius, 1793) (Entom. Syst. emend. auct., vol. 2, p. 449)

Rathbun, 1930, p. 133, pl. 57; Rathbun, 1933, p. 50, fig. 41.

Range: north coast of Cuba; Jamaica; Haiti; Puerto Rico; St. Thomas, Virgin Islands; Martinique.

Depth: 13 to 15 m (7 to 8.5 fm).

Habitat: mud bottoms.

Portunus Weber, 1795

(This genus has traditionally been divided into subgenera, based on morphological criteria. Modern studies of *Portunus* (Stephenson and Rees, 1967; Stephenson, Williams, and Lance, 1968) have raised serious doubts about the subgeneric relationships and they suggest that further study will be necessary to define the complex phylogenetic affinities of this group, including the closely related genera, *Arenaeus* and *Callinectes*. As in Williams (1965), subgeneric categories are deleted here.)

Portunus anceps (Saussure, 1858) (Mém. Soc. Phys. Hist. Nat. Geneve 14: 434)

Hay & Shore, 1918, p. 431, pl. 33, fig. 8; Rathbun, 1930, p. 42, pl. 15; Rathbun, 1933, p. 46; Williams, 1965, p. 163, fig. 145.

Range: North Carolina; Bermuda; Bahamas; Florida Keys; north and south coasts of Cuba; Caribbean coast of Yucatan; Jamaica; Puerto Rico; St. Thomas, Virgin Islands to Guadeloupe; Panama to Bahia, Brazil.

Depth: surface to 103 m (to 56 fm).

Habitat: mainly on sandy bottoms, often with weeds or grass; also on mud, shell, and stone substrates; on coral reefs; buries in sand; in shallow waters, brackish ponds.

Remarks: Rathbun (1930) reported ovigerous females from North Carolina in October and from Cuba in June. Listed from Brazil by Coelho (1971a) and by Coelho and Ramos (1972).

Portunus binoculus Holthuis, 1969 (Bull. Mar. Sci. 19: 409)

Holthuis, 1969, p. 409, fig. 1.

Range: Bahamas; Florida Straits; north coast of Cuba; east of Yucatan, in Caribbean Sea; off Caribbean coasts of Panama and Colombia.

Depth: 74 to 291 m (40 to 159 fm), possibly to a range of 63 to 467 m (34 to 255 fm).

Remarks: Holthuis (1969) notes that some of the specimens of *P. spinicarpus* in Rathbun (1930) may be *P. binoculus* and these may include a record from Tortugas at 37 m (20 fm).

Portunus depressifrons (Stimpson, 1859) (Ann. Lyc. Nat. Hist. New York 7: 58)

Hay & Shore, 1918, p. 430, pl. 33, fig. 7; Rathbun, 1930, p. 84, pl. 41; Rathbun, 1933, p. 47; Williams, 1965, p. 166, fig. 149.

Range: North Carolina; Bermuda; south Florida; Florida Keys and Dry Tortugas; Bahamas; Gulf of Campeche, off Yucatan; north coast of Cuba; Culebra; St. Thomas, Virgin Islands; Aruba, Netherlands Antilles; Old Providence Island (Carib.).

Depth: surface to 29 m (to 16 fm).

Habitat: shallow water coves and inlets with sandy bottoms; on coral, shell, and grass-covered sand substrates.

Remarks: Listed from northwest Florida by Wass (1955), Abele (1970), and Menzel (1971). Williams (1965) notes the lack of recent specimens from the Florida Keys. Ovigerous females have been reported from Florida in June and August, the Caribbean in August (Rathbun, 1930), and from Campeche in August (Williams, 1965). Chace (1956) recorded this species off the north coast of Yucatan. Abele (1970) provided detailed notes on the color of a live female and he listed it as common in St. Andrews Bay, Florida.

Portunus floridanus Rathbun, 1930 (Bull. U.S. Nat. Mus. 152: 82)

Rathbun, 1930, p. 82, pl. 40.

Range: Key West, Florida.

Depth: 82 m (45 fm).

Habitat: coral reefs.

Portunus gibbesii (Stimpson, 1859) (Ann. Lyc. Nat. Hist. New York 7: 57)

Hay & Shore, 1918, p. 428, pl. 33, fig. 1; Rathbun, 1930, p. 49, pls. 16-17; Williams, 1965, p. 164, fig. 146; Felder, 1973a, p. 60, pl. 8, fig. 16.

Range: Massachusetts to south Florida; Florida Keys and Straits; Dry Tor-

tugas; west coast of Florida to south Texas; Campeche, Mexico; Venezuela; Surinam.

Depth: surface to 88 m (to 48 fm), rarely deeper.

Habitat: mud, sand, and broken shell bottoms; usually in deeper off-shore waters of Gulf coast and in deeper parts of near-shore marine passes, inlets, and bays.

Remarks: Regional lists include Florida (Wass, 1955; Tabb and Manning, 1961; Rouse, 1970; Abele, 1970; Menzel, 1971), Mississippi (Richmond, 1968; Franks *et al.*, 1972; Christmas and Langley, 1973), Louisiana (Dawson, 1966; Hoese and Valentine, 1972), Texas (Gunter, 1950; Hildebrand, 1954; Parker, 1959; Leary, 1967), and from the Gulf (Chace, 1956; Fotheringham and Brunenmeister, 1975). Hildebrand (1954) reported this crab abundant at 6 to 10 fm at Campeche. Tabb and Manning (1961) found this crab feeding on concentrations of cyprinodont fishes in Coot Bay, Florida. Musick and McEachran (1972) listed it from Chesapeake Bight at depths of 18 to 49 m. Felder (1973a) notes that this species is often found in association with *P. spinimanus*. Rathbun (1930) reported ovigerous females from Florida in April; Williams (1965) cites other ovigerous females from North Carolina to Surinam, from February to June. Gray (1957) measured the gill area of this crab and compared it with habitat preferences.

Portunus ordwayi (Stimpson, 1860) (Ann. Lyc. Nat. Hist. New York 7: 224)

Hay & Shore, 1918, p. 431, pl. 33, fig. 6; Rathbun, 1930, p. 71, pl. 33; Rathbun, 1933, p. 46; Chace, 1940, p. 31; Williams, 1965, p. 166, fig. 148.

Range: Massachusetts; North Carolina; Bermuda; Bahamas; southeast Florida; Florida Keys and Dry Tortugas; west and northwest coasts of Florida; north and south coasts of Cuba; Gulf and Caribbean coasts of Yucatan; Jamaica; Puerto Rico; St. Thomas, Virgin Islands to Dominica; Old Providence Island (Carib.); Pará and Fernando de Noronha to Bahia, Brazil.

Depth: surface to 106 m (58 fm), rarely deeper.

Habitat: sand, gravel, broken shell, and coral substrates.

Remarks: Rathbun (1930) listed an ovigerous female from Florida in March. Chace (1956) recorded this species from off northwest Florida. Listed from Brazil by Coelho (1971a), Coelho and Ramos (1972), and Fausto Filho (1974).

Portunus sayi (Gibbes, 1850) (Proc. Amer. Assoc. Adv. Sci., 3rd Meeting, 7: 178)

Common Name: Sargassum Crab

Hay & Shore, 1918, p. 428, pl. 33, fig. 2; Rathbun, 1930, p. 37, text-figs. 6-7, pl. 14; Rathbun, 1933, p. 45, fig. 39; Chace, 1940, p. 31; Williams, 1965, p. 163, fig. 144; Felder, 1973a, p. 59, pl. 8, fig. 12.

Range: Nova Scotia to south Florida; Bermuda; Bahamas; Florida Keys and Dry Tortugas; west coast of Florida to south Texas; north and south coasts of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Trinidad; Guiana; Brazil; in eastern Atlantic Ocean; off Kerguelon Island in the Indian Ocean.

Depth: pelagic, at surface.

Habitat: normally among *Sargassum*, floating on surface; also on other flotsam; occasionally swimming freely.

Remarks: Larval stages from Bermuda were described by Lebour (1944) and Coventry (1944) described Caribbean collections. Williams (1965) listed ovigerous females in the northeast Gulf and in the West Indies from February to August and from off Massachusetts in September. Regional lists include Florida (Abele, 1970), Mississippi (Franks *et al.*, 1972), Louisiana (Behre, 1950), Texas (Leary, 1967), and the Gulf (Chace, 1956; Fotheringham and Brunenmeister, 1975). Autotomy and regeneration of chelae was studied by Zeleny (1908); Hartnoll (1971) described swimming behavior.

Portunus sebae (H. Milne Edwards, 1834) (Hist. Nat. Crust., vol. 1, p. 455)

Rathbun, 1930, p. 79, pls. 34-35; Rathbun, 1933, p. 46.

Range: Bermuda; Florida Keys and Straits; Dry Tortugas; south coast of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Dominica; Netherlands Antilles.

Depth: 4 to 18 m (2 to 10 fm).

Habitat: sand, rocky, and grass-covered sandy bottoms.

Portunus spinicarpus (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 148)

Hay & Shore, 1918, p. 429, pl. 33, fig. 3; Rathbun, 1930, p. 92., pl. 45; Rathbun, 1933, p. 47; Chace, 1940, p. 32; Williams, 1965, p. 167, fig. 150; Holthuis, 1969, p. 415, fig. 1; Felder, 1973a, p. 60, pl. 8, figs. 13-14.

Range: North and South Carolina; southeast Florida; Florida Keys and Dry Tortugas; west coast of Florida to south Texas; north and south coasts of Cuba; Puerto Rico; Trinidad; Caribbean coast of Colombia; Guianas to Santa Catarina, Brazil.

Depth: 9 to 550 m (5 to 300 fm).

Habitat: on sand, gravel, coral, broken shell, and mud substrates.

Remarks: Larval development was described by Bookhout and Costlow (1974). Williams (1965) listed ovigerous females from Texas in November. Rathbun (1930) noted that despite large numbers of this crab in the Florida Keys, fish stomachs do not contain this species, which she attributes to the Crab's extendable carpal spines. Pearse (1932a) reported the presence of the barnacle, *Dichelastis sinvata*, on crabs from North Carolina. Holthuis (1969) expanded the previously known depth range to 550 m and he reported ovigerous females from the Caribbean from January to September, with some females carrying sacculinid barnacles. Musick and McEachran (1972) collected this crab from a depth of 101 m in Chesapeake Bight. Regional lists included Florida (Hulings, 1961; Abele, 1970), Mississippi (Richmond, 1962; 1962; Franks *et al.*, 1972), Louisiana (Dawson, 1966), and Texas (Hildebrand, 1954; Chace, 1956; Leary, 1967). Listed from Brazil by Rodrigues da Costa (1968a), Coelho (1971a) and Coelho and Ramos (1972).

Portunus spinimanus Latreille, 1819 (Nouv. Dict. Hist. Nat., ed. 2, vol. 28, p. 47)

Hay & Shore, 1918, p. 429, pl. 33, fig. 4; Rathbun, 1930, p. 62, text-fig. 10, pls.

26-28; Rathbun, 1933, p. 46; Chace, 1940, p. 31; Williams, 1965, p. 165, fig. 147; Felder, 1973a, p. 59, pl. 8, fig. 15.

Range: New Jersey to south Florida; Bermuda; Bahamas; Florida Keys and Dry Tortugas; west Florida to south Texas; Campeche, off Mexico; south coast of Cuba; Jamaica; Puerto Rico; Hispaniola; Trinidad; Aruba, Netherlands Antilles; Bahia to Santa Catarina, Brazil.

Depth: surface to 91 m (50 fm).

Habitat: waters of inlets, canals, and harbors; on sand, gravel, broken shell, and mud bottoms; occasionally on *Sargassum*.

Remarks: Lebour (1950) raised and described larvae collected at Bermuda. Holthuis (1959) listed ovigerous females from Surinam in May, August, and September; Williams (1965) summarized other records of ovigerous females. Regional lists include Florida (Wass, 1955; Tabb and Manning, 1961; Abele, 1970; Menzel, 1971), Mississippi (Richmond, 1962), Louisiana (Behre, 1950), Texas (Hildebrand, 1954; Parker, 1959; Leary, 1967), and Campeche (Hildebrand, 1954). This crab was abundant at Campeche at depths of 6 to 10 fm, often in association with *P. gibbesii*. Reported from the Gulf by Chace (1956) and described generally by Fotheringham and Brunenmeister (1975). Musick and McEachran (1972) found this crab at 49 m depth in Chesapeake Bight. Gray (1957) measured gill area. Lobo de Mesquita (1972) reported biometrical data on Brazilian specimens; listed from Brazil by Coelho and Ramos (1972).

Portunus ventralis (A. Milne Edwards, 1879) (Crust. Rég. Mex., p. 215)

Rathbun, 1930, p. 43, pl. 13, figs. 1-2; Rathbun, 1933, p. 46; Chace, 1940, p. 31; Felder, 1973a, p. 59, pl. 8, fig. 11.

Range: Georgia to east coast of Florida; Dry Tortugas; ? Texas; north and west coasts of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Barbados; Rio Grande do Norte to Rio de Janeiro, Brazil.

Depth: shallow water, near shore, to 25 m (14 fm).

Habitat: sandy beaches; tide pools; on surface of open waters.

Remarks: Felder (1973a) questioned reports by Parker (1959) and Trott (unpublished) of this crab's occurrence in Texas; these records have not been verified by later collections. Rathbun (1930) reported ovigerous females from Dry Tortugas in August. Listed from Brazil by Coelho and Ramos (1972).

Portunus vocans (A. Milne Edwards, 1878) (Bull. Soc. Philom. Paris, ser. 7, vol. 2, p. 225)

Rathbun, 1930, p. 60, text-figs. 8-9, pl. 25.

Range: north coast of Cuba; between Jamaica and Haiti; Ascension Island, in the South Atlantic Ocean.

Depth: 37 to 309 m (20 to 169 fm).

Habitat: coral and broken shell bottoms.

Superfamily XANTHOIDEA Dana, 1851

Family POTAMIDAE Ortmann, 1896

The Freshwater Crabs

The familial name, based on the genus *Potamon* Savigny, 1816, was corrected from the original Potamonidae of Ortmann by an ICZN decision (Opinion 712, p. 342, in 1964). A revision of the freshwater crabs by Bott (1955b) split this family into two: the Pseudothelphusidae and the Trichodactylidae. Various other schemes for classification have been proposed (Pretzmann, 1965; Bott, 1968; Smalley, 1970). Gulf region species are confined to northern Cuba and southern Mexico, but insufficient material and information is available for most species to yield accurate ranges or to present a satisfactory organization of taxonomic relationships. Because these crabs are inhabitants of rivers, lakes, and caves, species ranges tend to be restricted to small areas; several are known only from a single type specimen. Species occurring in the center of a land mass, such as *Potamocarcinus* (*Typhlopseudothelphusa*) *mocinoi* in caves near Comitán (Chiapas) Mexico (Rioja, 1952) could be assigned to a coastal region based on watershed drainage patterns. The Cuban species listed by Chace and Hobbs (1969) are not localized to either the Gulf or the Caribbean side of the island in the records cited. For these reasons, these crabs are not included in the present work. Further references, including some older systematic papers, can be found in the papers cited above. There exists a clear need for more extensive work on this group of brachyurans.

Family XANTHIDAE Dana, 1851

This is a large family of crabs (about 1000 species and more than 130 genera) that has traditionally posed a number of taxonomic problems. Many of the species are small in size and appear morphologically similar. Individual variability and the large number of closely related species has often made definitive identification difficult, so that earlier collection records must be used cautiously. Rathbun (1930) did not subdivide her account of the xanthids into subfamilies. Guinot (1971) offers a number of systematic revisions and she comments at some length on affinities, but she also avoids listing the 51 genera that she treats under subfamilies. The same procedure is followed here, by arrangement of the 33 Gulf genera in alphabetical order, without regard for proposed affinity within the family. The xanthids are currently being revised by various workers around the world, so that a better organized and more accurate representation of this family should be forthcoming.

Actaea de Haan, 1833

Actaea acantha (H. Milne Edwards, 1834) (Hist. Nat. Crust., vol. 1, p. 379)

Rathbun, 1930, p. 261, pl. 105, fig. 5, pl. 106, figs. 1-2; Rathbun, 1933, p. 57.

Range: Bahamas; Florida Keys and Dry Tortugas; northwest coast of Cuba; Jamaica; Haiti; Puerto Rico; Guadeloupe; St. Bartholomew; Fernando de Noronha, Brazil.

Depth: surface to 22 m (to 12 fm).

Habitat: sand, shell, coral bottoms; from coral reefs; off mud and grassy bottoms.

Remarks: Habitat and color of specimens from Brazil were described by Fausto Filho (1974).

Actaea bifrons Rathbun, 1898 (Bull. Lab. Nat. Hist. State Univ. Iowa 4: 262)

Rathbun, 1930, p. 255, text-fig. 41, pl. 104, figs. 3-6; Rathbun, 1933, p. 56, fig. 48.

Range: Key West, Florida; Puerto Rico; Virgin Islands; St. Bartholomew; Barbados; Curaçao; Colon, Panama.

Depth: 18 to 73 m (10 to 40 fm).

Habitat: in coral, *Porites furcata*; on shoal banks, coral bottoms, sponge areas.

Actaea palmeri Rathbun, 1894 (Proc. U.S. Nat. Mus. 17: 85)

Rathbun, 1930, p. 260, pl. 106, figs. 3-6; Rathbun, 1933, p. 56.

Range: Bahamas; east coast of Florida; Florida Keys; north coast of Cuba; Haiti; Virgin Islands; Curaçao.

Depth: shallow water to 145 m (to 78 fm).

Habitat: from sponges and among coral (*Porites furcata*).

Actaea rufopunctata nodosa Stimpson, 1860.

Transferred to a new genus, *Paractaea*, by Guinot (1969b). See *Paractaea rufopunctata nodosa* (Stimpson, 1860).

Actaea setigera (H. Milne Edwards, 1834).

Transferred to a new genus, *Platyactaea*, by Guinot (1967b). See *Platyactaea setigera* (H. Milne Edwards, 1834).

Carpilius Leach, 1823

Carpilius corallinus (Herbst, 1783) (Natur. Krabben u. Krebse, vol. 1, p. 133)

Common Names: Queen Crab; Coral Crab

Rathbun, 1930, p. 240, pls. 97-99; Rathbun, 1933, p. 53; Chace, 1940, p. 33; Guinot, 1968b, p. 157, fig. 9; Guinot, 1968c, p. 321, figs. 10-11; Pequegnat & Ray, 1974, p. 237, figs. 13-15.

Range: Bermuda; Bahamas; West Flower Garden Bank, off Texas; north coast of Cuba; Jamaica; Puerto Rico; Virgin Islands; Guadeloupe; Dominica; Curaçao; Old Providence Island (Carib.); Pernambuco and Ceará, Brazil.

Depth: 2 to 46 m (1 to 25 fm).

Habitat: on coral reefs; sandy, coral, and stone substrates.

Remarks: This is the largest crab found in the Gulf of Mexico and Caribbean area and it is used for food in the West Indies. It reaches carapace widths of more than 15 cm. Garth (1965a) compared this species with *C. convexus* from the Pacific. Pequegnat and Ray (1974) reported some observations on mating and other behavior seen on a coral reef off the Texas coast. Saraiva da Costa (1968) described the biology and fishery of this species at Ceará, Brazil. Listed from Brazil by Fausto Filho (1968).

***Carpoporus* Stimpson, 1871**

***Carpoporus papulosus* Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 139)**

Rathbun, 1930, p. 269, pl. 110, figs. 3-6, pl. 111; Williams, 1965, p. 186, figs. 168, 183B; Guinot, 1967b, p. 551, figs. 18-19, 22.

Range: North Carolina; Florida Keys and Dry Tortugas; west and northwest coasts of Florida; Alabama; north of Yucatan, Mexico.

Depth: 33 to 110 m (18 to 60 fm).

Habitat: sand, broken shell, and coral bottoms.

Remarks: Wass (1955) listed this species from northwest Florida.

***Cataleptodius* Guinot, 1968**

***Cataleptodius floridanus* (Gibbes, 1850) (Proc. Amer. Assoc. Adv. Sci. 3: 175)**

As *Leptodius floridanus*—Rathbun, 1930, p. 297, pl. 137, figs. 1-2, pl. 138, fig. 1; Rathbun, 1933, p. 57.

As *Cataleptodius floridanus*—Guinot, 1968a, p. 706, figs. 20, 23, 29.

Range: Bermuda; Bahamas; Florida Keys and Dry Tortugas; northwest coast of Florida; north coast of Cuba; Jamaica; Puerto Rico; Virgin Islands; Antigua; Barbados; Curaçao; Panama to Colombia (Caribbean coasts; Abolhos Islands to São Paulo, Brazil.

Depth: shallow water to 33 m (to 18 fm).

Habitat: on coral and stone reefs; in *Sargassum*; in living sponges; from sand, shell, grassy, mud bottoms; intertidal pools in rocky areas; under rocks.

Remarks: Menzel (1971) listed this crab as rare on oyster reefs at Apalachee Bay, Florida. Listed from Florida by Abele (1970). Sulkin (1973) described larval depth regulation and Epifanio (1972) studied the effects of dieldrin on larval development. Hazlett (1976) described agonistic behavior in this crab. Listed from Brazil by Fausto Filho (1974), he provided notes on color of crab and its habitat.

***Chlorodiella* Rathbun, 1897**

***Chlorodiella longimana* (H. Milne Edwards, 1834) (Hist. Nat. Crust., vol. 1, p. 401)**

Rathbun, 1930, p. 462, pl. 186; Rathbun, 1933, p. 68, fig. 58.

Range: Bahamas; east coast of Florida; Florida Keys and Dry Tortugas; Jamaica; Puerto Rico; Virgin Islands; Martinique; Barbados; Curaçao.

Depth: 5 to 154 m (3 to 84 fm).

Habitat: coral reefs; rock and stone substrates; in sponges.

Remarks: Rathbun (1930) also listed this species from West Africa.

***Domecia* Eydoux & Souleyet, 1842**

***Domecia acanthophora acanthophora* (Desbonne & Schramm, 1867) (Crust. Guadeloupe, p. 35)**

As *D. hispida*—Rathbun, 1930, p. 554, pl. 227; Pequegnat & Ray, 1974, p. 237, figs. 16-17.

As *D. acanthophora acanthophora*—Guinot, 1964, p. 271, figs. 4–5, 7–8, 15; Williams, McCloskey & Gray, 1968, p. 52.

Range: North and South Carolina; east coast of Florida; Florida Keys and Dry Tortugas; West Flower Garden Bank, off Texas; Cuba; Jamaica; Puerto Rico; Barbados; Curaçao; Alagoas to Pernambuco, Brazil.

Depth: low tide mark to 146 m (to 80 fm).

Habitat: on coral reefs; among sponges; rocky and coral bottoms.

Remarks: The Pacific specimens of *D. hispida* listed by Rathbun (1930) remain with the species indicated; the Atlantic specimens were referred by Guinot (1964) to *D. acanthophora*. Patton (1967) studied the ecology of this species on coral reefs (*Acropora*) off Puerto Rico. A single male was taken from *Oculina* off North Carolina by Williams, McCloskey and Gray (1968).

Domecia hispida Eydoux & Souleyet, 1842.

Atlantic specimens were referred to *D. acanthophora* (Desbonne & Schramm, 1867) by Guinot (1964).

Eriphia Latreille, 1817

Eriphia gonagra (Fabricius, 1781) (Species Insectorum, p. 505)

Common Name: Calico Crab

Hay & Shore, 1918, p. 439, pl. 35, fig. 6; Rathbun, 1930, p. 545, text-fig. 83, pl. 222; Rathbun, 1933, p. 75, fig. 64; Williams, 1965, p. 182, figs. 164A–C, 165; Felder, 1973a, p. 64, pl. 9, fig. 5.

Range: North Carolina; Bermuda; Bahamas; Florida Keys and Dry Tortugas; south Texas; north coast of Cuba; Jamaica; Puerto Rico; Virgin Islands; Antigua; Barbados; Trinidad; Aruba, Netherlands Antilles; Caribbean coasts of Panama and Colombia; Parahyba, Brazil to Argentina.

Depth: intertidal to shallow water subtidal.

Habitat: coral and stone reefs; under rocks and in crevices of intertidal pools; on rock jetties; brackish ponds; in seaweed and sponges.

Remarks: Ovigerous females are known from south Florida and the West Indies during March to September (Williams, 1965). Furtado-Ogawa (1972) noted habitats in Brazil.

Etisus H. Milne Edwards, 1834

Etisus maculatus (Stimpson, 1860) (Ann. Lyc. Nat. Hist. New York 7: 210)

As *Phymodius maculatus*—Rathbun, 1930, p. 295, pl. 136; Rathbun, 1933, p. 57, fig. 49.

As *Etisus maculatus*—Guinot, 1969b, p. 234.

Range: Florida Keys and Dry Tortugas; Bahamas; north coast of Cuba; Puerto Rico; Virgin Islands.

Depth: low tide mark and shallow water.

Habitat: coral reefs; among rocks.

Remarks: In transferring this species to *Etisus*, Guinot (1969b) remarks that

it becomes the sole representative of this genus in the Atlantic, other members of the genus being Indo-Pacific.

Eucratodes A. Milne Edwards, 1880

Eucratodes agassizii A. Milne Edwards, 1880 (Crust. Rég. Mex., p. 347)

Rathbun, 1930, p. 471, pl. 190; Rathbun, 1933, p. 68, fig. 59; Guinot, 1969a, p. 722, figs. 145-146; Pequegnat, 1970, p. 188.

Range: off Mississippi; Yucatan Channel (Caribbean); Puerto Rico.

Depth: 156 to 315 m (85 to 172 fm).

Habitat: mud, sand, and shell bottoms.

Remarks: Pequegnat (1970) collected an ovigerous female from 100 fm off Mississippi in early December.

Eurypanopeus A. Milne Edwards, 1880

Eurypanopeus abbreviatus abbreviatus (Stimpson, 1860) (Ann. Lyc. Nat. Hist. New York 7: 211)

Rathbun, 1930, p. 405, text-fig. 63, pl. 172, figs. 1-2; Rathbun, 1933, p. 64, fig. 55; Williams, 1965, p. 194, figs. 178, 183K; Felder, 1973a, p. 68, pl. 9, fig. 13.

Range: South Carolina; Bahamas; Florida Keys; ? Louisiana; Texas; Jamaica; Haiti; Puerto Rico; Virgin Islands; Antigua; Barbados; Curaçao; Trinidad; Colombia to Venezuela; Parahyba to Santa Catarina, Brazil.

Depth: intertidal to shallow subtidal.

Habitat: under rocks; on stone and coral reefs; under sponges and bryozoans; on oyster beds.

Remarks: Behre (1950) listed a specimen of *E. crenatus* from Grand Isle, Louisiana. Felder (1973a) notes that *E. crenatus* is known primarily from the Pacific coast of South America and that Behre's record may be a mistaken identification of *E. abbreviatus*. Williams (1965) listed ovigerous females from the West Indies during April to November and in southern Brazil from August to November. Furtado-Ogawa (1972) commented on ecology of this species in Brazil.

Eurypanopeus abbreviatus ater Rathbun, 1930 (Bull. U.S. Nat. Mus. 152: 407)

Rathbun, 1930, p. 407, pl. 172, figs. 3-4.

Range: Vera Cruz, Mexico.

Remarks: Known only from the single male type specimen; no other data available.

Eurypanopeus depressus (Smith, 1869) (Proc. Boston Soc. Nat. Hist. 12: 283)

Hay & Shore, 1918, p. 437, pl. 34, fig. 4; Rathbun, 1930, p. 410, text-fig. 65, pl. 173, figs. 3-4; Williams, 1965, p. 195, figs. 179, 183L; Felder, 1973a, p. 67, pl. 9, fig. 17.

Range: Massachusetts to south Florida; Bermuda; west coast of Florida to Texas; St. Martin, Leeward Islands.

Depth: intertidal to 48 m (to 26 fm).

Habitat: very common on oyster bars; on muddy and stony shores, usually intertidal; on wharves and submerged pilings; in eel grass; brackish waters, to 4.5 ppt salinity.

Remarks: Developmental studies include descriptions of larval stages (Costlow and Bookhout, 1961b) and data on larval ecology in Chesapeake Bight (Sandifer, 1973). Life history information was provided by Ryan (1956). Hyman (1925) figured zoeal stages. Lunz (1937) observed the association of this crab with oysters; McDermott (1960) noted the threat to oyster spats in New Jersey. Abele (1970) noted that the presence of the red spot on the third maxillipeds was not consistent among specimens, but is found in crabs only associated with oysters. Records of ovigerous females are summarized in Williams (1965). Regional lists include Florida (Wass, 1955; Tabb and Manning, 1961; Dragovich and Kelly, 1964; Abele, 1970; Lyons *et al.*, 1971; Menzel, 1971), Mississippi (Christmas and Langley, 1973), Louisiana (Behre, 1950), Texas (Leary, 1967), and the northwestern Gulf of Mexico (Fotheringham and Brunenmeister, 1975). Ayres (1938) studied respiration in relation to habitat preferences.

Eurypanopeus dissimilis (Benedict & Rathbun, 1891) (Proc. U.S. Nat. Mus. 14: 366)

Rathbun, 1930, p. 411, text-fig. 66, pl. 173, figs. 1-2.

Range: west coast of Florida; north coast of Cuba; Jamaica; Nicaragua; Trinidad; Brazil.

Habitat: Listed from a salt water lagoon in Nicaragua and from harbors in Florida and Jamaica, no depths given.

Eurypanopeus turgidus (Rathbun, 1930).

This is a manuscript name used by Abele (1970) for *Panopeus turgidus* Rathbun, 1930 and listed by Menzel (1971, p. 80) for northwest Florida. Dr. Fenner A. Chace, Jr. examined the identification of Abele's specimens and compared them with species of *Eurypanopeus*. Until official revision of the two genera is complete, the present work will continue to recognize this species as a member of the genus *Panopeus*, while recognizing its probable affinity with *Eurypanopeus*. See *Panopeus turgidus*.

Eurytium Stimpson, 1859

Eurytium limosum (Say, 1818) (J. Acad. Nat. Sci., Philadelphia 1: 446)

Hay & Shore, 1918, p. 438, pl. 35, fig. 7; Rathbun, 1930, p. 423, pl. 176, figs. 1-2; Rathbun, 1933, p. 65, fig. 56; Chace, 1940, p. 34; Williams, 1965, p. 199, figs. 182, 1830; Chace & Hobbs, 1969, p. 153, figs. 45, 46b; Felder, 1973a, p. 65, pl. 9, fig. 4.

Range: Bermuda; Bahamas; New York to south Florida; Florida Keys and Dry Tortugas; west and northwest coasts of Florida; Louisiana; north and south coasts of Cuba; Jamaica; Hispaniola; Puerto Rico; Virgin Islands; Islas Los Roques; Curaçao; Belize; Caribbean coast of Panama; Maranhão to São Paulo, Brazil.

Depth: intertidal to shallow subtidal, near shore.

Habitat: muddy shores, especially among mangroves; burrows along tidal stream banks, burrows partially filled with water; under stones at high tide mark.

Remarks: Williams (1965) notes that modern records limit the northern extent of the range to South Carolina. Regional lists include Florida (Wass, 1955; Tabb and Manning, 1961; Abele, 1970; Menzel, 1971; Subrahmanyam *et al.*, 1976), and Louisiana (Behre, 1950; Hoese and Valentine, 1972). Manning (1961) compared growth of this crab with that in *Menippe mercenaria* and *Panopeus herbstii*. Warner (1969) provided ecological data on this species in Jamaica and Ryan (1956) described the life history of populations observed in Chesapeake Bay. Teal (1959) studied respiration of this crab under different field conditions.

^{SP}
Glyptoxanthosus A. Milne Edwards, 1879 *Glyptoxanthosus*

Glyptoxanthosus erosus (Stimpson, 1859) (Ann. Lyc. Nat. Hist. New York 7:51)

Rathbun, 1930, p. 263, pl. 107; Williams, 1965, p. 185, figs. 167, 183A; Guinot, 1967b, p. 551, fig. 30; Felder, 1973a, p. 60, pl. 9, fig. 9.

Range: Bahamas; North Carolina; east coast of Florida; Florida Keys and Dry Tortugas; west and northwest coasts of Florida; Louisiana and Texas (uncommon); Campeche Banks and off Yucatan, Mexico; Guadeloupe.

Depth: low tide mark to 68 m (to 37 fm).

Habitat: sand, broken shell, coral, and rock bottoms; from rocks and algal mats (*Halimeda*) in shallow water; sponges and coral reefs of deeper water.

Remarks: Ovigerous females are known from off northwest Florida in January (Williams, 1965). Listed from Florida by Wass (1955) and Chace (1956), from Louisiana by Behre (1950), and from Texas by Leary (1967).

Heteractaea Lockington, 1877

Heteractaea ceratopus (Stimpson, 1860) (Ann. Lyc. Nat. Hist. New York 7:215)

Rathbun, 1930, p. 530, pl. 212, figs. 5-8; pl. 213; Guinot, 1968a, p. 721, figs. 50, 56.

Range: Bahamas; east coast of Florida; Florida Keys and Dry Tortugas; north coast of Cuba; Curaçao; Trinidad; Barbados.

Depth: shallow water.

Habitat: on coral reefs; in sponges and coral.

Hexapanopeus Rathbun, 1898

Hexapanopeus angustifrons (Benedict & Rathbun, 1891) (Proc. U.S. Nat. Mus. 14: 373)

Hay & Shore, 1918, p. 436, pl. 34, fig. 7; Rathbun, 1930, p. 384, text-fig. 60, pl. 169, figs. 1-2; Williams, 1965, p. 188, figs. 170, 183D; Felder, 1973a, pl. 9, fig. 24.

Range: Massachusetts to South Carolina; Bahamas; west and northwest coasts of Florida; Mississippi to Texas; Jamaica.

Depth: near shore subtidal to 139 m (to 76 fm).

Habitat: most commonly on muddy bottoms; occasionally on sand, shell, and gravel substrates; on oyster beds.

Remarks: Developmental studies include data on growth conditions for zoeal stages (Chamberlain, 1961), descriptions of larval stages as reared in the laboratory (Costlow and Bookhout, 1966a), and notes on larval ecology in Chesapeake Bight (Sandifer, 1973). Williams (1965) summarized records of ovigerous females. Ryan (1956) described the general life history of this crab in Chesapeake Bay. Hazlett (1976) studied agonistic behaviors. Regional lists include Florida (Wass, 1955; Dragovich and Kelly, 1964; Abele, 1970; Menzel, 1971; Lyons *et al.*, 1971), Louisiana (Behre, 1950), and Texas (Hedgpeth, 1953; Leary, 1967).

Hexapanopeus hemphillii (Benedict & Rathbun, 1891) (Proc. U.S. Nat. Mus. 14: 374)

Rathbun, 1930, p. 400, pl. 171, figs. 1–2, 6; Rathbun, 1933, p. 63.

Range: Florida Keys and Dry Tortugas; west coast of Florida; north coast of Cuba; Puerto Rico; St. Thomas, Virgin Islands.

Depth: low tide mark to 9 m (to 5 fm).

Habitat: coral, sand, and gravel bottoms; off turtle grass flats.

Hexapanopeus lobipes (A. Milne Edwards, 1880) (Crust. Rég. Mex., p. 331)

As *Lophopanopeus lobipes*—Rathbun, 1930, p. 329, text-fig. 50, pl. 155, figs. 3–5.

As *Hexapanopeus lobipes*—Menzies, 1948, p. 23.

Range: Bahamas; off Key West, in Florida Straits; northwest of Dry Tortugas.

Depth: 68 m (37 fm); also recorded from shallow water.

Habitat: sand bottoms; coral banks.

Remarks: Menzies (1948) noted that this species did not fit the generic description of *Lophopanopeus* and he transferred it to *Hexapanopeus*, the American genus it most closely resembled.

Hexapanopeus paulensis Rathbun, 1930 (Bull. U.S. Nat. Mus. 152: 395)

Rathbun, 1930, p. 395, pl. 170, figs. 5–6; Williams, 1965, p. 189, figs. 171, 183E; Felder, 1973a, p. 70, pl. 9, fig. 23.

Range: South Carolina; northwest Florida; Texas; São Paulo, Brazil.

Depth: intertidal to 5 m (3 fm).

Habitat: sand, broken shell, and rock substrates; off rock jetties; among sponges, ascidians, and bryozoans.

Remarks: Originally listed from South Carolina by Lunz (1937). Additional records, from northwest Florida, were provided by Abele (1970), including ovigerous females taken in July.

Hexapanopeus quinquentatus Rathbun, 1901 (Bull. U.S. Fish Comm. for 1900, 2: 31)

Rathbun, 1930, p. 402, text-fig. 62.

Range: northwest Florida; Puerto Rico.

Depth: 15 to 22 m (8.5 to 12 fm).

Habitat: coral sand, sand-grass, rocky, and sticky mud bottoms.

Remarks: Rathbun (1930) noted the resemblance between this species and *H. sinaloensis*, a rare species from the west coast of Mexico. Abele (1970) provided the first Gulf records and noted a close resemblance to *H. paulensis*. Abele (1970) states that Fenner Chace, Jr., of the U.S.N.M. compared the Floridian specimens with the Puerto Rico holotype and noted the differences between them. The paucity of specimens of these three similar forms only permits a tentative designation of this crab; until further material and revision is available, this list follows the nomenclature of Abele (1970, p. 78).

Leptodius A. Milne Edwards, 1863

Leptodius agassizii A. Milne Edwards, 1880.

Transferred to a new genus by Guinot (1968a), see *Pseudomedeus agassizii*.

Leptodius floridanus (Gibbes, 1850).

Transferred to a new genus by Guinot (1968a), see *Cataleptodius floridanus*.

Leptodius parvulus (Fabricius, 1793) (Entom. Syst. Auct. et emend., vol. 2, p. 451) novo comb. Rathbun, 1930 (Bull. U.S. Nat. Mus. 152:305) ^{transferred}

Rathbun, 1930, p. 305, pl. 141, figs. 1-3; Rathbun, 1933, p. 58, fig. 50. _{by Guinot}

Range: Bermuda; Bahamas; Florida Keys; Jamaica; Haiti; Puerto Rico; Barbados; Curaçao; Fernando de Noronha, Brazil.

Habitat: shallow water, near shore; under rocks; in tide pools.

Remarks: Fausto Filho (1974) provided notes on color and habitat of Brazilian specimens.

Lobopilumnus A. Milne Edwards, 1880

Lobopilumnus agassizii (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 142)

Hay & Shore, 1918, p. 441, pl. 34, fig. 5; Rathbun, 1930, p. 526, pl. 211, figs. 1-5; Williams, 1965, p. 181, figs. 157G, 163.

Range: Bermuda; North Carolina; Florida Keys and Dry Tortugas; west and northwest coasts of Florida; north of Yucatan; north coast of Cuba; Trinidad.

Depth: low tide mark to 51 m (to 28 fm).

Habitat: sand, gravel, rock, coral, and broken shell substrates; under stones and dead corals; in sponges.

Remarks: Four environmental forms of this highly variable crab were recognized by Rathbun (1930): *L. a. typica*, *L. a. bermudensis*, *L. a. pulchella*, and *L. a. trinidadensis*. The distributions of the various forms overlap to some extent. Verrill (1908) described this species in Bermuda. Reported from the northeast Gulf by Chace (1956) and listed from northwest Florida by Abele (1970). Ovigerous females are known from Florida and Cuba during February to July (Williams, 1965). Pearse (1934) noted the presence of this crab inside the loggerhead sponge, *Spheciospongia vesparia*.

Lophopanopeus Rathbun, 1898

Lophopanopeus distinctus Rathbun, 1898.

Transferred to the genus *Micropanope* by Menzies (1948), see *Micropanope distincta* (Rathbun, 1898).

Lophopanopeus lobipes (A. Milne Edwards, 1880).

Transferred to the genus *Hexapanopeus* by Menzies (1948), see *Hexapanopeus lobipes* (A. Milne Edwards, 1880).

Melybia Stimpson, 1871

Melybia thalamita Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 144)

Rathbun, 1930, p. 562, pl. 230.

Range: Dry Tortugas; north coast of Cuba; Jamaica; Haiti; Culebra and Vieques Islands; Barbados; Curaçao; Colon, Panama; Abrolhos Islands, Brazil.

Depth: low tide mark to 368 m (to 201 fm), most common to 82 m (45 fm).

Habitat: coral, sand, rock, and broken shell bottoms.

Remarks: The deepest recorded specimens (192 and 201 fm) are from off Havana, Cuba.

Menippe de Haan, 1833

Menippe mercenaria (Say, 1818) (J. Acad. Nat. Sci. Philadelphia 1: 448)

Common Name: Stone Crab

Hay & Shore, 1918, p. 439, pl. 35, fig. 8; Rathbun, 1930, p. 472, text-fig. 78, pls. 191-193; Williams, 1965, p. 183, figs. 164D-E, 166; Felder, 1973a, p. 64, pl. 9, figs. 2-3.

Range: North Carolina to south Florida; Bahamas; Florida Keys; southwest Florida to south Texas; off Yucatan Gulf coast; north coast of Cuba; Jamaica.

Depth: surface and intertidal to 51 m (to 28 fm).

Habitat: in estuaries and bays of near-marine salinity; from sand, shell, clay, and mud substrates; in deeper waters on offshore reefs; in turtle grass (*Thalassia*) beds off northwest Florida; post-larval crabs are common in deeper channels and bays, under shell fragments; older juveniles and adults are among rocks, under stones, and on and among oyster bars.

Remarks: Except for the blue crab (*Callinectes sapidus*), this species is the most important crab commercially harvested on the Gulf coast, primarily in Florida. Williams (1965) summarizes much of the literature and accounts of the natural history of stone crabs in Texas are provided by Powell and Gunter (1968) and by Futch (1966) for Florida populations. Fotheringham and Brunenmeister (1975) offer general comments on the stone crab in the northwestern Gulf.

Developmental studies include data on fertilization (Binford, 1913), zoeal stage descriptions (Porter, 1960), studies of effects of temperature and salinity on larvae (Ong and Costlow, 1970), effects of Mirex on larvae (Bookhout *et al.*, 1972), and studies of larval energy budgets (Mootz and Epifanio, 1974). Studies

of behavior include those on mating (Hartnoll, 1969; Savage, 1971b), stridulation (Guinot-Dumortier and Dumortier, 1960), and shadow responses of larvae (Forward, 1977). Physiological research includes data on hormonal control of reproduction (Cheung, 1967, 1969), respiration in relation to habitat (Ayres, 1938), gill area measurements (Gray, 1957), respiration and metabolism (Leffler, 1973), respiration and osmoregulation (Karandieva and Lee, 1967), and digestion and energy balance (Suchenia and Claro Madruga, 1967).

Information on growth, molting, and fisheries includes reports by Manning (1961), Savage and McMahan (1968), Savage (1971a), and Savage, Sullivan and Kalman (1974, 1975). Cheung (1973, 1976) studied regeneration of claws in relation to molting. Menzel and Hopkins (1956) described predation of stone crabs on oysters in Louisiana. Iversen and Beardsley (1976) studied shell disease in south Florida populations. Humes (1941a) noted the presence of a parasitic barnacle.

Regional lists include Florida (Wass, 1955; Tabb and Manning, 1961; Dragovich and Kelly, 1964; Abele, 1970; Menzel, 1971; Lyons *et al.*, 1971), Mississippi (Richmond, 1962; Franks *et al.*, 1972; Christmas and Langley, 1973), Louisiana (Behre, 1950; Hoese and Valentine, 1972), Texas (Gunter, 1950; Whitten, Rosene and Hedgpeth, 1950; Hedgpeth, 1953; Hildebrand, 1954; Simmons, 1957; Parker, 1959; Hoese, 1960; Leary, 1967), and the northeastern Gulf (Chace, 1956).

Menippe nodifrons Stimpson, 1859 (Ann. Lyc. Nat. Hist. New York 7: 53)

Rathbun, 1930, p. 479, pl. 198, fig. 3, pl. 199. 2

Range: east coast of Florida; ? Louisiana; north and south coasts of Cuba; Jamaica; Virgin Islands; Trinidad; Caribbean coasts of Panama and Colombia; Paraiba to São Francisco do Sul, Brazil; Gabon, West Africa.

Depth: shallow water, near shore.

Habitat: tide pools; under rocks; near dock pilings; on sponges and among bryozoans.

Remarks: Rathbun's (1930) record of this crab from Louisiana has been questioned by Felder (1973a) because extensive collecting along the northwestern Gulf coast has produced no further specimens. The only other Gulf record would be that of a single male collected near Havana, Cuba (Rathbun, 1930).

Micropanope Stimpson, 1871

(As restricted by Guinot (1967a, 1968b), this genus would include only the type species, *M. sculptipes* Stimpson, and one other, leaving a number of species formerly in *Micropanope* without an apparent generic name. Following a suggestion of Garth (pers. comm.), this list retains these orphaned species, not included in other genera by Guinot, as members of *Micropanope*, pending further clarification of their taxonomic status.)

Micropanope barbadensis (Rathbun, 1921) (Bull. Lab. Nat. Hist. State Univ. Iowa 9: 73), *novo comb.* Rathbun, 1930 (Bull. U.S. Nat. Mus. 152: 446) 2

Rathbun, 1930, p. 446, text-fig. 72. 10

Range: Dry Tortugas; Barbados.

Depth: shallow water.

Habitat: from coral heads.

Micropanope distincta (Rathbun, 1898) (Bull. Lab. Nat. Hist. State Univ. Iowa 4: 272)

As *Lophopanopeus distinctus*—Rathbun, 1930, p. 331, pl. 155, figs. 1-2.

As *Micropanope distincta*—Menzies, 1948, p. 24.

Range: North Carolina; Florida Straits; Dry Tortugas; northwest Florida; Barbados.

Depth: 48 to 185 m (26 to 101 fm).

Habitat: sand and coral bottoms.

Remarks: Menzies (1948) transferred this species to *Micropanope* and he considered it to be close to *M. sculptipes* Stimpson, the type species of the genus. Cerame-Vivas and Gray (1966) extended the known range (listed as *Lophopanopeus*) to North Carolina.

Micropanope lobifrons A. Milne Edwards, 1880 (Crust. Rég. Mex., p. 327)

Rathbun, 1930, p. 429, pl. 178, figs. 4-6; Rathbun, 1933, p. 66.

Range: south Florida, in Gulf Stream; Dry Tortugas; off northwest Florida; off north coast of Cuba; Puerto Rico; Virgin Islands; Santa Cruz Island (Carib.); Grenada; Barbados; Colon, Panama.

Depth: 37 to 311 m (20 to 170 fm).

Habitat: sand, coral, rock, and broken shell substrates.

Remarks: Guinot (1968b) retained this species in *Micropanope* along with the type species, while revising the genus.

Micropanope nuttingi (Rathbun, 1898) (Bull. Lab. Nat. Hist. State Univ. Iowa 4: 271), novo comb. Rathbun, 1930 (Bull. U.S. Nat. Mus. 152: 450)

Rathbun, 1930, p. 450, text-fig. 74; Rathbun, 1933, p. 67, fig. 57; Williams, 1965, p. 194, figs. 177, 183J; Felder, 1973a, p. 66, pl. 9, fig. 22.

Range: North Carolina; east coast of Florida; Florida Keys and Dry Tortugas; west and northwest coasts of Florida; Texas; north coast of Yucatan; north coast of Cuba; Jamaica; Puerto Rico; Virgin Islands; Old Providence Island (Carib.); Rio Grande do Norte, Brazil.

Depth: shore to 183 m (to 100 fm).

Habitat: rock, sand, coral, and broken-shell substrates; from boulder-covered beaches; from clumps of *Porites* and *Halimede*.

Remarks: Felder (1973a) notes that this species may be eventually placed in a different genus on the basis of differences in male first pleopods. Williams (1965) listed ovigerous females from Florida in July.

Micropanope pusilla A. Milne Edwards, 1880 (Crust. Rég. Mex., p. 327)

Rathbun, 1930, p. 431, pl. 179, figs. 7-8; Rathbun, 1933, p. 66.

Range: Dry Tortugas; northwest of Key West; west and northwest coasts of Florida; Alabama; north coast of Cuba; Jamaica; Puerto Rico; Virgin Islands.

Depth: 31 to 311 m (17 to 170 fm).

Habitat: sand, gravel, coral, and broken shell bottoms.

Remarks: Listed from Florida by Wass (1955), Abele (1970), and Menzel (1971). Rathbun (1930) reported ovigerous females from the Gulf of Mexico during February-March and June-July. Abele (1970) noted that this species does not fit any of the genera reviewed or erected by Guinot (1968a).

Micropanope sculptipes Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 140)

Rathbun, 1930, p. 428, pl. 178, figs. 1-3; Rathbun, 1933, p. 66; Williams, 1965, p. 193, fig. 175; Felder, 1973a, p. 66, pl. 9, fig. 15.

Range: North and South Carolina; Florida Keys and Dry Tortugas; west and northwest coasts of Florida; Texas; St. Croix; Grenada; Barbados.

Depth: 9 to 311 m (5 to 170 fm).

Habitat: sand, gravel, coral, and broken shell bottoms.

Remarks: Listed from Florida by Wass (1955). Cerame-Vivas and Gray (1966) extended the known range of this crab to North Carolina.

Micropanope spinipes A. Milne Edwards, 1880 (Crust. Rég. Mex., p. 326)

Rathbun, 1930, p. 443, text-fig. 71, pl. 181, figs. 1-2; Pequegnat & Ray, 1974, p. 238, figs. 18-22.

Range: Bermuda; Bahamas; Florida Keys; West Flower Garden Bank, off Texas; Curaçao; Alagoas and off the Abrolhos Islands, Brazil.

Depth: low tide mark to 55 m (to 30 fm).

Habitat: sand and coral bottoms; off coral reefs; in sponges.

Micropanope truncatifrons Rathbun, 1898 (Bull. Lab. Nat. Hist. State Univ. Iowa 4: 274)

Rathbun, 1930, p. 433, text-fig. 68, pl. 178, figs. 7-8.

Range: off north coast of Cuba; off Caribbean coast of Yucatan.

Depth: 238 to 355 m (130 to 194 fm).

Habitat: coral sand bottoms.

Remarks: This species may be eventually transferred to the genus *Nanocasiope*. Guinot (1971, p. 1076) noted a resemblance between this species and *N. melanodactylus* (A. Milne Edwards), the type species of the genus recently erected by Guinot (1967a).

Micropanope urinator (A. Milne Edwards, 1881) (Crust. Rég. Mex., p. 289)

Rathbun, 1930, p. 451, pl. 182, figs. 3-4, pl. 183, figs. 1-3; Chace, 1940, p. 34; Williams, McCloskey & Gray, 1968, p. 51, fig. 7.

Range: off North Carolina; Florida Keys; north and south coasts of Cuba; St. Croix.

Depth: 146 to 457 m (80 to 250 fm).

Habitat: sand and coral bottoms.

Micropanope xanthiformis (A. Milne Edwards, 1880).

Transferred to a new genus, *Nanoplax*, by Guinot (1967a). See *Nanoplax xanthiformis*.

Nanoplax Guinot, 1967***Nanoplax xanthiformis*** (A. Milne Edwards, 1880) (Crust. Rég. Mex., p. 353)As *Micropanope xanthiformis*—Rathbun, 1930, p. 442, pl. 180, figs. 7–8; Rathbun, 1933, p. 67; Williams, 1965, p. 193, figs. 176, 183I.As *Nanoplax xanthiformis*—Guinot, 1967a, p. 362, fig. 16.

Range: North Carolina to off south Florida; Florida Keys and Dry Tortugas; northwest coast of Florida; north coast of Cuba; off Caribbean coast of Yucatan; Puerto Rico; Dominica; Barbados; Grenada; Curaçao; off Cape Frio, Brazil.

Depth: 9 to 333 m (5 to 182 fm).

Habitat: sand, broken shell, coral, and mud substrates.

Remarks: Guinot (1967a) discussed the affinities of this species and genus with certain genera of Goneplacidae. Listed from northwest Florida (as *Micropanope*) by Wass (1955) and Hulings (1961). Ovigerous females are known from Florida in June and August and from North Carolina in October (Williams, 1965).***Neopanope*** A. Milne Edwards, 1880***Neopanope packardii*** (Kingsley, 1879) (Proc. Boston Soc. Nat. Hist. 20: 152)

Rathbun, 1930, p. 380, text-fig. 59, pl. 168, figs. 5–6; Abele, 1972a, p. 269, figs. 1B, 3A.

Range: southeast and south Florida; Bahamas; Florida Keys and Dry Tortugas; west and northwest coasts of Florida; Louisiana; north coast of Cuba.

Depth: low tide mark to 74 m (to 135 fm).

Habitat: sand, gravel, rock, and coral substrates; occasionally in mud; grass beds, algal beds, marshes; on *Styela*.Remarks: A systematic review and morphological key for this genus was presented by Abele (1972a). Abele (1971) provided scanning electron micrographs of the gonopods of this crab and other *Neopanope*. Larval development was described by Costlow and Bookhout (1967); larvae of various *Neopanope* species were compared by McMahan (1967). Records of ovigerous crabs were listed by Rathbun (1930). Listed from Florida by Tabb and Manning (1961), Dragovich and Kelly (1964), Abele (1970), Rouse (1970), Menzel (1971), and Lyons *et al.* (1971), and from Louisiana by Hoese and Valentine (1972).***Neopanope texana*** (Stimpson, 1859) (Ann. Lyc. Nat. Hist. New York 7: 55)As *N. texana texana*—Rathbun, 1930, p. 367, text-fig. 57, pl. 168, figs. 1–2.As *N. texana*—Abele, 1972a, p. 266, figs. 1A, 2A, 2E, 3B, 3C; Felder, 1973a, p. 68, pl. 9, fig. 19; not Williams, 1965, p. 190 (= *N. sayi*).

Range: west coast of Florida (south as far as Charlotte County) to south Texas.

Depth: low tide mark to 51 m (to 28 fm).

Habitat: *Thalassia* grass flats; mud, sand, rock, and gravel substrates; among barnacles, clumps of ascidians.Remarks: Abele (1972a) reviewed the status of the two related species, *N. texana* and *N. sayi*. The latter form is restricted in distribution to the east coast

of the United States, whereas *N. texana* occurs only along the Gulf coast. Reports of *N. texana* in southern Florida (Tabb and Manning, 1961) are attributed by Abele to *N. packardii*. Williams (1965) listed this species from North Carolina, but Abele (1972a) states that the figures are of *N. sayi*. Florida listings include Wass (1955), Dragovich and Kelly (1964), Abele (1970), Lyons *et al.* (1971), and Menzel (1971). Other listings (some as *N. texana sayi* (= *N. sayi*), but all are referred to *N. texana*) include the northwestern Gulf (Fotheringham and Brunenmeister, 1975) and Texas (Simmons, 1957; Parker, 1959; Hoese, 1960; Breuer, 1962; Hoese and Jones, 1963; Keith and Hulings, 1965; Leary, 1967). McMahan (1967) described larvae reared in the laboratory. Landers (1954) noted predation by this species on clams. Ryan (1956) gave accounts of life history for *N. sayi*, which should be very similar to that of *N. texana*.

Panopeus H. Milne Edwards, 1834

Panopeus americanus Saussure, 1857 Rev. Mag. Zool., ser. 2, vol. 9: 502)

Rathbun, 1930, p. 357, pl. 164, figs. 3-4, 6; Rathbun, 1933, p. 62.

Range: Bahamas; Florida Keys; west coast of Florida; north coast of Cuba; Jamaica; Dominican Republic; Puerto Rico; St. Thomas, Virgin Islands; Guadeloupe; Trinidad; Caribbean coast of Colombia; Rio Parahyba do Norte to Santa Catarina, Brazil.

Depth: intertidal to 22 m (to 12 fm).

Habitat: under stones, on mud flats; on mangroves; sand, shell, and mud bottoms; grass flats; under sponges and bryozoans.

Remarks: Listed from south Florida by Tabb and Manning (1961).

Panopeus bermudensis Benedict & Rathbun, 1891 (Proc. U.S. Nat. Mus. 14: 376)

Rathbun, 1930, p. 360, text-fig. 56, pl. 165; Rathbun, 1933, p. 62; Garth, 1961, p. 149; Felder, 1973a, p. 69, pl. 9, fig. 20.

Range: Bermuda; Bahamas; west coast of Florida; ?Texas; north coast of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands; Trinidad; Old Providence Island (Carib.); Colombia to Santa Catarina, Brazil; in Pacific—from Magdalena Bay, Mexico to Peru.

Depth: low tide mark to 15 m (to 8 fm).

Habitat: oyster beds; rocky tide pools; under rocks, sponges, bryozoans, debris, and among fouling organisms; from bays and near-marine waters; on mangrove roots; from sand bottoms.

Remarks: Felder (1973a) listed this species from Texas on the basis of a tentative identification. Garth (1961) noted the variability of this species over its entire geographical range and pointed to the possibility of future segregation of this species into distinct Pacific and Atlantic forms.

Panopeus hartii Smith, 1869 (Proc. Boston Soc. Nat. Hist. 12: 280)

Rathbun, 1930, p. 355, pl. 164, figs. 1-2, 5; Rathbun, 1933, p. 62.

Range: Florida Keys and Dry Tortugas; Isla de Piños, Cuba (Carib.); Puerto

Rico; St. Thomas, Virgin Islands; Antigua; Barbados; Pernambuco to São Paulo, Brazil.

Depth: low tide mark and shallow waters.

Habitat: on rocks and coral reefs.

Remarks: Color and habitat in Brazil were described by Fausto Filho (1974).

Panopeus herbstii H. Milne Edwards, 1834 (Hist. Nat. Crust., vol. 1, p. 403)

Common Name: Common Mud Crab

Hay & Shore, 1918, p. 437, pl. 34, fig. 9; Rathbun, 1930, p. 335, text-figs. 52-53, pls. 156-157; Chace, 1940, p. 34; Ryan, 1956, p. 147, text-figs. 4B, 5B, 9A, pl. 1C; Edmondson, 1962, p. 277; Williams, 1965, p. 196, figs. 180, 183M; Chace & Hobbs, 1969, p. 154, figs. 46c, 47; Felder, 1973a, p. 69, pl. 9, fig. 21.

Range: Bermuda; Bahamas; Massachusetts to south Florida; Florida Keys and Dry Tortugas; west coast of Florida to Veracruz, Mexico; north and south coasts of Cuba; Jamaica; Haiti; Puerto Rico; Virgin Islands; Antigua to Barbados; Trinidad; Islas Los Roques; Netherlands Antilles; Belize; Caribbean coast of Panama to Venezuela; Rio Parahyba do Norte, Brazil to Uruguay; in Pacific—Hawaiian Islands.

Depth: intertidal to 22 m (12 fm).

Habitat: muddy bottoms of bays and estuaries; shell, rock, and stone bottoms; oyster beds; among mangroves and in banks of tidal streams; burrows in mud banks and under stones or shells; on coral and rock reefs; brackish tide pools; sandy beaches with rocks.

Remarks: Larval development has been described by Costlow and Bookhout (1961a) and Costlow, Bookhout and Monroe (1962). Sulkin (1973) reported on larval depth regulation and Sandifer (1973) gathered data on larval ecology in Chesapeake Bight. Schwartz and Cargo (1960) recorded this crab in Virginia and Maryland. Life history studies include those of Ryan (1956) in Chesapeake Bay and Warner (1969) in Jamaica. Rathbun (1930) listed three forms, in addition to the typical form, of this species and provided distribution records for each. Furtado-Ogawa (1972) noted individual variations and habitat differences for this species in Brazil. Williams (1965) provided a summary of the many studies on this crab. Relationships with oysters and role as a molluscan predator are included in reports by McDermott and Flower (1953), Ryan (1956), Menzel and Nichy (1958) and McDermott (1960). Perkins (1975) describes the fine structure of a haplosporid parasite of this crab and Humes (1941a) noted the presence of a parasitic barnacle. Forward (1977) studied shadow responses of the larvae. Physiological studies include reports on antennule chemosensitivity (Hazlett, 1971), coagulation (Morrison and Morrison, 1952), thoracic neurosection (Maynard, 1961a, 1961b; Maynard and Maynard, 1962), gill area (Gray, 1957), respiration in relation to habitat (Ayres, 1938; Teal, 1959), amino acid metabolism (Boone and Claybrook, 1977), and respiration and metabolism (Leffler, 1973). Regional lists include Florida (Wass, 1955; Dragovich and Kelly, 1964; Abele, 1970; Menzel, 1971; Subrahmanyam *et al.*, 1976), Mississippi (Richmond, 1968; Christmas and Langley, 1973), Louisiana (Behre, 1950; Hoese and Valentine, 1972), Texas (Leary, 1967), and the northwestern Gulf of Mexico (Fotheringham and Brunenmeister, 1975).

Panopeus occidentalis Saussure, 1857 (Rev. Mag. Zool., ser. 2, vol. 9: 502)

Rathbun, 1930, p. 348, text-fig. 55, pl. 161; Rathbun, 1933, p. 61; Williams, 1965, p. 198, figs. 181, 183N.

26 Range: Bermuda; Bahamas; North Carolina to southeast Florida; Florida Keys and Dry Tortugas; west coast of Florida; north coast of Cuba; Jamaica; Hispaniola; Puerto Rico; Virgin Islands; Guadeloupe; Old Providence Island (Carib.); Curaçao; Trinidad; Colon, Panama to Santa Catarina, Brazil.

Depth: intertidal to 18 m (to 10 fm).

Habitat: sand, shell, rock, and gravel bottoms; among ascidians, sponges, and seaweed; on mangrove roots; under rocks; on pilings and piers.

Remarks: Although both of the environmental forms listed by Rathbun (1930) were recorded from Louisiana by Behre (1950), Felder (1973a) doubts the validity of the Grand Isle records and believes that they may represent *P. herbstii*. Williams (1965) provided a good summary of data on this crab. Listed from Florida by Tabb and Manning (1961) and by Lyons *et al.* (1971). De Oliveira (1940) described specimens from Brazil and provided notes on life history; Furtado-Ogawa (1972) noted individual variations and habitat differences in Brazil.

Panopeus rugosus A. Milne Edwards, 1880 (Crust. Rég. Mex., p. 314)

Rathbun, 1930, p. 353, pls. 162-163.

26? Range: Florida Keys and Dry Tortugas; west and northwest coasts of Florida; north coast of Cuba; Haiti; Virgin Islands; Puerto Rico; Honduras to Nicaragua; Curaçao; Bahia to Santa Catarina, Brazil.

Depth: low tide mark to 51 m (to 28 fm).

Habitat: sand, shell, rock, and coral bottoms; coral reefs; on pilings.

Remarks: listed from northwest Florida by Wass (1955)

Panopeus turgidus Rathbun, 1930 (Bull. U.S. Nat. Mus. 152: 364)

Rathbun, 1930, p. 364, pl. 166; Felder, 1973a, p. 68, pl. 9, fig. 18.

27 Range: northwest coast of Florida to Texas.

Depth: near shore, shallow waters.

Habitat: bay and near-marine waters; found on or among rocks, shells, debris, and vegetation.

Remarks: Listed from northwest Florida by Wass (1955) and from Louisiana by Behre (1950). Abele (1970) collected a specimen from northwest Florida and Fenner Chace, Jr. of the USNM confirmed its identity by comparison with the type, suggesting it was similar to the genus *Eurypanopeus*. Abele (1970) listed it in his thesis as *Eurypanopeus turgidus* and this was repeated in the list by Menzel (1971). As indicated under this latter name, this present listing will continue to use *Panopeus turgidus* until a definitive study or revision is available.

Paractaea Guinot, 1969***Paractaea rufopunctata nodosa*** (Simpson, 1860) (Ann. Lyc. Nat. Hist. New York 7: 203)

As *Actaea rufopunctata nodosa*—Rathbun, 1930, p. 257, pl. 105, figs. 1-2;

Rathbun, 1933, p. 56; Holthuis & Gottlieb, 1956, p. 287; Williams, McCloskey & Gray, 1968, p. 51.

As *Paractaea rufopunctata* forma *nodosa*—Guinot, 1969b, p. 252, fig. 25.

Range: North Carolina; Bahamas; east coast of Florida; Florida Keys and Dry Tortugas; north coast of Cuba; Jamaica; Haiti; Puerto Rico; Virgin Islands; Barbados; Curaçao; Cape Frio, Brazil; Ascension Island, South Atlantic.

Depth: 5 to 212 m (3 to 115 fm).

Habitat: coral, broken shell, stone, and sand bottoms; in sponges.

Remarks: This genus was erected by Guinot (1969b) for several species of *Actaea*. She compared the differences and similarities of this form and forma *africana* and other forms of the *rufopunctata* complex, deciding that further research was needed to clarify the systematic relationships of the genus.

Paraliomera Rathbun, 1930

Paraliomera dispar (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 140)

Rathbun, 1930, p. 244, text-fig. 38, pl. 101, figs. 4-5; Rathbun, 1933, p. 54.

Range: Bermuda; Florida Keys and Dry Tortugas; north coast of Cuba; Jamaica; Haiti; Puerto Rico; Antigua; Barbados; Curaçao; Caribbean coast of Colombia.

Depth: shallow water to 154 m (to 84 fm).

Habitat: sand, shell, grassy, and mud bottoms; coral reefs and rocky areas.

Paraliomera longimana (A. Milne Edwards, 1865) (Nouv. Arch. Mus. Hist. Nat., Paris 1: 221).

Rathbun, 1930, p. 243, pl. 101, figs. 1-3; Rathbun, 1933, p. 53, fig. 46.

Range: Florida Keys and Dry Tortugas; Veracruz, Mexico; Puerto Rico; Virgin Islands; Barbados; Curaçao.

Depth: shallow water to 154 m (84 fm).

Habitat: coral reefs and rocky and grassy substrates.

Phymodius A. Milne Edwards, 1863

Phymodius maculatus (Stimpson, 1860).

This species was transferred to *Etisus* by Guinot (1969b), thus leaving the genus *Phymodius* unrepresented in the Gulf of Mexico.

Pilumnoides H. Milne Edwards & Lucas, 1843

Pilumnoides nudifrons (Stimpson, 1871) (Bull. Mus. Comp. Zool. 2: 143)

Rathbun, 1930, p. 538, pl. 218, figs. 1-2.

Range: Florida Straits and Keys; Barbados.

Depth: 128 to 556 m (70 to 304 fm).

Habitat: sand and rock substrates.

Pilumnus Leach, 1815***Pilumnus caribaeus*** Desbonne & Schramm, 1867 (Crust. Guadeloupe, p. 32)

Rathbun, 1930, p. 491, pl. 200, figs. 3-4; Rathbun, 1933, p. 71.

Range: Bahamas; Florida Keys; north coast of Cuba; Jamaica; Puerto Rico; Vieques and Culebra; Virgin Islands; Guadeloupe; Curaçao; Bahia to São Paulo, Brazil.

Depth: low tide mark to 29 m (to 16 fm).

Habitat: mud, sand, shell, grassy, and coral bottoms.

Pilumnus dasypodus Kingsley, 1879 (Proc. Boston Soc. Nat. Hist, 20: 155)

Rathbun, 1930, p. 493, pl. 200, figs. 5-6; Rathbun, 1933, p. 72; Williams, 1965, p. 178, figs. 157C, 159; Felder, 1973a, p. 61, pl. 9, fig. 7.

Range: North and South Carolina; Florida Keys and Dry Tortugas; south to northwest Florida; Mississippi; Texas; north and south coasts of Cuba; Jamaica; Puerto Rico; Culebra; Virgin Islands; Martinique; Curaçao; Pernambuco to Santa Catarina, Brazil.

Depth: 1 to 29 m (0.5 to 16 fm).

Habitat: sand, shell, rocky, and coral bottoms; from mangrove roots; with fouling materials on pilings, buoys and jetties; off sponges.

Remarks: Sandifer (1974) studied larval development. Williams (1965) reviewed records of ovigerous females. Ecological notes were provided by Lunz (1937), Pearse (1934), and Pearse and Williams (1951). Listed from Florida (Wass, 1955; Abele, 1970; Menzel, 1971; Lyons *et al.*, 1971), Mississippi (Christmas and Langley, 1973) and from Texas (Leary, 1967).

Pilumnus diomedae Rathbun, 1894 (Proc. U.S. Nat. Mus. 17: 85)

Rathbun, 1930, p. 501, pl. 202, figs. 2-3.

Range: north coast of Cuba; Caribbean coast of Yucatan.

Depth: 238 to 337 m (130 to 184 fm).

Habitat: coral and sand bottoms.

Pilumnus floridanus Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 141)

Rathbun, 1930, p. 507, pl. 205, figs. 3-4; Rathbun, 1933, p. 72; Williams, 1965, p. 179, figs. 157D, 160; Felder, 1973a, p. 61, pl. 9, fig. 8; Pequegnat & Ray, 1974, p. 238, figs. 23-24.

Range: North Carolina; Bahamas; Florida Keys and Dry Tortugas; west and northwest coasts of Florida; off Texas; north of Yucatan; Honduras; Jamaica; Puerto Rico; Culebra; Virgin Islands; Venezuela.

Depth: low tide mark to 146 m (to 80 fm).

Habitat: sand, shell, gravel, rock, and coral substrates; mud and grassy bottoms; inside sponges.

Remarks: Listed from the Gulf by Chace (1956) and from Florida by Wass (1955) and Lyons *et al.* (1971). Williams (1965) reported ovigerous females from Florida in March to August and from North Carolina in February. Pearse and Williams (1951) collected this crab from sponges in North Carolina waters.

Pilumnus gemmatus Stimpson, 1860 (Ann. Lyc. Nat. Hist. New York 7: 214)

Rathbun, 1930, p. 513, pl. 207, figs. 1-3; Rathbun, 1933, p. 72.

Range: Dry Tortugas; Culebra; Virgin Islands; Curaçao.

Depth: shore to 42 m (to 23 fm).

Habitat: shallow water lagoons; coral and rock bottoms; in corals; on seawoods.

Pilumnus holosericus Rathbun, 1898 (Bull. Lab. Nat. Hist. State Univ. Iowa 4: 268)

Rathbun, 1930, p. 519, text-fig. 81, pl. 207, figs. 8-9; Rathbun, 1933, p. 73, fig. 61.

Range: Bahamas; Dry Tortugas; Puerto Rico; Virgin Islands; Trinidad; Curaçao.

Depth: shallow water.

Habitat: near shore, under stones; coral reefs; rocky areas.

Pilumnus lacteus Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 142)

Hay & Shore, 1918, p. 440, pl. 35, fig. 3; Rathbun, 1930, p. 511, pl. 205, figs. 1-2; Williams, 1965, p. 180, figs. 157E, 161.

Range: North and South Carolina; Florida Keys and Dry Tortugas; west coast of Florida; north coast of Cuba.

Depth: low tide mark to 15 m (8 fm).

Habitat: sand, shell, rock, coral, and mud substrates; among sponges and seaweed; under stones; on buoys and pilings; in beds of *Thalassia*.

Remarks: Williams (1965) listed records of ovigerous females; Lunz (1937) provided notes on ecology of South Carolina populations, listed from Florida by Wass (1955), Tabb and Manning (1961) and Lyons *et al.* (1971). Rouse (1970) found this crab to be the most common pilumnid collected in Florida Bay.

Pilumnus longleyi Rathbun, 1930 (Bull. U.S. Nat. Mus. 152: 502)

Rathbun, 1930, p. 502, pl. 202, figs. 4-5.

Range: Bahamas; Florida Keys and Dry Tortugas.

Depth: shallow water.

Habitat: rocky and coral bottoms.

Remarks: Rathbun (1930) noted that this species is easily confused with *P. caribaeus* and *P. sayi*; she compared the three species morphologically and described the young of *P. longleyi*.

Pilumnus marshi Rathbun, 1901 (Bull. U.S. Fish. Comm. for 1900. vol. 20, pt. 2, p. 41)

Rathbun, 1930, p. 499, text-fig. 80; Rathbun, 1933, p. 72.

Range: Dry Tortugas; Virgin Islands.

Depth: shallow water to 37 m (to 20 fm).

Habitat: coral bottoms.

Pilumnus pannosus Rathbun, 1896 (Proc. U.S. Nat. Mus. 19: 142)

Rathbun, 1930, p. 514, pl. 207, figs. 4-5; Rathbun, 1933, p. 72; Williams, 1965, p. 181, figs. 157F, 162; Felder, 1973a, p. 64, pl. 9, fig. 12.

Range: North Carolina; Bahamas; southeast Florida; Florida Keys; west and northwest coasts of Florida; Texas; Puerto Rico; Jamaica; Virgin Islands.

Depth: 1 to 16 m (to 9 fm).

Habitat: sand, shell, rock, and coral substrates; on jetties and reefs with fouling material; with sponges and corals.

Remarks: Williams (1965) listed records of ovigerous females. Listed from Florida by Wass (1955) and from Texas by Leary (1967). Pearse and Williams (1951) collected this crab from submerged reefs off North Carolina.

Pilumnus sayi Rathbun, 1897 (Ann. Inst. Jamaica 1: 15)

Common Name: Hairy Crab

Hay & Shore, 1918, p. 440, pl. 35, fig. 4; Rathbun, 1930, p. 484, pl. 200, figs. 1-2, pl. 201, figs. 4-7; Rathbun, 1933, p. 71; Williams, 1965, p. 177, figs. 157A-B, 158; Felder, 1973a, p. 61, pl. 9, fig. 6.

Range: North Carolina to Georgia; Bahamas; east coast of Florida; Dry Tortugas; west and northwest coasts of Florida; Louisiana and Texas; Jamaica; Puerto Rico; Guadeloupe; Curaçao.

Depth: low tide mark to 90 m (to 49 fm).

Habitat: sand, shell, rock, coral, and gravel bottoms; on offshore reefs; among fouling material on jetties, pilings, and buoys.

Remarks: Listed from Florida by Wass (1955), Dragovich and Kelly (1964), Abele (1970), Menzel (1971) and Lyons *et al.* (1971), the latter group remarking on the considerable variation in morphology of this species. Lunz (1939) reported collecting *P. marshi* off a shoal in North Carolina, but Williams (1965) believed this to be an aberrant *P. sayi*. Pearse (1934) reported this crab from inside a sponge. Collected by Chace (1956) in the northeastern Gulf. Chamberlain (1961) studied the physiological ecology of larval and megalops stages.

Pilumnus spinosissimus Rathbun, 1898 (Bull. Lab. Nat. Hist. State Univ. Iowa 4: 265)

Rathbun, 1930, p. 494, text-fig. 79, pl. 200, figs. 7-8.

Range: Florida Keys and Dry Tortugas.

Depth: 5 to 11 m (3 to 6 fm).

Habitat: from rocks, shell, and coral substrates.

Platyactaea Guinot, 1967

Platyactaea setigera (H. Milne Edwards, 1834) (Hist. Nat. Crust., vol. 1, p. 390)

As *Actaea setigera*—Rathbun, 1930, p. 251, pl. 103; Rathbun, 1933, p. 55.

As *Platyactaea setigera*—Guinot, 1967b, p. 561, fig. 36.

Range: Bermuda; Bahamas; Florida Keys and Dry Tortugas; north coast of Cuba; Jamaica; Puerto Rico; Virgin Islands; Antigua; Barbados; Trinidad; Curaçao; Caribbean coast of Colombia.

Depth: intertidal and shallow water.

Habitat: coral reefs; among and under rocks.

Remarks: Guinot (1967b) placed this species in a new genus and discussed the systematic affinities of related species.

Platypodia Bell, 1835

Platypodia spectabilis (Herbst, 1794).

This species was transferred to a new genus, *Platypodiella*, by Guinot (1967b), leaving this genus without a representative in the Gulf of Mexico.

Platypodiella Guinot, 1967

Platypodiella spectabilis (Herbst, 1794) (Natur. Krabben u. Krebse, vol. 2, p. 153)

As *Platypodia spectabilis*—Rathbun, 1930, p. 247, text-fig. 39, pl. 102, fig. 4; Rathbun, 1933, p. 54, fig. 47.

As *Platypodiella spectabilis*—Guinot, 1967b, p. 562; Felder, 1973a, p. 65, pl. 9, fig. 10.

Range: Bermuda; Bahamas; Florida Keys; Texas; Veracruz, Mexico, Jamaica; Puerto Rico; Virgin Islands; Guadeloupe; Martinique; Barbados; Curaçao; Fernando de Noronha, Brazil.

Depth: 4 to 13 m (2 to 7.5 fm).

Habitat: coral reefs; under stones.

Remarks: This species is illustrated by Forest & Guinot (1966, fig. 7). Listed from Brazil and described by Fausto Filho (1974). Guinot (1967b) transferred this species from *Platypodia* to the newly-erected genus.

Pseudomadaeus Guinot, 1967

Pseudomadaeus agassizii (A. Milne Edwards, 1880) (Crust. Rég. Mex., p. 270)

As *Leptodius agassizii*—Hay & Shore, 1918, p. 441, pl. 34, fig. 6; Rathbun, 1930, p. 307, pl. 141, figs. 4-5; Williams, 1965, p. 192, figs. 174, 183H.

As *Pseudomadaeus agassizii*—Guinot, 1968a, p. 726, fig. 25; Felder, 1973a, p. 67, pl. 9, fig. 11.

Range: North Carolina; Florida Straits and Keys; Dry Tortugas; west and northwest coasts of Florida; Louisiana and Texas; Virgin Islands.

Depth: 7 to 82 m (4 to 45 fm).

Habitat: sand, shell, rock, and coral bottoms; from sponges.

Remarks: Costlow and Bookhout (1968a) described larval development. Williams (1965) noted morphological variability over the range of this species and reported ovigerous females from April to November. Listed from Florida by Wass (1955) and Abele (1970). Abele (1970) commented on variation in anteriolateral teeth and tuberculation of the chelae.

Rhithropanopeus Rathbun, 1898

Rhithropanopeus harrisii (Gould, 1841) (Rept. Invert. Massachusetts, p. 326)

Hay & Shore, 1918, p. 441, pl. 35, fig. 5; Rathbun, 1930, p. 456, text-fig. 75, pl. 183, figs. 7-8; Williams, 1965, p. 187, figs. 169, 183C; Christiansen, 1969, p. 81, fig. 33, map 27; Felder, 1973a, p. 67, pl. 9, fig. 14.

Range: New Brunswick to south Florida; west and northwest coasts of Florida;

Mississippi to Veracruz, Mexico; introduced to west coast of United States and to various parts of Europe in recent times.

Depth: intertidal to 37 m (to 20 fm), most common to 9 m (5 fm).

Habitat: freshwater and estuarine areas; upper bays with salinities usually less than 20 ppt; on oyster reefs; on living and dead vegetation; under rocks and in old cans and other debris.

Remarks: Ryan (1956) provided an account of the life history of this crab, based on studies in Chesapeake Bay. Williams (1965) reviewed much of the pertinent literature on this species. Studies of larval forms include those of Connolly (1925), Hood (1962), Chamberlain (1962), Costlow (1966), Costlow, Bookhout and Monroe (1966), Bookhout *et al* (1972), Ott and Forward (1976), and Sandifer (1973).

The European populations were considered a separate subspecies, *R. h. tridentatus* Maitland, by Buitendijk and Holthuis (1949), but this was questioned by Wolff (1954). Christiansen (1969) reviewed the distributional data for this species in Europe and treated the different populations as conspecific, with no distinct subspecies recognized. Reports of this crab on the U.S. Pacific coast include Jones (1940), Felice (1958), and Ricketts and Calvin (1968). Regional lists include Florida (Wass, 1955; Tabb and Manning, 1961; Abele, 1970; Menzel, 1971), Mississippi (Christmas and Langley, 1973), Louisiana (Behre, 1950), Texas (Hedgpeth, 1953; Leary, 1967), and the northwestern Gulf (Fotheringham and Brunenmeister, 1975).

Physiological studies include data on osmoregulation (Jones, 1941; Verway, 1957; Kalber and Costlow, 1966; R. I. Smith, 1967), effects of pesticides on larval development (Bookhout and Costlow, 1976), metabolism and larval development (Rosenberg and Costlow, 1976), effects of juvenile hormone on larvae (Christiansen, Costlow and Monroe, 1977a, b), sterol synthesis in larvae (Whitney, 1969), androgen glands (Payen, Costlow and Charniaux-Cotton, 1971), and eyestalk hormones (Skorkowski, 1972).

Tetraxanthus Rathbun, 1898

Tetraxanthus bidentatus (A. Milne Edwards, 1880) (Crust. Rég. Mex., p. 353)

As *T. rugosus* Rathbun—Rathbun, 1930, p. 459, pl. 185.

As *T. bidentatus*—Chace, 1939, p. 52; Chace, 1940, p. 36.

Range: Florida Keys; north and south coasts of Cuba; Grenada.

Depth: 168 to 293 m (92 to 160 fm).

Habitat: sand and coral bottoms.

Remarks: The species described and illustrated by Rathbun (1930) as *T. bidentatus* was actually a new species, which Chace (1939) named *T. rathbunae* (= *T. rathbuni*). The species Rathbun (1930) listed as *T. rugosus* n. sp. is *T. bidentatus*.

Tetraxanthus rathbunae Chace, 1939 (Mem. Soc. Cubana Hist. Nat. 13: 52)

As *T. bidentatus* (A. Milne Edwards)—Rathbun, 1930, p. 458, pl. 184.

As *T. rathbunae*—Chace, 1939, p. 52; Chace, 1940, p. 37.

As *T. rathbuni*—Pequenat, 1970, p. 195.

Range: North Carolina; southeast Florida; Florida Keys; west coast of Florida; north and south coasts of Cuba; off Mississippi; Campeche, off Yucatan; off Grenada; off Cape Frio, Brazil.

Depth: 108 to 476 m (59 to 260 fm).

Habitat: mud, coral, shell, sand, and rock bottoms.

Remarks: This is the species described by Rathbun (1930) as *T. bidentatus* (A. Milne Edwards), but it is not *Xanthodes bidentatus* A. Milne Edwards, the original designation of the species *Tetraxanthus bidentatus*. Chace (1956, p. 19) listed this species from the Gulf of Mexico.

***Tetraxanthus rugosus* Rathbun, 1930.**

This is a junior synonym of *Xanthodes bidentatus* (= *Tetraxanthus bidentatus*), therefore Chace (1939, p. 52) substituted *T. rathbunae* for Rathbun's *T. bidentatus* and *T. rugosus* is synonymized with *T. bidentatus* (A. Milne Edwards).

***Xantho* Leach, 1814**

***Xantho denticulata* White, 1847 (List Crustacea British Mus., p. 17)**

As *Xanthodius denticulatus*—Rathbun, 1930, p. 314, pl. 314, pl. 145, fig. 1, pl. 146.

As *Xantho denticulata*—Monod, 1956, p. 280, figs. 335–339; Forest & Guinot, 1961, p. 60, fig. 51.

Range: Bermuda; Bahamas; Florida Keys and Dry Tortugas; northwest Florida; Jamaica; Puerto Rico; Virgin Islands; Antigua; Barbados; Colon, Panama; Curaçao; Trinidad; Pernambuco to Abrolhos Islands, Brazil, Gulf of Guinea, west coast of Africa.

Depth: near shore to 12 m (to 7 fm).

Habitat: tide pools; coral reefs; under rocks.

Remarks: Guinot (1968a, p. 711) commented on the relationships between species of this genus and other related genera, but she did not discuss *X. denticulata*, other than to raise the question of differences between specimens from west Africa and South America. Monod (1956) compared this crab with species of other genera, especially *Cycloxanthops*. Listed from northwest Florida by Abele (1970).

***Xanthodius* Stimpson, 1859**

***Xanthodius denticulatus* (White, 1847).**

Monod (1956) listed synonymies. This species was transferred to the genus *Xanthodius* by Rathbun (1930, p. 314); other authors have continued to treat it as a species of *Xantho*. Not *Xantho denticulata* Stimpson, 1860, a west coast species (= *Xanthodius stimpsoni* (A. Milne Edwards, 1879) proposed by Rathbun, 1930).

Family GERYONIDAE Colosi, 1924

(The systematic status of this family and of the Goneplacidae is still subject to revision by Guinot and others. Balss (1957) placed this genus

between the Xanthidae and Carcinoplacinae as a link to the Goneplacidae. Bouvier (1940) included *Geryon* in the Xanthidae, but Rathbun (1937), Sakai (1939) and Barnard (1950) treated it as a genus of the Goneplacidae. In the present list, only the type genus, *Geryon*, is listed in the family; *Bathyplox* is retained in the Goneplacidae until further studies are available (refer to Guinot, 1969a, 1971).

***Geryon* Krøyer, 1837**

***Geryon quinquedens* Smith, 1879** (Trans. Connecticut Acad. Arts Sci. 5: 35)

Common Name: Deep Sea Red Crab

Rathbun, 1937, p. 271, not pls. 85–86; Chace, 1940, p. 38; Pequegnat, 1970, p. 189, fig. 6–7; Wigley, Theroux & Murray, 1975, p. 1.

Range: Nova Scotia to South Carolina; east coast of Florida; Florida Straits; off Key West; Dry Tortugas; off Alabama and Texas; off northeast Mexico; north coast of Cuba; off Cape Frio, Brazil.

Depth: 40 to 2153 m (22 to 1178 fm), most common at 320 to 914 m (175 to 500 fm).

Habitat: primarily on mud and ooze substrates, occasionally on sand or shell-mud bottoms.

Remarks: Chace (1940) compared this species with *G. affinis*, which is very similar in appearance. Plates 85 and 86 in Rathbun (1937) are of *G. affinis*. This latter species has also been caught off the coast of Florida, but it is more common in southern areas of the Atlantic and in the Indian Ocean. Le Loeuff *et al.* (1974) recently reported *G. quinquedens* from the Ivory Coast of Africa. When alive, there are color differences between these two species (Schroeder, 1959). Leone (1951) used serological techniques to investigate the taxonomic status of this crab. Musick and McEachran (1972) reported it from depths of 168 m in Chesapeake Bight. Accounts of the red crab fishery include those of Schroeder (1959), McRae (1961), Meade and Gray (1973), Holmsen and McAllister (1974), Ganz and Herrmann (1975), and Wigley, Theroux and Murray (1975). The latter study contains details of substrate and temperature data, a review of the life history of this crab, and *in situ* bottom photographs of the animal. Gray (1969) also provided an account of the biology of this species. Haefner and Musick (1974) reported its occurrence in Norfolk Canyon. Pequegnat (1970) indicated a center of depth range at about 914 m (500 fm) within the Gulf of Mexico, but all of the crabs taken by the R/V *Alaminos* at depths greater than 1170 m (640 fm) were juveniles. Haefner (1977) investigated reproduction in females.

Family GONEPLACIDAE Macleay, 1838 (sensu Balss, 1957)

(Guinot (1969a) has proposed several revisions in the systematic relationships of this family, most of them based on presumed affinities with genera of the Xanthidae, from which the Goneplacidae may have been derived. In the present list, the Gulf genera are listed alphabetically, as

they were for the Xanthidae, without regard to subfamily alignment. The genus *Geryon* is treated under a separate family, Geryonidae.)

Bathylax A. Milne Edwards, 1880

Bathylax typhla A. Milne Edwards, 1880 (Bull. Mus. Comp. Zool. 8: 16)

Rathbun, 1918, p. 19, text-fig. 4, pl. 2; Rathbun, 1933, p. 77, fig. 67; Chace, 1940, p. 43; Williams, McCloskey & Gray, 1968, p. 52, fig. 8; Pequegnat, 1970, p. 192, figs. 6-9, 6-10.

Range: off North Carolina; west coast of Florida; Mississippi; Texas and Mexico; west coast of Cuba; St. Croix; St. Lucia; off Recife, Brazil.

Depth: 402 to 878 m (220 to 480 fm); at 1106 m (605 fm) off Cuba. Pequegnat (1970) determined the highest densities of crabs at 512 m (280 fm) in the Gulf of Mexico.

Habitat: mainly from muddy substrates; also from coral bottoms. Pequegnat (1970) reported that blackened specimens were commonly collected, apparently due to contact with natural oil seepage.

Remarks: Recorded from the Gulf of Chace (1956). Pequegnat (1970) considers this crab to be the most common deep-water species in the Gulf of Mexico; ovigerous females were collected in August, November, and December. Guinot (1971) listed this species under the Xanthidae and under the Goneplacidae because of the uncertainty of its systematic affinities.

Chacellus Guinot, 1969

Chacellus filiformis Guinot, 1969 (Bull. Mus. Nation. Hist. Nat. 41: 722)

Guinot, 1969, p. 722, figs. 135-136, pl. V, fig. 4.

Range: between Bahamas and east coast of Florida; off northwest Florida.
Depth: 183 to 223 m (100 to 122 fm).

Chasmocarcinus Rathbun, 1898

Chasmocarcinus cylindricus Rathbun, 1901 (Bull. U.S. Fish. Comm. for 1900, vol. 2: 10)

Rathbun, 1918, p. 59, text-figs. 28-29; Rathbun, 1933, p. 80, fig. 73; Chace, 1940, p. 49; Pequegnat, 1970, p. 195.

Range: Mississippi; off Louisiana; Campeche, off Yucatan; north and south coasts of Cuba; Jamaica; Puerto Rico.

Depth: 13 to 1906 m (7 to 1075 fm).

Habitat: mud bottoms; sand, rock, and coral substrates.

Chasmocarcinus mississippiensis Rathbun, 1931 (Proc. Biol. Soc. Washington 44: 71)

Rathbun, 1931b, p. 71; Felder, 1973a, p. 70, pl. 10, fig. 2.

Range: off coasts of Mississippi, Louisiana, and Texas.

Depth: 4 to 91 m (2 to 50 fm).

Habitat: sand and mud bottoms.

Remarks: Dawson (1966) reported this species from off Grand Isle, Louisiana; Franks *et al.* (1972) obtained a single specimen from off Mississippi at 50 fm. Felder (1973a) reported specimens from Padre Island, Texas.

Chasmocarcinus obliquus Rathbun, 1898 (Bull. Lab. Nat. Hist. State Univ. Iowa 4: 286)

Rathbun, 1918, p. 58, text-fig. 27, pl. 14, figs. 1-2; Chace, 1940, p. 48.

Range: southeast of Bahamas; north and south coasts of Cuba.

Depth: 177 to 503 m (97 to 275 fm).

Habitat: mud and ooze substrates.

Eucratopsis Smith, 1869

Eucratopsis crassimanus (Dana, 1852) (Proc. Acad. Nat. Sci. Philadelphia, for 1851, vol. 5: 248)

Rathbun, 1918, p. 52, text-fig. 22, pl. 12, fig. 3, pl. 159, figs. 1-2; Guinot, 1969a, p. 258, figs. 6, 10, 25.

Range: Florida Keys; south and west coasts of Florida; Yucatan; Jamaica; Bahia to Rio de Janeiro, Brazil.

Depth: shallow water to 14 m (to 7.5 fm).

Habitat: sand, coral, and broken shell substrates.

Remarks: Tabb and Manning (1961) collected ovigerous females in October from Oyster Bay in south Florida.

Euphosynoplax Guinot, 1969

Euphosynoplax clausa Guinot, 1969 (Bull. Mus. Nation. Hist. Nat. 41: 720)

Guinot, 1969a, p. 720, figs. 127, 139, pl. IV, fig. 3; Pequegnat, 1970, p. 194.

Range: Dry Tortugas; off Alabama and Mississippi; Campeche, Yucatan.

Depth: 91 to 210 m (50 to 115 fm).

Euryplax Stimpson, 1859

Euryplax nitida Stimpson, 1859 (Ann. Lyc. Nat. Hist. New York 7: 60)

Rathbun, 1918, p. 34, pl. 7; Rathbun, 1933, p. 78, fig. 69; Williams, 1965, p. 202, fig. 185; Guinot, 1969a, p. 512, figs. 39, 41, 47, 56-57, pl. II, fig. 1; Felder, 1973a, p. 70, pl. 10, fig. 1.

Range: off North Carolina; Bermuda; Florida Keys; south and west coasts of Florida to Texas; Puerto Rico; St. Thomas, Virgin Islands.

Depth: shallow water to 90 m (to 49 fm).

Habitat: sand, shell, rock, and coral substrates; sandy grass flats.

Remarks: Menzel (1971) listed this crab as rare at Apalachee Bay and Abele (1970) collected only a single specimen in his study of the northeastern near-shore Gulf. Range reports for the western Gulf of Mexico are scant and questionable. Rathbun (1918) listed one specimen from New Orleans and Williams (1965) indicated a range extending to Texas. Felder (1973a) cites these reports but adds no new records.

Frevillea A. Milne Edwards, 1880***Frevillea barbata*** A. Milne Edwards, 1880 (Bull. Mus. Comp. Zool. 8: 15)As ***Goneplax barbata***—Rathbun, 1918, p. 26, pl. 4, figs. 1, 3, pl. 5.As ***Frevillea barbata***—Guinot, 1969a, p. 513, pl. II, fig. 2.

Range: west coast of Florida; Yucatan (Gulf); north coast of Cuba; off Grenada.

Depth: 55 to 168 m (30 to 92 fm).

Habitat: sand, broken coral bottoms.

Frevillea hirsuta (Borradaile, 1916) (Brit. Antarctic Exped., 1910, Zool., vol. 3, no. 2, p. 99)As ***Goneplax hirsuta***—Rathbun, 1918, p. 28, text-fig. 7; Williams, 1965, p. 201, fig. 184.As ***Frevillea hirsuta***—Guinot, 1969a, p. 513, text-figs. 33, 40, 58–59, pl. II, fig. 3.

Range: off North Carolina; off west and northwest Florida and Alabama; north of Yucatan; off Rio de Janeiro, Brazil.

Depth: 73 to 146 m (40 to 80 fm).

Remarks: Collected by the R/V *Oregon* from the Gulf of Mexico (Chace, 1956).***Frevillea tridentata*** A. Milne Edwards, 1880.Transferred to the genus *Trapezioplax* by Guinot (1969a). *Trapezioplax tridentata*.***Glyptoplax*** Smith, 1870***Glyptoplax smithii*** A. Milne Edwards, 1880 (Crust. Rég. Mex., p. 336)

Rathbun, 1918, p. 51, pl. 13, figs. 3–4; pl. 158, figs. 7–10; Milne Edwards & Bouvier, 1923, p. 328, pl. 5, fig. 5; Williams, McCloskey & Gray, 1968, p. 55, fig. 11; Guinot, 1969a, p. 259.

Range: Bermuda; off North Carolina; west coast of Florida; Cape Catoche, Yucatan (Gulf coast).

Depth: 24 to 55 m (13 to 30 fm).

Habitat: sand, coral, gravel and rock bottoms.

Remarks: Guinot (1969a) believed that this species should be excluded from the genus, based on differences from the type species, *G. pugnax*, a Pacific form from Central America.***Goneplax*** Leach, 1814(All three of the recognized Gulf species of this genus have been transferred to other genera by Guinot (1969a). *Goneplax barbata* and *G. hirsuta* are referred to *Frevillea* and *G. tridentata* is referred to *Trapezioplax*.)***Neopilumnoplax*** Serene, 1969***Neopilumnoplax americana*** (Rathbun, 1898) (Bull. Lab. Nat. Hist. State Univ. Iowa 4: 283)As ***Pilumnoplax americana***—Rathbun, 1918, p. 21, text-figs. 5–6; Williams,

McCloskey & Gray, 1968, p. 52, fig. 9.

As *Neopilumnoplax americana*—Guinot, 1969a, p. 689, figs. 83–84.

Range: off North Carolina and Georgia; Florida Keys and Straits; north coast of Cuba; Guadeloupe; Espirito Santo, Brazil; Arabian Sea.

Depth: 128 to 805 m (70 to 440 fm).

Habitat: sand, shell, coral, and rocky substrates.

Remarks: Chace (1940) recovered this crab from the stomach of a smooth dogfish (*Mustelus canis*) from off Havana, Cuba. Listed from Brazil by Rodrigues da Costa (1968a) and Coelho and Ramos (1972).

Panoplax Stimpson, 1871

Panoplax depressa Stimpson, 1871 (Bull. Mus. Comp. Zool. 2: 151)

Rathbun, 1918, p. 47, text-fig. 21, pl. 12, figs. 1–2; Rathbun, 1933, p. 80, fig. 72;

Guinot, 1969a, p. 264, figs. 3, 12, 28, ? 29; Bright & Pequegnat, 1974, p. 33.

Range: Dry Tortugas; west coast of Florida; West Flower Garden Bank, off Texas; north coast of Cuba; north coast of Yucatan; Puerto Rico.

Depth: shallow water to 101 m (to 55 fm).

Habitat: sand, coral, and broken shell bottoms.

Remarks: Listed by Chace (1956) from off the west coast of Florida and by Bright and Pequegnat (1974) from silty-sand bottoms at West Flower Garden coral reef, at 330 foot depth.

Pilumnoplax Stimpson, 1858

Pilumnoplax americana Rathbun, 1898.

Transferred to the genus *Neopilumnoplax* by Guinot (1969a). See *Neopilumnoplax americana*.

Pilumnoplax elata (A. Milne Edwards, 1880) (Bull. Mus. Comp. Zool. 8: 18)

As *Eucratoplax elata*—A. Milne Edwards, 1880, p. 18 (original type description, female holotype in Paris Museum, type locality is West Florida, 13 fms.). Not Rathbun, 1898, p. 281.

As *Pilumnoplax elata*—Guinot, 1969a, p. 688. Not Rathbun, 1918, p. 23.

Range: West Florida, type locality.

Depth: 24 m (13 fm).

Remarks: Only the original description of A. Milne Edwards (1880) is valid for this species. All of the other material described by Rathbun (1918, p. 23) has been referred by Guinot (1969a, p. 688, 716–717) to other genera. The male specimen description of Rathbun (1918, p. 23) was referred to *Robertsella mystica* (Guinot, 1969a, p. 716) and the females and juveniles were referred to *Thalassoplax angusta* (Guinot, 1969a, p. 717). This leaves the original type specimen to represent this poorly known species in the Gulf of Mexico.

Pilumnoplax nitida Chace, 1940 (Torreia 3: 44)

Chace, 1940, p. 44, figs. 17–18; Guinot, 1969a, p. 689.

Range: north coast of Cuba.

Depth: 348 to 476 m (190 to 260 fm).

Remarks: Guinot (1969a) retained this species in the genus, but she commented on the obscurity of its relationships.

Prionoplax H. Milne Edwards, 1852

Prionoplax atlantica Kendall, 1891.

Referred to *Frevillea tridentata* by Guinot (1969a), who later decided that *F. tridentata* was different enough from other *Frevillea* to establish a new genus, *Trapezioplax*, for this species. See *Trapezioplax tridentata*.

Robertsella Guinot, 1969

Robertsella mystica Guinot, 1969 (Bull. Mus. Nation. Hist. Nat. 41: 716)

As *Pilumnoplax elata* in Rathbun (not A. Milne Edwards)—Rathbun, 1918, p. 23 (part, male description only), pl. 3 (part, male only).

As *Robertsella mystica*—Guinot, 1969a, p. 716, figs. 132–133, pl. V, fig. 4.

Range: off southeast coast of Florida, Florida Straits.

Depth: 353 m (193 fm).

Habitat: sand bottom.

Remarks: Although this species has not been recorded from the Gulf of Mexico, it is included here because of the confusing nomenclatural history of *Pilumnoplax elata*. At present, this new species and genus includes only the mature male specimen described by Rathbun (1918, p. 23 and part of pl. 3).

Speocarcinus Stimpson, 1859

Speocarcinus carolinensis Stimpson, 1859.

This species was reported from the Gulf of Mexico, prior to revision by Guinot (1969a, p. 710), who referred the Gulf specimens to *S. lobatus*. The Carolinean specimens described in Williams (1965) are *S. carolinensis*.

Speocarcinus lobatus Guinot, 1969 (Bull. Mus. Nation, Hist. Nat. 41: 710)

As *S. carolinensis*—Rathbun, 1918, p. 39 (part, specimen from Dry Tortugas only).

As *S. lobatus*—Guinot, 1969a, p. 710, text-figs. 124–125, pl. IV, fig. 2; Felder 1973a, p. 70, pl. 10, fig. 3.

Range: Dry Tortugas; off Louisiana and Texas.

Depth: shallow water to 37 m (to 20 fm).

Habitat: probably inhabits burrows of polychaetes and crustaceans as does *S. carolinensis*.

Remarks: Dawson (1966) reported *S. carolinensis* from off Grand Isle, Louisiana and Felder (1973a) reported that specimens taken from that same area at later dates were *S. lobatus*.

Tetraplax Rathbun, 1901

Tetraplax quadridentata (Rathbun, 1898) ((Bull. Lab. Nat. Hist. State Univ. Iowa 4: 287)

Rathbun, 1918, p. 32, text-figs. 9–10, pl. 6, figs. 3–4; Rathbun, 1933, p. 78, fig. 68; Guinot, 1969a, p. 256, figs. 1, 14, 26.

Range: north coast of Cuba; Puerto Rico, Curaçao.

Depth: 8 to 22 m (4.5 to 12 fm).

Habitat: mud bottoms.

***Thalassoplax* Guinot, 1969**

***Thalassoplax angusta* Guinot, 1969** (Bull. Mus. Nation. Hist. Nat. 41: 717)

As *Pilumnoplax elata* in Rathbun (not A. Milne Edwards)—Rathbun, 1918, p. 23 (part, female and juveniles, includes female on pl. 3).

As *Thalassoplax angusta*—Guinot, 1969a, p. 717, figs. 131–132, pl. IV, fig. 2; Pequegnat, 1970, p. 192.

Range: east coast of Florida; off northwest Florida, Alabama and Mississippi; off east coast of Mexico; off Campeche, Yucatan.

Depth: 183 to 752 m (100 to 411 fm).

Habitat: mud, sand, broken shell bottoms.

Remarks: Pequegnat (1970) added several new records for the Gulf of Mexico and also commented on differences in morphological descriptions between the *Alaminos* specimens and that provided by Guinot (1969a, p. 717). The legends on Guinot's plates were transposed; the correct citations appear above and in Pequegnat (1970, p. 192). As previously discussed for *Robertsella mystica*, Rathbun's (1918, p. 23) description of *Pilumnoplax elata* does not conform to the type description of this species provided by A. Milne Edwards; thus all of the material included in Rathbun was transferred to new species and genera by Guinot (1969). *Thalassoplax angusta* includes the females and juveniles described by Rathbun, but not the mature male nor the original female holotype of *P. elata*.

***Trapezioplax* Guinot, 1969**

***Trapezioplax tridentata* (A. Milne Edwards, 1880)** (Bul. Mus. Comp. Zool. 8: 16)

As *Goneplax tridentata*—Rathbun, 1918, p. 29.

As *Prionoplax atlantica*—Rathbun, 1918, p. 30, text-fig. 8, pl. 6, figs. 1–2.

As *Trapezioplax tridentata*—Guinot, 1969a, p. 713, figs. 128–129, 142.

Range: Florida Keys and Dry Tortugas; west coast of Florida; Barbados.

Depth: 13 to 42 m (7 to 23 fm).

Habitat: sand, coral, shell, and mud bottoms.

Remarks: Guinot (1969a) established this new genus based on a separation of *Frevillea tridentata* from the other species of *Frevillea*. *Trapezioplax tridentata* includes the original type, *Goneplax tridentata*, and *Prionoplax atlantica*, listed as separate species by Rathbun (1918).

Family PALICIDAE Bouvier, 1898 (= CYMOPOLIDAE Faxon, 1895)

***Palicus* Philippi, 1838**

***Palicus affinis* A. Milne Edwards & Bouvier, 1899** (Bull. Mus. Hist. Nat. Paris 5: 122)

As *Cymopolia affinis*—Rathbun, 1918, p. 196, text-fig. 121, pl. 46, pl. 47, fig. 3;
Rathbun, 1933, p. 85.

Range: southeast and west coasts of Florida; Dry Tortugas; Virgin Islands;
Barbados; Guianas to Espirito Santo, Brazil.

Depth: 33 to 214 m (18 to 117 fm).

Habitat: sand, shell, and coral substrates.

Remarks: Listed from Brazil by Rodrigues da Costa (1968a), Coelho (1971c),
and Coelho and Ramos (1972).

Palicus alternatus Rathbun, 1897 (Proc. Biol. Soc. Washington 11: 95)

As *Cymopolia alternata*—Rathbun, 1918, p. 188, text-fig. 117, pls. 42–43.

As *Palicus alternatus*—Williams, 1965, p. 215, fig. 200.

Range: North Carolina; Florida Keys; west and northwest coasts of Florida.

Depth: 7 to 110 m (4 to 60 fm).

Habitat: sand, gravel, broken shell, coral, and sand-mud bottoms.

Remarks: Ovigerous females are known from Florida during January to
August and from North Carolina in October (Williams, 1965).

Palicus cursor (A. Milne Edwards, 1880) (Bull. Mus. Comp. Zool. 8: 29)

As *Cymopolia cursor*—Rathbun, 1918, p. 215, text-figs. 130–131, pl. 52, figs. 1–2;
Chace, 1940, p. 50.

Range: North Carolina; Florida Keys; northwest coast of Florida; north coast
of Cuba; St. Christopher; Dominica; Barbados.

Depth: 206 to 530 m (107 to 290 fm).

Habitat: sand ooze, sand-mud, sand, and broken shell bottoms.

Remarks: Rathbun (1918) reported ovigerous females from North Carolina
in October, from Florida in March, and from the Antilles in January–February.

Palicus dentatus (A. Milne Edwards, 1880) (Bull. Mus. Comp. Zool. 8: 28)

As *Cymopolia dentata*—Rathbun, 1918, p. 202, text-fig. 124.

As *Palicus dentatus*—Pequegnat, 1970, p. 197.

Range: Florida Keys; west coast of Florida; off Alabama; off Barbados.

Depth: 27 to 139 m (15 to 76 fm).

Habitat: coral and broken shell bottoms.

Palicus faxoni Rathbun, 1897 (Proc. Biol. Soc. Washington 11: 96)

As *Cymopolia faxoni*—Rathbun, 1918, p. 194, text-fig. 120, pl. 45, 2–3.

As *Palicus faxoni*—Williams, 1965, p. 216, fig. 201.

Range: North Carolina; east coast of Florida; northeast of Yucatan (Gulf); ?
off Cape Frio, Brazil.

Depth: 59 to 93 m (32 to 51 fm).

Habitat: sand substrates.

Palicus gracilipes (A. Milne Edwards, 1880) (Bull. Mus. Comp. Zool. 8: 29)

As *Cymopolia gracilipes*—Rathbun, 1918, p. 221, text-fig. 133, pl. 52, figs. 3–4;
Chace, 1940, p. 51.

Range: Bahamas; north of Yucatan; north coast of Cuba.

Depth: 112 to 545 m (61 to 298 fm).

Habitat: sand and shell bottoms.

Palicus gracilis (Smith, 1883) (Proc. U.S. Nat. Mus. 6: 20)

As *Cymopolia gracilis*—Rathbun, 1918, p. 218, text-fig. 132, pl. 50, pl. 51, fig. 1; Chace, 1940, p. 50.

As *Palicus gracilis*—Pequegnat, 1970, p. 195, fig. 6-11.

Range: off Massachusetts; east coast of Florida; northwest Florida; Louisiana to central east coast of Mexico; north coast of Cuba; Curaçao.

Depth: 183 to 686 m (100 to 375 fm).

Habitat: fine sand and mud substrates.

Remarks: Chace (1956) reported this species in the Gulf of Mexico and Pequegnat (1970) reported ovigerous females from the same area in August and November. He further noted that this crab may be able to swim, but is probably not pelagic.

Palicus obesus (A. Milne Edwards, 1880) (Bull. Mus. Comp. Zool. 8: 27)

As *Cymopolia obesa*—Rathbun, 1918, p. 205, text-fig. 125, pl. 49.

As *Palicus obesus*—Pequegnat, 1970, p. 197.

Range: off northwest Florida and Mississippi; Campeche, Mexico.

Depth: 24 to 220 m (13 to 120 fm).

Remarks: Collected by the R/V *Oregon* from the northeastern Gulf of Mexico (Chace, 1956).

Palicus sica (A. Milne Edwards, 1880) (Bull. Mus. Comp. Zool. 8: 29)

As *Cymopolia sica*—Rathbun, 1918, p. 208, text-fig. 127, pl. 40, figs. 3-4; Rathbun, 1933, p. 85, fig. 78; Chace, 1940, p. 49.

As *Palicus sicus*—Milne Edwards & Bouvier, 1902, p. 56, pl. 10, figs. 7-11, pl. 11, fig. 9; Pequegnat, 1970, p. 198.

Range: Florida Keys and Straits; west coast of Florida; north coast of Cuba; Puerto Rico; Virgin Islands; Barbados; Grenada.

Depth: 27 to 348 m (15 to 190 fm).

Habitat: sand, mud, shell, and coral bottoms.

Remarks: Pequegnat (1970) reported ovigerous females from the Gulf of Mexico in mid-July. Rathbun (1918) described the colors of freshly-preserved specimens. Chace (pers. comm.) notes that the specific name *sica* is used as a noun in opposition (L. = curved dagger) and thus should not be changed due to the transfer of the species to the masculine genus *Palicus*.

Family PINNOTHERIDAE de Haan, 1833

Subfamily PINNOTHERINAE de Haan, 1833

Dissodactylus Smith, 1870

Dissodactylus alcocki Rathbun, 1918 (Bull. U.S. Nat. Mus. 97: 124)

Rathbun, 1918, p. 124, text-figs. 70-71, pl. 28, figs. 3-4; Schmitt, McCain & Davidson, 1973, p. 16.

Range: off delta of Mississippi River.

Depth: 64 m (35 fm).

Habitat: sand-mud bottom.

Remarks: This species is known only from a female type and a damaged male paratype.

Dissodactylus borradailei Rathbun, 1918 (Bull. U.S. Nat. Mus. 97: 121)

Rathbun, 1918, p. 121, text-fig. 68, pl. 27, figs. 5–8; Schmitt, McCain & Davidson, 1973, p. 16.

Range: off southeast and southwest coasts of Florida; Jamaica.

Depth: 49 to 55 m (27 to 30 fm).

Habitat: fine white sand.

Dissodactylus calmani Rathbun, 1918 (Bull. U.S. Nat. Mus. 97: 125)

Rathbun, 1918, p. 125, text-figs. 72–73, pl. 28, figs. 5–6; Schmitt, McCain & Davidson, 1973, p. 16.

Range: east coast of Florida; northwest coast of Cuba.

Depth: 4 to 7 m (2 to 4 fm).

Habitat: coral, sand, gravel, and rock bottoms; near shore.

Dissodactylus crinitichelis Moreira, 1901 (Arch. Mus. Nac. Rio de Janeiro 11: 37)

As *D. encopei*—Rathbun, 1918, p. 119, text-fig. 67, pl. 27, figs. 1–4; Williams, McCloskey & Gray, 1968, p. 56, fig. 12.

As *D. crinitichelis*—Rathbun, 1933, p. 83, fig. 76; Schmitt, McCain & Davidson, 1973, p. 17.

Range: off North Carolina; northwest coast of Florida; Jamaica; Puerto Rico; Belize; Caribbean coast of Colombia; Paraiba to Rio Grande do Sul, Brazil.

Depth: shore to 52 m (to 28 fm).

Habitat: fine white sand, coral, and broken shell bottoms; on *Halodule* (sea grass); with the echinoids *Encope marginata*, *E. michelini* (sand dollars) and *Clypeaster subdepressus* (sea biscuit).

Remarks: L. H. Hyman (1955) commented on host relationships with echinoids (as *D. encopei*). Listed from Florida by Wass (1955), Abele (1970), and Menzel (1971); listed from Brazil by Coelho and Ramos (1972) and Rodrigues da Costa (1971).

Dissodactylus encopei Rathbun, 1901.

A junior synonym of *D. crinitichelis* Moreira, 1901.

Dissodactylus juvenilis Bouvier, 1917 (Bull. Mus. Nat. Hist. Natur. Paris 23: 397)

Milne Edwards & Bouvier, 1923, p. 349, text-figs. 11–12, pl. 9, figs. 3–4; Schmitt, McCain & Davidson, 1973, p. 17.

Range: north of Yucatan, Mexico.

Dissodactylus mellitae (Rathbun, 1900) (Amer. Natural. 34: 590)

Hay & Shore, 1918, p. 444, pl. 36, fig. 1; Rathbun, 1918, p. 117, text-fig. 66, pl. 28,

figs. 7-8; Williams, 1965, p. 209, fig. 192; Williams, McCloskey & Gray, 1968, p. 57; Rogers, 1968, p. 318; Schmitt, McCain & Davidson, 1973, p. 18.

Range: Massachusetts to South Carolina; northwest coast of Florida; Texas.
Depth: 9 to 52 m (5 to 28 fm).

Habitat: sand and broken shell bottoms; areas of scattered sponges and coral heads; with the echinoids *Mellita quinquesperforata*, *Encope michelini*, *Echinarachinius parma*, and *Clypeaster subdepressus*.

Remarks: Larval stages have been described by O. W. Hyman (1924), Lebour (1928), Aikawa (1937) and Costlow and Bookhout (1966b). Host relationships were described by Johnson (1952), L. H. Hyman (1955), Gray (1961), Gray, McCloskey and Weihe (1968), and MacGinitie and MacGinitie (1968, p. 314). Regional lists include Florida (Wass, 1955; Abele, 1970; Menzel, 1971). Not listed by Felder (1973a) for Texas, but see Rogers (1968) for a report on this species at Galveston.

Dissodactylus primitivus Bouvier, 1917 (Bull. Mus. Nat. Hist. Natur. Paris 23: 394)

Milne Edwards & Bouvier, 1923, p. 346, text-fig. 8, pl. 8, figs. 3-4, pl. 9, fig. 1; Schmitt, McCain & Davidson, 1973, p. 20.

Range: west of Tortugas, Florida.

Remarks: The above location is the only known record for this species.

Dissodactylus stebbingi Rathbun, 1918 (Bull. U.S. Nat. Mus. 97: 123)

Rathbun, 1918, p. 123, text-fig. 69, pl. 28, figs. 1-2; Schmitt, McCain & Davidson, 1973, p. 20.

Range: Virginia; west and northwest coasts of Florida.

Depth: 9 to 10 m (17 fm).

Habitat: on sea biscuits, *Clypeaster subdepressus*, in an area of scattered sponges and coral heads (northwest Florida).

Remarks: Wass (1955) and Menzel (1971) provide some ecological notes on this crab.

Fabia Dana, 1851

Fabia byssomiae (Say, 1818) (J. Acad. Nat. Sci. Philadelphia 1: 451)

Rathbun, 1918, p. 105, text-fig. 56, pl. 24, figs. 6, 8; Schmitt, McCain & Davidson, 1973, p. 22.

Range: west coast of Florida; northwest coast of Cuba.

Depth: 4 to 9 m (2 to 5 fm).

Habitat: in bivalve molluscs, *Hiatella arctica*; located on beds of *Alcyonium* and between individuals of aggregating ascidians.

Fabia tellinae Cobb, 1973 (Crustaceana 25: 70)

Cobb, 1973, p. 70, figs. 1-2.

Range: off northwest Florida to Alabama.

Depth: 5 to 18 m (3 to 10 fm).

Habitat: commensal in bivalves, *Tellina magna* Spengler (females in mantle cavity, males in excurrent siphon); from sandy bottoms.

***Orthoheres* Sakai, 1969**

***Orthoheres serrei* (Rathbun, 1909) (Bull. Mus. Hist. Nat. Paris 2: 69)**

As *Pinnotheres serrei*—Rathbun, 1918, p. 84, text-fig. 41, pl. 19, figs. 1–7; Rathbun, 1933, p. 82.

As *Orthoheres serrei*—Sakai, 1969, p. 275; Schmitt, McCain & Davidson, 1973, p. 27.

Range: northwest Cuba; Jamaica; Puerto Rico.

Habitat: at surface; on reef flats; in mantle cavity of *Strombus*.

***Orthoheres strombi* (Rathbun, 1905) (Proc. Acad. Nat. Sci. Philadelphia 1905: 371)**

As *Pinnotheres strombi*—Rathbun, 1918, p. 90, text-fig. 45, pl. 20, figs. 1–2.

As *Orthoheres strombi*—Sakai, 1969, p. 275; Schmitt, McCain & Davidson, 1973, p. 27.

Range: west and northwest coasts of Florida.

Habitat: commensal in the gastropods *Strombus pugilis*, *S. alatus*, and *Pleuroploca*.

Remarks: Listed from Florida by Wass (1955), Abele (1970), and Menzel (1971).

***Parapinnixa* Holmes, 1894**

***Parapinnixa bouvieri* Rathbun, 1918 (Bull. U.S. Nat. Mus. 97: 111)**

Rathbun, 1918, p. 111, text-fig. 60, pl. 25, figs. 4–10; Rathbun, 1933, p. 83, fig. 75; Williams, 1965, p. 208, fig. 191; Schmitt, McCain & Davidson, 1973, p. 31.

Range: South Carolina; northeast of Yucatan (Gulf); Puerto Rico.

Depth: 5 to 73 m (3 to 40 fm).

Habitat: coral and sand bottoms; among ventral spines of a sea urchin.

Remarks: Williams (1965) lists ovigerous females from Florida and notes the association of this crab with a sea urchin.

***Parapinnixa hendersoni* Rathbun, 1918 (Bull. U.S. Nat. Mus. 97: 109)**

Rathbun, 1918, p. 109, text-fig. 59, pl. 26, figs. 1–5; Schmitt, McCain & Davidson, 1973, p. 32.

Range: west coast of Florida; northwest Cuba; Curaçao; Maranhão to Bahia, Brazil.

Depth: 38 to 55 m (21 to 30 fm).

Habitat: free-swimming, pelagic (Cuba, in Rathbun, 1918); on sand and broken coral bottom in Florida.

Remarks: Recorded from Brazil by Righi (1967) and Coelho and Ramos (1972).

***Pinnaxodes* Heller, 1865**

***Pinnaxodes floridensis* Wells & Wells, 1961 (Bull. Mar. Sci. Gulf Carib. 11: 267)**

Wells & Wells, 1961, p. 267, figs. 1–2; Schmitt, McCain & Davidson, 1973, p. 34.

Range: west and northwest coasts of Florida.

Habitat: commensal in the cloaca and respiratory tree of the holothurian, *Theleothuria princeps* (Seleniza), which buries in sand; juvenile crabs are found in the anterior digestive tract of the host.

Remarks: Wells and Wells (1961) provided data on the natural history, ecology, and morphology of this crab and Pearce (1966) reviewed the biology and host relationships. Abele (1970) listed the angel wing mollusc, *Cyrtopleura costata*, as a host for a sexual intermediate form of this crab. Listed from north-west Florida by Menzel (1971).

***Pinnotheres* Bosc, 1801–1802**

***Pinnotheres geddesi* Miers, 1880 (J. Linn. Soc. London, Zool. 15: 86)**

Rathbun, 1918, p. 70, text-fig. 32, pl. 16, figs. 1–4; Rathbun, 1933, p. 82; Schmitt, McCain & Davidson, 1973, p. 45.

Range: Veracruz, Mexico; eastern Cuba (Atlantic); Puerto Rico; ? Jamaica.

Habitat: commensal in mangrove oysters (? *Crassostrea rhizophorae*) and *Ostrea*.

***Pinnotheres guerini* H. Milne Edwards, 1853 (Ann. Sci. Nat. Zool. Paris 20: 219)**

Rathbun, 1918, p. 101, text-fig. 52; Rathbun, 1933, p. 83; Schmitt, McCain & Davidson, 1973, p. 48.

Range: Cuba (location not specified); Puerto Rico.

Habitat: reported from oysters.

Remarks: The location of the type specimen in Cuba is unspecified, thus this species may not be present in the Gulf of Mexico.

***Pinnotheres hemphilli* Rathbun, 1918 (Bull. U.S. Nat. Mus. 97: 99)**

Rathbun, 1918, p. 99, text-fig. 51, pl. 23; Schmitt, McCain & Davidson, 1973, p. 48.

Range: Cedar Keys, Florida.

Habitat: intertidal.

Remarks: Known only from a single type specimen.

***Pinnotheres hirtimanus* H. Milne Edwards, 1853 (Ann. Sci. Nat. Zool. Paris 20: 219)**

Rathbun, 1918, p. 101; Schmitt, McCain & Davidson, 1973, p. 48.

Range: Cuba, location unspecified.

Remarks: Known only from the single type specimen.

***Pinnotheres maculatus* Say, 1818 (J. Acad. Nat. Sci. Philadelphia 1: 450)**

Common Names: Mussel Crab; Pea Crab

Hay & Shore, 1918, p. 443, pl. 35, fig. 10; Rathbun, 1918, p. 74, text-figs. 35–36, pl. 17, figs. 3–6; Rathbun, 1933, p. 82, fig. 74; Williams, 1965, p. 206, fig. 190; Felder, 1973a, p. 74, pl. 10, figs. 10–11; Schmitt, McCain & Davidson, 1973, p. 53.

Range: Massachusetts to south Florida; west coast of Florida to Texas; north-west Cuba; Jamaica; Puerto Rico; Virgin Islands; Uruguay and Argentina.

Depth: surface to 50 m (to 27 fm).

Habitat: commensal in a variety of bivalve molluscs; young of both sexes and often adult males are free-swimming; most common in the mantle cavities of mussels, *Mytilus edulis*; in tubes of the polychaetes *Arenicola* and *Chaetopterus* (*C. pergamentaceus* and *C. variopedatus*); from mud, sand, shell and gravel substrates. Other molluscan hosts include: *Atrina rigida*, *A. seminuda*, *A. serrata*, *Anomia simplex*, *Argopecten gibba*, *A. irradians*, *Cyrtopleura costata*, *Modiolus modiolus*, *M. tulipa*, *Mya arenaria*, and *Placopecten magellanica*.

Remarks: This species has a large literature, catalogued by Schmitt, McCain and Davidson (1973). Larval stages were described by O. W. Hyman (1924), Aikawa (1937) and Costlow and Bookhout (1966b). Life history data is provided in MacGinitie and MacGinitie (1968) and by Christensen and McDermott (1959). Caine (1975) studied feeding behavior and physiology and Kruczynski (1975) measured food intake and digestion. Pearce (1964) described reproductive aspects. Larval shadow responses were studied by Forward (1977) and behavior in relation to hosts was described by Sastry and Menzel (1962) and by Eidemiller (1969). The effects of this crab on the growth and biology of its scallop hosts were studied by Kruczynski (1971, 1972). Sandifer (1973) commented on larval ecology in Virginia and Fotheringham and Brunenmeister (1975) described this crab as it occurs in the Gulf of Mexico. Regional lists include Florida (Wass, 1955; Tabb and Manning, 1961; Abele, 1970; Menzel, 1971; Godcharles and Jaap, 1973) and Texas (Leary, 1967). Listed from Brazil by Rodrigues da Costa (1971) and Coelho and Ramos (1972).

Pinnotheres moseri Rathbun, 1918 (Bull. U.S. Nat. Mus. 97: 94)

Rathbun, 1918, p. 94, text-fig. 47, pl. 21, figs. 3-4; Schmitt, McCain & Davidson, 1973, p. 58.

Range: west coast of Florida.

Depth: 1.5 to 5.5 m (1 to 3 fm).

Habitat: commensal in sea squirts (tunicates) and from the brachial cavity of an ascidian, *Polycarpa obtecta*; off rocky bottoms with grass and thin layers of sand and mud.

Remarks: Rathbun (1918) did not list sea squirts as commensals, only as present in the dredges in which the crabs were found. Pearce (1966) provided information on life history and Hartnoll (1964a) described a larval stage. Listed from Florida by Godcharles and Jaap (1973), including data from the collection locality.

Pinnotheres ostreum Say, 1817 (J. Acad. Nat. Sci. Philadelphia 1: 67)

Common Names: Oyster Crab; Common Pea Crab

Hay & Shore, 1918, p. 543, pl. 35, fig. 9; Rathbun, 1918, p. 66, text-fig. 30, pl. 15, figs. 3-6; Williams, 1965, p. 203, figs. 187-189; Felder, 1973a, p. 75, pl. 10, figs. 12-14; Schmitt, McCain & Davidson, 1973, p. 61.

Range: Massachusetts to south Florida; Texas; northwest Cuba; Guadeloupe; Pernambuco to Santa Catarina, Brazil.

Habitat: parasitic in oysters and present in other bivalve molluscs, including: *Crassostrea virginica*, *C. rhizophorae*, *Anomia simplex*, *Mytilus edulis*, and

Pecten spp.; occasionally in polychaete (*Chaetopterus*) tubes; only the first crab ("invasive") stage is free-swimming; found primarily in shallow bays and other suitable oyster habitats.

Remarks: The large literature on this species was catalogued by Schmitt, McCain and Davidson (1973). Earlier biologists thought that this crab was a commensal of oysters, but its parasitic nature was definitely established, as summarized by Stauber (1945), Flower and McDermott (1953) and Haven (1958). Information on larval stages can be found in O. W. Hyman (1924), Lebour (1928), Aikawa (1937), Costlow and Bookhout (1966b) and in Sandoz and Hopkins (1947). Natural history of this species is reviewed by Christensen and McDermott (1959) and by MacGinitie and MacGinitie (1968); Williams (1965) summarized much of the current literature. Beach (1969) studied the life history of this crab in North Carolina. Hartnoll (1971) noted modifications for swimming activity. Listed from Texas by Hedgpeth (1953), Breuer (1962) and Leary (1967).

Pinnotheres serrei Rathbun, 1909.

Transferred to a new genus, *Orthotheres*, by Sakai (1969). Refer to *Orthotheres serrei*.

Pinnotheres shoemakeri Rathbun, 1918 (Bull. U.S. Nat. Mus. 97: 95)

Rathbun, 1918, p. 95, text-fig. 48, pl. 22, figs. 1-4; Rathbun, 1933, p. 83; Schmitt, McCain & Davidson, 1973, p. 86.

Range: west coast of Florida; St. Thomas, Virgin Islands.

Pinnotheres strombi Rathbun, 1905.

Transferred to a new genus, *Orthotheres*, by Sakai (1969). Refer to *Orthotheres strombi*.

Subfamily PINNOTHERELIINAE Alcock, 1900

Pinnixa White, 1846

Pinnixa chacei Wass, 1955 (Quart. J. Flor. Acad. Sci. 18: 160)

Wass, 1955, p. 160, figs. 5-9; Felder, 1973a, p. 71, pl. 10, fig. 5; Schmitt, McCain & Davidson, 1973, p. 104.

Range: northwest Florida; Louisiana and Texas.

Habitat: intertidal, commensal with burrowing shrimp, *Callinassa islagrande*, living in upper part of burrow; on sandy bottoms.

Remarks: Listed from Florida by Wass (1955) and Menzel (1971), from Louisiana by Behre (1950) as *Pinnixa* sp., and from Texas (Leary, 1967).

Pinnixa chaetoptera Stimpson, 1860 (Ann. Lyc. Nat. Hist. New York 7: 235)

Hay & Shore, 1918, p. 445, pl. 36, fig. 4; Rathbun, 1918, p. 151, text-figs. 93-94, pl. 33, figs. 3-6; Williams, 1965, p. 210, fig. 194; Felder, 1973a, p. 74, pl. 10, fig. 8; Schmitt, McCain & Davidson, 1973, p. 104.

Range: Massachusetts to Florida; northwest Florida to Texas; Pernambuco to Rio Grande do Sul, Brazil.

Depth: shore to 16 m (to 9 fm).

Habitat: mud, shell, and gravel bottoms; there are two forms of this crab along the northern Gulf coast: the larger is a commensal with the polychaetes *Amphitrite ornata* and *Chaetopterus variopedatus*, living inside the tubes of the hosts; the smaller form occupies the upper portion of burrows of *Callinassa jamaicensis louisianensis*.

Remarks: Larval stages were described by O. W. Hyman (1924), Lebour (1928), and Aikawa (1937). Sandifer (1973) noted aspects of larval ecology in Virginia. Williams (1965) summarized current literature on this species and MacGinitie and MacGinitie (1968) provided a general account of its life history. Johnson (1952) described a "host factor" for this crab. Behavioral studies include Pearse (1913) and Gray (1961), including notes on symbiotic relationships. Craig (1974) measured temperature tolerances and oxygen consumption. Listed from Florida by Wass (1955), Menzel (1971), and Godcharles and Jaap (1973) and from Mississippi by Richmond (1962) and Christmas and Langley (1973). Listed from Brazil by Righi (1967), Rodrigues da Costa (1971) and Coelho and Ramos (1972).

Pinnixa cristata Rathbun, 1900 (Amer. Natural. 34: 589)

Hay & Shore, 1918, p. 446, pl. 36, fig. 5; Rathbun, 1918, p. 134, text-fig. 78, pl. 29, figs. 8-9; Williams, 1965, p. 210, fig. 193; Felder, 1973a, p. 74, pl. 10, fig. 6; Schmitt, McCain & Davidson, 1973, p. 106.

Range: North and South Carolina; Louisiana and Texas.

Habitat: intertidal beaches; shallow sand and sand-mud substrates of brackish to marine waters; usually commensal with callinassid shrimps and other burrowers.

Remarks: Hedgpeth (1950) described these crab from salt flats that border the bays and intercoastal waterways of Texas. MacGinitie and MacGinitie (1968) included information on the ecology of this species. Listed from Louisiana by Behre (1950).

Pinnixa cylindrica (Say, 1818) (J. Acad. Nat. Sci. Philadelphia 1: 452)

Hay & Shore, 1918, p. 446, pl. 36, fig. 3; Rathbun, 1918, p. 159, text-fig. 99, pl. 35, figs. 5, 8; Milne Edwards & Bouvier, 1923, p. 345; Williams, 1965, p. 213, fig. 197; Schmitt, McCain & Davidson, 1973, p. 106.

Range: Massachusetts to South Carolina; west and northwest coasts of Florida.

Depth: shallow water to 37 m (to 20 fm).

Habitat: commensal with *Arenicola cristata* (lugworm) in the non-tubular burrows; young crabs occur near the intertidal zone of slimy shores.

Remarks: McDermott (1962) provided a general account of this species, which is also summarized by Williams (1965). Sandifer (1973) commented on larval ecology in Virginia. Listed by Menzel (1971) from northwest Florida.

Pinnixa floridana Rathbun, 1918 (Bull. U.S. Nat. Mus. 97: 138)

Rathbun, 1918, p. 138, text-fig. 82, pl. 30, figs. 4-7; Williams, McCloskey & Gray, 1968, p. 57, fig. 13; Schmitt, McCain & Davidson, 1973, p. 110.

Range: North Carolina; west and northwest coasts of Florida.

Habitat: shallow water, possibly in tubes of the polychaete, *Diopatra cuprea*; collected from a compound ascidian growing on a soft coral; from under rocks in 10 feet of water.

Remarks: Rathbun (1918) commented on morphological variation between the sexes of this species. Listed from Florida by Wass (1955) and Menzel (1971).

Pinnixa leptosynaptae Wass, 1968 (Tulane Stud. Zool. 14: 137)

Wass, 1968, p. 137, figs. 1-6; Schmitt, McCain & Davidson, 1973, p. 112.

Range: west coast of Florida.

Habitat: found on the surface of the holothurian *Leptosynapta crassipatina*.

Remarks: Listed from northwest Florida by Menzel (1971).

Pinnixa lunzi Glassell, 1937 (Charleston Mus. Leaflet 9: 3)

Glassell, 1937, p. 3, figs. 1-8; Williams, 1965, p. 214, figs. 198-199; Felder, 1973a, p. 71, pl. 10, fig. 4; Schmitt, McCain & Davidson, 1973, p. 114.

Range: Virginia to Georgia; Mississippi to Texas.

Depth: near shore to 22 m (to 12 fm).

Habitat: on beaches, under drift material; in burrows of echiurans (*Thalassema hartmani*) and possibly other burrowers.

Remarks: Boesch (1971) listed this crab from an echiuran burrow in Virginia. Felder (1973b) reported a specimen taken from a red snapper stomach from a reef off Texas.

Pinnixa pearsei Wass, 1955 (Quart. J. Flor. Acad. Sci. 18: 164)

Wass, 1955, p. 164, figs. 10-13; Schmitt, McCain & Davidson, 1973, p. 116.

Range: northwest coast of Florida.

Habitat: commensal in tubes of the polychaete, *Diopatra*, from sand-mud beaches.

Remarks: Listed from northwest Florida by Abele (1970) and Menzel (1971). Although Menzel (1971) listed this crab as a commensal of an undetermined annelid, Abele (1970) stated that the crab does not seem to be restricted to commensal relationships where it was common in the sand-mud intertidal zone of Alligator Harbor's south shore.

Pinnixa retinens Rathbun, 1918 (Bull. U.S. Nat. Mus. 97: 139)

Rathbun, 1918, p. 139, text-figs. 83-84, pl. 41, figs. 1-2; Williams, 1965, p. 212, fig. 196; Felder, 1973a, p. 74, pl. 10, fig. 7; Schmitt, McCain & Davidson, 1973, p. 118.

Range: Chesapeake Bay; west coast of Florida; Texas.

Depth: low tide mark to 37 m (to 20 fm).

Habitat: mud bottoms of estuarine and marine waters; from burrows of the callinassid shrimp, *Upogebia affinis*.

Remarks: Rathbun (1918) believed this crab to be allied with *P. floridana*. Listed from Florida by Wass (1955) and Menzel (1971). Williams (1965) included this species in the Carolina fauna, even though it hadn't yet been collected from that area. Based on the wide range of locales but paucity of specimens, it is probably a rare species.

Pinnixa sayana Stimpson, 1860 (Ann. Lyc. Nat. Hist. New York 7: 236)

Hay & Shore, 1918, p. 446, pl. 36, fig. 3; Rathbun, 1918, p. 156, text-fig. 98, pl. 34, figs. 2-4; Williams, 1965, p. 212, fig. 195; Felder, 1973a, p. 74, pl. 10, fig. 9; Schmitt, McCain & Davidson, 1973, p. 119.

Range: Massachusetts to North Carolina; west coast of Florida; Louisiana; Amapá to São Paulo, Brazil.

Depth: surface to 75 m (to 41 fm).

Habitat: free-swimming; on sandy beaches in drift material; from mud bottoms; in tubes of lugworm, *Arenicola cristata*.

Remarks: Larval descriptions include O. W. Hyman (1924), Lebour (1928), and Aikawa (1929, 1937). Regional lists include Louisiana (Behre, 1950). Schmitt, McCain and Davidson (1973) note that the host record of *Arenicola* may be due to synonymy of *P. sayana* with *P. cylindrica* by Hay and Shore (1918) and so may be in error. Listed from Brazil by Righi (1967), Rodrigues da Costa (1968a), Coelho (1971a) and Coelho and Ramos (1972).

Family GRAPSIDAE, Macleay, 1838

Subfamily GRAPSINAE Macleay, 1838

Geograpsus Stimpson, 1858

Geograpsus lividus (H. Milne Edwards, 1837) (Hist. Nat. Crust., vol. 2. p. 85)

Rathbun, 1918, p. 234, pl. 55; Rathbun, 1933, p. 87, fig. 80; Garth, 1965a, p. 26; Forest & Guinot, 1966, p. 91; Chace & Hobbs, 1969, p. 157, figs. 48, 52a-c.

Range: Bermuda; Florida Keys, north and south coasts of Cuba; Jamaica; Puerto Rico; Virgin Islands to Barbados; Netherlands Antilles to Trinidad; Old Providence Island (Carib.); Caribbean coast of Colombia to São Paulo, Brazil; eastern Atlantic, from Senegal to Angola; Cape Verde Islands; eastern Pacific, from southern part of Baja California to northern Chile; Clipperton Island; Galapagos Islands; Hawaiian Islands.

Habitat: supralittoral, near the splash zone of rocky areas and stone beaches; from middle to upper intertidal, beneath stones.

Remarks: Hartnoll (1965b) provided ecological notes on populations in Jamaica. Chace and Hobbs (1969) commented on ecology of this crab in Dominica. Listed from Brazil by Coelho and Ramos (1972) and Fausto Filho (1974).

Goniopsis de Haan, 1833

Goniopsis cruentata (Latreille, 1803) (Hist. Nat. Crust., vol. 6, p. 70)

Common Names: Mangrove Crab; Tree Crab

Rathbun, 1918, p. 237, text-fig. 136, pl. 57; Rathbun, 1933, p. 87, fig. 81; Chace, 1940, p. 52; Bott, 1955a, p. 62; Holthuis, 1959, p. 235, figs. 59-60; Forest & Guinot, 1966, p. 91; Chace & Hobbs, 1969, p. 160, figs. 49, 52d-f; Felder, 1973a, p. 78, pl. 11, figs. 8-9.

Range: Bermuda; Bahamas; northwest Florida (rare); Tampico, Mexico; north and south coasts of Cuba; Jamaica; Hispaniola; Puerto Rico; Virgin Is-

lands to Barbados; Netherlands Antilles; Belize; Old Providence Island (Carib.); Surinam to Rio de Janeiro, Brazil; eastern Atlantic, from Senegal to northern Angola.

Habitat: mangrove swamps, along roots and on trunks of trees; on wet muddy marine shores, along inlets and estuaries; intertidal to supratidal.

Remarks: Leary (1967) listed this species from Texas, based on a collection by Hildebrand in 1958, but its occurrence along the Gulf coast is scattered and rare. Ecological studies include field work in Jamaica (Hartnoll, 1965b; Warner, 1969) and Dominica (Chace and Hobbs, 1969). Behavioral data is provided by Schone and Schone (1963), Schone (1968), and Warner (1970). Physiological studies include data on thoracic neurosecretion (Maynard, 1961a, 1961b; Maynard and Maynard, 1962) and coagulation (Morrison and Morrison, 1952). Listed from Brazil by Coelho and Ramos (1972).

***Grapsus* Lamarck, 1801**

***Grapsus grapsus* (Linnaeus, 1758)** (Syst. Nat., ed. 10, vol. 1, p. 630)

Common Names: Rock Crab; Cliff Crab; Sally Lightfoot

Rathbun, 1918, p. 227, text-fig. 135, pls. 53-54; Rathbun, 1933, p. 86, fig. 79; Garth, 1965, p. 25; Forest & Guinot, 1966, p. 90; Chace & Hobbs, 1969, p. 163, figs. 50, 52g-i; Felder, 1973a, p. 78, pl. 11, fig. 15.

Range: Bermuda; Bahamas; southeast and south Florida; Texas; north and south coasts of Cuba; Jamaica; Puerto Rico; Hispaniola; Virgin Islands to Barbados; Netherlands Antilles to Trinidad; Old Providence Island and Swan Island (Carib.); Colombia to northern Brazil; eastern Atlantic, from Portugal to Angola; Cape Verde Islands and Azores; St. Helena Island; Ascension Island; eastern Pacific, from central Baja California to central Chile; Galapagos Islands; Clipperton Island.

Habitat: intertidal and supratidal zones of rocky areas, stone beaches, and sea walls; within reach of splash from surf and wave action; in crevices and cracks of rock cliffs near water's edge.

Remarks: Reports of this crab in the Gulf of Mexico are confined to the north coast of Cuba and from the rock jetties of Texas, where they are rare. Listed from Texas by Leary (1967), based on collections by Hildebrand. Also recorded from Brazil by Coelho and Ramos (1972) and Fausto Filho (1974). Ecological studies include those of Hartnoll (1965b) in Jamaica, Chace and Hobbs (1969) in Dominica, and Johnson (1965) on the relation of behavior to development and ecology. Social behavior was studied by Wright (1966, 1968), Schone and Eibl-Eibesfeldt (1965), Kramer (1967), Schone (1968), and Eibl-Eibesfeldt (film, 1963). Hartnoll (1971) noted the ability of this crab to swim. Physiological and anatomical studies include work on gill anatomy (Chen, 1933), coagulation (Morrison and Morrison, 1952), thoracic neurosecretion (Maynard, 1961a, 1961b; Maynard and Maynard, 1962), and neural fine structure (Skobe and Nunnemacher, 1970).

***Pachygrapsus* Randall, 1840**

Pachygrapsus gracilis (Saussure, 1858) (Mém. Soc. Phys. Hist. Nat. Genève 15: 443)

Common Name: Wharf Crab

Rathbun, 1918, p. 249, pl. 60, fig. 3, pl. 61, fig. 1; Rathbun, 1933, p. 89; Holthuis, 1959, p. 239, pl. 10, fig. 3; Forest & Guinot, 1966, p. 92; Chace & Hobbs, 1969, p. 167, figs. 51, 52j; Felder, 1973a, p. 79, pl. 11, figs. 3-4, 11.

Range: Bermuda; Bahamas; south Florida; Texas; north and south coasts of Cuba; Jamaica; Puerto Rico; Virgin Islands; Caribbean coast of Columbia; Pernambuco to Bahia, Brazil; eastern Atlantic, from Senegal to Zaire.

Habitat: mangrove roots; along river banks near the sea; on pilings, wharves, stone jetties; rocky areas, just above the water level; intertidal to near supra-tidal.

Remarks: Extensive notes on the natural history of this crab in Jamaica were provided by Hartnoll (1965b) and by Warner (1969). Felder (1973a) listed some collection localities in Texas, but it is absent from collection lists of west Florida, Mississippi, and Louisiana. Listed from Brazil by Coelho and Ramos (1972).

Pachygrapsus transversus (Gibbes, 1850) (Proc. Amer. Assoc. Adv. Sci. 3: 181.

Common Name: Mottled Shore Crab

Hay & Shore, 1918, p. 447, pl. 36, fig. 9; Rathbun, 1918, p. 244, pl. 61; Rathbun, 1933, p. 88, fig. 82; Williams, 1965, p. 217, fig. 202; Forest & Guinot, 1966, p. 91; Chace & Hobbs, 1969, p. 169, fig. 52k; Felder, 1973a, p. 79, pl. 11, figs. 5, 10.

Range: North Carolina; Bermuda; Bahamas; east coast of Florida; Florida Keys and Dry Tortugas; Louisiana to east coast of Mexico; north coast of Cuba; Jamaica; Hispaniola; Puerto Rico; St. Thomas, Virgin Islands to Barbados; Trinidad; Netherlands Antilles; Bahia, Brazil to Uruguay; eastern Atlantic, from Mediterranean Sea to Angola; eastern Pacific, from central California to Peru; Galapagos Islands; Easter Island.

Habitat: mainly in rocky areas near the tideline; beneath stones and on wharves and pilings; occasionally found among mangrove roots and on sandy beaches.

Remarks: Leobur (1944) figured some of the larval stages. Ecological studies include Hartnoll (1965b) in Jamaica and Verrill (1908) in Bermuda. Listed from Louisiana (Behre, 1950) and Texas (Whitten, Rosene and Hedgpeth, 1950; Leary, 1967). Pearse (1932a) reported a protozoan from the gill cavity of this crab. Hazlett (1971) studied antennule chemosensitivity and Alves (1974) tested salinity tolerances. Listed from Brazil by Coelho and Ramos (1972) and Fausto Filho (1974).

***Planes* Bowdich, 1825**

Planes cyaneus Dana, 1852 (Proc. Acad. Nat. Sci. Philadelphia 5: 250)

As *P. minutus*—Barnard, 1950, p. 120.

As *P. cyaneus*—Chace, 1951, pp. 65–103, figs. 1b, 2b, 2e, 2h, 2m–o, 3i–n; Chace 1966, p. 646; Sakai, 1965, p. 197, pl. 93, figs. 3–4; Felder, 1973a, p. 78, pl. 11, fig. 1.

Range: rare occurrence in Texas; south Atlantic, at St. Helena Island and off west coast of Africa; throughout eastern Pacific and westward to the north-western Pacific and Indian Ocean.

Habitat: pelagic, on floating objects, debris, and gulfweed; open ocean, but occasionally washed up with flotsam onto beaches.

Remarks: Chace (1951) provided a definitive taxonomic review, but this crab was considered at that time to be confined to the Pacific Ocean. In 1966, Chace reported it from St. Helena Island in the south Atlantic and noted that Barnard's (1950) citations of *P. minutus* from South Africa may have been partly or entirely records of *P. cyaneus*. Crosnier (1967) reported this crab from West Africa. Shirley (1974) found two specimens washed up on the beaches at Padre Island, Texas, in driftwood. This species should be considered extralimital for the Gulf of Mexico, although re-examination of *Planes* in collections may reveal additional specimens of this species.

Subfamily PLAGUSIINAE Dana, 1851

Percnon Gistel, 1848

Percnon gibbesi (H. Milne Edwards, 1853) (Ann. Sci. Nat. ser. 3, Zool. 20: 146 and 180)

Common Name: Spray Crab

Rathbun, 1918, p. 337, pl. 105; Rathbun, 1933, p. 93; fig. 88; Schmitt, 1939, p. 24; Garth, 1965, p. 34; Williams, 1965, p. 224.

Range: North Carolina, Bermuda; Bahamas, south Florida; Florida Keys; north coast of Cuba; Jamaica; Puerto Rico; Colon, Panama; Brazil; eastern Atlantic, from the Azores to South Africa; eastern Pacific, from Baja California to Chile; Galapagos Islands; Clipperton Island.

Habitat: low tide zone of rocky areas; surf zone, on rock and pebble bottoms, commensal with *Diadema* in Puerto Rico.

Remarks: Rathbun (1918) questioned the inclusion of the Pacific and Atlantic populations of this crab into one species, but subsequent authors have treated them as identical. Verrill (1908) commented on this crab in Bermuda and Hartnoll (1965b) described its biology and ecology in Jamaica. Garth (1946) described it from the Galapagos Islands. Schmitt (1939) provided the key characters for the genus. The habitat and color of Brazilian specimens was provided by Fausto Filho (1974).

Plagusia Latreille, 1806

Plagusia depressa (Fabricius, 1775) (Syst. entom., 1775), p. 406

Common Name: Cliff Crab

Rathbun, 1918, p. 332, text-fig. 154, pl. 101; Rathbun, 1933, p. 93, fig. 87; Monod,

1956, p. 455, figs. 614–617; Williams, 1965, p. 223, fig. 207; Forest & Guinot, 1966, p. 93; Chace & Hobbs, 1969, p. 192, figs. 62r–t, 63; Felder, 1973a, p. 75, pl. 11, fig. 13.

Range: North and South Carolina; Bermuda; Florida Keys and Dry Tortugas; Texas; Cuba; Jamaica; Hispaniola; Puerto Rico; Virgin Islands; Dominica to Barbados; Trinidad; Netherlands Antilles; Ceará to Pernambuco, Brazil; eastern Atlantic, from the Azores and Madeira and Senegal to Angola.

Habitat: in fissures and crevices of rocks; in tide pools; on jetties; intertidal.

Remarks: Chace and Hobbs (1969) provided notes on color patterns of living crabs. Haratnoll (1965b) studied the ecology of this crab in Jamaica and notes on swimming behavior were given in Hartnoll (1971). Physiological studies include Morrison and Morrison (1952) on coagulation and data on thoracic neurosecretion (Marynard, 1961a, 1961b; Marynard and Marynard, 1962). Listed from Brazil by Coelho (1971a), Coelho and Ramos (1972), and Fausto Filho (1974).

Subfamily SESARMINAE Dana, 1852

Aratus H. Milne Edwards, 1853

Aratus pisonii (H. Milne Edwards, 1837) (Hist. Nat. Crust., vol. 2: 76)

Common Names: Mangrove Crab; Tree Crab

Rathbun, 1918, p. 323, pl. 96; Rathbun, 1933, p. 92, fig. 85; Chace & Hobbs, 1969, p. 172, figs. 54, 58a.

Range: Bahamas; southeast to southwest Florida; north and south coasts of Cuba; New Providence Island (Atlantic); Jamaica; Puerto Rico; Virgin Islands to Guadeloupe; Netherlands Antilles; Belize; Rio Parahyba do Norte to São Paulo, Brazil; Nicaragua to Peru, in eastern Pacific.

Habitat: along shores of estuaries and near fresh, brackish or marine waters; on rocks, piles, and wharves; commonly in mangroves, on which this crab can easily climb.

Remarks: Warner (1968) described larval development. Hartnoll (1965b) provided extensive notes on the biology of this crab in Jamaica, including ecology, growth, feeding, behavior, and reproduction. Warner (1967, 1969, 1970) also studied this species in Jamaica. Hartnoll (1971) briefly commented on swimming activity. Listed from south Florida by Tabb and Manning (1961) and from Brazil by Coelho and Ramos (1972).

Cyclograpsus H. Milne Edwards, 1837

Cyclograpsus integer H. Milne Edwards, 1837 (Hist. Nat. Crust., vol. 2: 79)

Common Name: Marsh Crab

Rathbun, 1918, p. 326, pl. 97, figs. 1–2; Rathbun, 1933, p. 92, fig. 86; Monod, 1956, p. 451, figs. 609–612; Chace & Hobbs, 1969, p. 173, figs. 55, 58b–d; Felder, 1973a, p. 75, pl. 11, figs. 12, 14.

Range: Bermuda; Bahamas; south Florida; Florida Keys; Texas; Cuba; Jamaica; Hispaniola; Puerto Rico; St. Croix; Dominica; Islas Los Roques and

Caribbean coast of Colombia; Ceará to Pernambuco, Brazil; eastern Atlantic, from Senegal to Zaire.

Habitat: burrows in marshy marine areas; among rocky and stony areas of the intertidal zone and up to the high tide line.

Remarks: Felder (1973a) provided the only other specific Gulf record in addition to the previous record for the Florida Keys. Hartnoll (1965b) commented on the ecology of this crab in Jamaica. Listed from Brazil by Coelho and Ramos (1972) and Fausto Filho (1974).

Sesarma Say, 1817

Subgenus *Holometopus* H. Milne Edwards, 1853

Sesarma (Holometopus) americanum Saussure, 1858 (Mém. Soc. Hist. Nat. Genève 14: 441)

As *S. tampicense*—Rathbun, 1918, p. 307, text-fig. 151, pl. 88,

As *S. americanum*—Chace & Hobbs, 1969, p. 178, figs. 62a–f.

Range: Tampico, Mexico; St. Thomas, Virgin Islands.

Habitat: soft mud, along river banks.

Remarks: Chace and Hobbs (1969) determined that *S. tampicense* Rathbun was a junior synonym of *S. americanum* Saussure. Behre (1950) tentatively listed the species from Louisiana (as *S. tampicense*), but noted that Chace had examined the specimens and preferred not to record the species as indicated. Abele (1972b) mentions the similarities between several of the western Atlantic members of the genus; he notes the distinct differences between *S. angustipes* Dana and *S. americanum*.

Sesarma (Holometopus) angustipes Dana, 1852.

This species was restricted by Abele (1972b) to the specimens from Brazil and Trinidad; refer to his paper for a discussion of synonymy. Material listed under this name by Rathbun (1918, p. 331) was synonymized with *S. roberti* by Chace and Hobbs (1969, p. 184).

Sesarma (Holometopus) benedicti Rathbun, 1897 (Proc. Biol. Soc. Washington 11: 90)

Rathbun, 1918, p. 316, pl. 93; Holthuis, 1959, p. 248, fig. 62; Abele, 1973, p. 379, figs. 1A, 1G.

Range: Key West, Florida; Guyana and Surinam; Brazil.

Habitat: under wood and stones on banks of brackish and almost freshwater streams.

Remarks: The female specimen from Key West (MCZ 6236) listed by Rathbun (1918) is the only Gulf of Mexico record; all others are from South America. Ecological notes were provided by Holthuis (1959).

Sesarma (Holometopus) cinereum—(Bosc, 1802) (Hist. Nat. Crust., vol. 1, an X, p. 204)

Common Names: Square-backed Fiddler; Wharf Crab; Wood Crab; Friendly Crab

As *S. cinerea*—Hay & Short, 1918, p. 449, pl. 36, fig. 11.

As *S. cinereum*—Rathbun, 1918, p. 300, text-fig. 149, pl. 83; Williams, 1965, p. 222, fig. 206; Felder, 1973a, p. 78, pl. 11, fig. 6; Abele, 1973, p. 377, figs. 1B, 1H.

Range: Maryland to southeast Florida; southwest Florida to Vera Cruz, Mexico.

Habitat: on wharves, pilings, and other wooden objects; stone jetties and rocky areas; in *Spartina* marshes and along the edges of mangrove swamps; burrows from the high tide mark to well inland in mud and sand substrates; frequently found on boats and ships.

Remarks: Abele (1973) states that previous records of *S. cinereum* from the West Indies and elsewhere in the Caribbean were based on juvenile specimens of *S. ricordi* and *S. americanum*. Regional lists include Florida (Wass, 1955; Menzel, 1971; Subrahmanyam *et al.*, 1976), Mississippi (Richmond, 1968), Louisiana (Behre, 1950; Hoesel and Valentine, 1972) and Texas (Hedgpeth, 1953; Leary, 1967). Hedgpeth (1953) presented a map, showing the ranges of *Sesarma* in the northern Gulf of Mexico. Williams (1965) listed ovigerous females from North Carolina in May to November and from the mouth of the Potomac River in January. Abele (1973) collected ovigerous females from Florida in June and from Texas in July. Sandifer (1973) commented on larval abundance in Virginia.

Larval development was studied by Costlow, Bookhout and Monroe (1960) and Costlow and Bookhout (1960b). Ecological notes were provided by Williams (1965) and by Fotheringham and Brunenmeister (1975). Physiological studies include observations on gill area (Gray, 1957), oxygen consumption (Teal, 1959), and tolerance to dilute salt water (Pearse, 1929). Observations on the behavior of this crab in captivity were made by Oler (1941) and by Duncker (1934).

Sesarma (Holometopus) miersii Rathbun, 1897 (Proc. Biol. Soc. Washington 11: 91)

Rathbun, 1918, p. 303 (part), pl. 84; Chace & Hobbs, 1969, p. 180, figs. 59, 62g-i;

Abele, 1972b, p. 166, figs. 1B, 1C, 2B, 2C; Abele, 1973, p. 380, fig. 1I.

Range: Bahamas; Key West, Florida; south coast of Cuba; Swan Island (Carib.); Dominica.

Habitat: marshy tidal flats.

Remarks: Abele (1972b) reviewed the status of this crab and the confusion in nomenclature that existed from Rathbun's (1897) description of Mier's original material. Only the original specimens from the Bahamas became the type material for this species. Specimens from Brazil in Rathbun (1918) are now designated *S. angustipes* and the specimen from Uruguay is *Metasesarma rubripes*. Hartnoll (1965b) found no evidence of *S. miersii* in Jamaica and later authors agree that the observations of Andrews cited in Rathbun (1918, p. 304) refer to *S. roberti* (see Abele, 1973). The only Gulf of Mexico record is that from Key West (Abele, 1973).

Sesarma (Holometopus) ricordi H. Milne Edwards, 1853 (Ann. Sci. Nat., ser. 3, Zool. 20: 183)

Common Name: Beach Crab

Rathbun, 1918, p. 309 (part), pl. 89; Rathbun, 1933, p. 91; Holthuis, 1959, p. 246, pl. 11, fig. 3; Chace and Hobbs, 1969, p. 183, fig. 62k; Abele, 1973, p. 378, fig. 1J.

Range: Bermuda; Bahamas; southeast Florida; Florida Keys; west coast of Florida; north coast of Yucatan; Cuba; Jamaica; Hispaniola; Puerto Rico; Virgin Islands to Trinidad; Curaçao; Old Providence Island (Carib.); Yucatan to Surinam.

Habitat: from intertidal zone to about 50 meters inland; in low-lying pine woods; edges of mangrove swamps; burrows in grassy areas above sandy beaches; under driftwood and among rocks, along shorelines.

Remarks: Previous records of this species from Mississippi have been identified as *S. cinereum* by Chace (in Hedgpeth, 1953) and the material from Brazil was determined to be *S. angustipes* by Abele (1972b). Abele (1973) listed ovigerous females from Florida in May, June and August and from Panama in January. Larval development was studied by Diaz and Ewald (1968). The ecology and other aspects of biology of this crab in Jamaica were reported by Hartnoll (1965b), Warner (1969), and Standing (1972).

Sesarma (Holometopus) roberti H. Milne Edwards, 1853 (Ann. Sci. Nat. ser. 3, Zool. 20: 182)

Common Name: Brackish-water Crab

As *S. angustipes*—Rathbun, 1918, p. 311, pl. 90.

As *S. roberti*—Rathbun, 1918, p. 312, pl. 91; Rathbun, 1933, p. 91; Monod, 1956, p. 443, figs. 602–604; Chace & Hobbs, 1969, p. 184, figs. 60, 62l–n.

Range: Veracruz, Mexico to Nicaragua; north and south coasts of Cuba; Jamaica; Hispaniola; Puerto Rico; Virgin Islands to Trinidad; Venezuela.

Habitat: streams, rivers, and bays, including a wide range of freshwater and brackish environments; from marine shorelines to upland elevations of 1000 feet; burrows in steep muddy banks; among mangroves; on rocks in streams; among stony areas at bases of cliffs.

Remarks: Monod (1956) questioned the occurrence of *S. roberti* on Goree Island off West Africa, the indicated type-locality for this species. Hartnoll (1965b) reported on the biology of this crab in Jamaica (under the name *S. angustipes*, which he used as a senior synonym of *S. roberti*). Chace & Hobbs (1969) maintained the distinction between these two species, at least until a South American specimen with Caribbean characteristics is discovered. Abele (1972b) reviewed these reports and also regarded the two forms as separate species.

Sesarma (Holometopus) tampicense Rathbun, 1914.

This name was determined by Chace & Hobbs (1969) to be a junior synonym of *S. americanum*.

Subgenus *Sesarma* Say, 1817***Sesarma (Sesarma) curacaoense*** de Man, 1892 (Notes Leyden Mus. 14: 257)

Common Name: Mangrove Crab

Rathbun, 1918, p. 293, text-fig. 147, pl. 78, figs. 1-2, pl. 160, fig. 3; Rathbun, 1933, p. 90; Holthuis, 1959, p. 242; Chace & Hobbs, 1969, p. 188, figs. 61, 62p; Abele, 1973, p. 380, figs. 1C, 1F.

Range: Key West, Florida; south and southwest Florida; north coast of Cuba; Jamaica; Puerto Rico; Curaçao; Bahia, Brazil.

Habitat: muddy banks of rivers and ditches, including brackish water; in mangrove swamps, under rocks and litter; intertidal zone, in clumps of oysters and among rocks.

Remarks: Abele (1973) commented on sexual maturity and size ranges, but he did not report ovigerous females in the Florida material he studied. Tabb and Manning (1961) listed this crab from mangroves at Whitewater Bay in south Florida. Ecology of this species in Jamaica was studied by Hartnoll (1965b) and Warner (1969). Listed from Brazil by Coelho and Ramos (1972).

Sesarma (Sesarma) reticulatum (Say, 1817) (J. Acad. Nat. Sci. Philadelphia 1: 73)

Common Name: Marsh Crab

As *S. reticulata*—Hay & Shore, 1918, p. 448, pl. 36, fig. 12.

As *S. reticulatum*—Rathbun, 1918, p. 290, pl. 77; Williams, 1965, p. 221, fig. 205; Felder, 1973a, p. 78, pl. 11, fig. 7; Abele, 1973, p. 380, fig. 1D, 1E.

Range: Massachusetts to east coast of Florida; west coast of Florida to central Texas.

Habitat: *Spartina* salt marshes; burrows in soft muds and sand-mud; under rocks and litter of intertidal streams and near-marine to brackish waters.

Remarks: Larval descriptions were provided by O. W. Hyman (1924) and Costlow and Bookhout (1962). Sandifer (1973) reported on larval ecology in Virginia. Crichton (1960) gave a general account of life history as noted in Delaware marshes. Regional lists include Florida (Wass, 1955; Menzel, 1971; Subrahmanyam *et al.*, 1976), Louisiana (Behre, 1950), and Texas (Hedgpeth, 1953; Leary, 1967). The specimen listed by Tabb and Manning (1961) for this species was determined by Abele (1973) to be *S. curacaoense*. Humes (1941b) described a parasitic copepod in the gill chambers of this crab. Physiological studies include work on gill area (Gray, 1957), oxygen consumption (Teal, 1959), rhythmic activity (Palmer, 1967), antennule chemosensitivity (Hazlett, 1971), and melanophore hormones (Fingerman, Nagabhushanam and Philpott, 1961).

Subfamily VARUNINAE H. Milne Edwards, 1852

Euchirograpsus H. Milne Edwards, 1853***Euchirograpsus americanus*** A. Milne Edwards, 1880 (Bull. Mus. Comp. Zool. 8: 18)

Hay & Shore, 1918, p. 448, pl. 36, fig. 7; Rathbun, 1918, p. 282, text-fig. 144, pl.

74; Chace, 1940, p. 52; Williams, 1965, p. 220, fig. 204; Türkay, 1975, p. 114, figs. 6-7, 16b, 20, 24.

Range: North and South Carolina; south Florida; north and south coasts of Cuba; St. Lucia; Barbados; Colombia to Venezuela.

Depth: 31 to 508 m (17 to 278 fm).

Habitat: rocky, coral, and sand substrates.

Remarks: Listed from off the Carolinas by Cerame-Vivas, Williams and Gray (1963) and Cerame-Vivas and Gray (1966). Williams (1965) listed ovigerous females from Florida in March to September. Recorded off the Atlantic coast of Cuba by Chace (1956). The Pacific specimens listed by Garth (1946) were referred to a new species, *E. pacificus*, and the specimen (USNM 17672) listed by Rathbun (1918) from off Yucatan was referred to a new species, *E. antillensis*, by Türkay (1975).

Euchirograpsus antillensis Türkay, 1975 (Senckenbergiana Biol. 56: 112)

As *E. americanus*—Rathbun, 1918, p. 283 (part, Yucatan specimen only).

As *E. antillensis*—Türkay, 1975, p. 112, figs. 4-5, 16a, 19, 25.

Range: off Havana, Cuba; Arrowsmith Banks, between Cuba and Yucatan; south of Florida Keys; Bahamas.

Depth: 192 to 430 m (105 to 235 fm).

Remarks: Türkay (1975) compared this new species with *E. americanus* and the other species of the genus, which now number a total of six.

Platychoirapsus de Man, 1896

Platychoirapsus spectabilis de Man, 1896 (Zool. Anz. 19: 292)

Common Names: River Crab; Saber Crab

As *P. typicus*—Rathbun, 1918, p. 278, text-figs. 141-143, pl. 73; Bolívar y Pieltain, 1945, p. 267-270, figs. 1-5.

As *P. spectabilis*—Buitendijk, 1950, p. 280, fig. 1b; Monod, 1956, p. 426, text-figs. 584-588.

Range: Gulf coast of Mexico; west coast of Florida.

Habitat: burrows in clay banks, just above the water line, along rivers; shallow rocky areas of rivers; known from altitudes of greater than 100 feet, up to 140 miles from the sea.

Remarks: This species was described on the basis of only a few specimens, all from Tabasco, Mexico. Marchand (1946) discovered a large population of these crabs along the Hillsborough River in west Florida, which empties into the Gulf near Tampa. These crabs were found to have originated in Mexico from where they were transported on logs and in lumber for Tampa, beginning about 1915. Marchand (1946) provided notes on ecology, behavior, and feeding habits of the west Florida populations, under the name *P. typicus*. Buitendijk (1950) determined this latter name to be a junior synonym of *P. spectabilis*.

Platychoirapsus typicus Rathbun, 1914.

Determined by Buitendijk (1950) to be a junior synonym of *P. spectabilis* de Man, 1896.

Family GECARCINIDAE Macleay, 1838

Cardisoma Latreille, 1825

Cardiosoma guanhumi Latreille, 1825 (Encycl. Méth., Hist. Nat., Entom., vol. 10, p. 685)

Common Names: Great Land Crab; White Land Crab; Mulatto Land Crab; Juey; Tourlourou; Guanhumi; Guaiamu

Rathbun, 1918, p. 341, text-fig. 155, pls. 106-107; Rathbun, 1933, p. 94, fig. 89; Bright, 1966, p. 191, fig. 4-I; Forest & Guinot, 1966, p. 94; Chace & Hobbs, 1969, p. 195, figs. 64, 67a-c; Türkay, 1970, p. 345; Bright & Hogue, 1972; p. 16; Felder, 1973a, p. 79, pl. 12, figs. 1, 4.

Range: Bermuda; Bahamas; southeast Florida; Florida Keys; Louisiana and south Texas; eastern Mexico to Colombia; north and south coasts of Cuba; Jamaica; Puerto Rico; St. Thomas, Virgin Islands to Barbados; Trinidad; Netherlands Antilles; Colombia to São Paulo, Brazil.

Habitat: low-lying coastal areas, especially mangrove swamps; open fields; along rivers, streams, drainage canals and ditches; under buildings; saline soils with high water tables; primarily nocturnal, but diurnal in heavily shaded areas and on days when the sky is heavily overcast.

Remarks: This is the largest land crab in the Gulf of Mexico region. It is commercially harvested as a food item on some islands of the West Indies, especially Puerto Rico. In other areas, such as southern Florida, this crab is considered an agricultural pest because of the damage caused to fields by the large, extensive burrows and also due to the fondness of the crabs for young, growing shoots.

Regional lists include Louisiana (Behre, 1949, 1950) and Texas (Leary, 1967; Felder, 1973a). Listed from Cuba by Chace (1940) and from Brazil by Coelho and Ramos (1972). Accounts of natural history were provided by Gifford (1962b), Feliciano (1962), and Fotheringham and Brunenmeister (1975). Henning (1975a, 1975b) studied the biology of this crab in northern Columbia, including extensive observations on behavior and ecology. Wright (1968) described agonistic behavior, especially chela displays during social encounters. Herreid (1963) investigated feeding behavior and Herreid and Gifford (1963) reported on the burrow as a habitat and on ionic regulation by the crab.

Developmental stages were described by Moreira (1913) and Costlow and Bookhout (1968b). Costlow and Bookhout (1968c) studied the effects of various environmental factors on development. Physiological studies include work on calcium metabolism (Gifford and Johnson, 1962), growth and morphometrics (Herreid, 1967), various terrestrial adaptations (Bliss, 1963, 1968), aerial respiration (Cameron, 1975), respiratory pigments (Redmond, 1962), and uric acid metabolism (Gifford, 1968), osmoregulation by larval stages (Kalber and Costlow, 1968), neurobiology of autotomy and leg elevation (Moffett, 1975), pericardial organ neurosecretion (Cooke and Goldstone, 1970; Berlind and Cooke, 1970; Berlind, Cooke and Goldstone, 1970) neural control of walking (Barnes, Spirito and Evoy, 1972; Spirito, Evoy and Barnes, 1972; Evoy and Fournier,

1973; Fourtner and Evoy, 1973; Moffett, 1975) and biochemistry (Quinn and Lane, 1966, 1967).

Studies of economic and commercial impact include those of de Oliviera (1946) on the fishery and ecology of this crab in Brazil and Feliciano (1962) on the fishery in Puerto Rico. Humes (1958) described a copepod from the gill chambers of this crab.

***Cecarcinus* Leach, 1814**

Cecarcinus lateralis (Fremenville, 1835) (Ann. Sci. Nat., ser. 2, Zool. 3: 224)

Common Names: Black Land Crab; Common Land Crab

Rathbun, 1918, p. 355, text-fig. 161, pls. 119–120; Rathbun, 1933, p. 95, fig. 91; Chace & Holthuis, 1948, p. 26; Chace & Hobbs, 1969, p. 198, figs. 65, 67e–g; Türkay, 1970, p. 337, figs. 2a–c; Bright & Hogue, 1972, p. 21; Felder, 1973a, p. 82, pl. 12, figs. 2–3; Türkay, 1973, p. 974, fig. 2.

Range: Bermuda; Bahamas; southeast Florida; Florida Keys; south Texas to north coast of Yucatan; north and south coasts of Cuba; Jamaica; Hispaniola; Puerto Rico; St. Thomas, Virgin Islands to Barbados; Netherlands Antilles; Honduras to Costa Rica; Caribbean coast of Columbia to Surinam.

Habitat: burrows in dry, sandy areas; in back dunes and on dune ridges; up to 1000 foot elevation in Dominica (Chace and Hobbs, 1969); in wooded areas of dune ridges and back dunes of eastern Florida, under logs and leaf litter; primarily nocturnal, but diel in heavily wooded habitats.

Remarks: The status of this species and of *G. quadratus* have been unclear for some time. Türkay (1970) listed *G. quadratus* as a subspecies of *G. lateralis*, but an examination of types in the Paris Museum led Türkay (1973) to conclude that the two are synonymous. Most other workers have listed them as distinct and separate species, yet recognizing the complex distribution pattern of *G. quadratus* on both sides of Central America. Some specimens of *G. lateralis* have also been reported from the Pacific coast. Specimens collected in Texas (Ray, 1967; Britton, 1976; personal collections) are of *G. lateralis*.

Regional lists include Texas (Ray, 1967; Britton, 1976; Felder, 1973a; Fotheringham and Brunenmeister, 1975) and Mexico (Cabrera, 1965, zoea only). Listed from Costa Rica by Bright (1966). Reports on ecology and natural history of this species include Bliss and Sprague (1958b), Weitzman (1963), Bliss (1968), Chace and Hobbs (1969), and Klaassen (1975). This crab has been used for a variety of physiological studies, especially for research on molting and regeneration (Hodge, 1956a, 1956b; Bliss, 1960a, 1960b, 1966; Bliss and Boyer, 1964; Bliss *et al.*, 1972; Skinner, 1965, 1966; Skinner and Graham, 1972; Mason, 1970; Holland and Skinner, 1976; Yamaoka and Skinner, 1976). Other physiological reports include work on osmoregulation and water balance (Bliss, 1963; Bliss, Wang and Martinez, 1966; Mantel, 1968; Copeland, 1968), aerial respiration (Cameron, 1975), lipid metabolism (O'Connor and Gilbert, 1968), coagulation (Morrison and Morrison, 1952; Stutman and Dolliver, 1968), neuroendocrinology (Hodge and Chapman, 1958; Bliss and Sprague, 1958a; Maynard, 1961a, 1961b; Maynard and Maynard, 1962; Weitzman, 1969; Mantel *et al.*, 1975),

oxygen transport in hemolymph (Redmond, 1968), saline composition for lab experiments (Skinner, Marsh and Cook, 1965), neural fine structure (Skobe and Nunnemacher, 1970), and sensitivity to substrate vibrations (Klaassen, 1973).

Gecarcinus quadratus Saussure, 1853 (Rev. Mag. Zool., ser. 2, vol. 5, p. 360)

Common Names: Red Land Crab; Whitespot Crab

Rathbun, 1918, p. 358, text-fig. 162, pls. 121-122; Garth, 1948, p. 58; Bright, 1966, p. 190, fig. 4G; Türkay, 1970, p. 338, fig. 4; Bright & Hogue, 1972, p. 20; Türkay, 1973, p. 974.

Range: Veracruz, Mexico to Turbo, Columbia (Carib.); St. Croix; Jamaica; Barbados; in eastern Pacific, from Acapulco, Mexico to La Libertad, Ecuador.

Habitat: well above the high tide mark of sandy beaches; mangrove and other heavily-vegetated areas of marine shores; under debris and other litter.

Remarks: As indicated for *G. lateralis*, this species may be regarded as distinct, as a subspecies of *G. lateralis*, or as completely synonymous with the latter. Türkay (1970) published a biogeographical distribution map of the two forms which shows considerable overlap in the ranges of the two species. The two forms are listed here as separate species, but those workers accepting Türkay's (1973) synonymy of the two forms can combine the references and locality records.

Gecarcinus ruricola (Linnaeus, 1758) (Syst. Nat., ed. 10, vol. 1, p. 626)

Common names: Black Crab; Mountain Crab; Blue Land Crab; Red Tourlourou

Rathbun, 1918, p. 352, text-fig. 160, pls. 117-118; Chace & Holthuis, 1948, p. 26; Chace & Hobbs, 1969, p. 200, figs. 66, 67h-j; Türkay, 1970, p. 336, fig. 1a-f; Bright & Hogue, 1972, p. 20.

Range: Bahamas; southeast Florida; north and south coasts of Cuba; Cayman Islands; Jamaica; Navassa Island (Carib.); Hispaniola; Puerto Rico; St. Croix to Barbados; Curaçao; Old Providence and Swan Islands (Carib.).

Habitat: closer to the intertidal zone than other species of this genus; on low and marshy ground and on lower slopes of island mountains, up to elevations of 500 m; in wooded dune areas of southeast Florida (rare, personal observation).

Remarks: Descriptions of ecology and behavior are found in Rathbun (1918) and Chace and Hobbs (1969); a summary is provided by Bright and Hogue (1972). Listed from the south coast of Cuba by Chace (1940). This species occurs among dense populations of *G. lateralis* in southeast Florida, but is rather rare.

Superfamily OCYPODOIDEA Rafinesque, 1815

Family OCYPODIDAE Rafinesque, 1815

Subfamily OCYPODINAE Rafinesque, 1815

Ocypode Weber, 1795

Ocypode albicans Bosc, 1801-1802.

Junior synonym for *Ocypode quadrata*, used by Rathbun (1918) and others prior to revision by Holthuis (1959).

Ocypoda arenaria Say, 1817.

Junior synonym and invalid generic name for *Ocypode quadrata*, used by Cowles (1908) and some other early studies.

Ocypode quadrata (Fabricius, 1787) (Mantissa insect . . ., vol. 1, p. 315)

Common Names: Ghost Crab; Sand Crab; Racing Crab

As *O. albicans*—Rathbun, 1918, p. 367, pls. 127–128; Rathbun, 1933, p. 96, fig. 92.

As *O. quadrata*—Holthuis, 1959, p. 259; Williams, 1965, p. 225, fig. 208; Chace & Hobbs, 1969, p. 204, figs. 68–69; Felder, 1973a, p. 82, pl. 12, figs. 5, 8.

Range: Bermuda; Bahamas; Rhode Island to south Florida; Florida Keys and Dry Tortugas; west coast of Florida, around entire Gulf coast to Yucatan; north and south coasts of Cuba, through West Indies to Barbados; from Yucatan, along east coast of Central America and the north coast of South America to Estado de Santa Catarina, Brazil; most Caribbean Islands, including Netherlands Antilles, Old Providence Island, etc. Megalops have been collected as far north as Massachusetts.

Habitat: on sandy beaches, from high water line to back dunes areas; younger crabs burrow closer to water line and among beach vegetation; along wave-exposed shores, protected harbor beaches, bays, intracoastal canals, and lagoons; juveniles are mainly diel and older adults are primarily nocturnal, depending on degree of disturbance by man and various environmental factors. Adults usually burrow well back from the waterline, but often feed at the driftline.

Remarks: Regional lists include Florida (Wass, 1955; Menzel, 1971), Mississippi (Richmond, 1962), Louisiana (Behre, 1950; Hoese and Valentine, 1972), and Texas (Whitten, Rosene and Hedgpeth, 1950; Hedgpeth, 1953; Leary, 1967; Fotheringham and Brunenmeister, 1975). Bright and Hogue (1972) include this species in their world-list of land crabs; listed from Brazil by Coelho (1971a), Coelho and Ramos (1972), and Fausto Filho (1974).

Diaz and Costlow (1972) described and illustrated larval stages raised under laboratory conditions. Haley (1969) provided data on growth and morphometrics of Texas populations; reproductive cycling, female morphometrics, and population dynamics were covered in Haley (1967, 1972). Hughes (1973) described mating behavior in the laboratory and compared the mating functions of burrows in several ghost crab species. Population densities and interactions with man were studied by Teerling (1970). Accounts of general natural history were provided by Cowles (1908) for populations in the Tortugas (as *Ocypoda arenaria*) and by Milne and Milne (1946) for New Jersey. Williams (1965) summarized many of the recent studies. Chace and Hobbs (1969) described color phases of Dominican populations. Burrow construction and ecology in Texas was reported by Hill and Hunter (1973) and predatory behavior on mole crabs was noted by Fales (1976). Schone (1968) investigated agonistic displays and these were also presented in a film (Schone and Eibl-Eibesfeldt, 1965).

Physiological studies include work on oxygen consumption (Pearse, 1929; Ayres, 1938; Vernberg, 1956; Gray, 1957), water relations and the role of the pericardial sac (Blass, 1963, 1968), ionic regulation and respiration (Flemister

and Flemister, 1951; Flemister, 1958), ionic and osmotic regulation (Gifford, 1962a), gill and "kidney" histophysiology (Flemister, 1959), biochemistry of terrestrial adaptations (Vernberg and Vernberg, 1968), thoracic neurosecretion (Maynard, 1961a, 1961b; Maynard and Maynard, 1962), and visual perception (Schone and Schone, 1961). Studies of acoustic perception and related behavior include Horch and Salmon (1969), Horch (1971), and Salmon and Horch (1972).

Uca Leach, 1814

(This genus of intertidal ocypodids, along with a few other Australo-Asian genera, are commonly known as fiddler crabs. A number of subgenera have been proposed for this large and diverse genus. Bott (1973) split this genus into 10 genera and Crane (1975) also created a number of subgenera. Although Crane's (1975) monograph is a comprehensive and monumental work, taxonomic precedence must be given to Bott (1973) with regard to most of these proposed changes. For the present, and in agreement with a review by von Hagen (1976), this compilation will avoid the use of subgenera and will continue the use of *Uca*, with the species arranged alphabetically.)

Uca burgersi Holthuis, 1967 (Zool. Meded. Leiden 42: 52)

As *U. mordax*—Rathbun, 1918, p. 391 (part), not text-fig. 166, nor pl. 134, figs. 3-4; Maccagno, 1928, p. 46 (part); de Oliviera, 1939a, p. 138; Holthuis, 1959, p. 265.

As *U. burgersi*—Holthuis, 1967, p. 52; Chace & Hobbs, 1969, p. 207, figs. 70, 71a-d; Gibbs, 1974, p. 84; Crane, 1975, p. 168, figs. 26F, 31H, 54G, 66F, 100, pl. 24E-H, map 12.

Range: Bahamas; east coast of Florida; northeast (Gulf) coast of Yucatan; north and south coasts of Cuba; Jamaica; Hispaniola; Puerto Rico; St. Thomas, Virgin Islands to Trinidad; Curaçao; east coast of Yucatan to Guatamala; Caribbean coast of Panama; Venezuela to Rio de Janeiro, Brazil.

Habitat: sheltered mud flats; sloping mud banks and mud-sand areas at mouths of streams; along shores of lagoons and estuaries; often near mangroves; intertidal; above high tide mark in mangrove thickets of Florida, associated with *U. rapax*.

Remarks: Earlier references confused this species with *U. mordax*, to which it is similar in morphology, ecology, and geographical range. This species was also recorded as *U. affinis* by Holthuis (1959), when he distinguished it from *U. mordax* in Surinam. Records of this crab from west Africa are questioned by Crane (1975). Crane (1957) included data on waving displays as part of her description of *U. mordax*. Adaptation to intertidal zone habitats was reported by von Hagen (1970b) and Salmon (1967) obtained sound recordings of leg-wagging (as *U. mordax*). Gibbs (1974) investigated the ecology of this crab on Barbuda and Gibbs and Bryan (1972) studied cation composition of the exoskeleton.

Uca leptodactyla Rathbun, 1898 (Ann. New York, Acad. Sci. 11: 227)

Rathbun, 1918, p. 420, pl. 156; Maccagno, 1928, p. 41 (part), not text-fig. 25; Rathbun, 1933, p. 98; de Oliveira, 1939a, p. 126, pl. 5, text-figs. 25–28, pl. 6, fig. 29, pl. 8, fig. 47, pl. 13, figs. 61–62; Chace & Hobbs, 1969, p. 212, figs. 71g–h; von Hagen, 1970a, p. 227; Crane, 1975, p. 304, text-figs. 37M, 56F, 60N–O, 69K–L, 101, map 17, pl. 41A–D.

Range: Bahamas; west coast of Florida (not recently); east coast of Yucatan; north coast of Cuba; Jamaica; Puerto Rico; St. Croix; Curaçao; Venezuela to Santa Catarina, Brazil.

Habitat: relatively sandy tidal flats of marine waters; sometimes at supratidal levels, where burrows are covered only by spring tides; occasionally in mud or clay substrates, in partial shade of mangrove trees.

Remarks: Behavioral studies include Matthews (1930), Crane (1957), and Gerlach (1958b). Ecological data were provided by Matthews (1930), de Oliveira (1939a, 1939b, 1939c), and Crane (1957). Bott (1973) lists this species under his genus *Leptuca*; Crane (1975) lists it under her subgenus *Celuca*. Listed from Brazil by Coelho (1971a) and Coelho and Ramos (1972).

Uca longisignalis Salmon & Atsaiades, 1968 (Proc. Biol. Soc. Washington 81: 279)

As *U. pugnax rapax*—Rathbun, 1918, p. 397 (part), not. pl. 140.

As *U. longisignalis*—Salmon & Atsaiades, 1968b, p. 279, text-figs. 1–4, 6, 7.

As *U. rapax longisignalis*—Crane, 1975, p. 190, map 14.

Range: northwest Florida to south Texas.

Habitat: similar to that of *U. rapax*; often on exposed salt flats and algal beds; among marsh vegetation; substrates of mud, mud-sand, and sand-mud.

Remarks: Felder (1973a) comments on the distribution of *U. pugnax*, *U. virens* and *U. longisignalis* and retains the name of *U. pugnax* for all forms in the northwestern Gulf. Crane (1975) listed this form as a subspecies of *U. rapax*, which until recently was a subspecies of *U. pugnax*. In a review of Crane's (1975) monograph, von Hagen (1976) commented that an examination of Salmon's holotypes left "no doubt that *U. rapax longisignalis* is a synonym of *U. minax* (Le Conte)." The present list treats this crab as a separate species, mainly on the basis of Salmon and Atsaiades (1968b) evidence of behavioral separation and on the lack of agreement among other authorities as to which subspecies or species it is most closely allied with. Subrahmanyam *et al.* (1976) listed this crab from northwest Florida and Powers (1975) noted its occurrence in Texas. Included in the key to northwestern Gulf *Uca* by Fotheringham and Brunenmeister (1975). Abele (1970) provided habitat notes on populations at Alligator Harbor, Florida.

Uca minax (Le Conte, 1855) (Proc. Acad. Nat. Sci. Philadelphia 7: 403)

Common Names: Red-Jointed Fiddler Crab; Brackish Water Fiddle Crab

Hay & Shore, 1918, p. 451, pl. 37, fig. 3; Rathbun, 1918, p. 389, pl. 137; Maccagno, 1928, p. 48, text-fig. 31; Crane, 1943a, p. 220, text-fig. 1b; Williams, 1965, p. 227, figs. 209A, 210B; Felder, 1973a, p. 85, pl. 12, fig. 12; Crane, 1975, p. 176, figs. 67D, 81K, 100, pl. 25E–H, map 12.

Range: Massachusetts to northeast Florida; northwest Florida to Louisiana, possibly Texas.

Habitat: brackish to fresh waters of estuaries, bays and streams; drainage ditches and canals; usually located some distance from marine waters, but often subject to some tidal influence, particularly along the Atlantic coast; burrows in mud banks and among marsh vegetation, often supratidal; occasionally at edges of fields or woodlands.

Remarks: Crane (1975) questioned the presence of this species from Texas, stating that records listed by Rathbun (1918) from Texas were found to be *U. rapax longisignalis* and *U. pugnax virens*; von Hagen (1976) believes that *U. (rapax) longisignalis* is a synonym of *U. minax*. If the latter is true, then the range of *U. minax* would definitely include Texas, but the form described by Salmon and Atsaiades (1968b) as *U. longisignalis* does not inhabit freshwater and brackish areas. Listed from northwest Florida by Wass (1955), Abele (1970), and Menzel (1971), from Louisiana by Behre (1950) and Felder (1973a), and from Texas by Leary (1967), but the latter is probably based on Rathbun (1918).

Gray (1942) and Miller (1965) provided descriptions of natural history and Williams (1965) summarized much of the current literature. Hyman (1920, 1922) described post-larval development and behavior, including spawning. Ecological studies include Teal (1958) in Georgia, Miller and Maurer (1973) on distribution in relation to salinity, Whiting (1972) and Whiting and Moshiri (1974) on distribution in relation to substrate, and Kerwin (1971) on distribution in relation to marsh vegetation. Salmon (1967) studied distribution in Florida. Miller (1961) compared feeding adaptations in this and other *Uca* species. Physiological studies include work on gill area (Gray, 1957), osmotic and ionic regulation (Cole, 1971), the relationship between respiration and habitat (Teal, 1959), tolerance to desiccation (Pearse, 1929), acclimation to temperature (Vernberg, 1959), tidal rhythms of color change (Fingerman, Lowe and Moberly, 1958), and radiation sensitivity (Engel, 1973). Nimmo *et al.* (1971) studied PCB absorption from sediments. Behavioral studies include descriptions of waving displays (Crane, 1943a, 1957; Salmon, 1965), sound production (Salmon, 1965), and vibration reception (Salmon and Horch, 1973).

Uca mordax (Smith, 1870).

Although recorded from the Gulf of Mexico by Rathbun (1918, p. 391), these and earlier records had confused this species with *U. vocator*, *U. rapax*, and *U. burgersi*. Crane (1975, p. 173) restricts *U. mordax* to the continental coast, from Guatamala to Brazil, plus the island of Trinidad.

Uca panacea Novak & Salmon, 1974 (Proc. Biol. Soc. Washington 87: 313)

Common Name: Sand Fiddler Crab

As *U. pugilator*—Rathbun, 1918, p. 400 (part), not pl. 141 and pl. 169, fig. 2; Felder, 1973a, p. 83; Crane, 1975, p. 223 (part), not pl. 29E–H, part map 16, other figs. indet.

As *U. panacea*—Novak & Salmon, 1974, p. 316, figs. 1–8.

Range: northwest Florida to south Texas.

Habitat: sandy areas of marshes and tidal flats; often supratidal, intertidal in some areas; well inland on Texas barrier islands; similar to habitats of *U. pugilator* when intertidal.

Remarks: This species is morphologically similar to *U. pugilator*, but it has been separated from the latter on the basis of biochemical (Selander, Johnson and Avise, 1971) and behavioral studies (Novak and Salmon, 1974). Color variants of "*U. pugilator*" from Florida were noted by Rao & Fingerman (1968), a species-specific character present only in living specimens. It is likely that many of the studies reported as "*U. pugilator*," when collected from the central and western Gulf coasts, actually utilized *U. panacea*. *Uca pugilator* has been collected as far west as central Texas (Carl Thurman, pers. comm.; pers. observ. of author), indicating greater geographical overlap between the two species than reported by Novak and Salmon (1974). Other authors (Felder, 1973a; Crane, 1975) regarded the two forms as one species. The southern limits of *U. panacea* have not been defined yet, but may extend into northeastern Mexico. Hedgpeth (1950) commented on the ecology of this crab on salt flats in Texas and Powers (1973) provided data on burrow densities. Ecological and behavioral data on Texas barrier island populations were presented by Powers (1975) and Powers and Cole (1976). Studies prior to 1974, using the name *U. pugilator*, may include either or both species; regional lists and references are listed under *U. pugilator*, but many of these studies are undoubtedly of *U. panacea*.

Uca pugilator (Bosc, 1802) (Hist. Nat. Crust., vol. 1, an X, p. 197)

Common Name: Sand Fiddler Crab

Hay & Shore, 1918, p. 452, pl. 37, fig. 2; Rathbun, 1918, p. 400 (part), pl. 141, pl. 160, fig. 2; Maccagno, 1928, p. 44, text-fig. 28; Crane, 1943a, p. 220; Williams, 1965, p. 232, figs. 209C, 210C-D, 211; Crane, 1975, p. 223 (part), text-figs. 37K, 69F, 101, pl. 29E-H, map 16 (part).

Range: Bahamas; Massachusetts to south Florida; Florida Keys; west and northwest coasts of Florida; Mississippi to Texas; ? Santo Domingo; ? Old Providence Island (Carib.)

Habitat: sandy and sand-mud substrates; intertidal to supratidal marshes; burrows on open sand flats or among thick clumps of grasses and other vegetation.

Remarks: This species is listed by Bott (1973) in the genus *Planuca* and by Crane (1975) in the subgenus *Celuca*. The status of the Caribbean specimens needs to be re-examined with respect to *U. pugilator* and *U. panacea*. Because of the widespread use of this animal in experimental studies, the taxonomic relationships and variability of morphological, behavioral, and physiological features need to be defined and established for both of these closely-related forms. A partial listing of the large literature on *U. pugilator*, including some of *U. panacea*, follows.

Accounts of natural history include Pease (1914) in Massachusetts, Schwartz and Safir (1915) in New York, and studies by O. W. Hyman (1920, 1922) and Dembowski (1925, 1926). Developmental studies were reported by Hyman

(1920), Gray (1942) on transient prezoa, and by Hernkind (1968b). Miller (1968) investigated asymmetry during growth and Vernberg and Costlow (1966) studied handedness. Ecological studies include work on habitat preferences in Georgia (Teal, 1958), feeding efficiency (Miller, 1961), habitats in the Bahamas (Coventry, 1944), habitats in Massachusetts (Knopf, 1966), distribution in relation to thermal tolerance (Miller and Vernberg, 1968), thermal relations of crab and microhabitat (Smith and Miller, 1973), and capture-recapture methods (Hockett and Kritzler, 1972).

Sand Fiddlers have been the subjects of many behavioral studies: displays and courtships (Pease, 1914; Dembowski, 1925, 1926; Crane, 1943a, 1957), threat displays (Schone, 1968; Aspey, 1971), sound production and visual signals (Burkenroad, 1947; Salmon and Stout, 1962; Salmon, 1965, 1967; Salmon and Atsides, 1969; Salmon and Horch, 1972), burrowing activity (Teal, 1958; Coward, Gerhardt and Crockett, 1970), visual orientation (Herrnkind, 1968a, 1968c, 1972), feeding (Miller, 1961), locomotion (Baird and Burleson, 1970), and larval shadow responses (Forward, 1977).

Physiological studies include work on molting (Abramowitz and Abramowitz, 1940; Guyselman, 1953; Stewart and Green, 1969; Skinner and Graham, 1972; Fingerman and Fingerman, 1976; Weis, 1976a), regeneration (Weis, 1976b, 1976c, 1977a, 1977b; Weis and Mantel, 1976), color changes and chromatophores (Carlson, 1935, 1936; Brown and Sandeen, 1948; Brown and Webb, 1948; Brown, 1950; Guyselman, 1953; Webb, Bennett and Brown, 1954; Fingerman and Yamamoto, 1967; Barnwell, 1968a; Rao and Fingerman, 1968; Fingerman, Rao and Ring, 1969; Coohill and Fingerman, 1975), metabolism (W. B. Vernberg and Vernberg, 1972), rhythmical activity and physiology (Brown *et al.*, 1955; Fingerman, 1956, 1957; Fingerman, Lowe and Mobberly, 1958; Barnwell, 1966, 1968b), biochemistry (Eisen *et al.*, 1973), sensitivity to anemone toxin (Blanquet, 1968), reproduction and endocrinology (Darby, 1935; Brown and Jones, 1949; Sandeen, 1950; Fingerman and Fitzpatrick, 1956; Fingerman and Couch, 1967; Rao, Fingerman and Bartell, 1967; Rao and Fingerman, 1969, 1970; Fingerman, 1970, 1973; Bartell, Rao and Fingerman, 1971; Fielder, Rao and Fingerman, 1971), thermoregulation and temperature adaptations (Edwards, 1950; Orr, 1955; Demeusy, 1957; Wilkins and Fingerman, 1965; Vernberg, DeCoursey and Padgett, 1973), osmoregulation (Pease, 1929; Teal, 1958; Green *et al.*, 1959; Evans, Cooper and Bogan, 1976), toxicology (Nimmo *et al.*, 1971; DeCoursey and Vernberg, 1972; O'Hara, 1973), respiration (Gray, 1957; Teal, 1959; Wilkins and Fingerman, 1965; Smith and Miller, 1973; Silverthorn, 1975a, 1975b), sensory perception (Salmon and Atsides, 1969; Horch and Salmon, 1969; Salmon, 1971; Langdon, 1971; Avent, 1974; Hyatt, 1974, 1975; Salmon, Horch and Hyatt, 1977), neurobiology (Nunne-macher, 1965; Andrews, 1973), radiation sensitivity (Engel, 1973), and infection by bacteria (Spindler-Barth, 1976).

Regional lists include Florida (Wass, 1955; Tabb and Manning, 1961; Menzel, 1971), Louisiana (Behre, 1950; Hoese and Valentine, 1972), and Texas (Hedgpeth, 1950, 1953; Whitten, Rosene and Hedgpeth, 1950; Simmons, 1957; Leary, 1967), but the Louisiana and Texas records probably refer to *Uca panacea*.

Uca pugnax (Smith, 1870).

This is another species with a history of frequent nomenclatural changes. Most older references have included this crab as a Gulf species, but Salmon and Atsaiades (1968b) referred the Gulf populations of *U. pugnax* to new species, *U. virens* and *U. longisignalis*. Crane (1975) maintained *U. virens* as a subspecies of *U. pugnax*, but she placed *U. longisignalis* as a subspecies of *U. rapax*. Until a revision by Tashian and Vernberg (1958), *U. rapax* was considered a subspecies of *U. pugnax*; thus all four forms are closely related. However, von Hagen (1976) synonymizes *U. virens* with *U. rapax* and considers *U. longisignalis* to be synonymous with *U. minax*! The present list treats each form separately, maintaining each species presented by Salmon and Atsaiades (1968b) and excluding *U. pugnax* as a Gulf species.

Uca rapax (Smith, 1870) (Trans. Connecticut Acad. Arts Sci. 2: 134)

As *U. pugnax rapax*—Rathbun, 1918, p. 397 (part), pl. 140; Maccagno, 1928, p. 45, text-fig. 29; Rathbun, 1933, p. 97; de Oliveira, 1939a, p. 134.

As *U. rapax*—Tashian & Vernberg, 1958, Holthuis, 1959, p. 266, text-figs. 64d-f, 65, pl. 14, figs. 4-6, pl. 15, fig. 3; Chace & Hobbs, 1969, p. 214, figs. 73a-b; von Hagen, 1970a, p. 226; Crane, 1975, p. 190, figs. 52C-DD, 54F, 67C, 86, 91E-F, 100, pls. 27A-D, 45C-F, map 14.

Range: Bahamas; east coast of Florida; Florida Keys; southwest coast of Florida; northeast coast of Mexico to northeast Yucatan; north and south coasts of Cuba; Jamaica; Hispaniola; Puerto Rico; St. Thomas, Virgin Islands to Trinidad and Tobago; Netherlands Antilles; east coast of Yucatan to Guatamala; Caribbean coast of Panama to Santa Catarina, Brazil.

Habitat: mud, sand-mud, and mud-sand flats; edges of mangroves; along rivers and streams on flats and banks.

Remarks: This species may also occur infrequently along the northwestern Gulf coast, but Crane (1975) attributes records of this crab to *U. rapax longisignalis*. Felder (1973a) listed *U. rapax* from the same area, but past records may be erroneous with regard to the several similar species involved. Listed from Brazil by Coelho and Ramos (1972).

Behavioral studies include observations on waving displays (Crane, 1943a, 1957), combat between males (Crane, 1957, 1967), visual and acoustical signalling (Salmon and Atsaiades, 1968a), kinaesthetic orientation (von Hagen, 1967), orientation to burrows (von Hagen, 1970b), and feeding (Miller, 1965). Warner (1969) studied the natural history of this crab in Jamaica and Holthuis (1959) provided ecological notes and populations in Surinam. Smith and Miller (1973) measured thermal adaptations. Barnwell (1963) observed motor activity and the rhythmicity of color changes in populations in Brazil. Handedness and its relationship to development was analyzed by Vernberg and Costlow (1966). Adaptations to particular tidal levels were observed by von Hagen (1970b). Salmon (1971) measured vibration receptivity and van Delft (1968) studied daily rhythms of color changes.

Uca speciosa (Ives, 1891) (Proc. Nat. Acad. Sci. Philadelphia 1891: 179)

Rathbun, 1918, p. 408, pl. 145; Chace & Hobbs, 1969, p. 215, figs. 73c-d; von Hagen, 1970a, p. 227; Crane, 1975, p. 236, text-figs. 68G, 101, map 15, pl. 31A-D.

Range: southeast Florida; Florida Keys; west and northwest coasts of Florida; northeast Yucatan and northwest Cuba.

Habitat: wet, muddy substrates; mid to high intertidal zone; commonly found in mangroves.

Remarks: Specimens from Curaçao reported by Rathbun (1918) were referred to *U. cumulanta* by Chace and Hobbs (1969); Crane (1975) referred the Jamaican specimen of Chace and Hobbs (1969) to *U. cumulanta*. Crane (1957) provided some preliminary data on courtship displays and Salmon (1967) analyzed waving patterns of the crabs. Miller (1965) studied the distribution and ecology of this species. Listed from Florida by Wass (1955), Tabb and Manning (1961) and Subrahmanyam *et al.* (1976).

Uca spinicarpa Rathbun, 1900 (Amer. Natural. 34: 586)

As *U. spinicarpa*—Rathbun, 1918, p. 411, pl. 148; Felder, 1973a, p. 83, pl. 12, fig. 11.

As *U. speciosa spinicarpa*—Crane, 1975, p. 239, figs. 68K, 101, pl. 31E-H, map 15.

Range: Alabama to northeastern coast of Mexico.

Habitat: muddy banks of coastal freshwater ponds and streams; muddy, brackish beaches of the Gulf; grassy mud flats off bays (after Felder, 1973a).

Remarks: This crab was considered a separate, but allied species to *U. speciosa* by Rathbun (1918) and subsequent workers. Crane (1975) placed it as a subspecies of the latter. Felder (1973a) listed several personal collections in Louisiana and Mississippi. Listed from Texas by Leary (1967) and Fotheringham and Brunenmeister (1975) comment on its presence in the northwestern Gulf, providing a key for comparison with other *Uca* species. Bott (1973) placed this species and *U. speciosa* in his genus *Leptuca*; Crane's (1975) subgenus designation is *Celuca*.

Uca subcylindrica (Stimpson, 1859) (Ann. Lyc. Nat. Hist. New York 7: 63)

Rathbun, 1918, p. 419, pl. 155, pl. 160, fig. 5; Felder, 1973a, p. 83, pl. 12, fig. 10; Crane, 1975, p. 209, figs. 67, 100, pl. E-H, map 11.

Range: Texas to northeastern coast of Mexico.

Habitat: banks of freshwater streams; brackish water areas; on mud flats and algal beds, often some distance upstream from mouths of rivers and creeks.

Remarks: This species is uncommon and has a restricted range. Listed by Fotheringham and Brunenmeister (1975) for the northwestern Gulf. Very little is known about this crab's ecology, behavior, or other biological aspects.

Uca thayeri Rathbun, 1900 (Proc. Washington Acad. Sci. 2: 134)

Rathbun, 1918, p. 406, text-fig. 169, pl. 144; Rathbun, 1933, p. 98; Holthuis, 1959, p. 275, text-figs. 68b-c, pl. 16; Chace & Hobbs, 1969, p. 216, text-figs. 73e-f; von Hagen, 1970a, p. 226; Crane, 1975, p. 112, figs. 46K, 56E, 60H-I, 73A-B, 81I, 82I, 99, map 11, pl. 17.

Range: east and southwest coasts of Florida; north and south coasts of Cuba;

Jamaica; Hispaniola; Puerto Rico; Guadeloupe; Trinidad; Tobago; Guatamala and Panama (Caribbean coasts) Venezuela to São Paulo, Brazil.

Habitat: deep mud on banks of streams and estuaries, among mangrove swamps; burrows are often shaded by vegetation.

Remarks: Ecological studies include Gerlach (1958a) in Brazil Warner (1969) in Jamaica, and Salmon (1967) in Florida. Crane (1957) described daily behavioral displays and Barnwell (1963) reported on daily and tidal rhythms of activity. Bott (1973) placed this species in his genus *Planuca*; Crane (1975) designated the subgenus *Bcboruca*. Listed from Brazil by Coelho and Ramos (1972).

Uca virens Salmon & Atsoides, 1968 (Proc. Biol. Soc. Washington 81: 281)

As *U. pugnax*—Felder, 1973a, p. 84.

As *U. virens*—Salmon & Atsoides, 1968b, p. 281, figs. 2-3, 5-7.

As *U. pugnax virens*—Crane, 1975, p. 203, map 10.

As *U. rapax*—von Hagen, 1976, p. 224.

Range: Mississippi to Coatzacoalcos (central Gulf coast), Mexico.

Habitat: muddy sand, sand-mud, and mud substrates of salt marshes; algal flats, close to bays, estuaries and inlets; often among marsh vegetation.

Remarks: Studies that have recorded "*Uca pugnax*" from the Gulf coast may represent records of *U. virens* or *U. longisignalis*; the occasional presence of *U. rapax* along the Texas and Mexico coasts is also possible. Salmon and Atsoides (1968b) reported analyses of waving and acoustic signalling. Powers (1975) and Powers and Cole (1976) provided some data on habitats of this crab in Texas. See *Uca pugnax* and *Uca longisignalis* for a discussion of nomenclatural changes in the *pugnax-rapax* species group.

Uca vocator (Herbst, 1804) (Versuch. Natur. Krabben u. Krebse, vol. 1. pl. 59, fig. 1)

As *U. mordax*—Rathbun, 1918, p. 391 (part), pl. 134, figs. 3-4.

As *U. murifecenta*—Crane, 1943b, p. 38, text-figs. 1d-f, pl. 1, figs. 1-2.

As *U. vocator*—Holthuis, 1959, p. 269, text-figs. 66-67, pl. 14, fig. 1, pl. 15, fig. 1;

Chace & Hobbs, 1969, p. 217, figs. 73g-j, 74; von Hagen, 1970a, p. 225; Crane, 1975, p. 27, figs. 66D, 100, pl. 23E-G, pl. 24A-D, map 13.

Range: Tampico, Mexico; Belize to Guyana; Puerto Rico; Santo Domingo; Guadeloupe; Dominica; Trinidad and Tobago; Paraiba to Pernambuco, Brazil; ? Santa Catarina, Brazil.

Habitat: grassy marshes; mud flats; flat banks of streams and rivers; in damp mud among mangroves.

Remarks: This species was presented by Rathbun (1918) with *U. mordax*. Ecological data includes studies by Crane (1943b) in Venezuela, by Holthuis (1959) in Surinam, by Chace and Hobbs (1969) in Dominica, and by von Hagen (1970a, 1970c), who also commented on sound production and other aspects of behavior. The only record for this species from the Gulf of Mexico is that listed by Crane (1975) for Tampico, Mexico. Listed from Brazil by Coelho and Ramos (1972) for Paraiba and Pernambuco, but Crane (1975) questions the record from Santa Catarina.

Ucides Rathbun, 1897*Ucides cordatus* (Linnaeus, 1763) (Amoen. Acad., vol. 6, p. 414)

Common Names: Pagurus; Kaburi

Rathbun, 1918, p. 347, text-fig. 158, pls. 110–113, pl. 159, figs. 3–4; Rathbun, 1933, p. 95, fig. 90; Bott, 1955, p. 66; Bright, 1966, p. 191; Chace & Hobbs, 1969, p. 219, figs. 75–76; Türkay, 1970, p. 351, fig. 10a–d; Bright & Hogue, 1972, p. 14.

Range: Bahamas; southeast Florida; northeast Mexico to Panama; north and south coasts of Cuba; Jamaica; Hispaniola; Puerto Rico; St. Thomas, Virgin Islands to Grenada; Colombia to Santa Catarina, Brazil.

Habitat: swampy ground, among mangrove roots; in areas of standing brackish water; on mud flats, among *Uca* and *Cardisoma* burrows.

Remarks: Chace and Hobbs (1969) transferred this genus to the Ocypodidae from the Gecarcinidae and Türkay (1970) accepted this transfer. Although the genus did not fit conveniently into any of the existing subfamilies, Chace and Hobbs (1969) felt that it was more closely allied with the Ocypodinae than with others. Türkay (1970) placed the Pacific species, *U. occidentalis*, under *U. cordatus* as a subspecies. Manning and Provenzano (1961) comment on *Ucides* in Florida. Ecological and behavioral notes were provided by Chace and Hobbs (1969) and Bright and Hogue (1972). Warner (1969) discussed the ecology of this crab in Jamaica and de Oliveira (1946) studied its biology in Brazil. Listed from Brazil by Coelho and Ramos (1972). De Souza and Caland (1968) reported on bacterial infections in this species. Ogawa *et al.* (1973a, 1973b) described commercial processing of this crab for food in Brazil. Alves (1975) studied reproductive biology of Brazilian populations.

BIBLIOGRAPHY

- ABBOTT, W. 1967. Unusual climbing behavior by *Callinectes sapidus* Rathbun (Decapoda, Brachyura). *Crustaceana*. **13**: 128.
- ABELE, L. G. 1970. The marine decapod Crustacea of the northwestern Gulf of Mexico. Masters Thesis, Florida State University, Tallahassee. 137 pp.
- . 1971. Scanning electron photomicrographs of brachyuran genopods. *Crustaceana*. **21**: 218–220.
- . 1972a. A reevaluation of the *Neopanope texanasi* complex with notes, on *N. packardii* (Crustacea: Decapoda: Xanthidae) in the northwestern Atlantic. *Chesapeake Sci.* **13**: 263–271.
- . 1972b. The status of *Sesarma angustipes* Dana, 1852, *S. trapezium* Dana, 1852, and *S. miersii* Rathbun, 1897 (Crustacea: Decapoda: Grapsidae) in the western Atlantic. *Caribb. J. Sci.* **12**: 165–170.
- . 1973. Taxonomy, distribution and ecology of the genus *Sesarma* (Crustacea, Decapoda, Grapsidae) in eastern North America, with special reference to Florida. *Am. Midl. Nat.* **90**: 375–386.
- . 1974. Species diversity of decapod crustaceans in marine habitats. *Ecology*. **55**: 156–161.
- ABRAMOWITZ, R. K. and A. A. ABRAMOWITZ. 1940. Moulting, growth, and survival after eyestalk removal in *Uca pugnator*. *Biol. Bull.* **78**: 179–188.

- ADKINS, G. 1972a. Notes on the occurrence and distribution of the rhizocephalan parasite (*Loxothylacus texanus* Boschma) of the blue crabs (*Callinectes sapidus* Rathbun) in Louisiana estuaries. *Tech. Bull. La. Wildl. Fish. Comm.* **2**: 1-13.
- . 1972b. A study of the blue crab fishery in Louisiana. *Tech. Bull. La. Wildl. Fish. Comm.* **3**: 1-57.
- AIKAWA, H. 1929. On larval forms of some Brachyura. *Rec. oceanogr. Wks. Japan.* **2**: 17-55.
- . 1937. Further notes on brachyuran larvae. *Rec. oceanogr. Wks. Japan.* **9**: 87-162.
- ALDRICH, J. C. 1974. Allometric studies on energy relationships in the spider crab *Libinia emarginata* (Leach). *Biol. Bull.* **147**: 257-273.
- . 1976. The spider crab, *Libinia emarginata* Leach, 1815 (Decapoda, Brachyura), and the starfish, an unsuitable predator but a cooperative prey. *Crustaceana.* **31**: 151-156.
- ALVES, M. I. M. 1974. Resistencia a variacoes de salinidade apresentada por *Pachygrapsus transversus* (Gibbes, 1850)—Crustacea, Grapsidae. *Arquivos de Ciencias do Mar.* **14**: 91-93.
- . 1975. Sobre a reprodução do caranguejo-Uçá, *Ucides cordatus* (Linnaeus), em mangues do estado do Ceará (Brasil). *Arquivos de Ciencias do Mar.* **15**: 85-91.
- ANDREWS, P. M. 1973. Ultrastructural study of the pericardial organ-anterior ramifications complex neurosecretory terminals. *Z. Zellforsch. mikrosk. Anat.* **144**: 309-324.
- , D. E. COPELAND and M. FINGERMAN. 1971. Ultrastructural study of the neurosecretory granules in the sinus gland of the blue-crab. *Callinectes sapidus*. *Z. Zellforsch. mikrosk. Anat.* **113**: 461-471.
- ANTIONE, J. W. 1972. Structure of the Gulf of Mexico. In: R. Rezak and V. J. Henry (eds.), *Contributions on the Geological and Geophysical Oceanography of the Gulf of Mexico*. Gulf Publishing Co.: Houston. Pp. 1-34.
- ASPEY, W. P. 1971. Inter-species sexual discrimination and approach-avoidance conflict in two species of fiddler crabs, *Uca pugnax* and *Uca pugilator*. *Anim. Behav.* **19**: 669-676.
- AVENT, R. M. 1974. The effects of hydrostatic pressure on living aquatic organisms. VIII. Behavioral and metabolic responses of *Uca pugilator* to variations in hydrostatic pressure and temperature. *Int. Revue Ges. Hydrobiol. Hydrogr.* **59**: 219-238.
- AYERS, J. C. 1938. Relationship of habitat to oxygen consumption by certain estuarine crabs. *Ecology.* **19**: 523-527.
- BAIRD, J. L., JR. and A. L. BURLESON. 1970. An analysis of locomotor behavior in the fiddler crab *Uca pugilator*. *Am. Zoologist.* **10**: 500.
- BALLARD, B. S. and W. ABBOTT. 1969. Osmotic accommodation in *Callinectes sapidus* Rathbun. *Comp. Biochem. Physiol.* **29**: 671-687.
- BALSS, H. 1957. Decapoda. In: H. G. Broun, *Klassen und Ordnungen des Tierreichs*, volume 5, Crustacea. C. F. Winter and Akademische Verlagsgesellschaft (Leipzig). Lief. 12, pp. 1505-1672, figs. 1131-1199; Lief. 13, pp. 1673-1770, figs. 1200-1212.
- BARNARD, K. H. 1950. Descriptive catalogue of South African decapod Crustacea (Crabs and Shrimps). *Ann. S. Afr. Mus.* **38**: 1-837.
- BARNES, W. J. P., C. P. SPIRITO, and W. E. EVOY. 1972. Nervous control of walking in the crab, *Cardisoma guanhumi*. II. Role of resistance reflexes in walking. *Z. vergl. Physiol.* **76**: 16-31.
- BARNWELL, F. H. 1963. Observations on daily and tidal rhythms in some fiddler crabs from equatorial Brazil. *Biol. Bull.* **125**: 399-415.
- . 1966. Daily and tidal patterns of activity in individual fiddler crabs (genus *Uca*) from the Wood's Hole region. *Biol. Bull.* **130**: 1-13.
- . 1968a. Comparative aspects of the chromatophoric responses to light and temperature in fiddler crabs of the genus *Uca*. *Biol. Bull.* **134**: 221-234.

- . 1968b. The role of rhythmic systems in the adaptation of fiddler crabs to the intertidal zone. *Am. Zoologist*. **8**: 569-583.
- BARR, L. 1971. Observations on the biology of the arrow crab, *Stenorhynchus seticornis* (Herbst) in Lameshur Bay, St. John, Virgin Islands. pp. 213-220. In: J. Miller, J. van Derwalker and R. Walkers (eds.), *Scientists-in-the-sea*. Dept. of Interior, Washington, D.C.
- . 1975. Biology and behavior of the arrow crab, *Stenorhynchus seticornis* (Herbst), in Lameshur Bay, St. John, Virgin Islands. In: Results of the Tektite Program. *Bull. Mus. Nat. Hist. Los Angeles County*. **20**: 47-56.
- BARTELL, C. K., K. RANGA RAO and M. FINGERMAN. 1971. Comparison of the melanin-dispersing fractions in extracts prepared initially in ethanol, saline, or distilled water from eyestalks of the fiddler crab, *Uca Pugilator*. *Comp. Biochem. Physiol.* **38**: 17-36.
- BEACH, N. W. 1969. The oyster crab, *Pinnotheres ostreum* Say, in the vicinity of Beaufort, North Carolina. *Crustaceana*. **17**: 187-199.
- BEHRE, E. H. 1949. Notes on the occurrence of *Cardisoma guanhumi* Latreille at Grand Isle, Louisiana. *Proc. La. Acad. Sci.* **12**: 19-22.
- . 1950. Annotated list of the fauna of the Grand Isle region. *Occas. Papers mar. Lab. La. State Univ.* **6**: 1-66.
- . 1954. Decapoda of the Gulf of Mexico. *Fishery Bull. Fish Wildl. Serv. U.S.* **55**: 451-455.
- BERLIND, A. and I. M. COOKE. 1970. Release of a neurosecretory hormone as peptide by electrical stimulation of crab pericardial organs. *J. exp. Biol.* **53**: 679-686.
- , ——— and M. W. GOLDSTONE. 1970. Do the monoamines in crab pericardial organs play a role in peptide neurosecretion? *J. exp. Biol.* **53**: 669-677.
- BINFORD, R. 1913. The germ-cells and the process of fertilization in the crab, *Menippe mercenaria*. *J. Morphol.* **24**: 147-200.
- BLAND, C. E., D. G. RUCH, B. R. SALSER and D. V. LIGHTNER. 1976. Chemical control of *Lagenidium*, a fungal pathogen of marine Crustacea. Sea Grant Publ., Univ. North Carolina, **UNC-SG-76-02**. 38 pp.
- BLANQUET, R. 1968. Properties and composition of the nematocyst toxin of the sea anemone, *Aiptasia pallida*. *Comp. Biochem. Physiol.* **25**: 893-902.
- BLISS, D. E. 1960a. Autotomy and regeneration. In: T. H. Waterman (ed.), *The Physiology of Crustacea*. Vol. I. Academic Press. Pp. 561-589.
- . 1960b. Locomotor activity of land crabs during the premolt period. *Science*. **132**: 145-147.
- . 1963. The pericardial sacs of terrestrial Brachyura. In: H. B. Whitting and W. D. I. Rolfe (eds.), *Phylogeny and Evolution of Crustacea*. Mus. Comp. Zool. Spec. Publ., Harvard Univ. pp. 59-78.
- . 1966. Introduction: relation between reproduction and growth in decapod crustaceans. *Am. Zoologist*. **6**: 231-233.
- . 1968. Transition from water to land in decapod crustaceans. *Am. Zoologist*. **8**: 355-392.
- and J. R. BOYER. 1964. Environmental regulation of growth in the decapod crustacean *Gecarcinus lateralis*. *Gen. Comp. Endocrin.* **4**: 15-41.
- , J. ROUILLON BOYER, P. M. CONNELL and S. W. SHEEHAN. 1972. Bioassaying for crustacean limb growth-controlling factors. *Am. Zoologist*. **12**: abst. 105.
- and P. C. SPRAGUE. 1958a. The level of activity in *Gecarcinus lateralis* after eyestalk removal. *Anat. Rec.* **132**: 417.

- and P. C. SPRAGUE. 1958b. Diurnal locomotor activity in *Gecarcinus lateralis*. *Anat. Rec.* **132**: 416-417.
- , S. M. E. WANG and E. A. MARTINEZ. 1966. Water balance in the land crab, *Gecarcinus lateralis*, during the intermolt cycle. *Am. Zoologist*. **6**: 197-212.
- BOESCH, D. F. 1971. On the occurrence of *Pinnixa lunzi* Glassell (Decapoda, Pinnotheridae) off Virginia, U.S.A. *Crustaceana*. **20**: 219-220.
- BOLIVAR y PIELTAIN, C. 1945. Notas sobre *Platychirograpsus typicus* Rathb. (*Dec. Graps.*). *Ciencia, Mex.* **6**: 267-270.
- BONAVENTURA, C., B. SULLIVAN, J. BONAVENTURA and S. BOURNE. 1974. CO binding by hemocyanins of *Limulus polyphemus*, *Busycon carica* and *Callinectes sapidus*. *Biochemistry*. **13**: 4784-4789.
- BOOKHOUT, C. G. and J. D. COSTLOW, JR. 1974. Larval development of *Portunus spinicarpus* reared in the laboratory. *Bull. mar. Sci.* **24**: 20-51.
- , and ———. 1976. Effects of Mirex, Methoxychlor, and Malathion on development of crabs. Report, EPA-600-3-76-007 (NTIS PB-252 007/OST). 96 pp.
- , A. J. WILSON, JR., T. W. DUKE and J. I. LOWE. 1972. Effects of mirex on the larval development of two crabs. *Water, Air, Soil Pollut.* **1**: 165-180.
- BOONE, W. R. and D. L. CLAYBROOK. 1977. The effect of low salinity on amino acid metabolism in the tissues of the common mud crab, *Panopeus herbstii* (Milne-Edwards). *Comp. Biochem. Physiol.* **57A**: 99-106.
- BOTT, R. 1955a. Dekapoden (Crustacea) aus El Salvador. 2. Litorale Dekapoden ausser *Uca*. *Senckenberg. Biol.* **36**: 45-70.
- . 1955b. Die Susswasserkrabben von Afrika (Crust., Decap.) und ihre Stammesgeschichte. *Annls Mus. r. Congo beleg Ser., Zool.* **1**: 209-352.
- . 1968. Fluss-Krabben aus dem östlichen Mittel-Amerika und von dem Grossen Antillen (Crustacea, Decapoda). *Senckenberg. Biol.* **49**: 39-49.
- . 1973. Die verwandtschaftlichen Beziehungen der *Uca*-Arten (Decapoda: Ocypodidae). *Senckenberg. Biol.* **54**: 315-325.
- BOURNE, C. C. 1922. The Raninidae: a study in charcinology. *J. Linn. Soc., Zool.* **35**: 25-79.
- BOUVIER, E. L. 1940. Décapodes marcheurs. *Faune de France* **37**: 1-404.
- BREUER, J. P. 1962. An ecological survey of the lower Laguna Madre of Texas. *Publs Inst. mar. Sci. Univ. Tex.* **8**: 151-183.
- BRIGHT, D. B. 1966. The land crabs of Costa Rica. *Revta Biol. trop.* **14**: 183-203.
- and C. L. HOGUE. 1972. A synopsis of the burrowing land crabs of the world and list of their arthropod symbionts and burrow associates. *Los Angeles County Mus. Contrib. Sci.* **220**: 1-58.
- BRIGHT, T. J. and L. H. PEQUEGNAT. 1974. *Biota of the West Flower Garden Bank*. Gulf Publ. Co.: Houston. 435 pp.
- BRITTON, J. C. 1976. Additional reports of *Gecarcinus lateralis* (Fremerville) (Gecarcinidae) from the Texas coast. *Southwestern Naturalist* **21**: 251-252.
- BROWN, F. A., JR. 1950. Studies on the physiology of *Uca* red chromatophores. *Biol. Bull.* **98**: 218-226.
- and G. M. JONES. 1949. Ovarian inhibition by a sinus gland principle in the fiddler crab. *Biol. Bull.* **96**: 228-232.
- and M. I. SANDEEN. 1948. Responses of the chromatophores of the fiddler crab, *Uca*, to light and temperature. *Physiol. Zool.* **21**: 361-371.
- and H. M. WEBB. 1948. Temperature relations of an endogenous daily rhythmicity in the fiddler crab, *Uca*. *Physiol. Zool.* **21**: 371-381.

- , —————, M. F. BENNETT and M. I. SANDEEN. 1955. Evidence for an exogenous contribution to persistent diurnal and lunar rhythmicity under so-called constant conditions. *Biol. Bull.* **109**: 238-254.
- BROWN, G. G. 1966. Ultrastructural studies of sperm morphology and sperm-egg interaction in the decapod *Callinectes sapidus*. *J. Ultrastruct. Res.* **14**: 425-440.
- BROWNELL, W. N., A. J. PROVENZANO, JR. and M. MARTINEZ. 1977. Culture of the West Indian spider crab, *Mithrax spinosissimus* at Los Roques, Venezuela. Paper presented at 5th World Mariculture Conf., Costa Rica. Pp. A2-1 to A2-14.
- BRUES, C. T. 1927. Occurrence of the marine crab, *Callinectes ornatus*, in brackish and fresh water. *Am. Nat.* **61**: 566-568.
- BUITENDIJK, A. M. 1950. Note on a collection of Decapoda Brachyura from the coasts of Mexico, including the description of a new genus and species. *Zool. Meded., Leiden* **30**: 269-282.
- and L. B. HOLTHUIS. 1949. Note on the Zuiderzee crab, *Rithropanopeus harrisi* (Gould) subspecies *tridentatus* (Maitland). *Zool. Meded., Leiden.* **30**: 95-106.
- BULLIS, H. R. and J. R. THOMPSON. 1965. Collections by the exploratory fishing vessels Oregon, Silver Bay, Combat, and Pelican made during 1956-1960 in the southwestern North Atlantic. *Spec. scient. Rep. U.S. Fish Wildl. Serv. Fisheries.* **510**: 1-130.
- BURKENROAD, M. D. 1947. Production of sound by the fiddler crab, *Uca pugilator* Bosc with remarks on its nocturnal and mating behavior. *Ecology.* **28**: 458-462.
- CABRERA, J. A. 1965. Contribuciones carcinológicas I. El primer estadio zoea en *Gecarcinus lateralis* (Fremenville) (Brachyura Gecarcinidae) procedente de Veracruz, Mexico. *An. Inst. Biol. Univ. Mex.* **36**: 173-187.
- CAINE, E. A. 1974. Feeding of *Ovalipes guadulpensis* (Saussure) (Decapoda: Brachyura: Portunidae), and morphological adaptations to a burrowing existence. *Biol. Bull.* **147**: 550-559.
- , 1975. Feeding of *Pinnotheres maculatus* Say (Brachyura: Pinnotheridae). *Forma et Functio.* **8**: 395-404.
- CAMERON, J. N. 1975. Aerial gas exchange in the terrestrial Brachyura *Gecarcinus lateralis* and *Cardisoma guanhumi*. *Comp. Biochem. Physiol.* **52A**: 129-134.
- CARGO, D. G. 1960. A megalops of the blue crab, *Callinectes sapidus*, in the Patuxent River, Maryland. *Chesapeake Sci.* **1**: 110.
- CARLREN, O. and J. W. HEDGPETH. 1952. Actiniaria, Zoantharia and Ceriantharia from shallow water in the northwestern Gulf of Mexico. *Publs Inst. mar. Sci. Univ. Tex.* **2**: 141-172.
- CARLSON S. 1935. The color changes in *Uca pugilator*. *Proc. Natin. Acad. Sci. U.S.A.* **21**: 549-551.
- CARLSON, S. P. 1936. Color changes in brachyuran crustaceans, especially in *Uca pugilator*. *K. Fysiogr. Sallsk, i Lund Forhandl.* **6**: 63-80.
- CARRIKER, M. R. 1951. Observations of the penetration of tightly closing bivalves by *Busycon* and other predators. *Ecology.* **32**: 73-83.
- CERAME-VIVAS, M. J. and I. E. GRAY. 1966. The distributional pattern of benthic invertebrates of the continental shelf off North Carolina. *Ecology.* **47**: 260-270.
- CERAME-VIVAS, J. M., A. B. WILLIAMS and I. E. GRAY. 1963. New crustacean records for the coast of North Carolina. *Crustaceana.* **5**: 157-159.
- CHACE, F. A., JR. 1939. Reports on the scientific results of the first *Atlantis* expedition to the West Indies, under the joint auspices of the University of Havana and Harvard University. Preliminary descriptions of one new genus and seventeen new species of decapod and stomatopod Crustacea. *Mems. Soc. cub. Hist. nat. 'Felipe Poey'* **13**: 31-54.

- . 1940. Reports on the scientific results of the *Atlantis* expeditions to the West Indies, under the joint auspices of the University of Havana and Harvard University. The brachyuran crabs. *Torreta*. **3**: 3-67.
- . 1951. The oceanic crabs of the genera *Planes* and *Pachygrapsus*. *Proc. U.S. natn. Mus.* **101**: 65-103.
- . 1956. In: S. Springer and H. R. Bullis, Collections by the *Oregon* in the Gulf of Mexico, *Spec. scient. Rep. U.S. Fish Wildl. Serv. Fisheries*. **196**: 1-134.
- . 1966. Decapod crustaceans from St. Helena Island, South Atlantic. *Proc. U.S. natn. Mus.* **118**: 623-661.
- and L. B. HOLTHUIS. 1948. Land and fresh water decapod Crustacea from the Leeward Group and northern South America. *Studies on the Fauna of Curacao, Aruba, Bonaire and the Venezuelan Islands*. **3**: 21-28.
- and H. H. HOBBS, JR. 1969. The freshwater and terrestrial decapod crustaceans of the West Indies with special reference to Dominica. *Bull. U.S. natn. Mus.* **292**: 1-258.
- CHAMBERLAIN, N. A. 1961. Studies on the larval development of *Neopanope texana sayi* (Smith) and other crabs of the family Xanthidae (Brachyura). *Tech. Rept. Chesapeake Bay Inst.* **22**: 1-35.
- . 1962. Ecological studies of the larval development of *Rhithropanopeus harrisi* (Xanthidae, Brachyura). *Tech. Rept. Chesapeake Bay Inst.* **23**: 1-47.
- CHEN, P. S. 1933. Zur Morphologie und Histologie der Respirations-organe von *Grapsus grapsus* L. *Jena Z. naturwiss., Neue Folge*. **61**: 31-88.
- CHEUNG, T. S. 1967. Endocrine control of growth and reproduction in the stone crab, *Menippe mercenaria* (Say). *Am. Zoologist*. **7**: 200.
- . 1969. The environmental and hormonal growth and reproduction in the adult female stone crab, *Menippe mercenaria* (Say). *Biol. Bull.* **136**: 327-346.
- . 1973. The simultaneous regeneration of claws in the aged, male stone crab, *Menippe mercenaria* (Say), with special reference to the terminal molt. *Bull. Inst. Zool., Academia Sinica*. **12**: 1-11.
- . 1976. A biostatistical study of the functional consistency in the reversed claws of the adult male stone crabs, *Menippe mercenaria* (Say). *Crustaceana* **31**: 137-144.
- CHRISTENSEN, A. M. and J. J. McDERMOTT. 1959. Life-history and biology of the oyster crab, *Pinnotheres maculatus* Say. *Biol. Bull.* **114**: 146-179.
- CHRISTIANSEN, M. E. 1969. Crustacea Decapoda Brachyura. *Mar. Invertebr. Scand.* **2**: 1-143.
- , J. D. COSTLOW, JR., and R. J. MONROE. 1977a. Effects of the juvenile hormone mimic ZR-515 (Altosid^c) on larval development in the mud-crab *Rhithropanopeus harrisi* in various salinities and cyclic temperatures. *Mar. Biol.* **39**: 269-279.
- , J. D. COSTLOW, JR. and R. J. MONROE. 1977b. Effects of the juvenile hormone mimic ZR-512 (Altozar^c) on larval development of the mud-crab *Rhithropanopeus harrisi* at various cyclic temperatures. *Mar. Biol.* **39**: 281-288.
- CHRISTMAS, J. Y. and W. LANGLEY. 1973. Estuarine invertebrates, Mississippi. Section 4, Pp. 255-319. In: J. Y. Christmas (ed.), *Cooperative Gulf of Mexico Estuarine Inventory and Study, Mississippi*. Gulf Coast Research Lab.
- CHURCHILL, E. P. JR. 1919. Life history of the blue crab. *Bull. Bur. Fish., Wash.* **36**: 95-128.
- COBB, S. P. 1973. *Fabia tellinae*, a new species of commensal crab (Decapoda, Pinnotheridae) from the northeastern Gulf of Mexico. *Crustaceana*. **25**: 70-74.
- COCHRAN, D. M. 1935. The skeletal musculature of the blue crab, *Callinectes sapidus* Rathbun. *Smithson. misc. Collns.* **92**: 1-76.

- COELHO, P. A. 1971a. A distribuição dos crustáceos decápodos reptantes do Norte do Brasil. *Trab. Inst. Oceanogr. Univ. Fed. Pernambuco*, **7/8**: 71-90.
- . 1971b. Novas ocorrências de crustáceos decápodos em Pernambuco e Estados vizinhos (Brasil). *Trab. Inst. Oceanogr. Univ. Fed. Pernambuco*, **9/11**: 239-248.
- . 1971c. Nota prévia sobre os Majidae do norte e nordeste do Brasil. *Archos Mus. nac., Rio de J.*, **54**: 137-146.
- and M. de A. RAMOS. 1972. A constituição e a distribuição da fauna de decápodos do litoral leste da América do Sul entre as latitudes de 5°N e 39°S. *Trab. Oceanogr. Univ. Recife*, **13**: 133-236.
- COLE, T. J. 1971. Osmotic and ionic regulatory abilities of *Uca minax* in relation to its ecology. Masters Thesis, Univ. West Florida, Pensacola.
- CONNOLLY, C. J. 1925. The larval stages and megalops of *Rhithropanopeus harrisi* (Gould). *Contr. Can. Biol. Fish.* **2**: 329-334.
- CONTRERAS, F. 1930. Contribucion al conocimiento de las jaibas de Mexico. *An. Inst. Biol. Univ. Mex.* **1**: 227-241.
- COOHILL, T. P. and M. FINGERMAN. 1975. Relative effectiveness of ultraviolet and visible light in eliciting pigment dispersion in melanophores of the fiddler crab, *Uca pugilator*, through the secondary response. *Physiol. Zool.* **48**: 57-63.
- COOK, D. W. and S. R. LOFTON. 1973. Chitinoclastic bacteria associated with shell disease in *Penaeus* shrimp and the blue crab (*Callinectes sapidus*). *J. Wildl. Dis.* **9**: 154-159.
- COOKE, I. M. and M. W. GOLDSTONE. 1970. Fluorescence localization of monoamines in crab neurosecretory structures. *J. exp. Biol.* **53**: 651-668.
- COPELAND, B. J. 1965. Fauna of the Aransas Pass Inlet, Texas. I. Emigration as shown by tide trap collections. *Publ. Inst. mar. Sci. Univ. Tex.* **10**: 9-21.
- and T. J. BECHTEL. 1974. Some environmental limits of six Gulf coast estuarine organisms. *Contr. mar. Sci.* **18**: 169-204.
- COPEPLAND, D. E. 1968. Fine structure of salt and water uptake in the land crab, *Gecarcinus lateralis*. *Am. Zoologist*, **8**: 417-432.
- and A. T. FITZJARRELL. 1968. The salt absorbing cells in the gills of the blue crab (*Callinectes sapidus* Rathbun) with notes on modified mitochondria. *Z. Zellforsch. Mikrosk. Anat.* **92**: 1-22.
- CORRINGTON, J. D. 1927. Commensal association of a spider crab and a medusa. *Biol. Bull.* **53**: 346-350.
- COSTLOW, J. D., JR. 1963. The effect of eyestalk extirpation on metamorphosis of megalops of the blue crab, *Callinectes sapidus* Rathbun. *Gen. Comp. Endocrinol.* **3**: 120-130.
- . 1965. Variability in larval stages of the blue crab, *Callinectes sapidus*. *Biol. Bull.* **128**: 58-66.
- . 1966. The effect of eyestalk extirpation on larval development of the mud crab, *Rhithropanopeus harrisi* (Gould). *Gen. Comp. Endocrinol.* **7**: 255-274.
- . 1967. The effect of salinity and temperature on survival and metamorphosis of the blue crab, *Callinectes sapidus*. *Helgolander wiss. Meeresunters.* **15**: 84-97.
- and C. G. BOOKHOUT. 1959. The larval development of *Callinectes sapidus* Rathbun reared in the laboratory. *Biol. Bull.* **116**: 373-396.
- and ———. 1960a. A method for developing brachyuran eggs *in vitro*. *Limnol. Oceanogr.* **5**: 212-215.
- and ———. 1960b. The complete larval development of *Sesarma cinereum* (Bosc) reared in the laboratory. *Biol. Bull.* **118**: 203-214.

- and —————. 1961a. The larval stages of *Panopeus herbstii* Milne-Edwards reared in the laboratory. *J. Elisha Mitchell Sci. Soc.* **77**: 33-42.
- and —————. 1961b. The larval development of *Eurypanopeus depressus* (Smith) under laboratory conditions. *Crustaceana.* **2**: 6-15.
- and —————. 1962. The larval development of *Sesarma reticulatum* Say reared in the laboratory. *Crustaceana.* **4**: 281-294.
- and —————. 1966a. Larval development of the crab, *Hexapanopeus angustifrons*. *Chesapeake Sci.* **7**: 148-156.
- and —————. 1966b. Larval stages of the crab, *Pinnotheres maculatus*, under laboratory conditions. *Chesapeake Sci.* **7**: 157-163.
- and —————. 1967. The larval stages of the crab, *Neopanope packardii* (Kingsley), in the laboratory. *Bull. mar. Sci.* **17**: 52-63.
- and —————. 1968a. Larval development of the crab, *Leptodius agassizii* A. Milne Edwards in the laboratory (Brachyura, Xanthidae). *Crustaceana, Suppl.* **2**: 204-213.
- and —————. 1968b. The complete larval development of *Cardisoma guanhumi* Latreille in the laboratory (Brachyura, Gecarcinidae). *Crustaceana Suppl.* **2**: 259-270.
- and —————. 1968c. The effect of environmental factors on development of the land-crab, *Cardisoma guanhumi*. *Am. Zoologist* **8**: 399-410.
- COSTLOW, J. D., JR., C. G. BOOKHOUT and R. MONROE. 1960. The effect of salinity and temperature on larval development of *Sesarma cinereum* (Bosc) reared in the laboratory. *Biol. Bull.* **118**: 183-202.
- , ————— and —————. 1962. Salinity-temperature effects on the larval development of the crab, *Panopeus herbstii* Milne-Edwards, reared in the laboratory. *Physiol. Zool.* **35**: 79-93.
- , ————— and —————. 1966. Studies on the larval development of the crab, *Rhithropanopeus harristii* (Gould). I. The Effect of salinity and temperature on larval development. *Physiol. Zool.* **39**: 81-100.
- COUCH, J. N. 1942. A new fungus on crab eggs. *J. Elisha Mitchell Sci. Soc.* **58**: 158-161.
- COVENTY, G. A. 1944. Results of the Fifth George Vanderbilt Expedition (1941) (Bahamas, Caribbean Sea, Panama, Galapagos Archipelago, and Mexican Pacific islands). The Crustacea. *Monogr. Acad. Nat. Sci. Philadelphia*, no. **6**: pp. 531-544.
- COWARD, S. J., H. C. GERHARDT and D. T. CROCKETT. 1970. Behavioral variation in natural populations of two species of fiddler crabs (*Uca*) and some preliminary observations on directed modification. *J. Biol. Psychol.* **12**: 24-31.
- COWLES, R. P. 1908. Habits, reactions and associations of *Ocypoda arenaria*. *Pap. Tortugas Lab.* **2**: 1-41.
- CRAIG, W. J. 1974. Physiological ecology of the commensal crabs, *Polyonyx gibbesi* Haig and *Pinnixa chaetoptera* Stimpson. *Oecologia.* **15**: 235-244.
- CRANE, J. 1943a. Display, breeding and relationships of fiddler crabs (Brachyura, genus *Uca*) in the northeastern United States. *Zoologica.* **28**: 217-223.
- . 1943b. Crabs of the genus *Uca* from Venezuela. *Zoologica.* **28**: 33-44.
- . 1957. Basic patterns of display in fiddler crabs (Ocypodidae, Genus *Uca*). *Zoologica.* **42**: 69-82.
- . 1967. Combat and its ritualization in fiddler crabs (Ocypodidae) with special reference to *Uca rapax* (Smith). *Zoologica.* **52**: 49-76.
- . 1975. *Fiddler Crabs of the World*. Princeton University Press. 737 pp.
- CRICHTON, O. W. 1960. Marsh crab: intertidal tunnel maker and grass-eater. *Estuarine Bull.* **5**: 3-10.

- CRONIN, L. E. 1947. Anatomy and histology of the male reproductive system of *Callinectes sapidus* Rathbun. *J. Morph.* **81**: 209-239.
- , W. A. van ENGEL, D. G. CARGO and F. J. WOJCIK. 1957. A partial bibliography of the genus *Callinectes*. *Spec. Scient. Rept. Va. Fish. Lab.* no. **8**: 21 pp.
- CROSNIER, A. 1967. Remarques sur quelques Crustacés Décapodes benthiques ouest-africains. Description de *Heteropanope ocanthocarpus* et *Medaeus rectifrons* spp. nov. *Bull. Mus. Natn. Hist. Nat., Paris.* **39**: 320-344.
- DARBY, H. H. 1935. Intersexuality in the Crustacea. *Pap. Tortugas Lab.* **29**: 145-149.
- DARNELL, R. M. 1959. Studies of the life history of the blue crab in Louisiana waters. *Trans. Am. Fish. Soc.* **88**: 294-324.
- DAUGHERTY, F. M., JR. 1952a. The blue crab investigation, 1949-1950. *Tex. J. Sci.* **4**: 77-84.
- , 1952b. Notes on *Callinectes danae* Smith in Aransas Bay, Texas, and adjacent waters. *Tex. J. Sci.* **4**: 264-267.
- DAVIS, C. C. 1965. A study of the hatching process in aquatic invertebrates: XX. The blue crab, *Callinectes sapidus*, Rathbun, XXI. The nemertean. *Carcinonemertes carcinophila* (Kolliker). *Chesapeake Sci.* **6**: 201-208.
- DAWSON, C. E. 1966. Additions to the known marine fauna of Grand Isle, Louisiana. *Proc. La Acad. Sci.* **29**: 175-180.
- DECOURSEY, P. J. and W. B. VERNBERG. 1972. Effect of mercury on survival, metabolism and behavior of larval *Uca pugilator* (Brachyura). *Oikos.* **23**: 241-247.
- van DELFT, A. M. L. 1968. The daily colour rhythm of the fiddler crab *Uca repax* on Curaçao. *Stud. Fauna Curaçao.* **25**: 58-72.
- DEMBOWSKI, J. 1925. On the "speech" of the fiddler crab, *Uca pugilator*. *Pr. Inst. M. Nenci.* **3**: 1-7.
- , 1926. Notes on the behaviour of the fiddler crab. *Biol. Bull.* **50**: 179-200.
- DEMEUSY, N. 1957. Respiratory metabolism of the fiddler crab *Uca pugilator* from two different latitudinal populations. *Biol. Bull.* **113**: 245-253.
- DIAZ, H. and J. D. COSTLOW. 1972. Larval development of *Ocypode quadrata* (Brachyura: Crustacea) under laboratory conditions. *Mar. Biol.* **15**: 120-131.
- , and J. J. EWALD. 1968. A comparison of the larval development of *Metasesarma rubripes* (Rathbun) and *Sesarma ricodi* H. Milne Edwards (Brachyura, Grapsidae) reared under similar laboratory conditions. *Crustaceana, Suppl.* **2**: 225-248.
- DRAGOVICH, A. and J. A. KELLY, JR. 1964. Ecological observations of macroinvertebrates in Tampa Bay, Florida 1961-1962. *Bull. mar. Sci. Gulf Caribb.* **14**: 74-102.
- DUNCKER, G. 1934. Gefangenschaftsbeobachtungen an *Sesarma cinerea* Milne Edw. *Zool. Jahrbuch.. Syst. Geogr. Biol. Tiere, Jena.* **66**: 285-290.
- EDMONDSON, C. H. 1962. Xanthidae of Hawaii. *Occas. Pap. Bernice P. Bishop Mus.* **22**: 215-309.
- EDWARDS, G. A. 1950. The influence of eyestalk removal on the metabolism of the fiddler crab. *Physiol. Comp. Oecol.* **2**: 34-50.
- EIBL-EIBESFELDT, I. 1963. *Grapsus grapsus* (Brachyura), Drohen. E. 599, 2½ mins. (Film)..
- EIDEMILLER, A. 1969. Entry behavior of the crab *Pinnotheres maculatus* Say. *Q. Jl. Fla. Acad. Sci.* **32**: 266-274.
- EISEN, A. Z., K. O. HENDERSON, J. J. JEFFREY and R. A. BRADSHAW. 1973. A collagenolytic protease from the hepatopancreas of the fiddler crab *Uca pugilator*: Purification and properties. *Biochemistry.* **12**: 1814-1822.

- ENGEL, D. W. 1973. The radiation sensitivities of three species of fiddler crabs (*Uca pugilator*, *U. pugnax* and *U. minax*). *Chesapeake Sci.* **14**: 289-291.
- and L. D. EGGERT. 1974. The effect of salinity and sex on the respiration rates of excised gills of the blue crab, *Callinectes sapidus*. *Comp. Biochem. Physiol.* **47A**: 1005-1011.
- van ENGEL, W. A. 1958. The blue crab and its fishery in Chesapeake Bay. I. Reproduction, early development, growth, and migration. *Comm. Fish. Rev.* **20**: 6-17.
- EPIFANIO, C. E. 1972. Effects of dieldrin-contaminated food on the development of *Leptodius floridanus* larvae. *Mar. Biol.* **13**: 292-297.
- EVANS, D. H., K. COOPER and M. B. BOGAN. 1976. Sodium extrusion by the sea-water-acclimated fiddler crab *Uca pugilator*: comparison with other marine Crustacea and marine teleost fish. *J. exp. Biol.* **64**: 203-219.
- EVOY, W. H. and C. R. FOURTNER. 1973. Nervous control of walking in the crab, *Cardisoma guanhumi*. III. Proprioceptive influences on intra- and intersegmental coordination. *J. comp. Physiol.* **83**: 303-318.
- FALES, R. R. 1976. Apparent predation on the mole crab *Emerita talpoida* (Say) by the ghost crab *Ocypode quadrata* (Fabricius). *Chesapeake Sci.* **17**: 65.
- FAUSTO FILHO, J. 1967. Sobre os Calapideos do norte e nordeste do Brasil. *Arq. Estac. Biol. Mar., Univ. Fed, Ceará.* **7**: 31-62.
- . 1974. Stomatopod and decapod crustaceans of the archipelago of Fernando de Noronha, northeast Brazil. *Arq. Ciên. Mar.* **14**: 1-35.
- FELDER, D. L. 1973a. An Annotated Key to Crabs and Lobsters (Decapoda, Reptantia) from Coastal Waters of the Northwestern Gulf of Mexico. Pub. No. **LSU-SG-73-02** of the Center for Wetland Resources, Louisiana State University, Baton Rouge. 103 pp.
- . 1973b. A record of *Pinnixa lunzi* Glassell (Decapoda, Pinnotheridae) from off the coast of Texas, U.S.A. *Crustaceana.* **24**: 148-149.
- FELICE, E. P. 1958. Invertebrates from the estuarine portion of San Francisco Bay and some factors influencing their distributions. *Wasmann J. Biol.* **16**: 159-211.
- FELICIANO, C. 1962. Notes on the importance of the land crab *Cardisoma guanhumi*, Latreille of Puerto Rico, *Spec. Contrib. Inst. Mar. Biol. Univ. Puerto Rico.* 29 pp|
- FIELDER, R. H. 1930. Solving the question of crab migration. *Fishing Gazette.* **47**: 18-21.
- FIELDER, D. R., K. RANGA RAO, and M. FINGERMAN. 1971. A female-limited lipoprotein and the diversity of hemocyanin components in the dimorphic variants of the fiddler crab, *Uca pugilator*, as revealed by disc electrophoresis. *Comp. Biochem. Physiol.* **398**: 291-297.
- FINGERMAN, M. 1956. Phase difference in the tidal rhythms of color change in two species of fiddler crabs. *Biol. Bull.* **110**: 274-290.
- . 1957. Relation between position of burrows and tidal rhythm of *Uca*. *Biol. Bull.* **112**: 7-20.
- . 1970. Circadian rhythm of distal retinal pigment migration in the fiddler crab, *Uca pugilator*, maintained in constant darkness and its endocrine control. *J. Interdiscipl. Cycle Res.* **1**: 115-121.
- . 1973. Behavior of chromatophores of the fiddler crab *Uca pugilator* and the dwarf crayfish *Cambarellus shufeldtii* in response to synthetic *Pandalus* red pigment-concentrating hormone. *Gen. Comp. Endocrin.* **20**: 589-592.
- and E. F. COUCH. 1967. Differentiation of chromatophorotropins from the prawn, *Palaeomonetes vulgaris*, and the fiddler crab, *Uca pugilator*. *J. exp. Zool.* **165**: 183-194.
- and C. FITZPATRICK. 1956. An endocrine basis for the sexual difference in melanin dispersion in *Uca pugilator*. *Biol. Bull.* **110**: 138-143.

- , M. E. LOWE and W. C. MOBBERLY, JR. 1958. Environmental factors involved in setting the phases of activity of tidal rhythm of color change in the fiddler crabs *Uca pugilator* and *Uca minax*. *Limnol. Oceanogr.* **3**: 271-282.
- , R. NAGBHUSHANAM and L. PHILPOTT. 1961. Physiology of the melanophores of the crab *Sesarma reticulatum*. *Biol. Bull.* **120**: 337-347.
- , K. RANGA RAO and G. RING. 1969. Restoration of a rhythm of melanophore pigment dispersion in eyestalkless fiddler crabs, *Uca pugilator* (Bosc), at a low temperature. *Crustaceana.* **17**: 97-105.
- and Y. YAMAMOTO. 1967. Daily rhythm of melanophoric pigment migration in eyestalkless fiddler crabs, *Uca pugilator* (Bosc). *Crustaceana.* **12**: 303-319.
- FINGERMAN, S. W. and M. FINGERMAN. 1976. Effects of time of year and limb removal on rates of ecdysis of eyed and eyestalkless fiddler crabs, *Uca pugilator*. *Mar. Biol.* **37**: 357-362.
- FINNEGAN, S. 1931. Report on the Brachyura collected in Central America, the Gorgona and Galapagos Islands, by Dr. Crossland on the 'St. George' Expedition to the Pacific, 1924-25. *J. Linn. Soc. Zool.* **37**: 607-673.
- FLEMISTER, L. J. 1958. Salt and water anatomy, constancy and regulation in related crabs from marine and terrestrial habitats. *Biol. Bull.* **115**: 180-200.
- and S. C. FLEMISTER. 1951. Chloride ion regulation and oxygen consumption in the crab, *Ocypode albicans* (Bosc). *Biol. Bull.* **101**: 259-273.
- FLEMISTER, S. C. 1959. Histophysiology of gill and kidney of the crab *Ocypode albicans*. *Biol. Bull.* **116**: 37-48.
- FLOWER, F. B. and J. J. McDERMOTT. 1953. Observations on the occurrence of the oyster crab, *Pinnotheres ostreum*, as related to the oyster damage in Delaware Bay. *Nat. Shellfish Assoc., Conv. Addresses.* **1952**: 44-46.
- FOREST, J. and D. GUINOT. 1966. Campagne de la "Calypso" dans le Golfe de Guinée et aux îles Principe, São Tomé et Annobon (1956). 16. Crustaces. Decapodes. Brachycures. In: Res. Scient. Camp. "Calypso," fasc. 7. *Annls Inst. oceanogr, Monaco.* **44**: 23-124.
- FORWARD, R. B. 1977. Occurrence of a shadow response among brachyuran larvae. *Mar. Biol.* **39**: 331-341.
- FOTHERINGHAM, N. and S. BRUNENMEISTER. 1975. *Common Marine Invertebrates of the Northwestern Gulf Coast*. Gulf Publish. Co.: Houston. 197 pp.
- FOURTNER, C. R. and W. H. EVOY. 1973. Nervous control of walking in the crab, *Cardisoma quahumi*. IV. Effects of myochordotonal organ ablation. *J. comp. Physiol.* **33**: 319-329.
- FRANKS, J. S., J. Y. CHRISTMAS, W. L. SILER, R. COMBS, R. WALLER and C. BURNS. 1972. A study of nektonic and benthic faunas of the shallow Gulf of Mexico off the state of Mississippi as related to some physical, chemical and geological factors. *Gulf Res. Rep.* **4**: 1-148.
- FURTADO-OGAWA, E. 1972. Notas bioecológicas sobre a familia Xanthidae no estado do Ceará (Crustacea: Brachyura). *Arg. Ciên. Mar.* **12**: 99-104.
- FUTCH, C. R. 1965. The blue crab in Florida. *Flor. Bd. Conserv. Mar. Lab., Salt Wat. Fish. Leaflet. Ser.* **1**: 1-17.
- . 1966. The stone crab in Florida. *Flor. Bd. Conserv. Mar. Lab., Salt Wat. Fish. Leaflet. Ser.* **2**: 1-6.
- GANZ, A. R. and J. F. HERRMANN. 1975. Investigations into the southern New England red crab fishery. *R. I. Dept. Nat. Resour. Div. Fish Wildl. Mar. Fish. Sect.* 78 pp.
- GARTH, J. S. 1946. Littoral brachyuran fauna of the Galapagos Archipelago. *Allan Hancock Pac. Exped.* **5**: 341-601.

- . 1948. The Brachyura of the "Askoy" Expedition with remarks on carcinological collecting in the Panama Bight. *Bull. Am. Mus. nat. Hist.* **92**: 1-66.
- . 1958. Brachyura of the Pacific Coast of America, Oxyrhyncha. *Allan Hancock Pac. Exped.* **21**: 1-499.
- . 1961. Eastern Pacific Expeditions of the New York Zoological Society. XLV. Non-intertidal brachygnathous crabs from the west coast of tropical America. Part 2. Brachygnatha Brachyrhyncha. *Zoologica.* **46**: 133-159.
- . 1965a. The brachyuran decapod crustaceans of Clipperton Island. *Proc. Calif. Acad. Sci., Ser. 4,* **33**: 1-46.
- . 1965b. On the oceanic transport of crab larval stages. *Proc. Symp. Crustacea, Part I,* pp. 443-448.
- . 1966. Eastern Pacific expeditions of the New York Zoological Society. XLVI. Oxy stomatous and allied crabs from the west coast of tropical America. *Zoologica.* **51**: 1-16.
- and W. STEPHENSON. 1966. Brachyura of the Pacific coast of America. Brachyrhyncha: Portunidae. *Allan Hancock Monogr. Mar. Biol.* **1**: 1-154.
- GERARD, J. F. and R. GILLES. 1972. The free amino acid pool in *Callinectes sapidus* (Rathbun) tissues and its role in the intracellular osmotic regulation. *J. exp. mar. Biol. Ecol.* **10**: 125-136.
- GERLACH, S. A. 1958a. Die Mangroveregion tropischer Küsten als Lebensraum. *Z. Morph. Ökol. Tiere.* **46**:436-530.
- . 1958b. Beobachtungen über das Verhalten von Winkerkrabben (*Uca leptodactyla*). *Z. Tierpsychol.* **15**: 50-53.
- GIBBS, P. E. 1974. Notes on *Uca burgersi* Holthuis (Decapoda, Ocypodidae) from Barbuda, Leeward Islands. *Crustaceana.* **27**: 84-91.
- and G. W. BRYAN. 1972. A study of strontium, magnesium, and calcium in the environment and exoskeleton of decapod crustaceans, with special reference to *Uca burgersi* on Barbuda, West Indies. *J. exp. mar. Biol. Ecol.* **9**: 97-110.
- GIFFORD, C. A. 1962a. Some aspects of osmotic and ionic regulation in the blue crab, *Callinectes sapidus*, and the ghost crab, *Ocypode albicans*. *Publs Inst. mar. Sci. Univ. Tex.* **8**: 97-125.
- . 1962b. Some observations on the general biology of the land crab, *Cardisoma guanhumi* (Latreille), in South Florida. *Biol. Bull.* **123**: 207-223.
- . 1968. Accumulation of uric acid in the land crab, *Cardisoma guanhumi*. *Am. Zoologist* **8**: 521-528.
- and R. F. JOHNSON. 1962. Distribution of calcium in the land crab *Cardisoma guanhumi* during shell wound recalcification. *Comp. Biochem. Physiol.* **7**: 227-231.
- GILLES, R. 1970. Osmoregulation in the stenohaline crab *Libinia emarginata* Leech. *Arch. Int. Physiol. Biochem.* **78**: 91-99.
- GLAESSNER, M. F. 1969. Decapoda. In: R. C. Moore (ed.), *Treatise on Invertebrate Paleontology*. Part R. Arthropoda 4, Volume 2. pp. R399-532. Geol. Soc. Amer. and Univ. Kansas Press.
- GLASSELL, S. A. 1937. *Pinnixa lunzi*, a new commensal crab from South Carolina. *Charleston Mus. Leaflet* **9**: 3-8.
- GLEESON, R. A. and P. L. ZUBKOFF. 1977. The determination of hemolymph volume in the blue crab, *Callinectes sapidus*, utilizing ¹⁴C-thiocyanate. *Comp. Biochem. Physiol.* **56A**: 411-413.
- GODCHARLES, M. F. and W. C. JAAP. 1973. Fauna and flora in hydraulic clam dredge collections from Florida west and southeast coasts. *Spec. scient. Rep. Fla. Dept. Nat. Resources.* no. **40**, 89 pp.

- GOMES CORRÊA, M. M. 1970. Crustáceos braquiuros brasileiros da familia Raninidae. *Bolm. Mus. nac. Rio de J., Zool.* **276**: 1-21.
- GORDON, I. 1950. Crustacea: Dromiacea. I. Systematic account of the Dromiacea collected by the "John Murray" Expedition. II. The morphology of the spermatheca in certain Dromiacea. *Sci. Rept. John Murray Exped. 1933-1934.* **9**: 201-253.
- . 1963. On the relationships of Dromiacea, Tymolinae and Raninidae to the Brachyura. pp. 51-57. In: H. B. Whitting and W. D. I. Rolfe (eds.), "Phylogeny and Evolution of Crustacea." *Spec. Publ., Mus. Comp. Zool. Harvard Univ.*
- GORE, R. H. 1977. Studies on decapod Crustacea from the Indian River region of Florida. VI. The identity of *Parthenope (Platylambrus) seratta* (H. Milne Edwards, 1834) and *Parthenope (Platylambrus) granulata* (Kingsley, 1879). *Proc. biol. Soc. Washington* (in press).
- and R. E. GRIZZLE. 1974. Studies on decapod Crustacea from the Indian River region of Florida. III. *Callinectes bocourti* A. Milne Edwards, 1879 (Decapoda, Portunidae) from the central east coast of Florida. *Crustaceana.* **27**: 306-308.
- GRAY, E. H. 1942. Ecological and life history aspects of the red-jointed fiddler crab, *Uca minax* (Le Conte), region of Solomon Island, Maryland. *Contrib. Chesapeake Biol. Lab., Publ. no. 51*: 3-20.
- and C. L. NEWCOMBE. 1938. The relative growth of parts in the blue crab *Callinectes sapidus* Rathbun. *Growth.* **2**: 235-246.
- GRAY, G. W., JR. 1969. Investigation of the basic life history of the red crab (*Geryon quinquedens*). R. I. Div. Conserv. P.L. 88-309, Proj. 3-46-R Completion Rept., 36 pp.
- GRAY, I. E. 1957. A comparative study of the gill area of crabs. *Biol. Bull.* **112**: 34-42.
- . 1961. Changes in the abundance of the commensal crabs of *Chaetopterus*. *Biol. Bull.* **120**: 353-359.
- , L. R. McCLOSKEY and S. C. WEIHE. 1968. The commensal crab *Dissodactylus mellitae* and its reaction to sand dollar host-factor. *J. Elisha Mitchell Sci. Soc.* **84**: 472-481.
- GREEN, J. W., M. HARSCH, L. BARR and C. L. PROSSER. 1959. The regulation of water and salt by the fiddler crabs, *Uca pugnax* and *Uca pugnator*. *Biol. Bull.* **116**: 76-87.
- GUINOT, D. 1964. Les trois especes du genre *Domecia* (Decapoda, Brachyura): *D. hispida* Eydoux & Souleyet, *D. glabra* Alcock, et *D. acanthophora* (Desbonne & Schramm). *Crustaceana.* **7**: 267-283.
- . 1966. Recherches préliminaires sur les groupements naturels chez les Crustacés Décapodes Brachyours. I. Les affinités des genres *Aethra*, *Osachila*, *Hepatus*, *Hepatella* et *Actaemorpha*. *Bull. Mus. natn. Hist. Nat. Paris.* **38**: 744-762, 828-845.
- . 1967a. Recherches préliminaires sur les groupements naturels chez les Crustacés Décapodes Brachyours. II. Les anciens genres *Micropanope* Stimpson et *Medaeus* Dana. *Bull. Mus. natn. Hist. Nat., Paris.* **39**: 345-374.
- . 1967b. Recherches préliminaires sur les groupements naturels chez les Crustacés Décapodes Brachyours. III. A propos des affinités des genres *Dairoides* Stebbing et *Daira* de Haan. *Bull. Mus. natn. Hist. Nat., Paris.* **39**: 540-563.
- . 1968a. Recherches préliminaires sur les groupements naturels chez les Crustacés Décapodes Brachyours. IV. Observations sur quelques genres de Xanthidae. *Bull. Mus. natn. Hist. Nat., Paris.* **39**: 695-727.
- . 1968b. Recherches préliminaires sur les groupements naturels chez les Crustacés Décapodes Brachyours. V. Établissement d'un caractere evolutif: l'articulation ischiomerale des chelipedes. *Bull. Mus. natn. Hist. Nat., Paris.* **40**: 149-166.
- . 1968c. Recherches préliminaires sur les groupements naturels chez les Crustacés Décapodes Brachyours. VI. Les Carpilinae. *Bull. Mus. natn. Hist. Nat., Paris.* **40**: 320-334.

- . 1969a. Recherches préliminaires sur les groupements naturels chez les Crustacés Décapodes Brachyours. VII. Les Goneplacidae. *Bull. Mus. natn. Hist. Nat., Paris*. **41**: 241-265; 507-528; 688-714.
- . 1969b. Sur divers Xanthidae, notamment sur *Actaea* de Haan et *Paractaea* gen. nov. (Crustacea Decapoda Brachyura). *Cah. Pac.* **13**: 223-285.
- . 1971. Recherches préliminaires sur les groupements naturels chez les Crustacés Décapodes Brachyours. VIII. Synthèse et bibliographie. *Bull. Mus. natn. Hist. Nat., Paris*. **42**: 1063-1090.
- GUINOT-DUMORTIER, D. 1959. Sur une collection de Crustacés (Decapoda Reptantia) de Guyane Française. I. Brachyura (Oxyrhyncha exclus). *Bull. Mus. natn. Hist. Nat., Paris*. **31**: 423-434; 510-515.
- . 1960. Sur une collection de Crustacés (Decapoda Reptantia) de Guyane Française. II. Brachyura Oxyrhyncha et Macrura. *Bull. Mus. natn. Hist. Nat. Paris*. **32**: 177-187.
- and B. DUMORTIER. 1960. La stridulation chez les crabes. *Crustaceana*. **1**: 117-155.
- and ———. 1961. Description d'un appareil striduloire dans le genre *Cycloes* de Haan (Crustacea, Brachyura, Oxystemata, Calappidae). *Bull. Mus. natn. Hist. Nat. Paris, 2e Ser.* **32**: 558-561.
- GUNTER, G. 1938. The common blue crab in fresh waters. *Science*. **8**: 87-88.
- . 1950. Seasonal population changes and distributions as related to salinity, of certain invertebrates of the Texas coast, including the commercial shrimp. *Publs. Inst. mar. Sci. Univ. Tex.* **1**: 7-87.
- . 1954. Sagacity of a crab. *Science* **120**: 188-189.
- GUTSELL, J. S. 1928. The spider crab, *Libinia dubia*, and the jelly-fish, *Stomolophus meleagris*, found associated at Beaufort, North Carolina. *Ecology*. **9**: 358-359.
- GUYSELMAN, J. B. 1953. An analysis of the molting process in the fiddler crab, *Uca pugilator*. *Biol. Bull.* **104**: 115-137.
- HAEFNER, P. A., JR. 1977. Reproductive biology of the female deep-sea red crab, *Geryon quinquedens*, from the Chesapeake Bight. *Fishery Bull. Fish Wildl. Serv. U.S.* **75**: 91-102.
- and D. GARTEN. 1974. Methods of handling and shedding blue crabs, *Callinectes sapidus*. *Mar. Resources Advis. Ser 8* (Virginia Inst. Mar. Sci.), pp. 1-14.
- and J. A. MUSICK. 1974. Observations on distribution and abundance of red crabs in Norfolk Canyon and adjacent continental slope. *Mar. Fish. Rev.* **36**: 31-34.
- and C. N. SHUSTER, JR. 1964. Length increments during terminal molt of the female blue crab, *Callinectes sapidus*, in different salinity environments. *Chesapeake Sci.* **5**: 114-118.
- von HAGEN, H.-O. 1967. Nachweis einer kinasthetischen orientierung bei *Uca rapax*. *Z. Morph. Ökol, Tiere* **58**: 301-320.
- . 1970a. Verwandtschaftliche Gruppierung und Verbreitung der Karibischen Winterkrabben (Ocypodidae, Gattung *Uca*). *Zool. Meded. Leiden* **44**: 217-235.
- . 1970b. Anpassungen an das spezielle Gezeitenzonen-Niveau bei Ocypodiden (Decapoda, Brachyura). *Forma Functio*. **2**: 361-413.
- . 1970c. Die Balz von *Uca vocator* (Herbst) als ökologisches Problem. *Forma Functio*. **2**: 238-253.
- . 1976. Review: Jocelyn Crane, Fiddler Crabs of the World. Ocypodidae: Genus *Uca*. xxiv + 737 pp., 369 photographs, 101 figures, 21 maps. ISBN 08102-6, \$75.00. Princeton University Press, Princeton, New Jersey, 1975. *Crustaceana*. **31**: 221-224.

- HALEY, S. R. 1967. Reproductive biology of the Texas ghost crab *Ocypode albicans* Bosc (Decapoda: Ocypodidae). Ph.D. Dissertation, Univ. of Texas at Austin. 135 pp.
- . 1969. Relative growth and sexual maturity of the Texas ghost crab, *Ocypode quadrata* (Fabr.) (Brachyura, Ocypodidae). *Crustaceana*. **17**: 285–297.
- . 1972. Reproductive cycling in the ghost crab, *Ocypode quadrata* (Fabr.) (Brachyura, Ocypodidae). *Crustaceana*. **23**: 1–11.
- HAMILTON, P. V. 1976. Predation on *Littorina irrorata* (Mollusca: Gastropoda) by *Callinectes sapidus* (Crustacea: Portunidae). *Bull. mar. Sci.* **26**: 403–409.
- , R. T. NISHIMOTO and J. G. HALUSKY. 1976. Cheliped laterality in *Callinectes sapidus* (Crustacea: Portunidae). *Biol. Bull.* **150**: 393–401.
- HARTNOLL, R. G. 1964a. The freshwater grapsid crabs of Jamaica. *Proc. Linn. Soc. Lond.* **175**: 145–169.
- . 1964b. The zoeal stages of the spider crab *Microphrys bicornutus* (Latr.). *Ann. Mag. nat. Hist.* **7**: 241–246.
- . 1965a. The biology of spider crabs: a comparison of British and Jamaican species. *Crustaceana*. **9**: 1–16.
- . 1965b. Notes on the marine grapsid crabs of Jamaica. *Proc. Linn. Soc. Lond.* **176**: 113–147.
- . 1969. Mating in the Brachyura. *Crustaceana*. **16**: 161–181.
- . 1970. Swimming in the dromiid crab, *Homola barbata*. *Anim. Behav.* **18**: 588–591.
- . 1971. The occurrence, methods and significance of swimming in the Brachyura. *Anim. Behav.* **19**: 34–50.
- HAVEN, D. 1958. Effects of pea crabs, *Pinnotheres ostreum*, on oysters, *Crassostrea virginica*. *Proc. natl. Shellfish Assoc.* **49**: 77–86.
- and J. D. ANDREWS. 1957. Survival and growth of *Venus mercenaria*, *Venus campechiensis* and their hybrids in suspended trays and on natural bottoms. *Proc. natl. Shellfish Assoc.* **47**: 43–49.
- HAY, W. P. 1905. The life history of the blue crab (*Callinectes sapidus*). *Rept. U.S. Bur. Fish.* **1904**: 395–413.
- and C. A. SHORE. 1918. The decapod crustaceans of Beaufort, N.C., and surrounding region. *Bull. U.S. Bur. Fish.* **35**: 369–475.
- HAZLETT, B. A. 1971. Antennule chemosensitivity in marine decapod Crustacea. *J. Anim. Morph. Physiol.* **13**: 1–10.
- . 1972a. Responses to agonistic postures by the spider crab. *Microphrys bicornutus*. *Mar. Behav. Physiol.* **1**: 85–92.
- . 1972b. Stereotypy of agonistic movements in the spider crab *Microphrys bicornutus*. *Behaviour*. **42**: 270–278.
- . 1976. Agnostic behavior of two sympatric species of xanthid crabs, *Leptodius floridanus* and *Hexapanopeus angustifrons*. *Mar. Behav. Physiol.* **4**: 107–119.
- and G. F. ESTABROOK. 1974. Examination of agonistic behavior by character analysis. I. The spider crab *Microphrys bicornutus*. *Behaviour*. **44**: 131–144.
- HAZLETT, B. and D. RITTSCHOFF. 1975. Daily movements and home range in *Mithrax spinosissimus* (Majidae, Decapoda). *Mar. Behav. Physiol.* **3**: 101–118.
- HEDGPETH, J. W. 1950. Notes on the marine invertebrate fauna of salt flat areas in Aransas National Wildlife Refuge, Texas. *Publs. Inst. mar. Sci. Univ. Tex.* **1**: 103–119.
- . 1953. An introduction to the zoogeography of the northwestern Gulf of Mexico with reference to the invertebrate fauna. *Publs. Inst. mar. Sci. Univ. Tex.* **3**: 107–224.

- HENNING, H. G. 1975a. Ökologische, ethologische und sinnesphysiologische Untersuchungen an der Landkrabbe *Cardisoma guanhumi* Latreille (Decapoda, Brachyura) in Nordkolumbien. *Forma Functio*. **8**: 253-304.
- . 1975b. Kampf- Fortpflanzungs- und Hautungsverhalten; Wachstum und Geschlechtsreife von *Cardisoma guanhumi* Latreille. *Forma Functio*. **8**: 463-510.
- HERREID, C. F., II. 1963. Observations on the feeding behavior of *Cardisoma guanhumi* (Latreille) in southern Florida. *Crustaceana*. **5**: 176-180.
- HERREID, C. F. 1967. Skeletal measurements and growth of the land crab, *Cardisoma guanhumi* Latreille. *Crustaceana*. **13**: 39-44.
- and C. A. GIFFORD. 1963. The burrow habitat of the land crab, *Cardisoma guanhumi* (Latreille). *Ecology*. **44**: 773-775.
- HERRNKIND, W. F., 1968a. Adaptive visually-directed orientation in *Uca pugilator*. *Am. Zoologist*. **8**: 585-598.
- . 1968b. The breeding of *Uca pugilator* (Bosc) and mass rearing of the larvae with comments on the behavior of the larval and early crab stages (Brachyura, Ocypodidae). *Crustaceana, Suppl.* **2**: 214-224.
- . 1968c. Ecological and ontogenetic aspects of visual orientation in the sand fiddler crab, *Uca pugilator* (Bosc). Ph.D. Dissertation, Univ. of Miami, Miami, Florida.
- . 1972. Orientation of shore-living arthropods, especially the sand fiddler crab. pp. 1-59. In: H. E. Winn and B. L. Olla (eds.), *Behavior of Marine Invertebrates. Vol. 1, Invertebrates*. Plenum Press.
- HERRNKIND, W., G. STANTON and E. CONKLIN. 1976. Initial characterization of the commensal complex associated with the anemone, *Lebrunia danae*, at Grand Bahama. *Bull. mar. Sci.* **26**: 65-71.
- HILDEBRAND, H. H. 1954. A study of the fauna of the brown shrimp (*Penaeus aztecus* Ives) grounds in the western Gulf of Mexico. *Publs Inst. mar. Sci. Univ. Tex.* **3**: 229-366.
- . 1955. A study of the pink shrimp (*Penaeus duorarum* Burkenroad) grounds in the Gulf of Campeche. *Publs Inst. mar. Sci. Univ. Tex.* **4**: 171-232.
- . 1957. Estudios biológicos preliminares sobre la Laguna Madre de Tamaulipas. *Ciencia (Mexico)*. **17**: 151-173.
- HILL, G. W. and R. E. HUNTER. 1973. Burrows of the ghost crab *Ocypode quadrata* (Fabricius) on the barrier islands, south-central Texas coast. *J. Sediment. Petrol.* **43**: 24-30.
- HINSCH, G. W. 1968. Reproductive behavior in the spider crab, *Libinia emarginata* L. *Biol. Bull.* **135**: 273-278.
- . 1970. Some factors controlling reproduction in the spider crab, *Libinia emarginata*. *Biol. Bull.* **139**: 410.
- . 1973. Sperm structure of Oxyrhynga. *Can. J. Zool.* **51**: 421-426.
- and M. V. CONE. 1969. Ultrastructural observation of vitellogenesis in the spider crab, *Libinia emarginata* L. *J. Cell. Biol.* **40**: 336-342.
- HOCKETT, J. C. and H. KRITZLER. 1972. Capture-recapture methods with *Uca*. *Biol. Bull.* **142**: 49-56.
- HODGE, M. H. 1956a. Autotomy and regeneration in *Gecarcinus lateralis*. *Anat. Rec.* **125**: 633.
- . 1956b. Variations on the normal pattern of limb regeneration in *Gecarcinus lateralis*. *Anat. Rec.* **125**: 635-636.
- and G. B. CHAPMAN. 1958. Some observations on the fine structure of the sinus gland of a land crab, *Gecarcinus lateralis*. *J. biophys. biochem. Cytol.* **4**: 571-574.

- HOESE, H. D. 1960. Biotic changes associated with end of a drought. *Limnol. Oceanogr.* **5**: 326-336.
- , B. J. COPELAND, F. N. MOSELEY and E. D. LANE. 1968. Fauna of the Aransas Pass Inlet, Texas. III. Diel and seasonal variations in trawlable organisms of the adjacent area. *Tex. J. Sci.* **20**: 33-60.
- and R. S. JONES. 1963. Seasonality of larger animals in a Texas turtle grass community. *Publs Inst. mar. Sci. Univ. Tex.* **9**: 37-47.
- , and J. M. VALENTINE, JR., 1972. U.S.L. studies on the Chandeleur Islands. *Research Ser. no. 10. Southwest La.:* Lafayette, La. 60 pp.
- HOLLAND, C. A. and D. M. SKINNER. 1976. Interactions between molting and regeneration in the land crab. *Biol. Bull.* **150**: 222-240.
- HOLLAND, J. S., D. V. ALDRICH and K. STRAWN. 1971. Effects of temperature and salinity on growth, food conversion, survival and temperature resistance of juvenile blue crabs, *Callinectes sapidus* Rathbun. *Texas A&M University. Sea Grant Publ. TAMU-SG-71-222.* 166 pp.
- HOLMSEN, A. A. and H. McALLISTER. 1974. Technological and economic aspects of red crab harvesting and processing. *Tech. Rept. Univ. R.I. Mar.* **28**: 1-35.
- HOLTHUIS, L. B. 1958. West Indian crabs of the genus *Calappa*, with a description of three new species. *Stud. Fauna Curaçao.* **8**: 146-186.
- . 1959. The Crustacea Decapoda of Suriname (Dutch Guiana). *Zool. Verh., Leiden.* **44**: 1-296.
- . 1961. Report on a collection of Crustacea Decapoda and Stomatopoda from Turkey and the Balkans. *Zool. Verh., Leiden.* **47**: 1-67.
- . 1967. On a new species of *Uca* from the West Indian region (Crustacea, Brachyura, Ocypodidae). *Zool. Meded., Leiden.* **42**: 51-54.
- . 1969. *Portunus binoculus*, n. sp., a new deep-water swimming crab from the Caribbean region (Crustacea, Decapoda, Brachyura). *Bull. mar. Sci.* **19**: 409-427.
- and E. GOTTLIEB. 1956. Two interesting crabs (Crustacea Decapoda, Brachyura) from Mersin Bay, S.E. Turkey. *Zool. Meded., Leiden.* **34**: 287-299.
- HOOD, M. A. 1962. Studies on the larval development of *Rhithropanopeus harrisi* (Gould) of the family Xanthidae (Brachyura). *Gulf Res. Rept.* **1**: 122-130.
- HOPKINS, S. H. 1947. The nemertean *Carcinonemertes* as an indicator of the spawning history of the host, *Callinectes sapidus*. *J. Parasitol.* **33**: 146-150.
- HORCH, K. 1971. An organ for hearing and vibration sense in the ghost crab *Ocypode*. *Z. Vergl. Physiol.* **73**: 1-21.
- HORCH, K. W. and M. SALMON. 1969. Production, perception and reception of acoustic stimuli by semiterrestrial crabs (Genus *Ocypode* and *Uca*, family Ocypodidae). *Forma et Functio.* **1**: 1-25.
- HORN, E. C. and M. S. KERR. 1963. Hemolymph protein and copper concentrations of adult blue crabs (*Callinectes sapidus* Rathbun). *Biol. Bull.* **125**: 499-507.
- and ———. 1969. The hemolymph protein of the blue crab *Callinectes sapidus*. I. Hemocyanins and certain other major protein constituents. *Comp. Biochem. Physiol.* **29**: 493-508.
- HUGHES, D. A. 1973. On mating and the "copulation burrows" of crabs of the genus *Ocypode* (Decapoda, Brachyura). *Crustaceana.* **24**: 72-76.
- HULINGS, N. C. 1961. The barnacle and decapod fauna from the near shore area of Panama City, Florida. *Q. Jl. Fla. Acad. Sci.* **24**: 215-222.
- HUMES, A. G. 1941a. Notes on *Octolasmis mulleri* Coker, a barnacle commensal on crabs. *Trans. Am. microsc. Soc.* **60**: 101-103.

- . 1941b. A new harpacticoid copepod from the gill chambers of a marsh crab. *Proc. U.S. natn. mus.* **90**: 370-386.
- . 1958. *Antillessia cardisomae*, n. gen. and sp. (Copepoda: Harpacticoida) from the gill chambers of land crabs, with observations on the related genus *Cancricola*. *J. Wash. Acad. Sci.* **48**: 77-89.
- HYATT, G. W. 1974. Behavioural evidence for light intensity discrimination by the fiddler crab *Uca pugilator* (Brachyura, Ocypodidae). *Anim. Behav.* **22**: 796-801.
- . 1975. Physiological and behavioral evidence for colour discrimination by fiddler crabs (Brachyura, Ocypodidae, genus *Uca*). In: F. J. Vernberg (ed.) *Physiological Ecology of Estuarine Organisms*. Univ. South Carolina Press.
- HYMAN, L. H. 1955. *The Invertebrates. IV. Echinodermata: The Coelomate Bilateria*. McGraw-Hill: New York. 763 pp.
- HYMAN, O. W. 1920. The development of *Gelasimus* after hatching. *J. Morph.* **33**: 485-524.
- . 1922. Adventures in the life of a fiddler crab. *Rep. Smithsonian Inst. for 1920*. Pp. 443-460.
- . 1924. Studies on larvae of crabs of the family Grapsidae. *Proc. U.S. natn. mus.* **65**: 1-8.
- . 1925. Studies on the larvae of the crabs of the family Xanthidae. *Proc. U.S. natn. mus.* **67**: 1-22.
- IVERSON, E. S. and G. L. BEARDSLEY. 1976. Shell disease in crustaceans indigenous to South Florida. *Progr. Fish-Culturist.* **38**: 195-196.
- IVES, J. E. 1891. Crustacea from the northern coast of Yucatan, the harbor of Vera Cruz, the west coast of Florida and the Bermuda Islands. *Proc. Acad. Nat. Sci. Phil.* **1891**: 176-207.
- JACHOWSKI, R. 1963. Observations on the moon jelly, *Aurelia aurita*, and the spider crab, *Libinia dubia*. *Chesapeake Sci.* **4**: 195.
- JACHOWSKI, R. L. 1974. Agonistic behavior of the blue crab, *Callinectes sapidus* Rathbun. *Behaviour.* **50**: 232-253.
- JAHROMI, S. S. and C. K. GOVIND. 1976. Ultrastructural diversity in motor units of crustacean stomach muscles. *Cell Tiss. Res.* **166**: 159-166.
- JAWORSKI, E. 1970. Biogeography of the blue crab fishery, Barataria Bay, Louisiana, Ph.D. Dissertation, Louisiana State Univ., Baton Rouge.
- JOHNSON, G. E. 1965. An ethological study of the rock crab, *Grapsus grapsus* (family Grapsidae) with emphasis on behavioral observations during ontogeny and with habitat. *Am. Zoologist.* **5**: 632.
- JOHNSON, I. S. 1952. The demonstration of a "host-factor" in commensal crabs. *Trans. Kansas Acad. Sci.* **55**: 458-464.
- JOHNSON, P. T. 1976. Gas-bubble disease in the blue crab, *Callinectes sapidus*. *J. Invert. Pathol.* **27**: 247-253.
- JONES, H. G. 1968. Preliminary studies on the brachyuran Crustacea of Barbados. II. *J. Barbados Mus.* **32**: 187-189.
- JONES, L. L. 1940. An introduction of an Atlantic crab into San Francisco Bay. *Proc. 6th Pac. Sci. Congr.* **3**: 485-486.
- . 1941. Osmotic regulation in several crabs of the Pacific coast of North America. *J. cell. comp. Physiol.* **18**: 79-92.
- KALBER, F. A., JR. and J. D. COSTLOW, JR. 1966. The ontogeny of osmoregulation and its neurosecretory control in the decapod crustacean *Rhithropanopeus harrisi* (Gould). *Am. Zoologist.* **6**: 221-229.

- KALBER, F.A. and J. D. COSTLOW, JR. 1968. Osmoregulation in larvae of the land-crab, *Cardisoma guanhumii* Latreille. *Am. Zoologist*. **8**: 411-416.
- KARANDIEVA, O and A. S. LEE. 1967. Intensidad de respiracion y osmoregulacion del cangrejo comercial *Menippe mercenaria* (Say) de las aguas Cubanas. *Inst. Oceanol. Acad. Cien. Cuba, Estudios*. **2**: 5-19.
- KEITH, D. E. and N. C. HULINGS. 1965. A quantitative study of selected nearshore infauna between Sabine Pass and Bolivar Point, Texas. *Publs Inst. mar. Sci. Univ. Tex.* **10**: 33-40.
- KERWIN, J. A. 1971. Distribution of the fiddler crab (*Uca minax*) in relation to marsh plants within a Virginia estuary. *Chesapeake Sci.* **12**: 180-183.
- KIRCHER, A. B. 1970. The zoeal stages and glaucothoe of *Hypoconcha arcuata* Stimpson (Decapoda: Dromiidae) reared in the laboratory. *Bull. mar. Sci.* **20**: 769-792.
- KLAASSEN, F. 1973. Stridulation und Kommunikation durch Substratschall bei *Gecarcinus lateralis* (Crustacea Decapoda). *J. comp. Physiol.* **83**: 73-79.
- . 1975. Ökologische und ethologische Untersuchungen zue Fortpflanzungsbiologie von *Gecarcinus lateralis* (Decapoda, Brachyura). *Forma Functio*. **8**: 101-174.
- KNIGHT, M. D. 1968. The larval development of *Raninoides benedicti* Rathbun (Brachyura, Raninidae), with notes on the Pacific records of *Raninoides laevis* (Latreille). *Crustaceana, Suppl.* **2**: 145-169.
- KNOPF, G. N. 1966. Observations on behavioral ecology of the fiddler crab, *Uca pugilator* (Bosc). *Crustaceana*. **11**: 302-306.
- KRAMER, P. 1967. Beobachtungen zur Biologie und zum Verhalten der Klippenkrabbe *Grapsus grapsus* L. (Brachyura Grapsidae) auf Galapagos und am ekuadorianischen Festland. *Z. Tierpsychol.* **24**: 385-402.
- KRANTZ, G. E., R. R. COLWELL and E. LOVELACE. 1969. *Vibrio parahaemolyticus* from the Blue Crab *Callinectes sapidus* in Chesapeake Bay. *Science*. **164**: 1286-1287.
- KRUCZYNSKI, W. L. 1971. Relationship of the pea crab, *Pinnotheres maculatus* (Say), with the scallops, *Argopecten irradians concentricus* (Say) and *Argopecten gibbus* (Linne). Ph.D. Dissertation, Univ. North Carolina, Chapel Hill. 120 pp.
- . 1972. The effect of the pea crab, *Pinnotheres maculatus* Say, on growth of the bay scallop, *Argopecten irradians concentricus* (Say). *Chesapeake Sci.* **13**: 218-220.
- . 1975. A radioactive tracer study of food uptake by *Pinnotheres maculatus* in molluscan hosts. *Biol. Bull.* **148**: 60-67.
- LAIRD, C. E. and P. A. HAEFNER, JR. 1976. Effects of intrinsic and environmental factors on oxygen consumption in the blue crab, *Callinectes sapidus* Rathbun. *J. exp. mar. Biol. Ecol.* **22**: 171-178.
- LANDERS, W. S. 1954. Notes on the predation of the hard clam, *Venus mercenaria* by the mud crab, *Neopanope texana*. *Ecology*. **35**: 422.
- LANGDON, J. W. 1971. Shape discrimination and learning in the fiddler crab *Uca pugilator*. Ph.D. Dissertation. Florida State University, Tallahassee.
- LEARY, S. P. 1967. Crabs of Texas. Texas Parks and Wildlife Bulletin, no. **43**, 57 pp.
- LEBOUR, M. V. 1928. Studies of the Plymouth Brachyura. II. The larval stages of *Ebalia* and *Pinnotheres*. *J. mar. Biol. Assoc. U.K.* **15**: 109-123.
- . 1944. Larval crabs from Bermuda. *Zoologica*. **29**: 113-128.
- . 1950. Notes on some larval decapods (Crustacea) from Bermuda. *Proc. Zool. Soc. London*. **120**: 369-379.
- LEFFLER, C. W. 1972. Some effects of temperature on the growth and metabolic rate of juvenile blue crab, *Callinectes sapidus*, in the laboratory. *Mar. Biol.* **14**: 104-110.

- . 1973. Metabolic rate in relation to body size and environmental oxygen concentration in two species of xanthid crabs. *Comp. Biochem. Physiol.* **44A**: 1047-1052.
- LEIPPER, D. F. 1954. Physical oceanography of the Gulf of Mexico. In: Gulf of Mexico. Its origin, waters, and marine life. *Fishery Bull. Fish Wildl. Serv. U.S.* **55**: 119-137.
- LE LOEUFF, P., A. INTES and J. C. LE GUEN. 1974. Note sur les premiers essais de capture du crabe profond *Geryon quinquedens* en Côte d'Ivoire. *Doc. Sci. Cent. Rech. Océanogr. Abidjan.* **5**: 73-84.
- LEONE, C. A. 1951. A serological analysis of the systematic relationship of the brachyuran crab *Geryon quinquedens*. *Biol. Bull.* **100**: 44-48.
- LEWIS, E. G. 1976. Epizoots associated with *Bathynectes superbus* (Decapoda: Portunidae). *Fishery Bull. Fish Wildl. Serv. U.S.* **74**: 225-227.
- LEWIS, E. G. and P. A. HAEFNER, JR. 1976. Oxygen consumption of the blue crab, *Callinectes sapidus* Rathbun, from proecdysis to postecdysis. *Comp. Biochem. Physiol.* **54A**: 55-60.
- LOBO de MESQUITA, A. L. 1972. Dados biométricos do siri canela, *Portunus spinimanus* Latreille. 1819 (Decapoda-Brachyura-Portunidae). *Arq. Ciên. Mar.* **12**: 88-90.
- LUDOLPH, C., D. PAGNANALLI and M. I. MOTE. 1973. Neural control of migration of proximal screening pigment by reticular cells of the swimming crab *Callinectes sapidus*. *Biol. Bull.* **145**: 159-170.
- LUNZ, G. R., JR. 1937. Xanthidae (mud crabs) of the Carolinas. *The Charleston Mus., Leaflet.* **9**: 9-27.
- . 1939. New crustacean records for the Carolinas and Florida. *J. Elisha Mitchell Sci. Soc.* **55**: 335-338.
- . 1947. *Callinectes* versus *Ostrea*. *J. Elisha Mitchell Sci. Soc.* **63**: 81.
- LYNCH, M. P., K. L. WEBB and W. A. van ENGEL. 1973. Variation in serum constituents of the blue crab *Callinectes sapidus*: chloride and osmotic concentration. *Comp. Biochem. Physiol.* **44A**: 719-734.
- LYNCH, S. A. 1954. Geology of the Gulf of Mexico. In: Gulf of Mexico. Its origin, waters, and marine life. *Fishery Bull. Fish Wildl. Serv. U.S.* **55**: 67-86.
- LYONS, W. G., S. P. COBB, D. K. CAMP, J. A. MOUNTAIN, T. SAVAGE, L. LYONS and E. A. JOYCE. 1971. Preliminary inventory of marine invertebrates collected near the electrical generating plant, Crystal River, Florida, in 1969. *Prof. Pap. Fla. Dept. Nat. Res.* **14**.
- MACCAGNO, T. 1928. Crostacei decapodi. Le specie del genere *Uca* Leach conservate nel Regio Museo Zoologico di Torino. *Boll. Musei Zool. Anat. comp. R. Univ. Torino.* **41**: 1-52.
- MACGINITIE, G. E. and N. MACGINITIE. 1968. *Natural History of Marine Animals*. 2d Ed. McGraw-Hill: New York. 523 pp.
- MAHOOD, R., M. MCKENZIE, D. MIDDAGH, S. BOLLAR, J. DAVIS and D. SPITSBERGEN. 1970. A report on the cooperative blue crab study—South Atlantic states. *Georgia Game & Fish Comm., Coastal Fish. Div. Contrib. Ser.* **19**: 1-32.
- MANGUM, C. P. and L. M. AMENDE. 1972. Blood osmotic concentration of blue crabs (*Callinectes sapidus* Rathbun) found in fresh water. *Chesapeake Sci.* **13**: 318-320.
- , S. U. SILVERTHORN, J. L. HARRIS, D. W. TOWNE and A. R. KRALL. 1976. The relationship between blood pH, ammonia excretion and adaptation to low salinity in the blue crab *Callinectes sapidus*. *J. exp. Zool.* **195**: 129-136.
- MANGUM, C. and D. TOWLE. 1977. Physiological adaptation to unstable environments. *Am. Scient.* **65**: 67-75.
- MANGUM, C. P. and A. L. WEILAND. 1975. The function of hemocyanin in respiration of the blue crab *Callinectes sapidus*. *J. Exp. Zool.* **193**: 257-264.

- MANNING, R. B. 1961. Some growth changes in the stone crab, *Menippe mercenaria* (Say). *Q. Jl. Fla. Acad. Sci.* **23**: 273-277.
- . 1975. The identity of *Raninoides fossor* A. Milne Edwards & Bouvier, 1923 (Decapoda). *Crustaceana.* **29**: 297-298.
- and A. J. PROVENZANO, JR. 1961. The occurrence of *Ucides cordatus* (Linnaeus 1763) (Decapoda) in the United States. *Crustaceana.* **2**: 158-159.
- MANTEL, L. H. 1967. Asymmetry potentials, metabolism and sodium fluxes in gills of the blue crab, *Callinectes sapidus*. *Comp. Biochem. Physiol.* **20**: 743-753.
- . 1968. The foregut of *Gecarcinus lateralis* as an organ of salt and water balance. *Am. Zoologist.* **8**: 433-442.
- , D. E. BLISS, S. W. SHEEHAN and E. A. MARTINEZ. 1975. Physiology of hemolymph, gut fluid, and hepatopancreas of the land crab *Gecarcinus lateralis*. (Freminville) in various neuroendocrine states. *Comp. Biochem. Physiol.* **15A**: 663-671.
- MARCHAND, L. J. 1946. The saber crab, *Platychirograpsus typicus* Rathbun, in Florida: a case of accidental dispersal. *Q. Jl. Fla. Acad. Sci.* **9**: 93-100.
- MASON, C. A. 1970. Function of the pericardial sacs during the molt cycle in the land crab *Gecarcinus lateralis*. *J. exp. Zool.* **174**: 381-390.
- MATTHEWS, L. H. 1930. Notes on the fiddler-crab, *Uca leptodactyla*, Rathbun. *Ann. Mag. nat. Hist.* (ser. 10) **5**: 659-663.
- MAYNARD, D. M. 1961a. Thoracic neurosecretory structures in Brachyura. I. Gross anatomy. *Biol. Bull.* **121**: 316-329.
- . 1961b. Thoracic neurosecretory structures in Brachyura. II. Secretory neurons. *Gen. Comp. Endocrinol.* **1**: 237-263.
- and M. R. DANDO. 1974. The structure of the stomatogastric neuromuscular system in *Callinectes sapidus*, *Homarus americanus* and *Panulirus argus* (Decapoda Crustacea). *Phil. Trans. R. Soc. ser. B.* **263**: 161-220.
- and E. A. MAYNARD. 1962. Thoracic neurosecretory structures in Brachyura. III. Microanatomy of peripheral structures. *Gen. Comp. Endocrinol.* **2**: 12-28.
- McDERMOTT, J. J. 1960. The predation of oysters and barnacles by crabs of the family Xanthidae. *Proc. Pa. Acad. Sci.* **34**: 199-211.
- . 1962. The occurrence of *Pinnixa cylindrica* (Crustacea, Pinnotheridae) in the tubes of the lugworm, *Arenicola cristata*. *Proc. Penn. Acad. Sci.* **36**: 53-57.
- and F. B. FLOWER. 1953. Preliminary studies of the common mud crabs on oyster beds of Delaware Bay. *Nat. Shellfish. Assoc. 1952 Conv. Addr.* pp. 87-50.
- McMAHAN, M. R. 1967. The larval development of *Neopanope texana texana* (Stimpson) (Xanthidae). *Leaflet. Ser. Florida Bd. Conserv., Div. Salt Water Fish., Mar. Lab.*, no. **2**: pt. 1:1-16.
- McRAE, E. D. 1961. Red crab explorations off the northeastern coast of the United States. *Commer. Fish. Rev.* **23**(5): 5-10.
- MEADE, T. L. and G. W. GRAY, JR. 1973. The red crab. *Tech. Rept. Univ. R.I. Mar.* **11**: 1-21.
- MENDELSON, M. 1963. Some factors in the activation of crab movement receptors. *J. exp. Biol.* **40**: 157-169.
- . 1966. The site of impulse initiation in bipolar receptor neurons of *Callinectes sapidus* L. *J. exp. Biol.* **45**: 411-420.
- MENZEL, R. W. (ed.) 1971. Checklist of the marine fauna and flora of the Apalachee Bay and the St. George's Sound area. 3d edit. Dept. Oceanogr. Florida State Univ.: Tallahassee.

- MENZEL, R. and S. H. HOPKINS. 1956. Crabs as predators of oysters in Louisiana. *Proc. Nat. Shell Fish. Assoc.* **46**: 177-184.
- MENZEL, R. W. and F. W. NICHY. 1958. Studies of the distribution and feeding habits of some oyster predators in Alligator Harbor, Fla. *Bull. Mar. Sci. Gulf Carib.* **8**: 125-145.
- MENZIES, R. J. 1948. A revision of the brachyuran genus *Lophopanopeus*. *Occas. Pap. Allan Hancock.* **4**: 1-42.
- MILLER, D. C. 1961. The feeding mechanism of fiddler crabs, with ecological considerations of feeding adaptations. *Zoologica.* **46**: 89-100.
- . 1965. Studies of the systematics, ecology and geographical distribution of certain fiddler crabs. Doctoral Dissertation, Duke University. 240 pp.
- . 1968. Growth in *Uca*. 1. Ontogeny of asymmetry in *Uca pugilator* (Bosc) (Decapoda, Ocypodidae). *Crustaceana.* **24**: 119-131.
- and F. J. VERNBERG. 1968. Some thermal requirements of fiddler crabs of the temperate and tropical zones and their influence on geographic distribution. *Am. Zoologist.* **8**: 459-469.
- MILLER, K. G. and D. MAUER. 1973. Distribution of the fiddler crabs, *Uca pugnax* and *Uca minax*, in relation to salinity in Delaware rivers. *Chesapeake Sci.* **14**: 219-221.
- MILLER, R. E., S. D. SULKIN and R. L. LIPPSON. 1975. Composition and seasonal abundance of the blue crab, *Callinectes sapidus* Rathbun, in the Chesapeake and Delaware Canal and adjacent waters. *Chesapeake Sci.* **16**: 27-31.
- MILNE, L. J. and M. J. MILNE. 1946. Notes on the behavior of the ghost crab. *Am. Nat.* **80**: 362-380.
- MILNE EDWARDS, A. 1880. Reports on the results of dredging under supervision of Alexander Agassiz in the Gulf of Mexico, and in the Caribbean Sea, 1877, '78, '79, by the U.S. Coast Survey Steamer "Blake". VIII Études préliminaires sur les Crustacés. *Bull. Mus. Comp. Zool.* **8**: 1-68.
- and E. L. BOUVIER. 1902. Reports on the results of dredging, under the supervision of Alexander Agassiz, in the Gulf of Mexico (1877-78), in the Caribbean Sea (1878-79), and along the Atlantic coast of the United States (1880), by the U.S. Coast Survey Steamer "Blake". XXXIX. Les Dromiaces et Oxystomes. *Mem. Mus. Comp. Zool.* **27**: 1-127.
- and ———. 1923. Reports on the results of dredging under the supervision of Alexander Agassiz in the Gulf of Mexico (1877-78), in the Caribbean Sea (1878-79), and along the Atlantic coast of the United States (1880), by the U.S. Coast Survey Steamer Blake. . . . XLVII. Les Porcellanides et des Brachyures. *Mem. Mus. Comp. Zool.* **47**: 283-395.
- MOFFETT, S. 1975. Motor patterns and structural interactions of basi-ischiopodite levator muscles in routine limb elevation and production of autotomy in the land crab, *Cardisoma guanhumi*. *J. comp. Physiol.* **96**: 285-305.
- MONOD, TH. 1956. Hippidea et Brachyura ouest-africains. *Mem. Inst. Franc. Afr. No.* **45**: 1-674.
- MOORE, D. J. 1971. The uptake and concentration of fluoride by the blue crab, *Callinectes sapidus*. *Chesapeake Sci.* **12**: 1-13.
- MOOTZ, C. A. and C. E. EPIFANIO. 1974. An energy budget for *Menippe mercenaria* larvae fed *Artemia* nauplii. *Biol. Bull.* **146**: 44-55.
- MORE, W. R. 1969. A contribution to the biology of the blue crab, (*Callinectes sapidus* Rathbun) in Texas, with a description of the fishery. *Technical Ser. Texas Parks and Wildl. Dept.* **1**: 31 pp.
- MOREIRA, C. 1913. Embryologie de *Cardisoma guanhumi*, Latr. *Mem. Soc. Zool. France.* **25**: 155-161.

- MORRISON, P. R. and K. C. MORRISON. 1952. Bleeding and coagulation in some Bermudan Crustacea. *Biol. Bull.* **103**: 395-406.
- MURINA, V. V., V. D. CHUJCHIN, O. GOMEZ, and G. SUAREZ. 1969. Distribucion cuantitativa de la macrofauna bentonica del sublitoral superior de la plataforma cubana (region noroccidental). *Acad. Cienc. Cuba. Ser. Oceanol.* **6**: 1-14.
- MUSICK, J. A. and J. D. McEACHRAN. 1972. Autumn and winter occurrence of decapod crustaceans in Chesapeake Bight, U.S.A. *Crustaceana.* **22**: 190-200.
- NAYLOR, E. and M. J. ISAAC. 1973. Behavioural significance of pressure responses in megalopa larvae of *Callinectes sapidus* and *Macropipus* sp. *Mar. Behav. Physiol.* **1**: 341-350.
- NEWCORBE, C. L. and M. R. ROGERS. 1947. Studies of a fungus parasite that infects blue crab eggs. *Turtlex News.* **25**: 180-186.
- NIMMO, D. R., P. D. WILSON, R. R. BLACKMAN and A. J. WILSON, JR. 1971. Polychlorinated biphenyl absorbed from sediments by fiddler crabs and pink shrimp. *Nature.* **231**: 50-53.
- NOMURA, H. and J. FAUSTO FILHO. 1966. Alguns dados biométricos de dois crustáceos marinhos do nordeste Brasileiro. *Arq. Est. Biol. Mar. Univ. Fed. Ceará.* **6**: 119-121.
- NORSE, E. A. 1972. Preliminary study of six Jamaican blue crabs, genus *Callinectes* (Decapods: Portunidae). In: Marine studies on the north coast of Jamaica (ed. G. J. Bakus). *Atoll Res. Bull.* **152**: 5.
- NOVAK, A. and M. SALMON. 1974. *Uca panacea*, a new species of fiddler crab from the Gulf coast of the United States. *Proc. Biol. Soc. Wash.* **87**: 313-326.
- NUNNEMACHER, R. F. 1965. The fine structure of the optic tracts of Decapoda. In: "Proceedings of the International Symposium on the functional organization of the compound eye, 25-27 October, 1965, Stockholm, Sweden. Symp. Publ. Div., Pergamon Press. Vol. 7. pp. 363-375.
- O'CONNOR, J. D. and L. I. GILBERT. 1968. Aspects of lipid metabolism in crustaceans. *Am. Zoologist.* **8**: 529-539.
- OGAWA, M., T. T. ALVES, M. da C. CALAND-NORONHA, C. A. E. ARARIPE, and E. L. MAIA. 1973. Industrialização do caranguejo uçá, *Ucides cordatus* (Linnaeus). I.—Técnicas para o processamento da carne. *Arq. Ciên. Mar.* **13**: 31-37.
- , T. T. ALVES, R. B. FILHO, A. S. RODRIGUES and E. L. LIMA. 1973. Industrialização do caranguejo uçá, *Ucides cordatus* (Linnaeus). II.—Aproveitamento dos resíduos e carapaça. *Arq. Ciên. Mar.* **13**: 83-90.
- O'HARA, J. 1973. The influence of temperature and salinity on the toxicity of cadmium to the fiddler crab, *Uca pugilator*. *Fishery Bull. Fish Wildl. Serv. U.S.* **71**: 149-153.
- OLER, T. M. 1941. Some notes on the occurrence of a small land crab (*Sesarma cinereum* Bosc) on the Magorothy River, Md. *Bull. nat. Hist. Soc. Md.* **11**: 51-53.
- de OLIVEIRA, L. P. H. 1939a. Contribuição ao conhecimento dos crustáceos do Rio de Janeiro. Gênero *Uca* (Decapoda: Ocypodidae). *Mems. Inst. Oswaldo Cruz.* **34**: 115-148.
- . 1939b. Observações sobre a biologia dos adultos do gênero *Uca* Leach 1814. *Liv. Hom. Profs. A. e M. Ozorio de Almeida Rio de Janeiro*, pp. 490-497.
- . 1939c. Alguns fatores que limitam o habitat de varias espécies de carangueiros do gênero *Uca* Leach. *Mems. Inst. Oswaldo Cruz.* **34**: 519-526.
- . 1940. Observações preliminares sobre a biologia dos crustáceos do gênero *Panopeus* Milne Edwards, 1834. *Mems. Inst. Oswaldo Cruz.* **35**: 153-171.
- . 1946. Ecological studies on the edible crabs *Uca* and *Guaianu*, *Cardisoma guanhumi*, and *Ucides cordatus*. *Mems. Inst. Oswaldo Cruz.* **44**: 295-322.

- ONG, K.-S. and J. D. COSTDOW, JR. 1970. The effect of salinity and temperature on the larval development of the stone crab, *Menippe mercenaria* (Say), reared in the laboratory. *Chesapeake Sci.* **11**: 16-29.
- ORR, P. R. 1955. Heat death. 1. Time temperature relationship in marine animals. *Physiol. Zool.* **28**: 290-293.
- OTT, F. S. and R. B. FORWARD, JR. 1976. The effect of temperature on phototaxis and geotaxis by larvae of the crab *Rhithropanopeus harrisi* (Gould). *J. exp. mar. Biol. Ecol.* **23**: 97-107.
- OVERSTREET, R. and H. M. PERRY. 1972. A new microphallid trematode from the blue crab in the northern Gulf of Mexico. *Trans. Am. Mic. Soc.* **91**: 436-440.
- PALMER, J. D. 1967. Daily and tidal components in the persistent rhythmic activity of the crab, *Sesarma*. *Nature.* **215**: 64-66.
- PARK, J. R. 1969. A preliminary study of portunid crabs in Biscayne Bay. *Q. Jl. Fla. Acad. Sci.* **32**: 12-20.
- PARKER, R. H. 1959. Macro-invertebrate assemblages of central Texas coastal bays and Laguna Madre. *Bull. Am. Ass. Petrol. Geol.* **43**: 2100-2166.
- PATTON, W. K. 1967. Studies on *Domecia acanthophora*, a commensal crab from Puerto Rico, with particular reference to modifications of the coral host and feeding habits. *Biol. Bull.* **132**: 56-67.
- PAULEY, G. B., M. W. NEWMAN and E. GOULD. 1975. Serum changes in the blue crab, *Callinectes sapidus*, associated with *Paramoeba pernicioso*, the causative agent of Gray Crab Disease. *Mar. Fish. Rev.* **37**: 34-38.
- PAYEN, G., J. D. COSTLOW, JR. and H. CHARNIAUX-COTTON. 1971. Etude comparative de l'ultrastructure des glandes androgènes de Crabes normaux et pédonulectomisés pendant le vie larvaire ou apres la puberté chez les espèces: *Rhithropanopeus harrisi* (Gould) et *Callinectes sapidus* Rathbun. *Gen. comp. Endocrin.* **17**: 526-542.
- PEARCE, J. B. 1964. On reproduction in *Pinnotheres maculatus* (Decapoda: Pinnotheridae). *Biol. Bull.* **127**: 384.
- . 1966. On *Pinnixa faba* and *Pinnixa littoralis* (Decapoda: Pinnotheridae) symbiotic with the clam, *Tresus capax* (Pelecypoda: Mactridae). Pp. 565-589. In: Barnes, H. ed.), *Some Contemporary Studies in Marine Science*. Allen & Unwin, Ltd.
- PEARSE, A. S. 1913. On the habits of the crustaceans found in *Chaetopterus* tubes at Woods Hole, Mass. *Biol. Bull.* **24**: 102-114.
- . 1914. On the habits of *Uca pugnax* (Smith) and *U. pugilator* (Bosc). *Trans. Wisc. Acad. Sci.* **17**: 791-802.
- . 1929. The ecology of certain estuarine crabs at Beaufort, N. C. *J. Elisha Mitchell Scient. Soc.* **44**: 230-237.
- . 1932a. Observations on the parasites and commensals found associated with crustaceans of fishes at Dry Tortugas, Fla. *Pap. Tortugas Lab.* **28**: 103-115.
- . 1932b. Freezing points of bloods of certain littoral and estuarine animals. *Pap. Tortugas Lab.* **28**: 93-102.
- . 1934. Inhabitants of certain sponges at Dry Tortugas. *Pap. Tortugas Lab.* **28**: 117-124.
- . 1949. Observations on flatworms and nemertean collected at Beaufort, N.C. *Proc. U.S. natn mus.* **100**: 25-38.
- . 1952. Parasitic Crustacea from the Texas coast. *Publs Inst. mar. Sci. Univ. Tex.* **2**: 5-42.
- and L. G. WILLIAMS. 1951. The biota of the reefs off the Carolinas. *J. Elisha Mitchell Sci. Soc.* **67**: 133-161.

- PEQUEGNAT, L. H. and J. P. RAY. 1974. Crustacea and other arthropods. Pp. 232-288. In: T. J. Bright and L. H. Pequegnat (eds.), *Biota of the West Flower Garden Bank*. Gulf Publ. Co.: Houston.
- PEQUEGNAT, W. E. 1970. Deep-water brachyuran crabs. P. 171-204. In: F. A. Chace, Jr. and W. E. Pequegnat, eds. Texas A&M Univ. Oceanogr. Stud., I. *Contributions on the Biology of the Gulf of Mexico*. Gulf Publishing Co.: Houston.
- PERKINS, F. O. 1975. Fine structure of *Minichinia* sp. (Haplosporida) sporulation in the mud crab, *Panopeus herbstii*. *Mar. Fish. Rev.* **37**:46-60.
- PERRET, W. S. 1967. Occurrence, abundance and size distribution of the blue crab *Callinectes sapidus* taken with otter trawl in Vermillion Bay, Louisiana. *Proc. La Acad. Sci.* **30**: 63-69.
- PERRY, H. M. 1973. The occurrence of *Callinectes bocourti* (A. Milne Edwards, 1879) (Decapoda, Portunidae) in Biloxi Bay, Mississippi, U.S.A. *Crustaceana.* **25**: 110.
- . 1975. The blue crab fishery in Mississippi. *Gulf Res. Rept.* **5**: 39-57.
- PORTER, H. J. 1960. Zoal stages of the stone crab, *Menippe mercenaria* Say. *Chesapeake Sci.* **1**: 168-177.
- POWELL, E. H., JR. and G. GUNTER. 1968. Observations on the stone crab, *Menippe mercenaria* Say, in the vicinity of Port Aransas, Texas. *Gulf Res. Rep.* **2**: 285-299.
- POWERS, L.W. 1973. Ecological aspects of burrows and fiddler crab behavior. *Am. Zoologist.* **13**: 1271.
- . 1975. Fiddler crabs in a nontidal environment. *Contrib. mar. Sci.* **19**: 67-78.
- and J. F. COLE. 1976. Temperature variation in fiddler crab microhabitats. *J. exp. mar. Biol. Ecol.* **21**: 141-158.
- PRETZMANN, G. 1965. Verläufiger Bericht über die Familie Pseudothelphusidae. *Anz. öst. Akad. Wiss.* **1965**: 1-10.
- PROVENZANO, A. J. 1961. A North American record for *Callinectes boucourti* (A. Milne Edwards, 1879) (Decapoda, Portunidae). *Crustaceana.* **3**: 167.
- PYLE, R. and E. CRONIN. 1950. The general anatomy of the blue crab. *Publ. Ches. Biol. Lab.* **87**: 38 pp.
- QUINN, D. J. and C. E. LANE. 1966. Ionic regulation and Na⁺-K⁺-stimulated ATPase activity in the land crab, *Cardisoma guanhumii*. *Comp. Biochem. Physiol.* **19**: 533-543.
- and C. E. LANE. 1967. Na⁺- and K⁺-stimulated respiration in the excised gill of the land crab *Cardisoma guanhumii*. *Biol. Bull.* **133**: 245-254.
- RAO, K. RANGA and M. FINGERMAN. 1968. Dimorphic variants of the fiddler crab *Uca pugilator* and their chromatophore responses. *Proc. La. Acad. Sci.* **31**: 27-38.
- and ———. 1969. The influence of size on the response of melanophores in the fiddler crab, *Uca pugilator*, to eyestalk extracts. *Z. vergl. Physiol.* **62**: 86-96.
- and ———. 1970. Action of biogenic amines on crustacean chromatophores. II. Analysis of the responses of erythrophores in the fiddler crab, *Uca pugilator*, to indolealkylamines and an eyestalk hormone. *Comp. Gen. Pharmacol.* **1**: 117-126.
- , ——— and C. K. BARTELL. 1967. Physiology of the white chromatophores in the fiddler crab, *Uca pugilator*. *Biol. Bull.* **133**: 606-617.
- RATHBUN, M. J. 1894. Notes on crabs of the family Inachidae in the U.S. National Museum. *Proc. U.S. natn. mus.* **17**: 43-75.
- . 1897. List of the decapod Crustacea of Jamaica. *Ann. Inst. Jamaica, Kingston.* **1**: 1-46.
- . 1898. The Brachyura of the biological expedition to the Florida Keys and the Bahamas in 1893. *Bull. Lab. Nat. Hist. State Univ. Iowa.* **4**: 250-294.

- . 1901. The Brachyura and Macura of Porto Rico. *Bull. U.S. Fish Comm.* **20**: 1-127.
- . 1918. The grapsoid crabs of America. *Bull. U.S. natn. mus.* **97**: 1-461.
- . 1920. New species of spider crabs from the Straits of Florida and Caribbean Sea. *Proc. Biol. Soc. Washington.* **33**: 23-24.
- . 1925. The spider crabs of America. *Bull. U.S. natn. mus.* **129**: 1-613.
- . 1930. The cancroïd crabs of America of the families Euryalidae, Portunidae, Ateleyclidae, Cancridae and Xanthidae. *Bull. U.S. natn. mus.* **152**: 1-609.
- . 1931a. New crabs from the Gulf of Mexico. *J. Wash. Acad. Sci.* **21**: 125-129.
- . 1931b. Two new crabs from the Gulf of Mexico. *Proc. Biol. Soc. Wash.* **44**: 71-72.
- . 1933. Brachyuran crabs of Porto Rico and the Virgin Islands. In: *Scient. Surv. P. Rico.* **15(1)**: 1-121, fig. 1-107. New York Acad. Sciences.
- . 1937. The oxystomatous and allied crabs of America. *Bull. U.S. natn. mus.* **166**: 1-278.
- RAY, C. 1967. *Gecarcinus lateralis* Fremenville in Texas. *Texas J. Sci.* **19**: 109.
- RADMOND, J. R. 1962. Oxygen-hemocyanin relationships in the land crab, *Cardisoma guanhumi*. *Biol. Bull.* **122**: 252-262.
- . 1968. Transport of oxygen by the blood of the land crab, *Gecarcinus lateralis*. *Am. Zoologist.* **8**: 471-479.
- REED, C. T. 1941. *Marine Life in Texas Waters*. Pp. i-xii, 1-88, Texas Academy of Science, Publ. Nat. Hist.
- REINGOLD, S. C. 1975. Temperature effects on axonal conduction in *Callinectes sapidus* (Rathbun) and *Carcinus maenas* (L.). *Comp Biochem. Physiol.* **51A**: 195-199.
- REINHARD, E. G. 1950a. An analysis of the effects of a sacculinid parasite on the external morphology of *Callinectes sapidus* Rathbun. *Biol. Bull.* **98**: 277-288.
- . 1950b. The morphology of *Loxothylacus texanus* Boschma, a sacculinid parasite of the blue crab. *Tex. J. Sci.* **2**: 360-365.
- RICE, A. L. 1964. The metamorphosis of a species of *Homola* (Crustacea, Decapoda: Dromiacea). *Bull. mar. Sci. Gulf. Caribb.* **14**: 221-238.
- and A. J. PROVENZANO. 1966. The larval development of the West Indian sponge crab *Dromidia antillensis*. *J. Zool., Lond.* **149**: 297-319.
- and A. J. PROVENZANO, JR. 1970. The larval stages of *Homola barbata* (Fabricius) (Crustacea, Decapoda, Homolidae) reared in the laboratory. *Bull. mar. Sci.* **20**: 446-471.
- RICHMOND, E. A. 1962. The fauna and flora of Horn Island, Mississippi. *Gulf Res. Rept.* **1**: 59-106.
- . 1968. A supplement to the fauna and flora of Horn Island, Mississippi. *Gulf Res. Rept.* **2**: 213-254.
- RICKETTS, E. F. and J. CALVIN. 1968. *Between Pacific Tides*. 4th ed., rev. by J. W. Hedgpeth. Stanford Univ. Press. 614 pp.
- RIGHI, G. 1967. Sobre alguns Decápoda do Brasil (Crustacea, Brachyura: Pinnotheridae e Parthenopidae). *Pap. Dep. Zool. S. Paulo* **20**: 99-116.
- RIOJA, E. 1952. Descripción de un nuevo genero de potamonidos cavernícolas y ciegos de la cueva del Tío Ticho, Comitan, Chis. *An. Inst. Biol. Univ. Mex.* **23**: 217-225.
- ROBERT, M. and I. GRAY. 1972. Enzymatic mechanisms during temperature acclimation of the blue crab *Callinectes sapidus*. *Comp. Biochem. Physiol.* **42B**: 377-401.

- ROBERTS, M. H. JR. 1969. Larval development of *Bathynectes superba* (Costa) reared in the laboratory. *Biol. Bull.* **137**: 338-351.
- RODRIGUES DA COSTA, H. 1968a. Crustacea Brachyura récoltés par les draguages de la "Calypso" sur les cotes brésiliennes (1962). *Recl. Trav. Stn. mar. Endoume.* **43**: 333-343.
- . 1968b. Ocorrência do gênero "Cycloes" de Haan, 1837 no Brasil. Descrição de formas jovens de "Cycloes bairdii" Stimpson encontradas no litoral brasileiro (Brachyura, Oxystomata, Calappidae). *Atas Soc. Biol. Rio de J.* **12**: 29-30.
- . 1971. As espécies brasileiras da família Pinnotheridae (Crustacea, Reptantia) com descrição de uma nova espécie (*Fabia sebastianensis*). *Trab. Inst. Oceanogr. Univ. Fed. Pernambuco.* **9/11**: 255-264.
- ROGERS, B. G. 1968. An extension of the range of the pinnotherid crab, *Dissodactylus melittae* Rathbun. *Crustaceana.* **14**: 318.
- ROGERS-TALBERT, R. 1948. The fungus *Lagenidium callinectes* Couch (1942) on eggs of the blue crab in Chesapeake Bay. *Biol. Bull.* **95**: 214-228.
- ROSENBERG, R. and J. D. COSTLOW, JR. 1976. Synergistic effects of cadmium and salinity combined with constant and cycling temperatures on the larval development of two estuarine crab species. *Mar. Biol.* **38**: 291-303.
- ROUSE, W. L. 1970. Littoral Crustacea from southwest Florida. *Q. Jl. Fla. Acad. Sci.* **32**: 127-152.
- RUST, J. D. and F. CARLSON. 1960. Some observations on rearing blue crab larvae. *Chesapeake Sci.* **1**: 196-197.
- RYAN, E. P. 1956. Observations on the life histories and the distribution of the Xanthidae (mud crabs) of Chesapeake Bay. *Amer. Midl. Nat.* **56**: 138-162.
- SAKAI, T. 1939. *Studies on the crabs of Japan.* IV. Brachygnatha, Brachyrhyncha. Yokendo, Tokyo. pp. 365-741.
- . 1965. *The Crabs of Sagami Bay Collected by His Majesty The Emperor of Japan.* East-West Center Press: Honolulu. Text in English and Japanese, 100 color pl. + map.
- . 1969. Two new genera and twenty-two new species of crabs from Japan. *Proc. Biol. Soc. Wash.* **82**: 243-280.
- . 1976a. Notes from the carcinological fauna of Japan (VI). The occurrence of *Callinectes sapidus* Rathbun in Japan. *Res. Crust.* **7**: 29-30.
- . 1976b. *Crabs of Japan and the Adjacent Seas.* 3 vols. Kodansha Ltd., Tokyo. 773 pp., English text.
- SALMON, M. 1965. Waving display and sound production in the courtship behavior of *Uca pugilator*, with comparisons to *U. minax* and *U. pugnax*. *Zoologica.* **50**: 123-150.
- . 1967. Coastal distribution, display and sound production by Florida fiddler crabs (genus *Uca*). *Anim. Behav.* **15**: 449-459.
- . 1971. Signal characteristics and acoustic detection by the fiddler crabs, *Uca rapax* and *Uca pugilator*. *Physiol. Zool.* **44**: 210-224.
- and S. P. ATSAIDES. 1968a. Visual and acoustical signalling during courtship by fiddler crabs (genus *Uca*). *Am. Zoologist.* **8**: 623-639.
- and ———. 1968b. Behavioral, morphological and ecological evidence for two species of fiddler crabs (genus *Uca*) from the Gulf coast of the United States. *Proc. Biol. Soc. Wash.* **81**: 275-290.
- and ———. 1969. Sensitivity to substrate vibration in the fiddler crab, *Uca pugilator* Bosc. *Anim. Behav.* **17**: 68-76.
- and K. W. HORCH. 1972. Acoustic signalling and detection by semiterrestrial crabs of the family Ocypodidae. pp. 60-96. In: H. E. Winn and B. L. Olla (ed.), *Behavior of Marine Animals. Vol. 1—Invertebrates.* Plenum Press.

- _____ and _____. 1973. Vibration reception by the fiddler crab, *Uca minax*. *Comp. Biochem. Physiol.* **44A**: 527-541.
- _____, _____ and G. W. HYATT. 1977. Barth's myochordotonal organ as a receptor for auditory and vibrational stimuli in fiddler crabs (*Uca pugilator* and *U. minax*). *Mar. Behav. Physiol.* **4**: 187-194.
- _____ and J. F. STOUT. 1962. Sexual discrimination and sound production in *Uca pugilator* Bosc. *Zoologica.* **47**: 15-21.
- SANDEEN, M. I. 1950. Chromatophoretropins in the central nervous system of *Uca pugilator*, with special reference to their origins and actions. *Physiol. Zool.* **23**: 337-352.
- SANDIFER, P. A. 1973. Distribution and abundance of decapod crustacean larvae in the York River Estuary and adjacent Lower Chesapeake Bay, Virginia, 1968-1969. *Chesapeake Sci.* **14**: 235-257.
- _____. 1974. Larval stages of the crab, *Pilumnus dasypodus* Kingsley (Crustacea, Brachyura, Xanthidae), obtained in the laboratory. *Bull. mar. Sci.* **24**: 378-391.
- _____ and W. A. van ENGEL. 1971. Larval development of the spider crab, *Libinia dubia* H. Milne Edwards (Brachyura, Majidae, Pisinae), reared in laboratory culture. *Chesapeake Sci.* **12**: 18-25.
- _____ and _____. 1972. Larval stages of the spider crab, *Anasimus latus* Rathbun, 1894 (Brachyura, Majidae, Inachinae) obtained in the laboratory. *Crustaceana.* **23**: 141-151.
- SANDOZ, M. and S. H. HOPKINS. 1947. Early life-history of the oyster crab, *Pinnotheres ostreum* Say. *Biol. Bull.* **93**: 250-258.
- _____ and R. ROGERS. 1944. The effect of environmental factors on hatching, moulting, and survival of zoea larvae of the blue crab, *Callinectes sapidus* Rathbun. *Ecology.* **25**: 216-228.
- SARAIVA da COSTA, R. 1968. Estudo preliminar sôbre a biologia e a pesca do caranguejo *Carpilius corallinus* (Herbst), no estado do Ceará. *Arq. Estac. Biol. Mar. Univ. Ceará.* **3**: 211-219.
- SASTRY, A. N. and R. W. MENZEL. 1962. Influence of hosts on the behavior of the commensal crab *Pinnotheres maculatus* Say. *Biol. Bull.* **123**: 388-395.
- SAVAGE, T. 1971a. Effect of maintenance parameters on growth of the stone crab, *Menippe mercenaria* (Say). *Spec. Sci. Rept. Fla. Dept. Nat. Resources.* **28**: 1-19.
- _____. 1971b. Mating of the stone crab, *Menippe mercenaria* (Say) (Decapoda, Brachyura). *Crustaceana.* **20**: 315-316.
- _____ and M. R. McCAHAN. 1968. Growth of early juvenile stone crabs, *Menippe mercenaria* (Say, 1819). *Spec. Sci. Rept. Fla. Bd. Conserv.* **21**: 1-17.
- _____, J. R. SULLIVAN and C. E. KALMAN. 1974. Claw extraction during molting of a stone crab, *Menippe mercenaria* (Decapoda, Brachyura, Xanthidae). *Florida Mar. Res. Publ.* **4**: 1-5.
- _____, _____ and _____. 1975. An analysis of stone crab (*Menippe mercenaria*) landings on Florida's west coast, with a brief synopsis of the fishery. *Florida Mar. Res. Publ.* **13**: 1-37.
- SAWYER, T. K. 1969. Preliminary study of the epizootiology and host-parasite relationship of *Paramoeba* sp. in the blue crab, *Callinectes sapidus*. *Proc. Natl. Shellfish Assoc.* **59**: 60-64.
- _____, R. COX and M. HIGGINBOTTOM. 1970. Hemocyte values in healthy blue crabs, *Callinectes sapidus*, and crabs infected with the amoeba, *Paramoeba perniciosa*. *J. Invert. Pathol.* **15**: 440-446.
- SCHMITT, W. L. 1931. Some carcinological results of the deeper water trawlings of the *Anton Dohrn*, including description of two new species of Crustacea. *Carnegie Inst. Year Book* **30**: 389-394.

- . 1939. Decapod and other Crustacea collected on the Presidential Cruise of 1938 (with introduction and station data). *Smithson. Misc. Collect.* **98**: 1–29.
- , J. C. McCAIN and E. S. DAVIDSON. 1973. Crustaceorum Catalogus. Decapoda. Brachyura. Fam. Pinnotheridae. (ed. by H.-E. Grunner and L. B. Holthuis). Junk: Haag, Netherlands. 160 pp.
- SCHÖNE, H. 1968. Agonistic and sexual display in aquatic and semi-terrestrial brachyuran crabs. *Am. Zoologist*, **8**: 641–654.
- and I. EIBL-EIBESFELDT. 1965. *Grapsus grapsus* (Brachyura) Drohen. Encyclopaedia Cinematographica. *Publ. Wissenschaft. Film.* **1A**: 391–396.
- and H. SCHÖNE. 1961. Eyestalk movements induced by polarized light in the ghost crab, *Ocypode quadrata*. *Science*, **134**: 675–676.
- and ———. 1963. Balz und andere Verhaltensweisen der Mangrovekrabbe *Goniopsis cruentata* Latr. und das Winkverhalten der eulitoralen Brachyuren. *Z. Tierpsychol.* **20**: 642–656.
- SCHROEDER, W. C. 1959. The lobster, *Homarus americanus*, and the red crab, *Geryon quinquedens*, in the offshore waters of the western North Atlantic. *Deep-Sea Res.* **5**: 266–282.
- SCHWARTZ, B. and S. R. SAFIR. 1915. Habit formation in the fiddler crabs. *J. Anim. Behav.* **5**: 226–239.
- SCHWARTZ, F. J. and D. G. CARGO. 1960. Recent records of the xanthid crab, *Panopeus herbsti*, from Maryland and Virginia waters. *Chesapeake Sci.* **1**: 201–203.
- SELANDER, R. K., W. E. JOHNSON and J. C. AVISE. 1971. Biochemical population genetics of fiddler crabs (*Uca*). *Biol. Bull.* **141**: 402.
- SERENE, R. 1966. Note sur la taxonomie et la distribution géographique des Hapalocarcinidae (Decapoda—Brachyura). *Proc. Symp. Crustacea Ser. 2.* **1**: 395–398.
- SHERIDAN, P. F. 1975. Uptake, metabolism, and distribution of DDT in organs of the blue crab, *Callinectes sapidus*. *Chesapeake Sci.* **16**: 20–26.
- SHIRLEY, T. C. 1974. *Planes cyaneus* Dana, 1852 (Decapoda, Grapsidae) from Padre Island, Texas: a new record for the Gulf of Mexico and the North Atlantic. *Crustaceana*, **26**: 107–108.
- SHOUP, J. B. 1968. Shell opening by crabs of the genus *Calappa*. *Science*, **160**: 887–888.
- SILVERTHORN, S. U. 1975a. Hormonal involvement in thermal acclimation in the fiddler crab *Uca pugilator* (Bosc)—1. Effect of eyestalk extracts on whole animal respiration. *Comp. Biochem. Physiol.* **50A**: 281–283.
- . 1975b. Hormonal involvement in thermal acclimation in the fiddler crab *Uca pugilator* (Bosc)—2. Effects of extracts on tissue respiration. *Comp. Biochem. Physiol.* **50A**: 285–290.
- SIMMONS, E. G. 1957. An ecological survey of the Upper Laguna Madre of Texas. *Publ. Inst. mar. Sci. Univ. Tex.* **4**: 134–155.
- SIMS, H. W., JR. and E. A. JOYCE, JR. 1966. Partial albinism in a blue crab. *Q. Jl. Fla. Acad. Sci.* **28**: 373–374.
- SKINNER, D. M. 1965. Amino acid incorporation into protein during the molt cycle of the land crab, *Gecarcinus lateralis*. *J. exp. Zool.* **160**: 225–234.
- . 1966. Breakdown and reformation of somatic muscle during the molt cycle of the land crab, *Gecarcinus lateralis*. *J. exp. Zool.* **163**: 115–123.
- , and D. E. GRAHAM. 1972. Loss of limbs as a stimulus to ecdysis in Brachyura (true crabs). *Biol. Bull.* **143**: 222–233.
- , D. J. MARSH and J. S. COOK. 1965. Physiological salt solution for the land crab, *Gecarcinus lateralis*. *Biol. Bull.* **129**: 355–365.

- SKOBE, Z. and R. F. NUNNEMACHER. 1970. The fine structure of the circumesophageal nerve in several decapod crustaceans. *J. comp. Neurol.* **139**: 81-92.
- SKORKOWSKI, E. F. 1972. Separation of three chromatophorotropic hormones from the eyestalk of the crab *Rhithropanopeus harrisi* (Gould). *Gen. comp. Endocrin.* **18**: 329-334.
- SMALLEY, A. E. 1970. A new genus of freshwater crabs from Guatemala, with a key to the middle American genera (Crustacea Decapoda, Pseudothelphusidae). *Am. Midl. Nat.* **83**: 96-106.
- SMITH, R. I. 1967. Osmotic regulation and adaptive reduction of water permeability in a brackish water crab, *Rhithropanopeus harrisi* (Brachyura, Xanthidae). *Biol. Bull.* **133**: 643-658.
- SMITH, W. K. and P. C. MILLER. 1973. The thermal ecology of two south Florida fiddler crabs: *Uca rapax* Smith and *U. pugilator* Bosc. *Physiol. Zool.* **46**: 186-207.
- de SOUZA, T. T. and M. da C. CALAND. 1968. Estudo preliminar sobre a bacteriologia do caranguejo *Ucides cordatus* (Linnaeus). *Arq. Estac. Biol. Mar. Univ. Fed. Ceará.* **3**: 107-108.
- SPINDLER-BARTH, M. 1976. A bacterial infection in the common shore crab *Carcinus maenas* and the fiddler crab *Uca pugilator*. *Mar. Biol.* **36**: 1-4.
- SPIRITO, C. P. 1972. An analysis of swimming behavior in the portunid crab *Callinectes sapidus*. *Mar. Behav. Physiol.* **1**: 261-276.
- , W. R. EVOY and W. J. P. BARNES. 1972. Nervous control of walking in the crab, *Cardisoma guanhumi*. I. Characteristics of resistance reflexes. *Z. vergl. Physiol.* **76**: 1-15.
- SPRAGUE, V. 1965. *Nosema* sp. (Microsporida, Nosematidae) in the musculature of the crab, *Callinectes sapidus*. *J. Protozool.* **12**: 66-70.
- . 1966. Two new species of *Plistophora* (Microsporida, Nosematidae) in decapods, with particular reference to one in the blue crab. *J. Protozool.* **13**: 196-199.
- and R. BECKETT. 1966. A disease of blue crabs (*Callinectes sapidus*) in Maryland and Virginia. *J. Invert. Pathol.* **8**: 287-289.
- , R. L. BECKETT and T. K. SAWYER. 1969. A new species of *Paramoeba* (Amoebida, Paramoebidae) parasitic in the crab *Callinectes sapidus*. *J. Invert. Pathol.* **14**: 167-174.
- SPRINGER, S. and H. R. BULLIS. 1956. Collections by the Oregon in the Gulf of Mexico. *Spec. Scient. Rep. U.S. Fish Wildl. Serv.* **196**: 1-134.
- STANDING, J. D. 1972. Larval salinity acclimatization in the tropical shore crab, *Sesarma ricordi*. In: Marine studies on the north coast of Jamaica (ed., G. J. Bakus). *Atoll Res. Bull.* **152**: 6.
- STAUBER, L. A. 1945. *Pinnoheres ostreum*, parasitic on the American oyster, *Ostrea (Gryphea) virginica*. *Biol. Bull.* **88**: 269-291.
- STEINACKER, A. 1975. Perfusion of the central nervous system of decapod crustaceans. *Comp. Biochem. Physiol.* **52A**: 103-104.
- STEPHENSON, W. and M. REES. 1967. Some portunid crabs from the Pacific and Indian Oceans in the collections of the Smithsonian Institution. *Proc. U.S. natn. mus.* **120**: 1-114.
- and ———. 1968. A revision of the genus *Ovalipes* Rathbun, 1898 (Crustacea, Decapoda, Portunidae). *Rec. Aust. Mus.* **27**: 213-261.
- , W. T. WILLIAMS and G. N. LANCE. 1968. Numerical approaches to the relationships of certain American swimming crabs (Crustacea: Portunidae). *Proc. U.S. natn. mus.* **124**: 1-25.
- ŠTAVČIĆ, Z. 1971a. The pathways of brachyuran evolution. *Proc. 1st Symp. Biosyst. Yugoslav.* pp. 187-193.
- . 1971b. The main features of brachyuran evolution. *Sys. Zool.* **20**: 331-340.

- . 1973. The systematic position of the family Raninidae. *Syst. Zool.* **22**: 625–632.
- STEWART, J. R. and J. P. GREEN. 1969. Ecdysone mediated events in the molting of the fiddler crab, *Uca pugilator*. *Am. Zoologist.* **9**: 579.
- STUTMAN, L. J. and M. DOLLIVER. 1968. Mechanism of coagulation in *Gecarcinus lateralis*. *Am. Zoologist.* **8**: 481–489.
- SUBRAHMANYAM, C. B., W. L. KRUCZYNSKI and S. H. DRAKE. 1976. Studies on the animal communities in two North Florida salt marshes. Part II. Macroinvertebrate communities. *Bull. mar. Sci.* **26**: 172–195.
- SUCHENIA, L. M. and R. CLARO MADRUGA. 1967. Datos cuantitativos de la alimentacion del cangrejo comercial *Menippe mercenaria* (Say) y su relacion con el balance energetico del mismo. *Inst. Oceanogr. Acad. Cien. Cuba, Estudios.* **2**: 75–97.
- SULKIN, S. D. 1973. Depth regulation of crab larvae in the absence of light. *J. exp. mar. Biol. Ecol.* **13**: 73–82.
- and C. E. EPIFANIO. 1975. Comparison of rotifers and other diets for rearing early larvae of the blue crab, *Callinectes sapidus* Rathbun. *Estuarine Coastal mar. Sci.* **3**: 109–113.
- TABB, D. C. and R. B. MANNING. 1961. A checklist of the flora and fauna of northern Florida Bay and adjacent brackish water of the Florida mainland collected during the period July, 1957 through September, 1960. *Bull. mar. Sci.* **11**: 552–649.
- TAGATZ, M. E. 1967. Noncommercial crabs of the genus *Callinectes* in St. Johns River, Florida. *Chesapeake Sci.* **8**: 202–203.
- . 1969a. Some relations of temperature acclimation and salinity to thermal tolerances of the blue crab (*Callinectes sapidus*). *Trans. Am. Fish. Soc.* **98**: 713–716.
- . 1969b. Growth of juvenile blue crabs, *Callinectes sapidus* Rathbun, in the St. Johns River, Florida. *Fishery Bull. Fish Wildl. Serv. U.S.* **67**: 281–288.
- . 1971. Osmoregulatory ability of blue crabs in different temperature-salinity combinations. *Chesapeake Sci.* **12**: 14–17.
- and A. B. HALL. 1971. Annotated bibliography on the fishing industry and biology of the blue crab, *Callinectes sapidus*. *NOAA Tech Rept. MMFS SSRF-640*, 94 pp.
- TAN, E. C. and W. A. van ENGEL. 1966. Osmoregulation in the adult blue crab, *Callinectes sapidus* Rathbun. *Chesapeake Sci.* **7**: 30–35.
- TASHIAN, R. E. and F. J. VERNBERG. 1958. The specific distinctness of the fiddler crabs *Uca pugnax* (Smith) and *Uca rapax* (Smith) at their zone of overlap in northeastern Florida. *Zoologica.* **43**: 89–92.
- TEAL, J. M. 1958. Distribution of fiddler crabs in Georgia salt marshes. *Ecology.* **39**: 185–193.
- . 1959. Respiration of crabs in Georgia salt marshes and its relation to their ecology. *Physiol. Zool.* **32**: 1–14.
- TEERLING, J. 1970. The incidence of the ghost crab *Ocypode quadrata* on the forebeach of Padre Island and some of its responses to man. Master's Thesis, Texas A&I Univ., Kingsville. 71 pp.
- TEYTAUD, A. R. 1971. Laboratory studies of sex recognition in the blue crab *Callinectes sapidus* Rathbun. *Sea Grant Tech. Bull.* **15**, Univ. Miami.
- TOWLES, D. W., G. E. PALMER and J. L. HARRIS. 1976. Role of gill Na⁺ + K⁺-dependent ATPase in acclimation of blue crabs (*Callinectes sapidus*) to low salinity. *J. exp. Zool.* **196**: 315–321.
- TRENT, L., E. J. PULLEN and R. PROCTOR. 1976. Abundance of macrocrustaceans in a natural marsh and a marsh altered by dredging, bulkheading, and filling. *Fish. Bull.* **74**: 195–200.

- TRUITT, R. V. 1939. The blue crab. In: Our Water Resources and their Conservation. *Contr. Chesapeake Biol. Lab.* **27**: 10-38.
- TÜRKAY, M. 1968. Dekapoden von den Margarita-Inseln (Venezuela), *Senckenberg. Biol.* **49**: 249-257.
- . 1970. Die Gecarcinidae Amerikas. Mit einem Anhang über *Ucides* Rathbun. *Senckenberg. Biol.* **51**: 333-354.
- . 1971. Die Portunidae des Naturhistorischen Museums Genf, mit einem Anhang über die Typen von *Ovalipes ocellatus floridanus* Hay and Shore, 1918 (Crustacea, Decapoda). *Archs. Sci., Genève.* **24**: 111-143.
- . 1973. Bemerkungen zu einigen Landkrabben (Crustacea, Decapoda). *Bull. Mus. natn. Hist. Nat., Paris, Zool.* **106**: 969-979.
- . 1975. Zur Kenntnis der Gattung *Euchirograpsus* mit Bemerkungen zu *Brachygrapsus* und *Litocheira* (Crustacea: Decapoda). *Senckenberg. Biol.* **56**: 103-132.
- TYLER, A. V. and D. G. CARGO. 1963. Size relations of two instars of the blue crab, *Callinectes sapidus*. *Chesapeake Sci.* **4**: 52-54.
- VERNBERG, F. J. 1956. Study of the oxygen consumption of excised tissues of certain marine decapod Crustacea in relation to habitat. *Physiol. Zool.* **29**: 227-234.
- . 1959. Studies on the physiological variation between tropical and temperate zone fiddler crabs of the genus *Uca*, II. Oxygen consumption of whole organisms. *Biol. Bull.* **117**: 163-184.
- and J. D. COSTLOW, JR. 1966. Handedness in fiddler crabs. *Crustaceana.* **11**: 61-64.
- VERNBERG, W. B., P. J. DECOURSEY and W. J. PADGETT. 1973. Synergistic effects of environmental variables on larvae of *Uca pugilator*. *Mar. Biol.* **22**: 307-312.
- and F. J. VERNBERG. 1968. Physiological diversity in metabolism in marine and terrestrial Crustacea. *Am. Zoologist.* **8**: 449-458.
- and ———. 1972. The synergistic effects of temperature, salinity and mercury on survival and metabolism of the adult fiddler crab, *Uca pugilator*. *Fishery Bull. Fish Wildl. Serv. U.S.* **70**: 415-420.
- VERRILL, A. E. 1908. Brachyura and Anomura. Their distribution, variations, and habits. Decapod Crustacea of Bermuda I. *Trans. Conn. Acad. Arts Sci.* **13**: 299-474.
- VERWAY, J. 1957. A plea for the study of temperature influence on osmotic regulation. *L'Annee Biol.* **33**: 129-149.
- WARNER, G. F. 1967. The life history of the mangrove tree crab, *Aratus pisoni*. *J. Zool., London.* **153**: 321-335.
- . 1968. The larval development of the mangrove tree crab, *Aratus pisonii* (H. Milne Edwards), reared in the laboratory (Brachyura, Grapsidae). *Crustaceana, Suppl.* **2**: 249-258.
- . 1969. The occurrence and distribution of crabs in a Jamaican mangrove swamp. *J. Anim. Ecol.* **38**: 379-389.
- . 1970. Behaviour of two species of grapsid crab during intraspecific encounters. *Behaviour.* **36**: 9-19.
- WASS, M. L. 1955. The decapod crustaceans of Alligator Harbor and adjacent inshore areas of northwestern Florida. *Q. Jl. Fla. Acad. Sci.* **18**: 129-176.
- . 1968. A new pinnixed commensal with a holothurian (Crustacea: Decapoda). *Tulane Stud. Zool.* **14**: 137-139.
- WEBB, H. M., M. F. BENNETT and F. A. BROWN, JR. 1954. A persistent diurnal rhythm of chromatophoric response in eyestalkless *Uca pugilator*. *Biol. Bull.* **106**: 371-377.

- WEILAND, A. L. and C. P. MANGUM. 1975. The influence of environmental salinity on hemocyanin function in the blue crab, *Callinectes sapidus*. *J. exp. Zool.* **193**: 265-274.
- WEIS, J. S. 1976a. Effects of environmental factors on regeneration and molting in fiddler crabs. *Biol. Bull.* **150**: 152-162.
- . 1976b. Effects of mercury, cadmium, and lead salts on regeneration and ecdysis in the fiddler crab, *Uca pugilator*. *Fishery Bull. Fish Wildl. Serv. U.S.* **74**: 464-467.
- . 1976c. Regeneration in the fiddler crab, *Uca pugilator*, after injury or removal of limb buds. *J. exp. Zool.* **197**: 21-30.
- . 1977a. Limb regeneration in fiddler crabs: species differences and effects of methylmercury. *Biol. Bull.* **152**: 263-274.
- . 1977b. Regeneration of limbs autotomized at different times in the fiddler crab, *Uca pugilator*. *Can. J. Zool.* **55**: 656-660.
- and L. H. MANTEL. 1976. DDT as an accelerator of limb regeneration and molting in the fiddler crabs. *Estuarine Coastal mar. Sci.* **4**: 461-466.
- WEITZMAN, M. C. 1963. The biology of the tropical land crab, *Gecarcinus lateralis* (Fremerville). Ph.D. thesis, Albert Einstein College of Medicine, Yeshiva University.
- WEITZMAN, M. 1969. Ultrastructural study on the release of neurosecretory material from the sinus gland of the land crab, *Gecarcinus lateralis*. *Z. Zellforsch. Mikrosk. Anat.* **94**: 147-154.
- WELLS, H. and M. WELLS. 1961. Observations on *Pinnaxodes floridensis*, a new species of pinnotherid crustacean commensal in holothurians. *Bull. mar. Sci. Gulf Caribb.* **11**: 267-279.
- WHITE, A. Q. and C. P. SPIRITO. 1973. Anatomy and physiology of the swimming leg musculature in the blue crab, *Callinectes sapidus*. *Mar. Behav. Physiol.* **2**: 141-153.
- WHITING, N. H. 1972. The effect of substrate on the distribution of *Uca minax* (Crustacea: Decapoda). M. S. Thesis, Univ. West Florida, Pensacola.
- and G. A. MOSHIRI. 1974. Certain organism-substrate relationships affecting the distribution of *Uca minax*. *Hydrobiologia.* **44**: 481-493.
- WHITNEY, J. O. 1969. Absence of sterol synthesis in larvae of the mul crab *Rhithropanopeus harrisi* and of the spider crab *Libinia emarginata*. *Mar. Biol.* **3**: 134-135.
- WHITTEN, H. L., H. F. ROSENE and J. W. HEDGPETH. 1950. The invertebrate fauna of Texas coast jetties: a preliminary survey. *Publ. Inst. mar. Sci. Univ. Tex.* **1**: 53-87.
- WIGLEY, R. L. and J. L. MESSERSMITH. 1976. *Benthochascon schmitti* Rathbun (Decapoda, Brachyura) off southern New England. *Crustaceana.* **31**: 111-112.
- , R. B. THEROUX and H. E. MURRAY. 1975. Deep-sea red crab, *Geryon quinquegens*, survey off northeastern United States. *Mar. Fish. Rev.* **37**: 1-21.
- WILKINS, J. L. and M. FINGERMAN. 1965. Heat tolerance and temperature relationships of the fiddler crab, *Uca pugilator*, with reference to body coloration. *Biol. Bull.* **128**: 133-141.
- WILLIAMS, A. B. 1962. A re-examination of *Ovalipes* sp. in the Carolinas (Decapoda, Portunidae). *Crustaceana.* **4**: 39-41.
- . 1965. Marine decapod crustaceans of the Carolinas. *Fish. Bull.* **65**: 1-298.
- , 1966. The Western Atlantic swimming crabs *Callinectes ornatus*, *C. danae*, and a new, related species (Decapoda, Portunidae). *Tulane Stud. Zool.* **13**: 83-93.
- . 1974a. The swimming crabs of the genus *Callinectes* (Decapoda: Portunidae). *Fish. Bull.* **72**: 685-798.
- . 1974b. A new species of *Hypsophrys* (Decapoda: Homolidae) from the Straits of Florida, with notes on related crabs. *Proc. Biol. Soc. Wash.* **87**: 485-492.

- . 1976. Distinction between a Gulf of Mexico and a Carolinian Atlantic species of the swimming crab *Ovalipes* (Decapoda: Portunidae). *Proc. Biol. Soc. Wash.* **89**: 205-214.
- , L. R. McCLOSKEY and I. E. GRAY. 1968. New records of brachyuran decapod crustaceans from the continental shelf off North Carolina, U.S.A. *Crustaceana*. **15**: 41-66.
- and H. J. PORTER. 1964. An unusually large turtle barnacle (*Chelonibia p. patula*) on a blue crab from Delaware Bay. *Chesapeake Sci.* **5**: 150-151.
- WOLFF, T. 1954. Occurrence of two East American species of crabs in European waters. *Nature*. **174**: 188-189.
- WRIGHT, H. O. 1966. Comparative studies of social behavior in grapsoid crabs. Ph.D. Dissertation, Univ. Calif., Berkeley. 227 pp. (*Dissert. Abstr.* **27**: 4184-B).
- . 1968. Visual displays in brachyuran crabs: field and laboratory studies. *Am. Zoologist*. **8**: 655-665.
- YAMAOKA, L. H. and D. M. SKINNER. 1976. Free amino acid pools in muscle and hemolymph during the molt cycle of the land crab, *Gecarcinus lateralis*. *Comp. Biochem. Physiol.* **55A**: 129-134.
- YANG, W. T. 1967. A study of zoeal, megalopal, and early crab stages of some oxyrhynchous crabs (Crustacea, Decapoda). Doctoral Dissert. Univ. Miami, Coral Gables, Florida.
- . 1968. The zoeae, megalopa and first crab of *Epiplatys dilatatus* (Brachyura, Majidae) reared in the laboratory. *Crustaceana, Suppl.* **2**: 181-202.
- . 1976. Studies on the western Atlantic arrow crab genus *Stenorhynchus* (Decapoda Brachyura, Majidae). 1. Larval characters of two species and comparison with other larvae of Inachinae. *Crustaceana*. **31**: 157-177.
- ZELENY, C. 1908. Some internal factors concerned with the regeneration of the Gulf-weed crab (*Portunus sayi*). *Pap. Tortugas Lab.* **2**: 103-138.

INDEX

- Acanthocarpus* 29
 A. alexandri 29
 A. bispinosus 30
 Acanthonychinae 41
Acanthonyx 41
 A. petiverii 41
Actaea 87, 104
 A. acantha 87
 A. bifrons 88
 A. palmeri 88
 A. rufopunctata nodosa 88, 103
 A. setigera 88, 107
Acteomorpha 33
Aepinus 43
 A. septemspinus 43
Aethra 33
Aelhrinae 33
Anasimus 43
 A. latus 43
Anomalothir 43
 A. frontalis 43
 A. furcillatus 43
Arachnopsis 44
 A. filipes 44
 araña del mar 49
Aratus 132
 A. pisonii 132
Arenaeus 74, 82
 A. cribrarius 74
 arrow crab 49
 assemblages, faunal 13–14
 ATELECYCLIDAE 9, 11–13, 71, 72
Atelecyclinae 71

Bathynectes 72
 B. superba 72
Bathyplax 111, 112
 B. typhla 112
Batrachonotus 44
 B. fragosus 44
 beach crab 135
Benthochascon 73
 B. schmitti 73
 black crab 140
 black land crab 139
 blue crab 78–81, 96
 blue land crab 140
Boboruca 149
 box crab 30
 BRACHYRHYNCHA 8, 9, 72
 brackish-water crab 135
 brackish-water fiddler crab 143

Cabouca 58
Calappa 30
 C. angusta 30
 C. flammea 30, 31
 C. gallus 31
 C. ocellata 31
 C. springeri 32
 C. sulcata 32
 CALAPPIDAE 9, 11–13, 29
 Calappinae 29
 CALAPPOIDEA 9, 29
 Calico crab 33, 90
Callidactylus 37
 C. asper 37
Callinectes 18, 75, 82
 C. bocourti 75
 C. danae 75, 76, 77, 81
 C. exasperatus 76
 C. marginatus 77
 C. ornatus 75, 77–78, 81
 C. rathbunae 75, 78
 C. sapidus 75, 78–81, 96
 C. sapidus acutidens 78, 79
 C. similis 75, 76, 77, 78, 81
Cancer 71
 C. borealis 71
 C. irroratus 72
 CANCRIDAE 9, 11–13, 71, 72
 CANCRIDEA 9, 71, 72
 Cancrinae 71
 cancrivora crabs 72
 CANCROIDAE 9, 71
 cangrejo de la Santa Virgen 58
 Carcinoplacinae 111
Cardisoma 138, 150
 C. guanhumi 138
 Carolinean Province 11, 13–14, 73
Carpilius 88
 C. convexus 88
 C. corallinus 88
Carpoporus 89
 C. papulosus 89
Cataleptodius 89
 C. floridanus 89, 95

 Catametopa 72
Celuca 143, 145, 148
Chacellus 112
 C. filiformis 112
Chasmocarcinus 112
 C. cylindricus 112
 C. mississippiensis 112
 C. obliquus 113
Chlorodiella 89
 C. longimana 89
Chorinus 62
 C. heros 62
 cliff crab 129, 131
Clythrocerus 26
 C. nitidus 26
 C. stimpsoni 26
Coelocerus 50
 C. spinosus 50
Collodes 44
 C. armatus 44
 C. leptocheles 44
 C. trispinosus 45
 common edible crab 78
 common land crab 139
 common mud crab 102
 common pea crab 124
 common spider crab 64
 coral crab 56, 88
Corycodus 26
 C. bullatus 26
 Corystidae 9
Cronius 81
 C. ruber 81
 C. tumidulus 82
Cryptochirus 40
 C. corallicola 40
Cryptopodia 67
 C. concava 67
Cyclodorippe 26
 C. antennaria 26
 C. bowieri 26
 C. ornata 27
Cycloes 32
 C. bairdii 32
Cyclograpsus 132
 C. integer 132
 Cyclometopa 72
Cycloxanthops 110
Cymonomus 27
 C. caecus 27

- C. cubensis* 27
C. quadratus 27
C. rostratus 27
Cymopolia 27
C. affinis 118
C. alternata 118
C. cursor 118
C. dentata 118
C. faxoni 118
C. gracilipes 118
C. gracilis 119
C. obesa 119
C. sica 119
 CYMOPOLIDAE 117
Cymopolus 27
C. agassizi 27
- decorator crab 52, 58
 deep sea red crab 111
Dicranodromia 21
D. ovata 21
 dirty decorator crab 54
 disjunct distribution 14, 18
Dissodactylus 119
D. alcocki 119
D. borradalei 120
D. calmani 120
D. crinitichelis 120
D. encopei 120
D. juvenilis 120
D. mellitae 120
D. primitivus 121
D. stebbingi 121
 Dolly Varden crab 33
Domecia 89
D. acanthophora acanthophora 89–90
D. hispida 89–90
 DORIPPIDAE 9, 11–13, 26
 DORIPPOIDEA 9, 26
Dromia 19
D. erythropus 19
 DROMIACEA 9, 19, 22
Dromidia 19
D. antillensis 19
 DROMIIDAE 9, 11–13, 19
 DROMIOIDEA 9, 19
 Dynomenidae 9, 10
- Ebalia* 35
E. cariosa 35
E. simpsoni 35
 Ebalinae 35
- edible crab 78
 endemic species 11, 14
Epialtus 41
E. bituberculatus 41
E. dilatatus 42
E. dilatatus forma elongata 42
E. longirostris 42
Eriphia 90
E. gonagra 90
Ethusa 28
E. lata 28
E. mascarone americana 28
E. microphthalmia 28
E. tenuipes 28
E. truncata 29
Ethusina 29
E. abyssicola 29
E. favonii 29
Etisus 90, 104
E. maculatus 90
Euchirograpsus 136
E. americanus 136, 137
E. antillensis 137
E. pacificus 137
Eucratodes 91
E. agassizii 91
Eucratopsis 113
E. crassimanus 113
Euphosynoplax 113
E. clausa 113
Euprognatha 45
E. gracilipes 45
E. rastellifera acuta 45
E. rastellifera marthae 45
Eurypanopeus 91, 103
E. abbreviatus abbreviatus 91
E. abbreviatus ater 91
E. crenatus 91
E. depressus 91
E. dissimilis 92
E. turgidus 92, 103
Euryplax 113
E. nitida 113
Eurytium 92
E. limosum 92
- Fabia* 121
F. byssomiae 121
F. tellinae 121
 fiddler crab (see *Uca*)
 flame-streaked box crab 30
 freshwater crab 87
- Frevillea* 114
F. barbata 114
F. hirsuta 114
F. tridentata 114, 116, 117
 friendly crab 133
 frog crab 25
- GECARCINIDAE 9, 11–13, 138
Gecarcinus 139
G. lateralis 139, 140
G. quadratus 139, 140
G. ruricola 140
Geograpsus 128
G. lividus 128
Geryon 111, 112
G. affinis 111
G. quinquedens 111
 GERYONIDAE 9, 11–13, 110, 112
 ghost crab 141
Glyptoplax 114
G. pugnax 114
G. smithii 114
Glyptoxanthus 93
G. erosus 93
 GONEPLACIDAE 9, 11–13, 72, 100, 110, 111
Goneplax 114
G. barbata 114
G. hirsuta 114
G. tridentata 117
Goniopsis 128
G. cruentata 128
 GRAPSIDAE 9, 72, 128
 Grapsinae 128
 grapsoid crabs 72
Grapsus 129
G. grapsus 129
 grass crab 52
 great land crab 138
 guaiamu 138
 guanhumí 138
Gymnopleura 23
- hairy crab 107
 HAPALOCARCINIDAE 8, 9, 11–13, 40
 HAPALOCARCINIDEA 9, 40
 HAPALOCARCINOIDEA 9, 40
Hemus 50
H. cristulipes 50
Hepatus 32, 33
H. epheliticus 33

- H. princeps* 33, 34
H. pudibundus 33, 34
Heteractaea 93
H. ceratopus 93
Heterocrypta 67
H. granulata 67
Hexapanopeus 93, 96
H. angustifrons 93
H. hemphillii 94
H. lobipes 94, 96
H. paulensis 94, 95
H. quinquedentatus 94
H. sinaloensis 95
Holometopus 133
Holoplites 63
H. armata 63
Homola 22
H. barbata 22
H. vigil 22
HOMOLIDAE 9, 10, 11–13,
21, 23
Homolodromia 21
H. paradoxa 21
Homolodromiinae 10, 21
Homologenus 22
H. rostratus 22
HOMOLOIDEA 9, 21
Hypoconcha 20
H. arcuata 20
H. sabulosa 20
H. spinosissima 21
Hypsophrys 22–23
H. noar 22

Iliacantha 37
I. intermedia 37
I. liodactylus 37
I. sparsa 38
I. subglobosa 38
Inachinae 42
Inachoides 45
I. forceps 45, 46
I. laevis 45, 46
isozyme patterns 18

jonah crab 71
juey 138

kaburi 150

lady crab 74
land crab 138, 139, 140, 150
Latreillia 10, 23

L. elegans 23
LATREILLIDAE 10, 11–13,
23
Latreillopsis 23
lazy crab 58
Leiolambrus 68
L. nitidus 68
LEUCOSIIDAE 9, 11–13, 33,
35
Leucosiinae 36
leopard crab 33
Leptodius 95
L. agassizzi 95, 108
L. floridanus 89, 95
L. parvulus 95
Leptuca 143, 148
lesser blue crab 81
Iibinia 50, 63
L. dubia 63, 64, 65
L. emarginata 63, 64
L. erinacea 65
L. rhomboidea 65
Lithadia 35
L. cadaverosa 35
Lobopilumnus 95
L. agassizii 95
loop currents 15
Lophopanopeus 94, 96
L. distinctus 96, 98
L. lobipes 94, 96
Lupella 82
L. forceps 82
Lyreidus 23
L. bairdii 23

macca crab 58
Macrocoeloma 50
M. camptocerum 50
M. diplacanthum 51
M. eutheca 51
M. intermedium 51
M. laevigatum 51
M. septemspinatum 52
M. subparallelum 52
M. trispinosum trispinosum
52
M. trispinosum nodipes 52, 53
M. trispinosum variety
Rathbun 53
Macrocoelominae 50
Macropipinae 72
Macropipus 73
MAJIDAE 9, 11–13, 41
Majinae 49, 50
mangrove crab 128, 132, 136
marsh crab 132, 136
Matuta 33
Matutinae 32, 33
Melybia 96
M. thalamera 96
Menippe 96
M. mercenaria 93, 96–97
M. nodifrons 97
Mesorhoea 68
M. sexspinosa 68
Metasesarma rubripes 134
Metoporphaphis 46
M. calcarata 46
Micropanope 96, 97
M. barbadosis 97
M. distincta 96, 98
M. lobifrons 98
M. nuttingi 98
M. pusilla 98
M. sculptipes 97, 98, 99
M. spinipes 99
M. truncatifrons 99
M. urinator 99
M. xanthiformis 99, 100
Microphrys 53
M. antillensis 53
M. bicornutus 54
M. platysoma 53, 54
Mithracinae 50
Mithraculus 54
Mithrax 54, 56
M. (Mithraculus) cincti-
manus 54
M. (Mithraculus) coryphe 55
M. (Mithraculus) forceps
55, 57
M. (Mithraculus) ruber 55
M. (Mithraculus) sculptus
56
M. (Mithrax) acuticornis 56
M. (Mithrax) cornutus 56
M. (Mithrax) depressus 57
M. (Mithrax) hispidus 56
M. (Mithrax) holderi 57
M. (Mithrax) pilosus 57
M. (Mithrax) pleuracanthus
57
M. (Mithrax) spinosissimus
56, 58
M. (Mithrax) verrucosus 58
Mocosoa 42

- M. crebripunctata* 42
 mottled shore crab 130
 mountain crab 140
 mud crab 102
 mulatto land crab 138
 mussel crab 123
Myropsis 38
 M. quinquespinosa 38
- Nanocassiope* 99
 N. melanodactylus 99
Nanoplax 99, 100
 N. xanthiformis 99, 100
Neodoclea 50
Neopanope 100
 N. packardii 100, 101
 N. sayi 100, 101
 N. texana 100
 N. texana sayi 101
Neopilumnoplax 114, 115
 N. americana 114, 115
Nibilia 65
 N. antilocapra 65
 northern crab 71
Notosceles chummonis 25
- Ocypode* 140
 O. albicans 140
 O. arenaria 141
 O. quadrata 140, 141
 OCYPODIDAE 9, 10, 11–13, 140
 OCYPODOIDEA 9, 140
 Ocypodinae 140
 Ophthalmiinae 60
Orthotheres 122, 125
 O. serrei 122, 125
 O. strombi 122, 125
Osachila 32, 33, 34
 O. antillensis 34
 O. semilevis 34
 O. tuberosa 34
Ovalipes 13, 18, 72, 73
 O. floridanus 74
 O. guadulpensis 73, 74
 O. ocellatus 73
 O. stephensoni 74
 OXYRHYNCHA 9, 41
 OXYSTOMATA 8, 9, 23
 oyster crab 124
- Pachygrapsus* 130
 P. gracilis 130
- P. transversus* 130
 pagurus 150
 PALICIDAE 8, 9, 11–13, 27, 72, 117
Palicus 27, 117
 P. affinis 117
 P. alternatus 118
 P. cursor 118
 P. dentatus 118
 P. faxoni 118
 P. gracilipes 118
 P. gracilis 119
 P. obesus 119
 P. sica 119
Panopeus 101
 P. americanus 101
 P. bermudensis 101
 P. hartii 101
 P. herbstii 93, 102
 P. occidentalis 103
 P. rugosus 103
 P. turgidus 92, 103
Panoplax 115
 P. depressa 115
Paractaea 88, 103
 P. rufopunctata nodosa 88, 103
Paraliomera 104
 P. dispar 104
 P. longimana 104
Parapinnixa 122
 P. bouvieri 122
 P. hendersoni 122
 parrot crab 32
Parthenope 68
 P. (Parthenope) agonus 68
 (Platylambrus) fraterculus 68
 P. (Platylambrus) pourtalesii 69
 P. (Platylambrus) serrata 69
 PARTHENOPIDAE 9, 11–13, 33, 67
 Parthenopinae 67
 pea crab 123, 124
Pelia 65
 P. mutica 65
 pentagon crab 67
Percnon 131
 P. gibbesi 131
 Periceroida 50
Persephona 39
 P. aquilonaris 39
 P. crinita 39
 P. mediterranea 37, 39
 P. punctata 39
 P. punctata aquilonaris 39, 40
 Philyrinae 38
Phyfodius 90, 104
 P. maculatus 90, 104
Picroceroides 60
 P. tubularis 60
Pilumnoides 104
 P. nudifrons 104
Pilumnoplax 115
 P. americana 114, 115
 P. elata 115, 116, 117
 P. nitida 115
Pilumnus 105
 P. caribaeus 105
 P. dasypodus 105
 P. diomedea 105
 P. floridanus 105
 P. gemmatus 106
 P. holosericus 106
 P. lacteus 106
 P. longleyi 106
 P. marshi 106, 107
 P. pannosus 106
 P. sayi 106, 107
 P. spinosissimus 107
Pinnaxodes 122
 P. floridensis 122
Pinnixa 125
 P. chacei 125
 P. chaopterana 125
 P. cristata 126
 P. cylindrica 126, 128
 P. floridana 126, 127
 P. leptosynaptae 127
 P. lunzi 127
 P. pearsei 127
 P. retinens 127
 P. sayana 128
 Pinnothereliinae 125
Pinnotheres 123
 P. geddesi 123
 P. guerini 123
 P. hemphilli 123
 P. hirtimanus 123
 P. maculatus 123
 P. mosei 124
 P. ostreum 124
 P. serrei 122, 125
 P. shoemakeri 125
 P. strombi 122, 125

- PINNOTHERIDAE 9, 10, 11–13, 119
 Pinnotherinae 119
 Pisinae 50, 62
Pitho 61
P. aculeata 61
P. anisodon 61
P. laevigata 61
P. lherminieri 61
P. mirabilis 62
 Plagusiinae 131
Plagusia 131
P. depressa 131
Planes 130
P. cyaneus 130–131
P. minutus 130, 131
Planuca 145, 149
Platyactaea 88, 107
P. setigera 88, 107
Platychirograpsus 137
P. spectabilis 137
P. typicus 137
Platylambrus 68
Platypodia 108
P. spectabilis 108
Platypodiella 108
P. spectabilis 108
Podocheila 46
P. curvirostris 46
P. gracilipes 47
P. lamelligera 47
P. macrodera 47
P. riisei 47
P. sidneyi 48
 Polybiinae 72
 PORTUNIDAE 9, 11–13, 72
 Portuninae 74
 PORTUNOIDEA 9, 72
Portunus 82
P. anceps 82
P. binoculus 83
P. depressifrons 83
P. floridanus 83
P. gibbesii 83, 86
P. ordwayi 84
P. sayi 77, 84
P. sebae 85
P. spinicarpus 83, 85
P. spinimanus 84, 85
P. ventralis 86
P. vocans 86
 POTAMIDAE 8, 11–13, 72, 86
Potamocarcinus 87
Potamon 87
 Potamonidae 87
Prinoplax 116
P. atlantica 116, 117
 PROSOPIDAE 8, 9, 11–13, 21
Pseudomedeaeus 108
P. agassizii 95, 108
 Pseudothelphusidae 8, 87
 purse crab 39
Pyromaia 48
P. arachna 48
P. cuspidata 48
 queen crab 88
 racing crab 141
Ranilia 24
R. constricta 24
R. muricata 24
 RANINIDAE 9, 11–13, 23
 RANINOIDEA 9, 23
Raninoides 24
R. benedicti 25
R. fossor 24
R. lamarcki 24
R. loevis 25
R. louisianensis 25
 red crab, deep-sea 111
 red-jointed fiddler crab 143
 red land crab 140
 red spider crab 56
 red tourlourou 140
 Retroplumidae 9, 10
Rhithropanopeus 108
R. harrisii 108
R. harrissi tridentatus 109
 river crab 137
Robertella 116
R. mystica 115, 116, 117
Rochinia 66
R. crassa 66
R. hystrix 66
R. tanneri 66
R. umbonata 67
 rock crab 129
 saber crab 137
 sally lightfoot 129
 sand crab 74, 141
 sand fiddler crab 144, 145
 sargassum crab 84
Sesarma 18, 133
S. (Holometopus) ameri-
canum 133, 134, 135
S. (Holometopus) angustipes
 133, 134, 135
S. (Holometopus) benedicti
 133
S. (Holometopus) miersii 134
S. (Holometopus) cinereum
 18, 133–134, 135
S. (Holometopus) miersii 134
S. (Holometopus) ricordi 18,
 134, 135
S. (Holometopus) roberti
 133, 134, 135
S. (Holometopus) tampi-
cense 133, 135
S. (Sesarma) curacaoense 136
S. (Sesarma) reticulatum
 18, 136
 Sesarminae 132
 shame-faced crab 30, 32
 shore crab 130
Solenolambrus 70
S. decemspinus 70
S. tenellus 70
S. typicus 70
 speckled crab 74
Speloeophorus 36
S. elevatus 36
S. nodosus 36
S. pontifer 36
Speocarcimus 116
S. carolinensis 116
S. lobatus 116
 spider crab (see MAJIDAE)
 sponge crab 52
 spray crab 131
 square-backed fiddler crab 133
Stenocionops 50, 58
S. furcata coelata 59
S. furcata furcata 58
S. spinimana 59
S. spinosissima 59
Stenorhynchus 49
S. lanceolatus 49
S. seticornis 49
 stone crab 96
 swimming crab (see
 PORTUNIDAE)
Symethis 25
S. variolosa 25
Telephrys 60
T. ornatus 60
Temnonotus 49

- T. granulatus* 49–50
T. simplex 49, 50
Tetraplax 116
T. quadridentata 116
Tetrazanthus 109
T. bidentatus 109, 110
T. rugosus 109, 110
T. rathbunae 109
Thalassoplex 117
T. angusta 115, 117
Thelxiope 22
T. barbata 22
T. vigil 22
Thoe 60
T. puella 60
Thyrolambrus 70
T. astroides 70
 tourlourou 138
Trachycarcinus 71
T. spinulifer 71
Trapezioplax 114, 116, 117
T. tridentata 114, 116, 117
 tree crab 128, 132
 Trichodactylidae 8, 87
Trichopeltarion 71
T. nobile 71
Troglocarcinus 41
T. balssi 41
T. corallicola 41
Tutankhamen 71
T. cristatipes 71
Tyche 62
T. emarginata 62
T. lamellifrons 62
Typhlopseudothelphusa 87
Uca 10, 13, 18, 142, 150
U. affinis 142
U. burgersi 142, 144
U. cumulanta 148
U. leptodactyla 143
U. longisignalis 18, 143, 147, 149
U. minax 143, 144, 147
U. mordax 142, 144, 149
U. murifecenta 149
U. panacea 18, 144, 145
U. pugilator 18, 144, 145–146
U. pugnax 18, 143, 147, 149
U. pugnax repax 143, 144, 147
U. pugnax virens 144, 149
U. rapax 142, 143, 147, 149
U. rapax longisignalis 143, 144, 147
U. speciosa 148
U. speciosa spinicarpa 148
U. spinicarpa 148
U. subcylindrica 18, 148
U. thayeri 148
U. virens 18, 143, 147, 149
U. vocator 144, 149
Ucides 150
U. cordatus 150
U. occidentalis 150
Uhlias 36
U. limbatus 36
 Varuninae 136
 Virginian Province 11
 wharf crab 130, 133
 white land crab 138
 whitespot crab 140
 wood crab 133
 XANTHIDAE 9, 10, 11–13, 72, 87, 111, 112
Xantho 110
X. denticulata 110
Xanthodes bidentatus 110
Xanthodius 110
X. denticulatus 110
X. simpsoni 110
 XANTHOIDEA 8, 9, 72, 86
 yellow box crab 31, 32