

**Supporting Information.** Anthropogenic versus fish-derived nutrient effects on seagrass community structure and function. Jacob E. Allgeier, Craig A. Layman, Carmen G. Montana, Enie Hensel, Richard Appaldo, and Amy D. Rosemond. *Ecology*. 2018.

## Appendix S4

**Table S1** – List of all species found in benthic surveys of seagrass beds presented in descending order of mean abundance across surveys.

**Table S1**

<b>Producer Community</b>
<i>Thalassia testudinum</i>
<i>Syringodium filiforme</i>
<i>Batophora oerstedii</i>
<i>Avrainvillea sp.</i>
<i>Penicillis capitatus</i>
<i>Rhipocephalus phoenix</i>
<i>Udotea flabellum</i>
<i>Halimeda incrassata</i>
<i>Heterosiphonia gibbesii</i>
<i>Cyanobacterial mat</i>
<i>Laurencia sp.</i>
<i>Dictyosphaeria cavernosa</i>
<i>Acetabularia crenulata</i>
<i>Valonia ventricosa</i>
<i>Dictyopteris sp.</i>
<i>Halodule wrightii</i>
<i>Styopodium sp.</i>
<i>Cladosiphon occidentalis</i>
<i>Ernodesmis verticillata</i>
<i>Sargassum sp.</i>
<i>Padina sp.</i>