
The Cambrian System in Northwestern Argentina: stratigraphical and palaeontological framework

G. F. ACEÑOLAZA

INSUGEO – CONICET. Facultad de Ciencias Naturales e I.M.L., Universidad Nacional de Tucumán
Miguel Lillo 205, 4000 Tucumán, Argentina. E-mail: insugeo@unt.edu.ar

ABSTRACT

Cambrian sequences are widespread in the early Paleozoic of the Central Andean Basin. Siliciclastic sediments dominate these sequences although several minor occurrences of carbonates and volcanic rocks have been observed. The rocks assigned to the Cambrian System in NW Argentina are recognized in the Puna, Eastern Cordillera, Subandean Ranges and the Famatina System. This paper gives a general overview of the Cambrian formations outcropping in the northern provinces of Jujuy, Salta, Tucumán, Catamarca and La Rioja. Special emphasis has been given to the stratigraphical and biostratigraphical framework of the sequences. Late Precambrian-Early Cambrian thick sedimentary wackes dominate the basal Puncoviscana Formation (s.l.), characterized by a varied ichnofauna that includes the Precambrian-Cambrian transitional levels. Thick packages of sandstone facies characterize the overlying highly ichnofossiliferous Mesón Group. A rich shelly fauna included in sandstones, shales and volcanoclastics, dominate in the Cambrian-Ordovician transitional levels of the overlying Cambro-Ordovician Santa Victoria and Cachiuyuyo Groups. Palaeogeographic aspects dealing with the position and development of Cambrian basins are discussed. Several synthetic biostratigraphic tables are presented displaying the known record for the Cambrian System, including the transitional levels to the lowermost Ordovician in the above-mentioned regions.

KEYWORDS | Cambrian. Northwestern Argentina. Lithostratigraphy. Biostratigraphy. Palaeogeography.

INTRODUCTION

Widely distributed sedimentary rocks assigned to the Cambrian System are recorded in the early Paleozoic of the Central Andean Basin, cropping out from Southern Peru to northern Argentina. Thick sequences of over 7000 m are recognized in Jujuy, Salta, Tucumán, Catamarca and La Rioja provinces (Northern Argentina, Fig. 1). They were sedimented in large elongated marginal basins in the western border of Gondwana, connected to the north to other Bolivian, Peruvian, Paraguayan and Brazilian sedimentary systems.

The Cambrian sequences in northwestern Argentina display three clear depositional stages. The earlier one is recorded by the slates of the Puncoviscana Formation and equivalent units. The second stage resulted in the sedi-

mentation of the Mesón Group (only represented in the Eastern Cordillera, Puna and Subandean Ranges). Finally, the third stage is represented by the lowermost part of the Santa Victoria (Puna, Eastern Cordillera and Subandean Ranges) and Cachiuyuyo groups (Famatina System; Aceñolaza and Peralta, 2000). The first stage was developed during the Pampean Cycle of Aceñolaza and Toselli (1976), while the two others were deposited during the Famatinian Cycle (Aceñolaza et al., 2000; Fig. 2).

The Central Andean Basin is well known by its widespread Ordovician related economic deposits and fossiliferous content. Less attention has historically been given to the Cambrian sequences, as a consequence of their relative homogeneity and remarkable scarcity of fossiliferous shelly fauna.




FIGURE 1 | Location and distribution of Cambrian stratigraphic units (Precambrian-Cambrian Puncoviscana Formation (*s.l.*), the Cambrian Mesón Group and the Cambro-Ordovician Santa Victoria Group) in northwestern Argentina and location of significant outcrop zones mentioned in the text.

The stratigraphy, biostratigraphy and fossiliferous distribution of the Cambrian sequences in northwestern Argentina, as well as some paleopalaeogeographic aspects, will be summarized in this paper.

GEOLOGICAL AND PALAEOGEOGRAPHICAL SETTING

Cambrian sequences display a large geographic distribution in northwestern Argentina, covering an area of over




FIGURE 2 | Stratigraphic sketch for the Precambrian - Mid Paleozoic sequences in NW Argentina. The Puna, Eastern Cordillera, Subandean Ranges, and the Famatina System are considered (modified from Aceñolaza et al., 2000).

400000 km². They are noticeable in many sectors of the Puna, most of the Cordillera Oriental, the Subandean Ranges and the Famatina System of Argentina, from the international boundary with Bolivia to La Rioja Province (Fig. 1). In addition, some successions, which are considered to be Cambrian in age, have been recorded to the east in well-cores of the Pampas, while to the west, an imprecise boundary is mentioned on the basis of subsurface strata (Aceñolaza and Toselli, 1981; Finney et al. in press; Bordonaro, 1992; González-Bonorino and Llambías, 1996; Sánchez and Salfity, 1999).

To the south of this area, an extended carbonate platform was developed in the Precordillera of Mendoza and San Juan, but a different geotectonic history has been given to this sector, assuming an allochthonous to parautochthonous origin for the latter (see Bordonaro, this volume; Aceñolaza and Tosselli, 2000).

The outcrops analyzed in this paper have a N-S orientation and record depositional processes which

developed on the margin of Western Gondwana. On a broad outline, the sedimentary basins developed in this region had a border to the east with a more or less continuous belt of cratonic regions, which are regarded as the source areas for the basin infill. This palaeogeographic framework lasted, with only minor changes, from the Late Proterozoic to the early Silurian (Fig. 3A,B).

For the Puncoviscana Formation (s.l.), a rather long period of crustal stability resulted in very homogeneous sedimentary characteristics within a large area. A deep slope depositional setting, associated to large coalescent submarine fans, has been proposed for the facies assemblages included in this unit (Jezek et al., 1985; Jezek, 1990). Meanwhile, the deposition of The Mesón Group and of the lower part of the Santa Victoria Group took place mostly on shore lines, strongly influenced by the eustatic variations that characterised the Cambro-Ordovician transition.

Siliciclastic sediments dominate the sequences, while carbonates are scarce and occur mainly as local limestone successions, bioclastic beds and coquinas (Sánchez, 1994, 1999; Moya, 1988, 1998; Aceñolaza, 1996; Sánchez and Salfity, 1999).

LITHOSTRATIGRAPHY OF THE PRECAMBRIAN-EARLY ORDOVICIAN RECORD

The lithostratigraphic framework of the Early Paleozoic in the Eastern Cordillera, Puna, Subandean Ranges and the Famatina System is summarized in figure 2.

The Proterozoic-Early Cambrian Puncoviscana Formation (s.l.)

The precise lower boundary of the Cambrian System is not known in the provinces of northwestern Argentina (e.g., Durand, 1996). It has been considered that the boundary is included in the highly tectonised, dominantly clastic, thick sequences which are widely represented in the region. Different stratigraphic names such as Puncoviscana *s.l.* (Jujuy, Salta and Tucumán provinces), Suncho

(Catamarca Province) and Aguaditas/Negro Peinado (Famatina System, La Rioja Province) have been used to refer to these sequences.

Lithologically, they are made up by over 2000 meters of shales, slates, sandstones and quartzites, with minor conglomerates, limestones and some interstratified basic lavas and tuffs. The sequence is highly deformed and fractured, with a general N-S alignment. Structural complexity obscures the stratigraphy of these sediments, without a single known undisturbed section. Distinctive structural styles characterize approximately parallel N-S zones within the outcrops of the Puncoviscana Formation (Mon and Hongn, 1991; Mon and Salfity, 1995).

Aceñolaza and Durand (1986) recognised the existence of a SW/NE oriented basin during the Vendian/Tommotian. This basin was developed by an expansion of the Gondwanan margin between the Río de la Plata Craton (SE) and the Arequipa Massif (NW), both of Rifean age (approximately 1000 My). The opening of the basin is interpreted to be related to a triple junction point placed in the centre of Bolivia. The southern branch of this rift corresponds to the early above-mentioned Puncoviscana basin (Fig. 3A).




FIGURE 3 | Paleogeographic sketch of the Andean margin of South America for Precambrian-Cambrian and Cambro-Ordovician times. The Puncoviscana (Precambrian-Cambrian), Mesón (Cambrian) and Santa Victoria (Cambro-Ordovician) basins are shown related to the main structural elements in a Gondwanan framework

This Early Paleozoic basin had smooth floor morphology with large, low- gradient submarine fans which spread towards the axis of the basin. Six facies have been distinguished within this fan system, ranging from proximal to distal submarine fan deposits, whose relationships are quite complex. (Aceñolaza et al., 1988; Jezek, 1990). Widespread turbidites are recorded within the sections (Baldis and Omarini, 1984; Durand and Spalletti, 1986; Spalletti and Durand, 1987; Jezek, 1990; Durand, 1990). In addition, near-shore sediments were unconformably deposited upon the cratons on the marginal sectors of the basin (Jezek et al., 1985).

The Puncoviscana sequence was intruded by Middle Cambrian (517 My) post-tectonic granites with important outcrops in Tastil and La Quesera localities (Salta Province). These episodes were related to an important deformation process, represented by the Lower Cambrian (?Atdabanian) “Tilcaric” unconformity (Turner, 1960). Jezek (1990) states that deformation affecting the Puncoviscana Formation had multiple phases: the strongest one occurred at the end of the Lower Cambrian (F1), the second during the Ordovician (F2), and lastly the third is represented by the Andean deformation (Cenozoic fracturation – F3).

Trace fossils were used as the only biostratigraphic marker to date the above- mentioned unit (Fig. 4). Among the recognizable Vendian traces are: *Nereites saltensis* ACEÑOLAZA and DURAND, *Tasmanadia* and probably *Sekwia*. Higher levels assigned to the Tommotian are characterised by: *Oldhamia radiata* FORBES, *O. flabellata* ACEÑOLAZA and DURAND, *O. antiqua* KINAHAN, *Diplichnites* and *Dimorphichnus* among others (Aceñolaza and Durand, 1986; Aceñolaza et al., 1999; Aceñolaza and Tortello, 2000; Bua-tois et al., 2000; Aceñolaza and Alonso, 2000, Aceñolaza and Tortello, this volume. See Appendix).

The Cambrian Mesón Group

After the “Tilcaric” orogenic phase, a new depositional setting developed in northwestern Argentina and southern Bolivia. Sediments associated to this basinal evolution are characterised by over 3000 m thick marine siliciclastic sequences. Outcrops of the Mesón Group are widely distributed in the northern provinces of Jujuy, Salta and Tucumán (Puna, Eastern Cordillera and Subandean Ranges; with the southernmost outcrop at about 27°S; Fig. 1). In addition, thick packages of sandstones attributed to the group have been recorded to the east in cores from the provinces of Santiago del Estero (Árbol Blanco), Chaco (Charata) and Formosa (Mariano Boedo).

The Mesón basin was surrounded by the Río de La Plata and Arequipa cratons, and by some structural highs with a general SE/NW orientation (“Calchaquí Dorsal” *sensu* Auboin et al., 1973; or “Pampean Craton” *sensu*




FIGURE 4 | Litho and biostratigraphic sketch of the Puncoviscana Formation (s.l.) with distribution of the different recognized ichnofacies (based on Durand and Aceñolaza, 1990b). See Figs. 1 and 2 for location, and Appendix.

Braccacini, 1960). The development of this basin was strongly related to the first marine flooding of the Proto-Andean border. A wide corridor that involved areas between the Arequipa Massif and the Guaporé Craton (SW margin of the Brazilian shield) seems to have been developed from the Lower to Upper Cambrian, connecting the southern Mesón Group basin northwards to other Bolivian, Peruvian and Brazilian contemporary basins (Fig. 3B).

The sequence is dominated in certain areas by thousands of meters of thick massive sandstones and quartzites (Santa Victoria Range in Salta, and Tarija region in southern Bolivia). These are correlated as a whole with thinner sequences that were deposited in some marginal sectors of the same basin. These rocks were named as Mesón Group by Turner (1960), which can be clearly split from bottom to top into three formations: Lizoite, Campanario and Chalhualmayoc (see Moya, 1998; Sánchez and Salfity, 1999) (Fig. 5).

Lizoite Formation

White, pinkish and grayish quartzites and sandstones characterize this Formation. The thickness of this unit




FIGURE 5 | Stratigraphic characterization of the Cambrian Mesón Group with its three-fold stratigraphic subdivision: Lizoite, Campanario and Chalhualmayoc formations. The relative distribution of fossils is displayed by the symbols (Brachiopoda and Trilobita) and asterisks (ichnofossils). Modified after Sánchez and Salfity, 1999. See Figs. 1 and 2 for location, and Appendix.

ranges from 20 to 1100 m. The sequence starts after the Tilcarian unconformity with a conglomerate level. The successions of this unit show a fining upwards arrangement, and have been interpreted as intertidal and subtidal sandbars and channels (Moya, 1998; Sánchez, 1999; Sánchez and Salfity, 1999).

The only biostratigraphic markers for the Lizoite Fm. were recorded to the west, in the Puna Region (Potrerillos, Salta), where Aceñolaza (1973) and Aceñolaza and Bor-

donaro (1989) recognized fragmentary and poorly preserved trilobites (*Asaphiscus* sp.) and trace fossils (*Cruziana* isp.).

Campanario Formation

Two members have been differentiated within this unit: a lower “green member”, characterized by medium grained sandstones and quartzites, and a upper “purple member” with fine grained sandstones and subordinated quartzites. This formation is 30 to 1100 meters thick. It has been interpreted as an intertidal flat sequence, which records sporadic storm events (Sánchez, 1999; Sánchez and Salfity, 1999).

Abundant trace fossils are recognised within this unit. Well defined “pipe rocks” with *Skolithos*, various types of *Rusophycus*, *Cruziana* and the Lower Cambrian *Syringomorpha nilssoni* (TORELL) are among the most remarkable ones (Ramos, 1973; Alonso and Marquillas, 1981; Aceñolaza et al., 1982; Manca, 1986; Sánchez, 1994; Moya, 1998). *Lingula* sp. and *Lingulepis* sp. are the only records of shelly fauna in the unit (Sánchez and Herrera, 1994 with references).

Chalhualmayoc Formation

The Chalhualmayoc Formation mostly consists of a fining upwards sequence of white to pinkish quartzites and subordinate sandstones with minor interbedded fine conglomerates. Abundant trace fossils and even *Skolithos* “pipe rock” are recorded in the sequence. This succession records deposition developed in relation to intertidal to subtidal sand bars, with migrating dune deposition (Sánchez, 1999; Sánchez and Salfity, 1999).

The age of the Mesón Group has been established on the basis of the scarce fossils yielded by the Chalhualmayoc Fm. It ranges between the Lower and Upper Cambrian. The Middle Cambrian trilobite *Asaphiscus* sp. has been mentioned for the basal Lizoite Formation, while with some temporal incongruity, the Lower Cambrian trace fossil *Syringomorpha nilssoni* (TORELL) was found in the middle Campanario Formation. Lastly, a single occurrence of the Cambro-Ordovician trilobite *Parabolina* (*N.*) *frequens argentina* (KAYSER) at the Azul Pampa locality, Jujuy Province is the only suitable biostratigraphic datum for the Group. The Mesón Group is overlain by the Cambro-Ordovician Santa Victoria Group.

The Cambro-Ordovician Santa Victoria Group

A few thousands of meters of Upper Cambrian to Lower Ordovician shales, sandstones, quartzites and some volcanoclastics make up this unit. The Santa Victoria Group comprises most of the sequences which overlie the




FIGURE 6 | Stratigraphic sketch and biostratigraphic framework for the Cambro-Ordovician transition in the study area (lower part of Santa Victoria Group). The stratotype for the Cambrian-Ordovician boundary at the Sierra de Cajas is shown (based in Rao and Hünicken, 1995b). See Figs. 1 and 2 for location, and Appendix.

Mesón Group and crop out in a large area of the Puna, Eastern Cordillera and Subandean Ranges (Provinces of Jujuy and Salta; Aceñolaza, 1998).

The stratigraphic relation between the Mesón and Santa Victoria Group is a matter of discussion, due to the frequent occurrence of conglomerate beds in the transitional levels between both units within the uppermost part of the underlying Mesón Group. Some authors suggest an unconformable contact for the two groups (Turner and Méndez, 1975; Moya, 1988, 1998; Sánchez and Salfity, 1999). Nevertheless, the lithology mostly does not change below and above the conglomerates and quartzites and sandstones are dominant. Therefore, in this contribution the unconformity (named as “Tilcaric”) is considered earlier, and included in the underlying Chalhualmayoc Formation (Mesón Group).

These conglomerate layers have been interpreted as a channel fill within the platform, being related to the several eustatic episodes that characterised the Cambro-Ordovi-

cian transition (Aceñolaza and Aceñolaza, 1992; Aceñolaza, 1996).

The Santa Rosita Formation is the lower unit of the Santa Victoria Group and is composed of sandstones, shales and quartzites. The stratigraphic interval that includes the Cambrian-Ordovician transition reaches up to 900 m in thickness in the Santa Victoria Range (Salta Province), and is placed in the lower part of the Santa Rosita Formation (Figs. 6, 8).

The highly fossiliferous stratigraphic section cropping out at the Cajas Range has been proposed several times as the type section for the Cambrian – Ordovician transition in northwestern Argentina (Fig. 6).

The Cambro-Ordovician Volcancito Formation

To the south, in the Famatina System, neither the base nor the top of the Volcancito Formation is known. Esteban (1999) has recognized three members within the




FIGURE 7 | Stratigraphic section and biostratigraphic framework for the Cambro-Ordovician transition in the Famatina System (Cachiyuyo Group, lower member of the Volcancito Formation). Volcancito River section, La Rioja Province (modified after Tortello and Esteban, 1999. See Figs. 1 and 2 for location, and Appendix.




FIGURE 8 | Stratigraphical and biostratigraphical sketch for the Cambrian System in northwestern Argentina. The main fossiliferous groups have been integrated into a schematic stratigraphic section. See Figs. 5 to 7 and Appendix.

unit: a lower Filo Azul Member, the mid Peña Negra Member and an upper Bordo Atravesado Member. The Cambrian- Ordovician boundary has been placed within the upper part of the 170 m thick Filo Azul Member (Fig. 7).

The formation is constituted by 430 meters of highly fossiliferous slates, black shales, sandstones and few limestones. These lithofacies yielded graptolites, trilobites, conodonts, brachiopods and filocariids (Harrington, 1938; Harrington and Leanza, 1957; Turner, 1959; Toselli, 1977; Aceñolaza and Durand, 1984; Esteban and Gutiérrez-Marco, 1997; Tortello and Esteban, 1997; Albanesi et al., 1999, 2000). The Filo Azul Member has been interpreted to be deposited on shore zones (inner and outer shelf settings), reaching to shoreface in the uppermost part (Esteban, 1999). The lowermost part of the Volcancito Formation in the type locality displays a tectonic contact with the underlying Negro Peinado Formation (equivalent to the Puncoviscana Formation s.l.), while upwards, is followed by La Alumbreira Formation and the Arenig shales, sandstones and volcanoclastics of the Suri Formation (Fig. 2).

THE CAMBRIAN BIOSTRATIGRAPHIC RECORD

The biostratigraphic record of the above described sedimentary succession, which spans throughout the Cambrian, is analyzed in this section (Fig. 8).

The Cambrian System in Northwestern Argentina lacks a good biostratigraphical zonation compared to its counterpart in the Precordillera. Palaeoecological conditions related to the palaeogeography are interpreted to be the main cause of this contrast. The only fairly good biostratigraphical data come from the transitional levels to the Ordovician, in the uppermost part of the Cambrian sequences (Fig. 8). A para-autochthonous or autochthonous origin for the highly fossiliferous Cambrian Precordilleran sequences in the Gondwanan margin has been widely discussed during the last decade (see Bordonaro in this volume; Pankhurst and Rapela, 1998; Aceñolaza and Toselli, 2000; Aceñolaza et al., 2000; Peralta, 2000 with references; Finney et al. in press).

In Northwestern Argentina, low-grade metamorphism affected the lower Puncoviscana Formation (s.l.) and

equivalent units, dated only by means of trace fossils (Fig. 4). Quartzites and sandstones typify the Mesón Group, with remarkable good trace fossils but no significant shelly fauna (Fig. 5). Good biostratigraphic data can be obtained only from the shales and sandstones of the basal Santa Victoria and Cachiuyo Groups, which includes the Cambrian-Ordovician transition, with graptolites, trilobites, conodonts, brachiopods, molluscs, echinoderms and ichnofossils among others.

For the Eastern Cordillera, Puna and Subandean Ranges, the Santa Rosita Formation (basal unit of the Santa Victoria Group) is the one that includes the Cambrian-Ordovician boundary (Fig. 6). This boundary is included in the Filo Azul Member of the Volcancito Formation (basal part of the Cachiuyo Group) in the Famatina System of La Rioja Province (Fig. 7). Lithologically, both units are mainly characterized by shales and sandstones with a few quartzites of greyish, greenish and black colors, and displaying a mean thickness of a few hundred meters. The transition between the Cambrian and Ordovician is biostratigraphically well tied by graptolites (*Rhabdinopora flabelliformis* EICHWALD), conodonts (*Iapetognathus-Cordylodus*) and trilobites (*Jujuyaspis keideli* KOBAYASHI).

The tables provided in the Appendix to this paper show the known taxa for all the recorded palaeontological material yielded by the Precambrian-Cambrian Puncoviscana Formation, up to the Cambro-Ordovician Volcancito and Santa Rosita formations. The transitional levels to the Ordovician were also considered.

CONCLUDING REMARKS

The Cambrian sequences in the Andean margin of South America display a more or less shared, similar history. Nevertheless, the reduced biostratigraphic record does not provide sufficient insight into the studied sequences, to obtain a clear correlation with other Cambrian sequences and biostratigraphic records around the world.

The Cambrian biostratigraphic record displays a relatively low diversity in the Puna, Eastern Cordillera, Subandean Ranges and the Famatina System if compared to the record of the Precordillera terrane (see Bordonaro, in this volume). It is considered that the rather diverse palaeogeographic and palaeoenvironmental conditions under which both basins evolved during Cambrian account for the observed biostratigraphic record differences. While limestones are common in the western precordilleran sequences, siliciclastic rocks (sandstones, shales and conglomerates) are dominant in northwestern Argentina.

The transition from the Precambrian to the Lower Cambrian is located in the slightly metamorphosed, highly deformed Puncoviscana Formation. In spite of the scarcity of ichnofossil-bearing sections, the transition between the Precambrian and Cambrian is recognised by means of the trace fossils represented in the sequences.

The so called Cambrian explosion is recorded by means of the trace fossils represented in the slates of the Puncoviscana Formation (*s.l.*), with several ichnogenera as *Oldhamia*, *Protichnites*, *Nereites*, *Monomorphychnus*, *Dimorphichnus* and *Diplichnites* among others. During the deposition of the Mesón Group, little palaeontological information is displayed due to the special lithological characters (high-energy shelf quartzites and sandstones are dominant).

The Puncoviscana Formation is overlain by the almost barren, high-energy sandstones of the Mesón Group, which has delivered good palaeoichnological assemblages. The occurrence of the Middle Cambrian trilobite *Asaphiscus* sp. (which is associated to several other undetermined forms and brachiopods in the Lizoite Formation at the Locality of Potrerillos, Salta) causes some uncertainty into the dating of the unit. This occurrence also highlights the importance of the Mesón Group sections, which must be revisited for a better understanding of the Cambrian sequences in the region.

The second record of biodiversification in the study area is recognised in the finer sediments (shales and sandstones) of the lower part of the Cambro-Ordovician Santa Victoria Group, where most of the typical Early Paleozoic fossil groups are abundant. This transition to the Ordovician in the lower part of the Santa Victoria Group displays the most complete biostratigraphic data available in the studied sections. This “Cambro-Ordovician” biodiversification recorded in the Santa Victoria Group, may have started a short time before the flooding of the shelf during the early Tremadocian. This biodiversification is not well recorded because of the unfavorable lithology for fossil preservation (sandstone facies) that dominate the Mesón Group.

The number of taxa represented in the Puncoviscana Formation and Mesón Group is small compared to the material found in the transitional levels to the Ordovician in the Santa Victoria Group. Trace fossils are the most important elements within the older units, while a much more varied spectrum of taxa is represented in the younger ones (trilobites, graptolites, conodonts, brachiopods, and echinoderms, etc.).

Further biostratigraphic work is needed for a better understanding of the Cambrian System in Northwestern

Argentina. Research shall be focused in the fossiliferous levels of the Puncoviscana Formation (*s.l.*), and within marginal sectors of the Mesón basin, seeking softer environmental conditions that will help on a better fossil preservation within this high energy shallow Cambrian platform.

ACKNOWLEDGEMENTS

I wish to thank F. Aceñolaza, S. Esteban, J.C. Gutiérrez-Marco, J. Soler, D. Brown and L. Cabrera for the critical reading of this contribution. M.F. Tortello helped with many constructive discussions on some important biostratigraphic aspects of the different sections. D. Ruiz Holgado and E. Gómez are also thanked for drawing the figures. The Fundación Antorchas provided partial financial support to this research.

REFERENCES

- Aceñolaza, F.G., 1973. Sobre la presencia de trilobites en las cuarcitas del Grupo Mesón en Potrerillos, Provincia de Salta. *Revista de la Asociación Geológica Argentina*, 28, 309-311.
- Aceñolaza, F.G., 1978. Trazas de trilobites en los estratos del Ordovícico basal de la Puna argentina. *Acta Geológica Lilloana*, 15, 23-35.
- Aceñolaza, F.G., 1986. El género *Macrocystella* (Cystoidea) en el Tremadoc de Salta y Jujuy. *IV Congreso Argentino de Paleontología y Bioestratigrafía*, 1, 133-135.
- Aceñolaza, F.G., Aceñolaza G.F., 1992. The genus *Jujuyaspis* as a world reference fossil for the Cambrian-Ordovician boundary. In: Webby, B.D., Laurie, J.R. (eds.). *Global Perspectives on Ordovician Geology*, Elsevier, Rotterdam, 115-120.
- Aceñolaza, F.G., Alonso, R.N., 2000. La Formación Puncoviscana *s.l.* a partir de nuevos elementos icnológicos en la provincia de Salta. *Ameghiniana*, 37(4) Suplemento, 4R.
- Aceñolaza, F.G., Bordonaro, O., 1989. Presencia de *Asaphiscus* (Asaphiscidae – Trilobita) en la Formación Lizoite, Potrerillos, Salta y su significado geológico. *Serie Correlación Geológica*, 5, 21-28.
- Aceñolaza, F.G., Durand, F.R., 1984. Observaciones sobre la fauna graptolítica tremadociana del Famatina, provincia de La Rioja. *IX Congreso Geológico Argentino*, 4, 267-276.
- Aceñolaza, F.G., Durand, F.R., 1986. Upper Precambrian – Lower Cambrian biota from the northwest of Argentina. *Geological Magazine*, 123(4), 367-375.
- Aceñolaza, F.G., Durand, F.R., 1987. Paleontología del límite Precámbrico-Cámbrico de Argentina. *X Congreso Geológico Argentino*, 1, 315-320.
- Aceñolaza, F.G., Fernández, R.I., 1978. Trazas fósiles en el Ordovícico Inferior de la sierra de Cajas, Jujuy. *Acta Geologica Lilloana* (Suplemento), 14, 33-37.
- Aceñolaza, F.G., Tortello, M.F., 2003. El Alisal: a new locality with trace fossils of the Puncoviscana Formation (Late Precambrian-Early Cambrian) in Salta Province, Argentina. *Geologica Acta*, 1, 87-94.
- Aceñolaza, F.G., Toselli, A.J., 1976. Consideraciones estratigráficas y tectónicas sobre el Paleozoico Inferior del Noroeste Argentino. *Memorias del II Congreso Latinoamericano de Geología*, 2, 233-256.
- Aceñolaza, F.G., Toselli, A.J., 1981. *Geología del Noroeste Argentino*. Facultad de Ciencias Naturales e Instituto Miguel Lillo, Universidad Nacional de Tucumán, *Publicación Especial*, 1287, 212 pp.
- Aceñolaza, F.G., Toselli, A.J., 2000. Argentine Precordillera: allochthonous or autochthonous gondwanic?. *Zentralblatt für Geologie und Paläontologie*, 2000, 1(7/8), 743-756.
- Aceñolaza, F.G., Fernández, R.I., Manca, N., 1982. Caracteres bioestratigráficos y paleoambientales del Grupo Mesón (Cámbrico Medio-Superior), centro-oeste de América del Sur. *Estudios Geológicos*, 38, 385-392.
- Aceñolaza, F.G., Miller, H., Toselli, A.J., 1988. The Puncoviscana Formation (Late Precambrian – Early Cambrian), sedimentology, tectonometamorphic history and age of the oldest rocks of NW Argentina. In: Bahlburg, H., Breitenkreuz, Ch., Giese, P. (eds.). *The Southern Central Andes*. Springer Verlag, *Lecture Notes in Earth Sciences*, 17, 25-37.
- Aceñolaza, F.G., Aceñolaza, G.F., Esteban, S.B., 1999. Bioestratigrafía de la Formación Puncoviscana y unidades equivalentes en el NOA. In: González Bonorino, G., Omarini, R., Viramonte, J. (eds.). *Geología del Noroeste Argentino*, *Relatorio*, 1, 91-114.
- Aceñolaza, F.G., Miller, H., Toselli, A.J., 2000. The Pampean and Famatinian Cycles – Superimposed Orogenic Events in west Gondwana. In: Miller, H., Hervé, F. (eds.). *Geoscientific Cooperation with Latin America*. *Zeitschrift für Angewandte Geologie Sonderheft*, Hannover, SH 1, 337-344.
- Aceñolaza, G.F., 1996. Bioestratigrafía del límite Cámbrico-Ordovícico y Ordovícico Basal en la Quebrada de Humahuaca, Provincia de Jujuy, República Argentina. *Doctoral Thesis*. Facultad de Ciencias Naturales e IML, Universidad Nacional de Tucumán, 245 pp.
- Aceñolaza, G.F., 1998. Graptolites, conodonts and trilobites of uppermost Cambrian and Lower Ordovician age from Jujuy Province, northern Argentina. *Temas Geológico-Mineros ITGE*, 23, 145-147.
- Aceñolaza, G.F., 1999. *Macrocystella? durandi* sp. nov. (Echinodermata, Rhombifera) y el registro del género *Macrocystella* en la cuenca cambro-ordovícica del norte argentino. *Acta Geologica Hispanica*, 34(1), 89-101.
- Aceñolaza, G.F., Peralta, S., 2000. Cambrian from the southern edge. *Instituto Superior de Correlación Geológica, Miscelánea*, 6, 1-148.
- Aceñolaza, G.F., Tortello, M.F., 2000. El Alisal: a new locality with trace fossils of the Puncoviscana Formation (Late Precambrian-Early cambrian) in Salta Province. In: Aceñolaza, G., Peralta, S. (eds.). *Cambrian from the southern edge*. *Instituto Superior de Correlación Geológica, Miscelánea*, 6, 67-69.

- Albanesi, G.L., Astini, R., 2000. New conodont fauna from Suri Formation (Early-middle Ordovician), Famatina System, western Argentina. *Ameghiniana*, 37(4) Suplemento, 68R.
- Albanesi, G.L., Esteban, S.B., Barnes, C.R., 1999. Conodontes del límite Cámbrico-Ordovícico en la Formación Volcancito, Sistema de Famatina, Argentina. *Temas Geológico-Mineros ITGE*, 26, 521-526.
- Albanesi, G.L., Esteban, S.B., Hünicken, M.A., Barnes, C.R., 2000. Las biozonas de conodontes de la Formación Volcancito (Cámbrico tardío - Ordovícico temprano), Sistema de Famatina, noroeste de Argentina. *Ameghiniana*, 7(4) Suplemento, 5R.
- Alonso, R.N., Marquillas, R.A., 1981. Trazas fósiles de la Formación Campanario (Grupo Mesón, Cámbrico) en el norte argentino, consideraciones ambientales y geocronológicas. *Revista del Instituto de Geología y Minería de Jujuy*, 4, 95-111.
- Auboin, J., Borrello, A., Cecioni, G., Charrier, R., Chotin, P., Frutos, J., Thiele, R., Vicente, J., 1973. Esquisse paléogéographique et structural de Andes Meridionales. *Revue Geologique Physique Geologique Dynamique*, 15, 11-72.
- Baldís, B.A., Omarini, R., 1984. El Grupo Lerma (Precámbrico-Cámbrico) en la comarca central salteña y su posición en el borde pacífico sudamericano. *XI Congreso Geológico Argentino*, 1, 64-78.
- Bordonaro, O., 1992. El Cámbrico en Sudamérica. In: Gutiérrez-Marco, J.C., Saavedra Alonso, J., Rábano, I. (eds.). *El Paleozoico Inferior de Iberoamérica*. Publicación Especial, Universidad de Extremadura, 1, 69-84.
- Bordonaro, O., 2003. Review of the Cambrian stratigraphy of the Argentine Precordillera. *Geologica Acta*, 1, 9-18.
- Braccacini, I.O., 1960. Lineamientos principales de la evolución estructural de la Argentina. *Petrotecnica*, 25, 63-94.
- Buatois, L.A., Mángano, M.G., Aceñolaza, F.G., Esteban, S.B., 2000. The Puncoviscana ichnofauna of northwestern Argentina: a glimpse into the ecology of the Precambrian-Cambrian transition. In: Aceñolaza, G., Peralta, S. (eds.). *Cambrian from the southern edge*. Instituto Superior de Correlación Geológica, Miscelánea, 6, 82-84.
- Durand, F.R., 1990. Los conglomerados del Ciclo Pampeano en el noroeste Argentino. In: Aceñolaza, F.G., Miller, H., Toselli, A.J. (eds.). *El ciclo pampeano en el noroeste argentino*, Serie Correlación Geológica, 4, 61-69.
- Durand, F.R., 1996. La Transición Precámbrico-Cámbrico en el sur de sudamérica. In: Baldís, B.A., Aceñolaza, F.G. (eds.). *Early paleozoic Evolution in NW Gondwana*. Serie de Correlación Geológica, 12, 195-205.
- Durand, F.R., Aceñolaza, F.G., 1990a. Caracteres biofaunísticos, paleoecológicos y paleogeográficos de la Formación Puncoviscana (Precámbrico Superior – Cámbrico Inferior) del noroeste argentino. In: Aceñolaza, F.G., Miller, H., Toselli, A.J. (eds.). *El ciclo pampeano en el noroeste argentino*. Serie de Correlación Geológica, 4, 71-112.
- Durand, F.R., Aceñolaza, F.G., 1990b. Presencia de un artrópodo (?) primitivo en la Formación Puncoviscana (Precámbrico Superior – Cámbrico Inferior) de la Quebrada del Toro, Provincia de Salta, Argentina. *Serie de Correlación Geológica*, 7, 13-17.
- Durand, F.R., Spalletti, L., 1986. Las facies turbidíticas del Precámbrico Superior – Cámbrico Inferior en la zona de Corralito, Provincia de Salta. *I Reunión Argentina de Sedimentología*, Actas, 113-116.
- Esteban, S.B., 1996. Los primeros trilobites ciclopígidos en el Ordovícico de Argentina (Formación Volcancito, Sistema de Famatina). *Ameghiniana*, 33(1), 57-64.
- Esteban, S.B., 1999. Estratigrafía, geología sedimentaria y paleontología del Ordovícico basal del Sistema de Famatina. Unpublished Doctoral Thesis. Facultad de Ciencias Naturales e IML, Universidad Nacional de Tucumán. 189 pp.
- Esteban, S.B., Gutiérrez-Marco, J.C., 1997. Graptolitos del Tremadoc del Sistema de Famatina (Argentina). *V Reunión Internacional Proyecto 351 PICG, XIII Jornadas de Paleontología*. Publicación Especial Universidade da Coruña, 59-63.
- Finney, S., Gleason, J., Gehrels, G., Peralta, S., Aceñolaza, G. in press. Early Gondwanan Connection for the Argentine Precordillera Terrane. *Earth and Planetary Sciences Letters*.
- Gonzalez-Bonorino, G., Llanbías, E., 1996. Geology and paleogeographic development of southern South America (excluding Brazil) in the late Proterozoic and early Paleozoic. In: Moullade, M., Nairn, A.E.M. (eds.). *The Phanerozoic Geology of the World I, The Paleozoic B*, Elsevier, 265-338.
- Harrington, H.J., 1938. Sobre las faunas del Ordoviciano inferior del norte argentino. *Revista del Museo de La Plata, Nueva Serie, Sección Paleontología*, 1, 109-289.
- Harrington, H.J., Leanza, A.J., 1975. Ordovician trilobites of Argentina, University of Kansas, Special publication, 1, 259 pp.
- Hünicken, M.A., Suárez Riglos, M., Sarmiento, G.N., 1985. Conodontes tremadocianos de la Sierra de Cajas. Departamento Humahuaca, Provincia de Jujuy. *Boletín de la Academia Nacional de Ciencias, Córdoba*, 56(3-4), 333-347.
- Jezek, P., 1990. Análisis sedimentológico de la Formación Puncoviscana entre Tucumán y Salta. In: Aceñolaza, F.G., Miller, H., Toselli, A.J. (eds.). *El Ciclo Pampeano en el Noroeste Argentino*. Serie Correlación Geológica, 4, 9-36.
- Jezek, P., Willner, A.P., Aceñolaza, F.G., Miller, H., 1985. The Puncoviscana trough – a large basin of Late Precambrian to Early Cambrian age on the pacific edge of the Brazilian Shield. *Geologische Rundschau*, 74(3), 573-584.
- Koukharsky, M., Torres Claro, R., Etcheverría, M., Vaccari, N.E., Waisfeld, B.G., 1996. Episodios volcánicos del Tremadociano y del Arenigiano en Vega Pinato, Puna Salteña, Argentina. *XIII Congreso Geológico Argentino y III Congreso de Exploración de Hidrocarburos*, 5, 535-542.
- Malanca, S., 1996. Morfología y ontogenia de un nuevo shumardiidae (Trilobita) del Tremadociano de la Sierra de Mojotoro, Salta, Argentina, *Memorias del XII Congreso Geológico Boliviano*, 2, 319-399.
- Malanca, S., Monteros, J.A., Moya, M.C., 1998. Nuevos datos paleontológicos de la Formación Las Vicuñas (Tremadoc

- temprano), Puna occidental Argentina. VII Congreso Argentino de Paleontología y Bioestratigrafía (Bahía Blanca), Resúmenes, 90.
- Manca, N., 1986. Caracteres icnológicos de la Formación Campanario (Cámbrico Superior) en Salta y Jujuy. *Ameghiniana*, 23(1-2), 75-87.
- Mángano, M.G., Buatois, L.A., 2000. Ichnology, sedimentary dynamics, and sequence stratigraphy of the Mesón Group: A Cambrian macrotidal shallow-marine depositional system in northwestern Argentina. In: Aceñolaza, G., Peralta, S. (eds.). *Cambrian from the southern edge*. Instituto Superior de Correlación Geológica, Miscelánea, 6, 109-110.
- Mángano, M.G., Buatois, L., Aceñolaza, G.F., 1996. Trace fossils and sedimentary facies from a Late Cambrian - Early Ordovician tide dominated shelf (Santa Rosita Formation, northwest Argentina): Implications for ichnofacies models of shallow marine successions. *Ichnos*, 5(1), 53-88.
- Mon, R., Hongn, F.D., 1991. The structure of the Precambrian and Lower Paleozoic basement of the Central Andes between 22° and 32°. *Latinamerika Geologische Rundschau*, 80(3), 745-758.
- Mon, R., Salfity, J.A., 1995. Tectonic evolution of the Andes of Northern Argentina. In: Tankard, A.I., Suárez Soruco, R., Welsink, J.H. (eds.). *Petroleum Basins of South America*. American Association of Petroleum Geologists, Memoir, 62, 269-283.
- Moya, M.C., 1988. Lower Ordovician in the southern part of the Argentine Eastern cordillera. In: Bahlburg, H., Bretkreuz, Ch., Giese, P. (eds.). *The Southern Central Andes*. Springer Verlag, Lecture Notes in Earth Sciences, 17, 55-69.
- Moya, M.C., 1998. El Paleozoico Inferior en la Sierra de Mojotoro, Salta-Jujuy. *Revista de la Asociación Geológica Argentina*, 53(2), 219-238.
- Moya, M.C., Malanca S., Hongn, F.D., Bahlburg, H., 1993. El Tremadoc temprano en la Puna occidental Argentina. XII Congreso Geológico Argentino y II Congreso de Exploración de Hidrocarburos, Mendoza, 2, 20-30.
- Moya, M.C., Malanca, S., Monteros, J.A., Cuerda, A., 1994. Bioestratigrafía del Ordovícico Inferior en la Cordillera Oriental Argentina basada en graptolitos. *Revista Española de Paleontología*, 9, 91-104.
- Ortega, G., Rao, R.I., 1995. Nuevos hallazgos de graptolitos (Ordovícico Inferior) en la Sierra de Cajas. Departamento Humahuaca, Provincia de Jujuy, Argentina. *Boletín de la Academia Nacional de Ciencias, Córdoba*, 60(3-4), 293-316.
- Ortega, G., Albanesi, G., Rao, R.I., 1998. Lower Ordovician graptolites and conodonts from Cajas range and Parcha area, Eastern Cordillera, Northern Argentina. *Temas Geológico-Mineros ITGE*, 23, 236-240.
- Pankhurst, R.J., Rapela, C.W., 1998. The Proto-Andean Margin of Gondwana. *Geological Society, Special Publication*, 142, 383 pp.
- Peralta, S.H., 2000. An introduction to geology of the Precordillera, Western Argentina. In: Aceñolaza, G., Peralta, S. (eds.). *Cambrian from the southern edge*. Instituto Superior de Correlación Geológica, Miscelánea, 6, 14-20.
- Ramos, V.A., 1973. Trazas de trilobites de la Formación Campanario (Cámbrico) en la Puna Salteña y otros fósiles problemáticos del Paleozoico Inferior. *Ameghiniana*, 10(3), 229-234.
- Rao, R.I., 1992. Upper Cambrian conodonts from Sierra de Cajas, Cordillera Oriental of Argentina. Contribución al proyecto IGCP 271, LACON II. Publicación Especial de la Academia Nacional de Ciencias, Córdoba, 89 pp.
- Rao, R.I., 1994. Conodontes Ordovícicos de la Sierra de Cajas y del Espinazo del Diablo, Departamento Humahuaca, Provincia de Jujuy, República Argentina. Unpublished Doctoral Thesis. Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba, 89 pp.
- Rao, R.I., 1999. Los conodontes Cambro-Ordovícicos de la Sierra de Cajas y Espinazo del diablo, Cordillera Oriental, República Argentina. *Revista Española de Micropaleontología*, 31(1), 23-51.
- Rao, R.I., Hünicken, M.A., 1995a. Conodontes del Cámbrico Superior-Ordovícico Inferior en el área de Purmamarca. Cordillera Oriental, Provincia de Jujuy, Argentina. *Boletín de la Academia Nacional de Ciencias, Córdoba*, 60(3-4), 249-266.
- Rao, R.I., Hünicken, M.A., 1995b. Conodont biostratigraphy of the Cambrian-Ordovician boundary in northwestern Argentina. In: S.C. Finney, M. Droser (eds.). *Ordovician Odyssey*. Seventh International Symposium on the Ordovician System, Pacific Section Society for Sedimentary Geology, 77, 124-127.
- Rao, R.I., Tortello, M.F., 1998. Tremadoc conodonts and trilobites from Cardonal Formation, Incamayo Creek, Salta Province, Northwestern Argentina. *Paleontologia Polonica*, 58, 31-45.
- Rao, R.I., Ortega, G., Hünicken, M.A., 1994. Cambrian Ordovician Boundary in the Sierra de Cajas, Cordillera Oriental, Jujuy Province, Argentina. *Serie de Correlación Geológica*, 12, 22-23.
- Sánchez, M.C., 1994. El Grupo Mesón (Cámbrico) en el tramo austral de la Cordillera Oriental Argentina. Unpublished Doctoral Thesis. Universidad Nacional de Salta. 248 pp.
- Sánchez, M.C., 1999. Sedimentología y paleogeografía del Grupo Mesón (Cámbrico). In: González Bonorino, G., Omarini, R., Viramonte, J. (eds.). *Geología del Noroeste Argentino, Relatorio del XIV Congreso Geológico Argentino*, 126-133.
- Sánchez, M.C., Herrera, Z.A., 1994. Braquiópodos inarticulados cámbricos en la Formación Campanario (Grupo Mesón), Rio Reyes, Provincia de Jujuy. *Resúmenes, VI Congreso Argentino de Paleontología y Bioestratigrafía, Trelew*, 69.
- Sánchez, M.C., Salfity, J.A., 1999. La cuenca cámbrica del Grupo Mesón en el Noroeste Argentino: Desarrollo estratigráfico y paleogeográfico. *Acta Geologica Hispanica*, 34(2-3), 123-139.
- Sánchez, T.M., 2000. The oldest record of Rostroconchia (Mollusca) from western Gondwana (Cordillera Oriental, northwestern Argentina). *Ameghiniana*, 37(3), 265-269.

- Spalletti, L., Durand, F.R., 1987. Estudio de las facies sedimentarias del basamento de la región de Corralito, Provincia de Salta, Argentina. X Congreso Geológico Argentino, 1, 311-314.
- Tortello, M.F., Aceñolaza, G.F., 1993. Trilobites agnóstidos del límite Cámbrico-Ordovícico de la Formación Lampazar, Sierra de Cajas, Provincia de Jujuy, Argentina. Revista Española de Paleontología, 8(2), 177-184.
- Tortello, M.F., Aceñolaza, G.F., 1999. Trilobites agnóstidos del Ordovícico basal en la localidad de Purmamarca, Provincia de Jujuy, Argentina. Temas Geológico-Mineros ITGE, 26, 585-588.
- Tortello, M.F., Esteban, S.B., 1995. Un trilobite agnóstido en el Ordovícico de la región de Bordo Atravesado (Cuesta de Miranda), Provincia de La Rioja, Argentina. VI Congreso Argentino de Paleontología y Bioestratigrafía, 271-275.
- Tortello, M.F., Esteban, S.B., 1997. Significado bioestratigráfico de una asociación de trilobites del tramo basal de la Formación Volcancito (Sistema de Famatina, La Rioja, Argentina). Ameghiniana, 34(3), 265-270.
- Tortello, M.F., Esteban, S.B., 1999. La transición Cámbrico-Ordovícico en la Formación Volcancito (Sierra de Famatina, La Rioja, Argentina). Ameghiniana, 36(4), 371-387.
- Tortello, M.F., Esteban, S.B., Lech, R.R., 1996. Paleontología del Sistema de Famatina. Münchner Geologische Hefte, 19 (Reihe A), 125-136.
- Tortello, M.F., Rábano, I., Rao, R.I., Aceñolaza, F.G., 1999. Los trilobites de la transición Cámbrico-Ordovícico en la Quebrada Amarilla (Sierra de Cajas, Jujuy, Argentina). Boletín Geológico y Minero, 110(5), 555-572.
- Toselli, G., 1977. El Paleozoico Inferior y Medio de la región de Volcancito, Sierra de Famatina, Provincia de La Rioja, República Argentina. Acta Geologica Lilloana, 14, 83-104.
- Turner, J.M.C., 1959. La fauna graptolítica de América del Sur. Revista de la Asociación Geológica Argentina, 15(1-2), 5-160.
- Turner, J.M.C., 1960. Estratigrafía de la Sierra de Santa Victoria y adyacencias. Boletín de la Academia Nacional de Ciencias, Córdoba, 41, 163-196.
- Turner, J.M.C., 1964. Descripción geológica de la hoja 2c, Santa Victoria (Provincias de Salta y Jujuy). Instituto Nacional de Geología y Minería., Boletín 104, 135 pp.
- Turner, J.M.C., Méndez, V., 1975. Geología del sector oriental de los departamentos de Santa Victoria e Iruya, Provincia de Salta, República Argentina. Boletín de la Academia Nacional de Ciencias, 51(1-2), 11-24.

Manuscript received October 2001;
revision accepted July 2002.

APPENDIX

Biostratigraphic record of the Cambrian in NW Argentina

The taxa included in this appendix are considered significant for the definition of the Lower Paleozoic boundaries.

PHYLUM CONODONTA

The scheme given by Rao (1999) for the proposed Cambro-Ordovician stratotype in northwestern Argentina is followed (Sierra de Cajas, Jujuy Province). Summarised data from Hünicken et al. (1985), Rao (1992, 1994, 1999), Rao et al. (1994), Rao and Hünicken (1995a, 1995b), Aceñolaza (1998) and Rao and Tortello (1998). Additional data from the Volcancito Formation by Albanesi et al. (1999, 2000) and Albanesi and Astini (2000). Asterisk indicates the first Ordovician taxa. See Figs. 7 to 8.

***Cordylodus proavus* Zone**

***Hirsutodontus hirsutus* Subzone (Late Cambrian)**

Cordylodus primitivus

C. proavus

Hertzina elongata

Eoconodontus notchpeakensis

Eoconodontus sp.

Phakelodus elongatus

Furnishina furnishi

F. gladiata

F. primitiva

Teridontus nakamurai

Ph. simplex

Ph. tenuis

Fryxellodontus sp.

Hirsutodontus aff. *hirsutus*

Teridontus sp.

Proconodontus posterocostatus

Prooneotodus gallatini

P. rotundatus

Gen. et sp. indet. 2

***Cordylodus caboti* Zone
(Late Cambrian)**

Cordylodus caboti
Teridontus gracilis
T. nakamurai

***Cordylodus intermedius* Zone
(Late Cambrian)**

Cordylodus intermedius
C. caboti
C. durcei
Teridontus gracilis

T. nakamurai
Cordylodus proavus
Cordylodus sp.
 Gen. et sp. indet.1

***Cordylodus lindstromi* Zone *
(Base of Ordovician)**

Cordylodus caboti
Cordylodus aff. deflexus
C. intermedius
C. lindstromi
Semiacontiodus sp.

Teridontus gracilis
Eoconodontus? sp.
C. proavus
C. viruanus
Cordylodus sp.
Drepanodus cf. simplex
T. nakamurai

***Iapetognathus aengensis* Zone *
(Base of Ordovician)**

Iapetognathus aengensis

CLASS GRAPTOLITHINA

Data from Harrington and Leanza (1957), Turner (1959, 1960, 1964), Toselli (1977), Aceñolaza and Durand (1984), Moya et al. (1994), Ortega and Rao (1995), Tortello et al. (1996), Esteban and Gutiérrez-Marco (1997), Aceñolaza (1998), Ortega et al. (1998).

Aspidograptus cf. implicatus *
Callograptus cf. salteri *
Rhabdinopora flabelliformis “rustica”
R. flabelliformis flabelliformis *
R. flabelliformis cf. socialis *
R. flabelliformis cf. famatinense *
R. flabelliformis cf. norvegica *
R. flabelliformis cf. scitulun *
R. flabelliformis parabola
R. flabelliformis anglica *
Bryograptus sp. aff. kjerulfi *
Bryograptus sp. *
Paradelograptus sp. *

Anisograptus cf. richardsoni *
Anisograptus cf. flexuosus *
Aspidograptus cf. implicatus *
Aspidograptus? cf. *minor* *
Rhabdinopora n. sp.1 *
Rhabdinopora n. sp. 2 *
R. flabelliformis “patula” *
Staurograptus sp. *
Aspidograptus? cf. *minor* *
Staurograptus cf. dichotomus *
Adelograptus tenellus *
Aspidograptus cf. implicatus *
Callograptus cf. salteri *

CLASS TRILOBITA

Data from Harrington and Leanza (1957), Aceñolaza (1973), Aceñolaza and Bordonaro (1989), Moya et al. (1993), Tortello and Aceñolaza (1993), Tortello and Esteban (1995, 1997, 1999), Koukharsky et al. (1996), Esteban (1996, 1999), Malanca (1996), Malanca et al. (1998), Tortello and Aceñolaza (1999), Tortello et al. (1999). Asterisk indicates the first Ordovician taxa. See Figs. 6 to 8.

Cambrian fauna

Asaphiscus sp.
Parabolina (N.) frequens argentina
Lotagnostus (Lotagnostus) sp.
Lotagnostus sp.
Lotagnostus (Semagnostus) zuninoi

Leiostegium douglasi
Shumardia alata
Micragnostus calviformis,
Asaphellus riojanus
Gymnagnostus bolivianus
Kainella? conica

Leiagnostus turgidulus
Micragnostus vilonii
Onychopyge riojana
Rhadinopleura eurycephala
Rossaspis sp.

Strictagnostus? *micropeltis*
Trilobagnostus chiushuensis
 “*Bodenbenderia longifrons*”
 “*Sphaerocare globifrons*”

Lower Ordovician *

Jujuyaspis keideli *
Parabolina (N.) *frequens argentina*
Parabolinella argentinensis
Anglagnostus? n. sp. A
Micragnostus sp.
Cyclopyge? sp.
Corrugatagnostus? sp.
Micragnostus sp.
Trilobagnostus chiushuensis

?*Gymnagnostus bolivianus*
 “*Geragnostus*” sp.
Angelina hyeronimi
Anglagnostus? sp.
Kainella meridionalis *
Bienvillia tetragonalis *
Degamella famatinensis
Shumardia minutula *

Other trilobite taxa recorded for Cambrian - lowermost Ordovician strata

Shumardia alata
Shumardia sp.
Asaphellus catamarcensis
Pricyclopyge sp.
Pseudoperonopsis zuninoi
Rossaspis rossi
Bellaspidela sp.
Ceratopyge forficuloides
Cyclopyge sp.
Micragnostus hoeki
Neoagnostus (*Machairagnostus*) *tmetus*
Notopeltis orthometopa
Orometopus pyriformis

Parabolinella triarhroides
Basiliela carinata
Beltella ulrichi
Leptoplastides marianus
Hapalopleura clavata
Kainella? sp.
Leiagnostus turgidulus
Leiostegium sp.
Plicatolina scalpta
Acerocarina glaber
Apatokephalus serratus
Shumardia (*Conophrys*) *erquensis*
Shumardia (*Conophrys*) *sulcatus*
 “*Trinodus*” *jujuyensis*

PHYLLUM BRACHIOPODA

Data from Sánchez and Herrera (1994, with references). Both samples come from the Cambrian Mesón Group. See Fig. 5.

Lingula sp.
Lingulepis sp.

PHYLLUM MOLLUSCA

Data from Sánchez (2000).

Ribeiria francae

PHYLLUM ECHINODERMATA

Data from Aceñolaza (1986, 1999).

Macrocystella sp. *
Macrocystella? *durandi*

TRACE FOSSILS

Data from Ramos (1973), Aceñolaza (1978), Aceñolaza and Fernández (1978), Alonso and Marquillas (1981), Manca (1986), Aceñolaza and Durand (1987), Durand and Aceñolaza (1990a), Mángano et al. (1996), Aceñolaza et al. (1999, with references), Aceñolaza and Tortello (2000), Buatois et al. (2000), Mángano and Buatois (2000). Asterisk indicates the first Ordovician taxa. See Figs. 4 and 5.

<i>Alcyonidiopsis pharmaceus</i> *	<i>Monomorphichnus</i> isp.
<i>Arenicolites</i> isp.	<i>Multipodichnus</i> isp.
<i>Asaphoidichnus trifidus</i>	<i>Neonereites uniserialis</i>
<i>Asaphoidichnus</i> isp.	<i>Neonereites</i> isp.
<i>Bergaueria</i> isp.	<i>Nereites saltensis</i>
<i>Cochlichnus</i> isp.	<i>Nereites</i> isp.
<i>Alcyonidiopsis pharmaceus</i> *	<i>Oldhamia antiqua</i>
<i>Arenicolites</i> isp.	<i>Oldhamia flabellata</i>
<i>Asaphoidichnus trifidus</i>	<i>Oldhamia radiata</i>
<i>Asaphoidichnus</i> isp.	<i>Palaeophycus tubularis</i>
<i>Bergaueria</i> isp.	<i>Palaeophycus</i> isp.
<i>Cochlichnus</i> isp.	<i>Parahaentschelinia</i> isp.
<i>Cochlichnus anguineus</i>	<i>Phycodes pedum</i> *
<i>Cruziana</i> cf. <i>furcifera</i> *	<i>Planocraterion carbajalis</i> *
<i>Cruziana</i> cf. <i>breadstoni</i> *	<i>Planolites</i> isp.
<i>Cruziana goldfussi</i> *	<i>Protichnites</i> isp.
<i>Cruziana omanica</i> *	<i>Protovirgularia</i> isp.
<i>Cruziana problematica</i>	<i>Rusophycus carbonarius</i>
<i>Cruziana semiplicata</i>	<i>Rusophycus jenningsi</i>
<i>Cruziana</i> isp.*	<i>Rusophycus latus</i> *
<i>Didymaulichnus</i> isp.	<i>Rusophycus leifeirikssoni</i>
<i>Dimorphichnus</i> isp.	<i>Rusophycus</i> isp. A
<i>Diplichnites</i> isp.	<i>Rusophycus</i> isp. B
<i>Diplocraterion</i> isp.	<i>Rusophycus</i> isp. C
<i>Glockerichnus</i> isp.	<i>Rusophycus</i> isp.
<i>Gordia</i> isp.	<i>Scolicia</i> isp.
<i>Helminthoidichnites tenuis</i>	<i>Skolithos linearis</i>
<i>Helminthoidichnites</i> isp.	<i>Syringomorpha nilssoni</i>
<i>Helminthopsis</i> aff. <i>tenuis</i>	<i>Syringomorpha</i> isp.
<i>Helminthopsis tenuis</i>	<i>Tasmanadia cachii</i>
<i>Helminthopsis</i> isp.	<i>Torrowangea</i> isp.
<i>Helminthorhapse</i> isp.	<i>Tomaculum problematicum</i> *
<i>Helmintoida</i> cf. <i>miocenica</i>	<i>Treptichnus</i> isp.
<i>Helmintoida</i> isp.	cf. <i>Aulichnites</i> isp.
<i>Laevicyclus</i> isp.	cf. <i>Phycodes</i>
<i>Monocraterion</i> isp.	cf. <i>Protopaleodictyon</i> isp.
<i>Monomorphichnus bilinearis</i>	
<i>Monomorphichnus lineatus</i>	
<i>Monomorphichnus multilineatus</i>	