

# cryptogamie

*Algologie*

2021 • 42 • 7

*Melanothamnus macaronesicus*  
Rodríguez-Buján & Díaz-Tapia, sp. nov.  
(Rhodomelaceae, Rhodophyta): a new turf-forming  
species from the Azores and the Canary Islands

Iván RODRÍGUEZ-BUJÁN, Manuel PIMENTEL & Pilar DÍAZ-TAPIA

DIRECTEUR DE LA PUBLICATION / PUBLICATION DIRECTOR: Bruno DAVID  
Président du Muséum national d'Histoire naturelle

RÉDACTRICE EN CHEF / EDITOR-IN-CHIEF: Line LE GALL  
Muséum national d'Histoire naturelle

ASSISTANTE DE RÉDACTION / ASSISTANT EDITOR: Marianne SALAÜN ([algo@cryptogamie.com](mailto:algo@cryptogamie.com))

MISE EN PAGE / PAGE LAYOUT: Marianne SALAÜN

RÉDACTEURS ASSOCIÉS / ASSOCIATE EDITORS

**Ecoevolutionary dynamics of algae in a changing world**

Stacy KRUEGER-HADFIELD

Department of Biology, University of Alabama, 1300 University Blvd, Birmingham, AL 35294 (United States)

Jana KULICHOVA

Department of Botany, Charles University, Prague (Czech Republic)

Cecilia TOTTI

Dipartimento di Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, Via Brecce Bianche, 60131 Ancona (Italy)

**Phylogenetic systematics, species delimitation & genetics of speciation**

Sylvain FAUGERON

UMI3614 Evolutionary Biology and Ecology of Algae, Departamento de Ecología, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Av. Bernardo O'Higgins 340, Santiago (Chile)

Marie-Laure GUILLEMIN

Instituto de Ciencias Ambientales y Evolutivas, Universidad Austral de Chile, Valdivia (Chile)

Diana SARNO

Department of Integrative Marine Ecology, Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Napoli (Italy)

**Comparative evolutionary genomics of algae**

Nicolas BLOUIN

Department of Molecular Biology, University of Wyoming, Dept. 3944, 1000 E University Ave, Laramie, WY 82071 (United States)

Heroen VERBRUGGEN

School of BioSciences, University of Melbourne, Victoria, 3010 (Australia)

**Algal physiology & photosynthesis**

Janet KÜBLER

California State University Northridge, Department of Biology, California State University, Northridge, CA 91330-8303 (United States)

**Prokaryotic algae**

Nico SALMASO

IASMA Research and Innovation Centre, Fondazione Mach-Istituto Agrario di S. Michele all'Adige, Limnology and River Ecology, Via E. Mach, 1, 38010 San Michele all'Adige, Trento (Italy)

Vitor VASCONCELOS

Faculdade de Ciências da Universidade do Porto and CIIMAR, Rua do Campo Alegre, s/n, 4169-007 Porto (Portugal)

COUVERTURE / COVER:

Extraits d'éléments de la Figure 3 / Extracts of the Figure 3

*Cryptogamie, Algologie* est indexé dans / *Cryptogamie, Algologie* is indexed in:

- Aquatic Sciences & Fisheries Abstracts Part I.
- Biological Abstracts
- Chemical Abstracts
- Current Contents
- Marine Science Contents Tables (FAO)
- Science Citation Index
- Publications bibliographiques du CNRS (Pascal).

*Cryptogamie, Algologie* est distribué en version électronique par / *Cryptogamie, Algologie* is distributed electronically by:

- BioOne® (<http://www.bioone.org/loi/crya>)

**Cryptogamie, Algologie** est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris  
*Cryptogamie, Algologie* is a fast track journal published by the Museum Science Press, Paris

Les Publications scientifiques du Muséum publient aussi / The Museum Science Press also publishes: *Adansonia*, *Geodiversitas*, *Zoosystema*, *Anthropozoologica*, *European Journal of Taxonomy*, *Naturae*, *Comptes Rendus Palévol*, *Cryptogamie* sous-sections *Bryologie*, *Mycologie*.

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle

CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)

Tél. : 33 (0)1 40 79 48 05 / Fax : 33 (0)1 40 79 38 40

[diff.pub@mnhn.fr](mailto:diff.pub@mnhn.fr) / <http://sciencepress.mnhn.fr>

© Publications scientifiques du Muséum national d'Histoire naturelle, Paris, 2021

ISSN (imprimé / print): 0181-1568 / ISSN (électronique / electronic): 1776-0984

# ***Melanothamnus macaronesicus* Rodríguez-Buján & Díaz-Tapia, sp. nov. (Rhodomelaceae, Rhodophyta): a new turf-forming species from the Azores and the Canary Islands**

**Iván RODRÍGUEZ-BUJÁN**

Coastal Biology Research Group, Faculty of Sciences and Centre for Advanced Scientific Research (CICA), University of A Coruña, 15071, A Coruña (Spain)

**Manuel PIMENTEL**

Evolutionary Biology Research Group, Faculty of Sciences and Centre for Advanced Scientific Research (CICA), University of A Coruña, 15071, A Coruña (Spain)

**Pilar DÍAZ-TAPIA**

Coastal Biology Research Group, Faculty of Sciences and Centre for Advanced Scientific Research (CICA), University of A Coruña, 15071, A Coruña (Spain) and Instituto Español de Oceanografía (IEO), Centro Oceanográfico de A Coruña, Aptdo. 130, 15080, A Coruña (Spain)  
[pdiaz@udc.es](mailto:pdiaz@udc.es) (corresponding author)

Submitted on 2 November 2020 | Accepted on 18 February 2021 | Published on 12 May 2021

Rodríguez-Buján I., Pimentel M. & Díaz-Tapia P. 2021. — *Melanothamnus macaronesicus* Rodríguez-Buján & Díaz-Tapia, sp. nov. (Rhodomelaceae, Rhodophyta): a new turf-forming species from the Azores and the Canary Islands. *Cryptogamie, Algologie* 42 (7): 77-91. <https://doi.org/10.5252/cryptogamie-algologie2021v42a7>. <http://cryptogamie.com/algologie/42/7>

## **ABSTRACT**

The use of molecular tools in red algal diversity surveys often reveals the existence of undescribed species. Here, we report a new Macaronesian turf-forming red alga in the otherwise mostly Pacific genus *Melanothamnus* Bornet & Falkenberg. This new taxon, *Melanothamnus macaronesicus* Rodríguez-Buján & Díaz-Tapia, sp. nov. is described based on morphological and molecular (*rbcL* gene) evidence. Morphologically, it differs from other *Melanothamnus* taxa by a combination of characters that includes its decumbent habit with an extensive system of prostrate axes, small size ( $\leq 2.5$  cm), four ecorticate pericentral cells, unbranched to once-branched trichoblasts and spermatangial branches formed on the first dichotomy of trichoblasts that have a sterile apical cell. Molecularly, this new species differs from its congeners by sequence divergence  $\geq 3.3\%$  in the *rbcL* gene. Current known distribution of *M. macaronesicus* Rodríguez-Buján & Díaz-Tapia, sp. nov. is restricted to Macaronesia, suggesting that it could be endemic to this archipelago. Our findings suggest that endemism among red algae might be more common than previously thought in this bioregion, particularly among easily overlooked turf-forming species.

**KEY WORDS**  
Algal turfs,  
endemism,  
Macaronesia,  
morphology,  
overlooked diversity,  
phylogeny,  
*Polysiphonia*,  
*rbcL*,  
Streblocladiae,  
new species.

## **RÉSUMÉ**

*Melanothamnus macaronesicus* Rodríguez-Buján & Díaz-Tapia, sp. nov. (Rhodomelaceae, Rhodophyta) : une nouvelle espèce gazonnante dans les Açores et les îles Canaries.  
L'utilisation d'outils moléculaires dans les suivis de diversité des algues rouges révèle souvent l'existence d'espèces non décrites. Ici, nous rapportons une nouvelle algue rouge de Macaronésie formant un gazon du genre *Melanothamnus* Bornet & Falkenberg, dont la distribution est principalement Pacifique. Ce nouveau taxon, *Melanothamnus macaronesicus* Rodríguez-Buján & Díaz-Tapia, sp. nov.



**MOTS CLÉS**  
Gazons algaux,  
endémisme,  
Macaronésie,  
morphologie,  
diversité négligée,  
phylogénie,  
*Polysiphonia*,  
*rbcL*,  
Streblocladiaeae,  
espèce nouvelle.

est décrit sur la base de preuves morphologiques et moléculaires (gène *rbcL*). Morphologiquement, il diffère des autres taxons de *Melanothamnus* par une combinaison de caractères qui incluent son port décombant avec un système étendu d'axes prostrés, sa petite taille ( $\leq 2,5$  cm), quatre cellules péricentrales écourtées, des trichoblastes non ramifiés à une fois ramifiés et des rameaux spermatangiaux formés sur la première dichotomie des trichoblastes qui ont une cellule apicale stérile. Moléculairement, cette nouvelle espèce diffère de ses congénères par une divergence de séquence  $\geq 3,3$  % dans le gène *rbcL*. La distribution actuelle connue de *M. macaronesisus* Rodríguez-Buján & Díaz-Tapia, sp. nov. est limitée à la Macaronésie, ce qui laisse à penser qu'elle pourrait être endémique à cet archipel. Nos résultats suggèrent que l'endémicité parmi les algues rouges pourrait être plus fréquente qu'on ne le pensait auparavant dans cette biorégion, en particulier parmi les espèces formant des gazons, facilement négligés.

## INTRODUCTION

More than 7000 red algal species have been described (Guiry & Guiry 2020), representing only an estimated *c.* 50% of the total diversity (Guiry 2012). Cryptic species are common among the Rhodophyta with simpler morphologies (Van Oppen *et al.* 1996), especially in lineages displaying a strong morphological convergence that breaks the positive correlation between genetic and morphological differentiation (Mineur *et al.* 2012; Verbruggen 2014).

The Rhodomelaceae is the most diverse family of the Rhodophyta, with the tribes Polysiphoniae and Streblocladiae covering much of that diversity (Díaz-Tapia *et al.* 2017a). Most members of these tribes are small in size (<3 cm) and morphologically similar, and several groups of species share many of the diagnostic characters that are relevant for the delineation of taxa (Stuercke & Freshwater 2008; Savoie & Saunders 2015, 2018; Díaz-Tapia *et al.* 2020a). The identification of species in these rhodomelacean tribes usually requires detailed micro-morphological studies as well as molecular analyses, essential when taxa are morphologically indistinguishable. High diversity, small size and phenotypic conservatism render the tribes Polysiphoniae and Streblocladiae good candidates for hosting undiscovered and cryptic species. The combination of molecular and morphological studies provides the best tool to explore taxonomic diversity within morphologically similar red algal lineages (e.g. Stuercke & Freshwater 2010; Saunders *et al.* 2017; Soares *et al.* 2019; Díaz-Tapia *et al.* 2020b; Rodríguez-Prieto *et al.* 2020).

Genus *Melanothamnus* Bornet & Falkenberg (Rhodomelaceae; Streblocladiae) was originally defined as monotypic, including only *M. somalensis* Bornet & Falkenberg. Díaz-Tapia *et al.* (2017b) proposed a new circumscription for this genus based on phylogenetic studies, and 46 species formerly placed in *Kintarosiphonia* S.Uwai & M.Masuda, *Fernandosiphonia* Levring, *Neosiphonia* M.S.Kim & I.K.Lee and *Polysiphonia* Greville were transferred to *Melanothamnus*. This genus is differentiated from other similar genera by two synapomorphies: the presence of three cells in the carpogonial branch (other Rhodomelaceae have four); and the arrangement of plastids only on the radial walls of pericentral cells (other Rhodomelaceae have plastids also on the outer cell walls)

(Díaz-Tapia *et al.* 2017b). Eleven (20%) out of the 56 currently recognized species in *Melanothamnus* were discovered in the last 11 years (Mamoozadeh & Freshwater 2012; Bustamante *et al.* 2012, 2013a, b; Muangmai *et al.* 2014; Kim & Kim 2016; Díaz-Tapia *et al.* 2017c; Huisman 2018; Woodworth *et al.* 2019; Serio *et al.* 2020), suggesting that its diversity has been greatly underestimated.

*Melanothamnus* is distributed in most coastal areas worldwide, but it is far more diverse in the northwestern Pacific (Díaz-Tapia *et al.* 2017b). In Macaronesia, a group of volcanic archipelagos in the North Atlantic, this genus is represented by five species (Haroun *et al.* 2002; Freitas *et al.* 2019). Two of these taxa [*M. collabens* (C.Agardh) Díaz-Tapia & Maggs and *M. harveyi* (Bailey) Díaz-Tapia & Maggs] are considered as cryptogenic or introduced in the Atlantic (McIvor *et al.* 2001; Díaz-Tapia *et al.* 2017b; but for alternative interpretations of *M. harveyi*/ *japonicus* distribution see Kim & Yang 2006; Savoie & Saunders 2015; Piñeiro-Corbeira *et al.* 2020b). *Melanothamnus ferulaceus* (Suhr ex J.Agardh) Díaz-Tapia & Maggs and *M. sphaerocarpus* (Børgesen) Díaz-Tapia & Maggs have been widely recorded in the Atlantic and Pacific Oceans, with the former also distributed in the Indian Ocean, while *M. gorgoniae* (Harvey) Díaz-Tapia & Maggs is predominantly known in the western Atlantic and has also been recorded in Cape Verde (Guiry & Guiry 2020). In addition to these five species, the recently described *M. testudinis* Serio, G.Furnari, I.Moro & K.Sciuto that lives on turtles in the Mediterranean is also presumed to be found in the Canary Islands (Serio *et al.* 2020). The diversity of species of *Melanothamnus* in Macaronesia has been characterized based on morphological studies (Rojas-González 1997; Neto *et al.* 2001; both as *Polysiphonia*), whereas surveys based on molecular data have not been performed.

In 2018, we conducted a diversity survey of the family Rhodomelaceae in the islands of Lanzarote (Canary Islands, Spain) and São Miguel (Azores, Portugal). The molecular characterization of samples using the *rbcL* gene led to the discovery of several species that differ from others molecularly characterized elsewhere and that morphologically do not conform to the species previously described in the study area. The objective of this work is to clarify the taxonomic identity of one of the discovered species analyzing its phylogenetic position and characterizing its morphology.




FIG. 1. — Distribution of *Melanothamnus macaronesicus* Rodríguez-Buján & Díaz-Tapia, sp. nov. in Lanzarote (Canary Islands) and São Miguel (Azores).

## MATERIAL AND METHODS

Material was collected in 2018 in the intertidal of São Miguel (Azores) and Lanzarote (Canary Islands) during diversity surveys of the family Rhodomelaceae (Fig. 1; Appendix 1). Specimens were carefully cleaned and isolated using a stereomicroscope. For each species and sample, part of the material was dried in silica gel desiccant for DNA extraction and the remaining material was preserved in 4% formalin seawater at 4°C and stored in the dark for morphological studies. Some specimens were mounted in 20% Karo® Syrup solution (ACH Foods, Memphis, Tennessee, United States) or in a solution composed of 50% Karo, 1% aqueous aniline blue and 1% acetic acid. Sections for microscopic observations were made by hand using a razor blade. Pressed herbarium specimens were deposited at SANT (Herbarium of the University of Santiago de Compostela, Spain). Herbarium abbreviations follow Thiers (2021).

DNA was extracted from silica gel-dried material following an adapted cetyltrimethylammonium bromide (CTAB) protocol (Doyle & Doyle 1987). PCR amplification of the plastid *rbcL* gene was carried out using primers and condi-

tions specified in Piñeiro-Corbeira *et al.* (2020a). The PCR products were purified and sequenced by Macrogen (Korea).

In order to establish the phylogenetic position of the studied species in the tribes Polysiphoniae and Streblocladiae, we constructed a matrix for the *rbcL* plastid gene including a sequence of each haplotype of the studied species along with 81 sequences belonging to other species of these tribes (Appendix 1, 2). Three sequences of *Symphyocladia* Falkenberg (Pterosiphoniae) were also added to the matrix as outgroups (Appendix 2) based on Díaz-Tapia *et al.* (2017a). All sequences were downloaded from GenBank except those belonging to the species under study and the South African sample of *M. incomptus* (Harvey) Díaz-Tapia & Maggs (MW567797).

Sequences were aligned using MAFFT (Katoh *et al.* 2019). We used Mr. Model test v.2.3.7 (Nylander 2004) in order to determine the evolutionary model that best fitted our data. Phylogenetic trees were obtained using Maximum Parsimony (MP), Maximum Likelihood (ML) and Bayesian Inference (BI) strategies, but only the ML tree is shown. MP and BI tests were conducted according to Torrecilla *et al.* (2003) and Pimentel *et al.* (2013), respectively. ML analyses were performed according to Díaz-Tapia *et al.* (2017b). Paup v.4.a165




FIG. 2. — Phylogenetic tree estimated with ML analysis of *rbcL* sequences. Values at nodes indicate bootstrap support (only shown if >60). Species names shaded and printed in bold correspond to the new species, haplotype numbers (H1-H3 as in Appendix 1) and their distribution (AZ = Azores, CI = Canary Islands) are indicated. **Asterisks** indicate the species of *Melanothamnus* Bornet & Falkenberg that have been previously reported in Macaronesia.

(Swofford 2000), MrBayes 3.2.7a (Huelsenbeck & Ronquist 2001) and RAxML 8.1.X (Stamatakis 2014) were used for MP, BI and ML analyses, respectively. BI and ML tests were run in the *Cipres Science Gateway* server (<https://www.phylo.org/portal2/home.action>).

## RESULTS

### PHYLOGENY

The *rbcL* aligned matrix consisted of 1467 base pairs (length of new sequences 1183-1368 bp), of which 37.83% (555/1467) were variable. The GTR+I+G model was selected for all analyses. The phylogenetic analysis of the *rbcL* gene in the tribes Polysiphoniae and Streblocadidae placed the studied species in the tribe Streblocadidae with full support, together with the other *Melanothamnus* taxa (Fig. 2). The studied species was resolved as sister to *M. incomptus* (Harvey) Díaz-Tapia & Maggs from South Africa with high support (96%), although relationships with other members of *Melanothamnus* remained unresolved.

Eight *rbcL* sequences were newly obtained for the studied species, corresponding to three different haplotypes that were resolved in a fully supported clade. One haplotype was exclusively found in the Canary Islands, while the other two haplotypes were found only in the Azores (Fig. 2). Sequence divergence among haplotypes was up to 0.3% (3 base pairs). Sequence divergence between the studied species and the most closely related species (*M. incomptus*) was 3.3-3.4% (41-43 base pairs out of 1278). Thus, molecular analyses evidenced that the studied species differs from the other sequenced species in the genus, and *M. macaronesicus* Rodríguez-Buján & Díaz-Tapia, sp. nov. is described below.

### MORPHOLOGICAL OBSERVATIONS

Family RHODOMELACEAE Horaninow  
Genus *Melanothamnus* Bornet & Falkenberg

*Melanothamnus macaronesicus*  
Rodríguez-Buján & Díaz-Tapia, sp. nov.  
(Figs 3; 4)

DIAGNOSIS. — Thalli up to 2.5 cm in height, forming dense turfs composed of decumbent axes that produce extensive prostrate systems. Axes with four pericentral cells, ecorticate. Erect axes 130-200 µm in basal parts, with exogenous branches that replace trichoblasts, alternately or pseudodichotomously branched. Trichoblasts short, simple or once branched, formed on every segment. Spermatangial branches growing on the first dichotomy of fertile trichoblasts, with 1-2 apical sterile cells at maturity. Cystocarps globose. Fertile parts of tetrasporangial branches bifurcated or bearing alternate branches, with tetrasporangia forming markedly spiral series. *rbcL* sequence of the holotype: MW567802.

HOLOTYPE. — SANT 33663 (male and female gametophytes, tetrasporophytes).

ADDITIONAL STUDIED SPECIMENS. — PD4116 (sterile), PD4121 (tetrasporophytes), PD4129 (sterile), PD4130 (tetrasporophytes),

Puerto del Carmen, Lanzarote, Canary Islands, Spain; PD3692 (sterile), Praia do Populo, São Miguel, Azores, Portugal; PD3843 (sterile), PD3847 (sterile), Praia San Roque, São Miguel, Azores, Portugal.

TYPE LOCALITY. — Puerto del Carmen, Lanzarote, Spain.

ETYMOLOGY. — “*macaronesicus*” refers to the name of the region where the species was found.

HABITAT AND DISTRIBUTION. — *Melanothamnus macaronesicus* Rodríguez-Buján & Díaz-Tapia, sp. nov. was collected from turfs mixed with other small species in intertidal sand-covered rocks from the Azores and the Canary Islands.

### VEGETATIVE MORPHOLOGY

Thalli up to 2.5 cm high, forming turfs of densely entangled axes. Thalli formed by numerous decumbent axes, forming extensive prostrate systems. Prostrate axes bear erect axes dorsally that branch alternately or pseudodichotomously (Fig. 3A). Thallus brownish-red in colour, texture flaccid and delicate.

Axes ecorticate consisting of an axial cell surrounded by four pericentral cells (Fig. 3B). Pericentral cells with discoid plastids, lying exclusively on radial walls of pericentral cells (Fig. 3C), so outer walls appear transparent. Prostrate axes (130-) 150-340 µm in diameter, composed of segments Length/Diameter (L/D) 0.3-1.1 (-1.6). Rhizoids arising in the middle or in proximal parts of the pericentral cells, up to two per segment, cut off from the pericentral cells, unicellular, up to 1 mm in length and consisting of a filament 25-130 µm in diameter, occasionally branched and often terminated in digitate haptera (Fig. 3D). Prostrate axes bear erect axes dorsally at regular intervals and branch laterally forming further prostrate axes (Fig. 3E). Erect axes growing from obtuse apical cells of about 12.5 µm in diameter. Branches predominantly exogenous, replacing trichoblasts and formed every six segments, although larger or smaller distances are common (Fig. 3F). Adventitious branches occasional, growing from trichoblast scar cells. Basal segments in erect axes 130-200 µm in diameter; L/D 0.6-1.1. Segments in the middle part of the erect axes 80-120 µm in diameter, L/D 0.7-1.5.

Trichoblasts short, <260 µm in length with basal cells 7.5-10 (12.5) µm in diameter, simple or once branched. Trichoblasts formed on every segment, in a ¼ spiral pattern, deciduous and leaving conspicuous scar cells when shed (Fig. 3F, G).

### REPRODUCTIVE MORPHOLOGY

Gametophytes dioecious. Spermatangial branches growing at the apices of erect axes on the first dichotomy of a trichoblast, one per segment in a ¼ spiral (Fig. 4A). Spermatangial axes cylindrical, 90-190 µm in length and 25-40 µm in diameter, with 1-2 apical sterile cells at maturity (Fig. 4B).

Cystocarps formed at the apices of erect axes, globose, up to 550 µm in height and diameter, cells around the ostiole are slightly larger than cells of the pericarp immediately below (Fig. 4C, D).

Tetrasporangia developing at the apices of alternate branches whose fertile parts are bifurcated or alternately branched






FIG. 3. — *Melanothamnus macaronensis* Rodríguez-Buján & Díaz-Tapia, sp. nov.: vegetative morphology: **A**, habit, prostrate axis with many rhizoids (arrows) and erect axes alternately branched; **B**, cross section with a small axial cell and four pericentral cells; **C**, surface view of pericentral cells with plastids lying only on radial walls, as well as a scar cell of a trichoblast (arrow); **D**, rhizoid cut off from pericentral cell; **E**, prostrate axes branched laterally; **F**, apices of erect axes with trichoblasts, alternately branched every 5-6 segments and showing scar cells of trichoblasts (arrows); **G**, erect axis showing the spiral arrangement of the scar cells of trichoblasts (arrows). A, E, G correspond to the holotype (PD4111); B (PD4116); C (PD4121); D, F (PD4129). Scale bars: A, 1 mm; B, C, G, 50  $\mu$ m; D, 200  $\mu$ m; E, 500  $\mu$ m; F, 100  $\mu$ m.

(Fig. 4E, F). Tetrasporangia forming markedly spiral series, subspherical, 37.5-67.5  $\mu$ m in diameter, tetrahedrally divided (Fig. 4F, G).

## DISCUSSION

In this work the new species *Melanothamnus macaronensis* Rodríguez-Buján & Díaz-Tapia, sp. nov. is described based on both molecular and morphological evidence. The new species was collected in two Macaronesian archipelagos (the Azores and the Canary Islands), growing in intertidal algal turfs mixed with other turf-forming species.

Our phylogenetic analyses evidenced that *Melanothamnus macaronensis* Rodríguez-Buján & Díaz-Tapia, sp. nov. is placed with full support in the genus *Melanothamnus* of the tribe Streblocladiae. Morphological observations support this placement, as *M. macaronensis* Rodríguez-Buján & Díaz-Tapia, sp. nov. has unicellular rhizoids cut off from the pericentral cells, a character distinctive of the tribe (Díaz-Tapia *et al.* 2017a). Moreover, plastids are arranged lying only on radial walls of pericentral cells, a synapomorphic character

of *Melanothamnus* (Díaz-Tapia *et al.* 2017b). However, the uniformity of this character in *Melanothamnus* has been recently discussed (Huisman 2018; Serio *et al.* 2020). Three Australian species with plastids lying on all walls of pericentral cells were included tentatively in *Melanothamnus* (Huisman 2018). Similarly, this character was described in Australian specimens identified based on morphology as *M. tongatensis* (Harvey *ex* Kützing) Díaz-Tapia & Maggs and *M. nudus* (Mamoozadeh & Freshwater) Freshwater, whose type localities are in Tonga and the Caribbean, respectively (Huisman 2018). Thus, further studies using molecular data are needed to test whether these Australian species or specimens actually belong to the genus *Melanothamnus* or if this character can exhibit some variability in the genus. Moreover, the recently described species *M. testudinis* has pericentral cells with plastids arranged either on all or only on radial walls (Serio *et al.* 2020). Our study of *M. macaronensis* Rodríguez-Buján & Díaz-Tapia, sp. nov. shows that this character can be easily observed in most formalin preserved specimens, but it can occasionally not be adequately studied in some specimens or some parts of them depending on the preservation conditions of plastids. The other synapomorphic character of *Melanoth-*






FIG. 4. — *Melanothamnus macaroneticus* Rodríguez-Buján & Díaz-Tapia, sp. nov.: reproductive morphology: **A**, erect axis with spermatangial branches; **B**, mature spermatangial branch with a sterile apical cell (arrow); **C**, erect axis with cystocarps; **D**, cystocarp and carpospores; **E**, tetrasporophyte; **F**, tetrasporangia arranged in spiral series; **G**, detail of tetrasporangia. A-E, G correspond to the holotype (PD4111); F (PD3843). Scale bars: A, 400  $\mu$ m; B, G, 50  $\mu$ m; C, 1 mm; D, F, 200  $\mu$ m; E, 500  $\mu$ m.

*amnus*, having 3-celled carpogonial branches (Díaz-Tapia *et al.* 2017b), could not be observed in *M. macaroneticus* Rodríguez-Buján & Díaz-Tapia, sp. nov. Other relevant characters that are common in *Melanothamnus* and that were present in the new species are the spiral arrangement of tetrasporangia, the development of spermatangial branches on the first dichotomy of trichoblasts and the formation of trichoblasts or branches on successive segments (Kim & Lee 1999, as *Neosiphonia*; Díaz-Tapia *et al.* 2017b).

*Melanothamnus macaroneticus* Rodríguez-Buján & Díaz-Tapia, sp. nov. differs from other sequenced congeners by *rbcL* sequence divergence  $\geq 3.3\%$ , supporting its recognition as a new species. Moreover, the new species can be morphologically distinguished from other species by its unique combination of the characters habit, number of pericentral cells, cortication, trichoblasts and number of apical sterile cells in spermatangial branches (Table 1). Molecular and/or morphological distinction of the new species regarding the other species of the genus includes the five formerly reported

species in Macaronesia (Fig. 2; Table 1). As most species of *Melanothamnus*, the new species has four pericentral cells, differing from the 12 species that have a higher number such as *M. forfex* (Harvey) Díaz-Tapia & Maggs or *M. collabens* (Womersley 1979, as *Polysiphonia*; Díaz-Tapia & Bárbara 2013, as *Streblocladia*). This trait has a high diagnostic value for distinguishing species in the Polysiphonieae and Streblocladieae (Stuercke & Freshwater 2008, as *Polysiphonia sensu lato*) and, therefore, we restricted our comparisons of other morphological characters to the new species and the other *Melanothamnus* spp. that have four pericentral cells.

*Melanothamnus macaroneticus* Rodríguez-Buján & Díaz-Tapia, sp. nov. differs morphologically from most *Melanothamnus* spp. by its decumbent habit with an extensive system of profusely branched prostrate axes. Only 13 species of the genus have a similar decumbent habit, which contrasts with most species that are erect or have short prostrate axes (Table 1). Another character that distinguishes *M. macaroneticus* Rodríguez-Buján & Díaz-Tapia, sp. nov. from most congeners

TABLE 1. — Comparison of selected morphological characters for species currently assigned to *Melanothamnus* Bornet & Falkenberg that have four pericentral cells.

Species	Type locality	Habit and length (cm)	Cortication	Trichoblasts: Arrangement, maximum length (µm) and number of times branched	Apical sterile cells in spermatangial branches	Diameter (µm) of erect axes	References
<i>M. macaronesicus</i> Rodríguez-Buján & Díaz-Tapia, sp. nov.	Canary Islands, Spain	Decumbent, < 2.5	–	On every segment. 220. Simple or once branched	1-2	Base: 130-200 Middle: 80-120	This work
<i>M. afaqhussainii</i> Shameel	Pakistan	Erect	+	Absent (old apices). Several segments apart (young apices)	–	<4000	Shameel (1999); Afaq-Husain & Shameel (2000)
<i>M. apiculatus</i> (Hollenberg) Díaz-Tapia & Maggs	Hawaii, United States	Erect	–	On every segment. 740. 3-4 times branched	1	150-250 (-320)	Hollenberg (1968); Kim & Abbott (2006)
<i>M. bajacali</i> (Hollenberg) Díaz-Tapia & Maggs	Baja California, Mexico	Erect	Light in some old apices	On every segment. 750. Branched many times	0	75-150	Hollenberg (1961); Mamoozadeh & Freshwater (2011)
<i>M. blandii</i> (Harvey) Díaz-Tapia & Maggs	Victoria, Australia	Decumbent	–	On every segment. 1-3 times branched	(1-) 2-3	300-550	Womersley (1979)
<i>M. cheloniae</i> (Hollenberg & J.N.Norris) Díaz-Tapia & Maggs	Gulf of California, Mexico	Erect	–	On every segment. 400. 1-3 times branched	0	300	Hollenberg & Norris (1977); Norris (2014)
<i>M. decumbens</i> (T.Segi) Díaz-Tapia & Maggs	Japan	Erect	+	On every segment. 1-3 times branched	1-2	Base: 600-1000 Middle: 300-600	Segi (1951); Kim (2003)
<i>M. delicatulus</i> (Hollenberg) Huisman	Hawaii, United States	Decumbent	–	On every segment. 230. 2-4 times branched	–	Middle: 25-30	Hollenberg (1968)
<i>M. eastwoodiae</i> (Setchell & N.L.Gardner) Díaz-Tapia & Maggs	Baja California, Mexico	Erect, 5-12	–	On every segment. Once branched	–	Base: 300-400	Norris (2014)
<i>M. ecorticatus</i> (R.E.Norris) Díaz-Tapia & Maggs	Hawaii, United States	Erect, <2	–	On every segment. Branched many times	>1	<2000	Norris (1994)
<i>M. ferulaceus</i> (Suhr ex J.Agardh) Díaz-Tapia & Maggs	East coast of Mexico; North America; Guadeloupe; Australia; Marquesas Islands; Hawaiian Islands.	Erect, <4 (-15)	–	On every segment. 90 µm. Branched many times	1-2	(150-) 250-350	Mamoozadeh & Freshwater (2012)
<i>M. flavimarinus</i> (M. S.Kim & I.K.Lee) Díaz-Tapia & Maggs	Korea	Erect, 4-5	–	On every segment. 150. 1-2 times branched	1-2	Base: 85-170 Middle: 100-130	Kim & Lee (1999)
<i>M. gigas</i> Huisman	Western Australia	Erect, 40	+	On every segment	Several		Huisman (2018)
<i>M. gorgoniae</i> (Harvey) Díaz-Tapia & Maggs	Florida, United States	Erect	–	Scarce. On every segment	1	Base: 160-270 Middle: 130-230	Kapraun (1979); Guimarães <i>et al.</i> (2004);
<i>M. hancockii</i> (E.Y.Dawson) Díaz-Tapia & Maggs	Baja California, Mexico	Erect, <2	+	On every segment. Branched	–	Base: <1000	Dawson (1944); Norris (1994)
<i>M. harlandii</i> (Harvey) Díaz-Tapia & Maggs	Hong Kong, China	Erect, 4-8	+	On every segment. 700. 3-4 times branched	1-2	Base: 800-1000 Middle: 500-800	Yoon (1986); Kim (2003)
<i>M. harveyi</i> (Bailey) Díaz-Tapia & Maggs	Connecticut, United States	Erect	+	On every segment	1-2	400-500	Maggs & Hommersand (1993); Kapraun (1977)
<i>M. hawaiiensis</i> (Hollenberg) Díaz-Tapia & Maggs	Hawaii, United States	Erect	–	On every segment. 500. Branched many times	1	300-500	Hollenberg (1968); Kim & Abbott (2006)

TABLE 1. — Continuation

Species	Type locality	Habit and length (cm)	Cortication	Trichoblasts: Arrangement, maximum length (µm) and number of times branched	Apical sterile cells in spermatangial branches	Diameter of erect axes (µm)	References
<i>M. incomptus</i> (Harvey) Díaz-Tapia & Maggs	South Africa	Decumbent, 2-3 (-6)	-	On every segment. Simple or once branched	0	<400	Anderson <i>et al.</i> (2016)
<i>M. infestans</i> (Harvey) Huisman	Western Australia	Erect, 16	-	On every segment	0-1	-	Huisman (2018)
<i>M. japonicus</i> (Harvey) Díaz-Tapia & Maggs	Japan	Erect, 1-20	+	On every segment. Once 1 or more branched	1	700-800	Segi (1951); Kim (2003)
<i>M. masonii</i> (Setchell & N.L.Gardner) Díaz-Tapia & Maggs	Baja California, Mexico	Erect, 2-3	-	On every segment. 480. 2-3 times branched	-	350-400	Hollenberg & Norris (1977)
<i>M. minutissimus</i> (Hollenberg) Díaz-Tapia & Maggs	Baja California, Mexico	Erect, <0.3	-	On every segment. 80	0	<145	Hollenberg (1942)
<i>M. nanus</i> (A.J.K.Millar) Díaz-Tapia & Maggs	New South Wales, Australia	Erect, <1	+	On every segment. 26	2	Base: <400 Middle: <180	Millar (1990)
<i>M. nudus</i> (N.R.Mamoozadeh & D.W.Freshwater) D.W.Freshwater	Panama	<0.7	-	Absent	-	70-140	Mamoozadeh & Freshwater (2012)
<i>M. platycarpus</i> (Børgesen) Díaz-Tapia & Maggs	India	Decumbent	-	On every segments. Branched many times	-	70-80	Børgesen (1934)
<i>M. pseudoforcipatus</i> Díaz-Tapia	Galicia, Spain	Erect	-	Scarce irregularly	-	Base: 170-190 (-220)	Díaz-Tapia <i>et al.</i> (2017b)
<i>M. pseudovillum</i> (Hollenberg) Díaz-Tapia & Maggs	Johnson Islands	Decumbent, 0.1-0.27	-	Most segments. 1000. 4 times branched	4	40-60	Hollenberg (1968)
<i>M. quadratus</i> (Hollenberg) Huisman	Gilbert Islands	Decumbent, 0.8	-	Irregular intervals of 4-8. 2-3 times branched	-	100-160	Hollenberg (1968)
<i>M. ramireziae</i> (D.E.Bustamante, B.Y.Won & T.O.Cho) Díaz-Tapia & Maggs	Peru	Erect, 2.4-5	-	On every segment. 31. Simple or once branched	-	86-103	Bustamante <i>et al.</i> (2013a)
<i>M. savatieri</i> (Hariot) Díaz-Tapia & Maggs	Japan	Erect, 0.5-1	+	On every segment. Twice 1-2 or more branched	1-2	100-200	Segi (1951); Kim (2005)
<i>M. silvae</i> (D.E.Bustamante, B.Y.Won & T.O.Cho) Díaz-Tapia & Maggs	Bali, Indonesia	Decumbent, 1.0-2.9	-	On every segment. 125. 1-2 times branched	-	61-73	Bustamante <i>et al.</i> (2013b)
<i>M. simplex</i> (Hollenberg) Díaz-Tapia & Maggs	California, United States	Decumbent	-	On every segment. 1-3 times branched	0	160-250	Hollenberg (1942)
<i>M. somalensis</i> Bornet & Falkenberg	Somalia	Erect, <36	+	Several segments apart	-	-	Kylin (1956); Wynne & Banaimoon (1990)
<i>M. sparsus</i> (Setchell) Díaz-Tapia & Maggs	Tahiti	Prostrate, < 1	-	On every segment. 650. 3-5 times branched	0	-	Hollenberg (1968)
<i>M. sphaerocarpus</i> (Børgesen) Díaz-Tapia & Maggs	Virgin Islands	Erect	-	On every segment. 875. Many times branched	0-1	100-180	Hollenberg (1968)
<i>M. strictissimus</i> (J.D.Hooker & Harvey) Díaz-Tapia & Maggs	New Zealand	Decumbent, (1)2-12(15)	+	On every segment. Branched many times.	1-2	500-1000	Adams (1991)
<i>M. testudinis</i> Serio, G.Furnari, I.Moro, K. Sciuto	Lampedusa, Italy	Decumbent, 2-3	-	On every segment. 500. 1-2 times branched	-	100-140	Serio <i>et al.</i> (2020)
<i>M. thailandicus</i> (N.Muangmai & C.Kaewsuralikhit)	Thailand	Erect, 5-15	-	Scarce. Associated with 1 reproductive structure. 100. 2-3 times branched	1	100-500	Muangmai <i>et al.</i> (2014)
<i>M. tongatensis</i> (Harvey ex Kützing) Díaz-Tapia & Maggs	Panama	Erect, 1.5-3(8)	-	On every segment. 175. Branched many times	0-1	50-150	Mamoozadeh & Freshwater (2012)



TABLE 1. — Continuation

Species	Type locality	Habit and length (cm)	Cortication	Trichoblasts: Arrangement, maximum length (µm) and number of times branched	Apical sterile cells in spermatangial branches	Diameter (µm) of erect axes	References
<i>M. unilateralis</i> (Levring) Díaz-Tapia & Maggs	Juan Fernández Islands, Chile	Erect	+	On every segment	–	750-1250 (-1500)	Levring (1941); Díaz-Tapia <i>et al.</i> (2017a)
<i>M. upolensis</i> (Grunow) Díaz-Tapia & Maggs	Samoa	Decumbent, 1-2-(3)	–	Every 2 or 3 segments. 340-(700). 1-3(4) times branched	1-2	100-120	Hollenberg (1968)
<i>M. yendoi</i> (T.Segi) Díaz-Tapia & Maggs	Japan	Decumbent, 1.5-4	–	On every segment. 200-300. 2-3 times branched	1	Base: 190-300 Middle: 100-140	Segi (1951); Yoon (1986)
<i>M. yongpili</i> (Kim & Kim) Díaz-Tapia & Maggs	Korea	Erect, 5-12	–	On every segment. 420. 1-3 times branched	2-3	180-340	Kim & Kim (2016)

and that has been uniformly observed in all studied specimens is that trichoblasts are short (<260 µm) and unbranched or branched only once. The only species with similar trichoblasts are *M. incomptus* and *M. ramireziae* (D.E.Bustamante, B.Y. Won & T.O.Cho) Díaz-Tapia & Maggs (Bustamante *et al.* 2012; Anderson *et al.* 2016), while trichoblasts are more profusely branched in other species.

*Melanothamnus incomptus* is the only currently recognized species that shares most of the diagnostic characters with *M. macaronesicus* Rodríguez-Buján & Díaz-Tapia, sp. nov. (decumbent, four pericentral cells, ecorticate, short trichoblasts). Interestingly, both species have been resolved as sister with high support in our phylogeny. Despite morphological similarities, they clearly constitute different molecular entities, as their divergence in the *rbcl* gene was 3.3-3.4%. It is worth mentioning that two sequences assigned to *M. incomptus* with divergent phylogenetic affinities were included in our phylogeny, suggesting the existence of cryptic diversity in this South African species. Regardless of which of the two sequences corresponds to the genuine *M. incomptus*, our phylogeny evidenced that both entities clearly differ from *M. macaronesicus* Rodríguez-Buján & Díaz-Tapia, sp. nov.

In addition to the molecular divergence between *M. incomptus* and *M. macaronesicus* Rodríguez-Buján & Díaz-Tapia, sp. nov., they differ from each other in the number of sterile cells in spermatangial branches (0 vs 1-2, respectively). However, even if this trait has been often used for species delineation, its reliability has been questioned as it can be variable in some species as well as depending on the maturity of spermatangial branches (Kim & Lee 1999; Stuercke & Freshwater 2008; Díaz-Tapia *et al.* 2021). Moreover, *M. incomptus* is larger in size than *M. macaronesicus* Rodríguez-Buján & Díaz-Tapia, sp. nov. as indicated by both thallus height (up to 6 vs up to 2.5 cm) and diameter of the prostrate and erect axes (up to 400 vs 150-340 µm; 200-240 µm vs 80-120 µm) (Rull Lluçh 2002; Anderson *et al.* 2016). Furthermore, *M. incomptus* may present a greater number of pericentral cells in some prostrate axes (6), a character never observed in *M. macaronesicus* Rodríguez-Buján & Díaz-Tapia, sp. nov. However, this

character was only observed in a North Namibian population (Rull Lluçh 2002) and it might be absent in the South African populations from which we obtained the sequences included in our phylogeny.

The current known distribution of *Melanothamnus macaronesicus* Rodríguez-Buján & Díaz-Tapia, sp. nov. and our previous work on the Streblocladieae suggest that this species could be endemic to Macaronesia. Extensive surveys of the Streblocladieae in the Atlantic Iberian Peninsula and the Mediterranean that included the sequencing of more than 250 specimens (PD, pers. obs.) failed to detect this taxon. Likewise, the new species was not found in molecular diversity surveys of this group in Central American coasts (Stuercke & Freshwater 2008, 2010; Mamoozadeh & Freshwater 2011, 2012), even if this region and Macaronesia often share the occurrence of species (Haroun & Prud'homme van Reine 1993; Machín-Sánchez *et al.* 2018). While we cannot rule out that the new species could have remained unnoticed in these or other undersampled regions (e.g. northern Africa), endemism would not be surprising. The Macaronesian archipelagos, of recent volcanic origin, host a high diversity of endemic animals and plants, which contrasts the low number (19) of endemic macroalgae (Freitas *et al.* 2019). Our finding suggests that further biodiversity studies of seaweeds using molecular data could reveal a higher number of endemisms than previously thought; particularly in algal turfs, the most common type of algal assemblage in Macaronesia (Tuya & Haroun 2006; Waltenstein *et al.* 2009) whose diversity has been barely studied using molecular tools (but see Díaz-Tapia *et al.* 2021). Turfs are composed by small species with a similar habit including prostrate and erect axes, and species identification often requires the observation of microscopic characters (Price & Scott 1992; Díaz-Tapia & Bárbara 2013, 2014). These traits render turf-forming species easy to overlook, as probably has occurred with *M. macaronesicus* Rodríguez-Buján & Díaz-Tapia, sp. nov. Our work provides further evidence that the species diversity in algal turfs has been largely underestimated and more effort is yet required in order to improve our understanding of these complex assemblages.

**Acknowledgements**

We warmly thank Ana Neto for providing assistance during field and lab work. This work was supported by Xunta de Galicia (“Axudas de apoio á etapa de formación posdoutoral” [grant ED481D/2017/011] and “Talento Senior” [grant 03IN858A2019-1630129] to PDT)).

**REFERENCES**

ADAMS N. M. 1991. — The New Zealand species of *Polysiphonia* Greville. *New Zealand Journal of Botany* 29: 411-427. <https://doi.org/10.1080/0028825X.1991.10415494>

AFAQ-HUSAIN S. & SHAMEEL M. 2000. — Further investigations on red alga *Melanothamnus afaqhusainii* (Ceramiales) from the coast of Pakistan. *Pakistan Journal of Botany* 31: 211-214.

ANDERSON R. J., STEGENGA H. & BOLTON J. J. 2016. — Seaweeds of the South African South Coast. World Wide Web electronic publication, University of Cape Town, <http://southafrseaweeds.uct.ac.za>. (Accessed on 16 Aug. 2020).

BÁRBARA I., CHOI H. G., SECILLA A., DÍAZ-TAPIA P., GOROSTIAGA J. M., SEO T. K., JUNG M. Y. & BERECIBAR E. 2013. — *Lampisiphonia iberica* gen. et sp. nov. (Ceramiales, Rhodophyta) based on morphology and molecular evidence. *Phycologia* 52: 137-155.

BØRGENSEN F. 1934. — Some Indian Rhodophyceae especially from the shores of the Presidency of Bombay – IV. *Bulletin of Miscellaneous Information, Royal Botanic Gardens, Kew* 1934: 1-30. <https://doi.org/10.2307/4111648>

BUSTAMANTE D. E., WON B. Y., RAMIREZ M. E. & CHO T. O. 2012. — *Neosiphonia peruviana* sp. nov. (Rhodomelaceae, Rhodophyta) from the Pacific coast of South America. *Botanica Marina* 55: 359-366. <https://doi.org/10.1515/bor-2012-0146>

BUSTAMANTE D. E., WON B. Y. & CHO T. O. 2013a. — *Neosiphonia ramirezii* sp. nov. (Rhodomelaceae, Rhodophyta) from Peru. *Algae* 28: 73-82. <https://doi.org/10.4490/algae.2013.28.1.073>

BUSTAMANTE D. E., WON B. Y. & CHO T. O. 2013b. — *Neosiphonia baliana* sp. nov. and *N. silvae* sp. nov. (Rhodomelaceae, Rhodophyta) from Bali, Indonesia. *Botanica Marina* 56: 515-524.

BUSTAMANTE D. E., WON B. Y. & CHO T. O. 2015. — *Polysiphonia freshwateri* sp. nov. and *Polysiphonia koreana* sp. nov.: two new species of *Polysiphonia* (Rhodomelaceae, Rhodophyta) from Korea. *European Journal of Phycology* 50: 330-342.

CARLILE A. L. 2009. — *Molecular systematics of North Pacific Ceramiales (Rhodophyta): Phylogeny, taxonomy, and phylogeography*. PhD, University of Washington.

D’ARCHINO R., NEILL K. & NELSON W. A. 2013. — Recognition and distribution of *Polysiphonia morrowii* (Rhodomelaceae, Rhodophyta) in New Zealand. *Botanica Marina* 56: 41-47.

DAWSON E. Y. 1944. — The marine algae of the Gulf of California. *Allan Hancock Pacific Expeditions* 3: 189-432.

DÍAZ-TAPIA P. & BÁRBARA I. 2013. — Seaweeds from sand-covered rocks of the Atlantic Iberian Peninsula. Part 1. The Rhodomelaceae (Ceramiales, Rhodophyta). *Cryptogamie Algologie* 34: 325-422. <https://doi.org/10.7872/crya.v34.iss4.2013.325>

DÍAZ-TAPIA P. & BÁRBARA I. 2014. — Seaweeds from sand-covered rocks of the Atlantic Iberian Peninsula. Part 2. Palmariales, Ceramiales (excluding Rhodomelaceae), Gelidiales, Gigartinales, Plocamiales, Rhodymeniales and Scytothamniales. *Cryptogamie Algologie* 35: 157-199. <https://doi.org/10.7872/crya.v35.iss2.2014.157>

DÍAZ-TAPIA P., KIM M. S., SECILLA A., BÁRBARA I. & CREMADES J. 2013. — Taxonomic reassessment of *Polysiphonia foetidissima* (Rhodomelaceae, Rhodophyta) and similar species, including *P. schneideri*, a newly introduced species in Europe. *European Journal of Phycology* 48: 345-362.

DÍAZ-TAPIA P., MAGGS C. A., WEST J. A. & VERBRUGGEN H.

2017a. — Analysis of chloroplast genomes and a supermatrix inform reclassification of the Rhodomelaceae (Rhodophyta). *Journal of Phycology* 53: 920-930. <https://doi.org/10.1111/jpy.12553>

DÍAZ-TAPIA P., MCIVOR L., FRESHWATER D. W., VERBRUGGEN H., WYNNE M. J. & MAGGS C. A. 2017b. — The genera *Melanothamnus* Bornet & Falkenberg and *Vertebrata* S.F. Gray constitute well-defined clades of the red algal tribe Polysiphoniae (Rhodomelaceae, Ceramiales). *European Journal of Phycology* 52: 1-31. <https://doi.org/10.1080/09670262.2016.1256436>

DÍAZ-TAPIA P., BÁRBARA I., CREMADES J., VERBRUGGEN H. & MAGGS C. A. 2017c. — Three new cryptogenic species in the tribes Polysiphoniae and Streblodidae (Rhodomelaceae, Rhodophyta). *Phycologia* 56: 605-623. <https://doi.org/10.2216/17-17.1>

DÍAZ-TAPIA P., LY M. & VERBRUGGEN H. 2020a. — Extensive cryptic diversity in the widely distributed *Polysiphonia scopulorum* (Rhodomelaceae, Rhodophyta): Molecular species delimitation and morphometric analyses. *Molecular Phylogenetics and Evolution* 152: 106909. <https://doi.org/10.1016/j.ympev.2020.106909>

DÍAZ-TAPIA P., MAGGS C. A., NELSON W., MACAYA E. C. & VERBRUGGEN H. 2020b. — Reassessment of the genus *Lophurella* (Rhodomelaceae, Rhodophyta) from Australia and New Zealand reveals four cryptic species. *European Journal of Phycology* 55: 113-128. <https://doi.org/10.1080/09670262.2019.1659419>

DÍAZ-TAPIA P., MUÑOZ-LUQUE, L., PIÑEIRO-CORBEIRA, C. & MAGGS C. A. 2021. — Phylogenetic analyses of Macaronesian turf-forming species reveal cryptic diversity and resolve *Stichothamnion* in the *Vertebrata* clade (Rhodomelaceae, Rhodophyta). *European Journal of Phycology* in press.

DOYLE, J. J. & DOYLE, J. L. 1987. — A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin* 19: 11-15.

FREITAS R., ROMEIRAS M., SILVA L., CORDEIRO R., MADEIRA P., GONZÁLEZ J. A., WIRTZ P., FALCÓN J. M., BRITO A. & FLOETER S. R. 2019. — Restructuring of the ‘Macaronesia’ biogeographic unit: A marine multi-taxon biogeographical approach. *Scientific Reports* 9: 15792. <https://doi.org/10.1038/s41598-019-51786-6>

FUJII M. T., GUIMARÃES S. M. P. D. B., GURGEL C. F. D. & FREDERICQ S. 2006. — Characterization and phylogenetic affinities of the red alga *Chondrophycus flagelliferus* (Rhodomelaceae, Ceramiales) from Brazil on the basis of morphological and molecular evidence. *Phycologia* 45: 432-441.

GUIMARÃES S. M. P. D. B., FUJII M. T., PUPO M. T. & YOKOYA N. S. 2004. — Reavaliação das características morfológicas e suas implicações taxonômicas no gênero *Polysiphonia sensu lato* (Ceramiales, Rhodophyta) do litoral dos Estados de São Paulo e Espírito Santo, Brasil. *Revista Brasileira de Botânica* 27: 163-183.

GUIRY M. D. 2012. — How many species of algae are there? *Journal of Phycology* 48: 1057-1063. <https://doi.org/10.1111/j.1529-8817.2012.01222.x>

GUIRY M. D. & GUIRY G. M. 2020. — *AlgaeBase*. World-wide electronic publication, National University of Ireland, Galway. <http://www.algaebase.org> (Accessed on 12 June 2020).

HAROUN R. J. & PRUD’HOMME VAN REINE W. F. 1993. — A biogeographical study of *Laurencia* and *Hypnea* species of the Macaronesian region. *Courier Forschungsinstitut Senckenberg* 159: 119-125.

HAROUN R. J., GIL-RODRÍGUEZ M. C., DÍAZ DE CASTRO J. & PRUD’HOMME VAN REINE W. F. 2002. — A checklist of the marine plants from the Canary Islands (central eastern Atlantic Ocean). *Botanica Marina* 45: 139-169. <https://doi.org/10.1515/BOT.2002.015>

HOLLENBERG G. J. 1942. — An account of the species of *Polysiphonia* on the Pacific coast of North America. I. *Oligosiphonia*. *American Journal of Botany* 29: 772-785. <https://doi.org/10.1002/j.1537-2197.1942.tb10279.x>

HOLLENBERG G. J. 1961. — Marine red algae of Pacific Mexico, Part 5: The genus *Polysiphonia*. *Pacific Naturalist* 2: 345-375.



- HOLLENBERG G. J. 1968. — An account of the species of *Polysiphonia* of the central and western tropical Pacific ocean. I *Oligosiphonia*. *Pacific Science* 22: 56-98.
- HOLLENBERG G. J. & NORRIS J. N. 1977. — The red alga *Polysiphonia* (Rhodomelaceae) in the Northern Gulf of California. *Smithsonian Contributions to the Marine Sciences* 1: 1-21.
- HUELSENBECK J. P. & RONQUIST F. 2001. — MrBayes: Bayesian Inference of phylogenetic trees. *Bioinformatics* 17: 754-755. <https://doi.org/10.1093/bioinformatics/17.8.754>
- HUISMAN J. M. 2018. — *Algae of Australia. Marine benthic algae of north-western Australia. 2. Red algae*. ABRIS & CSIRO Publishing, Canberra & Melbourne, 672 p.
- KAPRAUN D. F. 1977. — The Genus *Polysiphonia* in North Carolina, USA. *Botanica Marina* 20: 5. <https://doi.org/10.1515/botm.1977.20.5.313>
- KAPRAUN D. F. 1979. — The genus *Polysiphonia* (Ceramiales, Rhodophyta) in the vicinity of Port Aransas, Texas. *Contributions in Marine Sciences* 22: 105-120.
- KATOH K., ROZEWICKI J. & YAMADA K. D. 2019. — MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* 20: 1160-1166. <https://doi.org/10.1093/bib/bbx108>
- KIM M. S. & LEE I. K. 1999. — *Neosiphonia flavimarina* gen. et sp. nov. with a taxonomic reassessment of the genus *Polysiphonia* (Rhodomelaceae, Rhodophyta). *Phycological Research* 47: 271-281. <https://doi.org/10.1111/j.1440-1835.1999.tb00308.x>
- KIM M. S. 2003. — Taxonomic reassessment of the genus *Polysiphonia* (Rhodomelaceae, Rhodophyta): morphology and reproduction of *Neosiphonia decumbens* and *N. harlandii*. *Algae* 18: 107-119. <https://doi.org/10.4490/ALGAE.2003.18.2.107>
- KIM M. S. 2005. — Taxonomy of a poorly documented alga, *Neosiphonia savatieri* (Rhodomelaceae, Rhodophyta) from Korea. *Nova Hedwigia* 81: 163-176. <https://doi.org/10.1127/0029-5035/2005/0081-0163>
- KIM M. S. & YANG E. C. 2005. — Taxonomic note of *Polysiphonia pacifica* (Ceramiales, Rhodophyta) complex with focus on Pacific isolates. *Algae* 20: 15-23
- KIM M. S. & ABBOTT I. A. 2006. — Taxonomic notes on Hawaiian *Polysiphonia*, with transfer to *Neosiphonia* (Rhodomelaceae, Rhodophyta). *Phycological Research* 54: 32-39. <https://doi.org/10.1111/j.1440-1835.2006.00406.x>
- KIM M. S. & YANG E. C. 2006. — Taxonomy and phylogeny of *Neosiphonia japonica* (Rhodomelaceae, Rhodophyta) based on *rbcL* and *cpeA/B* gene sequences. *Algae* 21: 287-294.
- KIM B. & KIM M. S. 2014. — Three new species of *Polysiphonia* sensu lato (Rhodophyta) based on the morphology and molecular evidence. *Algae* 29: 183-195.
- KIM B. & KIM M. S. 2016. — *Neosiphonia yongpili* sp. nov. (Rhodomelaceae, Rhodophyta), known as *Neosiphonia simplex* from Korea, with an emphasis on cystocarp development. *Phycologia* 55: 323-332. <https://doi.org/10.2216/15-132.1>
- KIM M. S., KIM S. Y. & NELSON W. A. 2010. — *Symphyocladia lithophylla* sp. nov. (Rhodomelaceae, Ceramiales), a new Korean red algal species based on morphology and *rbcL* sequences. *Botanica Marina* 53: 233-241.
- KYLIN H. 1956. — *Die Gattungen der Rhodophyceen*. C.W.K. Gleerups, Lund, 673 p.
- LAM W. D., GARCÍA-FERNÁNDEZ M. E., ABOAL M. & VIS M. L. 2013. — *Polysiphonia subtilissima* (Ceramiales, Rhodophyta) from freshwater habitats in North America and Europe is confirmed as conspecific with marine collections. *Phycologia* 52: 156-160.
- LIN S. M., FREDERICQ S. & HOMMERSAND M. H. 2001. — Systematics of the Delesseriaceae (Ceramiales, Rhodophyta) based on large subunit rDNA and *rbcL* sequences, including the Phycodryioideae, subfam. nov. *Journal of Phycology* 37: 881-899.
- LEVRING T. 1941. — Die Meeresalgen der Juan Fernandez-Inseln. Die Corallinaceen der Juan Fernandez-Inseln, in Skottsberg C. (ed), *The natural history of Juan Fernandez and Easter Island*. Vol 2. Almqvist & Wiksells Boktryckeri, Uppsala: 601-670; 753-757.
- MACHÍN-SÁNCHEZ M., GIL-RODRÍGUEZ M. C. & HAROUN R. 2018. — Phylogeography of the red algal *Laurencia* complex in the Macaronesia region and nearby coastal areas: recent advances and future perspectives. *Diversity* 10: 10. <https://doi.org/10.3390/d10010010>
- MAGGS C. A. & HOMMERSAND M. H. 1993. — *Seaweeds of the British Isles. Volume 1. Rhodophyta, Part 3A Ceramiales*. Natural History Museum/HMSO, London, 444 p.
- MAMOOZADEH N. R. & FRESHWATER D. W. 2011. — Taxonomic notes on Caribbean *Neosiphonia* and *Polysiphonia* (Ceramiales, Florideophyceae): five species from Florida, USA and Mexico. *Botanica Marina* 54: 269-292. <https://doi.org/10.1515/bot.2011.036>
- MAMOOZADEH N. R. & FRESHWATER D. W. 2012. — *Polysiphonia sensu lato* (Ceramiales, Florideophyceae) species of Caribbean Panama including *Polysiphonia lobophoralis* sp. nov. and *Polysiphonia nuda* sp. nov. *Botanica Marina* 55: 317-347. <https://doi.org/10.1515/bot-2012-0147>
- MCIVOR L., MAGGS C. A., PROVAN J. & STANHOPE M. J. 2001. — *rbcL* sequences reveal multiple cryptic introductions of the Japanese red alga *Polysiphonia harveyi*. *Molecular Ecology* 10: 911-919. <https://doi.org/10.1046/j.1365-294X.2001.01240.x>
- MILLAR A. J. K. 1990. — Marine red algae of the Coffs Harbour region, northern New South Wales. *Australian Systematic Botany* 3: 293-593. <https://doi.org/10.1071/SB9900293>
- MINEUR F., COOK E. J., MINCHIN D., BOHN K., MACLEOD A. & MAGGS C. A. 2012. — Changing coasts: marine aliens and artificial structures. *Oceanography and Marine Biology: An Annual Review* 50: 187-232.
- MUANGMAI N., YAMAGISHI Y., MANEEKAT S. & KAEWSURALIKHIT C. 2014. — The new species *Neosiphonia thailandica* sp. nov. (Rhodomelaceae, Rhodophyta) from the Gulf of Thailand. *Botanica Marina* 57: 459-467. <https://doi.org/10.1515/bot-2014-0042>
- NETO A. I., CRAVO D. C. & HAROUN R. T. 2001. — Checklist of the benthic marine plants of the Madeira Archipelago. *Botanica Marina* 44: 391-414.
- NORRIS R. E. 1994. — Some cumophytic Rhodomelaceae (Rhodophyta) occurring in Hawaiian surf. *Phycologia* 33: 434-443. <https://doi.org/10.2216/i0031-8884-33-6-434.1>
- NORRIS J. N. 2014. — *Marine algae of the northern Gulf of California, II: Rhodophyta*. Smithsonian Institution Scholarly Press, Washington, 96 p.
- NYLANDER J. A. A. 2004. — MrModeltest v2. Program distributed by the author. Uppsala: Uppsala University. <https://github.com/nylander/MrModeltest2>.
- PIMENTEL M., SAHUQUILLO E., TORRECILLA Z., POPP M., CATALÁN P. & BROCHMANN P. 2013. — Hybridization and long-distance colonization at different time scales: towards resolution of long-term controversies in the sweet vernal grasses (*Anthoxanthum*). *Annals of Botany* 112: 1015-1030. <https://doi.org/10.1093/aob/mct170>
- PIÑEIRO-CORBEIRA C., MAGGS C. A., RINDI F., BUNKER F., BALDOCK L. & DÍAZ-TAPIA P. 2020a. — Molecular assessment of the tribes Strebloladiaceae and Polysiphoniaceae (Rhodomelaceae, Rhodophyta) in the British Isles reveals new records and species that require taxonomic revision. *Cryptogamie, Algologie* 41: 55-72. <https://doi.org/10.5252/cryptogamie-algologie2020v41a8>
- PIÑEIRO-CORBEIRA C., VERBRUGGEN H. & DÍAZ-TAPIA P. 2020b. — Molecular survey of the red algal family Rhodomelaceae (Ceramiales, Rhodophyta) in Australia reveals new introduced species. *Journal of Applied Phycology* 32: 2535-2547. <https://doi.org/10.1007/s10811-019-01932-4>
- PRICE I. R. & SCOTT F. J. 1992. — *The turf algal flora of the Great Barrier Reef. Part I. Rhodophyta*. Botany Department, James Cook University, Townsville, Australia, 266 p.
- RODRÍGUEZ-PRÍETO C., AFONSO-CARRILLO J., DE CLERCK O., HUISMAN J. M. & LIN S. M. 2020. — Systematic revision of the



- foliose Halymeniaceae (Halymeniales, Rhodophyta) from Europe, with the description of *Halymenia ballesterosii* sp. nov. from the Mediterranean Sea and *Nesoa hommersandii* from the Canary Islands. *European Journal of Phycology* 55: 454-466. <https://doi.org/10.1080/09670262.2020.1749888>
- ROJAS-GONZÁLEZ B. 1997. — *Estudio de las especies de la familia Rhodomelaceae (Rhodophyta), con exclusión de las tribus Chondrieae y Laurencieae, en las Islas Canarias*. Universidad de La Laguna, La Laguna, 647 p.
- RULL LLUCH J. R. 2002. — Marine benthic algae of Namibia. *Scientia Marina* 66 (Suppl 3): 1-258.
- SAVOIE A. M. & SAUNDERS G. W. 2015. — Evidence for the introduction of the Asian red alga *Neosiphonia japonica* and its introgression with *Neosiphonia harveyi* (Ceramiales, Rhodophyta) in the Northwest Atlantic. *Molecular Ecology* 24: 5927-5937. <https://doi.org/10.1111/mec.13429>
- SAVOIE A. M. & SAUNDERS G. W. 2016. — A molecular phylogenetic and DNA barcode assessment of the tribe Pterosiphoniae (Ceramiales, Rhodophyta) emphasizing the Northeast Pacific. *Botany* 94: 917-939.
- SAVOIE A. M. & SAUNDERS G. W. 2018. — A molecular assessment of species diversity and generic boundaries in the red algal tribes Polysiphoniae and Streblodactylidae (Rhodomelaceae, Rhodophyta) in Canada. *European Journal of Phycology* 54: 1-25. <https://doi.org/10.1080/09670262.2018.1483531>
- SAUNDERS G. W., HUISMAN J. M., VERGÉS A., KRAFT G. T. & LE GALL L. 2017. — Phylogenetic analyses support recognition of ten new genera, ten new species and 16 new combinations in the family Kallymeniaceae (Gigartinales, Rhodophyta). *Cryptogamie, Algologie* 38: 79-132. <https://doi.org/10.7872/cryalv38.iss2.2017.79>
- SOARES L. P., FUJII M. T., DIAZ-TAPIA P. & HOMMERSAND M. H. 2019. — A molecular and morphological reinvestigation of topotype specimens of *Calliblepharis occidentalis* (Cystocloniaceae, Rhodophyta). *Phycologia* 58: 169-179. <https://doi.org/10.1080/00318884.2018.1535741>
- SHAMEEL M. 1999. — *Melanothamnus afaqhusainii*, a new red alga from the coast of Karachi. *Pakistan Journal of Botany* 31: 211-214.
- SEGI T. 1951. — Systematic study of the genus *Polysiphonia* from Japan and its vicinity. *Journal of the Faculty of Fisheries, Prefectural University of Mie* 1: 169-272.
- SERIO D., FURNARI G., MORO I., & SCIUTO K. 2020. — Molecular and morphological characterisation of *Melanothamnus testudinis* sp. nov. (Rhodophyta, Rhodomelaceae) and its distinction from *Polysiphonia caretta*. *Phycologia* 59: 281-291. <https://doi.org/10.1080/00318884.2020.1752531>
- SWOFFORD D. L. 2000. — *PAUP\*: Phylogenetic analysis using parsimony*. Version 4.0. Washington DC, Smithsonian Institution
- STAMATAKIS A. 2014. — RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1312-1313.
- STUERCKE B. & FRESHWATER D. W. 2008. — Consistency of morphological characters used to delimit *Polysiphonia sensu lato* species (Ceramiales, Florideophyceae): analyses of North Carolina, USA specimens. *Phycologia* 47: 541-559. <https://doi.org/10.2216/08-09.1>
- STUERCKE B. & FRESHWATER D. W. 2010. — Two new species of *Polysiphonia* (Ceramiales, Florideophyceae) from the western Atlantic. *Botanica Marina* 53: 301-311. <https://doi.org/10.1515/BOT.2010.036>
- THIERS B. 2021. — *Index Herbariorum*: A Global Directory of Public Herbaria and Associated Staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/> (Accessed on 01 Feb. 2021).
- TORRECILLA P., LÓPEZ RODRÍGUEZ J.A., STANCIK D. & CATALÁN P. 2003. — Systematics of *Festuca* L. sects. *Eskia* Willk., *Pseudotropis* Kriv., *Amphigenes* (Janka) Tzvel., *Pseudoscariosa* Kriv. and *Scariosae* Hack. based on analysis of morphological characters and DNA sequences. 2003. *Plant Systematics and Evolution* 239: 113-139. <https://doi.org/10.1007/s00606-002-0265-2>
- TUYA F. & HAROUN R. 2006. — Spatial patterns and response to wave exposure of shallow water algal assemblages across the Canarian Archipelago: a multi-scaled approach. *Marine Ecology Progress Series* 311: 15-28.
- VAN-ÖPPEN M. J. H., KLERK H., OLSEN J. L., & STAM W. T. 1996. — Hidden diversity in marine algae: some examples of genetic variation below the species level. *Journal of the Marine Biological Association of the United Kingdom* 76: 239-242. <https://doi.org/10.1017/S0025315400029192>
- VERBRUGGEN H. 2014. — Morphological complexity, plasticity, and species diagnosability in the application of old species names in DNA-based taxonomies. *Journal of Phycology* 50: 26-31. <https://doi.org/10.1111/jpy.12155>
- WALLENSTEIN F. M., TERRA M. R., POMBO J. & NETO A. I. 2009. — Macroalgal turfs in the Azores. *Marine Ecology* 30: 113-117. <https://doi.org/10.1111/j.1439-0485.2009.00311.x>
- WOMERSLEY H. B. S. 1979. — Southern Australian species of *Polysiphonia* Greville (Rhodophyta). *Australian Journal of Botany* 27: 459-528. <https://doi.org/10.1071/BT9790459>
- WOODWORTH K. A., FRANKOVICH T. A. & FRESHWATER D. W. 2019. — *Melanothamnus maniticolus* sp. nov. (Ceramiales, Rhodophyta): an epizoic species evolved for living on the West Indian Manatee. *Journal of Phycology* 55: 1239-1245. <https://doi.org/10.1111/jpy.12912>
- WYNNE M. J. & BANAIMOON S. A. 1990. — The occurrence of *Jolyna laminarioides* (Phaeophyta) in the Arabian Sea and the Indian Ocean and a new report of *Melanothamnus somalensis* (Rhodophyta). *Botanica Marina* 33: 213-218. <https://doi.org/10.1515/botm.1990.33.2.213>
- YANG E. C., BOO S. B., BHATTACHARYA D., SAUNDERS G. W., KNOLL A. H., FREDERICQ S., GRAF L. & YOON H. S. 2016. — Divergence time estimates and the evolution of major lineages in the florideophyte red algae. *Scientific Reports* 6: 21361.
- YOON H. Y. 1986. — A taxonomic study of Genus *Polysiphonia* (Rhodophyta) from Korea. *The Korean Journal of Phycology* 1: 3-86.

Submitted on 2 November 2020;  
accepted on 18 February 2021;  
published on 12 May 2021.

APPENDICES

APPENDIX 1. — Materials collected during this study, including collection information and GenBank accession numbers.

Species	Haplotype (Fig. 2)	Collection information and herbarium vouchers for studied specimens	GenBank accession number
<i>Melanothamnus macaronesicus</i> Rodríguez-Buján & Díaz-Tapia, sp. nov.	H1	Puerto del Carmen, Lanzarote, Canary Islands; 11.VI.2018; SANT 33663; PD4111; 28°55'10"N, 13°39'47"W	MW567802
		Puerto del Carmen, Lanzarote, Canary Islands; 11.VI.2018; PD4116; 28°55'10"N, 13°39'47"W	MW567803
		Puerto del Carmen, Lanzarote, Canary Islands; 11.VI.2018; SANT 33664; PD4121; 28°55'10"N, 13°39'47"W	MW567804
		Puerto del Carmen, Lanzarote, Canary Islands; 11.VI.2018; SANT 33665; PD4129; 28°55'10"N, 13°39'47"W	MW567805
		Puerto del Carmen, Lanzarote, Canary Islands; 11.VI.2018; SANT 33666; PD4130; 28°55'10"N, 13°39'47"W	MW567801
	H2	Praia do Populo, São Miguel, Azores; 12.VI.2018; SANT 33649; PD3692; 37°45'00"N, 25°37'05"W	MW567800
	H3	Praia San Roque, São Miguel, Azores; 15.VI.2018; SANT 33656; PD3843; 37°45'00"N, 25°37'05"W	MW567798
		Praia San Roque, São Miguel, Azores; 15.VI.2018; SANT 33657; PD3847; 37°45'00"N, 25°37'05"W	MW567799

APPENDIX 2. — Sample information for the species included in the phylogenetic analysis.

Species	Publication	GenBank accession number
<i>Acanthosiphonia echinata</i> (Harvey) Savoie & G.W.Saunders	Mamoozadeh & Freshwater (2011), as <i>Polysiphonia</i>	HM573559
<i>Bryocladia cuspidata</i> (J.Agardh) De Toni	Lin <i>et al.</i> (2001)	AF259498
<i>Carradoriella denudata</i> (Dillwyn) Savoie & G.W.Saunders	Bárbara <i>et al.</i> (2013), as <i>Polysiphonia</i>	JX828147
<i>Carradoriella elongata</i> (Hudson) Savoie & G.W.Saunders	Bárbara <i>et al.</i> (2013), as <i>Polysiphonia</i>	JX828162
<i>Carradoriella elongella</i> (Harvey) Savoie & G.W.Saunders	Mclvor <i>et al.</i> (2001), as <i>Polysiphonia</i>	AF342913
<i>Carradoriella virgata</i> (C. Agardh) P.C.Silva	Bárbara <i>et al.</i> (2013), as <i>Polysiphonia</i>	JX828152
<i>Eutrichosiphonia sabulosia</i> (B.Kim & M.S.Kim) Savoie & G.W.Saunders	Kim & Kim (2014), as <i>Polysiphonia</i>	KF479250
<i>Kapraunia amplacapiilli</i> (B.Kim & M.S.Kim) Savoie & G.W.Saunders	Kim & Kim (2014), as <i>Polysiphonia</i>	KF479253
<i>Kapraunia morroides</i> (B.Kim & M.S.Kim) Savoie & G.W.Saunders	Kim & Kim (2014), as <i>Polysiphonia</i>	KF479257
<i>Kapraunia pentamera</i> (Hollenberg) Savoie & G.W.Saunders	Mamoozadeh & Freshwater (2011), as <i>Polysiphonia</i>	HM573564
<i>Kapraunia schneideri</i> (B.Stuercke & D.W.Freshwater) Savoie & G.W.Saunders	Stuercke & Freshwater (2010), as <i>Polysiphonia</i>	GU385836
<i>Lampisiphonia iberica</i> Bárbara, Secilla, Díaz Tapia & H.-G. Choi	Bárbara <i>et al.</i> (2013)	JX828131
<i>Leptosiphonia brodiei</i> (Dillwyn) Savoie & G.W.Saunders	Mclvor <i>et al.</i> (2001), as <i>Polysiphonia</i>	AF342916
<i>Leptosiphonia fibrata</i> (Dillwyn) Savoie & G.W.Saunders	Mclvor <i>et al.</i> (2001), as <i>Polysiphonia</i>	AF342915
<i>Leptosiphonia fibrillosa</i> (Dillwyn) Savoie & G.W.Saunders	Mclvor <i>et al.</i> (2001), as <i>Polysiphonia</i>	AF342912
<i>Leptosiphonia schousboei</i> (Thuret) Kylin	Bárbara <i>et al.</i> (2013)	JX828133
<i>Melanothamnus afaqhusainii</i> Bornet & Falkenberg	Savoie & Saunders (2016)	KU564460
<i>Melanothamnus bajacali</i> (Hollenberg) Díaz-Tapia & Maggs	Mamoozadeh & Freshwater (2011), as <i>Neosiphonia</i>	HM573572
<i>Melanothamnus balianus</i> (D.E.Bustamante, B.Y.Won & T.O.Cho) Díaz-Tapia & Maggs	Bustamante <i>et al.</i> (2013b), as <i>Neosiphonia</i>	KF146879
<i>Melanothamnus blandii</i> (Harvey) Díaz-Tapia & Maggs	Díaz-Tapia <i>et al.</i> (2017a), as <i>Polysiphonia</i>	KX499556
<i>Melanothamnus collabens</i> (C.Agardh) Díaz-Tapia & Maggs	Bárbara <i>et al.</i> (2013), as <i>Streblocladia</i>	JX828157
<i>Melanothamnus ferulaceus</i> (Suhr ex J.Agardh) Díaz-Tapia & Maggs	Mamoozadeh & Freshwater (2011), as <i>Neosiphonia</i>	HM573584
<i>Melanothamnus flavimarinus</i> (M.-S.Kim & I.K.Lee) Díaz-Tapia & Maggs	Kim & Yang (2006), as <i>Neosiphonia</i>	DQ787481
<i>Melanothamnus forfex</i> (Harvey) Díaz-Tapia & Maggs	Díaz-Tapia <i>et al.</i> (2017a), as <i>Polysiphonia</i>	KX499565
<i>Melanothamnus hancockii</i> (E.Y.Dawson) Díaz-Tapia & Maggs	Díaz-Tapia <i>et al.</i> (2017a), as <i>Polysiphonia</i>	KX499548
<i>Melanothamnus harveyi</i> (J. Bailey) Díaz-Tapia & Maggs	Bárbara <i>et al.</i> (2013), as <i>Polysiphonia</i>	JX828134
<i>Melanothamnus incomptus</i> (Harvey) Díaz-Tapia & Maggs	This study, collected in Knysna, Western Cape, South Africa by J. Costa and K. Dixon in December 2014	MW567797
<i>Melanothamnus cf. minutissimus</i> (Hollenberg) Díaz-Tapia & Maggs	Díaz-Tapia <i>et al.</i> (2017a), as <i>Polysiphonia</i>	KX499557
<i>Melanothamnus nudus</i> (Mamoozadeh & Freshwater) Freshwater	Mamoozadeh & Freshwater (2012), as <i>Polysiphonia</i>	HM573571
<i>Melanothamnus peruviansis</i> (D.E.Bustamante, B.Y.Won, M.E.Ramirez & T.O.Cho) Díaz-Tapia & Maggs	Bustamante <i>et al.</i> (2012), as <i>Neosiphonia</i>	JN989969
<i>Melanothamnus pseudofoeniculatus</i> Díaz-Tapia	Díaz-Tapia <i>et al.</i> (2017b)	KY620089
<i>Melanothamnus pseudovillum</i> (Hollenberg) Díaz-Tapia & Maggs	Mamoozadeh & Freshwater (2011), as <i>Polysiphonia</i>	HM573568
<i>Melanothamnus savatieri</i> (Hariot) Díaz-Tapia & Maggs	Díaz-Tapia <i>et al.</i> (2017a), as <i>Polysiphonia</i>	KX499558
<i>Melanothamnus silvae</i> (D.E.Bustamante, B.Y.Won & T.O.Cho) Díaz-Tapia & Maggs	Bustamante <i>et al.</i> (2013b), as <i>Neosiphonia</i>	KF146878

APPENDIX 2. — Continuation.

Species	Publication	GenBank accession number
<i>Melanothamnus simplex</i> (Hollenberg) Díaz-Tapia & Maggs	Mclvor <i>et al.</i> (2001), as <i>Polysiphonia</i>	AF342909
<i>Melanothamnus somalensis</i> Bornet & Falkenberg	Díaz-Tapia <i>et al.</i> (2017a)	KX499555
<i>Melanothamnus</i> sp.	Carlile (2009), as <i>Polysiphonia</i>	GQ252567
<i>Melanothamnus</i> sp.	Mamoozadeh & Freshwater (2011), as <i>Neosiphonia</i>	HM573573
<i>Melanothamnus</i> sp.	Díaz-Tapia <i>et al.</i> (2017a), as <i>Polysiphonia</i>	KX499561
<i>Melanothamnus</i> sp.	Díaz-Tapia <i>et al.</i> (2017a), as <i>Polysiphonia</i>	KX499562
<i>Melanothamnus sphaerocarpus</i> (Børgesen) Díaz-Tapia & Maggs	Mamoozadeh & Freshwater (2011), as <i>Neosiphonia</i>	HM573569
<i>Melanothamnus strictissimus</i> (J.D.Hooker & Harvey) Díaz-Tapia & Maggs	Stuercke & Freshwater (2010), as <i>Polysiphonia</i>	GU385833
<i>Melanothamnus teradomariensis</i> (M.Noda) Díaz-Tapia & Maggs	Bárbara <i>et al.</i> (2013), as <i>Neosiphonia</i>	JX828136
<i>Melanothamnus thailandicus</i> (N.Muangmai & C.Kaewsuralikhit) Díaz-Tapia & Maggs	Muangmai <i>et al.</i> (2014), as <i>Neosiphonia</i>	KM502785
<i>Melanothamnus tongatensis</i> (Harvey ex Kützing) Díaz-Tapia & Maggs	Mamoozadeh & Freshwater (2011), as <i>Neosiphonia</i>	HM573570
<i>Melanothamnus unilateralis</i> (Levring) Díaz-Tapia & Maggs	Díaz-Tapia <i>et al.</i> (2017a), as <i>Fernandosiphonia</i>	KX499549
<i>Melanothamnus yendoii</i> (Segi) Díaz-Tapia & Maggs	Bárbara <i>et al.</i> (2013), as <i>Neosiphonia</i>	JX828137
<i>Melanothamnus yongpili</i> (B.Kim & M.-S.Kim) Díaz-Tapia & Maggs	Kim & Kim (2014), as <i>Polysiphonia</i>	KT964456
<i>Polysiphonia atlantica</i> Kapraun & J.N. Norris	Bárbara <i>et al.</i> (2013)	JX828141
<i>Polysiphonia binneyi</i> Harvey	Mamoozadeh & Freshwater (2011)	HM573554
<i>Polysiphonia delicata</i> Díaz-Tapia	Díaz-Tapia <i>et al.</i> (2017b)	KY620062
<i>Polysiphonia havanensis</i> Montagne	Mamoozadeh & Freshwater (2011)	HM573554
<i>Polysiphonia kapraunii</i> B.Stuercke & D.W.Freshwater	Stuercke & Freshwater (2008)	EU492920
<i>Polysiphonia koreana</i> D.Bustamante, B.Y.Won & T.O.Cho	Bustamante <i>et al.</i> (2015)	KJ957811
<i>Polysiphonia macrocarpa</i> (C.Agardh) Sprengel	Mamoozadeh & Freshwater (2011)	HM573545
<i>Polysiphonia morrowii</i> Harvey	D'Archino <i>et al.</i> (2013)	KC152488
<i>Polysiphonia pacifica</i> Hollenberg	Kim <i>et al.</i> (2005)	AY958162
<i>Polysiphonia scopulorum</i> Harvey	Bárbara <i>et al.</i> (2013), as <i>P. caespitosa</i>	JX828149
<i>Polysiphonia</i> sp.	Mamoozadeh & Freshwater (2011, as <i>P. pernacola</i> )	HM573576
<i>Polysiphonia stricta</i> (Dillwyn) Greville	Bárbara <i>et al.</i> (2013)	JX828151
<i>Polysiphonia subtilissima</i> Montagne	Lam <i>et al.</i> (2013)	JX294918
<i>Streblocladia glomerulata</i> (Montagne) Papenfuss	Díaz-Tapia <i>et al.</i> (2017a)	KX499569
<i>Streblocladia muelleriana</i> J.Agardh	Fujii <i>et al.</i> (2006), as <i>Polysiphonia</i>	AY588412
<i>Symphyocladia latiuscula</i> (Harvey) Yamada	Kim <i>et al.</i> (2010)	GQ867072
<i>Symphyocladia linearis</i> (Okamura) Falkenberg	Bárbara <i>et al.</i> (2013)	JX828158
<i>Symphyocladia lithophila</i> M.-S.Kim	Kim <i>et al.</i> (2010)	GQ867078
<i>Vertebrata australis</i> (C.Agardh) Kuntze	Díaz-Tapia <i>et al.</i> (2017a)	KX499546
<i>Vertebrata byssoides</i> (Goodenough & Woodward) Kuntze	Yang <i>et al.</i> (2016), as <i>Brongniartella</i>	DQ787584
<i>Vertebrata constricta</i> (Womersley) Díaz-Tapia & Maggs	Stuercke & Freshwater (2010), as <i>Polysiphonia</i>	GU385832
<i>Vertebrata foetidissima</i> (Cocks ex Bornet) Díaz-Tapia & Maggs	Díaz-Tapia <i>et al.</i> (2013), as <i>Polysiphonia</i>	JQ653284
<i>Vertebrata fucoides</i> (Hudson) Kuntze	Bárbara <i>et al.</i> (2013), as <i>Polysiphonia</i>	JX828163
<i>Vertebrata hypnoides</i> (Welwitsch ex J.Agardh) Kuntze	Díaz-Tapia <i>et al.</i> (2017a)	KX499547
<i>Vertebrata lanosa</i> (Linnaeus) T. Christensen	Stuercke & Freshwater (2008)	EU492914
<i>Vertebrata lobophoralis</i> (N.R.Mamoozadeh & D.W.Freshwater)	Mamoozadeh & Freshwater (2011), as <i>Polysiphonia</i>	HM573551
<i>Vertebrata martensiana</i> (Kützing) Piñeiro-Corbeira, Maggs & Díaz-Tapia	Bárbara <i>et al.</i> (2013), as <i>Boergeseniella fruticulosa</i>	JX828161
<i>Vertebrata nigra</i> (Hudson) Díaz-Tapia & Maggs	Bárbara <i>et al.</i> (2013), as <i>Polysiphonia</i>	JX828164
<i>Vertebrata reptabunda</i> (Suhr) Díaz-Tapia & Maggs	Díaz-Tapia <i>et al.</i> (2017a)	KX499554
<i>Vertebrata simulans</i> (Harvey) Kuntze	Díaz-Tapia <i>et al.</i> (2017a)	KX499568
<i>Vertebrata subulifera</i> (C.Agardh) Kuntze	Díaz-Tapia <i>et al.</i> (2017a)	KX499564
<i>Vertebrata thuyoides</i> (Harvey) Kuntze	Bárbara <i>et al.</i> (2013), as <i>Polysiphonia</i>	JX828125