



Title	A Taxonomic Catalogue of Japanese Nemerteans (Phylum Nemertea)
Author(s)	Kajihara, Hiroshi
Citation	Zoological Science, 24(4), 287-326 https://doi.org/10.2108/zsj.24.287
Issue Date	2007-04
Doc URL	http://hdl.handle.net/2115/39621
Rights	(c) Zoological Society of Japan / 本文献の公開は著者の意思に基づくものである
Type	article
Note	REVIEW
File Information	zsj24p287.pdf



[Instructions for use](#)

[REVIEW]

A Taxonomic Catalogue of Japanese Nemerteans (Phylum Nemertea)

Hiroshi Kajihara*

Department of Natural History Sciences, Faculty of Science, Hokkaido University,
Sapporo 060-0810, Japan

A literature-based taxonomic catalogue of the nemertean species (Phylum Nemertea) reported from Japanese waters is provided, listing 19 families, 45 genera, and 120 species as valid. Applications of the following species names to forms previously recorded from Japanese waters are regarded as uncertain: *Amphiporus cervicalis*, *Amphiporus depressus*, *Amphiporus lactifloreus*, *Cephalothrix filiformis*, *Cephalothrix linearis*, *Cerebratulus fuscus*, *Lineus vegetus*, *Lineus bilineatus*, *Lineus gesserensis*, *Lineus grubei*, *Lineus longifissus*, *Lineus mcintoshii*, *Nipponnemertes pulchra*, *Oerstedia venusta*, *Prostoma graecense*, and *Prostoma grande*. The identities of the taxa referred to by the following four nominal species require clarification through future investigations: *Cosmocephala japonica*, *Dicelis rubra*, *Dichilus obscurus*, and *Nareda serpentina*. The nominal species established from Japanese waters are tabulated. In addition, a brief history of taxonomic research on Japanese nemerteans is reviewed.

Key words: checklist, Pacific, classification, ribbon worm, Nemertinea

INTRODUCTION

The phylum Nemertea comprises about 1,200 species (Gibson, 1995). Nemerteans are distributed worldwide, mostly in marine benthic habitats, though some species have adapted to live in pelagic, freshwater, and land habitats. Nemerteans are basically carnivorous, feeding on small crustaceans, polychaetes, and mollusks (McDermott and Roe, 1985). They are distinct from other metazoans by possessing a unique organ, the proboscis, which is housed in a fluid-filled cavity, the rhynchocoel, and eversible when used in prey capture. The phylum is currently regarded as the sister taxon to the Neotrochozoa (comprising the Annelida, Echiura, Mollusca, and Sipuncula) (Jenner, 2004). Morphological characters supporting the close relationship between the Nemertea and Neotrochozoa include: 1) modified coelomic cavities derived by schizocoely (Turbeville, 1986) and lined by mesothelium, with at least some cells bearing rudimentary cilia (Turbeville and Ruppert, 1985; Turbeville, 1991, 2002); 2) a glialinterstitial cell system (Turbeville and Ruppert, 1985; Turbeville, 1991, 2002); and 3) the prototroch, a transitory larval structure consisting of a preblastoporal belt of specialized cells derived from the trophoblast cell lineage (Maslakova *et al.*, 2004a, b).

Despite considerable efforts by previous researchers, a number of undescribed nemerteans remain in Japanese waters, especially those in the southwestern part (Kajihara,

2001). The only recent listing of previously described Japanese species is the checklist of Crandall *et al.* (2002), but the relevant literature is scattered. The present catalogue, identifying 19 families, 45 genera, and 120 species so far reported from Japanese waters as valid, has been compiled to integrate this scattered nemertean literature and to point out taxonomic issues to be resolved for the species already described, in order to offer a perspective for future studies.

Generally, the ideal situation for taxonomic studies is that name-bearing type specimens for every nominal species are extant and available. The present study revealed, however, that type specimens are either unavailable or unlocated for 45 out of 101 nominal species established from Japanese waters (Tables 1, 2).

While modern nemertean taxonomy depends upon the examination of internal structures from serially sectioned material (Gibson, 1985), virtually all of the original descriptions made by Stimpson and Takakura lack such information (Stimpson, 1855, 1857; Takakura, 1898). Fortunately, most of these species can be identified by their external features, but their systematic position requires reappraisal based on their internal morphology. Since most of the type material of early researchers, viz., Stimpson, Takakura, and Yamaoka, is unavailable (Nishimura, 1992; Kajihara, 2004; see the following section, "Brief History..."), re-collection of the species established by them is essential to correctly determine taxonomic identity.

Throughout the text and tables, "Code" and "ICZN" refer to the International Code for Zoological Nomenclature and the International Commission on Zoological Nomenclature, respectively.

* Corresponding author. Phone: +81-11-706-2755;

Fax : +81-11-746-0862;

E-mail: kazi@sci.hokudai.ac.jp

Table 1. List of the nominal species established from Japanese waters arranged by their type locality from north to south. *Type specimen depository: FI Dr. Iwata's collection; LBM Lake Biwa Museum, Shiga, Japan; NHMW-EV Naturhistorisches Museum Wien, Evertebrata-Varia, Wien, Austria; U unavailable; ZIHU Hokkaido University Museum, Sapporo, Japan; ? unlocated.

Locality	Nominal species	Type specimen*	Further detailed locality and comments.
HOKKAIDO			
Rishiri Island	<i>Lineus fulvus</i> Iwata, 1954	FI	
	* <i>Oerstedia venusta</i> Iwata, 1954	FI	Also based on the material from Muroran; a <i>nomen dubium</i> .
Akkeshi	<i>Amphiporus antifuscus</i> Iwata, 1954	FI	
	<i>Amphiporus parvus</i> Yamaoka, 1940	U	
	<i>Cephalothrix notabilis</i> Iwata, 1954	FI	
	<i>Lineus spatiiosus</i> Iwata, 1954	FI	
	* <i>Micrura akkeshiensis</i> Yamaoka, 1940	U	Also based on the material from Abashiri.
	<i>Micrura magna</i> Yamaoka, 1940	U	Daikokujima Island.
	<i>Nemertellina yamaokai</i> Kajihara et al., 2000	ZIHU-1260	
	<i>Oerstedia polyorbis</i> Iwata, 1954	FI	
	* <i>Tetrastemma pinnatum</i> Iwata, 1954	FI	
	* <i>Tetrastemma stimpsoni</i> Chernyshev, 1992	U	Also based on the material from Abashiri.
	<i>Tubulanus ezoensis</i> Yamaoka, 1940	U	Daikokujima Island.
	<i>Zygonemertes glandulosa</i> Yamaoka, 1940	U	
	<i>Zygonemertes jamsteci</i> Kajihara, 2002	ZIHU-1928	
Abashiri	* <i>Micrura akkeshiensis</i> Yamaoka, 1940	U	Also based upon the material from Akkeshi.
	* <i>Tetrastemma stimpsoni</i> Chernyshev, 1992	U	Also based upon the material from Akkeshi.
Oshoro	<i>Amphiporus musculus</i> Iwata, 1954	FI	
	<i>Tetrastemma yamaokai</i> Iwata, 1954	FI	
	<i>Tetrastemma verinigrum</i> Iwata, 1954	FI	
	<i>Zygonemertes shintai</i> Kajihara, 2002	ZIHU-1296	
Shizunai	<i>Potamostoma shizunaiense</i> Kajihara et al., 2003	ZIHU-2037	The mouth of the River Shizunai.
Muroran	<i>Amphiporus regius</i> Iwata, 1954	FI	
	<i>Micrura uchidai</i> Yamaoka, 1940	U	Now <i>Nipponomicrura uchidai</i> .
	* <i>Oerstedia venusta</i> Iwata, 1954	FI	Also based on the material from Rishiri; a <i>nomen dubium</i> .
Hakodate	<i>Tatsnoskia depressa</i> Stimpson, 1857	U	A <i>nomen dubium</i> .
	<i>Tetrastemma stigmatum</i> Stimpson, 1857	U	
Further detail of locality not specified	<i>Cerebratulus bellus</i> Stimpson, 1857	U	
	<i>Cerebratulus fasciatus</i> Stimpson, 1857	U	Now <i>Micrura bella</i> .
HONSHŪ			
AOMORI PREFECTURE	<i>Emplectonema kandai</i> Kato, 1939	U	Asamushi.
IWATE PREFECTURE	<i>Diopsonemertes acanthocephala</i> Kajihara et al., 2001	ZIHU-1290	Ōtsuchi.
	<i>Ototyphlonemertes dolichobasis</i> Kajihara, 2007	ZIHU-3200	Ōtsuchi.
IBARAKI PREFECTURE	<i>Hinumanemertes kikuchii</i> Iwata, 1970	FI	Lake Hinuma.
	<i>Sacconemertella lutulenta</i> Iwata, 1970	FI	Lake Hinuma.
	<i>Sacconemertopsis olivifera</i> Iwata, 1970	FI	Lake Hinuma.
CHIBA PREFECTURE	<i>Carcinonemertes mitsukurii</i> Takakura, 1910	U	Among egg masses of <i>Eriocheir japonicus</i> (Decapoda) at the mouth of the River Minatogawa.
	<i>Coeia ijimai</i> Takakura, 1922	U	Tateyama; now <i>Hubrechtelea ijimai</i> .
	* <i>Lineus fuscoviridis</i> Takakura, 1898	U	Sunosaki, Tateyama, also based on the material from Misaki.
	* <i>Lineus mitellatus</i> Takakura, 1898	U	Sunosaki, Tateyama, also based on the material from Misaki; now <i>Notospermus geniculatus</i> .
	<i>Malacobdella japonica</i> Takakura, 1897	U	Kujūkuri, in the mantle cavity of <i>Mactra sachalinensis</i> (Bivalvia).
KANAGAWA PREFECTURE			
Misaki and its vicinity	<i>Carinella punctata</i> Takakura, 1898	U	Now <i>Tubulanus punctatus</i> .
	<i>Cerebratulus carnosus</i> Takakura, 1898	U	
	<i>Cerebratulus communis</i> Takakura, 1898	U	
	<i>Lineus alborostratus</i> Takakura, 1898	U	
	<i>Lineus bipunctatus</i> Takakura, 1898	U	
	<i>Lineus caputornatus</i> Takakura, 1898	U	
	* <i>Lineus fuscoviridis</i> Takakura, 1898	U	Also based on the material from Tateyama.
	* <i>Lineus mitellatus</i> Takakura, 1898	U	Also based on the material from Tateyama; now <i>Notospermus geniculatus</i> .
	<i>Lineus subcingulatus</i> Takakura, 1898	U	
	<i>Micrura dorsovittata</i> Takakura, 1898	U	
	<i>Micrura festiva</i> Takakura, 1898	U	Now <i>Micrura bella</i> .
Sagami Bay	<i>Amphiporus nagaiensis</i> Iwata, 1957	FI	Now <i>Sagaminemertes nagaiensis</i> .
	<i>Amphiporus parmiornatus</i> Iwata, 1957	FI	Now <i>Kameginemertes parmiornata</i> .

To be continued.

Tabel. 1. continued.

<i>Amphiporus redundcus</i> Iwata, 1957	FI	
<i>Amphiporus retrotrumidus</i> Iwata, 1957	FI	
<i>Cephalomastax brevis</i> Iwata, 1957	FI	
<i>Cerebratulus albocirculus</i> Iwata, 1957	FI	
<i>Cerebratulus formosus</i> Iwata, 1957	FI	
<i>Cerebratulus macrone</i> Hubrecht, 1887	?	
<i>Cerebratulus penniger</i> Iwata, 1957	FI	
<i>Cerebratulus superniger</i> Iwata, 1957	FI	
<i>Drepanophorus longiceps</i> Iwata, 1957	FI	
<i>Euborlasia proteres</i> Iwata, 1957	FI	
<i>Eupolia nipponensis</i> Hubrecht, 1887	?	Now <i>Baseodiscus nipponensis</i> .
<i>Micrura multinotata</i> Iwata, 1957	FI	
<i>Nectonemertes japonica</i> Foshay, 1912	?	
<i>Paranemertes plana</i> Iwata, 1957	FI	
<i>Pelagonemertes moseleyi</i> Bürger, 1895	?	
<i>Tetramys ramicerebrus</i> Iwata, 1957	FI	
SHIZUOKA PREFECTURE		
Shimoda		
<i>Amphiporus ogumai</i> Yamaoka, 1947	U	Now <i>Nipponnemertes ogumai</i> .
<i>Cosmocephalia japonica</i> Stimpson, 1857	U	Now <i>Amphiporus japonicus</i> , a nomen dubium.
<i>Emplectonema mitsuii</i> Yamaoka, 1947	U	
<i>Paranemertes katoi</i> Yamaoka, 1947	U	
<i>Polina cervicalis</i> Stimpson, 1857	U	A nomen dubium.
<i>Prostoma roseocephalum</i> Yamaoka, 1947	U	Now <i>Tetrastemma roseocephalum</i> .
Lake Hamanako		
<i>Callinera nishikawai</i> Kajihara, 2006	ZIHU-3133	
<i>Carinina plecta</i> Kajihara, 2006	ZIHU-3123	
<i>Hubrechtiella kimuraorum</i> Kajihara, 2006	ZIHU-3127	
MIE PREFECTURE		
<i>Uchidana parasita</i> Iwata, 1967	FI	In mantle cavities of <i>Mactra sulcatoria</i> in muddy sand at the mouth of Aikawa River.
WAKAYAMA PREFECTURE		
<i>Amphiporus insolitus</i> Iwata, 1954	FI	Kushimoto.
<i>Lineus cancelli</i> Iwata, 1954	FI	Shirahama.
<i>Nemertopsis mitellicola</i> Kajihara, 2007	ZIHU-3204	Shirahama.
SHIGA PREFECTURE		
<i>Prostoma ohmiense</i> Chernyshev et al., 1998	LBM	Lake Biwa.
HIROSHIMA PREFECTURE		
<i>Paralineopsis taki</i> Iwata, 1993	FI	Mukaishima.
<i>Stichostemma grandis</i> Ikeda, 1913	U	Hiroshima, a nomen dubium.
KYŪSHŪ and OKINAWA		
KUMAMOTO PREFECTURE		
Tomioka		
* <i>Micrura japonica</i> Iwata, 1952	FI	Also based on the material from Fukue.
<i>Paranemertes incola</i> Iwata, 1952	FI	
<i>Procephalothrix fasciculus</i> Iwata, 1952	FI	Now <i>Cephalothrix fasciculus</i> .
* <i>Tetrastemma insolens</i> Iwata, 1952	FI	Also based on the material from Fukue.
NAGASAKI PREFECTURE		
Fukue		
<i>Carinesta uchidai</i> Iwata, 1952	FI	
<i>Euborlasia gotoensis</i> Iwata, 1952	FI	
* <i>Micrura japonica</i> Iwata, 1952	FI	Also based on the material from Tomioka.
<i>Procephalothrix simula</i> Iwata, 1952	FI	Now <i>Cephalothrix simula</i> .
* <i>Tetrastemma insolens</i> Iwata, 1952	FI	Also based on the material from Tomioka.
<i>Tubulanus lucidus</i> Iwata, 1952	FI	
KAGOSHIMA PREFECTURE		
Kagoshima Bay		
<i>Diplopleura japonica</i> Stimpson, 1857	U	Kagoshima Bay.
Tanegashima		
<i>Dicelis rubra</i> Stimpson, 1857	U	Tanegashima, a nomen dubium.
Nakanoshima		
<i>Lineus nigrostriatus</i> Iwata, 1954	FI	Nakanoshima, Tokara Islands.
Amamiōshima		
<i>Cerebratulus nigrofuscus</i> Stimpson, 1857	U	Amamiōshima, now <i>Lineus nigrofuscus</i> .
<i>Dichilus obscurus</i> Stimpson, 1857	U	Amamiōshima, a nomen dubium.
<i>Taeniosoma aequale</i> Stimpson, 1857	U	Amamiōshima, now <i>Baseodiscus quinquelineatus</i> .
Kikajima		
<i>Meckelia piperata</i> Stimpson, 1855	U	Kikajima, now <i>Iwatanemertes piperata</i> .
OKINAWA PREFECTURE		
<i>Meckelia albovittata</i> Stimpson, 1855	U	Now <i>Lineus albovittatus</i> .
<i>Meckelia subacuta</i> Stimpson, 1857	U	Naha, a nomen dubium.
<i>Nareda serpentina</i> Stimpson, 1855	U	A nomen dubium.
<i>Prostoma specutaculum</i> Yamaoka, 1940	U	Naha and Chinen; now <i>Pantinonemertes specutacula</i> .
TYPE LOCALITY UNKNOWN		
<i>Lineus nipponensis</i> Senz, 2001	NHMW-EV 17026/3990	
<i>Tubulanus roretzi</i> Senz, 1997	NHMW-EV 3565/1886	

Table 2. List of the nominal species established from Japanese waters arranged taxonomically. *Type specimen depository: FI Professor Fumio Iwata's collection; LBM Lake Biwa Museum, Shiga, Japan; NHMW-EV Naturhistorisches Museum Wien, Evertebrata-Varia, Wien, Austria; U unavailable; ZIHU Hokkaido University Museum, Sapporo, Japan; ? unlocated.

Higer Taxa	Nominal species	Type specimen*	Type locality and comments.
PALAEONEMERTEA			
	<i>Callinera nishikawai</i> Kajihara, 2006	ZIHU-3133	Lake Hamanako, Shizuoka Prefecture.
	<i>Carinella punctata</i> Takakura, 1898	U	Misaki, Kanagawa Prefecture; now <i>Tubulanus punctatus</i> .
	<i>Carinesta uchidai</i> Iwata, 1952	FI	Fukue, Nagasaki Prefecture.
	<i>Carinina plecta</i> Kajihara, 2006	ZIHU-3123	Lake Hamanako, Shizuoka Prefecture.
	<i>Cephalothrix notabilis</i> Iwata, 1954	FI	Akkeshi, Hokkaidō Prefecture.
	<i>Procephalothrix fasciculus</i> Iwata, 1952	FI	Tomioka, Kumamoto Prefecture; now <i>Cephalothrix fasciculus</i> .
	<i>Procephalothrix simula</i> Iwata, 1952	FI	Fukue, Nagasaki Prefecture; now <i>Cephalothrix simula</i> .
	<i>Tubulanus ezoensis</i> Yamaoka, 1940	U	Daikokujima Island, Akkeshi, Hokkaidō Prefecture.
	<i>Tubulanus lucidus</i> Iwata, 1952	FI	Fukue, Nagasaki Prefecture.
	<i>Tubulanus roretzi</i> Senz, 1997	NHMW-EV 3565/1886	Precise type locality unknown.
PILIDIOPHORA			
	<i>Cephalomastax brevis</i> Iwata, 1957	FI	Sagami Bay, Kanagawa Prefecture.
	<i>Cerebratulus albocirculus</i> Iwata, 1957	FI	Sagami Bay, Kanagawa Prefecture.
	<i>Cerebratulus bellus</i> Stimpson, 1857	U	Hokkaidō; now <i>Micrura bella</i> .
	<i>Cerebratulus carnosus</i> Takakura, 1898	U	Misaki, Kanagawa Prefecture.
	<i>Cerebratulus communis</i> Takakura, 1898	U	Misaki, Kanagawa Prefecture.
	<i>Cerebratulus fasciatus</i> Stimpson, 1857	U	Hokkaidō.
	<i>Cerebratulus formosus</i> Iwata, 1957	FI	Sagami Bay, Kanagawa Prefecture.
	<i>Cerebratulus macrone</i> Hubrecht, 1887	?	Sagami Bay, Kanagawa Prefecture.
	<i>Cerebratulus nigrofuscus</i> Stimpson, 1857	U	Amamiōshima, Kagoshima Prefecture; now <i>Lineus nigrofuscus</i> .
	<i>Cerebratulus penniger</i> Iwata, 1957	FI	Sagami Bay, Kanagawa Prefecture.
	<i>Cerebratulus superniger</i> Iwata, 1957	FI	Sagami Bay, Kanagawa Prefecture.
	<i>Coeia ijimai</i> Takakura, 1922	U	Tateyama, Chiba Prefecture; now <i>Hubrechtella ijimai</i> .
	<i>Diplopleura japonica</i> Stimpson, 1857	U	Kagoshima Bay, Kagoshima Prefecture.
	<i>Euborlasia gotoensis</i> Iwata, 1952	FI	Fukue, Nagasaki Prefecture.
	<i>Euborlasia proteres</i> Iwata, 1957	FI	Sagami Bay, Kanagawa Prefecture.
	<i>Eupolia nipponensis</i> Hubrecht, 1887	?	Sagami Bay, Kanagawa Prefecture; now <i>Baseodiscus nipponensis</i> .
	<i>Hinumanemertes kikuchi</i> Iwata, 1970	FI	Lake Hinuma, Ibaraki Prefecture.
	<i>Hubrechtella kimuraorum</i> Kajihara, 2006	ZIHU-3127	Lake Hamanako, Shizuoka Prefecture.
	<i>Lineus alborostratus</i> Takakura, 1898	U	Misaki, Kanagawa Prefecture.
	<i>Lineus bipunctatus</i> Takakura, 1898	U	Misaki, Kanagawa Prefecture.
	<i>Lineus cancelli</i> Iwata, 1954	FI	Shirahama, Wakayama Prefecture.
	<i>Lineus caputornatus</i> Takakura, 1898	U	Misaki, Kanagawa Prefecture.
	<i>Lineus fulvus</i> Iwata, 1954	FI	Rishiri Island, Hokkaidō Prefecture.
	<i>Lineus fuscoviridis</i> Takakura, 1898	U	Sunosaki, Tateyama, Chiba Prefecture; Misaki, Kanagawa Prefecture.
	<i>Lineus mitellatus</i> Takakura, 1898	U	Sunosaki, Tateyama, Chiba Prefecture; Misaki, Kanagawa Prefecture; now <i>Notospermus geniculatus</i> .
	<i>Lineus nigrostriatus</i> Iwata, 1954	FI	Nakanoshima, Tokara Islands, Kagoshima Prefecture.
	<i>Lineus nipponensis</i> Senz, 2001	NHMW-EV 17026/3990	Precise type locality unknown.
	<i>Lineus spatirosus</i> Iwata, 1954	FI	Akkeshi, Hokkaidō Prefecture.
	<i>Lineus subcingulatus</i> Takakura, 1898	U	Misaki, Kanagawa Prefecture.
	<i>Meckelia albovittata</i> Stimpson, 1855	U	Okinawa; now <i>Lineus albovittatus</i> .
	<i>Meckelia piperata</i> Stimpson, 1855	U	Kikajima, Kagoshima Prefecture; now <i>Iwatanelemertes piperata</i> .
	<i>Meckelia subacuta</i> Stimpson, 1857	U	Naha, Okinawa; <i>nomen dubium</i> .
	<i>Micrura akkeshiensis</i> Yamaoka, 1940	U	Abashiri and Akkeshi, Hokkaidō Prefecture.
	<i>Micrura dorsovittata</i> Takakura, 1898	U	Misaki, Kanagawa Prefecture.
	<i>Micrura festiva</i> Takakura, 1898	U	Misaki, Kanagawa Prefecture; now <i>Micrura bella</i> .
	<i>Micrura japonica</i> Iwata, 1952	FI	Tomioka, Kumamoto Prefecture; Fukue, Nagasaki Prefecture.
	<i>Micrura magna</i> Yamaoka, 1940	U	Daikokujima Island, Akkeshi, Hokkaidō Prefecture.
	<i>Micrura multinotata</i> Iwata, 1957	FI	Sagami Bay, Kanagawa Prefecture.
	<i>Micrura uchidai</i> Yamaoka, 1940	U	Muroran, Hokkaidō Prefecture; now <i>Nipponomicrura uchidai</i> .
	<i>Paralineopsis taki</i> Iwata, 1993	FI	Mukaishima, Hiroshima Prefecture.
	<i>Taeniosoma aequale</i> Stimpson, 1857	U	Amamiōshima, Kagoshima Prefecture; now <i>Baseodiscus quinquelineatus</i> .
	<i>Tetramys ramicerebrus</i> Iwata, 1957	FI	Sagami Bay, Kanagawa Prefecture.
	<i>Uchidana parasita</i> Iwata, 1967	FI	Mouth of Aikawa River, Mie Prefecture.
HOPLONEMERTEA			
	<i>Amphiporus antifuscus</i> Iwata, 1954	FI	Akkeshi, Hokkaidō Prefecture.
	<i>Amphiporus insolitus</i> Iwata, 1954	FI	Kushimoto, Wakayama Prefecture.
	<i>Amphiporus musculus</i> Iwata, 1954	FI	Oshoro, Hokkaidō Prefecture.
	<i>Amphiporus nagaiensis</i> Iwata, 1957	FI	Sagami Bay, Kanagawa Prefecture; now <i>Sagaminemertes nagaiensis</i> .
	<i>Amphiporus ogumai</i> Yamaoka, 1947	U	Shimoda, Shizuoka Prefecture; now <i>Nipponnemertes ogumai</i> .
	<i>Amphiporus parmiornatus</i> Iwata, 1957	FI	Sagami Bay, Kanagawa Prefecture; now <i>Kameginemertes parmiornata</i> .

To be continued.

Table 2. continued.

<i>Amphiporus parvus</i> Yamaoka, 1940	U	Akkeshi, Hokkaidō Prefecture.
<i>Amphiporus reducens</i> Iwata, 1957	FI	Sagami Bay, Kanagawa Prefecture.
<i>Amphiporus regius</i> Iwata, 1954	FI	Muroran, Hokkaidō Prefecture.
<i>Amphiporus retrotumidus</i> Iwata, 1957	FI	Sagami Bay, Kanagawa Prefecture.
<i>Carcinonemertes mitsukurii</i> Takakura, 1910	U	Mouth of River Minatogawa, Chiba Prefecture.
<i>Cosmocephala japonica</i> Stimpson, 1857	U	Shimoda, Shizuoka Prefecture; <i>nomen dubium</i> .
<i>Dicelis rubra</i> Stimpson, 1857	U	Tanegashima Island, Kagoshima Prefecture; <i>nomen dubium</i> .
<i>Diopsonemertes acanthocephala</i> Kajihara et al., 2001	ZIHU-1290	Ōtsuchi, Iwate Prefecture.
<i>Drepanophorus longiceps</i> Iwata, 1957	FI	Sagami Bay, Kanagawa Prefecture.
<i>Emplectonema kandai</i> Kato, 1939	U	Asamushi, Aomori Prefecture.
<i>Emplectonema mitsuii</i> Yamaoka, 1947	U	Shimoda, Shizuoka Prefecture.
<i>Malacobdella japonica</i> Takakura, 1897	U	Kujūkuri, Chiba Prefecture.
<i>Nareda serpentina</i> Stimpson, 1855	U	Okinawa Prefecture; <i>nomen dubium</i> .
<i>Nectonemertes japonica</i> Foshay, 1912	?	Sagami Bay, Kanagawa Prefecture.
<i>Nemertellina yamaokai</i> Kajihara et al., 2000	ZIHU-1260	Akkeshi, Hokkaidō Prefecture.
<i>Nemertopsis mitellicola</i> Kajihara, 2007	ZIHU-3204	Shirahama, Wakayama Prefecture.
<i>Oerstedia polyorbis</i> Iwata, 1954	FI	Akkeshi, Hokkaidō Prefecture.
<i>Oerstedia venusta</i> Iwata, 1954	FI	Rishiri Island and Muroran, Hokkaidō Prefecture; <i>nomen dubium</i> .
<i>Ototyphlonemertes dolichobasis</i> Kajihara, 2007	ZIHU-3200	Ōtsuchi, Iwate Prefecture.
<i>Paranemertes incola</i> Iwata, 1952	FI	Tomioka, Kumamoto Prefecture.
<i>Paranemertes katoi</i> Yamaoka, 1947	U	Shimoda, Shizuoka Prefecture.
<i>Paranemertes plana</i> Iwata, 1957	FI	Sagami Bay, Kanagawa Prefecture.
<i>Pelagonemertes moseleyi</i> Bürger, 1895	?	Sagami Bay, Kanagawa Prefecture.
<i>Polina cervicalis</i> Stimpson, 1857	U	Shimoda, Shizuoka Prefecture; <i>nomen dubium</i> .
<i>Potamostoma shizunaiense</i> Kajihara et al., 2003	ZIHU-2037	The mouth of the River Shizunai, Hokkaidō Prefecture.
<i>Prostoma ohmienne</i> Chernyshev et al., 1998	LBM	Lake Biwa, Shiga Prefecture.
<i>Prostoma roseocephalum</i> Yamaoka, 1947	U	Shimoda, Shizuoka Prefecture; now <i>Tetrastemma roseocephalum</i> .
<i>Prostoma specutaculum</i> Yamaoka, 1940	U	Naha and Chinen, Okinawa; now <i>Pantinonemertes specutacula</i> .
<i>Sacconemertella lutulenta</i> Iwata, 1970	FI	Lake Hinuma, Ibaraki Prefecture.
<i>Sacconemertopsis olivifera</i> Iwata, 1970	FI	Lake Hinuma, Ibaraki Prefecture.
<i>Stichostemma grandis</i> Ikeda, 1913	U	Hiroshima Prefecture; <i>nomen dubium</i> .
<i>Tatsnoskia depressa</i> Stimpson, 1857	U	Hakodate, Hokkaidō Prefecture; <i>nomen dubium</i> .
<i>Tetrastemma insolens</i> Iwata, 1952	FI	Tomioka, Kumamoto Prefecture; Fukue, Nagasaki Prefecture.
<i>Tetrastemma pinnatum</i> Iwata, 1954	FI	Akkeshi, Hokkaidō Prefecture.
<i>Tetrastemma stigmatum</i> Stimpson, 1857	U	Hakodate, Hokkaidō Prefecture.
<i>Tetrastemma stimpsoni</i> Chernyshev, 1992	U	Abashiri and Akkeshi, Hokkaidō Prefecture.
<i>Tetrastemma yamaokai</i> Iwata, 1954	FI	Oshoro, Hokkaidō Prefecture.
<i>Tetrastemma verinigrum</i> Iwata, 1954	FI	Oshoro, Hokkaidō Prefecture.
<i>Zygonemertes glandulosa</i> Yamaoka, 1940	U	Akkeshi, Hokkaidō Prefecture.
<i>Zygonemertes jamsteci</i> Kajihara, 2002	ZIHU-1928	Akkeshi, Hokkaidō Prefecture.
<i>Zygonemertes shintai</i> Kajihara, 2002	ZIHU-1296	Oshoro, Hokkaidō Prefecture.
INCERTAE CEDIS		
<i>Dichilus obscurus</i> Stimpson, 1857	U	Amamiōshima, Kagoshima Prefecture; <i>nomen dubium</i> .

Brief History of Taxonomic Research on Japanese Nemerteans

William Stimpson (1832–1872) first reported nemerteans from Japanese waters (Stimpson, 1855). During the cruise of the North Pacific Exploring Expedition (1852–1856), in which Stimpson participated as a naturalist in zoology at the age of 21, he established 15 nominal species of nemerteans from Naha (Okinawa), Kikaishima (Kagoshima), Amamiōshima (Kagoshima), Shimoda (Shizuoka), and Hakodate (Hokkaidō) (Stimpson, 1855, 1857). Regrettably, his nemertean samples were lost in the Great Chicago Fire of 1871, together with his manuscripts, drawings, and other invertebrate collections obtained by the expedition (Nishimura, 1992). Even though Stimpson's descriptions are brief and composed of only a few lines, eight of 15 species of nemerteans he described from Japanese waters are still regarded as valid.

Dr. Albrecht von Roretz (1846–1884), a medical doctor who graduated from the University of Wien, came to Japan towards the end of 1874 as Consultats-Arzt for the Austro-Hungarian embassy in East Asia. From 1875 to 1882, when

Roretz returned to Austria, he made sampling trips to the four major islands in Japan, namely Hokkaidō, Honshū, Shikoku, and Kyūshū. Roretz's Japanese animal collection, now deposited in the Naturhistorisches Museum Wien, Austria, consists of more than 1,450 individuals ranging from sponges to mammals and comprises about 360 species (Nishikawa and Sattmann, 2001). Senz (1997a, 2001) recently described two new species based on nemertean specimens in Roretz's collection.

In 1875 the British Naval research vessel H.M.S. *Challenger* dropped into Japanese harbors in the course of her round-the-world scientific voyage, making collections at about 10 locations in Japanese waters (Tizard et al., 1885). Ambrosius Arnold Willem Hubrecht (1853–1915), Professor of Zoology at Utrecht University, reported the nemerteans obtained during the cruise; his report includes two species of nemerteans collected from Japanese waters (Hubrecht, 1887). One of the major zoological findings made by the *Challenger* expedition is the discovery of bathypelagic nemerteans. The first specimen was found on 7 March 1874 near the southern verge of the South-Australian current and

named *Pelagonemertes rollestoni* by Moseley (1875a). A second specimen, collected on 5 June 1875 off Sagami Bay, was first considered to be a young individual of the same species (Moseley, 1875b), but was later deemed to represent a different species and named *Pelagonemertes moseleyi* by Bürger (1895).

In 1906 the United States Fisheries Commission Steamer *Albatross* visited Japanese waters, with the ichthyologist Charles Henry Gilbert as the Naturalist-in-Charge. The nemerteans among the numerous specimens collected were studied by Coe (1944) and included several new locality records for species already known.

In the same year, 1906, Dr. Harold Heath secured from Mr. Alan Owston six specimens of pelagic nemertean species, which Foshey (1912) later described as *Nectonemertes japonica*.

Usamaro Takakura (1867–1944), Professor of Zoology first at the Higher Normal School (later renamed Tokyo Higher Normal School), then concurrently at Tokyo University of Literature and Science, was the first Japanese expert on nemerteans. Takakura reported 25 species and established 14 new species and one new genus, based primarily upon material from the Pacific coast of Honshū (Takakura, 1897, 1898, 1910, 1922, 1933). Takakura's principal work is his 1898 paper, which contains descriptions of 21 anoplan nemerteans from the vicinity of Misaki. Takakura's nemertean collection appears to have been lost during the relocation of Tokyo University of Education from Tokyo to Tsukuba (Kajihara, 2004).

Teiichi Yamaoka (ca. 1918–1945?) carried out taxonomic studies on nemerteans in his graduate studies under the guidance of Professor Tohru Uchida at Hokkaido Imperial University. After graduating in 1939, he became a researcher at Mitsui Marine Biological Station at Izu before he moved to the capital of Manchuria as a teacher at Shinkyō First Junior High School in 1940. Before he moved to Manchuria, he published two papers in which he reported 23 species (including seven new species) of nemerteans from Japanese waters (Yamaoka, 1940a, b). In addition, he reported two species from Taiwan, of which one was new to science, based on the specimens obtained by Dr. Shirō Okuda (Yamaoka, 1939). Yamaoka also published a paper on the entocommensal species *Malacobdella japonica* Takakura, 1897 in collaboration with Saburō Kawai (Kawai and Yamaoka, 1940) and wrote a chapter on nemerteans in a treatise on systematic zoology (Yamaoka, 1943). When Yamaoka had been a researcher at Mitsui Marine Biological Station, he prepared a manuscript that contained descriptions of several 'new' species, but the manuscript was not published before his death. Later, Dr. Okuda included four of these 'new' species in the *Revised Edition Illustrated Encyclopedia of the Fauna of Japan (Exclusive of Insects)*, with each of the four entries accompanied by a brief description. Crandall et al. (2001) argued that these four names are available in terms of the Code (ICZN, 1999), with Yamaoka as the naming authority and the date of publication 1947, the date when the *Encyclopedia* was published. Crandall et al. (2001) provided additional information on these four species in Yamaoka's unpublished manuscript, which had been in the care of Dr. Fumio Iwata. Quite recently, the manuscript was posthumously published (Yamaoka, 2005), sub-

mitted by Professor Iwata.

Much of our knowledge of the Japanese nemertean fauna depends upon the works by Dr. Fumio Iwata (b. 1925), Professor Emeritus of Hokkaido University. As with Yamaoka, Fumio Iwata began his nemertean studies under the guidance of Professor Uchida at Hokkaido University. After graduation in 1950, he was appointed as Assistant Professor at Akkeshi Marine Biological Station (Moriyama, 1995). He energetically investigated the nemertean fauna in various regions of Japan, as well as engaged in embryological studies on nemerteans. He published nine nemertean papers (Iwata, 1951, 1952, 1954a, b, c, 1957a, b, 1958, 1960a) before obtaining his doctorate in 1959 with a dissertation on the comparative embryology of nemerteans (Iwata, 1960b), in which he proposed the new order Archinemertea that he considered to be the most primitive group in the phylum. Although adopted by some researchers (e.g., Gibson, 1994), Iwata's (1960b) hypothesis was later questioned by Sundberg and Hylbom (1994), who found no morphological support for the Archinemertea. This group has lost acceptance among other nemertean researchers, since it is now regarded to be a group within the palaeonemerteans (Thollesson and Norenburg, 2003). However, Iwata's (1960b) embryological observations themselves are highly valued and frequently cited by modern researchers (e.g., Maslakova et al., 2004a, b; Nielsen, 2005). After receiving an Associate Professorship in Sapporo in 1963, Iwata wrote the chapter on nemerteans in a treatise on systematic zoology (Iwata, 1965a), descriptions of a parasitic nemertean in bivalves (Iwata, 1967) and three brackish-water nemerteans (Iwata, 1970a), another embryological paper (Iwata, 1972), and the chapter on nemerteans in a treatise on freshwater biology (Iwata, 1973). He was appointed to a Professorship in 1974. Iwata has attended all the international meetings on nemertean biology, held in 1983 (Philadelphia, USA), 1986 (Tjärnö, Sweden), 1991 (Bangor, UK), 1995 (Asilomar, California, USA), 2000 (Alcalá de Henares, Spain), and 2004 (Ogden, Utah, USA), and presented papers (Iwata, 1985, 1988, 1993, 2006). Even after retiring in 1988, Dr. Iwata is still quite active in research (Iwata, 2001).

Other taxonomic works on Japanese nemerteans include papers on freshwater nemerteans by Ikeda (1913), Ishizuka (1933), Sudzuki (1953), and Chernyshev et al. (1998); a report on the pelagic species *Pelagonemertes moseleyi* Bürger, 1895 by Kato and Tanaka (1938); Kato's (1939) description of the luminescent nemertean *Emplectonema kandai* Kato, 1939; Oki et al.'s (1987) report on the land nemertean *Geonemertes pelaensis* Semper, 1863; and reports on some marine benthic (Kajihara, 2002, 2006, 2007a, b; Kajihara et al., 2000, 2001) and a brackish-water (Kajihara et al., 2003) species.

Classification and Checklist of the Valid Japanese Nemertean Species

The higher classification system adopted here is based on Gibson (1982a, b, 1994), Chernyshev (1995, 2003), and Thollesson and Norenburg (2003).

Phylum NEMERTEA

- 1) Class PALAEONEMERTEA Hubrecht, 1879
- 1) Family CALLINERIDAE Bergendal, 1901

- 1) *Callinera nishikawai* Kajihara, 2006
 2) Family CEPHALOTRICHIDAE McIntosh, 1874
 2) *Cephalothrix fasciculus* (Iwata, 1952)
 3) *Cephalothrix notabilis* Iwata, 1954
 4) *Cephalothrix simula* (Iwata, 1952)
 3) Family TUBULANIDAE Bürger, 1904 (1874)
 5) *Carinesta uchidai* Iwata, 1952
 6) *Carinina plecta* Kajihara, 2006
 7) *Tubulanus capistratus* (Coe, 1901)
 8) *Tubulanus ezoensis* Yamaoka, 1940
 9) *Tubulanus lucidus* Iwata, 1952
 10) *Tubulanus punctatus* (Takakura, 1898)
 11) *Tubulanus roretzi* Senz, 1997
 2) Class PILIDIOPHORA Thollesson and Norenburg, 2003
 4) Family HUBRECHTELLIDAE Chernyshev, 2003
 12) *Hubrechtella ijimai* (Takakura, 1922)
 13) *Hubrechtella kimuraorum* Kajihara, 2006
 14) *Tetramys ramicerebrus* Iwata, 1957
 5) Family LINEIDAE McIntosh, 1874
 15) *Cerebratulus albocirculus* Iwata, 1957
 16) *Cerebratulus carnosus* Takakura, 1898
 17) *Cerebratulus communis* Takakura, 1898
 18) *Cerebratulus fasciatus* Stimpson, 1857
 19) *Cerebratulus formosus* Iwata, 1957
 20) *Cerebratulus longiceps* Coe, 1901
 21) *Cerebratulus macroren* Hubrecht, 1887
 22) *Cerebratulus marginatus* Renier, 1804
 23) *Cerebratulus montgomeryi* Coe, 1901
 24) *Cerebratulus penniger* Iwata, 1957
 25) *Cerebratulus subacutus* (Stimpson, 1857)
 26) *Cerebratulus superniger* Iwata, 1957
 27) *Cerebratulus zebra* Punnett and Cooper, 1909
 28) *Diplopleura japonica* Stimpson, 1857
 29) *Euborlasia gotoensis* Iwata, 1952
 30) *Euborlasia proteres* Iwata, 1957
 31) *Hinumanemertes kikuchii* Iwata, 1970
 32) *Iwatanelemertes piperata* (Stimpson, 1855)
 33) *Lineus alborostratus* Takakura, 1898
 34) *Lineus albovittatus* (Stimpson, 1855)
 35) *Lineus bipunctatus* Takakura, 1898
 36) *Lineus cancelli* Iwata, 1954
 37) *Lineus caputornatus* Takakura, 1898
 38) *Lineus fulvus* Iwata, 1954
 39) *Lineus fuscoviridis* Takakura, 1898
 40) *Lineus nigrofuscus* (Stimpson, 1857)
 41) *Lineus nigrostriatus* Iwata, 1954
 42) *Lineus nipponensis* Senz, 2001
 43) *Lineus spatirosus* Iwata, 1954
 44) *Lineus subcingulatus* Takakura, 1898
 45) *Lineus torquatus* Coe, 1901
 46) *Micrura akkeshiensis* Yamaoka, 1940
 47) *Micrura alaskensis* Coe, 1901
 48) *Micrura bella* (Stimpson, 1857)
 49) *Micrura dorsovittata* Takakura, 1898
 50) *Micrura japonica* Iwata, 1952
 51) *Micrura magna* Yamaoka, 1940
 52) *Micrura multinotata* Iwata, 1957
 53) *Nipponomicrura uchidai* (Yamaoka, 1940)
 54) *Notospermus geniculatus* (Delle Chiaje, 1828)
 55) *Paralineopsis taki* Iwata, 1993
 56) *Uchidana parasita* Iwata, 1967
 6) Family VALENCINIIDAE Hubrecht, 1879
 57) *Baseodiscus curtus* (Hubrecht, 1879)
 58) *Baseodiscus delineatus* (Delle Chiaje, 1825)
 59) *Baseodiscus hemprichii* (Ehrenberg, 1831)
 60) *Baseodiscus nipponensis* (Hubrecht, 1887)
 61) *Baseodiscus princeps* (Coe, 1901)
 62) *Baseodiscus quinque-lineatus* (Quoy and Gaimard, 1833)
 63) *Cephalomastax brevis* Iwata, 1957
 3) Class HOPLONEMERTEA Hubrecht, 1879
 1) Subclass MONOSTILIFERA Brinkmann, 1917
 7) Family AMPHIPORIDAE McIntosh, 1874
 64) *Amphiporus antifuscus* Iwata, 1954
 65) *Amphiporus formidabilis* Griffin, 1898
 66) *Amphiporus gelatinosus* Coe, 1905
 67) *Amphiporus imparispinosus* Griffin, 1898
 68) *Amphiporus insolitus* Iwata, 1954
 69) *Amphiporus musculus* Iwata, 1954
 70) *Amphiporus parvus* Yamaoka, 1940
 71) *Amphiporus reduncus* Iwata, 1957
 72) *Amphiporus regius* Iwata, 1954
 73) *Amphiporus retrotumidus* Iwata, 1957
 74) *Potamostoma shizunaiense* Kajihara, Gibson and Mawatari, 2003
 75) *Zygonemertes glandulosa* Yamaoka, 1940
 76) *Zygonemertes jamsteci* Kajihara, 2002
 77) *Zygonemertes shintai* Kajihara, 2002
 8) Family CARCINONEMERTIDAE Sumner, Osburn and Cole, 1913
 78) *Carcinonemertes mitsukurii* Takakura, 1910
 9) Family CRATENEMERTIDAE Friedrich, 1968
 79) *Nipponnemertes bimaculata* (Coe, 1901)
 80) *Nipponnemertes ogumai* (Yamaoka, 1947)
 81) *Nipponnemertes punctatula* (Coe, 1905)
 10) Family EMPLECTONEMATIDAE Bürger, 1904
 82) *Emplectonema buergeri* Coe, 1901
 83) *Emplectonema gracile* (Johnston, 1837)
 84) *Emplectonema kandai* Kato, 1939
 85) *Emplectonema mitsuii* Yamaoka, 1947
 86) *Nemertopsis mitellicola* Kajihara, 2007
 87) *Nemertopsis quadripunctata* (Quoy and Gaimard, 1833)
 88) *Paranemertes incola* Iwata, 1952
 89) *Paranemertes katoi* Yamaoka, 1947
 90) *Paranemertes peregrina* Coe, 1901
 91) *Paranemertes plana* Iwata, 1957
 11) Family MALACOBDELLIDAE Blanchard, 1847
 92) *Malacobdella japonica* Takakura, 1897
 12) Family OTOTYPHLONEMERTIDAE Bürger, 1895
 93) *Ototyphlonemertes dolichobasis* Kajihara, 2007
 94) *Ototyphlonemertes martynovi* Chernyshev, 1993
 95) *Ototyphlonemertes nikolaii* Chernyshev, 1998
 13) Family POSEIDONEMERTIDAE Chernyshev, 2002
 96) *Diopsonemertes acanthocephala* Kajihara, Gibson and Mawatari, 2001
 14) Family PROSORHOCHMIDAE Bürger, 1895
 97) *Geonemertes pelaensis* Semper, 1863
 98) *Pantinonemertes spectacula* (Yamaoka, 1940)

- 15) Family TETRASTEMMATIDAE Hubrecht, 1879
 99) *Nemertellina yamaokai* Kajihara, Gibson and Mawatari, 2000
- 100) *Oerstedia dorsalis* (Abildgaard, 1806)
 101) *Oerstedia polyorbis* Iwata, 1954
 102) *Prostoma ohmense* Chernyshev, Timoshkin and Kawakatsu, 1998
 103) *Quasitetrastemma nigritrons* (Coe, 1904)
 104) *Quasitetrastemma stimpsoni* (Chernyshev, 1992)
 105) *Sacconemertella lutulenta* Iwata, 1970
 106) *Sacconemertopsis olivifera* Iwata, 1970
 107) *Tetrastemma candidum* (Müller, 1774)
 108) *Tetrastemma insolens* Iwata, 1952
 109) *Tetrastemma melanocephalum* (Johnston, 1837)
 110) *Tetrastemma pinnatum* Iwata, 1954
 111) *Tetrastemma pseudocoronatum* Chernyshev, 1998
 112) *Tetrastemma roseocephalum* (Yamaoka, 1947)
 113) *Tetrastemma stigmatum* Stimpson, 1857
 114) *Tetrastemma verinigrum* Iwata, 1954
 115) *Tetrastemma yamaokai* Iwata, 1954
- 2) Subclass POLYSTILIFERA Brinkmann, 1917
- 1) Order REPTANTIA Brinkmann, 1917
 16) Family DREPANOPHORIDAE Verrill, 1892
 116) *Drepanophorus longiceps* Iwata, 1957
 117) *Kameginemertes parmiornata* (Iwata, 1957)
- 17) Family SAGAMINEMERTIDAE Chernyshev, 2003
 118) *Sagaminemertes nagaiensis* (Iwata, 1957)
- 2) Order PELAGICA Brinkmann, 1917
 18) Family NECTONEMERTIDAE Verrill, 1892
 119) *Nectonemertes japonica* Foshay, 1912
 19) Family PELAGONEMERTIDAE Moseley, 1875
 120) *Pelagonemertes moseleyi* Bürger, 1895

Taxonomic Catalogue of Japanese Nemerteans

The entries in the synonymy for each species are arranged chronologically. For convenience, each authority in the synonymy is indicated in bold letters. Bibliographic information is given in the synonymy for both primary and secondary literature. For primary literature, such information as locality, habitat, and date of collection are also provided. The prefecture of each locality is given, to facilitate relocation and avoid confusion by synonymy of place names. Where applicable, a long vowel in place names is marked by a macron (e.g., È, â), to avoid confusion between, e.g., "Ôshima" and "Oshima." The literature covered includes not only taxonomic papers, but also those on ecology and biochemistry, pictorial books, faunal reports, and field guides. Japanese common names that have previously been assigned are indicated for a number of species, but no attempt has been made to create new Japanese names for the remaining species. The following abbreviations are used to indicate museum depositories of specimens:

NHMW-EV: Naturhistorisches Museum Wien, Evertebrata-Varia, Wien, Austria.

USNM: National Museum of Natural History, Smithsonian Institution, Washington, D.C., USA.

ZIHU: Hokkaido University Museum, Sapporo, Japan.

Phylum NEMERTEA

Class PALAEONEMERTEA Hubrecht, 1879

Family CALLINERIDAE Bergendal, 1901

Genus *Callinera* Bergendal, 1900

Callinera Bergendal, 1900: 313.

TYPE SPECIES: *Callinera buergeri* Bergendal, 1900, by monotypic designation.

Callinera nishikawai Kajihara, 2006

Callinera nishikawai Kajihara, 2006: 17–27, figs. 11–14; sandy to muddy tidal flat, 34°41'04"N, 137°35'59"E, Ikarise, Hamanako Lake, Shizuoka Prefecture.

TYPE MATERIAL: Holotype, ZIHU-3133, 12 May 2002, collected by Taeko Kimura, Sho-ichi Kimura, and Teruaki Nishikawa, female, 6-µm serial transverse sections of an anterior body fragment about 1.5 cm in length and 0.8–1.0 mm in width, fixed in Bouin's fluid without anaesthetization.

NOTE: The species was described on the basis of a fixed anterior fragment of the body, and the shape of the living animal is unknown.

Family CEPHALOTRICHIDAE McIntosh, 1874

NOTE: Although the family name was incorrectly spelled as "Cephalothricidae" when established, the correct spelling of the family name should be "Cephalotrichidae" under Article 29.3 of the Code (ICZN, 1999), for the name of its type genus *Cephalothrix* gives the genitive singular "Cephalotrichos," and thus the stem "Cephalotrich-." Article 29.5 of the Code (ICZN, 1999) is not applicable, since the correct spelling has been widely used, e.g., by Bürger (1895: 533, 1904: 16), Wijnhoff (1913: 294), Coe (1930: 97, 1940: 257), Yamaoka (1940a: 215), Hylbom (1957: 553, 1993: 173), Moretto (1974: 9), Chernyshev (2004a: 152) and Tanu et al. (2004: 515).

Genus *Cephalothrix* Örsted, 1843

Cephalothrix Örsted, 1843: 573.

Procephalothrix Wijnhoff, 1913: 294; synonymized by Sundberg et al. (2003).

Cephalotrichella Wijnhoff, 1913: 298; synonymized by Sundberg et al. (2003).

TYPE SPECIES: *Cephalothrix coeca* Örsted, 1843, now regarded as a junior synonym of *Planaria linearis* Rathke, 1799 (Bürger, 1904), by subsequent designation.

NOTE: Gibson's (1995) assignment and Sundberg et al.'s (2003) statement of *Cephalothrix linearis* (Rathke, 1799) as the type species of the genus are irrelevant in terms of Articles 67.1.2 and 67.2 of the Code (ICZN, 1999), because the genus *Cephalothrix* contained only two nominal species, *Cephalothrix bioculata* and *Cephalothrix coeca*, when it was originally established.

Cephalothrix fasciculus (Iwata, 1952)

Procephalothrix fasciculus Iwata, 1952: 130, figs. 2, 8; under stones on stony beach near low-water level, Tomioka, Amakusa, Kumamoto Prefecture.

Procephalothrix fasciculasic [sic]: Crandall et al., 2002: 14, 16, 29, 36, 41.

NOTE: The species was originally classified in the genus *Procephalothrix*, which has been synonymized with *Cephalothrix* by Sundberg et al. (2003: 292); now it should be

known as *Cephalothrix fasciculus*. The species is characterized by the posterior end of its rhynchocoel reaching the hind end of the body. This character state is absent among its congeners and has only been recorded in this species. At the same time, however, this character state can be erroneously identified by misinterpretation of a body fragment as an intact specimen. If the anus cannot be confirmed in the holotype specimen, the name of the nominal species *Procephalothrix fasciculus* should be regarded as a *nomen dubium*.

***Cephalothrix notabilis* Iwata, 1954**

[Japanese name: shirayuki-himomushi or kita-hoso-himomushi]
Cephalothrix notabilis Iwata, 1954a: 8, fig. 1C, E, F; under stones near low-water level on stony beach, Akkeshi, Hokkaidō Prefecture. Yamaguchi and Yamada, 1955: 65. Uchida et al., 1963: 17. Iwata, 1965a: 201. Iwata, 1992: 196, fig. 7-2A. Iwata, 1997: 55. Crandall et al., 2002: 10, 16, 26, 33.

***Cephalothrix simula* (Iwata, 1952)**

[Japanese name: akahana-himomushi]

Procephalothrix simulus [sic] Iwata, 1952: 132; under stones near low-water level, Fukue, Gotō Islands, Nagasaki Prefecture; non *Procephalothrix simula* sensu Iwata (1954a), nec *Cephalothrix linearis* sensu Yamaoka (1940).

NOTE: The species was originally classified in the genus *Procephalothrix*, which was synonymized with *Cephalothrix* by Sundberg et al. (2003: 292); now it should be known as *Cephalothrix simula*. Iwata (1954a) synonymized *Cephalothrix linearis* sensu Yamaoka (1940) and *Procephalothrix simula* sensu Iwata (1954a) with *Procephalothrix simula* sensu Iwata (1952). *Cephalothrix* (=*Procephalothrix*) *simula* sensu Iwata (1952) is characterized by the absence between the rhynchocoel and alimentary canal of a longitudinal muscle plate (Iwata, 1952), which, however, is present in *Cephalothrix linearis* sensu Yamaoka (1940a) and *Procephalothrix simula* sensu Iwata (1954a). Therefore, I hesitate to apply the name *Cephalothrix simula* to the taxon to which the latter two authors referred. See NOTE under *Cephalothrix linearis*.

Family TUBULANIDAE Bürger, 1904 (1874)

NOTE: In response to Melville's (1986) proposal, ICZN (1988) ruled under Article 40b of the third edition of the Code (ICZN, 1985) that the name Tubulanidae has precedence over, but takes the date of, its senior subjective synonym Carinellidae. In his proposal, Melville (1986) stated that the name *Carinella trilineata* "has been regarded as a synonym of *Tubulanus polymorphus* since at least 1905," referring to Bürger (1897–1907), and that the family Tubulanidae "should be cited with the date '1905 (1874)'." The ICZN's ruling, basically following Melville's proposal, states that "the name *Tubulanidae* Bürger, 1905 (1874) ... is hereby placed on the Official List of Family-Group Names in Zoology." Bürger's (1897–1907) book was published in six different parts, and Melville (1986) was quite correct in that the relevant part about the replacement of Carinellidae with Tubulanidae was published in 1905 (pp. 401, 402, 405). However, the name Tubulanidae had already appeared prior to 1905 in Bürger (1904). Thus the family

name should be cited as "Tubulanidae Bürger, 1904 (1874)," with the date of priority being enclosed in parentheses in accordance with Recommendation 40A of the Code (ICZN, 1985, 1999).

Genus *Carinesta* Punnett, 1900

Carinesta Punnett, 1900: 569.

TYPE SPECIES: *Carinesta orientalis* Punnett, 1900 by monotypic designation.

***Carinesta uchidai* Iwata, 1952**

[Japanese name: kensaki-himomushi]

Carinesta uchidai Iwata, 1952: 128, fig. 7; under stones near low-water level, Fukue, Gotō Islands, Nagasaki Prefecture. Iwata, 1960c: 166, pl. 83, fig. 2. Iwata, 1965a: 201. Iwata, 1965b: 391, figs. a, b. Iwata, 1992: 197. Crandall et al., 2002: 10, 16, 26, 33.

NOTE: Sundberg and Hylbom's (1994) cladistic analysis based on morphological characters shows that the genus *Carinesta* is a polyphyletic group and that *Carinesta uchidai* comprises a monophyletic group together with members of the family Cephalotrichidae. The generic placement of this species, as well as the taxonomic status of the genus *Carinesta*, requires reassessment.

Genus *Carinina* Hubrecht, 1885

Carinina Hubrecht, 1885: 830.

Procarinina Bergendal, 1902: 422; synonymized by Hylbom (1957).

TYPE SPECIES: *Carinina grata* Hubrecht, 1887, by monotypic designation; Hubrecht (1885) did not designate the type species when he erected the genus *Carinina*.

***Carinina plecta* Kajihara, 2006**

Carinina plecta Kajihara, 2006: 5–16, figs. 3–10; sandy to muddy tidal flat, 34°41'04"N, 137°35'59"E, Ikarise, Hamanako Lake, Shizuoka Prefecture.

TYPE MATERIAL: Holotype, ZIHU-3123, 31 July 2003, female; serial transverse (6 µm thick) and longitudinal (10 µm thick) sections of a fragment containing the head; total 116 slides.

Genus *Tubulanus* Renier, 1804

Tubulanus Renier, 1804: 20.

Carinella Johnston, 1833: 232; synonymized by Bürger (1904).

TYPE SPECIES: *Tubulanus polymorphus* Renier, 1804 by monotypic designation.

***Tubulanus capistratus* (Coe, 1901)**

Tubulanus capistratus: Coe, 1944: 27, "One specimen nearly a meter in length was collected by the Albatross in 1906 near Hakodate, Japan." Crandall et al., 2002: 15, 16, 30, 37, 42.

NOTE: Originally described as *Carinella capistrata* by Coe (1901: 16) from Orca and Virgin Bay in Prince William Sound, Alaska; transferred to the genus *Tubulanus* by Coe (1940: 255). Apart from the record from Japan, the species is known to be distributed along the Pacific coast of North America (Gibson, 1995: 316).

Tubulanus ezoensis Yamaoka, 1940

[Japanese name: ezo-himomushi]

Tubulanus ezoensis Yamaoka, 1940a: 212–215, pl. XIV, figs. 3, 4, text figs. 3, 4; lower intertidal under stones, Daikokujima, Akkeshi, Hokkaidō Prefecture. **Okuda**, 1947: 1474, fig. 4158. **Iwata**, 1954a: 6. **Yamaguchi and Yamada**, 1955: 64. **Uchida et al.**, 1963: 17. **Okuda and Iwata**, 1965: 390; figs. a–c. **Crandall et al.**, 2002: 15, 16, 30, 37, 42.

NOTE: *Tubulanus ezoensis* has long been known only by Yamaoka's (1940) original description (Iwata, 1954a). Although the type material appears to have been lost (see "Brief History..."), some additional specimens referable to *T. ezoensis* were recently collected from the type locality (Kajihara, pers. obs.).

Tubulanus lucidus Iwata, 1952

Tubulanus lucidus Iwata, 1952: 126–128, figs. 1, 6; lower intertidal under stones, Fukue Island, Gotō Islands, Nagasaki Prefecture. **Crandall et al.**, 2002: 15, 16, 30, 37.

NOTE: Sundberg and Hylbom's (1994) cladistic analysis based on morphological characters indicates that this species is the sister taxon to hubrechtids, appearing in a clade which is different from that containing other *Tubulanus*, including the type species. The generic placement of this taxon thus needs reassessment.

Tubulanus punctatus (Takakura, 1898)

[Japanese name: kurige-himomushi or kugi-himomushi]

Carinella punctata **Takakura**, 1898: 117–118, fig. 3; sublittoral from 2–3 fathoms depth, Jōgashima, Kanagawa Prefecture.

Tubulanus punctatus: **Kaburaki**, 1927: 1662, fig. 3180.

Yamaoka, 1940a: 208–212, pl. XIV, figs. 1, 2, text figs. 1, 2; from lower intertidal under stones and in rock crevices to sublittoral among mud, Hokkaidō Prefecture (Akkeshi, Ochiishi, Nemuro, Abashiri, Usu and Muroran). **Kaburaki**, 1947: 1474, fig. 4157. **Iwata**, 1954a: 5; lower intertidal under stones, Hokkaidō Prefecture (Akkeshi, Muroran, Oshoro, Rishiri Island and Monbetsu). **Iwata**, 1954b: 34; habitat not recorded, northern coast near the Seto Marine Biological Laboratory, Wakayama Prefecture. **Yamaguchi and Yamada**, 1955: 64. **Utinomi**, 1956: 31; pl. 16, fig. 1. **Iwata**, 1960a: 96, fig. 2. **Iwata**, 1960b: 19–25, figs. 62–91; intertidal, Akkeshi, Hokkaidō Prefecture. **Iwata**, 1960c: 166, pl. 83, fig. 1. **Utinomi**, 1960: 31, pl. 16, fig. 1. **Satō and Itō**, 1961: 187, fig. 7.1.1. **Uchida et al.**, 1963: 17. **Iwata**, 1965a: 169, 201. **Iwata**, 1965b: 390, one figure. **Shiino**, 1969: 94, fig. 9-3A. **Utinomi**, 1969: 31, pl. 16, fig. 1. **Okada et al.**, 1971: 62. **Uchida et al.**, 1972: 62. **Kito**, 1975: 149; among the holdfasts of *Sargassum confusum*, Oshoro, Hokkaidō Prefecture. **Honma and Kitami**, 1978: 14; Sado Island, Nîgata Prefecture. **Iwata**, 1983: 181, 182, figs. 8–5g, h, i, 8–13b. **Hieda and Takahashi**, 1986: 41, with two color photographs of a specimen taken in Yakumo, Hokkaidō Prefecture. **Inaba**, 1988: 225; lower intertidal to shallow sublittoral, under stones on gravelly to rocky shores, found rarely in the Inland Sea of Seto. **Miyazawa et al.**, 1988: 867–874; intertidal, habitat unknown, Hiroshima Prefecture (Iwashijima and Mukaishima), Yugeshima, Ehime Prefecture, identified by Dr. T.

Hoshino, Mukaishima Marine Biological Station, Hiroshima University. **Iwata**, 1992: 196, pl. 44-1, fig. 7-2B. **Iwata**, 1997: 53 (with a color photograph take in life by Fumio Iwata), 55. **Shimomura et al.**, 2001: 46; dredged sublittorally from two sites in Ôtsuchi Bay, Iwate Prefecture; 39°21'01"N, 141°58'31"E, 58.7 m depth, mixture of sand and shell debris; 39°20'43"N, 141°57'43"E, 49.0 m depth, muddy sand. **Crandall et al.**, 2002: 15, 16, 30, 37, 42.

NOTE: Apart from the records from Japanese waters, *Tubulanus punctatus* is also known from Posjet Bay (Peter the Great Bay) (Korotkevitsch, 1971), Sakhalin, the Kuril Islands and Kamchatka Peninsula (Korotkevitsch, 1982), Vostok Bay (Kulikova, 1988), and Shandon Province (Qingdao, Huangdao, and Jiaonan), China (Yin et al., 1988).

Tubulanus roretzi Senz, 1997

Tubulanus roretzi **Senz**, 1997a: 424–430, figs. 1–4; locality and habitat unknown.

TYPE MATERIAL: Holotype, NHMW-EV 3565/1886; paratype, NHMW-EV 3566–3573.

NOTE: The material was collected by Dr. Albrecht von Roretz during his stay in Japan from 1874–1882.

Class PILIDIOPHORA Thollesson and Norenburg, 2003**Family HUBRECHTELLIDAE Chernyshev, 2003****Genus *Hubrechtella* Bergendal, 1902**

Hubrechtella Bergendal, 1902: 9.

Coeia **Takakura**, 1922: 419; synonymized by Kajihara (2006).

TYPE SPECIES: *Hubrechtella dubia* Bergendal, 1902, by monotypic designation.

Hubrechtella ijimai (Takakura, 1922)

[Japanese name: ijima-himomushi]

Coeia ijimai **Takakura**, 1922: 419–422, two figs.; among sandy gravel on beach, Enoura Bay, Shizuoka Prefecture and Tateyama Bay, Chiba Prefecture. **Crandall et al.**, 2002: 10, 16, 27, 34.

Coia* [sic] *ijimai: **Iwata**, 1960c: 166, pl. 83, fig. 3. **Iwata**, 1965a: 201. **Iwata**, 1965b: 391, figs. 6a, b; under stones on sandy beach in Asamushi, Aomori Prefecture, Onagawa, Miyagi Prefecture, Usa, Kôchi Prefecture, and Fukue, Nagasaki Prefecture. **Tsuchiya**, 1979: 82; intertidal, Hadakajima Island, Asamushi Aomori Prefecture. **Iwata**, 1992: 197, fig. 7-3B.

Hubrechtella ijimai: **Kajihara**, 2006: 28–37, figs. 15–19; sandy to muddy tidal flat, 34°41'04"N, 137°35'59"E, Ikarise, Hamanako Lake, Shizuoka Prefecture.

Hubrechtella kimuraorum Kajihara, 2006

Hubrechtella kimuraorum **Kajihara**, 2000: 37–43, figs. 20–23; sandy to muddy tidal flat, 34°41'04"N, 137°35'59"E, Ikarise, Hamanako Lake, Shizuoka Prefecture.

TYPE MATERIAL: Holotype, ZIHU-3127, male, 1 August 2003, 72 slides, 6-µm serial transverse sections of the body except in the middle portion.

Genus *Tetramys* Iwata, 1957

Tetramys Iwata, 1957a: 2.

TYPE SPECIES: *Tetramys ramicerebrus* Iwata, 1957 by monotypic designation.

NOTE: Cladistic analyses by Sundberg and Hylbom (1994) and Sundberg *et al.* (2003) based on morphological characters indicate that the genus appears to be synonymous with *Hubrechtella*.

Tetramys ramicerebrus Iwata, 1957

[Japanese name: miura-himomushi]

Tetramys ramicerebrum [sic] **Iwata**, 1957a: 3–5, pl. I, fig. 1, pl. II, figs. 1–6; dredged sublitorally from 20 m depth on 9 February 1955 by His Majesty Emperor Shōwa, sediment type not recorded, Higashioine, Sagami Bay, Kanagawa Prefecture. **Iwata**, 1965a: 201. **Iwata**, 1965b: 391, figs. a–d. *Tetramys ramicerebrus*: **Crandall et al.**, 2002: 14, 16, 29, 41.

Family LINEIDAE McIntosh, 1874

Genus Cerebratulus Renier, 1804

Cerebratulus Renier, 1804: 21.

Meckelia Leuckart, 1828: 17; synonymized by Hubrecht (1879) (in part).

TYPE SPECIES: *Cerebratulus marginatus* by subsequent designation of Gibson (1995).

Cerebratulus albocirculus Iwata, 1957

Cerebratulus albocirculus **Iwata**, 1957a: 17–18, pl. I, fig. 7, pl. V, figs. 6, 7; “Mosaki at Kamezyo,” dredged sublitorally from 10 m depth on 23 July 1956 by His Majesty Emperor Shōwa, Sagami Bay, off Kanagawa Prefecture. **Crandall et al.**, 2002: 10, 17, 26, 38.

NOTE: The place indicated by the name “Mosaki at Kamezyo” in Iwata’s (1957a) original description is uncertain. One possible candidate in Sagami Bay is “Kamegishō,” a bank located at approximately 35°35'N, 35°12'E.

Cerebratulus carnosus Takakura, 1898

Cerebratulus L.C. *arnosus* [sic] **Takakura**, 1898: 426, fig. 23; mud, Misaki Harbour and Koajiro Bay, Kanagawa Prefecture.

Cerebratulus arnosus: **Crandall et al.**, 2002: 10, 17, 26, 33.

NOTE: The abbreviation “L.C.”, most probably denoting “lower-case letters,” was mistakenly inserted between the generic and specific names in the original publication, with the initial “c” dropped from “carnosus” (=Latin, meaning “fleshy”), which refers to the body coloration of the species treated in the original description. The original spelling *arnosus* is incorrect, due to a printer’s error, according to Article 32.5 of the Code (ICZN, 1999) (Crandall, pers. comm.).

Cerebratulus communis Takakura, 1898

[Japanese name: nami-himomushi]

Cerebratulus L.C. *ommunis* [sic] **Takakura**, 1898: 425, fig. 22 (originally numbered as fig. 20); intertidal in sandy mud, Misaki Harbour, Koajiro Bay, Bishamon Bay, and Matsuwa Bay, Kanagawa Prefecture.

Cerebratulus communis: **Kaburaki**, 1927: 1666, fig. 3189. **Kaburaki**, 1947: 1471, fig. 4149. **Iwata**, 1952: 140–141, fig. 5; lower intertidal among sandy mud, Tomioka, Amakusa, Kumamoto Prefecture. **Utinomi**, 1956: 31, pl. 16, fig. 6. **Iwata**, 1960c: 169, pl. 84, fig. 6; Asamushi,

Aomori Prefecture. **Utinomi**, 1960: 31, pl. 16, fig. 6. **Inaba**, 1963: 228; lower intertidal on sandy to muddy shores or under stones on gravelly to rocky shore, commonly found in the Inland Sea of Seto. **Kikuchi**, 1968: 167; among *Zostera marina*, Tomioka Bay, Amakusa, Kumamoto Prefecture. **Utinomi**, 1969: 31, pl. 16, fig. 6. **Inaba**, 1988: 225; lower intertidal on sandy to muddy shores or under stones on gravelly to rocky shores, commonly found in the Inland Sea of Seto. **Crandall et al.**, 2002: 10, 17, 24, 26, 33.

Cerebratulus comunis [sic]: **Kaburaki and Iwata**, 1965: 395, with one figure.

NOTE: As in the case with *Cerebratulus arnosus*, a nomenclatural consideration is required as to the correct spelling of the species name. The species has been also recorded from Alaid Island, northern Kurile Islands (Takakura, 1933).

Cerebratulus fasciatus Stimpson, 1857

Cerebratulus fasciatus **Stimpson**, 1857: 161; obtained sublitorally from sandy to stony bottoms of about 7.4 m depth, Hokkaidō Prefecture. Originally recorded as “Apud oras insulae ‘Jesso’ Japoniae Borealis; in fundo arenoso-limoso profunditatis 4 orgyiarum” (near the region of Hokkaidō Island, northern Japan; in a bottom of muddy sand at a depth of 7.4 m). **Iwata**, 1954a: 14. **Yamaguchi and Yamada**, 1955: 67. **Crandall et al.**, 2002: 10, 17, 26, 38.

Cerebratulus formosus Iwata, 1957

Cerebratulus formosus **Iwata**, 1957a: 15–17, pl. I, fig. 4, pl. V, figs. 1–5; dredged sublitorally from 100 m depth on 13 December 1952 by His Majesty Emperor Shōwa, Nakafukari at Hayama, Sagami Bay, off Kanagawa Prefecture. **Crandall et al.**, 2002: 10, 17, 26, 38.

Cerebratulus longiceps Coe, 1901

Cerebratulus longiceps: **Coe**, 1944: 29, obtained by the United States Bureau of Fisheries Steamer *Albatross*, 250 m depth, off Ōshima, the Metropolis of Tōkyō. **Crandall et al.**, 2002: 10, 17, 27, 38.

NOTE: *Cerebratulus longiceps* was originally described from Yakutat, Alaska, USA, by Coe (1901: 77). The species is so far only known from Coe’s (1901, 1944) records.

Cerebratulus macroren Hubrecht, 1887

Cerebratulus macroren **Hubrecht**, 1887: 46–47, pl. I, figs. 13, 14, 18, 19, pl. X, figs. 8, 9, pl. XI, fig. 11, pl. XII, figs. 1, 2, 7, 8, pl. XIII, figs. 7–9, pl. XIV, figs. 7, 8, 11, pl. XV, figs. 2, 3, 19, text fig. 4; sublitoral from 345 fathoms (about 640 m) depth on green mud, collected on 12 May 1875 by H.M.S. *Challenger*, 35°11'00"N, 139°28'00"E, Sagami Bay, off Kanagawa Prefecture.

Cerebratulus marginatus Renier, 1804

[Japanese name: orochi-himomushi]

Cerebratulus marginatus: **Yamaoka**, 1940a: 222–224, pl. XV, figs. 6–8, text fig. 9; lower intertidal or sublitoral in soft mud or among fine sand under stones, Akkeshi, Hokkaidō Prefecture. **Coe**, 1944: 29. **Okuda**, 1947: 1472, fig. 4150. **Iwata**, 1954a: 14; sublitoral among soft mud, Hokkaidō Prefecture (Akkeshi, Kushiro and Nemuro); sublitoral among soft mud, Misaki, Kanagawa Prefecture. **Iwata**,

1954b: 37; sublittoral, obtained by trawling off the coast of the Kii Peninsula, Wakayama Prefecture. **Yamaguchi and Yamada**, 1955: 66. **Iwata**, 1957a: 11–12, pl. I, fig. 6; dredged sublitorally from 200–280 m depth on 5 February 1955 by His Majesty Emperor Shōwa, “Southern Maruyamadashi at Aamadaiba” [sic], Sagami Bay, Kanagawa Prefecture. **Iwata**, 1960c: 169, pl. 84, fig. 5. **Uchida et al.**, 1963: 17. **Iwata**, 1965a: 169, 201. **Satō and Itō**, 1961: 187, fig. 7.1.5. **Okuda and Iwata**, 1965: 385, figs. a, b. **Shiino**, 1969: 94, fig. 9-3B-D. **Honma and Kitami**, 1978: 15; Sado Island, Niigata Prefecture. **Iwata**, 1992: 198, fig. 7-2D. **Iwata**, 1997: 55. **Shimomura et al.**, 2001: 46; obtained from a sediment of mixed sand, mud, and shell debris at a depth of 64 m, 39°23.363'N, 141°58.971'E, Ōtsuchi Bay, Iwate Prefecture. **Crandall et al.**, 2002: 10, 17, 24, 27, 38.

NOTE: *Cerebratulus marginatus* Renier, 1804 was originally described from the Adriatic Sea, Italy (presumably Padua). Apart from the records in Japanese waters, the species is also reported from the Pacific coast of North America (Alaska to California), the western North Atlantic (Greenland, Labrador, and Cape Cod southwards under the offshore Arctic current), the Arctic (King Charles Land, Bremer Sound, Hinlopen Strait, Spitzbergen), Europe (Norway, the British Isles, the Mediterranean), and south to Madeira (Gibson, 1995: 340).

***Cerebratulus montgomeryi* Coe, 1901**

Cerebratulus montgomeryi: **Coe**, 1944: 29; obtained by the United States Bureau of Fisheries Steamer Albatross, 600 m depth, off Hokkaidō. **Crandall et al.**, 2002: 10, 17, 24, 27, 34, 38.

NOTE: Originally described from Alaska by Coe (1901: 80), *Cerebratulus montgomeryi* is distributed along Pacific coast of North America, the Aleutian Islands, Bering Sea, coast of Siberia, Japan Sea coasts of Russia, and Japan (Kulikova, 1988; Gibson, 1995: 341).

***Cerebratulus penniger* Iwata, 1957**

Cerebratulus penniger **Iwata**, 1957a: 13–14, pl. I, fig. 3, pl. IV, figs. 4–6; dredged sublitorally from 380 m depth on 28 September 1953 by His Majesty Emperor Shōwa, Nakafukari at Hayama, Sagami Bay, off Kanagawa Prefecture. **Crandall et al.**, 2002: 10, 17, 27, 38.

***Cerebratulus subacutus* (Stimpson, 1857)**

Meckelia subacuta **Stimpson**, 1857: 161; intertidal in mud, Naha, Okinawa Prefecture. Originally recorded as “In portu ‘Napa’ insulae ‘Loo Choo;’ littoralis in limo”; transferred to *Baseodiscus* by Bürger (1904: 120).

Cerebratulus subacutus: **Crandall et al.**, 2002: 10, 17, 30, 34.

***Cerebratulus superniger* Iwata, 1957**

Cerebratulus superniger **Iwata**, 1957a: 14–15, pl. I, fig. 5, pl. IV, figs. 7, 8; collected by His Majesty Emperor Shōwa from a depth of 10 m on 10 January 1930, “Ithishiki at Hayama” [sic], off Kanagawa Prefecture. **Crandall et al.**, 2002: 10, 17, 27, 39.

***Cerebratulus zebra* Punnett and Cooper, 1909**

Cerebratulus zebra: **Iwata**, 1957a: 12–13; dredged sublit-

torally from 410 m depth on 16 July 1940 by His Majesty Emperor Shōwa, “Aamadaiba” [sic], Sagami Bay, off Kanagawa Prefecture. **Crandall et al.**, 2002: 10, 17, 27, 39.

NOTE: *Cerebratulus zebra* was originally described from Sri Lanka (Punnett and Cooper, 1909: 11). The species is currently known only from the two localities, Sri Lanka and Japan.

Genus *Diplopleura* Stimpson, 1857

Diplopleura Stimpson, 1857: 162.

Langia Hubrecht, 1879: 220; synonymized by Verrill (1895: 528).

TYPE SPECIES: *Diplopleura japonica* Stimpson, 1857 by monotypic designation.

NOTE: The genus *Diplopleura*, currently containing five nominal species, was established only on the basis of external characters, in which the lateral margins of the body are dorsally curled up.

***Diplopleura japonica* Stimpson, 1857**

[Japanese name: hida-himomushi]

Diplopleura japonica **Stimpson**, 1857: 162; shallow sublitoral in sandy bottom of 5 orgya (=5 fathoms=9.3 m) depth, Kagoshima Bay, Kagoshima Prefecture. **Iwata**, 1965a: 201. **Crandall et al.**, 2002: 11, 17, 27, 39.

NOTE: *Diplopleura japonica* has not been reported since its original description. Stimpson’s specimen was light yellowish chestnut in color, measured 1.5 “pollex” (=1.5 inch=3.8 cm) in length and 0.12 “pollex” (=0.12 inch=0.3 cm) in width.

Genus *Euborlasia* Vaillant, 1890

Euborlasia Vaillant, 1890: 616.

TYPE SPECIES: *Borlasia elizabethae* McIntosh, 1874 by monotypic designation.

***Euborlasia gotoensis* Iwata, 1952**

[Japanese name: gotō-himomushi]

Euborlasia gotoensis **Iwata**, 1952: 133–134, figs. 3, 9, 10; lower intertidal under stones, Fukue, Gotō Islands, Nagasaki Prefecture. **Iwata**, 1960c: 166, pl. 83, fig. 8. **Iwata**, 1965a: 169, 201. **Iwata**, 1965b: 392, figs. a–c. **Honma and Kitami**, 1978: 15; Sado Island, Niigata Prefecture. **Iwata**, 1992: 198, pl. 44-3. **Crandall et al.**, 2002: 11, 17, 27, 34.

***Euborlasia proteres* Iwata, 1957**

Euborlasia proteres **Iwata**, 1957a: 8–9, pl. I, fig. 2, pl. IV, figs. 1–3; dredged sublitorally from 380 m depth on 28 September 1953 by His Majesty Emperor Shōwa, Nakafukari at Hayama, Sagami Bay, off Kanagawa Prefecture. **Crandall et al.**, 2002: 11, 17, 27, 39.

Genus *Hinumanemertes* Iwata, 1970

Hinumanemertes Iwata, 1970a: 134.

TYPE SPECIES: *Hinumanemertes kikuchii* Iwata, 1970 by original designation.

***Hinumanemertes kikuchii* Iwata, 1970**

[Japanese name: hinuma-himomushi]

Hinumanemertes kikuchii **Iwata**, 1970a: 136–142, fig. 1A–C,

pl. 1, figs. 1–8, pl. 2, figs. 9–17; obtained sublitorally in mud from a brackish-water lake, Lake Hinuma, Ibaraki Prefecture. **Iwata**, 1970b: 128. **Iwata**, 1973: 264. **Iwata**, 1992: 198, fig. 7-3E. **Crandall et al.**, 2002: 11, 17, 27, 43.

Genus *Iwatanemertes* Gibson, 1990

Iwatanemertes Gibson, 1990a: 75.

TYPE SPECIES: *Meckelia piperata* Stimpson, 1855 by original designation.

Iwatanemertes piperata (Stimpson, 1855)

[Japanese name: ryûkyû-himomushi]

Meckelia piperata Stimpson, 1855: 381; habitat not recorded, Kikaishima, Kagoshima Prefecture.

Lineus piperata [sic]: **Takakura**, 1898: 186–187, fig. 9; intertidal among algae, Enoshima and Misaki, Kanagawa Prefecture.

Lineus piperatus: **Stimpson**, 1857: 160; sublitoral between stones and among algae, Kikaishima, Kagoshima Prefecture; originally recorded as “In portu insulae ‘Kikaishima’ Japoniae Australis; sublitoralis inter lapillus et algas.” **Yamaoka**, 1940b: 13–15, figs. 1, 2; habitat not recorded, Shitaru, Minami-Izu, Shizuoka Prefecture; habitat not recorded, Naha, Okinawa Island, Okinawa Prefecture. **Iwata**, 1954b: 34; intertidal under stones, Shirahama, Wakayama Prefecture. **Iwata**, 1960c: 169, pl. 84, fig. 1; Aikawa, Sado Island, Niigata Prefecture. **Honma and Kitami**, 1978: 15; Sado Island, Niigata Prefecture. **Iwata**, 1965b: 394, with one figure.

Iwatanemertes piperata: **Crandall et al.**, 2002: 11, 17, 27, 31, 34, 39.

NOTE: Transferred to *Iwatanemertes* by Gibson (1990a: 75), based upon a study of material collected in Hong Kong. Apart from the records from Japanese waters, the species is also known from Hong Kong (Gibson, 1990a) and the Taiwan Straits (Sun, 1995).

Genus *Lineus* Sowerby, 1806

Lineus Sowerby, 1806: 15.

Heterolineus Friedrich, 1935a: 310; synonymized by Corrêa (1963: 43).

TYPE SPECIES: *Ascaris longissima* Gunnerus, 1770 by monotypic designation.

Lineus alborostratus Takakura, 1898

[Japanese name: takakura-himomushi]

Lineus alborostratus **Takakura**, 1898: 332, fig. 12; habitat not recorded, Misaki, Kanagawa Prefecture. **Yamaoka**, 1940a: 220–222, pl. XV, figs. 1–5, text fig. 8; lower intertidal under stones, Akkeshi and Ochiishi, Hokkaidô Prefecture. **Iwata**, 1951: 136–137, figs. 2, 3c; intertidal under stones, Mukaishima, Hiroshima Prefecture. **Iwata**, 1952: 138–139; lower intertidal under stones, Fukue, Gotô Islands, Nagasaki Prefecture. **Iwata**, 1954a: 12; lower intertidal under stones, Hokkaidô Prefecture (Akkeshi, Nemuro, Hiroo, Muroran, Monbetsu and Rishiri Island). **Yamaguchi and Yamada**, 1955: 66. **Iwata**, 1957c: 102. **Iwata**, 1960b: 26–27, figs. 92–96; intertidal, Akkeshi, Hokkaidô Prefecture. **Iwata**, 1960c: 166, pl. 83, fig. 9. **Inaba**, 1963: 227; lower intertidal to shallow sublitoral under stones on rocky to gravelly shores, commonly found in the Inland Sea of Seto.

Uchida et al., 1963: 17. **Iwata**, 1965b: 393, figs. a–c. **Iwata**, 1983: 182. **Inaba**, 1988: 225; lower intertidal to shallow sublitoral under stones on rocky to gravelly shores, commonly found in the Inland Sea of Seto. **Iwata**, 1992: 198, pl. 44–6. **Crandall et al.**, 2002: 11, 17, 27, 31, 34, 39.

NOTE: Apart from the records from Japanese waters, *Lineus alborostratus* is also known from Vostok Bay, Russia (Kulikova, 1988) and Shandong Province (Qingdao and Yantai), China (Yin et al., 1988). A similar-looking species, *Lineus hiatti* Coe, 1947, is known from Hawaii; the latter can be distinguished from *L. alborostratus* by the head having less distinct anterior white or colorless margins (Coe, 1947: 104).

Lineus albovittatus (Stimpson, 1855)

Meckelia albo-vittata [sic] **Stimpson**, 1855: 382; habitat not recorded, Okinawa Prefecture; originally recorded as “Loo Choo.”

Cerebratulus albovittatus: **Stimpson** 1857: 160; intertidal among algae and in rock crevices, Okinawa Prefecture.

Non Cerebratulus albo-vittatus: Bürger, 1890: 11, pl. 1, fig. 1, pl. 2, figs. 1–8, pl. 8, figs. 153–154 (from Ambon, Indonesia).

Lineus albovittatus: **Iwata**, 1954c: 27–29, figs. 1A, 2; habitat not recorded, Nakanoshima, Tokara Islands, Kagoshima Prefecture. **Crandall et al.**, 2002: 11, 17, 27, 34, 39. *Non Lineus albo-vittatus* [sic]: Punnett, 1900: 578, pl. LXI, figs. 46–47 (from Lifou, New Caledonia, South Pacific); *nec Lineus albovittatus*: Punnett and Cooper, 1909: 7, 14, pl. 1, fig. 10, pl. 2, fig. 14 (from Salomon, Indian Ocean).

NOTE: Two forms have been known by the specific name of either *albovittatus* or *albo-vittatus*. These differ in the shape of the white line across the dorsal surface of the head. The transverse line in one form, reported from Japanese waters, is straight (Stimpson, 1855, 1857; Iwata, 1954c), while it is W-shaped in the other (Bürger, 1890; Punnett, 1900; Punnett and Cooper, 1909). The latter was synonymized with *Lineus tricuspidatus* (Quoy and Gaimard, 1833) by Gibson (1981: 206) and later transferred to *Notospermus* by Riser (1991: 435), and is now known as *Notospermus tricuspidatus*, a species that has not been reported from Japanese waters.

Lineus bipunctatus Takakura, 1898

Lineus bipunctatus **Takakura**, 1898: 335–336, fig. 18: Jôgashima, Kanagawa Prefecture; collected sublitorally among thecatid hydroids from 2–3 fathoms depth. **Crandall et al.**, 2002: 11, 17, 27, 34, 39.

Lineus cancelli Iwata, 1954

Lineus cancelli **Iwata**, 1954b: 34–35, fig. 1; intertidal under stones, Shirahama, Wakayama Prefecture. **Crandall et al.**, 2002: 11, 17, 27, 34, 39.

Lineus caputornatus Takakura, 1898

Lineus caputornatus **Takakura**, 1898: 334–335, fig. 15; obtained sublitorally among thecate hydroids from 2–3 fathoms depth, Jôgashima, Kanagawa Prefecture. **Crandall et al.**, 2002: 11, 17, 27, 34, 39.

Lineus fulvus Iwata, 1954

Lineus fulvus **Iwata**, 1954a: 13, fig. 2C; intertidal among

laminarian holdfasts, Rishiri Island, Hokkaidô Prefecture. **Yamaguchi and Yamada**, 1955: 66, fig. 17-3. **Crandall et al.**, 2002: 11, 18, 27, 34, 39.

Lineus fuscoviridis Takakura, 1898

[Japanese name: midori-himomushi or midori-hera-himomushi]
Lineus fuscoviridis **Takakura**, 1898: 332–333, fig. 13; intertidal under stones on muddy sand, sublittoral between rock crevices from 2–3 fathoms depth, Misaki and Jôgashima, Kanagawa Prefecture and Sunosaki, Tateyama, Chiba Prefecture. **Kaburaki**, 1927: 1665, fig. 3186. **Ohuye**, 1942: 187–191, figs. 1–3. **Kaburaki**, 1947: 1470, fig. 4146. **Ohuye**, 1950: 22. **Iwata**, 1952: 134–136; lower intertidal under stones, Tomioka, Amakusa, Kumamoto Prefecture. **Iwata**, 1954b: 36–37; habitat not recorded, Kada, Wakayama Prefecture. **Utinomi**, 1956: 31, pl. 16, fig. 5. **Iwata**, 1957a: 10–11; habitat not recorded, obtained by His Majesty Emperor Shôwa on 5 June 1928, Koajiro at Misaki, Kanagawa Prefecture. **Iwata**, 1960c: 166, pl. 83, fig. 11. **Utinomi**, 1960: 31, pl. 16, fig. 5. **Satô and Itô**, 1961: 187, fig. 7.1.6. **Inaba**, 1963: 228; sandy to muddy sediment or under stones on gravelly to rocky shores, lower intertidal to shallow sublittoral, commonly found in the Inland Sea of Seto; a specimen collected 21 May 1936 at Hakanjima, Hiroshima Prefecture, is deposited in Mukaishima Marine Biological Station, Hiroshima University, under registration number 70-1. **Iwata**, 1965a: 169, 201. **Kaburaki and Iwata**, 1965: 393, with one figure. **Shiino**, 1969: 93. **Utinomi**, 1969: 31, pl. 16, fig. 5. **Saito and Suzuki**, 1974: 38; intertidal, Niisaki Beach, Kanagawa Prefecture; identified by Dr. Iwata. **Honma and Kitami**, 1978: 15; Sado Island, Nîgata Prefecture. **Ochi**, 1979: 640. **Inaba**, 1988: 225; sandy to muddy sediment or under stones on gravelly to rocky shores, lower intertidal to shallow sublittoral, commonly found in the Inland Sea of Seto. **Miyazawa et al.** 1988: 867–874; intertidal, Iwashijima and Mukaishima, Hiroshima Prefecture; intertidal, Yugeshima, Ehime Prefecture; identified by Dr. T. Hoshino, Mukaishima Marine Biological Station, Hiroshima University. **Iwata**, 1992: 198, pl. 44-7. **Uchida**, 1994: 88, with a color photograph taken in life by Mr. Isamu Soyama in Ôsezaki, Shizuoka Prefecture. **Iwata**, 1997: 55. **Crandall et al.**, 2002: 11, 18, 27, 34, 39.
Lineus fuscoviridis [sic]: **Kikuchi**, 1968: 167; among *Zostera marina*, Tomioka Bay, Amakusa, Kumamoto Prefecture.

Lineus nigrofuscus (Stimpson, 1857)

Cerebratulus nigrofuscus **Stimpson**, 1857: 161; intertidal between stones, Amamiôshima, Kagoshima Prefecture. Originally reported as "Ad insulam 'Ousima' Japoniae Australis; littoralis inter lapillus"; transferred to *Lineus* by Bürger (1904: 102). **Crandall et al.**, 2002: 11, 27, 34, 39.

Lineus nigrostriatus Iwata, 1954

Lineus nigrostriatus **Iwata**, 1954c: 30, fig. 1B; habitat not recorded, Nakanoshima, Tokara Islands, Kagoshima Prefecture. **Crandall et al.**, 2002: 11, 18, 27, 34, 39.

Lineus nipponeensis Senz, 2001

Lineus nipponeensis **Senz**, 2001: 5–13, figs. 1–9; habitat and locality unknown.

TYPE MATERIAL: Holotype, NHMW-EV 17026/3990; paratype, NHMW-EV 17027/3991.

NOTE: The material was collected by Dr. Albrecht von Roretz during his stay in Japan from 1874–1882.

Lineus spatirosus Iwata, 1954

Lineus spatirosus **Iwata**, 1954a: 11–12, fig. 2B; lower intertidal under stones, Akkeshi, Hokkaidô Prefecture.

Yamaguchi and Yamada, 1955: 65, fig. 17-2. **Uchida et al.**, 1963: 17. **Crandall et al.**, 2002: 11, 18, 27, 34, 39.

Lineus subcingulatus Takakura, 1898

[Japanese name: koajiro-himomushi]

Lineus subcingulatus **Takakura**, 1898: 335, fig. 16; intertidal among *Sargassum thunbergii*, Koajiro Bay, Kanagawa Prefecture. **Kaburaki**, 1927: 1665, fig. 3187. **Kaburaki**, 1947: 1471, fig. 4147. **Inaba**, 1963: 228; lower intertidal to shallow sublittoral, commonly occurring on algae, or on gravelly to rocky shores, the Inland Sea of Seto. **Kaburaki and Iwata**, 1965: 394. **Inaba**, 1988: 225; lower intertidal to shallow sublittoral, commonly occurring on algae, or on gravelly to rocky shores, the Inland Sea of Seto. **Crandall et al.**, 2002: 11, 18, 27, 34, 39.

Lineus torquatus Coe, 1901

[Japanese name: kasuri-himomushi or kasuri-hera-himomushi]

Lineus torquatus: **Yamaoka**, 1940a: 218–220, pl. XIV, figs. 9–14, text fig. 7; Hokkaidô Prefecture (Akkeshi, Ochiishi, and Nemuro), lower intertidal under stones. **Okuda**, 1947: 1471, fig. 4148 (1–3). **Iwata**, 1954a: 12; lower intertidal under stones or among laminarian holdfasts, Hokkaidô Prefecture (Akkeshi, Nemuro and Hiroo). **Yamaguchi and Yamada**, 1955: 66. **Utinomi**, 1956: 31, pl. 16, fig. 4. **Iwata**, 1957b: 54–57, figs. 1–10; intertidal under stones, Akkeshi, Hokkaidô Prefecture. **Iwata**, 1957c: 102, figs. 5.1a–c, 5.2. **Iwata**, 1960b: 26. **Iwata**, 1960c: 166, pl. 83, fig. 13. **Utinomi**, 1960: 31, pl. 16, fig. 4. **Uchida et al.**, 1963: 17. **Okuda and Iwata**, 1965: 394, figs. a–c. **Utinomi**, 1969: 31, pl. 16, fig. 4. **Okada et al.**, 1971: 62. **Uchida et al.**, 1972: 62. **Iwata**, 1983: 182, 183, 187, 189, figs. 8-1, 8-2, 8-6. **Crandall et al.**, 2002: 11, 18, 27, 35, 39.

NOTE: Originally reported from Alaska (Coe, 1901: 66); also known to occur in San Francisco Bay, USA (Corrêa, 1964: 528) and Santa Maria Basin, California, USA (Blake, 1993: 118), on the coasts around northern China (Sun and Pan, 1994: 328), and the Japan Sea coast of Russia (Korotkevitsch, 1955, 1971; Kulikova, 1988). Manchenko and Kulikova (1996a) demonstrated on the basis of allozyme analyses that the species is a mixture of at least two cryptic species, which have been recognized since the original description by Coe (1901) as brown and reddish color morphs. A nomenclatural procedure, such as neotypification, may be required in the future to make it clear to which taxon the name *torquatus* will be applied. Even if some syntypes were extant, designating a lectotype would not solve the problem, for the body color might have been changed or not preserved by fixation.

Genus Micrura Ehrenberg, 1831

Micrura Ehrenberg, 1831: 57.

TYPE SPECIES: *Micrura fasciolata* Ehrenberg, 1831, by

monotypic designation.

Micrura akkeshiensis Yamaoka, 1940

[Japanese name: akkeshi-himomushi]

Micrura akkeshiensis Yamaoka, 1940a: 227–228, pl. XV, figs. 11, 12, pl. XVI, fig. 1, text fig. 11; upper intertidal under stones on sandy beaches, Akkeshi and Abashiri, Hokkaidō Prefecture. Okuda, 1947: 1473, fig. 4153. Iwata, 1954a: 14–15; intertidal under stones, Akkeshi and Muroran, Hokkaidō Prefecture. Yamaguchi and Yamada, 1955: 67. Iwata, 1957c: 106. Iwata, 1958: 104–127, figs. 1–60; in July 1955, intertidal, Daikokujima Island, Akkeshi, Hokkaidō Prefecture. Iwata, 1960a: 96, figs. 3, 4. Iwata, 1960b: 27. Iwata, 1960c: 169, pl. 84, fig. 2. Satō and Itō, 1961: 187, fig. 7.1.4. Uchida et al., 1963: 17. Iwata, 1965a: 169, 201. Okuda and Iwata, 1965: 395, figs. a, b. Iwata, 1983: 181, 182, 185, 191, figs. 8–12. Crandall et al., 2002: 12, 18, 28, 35, 40.

NOTE: Apart from the records from Hokkaido, *Micrura akkeshiensis* is also known from Vostok Bay, Russia (Kulikova, 1988).

Micrura alaskensis Coe, 1901

Micrura alaskensis: Yamaoka, 1940a: 225–226, pl. XV, figs. 9, 10, text fig. 10; intertidal under stones on beaches, Akkeshi, Hokkaidō Prefecture. Coe, 1944: 29. Iwata, 1954a: 14; lower intertidal under stones, Akkeshi, Hokkaidō Prefecture. Yamaguchi and Yamada, 1955: 67. Uchida et al., 1963: 17. Crandall et al., 2002: 12, 18, 28, 35, 40.

NOTE: Originally described from Alaska (at New Metlakahtla on Annette Island, Glacier Bay, Sitka, Yakutat, and at Orca and Virgin Bay in Prince William Sound) by Coe (1901: 71), *Micrura alaskensis* is also reported from San Pedro and Monterey Bay, California, USA (Coe, 1904: 118), British Columbia, Canada (Coe, 1940: 271), and Ensenada, Mexico (Coe, 1940: 271); the record from Santa Maria Basin, California, USA (Blake, 1993: 119), based on fixed fragments of bodies collected from depths of 297 m and 591 m, might represent different species.

Micrura bella (Stimpson, 1857)

[Japanese name: kuchibeni-himomushi]

Cerebratulus bellus Stimpson, 1857: 161; obtained from an empty shell from a muddy bottom, 11 m depth, Hokkaidō Prefecture; originally reported as "Prope oras insulae 'Jesso,' in conchis desertis e fundo limoso profunditatis sex orgyiarum."

Micrura festiva Takakura, 1898: 336, fig. 20; sublittoral 2–3 fathoms (=3.7–5.6 m) depth, Jōgashima, Kanagawa Prefecture. Kaburaki, 1927: 1666, fig. 3188. Crandall et al., 2002: 12, 18, 28, 35, 40.

Micrura bella: Yamaoka, 1940a: 230–232, pl. XV, figs. 13–15; on floating rotten wood perforated by the shipworm *Teredo hibicola*, Usu, Hokkaidō Prefecture; habitat not recorded, Muroran, Hokkaidō Prefecture; habitat not recorded, Kushimoto, Wakayama Prefecture. Okuda, 1947: 1472, fig. 4151. Iwata, 1954a: 14; lower intertidal under stones or among holdfasts of seaweeds, Muroran and Oshoro, Hokkaidō Prefecture, Misaki, Kanagawa Prefecture, and Kushimoto, Wakayama Prefecture. Iwata,

1954b: 37; intertidal under stones, Yuzaki, Shirahama, Wakayama Prefecture. Yamaguchi and Yamada, 1955: 67. Utinomi, 1956: 32, pl. 16, fig. 10. Iwata, 1960c: 169, pl. 84, fig. 3; habitat not recorded, Usa, Kōchi Prefecture and Aikawa, Sado Island, Nigata Prefecture. Utinomi, 1960: 32, pl. 16, fig. 10. Okuda and Iwata, 1965: 395, figs. a, b. Utinomi, 1969: 32, pl. 16, fig. 10. Okada et al., 1971: 62. Uchida et al., 1972: 62. Honma and Kitami, 1978: 15; Sado Island, Nigata Prefecture. Crandall et al., 2002: 12, 18, 28, 35, 40.

NOTE: *Micrura bella* possesses a striking body color pattern, which is an off-white background with a deep vermillion tip of the head and a longitudinal dorsal band composed of broad purple rectangles separated by narrow spaces. There are at least three different names for similar-looking forms: *Lineus striatus* Griffin, 1898, *Micrura impressa* Stimpson, 1857, and *Micrura verrilli* Coe, 1901. Crandall et al. (2002) pointed out that the difference among these forms is whether or not the head marking encircles the tip. Crandall et al. (2002) suggested that the head marking is complete in the western-Pacific forms, while it is restricted to the dorsal half in the eastern-Pacific forms. Future assessment will be required to delineate these forms for the proper application of their names.

Micrura dorsovittata Takakura, 1898

Micrura dorsovittata Takakura, 1898: 337, fig. 21; habitat not recorded, Jōgashima, Kanagawa Prefecture. Crandall et al., 2002: 12, 18, 28, 35, 40.

NOTE: *Micrura dorsovittata* is so far known only by its original description. It resembles *Micrura kulikovae* Chernyshev, 1992, a new name given to a form identified as *Micrura bella* by Kulikova and Kutishchev (1984), in which the color of the dorsal band is brownish.

Micrura japonica Iwata, 1952

[Japanese name: kuro-himomushi]

Micrura japonica Iwata, 1952: 139–140, figs. 4, 11, 12; lower intertidal under stones on sandy beaches, Fukue, Gotō Islands, Nagasaki Prefecture; upper intertidal under stones in tide pools on a rocky shore, Tomioka, Amakusa, Kumamoto Prefecture; habitat not recorded, Shimoda, Shizuoka Prefecture. Utinomi, 1956: 32, pl. 16, fig. 9. Iwata, 1957a: 18–19; collected intertidally on 18 January 1930 by His Majesty Emperor Shōwa, Samejima at Hayama, Kanagawa Prefecture. Iwata, 1960c: 169; pl. 84, fig. 4; habitat not recorded, Sakurajima, Kagoshima Prefecture; habitat not recorded, Aikawa, Sado Island, Nigata Prefecture. Utinomi, 1960: 32, pl. 16, fig. 9. Satō and Itō, 1961: 187, fig. 7.1.3. Iwata, 1965b: 396, figs. a, b. Utinomi, 1969: 32, pl. 16, fig. 9. Saito and Suzuki, 1974: 38; intertidal, Niisaki Beach, Kanagawa Prefecture; identified by Dr. Iwata. Honma and Kitami, 1978: 15; Sado Island, Nigata Prefecture. Iwata, 1992: 198–199, fig. 7-3F. Crandall et al., 2002: 12, 18, 28, 35, 40.

NOTE: The distinction between *Micrura japonica* Iwata, 1952 and *Micrura formosana* Yamaoka, 1939 will require future verification; the latter species, described from the northeastern coast of Taiwan, differs from the former by having a rhynchocoel diverticulum protruding ventrally into the lumen of the foregut, a character state that can be

interpreted as an artifact induced during fixation. Yamaoka's (1939) illustration of the external appearance of the preserved specimen shows strong shrinkage on the surface of the body, which suggests that the specimen was not, or not adequately, anaesthetized before fixation. This would further argue for conspecificity of these two nominal species, reinforced by the close proximity of their localities. Furthermore, *M. japonica* and *M. formosana* might be synonymous with *Cerebratulus niger* (Stimpson, 1855), described from Hong Kong, which is similar in having a truncated anterior end, lateral cephalic slits extending back to the mouth region, black body coloration, and white margins around the mouth.

Micrura magna Yamaoka, 1940

Micrura magna Yamaoka, 1940a: 227–228, pl. XVI, figs. 2–4; lower intertidal under stones, Daikokujima, Akkeshi, Hokkaidō Prefecture. Iwata, 1954a: 14. Yamaguchi and Yamada, 1955: 67. Uchida et al., 1963: 17. Crandall et al., 2002: 12, 18, 28, 35, 40.

NOTE: Besides its original description, *Micrura magna* is also known from Vostok Bay, Russia (Kulikova, 1988).

Micrura multinotata Iwata, 1957

Micrura multinotarum [sic] Iwata, 1957a: 19–20, pl. I, fig. 8, pl. V, figs. 8, 9; dredged from 30–40 m depth on 20 January 1949 by His Majesty Emperor Shōwa, Tateishi at Ogashima, Kanagawa Prefecture.

Micrura multinotata: Crandall et al., 2002: 12, 18, 28, 40.

Genus *Nipponomicrura* Chernyshev, 1995

Nipponomicrura Chernyshev, 1995: 15.

TYPE SPECIES: *Micrura uchidai* Yamaoka, 1940, by original designation.

Nipponomicrura uchidai (Yamaoka, 1940)

[Japanese name: uchida-himomushi]

Micrura uchidai Yamaoka, 1940a: 232–234, pl. XVI, figs. 5–7, text fig. 12; lower intertidal under stones, Muroran, Hokkaidō Prefecture. Okuda, 1947: 1472, fig. 4152 (1–3). Iwata, 1954a: 14. Yamaguchi and Yamada, 1955: 67. Okuda and Iwata, 1965: 396, figs. a–c.

Nipponomicrura uchidai: Chernyshev, 1995: 15. Crandall et al., 2002: 13, 18, 28, 35, 40.

NOTE: Besides the original record from Muroran, *Nipponomicrura uchidai* is also known from Vostok Bay, Russia (Kulikova, 1988).

Genus *Notospermus* Huschke, 1830

Notospermus Huschke, 1830: 682.

TYPE SPECIES: *Notospermus drepanensis* Huschke, 1830 (from Sicilia, Italy; now regarded as a junior synonym of *Polia geniculata* Delle Chiaje, 1828) by monotypic designation.

Notospermus geniculatus (Delle Chiaje, 1828)

[Japanese name: misaki-himomushi or kurohera-himomushi]
Lineus mitellatus Takakura, 1898: 333–334, fig. 14; intertidal under stones on muddy sand, sublittoral in rock crevices from 2–3 fathoms depth, Misaki and Jōgashima, Kanagawa Prefecture and Sunosaki, Tateyama, Chiba

Prefecture. Iwata, 1952: 136–137, fig. 13; lower intertidal under stones, Tomioka, Amakusa, Kumamoto Prefecture. *Lineus geniculatus*: Iwata, 1954b: 35–36; intertidal under stones, Kushimoto and Shirahama, Wakayama Prefecture. Utinomi, 1956: 31, pl. 16, fig. 3. Iwata, 1957a: 9–10; habitat not recorded, collected by His Majesty Emperor Shōwa on 16 July, 1939 Najima at Hayama, Kanagawa Prefecture. Iwata, 1960c: 166, pl. 83, fig. 10. Utinomi, 1960: 31, pl. 16, fig. 3. Inaba, 1963: 228; sandy to muddy sediment or under stones on gravelly to rocky shores, lower intertidal to shallow sublittoral, commonly found in the Inland Sea of Seto; a specimen collected on 20 June 1963 at Mukaishima, Hiroshima Prefecture, is deposited in Mukaishima Marine Biological Station, Hiroshima University, under registration number 70-2. Iwata, 1965b: 393, figs. a–c. Utinomi, 1969: 31, pl. 16, fig. 3. Saito and Suzuki, 1974: 38; intertidal, Niisaki Beach, Kanagawa Prefecture; identified by Dr. Iwata. Honma and Kitami, 1978: 15; Sado Island, Niigata Prefecture. Inaba, 1988: 225; sandy to muddy sediment or under stones on gravelly to rocky shores, lower intertidal to shallow sublittoral, commonly found in the Inland Sea of Seto. Iwata, 1992: 198, pl. 44–4, 5. Uchida, 1994: 88–89, with a color photograph taken in life by Mr. Isamu Soyama at Ōsezaki, Shizuoka Prefecture.

?*Lineus genicalatus* [sic]: Iwata, 1997: 53, a color photograph taken in life by Eiichi Kurasawa, possibly depicting a species of the genus *Cerebratulus* that lost its tail.

Notospermus geniculatus: Crandall et al., 2002: 13, 18, 28, 35, 40. Thollesson and Norenburg, 2003: 408; Shirahama, Wakayama Prefecture.

NOTE: Iwata (1954b) synonymized *Lineus mitellatus* sensu Takakura (1898) and sensu Iwata (1952) with *Lineus geniculatus*, which was originally described as *Polia geniculata* by Delle Chiaje (1828: 177) from Naples, Italy. The species has been redescribed as *Notospermus geniculatus* by Riser (1991: 428–434). Outside Japanese waters it is also known from the Black Sea, Mediterranean (France, Italy, Greece, Malta), Canary Is., Gulf of Guinea (West Africa), Australia, New Zealand, and the western coasts of tropical America (Gulf of California, Panama, and Peru) (Gibson, 1995: 480).

Genus *Paralineopsis* Iwata, 1993

Paralineopsis Iwata, 1993: 186.

TYPE SPECIES: *Paralineopsis taki* Iwata, 1993, by original designation.

Paralineopsis taki Iwata, 1993

Zygeupolia littoralis: Iwata, 1951: 135–136; lower intertidal under stones on sand, Mukaishima, Hiroshima Prefecture.

Inaba, 1963: 227. Mukaishima, Hiroshima Prefecture.

Zyreupolia littoralis [sic]: Inaba, 1988: 225. Mukaishima, Hiroshima Prefecture.

Paralineopsis taki Iwata, 1993: 186–199, figs. 1–6.

Paralinoepsis taki [sic]: Crandall et al., 2002: 13, 18, 28, 36.

NOTE: Iwata (1951) originally identified his material as *Zygeupolia littoralis* Thompson, 1900 on the basis of its external features. Later he thoroughly redescribed the taxon as a new genus and species (Iwata, 1993).

Genus *Uchidana* Iwata, 1967*Uchidana* Iwata, 1967: 123.TYPE SPECIES: *Uchidana parasita* Iwata, 1967 by original designation.***Uchidana parasita* Iwata, 1967**

[Japanese name: uchida-kisei-himomushi]

Uchidana parasita Iwata, 1967: 124–136, text fig. 1, pl. 1, figs. 1–8, pl. 2, figs. 9–16, pl. 3, figs. 17–24; in the mantle cavity of *Mactra sulcatoria* in muddy sand, mouth of the River Aikawa, Tsu, Mie Prefecture. Iwata, 1970b: 128, 132. Iwata, 1973: 264. Iwata, 1992: 198, fig. 7–3D. Crandall et al., 2002: 15, 18, 30, 44.NOTE: *Uchidana parasita* is the only heteronemertean species parasitizing bivalves.**Family VALENCINIIDAE Hubrecht, 1879****Genus *Baseodiscus* Diesing, 1850***Polia* Delle Chiaje, 1825: 406; non *Polia* Ochsenheimer, 1816: 73 (Lepidoptera: Noctuidae).*Baseodiscus* Diesing, 1850: 243.*Eupolia* Hubrecht, 1887: 10; synonymized by Bürger (1904).TYPE SPECIES: *Polia delineata* Delle Chiaje, 1825 by monotypic designation.***Baseodiscus curtus* (Hubrecht, 1879)**

[Japanese name: tatejima-himomushi]

Eupolia curta: Takakura, 1898: 185, fig. 7; sublittoral from 2–3 fathoms, Jōgashima, Kanagawa Prefecture; intertidal, Matsuwa Bay, Miura, Kanagawa Prefecture.*Baseodiscus curtus*: Kaburaki, 1927: 1664, fig. 3185.*Kaburaki*, 1947: 1473, fig. 4154. Iwata, 1952: 141–142, fig. 14; lower intertidal under stones, Fukue, Gotō Islands, Nagasaki Prefecture. Utinomi, 1956: 31, pl. 16, fig. 7. Iwata, 1960c: 166, pl. 83, fig. 6. Iwata, 1965a: 169, 201. Iwata, 1965b: 391, one fig. Utinomi, 1969: 31, pl. 16, fig. 7. Crandall et al., 2002: 9, 16, 26, 33, 38.*Baseodiscus delineatus* var. *curtus*: Utinomi, 1960: 31, pl. 16, fig. 7. Saito and Suzuki, 1974: 38; intertidal, Niisaki Beach, Kanagawa Prefecture; identified by Dr. Fumio Iwata.NOTE: *Baseodiscus curtus*, originally described as *Polia curta* Hubrecht, 1879, from Naples, Italy, was transferred to the genus *Baseodiscus* by Bürger (1904). *Baseodiscus curtus* was synonymized with *B. delinatus* by Gibson (1979). However, *B. curtus* can be distinguished from *B. delineatus* by lacking stripes on the ventral surface of the body (Hubrecht, 1879: 209). Although these two species have been regarded as conspecific (e.g., Gibson, 1995), they are treated as distinct species in the present paper. These two species appear to occur globally (Gibson, 1995), with virtually completely overlapping ranges of distribution (Coe, 1944).***Baseodiscus delineatus* (Delle Chiaje, 1825)**

[Japanese name: iso-himomushi]

?*Eupolia* sp. Takakura, 1898: 185, fig. 8; intertidal, Moroiso and Mwatsuwa, Kanagawa Prefecture.*Baseodiscus delineatus*: Coe, 1944: 28. Crandall et al., 2002: 9, 16, 26, 33, 38.?*Baseodiscus takakurai* Gibson, 1995: 305.NOTE 1: *Baseodiscus delineatus* was originally described as *Polia delineata* Delle Chiaje, 1825 from Naples, Italy, then transferred to *Baseodiscus* by Diesing (1850: 243). The species can be distinguished from *B. curtus* by its either striped or mottled ventral body surface. The species shows a circumglobal distribution; apart from the records from Japan, *Baseodiscus delineatus* is also known from the Mediterranean, the Adriatic and Atlantic coasts of Europe, Cape Verde Is., Bermuda, Barbados, southern Florida, USA, Puerto Rico, Gulf of California, Fiji Is., Mariana Is., Java, Torres Straits, Australia (the Great Barrier Reef and southern coast of Western Australia), Mauritius, Zanzibar, Brazil, and Chile (Gibson, 1995: 479).NOTE 2: Takakura (1898) recorded a form as *Eupolia* sp. that possessed black mottles on both the dorsal and ventral surfaces of the body. Similar specimens collected on Kakeroma Island (Kagoshima Prefecture) and Ishigaki Island (Okinawa Prefecture) show mottles becoming gradually fused together to form incomplete longitudinal stripes in the middle region of the body (Kajihara, pers. obs.). Takakura's *Eupolia* sp. is herein tentatively regarded as *Baseodiscus delineatus*, though future study must confirm this identification. Takakura (1898) did not identify his material to species, mentioning that it resembled *Eupolia antillensis* Bürger, 1895. Gibson (1995) misinterpreted this as Takakura's (1898) having established a new taxon with the specific name *antillensis*, and superfluously gave the new name *Baseodiscus takakurai* for Takakura's (1898) *Eupolia* sp.***Baseodiscus hemprichii* (Ehrenberg, 1831)**

[Japanese name: sanada-himomushi]

Baseodiscus hemprichi [sic]: Iwata, 1954b: 37; habitat not recorded, Tōshima, Shirahama, Wakayama Prefecture. Iwata, 1954c: 30–31; habitat not recorded, Nakanoshima, Tokara Islands, Kagoshima Prefecture. Utinomi, 1956: 31, pl. 16, fig. 8. Utinomi, 1960: 31, pl. 16, fig. 8. Utinomi, 1969: 31, pl. 16, fig. 8.*Baseodiscus hemprichii*: Iwata, 1960c: 166, pl. 83, fig. 7. Ooishi, 1964: 193; intertidal among dead reef corals, Ushuku, Amamiōshima, Kagoshima Prefecture. Okuda and Iwata, 1965: 392, figs. a, b. Crandall et al., 2002: 10, 16, 24, 26, 30, 33, 38.NOTE: Originally described as *Nemertes hemprichii* Ehrenberg, 1831 from the Red Sea, subsequently transferred to *Baseodiscus* by Bürger (1904: 83). Apart from the records from Japanese waters and its type locality, this species is widely known from India, Pakistan, East Africa (off Mozambique and Zanzibar), Maldives, Laccadive Islands, Coetivy Island, Mauritius, Malay Peninsula, Java, Ambon, Taiwan, Australia (Great Barrier Reef), Papua New Guinea, New Britain (Solomon Is.), Loyalty Is., Caroline Is., Wake Is., West Samoa, and Hawaiian Islands (Gibson, 1995: 432–433).***Baseodiscus nippensis* (Hubrecht, 1887)***Eupolia nippensis* Hubrecht, 1887: 14–15, pl. I, figs. 4, 5, 10, pl. VII, figs. 6, 11, 12; dredged from 345 fathoms (about 640 m) depth on green mud, collected on 12 May 1875 by H.M.S. Challenger, 35°11'00"N, 139°28'00"E, Sagami Bay, off Kanagawa Prefecture; transferred to

Baseodiscus by Bürger (1904: 84).

Baseodiscus nippensis: Crandall et al., 2002: 10, 16, 26, 38.

NOTE: *Baseodiscus nippensis* has not been reported since its original description. The species was described from preserved body fragments.

***Baseodiscus princeps* (Coe, 1901)**

[Japanese name: arasuka-himomushi]

Baseodiscus curtus: Yamaoka, 1940a: 234–236, pl. XVI, fig. 8–11, text fig. 13; lower intertidal between or under stones, Daikokujima, Akkeshi, Hokkaidō Prefecture; synonymized by Iwata (1954a: 15).

Baseodiscus princeps: Coe, 1944: 28, obtained by the United States Bureau of Fisheries Steamer *Albatross*, 340 m depth, south of Hokkaidō Island; 250 m depth, off Ōshima, the Metropolis of Tōkyō; 260 m depth, off Ōsezaki, Izu Peninsula, Shizuoka Prefecture; 135–290 m depth, Sea of Japan. Iwata, 1954a: 15; lower intertidal under stones, Akkeshi, Hokkaidō Prefecture. Yamaguchi and Yamada, 1955: 68. Uchida et al., 1963: 17. Iwata, 1965b: 392, figs. a–c. Crandall et al., 2002: 10, 16, 26, 33, 38.

NOTE: Originally described as *Taeniosoma princeps* Coe, 1901 from Alaska (Cape Fox, Yakutat, and Orca in Prince William Sound), transferred to *Baseodiscus* by Coe (1940: 262). Also known from Puget Sound, Washington, USA (Coe, 1944: 28).

***Baseodiscus quinque-lineatus* (Quoy and Gaimard, 1833)**

[Japanese name: kurosuji-himomushi]

Taeniosoma aequale Stimpson, 1857: 162; intertidal under stones, Amamiōshima, Kagoshima Prefecture. The locality and habitat were originally recorded as “in situ insulae ‘Ousima;’ littorale sub lapidibus”; synonymized by Bürger (1904: 83).

Baseodiscus quinque-lineatus: Iwata, 1992: 197, pl. 44-2. Crandall et al., 2002: 10, 16, 26, 33, 38.

NOTE: Originally described as *Borlasia quinque-lineata* by Quoy and Gaimard (1833: 285) from Dorey, New Guinea, transferred to *Baseodiscus* by Bürger (1904: 83). *Baseodiscus quinque-lineatus* is distributed in the western Pacific (Japan, Singapore, Indonesia [Java, Timor, Ambon], New Guinea, Solomon Is., Loyalty Is., Torres Straits, and east coast of Australia, including the Great Barrier Reef) (Gibson, 1995: 310).

Genus *Cephalomastax* Iwata, 1957

Cephalomastax Iwata, 1957a: 5.

TYPE SPECIES: *Cephalomastax brevis* Iwata, 1957 by monotypic designation.

***Cephalomastax brevis* Iwata, 1957**

[Japanese name: amadaiba-himomushi]

Cephalomastax brevis Iwata, 1957a: 5–7, pl. I, fig. 9, pl. II, fig. 7, pl. III, figs. 1–7; dredged from 200–300 m depth on 7 August 1953 by His Majesty Emperor Shōwa, “Southern Maruyamadashi at Aamadaiba” [sic], Sagami Bay, off Kanagawa Prefecture. Iwata, 1965a: 201. Iwata, 1965b: 392, figs. a, b. Iwata, 1992: 197, fig. 7-3C. Crandall et al., 2002: 10, 17, 26, 38.

Class HOPLONEMERTEA Hubrecht, 1879

Subclass MONOSTILIFERA Brinkmann, 1917

Family AMPHIPORIDAE McIntosh, 1874

Genus *Amphiporus* Ehrenberg, 1831

Amphiporus Ehrenberg, 1831: 63.

TYPE SPECIES: *Planaria lactiflorea* Johnston, 1828, designated under the plenary power of the ICBN (ICZN, 1992); all previous designations of *Amphiporus albicans* as the type species for *Amphiporus*, including that of Friedrich (1955: 154), have thereby been set aside.

***Amphiporus antifuscus* Iwata, 1954**

Amphiporus antifuscus Iwata, 1954a: 24–25, fig. 6A; lower intertidal among algae, Akkeshi, Hokkaidō Prefecture.

Yamaguchi and Yamada, 1955: 70. Uchida et al., 1963: 17. Crandall et al., 2002: 9, 19, 25, 33.

NOTE: Gibson and Crandall (1989) listed this form as a species inquirenda.

***Amphiporus formidabilis* Griffin, 1898**

Amphiporus cervicalis: Yamaoka, 2005: 143, pl. 1, figs. 2, 4, text fig. 2; Susaki, Sotoura, and Manazuru, near Shimoda, Shizuoka Prefecture; Muroran, Hokkaidō Prefecture.

Amphiporus formidabilis: Iwata, 1952: 144–146, figs. 15, 16; intertidal under stones on sandy beaches, Tomioka, Amakusa Islands, Kumamoto Prefecture and Fukue, Gotō Islands, Nagasaki Prefecture. Okuda, 1947: 1467, fig. 4137. Crandall et al., 2002: 9, 19, 26, 33.

NOTE: *Amphiporus formidabilis* Griffin, 1898 was originally described from Puget Sound and Alaska, USA. Coe (1904: 115) synonymized *Amphiporus exilis* Coe, 1901 with *A. formidabilis*, but Gibson and Crandall (1989) regarded these taxa as species inquirendae, retaining them as separate species. *Amphiporus formidabilis* has also been reported from the Aleutian Islands (Coe, 1905: 252). Some earlier records under the name *Amphiporus cervicalis* (Stimpson, 1857) from Japanese waters may represent *Amphiporus formidabilis*.

***Amphiporus gelatinosus* Coe, 1905**

Amphiporus gelatinosus: Coe, 1944: 30; obtained by the United States Bureau of Fisheries Steamer *Albatross*, at 130 m depth in Uraga Strait, between the Metropolis of Tōkyō and Chiba Prefecture. Crandall et al., 2002: 9, 19, 26, 37.

NOTE: The original description by Coe (1905: 259) was based on a single specimen dredged by *Albatross* on 9 August 1888 at Station 2853, 56°00'N, 154°20'W, southwest of Kadiak Island, Alaska, at a depth of 159 fathoms (=290 m). Known to occur from Alaska to Puget Sound, Washington State, North America (Gibson, 1995: 283). Gibson and Crandall (1989) included it as a species inquirenda.

***Amphiporus imparispinosus* Griffin, 1898**

Amphiporus imparispinosus: Yamaoka, 2005: 145, text figs. 3, 4a; Sotoura and Mikimoto Island, near Shimoda, Shizuoka Prefecture.

NOTE: *Amphiporus imparispinosus* was originally described from Port Townsend, Washington and Sitka, Alaska, USA by Griffin (1898). The species is distributed from San

Pedro, California, to Puget Sound, Alaska, to the Commander Islands off the coast of Kamchatka, to the Bering Strait (Coe, 1905: 249). Some forms reported as *Amphiporus cervicalis* (Stimpson, 1857), *Amphiporus depressus* (Stimpson, 1857), and *Amphiporus lactifloreus* (Johnston, 1828) from Japanese waters appear to represent *A. imparispinosus*. *Pantinonemertes daguillarensis* Gibson and Sundberg, 1992, described from Hong Kong, might be conspecific with *Amphiporus imparispinosus*. The taxonomic identity of this species should be delineated by future studies.

Amphiporus insolitus Iwata, 1954

Amphiporus insolitus Iwata, 1954b: 39–41; lower intertidal under stones, Kushimoto, Wakayama Prefecture.

Crandall et al., 2002: 9, 19, 26, 33.

NOTE: Gibson and Crandall (1989) regarded this form as a species *inquirenda*. *Amphiporus insolitus* resembles *Diplomma serpentina* (Stimpson, 1855) in body coloration, shape of the head, and arrangement of the eyes; these two may be conspecific.

Amphiporus musculus Iwata, 1954

Amphiporus musculus Iwata, 1954a: 26–27, fig. 6B; intertidal among algal holdfasts, Oshoro, Hokkaidō Prefecture.

Yamaguchi and Yamada, 1955: 70. Okada et al., 1971: 62. Uchida et al., 1972: 62. Crandall et al., 2002: 9, 19, 26, 38.

NOTE: Listed as a species *inquirenda* by Gibson and Crandall (1989).

Amphiporus parvus Yamaoka, 1940

Amphiporus parvus Yamaoka, 1940a: 243–244, pl. XVII, fig. 7, text figs. 20, 21; intertidal on sandy beaches, Akkeshi, Hokkaidō Prefecture. Iwata, 1954a: 19. Yamaguchi and Yamada, 1955: 69. Uchida et al., 1963: 17. Crandall et al., 2002: 9, 19, 26, 33.

NOTE: Listed as a species *inquirenda* by Gibson and Crandall (1989).

Amphiporus reduncus Iwata, 1957

Amphiporus reduncus Iwata, 1957a: 23–24; dredged subtidally from 100–130 m depth on 7 August 1950 by His Majesty Emperor Shōwa, northeastern Nakafukari near Nagai, Sagami Bay, off Kanagawa Prefecture. Crandall et al., 2002: 9, 19, 26, 33, 38.

NOTE: Gibson and Crandall (1989) regarded this form as a species *inquirenda*, with the comment that it may be related to the genus *Nipponnemertes* or some similar taxon. It might also prove to be a reptantic polystiliferan (Crandall, pers. comm.).

Amphiporus regius Iwata, 1954

Amphioprus regius Iwata, 1954a: 27–29, fig. 7; lower intertidal under stones on rocky shores, Muroran, Hokkaidō Prefecture. Yamaguchi and Yamada, 1955: 70. Crandall et al., 2002: 9, 19, 26, 33.

NOTE: Listed as a species *inquirenda* by Gibson and Crandall (1989). The presence of four well-developed eyes and a cephalic patch on the dorsal surface of the head indicates that this species may belong to the genus *Tetrastramma* or a related taxon.

Amphiporus retrotumidus Iwata, 1957

Amphiporus retrotumidus Iwata, 1957a: 25–27, pl. I, fig. 13; dredged from 30–55 m depth on 6 August 1953 by His Majesty Emperor Shōwa, "Shuragane at Hayama" [sic], Kanagawa Prefecture. Crandall et al., 2002: 9, 19, 26, 38.

NOTE: Gibson and Crandall (1989) regarded this form as a species *inquirenda*, with the comment that it may be related to the genus *Nipponnemertes* or some similar taxon. It might also prove to be a reptantic polystiliferan (Crandall, pers. comm.).

Genus *Potamostoma* Kajihara, Gibson, and Mawatari, 2003

Potamostoma Kajihara et al., 2003: 492.

TYPE SPECIES: *Potamostoma shizunaiense* Kajihara, Gibson, and Mawatari, 2003, by original designation.

NOTE: The familial affiliation of the genus, not referred to in its original description, is here provisionally designated as the Amphiporidae because of similarities in eye pattern (multiple and grouped) and the nature of the rhynchocoel wall (two-layered). Confirmation of this familial placement will have to be resolved by future studies, hopefully involving molecular data.

Potamostoma shizunaiense Kajihara, Gibson, and Mawatari, 2003

Potamostoma shizunaiense Kajihara et al., 2003: 491–500, figs. 1–7, tabs. 1–2; under stones on a sandy bottom, salinity at ebb tide 2 psu, mouth of the River Shizunai, 42°20'N, 142°22'E, Shizunai, Hokkaidō Prefecture.

TYPE MATERIAL: Holotype ZIHU-2037, immature female, complete series of transverse sections, 7 µm, 81 slides. Paratypes: ZIHU-1930, immature male, transverse sections of the anterior body region, 6 µm, 39 slides; ZIHU-2040, immature female, complete series of transverse sections, 6 µm, 84 slides. Eight sectioned voucher specimens are also accessioned as ZIHU-1931, -1932, -2038, -2039, -2041, -2042, -2043, and -2044. One voucher specimen, longitudinal sections of the anterior body region, 10 slides, is deposited under NHMW-EV 19875.

Genus *Zygonemertes* Montgomery, 1897

Zygonemertes Montgomery, 1897: 2.

TYPE SPECIES: *Amphiporus virescens* Verrill, 1879 by monotypic designation.

Zygonemertes glandulosa Yamaoka, 1940

[Japanese name: fujikasa-himomushi]

Zygonemertes glandulosa Yamaoka, 1940a: 244–247, pl. XVII, fig. 8, text figs. 22–24; lower intertidal under stones, Akkeshi, Hokkaidō Prefecture. Okuda, 1947: 1468, fig. 4139 (1–4). Iwata, 1954a: 19. Yamaguchi and Yamada, 1955: 69. Uchida et al., 1963: 17. Crandall et al., 2002: 15, 22, 30, 37, 42.

Zygonemertes grandulosa [sic]: Iwata, 1960c: 169, pl. 84, fig. 12. Iwata, 1965a: 217. Okuda and Iwata, 1965: 399, figs. a–d. Iwata, 1992: 202, fig. 7–5C.

Zygonemertes jamsteci Kajihara, 2002

Zygonemertes jamsteci Kajihara, 2002: 131–140, figs. 6–9; about 1 m deep, among eelgrass (*Zostera marina*),

Akkeshi Bay, Hokkaidô Prefecture.

TYPE MATERIAL: Holotype, ZIHU-1928, mature male, full series of transverse sections, 43 slides. Paratypes: ZIHU-1929, mature female, serial longitudinal sections of anterior portion of the body, nine slides; ZIHU-2045, mature male, full series of transverse sections, 41 slides; ZIHU-2046, mature male, full series of transverse sections, 36 slides; ZIHU-2047, mature female, serial transverse sections of anterior portion of the body, three slides; ZIHU-2048, mature female, serial transverse sections of the anterior portion of the body, 14 slides. All collected on 8 July 1997.

Zygonemertes shintai Kajihara, 2002

Zygonemertes shintai Kajihara, 2002: 122–131, figs. 1–5; intertidal, among blue mussels (*Mytilus trossulus* Gould), Oshoro, Hokkaidô Prefecture.

TYPE MATERIAL: Holotype, ZIHU-1296, immature, full series of transverse sections, 16 slides, collected on 2 July 1998. Paratypes: ZIHU-1927, immature, full series of transverse sections, 31 slides, collected on 2 July 1998, ZIHU-2105, -2106; whole specimens, preserved in 100% EtOH for molecular analyses, collected on 3 March 2001.

Family CARCINONEMERTIDAE Sumner, Osburn and Cole, 1913

Genus *Carcinonemertes* Coe, 1902

Carcinonemertes Coe, 1902: 440.

TYPE SPECIES: *Nemertes carcinophilus* [originally spelled *cartinophilus*] Kölliker, 1845, by subsequent designation of Friedrich (1955: 176).

Carcinonemertes mitsukurii Takakura, 1910

[Japanese name: kani-himomushi]

Carcinonemertes mitsukurii Takakura, 1910: 111–116, figs. 1–4; obtained from egg masses of the crab *Eriocheir japonicus* De Haan, mouth of the River Minatogawa, Tateyama Bay, Chiba Prefecture. **Satô and Itô**, 1961: 187, fig. 7.1.12. **Kaburaki and Iwata**, 1965: 397, with one figure. **Shiino**, 1969: 94. **Iwata**, 1992: 203. **Crandall et al.**, 2002: 10, 19, 26, 30.

Emplectonema mitsukurii: **Kaburaki**, 1927: 1663, fig. 3182. **Kaburaki**, 1947: 1466, fig. 4132.

Carcinonemertes mitukuri [sic]: **Iwata**, 1965a: 218.

NOTE: Apart from the Japanese record, *Carcinonemertes mitsukurii* is also known from San Andreas Is. (between Marinduque and Luzon), Hong Kong, Hawaiian Is., Kingsmill Is., Society Is., and Singapore (Humes, 1942).

Family CRATENEMERTIDAE Friedrich, 1968

Genus *Nipponnemertes* Friedrich, 1968

Nipponnemertes Friedrich, 1968: 34.

TYPE SPECIES: Friedrich (1968) did not designate the type species, and the name *Nipponnemertes* was thus unavailable. This introduced nomenclatural confusion. Gibson and Crandall (1989: 463) designated *Amphiporus drepanophoroides* Griffin, 1898 as the type species. Gibson (1995: 442) later indicated *Nipponnemertes pulchra* (Johnston, 1837) as the type species. Finally, Crandall (2001: 106) designated *Amphiporus punctatulus* Coe, 1905. Crandall (2001: 106) was correct in that Gibson's (1995) listing of

Nipponnemertes pulchra as the type species of the genus was invalid, since the nominal species *Nemertes pulchra* Johnston, 1837 was not included when the genus was established. The nominal species *Amphiporus drepanophoroides*, first designated by Gibson and Crandall (1989), is eligible for the type species of the genus, but Crandall (2001: 106) discussed that its original description is too brief and its type specimen has been lost by shipwreck. Confusion remains as to whether the author of the name is Gibson and Crandall (1989) or Crandall (2001) (Chernyshev, pers. comm.). Here, I follow the prevailing usage of the authority and date of the name *Nipponnemertes* as Friedrich (1968).

Nipponnemertes bimaculata (Coe, 1901)

[Japanese name: rishiri-himomushi]

Amphiporus bimaculatus: **Iwata**, 1954a: 21–22, fig. 5D; sublittoral among laminarian holdfasts, Rishiri Island, Hokkaidô Prefecture. **Yamaguchi and Yamada**, 1955: 69. **Iwata**, 1960c: 169, pl. 84, fig. 14. **Iwata**, 1965b: 399, figs. a–e.

Nipponnemertes bimaculata **Crandall et al.**, 2002: 12, 20, 28, 35, 40.

NOTE: The species, originally described as *Amphiporus bimaculatus* Coe, 1901 from Victoria, B.C., Canada, and Sitka, Alaska and Puget Sound, Washington, USA, was later transferred to the genus *Nipponnemertes* by Friedrich (1968). Crandall et al. (2002) noted that the form identified by Iwata (1954a) differs from Coe's (1901) taxon in having a pair of quadrangular head markings, rather than the long-triangular markings of Coe's form, and a quite different proboscis central armature.

Nipponnemertes ogumai (Yamaoka, 1947)

[Japanese name: oguma-himomushi]

Amphiporus ogumai **Yamaoka**, 1947: 1468, fig. 4138 (1–3). **Okuda and Iwata**, 1965: 399, figs. a–c.

Nipponnemertes ogumai: **Crandall et al.**, 2001: 179–180, pl. 1, figs. 8–10, pl. 2, fig. 18. **Crandall et al.**, 2002: 12, 20, 28, 35, 40.

NOTE: Crandall et al. (2001) mentioned that the type locality for this species was not specified in the original manuscript; Yamaoka (1947) obtained specimens on sandy beaches at Itado, near Shimoda, Shizuoka Prefecture and Seto, Kishû (probably Shirahama, Wakayama Prefecture).

Nipponnemertes punctatula (Coe, 1905)

[Japanese name: madara-himomushi]

Amphiporus nebulosus: **Takakura**, 1933: 226–227; Kitaura, Alaid Island (=Atlasova), Kurile Islands. **Non Crandall et al.**, 2002: 9, 19, 24, 26, 33.

Amphiporus punctatulus: **Iwata**, 1951: 137–138, figs. 1, 2D; intertidal under stones, Mukaishima, Hiroshima Prefecture.

Iwata, 1952: 143–144; lower intertidal under stones, Tomioka, Amakusa Islands, Kumamoto Prefecture and Fukue, Gotô Islands, Nagasaki Prefecture. **Iwata**, 1954a: 22–23; intertidal among algal holdfasts, Oshoro and Rishiri Island, Hokkaidô Prefecture. **Iwata**, 1954b: 39; intertidal under stones, Shirahama, Wakayama Prefecture. **Yamaguchi and Yamada**, 1955: 70. **Utinomi**, 1956: 32, pl. 16, fig. 12. **Iwata**, 1957a: 21–22; obtained by a hand-

reeled net on 16 March 1956 by His Majesty Emperor Shōwa, Samejima at Hayama, Kanagawa Prefecture. **Iwata**, 1960c: 169, pl. 84, fig. 13. **Utinomi**, 1960: 32, pl. 16, fig. 12. **Inaba**, 1963: 228; lower intertidal to shallow sublittoral, under stones on gravelly to rocky shores, commonly found in the Inland Sea of Seto. **Iwata**, 1965a: 169. **Iwata**, 1965b: 400, figs. a–c. **Utinomi**, 1969: 32, pl. 16, fig. 12. **Saito and Suzuki**, 1974: 38; intertidal, Niisaki Beach, Kanagawa Prefecture; identified by Dr. Iwata. **Inaba**, 1988: 226; lower intertidal to shallow sublittoral, under stones on gravelly to rocky shores, commonly found in the Inland Sea of Seto.

Amphiporus punctatus [sic]: **Okada et al.**, 1971: 62. **Kikuchi**, 1968: 167; among *Zostera marina*, Tomioka Bay, Amakusa, Kumamoto Prefecture.

Amphiporus punctatulus [sic]: **Honma and Kitami**, 1978: 15; Sado Island, Niigata Prefecture.

Cratenemertes punctatulus: **Iwata**, 1992: 202, pl. 44–9, fig. 7–E.

Nipponnemertes punctatulus [sic]: **Uchida et al.**, 1972: 55; habitat not recorded, Horomui, Hokkaidō Prefecture.

Iwata, 1997: 53 (with a color drawing), 55. **Shimomura et al.**, 2001: 46; intertidal on a rocky shore, Akahama, Ōtsuchi Bay, Iwate Prefecture. **Thollesson and Norenburg**, 2003: 408; Oshoro, Hokkaidō Prefecture.

Nipponnemertes punctatula: **Crandall et al.**, 2002: 12, 20, 25, 28, 35, 40.

NOTE: Takakura's (1933) record of *Amphiporus nebulosus* Coe, 1901 from the Kurile Islands was regarded as *Amphiporus punctatulus* by Iwata (1951), whereas *Amphiporus nebulosus* s. str., known only from its type locality Kukak Bay, Alaska Peninsula, was regarded as a species *inquirenda* by Gibson and Crandall (1989). Crandall et al. (2002) noted that there are two coexisting species of cratenemertids in Japanese waters that possess a brown dorsal blotch pattern, *Nipponnemertes arenaria* (Uschakov, 1927) and *Nipponnemertes punctatula* (Coe, 1905), and the records of the latter by Iwata in the 1950s were probably of *Nipponnemertes arenaria*.

Family EMPLECTONEMATIDAE Bürger, 1904

Genus *Emplectonema* Stimpson, 1857

Emplectonema Stimpson, 1857: 163.

TYPE SPECIES: *Emplectonema viride* Stimpson, 1857, was originally described from San Francisco, USA, and now is regarded as a junior synonym of *Nemertes gracilis* Johnston, 1837, by subsequent designation of Friedrich (1955: 172).

Emplectonema buergeri Coe, 1901

Emplectonema buergeri [sic]: **Coe**, 1944: 29, obtained by the United States Bureau of Fisheries Steamer Albatross, 250 m depth, off Ōshima, the Metropolis of Tōkyō.

Emplectonema buergeri: **Crandall et al.**, 2002: 11, 20, 24, 27, 34, 39.

NOTE: Originally described from Sitka and Glacier Bay, Alaska by Coe (1901: 28), known to be distributed in North Pacific (Japan, Pribilof Islands, Bering Sea, and the coast of North America from Alaska to California) (Gibson, 1995: 362).

Emplectonema gracile (Johnston, 1837)

[Japanese name: hoso-midori-himomushi]

Emplectonema gracile: **Yamaoka**, 1940a: 237–238, pl. XVII, figs. 1, 2, text fig. 14; lower intertidal on the surfaces of stones, Daikokujima, Akkeshi, Hokkaidō Prefecture. **Iwata**, 1954a: 15; intertidal on the surfaces of stones or in rock crevices, Akkeshi and Muroran, Hokkaidō Prefecture. **Yamaguchi and Yamada**, 1955: 68. **Iwata**, 1957c: 102. **Iwata**, 1960a: 96, fig. 5. **Iwata**, 1960b: 27–35, figs. 97–122; intertidal, Akkeshi, Hokkaidō Prefecture. **Iwata**, 1960c: 169, pl. 84, fig. 7; habitat not recorded, Asamushi, Aomori Prefecture. **Uchida et al.**, 1963: 17. **Iwata**, 1965a: 169, 218. **Iwata**, 1965b: 396, figs. a, b. **Tsuchiya**, 1979: 82; intertidal, Hadakajima Island, Asamushi, and Aomori Harbor, Aomori Prefecture. **Iwata**, 1983: 181, 182, 184, figs. 8–5j, i, 8–13c. **Iwata**, 1992: 202–203, fig. 7–4K, 7–5D. **Crandall et al.**, 2002: 11, 20, 27, 34, 39.

NOTE: Originally described as *Nemertes gracilis* from the British Isles by Johnston (1837); transferred to *Emplectonema* by Verrill (1895: 528). Other than Japanese waters, the species is widely distributed in the northern hemisphere: Peter the Great Bay, Aleutian Islands, Pacific coast of North America, northern coast of Europe, Mediterranean, Rumanian coast of the Black Sea, and Madeira (Gibson, 1995: 432).

Emplectonema kandai Kato, 1939

[Japanese name: hikari-himomushi]

Emplectonema kandai **Kato**, 1939: 251–253, pl. XXXII, figs. 1–6; sublittoral on the tunic of *Chelyosoma siboga* collected from sandy or muddy bottom, 30–40 m depth, near Asamushi Marine Biological Station, Aomori Bay, Aomori Prefecture. **Kanda**, 1939: 166–173, figs. 1–4. **Kato**, 1947: 1466, fig. 4134. **Satō and Itō**, 1961: 187, fig. 7.1.13. **Iwata**, 1965a: 218. **Kato and Iwata**, 1965: 396, figs. a, b. **Crandall et al.**, 2002: 11, 20, 27, 39.

Emplectonema candai [sic]: **Iwata**, 1970b: 129.

NOTE: *Emplectonema kandai* is so far the only known luminescent species in the phylum.

Emplectonema mitsuii Yamaoka, 1947

[Japanese name: mitsui-himomushi]

Emplectonema mitsuii **Yamaoka**, 1947: 1466, fig. 4133; intertidal among algae on rocky shores in southern Japan. **Satō and Itō**, 1961: 187, fig. 7.1.7. **Okuda and Iwata**, 1965: 397, figs. a–d. **Crandall et al.**, 2001: 177–178, pl. 1, figs. 1–4, pl. 2, figs. 16, 16a, 16b. **Crandall et al.**, 2002: 11, 20, 27, 34.

NOTE: Crandall et al. (2001) introduced Yamaoka's original data on the habitat and locality of this species as intertidal among rockweeds at Susaki, Sotoura, and Mikimoto Island, near Shimoda, Shizuoka Prefecture.

Genus *Nemertopsis* Bürger, 1895

Nemertopsis Bürger, 1895: 548.

TYPE SPECIES: *Nemertes peronea* Quatrefages, 1846 (now regarded as a junior synonym of *Polia bivittata* Delle Chiaje, 1841) by subsequent designation of Friedrich (1955: 173).

NOTE: Chernyshev (pers. comm.) indicated that the genus *Nemertopsis* Bürger, 1895 has a senior subjective syn-

onym, *Colpocephalus* Diesing, 1850 (type species *Borlasia quadripunctata* Quoy and Gaimard, 1833). As far as I am aware, the name *Colpocephalus* Diesing, 1850 has not been used as valid since the year 1899, meeting the condition in Article 23.9.1.1 of the Code (ICZN, 1999). The junior subjective synonym *Nemertopsis* was used during the decade from 1989 to 1998 in the following 27 works, published by 25 authors, and thus meets the condition in Article 23.9.1.2 of the Code (ICZN, 1999): Fish and Fish (1989), Morton (1989), Riser (1989), Britton (1990), Gibson (1990a, b, 1997a,b, 1998), Gibson and Knight-Jones (1990), Turbeville (1991), Iwata (1992), Roe (1993), Hansson (1994), Henry and Martindale (1994, 1996, 1997a, b), Sun and Pan (1994), Walker (1994), Martindale and Henry (1995), Senz (1997b), Boyer and Henry (1998), Envall (1998), Hochberg and Lunianski (1998), Norenburg and Roe (1998), and Stricker and Folsom (1998). The name *Nemertopsis* Bürger, 1895 is herein regarded to have precedence over *Colpocephalus* Diesing, 1850, whenever the two names are considered to be synonymous, according to Article 23.9.2 of the Code (ICZN, 1999).

***Nemertopsis mitellicola* Kajihara, 2007**

Nemertopsis mitellicola **Kajihara**, 2007a: 51–57, figs. 7–11; among the gooseneck barnacle, *Capitulum mitella* (Linnaeus), Shirahama, Wakayama Prefecture.
TYPE MATERIAL: Holotype, ZIHU-3204, serial transverse sections of the complete body, total 52 slides: 6 µm, anterior end of body (1 cm long), 12 slides; 8 µm, rest of the body, 40 slides. Paratypes: ZIHU-3205, serial transverse sections of head (1.5 cm long), 8 µm, 15 slides; ZIHU-3206, serial longitudinal sections, 12 µm, 12 slides.

***Nemertopsis quadripunctata* (Quoy and Gaimard, 1833)**

[Japanese name: yotsume-himomushi]

Nemertopsis gracilis: **Iwata**, 1954b: 38–39, fig. 2A; in the mantle cavity of *Capitulum mitella* (Linnaeus), Shirahama, Wakayama Prefecture. **Utinomi**, 1956: 32, pl. 16, fig. 14. **Iwata**, 1960c: 169, pl. 84, fig. 8; Onomichi, Hiroshima Prefecture; Tomioka, Amakusa Islands, Kumamoto Prefecture; Cape Muroto, Kōchi Prefecture. **Utinomi**, 1960: 32, pl. 16, fig. 14. **Inaba**, 1963: 228; upper to mid intertidal, in the mantle cavity of *Capitulum mitella* (Linnaeus), the Inland Sea of Seto. **Iwata**, 1965a: 218. **Iwata**, 1965b: 397; figs. a–c. **Shiino**, 1969: 94. **Utinomi**, 1969: 32, pl. 16, fig. 14. **Inaba**, 1988: 226; upper to mid intertidal, in the mantle cavity of *Capitulum mitella* (Linnaeus), the Inland Sea of Seto. **Iwata**, 1992: 203, fig. 7–5F.

Nemertopsis quadripunctata: **Crandall et al.**, 2002: 12, 20, 28, 31, 35, 40. **Kajihara**, 2007a: 45–51, figs. 2–6; from *Capitulum mitella* (Linnaeus), Shirahama, Wakayama Prefecture.

NOTE: *Nemertopsis quadripunctata* was originally described as *Borlasia quadripunctata* Quoy and Gaimard, 1833 from Ambon, Indonesia. The Japanese taxon identified as *Nemertopsis gracilis* Coe, 1904 was regarded as conspecific with *Nemertopsis quadripunctata* by Gibson (1990a). Apart from the records from Japanese waters, the species is currently known from Ambon (Quoy and Gaimard, 1833) and Hong Kong (Gibson, 1990a).

Genus *Paranemertes* Coe, 1901

Paranemertes Coe, 1901: 32.

TYPE SPECIES: *Paranemertes peregrina* Coe, 1901, by subsequent designation of Friedrich (1955: 173).

***Paranemertes incola* Iwata, 1952**

Paranemertes incola **Iwata**, 1952: 142–143; lower intertidal under stones, Tomioka, Amakusa Islands, Kumamoto Prefecture. **Crandall et al.**, 2002: 13, 21, 28, 36, 41.

***Paranemertes katoi* Yamaoka, 1947**

[Japanese name: katō-himomushi]

Paranemertes katoi **Yamaoka**, 1947: 1467, fig. 4135. **Okuda and Iwata**, 1965: 397, figs. a–c. **Crandall et al.**, 2001: 178–179, pl. 1, figs. 5–7, pl. 2, fig. 15. **Crandall et al.**, 2002: 13, 21, 28, 36, 41.

NOTE: Crandall et al. (2001) noted that Yamaoka's manuscript reported the species as intertidally abundant from May to July under stones or on seaweeds at Susaki and Sotoura, near Shimoda, Shizuoka Prefecture.

***Paranemertes peregrina* Coe, 1901**

[Japanese name: onando-himomushi]

Paranemertes peregrina: **Yamaoka**, 1940a: 240–243, pl. XVII, figs. 3–5, text figs. 17–19; intertidal under or between stones; Akkeshi, Abashiri, and Muroran, Hokkaidō Prefecture. **Coe**, 1944: 29. **Okuda**, 1947: 1467, fig. 4136. **Iwata**, 1954a: 15; lower intertidal under stones or among laminarian holdfasts, Hokkaidō Prefecture (Akkeshi, Monbetsu, Muroran, Nemuro, Oshoro and Rishiri Island). **Yamaguchi and Yamada**, 1955: 68. **Utinomi**, 1956: 32, pl. 16, fig. 11. **Iwata**, 1960c: 169, pl. 84, fig. 9. **Utinomi**, 1960: 32, pl. 16, fig. 11. **Uchida et al.**, 1963: 17. **Iwata**, 1965a: 169. **Okuda and Iwata**, 1965: 398, figs. a–d. **Utinomi**, 1969: 32, pl. 16, fig. 11. **Okada et al.**, 1971: 62. **Uchida et al.**, 1972: 62. **Hieda and Takahashi**, 1986: 42, with two color photographs of a specimen taken at Yakumo, Hokkaidō Prefecture. **Iwata**, 1992: 203, figs. 7–4J, 7–5E. **Crandall et al.**, 2002: 13, 21, 25, 28, 36, 41.

NOTE: *Paranemertes peregrina* Coe, 1901 was originally described from Alaska. Yamaoka's (1940a) illustration in pl. XVII, fig. 6 depicts *Amphiporus parvus*, though the figure legend indicates *Paranemertes peregrina*. Besides the records from Japanese waters, the species is also known from the Commander Islands, Kamchatka Peninsula, Aleutian Islands, and the Pacific coast of North America from Alaska to Ensenada, Mexico (Gibson, 1995: 460).

***Paranemertes plana* Iwata, 1957**

[Japanese name: sagami-himomushi]

Paranemertes plana **Iwata**, 1957a: 20–21, pl. I, fig. 10, pl. VI, figs. 1–5; dredged sublitorally from 250–300 m depth on 16 July 1955 by His Majesty Emperor Shōwa, near “Goromo of southern Minamiaamadaiba” [sic], Sagami Bay, off Kanagawa Prefecture. **Iwata**, 1960c: 169, pl. 84, fig. 10. **Iwata**, 1965b: 398, figs. a, b. **Crandall et al.**, 2002: 13, 21, 29, 41.

Family MALACOBDELLIDAE Blanchard, 1847

Genus *Malacobdella* Blainville, 1827

Malacobdella Blainville, 1827: 270.

TYPE SPECIES: *Hirudo glossa* Müller, 1776, by monotypic designation.

Malacobdella japonica Takakura, 1897

[Japanese name: himobiru]

Malacobdella japonica **Takakura**, 1897: 105–112, pl. VII, figs. 1–6; in the mantle cavity of *Mactra sachalinensis*, Kujūkuri, Chiba Prefecture. **Kaburaki**, 1927: 1664, fig. 3184. **Yamaoka**, 1940a: 253–258, pl. XVII, figs. 14–16, text figs. 32, 33; in the mantle cavity of *Mactra sachalinensis*, Akkeshi, Hokkaidō Prefecture. **Kawai and Yamaoka**, 1940: 255–259, figs. 1–6; in the mantle cavity of *Mactra sachalinensis*, Akkeshi, Hokkaidō Prefecture. **Kaburaki**, 1947: 1470, fig. 4145. **Iwata**, 1954a: 36; in the mantle cavity of *Mactra sachalinensis*, Akkeshi, Hokkaidō Prefecture and Shimoda, Shizuoka Prefecture. **Yamaguchi and Yamada**, 1955: 73. **Iwata**, 1960c: 169, pl. 84, fig. 19. **Satô and Itô**, 1961: 187, fig. 7.1.11. **Uchida et al.**, 1963: 17. **Iwata**, 1965a: 169, 218. **Kaburaki and Iwata**, 1965: 401, figs. a, b. **Shiino**, 1969: 94, fig. 9-3F. **Okada et al.**, 1971: 63. **Uchida et al.**, 1972: 62. **Iwata**, 1992: 204, fig. 7-4M. **Iwata**, 1997: 55. **Crandall et al.**, 2002: 12, 23, 28, 44.

NOTE: Apart from the records from Japanese waters, *Malacobdella japonica* has also been reported from Sakhalin (Steksova, 2004).

Family OTOTYPHLONEMERTIDAE Bürger, 1895

Genus *Otopythonemertes* Diesing, 1863

Otopythonemertes Diesing, 1863: 180.

TYPE SPECIES: *Oerstedia pallida* Keferstein, 1862, by monotypic designation.

Otopythonemertes dolichobasis Kajihara, 2007

Otopythonemertes sp. **Shimomura et al.**, 2001: 47.

Otopythonemertes dolichobasis **Kajihara**, 2007b: 57–66, figs. 1–4; intertidal among coarse sand, Hakozaki, Ôtsuchi Bay, Iwate Prefecture.

TYPE MATERIAL: Holotype, ZIHU-3200, 22 July 1998, 23 slides. Paratypes: ZIHU-3199, 22 July 1998, 7 slides; ZIHU-3201, 25 September 1997, 16 slides; ZIHU-3202, 25 September 1997, 42 slides; ZIHU-3203, 25 September 1997, 31 slides; ZIHU-3208, 3209, 25 May 1998, unsectioned, fixed in Bouin's fluid, preserved in 70% EtOH; ZIHU-3210, 3211, 3212, 3214, 3215, 21 July 1998, unsectioned, fixed in Bouin's fluid, preserved in 70% EtOH; ZIHU-3213, 21 July 1998, fixed and preserved in 99% EtOH; ZIHU-3217, 3218, 22 July 1998, unsectioned, fixed in Bouin's fluid, preserved in 70% EtOH.

Otopythonemertes martynovi Chernyshev, 1993

Otopythonemertes martynovi: **Kajihara**, 1998: 11, pls 1–2; intertidal in coarse sand, Oshoro, Hokkaidō Prefecture; Ôtsuchi, Iwate Prefecture; Shirahama, Wakayama Prefecture; Sugashima, Mie Prefecture; Mukaishima, Hiroshima Prefecture. **Shimomura et al.**, 2001: 46; intertidal in coarse sand, Hakozaki, Ôtsuchi Bay, Iwate Prefecture. **Crandall et al.**, 2002: 13, 20, 28, 43.

NOTE: *Otopythonemertes martynovi* Chernyshev, 1993 was originally described from Peter the Great Bay, Russia, and appears to have a wide range of distribution in Japanese waters.

Otopythonemertes nikolaii Chernyshev, 1998

Otopythonemertes nikolaii: **Shimomura et al.**, 2001: 47; intertidal among coarse sand Hakozaki, Ôtsuchi Bay, Iwate Prefecture. **Crandall et al.**, 2002: 13, 20, 28, 43.

NOTE: *Otopythonemertes nikolaii* Chernyshev, 1998 was originally described from Peter the Great Bay, Russia. The species is currently known from Russia and Japan.

Family POSEIDONEMERTIDAE Chernyshev, 2002

Genus *Diopsonemertes* Kajihara, Gibson and Mawatari, 2001

Diopsonemertes Kajihara et al., 2001: 187.

TYPE SPECIES: *Diopsonemertes acanthocephala* Kajihara, Gibson and Mawatari, 2001, by original designation.

Diopsonemertes acanthocephala Kajihara, Gibson and Mawatari, 2001

Diopsonemertes acanthocephala **Kajihara et al.**, 2001: 187–198, figs. 1–23; Ôtsuchi Bay, Iwate Prefecture; sublittoral, 59 m depth on shell gravel. **Crandall et al.**, 2002: 11, 20, 27, 34.

TYPE MATERIAL: Holotype, ZIHU-1290, immature male, complete series of transverse sections, 71 slides.

Family PROSORHOCHMIDAE Bürger, 1895

Genus *Geonemertes* Semper, 1863

Geonemertes Semper, 1863: 559.

TYPE SPECIES: *Geonemertes pelaensis* Semper, 1863, by monotypic designation.

Geonemertes pelaensis Semper, 1863

[Japanese name: ogasawara-riku-himomushi]

Geonemertes pelaensis: **Oki et al.**, 1987: 69–75, figs. 1–4; among roadside bushes and under a flowerpot in a garden, Chichijima Island, Ogasawara Islands, the Metropolis of Tôkyô. **Kawakatsu**, 1991: 2, fig. 15. **Kawakatsu**, 1999: 11–12, figs. 1–5.

NOTE: Semper's (1863) original description of *Geonemertes pelaensis* was based upon material collected from the Palau Islands, Republic of Palau. The species is also known from Papua New Guinea, Sulawesi, Seychelle Islands, Peradeniya (Sri Lanka), Kei Island, Upolu Island (Samoa), Mauritius, Florida, Dominica (West Indies), Jamaica, Mangareva Island, Oahu (Hawaiian Islands), and Réunion (Gibson and Moore, 1998: 159).

Genus *Pantinonemertes* Moore and Gibson, 1981

Pantinonemertes Moore and Gibson, 1981: 176.

TYPE SPECIES: *Pantinonemertes winsori* Moore and Gibson, 1981, by original designation.

NOTE: The genus *Pantinonemertes* Moore and Gibson, 1981 now contains nine species (Sun, 2001), but the generic name has a subjective senior synonym *Neonemertes* Girard, 1893 (Chernyshev, pers. comm.). Moore and Gibson (1981) recognized the genus *Pantinonemertes* as including three nominal species: *Pantinonemertes winsori* Moore and Gibson, 1981, *Pantinonemertes enalios* Moore and Gibson, 1981, and *Tetrastemma agriculta* Willemoes-Suhm, 1874 (the name-bearing type of the nominal genus *Neonemertes* Girard, 1893), while Moore

and Gibson (1981) designated *Pantinonemertes winsori* Moore and Gibson, 1981 as the type species of the genus *Pantinonemertes* Moore and Gibson, 1981. The name *Neonemertes* has been used as valid by six works, including Girard (1893: 238), Joubin (1894: 193), Friedrich (1955: 142, 143, 161, 1958: 22), Corrêa (1966: 365), and Riser (1974: 363, 364), whereas *Pantinonemertes* has been used in at least 42 works since the year 1981. It is thus reasonable to conclude that the name *Pantinonemertes* has been adopted as the prevailing usage, and that the senior synonym *Neonemertes* should be suppressed by plenary power by the ICZN under Article 23.9.3 of the Code (ICZN, 1999). Recently, Maslakova (2005) concluded that these two genera should be synonymized due to lack of morphological differences between them, on the basis of a reinvestigation of all available type and voucher material of species of *Pantinonemertes* Moore and Gibson 1981 and *Prosadenoporus* Bürger, 1890. The name *Prosadenoporus* Bürger, 1890 has precedence over both *Neonemertes* Girard, 1893 and *Pantinonemertes* Gibson and Moore, 1981. However, since Maslakova (2005) disclaimed nomenclatural acts, the name *Pantinonemertes* Gibson and Moore 1981 is here used as valid.

Pantinonemertes spectacula (Yamaoka, 1940)

Prostoma spectaculum Yamaoka, 1940b: 16–17, fig. 3; habitat not recorded, Naha and Chinen, Okinawa Prefecture.

Pantinonemertes speculacula: Crandall et al., 2002: 13, 21, 31, 43.

NOTE: Gibson (1990a) redescribed the material from Hong Kong and transferred this species to the genus *Pantinonemertes*. Currently known from Okinawa and Hong Kong.

Family TETRASTEMMATIDAE Hubrecht, 1879

NOTE: The correct spelling of the family name is “Tetramermatidae” under Article 29.3. of the Code (ICZN, 1999), since the name of its type genus *Tetramermannum* (neuter gender) gives the genitive singular “Tetramermatos” and the stem “Tetramermat-.”

Genus *Nemertellina* Friedrich, 1935

Nemertellina Friedrich, 1935b: 10.

TYPE SPECIES: *Nemertellina oculata* Friedrich, 1935 by subsequent designation of Friedrich (1955: 164).

Nemertellina yamaokai Kajihara, Gibson and Mawatari, 2000

[Japanese name: yamaoka-himomushi]

Nemertellina minuta: Yamaoka, 1940a: 239–240, text figs. 15, 16; sublittoral from several meters depth; in the canals of sponges; Akkeshi Bay, Hokkaidô Prefecture. Iwata, 1954a: 15. Yamaguchi and Yamada, 1955: 68. Uchida et al., 1963: 17. Crandall et al., 2002: 12, 20, 28, 35, 40. Non Friedrich, 1935a: 320.

Nemertellina yamaokai Kajihara et al., 2000: 265–276, figs. 1–33; sublittoral, 6–8 m depth among sponges, seaweeds, rocks, and mollusks (*Patinopecten* sp. and oysters), 43°00'N, 144°46'E and 43°42'N, 144°51'E, Akkeshi Bay, Hokkaidô Prefecture. Crandall et al. 2002: 12, 20, 28, 35. Thollesson and Norenburg, 2003: 408; Akkeshi Bay,

Hokkaidô Prefecture.

TYPE MATERIAL: Holotype, ZIHU-1260, immature male, complete series of transverse sections, 26 slides. Paratypes: ZIHU-1261, female, series of transverse sections, 24 slides; ZIHU-1262, male, complete series of longitudinal sections, 14 slides; USNM 186063, female, complete series of transverse sections, 17 slides. Three unsectioned voucher specimens are also deposited under ZIHU-1271, ZIHU-1272, and ZIHU-1273.

Genus *Oerstedia* Quatrefages, 1846

Oerstedia Quatrefages, 1846: 221.

TYPE SPECIES: *Oerstedia maculata* Quatrefages, 1846, now regarded as a junior synonym of *Planaria dorsalis* Abildgaard, 1806.

NOTE: The genus *Oerstedia* Quatrefages, 1846 had long been classified into the family Prosorhochmidae, before Moore and Gibson (1988) argued that the genus could no longer be retained in that taxon. The familial affiliation of the genus has been treated as uncertain (Gibson, 1994). A recent molecular phylogenetic study (Thollesson and Norenburg, 2003) indicated that members of the genus are closely related to tetrastematids, although Strand and Sundberg's (2005b) molecular phylogenetic analyses were not decisive about the familial classification. The genus is here provisionally included in the family Tetrastematidae.

Oerstedia dorsalis (Abildgaard, 1806)

[Japanese name: botan-himomushi]

Oerstedia dorsalis: Iwata, 1954a: 17, fig. 4A; lower intertidal among algae or the hydrozoan *Eudendrium annulatum*, Akkeshi, Hokkaidô Prefecture. Yamaguchi and Yamada, 1955: 68. Iwata, 1960a: 96, fig. 6. Iwata, 1960c: 169; pl. 84, fig. 11. Uchida et al., 1963: 17. Iwata, 1965a: 169, 218.

Iwata, 1965b: 398; figs. a–e. Iwata, 1983: 181. Iwata, 1992: 203, fig. 7–5H. Shimomura et al., 2001: 47; intertidal, Akahama, Ôtsuchi Bay, Iwate Prefecture. Crandall et al., 2002: 13, 20, 28, 35, 40.

Oerstedia dorsalis var. *aqualis* Iwata, 1954a: 17, fig. 4A–4; lower intertidal among algae or the hydrozoan *Eudendrium annulatum*, Akkeshi, Hokkaidô Prefecture. Crandall et al., 2002: 10, 28.

Oerstedia dorsalis var. *albolineata*: Iwata, 1954a: 17, fig. 4A–2, 4A–3 lower intertidal among algae or the hydrozoan *Eudendrium annulatum*, Akkeshi, Hokkaidô Prefecture.

Iwata, 1960b: 35–40, figs. 123–142; intertidal, Akkeshi, Hokkaidô Prefecture. Iwata, 1983: 182, 187, 193, fig. 8–14C.

Oerstedia dorsalis var. *viridis*: Iwata, 1954a: 17, fig. 4A–1 lower intertidal among algae or the hydrozoan *Eudendrium annulatum*, Akkeshi, Hokkaidô Prefecture.

NOTE: *Oerstedia dorsalis*, originally described as *Planaria dorsalis* by Abildgaard (1806) from Denmark and Norway, was transferred to *Oerstedia* by Bürger (1895: 592). The species is known to exhibit a high degree of polymorphism in body color pattern (Bürger, 1895; Iwata, 1954a; Brunberg, 1964). A series of studies based on morphological (Sundberg, 1984) and molecular (Sundberg and Janson, 1988; Sundberg and Andersson, 1995) evidence have revealed the existence of a cryptic species, *Oerstedia striata*, that can be distinguished from *Oerstedia dorsalis* by enzyme differences, external pigmentation, and the

general appearance of the body (Sundberg, 1988). The forms reported under the name *Oerstedia dorsalis* are known from the coast of North America (from Puget Sound, Washington to Mexico), Gulf of Mexico, Atlantic coast of North America (Nova Scotia to Florida), western Baltic Sea, North Sea, Mediterranean Sea, Black Sea, northwestern Spain, and Madeira (Gibson, 1995: 467). Due to the high polymorphism, different taxa may be contained among these forms. Numerous varieties have been named, including three reported from Japanese waters: var. *aequalis* Iwata, 1954a, var. *albolineata* Bürger, 1895, and var. *viridis* Bürger, 1895. Until future studies determine whether or not these varieties warrant separate taxonomic status, these are regarded as synonymous with *Oerstedia dorsalis*.

Oerstedia polyorbis Iwata, 1954

Oerstedia polyorbis Iwata, 1954a: 18–19, fig. 4B; lower intertidal among the hydrozoan *Eudendrium annulatum* Norman, Daikokujima Island, Akkeshi, Hokkaidō Prefecture. Yamaguchi and Yamada, 1955: 69. Uchida et al., 1963: 17. Crandall et al., 2002: 13, 20, 28, 35, 40.

?*Oerstedia zebra*: Thollesson and Norenburg, 2003: 408; Akkeshi Bay, Akkeshi, Hokkaidō Prefecture.

?*Oerstedia venusta*: Thollesson and Norenburg, 2003: 408; Akkeshi Bay, Akkeshi, Hokkaidō Prefecture.

NOTE 1: Iwata (1954: 18) established *Oerstedia polyorbis*, which is about 5 mm in body length, with about 30 transverse dorsal bands and cephalic glands that are not well developed and limited only to the anterior portion of the head. Later, Chernyshev (1993: 13) described a similar form, *Oerstedella (Paroerstedella) zebra* (now *Oerstedia zebra*), which differs from *Oerstedia polyorbis* in body length (8–13 mm), the number of the transverse dorsal bands (10–18), and in having cephalic glands extending behind brain. However, ten specimens obtained from the same population as those that Thollesson and Norenburg (2003) identified as *Oerstedia zebra*, possessed 9–16 transverse bands, with body length varying from 2–4 mm and cephalic glands extending behind brain; there was a pair of pores on the ventral surface of the head, which represented the openings of the cerebral organ ducts, but there were no distinct anterior cephalic furrows (Kajihara, pers. obs.). This overlap in characters indicates that *Oerstedia zebra* might be a junior synonym of *Oerstedia polyorbis*.

NOTE 2: Thollesson and Norenburg (2003) identified their material from Akkeshi as *Oerstedia venusta*. Specimens from the same locality (n=10) were almost identical with what these authors identified as *Oerstedia zebra* in both external and internal morphology, except for the transverse dorsal bands that were present in the latter. However, the form identified as *Oerstedia venusta* by Thollesson and Norenburg (2003) differs from Iwata's (1954a) original description in not having distinct anterior cephalic furrows. It remains uncertain whether *Oerstedia venusta* sensu Thollesson and Norenburg (2003) represents the same taxon as Iwata's (1954a) form. *Oerstedia venusta* sensu Thollesson and Norenburg (2003) also resembles *Oerstedia oculata* (Kulikova, 1987) in external characters.

NOTE 3: Strand and Sundberg (2005a: 210) regarded *Oer-*

stedia zebra (Chernyshev, 1993) *sensu* Thollesson and Norenburg (2003) and *Oerstedia venusta* Iwata, 1954 *sensu* Thollesson and Norenburg (2003) as synonymous, on the basis of genetic similarity. However, as the taxonomic identity of the latter is unclear, Strand and Sundberg's (2005a) synonymization may require additional topotypic data before it is substantiated.

Genus *Prostoma* Dugès, 1828

Prostoma Dugès, 1828: 140.

Stichostemma Montgomery, 1894: 8; synonymized by Bürger (1904: 53).

TYPE SPECIES: The genus *Prostoma* was long used for species of *Tetrastremma*, until Stiasny-Wijnhoff (1938) circumscribed *Prostoma* to include only freshwater species. The single species included in the nominal genus *Prostoma* when it was established was *Prostoma clepsinoides* Dugès, 1828, which was the only nominal species eligible to be the type species of the genus. However, Friedrich (1955: 162) indicated "*Prostoma lumbricoideum* Dugès (1828) [sic]" (correctly *Prostoma lombricoideum* Dugès, 1830) as the type species of *Prostoma*, and recently Gibson (1995: 495) indicated *Prostoma graecense* (Böhmig, 1892). These nomenclatural acts cannot be regarded as valid designations of the type species, according to Article 67.2 of the Code (ICZN, 1999). Meanwhile, the taxonomic identity indicated by the name *Prostoma clepsinoides* has been regarded as vague (Stiasny-Wijnhoff, 1938; Gibson and Moore, 1976). When it becomes necessary to delineate the identity of *Prostoma*, especially in comparison with similar genera like *Limnemertes*, nomenclatural actions will be required, such as either 1) removing the name-bearing function from *Prostoma clepsinoides* and bestowing it on a well-known species like *Prostoma graecense*, or 2) designating a neotype for *Prostoma clepsinoides*, ideally obtained from the type locality, probably Montpellier, France.

Prostoma ohmiense Chernyshev, Timoshkin, and Kawakatsu, 1998

Prostoma ohmiense Chernyshev et al., 1998: 53–60, figs. 2–6; on rocks with overgrowing algae, 2 m depth, Lake Biwako, off Kitakomatsu, Shiga-chō, Shiga-gun, and off Shin-asahi-chō, Takashima-gun, Shiga Prefecture. Crandall et al., 2002: 14, 21, 29, 43.

TYPE MATERIAL: The holotype and two paratypes are supposed to be deposited in Biwako Museum, according to the original description. However, due to confusion arising during transportation of the specimens, the holotype cannot be identified among the specimens in the museum (Dr. Mark J. Grygier, pers. comm.).

Genus *Quasitetrastemma* Chernyshev, 2004

Quasitetrastemma Chernyshev, 2004b: 152.

TYPE SPECIES: *Tetrastremma nigrifrons* Coe, 1904, by original designation.

Quasitetrastemma nigrifrons (Coe, 1904)

[Japanese name: menoko-himomushi]

Prostoma nigrifrons: Yamaoka, 1940a: 249–251, pl. XVI, fig. 14, pl. XVII, figs. 9–12, text figs. 26–29; lower intertidal

under stones, Akkeshi and Abashiri, Hokkaidô Prefecture; sublittoral, among the canal system of sponges attached to gastropod shells collected from a depth of several meters, Akkeshi, Hokkaidô Prefecture. **Okuda**, 1947: 1469, fig. 4142 (1–8).

Tetrastremma nigrifrons: **Iwata**, 1954a: 30–32, fig. 8B. **Yamaguchi and Yamada**, 1955: 71, fig. 18–2. **Utinomi**, 1956: 32, pl. 16, fig. 15. **Iwata**, 1960c: 169, pl. 84, fig. 16. **Utinomi**, 1960: 32, pl. 16, fig. 15. **Uchida et al.**, 1963: 17. **Iwata**, 1965a: 169, 218. **Okuda and Iwata**, 1965: 400, figs. a–h. **Utinomi**, 1969: 32, pl. 16, fig. 15. **Okada et al.**, 1971: 62. **Uchida et al.**, 1972: 55; habitat not recorded, Horomui, Hokkaidô Prefecture. **Iwata**, 1992: 203, fig. 7–41. **Shimomura et al.**, 2001: 47; shallow sublittoral, among sessile organisms on mooring floats, Akahama, Ôtsuchi Bay, Iwate Prefecture. **Crandall et al.**, 2002: 14, 21, 29, 36, 41. **Tetrastremma nigrifrons** var. *bilineatum*: **Iwata**, 1957a: 27, pl. I, fig. 14; collected subtidally from 4–6 m depth on 8 December 1953 by His Majesty Emperor Shôwa, Samejima at Hayama, Kanagawa Prefecture. **Crandall et al.**, 2002: 14, 21, 29, 36, 41.

Tetrastremma nigrifrons var. *punctatum*: **Crandall et al.**, 2002: 14, 21, 29, 36, 41. **Tetrastremma nigrifrons** var. *spadix*: **Crandall et al.**, 2002: 14, 21, 29, 37, 41.

NOTE: Originally described as *Tetrastremma nigrifrons* by Coe (1904: 159) from Pacific Grove (36°38'N 121°56'W) and San Pedro, California, USA, the species was recently transferred to *Quasitetrastremma* by Chernyshev (2004b). Known from the Pacific coasts of North and Central America (Puget Sound, Washington, to Salinas Bay, Costa Rica) (Gibson, 1995: 520), the species shows a high degree of polymorphism in color pattern (Coe, 1940: 305). The following varieties have been named: var. *albino* Manchenko and Kulikova, 1996b; var. *bicolor* Coe, 1904; var. *bilineatum* Iwata, 1954a; var. *pallidum* Coe, 1904; var. *punctata* Iwata, 1954a; var. *purpureum* Coe, 1904; var. *spadix* Iwata, 1954a; and var. *zonatum* Coe, 1940. Manchenko and Kulikova (1996b) demonstrated by isozyme analyses that the five sympatric varieties *albino*, *bicolor*, *pallidum*, *punctata*, and *purpureum* are conspecific. Incidentally, Manchenko and Kulikova's (1996b) description of their var. *albino* that possesses no pigmentation gives the impression that it might represent *Quasitetrastremma stimpsoni*.

Quasitetrastremma stimpsoni (Chernyshev, 1992)

Prostoma stigmatum: **Yamaoka**, 1940a: 251–253, pl. XVII, fig. 13, text figs. 30, 31; intertidal under stones or among algae, Akkeshi and Abashiri, Hokkaidô Prefecture.

Tetrastremma stigmatum: **Iwata**, 1954a: 35; intertidal under stones and among algae, Hokkaidô Prefecture (Abashiri, Akkeshi and Hiroo). **Yamaguchi and Yamada**, 1955: 72. **Uchida et al.**, 1963: 17. *Non* Stimpson, 1857: 163.

Tetrastremma stimpsoni **Chernyshev**, 1992: 135. **Crandall et al.**, 2002: 15, 22, 25, 29, 37, 42.

Quasitetrastremma stigmatum: **Chernyshev**, 2004b: 154.

NOTE 1: Although Stimpson's (1857) original description of *Tetrastremma stigmatum* was brief and accompanied by no illustration, Yamaoka (1940a) and Iwata (1954a) considered their material as conspecific with Stimpson's.

Based upon literature, Chernyshev (1992) regarded *Prostoma stigmatum* sensu Yamaoka (1940a) as different from *Tetrastremma stigmatum* Stimpson, 1857 and gave to Yamaoka's taxon a new name, *Tetrastremma stimpsoni* Chernyshev, 1992, while he considered the name *Tetrastremma stigmatum* to be a *nomen dubium*. Later, Chernyshev (2004b) transferred *Tetrastremma stigmatum* (=*Prostoma stigmatum*) sensu Yamaoka (1940a) into *Quasitetrastremma* Chernyshev, 2004, ascribing "Quasitetrastremma stigmatum" (Yamaoka, 1940)." Chernyshev's (1992, 2004b) treatment of the names raises the following two issues: 1) Homonymy. Chernyshev (2004b) states "*Prostoma stigmatum* Yamaoka, 1940 was replaced by a new name *Tetrastremma stimpsoni*," although Yamaoka (1940a) did not establish any nominal species bearing the epithet *stigmatum*. Accordingly, there was no homonymy when Chernyshev (1992) created a new name. 2) Authorship. As mentioned, Chernyshev (2004b) appears to misinterpret Yamaoka (1940a) as establishing a new nominal species *Prostoma stigmatum*. Since Stimpson's material is deemed to belong to a different species from Yamaoka's, the latter taxon name should be ascribed as *Quasitetrastremma stimpsoni* (Chernyshev, 1992).

NOTE 2: It seems likely that Yamaoka did not have access to a copy of Stimpson's 1857 paper, and probably had to refer to Bürger (1904) for the identification of his material as *Prostoma stigmatum*. Unfortunately, the German translation of an excerpt of Stimpson's (1857) Latin description of the species in Bürger (1904) lacked an important sentence for the identification of tetrastemmatids, namely, the presence and coloration of the cephalic patch. Nothing equivalent to the sentence in Stimpson (1857: 163) "pone ocellos anteriores fascia transversa obscure rubra" [behind the anterior eyes there is a dark red transverse band] can be found in Bürger (1904). This could account for why Yamaoka (1940a) identified his material without a cephalic patch as *Tetrastremma stigmatum*, and also why he later established a new species that possessed a red cephalic patch as *Tetrastremma rosecephalum*.

Genus Sacconemertella Iwata, 1970

Sacconemertella Iwata, 1970a: 147.

TYPE SPECIES: *Sacconemertella lutulenta* Iwata, 1970 by original designation.

Sacconemertella lutulenta Iwata, 1970

[Japanese name: chibi-kisui-himomushi]

Sacconemertella lutulenta Iwata, 1970a: 148–151, fig. 1G–I, pl. 4, figs. 26–33; sublittoral in mud, brackish Lake Hinuma, Ibaraki Prefecture. **Iwata**, 1973: 264. **Crandall et al.**, 2002: 14, 21, 29, 43.

Genus Sacconemertopsis Iwata, 1970

Sacconemertopsis Iwata, 1970a: 142.

TYPE SPECIES: *Sacconemertopsis olivifera* Iwata, 1970 by original designation.

Sacconemertopsis olivifera Iwata, 1970

[Japanese name: hime-kisui-himomushi]

Sacconemertopsis olivifera Iwata, 1970a: 143–147, fig. 1D–

F, pl. 3, figs. 18–25; sublittoral in mud, brackish Lake Hinuma, Ibaraki Prefecture. **Iwata**, 1973: 264. **Crandall et al.**, 2002: 14, 21, 29, 43.

Genus *Tetrastemma* Ehrenberg, 1831

Tetrastemma Ehrenberg, 1831: 61.

TYPE SPECIES: *Tetrastemma flavidum* Ehrenberg, 1831, by monotypic designation.

***Tetrastemma candidum* (Müller, 1774)**

Tetrastemma candidum: **Iwata**, 1954a: 35–36; lower intertidal among algae, Akkeshi, Hokkaidō Prefecture.

Yamaguchi and Yamada, 1955: 72. **Uchida et al.**, 1963: 17. **Crandall et al.**, 2002: 14, 21, 29, 36, 41.

NOTE: Originally described as *Fasciola candida* Müller, 1774 from Norway, this species was transferred to *Tetrastemma* by Örsted (1844: 88). The species has a circum polar distribution in the northern hemisphere (British Isles, coasts of Scandinavia, North Sea, Mediterranean, Madeira, Faroe Islands, Iceland, Greenland, Caribbean, Atlantic and Pacific coasts of North America) (Gibson, 1995: 372).

***Tetrastemma insolens* Iwata, 1952**

Tetrastemma insolens **Iwata**, 1952: 146–147, figs. 17, 18; intertidal under stones, Tomioka, Amakusa, Kumamoto Prefecture and Fukue, Gotō Islands, Nagasaki Prefecture.

Crandall et al., 2002: 14, 21, 29, 36, 41.

***Tetrastemma melanocephalum* (Johnston, 1837)**

Tetrastemma melanocephalum: **Yamaoka**, 2005: 153, pl. 1, fig. 5, pl. 2, fig. 5, text fig. 9a–c; intertidal under stones and among algal holdfasts, Shimoda, Shizuoka Prefecture.

NOTE: *Tetrastemma melanocephalum* was originally described as *Nemertes melanocephala* by Johnston (1837). The species is reported from west coast of Sweden, Baltic Sea coasts of Germany, Denmark, British Isles, Mediterranean, Adriatic and Black Sea coasts, northern Spain, Madeira, and the Canary Islands (Gibson, 1995).

***Tetrastemma pinnatum* Iwata, 1954**

Tetrastemma pinnatum **Iwata**, 1954a: 34–35, fig. 9C; sublittoral among algae about 4 m deep, Akkeshi, Hokkaidō Prefecture. **Yamaguchi and Yamada** 1955: 72. **Uchida et al.** 1963: 17. **Crandall et al.**, 2002: 15, 21, 29, 37, 42.

***Tetrastemma pseudocoronatum* Chernyshev, 1998**

Prostoma coronatum: **Yamaoka** 1940a: 247–249, pl. XVI, figs. 12, 13, text fig. 25; lower intertidal on stones, Akkeshi and Abashiri, Hokkaidō Prefecture.

Tetrastemma coronatum: **Iwata**, 1954a: 32; lower intertidal, under stones and among algae, Akkeshi and Abashiri, Hokkaidō Prefecture. **Yamaguchi and Yamada**, 1955: 72. **Uchida et al.**, 1963: 17. **Shimomura et al.**, 2001: 47; intertidal, coarse sand, Akahama, Ōtsuchi Bay, Iwate Prefecture. **Crandall et al.**, 2002: 14, 21, 29, 36, 41.

Tetrastemma pseudocoronatum: **Crandall et al.**, 2002: 15, 21, 29, 37, 42.

NOTE: *Tetrastemma coronatum* was originally described as *Polia coronata* from Bréhat, France, by Quatrefages (1846: 213) and was later transferred to *Tetrastemma* by

Hubrecht (1879: 228). Gibson (1995: 478) stated that the species has been reported from the British Isles, Scandinavia, the Atlantic coast of France, the Mediterranean, the Adriatic and Black Seas, and Madeira, but questioned the validity of the records by Yamaoka (1940a) and Iwata (1954a). Chernyshev (1998) described *Tetrastemma pseudocoronatum* based upon material obtained from Kunashiri Island and regarded *Prostoma coronatum* sensu Yamaoka (1940a) as conspecific.

***Tetrastemma roseocephalum* (Yamaoka, 1947)**

Prostoma roseocephalum **Yamaoka**, 1947: 1469, fig. 4141 (1–4); under stones and among algal holdfasts; Shimoda, Shizuoka Prefecture.

Tetrastemma roseocephalum: **Okuda and Iwata**, 1965: 400; figs. a–d. **Crandall et al.**, 2001: 180, pl. 1, figs. 11–14, pl. 2, figs. 17, 17a. **Crandall et al.**, 2002: 15, 22, 29, 37, 42. **Yamaoka**, 2005: 155.

NOTE: *Tetrastemma roseocephalum* is potentially a junior synonym of *Tetrastemma stigmatum*. See NOTE under *Tetrastemma yamaokai*.

***Tetrastemma stigmatum* Stimpson, 1857**

Tetrastemma stigmatum **Stimpson**, 1857: 163; sublittoral, under stones or among algae at a depth of about 11 m, Hakodate, Hokkaidō Prefecture. **Crandall et al.**, 2002: 15, 22, 29, 37, 42.

NOTE: *Tetrastemma stigmatum* is potentially a senior synonym of the two nominal species *Prostoma roseocephalum* and *Tetrastemma yamaokai*. See NOTE for *Tetrastemma yamaokai*.

***Tetrastemma verinigrum* Iwata, 1954**

Tetrastemma verinigrum **Iwata**, 1954a: 32–33, fig. 9A; intertidal among algal holdfasts, Oshoro, Hokkaidō Prefecture.

Yamaguchi and Yamada, 1955: 72. **Okada et al.**, 1971: 62. **Uchida et al.**, 1972: 62. **Crandall et al.**, 2002: 15, 22, 30, 31, 37, 42.

Tetrastemma verinigrum var. *meridianum* **Iwata**, 1954b: 41, fig. 2C; lower intertidal under stones, Kushimoto, Wakayama Prefecture. **Crandall et al.**, 2002: 15, 22, 30, 37, 42.

NOTE: Apart from the records from Japanese waters, *Tetrastemma verinigrum* is also known from Hong Kong (Gibson, 1990a).

***Tetrastemma yamaokai* Iwata, 1954**

[*sarinuri-himomushi*]

Tetrastemma yamaokai **Iwata**, 1954a: 33, fig. 9B; Oshoro, Hokkaidō Prefecture, intertidal among algal holdfasts.

Yamaguchi and Yamada, 1955: 72. **Okada et al.**, 1971: 63. **Uchida et al.**, 1972: 62. **Crandall et al.**, 2002: 15, 22, 30, 37, 42.

NOTE: Judging from the original description of *Tetrastemma yamaokai* Iwata, 1954, the taxonomic identity of this species can be regarded to be encompassed by that of *Tetrastemma roseocephalum* Yamaoka, 1947; thus the former name may possibly be a junior synonym of the latter. *Tetrastemma stigmatum* Stimpson, 1857 was regarded by Chernyshev (1992) as a *nomen dubium*, but Stimpson's (1857) original description contains such

important taxonomic characters for identifying tetrastemmatids as the coloration of the body and cephalic patch. These character states in *Tetraستemma stigmatum* Stimpson, 1857 also apply to *Tetraستemma yamaokai* Iwata, 1954 and *Tetraستemma roseocephalum* Yamaoka, 1947. These might be synonymized by future studies.

Subclass POLYSTILIFERA Brinkmann, 1917

Order REPTANTIA Brinkmann, 1917

Family DREPANOPHORIDAE Verrill, 1892

Genus Drepanophorus Hubrecht, 1874

Drepanophorus Hubrecht, 1874: 42.

TYPE SPECIES: *Drepanophorus rubrostriatus* Hubrecht, 1874 by subsequent designation of Gibson (1995: 360).

***Drepanophorus longiceps* Iwata, 1957**

[Japanese name: mikado-himomushi or tsurugi-himomushi] *Drepanophorus longiceps* Iwata, 1957a: 27–30, pl. I, fig. 15, pl. VI, figs. 9, 10, pl. VII, figs. 1–8; dredged from a depth of 50 m on 7 November 1954 by His Majesty Emperor Shōwa, Shimoda, Shizuoka Prefecture. Iwata, 1960c: 169, pl. 84, fig. 17. Iwata, 1965a: 216. Iwata, 1965b: 401, figs. a, b. Iwata, 1992: 199, fig. 7–5B.

Hirohitonemertes longiceps [nomen nudum]: Crandall et al., 2002: 11, 22, 27, 39.

Genus Kameginemertes Iwata, 1998

Kameginemertes Iwata, 1998: 199.

TYPE SPECIES: *Amphiporus parmiornatus* Iwata, 1957 by original designation.

***Kameginemertes parmiornata* (Iwata, 1957)**

Amphiporus parmiornatus Iwata, 1957a: 24–25, pl. I, fig. 12, pl. VI, fig. 6; dredged from 50–55m depth on 6 December 1951 by His Majesty Emperor Shōwa, “Kamejyo” [sic], Sagami Bay, off Kanagawa Prefecture.

Kameginemertes parmiornatus Iwata, 1998: 199–213, figs. 1–9.

Kameginemertes parmiornata: Crandall et al., 2002: 11, 22, 27, 39.

NOTE: Iwata (1998) redescribed *Amphiporus parmiornatus* Iwata, 1957 based on the original material, establishing a new genus; in his 1998 paper, the locality is noted as “On off-shore reef at Kamegisho, near Nagai.”

Family SAGAMINEMERTIDAE Chernyshev, 2003

Genus Sagaminemertes Friedrich, 1968

Sagaminemertes Friedrich, 1968: 34.

TYPE SPECIES: *Amphiporus nagaiensis* Iwata, 1957 by monotypic designation.

***Sagaminemertes nagaiensis* (Iwata, 1957)**

[Japanese name: nagai-himomushi]

Amphiporus nagaiensis Iwata, 1957a: 23–24, pl. I, fig. 11, pl. VI, figs. 6, 7; dredged sublitorally from 100–130 m depth on 7 August 1950 by His Majesty Emperor Shōwa, northern Nakafukari near Nagai, Sagami Bay, Kanagawa Prefecture.

Sagaminemertes nagaiensis Iwata, 1988: 115–123, figs. 1–7; Iwata, 1992: 199, fig. 7–5A. Crandall et al., 2002: 14,

22, 29, 41.

NOTE: Iwata (1988) redescribed *Amphiporus nagaiensis* Iwata, 1957 based on the original material, establishing a new genus. Iwata (1988) indicated the depth as 100–110 m, whereas Iwata (1957) gave it as 100–130 m.

Order PELAGICA Brinkmann, 1917

Family NECTONEMERTIDAE Verrill, 1892

Genus Nectonemertes Verrill, 1892

Nectonemertes Verrill, 1892: 447.

TYPE SPECIES: *Nectonemertes mirabilis* Verrill, 1892 by monotypic designation.

***Nectonemertes japonica* Foshay, 1912**

[Japanese name: hoso-oyogi-himomushi]

Nectonemertes japonica Foshay, 1912: 50–53, fig. 1; off Misaki, Kanagawa Prefecture, “taken in the vicinity of Misaki..., but no depth is recorded.” Crandall et al., 2002: 12, 22, 28, 31, 42.

Nectonemertes mirabilis: Komai, 1919: 295, fig. 2. Kato, 1947: 1469, fig. 4143. Satô and Itô, 1961: 187, fig. 7.1.9.

Kato and Iwata, 1965: 401; with one figure. Shiino, 1969: 94, fig. 9–3E. Iwata, 1992: 201, fig. 7–4E. Iwata, 1997: 55.

NOTE: Brinkmann (1917: 9) synonymized *Nectonemertes japonica* Foshay, 1912 with *Nectonemertes mirabilis* Verrill, 1892. Coe (1926: 174) proposed separating these species, but later synonymized *Nectonemertes japonica* with *Nectonemertes mirabilis* (Coe, 1954: 259). Korotkevitsch (1955: 72, 81–82; 1977: 17) retained *Nectonemertes japonica* under its original name. Gibson (1995: 425) listed *Nectonemertes japonica* as a valid species name. Future studies must settle the problem of which name should be applied to the Japanese species. *Nectonemertes mirabilis* was originally described from the Atlantic, but is known to be distributed in the North, equatorial and South Atlantic and North Pacific (Gibson, 1995: 426), while *Nectonemertes japonica* is only known from Japanese waters.

Family PELAGONEMERTIDAE Moseley, 1875

Genus Pelagonemertes Moseley, 1875

Pelagonemertes Moseley, 1875a: 168.

TYPE SPECIES: *Pelagonemertes rollestoni* Moseley, 1875 by monotypic designation.

***Pelagonemertes moseleyi* Bürger, 1895**

[Japanese name: oyogi-himomushi]

Pelagonemertes rollestoni Moseley, 1875b: 377–383, pl. XI, figs. 1–5; trawled from 420–755 fathoms by H.M.S. Challenger on 5 June 1875, obtained by Willemoes-Suhm, 34°58'N, 139°30'E, about halfway between Ôshima (the Metropolis of Tôkyô) and Cape Sagami (Kanagawa Prefecture).

Pelagonemertes moseleyi: Komai, 1919: 294, fig. 1. Kato and Tanaka, 1938: 595–598, pl. XL, figs. A–F, text figs. 1 and 2; “In the middle of November, 1937, one of the writers, Otohiko Tanaka, obtained several specimens of pelagic nemerteans along with a large number of deep-sea medusae, copepods, arrow-worms, etc., by the vertical net from about 1,000 meters to the surface, at a station 3 miles off Hasima in Sagami Bay.” Kato, 1940: 101, two

figs. **Kato**, 1947: 1470, fig. 4144. **Iwata**, 1960c: 169; pl. 84, fig. 18. **Satô and Itô**, 1961: 187, fig. 7.1.8. **Iwata**, 1965a: 169, 217. **Kato and Iwata**, 1965: 401, one figure. **Iwata**, 1992: 201, fig. 7-4F. **Crandall et al.**, 2002: 13, 22, 29, 32.

NOTE: *Pelagonemertes moseleyi* was first reported by Moseley (1875b) as a young individual of *Pelagonemertes rollestoni*, which was also obtained during the scientific cruise of H.M.S. *Challenger* (Moseley, 1875a). Later, Bürger (1895: 596) regarded the former as different from the latter and gave it a new name, *Pelagonemertes moseleyi*; this species has been found in the North and tropical Atlantic and the North Pacific (Gibson, 1995: 463).

Records for Which Application of the Species Name is Doubtful

Class PALAEONEMERTEA Hubrecht, 1879
Family CEPHALOTRICHIDAE McIntosh, 1874
Cephalothrix filiformis (Johnston, 1829)

[Japanese name: daikoku-hoso-himomushi]

Procephalothrix filiformis: **Iwata**, 1954a: 7, fig. 1B, D; under stones near the low-water level on a stony beach, Daikokujima Island, Akkeshi, Hokkaidô Prefecture. **Yamaguchi and Yamada**, 1955: 65. **Iwata**, 1960b: 14–18, figs. 46–61; intertidal, Akkeshi, Hokkaidô Prefecture. **Uchida et al.**, 1963: 17. **Iwata**, 1983: 181, 182, 188, fig. 8-5d, e, f. **Crandall et al.**, 2002: 14, 16, 29, 36, 41. **Thollesson and Norenburg**, 2003: 409; Akkeshi Bay, Hokkaidô Prefecture.

NOTE: *Cephalothrix filiformis* was originally described as *Planaria filiformis* by Johnston (1828) from the British Isles. Johnston's type material is presumably not extant. The taxonomic identity of *Cephalothrix filiformis sensu* Iwata (1954a) requires further investigation, since the occurrence in Japanese waters is quite outside the range of this species based on other records from the British Isles, the coast of France, and northern Spain (Gibson, 1994: 60, 1995: 467).

Cephalothrix linearis (Rathke, 1799)

[Japanese name: hoso-himomushi]

Cephalothrix linearis: **Takakura**, 1898: 119, fig. 4; intertidal among *Sargassum thunbergii*, Jôgashima, Kanagawa Prefecture. **Kaburaki**, 1927: 1662, fig. 3181. **Kaburaki**, 1947: 1474, fig. 4156. **Iwata**, 1951: 135; habitat not recorded, Onomichi, Hiroshima Prefecture. **Iwata**, 1952: 132; intertidal under stones, Tomioka, Amakusa, Kumamoto Prefecture and Fukue, Gotô Islands, Nagasaki Prefecture. **Yamaoka**, 1940a: 215, pl. XIV, figs. 5–8, text figs. 5, 6; under stones on sandy beach near high-water level, Akkeshi, Hokkaidô Prefecture. **Utinomi**, 1956: 31, pl. 16, fig. 2. **Iwata**, 1960c: 166, pl. 83, fig. 4. **Utinomi**, 1960: 31, pl. 16, fig. 2. **Satô and Itô**, 1961: 187, fig. 7.1.2. **Inaba**, 1963: 227; lower intertidal to shallow sublittoral under stones on rocky to gravelly shores; commonly found in the Inland Sea of Seto. **Iwata**, 1965a: 169. **Kaburaki and Iwata**, 1965: 390, figs. 1a, b. **Utinomi**, 1969: 31, pl. 16, fig. 2. **Honma and Kitami**, 1978: 14; Sado Island, Niigata Prefecture. **Inaba**, 1988: 225; lower intertidal to shallow sublittoral under stones on rocky to gravelly shores; commonly found in the Inland Sea of Seto. **Ali et al.**, 1990: 1083; intertidal, Shimoda, Shizuoka Prefecture, identified by Dr.

Minoru Imajima. **Noguchi et al.**, 1991: 846; Shimoda, Shizuoka Prefecture. **Iwata**, 1992: 195, fig. 7-3A. **Asakawa et al.**, 2000: 764; among shells of the oyster, *Crassostrea gigas*, Hiroshima Bay, Hiroshima Prefecture, identified by Prof. Iwata. **Crandall et al.**, 2002: 10, 16, 24, 26, 33, 38.

Procephalothrix simulus [sic]: **Iwata**, 1954a: 6, fig. 1A; under stones or among laminarian holdfasts, Hokkaidô Prefecture (Akkeshi, Muroran, Hiroo, Nemuro and Oshoro). **Yamaguchi and Yamada**, 1955: 65. **Iwata**, 1957c: 108. **Iwata**, 1960a: 96, fig. 1. **Iwata**, 1960b: 3–14, figs. 1–45; intertidal, Akkeshi, Hokkaidô Prefecture. **Iwata**, 1960c: 166, pl. 83, fig. 5. **Uchida et al.**, 1963: 17. **Iwata**, 1965a: 169, 201. **Iwata**, 1965b: 390, figs. a, b. **Okada et al.**, 1971: 62. **Honma and Kitami**, 1978: 14; Sado Island, Niigata Prefecture. **Iwata**, 1983: 181, 182, 193, figs. 8-5a, b, c, 8-13a. **Iwata**, 1992: 196. **Iwata**, 1997: 55. **Thollesson and Norenburg**, 2003: 409; Akkeshi Bay, Hokkaidô Prefecture.

Procephalothrix simulus [sic]: **Uchida et al.**, 1972: 62.

Procephalothrix simula: **Crandall et al.**, 2002: 14, 16, 29, 36, 41.

Cephalothrix sp. **Asakawa et al.**, 2003: 748; among shells of the oyster, *Crassostrea gigas*, Hiroshima Bay, Hiroshima Prefecture. **Tanu et al.**, 2004: 516; Hiroshima Bay, Hiroshima Prefecture.

NOTE: *Cephalothrix linearis* was originally described as *Planaria linearis* by Rathke (1799) from the North Sea coast of Denmark, based on two specimens. The original description was so brief and uninformative that Jensen (1878) even suspected that Rathke's two specimens represented two different species. Because of the vagueness of the taxonomic identity of this species, determining whether or not the Japanese population identified as *C. linearis* can be included in the same species will require further investigation. Comparative toxicological (Dr. Manabu Asakawa, unpublished) and molecular (Kajihara, unpublished) data from Hiroshima Bay, Ôtsuchi Bay, and Akkeshi Bay indicate that the species previously recorded as *Cephalothrix linearis* from Japanese waters appears to be conspecific with *Cephalothrix simula* sensu Iwata (1954a). The species possesses strong toxicity due to a high concentration of tetrodotoxin and/or related chemicals (Ali et al., 1990; Asakawa et al., 2000, 2003). Possible nominal species contained in this taxon include *Procephalothrix fasciculus* Iwata, 1952 and *Procephalothrix arenaria* Gibson, 1990.

Class PILIDIOPHORA Thollesson and Norenburg, 2003

Family LINEIDAE McIntosh, 1874

Cerebratulus fuscus (McIntosh, 1874)

Cerebratulus fuscus: **Takakura**, 1898: 426–427, fig. 24; on the surfaces of rocks obtained sublitorally from a depth of several fathoms on a muddy sand substrate, Jôgashima, Kanagawa Prefecture. **Crandall et al.**, 2002: 10, 17, 27, 38.

NOTE: *Cerebratulus fuscus* was originally described as *Micrura fusca* by McIntosh (1873–1874) from the British Isles and later transferred to *Cerebratulus* by Hubrecht (1879: 219). The species is distributed in European waters, including the Mediterranean (Gibson, 1994: 78). Records from North American and Greenland are related to *Cerebratulus marginatus* Renier, 1804 (Coe, 1940: 276, 1943: 255). Gibson

(1995: 417) doubted the validity of Wheeler's (1934: 232, 1940: 32) records from South Africa. Takakura's (1898) record of *C. fuscus* from Japan is based on external characters and thus requires further investigation.

Lineus vegetus Coe, 1931

Lineus cf. vetatus [sic]: **Inaba**, 1988: 225; lower intertidal to shallow sublittoral, sandy to muddy sediment; a specimen collected on May 1976 in Bizen, Okayama Prefecture, is deposited in Mukaishima Marine Biological Station, Hiroshima University.

Lineus vegetus: **Iwata**, 1997: 53, species name appearing as the caption of a color photograph taken by Fumio Iwata, locality not indicated.

NOTE: *Lineus vegetus* was originally described by Coe (1931) from California, USA, as possessing a strong capacity for regeneration. It was later synonymized with the nominal species *Ramphogordius sanguineus* Rathke, 1799 by Riser (1994), who established a new genus *Myiosiphagos* to accommodate the species, along with the two nominal species *Planaria sanguinea* Rathke, 1799 and *Lineus pseudolacteus* Gontcharoff, 1951. However, as the genus *Myiosiphagos* Riser, 1994 constitutes a junior synonym of *Ramphogordius* Rathke, 1843, the species should now called *Ramphogordius sanguineus* Rathke, 1799 (Riser, 1998). There is no taxonomic account of any material of this species from Japanese waters, and the use of the species name by Inaba (1988) and Iwata (1997) should be regarded as questionable.

Lineus bilineatus (Renier, 1804) *sensu* Iwata (1954a)

[Japanese name: hutasuji-himomushi]

Lineus bilineatus: **Iwata**, 1954a: 9–10, fig. 2A; lower intertidal under stones, Akkeshi, Hokkaidō Prefecture. **Yamaguchi and Yamada**, 1955: 65, fig. 17-1. **Uchida et al.**, 1963: 17. **Iwata**, 1965b: 393, figs. a–c. **Crandall et al.**, 2002: 11, 17, 27, 34, 39.

NOTE: *Lineus bilineatus*, originally described as *Cerebratulus bilineatus* Renier, 1804 from the Adriatic Sea (presumably near Padua), has a color pattern and number of eyes different from the Japanese form identified under this name (Gibson, 1995: 330–331).

Lineus gesserensis (Müller, 1774) *sensu* Takakura, 1898

Lineus gesserensis: **Takakura**, 1898: 335, fig. 17; intertidal among algae, Koajiro Bay, Kanagawa Prefecture.

NOTE: *Lineus gesserensis*, originally described by Müller (1788: 32) as *Planaria gesserensis* from Denmark, was subsequently synonymized with *Lineus ruber* (Müller, 1774) by Bürger (1904: 101). However, Gibson (1982a: 90; 1994: 94) noted that forms identified as *Lineus gesserensis* also contain what should now be referred to as *Lineus viridis* (Müller, 1774). Takakura's (1898) description of the external features of what he called *Lineus gesserensis* equally applies to both *Lineus ruber* and *Lineus viridis*, so the identity of Takakura's material cannot be determined. *Lineus ruber* and *Lineus viridis* were recently transferred to the genus *Poseidon* Girard, 1852 by Chernyshev (2004c).

Lineus grubei (Hubrecht, 1879) *sensu* Takakura, 1898

Lineus grubei: **Takakura**, 1898: 331–332, fig. 11; among

algae from 2–3 fathoms depth, Misaki and Jōgashima, Kanagawa Prefecture.

NOTE: Originally reported from Naples as *Cerebratulus grubei* by Hubrecht (1879: 215–216), this species was transferred to *Lineus* by Bürger (1892: 160). Gibson (1995: 335) stated that Takakura's (1898) report of this species from Japan "cannot be substantiated."

Lineus longifissus (Hubrecht, 1887) *sensu* Takakura (1898) and Iwata (1952)

[Japanese name: murasaki-himomushi]

Lineus longifissus: **Takakura**, 1898: 336, fig. 19; obtained sublitorally from muddy sediment, Moroiso Bay, Kanagawa Prefecture. **Iwata**, 1952: 137–138; collected sublitorally among sandy mud at 30 cm depth; Tomioka, Amakusa, Kumamoto Prefecture. **Iwata**, 1960c: 166, pl. 83, fig. 12. **Iwata**, 1965b: 394, one figure. **Saito and Suzuki**, 1974: 38; intertidal, Niisaki Beach, Kanagawa Prefecture; identified by Dr. Iwata. **Crandall et al.**, 2002: 11, 18, 27, 34, 39.

NOTE: *Lineus longifissus* was originally described as *Cerebratulus longifissus* Hubrecht, 1887, based on material from Marion Island, South Africa obtained during the cruise of H.M.S. *Challenger*. It was later transferred to *Lineus* by Wheeler (1934: 255), and more recently to *Heteronemertes* by Chernyshev (1995: 15). *Lineus longifissus* differs from Takakura's (1898) and Iwata's (1952) descriptions in the degree of posterior extension of the lateral cephalic grooves; the Japanese form belongs to a different species and will be given a different name when it is redescribed.

Lineus mcintoshii (Langerhans, 1880) *sensu*

Takakura (1898)

Lineus Mcintoshii [sic]: **Takakura**, 1898: 187, fig. 10; intertidal, Koajiro, Kanagawa Prefecture.

NOTE: Collected among algae on a rocky shore in Madeira and originally described as *Cerebratulus mcintoshii* by Langerhans (1880), this species was later transferred to *Lineus* by Bürger (1904: 95). It appears to differ from Takakura's (1898) material in the color pattern of the cephalic region. The taxon recognized by Takakura (1898) must be given a different name when additional material has been found and redescribed.

Class HOPLONEMERTEA Hubrecht, 1879

Subclass MONOSTILIFERA Brinkmann, 1917

Family AMPHIPORIDAE McIntosh, 1874

Amphiporus cervicalis (Stimpson, 1857)

[Japanese name: yajirobei-himomushi]

Polina cervicalis Stimpson, 1857: 165; intertidal under stones, Shimoda, Shizuoka Prefecture; transferred to *Amphiporus* by Bürger (1904: 39).

Amphiporus cervicalis: **Iwata**, 1954a: 25–26; intertidal among mussels, Muroran and Rishiri Island, Hokkaidō Prefecture; intertidal among mussels, Kominato, Chiba Prefecture; intertidal among mussels, Shimoda, Shizuoka Prefecture.

Yamaguchi and Yamada, 1955: 70. **Utinomi**, 1956: 32, pl. 16, fig. 13. **Iwata**, 1960c: 169, pl. 84, fig. 15; habitat not recorded, Asamushi, Aomori Prefecture; habitat not recorded, Cape Muroto, Kōchi Prefecture. **Utinomi**, 1960:

32, pl. 16, fig. 13. **Iwata**, 1965a: 217. **Okuda and Iwata**, 1965: 399, figs. a–c. **Shiino**, 1969: 94. **Utinomi**, 1969: 32, pl. 16, fig. 13. **Tsuchiya**, 1979: 82; intertidal, Hadakajima Island, Asamushi, and Aomori Harbor, Aomori Prefecture. **Uchida et al.**, 1972: 55; habitat not recorded, Horomui, Hokkaidō Prefecture. **Iwata** 1992: 202, pl. 44–8. **Crandall et al.**, 2002: 9, 19, 25, 33.

Amphiporus cervicalis [sic]: **Inaba**, 1988: 226; lower intertidal to shallow sublittoral, on algae or under stones on gravelly to rocky shores; a specimen collected 18 July 1977 at Shijūshima, Hiroshima Prefecture, is deposited in Mukaishima Marine Biological Station, Hiroshima University.

NOTE: The original description of *Polina cervicalis* contains little information about the arrangement of the ocelli, and the description can be applied as well to the conditions in both *Amphiporus formidabilis* Griffin, 1898 and *A. imparispinosus* Griffin, 1898. Records of nemerteans under the name *Amphiporus cervicalis* from Japanese waters probably represent either *A. formidabilis* or *A. imparispinosus*, or even another taxon.

Amphiporus depressus (Stimpson, 1857)

Tatsnoskia depressa **Stimpson**, 1857: 165; sublittoral on sandy bottom at a depth of about 3–5 m, Hakodate, Hokkaidō Prefecture; originally recorded as “In portu ‘Hakodadi’ insulae ‘Jesso;’ in fundo arenoso, e 6–10 org. profundo accepta”; transferred to *Amphiporus* by Bürger (1904: 44).

Amphiporus depressus: **Iwata**, 1954a: 19–21, fig. 5C; lower intertidal under stones, Muroran, Hokkaidō Prefecture. **Yamaguchi and Yamada**, 1955: 69. **Crandall et al.**, 2002: 9, 19, 26, 33, 37.

NOTE: Gibson and Crandall (1989: 458) regarded *Amphiporus depressus* sensu Stimpson (1857) as a *nomen dubium*, but *Amphiporus depressus* sensu Iwata (1954a) as a different taxon. The illustration of Iwata’s (1954a) taxon resembles *Amphiporus imparispinosus*.

Amphiporus lactifloreus (Johnston, 1828)

Amphiporus lactifloreus: **Iwata**, 1954a: 23–24; intertidal under stones, Akkeshi and Muroran, Hokkaidō Prefecture. **Yamaguchi and Yamada**, 1955: 70. **Uchida et al.**, 1963: 17. **Crandall et al.**, 2002: 9, 19, 26, 33, 38.

NOTE: *Amphiporus lactifloreus* was originally described from the British Isles as *Planaria lactiflorea* by Johnston (1828: 489). Gibson (1995: 469) questioned the conspecificity between Iwata’s material and Johnston’s taxon. Iwata’s (1954a) taxon may represent *Amphiporus imparispinosus*.

Family CRATENEMERTIDAE

Nipponnemertes pulchra (Johnston, 1837)

Nipponnemertes pulchra: **Yamaoka**, 2005: 147, pl. 2, fig. 2, text fig. 5a–d; subtidal, 50 m depth, muddy sediment, off Kawazu, near Shimoda, Shizuoka Prefecture; habitat not recorded, Hashima Island, near Itō, Shizuoka Prefecture.

NOTE: *Nipponnemertes pulchra* was originally described as *Nemertes pulchra* from Berwickshire, UK, by Johnston (1837: 536). The species has been reported in the northern hemisphere from the east coast of North America, Greenland, the Faroe Islands, the White Sea, and northern

Europe from the Atlantic coast of France to Scandinavia; also reported in the southern hemisphere from Chile, South Africa, and Antarctica (Gibson, 1995). Yamaoka’s (2005) material differs from the other records of *N. pulchra* in possessing a white head, and probably represents a different species.

Family TETRASTEMMATIDAE Hubrecht, 1879

Oerstedia venusta Iwata, 1954

[Japanese name: hime-himomushi]

Oerstedia venusta **Iwata**, 1954a: 15–16, fig. 3; intertidal among algal holdfasts, Muroran and Rishiri Island, Hokkaidō Prefecture. **Yamaguchi and Yamada**, 1955: 68. **Iwata**, 1965b: 398, figs. a–f. **Crandall et al.**, 2002: 13, 20, 28, 35, 40.

NOTE: Envall and Sundberg (1993: 313) stated, “It is not possible from the brief description of this species to identify it to the genus *Oerstedia*.” Gibson (1995: 447) regarded the name *Oerstedia venusta* as a *nomen dubium*.

Prostoma graecense (Böhmig, 1892)

[Japanese name: mamizu-himomushi]

Prostoma graecense: **Ishizuka**, 1933: 215–218, figs. A, B; paddy fields and ponds, especially in chalybeate water, Sapporo, Hokkaidō Prefecture. **Sudzuki**, 1953: 218; in paddy fields around Urawa, Saitama Prefecture; in paddy fields around Sugashima, Mie Prefecture. **Iwata**, 1954a: 29. **Iwata**, 1970b: 128, fig. 91; Urawa, Saitama Prefecture. **Iwata**, 1973: 262. **Iwata**, 1992: 203, fig. 7–5G; Sapporo, Urawa, Sugashima.

Prostoma hokkaidoensis [sic] **Stiasny-Wijnhoff**, 1938: 222.

Prostoma lacstre [sic]: **Sudzuki**, 1953: 218; habitat not recorded, Sapporo, Hokkaidō Prefecture.

Prostoma graecens [sic]: **Iwata**, 1965a: 217; Urawa, Saitama Prefecture; Sapporo, Hokkaidō Prefecture.

Prostoma gracense [sic]: **Crandall et al.**, 2002: 14, 21, 29, 43.

Prostoma hokkaidoense: **Crandall et al.**, 2002: 14, 21, 29, 43.

NOTE: Ishizuka (1933) identified his material from Sapporo as *Prostoma graecense* (type locality: a local botanic garden in Graz, Austria). Stiasny-Wijnhoff (1938) created a new name, *Prostoma hokkaidoense*, to refer to the Sapporo form, attributing the naming authority to Ishizuka. Sudzuki (1953) regarded the Sapporo form as *Prostoma lacstre* (du Plessis, 1892) (type locality: under pebbles on a beach of Lac Léman at Anière, near Genève, Switzerland), while he identified the form from Urawa and Sugashima as *Prostoma graecense* (Böhmig, 1892). Iwata (1954a) listed *Prostoma hokkaidoense* and *Prostoma lacstre* sensu Sudzuki (1953) as synonymous with *Prostoma graecense*. Chernyshev et al. (1998: 62) argued that since previous records of *Prostoma* from Japan lack histological information about internal morphology, the specimens involved cannot be identified with certainty, and concluded that “all previous records of *Prostoma* from Japan should be cited as *Prostoma* sp. (or spp.?.”

Prostoma grande (Ikeda, 1913)

[Japanese name: mimizu-himomushi]

Stichostemma grandis **Ikeda**, 1913: 239–256, pl. IV, figs. 1–5;

a vessel planted with the aquatic plant *Lisichiton kamtschatense* Schott in the Botanic Garden of the Hiroshima Normal School, Hiroshima Prefecture. **Iwata**, 1957c: 101.

Prostoma grandis: **Kaburaki**, 1927: 1663, fig. 3183. **Miyashita**, 1932: 328; among *Potamogeton cristatus* covering the bottom of a small river, Setagaya, the Metropolis of Tôkyô. **Kaburaki**, 1947: 1468, fig. 4140. **Satô and Itô**, 1961: 187, fig. 7.1.10. **Iwata**, 1965a: 217. **Kaburaki and Iwata**, 1965: 400, figs. a, b. **Iwata**, 1973: 262. **Iwata**, 1983: 181.

Prossoma grandis [sic]: **Okugawa**, 1932: 70; in paddy fields over much of Japan, including Hokkaidô, but excepting Kyushu and Shikoku.

Prostoma lubricoideum [sic]: **Sudzuki**, 1953: 217–218.

Prostoma grande: **Kawakatsu et al.**, 1989: 47. **Crandall et al.**, 2002: 14, 21, 29, 43.

NOTE: Described from Hiroshima by Ikeda (1913), *Prostoma grande* has been widely reported from Japan by various authors. Sudzuki (1953) regarded *Prostoma grande* (Ikeda, 1913) as synonymous with *Prostoma lombricoideum* Dugès, 1830 (type locality: Montpellier [?], France). The comments of Chernyshev et al. (1998: 62) above (see 'NOTE' under *Prostoma graecense*) equally apply to *Prostoma grande*. Chernyshev et al. (1998: 62) further commented, "Future taxonomic studies on the comparative morphology and histology of *Prostoma* samples from elsewhere in Japan, including the type localities of the nominal species *P. grande* and *P. hokkaidoense*, are necessary."

Species That Cannot With Certainty Be Assigned to Valid Genera

Cosmocephala japonica Stimpson, 1857

Cosmocephala japonica **Stimpson**, 1857: 165; intertidal under stones and in rock crevices, Shimoda, Shizuoka Prefecture.

?*Amphiporus angulatus* [in part]: **Coe**, 1944: 30. **Crandall et al.**, 2002: 9, 19, 24, 33, 37.

Amphiporus japonicus: **Crandall et al.**, 2002: 9, 19, 26, 33.

NOTE: Bürger (1904: 48) regarded this form as a subspecific taxon, *Amphiporus angulatus japonicus*. Iwata (1952: 144) mentioned the similarity between the cephalic marking of *Amphiporus angulatus japonicus* and those of *Amphiporus punctatulus* Coe, 1905 (now *Nipponnemertes punctatula*); the former, having a uniform dorsal body coloration, can be differentiated from the latter, in which the dorsal coloration is mottled. Gibson and Crandall (1989: 460) regarded *Amphiporus japonicus* as a *nomen dubium*. The external appearance of a specimen I recently collected in Hiroshima Bay agrees with the original description of this species, but also resembles that of the *Nipponnemertes* species. This might mean that *Cosmocephala* is a senior synonym of *Nipponnemertes*, which must be determined by future studies. If this proves to be the case, however, the nomenclature of Cratene-mertidae will have to be altered to a large extent.

ORIGINAL DESCRIPTION: Stimpson (1857) gave the following diagnosis: "Corpus subelongatum, utrinque obtusum; lateribus in extentione fere parallelis. Color

supra brunnea, subtus alba; caput linea mediana et maculis minutis irregularibus incoloratis; fronte, et maculis cervicalibus triangularibus, albis. Caput breve subdiscretum fronte rotundata, ad aperturam profunde fissa. Cervix utrinque pseudorima obliqua, antrorum curvata. Ocelli sat magni, in capitib marginibus antero-lateralibus, utrinque 10–15. Long. 4; lat. 0.18 poll." [Free translation: Body somewhat elongated, sometimes blunt; when extended the lateral margins are parallel. Dorsally brown, ventrally white; head with a white median line and irregularly-shaped small white dots; anterior end of the head and triangle-shaped neck spots are white. Head wide, somewhat discrete, anteriorly rounded, deeply splits toward proboscis pore. Neck on each side with pseudo-crevices antero-obliquely curved. Eyes sufficiently large, arranged on the antero-lateral margins of the head, 10–15 on each side. 10 cm long, 4.5 mm wide.]

Dicelis rubra Stimpson, 1857

Dicelis rubra **Stimpson**, 1857: 164; sublittoral, between barnacles and sponges at a depth of about 7–8 m, Tanegashima, Kagoshima Prefecture.

NOTE: Bürger (1904) included this species in a group of dubious nemertean taxa. Gibson (1995) regarded the name as invalid.

ORIGINAL DESCRIPTION: Stimpson (1857: 164) gave the following diagnosis for the genus *Dicelis*: "Corpus lineare, depressiusculum, utrinque obtusum. Caput continuum vel subdiscretum, fronte emarginata, apertura proboscidis terminali. Ocelli duo simplices, rotundati, subterminales. Maricolae." [Free translation: Body filiform, dorsoventrally flattened, blunt on both ends. Head continuous to, or somewhat discrete from, body; frontally convex, proboscis pore terminal. Two rounded eyes subterminally. Marine]. The diagnosis for the species was given as: "Subfiliformis, depressiuscula, antice subattenuata; colore rubra vel purpurea. Cervix quam caput vix angustior. Caput antice rotundata et emarginata. Ocelli duo parvi subterminales. Long. 1.5; lat. 1.03 [sic. probably 0.03] poll." [Free translation: somewhat filiform, dorsoventrally flattened, anteriorly somewhat tapered; red or purple in color. Neck hardly narrower than head. Head anteriorly rounded and convex. Two eyes slightly subterminally. 3.75 cm long, 0.75 mm wide.]

Dichilus obscurus Stimpson, 1857

Dichilus obscurus **Stimpson**, 1857: 163; intertidal between stones, Amamiôshima, Kagoshima Prefecture; originally recorded as "In portu insulae 'Ousima;' littoralis inter lapillus."

NOTE: Bürger (1904) included this species in a group of dubious nemertean taxa. Gibson (1995) regarded the name as invalid.

ORIGINAL DESCRIPTION: Stimpson (1857: 163) gave the following diagnosis for the genus *Dichilus*: "Corpus lineare depresso, longitudine mediocre. Caput corpori continuum subquadratum, plica transversa terminali bilabiatum; labio inferiore emarginato. Ocelli duo subterminales. Cervix supra rimis obsoletis (pseudorimis) impressa. Maricolae." [Free translation: head somewhat rectangular, continuous to body, transverse fold terminally bilobed; lower lip concave. Two ocelli subterminally. Neck dorsally with

pseudo-crevices. Marine.]. The diagnosis for the species is given as: "Corpus supra pallide rubro-fulvum, maculis duabus oblongis in capite. Ocelli fusi, sat magni, subdistantes, in maculis siti. Pseudorimae cervicales tres; una mediana longitudinalis, ex cujus media aliae versus marginem utrinque oblique extendunt. Long. 3; lat. 0.08 poll." [Free translation: Body dorsally pale reddish brown, with two oblong cephalic patches. Ocelli brown, large, rather separately situated in the cephalic patch. Three pseudo-crevices on neck; one medioligitudinal, from which other median ones extend obliquely towards the margins respectively. 7.5 cm long, 2mm wide.]

Diplomma serpentina (Stimpson, 1855)

Nareda serpentina **Stimpson**, 1855: 381; habitat not recorded, Okinawa Prefecture.

Diplomma serpentina: **Stimpson**, 1857: 164; intertidal under stones on muddy sand, Okinawa Prefecture.

NOTE: Bürger (1904) included this species in a group of dubious nemertean taxa. Gibson (1995) regarded the name as invalid. It appears that *Diplomma serpentina* is conspecific with *Amphiporus insolitus* Iwata, 1954 and *Paranemertes* sp. *sensu* Yamaoka (2005). The identity of this taxon should be clarified by future studies.

ORIGINAL DESCRIPTION: "Elongated, somewhat flattened, brownish; head broader than the body, emarginate in front; neck well contracted; eyes two, rather large, bilobate, placed one on each side at the middle of the head. Length 2 1/2 inches" (Stimpson, 1855: 381).

Records of Specimens Not Identified to Species

Class PALAEONEMERTEA Hubrecht, 1879

Family TUBULANIDAE Bürger, 1904 (1874)

Carinella sp.

Carinella sp. **Takakura**, 1898: 119; Misaki, Kanagawa Prefecture.

NOTE: The genus *Carinella* Johnston, 1833 was synonymized with *Tubulanus* Renier, 1804, by Bürger (1904). This form differs from *Tubulanus punctatus* (Takakura, 1898) in possessing continuous longitudinal stripes on the mid-dorsal and lateral surfaces of the body.

Class HOPLONEMERTEA Hubrecht, 1879

Subclass MONOSTILIFERA Brinkmann, 1917

Family AMPHIPORIDAE McIntosh, 1874

Amphiporus sp.

Amphiporus sp. **Yamaoka**, 2005: 147, pl. 1, fig. 3; text fig. 4b, c; among shelly bottom, several meters depth, Susaki, Shizuoka Prefecture.

NOTE: *Amphiporus* sp. *sensu* Yamaoka (2005) is well illustrated, which will facilitate identification when this species is again encountered; the external characters of this form include the ovoid-shaped head, the eyes arranged on the edges of the head in front of the anterior cephalic furrows, and the uniformly pinkish-red body coloration. The generic placement of this form may require further assessment based on new material in future studies.

Zygonemertes sp.

Zygonemertes sp. **Iwata**, 1954a: 19, fig. 5A; under a stone

near low tide mark, Muroran, Hokkaidō Prefecture.
Yamaguchi and Yamada, 1955: 69.

NOTE: This form can be identified as a member of the genera *Zygonemertes* or *Pheroneonemertes* by possessing post-cerebral ocelli; proper generic identification of this form will require histological examination of its internal morphology. The body is about 3 cm long, 1 mm wide, pale blue in color without any marking. There are short, double longitudinal lines on the mid-dorsal surface of the head (Iwata, 1954a).

Family CRATENEMERTIDAE Friedrich, 1968

Nipponnemertes sp. 1

Nipponnemertes sp. 1. **Yamaoka**, 2005: 151, pl. 2, fig. 3; text figs. 6b, c, 7; intertidal, under stones, Shitaru, near Shimoda, Shizuoka Prefecture; shelly bottom, several meters depth, Susaki, Shimoda, Shizuoka Prefecture.

NOTE: The form can be identified as *Nipponnemertes* by the interwoven longitudinal and circular muscle fibers in the rhynchocoel wall. Detailed external features illustrated by Yamaoka (2005) will suffice for identification when this species is again encountered. The body is 2.5 cm long, 0.8 mm wide, yellowish brown in color, rarely with small brown dots; about 15 eyes are irregularly arranged on either side of head. Remarkably, Yamaoka's (2005) specimen possessed only one accessory stylet pouch.

Nipponnemertes sp. 2

Nipponnemertes sp. 2. **Yamaoka**, 2005: 152, text fig. 8; subtidal, several meters depth, Susaki, Shimoda, Shizuoka Prefecture.

NOTE: Yamaoka's (2005) specimens were 10–12 mm long, 0.7 mm wide; anterior cephalic furrows ventrally forming M-shape; posterior cephalic furrows encircling body in esophageal region, curving forward on both dorsal and ventral surfaces; body color pure white, yellowish white, or yellow, with scattered small brown patches. The large cerebral organs extending behind the brain in this form is characteristic of the Cratenemertidae.

Family EMPLECTONEMATIDAE Bürger, 1904

Paranemertes sp.

Paranemertes sp. **Yamaoka**, 2005: 142, pl. 1, fig. 6; text fig. 1; Itado, near Shimoda, Shizuoka Prefecture.

NOTE: Body 6 cm long, 0.7 mm wide; uniformly bright chestnut-brown in color, except for white margins on cephalic tip. This form may be conspecific with *Diplomma serpentina* (Stimpson, 1855) and *Amphiporus insolitus* Iwata, 1954. Future study should clarify the identity of this taxon.

Family TETRASTEMMATIDAE Hubrecht, 1879

Prostoma sp.

[Japanese name: toyama-mamizu-himomushi]

Prostoma sp. **Iwata**, 1997: 53, with two color photographs taken in life by Dr. Fumio Iwata; under stones near a spring in a small pond near a paddy field, Asahi-chō, Toyama Prefecture.

NOTE: The body is about 1 cm long, 0.5 mm wide, pale orange in color. The proboscis retractor muscle is well developed (Iwata, 1997).

Tetrastemma sp.

Tetrastemma sp. **Iwata**, 1954a: 30, fig. 8A; among seaweeds, Akkeshi, Hokkaidō Prefecture. **Yamaguchi and Yamada**, 1955: 71, fig. 18-1.

NOTE: Body 2 cm long, 1 mm wide, yellowish green in basement body color, with four darker longitudinal stripes on the dorsal surface; with four eyes (Iwata, 1954a).

Tetrastemma sp.

Tetrastemma sp. **Yamaoka**, 2005: 155, pl. 2, fig. 1, text figs. 9d, e, 10; subtidal, about 5 fathoms deep, Shirahama, near Shimoda, Shizuoka Prefecture.

NOTE: Body 1 cm long, 2 mm wide; dorsally reddish brown, paler ventrally, with dark brown mid-dorsal stripe (Yamaoka, 2005).

ACKNOWLEDGMENTS

Thanks are due to Dr. Mark J. Grygier, Lake Biwa Museum, for information about the type specimens of *Prostoma ohmienne*, and to Dr. Manabu Asakawa, Hiroshima University, for his unpublished toxicological data on nemerteans from some places in Japan. I wish to express my gratitude to Dr. Alexei V. Chernyshev (Institute of Marine Biology, Far East Division, Russian Academy of Sciences, Russia), Dr. Frank B. Crandall (Department of Invertebrate Zoology, National Museum of Natural History, USA), Dr. Ray Gibson (Professor Emeritus of Liverpool John Moores University, UK), Dr. Svetlana A. Maslakova (University of Washington, USA), Dr. Jon L. Norenburg (Department of Invertebrate Zoology, National Museum of Natural History, USA) and Dr. Sun Shichun (Mariculture Research Lab, Ocean University of Qingdao, China) for their valuable comments on an earlier version of the manuscript. This study was supported by Grants-in-Aid from the Japan Society for the Promotion of Science (research grant numbers 16770059 and 16207005).

REFERENCES

- Abildgaard PC (1806) *Planaria dorsalis*. In "Zoologia Danica Seu Animalium Daniae et Norvegiae Rariorum ac Minus Notorum Descriptiones et Historia Vol 4" Ed by OF Müller, N Christensen, N Möller et Fillii, Havniae, p 25
- Ali AE, Arakawa O, Noguchi T, Miyazawa K, Shida Y, Hashimoto K (1990) Tetrodotoxin and related substances in a ribbon worm *Cephalothrix linearis* (Nemertean). Toxicon 28: 1083–1093
- Asakawa M, Toyoshima T, Shida Y, Noguchi T, Miyazawa K (2000) Paralytic toxins in a ribbon worm *Cephalothrix* species (Nemertean) adherent to cultured oysters in Hiroshima Bay, Hiroshima Prefecture, Japan. Toxicon 38: 763–773
- Asakawa M, Toyoshima T, Ito K, Bessho K, Yamaguchi C, Tsunetsugu S, Shida Y, Kajihara H, Mawatari SF, Noguchi T, Miyazawa K (2003) Paralytic toxicity in a ribbon worm *Cephalothrix* species (Nemertea) in Hiroshima Bay, Hiroshima Prefecture, Japan and the isolation of tetrodotoxin as a main component of its toxins. Toxicon 41: 747–753
- Bergendal D (1900) Über ein Paar sehr eigenthümliche nordische Nemertinen. Zool Anz 23: 313–328
- Bergendal D (1902) Zur Kenntnis der nordischen Nemertinen. 2. Eine der construierten Urnemertine entsprechende Palaeonemertine aus dem Meere der schwedischen Westküste. Zool Anz 25: 421–432
- Blainville HM de (1827) Vers. In "Dictionnaire des Sciences Naturelles Vol 57" Ed by FG Levrault, Normant, Paris, pp 270–271
- Blake JA (1993) Phylum Nemertea. In "Taxonomic Atlas of the Benthic Fauna of the Santa Maria Basin and Western Santa Barbara Channel Vol 1 Introduction, Benthic Ecology, Oceanology, Platyhelminthes, and Nemertea" Ed by JA Blake, A Lissner, Santa Barbara Museum of Natural History, California, pp 95–132
- Boyer BC, Henry JQ (1998) Evolutionary modifications of the spiralian developmental program. Am Zool 38: 621–633
- Brinkmann A (1917) Die pelagischen Nemertinen. Bergens Mus Skr 3: 1–180
- Britton JC (1990) The intertidal crevice fauna of Tolo Channel and Harbour, New Territories, Hong Kong. In "Proceedings of the Second International Marine Biological Workshop: The Marine Flora and Fauna of Hong Kong and Southern China, Hong Kong, 1986, Vol 2" Ed By B Morton, Hong Kong University Press, Hong Kong, pp 803–835
- Brunberg L (1964) On the nemertean fauna of Danish waters. Ophelia 1: 77–111
- Bürger O (1890) Untersuchungen über die Anatomie und Histologie der Nemertinen nebst Beiträgen zur Systematik. Z Wiss Zool 50: 1–277
- Bürger O (1892) Zur Systematik der Nemertinenfauna des Golfs von Neapel. Nachr Königl Gessels Wiss Georg
- Bürger O (1895) Nemertinen. F F Golf Neapel 22: 1–743
- Bürger O (1904) Nemertini. Tierreich 20: 1–151
- Bürger O (1897–1907) Nemertini (Schnurwürmer). In "Klassen und Ordnungen des Tier-Reichs, wissenschaftlich dargestellt in Wort und Bild Vol 4 Supplement" Ed by HG Bronn, CF Winter'sche Verlagshandlung, Leipzig, pp 1–542 [1897, pp 1–176; 1898, pp 177–240; 1899, pp 241–288; 1903, pp 289–384; 1905, pp 385–480; 1907, pp 481–542]
- Chernyshev AV (1992) On the names of some nemertines. Zool Zh 71: 134–136 [In Russian with English abstract]
- Chernyshev AV (1993) A review of the genera of nemerteans allied to *Oerstedia* (Monostilifera, Tetrastematidae) with description of four new species. Zool Zh 72: 11–20 [In Russian with English abstract]
- Chernyshev AV (1995) On the higher taxa of the phylum Nemertea with the taxonomic review of the subclass Anopla. Zool Zh 74: 7–18
- Chernyshev AV (1998) Nemerteans of the genus *Tetrastemma* (Enopla, Monostilifera) from the Far East seas of Russia. Zool Zh 77: 995–1002 [In Russian with English abstract]
- Chernyshev AV (2003) Classification system of the higher taxa of enoplan nemerteans (Nemertea, Enopla). Russ J Mar Biol 29: 57–65
- Chernyshev AV (2004a) Nemertini. In "Dalnevostochnyi Morskoi Biosfernyi Zadovednik Biota Vol 2" [Far Eastern Marine Biosphere Nature Reserve Biota] Ed by AN Tyurin, Dalnauka, Vladivostok, pp 151–155
- Chernyshev AV (2004b) Two new genera of nemertean worms of the family Tetrastematidae (Nemertea: Monostilifera). Zool Syst Russ 12: 151–156
- Chernyshev AV (2004c) Problems of taxonomy of the "*Lineus ruber*" heteronemertean complex (Nemertea, Anopla). Zool Zh 88: 788–794 [In Russian with English abstract]
- Chernyshev AV, Timoshkin OA, Kawakatsu M (1998) *Prostoma ohmienne* sp. nov., a new species of freshwater nemertean from Lake Biwa-Ko, central Japan, with special reference to the taxonomy and distribution of the known species in the genus *Prostoma* Dugès, 1828 (Enopla, Hoplonemertea, Monostilifera, Tetrastematidae). Bull Fuji Women's Coll Ser II 36: 51–66
- Coe WR (1901) Papers from the Harriman Alaska Expedition. XX. The nemerteans. Proc Wash Acad Sci 3: 1–84
- Coe WR (1902) The nemertean parasites of crabs. Amer Nat 36: 431–450
- Coe WR (1904) Nemerteans of the Pacific coast of North America. Part II. Harriman Alaska Ser 11: 111–220
- Coe WR (1905) Nemerteans of the west and northwest coasts of America. Bull Mus Comp Zool Harvard Coll 47: 1–318
- Coe WR (1926) The pelagic nemerteans. Mem Mus Comp Zool Harvard Coll 49: 1–244

- Coe WR (1930) Two new species of nemerteans belonging to the family Cephalotrichidae. Zool Anz 89: 97–103
- Coe WR (1931) A new species of nemertean (*Lineus vegetus*) with asexual reproduction. Zool Anz 94: 54–60
- Coe WR (1940) Revision of the nemertean fauna of the Pacific coasts of North, Central, and northern South America. Allan Hancock Pac Exped 2: 247–322
- Coe WR (1943) Biology of the nemerteans of the Atlantic coast of North America. Trans Conn Acad Arts Sci 35: 129–328
- Coe WR (1944) Geographical distribution of the nemerteans of the Pacific coast of North America, with descriptions of two new species. J Wash Acad Sci 34: 27–32
- Coe WR (1947) Nemerteans of the Hawaiian and Marshall Islands. Occ Pap Bernice P Bishop Mus 19: 101–106
- Coe WR (1954) Bathypelagic nemerteans of the Pacific Ocean. Bull Scripps Inst Oceanogr Univ Califor 6: 225–286
- Corrêa DD (1963) Nemerteans from Curaçao. Stud Faun Curaçao Carib Is 17: 41–56
- Corrêa DD (1964) Nemerteans from California and Oregon. Proc Calif Acad Sci 31: 515–558
- Corrêa DD (1966) A new hermaphroditic nemertean. An Acad Brasil Ciênc 38: 365–369
- Crandall FB (2001) A cladistic view of the Monostilifera (Hoplonephentea) with interwoven rhynchocoel musculature: a preliminary assessment. Hydrobiologia 456: 87–110
- Crandall FB, Kajihara H, Mawatari SF, Iwata F (2001) The status of four Japanese nemertean species of Yamaoka. Hydrobiologia 456: 175–185
- Crandall FB, Norenburg JL, Chernyshev AV, Maslakova S, Schwartz M, Kajihara H (2002) Checklist of the Nemertean Fauna of Japan and Northeastern Asia. The Smithsonian Institution, Washington, D.C. [PDF available from <http://nemertes.si.edu/PDFs/epub2917.pdf>]
- Delle Chiaje S (1825) Memorie Sulla Storia e Notomia Degli Animali Senza Vertebre del Regno di Napoli Vol 2. Societa' Tipografica, Napoli
- Delle Chiaje S (1828) Memorie Sulla Storia e Notomia Degli Animali Senza Vertebre del Regno di Napoli Vol 3. Societa' Tipografica, Napoli
- Diesing CM (1850) Systema Helminthum Vol I. W Braumuller, Vindobonae
- Diesing KM (1863) Nachträge zur Revision der Turbellarien. Sitzungsber Akad Wiss Wien Math Naturwiss Kl 46: 173–188
- Dugès A (1828) Recherches sur l'organisation et les moeurs des Planariées. Ann Sci Nat 15: 139–183
- Ehrenberg CG (1828–1831) Phytozoa turbellaria Africana et Asiatica in Phytozoorum Tabula IV et V delineata. In "Symbolae physicae, seu icones et descriptiones corporum naturalium novorum aut minus cognitorum quae ex itineribus per Libyam, Aegyptium, Nubiam, Dongalam Syriam, Arabiam et Habessiniam, pars zoologica II, animalia evertebrata exclusis insectis" Ed by FG Hemprich, CG Ehrenberg, Officina Academica, Berolina, pp 53–67, pls IV–V [Plates published in 1828, text in 1831]
- Envall M (1998) General problems in estimating nemertean relationships on ribosomal sequence data — an example using six monostiliferous species and mitochondrial 16S rDNA. Hydrobiologia 365: 19–31
- Envall M, Sundberg P (1993) Intraspecific variation in nemerteans (Nemertea): synonymization of the genera *Paroerstedia* and *Oerstediella* with *Oerstedia*. J Zool Lond 230: 293–318
- Fish JD, Fish S (1989) A Student's Guide to the Seashore. Unwin Hyman, London
- Foshay EA (1912) *Nectonemertes japonica*, a new nemertean. Zool Anz 40: 50–53
- Friedrich H (1935a) Studien zur Morphologie, Systematik und Ökologie der Nemertinen der Kieler Bucht. Arch Naturg NF 4: 293–375
- Friedrich H (1935b) Neue Hoplonemertinen der Kieler Bucht. Schrift Naturw Ver Schleswig-Holst 21: 10–19
- Friedrich H (1955) Beiträge zu einer Synopsis der Gattungen der Nemertini monostilifera nebst Bestimmungsschlüssel. Z Wiss Zool 158: 133–192
- Friedrich H (1958) Nemertini. Zool Iceland 2: 1–24
- Friedrich H (1968) *Sagaminemertes*, eine bemerkenswerte neue Gattung der Hoplonemertinen und ihre systematische Stellung. Zool Anz 180: 33–36
- Gibson R (1979) Nemerteans of the Great Barrier Reef. 1. Anopla Palaeonemertea. Zool J Linn Soc 65: 305–337
- Gibson R (1981) Nemerteans of the Great Barrier Reef 3. Anopla Heteronemertea (Lineidae). Zool J Linn Soc 71: 171–235
- Gibson R (1982a) British Nemerteans. University Press, Cambridge
- Gibson R (1982b) Nemertea. In "Synopsis and Classification of Living Organisms Vol 1" Ed by SP Parker, McGraw-Hill, New York, pp 823–846
- Gibson R (1985) The need for a standard approach to taxonomic descriptions of nemerteans. Am Zool 25: 5–14
- Gibson R (1990a) The macrobenthic nemertean fauna of Hong Kong. In "Proceedings of the Second International Marine Biological Workshop: The Marine Flora and Fauna of Hong Kong and Southern China, Hong Kong, 1986 Vol 1" Ed by B Morton, Hong Kong University Press, Hong Kong, pp 33–212
- Gibson R (1990b) The macrobenthic nemertean fauna of the Albany region, Western Australia. In "Proceedings of the Third International Marine Biological Workshop: the Marine Flora and Fauna of Albany, Western Australia Vol 1" Ed by FE Wells, DI Walker, H Kirkman, R Lethbridge, Western Australian Museum, Perth, pp 89–194
- Gibson R (1994) Nemerteans. Field Studies Council, Shrewsbury
- Gibson R (1995) Nemertean genera and species of the world: an annotated checklist of original names and description citations, synonyms, current taxonomic status, habitats and recorded zoogeographic distribution. J Nat Hist 29: 271–562
- Gibson R (1997a) An annotated checklist of the Nemertea recorded from the Cape d'Aguilar Marine Reserve, Hong Kong. In "The Marine Flora and Fauna of Hong Kong and Southern China Vol 4" Ed by B Morton, Hong Kong University Press, Hong Kong, pp 31–36
- Gibson R (1997b) Nemertea. In "The Species Directory of the Marine Fauna and Flora of the British Isles and Surrounding Seas" Ed by C M Howson, BE Picton, Ulster Museum and The Marine Conservation Society, Belfast and Ross-on-Wye, pp 49–54
- Gibson R (1998) Epilogue - one hundred years of nemertean research: Bürger (1895) to the present. Hydrobiologia 365: 301–310
- Gibson R, Crandall FB (1989) The genus *Amphiporus* Ehrenberg (Nemertea, Enopla, Monostiliferoidea). Zool Scr 18: 453–470
- Gibson R, Knight-Jones EW (1990) Platyhelminthes, Nematoda, Nemertea. In "The Marine Fauna of the British Isles and Northwest Europe Vol 1" Ed by PJ Hayward, JS Ryland, Clarendon Press, Oxford, pp 181–200
- Gibson R, Moore J (1976) Freshwater nemerteans. Zool J Linn Soc 58: 177–218
- Gibson R, Moore J (1998) Further observations on the genus *Geonemertes* with a description of a new species from the Philippine Islands. Hydrobiologia 365: 157–171
- Girard C (1893) Recherches sur les Planariés et les Némertiens de l'Amérique du Nord. Ann Sci Nat Sér 7 15: 145–310
- Hansson HG (1994) Sydskandinaviska marina flercelliga evertebrater. Lanstryckeriet i Göteborg, Göteborg
- Henry JQ, Martindale MQ (1994) Establishment of the dorsoventral axis in nemertean embryos: evolutionary considerations of spiralian development. Dev Gen 15: 64–78
- Henry JQ, Martindale MQ (1996) The establishment of embryonic

- axial properties in the nemertean, *Cerebratulus lacteus*. Dev Biol 180: 713–721
- Henry J, Martindale MQ (1997a) Nemerteans, the ribbon worms. In “Embryology: Constructing the Organism” Ed by SF Gilbert, AM Raunio, Sinauer Associates, Sunderland, pp 151–166
- Henry JQ, Martindale MQ (1997b) Regulation and the modification of axial properties in partial embryos of the nemertean, *Cerebratulus lacteus*. Dev Gen Evol 207: 42–50
- Hieda K, Takahashi Y (1986) Hokkaido Umibenoikimono. [Sea Shore Animals Around Hokkaido] Hokkaido Shimbun, Sapporo [In Japanese]
- Hochberg FG, Lunianski DN (1998) Nemertean collections at the Santa Barbara Museum of Natural History: type specimens and vouchers for Wesley R. Coe's 1940 publication. Hydrobiologia 365: 291–300
- Honma Y, Kitami T (1978) Fauna and flora in the waters adjacent to the Sado Marine Biological Station, Niigata University. Ann Rep Sado Mar Biol Stat Niigata Univ 8: 7–81
- Hubrecht AAW (1874) Aanteekeningen Over de Anatomie, Histologie en Ontwikkelingsgeschiedenis van Eneige Nemertinen. JL Beijer, Utrecht
- Hubrecht AAW (1879) The genera of European nemerteans critically revised, with description of several new species. Note Leyden Mus 1: 193–232
- Hubrecht AAW (1885) The Nemertea. Rep Sci Res Challenger Exped, Narrative 1: 830–833
- Hubrecht AAW (1887) Report on the Nemertea collected by H.M.S. Challenger during the years 1873–76. Rep Sci Res Challenger Exped, Zoology 19: 1–150
- Humes AG (1942) The morphology, taxonomy, and bionomics of the nemertean genus *Carcinonemertes*. Illinois Biol Monogr 18: 1–105
- Huschke E (1830) Beschreibung und Anatomie eines neuen an Sicilien gefundenen Meerwurms, *Notospermus drepanensis*. Isis 23: 681–683
- Hylbom R (1957) Studies on palaeonemerteans of the Gullmar Fiord area (West Coast of Sweden). Arkiv Zool 10: 539–582
- Hylbom R (1993) A proposal for a check-list of characteristics to be used in the descriptions of palaeonemertean species. Hydrobiologia 266: 169–174
- Ikeda I (1913) A new fresh-water nemertine from Japan (*Stichostemma grandis*). Ann Zool Jpn 8: 239–256
- Inaba A (1963) Fauna and Flora of the Inland Sea of Seto. Mukaishima Marine Biological Station, Faculty of Science, Hiroshima University, Hiroshima
- Inaba A (1988) Fauna and Flora of the Seto Inland Sea. 2nd ed, Mukaishima Marine Biological Station, Faculty of Science, Hiroshima University, Hiroshima
- International Commission on Zoological Nomenclature (1985) International Code of Zoological Nomenclature. 3rd ed, International Trust for Zoological Nomenclature, London
- International Commission on Zoological Nomenclature (1988) Opinion 1486, *Tubulanus* Renier, [1804] and *T. polymorphus* Renier, [1804] (Nemertea): reinstated and made available. Bull Zool Nom 45: 157–158
- International Commission on Zoological Nomenclature (1992) OPINION 1675 *Amphiporus* Ehrenberg, 1831 (Nemertea): *Planaria lactiflorea* Johnston, 1828 designated as the type species. Bull Zool Nom 49: 157
- International Commission on Zoological Nomenclature (1999) International Code of Zoological Nomenclature. 4th ed, International Trust for Zoological Nomenclature, London
- Ishizuka H (1933) On a fresh-water nemertean from Hokkaido. Proc Imp Acad Jpn 9: 215–218
- Iwata F (1951) Nemerteans in the vicinity of Onomichi. J Fac Sci Hokkaido Univ Ser 6 Zool 10: 135–138
- Iwata F (1952) Nemertini from the coasts of Kyusyu. J Fac Sci Hokkaido Univ Ser 6 Zool 11: 126–148
- Iwata F (1954a) The fauna of Akkeshi Bay. XX. Nemertini in Hokkaido (revised report). J Fac Sci Hokkaido Univ Ser 6 Zool 12: 1–39
- Iwata F (1954b) Some nemerteans from the coasts of the Kii Peninsula. Publ Seto Mar Biol Lab 4: 33–42
- Iwata F (1954c) Invertebrate fauna of the intertidal zone of the Tokara Islands. X. Nemertini. Publ Seto Mar Biol Lab 4: 27–31
- Iwata F (1957a) Nemerteans from Sagami Bay. Publ Akkeshi Mar Biol Stat 7: 1–31
- Iwata F (1957b) On the early development of the nemertine, *Lineus torquatus* Coe. J Fac Sci Hokkaido Univ Ser 6 Zool 13: 54–58
- Iwata F (1957c) Nemertini. In “Musekitsui-dōbutsu-hasseigaku” [Developmental Biology of Invertebrates] Ed by M Kume, K Dan, Baifūkan, Tokyo, pp 101–110 [In Japanese]
- Iwata F (1958) On the development of the nemertean *Micrura akkeshiensis*. Embryologia 4: 103–131
- Iwata F (1960a) The life history of the Nemertea. Bull Mar Biol Stat Asamushi 10: 95–97
- Iwata F (1960b) Studies on the comparative embryology of nemerteans with special reference to their interrelationships. Publ Akkeshi Mar Biol Stat 10: 1–51
- Iwata F (1960c) Nemertini. In “Encyclopaedia Zoologica Illustrated in Colours Vol 4” Ed by T Okada, T Uchida, Hokuryu-kan Publishing Co., Ltd., Tokyo, 166–169 [In Japanese]
- Iwata F (1965a) Nemertinea. In “Dōbutsu-keitō-bunruigaku Vol 3” [Systematic Zoology Vol 3] Ed by T Uchida, Nakayama-Shoten Co., Ltd, Tokyo, pp 167–219 [In Japanese]
- Iwata F (1965b) *Procephalothrix simulus* Iwata, *Tubulanus punctatus* Takakura, *Carinesta uchidai* Iwata, *Coia ijimai* Takakura, *Tetramys ramicerebrum* Iwata, *Baseodiscus curtus* Hubrecht, *Baseodiscus princeps* Coe, *Cephalomastax brevis* Iwata, *Euborlasia gotoensis* Iwata, *Lineus alborostratus* Takakura, *Lineus bilineatus* Renier, *Lineus geniculatus* Delle Chiaje, *Lineus longifissus* Hubrecht, *Lineus piperatus* Stimpson, *Micrura japonica* Iwata, *Emplectonema gracile* Johnston, *Nemertopsis gracilis* Coe, *Paranemertes plana* Iwata, *Oerstedia dorsalis* Abildgaard, *Oerstedia venusta* Iwata, *Amphiporus bimaculatus* Coe, *Amphiporus punctatulus* Coe, *Drepanophorus longiceps* Iwata. In “New Illustrated Encyclopedia of the Fauna of Japan Vol 1” Ed by Y Okada, S Uchida, T Uchida, Hokuryu-kan Publishing Co., Ltd., Tokyo, pp 390–401 [In Japanese]
- Iwata F (1967) *Uchidana parasita* nov. gen. et nov. sp., a new parasitic nemertean from Japan with peculiar morphological characters. Zool Anz 178: 122–136
- Iwata F (1970a) On the brackish water nemerteans from Japan, provided with special circulatory and nephridial organs useful for osmoregulation. Zool Anz 184: 133–154
- Iwata F (1970b) Nemertinea. In “Gendai-seibutugaku-taikei Vol 1 musekitui-dōbutsu” [Contemporary Biology Series Vol 1 Invertebrates] Ed by T Uchida, S Mawatari, Nakayama-shoten, Tokyo, pp 125–132 [In Japanese]
- Iwata F (1972) Axial changes in the nemertean egg and embryo during development and its phylogenetic significance. J Zool Lond 168: 521–526
- Iwata F (1973) Nemertinea. In “Freshwater Biology of Japan Enlarged and Revised Edition” Ed by M Uéno, Hokuryukan Publishing Co., Ltd., Tokyo, pp 262–264 [In Japanese]
- Iwata F (1983) Nemertinea. In “Musekitui-dōbutsu-no-hassei-jō” [Development of Invertebrates Vol 1] Ed by K Dan, K Sekiguchi, Y Andō, H Watanabe, Baifūkan, Tokyo, pp 181–196 [In Japanese]
- Iwata F (1985) Foregut formation of the nemerteans and its role in nemertean systematics. Amer Zool 25: 23–36
- Iwata F (1988) On the hoplonemertean *Sagaminemertes nagaiensis* (Iwata, 1957), with consideration of its systematics. Hydrobiologia 156: 115–124

- Iwata F (1992) Nemertinea. In "Guide to Seashore Animals of Japan with Color Pictures and Keys Vol 1" Ed by S Nishimura, Hoikusha Publishing Co., Ltd., Osaka, pp 194–204 [In Japanese]
- Iwata F (1993) *Paralineopsis taki* gen. et sp. nov., a littoral heteronemertean from Japan, provided with special proboscideal, circulatory and sensory organs of significance to nemertean systematics. *Hydrobiologia* 266: 185–201
- Iwata F (1997) Himogatadōbutsu (Himomushirui). In "The Encyclopaedia of Animals in Japan Vol 7 Invertebrates" Ed by T Okutani, M Takeda, M Imafuku, Heibonsha Ltd., Tokyo, pp 50–55 [In Japanese]
- Iwata F (1998) On the hoplonemertean, *Kameginemertes parmior-natus* (Iwata, 1957) gen. n., comb. n. from Sagami Bay, Japan. *Hydrobiologia* 365: 199–213
- Iwata F (2001) *Nipponnemertes fernaldi*, a new species of swimming monostiliferous hoplonemertean from the San Juan Archipelago, Washington, U.S.A. *Proc Biol Soc Wash* 114: 833–857
- Iwata F (2006) On four monostiliferous hoplonemerteans, including three new genera and species from Washington state and British Columbia. *J Nat Hist* 40: 873–913
- Jenner RA (2004) Towards a phylogeny of the Metazoa: evaluating alternative phylogenetic positions of Platyhelminthes, Nemertea, and Gnathostomulida, with a critical reappraisal of cladistic characters. *Contr Zool* 73: 1–216 [Available from <http://dpc2.uba.uva.nl/ctz/vol73/nr01/art01>]
- Jensen OS (1878) Turbellarier ved Norges Vestkyst. JW Eides, Bergen
- Johnston G (1828) Contributions to the British fauna. *Zool J* 4: 52–57
- Johnston G (1833) Illustrations in British Zoology. *Mag Nat Hist* 6: 232–235
- Johnston G (1837) A description of some planarian worms. *Mag Zool Bot* 1: 529–538
- Joubin L (1894) Les Némertiens. In "Fauna Française" Ed by R Blanchard, J de Guerne, Société d'Éditions Scientifiques, Paris, pp 1–235
- Kaburaki T (1927) Nemertea. In "Figuraro de Japanaj Bestoj" Ed by S Uchida, Hokuryukan & Co., Ltd., Tokyo, pp 1662–1666 [In Japanese]
- Kaburaki T (1947) *Emplectonema mitsukurii* (Takakura), *Prostoma grandis* (Ikeda), *Malacobdella japonica* Takakura, *Lineus fuscoviridis* Takakura, *Lineus subcingulatus* Takakura, *Cerebratulus communis* Takakura, *Baseodiscus curtus* (Hubrecht), *Cephalothrix linearis* (Rathke), *Tubulanus punctatus* (Takakura). In "Revised Edition Illustrated Encyclopedia of the Fauna of Japan (Exclusive of Insects)" Ed by S Uchida et al. The Hokuryukan Co., Ltd., Tokyo, pp 1466–1474 [In Japanese]
- Kaburaki T, Iwata F (1965) *Cephalothrix linearis* Rathke, *Lineus fuscoviridis* Takakura, *Lineus subcingulatus* Takakura, *Cerebratulus communis* Takakura, *Carcinonemertes mitsukurii* Takakura, *Prostoma grandis* Ikeda, *Malacobdella japonica* Takakura. In "New Illustrated Encyclopedia of the Fauna of Japan Vol 1" Ed by Y Okada, S Uchida, T Uchida, Hokuryu-kan Publishing Co., Ltd., Tokyo, pp 390–401 [In Japanese]
- Kajihara H (1998) Kangekisei-himomushirui — sunano-sukima-ni-sumu-himomushi [Interstitial nemerteans]. *Umiushi Tsūshin* 19: 10–11 [In Japanese]
- Kajihara H (2001) Taxonomic study on nemerteans from South West Islands, Japan. In "Toyoshio-maru Cruise Report No2001-6" Ed by A Go, Faculty of Applied Biological Science, Hiroshima University, Hiroshima, pp 10–11 [In Japanese]
- Kajihara H (2002) Two new species of *Zygonemertes* (Nemertea: Enopla: Monostilifera) from Hokkaido, Japan. *Spec Div* 7: 121–143
- Kajihara H (2004) Usamaro Takakura (1867–1944), Japanese pioneer nemertean researcher. *Arch Nat Hist* 31: 208–213
- Kajihara H (2006) Four palaeonemerteans (Nemertea: Anopla) from a tidal flat in middle Honshu, Japan. *Zootaxa* 1163: 1–47
- Kajihara H (2007a) Two species of *Nemertopsis* (Nemertea: Hoplonemertea: Monostilifera) living in association with *Capitulum mitella* (Crustacea: Cirripedia: Thoracica: Lepadomorpha). *Zootaxa* 1446: 43–58
- Kajihara H (2007b) *Ototyphlonemertes dolichobasis* sp. nov. (Nemertea: Hoplonemertea: Monostilifera: Ototyphlonemertidae), a new species of interstitial nemertean from Japan. *Spec Div* 12: 57–66
- Kajihara H, Gibson R, Mawatari SF (2000) Redescription and taxonomic reassessment of *Nemertellina minuta* Friedrich, 1935, *sensu* Yamaoka, 1940 (Nemertea, Hoplonemertea, Monostilifera). *Zool Sci* 17: 265–276
- Kajihara H, Gibson R, Mawatari SF (2001) A new genus and species of monostiliferous hoplonemertean (Nemertea: Enopla: Monostilifera) from Japan. *Hydrobiologia* 456: 187–198
- Kajihara H, Gibson R, Mawatari SF (2003) *Potamostoma shizunaiense* gen. et sp. nov. (Nemertea: Hoplonemertea: Monostilifera): a new brackish-water nemertean from Japan. *Zool Sci* 20: 491–500
- Kanda S (1939) The luminescence of a nemertean, *Emplectonema kandai* Kato. *Biol Bull* 77: 166–173
- Kato K (1939) A new luminous species of the nemertea, *Emplectonema kandai* sp. nov. *Jpn J Zool* 8: 251–254
- Kato K (1940) Oyogi-himomushi. *Saishū to Shiiku* 2: 101 [In Japanese]
- Kato K (1947) *Emplectonema kandai* Kato, *Nectonemertes mirabilis* Verrill, *Pelagonemertes moseleyi* Bürger. In "Revised Edition Illustrated Encyclopedia of the Fauna of Japan (Exclusive of Insects)" Ed by S Uchida, The Hokuryukan Co., Ltd., Tokyo, pp 1466–1470 [In Japanese]
- Kato K, Iwata F (1965) *Emplectonema kandai* Kato, *Pelagonemertes moseleyi* Bürger, *Nectonemertes mirabilis* Verrill. In "New Illustrated Encyclopedia of the Fauna of Japan Vol 1" Ed by Y Okada, S Uchida, T Uchida, Hokuryu-kan Publishing Co., Ltd., Tokyo, pp 396–401 [In Japanese]
- Kato K, Tanaka O (1938) Notes on *Pelagonemertes moseleyi* Bürger. *Jpn J Zool* 7: 595–598
- Kawai S, Yamaoka T (1940) On *Malacobdella japonica* Takakura. *Zool Mag* 52: 255–259 [In Japanese with English summary]
- Kawakatsu M (1991) Nemertinea. In "Pictorial Keys to Soil Animals of Japan" Ed by J Aoki, Tōkaidaiigaku Shuppan Kai, Tokyo, p 2 [In Japanese]
- Kawakatsu M (1999) Nemertinea. In "Pictorial Keys to Soil Animals of Japan" Ed by J Aoki, Tōkaidaiigaku Shuppan Kai, Tokyo, p 11 [In Japanese]
- Kawakatsu M, Nunomura N, Suzuki H (1989) Corrections and emendations of specific names of several Japanese species of turbellarians (Acoela, Tricladida-Maricola, Polycladida-Cotylea, and Tremocephalida-Scutariellida) and Nemertinea (Enopla-Hoplonemertea-Monostilifera). *Bull Fuji Women's Coll Ser II* 27: 41–51
- Kikuchi T (1968) Faunal list of the *Zostera marina* belt in Tomioka Bay, Amakusa, Kyushu. *Publ Amakusa Mar Biol Lab* 1: 163–192
- Kito K (1975) Preliminary report on the phytal animals in the *Sargassum confusum* region in Oshoro Bay, Hokkaido. *J Fac Sci Hokkaido Univ Ser 6 Zool* 20: 141–158
- Komai T (1919) Huyūsei-himomushi [Pelagic nemerteans]. *Zool Mag* 31: 294–296 [In Japanese]
- Korotkevitsch VS (1955) Pelagicheskie nemertiny dalnevostochnykh morei SSSR. Izd Akad Nauk SSSR 58: 1–131 [In Russian]
- Korotkevitsch VS (1971) K ekologii i sistematike nemertin saliva noseta yaponskogo morya. *Issl Faun Mor* 8: 109–122 [In Russian]
- Korotkevitsch VS (1977) K sistematike nemertin tribui Pelagica. Issl

- Faun Mor 20: 13–19 [In Russian]
- Korotkevitsch VS (1982) Novye i redkie vidy nemertin (Nemertini) s iuzhnogo shel'fa ostrova sakhalin. Issledv Faun Mor 29: 12–26 [In Russian]
- Kulikova VI (1988) Raspredelenie nemertin v agregatsiakh dvustvorchatykh molliuskov. Dokl Akad Nauk SSSR 302: 1013–1016 [In Russian]
- Kulikova VI, Kutishchev AA (1984) Nemertinui druz mitilid zaliva vostok yaponskogo morya. Izv Akad Nauk SSSR Ser Biol 4: 615–619
- Langerhans P (1880) Die Wurmfauna von Madeira III. Z Wiss Zool 34: 87–143
- Leuckart FS (1828) Breves animalium quorundam maxima ex parte marinorum descriptiones. Augusti Osswaldi, Heidelbergae
- Manchenko GP, Kulikova VI (1996a) Allozyme and colour differences between two sibling species of the heteronemertean *Lineus torquatus* from the Sea of Japan. Mar Biol 125: 687–691
- Manchenko GP, Kulikova VI (1996b) Enzyme and colour variation in the hoplonemertean *Tetrastemma nigrifrons* from the Sea of Japan. Hydrobiologia 337: 69–76
- Martindale MQ, Henry JQ (1995) Modifications of cell fate specification in equal-cleaving nemertean embryos: alternate patterns of spiralian development. Development 121: 3175–3185
- Maslakova SA (2005) Systematics and evolution of smiling worms family Prosorhochmidae (Hoplonemertea; Nemertea). Ph.D thesis, George Washington University, Washington DC, USA
- Maslakova SA, Martindale MQ, Norenburg JL (2004a) Fundamental properties of the spiralian developmental program are displayed by the basal nemertean *Carinoma tremaphoros* (Palaeonemertea, Nemertea). Dev Biol 267: 342–360
- Maslakova SA, Martindale MQ, Norenburg JL (2004b) Vestigial prototroch in a basal nemertean, *Carinoma tremaphoros* (Nemertea; Palaeonemertea). Evol Dev 6: 219–226
- McDermott JJ, Roe P (1985) Food, feeding behavior and feeding ecology of nemerteans. Amer Zool 25: 113–125
- McIntosh WC (1873–1874) A Monograph of the British Annelids Part I The Nemerteans. Ray Society, London [pp 1–96, pls I–X in 1873; pp 97–214, pls XI–XXIII in 1874]
- Melville RV (1986) *Tubulanus* Renier, [1804] and *T. polymorphus* Renier, [1804] (Polychaeta): proposed reinstatement under the plenary powers. Bull Zool Nom 43: 112–114
- Miyashita Y (1932) Kansatsu-zakkai. [Miscellaneous notes of observation]. Zool Mag 44: 327–331 [In Japanese]
- Miyazawa K, Higashiyama M, Ito K, Noguchi T, Arakawa O, Shida Y, Hashimoto K (1988) Tetrodotoxin in two species of ribbon worms (Nemertini), *Lineus fuscoviridis* and *Tubulanus punctatus*. Toxicon 26: 867–874
- Montgomery TH (1894) Kritische Uebersicht der Süßwasser-Nemertinen. Inaugural Dissertation, University of Berlin, Berlin
- Montgomery TH (1897) Descriptions of new metanemerteans, with notes on other species. Zool Jb Abt Syst 10: 1–14
- Moore J, Gibson R (1981) The Geonemertes problem (Nemertea). J Zool Lond 194: 175–201
- Moore J, Gibson R (1988) Marine relatives of terrestrial nemerteans: the genus *Prosadenoporus* Bürger, 1890 (Hoplonemertea). Hydrobiologia 156: 75–86
- Moretto HJA (1974) Un paleonemertino de Quequen. Neotropica 20: 9–13
- Moriyama H (1995) Nakamura Kazuhiko sensei Okazaki Yoshiro sensei Iwata Fumio sensei wo ookurisuru. Mem Kushiro Publ Univ Econom 7: 1–18 [In Japanese]
- Morton B (1989) Partnerships in the sea: Hong Kong's marine symbioses. Hong Kong University Press, Hong Kong
- Moseley HIN (1875a) On *Pelagonemertes rollestoni*. Ann Mag Nat Hist Ser 4 15: 165–169
- Moseley HIN (1875b) On a young specimen of *Pelagonemertes rollestoni*. Ann Mag Nat Hist Ser 4 16: 377–383
- Müller OF (1788) Zoologica Danica Seu Animalium Daniae et Norvegiae Rariorum ac Minus Notorum Descriptiones et Historia Vol 2. N Mölleri et Fillii, Havniae
- Nielsen C (2005) Trochophora larvae: cell-lineages, ciliary bands and body regions. 2. Other groups and general discussion. J Exp Zool 304b: 401–447
- Nishikawa T, Sattmann, H (2001) List of Dr. Albrecht von Roretz's collection of Japanese animals made about 120 years ago, compiled from the catalogues of Naturhistorisches Museum Wien. Bull Nagoya Univ Mus 17: 33–44
- Nishimura S (1992) Gaisetsu [Overview]. In "Guide to Seashore Animals of Japan with Color Pictures and Keys Vol I" Ed by S Nishimura, Hoikusha, Osaka, pp xi–xxxv [In Japanese]
- Noguchi T, Ali AE, Arakawa O, Miyazawa K, Kanoh S, Shida Y, Nishio S, Hashimoto K (1991) Tetrodonic acid-like substance; a possible precursor of tetrodotoxin. Toxicon 29: 845–855
- Norenburg JL, Roe P (1998) Reproductive biology of several species of recently collected pelagic nemerteans. Hydrobiologia 365: 73–91
- Ochi O (1979) Ultrastructures of the blood vessels and the haemocytes in a nemertean, *Lineus fuscoviridis*. Zool Mag 88: 640
- Ochsenheimer F (1816) Die Schmetterlinge von Europa Vol 4. Gerhard Fleischer, Leipzig
- Ohuye T (1942) On the blood corpuscles and the hemopoiesis of a nemertean, *Lineus fuscoviridis*, and of a sipunculus, *Dendrostoma minor*. Sci Rep Tōhoku Imp Univ 4th Ser Biol 17: 187–196
- Ohuye T (1950) Preliminary note on the survey of peroxidase distribution in the blood and coelomic cells of various invertebrates. Mem Ehime Univ Sect II Sci 1: 21–26
- Okada S, Igarashi T, Kobayashi K (1971) Invertebrates and fishes of Oshoro Bay and neighbouring Area. Bull Plankton Soc Jpn 18: 59–72
- Oki I, Tamura S, Ogren RE, Kitagawa K, Kawakatsu M (1987) The karyotype and a new locality for the land nemertine *Geonemertes palaensis* Semper, 1863. Bull Fuji Women's Coll Ser 2 25: 66–77
- Okuda S (1947) *Paranemertes peregrina* Coe, *Amphiporus formidabilis* Griffin, *Zygonemertes glandulosa* Yamaoka, *Prostoma nigrifrons* (Coe), *Lineus torquatus* Coe, *Cerebratulus marginatus* Renier, *Micrura bella* (Stimpson), *Micrura uchidai* Yamaoka, *Micrura akkeshiensis* Yamaoka, *Baseodiscus hemprichi* (Ehrenberg), *Tubulanus ezoensis* Yamaoka. In "Revised Edition Illustrated Encyclopedia of the Fauna of Japan (Exclusive of Insects)" Ed by S Uchida, The Hokuryukan Co., Ltd, Tokyo, pp 1467–1474 [In Japanese]
- Okuda S, Iwata F (1965) *Tubulanus ezoensis* Yamaoka, *Baseodiscus hemprichi* Ehrenberg, *Lineus torquatus* Coe, *Cerebratulus marginatus* Renier, *Micrura akkeshiensis* Yamaoka, *Micrura bella* Stimpson, *Micrura uchidai* Yamaoka, *Emplectonema mitsuii* Yamaoka, *Paranemertes katoi* Yamaoka, *Paranemertes peregrina* Coe, *Zygonemertes grandulosa* Yamaoka, *Amphiporus ogumai* Yamaoka, *Amphiporus cervicalis* (Stimpson), *Tetraستema roseocephalum* Yamaoka, *Tetraستema nigrifrons* Coe. In "New Illustrated Encyclopedia of the Fauna of Japan Vol 1" Ed by Y Okada, S Uchida, T Uchida, Hokuryukan Publishing Co., Ltd., Tokyo, pp 390–400 [In Japanese]
- Okugawa K (1932) An outline of the Japanese freshwater rhabdocoelids. Zool Mag 44: 69–70. [In Japanese]
- Ooishi S (1964) Results of Amami Expedition, no. 3, invertebrates. Rep Fac Fish Pref Univ Mie 5: 189–215
- Ørsted AS (1843) Forsog til en ny classification af Planarieerne (Planariae Duges) grundet paa mikroskopisk-anatomiske Undersøgelser. Naturhist Tidsskr 4: 519–581
- Ørsted AS (1844) Entwurf Einer Systematischen Eintheilung und Speciellen Beschreibung der Plattwürmer, auf Microscopische Untersuchungen. CA Reitzel, Copenhagen

- Punnett RC (1900) On some South Pacific nemertines collected by Dr Willey. Zool Resul New Brit New Guinea, Loyalty Is Coll 1895 1896 1897 Arthur Willey 5: 569–584
- Punnett RC and Cooper CF (1909) On some nemertines from the eastern Indian Ocean. Trans Linn Soc Lond Ser 2 Zool 13: 1–15
- Quatrefages A de (1846) Étude sur les types inférieurs de l'embranchement des annelés. Ann Sci Nat Sér 3 Zool 6: 173–303
- Quoy JRC, Gaimard JP (1833) Voyage de découvertes de l'Astrolabe exécuté par ordre du Roi, pendant les années 1826–1827–1828–1829, sous le commandement de M. J. Dumont d'Urville, Zoologie Vol 4. J Tastu, Paris, pp 284–292
- Rathke J (1799) Jagttagelser henhorende til Indvoldormenes og Bloddyrenes Naturhistorie. Skr naturh Selsk Khb 5: 61–148
- Renier SA (1804) Prospetto Della Classe dei Vermi, Nominati e Ordinati Secondo il Sistema di Bosc. pp xv–xxvii [A work commonly attributed to the year 1804, rejected for nomenclatural purposes by ICZN with Opinion 316 as not duly published within the meaning of Article 25 of the Régule Internationales de la Nomenclature Zoologique, thereby placed on the Official Index of Rejected and Invalid Works in Zoological Nomenclature (Work No. 25) in 1954; names of animals contained in this work were subsequently ruled to be unavailable by ICZN with Opinions 427 and 436, thereby placed on the Official Index of Rejected and Invalid Generic Names in Zoology and Official Index of Rejected and Invalid Specific Names in Zoology in 1956 and 1957, respectively; later, a certain number of these names were reinstated by the plenary power of ICZN, including the nemertean names *Cerebratulus marginatus* (Opinion 477) in 1957 and *Tubulanus polymorphus* (Opinion 1486) in 1988]
- Riser NW (1974) Nemertinea. In “Reproduction of Marine Invertebrates Vol 1” Ed by JS Pearse, AC Giese, Academic Press, New York, pp 359–389
- Riser NW (1989) Speciation and time — relationships of the nemertines to the acelomate metazoan Bilateria. Bull Mar Sci 45: 531–538
- Riser NW (1991) New Zealand nemertines from kelp holdfasts: Heteronemertea II. *Notospermus geniculatus* (Delle Chiaje, 1828) n. comb. N Z J Zool 18: 427–438
- Riser NW (1994) The morphology and generic relationships of some fissiparous heteronemertines. Proc Biol Soc Wash 107: 548–556
- Riser NW (1998) The morphology of *Micrura leidyi* (Verrill, 1892) with consequent systematic revaluation. Hydrobiologia 365: 149–156
- Roe P (1993) Aspects of the biology of *Pantinonemertes californiensis*, a high intertidal nemertean. Hydrobiologia 266: 29–44
- Saito M, Suzuki H (1974) Fauna in the littoral zone of Niisaki Beach in the western part of Sagami Bay I. Sci Rep Yokohama Natn Univ (Sect 2) 21: 31–48
- Satō H, Itō T (1961) Musekitsui Dōbutsu Saishū Shiiku Jikkenhō [Methods of Sampling, Raring, and Experiments on Invertebrates]. Hokuryūkan, Tokyo [In Japanese]
- Semper C (1863) Reisebericht. Briefliche Mittheilung an A. Kölliker. Z Wiss Zool 13: 558–570
- Senz W (1997a) Morphologie und klassifikatorische Position einiger anopler Nemertinen (Nemertini: Anopla). Ann Naturhist Mus Wien 99B: 423–496
- Senz W (1997b) Über die phylogenetische Herkunft und systematische Stellung der Gattung *Malacobdella* Blainville, 1827 (Nemertini, Bdellonemertini). Sitzungsber Math naturwiss Kl Abt I 204: 39–62
- Senz W (2001) Eine neue Heteronemertine von der Küste Japans (Nemertini). Spixiana 24: 5–12
- Shiino S (1969) Suisan-musekitsui-dōbutsugaku [Aquatic Invertebrate Zoology]. Baifūkan, Tokyo
- Shimomura M, Kato T, Kajihara H (2001) Records of some marine invertebrates (nemertines, asellotes and phylloocids) from the coast around Otsuchi Bay. Otsuchi Mar Sci 26: 46–50
- Sowerby J (1806) The British Miscellany: Or Coloured Figures of New, Rare, or Little Known Animal Subjects; Many Not Before Ascertained to Be Inhabitants of the British Isles. Taylor, London
- Steksova VV (2004) Invaziia dvustvorchatogo molliuska *Spisula sachalinensis* v zal. aniva o. sakhalin nemertinoi *Malacobdella japonica* (Takakura, 1897). Trudy Sakhin Nauch-Issled Inst Rybn Khoz Okean 6: 297–300 [In Russian]
- Stiasny-Wijnhoff G (1938) Das Genus *Prostoma* Dugès, eine Gattung von Süßwasser-nemertinen. Arch Néerland Zool 3: 219–230
- Stimpson W (1855) Descriptions of some of the new marine Invertebrata from the Chinese and Japanese seas. Proc Acad Nat Sci Philadelphia 7: 375–384
- Stimpson W (1857) Prodromus descriptionis animalium evertebratorum quae in Expeditione ad Oceanum Pacificum Septentrionalem, a Republica Federata missa, Dadwaladaro Ringgold et Johanne Rodgers Ducibus, observavit et descripsit. Pars II. Turbellarieorum Nemertineorum generum et specierum adhuc ineditarum descriptions; adjunctis notis de generibus jam constitutes. Proc Acad Nat Sci Philadelphia 9: 159–165
- Strand M, Sundberg P (2005a) Delimiting species in the hoplonemertean genus *Tetrastremma* (phylum Nemertea): morphology is not concordant with phylogeny as evidenced from mtDNA sequences. Biol J Linn Soc 86: 201–212
- Strand M, Sundberg P (2005b) Genus *Tetrastremma* Ehrenberg, 1831 (phylum Nemertea) — a natural group? Phylogenetic relationships inferred from partial 18S rRNA sequences. Mol Phylogenet Evol 37: 144–152
- Stricker SA, Folsom MW (1998) A comparative ultrastructural analysis of spermatogenesis in nemertean worms. Hydrobiologia 365: 55–72
- Sudzuki M (1953) On Japanese fresh-water nemertines. Zool Mag 62: 212–219 [In Japanese with English summary]
- Sun S (1995) The preliminary report of nemertines from Taiwan Channel. Mar Sci 5: 45–48
- Sun S (2001) A new mangrove-dwelling nemertean from China. Hydrobiologia 456: 199–209
- Sun S, Pan H (1994) Nemertinea. In “Marine Species and Their Distributions in China’s Seas” Ed by Z Huang, China Ocean Press, Beijing, pp 328–330 [In Chinese]
- Sundberg P (1984) Multivariate analysis of polymorphism in the hoplonemertean *Oerstedia dorsalis* (Abildgaard, 1806). J Exp Mar Biol Ecol 78: 1–22
- Sundberg P (1988) A new monostiliferous hoplonemertean (Nemertea), *Oerstedia striata* sp. n., from the west coast of Sweden. Zool Scr 17: 135–139
- Sundberg P, Andersson S (1995) Random amplified polymorphic DNA (RAPD) and intraspecific variation in *Oerstedia dorsalis* (Hoplonemertea, Nemertea). J Mar Biol Ass UK 75: 483–490
- Sundberg P, Gibson R (1995) The nemertines (Nemertea) of Rottnest Island, Western Australia. Zool Scr 24: 101–141
- Sundberg P, Gibson R, Olsson U (2003) Phylogenetic analysis of a group of palaeonemertians (Nemertea) including two new species from Queensland and the Great Barrier Reef, Australia. Zool Scr 32: 279–296
- Sundberg P, Hylbom R (1994) Phylogeny of the nemertean subclass Palaeonemertea (Anopla, Nemertea). Cladistics 10: 347–402
- Sundberg P, Janson K (1988) Polymorphism in *Oerstedia dorsalis* (Abildgaard, 1806) revised — electrophoretic evidence for a species complex. Hydrobiologia 156: 93–98
- Takakura U (1897) On a new species of *Malacobdella* (*M. japonica*). Ann Zool Jpn 1: 105–112

- Takakura U (1898) Misaki-kinbōsan-himomushirui-(Nemertine)-nobunrui [A classification of the nemerteans of the Misaki region]. Zool Mag 10: 38–44, 116–120, 184–187, 331–337, 424–429 [In Japanese]
- Takakura U (1910) Kisei-himomushi-no-ichi-shinshu [A new species of parasitic nemertean]. Zool Mag 22: 111–116 [In Japanese]
- Takakura U (1922) Honpōsan-himomushi-no-ichi-shinzoku-ni-tsukite [On a new genus of nemertean from Japan]. Zool Mag 34: 419–422 [In Japanese]
- Takakura U (1933) Kitachishima-no-himomushi-rui [Nemerteans of the Northern Kurile Islands]. Bull Biogeogr Soc Jpn 4: 226–227 [In Japanese]
- Tanu MB, Mahmud Y, Arakawa O, Takatani T, Kajihara H, Kawatsu K, Hamano Y, Asakawa M, Miyazawa K, Noguchi T (2004) Immunoenzymatic visualization of tetrodotoxin (TTX) in *Cephalothrix* species (Nemertea: Anopla: Palaeonemertea: Cephalotrichidae) and *Planocera reticulata* (Platyhelminthes: Turbellaria: Polycladida: Planoceridae). Toxicon 44: 515–520
- Thollesson M, Norenburg JL (2003) Ribbon worm relationships: a phylogeny of the phylum Nemertea. Proc R Soc Lond B 270: 407–414
- Tizard TH, Moseley HN, Buchanan JY, Murray J (1885) Japan. Rep Sci Challenger Exped, Narrative 1: 744–750
- Tsuchiya M (1979) Quantitative survey of intertidal organisms on rocky shores in Mutsu Bay, with special reference to the influence of wave action. Bull Mar Biol St Asamushi Tohoku Univ 16: 69–86
- Turbeville JM (1986) An ultrastructural analysis of coelomogenesis in the hoplonemertine *Prosorhochmus americanus* and the polychaete *Magelona* sp. J Morph 187: 51–60
- Turbeville JM (1991) Nemertinea. In "Microscopic Anatomy of Invertebrates Vol 3 Platyhelminthes and Nemertinea" Ed by FW Harrison, BJ Bogitsh, Wiley-Liss, Inc., New York, pp 285–328
- Turbeville JM (2002) Progress in nemertean biology: development and phylogeny. Integ Comp Biol 42: 692–703
- Turbeville JM, Ruppert EE (1985) Comparative ultrastructure and the evolution of nemertines. Amer Zool 25: 53–71
- Uchida H (1994) Nagamushirui [Vermes]. In "Yama-Kei Field Books Vol 8 Umibenoikimono" [Sea Shore Animals] Ed by T Okutani, Yamatokeikokusha, Tokyo, pp 85–90 [In Japanese]
- Uchida T, Iwata F, Nagao Z (1972) Shakotan-hantō-kaichū-kōen-kōho-chi-no-musekutsui-dōbutsu [Invertebrates in a site proposed for Shakotan Marine Park]. In "Hokkaidō-shakotan-hantō-otaru-kaigan-kaichū-kōen-chōsa-sōkokusho" [Report on Hokkaido Shakotan Peninsula and Otaru Beach Marine Park] Ed by Hokkaido Wildlife Conservation Section, Hokkaido, pp 45–66
- Uchida T, Yamada M, Iwata F, Oguro C, Nagao Z (1963) The zoological environs of the Akkeshi Marine Biological Station. Publ Akkeshi Mar Biol St 13: 1–36
- Utinomi H (1956) Coloured Illustrations of Sea Shore Animals of Japan. 1st ed, Hoikusha, Osaka
- Utinomi H (1960) Coloured Illustrations of Sea Shore Animals of Japan. 1st ed, 4th copy, Hoikusha, Osaka
- Utinomi H (1969) Coloured Illustrations of Sea Shore Animals of Japan. 2nd ed, Hoikusha, Osaka
- Vaillant L (1890) Histoire Naturelle des Annelés Marins et d'Eau Douce. Lombriciens, Hirudiniens, Bdellomorphes, Térétulaires et Planariens. Librairie Encyclopédique de Roret, Paris
- Verrill AE (1895) Supplement to the marine nemerteans and planarians of New England. Trans Connecticut Acad Art Sci 9: 523–534
- Walker G (1994) Preliminary observations on the association of the nemertean, *Nemertopsis quadripunctatus*, with the stalked barnacle, *Capitulum mitella*, on an exposed Hong Kong shore. Asian Mar Biol 11: 125–136
- Wheeler JFG (1934) Nemerteans from the South Atlantic and southern oceans. Discovery Rep 9: 215–294
- Wheeler JFG (1940) Some nemerteans from South Africa and a note on *Lineus corrugatus* M'Intosh. J Linn Soc Lond Zool 41: 20–49
- Wijnhoff G (1913) Die Gattung *Cephalothrix* und ihre Bedeutung für die Systematik der Nemertinen. II. Systematischer Teil. Zool Jb Abt Syst Ökol Geogr Tiere 34: 291–320
- Yamaguchi E, Yamada M (1955) Kitanihon-kaisan-musekutsuidōbutsu [Marine Invertebrates in Northern Japan]. Hokkai-kyōiku-hyōron-sha, Sapporo
- Yamaoka T (1939) Two nemerteans from Formosa. Ann Zool Jpn 18: 283–289
- Yamaoka T (1940a) The fauna of Akkeshi Bay. IX. Nemertini. J Fac Sci Hokkaido Imp Univ Ser 6 Zool 7: 205–263
- Yamaoka T (1940b) Two nemerteans from the Riukiu Islands. Ann Zool Jpn 19: 13–18
- Yamaoka T (1943) Nemertini. In "Keitō-dōbutsugaku" [Systematic Zoology] Ed by H Ohshima, Y Okada, Yōkenrō, Tokyo, pp 773–815 [In Japanese]
- Yamaoka T (1947) *Emplectoneema mitsuii* Yamaoka, *Paranemertes katoi* Yamaoka, *Amphiporus ogumai* Yamaoka, *Prostoma roseocephalum* Yamaoka, as submitted by S Okuda. In "Revised Edition Illustrated Encyclopedia of the Fauna of Japan (Exclusive of Insects)" Ed by S Uchida, The Hokuryukan Co. Ltd., Tokyo, pp 1466–1469 [In Japanese]
- Yamaoka T (2005) On fourteen monostiliferous hoplonemerteans from the Izu Peninsula, Middle Japan. Publ Seto Mar Biol Lab 40: 141–158 [A posthumous publication submitted by Prof. Fumio Iwata]
- Yin Z, Shi J, Li N (1988) A preliminary study of the intertidal nemerteans along the coast of Shandong Province. Mar Sci Bull 1: 68–74

(Received September 1, 2006 / Accepted November 8, 2006)