

**DIRECTIONS FOR USE**

**Restrictions:** DO NOT apply granules when the turf foliage is wet.  
DO NOT use on putting greens, low cut tees or bowling greens.  
DO NOT apply to newly seeded turf areas.

Situation	Target	Turf Species	Rate	Critical Comments
Warm season turfgrass areas, including fairways, golf tees, municipal sports fields, industrial and commercial lawns	Pre-emergent control of Summer Grass ( <i>Digitaria ciliaris</i> ), Crowsfoot (Crabgrass) ( <i>Eleusine indica</i> )	Kikuyu ( <i>Pennisetum clandestinum</i> ), Buffalo Grass ( <i>Stenotaphrum secundatum</i> ), Common Couch	300 kg/ha (3 kg/100m <sup>2</sup> )	This product is ready to apply. No mixing or dilution is required. Apply product evenly to dry foliage and bare areas using a granular spreader able to give even coverage. Application must be made before weeds have begun to germinate. Established weeds will not be controlled. After application water in with at least 10mm of sprinkler irrigation. Control of germinating weed seedlings can be expected for up to 10 weeks after application depending upon temperature, rainfall, soil type and other factors. If longer control is required, a repeat application should be made 7-10 weeks after initial application. DO NOT mow treated area until product has been watered in and turf foliage has dried. Avoid application when daytime temperatures are consistently above 32°C. This product is not recommended for use on putting greens. DO NOT apply on turf containing Carpet Grass ( <i>Axonopus compressus</i> ) and Dichondra unless some damage is acceptable.
	Pre-emergent control of Winter Grass ( <i>Poa annua</i> ), Creeping Oxalis ( <i>Oxalis corniculata</i> )	( <i>Cynodon dactylon</i> , <i>C. transvaalensis</i> ), Grass (not Santa Anna), Tiff Hybrid Couch Grass, Saltwater Couch ( <i>Paspalum vaginatum</i> ), Queensland Blue Couch ( <i>Digitaria didactyla</i> ), Bahia Grass ( <i>Paspalum notatum</i> ), Zoysia Grass ( <i>Zoysia matrella</i> )	400 kg/ha (4 kg/100m <sup>2</sup> )	
	Turf nutrition – provides slow release nitrogen for 8 weeks feeding		300-400 kg/ha (3-4 kg/100m <sup>2</sup> )	

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**GENERAL INSTRUCTIONS**

Ox Herbicide Plus Fertiliser 18-10-9 is a convenient ready to use formulation for the control of certain annual grass weeds in turf and also for fertilisation of turf where nitrogen, potassium and sulphur are required. The product should only be used when both functions are required.

**APPLICATION**

For a granular spreader which can be calibrated for even output and capable of giving even application. Use a double application technique: i.e. half rate width, to achieve a full rate application. Apply under calm conditions, thereby reducing drift to non-target areas and optimizing particle distribution for good weed control. Ensure turf area is free of leaves and other debris, mowing if necessary so that the granules reach the soil surface. Delay reseeding of Ryegrass or Bentgrass into treated areas for 4-5 months (Ryegrass 4 months & Bentgrass 5 months). DO NOT over apply, especially under the branch spread of trees and shrubs. Avoid contact with flowers and shrubs.

Suggested Spreader Settings				
The suggested spreader settings are not intended to replace calibration. Calibrate your spreader before applying product.			SPREADER SETTINGS	
Spreader	Ground Speed	Width of Coverage	300 kg/ha	400 kg/ha
AccuPro, SR-2000 (cone 4)	4.8 km/hr	2.7m	0 1/2	Q
Model 2000 (cone 4)	4.8 km/hr	2.7m	0 1/2	Q
Scotts R8A (cone 4)	4.8 km/hr	2.7m	N 1/2	P 1/2
Lesco	4.8 km/hr	2.1m	17	19
Spyker Spreader	4.8 km/hr	2.1m	4	4 1/2
Vicon (03 Series- Standard Spout)	7.3 km/hr	8.5m	37	44

**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**OXaMAX**

**Ox Herbicide *plus* Fertiliser 18-10-9**

**ACTIVE CONSTITUENT: 10 g/kg OXADIAZON**  
**SOLVENT: 10.5 g/kg POLYBUTENE**

**GROUP G HERBICIDE**

Combination Herbicide and Controlled Release Fertiliser for pre-emergent control of Summer Grass, Crabgrass and Winter Grass in turf and feeding of turf with slow release nitrogen with phosphorus and potassium.

**GUARANTEED MINIMUM ANALYSIS**

- 18.00% Total Nitrogen (N)**  
9.39% Ammoniacal Nitrogen  
8.61% Urea Nitrogen
- 10.40% Phosphorus (P) from ammonium phosphate**
- 9.90% Potassium (K) from potassium chloride**
- 0.80% Total Sulphur (S) as elemental**
- 10.47% Chlorine (Cl) Maximum**

\*8.61% Slowly Available Nitrogen from Polymer Sulphur Coated Urea

Nutrients derived from: ammonium phosphate, polymer sulphur coated urea, potassium chloride.

NET CONTENTS  
**20KG**  
**WARNING**

**RESISTANCE WEEDS WARNING**

Ox Herbicide plus Fertiliser 18-10-9 is a member of the Oxadiazole Group of herbicides. The herbicide is an inhibitor of protoporphyrinogen oxidase. For weed resistance management, the product is a Group G Herbicide. Some naturally occurring weed biotypes resistant to the herbicide may exist through normal genetic variability in weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other oxadiazole group herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Amgrow Australia Pty Ltd does not accept liability for any losses that may result from failure of this product to control resistant weeds.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT apply product if rapid storm run-off from treated areas into water bodies is likely to occur within several days of application.

**STORAGE & DISPOSAL**

Store in the closed original container, in a dry, cool, well-ventilated area out of direct sunlight. DO NOT store with seed, feed, foodstuffs, insecticides, fungicides and other fertilisers. Shake bag empty into applicator. Puncture and shred empty container and bury in a local authority landfill. If no landfill is available bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. DO NOT dispose of damaged material on-site. DO NOT re-use this container for any purpose.

**SPILLS**

Sweep up spilled materials and return to original container. Wash contaminated equipment taking care the washings. DO NOT enter sewers, drains and other waterways.

**SAFETY DIRECTIONS**

Avoid contact with eyes. DO NOT inhale dust. Wash hands after use.

**FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. Give a glass of water.

**SAFETY DATA SHEET**

Additional information is listed on the Safety Data Sheet

APVMA Approval No.: 69594/61193

**MADE IN USA**

Additional information required under the Globally Harmonised System (GHS) classification of the substance/mixture: **Very toxic to aquatic life with long lasting effects.** Do not get in eyes, on skin, or on clothing. Wash contacted areas thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. Wear protective gloves, protective clothing and eye or face protection. Take off contaminated clothing and wash before reuse. IF SWALLOWED: Immediately call a POISON CENTRE or doctor. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. IF ON SKIN (or hair): Remove immediately all contaminated clothing. Rinse skin with water. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. If experiencing respiratory symptoms: Call a POISON CENTRE or doctor. Collect spillage. In case of fire, use carbon dioxide, dry chemical, foam, water fog. Alcohol resistant foam is the preferred firefighting medium but, if it is not available, normal foam can be used.

