



**MARJORIE HARRIS CARR  
CROSS FLORIDA GREENWAY  
STATE RECREATION AND  
CONSERVATION AREA**

**UNIT MANAGEMENT PLAN  
(2017-2027)**

State of Florida  
Department of Environmental Protection  
Division of Recreation and Parks





## **Florida Park Service**

The Florida Park Service is one of the largest park systems in the country with a total of 163 state parks and 11 state trails spanning nearly 800,000 acres, 100 miles of sandy white beach, and more than 1,500 miles of multi-use trails. From swimming and diving in Florida's rivers and springs to birding and fishing or hiking and riding on natural scenic trails, Florida's state parks offer year-round outdoor activities for all ages. The Florida Park Service is the proud recipient of three National Gold Medals for Excellence in Park and Recreation Management, making Florida America's first three-time Gold Medal winner. Offering opportunities throughout the state, Florida State Parks welcomed more than 31.8 million visitors in 2015-2016.

### **Mission Statement**

To provide resource-based recreation while preserving, interpreting, and restoring natural and cultural resources.

### **Vision**

The Florida State Park System creates a sense of place and is recognized as containing the best of Florida's diverse natural and cultural heritage sustained for future generations while providing quality and appropriate resource-based recreational opportunities, interpretation, and education that help visitors connect to "The Real Florida."





# TABLE OF CONTENTS

<b>INTRODUCTION AND GENERAL INFORMATION .....</b>	<b>1</b>
Background .....	1
<b>PURPOSE AND SIGNIFICANCE OF THE PARK .....</b>	<b>9</b>
Park Significance.....	9
<b>PURPOSE, SCOPE, AND REVIEW OF THE PLAN .....</b>	<b>10</b>
<b>MANAGEMENT PROGRAM REVIEW.....</b>	<b>12</b>
Management Authority and Responsibility.....	12
Park Management Goals .....	13
Management Coordination.....	14
Public Participation .....	15
Preliminary Public Workshops.....	15
Public Hearings and Advisory Group Meeting.....	16
Other Designations.....	16
<b>RESOURCE MANAGEMENT COMPONENT .....</b>	<b>18</b>
Introduction.....	18
Resource Descriptions and Assessment.....	19
Natural Resources .....	19
Cultural Resources .....	123
Resource Management Program.....	141
Management Goals, Objectives, and Actions .....	141
Natural Resource Management.....	141
Special Management Considerations.....	147
Cultural Resources Management .....	152
Resource Management Schedule.....	154
Land Management Review.....	154
<b>LAND USE COMPONENT.....</b>	<b>155</b>
Introduction.....	155
CFG Acquisition History .....	156
Subleases, Agreements, and Easements.....	157
External Conditions .....	159
Past Uses.....	162
Future Land Use and Zoning.....	162

Current Recreation Use and Visitor Programs .....	162
Property Analysis by County .....	164
Citrus County.....	164
Levy County.....	175
Marion County.....	184
Putnam County .....	203
Conceptual Land Use Plan.....	214
Potential Uses.....	214
Proposed Facilities.....	219
Facilities Development.....	220
Other Potential Future Trail/Facilities Developments .....	220
Existing Use and Recreational Carrying Capacity .....	226
Optimum Boundary .....	228
<b>IMPLEMENTATION COMPONENT.....</b>	<b>237</b>
Management Progress.....	237
Park Administration and Operations .....	237
Resource Management.....	237
Recreational Facilities and Visitor Services.....	238
Management Plan Implementation.....	239

## List of Tables

Table 1. CFG Western Management Zones, Acreage, and Fire Type .....	19
Table 2. CFG Central Management Zones, Acreage, and Fire Type.....	21
Table 3. CFG Eastern Management Zones, Acreage, and Fire Type.....	25
Table 4. Flooding Frequency Summary for Soils on the CFG.....	33
Table 5. Hydric Soils Summary for Soils on the CFG.....	33
Table 6. Soil Drainage Class Summary for Soils on the CFG.....	33
Table 7. Soil Series Class Summary for the Commonly Occurring Soils on the CFG .....	34
Table 8. CFG Community Type, Acreage, and Management.....	39
Table 9. Imperiled Species on the CFG.....	87
Table 10. Nesting Data on the CFG Dredge Spoil Islands .....	94
Table 11. CFG Large Management Zones and Degree of Infestation.....	103
Table 12. Inventory of FLEPPC Category I and II Invasive Exotic Plant Species.....	104
Table 13. Cultural Sites Listed in the Florida Master Site File .....	126
Table 14. Prescribed Fire Management .....	143
Table 15. CFG Prescribed Fire History.....	144
Table 16. Subleases on the CFG .....	157
Table 17. Conservation Lands and Activities within 15 Miles of the CFG.....	160
Table 18. CFG Visitor Estimates (FY 09 to FY 17).....	163
Table 19. Historic, Current and Projected Population through 2045—Citrus County.....	165
Table 20. Population Change—Citrus County .....	165
Table 21. Population by Age—Citrus County .....	166
Table 22. Trails and Mileage on the CFG—Citrus County.....	168
Table 23. Natural Communities and Acreage on the CFG—Citrus County .....	168
Table 24. Historic, Current and Projected Population through 2045—Levy County.....	176
Table 25. Population Change—Levy County.....	176
Table 26. Population by Age—Levy County.....	176
Table 27. Trails and Mileage on the CFG—Levy County .....	179
Table 28. Natural Communities and Acreage on the CFG—Levy County.....	179
Table 29. Historic, Current, and Projected Population through 2045—Marion County .....	185
Table 30. Population Change—Marion County.....	185
Table 31. Population by Age—Marion County.....	186
Table 32. Roadway Projects—Marion County.....	187
Table 33. Trails and Mileage on CFG—Marion County.....	188
Table 34. Natural Communities and Acreage on CGF—Marion County .....	190
Table 35. Historic, Current, and Projected Population through 2045—Putnam County .....	204
Table 36. Population Change—Putnam County.....	204
Table 37. Population by Age—Putnam County.....	205
Table 38. Trails and Mileage on CFG—Putnam County.....	207
Table 39. Natural Communities and Acreage on CFG—Putnam County .....	207
Table 40. Recreational Carrying Capacity .....	227
Table 41. Marjorie Harris Carr Cross Florida Greenway State Recreation and Conservation Area 10-Year Implementation Schedule and Cost Estimates Sheet .....	240

## List of Figures

Figure 1. Cross Florida Greenway Vicinity Map 1—West Section .....	3
Figure 2. Cross Florida Greenway Vicinity Map 2—Central Section.....	4
Figure 3. Cross Florida Greenway Vicinity Map 3—East Section .....	5
Figure 4. Cross Florida Greenway Reference Map 1—West Section.....	6
Figure 5. Cross Florida Greenway Reference Map 2—Central Section .....	7
Figure 6. Cross Florida Greenway Reference Map 3—East Section .....	8
Figure 7. Citrus County Generalized Existing Land Use .....	173
Figure 8. Citrus County Generalized Future Land Use .....	174
Figure 9. Levy County Generalized Existing Land Use .....	182
Figure 10. Levy County Generalized Future Land Use .....	183
Figure 11. Marion County Generalized Existing Land Use, Page 1.....	199
Figure 12. Marion County Generalized Existing Land Use, Page 2.....	200
Figure 13. Marion County Generalized Future Land Use, Page 1 .....	201
Figure 14. Marion County Generalized Future Land Use, Page 2 .....	202
Figure 15. Putnam County Generalized Existing Land Use .....	212
Figure 16. Putnam County Generalized Future Land Use.....	213
Figure 17. Cross Florida Greenway Conceptual Land Use Map—West Section .....	223
Figure 18. Cross Florida Greenway Conceptual Land Use Map—Central Section.....	224
Figure 19. Cross Florida Greenway Conceptual Land Use Map—East Section.....	225
Figure 20. Cross Florida Greenway Optimum Boundary Map, Page 1 of 7 .....	230
Figure 21. Cross Florida Greenway Optimum Boundary Map, Page 2 of 7 .....	231
Figure 22. Cross Florida Greenway Optimum Boundary Map, Page 3 of 7 .....	232
Figure 23. Cross Florida Greenway Optimum Boundary Map, Page 4 of 7 .....	233
Figure 24. Cross Florida Greenway Optimum Boundary Map, Page 5 of 7 .....	234
Figure 25. Cross Florida Greenway Optimum Boundary Map, Page 6 of 7 .....	235
Figure 26. Cross Florida Greenway Optimum Boundary Map, Page 7 of 7 .....	236

## **List of Addenda**

Addendum 1. Acquisition History

Addendum 2. References

Addendum 3. Preliminary Public Workshop Summary

Addendum 4. Timber Management Analysis

Addendum 5. Advisory Group Members and Summary Report

Addendum 6. Manatee Management

Addendum 7. Land Management Review



1                   **INTRODUCTION AND**  
2                   **GENERAL INFORMATION**

3                   **BACKGROUND**

4                   The Marjorie Harris Carr Cross Florida Greenway (CFG)  
5                   State Recreation and Conservation Area spans 110 miles  
6                   from Yankeetown on Florida’s west coast to south of  
7                   Palatka on the St. Johns River, near the east coast of Florida.  
8                   The CFG ranges from 300 yards wide to one mile wide  
9                   and includes portions of four counties in the upper Florida  
10                  peninsula: Citrus, Levy, Marion and Putnam. Over the 110  
11                  miles, the CFG traverses numerous natural areas,  
12                  physiographic areas, and human features. Many access  
13                  points provide residents and visitors with diverse  
14                  recreation opportunities. Nearby communities include  
15                  Yankeetown, Inglis, Dunnellon, Ocala, Belleview, Silver  
16                  Springs, Interlachen, and Palatka. The major highways  
17                  along the CFG include United States Highway (US) 19/98,  
18                  US 41, State Road (SR) 200, US 27/301/441, SR 40, and  
19                  SR 19. In addition, the CFG crosses Interstate 75 (I-75) just  
20                  south of Ocala, via the first land bridge constructed over a  
21                  major interstate roadway in the United States. This land  
22                  bridge provides both pedestrian and wildlife crossing  
23                  opportunities.

24                 Vicinity maps of the three sections of the CFG are  
25                 presented below (see Figures 1 through 3), followed by  
26                 reference maps of the three sections (see Figures 4  
27                 through 6).

28                 The CFG has an unusual land acquisition history  
29                 (Addendum 1). The majority of CFG lands were acquired  
30                 for the purpose of constructing and operating a  
31                 commercial shipping channel across the state. There  
32                 were two major efforts to construct a canal across the  
33                 Florida peninsula along this corridor alignment: the Gulf-  
34                 Atlantic Ship Canal (1933–1935), and the Cross Florida  
35                 Barge Canal (1964–1990). Thousands of acres of land  
36                 were acquired and millions of dollars were spent  
37                 designing, engineering, and building various segments of  
38                 the project until it was halted by Presidential order due,  
39                 primarily, to environmental concerns. In 1990, the Cross



*State Archives of Florida, Florida Memory*

Construction photo of the Cross Florida Barge Canal construction in the 1960s



1 Florida Barge Canal project was de-authorized by  
2 Congress and all lands and structures were transferred to  
3 the State of Florida for management as a Public  
4 Recreation and Conservation Area. In 1991, the  
5 Governor and Cabinet of the State of Florida signed a  
6 resolution agreeing to the terms of the federal de-  
7 authorization bill, thereby officially de-authorizing the  
8 Cross Florida Barge Canal project.

9 Ultimately, this action led to the creation of the Cross  
10 Florida Greenway State Recreation and Conservation  
11 Area. The Cross Florida Greenway was officially renamed  
12 the Marjorie Harris Carr Cross Florida Greenway in honor  
13 of Marjorie Harris Carr, who led the fight to stop the  
14 Cross Florida Barge Canal project.

15 The Florida Department of Environmental Protection  
16 (DEP), Division of Recreation and Parks (DRP), manages  
17 these lands under Lease #4013, dated October 27, 1993.  
18 The initial lease for the former canal lands has been  
19 amended 20 times through 2006 to include additional  
20 lands, mostly acquired with Preservation 2000 and  
21 Florida Forever funds. The lease extends 50 years,  
22 expiring on October 26, 2043 (Addendum 1). As indicated  
23 above, the CFG begins on the west coast of Florida at the  
24 Gulf of Mexico within Section 036, Township 16 South and  
25 Range 17 East within Citrus County, Florida, extending to  
26 Section 38, Township 11 South, Range 26 East in Putnam  
27 County, Florida.

28 The CFG is designated multiple-use in accordance with  
29 253.034(2)a, Florida Statute (F.S.), to provide public  
30 outdoor recreation and other park-related uses. There are  
31 no legislative or executive directives that constrain the use  
32 of this property (see Addendum 2). DRP intends to manage  
33 the CFG as such. Other than proposed facilities described  
34 in this plan, no other user-oriented facilities are  
35 anticipated to be developed and maintained by local  
36 governments.





Figure 1. Cross Florida Greenway Vicinity Map 1—West Section





Figure 2. Cross Florida Greenway Vicinity Map 2—Central Section






Figure 3. Cross Florida Greenway Vicinity Map 3—East Section






Figure 4. Cross Florida Greenway Reference Map 1—West Section






Figure 5. Cross Florida Greenway Reference Map 2—Central Section







Figure 6. Cross Florida Greenway Reference Map 3—East Section





# PURPOSE AND SIGNIFICANCE OF THE PARK

## *Marjorie Harris Carr Cross Florida Greenway State Recreation and Conservation Area*

The CFG was acquired to provide exceptional resource-based public outdoor recreation opportunities and cultural resource-based opportunities to Florida residents and visitors while ensuring the conservation and protection of valuable natural resources, including species and unique systems.

### **PARK SIGNIFICANCE**

The 110-mile long CFG offers outstanding opportunities for a variety of land and water-based activities. The CFG protects exceptional natural areas and provides important strategic ecological connectors for significant north-south and east-west ecological networks.

Assisting with this ecological connection is the CFG land bridge—America’s first land bridge—which spans I-75 just south of Ocala. The land bridge is one of the first multi-use bridges of its kind, landscaped with native vegetation, including longleaf pine, live oak, and sand live oak.

The CFG provides a cross-sectional view of the natural areas of the northern portion of the Florida peninsula. Two of Florida’s top first magnitude springs, Rainbow Springs and Silver Springs, feed into the CFG riverine ecosystems. The CFG offers the potential to serve as a critical ecological connector for the Ocala National Forest/Osceola National Forest/Pinhook Swamp/Okefenokee Swamp complex on its east end and Chassahowitzka/Crystal River/Lower Suwanee/Big Bend Florida Fish and Wildlife Conservation Commission (FWC) Wildlife Management Area (WMA) complex on its west end.

Further, the FWC indicates that almost 22,000 acres of the CFG are identified as Strategic Habitat Conservation Areas (SHCAs). SHCAs are essential to the enhancement of the long-term protection of many plants, animals, and natural communities that constitute essential components of Florida’s natural diversity. Within the CFG, the FWC does identify much of the Ocklawaha River area, Marshall Swamp/Adams Marsh, and the Ross

- The Cross Florida Greenway contains America’s first land bridge—which spans I-75 just south of Ocala, Florida. The land bridge is one of the first multi-use bridges of its kind, landscaped with native vegetation, including longleaf pine, live oak, and sand live oak.)
- Approximately 71,000 acres under management
- 254 cultural resource sites
- 290+ miles of trails (paved and unpaved)
- 42 miles of Florida National Scenic Trail (FNST) on the CFG


1 Prairie area as biodiversity hot spots for seven or more  
2 focal species.

3 Culturally, the CFG is significant given its history of  
4 ship/barge canal activities and the unique opportunities it  
5 provides to experience the largest de-authorization of a  
6 federal civil works project in the United States. Given its  
7 original purpose, CFG lands were acquired through a  
8 variety of programs. While most of the lands are former  
9 ship and barge canal lands, a significant amount of  
10 property was acquired with Preservation 2000 and  
11 Florida Forever funds. A portion of the CFG was acquired  
12 with Greenways and Trails Florida Forever funds and other  
13 parcels as part of Etoniah-Cross Florida Greenway, Longleaf  
14 Pine Ecosystem and Crystal River-area projects. Some  
15 lands were donated, and some are released from the Felburn  
16 Foundation. CFG also manages some St. Johns River Water  
17 Management District (SJRWMD) and Southwest Florida  
18 Water Management District (SWFWMD) land through  
19 various management agreements.

20 The CFG is classified as a state recreation area in the DRP's  
21 unit classification system. In the management of a state  
22 recreation area, a major emphasis is placed on maximizing  
23 the recreational potential of the unit. However, preservation  
24 of the park's natural and cultural resources remains  
25 important. Depletion of a resource by any recreational  
26 activity is not permitted. To realize the park's recreational  
27 potential, the development of appropriate park facilities is  
28 undertaken with the goal to provide amenities that are  
29 accessible, convenient, and safe; and to support public  
30 recreational use or appreciation of the park's natural,  
31 cultural, aesthetic, and educational attributes.

## 32 **PURPOSE, SCOPE, AND** 33 **REVIEW OF THE PLAN**

34 This Unit Management Plan (UMP) serves as the basic  
35 statement of policy and direction for the management of the  
36 CFG as a unit of Florida's state park system from 2017–2027.  
37 It provides information about past usage, conservation  
38 acquisition history, and descriptions of the natural and  
39 cultural resources found on the CFG. Furthermore, this UMP  
40 identifies the goals, objectives, actions, and criteria or  
41 standards that guide each aspect of park administration,  
42 and sets forth the specific measures that will be



## Components of the Unit Management Plan

### ***The Resource Management***

**Component** (RMC) provides a detailed inventory and assessment of the natural and cultural resources of the park. Resource management permits and issues are identified and measurable management objectives are established for each of the park's management goals and resource types. THE RMC provides guidance on the applications of such measures as prescribed burning and exotic species removal, imperiled species management, cultural resource management, and restoration of natural conditions.

***The Land Use Component*** is the recreational resource allocation plan for the park. Based on considerations such as access, population, adjacent land uses, the natural and cultural resources of the park, current public uses, and existing development, measurable objectives are set to achieve the ideal allocation of the physical space of the park. These objectives identify use areas and propose the types of facilities and programs, as well as the volume of public use to be provided.

***The Implementation Component*** consolidates the measurable objectives and actions for each of the park's management goals. An implementation schedule and cost estimates are included for each objective and action.

1 implemented to meet management objectives and provide  
2 balanced public utilization. The plan is intended to meet the  
3 requirements of Sections 253.034 and 259.032, F.S., Chapter  
4 18-2, Florida Administrative Code (FAC), and is intended to  
5 be consistent with the State Lands Management Plan. With  
6 approval, this management plan will replace the 2007  
7 approved plan. The UMP consists of three interrelated  
8 components: Resource Management Component, Land Use  
9 Component and Implementation Component.

10 All development and resource alteration proposed in this  
11 UMP is subject to the granting of appropriate permits,  
12 easements, licenses, and other required legal instruments.  
13 Approval of the UMP does not constitute an exemption from  
14 complying with the appropriate local, state, or federal  
15 agencies. This plan also is intended to meet the requirements  
16 for beach and shore preservation, as defined in Chapter 161,  
17 Florida Statutes, and Chapters 62B-33, 62B-36 and 62R49,  
18 FAC.

19 In the development of this UMP, the potential of the park to  
20 accommodate secondary management purposes was  
21 analyzed. These secondary purposes were considered  
22 within the context of the DRP's statutory responsibilities  
23 and the resource needs and values of the park. This analysis  
24 considered the park's natural and cultural resources,  
25 management needs, aesthetic values, and visitation and  
26 visitor experiences. For the CFG, it was determined that  
27 timber management activities for restoration could be  
28 accommodated in a manner that would be compatible and  
29 not interfere with the primary purpose of resource-based  
30 outdoor recreation and conservation. This compatible  
31 secondary management purpose is addressed in the  
32 Resource Management Component of the plan. Uses such as  
33 water resource development projects, water supply  
34 projects, stormwater management projects, linear facilities,  
35 and sustainable agriculture and forestry (other than those  
36 forest management activities specifically identified in this  
37 plan) are not consistent with this plan or the management  
38 purposes of the CFG.

39 The potential for generating revenue to enhance  
40 management also was analyzed. Visitor fees and charges  
41 are the principal sources of revenue generated by the park.  
42 It was determined that timber management and hunting  
43 activities would be appropriate at this park as an additional

1 source of revenue for land management since it is  
2 compatible with the park's primary purpose of resource-  
3 based outdoor recreation and conservation.

4 DRP may provide the services and facilities outlined in  
5 this plan either with its own funds and staff, by  
6 building partnerships, or through an outsourcing  
7 contract. Private contractors may assist with natural  
8 resource management and restoration activities or a  
9 concessionaire may provide services to park visitors to  
10 enhance the visitor experience. For example, a  
11 concessionaire could be authorized to sell merchandise  
12 and food and to rent recreational equipment for use in  
13 the park. A concessionaire also may be authorized to  
14 provide specialized services, such as interpretive tours,  
15 or overnight accommodations when the required  
16 capital investment exceeds that which DRP can elect to  
17 incur. Decisions regarding outsourcing, contracting with  
18 the private sector, the use of concessions, etc., are made on  
19 a case-by-case basis in accordance with the policies set  
20 forth in DRP's Operations Manual (OM).

## 21 **MANAGEMENT PROGRAM REVIEW**

### 22 **MANAGEMENT AUTHORITY AND** 23 **RESPONSIBILITY**

24 In accordance with Chapter 258, Florida Statutes, and  
25 Chapter 62D-2, Florida Administrative Code, DRP is charged  
26 with the responsibility of developing and operating Florida's  
27 recreation and parks system. These responsibilities are  
28 administered in accordance with the following policy:

29 *It shall be the policy of the Division of Recreation and*  
30 *Parks to promote the state park system for the use,*  
31 *enjoyment, and benefit of the people of Florida and*  
32 *visitors; to acquire typical portions of the original*  
33 *domain of the state which will be accessible to all of the*  
34 *people, and of such character as to emblemize the*  
35 *state's natural values; conserve these natural values*  
36 *for all time; administer the development, use and*  
37 *maintenance of these lands and render such public*  
38 *service in so doing, in such a manner as to enable the*  
39 *people of Florida and visitors to enjoy these values*  
40 *without depleting them; to contribute materially to the*  
41 *development of a strong mental, moral, and physical*



*The Office of Park Planning is responsible for the development of park management plans, such as this UMP.*



1           *fiber in the people; to provide for perpetual*  
2           *preservation of historic sites and memorials of*  
3           *statewide significance and interpretation of their*  
4           *history to the people; to contribute to the tourist*  
5           *appeal of Florida.*

6       For the purposes of administering the Parks, the DRP is  
7       divided into five Districts covering the northwest, northeast,  
8       central, southeast, and southwest areas of the state. CFG is  
9       located entirely within DRP District 3. Each District Bureau  
10      Chief is responsible for the overall development and  
11      maintenance of comprehensive multi-purpose outdoor  
12      recreation and natural and cultural resource conservation  
13      programs for the District. A Park Manager—typically one is  
14      assigned per park—is responsible for the day-to-day park  
15      operations and reports to applicable District staff. DRP staff  
16      in Tallahassee support state parks and District offices.

17     Many operating procedures are standardized system-wide  
18     and are set by internal direction and documented in the  
19     DRP’s OM. The OM serves as a guide to provide quality  
20     assurance and consistency in Florida State Park operational  
21     procedures and provides the DRP with a defined course of  
22     action that guides present and future decisions. OM  
23     guidance is based on applicable Florida State Statutes,  
24     Florida Administrative Codes/Rules, Department directives,  
25     and policy and direction of the Division Management Team.  
26     The OM covers areas such as personnel management,  
27     uniforms and personal appearance, training, signs,  
28     communications, fiscal procedures, interpretation,  
29     concessions, public use regulations, resource management,  
30     law enforcement, protection, safety, and maintenance.

31     **PARK MANAGEMENT GOALS**

32     The following park goals express DRP’s long-term intent in  
33     managing the state park:

- 34           • Provide administrative support for all park  
35           functions.
- 36           • Protect water quality and quantity in the park,  
37           restore hydrology to the greatest extent feasible, and  
38           maintain the restored condition.
- 39           • Restore and maintain the natural  
40           communities/habitats of the park.
- 41           • Maintain, improve, or restore imperiled species  
42           populations and habitats in the park.

- 1 • Remove exotic and invasive plants and animals from
- 2 the park and conduct needed maintenance/control.
- 3 • Protect, preserve, and maintain the cultural
- 4 resources of the park.
- 5 • Provide public access and recreational opportunities
- 6 in the park.
- 7 • Develop and maintain the capital facilities and
- 8 infrastructure necessary to meet the goals and
- 9 objectives of this UMP.

*United States Code (USC)  
460tt provides terms and  
conditions for the de-  
authorization of the federal  
barge canal project.*

*Volunteers provide valuable  
assistance to the  
Florida Park Service:*

- 38 percent of the DRP workforce is volunteers
- 1.2 million volunteer hours were logged in Florida state parks last year
- The total value of volunteer hours is \$28 million per year.

## 10 MANAGEMENT COORDINATION

11 The CFG is managed in accordance with all applicable laws  
12 and administrative rules. Included are the federal de-  
13 authorization terms and conditions agreed upon by the state  
14 of Florida by adoption of the resolution to de-authorize the  
15 barge canal project. These terms and conditions include a  
16 federal oversight role in the management of the CFG to  
17 ensure that the land is maintained as a state park or  
18 conservation/recreation area. By doing this, the state agrees  
19 to preserve, enhance, interpret, and manage the water and  
20 related land resources of the area containing cultural, fish  
21 and wildlife, scenic, and recreational values for the benefit  
22 and enjoyment of present and future generations of people  
23 and the development of outdoor recreation.

24 Several state agencies have a major or direct role in the  
25 management of the park are discussed in this UMP. The  
26 Florida Department of Agriculture and Consumer Services  
27 (FDACS), FFS, assists DRP staff in the development of wildfire  
28 emergency plans and provides assistance with prescribed  
29 burning activities. The FWC provides staff in the enforcement  
30 of state laws pertaining to wildlife, freshwater fish and other  
31 aquatic life existing within the park. In addition, the FWC aids  
32 DRP with wildlife management programs, including  
33 imperiled species management. The Division of Historical  
34 Resources (DHR) assists staff with the management and  
35 protection of archaeological and historical sites.

36 Volunteers provide an extension of the DRP work force to  
37 accomplish the agency mission. The DRP provides volunteers  
38 an opportunity to work in areas such as resource  
39 management, and environmental education/ interpretation.  
40 In 2014-15, 3,201 volunteers donated approximately 32,997  
41 volunteer hours on the CFG. Volunteers will continue to  
42 provide vital assistance in managing the CFG in the future.



*Public participation is a hallmark of the Florida Park Service unit management plan development process.*



1 Friends groups, also referred to as Citizen Support  
2 Organizations (CSOs), provide support to individual parks  
3 by volunteering, educating visitors, hosting events, and  
4 raising funds for specific park projects. In accordance with  
5 its bylaws, the Florida Greenways and Trails Foundation  
6 (FGTF) serves as the official overall CSO to the CFG. There are  
7 currently 86 friends groups supporting state parks  
8 throughout Florida. These groups and the citizens that  
9 pledge their time to them play a vital role in supporting  
10 Florida's state parks, ensuring that our natural, cultural,  
11 recreational, and historical resources are protected for  
12 future generations.

### 13 PUBLIC PARTICIPATION

14 Public participation is a process, not a single event. It consists  
15 of a series of activities and actions over the lifespan of a  
16 project to both inform the public and obtain input from them.  
17 Public participation affords stakeholders (those that have an  
18 interest or stake in an issue, such as individuals, interest  
19 groups, communities) the opportunity to influence decisions  
20 that affect their lives. As such, DRP offered multiple  
21 opportunities for input throughout the development of this  
22 UMP.

### 23 Preliminary Public Workshops

24 Initial public input was solicited by DRP through three  
25 preliminary public workshops from December 6-8, 2016.  
26 These workshops were held in Inglis, Ocala, and Palatka,  
27 Florida. The purpose of these meetings was to gather  
28 public input at the beginning of the management planning  
29 process. Meeting notices were published in the Florida  
30 Administrative Register, Volume 42, Issue 46, included on  
31 the Department Internet Calendar, posted in clear view at  
32 the park, and promoted locally.

33 As part of the public outreach during the December 2016  
34 public workshops, an online survey was employed to obtain  
35 feedback from the public concerning their use and  
36 perceptions of the CFG.

37 There were 26 respondents who took the survey, and  
38 the following is a summary of their responses:



- 1 • Almost two-thirds of the respondents were
- 2 from Putnam County, followed by a mixture
- 3 of respondents from the other counties.
- 4 • An equal number of males and females were
- 5 reported on the survey (13 male/13 female)
- 6 and all respondents, except one, reported an
- 7 age over 40.
- 8 • More than 50 percent of the respondents
- 9 reported travel of more than 10 miles (one
- 10 way) to the Greenway, with two-thirds of all
- 11 respondents stating that they visit the
- 12 Greenway at least once a month.
- 13 • The most common reported access points to
- 14 the Greenway were Rodman Recreation Area
- 15 (62 percent), Kenwood Recreation Area (37
- 16 percent), followed by the Landbridge
- 17 Trailhead, Santos Trailhead, Eureka, Orange
- 18 Springs, and St. Johns Loop North Trailhead
- 19 (all at 25 percent).
- 20 • Hiking (60 percent), boating (40 percent), and
- 21 wildlife watching (40 percent) were the top
- 22 three reported recreational activities.
- 23 • Of the total respondents, 42 percent were
- 24 satisfied with the current level of recreational
- 25 opportunities offered on the CFG, while 36
- 26 percent were not satisfied.
- 27 • 75 percent of respondents had an “Excellent”
- 28 or “Very Good” opinion of the CFG.

29 Opportunities also were provided within the survey for  
30 respondents to provide general comments. Ten comments  
31 were received concerning removing Kirkpatrick Dam and  
32 restoring the Ocklawaha River and eight comments were  
33 received expressing desires to expand mountain biking  
34 trails within the Greenway. The following comments were  
35 expressed once per issue: proper land management,  
36 reopening camping at Kenwood Recreation Area, and the  
37 desire for more horse trails.

### 38 **Public Hearings and Advisory Group Meeting**

39 DRP also will conduct three public hearings to gather  
40 recommendations and comments on the draft UMP.

## 41 **OTHER DESIGNATIONS**

42 The CFG is not located within or adjacent to an Area of  
43 Critical State Concern as defined in Section 380.05, F.S., and

1 it is not presently under study for such designation. The park  
2 is a component of the Florida Greenways and Trails System,  
3 administered by the DRP.



***“Special Waters”  
OFWs include  
41 of Florida’s  
1,700 rivers, lakes, and  
lake chains, several  
estuarine areas, and  
the Florida Keys.***

4 As noted during a review of the eligible nearby  
5 conservation areas, many Outstanding Florida Waters  
6 (OFWs) are near the CFG. Section 403.061(27), F.S., grants  
7 the DEP the power to establish rules that provide for a  
8 special category of waterbodies within the state, to be  
9 referred to as “Outstanding Florida Waters,” which shall be  
10 worthy of special protection because of their natural  
11 attributes. In addition to those areas that are OFWs by virtue  
12 of being state parks, aquatic preserves, or acquisitions  
13 through the state’s environmental land acquisition  
14 programs, four areas within or contiguous to the CFG are  
15 designated as Special Waters OFWs: Ocklawaha, Rainbow  
16 River, Silver River, and the Withlacoochee Riverine and  
17 Lake System.

# RESOURCE MANAGEMENT COMPONENT

## CFG Natural Resource Management Accomplishments 2007-2016

### INTRODUCTION

The DRP, in accordance with Chapter 258, F.S., has implemented resource management programs for preserving the representative examples of natural and cultural resources of statewide significance under its administration. The Resource Management Component (RMC) of this UMP details the CFG's natural and cultural resources, while also identifying management methods that are consistent with the DEP's overall mission of ecosystem management.

DRP's philosophy of resource management is *natural systems management*. Primary emphasis is placed on restoring and maintaining, to the greatest degree possible, the natural processes that originally shaped the structure, function, and species composition of Florida's diverse natural communities. Single-species management for imperiled species is appropriate in state parks when the maintenance, recovery, or restoration of a species or population is complicated due to constraints associated with long-term restoration efforts, unnaturally high mortality, or insufficient habitat. Single-species management should be compatible with the maintenance and restoration of natural processes, and should not imperil other native species or seriously compromise park values.

DRP's management goal for cultural resources is to preserve sites and objects that represent Florida's cultural periods, significant historic events, or persons. This goal often entails active measures to stabilize, reconstruct, or restore resources, or to rehabilitate them for appropriate public use.

Within the CFG, there are biotic (living) communities, which include all the plant and animal populations living within the greenway. These communities interact with the abiotic (non-living) resources (soil, air, water, and sunlight) to form what is known as an *ecosystem*. The size of the area involved when defining communities or ecosystems can vary. For instance, given that the CFG

#### Prescribed Fire

15,977 acres burned

- 66 burn zones in rotation (7,325 acres)
- 29 new burn zones since 2007
- All the CFG fire type acreage now divided into burn zones
- Significantly increased utilization of prescribed burn contractors to increase annual acreage burned

#### Natural Community Restoration

- 3,925 acres in timber harvests for restoration of natural communities with the side benefit of earning \$1.065 million in revenue
- 498 acres of trees planted
- 157 acres of groundcover planted
- 540 acres of mechanical treatments
- A timber inventory and management plan for the 9,000 acre Etoniah addition

#### Endangered Species

- Coordinated with Audubon Florida to join their annual Jay Watch program.
- Restored 523 additional acres of scrub
- Contracted to band Jays to better manage responses to management measures
- Mechanically treated 538 acres
- Increased from 46 birds in 2009 to 111 in 2015
- Coordinated with FWC to set up boxes for kestrel recruitment.

1 incorporates a larger ecosystem, management often is affected by conditions and events occurring  
 2 well beyond park boundaries. Therefore, DRP uses *ecosystem management* through a resource  
 3 management evaluation program to assess resource conditions, evaluate management activities,  
 4 refine management actions, and review local comprehensive plans and development permit  
 5 applications for park/ecosystem impacts.

## 6 RESOURCE DESCRIPTIONS AND ASSESSMENT

### 7 Natural Resources

8 The CFG is 110 miles long and contains 70,640 acres, which are not uniform in the distribution of  
 9 plant and wildlife species. To assist with the identification of natural resources management  
 10 activities at the CFG, the entire CFG has been subdivided into the western, central, and eastern  
 11 sections. Within each of these, the greenway has been further subdivided into 740 management  
 12 zones that delineate areas on the ground used to reference management activities (see: Management  
 13 Zones Map). These management zones range in shape and size based on natural community type,  
 14 burn zone boundaries, and the location of existing roads and natural fire breaks. It is important to  
 15 note that all burn zones are management zones; however, not all management zones include fire-  
 16 dependent natural communities.

17 The westernmost portion of the greenway, which is considered the section from the Gulf of Mexico  
 18 to SR 200, contains 140 management zones that total 16,088 acres. The largest management zone  
 19 within the western portion of the CFG is 3,156 acres, while the smallest management zone is 0.001  
 20 acre. The average size management zone within the western portion of the CFG is 115 acres. There  
 21 are seven management zones that exceed 500 acres. Of the 140 management zones, 49 are  
 22 considered fire dependent (Table 1).

23 The central portion of the greenway, which is considered the section from SR 200 to CR 316, contains  
 24 281 management zones that total 27,270 acres. The largest management zone within the central  
 25 portion of the CFG is 2,378 acres and the smallest management zone is 0.30 acre. The average size  
 26 management zone within the central portion of the CFG is 96 acres. There are seven management  
 27 zones that exceed 500 acres. Of the 281 management zones, 142 are considered fire dependent  
 28 (Table 2).

29 The eastern portion of the greenway, which is considered the section from CR 316 to the St. Johns  
 30 River, contains 309 management zones that total 27,282 acres. The largest management zone within  
 31 the eastern portion of the CFG is 2,765 acres and the smallest management zone is 0.005 acre. The  
 32 average size management zone within the eastern portion of the CFG is 88 acres. There are seven  
 33 management zones that exceed 500 acres. Of the 309 management zones, 166 are considered fire  
 34 dependent (Table 3).

35 **Table 1. CFG Western Management Zones, Acreage, and Fire Type**

Management Zone ID	Acres	Managed by Fire	Management Zone ID	Acres	Managed by Fire
W001	51.78	Y	W071	9.77	N
W002	15.84	N	W072	0.97	N

Management Zone ID	Acres	Managed by Fire	Management Zone ID	Acres	Managed by Fire
W003	120.35	Y	W073	0.27	N
W004	202.90	Y	W074	0.082	N
W005	99.90	Y	W075	3.89	N
W006	190.08	Y	W076	5.25	N
W007	293.25	Y	W077	60.0	Y
W008	299.56	Y	W078	70.68	Y
W009	65.31	Y	W079	3,155.50	N
W010	24.38	Y	W080	60.76	Y
W011	90.86	Y	W081	15.28	N
W012	55.04	Y	W082	12.58	N
W013	90.91	Y	W083	62.90	N
W014	15.03	N	W084	27.52	Y
W015	63.12	Y	W085	40.75	N
W016	71.46	Y	W086	12.45	N
W017	95.95	N	W087	155.10	N
W018	147.24	Y	W088	39.58	N
W019	12.24	Y	W089	20.49	N
W020	67.18	Y	W090	16.24	N
W021	151.51	Y	W091	26.44	N
W022	10.11	N	W092	71.29	Y
W023	39.95	Y	W093	33.10	N
W024	147.28	N	W094	135.92	Y
W025	48.35	N	W095	100.99	Y
W026	67.41	N	W096	158.68	Y
W027	87.96	N	W097	47.34	N
W028	107.95	Y	W098	29.49	N
W029	5.58	N	W099	15.08	N
W030	262.98	Y	W100	12.72	N
W031	68.75	N	W101	1.66	N
W032	56.73	N	W102	13.91	N
W033	153.60	Y	W103	1.57	N
W034	19.33	N	W104	90.40	N
W035	89.87	Y	W105	27.84	N
W036	45.79	Y	W106	87.56	N
W037	24.03	N	W107	104.12	N
W038	0.82	N	W108	107.29	N
W039	91.08	Y	W109	68.33	N
W040	20.22	N	W110	43.21	Y
W041	28.45	N	W111	27.88	N

Management Zone ID	Acres	Managed by Fire	Management Zone ID	Acres	Managed by Fire
W042	9.33	N	W112	3.90	N
W043	66.46	N	W113	1.58	N
W044	67.20	N	W114	5.32	N
W045	8.34	N	W115	4.22	N
W046	6.61	N	W116	19.68	N
W047	28.44	N	W117	11.96	N
W048	0.268	N	W118	38.23	N
W049	0.003	N	W119	3.58	N
W050	0.001	N	W120	3.44	N
W051	2.17	N	W121	112.30	N
W052	7.32	N	W122	97.82	N
W053	2.30	N	W123	133.83	N
W054	8.00	N	W124	553.81	Y
W055	68.85	N	W125	126.02	Y
W056	55.67	N	W126	1,831.33	Y
W057	0.37	N	W127	42.26	N
W058	9.41	N	W128	46.32	N
W059	3.52	N	W129	13.20	N
W060	92.06	Y	W130	83.70	N
W061	19.40	N	W131	31.94	N
W062	0.93	N	W132	84.73	N
W063	0.27	N	W133	75.81	N
W064	0.14	N	W134	15.80	N
W065	0.88	N	W135	1,161.54	N
W066	6.31	N	W136	2.62	N
W067	42.43	N	W137	11.89	N
W068	0.79	N	W138	195.74	N
W069	4.30	N	W139	2,272.99	N
W070	1.93	N	W140	14.03	N
<b>Total Acreage, Western CFG Zones: 16,088</b>					

1

2 **Table 2. CFG Central Management Zones, Acreage, and Fire Type**

Management Zone ID	Acres	Managed by Fire	Management Zone ID	Acres	Managed by Fire
C001	3.9	N	C142	278.2	Y
C002	12.4	N	C143	19.2	N
C003	5.1	N	C144	34.8	Y
C004	17.1	N	C145	10.4	Y

Management Zone ID	Acres	Managed by Fire	Management Zone ID	Acres	Managed by Fire
C005	26.5	N	C146	10.5	N
C006	23.0	N	C147	34.4	N
C007	225.5	N	C148	24.4	Y
C008	65.2	N	C149	45.9	Y
C009	71.3	Y	C150	1.5	N
C010	181.2	Y	C151	3.7	N
C011	113.0	N	C152	9.7	N
C012	431.8	Y	C153	13.7	N
C013	1481.2	N	C154	3.1	N
C014	168.3	N	C155	0.3	N
C015	110.1	N	C156	30.8	Y
C016	33.8	N	C157	37.8	Y
C017	27.3	Y	C158	17.3	Y
C018	15.3	N	C159	3.5	N
C019	27.9	N	C160	12.4	Y
C020	140.5	Y	C161	1.9	N
C021	39.7	N	C162	15.9	N
C022	42.2	N	C163	21.7	N
C023	8.0	N	C164	17.8	N
C024	3.4	N	C165	7.5	N
C025	96.3	N	C166	8.9	N
C026	61.2	N	C167	34.7	N
C027	60.7	N	C168	466.0	N
C028	0.5	N	C169	13.9	Y
C029	21.6	N	C170	137.5	Y
C030	2.4	N	C171	17.6	N
C031	18.0	N	C172	12.1	N
C032	8.6	N	C173	2.1	N
C033	3.4	N	C174	9.6	Y
C034	1.0	N	C175	59.7	Y
C035	5.6	N	C176	30.3	Y
C036	4.5	N	C177	28.1	N
C037	1.7	N	C178	79.0	N
C038	3.1	N	C179	25.6	N
C039	5.9	N	C180	21.7	N
C040	7.0	N	C181	156.0	N
C041	2.8	N	C182	9.5	N
C042	0.4	N	C183	27.9	Y
C043	1.5	N	C184	130.5	Y



Management Zone ID	Acres	Managed by Fire	Management Zone ID	Acres	Managed by Fire
C044	14.0	Y	C185	25.5	Y
C045	15.2	Y	C186	11.6	Y
C046	145.2	Y	C187	4.3	N
C047	3.0	N	C188	8.1	N
C048	400.6	Y	C189	33.1	Y
C049	45.0	N	C190	160.0	Y
C050	47.5	N	C191	6.4	Y
C051	393.1	N	C192	11.7	Y
C052	287.8	N	C193	31.3	Y
C053	133.0	N	C194	42.2	Y
C054	6.6	N	C195	50.3	Y
C055	37.2	N	C196	18.0	Y
C056	31.7	N	C197	29.5	Y
C057	3.4	N	C198	15.6	Y
C058	852.4	Y	C199	17.3	Y
C059	22.4	N	C200	44.7	Y
C060	137.9	Y	C201	15.6	Y
C061	167.3	Y	C202	15.4	Y
C062	154.4	Y	C203	29.1	Y
C063	131.0	Y	C204	34.1	Y
C064	65.4	Y	C205	287.2	Y
C065	24.1	Y	C206	132.7	Y
C066	38.8	Y	C207	135.3	Y
C067	59.6	Y	C208	3.7	N
C068	92.1	Y	C209	6.4	N
C069	24.8	Y	C210	18.9	N
C070	57.8	Y	C211	75.7	Y
C071	45.5	Y	C212	68.4	N
C072	96.4	Y	C213	20.3	N
C073	901.3	N	C214	65.0	N
C074	170.1	Y	C215	26.9	N
C075	45.0	Y	C216	28.2	N
C076	11.8	Y	C217	586.4	Y
C077	127.0	Y	C218	61.4	Y
C078	40.0	Y	C219	68.4	N
C079	236.1	Y	C220	40.5	N
C080	30.3	Y	C221	75.3	Y
C081	54.9	Y	C222	119.3	Y
C082	1.4	N	C223	286.5	N

Management Zone ID	Acres	Managed by Fire	Management Zone ID	Acres	Managed by Fire
C083	4.0	N	C224	115.1	Y
C084	24.0	N	C225	0.4	N
C085	23.8	N	C226	4.2	N
C086	61.5	N	C227	9.9	N
C087	25.7	N	C228	30.9	N
C088	11.5	N	C229	22.0	Y
C089	28.3	N	C230	26.4	Y
C090	4.2	N	C231	15.3	Y
C091	7.6	N	C232	16.9	Y
C092	9.0	N	C233	28.3	Y
C093	2.7	N	C234	17.1	Y
C094	14.9	N	C235	28.9	Y
C095	23.7	N	C236	26.6	Y
C096	7.8	N	C237	24.3	Y
C097	5.2	N	C238	36.2	Y
C098	309.0	N	C239	44.8	Y
C099	337.5	N	C240	24.4	Y
C100	4.4	N	C241	13.6	Y
C101	17.0	N	C242	69.2	Y
C102	56.5	N	C243	25.1	N
C103	363.7	N	C244	72.0	Y
C104	2262.1	N	C245	20.4	Y
C105	2378.3	N	C245A	27.1	Y
C106	93.4	Y	C246	13.1	Y
C107	15.9	N	C247	12.9	Y
C108	12.0	N	C248	13.5	Y
C109	7.8	Y	C249	14.4	Y
C110	20.7	N	C250	17.7	Y
C111	31.9	Y	C251	4.1	Y
C112	22.8	Y	C252	31.8	Y
C113	5.9	N	C253	5.9	Y
C114	74.8	Y	C254	118.4	Y
C115	273.9	N	C255	561.4	Y
C116	165.7	Y	C256	159.1	Y
C117	9.5	N	C257	149.4	Y
C118	104.2	Y	C258	146.7	Y
C119	468.8	N	C259	140.2	Y
C120	47.6	Y	C260	12.1	N
C121	30.7	Y	C261	301.3	Y

Management Zone ID	Acres	Managed by Fire	Management Zone ID	Acres	Managed by Fire
C122	152.3	Y	C262	58.9	N
C123	70.3	Y	C263	299.7	Y
C124	51.8	Y	C264	201.6	Y
C125	67.1	Y	C265	212.6	Y
C126	77.4	N	C266	394.6	Y
C127	20.2	N	C267	186.7	N
C128	57.8	Y	C268	46.2	Y
C129	2.1	N	C269	12.6	Y
C130	15.4	N	C270	14.8	N
C131	0.5	N	C271	20.4	N
C132	59.5	N	C272	181.1	Y
C133	0.4	N	C273	171.5	Y
C134	0.6	N	C274	230.5	Y
C135	5.8	N	C275	475.4	Y
C136	77.1	N	C276	366.9	Y
C137	24.9	Y	C277	182.6	Y
C138	61.6	Y	C278	107.5	Y
C139	74.7	Y	C279	28.9	N
C140	54.1	N	C280	17.3	N
C141	37.0	Y	C281	11.4	N
<b>Total Acreage, Central CFG Zones: 27,269.9</b>					

1

2 **Table 3. CFG Eastern Management Zones, Acreage, and Fire Type**

Management Zone ID	Acres	Managed by Fire	Management Zone ID	Acres	Managed by Fire
E001	61.2	N	E155	69.1	Y
E002	130.1	N	E156	23.4	Y
E003	108.6	N	E157	19.1	N
E004	185.7	N	E158	11.4	N
E005	5.1	N	E159	22.9	N
E006	12.4	Y	E160	5.8	N
E007	2.8	N	E161	420.1	Y
E008	125.4	Y	E162	107.2	Y
E009	21.1	N	E163	105.2	Y
E010	42.3	N	E164	1.9	N
E011	81.5	N	E165	32.8	N
E012	135.4	Y	E166	50.5	N
E013	46.1	N	E167	68.9	N

Management Zone ID	Acres	Managed by Fire	Management Zone ID	Acres	Managed by Fire
E014	155.2	Y	E168	151.4	Y
E015	31.7	N	E169	108.3	Y
E016	16.9	N	E170	86.1	Y
E017	31.0	N	E171	74.4	N
E018	80.4	Y	E172	39.4	Y
E019	32.4	N	E173	12.2	N
E020	7.5	N	E174	7.8	N
E021	26.0	N	E175	102.8	N
E022	1.6	N	E176	93.8	Y
E023	2.6	N	E177	15.4	Y
E024	0.9	N	E178	122.5	Y
E025	6.9	N	E179	151.3	Y
E026	10.0	N	E180	153.1	Y
E027	7.1	N	E181	146.0	Y
E028	16.9	N	E182	299.7	Y
E029	44.8	Y	E183	61.3	Y
E030	53.0	Y	E184	16.8	Y
E031	13.3	N	E185	43.1	Y
E032	53.4	Y	E186	79.5	Y
E033	6.0	Y	E187	29.0	Y
E034	156.0	Y	E188	164.2	Y
E035	53.3	Y	E189	50.1	Y
E036	200.6	N	E190	53.7	Y
E037	85.4	Y	E191	37.3	Y
E038	32.2	Y	E192	75.3	Y
E039	134.7	Y	E193	28.0	Y
E040	47.1	Y	E194	74.9	N
E041	23.1	Y	E195	110.1	Y
E042	7.6	Y	E196	55.0	N
E043	28.6	Y	E197	51.7	Y
E044	448.1	Y	E198	14.1	Y
E045	97.6	Y	E199	210.2	N
E046	67.7	Y	E200	28.8	N
E047	15.7	Y	E201	51.7	Y
E048	10.9	Y	E202	181.1	Y
E049	116.6	Y	E203	19.4	Y
E050	64.5	Y	E204	128.0	Y
E051	4.6	N	E205	126.6	N
E052	11.8	Y	E206	89.0	N

Management Zone ID	Acres	Managed by Fire	Management Zone ID	Acres	Managed by Fire
E053	49.8	Y	E207	151.0	Y
E054	50.5	Y	E208	12.5	N
E055	22.7	Y	E209	88.8	Y
E056	73.6	Y	E210	96.7	Y
E057	14.8	Y	E211	75.8	Y
E058	61.5	Y	E212	13.8	N
E059	46.8	Y	E213	3.2	N
E060	12.5	N	E214	26.7	Y
E061	55.2	Y	E215	173.2	Y
E062	112.0	Y	E216	136.8	Y
E063	35.3	Y	E217	129.9	Y
E064	91.9	Y	E218	35.9	Y
E065	474.9	Y	E219	84.1	Y
E066	105.7	N	E220	96.6	N
E067	9.0	Y	E221	271.7	N
E068	57.4	Y	E222	65.2	N
E069	12.8	Y	E223	200.3	Y
E070	745.3	N	E224	157.6	Y
E071	10.1	N	E225	117.3	Y
E072	35.1	Y	E226	30.8	Y
E073	115.3	Y	E227	61.1	Y
E074	190.7	Y	E228	7.6	Y
E075	36.6	Y	E229	40.3	Y
E076	21.7	N	E230	52.5	Y
E077	58.5	Y	E231	31.1	Y
E078	160.6	Y	E232	15.2	N
E079	136.0	Y	E233	20.0	N
E080	42.0	N	E234	69.4	Y
E081	35.6	Y	E235	183.1	Y
E082	25.9	N	E236	155.7	Y
E083	23.5	N	E237	30.8	N
E084	10.4	N	E238	126.9	N
E085	163.6	Y	E239	10.0	N
E086	41.2	N	E240	36.0	Y
E087	7.3	N	E241	2,765.1	N
E088	260.2	Y	E242	110.3	N
E089	18.4	N	E243	63.4	N
E090	35.5	Y	E244	147.0	N
E091	13.3	Y	E245	175.8	N

Management Zone ID	Acres	Managed by Fire	Management Zone ID	Acres	Managed by Fire
E092	79.0	Y	E246	14.6	N
E093	45.2	Y	E247	37.1	N
E094	55.0	Y	E248	4.1	N
E095	18.4	Y	E249	84.9	N
E096	7.2	Y	E250	59.5	N
E097	232.4	Y	E251	64.6	N
E098	28.3	Y	E252	170.3	N
E099	1,018.2	Y	E253	3.9	N
E100	113.7	Y	E254	20.6	N
E101	90.8	Y	E255	152.4	N
E102	3.2	N	E256	4.0	N
E103	1,139.2	N	E257	194.1	N
E104	26.2	Y	E258	3.2	N
E105	60.0	Y	E259	18.7	N
E106	24.9	N	E260	29.6	N
E107	13.0	N	E261	2.1	N
E108	31.9	Y	E262	13.3	N
E109	4.1	N	E263	2.3	N
E110	4.7	N	E264	2.3	N
E111	16.2	N	E265	51.5	N
E112	18.0	Y	E266	13.1	N
E113	23.7	Y	E267	2.2	N
E114	41.0	Y	E268	0.0	N
E115	10.9	N	E269	0.1	N
E116	5.3	N	E270	15.0	N
E117	42.4	Y	E271	34.6	N
E118	16.1	N	E272	16.5	N
E119	188.2	Y	E273	35.5	N
E120	142.8	Y	E274	6.1	N
E121	12.7	Y	E275	25.3	N
E122	39.3	Y	E276	7.0	N
E123	11.1	Y	E277	14.4	N
E124	30.3	Y	E278	28.1	N
E125	4.9	Y	E279	340.4	Y
E126	230.9	Y	E280	46.6	N
E127	47.2	Y	E281	630.3	N
E128	35.0	Y	E282	308.3	N
E129	59.2	Y	E283	57.6	N
E130	40.0	Y	E284	2.0	N

Management Zone ID	Acres	Managed by Fire	Management Zone ID	Acres	Managed by Fire
E131	78.9	N	E285	203.5	N
E132	45.5	Y	E286	68.0	Y
E133	53.8	Y	E287	325.9	N
E134	21.4	N	E288	55.2	Y
E135	13.6	Y	E289	0.8	N
E136	99.7	Y	E290	8.1	N
E137	35.2	Y	E291	38.3	N
E138	102.0	Y	E292	579.8	N
E139	26.3	Y	E293	32.1	N
E140	62.3	Y	E294	29.6	N
E141	168.3	Y	E295	57.8	N
E142	87.9	Y	E296	24.1	N
E143	14.6	Y	E297	13.7	N
E144	23.1	N	E298	534.4	N
E145	24.3	Y	E299	26.7	N
E146	9.1	N	E300	54.5	Y
E147	27.8	Y	E301	55.1	Y
E148	15.3	Y	E302	47.5	N
E149	29.7	Y	E303	69.4	N
E150	7.9	Y	E304	1.7	N
E151	6.4	N	E305	4.9	N
E152	15.4	Y	E306	50.1	N
E153	11.9	Y	E307	109.6	Y
E154	114.2	N	E308	143.4	Y
			E309	141.4	N
<b>Total Acreage, Eastern Zones: 27,282.0</b>					

1

2 **Topography and Physiography**

3 The topography and physiography of the CFG reflects the underlying geology and the effect of large-  
4 scale man-made alterations to the landscape. The western half of the CFG is in the Ocala Uplift  
5 District, so named for the underlying subsurface geologic structure. In this section, the limestone is  
6 nearer to the surface and not covered by thick layers of old sediments. The sediments are thicker  
7 near I-75 and tend to be composed of sand, which erodes with slightly steeper slopes. The extreme  
8 western section from the Gulf of Mexico coast to Dunnellon is primarily composed of three  
9 physiographic provinces within the Ocala Uplift District (Brooks, 1981).

10 From the coast to U.S. Highway 19/98 (US 19/98), the area north of the Barge Canal is in the  
11 Waccasassa Coastal Strip. This province is a low limestone plain with some rockland upon which has  
12 developed a hardwood forest mixed with flatwoods and swamps. The similar Chassahowitzka

1 Coastal Strip occurs on the south side of the canal. It is characterized as a very low coastal strip of  
2 limestone rocklands mostly covered by hardwoods and swamps, and some flatwoods. Elevations  
3 typically are 10 feet and less (Brooks, 1981). The landscape here has been drastically altered by  
4 damming the Withlacoochee River and partially diverting its flow into a canal dredged into the Gulf  
5 of Mexico. The topography of the CFG is driven not only by the underlying geology, and the effects of  
6 river systems, but also by the hand of man on the landscape. The natural topography of the CFG  
7 generally is flat, but the levees along the barge canal can rise 25 feet to 30 feet higher than the natural  
8 grade.

9 Between US 19/98 and the town of Dunnellon is the Waccasassa Flats. Here, thin surficial clastic  
10 sediments overlie the limestone, and the elevation is less than 56 feet. Historically, this was flatwoods  
11 terrain (Brooks, 1981). Now, CFG property is relatively close to the shores of Lake Rousseau, which  
12 affects the hydrology of the area and does not provide much unflooded land for flatwoods in today's  
13 landscape.

14 The CFG property on the south side of the Withlacoochee River across from Dunnellon is in the  
15 Hernando Hammock province. The most characteristic features of the province are the thick, deeply  
16 weathered deposits of sand and clayey sand (Brooks, 1981). Typically, the elevation range is 100 feet  
17 to 160 feet in the province, but the CFG property is in the river floodplain and does not rise much  
18 beyond 50 feet.

19 From Dunnellon to County Road 484 (CR 484), the CFG is in the Tsala Apopka Basin province. Tsala  
20 Apopka is an erosional valley in the limestone terrain of the Ocala Uplift. The surficial sands generally  
21 are thin. In some areas, recent freshwater marls and peat have been deposited. The area is a maze of  
22 islands, swamps, marshes, and lakes. Flatwoods are typical on the land portions (Brooks, 1981). This  
23 portion of the CFG has been altered significantly by the digging of the old Ship Canal, then the Barge  
24 Canal. Five dig sites exist in the Tsala Apopka. The dig sites in the canal area can be up to 50 feet  
25 above the surrounding landscape and the pits from which material was removed can be another 20  
26 feet lower.

27 The CFG east of CR 484 to Marshall Swamp is composed of Newberry Sand Hills, Anthony Hills, and  
28 Ocala Hills. These provinces are deeply weathered and leached sands that rest directly on the Ocala  
29 Limestone. The sediments here are relatively thicker compared to the provinces to the west, and their  
30 predominantly sand composition accounts for the hilly landscape. Drainage is primarily internal.  
31 Historically, the xeric sandhills were woodland of longleaf pine (*Pinus palustris*) and turkey oak  
32 (*Quercus laevis*). Elevations generally are between 70 feet and 110 feet on the CFG (Brooks, 1981).

33 North of Marshall Swamp to the beginnings of Rodman Reservoir, the CFG is predominantly in the  
34 Ocklawaha Valley province. Ocklawaha Valley is an erosional valley partially backfilled with older  
35 estuarine deposits. A poorly drained flatwoods terrace (elevations about 80 feet) borders the river  
36 swamp (Brooks, 1981). The valley consists of relatively broad and flat floodplain.

37 A few areas—mostly outlying properties to the east of the Ocklawaha River and adjacent to the Ocala  
38 National Forest—are in the Central Lake District, and the Lynne Karst province. While this province  
39 is known for sandhills over much of its area, the CFG properties are in low-lying local karst areas that  
40 are flooded periodically.



1 The remainder of the CFG property—except for the northernmost extent, a few hundred acres just  
2 southeast of Hollister, Florida—is in the St. Johns Offset province. This portion of the St. Johns River  
3 Valley is very ancient. It is partially filled with estuarine deposits from when the Atlantic coastline  
4 extended this far inland up the St. Johns River. The underlying limestone is very near the surface and  
5 contributes to the development of the broad valley. Flatwoods occur on the Pleistocene terraces and  
6 a river swamp forest, generally with many cabbage palms (*Sabal palmetto*), occurs on the floodplain  
7 (Brooks, 1981).

## 8 **Geology**

9 Florida is composed of a large, relatively tectonically stable plateau. The deepest of the underlying  
10 basement rocks of the Floridan Plateau lie at least 4,000 feet below the land that comprises the state  
11 of Florida. These deep rocks are covered by layers of limestone and dolomite accumulated during the  
12 Cretaceous time period, 145 million years ago (mya) to 65 mya, when Florida was surrounded by  
13 ocean and covered by a warm sea.

14 Similar to the Bahama Islands now, Florida was an underwater plateau cut off from the mainland by  
15 an ocean current (Gulf Trough) until around 23 mya, when sea levels dropped and Florida emerged  
16 from the sea. The Ocala Uplift, a dominant surface feature, was created during this time. The Ocala  
17 Uplift is a very important geologic feature in Florida. It stretches all along West Central Florida from  
18 Brooksville in the south to Live Oak in the north. It runs parallel to and west of I-75 and is  
19 characterized by high, rolling hills.

20 As the Gulf Trough was closed off, sediments from erosion in the Appalachian Mountains were no  
21 longer carried away by ocean currents, and began to deposit on Florida. For the next 23 million years,  
22 the limestone of Florida was covered with these sediments, which have themselves eroded, forming  
23 Florida's current topography.

24 The geology of the CFG west of US 441 is dominated by the Ocala Uplift, and is composed primarily  
25 of two formations: the Undifferentiated Quaternary Sediments and the Ocala Limestone. Most of the  
26 sediments here are thinner and date, for the most part, from 25 mya to 11,700 years ago, while the  
27 underlying Ocala Limestone dates from 56 mya to 34 mya. The underlying geology gives rise to the  
28 distinctive topography of the western part of the CFG.

29 The Undifferentiated Quaternary Sediments are where these sediments exceed 20 feet (6.1 meters)  
30 in thickness, and were mapped as discrete units in the geologic map (Scott, 2001). The Ocala  
31 Limestone consists of nearly pure limestones and occasional dolostones. It can be subdivided into  
32 lower and upper facies. The lower member is composed of a white to cream-colored, fine to medium-  
33 grained limestone, and it is not encountered often. The upper facies is a white limestone with fossils  
34 and chert commonly occurring throughout (Scott, 2001).

35 The Ocala Limestone is at or near the surface on the western side of the CFG. In these areas, the Ocala  
36 Limestone exhibits extensive karstification. These karst features often have tens of feet (meters) of  
37 relief and may contain disappearing streams and springs within these areas. The permeable, highly  
38 transmissive carbonates of the Ocala Limestone form an important part of the Florida Aquifer System  
39 (FAS). It is one of the most permeable rock units in the FAS (Miller, 1986).

1 East of I-75, the CFG is dominated by geologically recent sediments associated with the Ocklawaha  
2 River Valley. The underlying limestones are much deeper, with a thicker layer of sediments.  
3 Primarily, these sediments are classified in geologic maps as Undifferentiated Quaternary Sediments  
4 and Holocene Sediments. Undifferentiated Quaternary Sediments consist of siliciclastic rocks,  
5 organics, and freshwater carbonates. Where these sediments exceed 20 feet (6.1 meters) in  
6 thickness, they were mapped as discrete units (Scott, 2001). In this area of the CFG, these deposits  
7 were created by nearshore sedimentation during times of higher sea levels. They tend to occur  
8 farther from the Ocklawaha River (Faulkner, 1973). More recent Holocene Sediments occur nearer  
9 to the river and were deposited since the area has been continuously above sea level. They consist of  
10 comparatively thin beds of alluvium; freshwater marl; peats and muds in stream and lake bottoms;  
11 and some windblown sand deposits (Faulkner, 1973).

12 There are smaller geologic areas of this portion of the CFG, which are less than 750 acres each. The  
13 narrow area between US 441 and Marshall Swamp comprises the Cypresshead Formation, which  
14 consists of reddish brown to reddish orange, unconsolidated to poorly consolidated, fine to very  
15 coarse grained, clean to clayey sands. Cross bedded sands are common within the formation. The  
16 outlying parcels of the CFG that are north of Silver Springs and east of the Ocklawaha River are made  
17 up of the Coosawhatchie Formation, which varies from a light gray to olive gray, poorly consolidated,  
18 variably clayey and phosphatic sand with few fossils, to an olive gray, poorly to moderately  
19 consolidated, slightly sandy, silty clay.

## 20 **Soils**

21 The CFG is such a large area and crosses so much of the central part of the state that it encompasses  
22 many of the soil series mapping units that make up Central Florida (see Addendum 7). There are 146  
23 different Natural Resources Conservation Service (NRCS) soil mapping units identified as being  
24 within the CFG. While the NRCS data about individual soil mapping units contain valuable  
25 information about the capabilities and limitations of these discrete units, it is difficult to adequately  
26 assess the current soil conditions of the CFG because of the drastic man-made alterations to the  
27 landscape.

28 Construction of levees and canals on the far western and eastern sections of the CFG have altered the  
29 soils. Soils were excavated, water regimes changed, excavation spoil distributed, and new materials  
30 brought in for engineering reasons. In addition, just off the coast, material was dredged up to make a  
31 channel and distributed in numerous soil piles, creating new above-ground areas.

32 In the central canal section, the abandoned dig sites not only disturbed the soil in the actual dig site,  
33 but the spoil piles of excavated material cover the archetypal soils near the sites. The hydrology of  
34 the soils also has been altered by inundating Lake Rousseau and Rodman Reservoir.

35 Before the land was obtained to create the Barge Canal or the CFG, various agricultural activities had  
36 been practiced throughout the area. Some of these activities date back more than a century. For  
37 instance, parts of Marshall Swamp were an active sugar plantation before the Civil War. Also,  
38 historically, there was phosphate mining around the town of Dunnellon during the turn of the 20th  
39 century.

40 Despite past impacts to soils, there are some general conclusions that can be made about the soils of  
41 the CFG. Generally, the soils reflect the sediments that are the parent material and the hydrologic

1 regime. Hydric soils are formed under conditions of saturation, flooding, or ponding for a long-  
 2 enough period during the growing season to develop anaerobic conditions in the upper part of the  
 3 soil. Soil drainage class represents the moisture condition of the soil in its natural condition  
 4 throughout the year, and flooding frequency, shown in Table 4, is the possibility of a soil flooding in  
 5 any given year.

6 **Table 4. Flooding Frequency Summary for Soils on the CFG**

<b>Flooding Frequency</b>	<b>Acres</b>
Very frequent	1,607.7
Frequent	7,955.9
Rare	1,434.3
None	48,142.6
<b>Total</b>	<b>57,532.8</b>

7  
 8 On the CFG, hydric soils exist in large areas from Dunnellon west, except for the canal levees; along  
 9 the Ocklawaha River and the far western area, except canal levees; and on Ross Prairie. The areas  
 10 along the canal east of Ross Prairie are not hydric soils. Table 5, below, shows the division between  
 11 soils that are hydric and not hydric.

12 **Table 5. Hydric Soils Summary for Soils on the CFG**

<b>Hydric</b>	<b>Acres</b>
Not	15,592.6
Hydric	43,547.9
<b>Total</b>	<b>59,140.6</b>

13  
 14 These same areas are in the poorly drained soil drainage classes, but flooding frequency is different.  
 15 The only large areas with a possibility of being flooded in any given year are next to the Gulf of Mexico  
 16 (GOM), areas next to the Ocklawaha River, and low wet areas to the north. Table 6 shows a summary  
 17 of soil drainage.

18 **Table 6. Soil Drainage Class Summary for Soils on the CFG**

<b>Drainage Class</b>	<b>Acres</b>
Excessively drained	8,144.6
Very poorly drained	23,110.4
Moderately well drained	1,474.2
Well drained	5,243.2
Somewhat poorly drained	3,118.8
Poorly drained	18,049.3
<b>Total</b>	<b>59,140.6</b>

19  
 20 Table 7, below, displays the Soil Series classes with more than 1,000 acres on the CFG. This reflects  
 21 78 percent of the soils on the CFG.

1 **Table 7. Soil Series Class Summary for the Commonly Occurring Soils on the CFG**

Series Name	Flooding Frequency	Drainage Class	Hydric	Acres
Terra Ceia	None	Very poorly drained	Yes	10,784.7
Candler	None	Well drained	No	6,776.8
Pomona	None	Poorly drained	Yes	6,111.8
Holopaw	None	Poorly drained	Yes	3,884.7
Arredondo	None	Well drained	No	2,317.9
Placid	None	Very poorly drained	Yes	2,267.3
Palmetto	None	Poorly drained	Yes	2,214.4
Bluff	Frequent	Very poorly drained	Yes	2,190.1
Riviera	None	Poorly drained	Yes	1,719.9
Homossa	Very frequent	Very poorly drained	Yes	1,564.3
Arents	None	Somewhat poorly drained	No	1,467.9
Myakka	None	Very poorly drained	Yes	1,340.6
Boca	None	Poorly drained	Yes	1,187.1
Anclote	None	Very poorly drained	Yes	1,156.3
Astatula	None	Excessively drained	No	1,110.9
<b>Total</b>				<b>46,094.7</b>

2

3 **Minerals**

4 The close proximity of the Ocala Limestone to the surface in the western half of the CFG makes it  
 5 feasible in areas to mine limestone. There are three limestone quarries adjacent to the CFG: two are  
 6 west of US 19/98 and one is on the north side of the CFG, midway between I-75 and US 441. In  
 7 addition, there are four sand mines within three miles of the CFG. It is reasonable to assume there  
 8 are mineable sand deposits on the CFG, given the nature of sediments and the close proximity of other  
 9 sand mining operations.

10 **Hydrology**

11 The CFG is hydrologically complex, highly manipulated, and diverse. It is a unique park in many  
 12 respects, especially since surface hydrology of major rivers can be actively managed/manipulated  
 13 via the lock and dam systems that exist on the east and west ends of the CFG.

14 The CFG crosses through portions of four river basins: Withlacoochee (HUC 03100208), Crystal-  
 15 Pithlachascotee (HUC 03100207), Ocklawaha (03080102), and Lower St. Johns (HUC 03080103).  
 16 Most of the CFG intersects with the Ocklawaha River, Withlacoochee River, and below ground,  
 17 portions of the Floridan Aquifer.

18 From jurisdictional and management perspectives, the St. Johns River Water Management District  
 19 (SJRWMD) covers the eastern half of the CFG, while the SWFWMD covers the remaining areas. Florida

1 designates its surface waters into one of five categories, each with a designated use. The majority of  
2 the CFG is considered Class III: recreation, propagation, and maintenance of a healthy, well-balanced  
3 population of fish and wildlife. At the western end of the CFG, two areas are designated as Class II  
4 (shellfish propagation or harvesting).

5 Construction of the Cross Florida Barge Canals resulted in significant alterations of and impacts to  
6 the immediate and adjacent landscape. These land-use actions, which occurred in the mid to latter  
7 parts of the 20<sup>th</sup> Century, have had a lasting effect on aquatic resources throughout the CFG. Despite  
8 the degree of immediate impact, portions of the CFG still are characterized by relatively intact aquatic  
9 ecosystems. For example, the Cross Florida Barge Canal was cleared and grubbed along its entire  
10 alignment, but not dredged upstream of Kenwood. Conversely, the Ocklawaha River still flows along  
11 the original channel in most locations.

12 The Ocklawaha River system arises from the Green Swamp, Lake Apopka and the Harris Chain of  
13 Lakes (Lake Griffin, Lake Harris, Little Lake Harris Lake Dora and Lake Beauclair) and flows north,  
14 forming the Upper Ocklawaha River. Other aquatic ecosystems that contribute to the headwaters of  
15 the Upper Ocklawaha River include Marshall Swamp and Adams Marsh. The CFG encompasses the  
16 Ocklawaha River from this point downstream to the Kirkpatrick Dam. The Ocklawaha River flows  
17 north, where it is joined by the Silver River from the west near SR 40. The Silver River is a spring-run  
18 stream flowing from Silver Springs and it represents the largest tributary of the Ocklawaha River.  
19 Orange Creek, Deep Creek, and Camp Branch Creek are other important surface waters that flow into  
20 the Ocklawaha River. The Ocklawaha River ultimately flows into the St. Johns River near Satsuma.

21 Construction of the Cross Florida Barge Canal and other related activities radically changed parts of  
22 the Ocklawaha River and associated riparian areas; 7,500 acres of floodplain forest associated with  
23 the Ocklawaha River were flooded, along with one second-magnitude (Blue Spring) and 19 third-  
24 magnitude springs and associated habitats along the river. The 3,000-acre (relatively intact) Adams  
25 Marsh (within Marshall Swamp) is impounded by a perimeter dike about nine miles long. From Silver  
26 Springs north to the Eureka Lock, the Ocklawaha River is essentially unaltered. The Eureka Dam and  
27 Lock are still in place, but have never been operational. North of Eureka, the Ocklawaha River retains  
28 much of its natural characteristics. Approaching Orange Springs, the Ocklawaha River displays more  
29 of the characteristics of Rodman Reservoir (also known as Lake Ocklawaha), a consequence of the  
30 Kirkpatrick Dam. As the Rodman Reservoir makes its turn to the east, Orange Creek flows in from the  
31 west. Farther east, Deep Creek and Sweetwater Creek flow into the reservoir. Much of Sweetwater  
32 Creek and its associated riparian areas are found within the CFG. East of the reservoir, the east Barge  
33 Canal extends about nine miles to the east-northeast, where it joins the St. Johns River. This part of  
34 the canal bisected the Camp Branch Creek, disrupted the natural surface flows, and altered the Cow  
35 Heaven Bay Swamp connection with the St. Johns River.

36 Buckman Lock (still operational) is located within the eastern canal. From this point, CFG  
37 landholdings extend about five-miles northeast along the St. Johns River. Historically, the Ocklawaha  
38 flowed east-southeast from the location of the Kirkpatrick Dam for approximately nine miles, where  
39 it joined the St. Johns River. The flow has been disrupted by the dam and water is released by a  
40 spillway into the man-made tailrace and the historical Ocklawaha River channel.

41 CFG staff maintain the Buckman Lock and Kirkpatrick Dam and Spillway. The Buckman Lock controls  
42 access to Rodman Reservoir from the St. Johns River through the east barge canal. The Kirkpatrick

1 Dam spillway controls the level of Rodman Reservoir. Generally, the water level of the reservoir is  
2 kept at 18 feet to 20 feet National Geodetic Vertical Datum (NGVD) 1929 level. Drawdown occurs  
3 every three to four years, which de-waters approximately 65 percent of the reservoir.

4 The Withlacoochee River is the main river system on the western end of the CFG. A small portion of  
5 the CFG abuts the Withlacoochee River near Dunnellon, where the Rainbow River, arising from a first-  
6 magnitude spring, flows into the river. Downstream of the confluence with the Rainbow River, the  
7 Withlacoochee River is dammed. Dam construction was completed in the early 1900's and resulted  
8 in Lake Rousseau (also known as Withlacoochee Backwaters)—a waterbody utilized for generating  
9 electric power. Lake Rousseau is no longer used for power generation. Downstream of Lake  
10 Rousseau, the CFG borders the south bank of the Withlacoochee River in several places all the way to  
11 the Gulf of Mexico.

12 Current water control structures associated with Lake Rousseau include the Inglis Dam and Spillway,  
13 Inglis Lock (as part of the western Barge Canal), and the Inglis Bypass Canal and Spillway. The bypass  
14 canal funnels water from Lake Rousseau just east of the lock to a spillway that provides water to the  
15 lower reaches of the Withlacoochee River. The Inglis Lock is no longer operational due to its  
16 deteriorated condition. The SWFWMD operates the western Barge Canal water control structures  
17 (dams and spillways) under a contract with and funding through FDEP. FDEP is currently responsible  
18 for maintenance and operation of the Inglis Lock. The Lake Rousseau water level generally is  
19 maintained at a fixed elevation of 27.5 NGVD. In times of heavy rainfall, additional water can be  
20 released to prevent or minimize localized flooding.

21 The western Barge Canal is about nine miles long and extends from near the western end of Lake  
22 Rousseau into the Gulf of Mexico for about 10.5 miles. The western Barge Canal cuts through the  
23 lower reaches of the Withlacoochee River between the Inglis Bypass Spillway and the western end of  
24 Lake Rousseau. Inglis Island, formerly bordered on the north side of Lake Rousseau and the  
25 Withlacoochee River, was surrounded by water by the canal being cut through on the north side of  
26 the land mass.

27 The CFG is not directly associated with any riverine systems from east of Dunnellon to Marshall  
28 Swamp. However, some significant wetlands are present in the Ross Prairie/Halpata Tastanaki  
29 Preserve area. Segments of old ship canal diggings also are in the stretch from the Pruitt Trailhead  
30 east to I-75.

### 31 *Water Quality*

32 Water quality within the CFG is influenced to a large degree by sources and land uses occurring  
33 outside of its boundaries. Water quality is tracked and managed at multiple levels within the state.  
34 For example, at the macro-scale, Basin Management Action Plans (BMAPs) have been established  
35 across Florida and serve as a "blueprint" for restoring impaired waters by reducing pollutant  
36 loadings to meet the allowable loadings established in a Total Maximum Daily Load (TMDL).  
37 Individual BMAPs represent a comprehensive set of strategies such as permit limits on wastewater  
38 facilities, urban and agricultural best management practices (BMPs), and conservation programs.  
39 Within and proximal to the CFG, six BMAPS have been established (Silver Springs, Kings Bay, Lower  
40 St. Johns main stem, Silver River, Rainbow Springs, and Rainbow River). Also within this same  
41 geographic area, seven impaired waterbodies have been identified (Withlacoochee River, Lake  
42 Rousseau Drain, Rainbow River Run, Ocklawaha River above Daisy Creek, Daisy Creek, and Little



1 Orange Creek) and two TMDLs have been established for the St. Johns River above Dunns Creek and  
2 Upper Silver River.

3 At the local scale, FDEP, USGS, and the SJRWMD routinely collect water quality samples in the  
4 Ocklawaha River Basin. Water quality at the SR 316 and SR 40 stations in Marion County is  
5 considered good according to the SJRWMD's water quality index. Orange Creek and the Silver River  
6 are each monitored near their confluences with the Ocklawaha River. Water quality at both of these  
7 monitoring sites is considered good according to the water quality index. Additional water quality  
8 information for the Ocklawaha River is available via multiple websites (need to verify and/or update  
9 links): and for Orange Creek.

10 Construction, shoreline alterations, and failing septic tanks contribute sediment, nutrient, and  
11 bacteria inputs to Lake Rousseau. Aquatic weed growth within Lake Rousseau is a problem, which is  
12 managed primarily via herbicide treatments by FWC. Downstream of Lake Rousseau, the  
13 Withlacoochee River has similar pollution sources.

14 The SJRWMD has several groundwater observation wells located both on and within one mile of the  
15 CFG. Selected wells provide daily observations, and the district and the USGS conduct potentiometric  
16 readings twice a year. The SJRWMD portion of the CFG has little groundwater development,  
17 especially in river areas. Sampling in the Summit Reach area near Ocala shows that nitrates have been  
18 steadily increasing over the years. The Ocala area is an important water recharge area for the  
19 Floridan Aquifer. The western end of the CFG is in an area generally with an unconfined Floridan  
20 Aquifer, except near the Withlacoochee River. Groundwater recharge rates are high, water  
21 withdrawal is low, and the level of nitrates in groundwater has increased over time.

#### 22 *Other Supporting Information*

23 Starting in the 1970's, numerous public interest groups have recommended the removal of the  
24 Kirkpatrick Dam due to impacts the Rodman Reservoir has had on the Ocklawaha River, its  
25 floodplain, and associated ecosystems.

26 Creation of the Kirkpatrick Dam has caused flooding within a portion of Ocala National Forest.  
27 Incidental flooding has been allowed under a special-use permit from the U.S. Department of  
28 Agriculture, Forest Service (first issued in 1994). The U.S. Forest Service included conditions about  
29 the issuance of the new permit, containing a schedule for the reservoir to be drawn down and the  
30 dam to be breached/disabled.

31 In addition to obvious hydrological disruptions resulting from Barge Canal activities, smaller-scale,  
32 but significant, hydrological disruptions are present on the CFG. Roads, planting beds, ditches, and  
33 firelines are responsible for many of these other disturbances.

#### 34 *Management Recommendations*

35 Managing water levels within Rodman Reservoir follows procedures and protocols established by  
36 professionals, such as civil engineers, fisheries and aquatic plant scientists, hydrologists, and  
37 geologists. A discussion of the technical aspects of those procedures and protocols is not germane to  
38 this UMP. With respect to other aspects of hydrologic/aquatic resources within the CFG, it is  
39 recommended that resource staff continue to follow Florida Silviculture Best Management Practices,  
40 as well as agricultural BMPs (research has consistently shown that properly applied BMPs protect

1 water quality). Aspects of land management having an immediate and recurring impact include soil  
2 erosion/sedimentation control measures related to dirt roads and firebreaks and proper use of  
3 herbicides in aquatic environments.

#### 4 **Natural Communities**

##### 5 *Introduction*

6 This section of the unit management plan describes and assesses each of the natural communities or  
7 land cover found on the CFG. Information on the CFG natural communities was obtained from the  
8 *2007-2008 FNAI Natural Communities Survey Report*. Land management and restoration is best  
9 guided by clear and specific ecological goals and/or desired future conditions. This section describes  
10 the desired future condition of each natural community or provides a description of how each  
11 community should look in the future. There are many values of identifying a desired future condition.  
12 These include: (1) providing a vision of future conditions that can be communicated to staff,  
13 stakeholders and the public, (2) guiding conservation and management actions, (3) providing spatial  
14 and temporal priorities for management and conservation, and (4) integrating monitoring and  
15 adaptive management into natural resource management. This section also discusses the  
16 management actions required to bring each community to its desired future condition. Specific  
17 management objectives and actions for natural community management, exotic species  
18 management, imperiled species management, and restoration are discussed in the Resource  
19 Management Program section of this component.

20 The system of classifying natural communities utilized in this plan was developed by the Florida  
21 Natural Areas Inventory (FNAI). Natural Communities are characterized and defined by a  
22 combination of physiognomy, vegetation structure and composition, topography, land form,  
23 substrate, soil moisture condition, climate, and fire. They are named for their most characteristic  
24 biological or physical feature (FNAI and FDEP, 2010). FNAI uses several criteria, including area  
25 covered and number of occurrences, to determine the relative rarity and threat to each community  
26 type; these are summarized into a global rank and a state rank, the G and S ranks listed for each  
27 community, respectively. Table 8 provides the ranking for each vegetative community on the CFG.

28 When a natural community within a park reaches the ideal future condition, it is considered to be in  
29 a “maintenance condition.” Required actions for sustaining a community’s maintenance condition  
30 may include maintaining optimal fire return intervals for fire-dependent communities, ongoing  
31 control of non-native plant and animal species, maintaining natural hydrological functions (including  
32 historic water flows and water quality), preserving a community’s biodiversity and vegetative  
33 structure, protecting viable populations of plant and animal species (including those that are  
34 imperiled or endemic), and preserving intact ecotones linking natural communities across the  
35 landscape.

36 The CFG extends from the Gulf of Mexico on the west to the St. Johns River on the east, and it  
37 crosses four major landscapes, including: Withlacoochee Coastal Lowlands, Ocala Uplands,  
38 Ocklawaha River Valley, and the Etoniah Basin.

39 The key ecological associations for each of these landscapes are:



- 1 • *Withlacoochee Coastal Lowlands*—associations include hydric hammock/tidal  
2 marsh/freshwater tidal swamp/wet flatwoods, wet flatwoods/tidal marsh, wet  
3 flatwoods/sandhill, and mesic flatwoods/ sandhill.
- 4 • *Ocala Uplands*—associations include sandhill/ scrub/upland pine forest (red oak),  
5 sandhill/xeric hammock/wet prairie, and wet flatwoods/mesic flatwoods/depression  
6 marsh.
- 7 • *Ocklawaha River Valley*—associations include floodplain swamp/bottomland  
8 forest/depression marsh/basin marsh/mesic flatwoods/ wet flatwoods, and  
9 scrub/sandhill.
- 10 • *Etoniah Basin*—associations include floodplain swamp/basin swamp/mesic  
11 flatwoods/wet flatwoods, and scrub/sandhill.

12 Outstanding natural systems of the greenway include springs and rivers, lakes, swamps, prairies,  
13 freshwater and saltwater marshes, and uplands. Many of the natural communities on the greenway  
14 represent a true cross section of natural Florida, with a possible 26 of FNAI's 82 natural communities  
15 in the region.

16 The greenway is home to a variety of wildlife, including indicator species such as the Florida black  
17 bear, gopher tortoise, Bald Eagle, Florida Scrub-Jay, and Wood Stork. Important black bear corridors  
18 exist throughout the greenway. The greenway affords the opportunity to study exotic species  
19 throughout its length; to evaluate disturbed hydrologic systems in the canals and river corridors; and  
20 to analyze impacts from urban and development pressures—for example, from domestic pets—on  
21 the natural systems in the Ocala Upland, and from development pressures on the Ocklawaha and  
22 Withlacoochee Rivers.

23 As illustrated below in Table 8, a total of 64 percent—or 45,663 acres—of the CFG can be  
24 classified as natural communities, while 24,868 acres—or 36 percent—can be classified as an  
25 altered land cover type. Major natural communities represented based on acreage on the CFG  
26 include sandhills, mesic flatwoods, wet flatwoods, scrub, upland mixed forest, and  
27 bottomlands. These communities total approximately 50 percent of the CFG acreage, and two-  
28 thirds of the natural community coverage on the CFG.

29 **Table 8. CFG Community Type, Acreage, and Management**

Community Type	FNAI Community Ranking		CFG Amount		Fire Return Interval
	Global	State	Acres	Percent of Total	
<b>Fire-Dependent Communities</b>					
Sandhill	G3	S2	6,408	9%	1–3 years (3)
Mesic Flatwoods	G4	S4	5,041	7%	1–4 years (3)
Wet Flatwoods	G4	S4	1,773	3%	5–10 years (7)
Scrub	G2	S2	1,630	2%	5–20 years (5)
Scrubby Flatwoods	—	—	683	1%	3–14 years (10)
Floodplain Marsh	—	—	245	0.3%	2–5 years

Community Type	FNAI Community Ranking		CFG Amount		Fire Return Interval
	Global	State	Acres	Percent of Total	
Basin Marsh	—	—	14	0.1%	2–10 years (5)
Upland Pine Forest	—	—	6	0.1%	1–3 years
<b>Total Fire-Dependent Communities</b>			<b>15,800</b>	<b>22%</b>	
<b>Altered Land Cover Fire-Type Acres</b>					
Clear Cut Pine Plantations	—	—	161	0.2%	3 years
Pine Plantation	—	—	9,071	13%	3 years
<b>Total Altered Land Cover Fire-Type Acres</b>			<b>9,232</b>	<b>13%</b>	
<b>Total Fire-Dependent/ Altered Land Cover Fire-Type Acres</b>			<b>25,032</b>	<b>35%</b>	
<b>Other Natural Communities (Non-Fire Type)</b>					
Basin Swamp	—	—	3,388	5%	
Baygall	G4	S4	549	1%	
Blackwater Stream	—	—	192	0.2%	
Bottomland Forest	—	—	1,258	2%	
Depression Marsh	G4	S4	735	1%	
Dome Swamp	G4	S4	298	0.3%	
Floodplain Swamp	—	—	10,497	15%	
Hydric Hammock	—	—	3,663	5%	
Upland Hardwood Forrest	—	—	350	0.3%	
Xeric Hammock	G3	S3	307	0.3%	
Mesic Hammock	—	—	4,313	6%	
Swamp Lake	—	—	28	0.1%	
Salt Marsh	—	—	4,285	6%	
<b>Total Other Natural Communities (Non-Fire Type)</b>			<b>29,863</b>	<b>42%</b>	
<b>Other Altered Land Cover Types</b>					
Abandoned Field	—	—	362	0.3%	
Abandoned Pasture	—	—	116	0.1%	
Agriculture	—	—	47	0.1%	
Canal/Ditch	—	—	3,423	5%	
Clearing	—	—	1,516	2%	
Developed	—	—	756	1%	
Impound-ment/ Artificial Ponds	—	—	8,270	12%	
Invasive Exotic Subculture	—	—	35	0.1	

Community Type	FNAI Community Ranking		CFG Amount		Fire Return Interval
	Global	State	Acres	Percent of Total	
Pasture Improved	—	—	313	0.3%	
Pasture Semi-Improved	—	—	85	0.1%	
Road	—	—	250	0.2%	
Spoil Area	—	—	204	0.2%	
Utility Corridor	—	—	259	0.2%	
<b>Total Altered Land Cover Types</b>			<b>15,636</b>	<b>22%</b>	
<b>Total CFG</b>			<b>70,531</b>	<b>100%</b>	

Source: CFG, 2016.

1

# Sandhills

**Global and State Rank: G2/S2**


## Desired Future Condition

Dominant pines usually will be longleaf pine (*Pinus palustris*). Herbaceous cover is 80 percent or greater, typically of wiregrass (*Aristida stricta*), and is less than three feet in height. In addition to groundcover and pine characteristics, there will be scattered individual trees, clumps, or ridges of onsite oak species (usually turkey oaks (*Quercus laevis*), sand post oak (*Quercus margaretta*), and blue-jack oak (*Quercus incana*)). In old-growth conditions, sand post oaks commonly are 150 to 200 years old, and some turkey oaks are more than 100 years old. The optimal fire return interval for this community is one year to three years.

## Description and Assessment

In total, as demonstrated in Table 8, the Sandhill community at CFG is comprised of 6,408 acres, or 9 percent of CFG total area. The highest-quality sandhills on the CFG are found south of CR 484 and east and west of SR 200. Other good examples of sandhills are located in the Caravelle Wildlife Management Area; north and south of the Buckman Lock; west of the Ocklawaha River in the vicinity of Country Club Boulevard; Baseline Road to Marshall Swamp Trailhead; the western edge of Marshall Swamp; north and south of CR 464, from US 301 west to I-75; in the Diggings from I-75 west of SR 200; in the area east of US 41 in Dunnellon; and on Inglis Island.

The overstory of sandhills on the CFG varies from open to semi-closed depending on history and frequency of fire. The canopy is primarily longleaf pine. In some areas, sand pine (*Pinus clausa*), turkey oak, laurel oak (*Quercus hemisphaerica*), or sand live oak (*Quercus geminata*) can reach into the canopy as well. Many areas on the CFG have experienced long-term fire exclusion, and a dense midstory of hardwoods has developed. Typically, these are the aforementioned oaks and pines, as well as common persimmon (*Diospyros virginiana*), black cherry (*Prunus serotina*), and sand post oak. Typical shrubs are slimleaf pawpaw (*Asimina angustifolia*), netted pawpaw (*Asimina*



**Global and State Rank: G2/S2**



**Fire Interval: 1–3 years**

## Listed Species:

Eastern indigo snake, gopher tortoise, longspurred mint, Garberia, sandhill spiny pod, giant orchid, angle pod, leafless beaked orchid, scrub stylisma, spiny pod, incised groove-bur, sand butterfly pea, scrub buckwheat, Lewton’s polygala

## Invasive Exotic Species:

Air-potato, camphor tree, Chinaberry, Chinese Brake Fern, Chinese Tallow tree, Chinese Wisteria, cogon grass, confederate jasmine, crapemyrtle, English ivy, flamegold, formosa firethorn, golden bamboo, Japanese climbing fern, Japanese honeysuckle, lantana, mimosa, natal grass, pampas grass, paper mulberry, ravenna grass, sago palm, silverthorn, sweet viburnum

## Management Practices:

- Prescribed fire
- Invasive exotic treatment/control



1 *reticulata*), Michaux's hawthorn (*Crataegus michauxii*), dwarf huckleberry (*Gaylussacia dumosa*),  
2 sand holly (*Ilex ambigua*), gopher apple (*Licania michauxii*), Carolina laurelcherry (*Prunus*  
3 *caroliniana*), flatwoods plum (*Prunus umbellata*), Chapman's oak (*Quercus chapmanii*), bluejack oak  
4 (*Quercus incana*), winged sumac (*Rhus copallinum*), sassafras (*Sassafras albidum*), saw palmetto  
5 (*Serenoa repens*), and dwarf live oak (*Quercus minima*).

6 A few areas of sandhills on the CFG contain southern red oak (*Quercus falcata*) and flowering  
7 dogwood (*Cornus florida*). Typically, these species are associated with upland pine forest  
8 communities, which represent a similar community that occurs on slightly richer, more clayey soils,  
9 (primarily occurring in northern Florida). Only one small area (six acres) is delineated for the upland  
10 pine forest (red oak) on the CFG, although a larger area is noted in the historic vegetation map (320  
11 acres). Possible remnants of upland pine forests are within the sandhills west of US 441 and within a  
12 ruderal area (historic sandhill) north of CR 464 in the Baseline Road/Marshall Swamp Recreation  
13 Trail.

14 The herbaceous layer of the sandhills at CFG offers a high diversity of species. Grasses include Elliott's  
15 bluestem (*Andropogon gyrans*), splitbeard bluestem (*Andropogon ternarius*), broomsedge bluestem  
16 (*Andropogon virginicus*), arrowfeather threeawn (*Aristida purpurascens*), bottlebrush threeawn  
17 (*Aristida spiciformis*), wiregrass, bearded skeletongrass (*Gymnopogon ambiguus*), pinewoods  
18 dropseed (*Sporobolus junceus*), little bluestem (*Schizachyrium scoparium*), needleleaf witchgrass  
19 (*Dichanthelium aciculare*), perennial sandgrass (*Triplasis americana*) and lopsided indian grass  
20 (*Sorghastrum secundum*).

21 Typical forbs in the Sandhills community include yellow false foxglove (*Aureolaria pedicularia* var.  
22 *pectinata*), coastalplain honeycomb-head (*Balduina angustifolia*), oblongleaf twinflower (*Dyschoriste*  
23 *oblongifolia*), Florida greeneyes (*Berlandiera subacaulis*), Elliott's milkpea (*Galactia elliottii*),  
24 shortleaf gayfeather (*Liatris tenuifolia*), skyblue lupine (*Lupinus diffusus*), sweet goldenrod (*Solidago*  
25 *odora*), chaffhead (*Carphephorus corymbosus*), Florida alicia (*Chapmannia floridana*), devil's  
26 grandmother (*Elephantopus tomentosus*), wild buckwheat (*Eriogonum tomentosum*), narrowleaf  
27 silkgrass (*Pityopsis graminifolia*), bracken fern (*Pteridium aquilinum*), manyflower beardtongue  
28 (*Penstemon multiflorus*), Feay's palafox (*Palafoxia feayi*), sandyfield beaksedge (*Rhynchospora*  
29 *megalocarpa*), Carolina wild petunia (*Ruellia caroliniensis*), Piedmont blacksenna (*Seymeria*  
30 *pectinata*), tread softly (*Cnidioscolus stimulosus*), silver croton (*Croton argyranthemus*), pineland  
31 croton (*Croton linearis*), ticktrefoil (*Desmodium* sp.), whitetop aster (*Sericocarpus tortifolius*), and  
32 camphorweed (*Heterotheca subaxillaris*). Vines tend to have a patchy distribution and primarily  
33 include Virginia creeper (*Parthenocissus quinquefolia*), earleaf greenbrier (*Smilax auriculata*), and  
34 muscadine (*Vitis rotundifolia*). In areas where fire has been absent for long periods, ground lichens  
35 (*Cladina evansii*, *Cladina subtenuis*, and *Cladonia leporina*) are common.

36 Fire exclusion and the resulting woody encroachment represent the most prevalent departure from  
37 desired future conditions in the sandhills throughout the CFG. Invasive exotic plants and  
38 disturbances (historic or routinely recurring) including clearing, ditches and canals (“the Diggings”),  
39 forestry operations, agriculture operations to include cattle grazing and watering, roads, off-road  
40 vehicle (ORV) trails, firebreaks and development account for most of the remaining impacts.

41 Two listed animal species—eastern indigo snake (*Drymarchon couperi*) and gopher tortoise  
42 (*Gopherus polyphemus*)—were observed in the sandhills on CFG during a survey conducted by FNAI

1 from 2007 to 2008. The following listed plants were identified during the surveys: longspurred mint  
2 (*Dicerandra cornutissima*), garberia (*Garberia heterophylla*), giant orchid, angle pod, leafless beaked  
3 orchid, scrub stylisma, spiny pod, and sandhill spiny pod (*Matelea pubiflora*).

4 Invasive plant species observed within the sandhills during these surveys conducted by FNAI were:  
5 air-potato (*Dioscorea bulbifera*), camphor tree (*Cinnamomum camphora*), Chinaberry (*Melia*  
6 *azedarach*), Chinese brake fern (*Pteris vittata*), Chinese tallow tree (*Triadica sebifera*), Chinese  
7 wisteria, cogon grass (*Imperata cylindrica*), confederate jasmine, crapemyrtle, English ivy, flamegold,  
8 formosa firethorn, golden bamboo, Japanese climbing fern, Japanese honeysuckle, lantana, mimosa  
9 (*Albizia julibrissin*), natal grass (*Rhynchelytrum repens*), pampas grass, paper mulberry, ravenna  
10 grass, sago palm, silverthorn, sweet viburnum.

## 11 **Fire Regime**

12 Frequent ground fires reduce hardwood competition and promote the growth and development of  
13 forbs, grasses, and predominant (or preferred) overstory species—longleaf pine. The natural fire  
14 frequency for the sandhills is every one year to three years, although some sites may be maintained  
15 with fire intervals up to five years. Naturally, fires would ignite principally during the early summer  
16 (April to June) when lightning strikes are frequent and fuels are not yet saturated by afternoon rains.  
17 Without frequent fires, the sandhills may eventually succeed to xeric hammock.

## 18 **General Management Measures**

19 Management activities in the sandhills on CFG should focus on prescribed burning, minimizing  
20 practices that disturb the soil, and restoring areas that are densely encroached with oaks or sand  
21 pines. When conditions allow, prescribed burning alone is the preferred method to reduce woody  
22 species abundance in the understory. Roller-chopping, particularly in xeric soil types, can be  
23 detrimental to herbaceous species, especially wiregrass. A chemical treatment may be appropriate  
24 in some areas of extreme oak encroachment. Hand felling and removal of very dense oaks prior to  
25 burning may be most effective and create the least amount of soil disturbance. In many situations,  
26 prescribed burning during early summer should provide the greatest benefit in reducing woody  
27 species abundance. Diligence in monitoring and controlling invasive exotic plants in the sandhills  
28 also is necessary.

29 During all management activities, every effort should be made to minimize detrimental effects to the  
30 gopher tortoise population (and its burrows) and to existing longspurred mint growth.



# 1 Scrub

2 **Global and State Rank: G2/S2**


3 **Desired Future Condition**

4 Dominant species over the vast majority of scrub include  
5 sand live oak (*Quercus geminata*), myrtle oak (*Quercus*  
6 *myrtifolia*), Chapman's oak (*Quercus chapmanii*), saw  
7 palmetto (*Serenoa repens*) and rusty staggerbush (*Lyonia*  
8 *ferruginea*). Scrub oak canopy will vary in height from  
9 three feet to eight feet. There is a variety of oak age classes  
10 and heights between different scrub patches. There are  
11 scattered openings in the canopy, with bare patches of  
12 sand that support many imperiled or endemic plant  
13 species; these species are regularly flowering and  
14 replenishing their seed banks. Sand pine (*Pinus clausa*),  
15 where present, usually is not dominant in abundance,  
16 percent cover, or height. The optimal fire return interval  
17 for this community is regionally variable; typically, five  
18 years to 20 years when aiming to achieve a mosaic of  
19 burned and unburned areas.

20 **Description and Assessment**

21 In total, as shown in Table 8, the scrub communities on the  
22 CFG are comprised of 1,630 acres, or 2 percent of the CFG  
23 total area. Scrub is centered in Florida, but extends  
24 westward on barrier islands to Alabama and Mississippi,  
25 and small patches are found northward into southeastern  
26 Georgia. In Florida, scrub tends to be distributed in long,  
27 narrow ridges parallel to coastlines and is scarce or absent  
28 from the limestone-dominated southernmost portion of  
29 the state.

30 The highest-quality scrub on the CFG occurs within the  
31 Triangle and the Diggings, within close proximity to  
32 I-75. Scrub also occurs in the Caravelle Wildlife  
33 Management Area, specifically, north of the Buckman Lock;  
34 north of the Rodman Reservoir; and east of Deep Creek.  
35 Other areas include the Eureka East Recreation site, in the  
36 area north of CR 314, west end of the CFG south of the  
37 Withlacoochee River, south of the Ocklawaha River, in the  
38 area of I-75 west (the Triangle), and in the Diggings east of  
39 SR 200.



**Global and State Rank:  
G2/S2**



**Fire Interval: 5–20 years**

### Listed Species:

Florida Scrub-Jay, gopher tortoise, longspurred mint, garberia, giant orchid, Chapman's skeletongrass, sand butterfly pea

### Invasive Exotic Species:

Mimosa, cogon grass, natal grass, Caesar's weed, Chinese tallow tree, pampas grass

### Management Practices:

- Prescribed fire
- Mechanical treatments
- Invasive exotic treatment/control



1 The scrub communities on the CFG typically have an open canopy of sand pine and sand live oak. The  
2 subcanopy is moderately dense and includes sand pine, sand live oak, rusty staggerbush, Chapman's  
3 oak, laurel oak (*Quercus hemisphaerica*), and myrtle oak. Shrubs are dense and diverse, but at times  
4 can be patchy, with white sand between the patches. Shrubs include the aforementioned overstory  
5 species and Florida rosemary (*Ceratiola ericoides*), garberia (*Garberia heterophylla*), blue  
6 huckleberry (*Gaylussacia frondosa* var. *tomentosa*), sand holly (*Ilex ambigua*), gopher apple (*Licania*  
7 *michauxii*), rusty staggerbush, fetterbush (*Lyonia lucida*), scrub wild olive (*Osmanthus megacarpus*),  
8 silk bay (*Persea borbonia* var. *humilis*), black cherry (*Prunus serotina*), turkey oak (*Quercus laevis*),  
9 scrub palmetto (*Sabal etonia*), saw palmetto, gum bully (*Sideroxylon lanuginosum*), tough bully  
10 (*Sideroxylon tenax*), sparkleberry (*Vaccinium arboreum*), Darrow's blueberry (*Vaccinium darrowii*),  
11 shiny blueberry (*Vaccinium myrsinites*), and deerberry (*Vaccinium stamineum*).

12 Herbs are sparse and occur primarily in the open sandy patches. Typically seen are broomsedge  
13 bluestem (*Andropogon virginicus*), bottlebrush threeawn (*Aristida spiciformis*), coastalplain  
14 honeycomb-head (*Balduina angustifolia*), capillary hairsedge (*Bulbostylis ciliatifolia*), Florida alicia  
15 (*Chapmannia floridana*), Feay's prairie clover (*Dalea feayi*), summer farewell (*Dalea pinnata*), rough  
16 hedgehyssop (*Gratiola hispida*), shortleaf gayfeather (*Liatis tenuifolia*), prickly pear (*Opuntia*), wild  
17 pennyroyal (*Piloblephis rigida*), narrowleaf silkgrass (*Pityopsis graminifolia*), racemed milkwort  
18 (*Polygala polygama*), sandyfield beaksedge (*Rhynchospora megalocarpa*), sand spike-moss  
19 (*Selaginella arenicola*), and pineland scalypink (*Stipulicida setacea*). Vines include earleaf greenbrier  
20 (*Smilax auriculata*) and muscadine (*Vitis rotundifolia*). Ground lichens (*Cladonia* spp. and *Cladina*  
21 spp.) are present. Fire exclusion and resulting woody encroachment represent the most serious  
22 disturbances in the scrub communities on the CFG. Other more-minor disturbances include clearing,  
23 forestry operations, ORV trails, firebreaks, and exotic plant introduction.

24 Listed species occur in the scrub communities of the CFG. Two listed animal species—the Florida  
25 Scrub-Jay (*Aphelocoma coerulescens*) and gopher tortoise (*Gopherus polyphemus*)—were  
26 documented during the 2007 and 2008 FNAI survey. Five listed plants—longspurred mint  
27 (*Dicerandra cornutissima*), garberia (*Garberia heterophylla*), Chapman's skeletongrass (*Gymnopogon*  
28 *chapmanianus*), sand butterfly pea (*Centrosema arenicola*), and giant orchid (*Pteroglossaspis*  
29 *ecristata*)—also were documented.

30 Six invasive plant species were recorded within the scrubs on the CFG: mimosa (*Albizia julibrissin*),  
31 cogon grass (*Imperata cylindrica*), natal grass (*Rhynchelytrum repens*), Caesar's weed (*Urena lobate*),  
32 Chinese tallow (*Triadica sebifera*) and pampas grass (*Cortaderia selloana*).

### 33 **Fire Regime**

34 Oak scrub, which has been the subject of considerable research in the peninsula, is thought to have a  
35 range of natural fire return intervals that are considerably shorter (five years to 20 years) than those  
36 of sand pine (20 years to 80 years) or rosemary scrub (10 years to 40 years). Sand pine, if present in  
37 the area at all, will invade and overtop oak scrub if left unburned for long periods of time.

## 1 **General Management Measures**

2 The primary management of scrub on the CFG should be guided by habitat requirements of the  
3 Florida Scrub-Jay. On a case-by-case basis, some special management considerations may be added  
4 to this goal to protect and maintain longspurred mint and the gopher tortoise.

5 Optimal Florida Scrub-Jay habitat consists of low oak shrubs (three feet to six feet tall) interspersed  
6 with numerous patches of exposed sand representing greater than 15 percent of the area. Jays need  
7 the bare sandy soil to bury and recover their annual cache of acorns. A mosaic of scrubs with various  
8 heights is ideal, but across all scrub in an area where Florida Scrub-Jays reside or are wanted, the  
9 general goal should be maintenance of 70 percent of all territories (or potential territories) at an  
10 optimal height for Jays of approximately three feet to six feet. The majority of the remaining 30  
11 percent should be in the lower height range (i.e., just burned and re-sprouting scrub vegetation less  
12 than four feet in height), with a very small percentage, if any, of the scrub landscape taller than six  
13 feet.

14 Florida Scrub-Jays are permanently territorial and do not abandon a breeding territory except under  
15 rare circumstances. To avoid burning all of one territory in one fire, it is important to conduct mosaic  
16 burns in occupied territories. A specific Florida Scrub-Jay habitat management plan may be needed  
17 to effectively implement and guide the timing and extent of management actions. Effective burning  
18 may require low humidity conditions, which presents challenges for the prescribed burner.  
19 Mechanical treatments in scrub can be used to facilitate burning under more controlled weather  
20 conditions. Mechanical reduction of shrub stature and/or removal of sand pines prior to burning are  
21 sometimes needed to reduce fuel levels.

22 Scrub fire regimes are highly variable, depending on landscape settings. Some older references  
23 recommend a fire return interval of 20 years or more; however, more current scientific research  
24 suggests most scrubs would have naturally burned more frequently. A return interval between four  
25 years and 10 years may be necessary to maintain shrub heights within the range favorable to the  
26 Florida Scrub-Jay. Intervals vary depending on site conditions, which will affect vegetation growth  
27 rates, and on the ideal vegetation structure.

28 The invasive exotic plant cogon grass threatens the state-listed longspurred mint in some areas;  
29 vigilant treatment and monitoring of cogon grass in these areas will be required to protect this rare  
30 mint. In certain locations, the source for this invasive exotic is on adjacent Department of  
31 Transportation lands, and, thus, effective treatment will require coordination with that agency.

# 1 Xeric Hammock

2 **Global and State Rank: G3/S3**


3 **Desired Future Condition**

4 Typically considered a late successional stage of scrub or  
5 sandhill that generally occurs in small isolated patches on  
6 excessively well drained soils. Vegetation will consist of a  
7 low, closed canopy dominated by sand live oak (*Quercus*  
8 *geminata*), which provides shady conditions. Typical plant  
9 species also may include Chapman's oak (*Quercus*  
10 *chapmanii*) and laurel oak (*Quercus laurifolia*). Slash pine  
11 or longleaf pine (*Pinus elliottii*, *Pinus palustris*,  
12 respectively) also may be a minor component. The  
13 understory species will include saw palmetto (*Serenoa*  
14 *repens*), fetterbush (*Lyonia lucida*), rusty staggerbush  
15 (*Lyonia ferruginea*), myrtle oak (*Quercus myrtifolia*),  
16 yaupon holly (*Ilex vomitoria*), Hercules' club (*Zanthoxylum*  
17 *clava-herculis*), and Florida rosemary (*Ceratiola ericoides*).  
18 A sparse groundcover layer of wiregrass (*Aristida stricta*)  
19 and other herbaceous species may exist, but typically will  
20 be absent. A continuous leaf litter layer may be present.  
21 Overgrown scrub in need of fire and/or mechanical  
22 treatment should not be confused with true xeric  
23 hammock.

24 **Description and Assessment**

25 In total, as illustrated in Table 8, the Xeric Hammock  
26 communities on the CFG are comprised of 307 acres, or  
27 0.3 percent of CFG total area. Xeric hammock on the CFG  
28 occurs north, south, and west of the Rodman Reservoir; in  
29 the Orange Springs Recreation Area and Eureka East  
30 Recreation Area; west of the Ocklawaha River from  
31 Country Club Boulevard south to Gores Landing; north of  
32 US 40; along the western edge of Marshall Swamp from  
33 Baseline Road to Marshall Swamp Trailhead; in the  
34 Diggings east and west of SR 200 and Ross Prairie.

35 The canopy and subcanopy are dominated by sand live oak,  
36 live oak (*Quercus virginiana*), and laurel oak (*Quercus*  
37 *hemisphaerica*), and sometimes sand pine (*Pinus clausa*) or  
38 southern magnolia (*Magnolia grandiflora*) are common.  
39 Shrubs include saw palmetto, scrub palmetto (*Sabal*



**Global and State Rank: G3/S3**

**Fire Interval: NA**

**Listed Species:**

Gopher tortoise, garberia, coontie  
longspurred mint, needle palm,  
pygmy pipes

**Invasive Exotic Species:**

Cogon grass, camphor tree,  
mimosa

**Management Practices:**

- Returning fire frequency to surrounding communities
- Invasive exotic treatment/control
- Oak management in ecotones





1 *etonia*), sparkleberry (*Vaccinium arboreum*), and deerberry (*Vaccinium stamineum*).

2 Herbs are sparse and include witchgrass (*Dichanthelium* sp.), Elliott's milkpea (*Galactia elliottii*),  
3 gayfeather (*Liatris* sp.), partridgeberry (*Mitchella repens*), narrowleaf silkgrass (*Pityopsis*  
4 *graminifolia*), bracken fern (*Pteridium aquilinum*), and sandyfield beaksedge (*Rhynchospora*  
5 *megalocarpa*).

6 Ground lichens (*Cladina evansii* and *Cladina subtenuis*) can be common. A few Spanish moss  
7 (*Tillandsia usneoides*) can be found in the canopy.

8 Vines are sparse, and include muscadine (*Vitis rotundifolia*) and earleaf greenbrier (*Smilax*  
9 *auriculata*).

10 Primary disturbances that were documented within xeric hammocks on the CFG include clearing, fire  
11 suppression, ORV and all-terrain vehicle (ATV) trails, trash dumping, ditching and canal construction,  
12 bike trails, and woody encroachment.

13 Two rare species were documented within the xeric hammock. There is one plant species: garberia,  
14 and one animal species: gopher tortoise (*Gopherus polyphemus*). In addition, three invasive species  
15 were noted: cogon grass (*Imperata cylindrica*), Camphor tree (*Cinnamomum camphora*), and Mimosa  
16 (*Albizia julibrissin*).

## 17 **Fire Regime**

18 Xeric hammocks typically develop after 30 or more years of fire exclusion and rarely burn, although  
19 occasional ground fires from adjacent uplands may pass through when leaf litter is dry. Catastrophic  
20 crown fires can result in reversion of the xeric hammock to sandhill, scrub, or another community  
21 type from which the hammock originated.

## 22 **General Management Measures**

23 Management activities in the xeric hammock communities on the CFG should focus on returning fire  
24 frequency to the surrounding natural communities and allowing fires to burn into the edges of the  
25 xeric hammock. In some areas, xeric hammock has replaced historic sandhills; in these areas, if  
26 restoration to sandhills is desired, mechanical removal of oaks prior to burning may be appropriate.  
27 Invasive exotic species monitoring and control also is needed.

# 1 Scrubby Flatwoods

2 **Global and State Rank: NA**


## 3 **Desired Future Condition**

4 Dominant tree species of the interior usually will be  
5 longleaf pine (*Pinus palustris*). Sand pines (*Pinus clausa*)  
6 typically are present. There will be a diverse shrubby  
7 understory often with patches of bare white sand. A scrub-  
8 type oak “canopy” will vary in height from three feet to  
9 eight feet and there will be a variety of oak age  
10 classes/heights across the landscape. Dominant shrubs  
11 include sand live oak (*Quercus geminata*), myrtle oak  
12 (*Quercus myrtifolia*), Chapman’s oak (*Quercus chapmanii*),  
13 saw palmetto (*Serenoa repens*), rusty staggerbush (*Lyonia*  
14 *ferruginea*), and tarflower (*Bejaria racemosa*). Cover by  
15 herbaceous species often is well below 40 percent. The  
16 optimal fire return interval for this community is  
17 regionally variable. Areas may be burned as frequently as  
18 every three years to 14 years when burn prescriptions are  
19 designed to achieve a mosaic of burned and unburned  
20 areas.

## 21 **Description and Assessment**

22 In total, as shown in Table 8, the Scrubby Flatwoods  
23 community on the CFG is comprised of 683 acres, or 1  
24 percent of the CFG total area. Scrubby flatwoods on the  
25 CFG are located in the Caravelle Wildlife Management  
26 Area; north of the Buckman Lock; north of the Rodman  
27 Reservoir; and along Deep Creek. Other scrubby flatwoods  
28 are located west of the Ocklawaha River from Country Club  
29 Boulevard south to Gores Landing; north of CR 314 and  
30 south of the Ocklawaha River; northeast of Marshall  
31 Swamp, the Triangle, and Inglis Island; and north of the  
32 Inglis Spillway.

33 The canopy of the scrubby flatwoods on the CFG is  
34 composed primarily of scattered sand pine, slash pine  
35 (*Pinus elliottii*), or longleaf pine. The subcanopy includes  
36 sand live oak, laurel oak (*Quercus hemisphaerica*), myrtle  
37 oak, water oak (*Quercus nigra*), live oak (*Quercus*  
38 *virginiana*), and cabbage palm (*Sabal palmetto*). Shrubs are  
39 dense and many of the overstory species occur in this layer  
40 as well.



**Global and State Rank: NA**



**Fire Interval: 3–14 years**

### **Listed Species:**

Gopher tortoise, garberia,  
Chapman’s skeletongrass

### **Invasive Exotic Species:**

Cogon grass, jelly palm

### **Management Practices:**

- Prescribed fire
- Invasive exotic treatment/control
- Hydrologic restoration



1 Additional species include tarflower, dwarf huckleberry (*Gaylussacia dumosa*), blue huckleberry  
2 (*Gaylussacia frondosa* var. *tomentosa*), sand holly (*Ilex ambigua*), myrtle dahoon (*Ilex cassine* var.  
3 *myrtifolia*), gallberry (*Ilex glabra*), gopher apple (*Licania michauxii*), rusty staggerbush, coastalplain  
4 staggerbush (*Lyonia fruticosa*), fetterbush (*Lyonia lucida*), wax myrtle (*Myrica cerifera*), wild olive  
5 (*Osmanthus americanus*), red bay (*Persea borbonia*), turkey oak (*Quercus laevis*), winged sumac (*Rhus*  
6 *copallinum*), scrub palmetto (*Sabal etonia*), saw palmetto, sparkleberry (*Vaccinium arboreum*),  
7 Darrow's blueberry (*Vaccinium darrowii*), shiny blueberry (*Vaccinium myrsinites*), and deerberry  
8 (*Vaccinium stamineum*).

9 The herbaceous, epiphyte, and vine strata of the scrubby flatwoods typically are sparse. Herbs  
10 include broomsedge bluestem (*Andropogon virginicus*), bottlebrush threeawn (*Aristida spiciformis*),  
11 wiregrass (*Aristida stricta* var. *beyrichiana*), coastalplain chaffhead (*Carphephorus corymbosus*),  
12 vanillaleaf (*Carphephorus odoratissimus*), Elliott's milkpea (*Galactia elliotii*), rough hedgehyssop  
13 (*Gratiola hispida*), shortleaf gayfeather (*Liatris tenuifolia*), narrowleaf silkgrass (*Pityopsis*  
14 *graminifolia*), bracken fern (*Pteridium aquilinum*), sandyfield beaksedge (*Rhynchospora*  
15 *megalocarpa*), sand spike-moss (*Selaginella arenicola*), sweet goldenrod (*Solidago odora*), lopsided  
16 indian grass (*Sorghastrum secundum*), and yellow hatpins (*Syngonanthus flavidulus*). Bartram's air-  
17 plant (*Tillandsia bartramii*), and Spanish moss (*Tillandsia usneoides*) are common epiphytes. Vines  
18 include earleaf greenbrier (*Smilax auriculata*), saw greenbrier (*Smilax bona-nox*), sarsaparilla vine  
19 (*Smilax pumila*), and muscadine (*Vitis rotundifolia*). The ground lichens (*Cladina evansii* and *Cladina*  
20 *subtenuis*) are present.

21 Historical and current disturbances that have impacted scrubby flatwoods include fire exclusion,  
22 clearing, forestry operations, ditching and canal construction, off-road trails, trash dumping, and  
23 utility corridors.

24 Three rare species—gopher tortoise (*Gopherus polyphemus*), garberia (*Garberia heterophylla*), and  
25 Chapman's skeletongrass (*Gymnopogon chapmanianus*)—occur within the scrubby flatwoods.

26 Two invasive species were noted: cogon grass (*Imperata cylindrica*) and jelly palm (*Butia capitata*).

## 27 **Fire Regime**

28 Although scrubby flatwoods occupy a drier environment than the surrounding mesic flatwoods, this  
29 community type does not burn as often. Natural fire frequency is three years to 14 years, with most  
30 burns occurring during late spring and early summer (April to June).

## 31 **General Management Measures**

32 Fire return intervals should be frequent enough to maintain shrub heights (less than six feet) within  
33 the ranges required by the Florida Scrub-Jay. If necessary to maintain ideal structure, effective fires  
34 in scrubby flatwoods should occur during hot, low-humidity conditions.

35 If embedded within or near to mesic flatwoods, scrubby flatwoods should be allowed to burn along  
36 with the mesic flatwoods on a two-year to five-year prescribed fire cycle, primarily during late spring  
37 and early summer (April to June). Scrubby flatwoods may not ignite as often as mesic flatwoods due  
38 to the relative incombustibility of the oak litter. In areas that have been converted to pine plantations,



- 1 thinning and reintroduction of prescribed fire are needed for restoration, and, in some cases, removal
- 2 of off-site pines and replanting with longleaf pine may be desirable.
  
- 3 Hydrologic alterations, such as ditching and other drainage channels, may be affecting scrubby
- 4 flatwoods vegetation in a few places. In these areas, restoration of natural hydrology is warranted.
- 5 Avoiding heavy ground disturbances is important to prevent elimination of the natural groundcover
- 6 and the establishment of weedy species.

# Mesic Flatwoods

**Global and State Rank: G4/S4**


## Desired Future Condition

Dominant pines usually will be longleaf pine (*Pinus palustris*). Native herbaceous groundcover should be more than 50 percent of the area and less than three feet in height. Saw palmetto (*Serenoa repens*) will comprise no more than 50 percent of total shrub species cover, and will be less than three feet in height. Shrub species include saw palmetto, gallberry (*Ilex glabra*), fetterbush (*Lyonia lucida*), runner oak (*Quercus elliotii*), dwarf live oak (*Quercus minima*), shiny blueberry (*Vaccinium myrsinites*), and dwarf huckleberry (*Gaylussacia dumosa*). Shrubs generally are knee-high or less, and there are few if any large trunks of saw palmetto along the ground. The optimal fire return interval for this community is one year to four years.

## Descriptions and Assessment

Within the CFG, there are 5,041 total acres of mesic flatwoods, or 7 percent of the total CFG area. Mesic flatwoods on the CFG are in the Caravelle Wildlife Management Area; north and south of the Buckman Lock; north, south, and east of the Rodman Reservoir; and along Deep Creek. Other areas of mesic flatwoods are located in the vicinity of the Orange Springs Recreation Areas; west of the Ocklawaha River from Country Club Boulevard south to Gores Landing; in Marshall Swamp and on Inglis Island; and north and south of the Inglis Spillway. Some of the higher-quality examples of mesic flatwoods occur south and north of Rodman Road and in the Country Club Boulevard area west of the Ocklawaha River.

The canopy and subcanopy layers of mesic flatwoods within the CFG typically are open and dominated by slash pine (*Pinus elliotii*) or longleaf pine, but a few loblolly pine (*Pinus taeda*) areas also occur. Many of the flatwoods on the CFG have a moderately dense to very dense midstory, which includes loblolly bay (*Gordonia lasianthus*), sweetgum (*Liquidambar styraciflua*), swamp bay (*Persea palustris*), laurel oak (*Quercus hemisphaerica*), water oak



**Global and State Rank: G4/S4**



**Fire Interval: 2–4 years**

## Listed Species:

Gopher tortoise, hooded pitcher plant, blueflower butterwort, giant orchid, pine lily, plume polypody, widespread polypody

## Invasive Exotic Species:

Cogon grass, camphor tree, torpedo grass, Chinese brake fern, grapefruit, Japanese climbing fern, Peruvian primrose-willow, sour orange, sword fern

## Management Practices:

- Prescribed fire
- Invasive exotic treatment/control
- Hydrologic restoration
- Timber treatments



1 (*Quercus nigra*), live oak (*Quercus virginiana*) and cabbage palm (*Sabal palmetto*).

2 On the CFG, shrubs are dense within many of the mesic flatwoods areas, evidence of long-term fire  
3 exclusion and lack of growing-season fires. Typical shrubs include saw palmetto, gallberry, shiny  
4 blueberry, fetterbush, huckleberry (*Gaylussacia frondosa* var. *tomentosa*), yaupon (*Ilex vomitoria*),  
5 rusty staggerbush (*Lyonia ferruginea*), wax myrtle (*Myrica cerifera*), red bay (*Persea borbonia*), dwarf  
6 live oak, dwarf pawpaw (*Asimina pygmea*), netted pawpaw (*Asimina reticulata*), scrub palmetto  
7 (*Sabal etonia*), highbush blueberry (*Vaccinium corymbosum*), Darrow's blueberry (*Vaccinium*  
8 *darrowii*), shiny blueberry (*Vaccinium myrsinites*), deerberry (*Vaccinium stamineum*), and coontie  
9 (*Zamia pumila*).

10 The herbaceous groundcover typically is dominated by wiregrass (*Aristida beyrichiana* var. *stricta*),  
11 bracken fern (*Pteridium aquilinum*), broomsedge bluestem (*Andropogon virginicus*), bottlebrush  
12 wiregrass (*Aristida spiciformis*), blue maidencane (*Amphicarpum muhlenbergianum*), switchgrass  
13 (*Panicum virgatum*), and tapered witchgrass (*Dichantherium acuminatum*).

14 Forbs include vanilla leaf (*Carphephorus odoratissimus*), yankeeweed (*Eupatorium compositifolium*),  
15 slender flattop goldenrod (*Euthamia caroliniana*), Elliott's milkpea (*Galactia elliotii*), shortleaf  
16 gayfeather (*Liatris tenuifolia*), narrowleaf silkgrass (*Pityopsis graminifolia*), orange milkwort  
17 (*Polygala lutea*), blackroot (*Pterocaulon pycnostachyum*), and pale meadow beauty (*Rhexia mariana*).

18 Epiphyte and vine abundances vary. Epiphytes include resurrection fern (*Pleopeltis polypodioides*  
19 var. *michauxiana*), Bartram's air-plant (*Tillandsia bartramii*), and Spanish moss (*Tillandsia*  
20 *usneoides*). Vines are abundant, especially in fire-excluded sites. Peppervine (*Ampelopsis arborea*),  
21 earleaf greenbrier (*Smilax auriculata*), saw greenbrier (*Smilax bona-nox*), cat greenbrier (*Smilax*  
22 *glauca*), eastern poison ivy (*Toxicodendron radicans*), and muscadine (*Vitis rotundifolia*) are common.

23 Listed species observed within the mesic flatwoods include: gopher tortoise (*Gopherus polyphemus*),  
24 hooded pitcher plant (*Sarracenia minor*), blueflower butterwort (*Pinguicula caerulea*), giant orchid  
25 (*Pteroglossaspis ecristata*), lilly plume, and the widespread polypody (*Pecluma dispersa*). Invasive  
26 species observed during the FNAI survey included cogon grass (*Imperata cylindrica*) camphor tree  
27 (*Cinnamomum camphora*), torpedo grass (*Panicum repens*), Chinese brake fern (*Pteris vittata*),  
28 grapefruit (*Citrus paradisi*), Japanese climbing fern (*Lygodium japonicum*), Peruvian primrose willow  
29 (*Ludwigia peruviana*), sour orange (*Citrus aurantium*), and sword fern (*Nephrolepis cordifolia*).

30 Historical and current disturbances that have impacted mesic flatwoods include fire exclusion,  
31 clearing, forestry operations, ditching and canal construction, hydrological alteration, off-road trails,  
32 trash dumping, and exotic plant invasion.

### 33 **Fire Regime**

34 Within mesic flatwood communities, it is important to conduct dormant season fires until heavy fuel  
35 loads are diminished enough to conduct growing season fires without undue tree mortality. Mesic  
36 flatwoods require application of growing-season fires on a one-year to four-year cycle. Nearly all  
37 plants and animals inhabiting this community are adapted to periodic fires; several species depend  
38 on fire for their continued existence. Without relatively frequent fires, mesic flatwoods succeed into  
39 hardwood-dominated forests, shading out and eliminating the diverse groundcover.

1 **General Management Measures**

2 Implementing prescribed fires on a two-year to four-year cycle is critical for restoration and  
3 maintenance of mesic flatwoods. Priority should be given to burning mesic flatwoods with higher-  
4 quality groundcover, using frequent growing-season fires to encourage herbaceous species,  
5 especially wiregrass, to reproduce naturally. Many of the mesic flatwoods on the CFG have been  
6 converted to pine plantations. In these areas, tree thinning also may be needed to open up the canopy  
7 and allow more light to reach the ground.

8 In some areas, ditching and other water control efforts have likely changed the natural hydrology of  
9 the mesic flatwoods. Restoration of hydrology may be needed in some of these areas. Restricting the  
10 use of heavy machinery to dry periods when the soil is not saturated will help with avoiding  
11 unnecessary vegetation and soil disturbances.

# Hydric Hammock

**Global and State Rank: G4/S4**

## Desired Future Condition


Hydric hammock is a closed canopy, evergreen hardwood and/or palm forest with a variable understory dominated by palms, with sparse to moderate ground cover of grasses and ferns. Typical canopy species will include laurel oak (*Quercus laurifolia*), cabbage palm (*Sabal palmetto*), live oak (*Quercus virginiana*), sweet bay (*Magnolia virginiana*), swamp tupelo (*Nyssa sylvatica biflora*), American elm (*Ulmus americana*), red maple (*Acer rubrum*) and other hydrophytic tree species. Soils are poorly drained, with a normal hydroperiod that is seldom more than 60 days per year. Hydric hammock should occasionally burn by allowing fires to naturally burn across ecotones from fires originating in adjacent upland natural communities.

## Description and Assessment

In total, as illustrated in Table 8, hydric hammocks account for 3,663 acres of the CFG, or 5 percent of CFG total area. Outstanding examples of hydric hammock within the CFG occur along Deep Creek; north of CR 310; and in several areas that join the Ocklawaha River, such as Marshall Swamp and north of US 40. Other areas include the Caravelle Wildlife Management Area and areas within the WMA such as north and south of the Buckman Lock and north and east of the Rodman Reservoir. Hydric hammocks also exist in the vicinity of the Kenwood Recreation Area and in various sections along the Ocklawaha River from NE 150 Avenue south to CR 314, from Turkey Landing south to US 40, and north and south of the Inglis Spillway.

The canopy and subcanopy layers within the hydric hammock are semi-closed to closed and contain a good diversity of trees. Dormant-season fires should be conducted until heavy fuel loads are diminished enough to conduct growing-season fires without undue tree mortality (at this time, no growing-season fires occur within the CFG flatwoods).

Dominant species are diamond leaf oak, sweetgum (*Liquidambar styraciflua*), live oak, cabbage palm, and red maple (*Acer rubrum*). A great variety of other canopy and



**Global and State Rank: G4/S4**

**Fire Interval: NA**

## Listed Species:

Royal fern, variable-leaved indian plantain, large-leaved grass of parnassus, pinkroot, Treat's zephyrlily, toothpetal false rein orchid, spiny-pod, southern twayblade, angle pod, buckthorn, Chapman's sedge, Florida willow, hairy shadow-witch, needle palm, palegreen orchid

## Invasive Exotic Species:

Paper mulberry, camphor tree, wild taro, Japanese climbing fern, cat's claw vine, Chinaberry, tropical soda apple, Caesar's weed, Chinese brake fern, cogon grass, coral ardisia, gardenia, glossy privet, Japanese honeysuckle, mimosa, Peruvian rimrose-willow, skunk vine, sword fern

## Management Practices:

- Maintaining natural hydrology
- Invasive exotic treatment/control





1 small trees can be found, including American hornbeam (*Carpinus caroliniana*), Atlantic white cedar  
2 (*Chamaecyparis thyoides*), Carolina ash (*Fraxinus caroliniana*), green ash (*Fraxinus pennsylvanica*),  
3 red cedar (*Juniperus virginiana*), tulip tree (*Liriodendron tulipifera*), southern magnolia (*Magnolia*  
4 *grandiflora*), sweet bay, swamp bay (*Persea palustris*), slash pine (*Pinus elliottii*), loblolly pine (*Pinus*  
5 *taeda*), swamp chestnut oak (*Quercus michauxii*), swamp dogwood (*Cornus foemina*), cypress  
6 (*Taxodium* spp.), basswood (*Tilia americana*), and American elm.

7 Typical shrubs include Florida hobblebush (*Agarista populifolia*), common buttonbush  
8 (*Cephalanthus occidentalis*), wax myrtle (*Myrica cerifera*), bluestem palmetto (*Sabal minor*),  
9 mountain azalea (*Rhododendron canescens*), highbush blueberry (*Vaccinium corymbosum*), and  
10 cabbage palm.

11 Herbs can be sparse or patchy, and include variable-leaved indian plantain (*Arnoglossum*  
12 *diversifolium*), false nettle (*Boehmeria cylindrica*), clustered sedge (*Carex glaucescens*), spadeleaf  
13 (*Centella asiatica*), woodoats (*Chasmanthium* spp.), cypress witchgrass (*Dichantherium dichotomum*),  
14 whorled marsh pennywort (*Hydrocotyle verticillata*), needle rush (*Juncus roemerianus*), redtop  
15 panicum (*Panicum rigidulum*), swamp smartweed (*Polygonum hydropiperoides*), millet beaksedge  
16 (*Rhynchospora miliacea*), and lizard's tail (*Saururus cernuus*).

17 Ferns are diverse and can be abundant. These include cinnamon fern (*Osmunda cinnamomea*), royal  
18 fern (*Osmunda regalis*), southern wood fern (*Dryopteris ludoviciana*), southern shield fern  
19 (*Thelypteris kunthii*), marsh fern (*Thelypteris palustris* var. *pubescens*), netted chain fern  
20 (*Woodwardia areolata*), and Virginia chain fern (*Woodwardia virginica*).

21 Epiphytes can be abundant, and include green fly orchid (*Epidendrum conopseum*), golden polypody  
22 (*Phlebodium aureum*), resurrection fern (*Pleopeltis polypodioides* var. *michauxiana*), Bartram's air-  
23 plant (*Tillandsia bartramii*), and Spanish moss (*Tillandsia usneoides*).

24 Vine diversity also is impressive, with climbing hempvine (*Mikania scandens*), Virginia creeper  
25 (*Parthenocissus quinquefolia*), peppervine (*Ampelopsis arborea*), rattan vine (*Berchemia scandens*),  
26 crossvine (*Bignonia capreolata*), trumpet creeper (*Campsis radicans*), climbing hydrangea  
27 (*Decumaria barbara*), yellow jessamine (*Gelsemium sempervirens*), earleaf greenbrier (*Smilax*  
28 *auriculata*), saw greenbrier (*Smilax bona-nox*), laurel greenbrier (*Smilax laurifolia*), bristly  
29 greenbrier (*Smilax tamnoides*), poison ivy (*Toxicodendron radicans*), summer grape (*Vitis aestivalis*),  
30 muscadine (*Vitis rotundifolia*), and frost grape (*Vitis vulpina*).

31 Previous and current disturbances include clearing, forestry operations, ditching and canal  
32 construction, hydrological alteration, impoundments, roads and off-road vehicle trails, trash  
33 dumping, and introduction of exotic plants.

34 The rare plants variable-leaved indian plantain, Florida milk pod (*Matelea floridana*), grass-of-  
35 Parnassus (*Parnassia grandifolia*), and pinkroot (*Spigelia loganioides*) were observed during the  
36 FNAI survey.

37 The following species of invasive plants occur in the hydric hammocks: paper mulberry (*Broussonetia*  
38 *papyrifera*), camphor tree (*Cinnamomum camphora*), wild taro (*Colocasia esculenta*), Japanese  
39 climbing fern (*Lygodium japonicum*), cat's claw vine (*Macfadyena unguis-cati*), Chinaberry (*Melia*

1 *azedarach*), tropical soda apple (*Solanum viarum*), Caesar's weed (*Urena lobata*), Chinese brake fern  
2 (*Pteris vittata*), cogon grass (*Imperata cylindrica*), coral ardisia (*Ardisia crenata*), gardenia (*Gardenia*  
3 *jasminoides*), glossy privet (*Ligustrum lucidum*), Japanese honeysuckle (*Lonicera japonica*), mimosa  
4 (*Mimosa pudica*), Peruvian primrose-willow (*Ludwigia peruviana*), skunk vine (*Paederia foetida*), and  
5 sword fern (*Nephrolepis cordifolia*).

## 6 **Fire Regime**

7 Hydric hammocks usually are too wet to support fires due to their saturated soils and the sparsity of  
8 fine fuels, so they are not considered a pyrogenic natural community.

## 9 **General Management Measures**

10 Effective management of hydric hammocks primarily consists of maintaining natural hydrology and  
11 monitoring and treating invasive/exotic plants.

12 Feral hogs (*Sus scrofa*) can cause significant vegetation damage. They should be monitored and  
13 efforts should be made to control numbers, if needed.



# Wet Flatwoods

**Global and State Rank: G4/S4**


## Desired Future Condition

Depending on the region of the state, dominant pines usually will be longleaf pine (*Pinus palustris*), slash pine (*Pinus elliottii*), pond pine (*Pinus serotina*), and/or loblolly pine (*Pinus taeda*). Pond cypress (*Taxodium ascendens*) may reach the canopy in some locations. The canopy will be open, with pines being widely scattered and of at least three age classes. The subcanopy may include scattered sweetbay (*Magnolia virginiana*), swamp bay (*Persea palustris*), and loblolly bay (*Gordonia lasianthus*). Common shrubs will include fetterbush (*Lyonia lucida*), large gallberry (*Ilex coriacea*), titi (*Cyrilla racemiflora*), and wax myrtle (*Myrica cerifera*). The optimal fire return interval for this community is five years to 10 years.

## Description and Assessment

Wet flatwoods occur on relatively flat, poorly drained, acidic sands overlying an organic hardpan or clay layer. The hardpan substantially reduces the percolation of water, so wet flatwoods can be inundated or saturated for one or more months per year. In total, as shown in Table 8, the wet flatwoods communities accounts for 1,773 acres of the CFG, or 3 percent of CFG total area. Wet flatwoods on the CFG occur in the Caravelle Wildlife Management Area and in areas within the WMA such as north and south of the Buckman Lock and north, west, and east of the Rodman Reservoir. Wet flatwoods also exist in the vicinity of the Kenwood Recreation Area; west of the Ocklawaha River from Country Club Boulevard south to Gores Landing; on the western edge of Marshall Swamp; on Inglis Island; and north and south of the Inglis Spillway.

The vegetative structure of wet flatwoods within the CFG is highly variable from site to site and dependent on fire history, hydroperiod, and land-use history, such as silviculture. The canopy typically is slash pine, longleaf pine, loblolly pine, and pond pine in a few locations. In long-term fire-excluded sites, loblolly bay (*Gordonia lasianthus*), sweetgum (*Liquidambar styraciflua*), water oak (*Quercus nigra*), and swamp bay (*Persea palustris*) may



**Global and State Rank: G4/S4**



**Fire Interval: 2-5 years**

## Listed Species:

Pineland plantain, royal fern, blueflower butterwort, hooded pitcher plant, cinnamon fern, mountain azalea

## Invasive Exotic Species:

Caesar's weed, camphor tree, Chinese brake fern, cogon grass, torpedo grass

## Management Practices:

- Prescribed fire
- Invasive exotic treatment/control



1 reach the canopy and dominate the subcanopy. These species also are found in the shrub layers, as  
2 are gallberry (*Ilex glabra*), saw palmetto (*Serenoa repens*), fetterbush, dwarf huckleberry  
3 (*Gaylussacia dumosa*), blue huckleberry (*Gaylussacia frondosa* var. *tomentosa*), Darrow's blueberry  
4 (*Vaccinium darrowii*), and shiny blueberry (*Vaccinium myrsinites*).

5 Due to the dense shrub layer, herbs typically are sparse in wet flatwoods on the CFG, although a few  
6 locations have herb cover greater than 50 percent due to recent fires. Typical species in the herb  
7 layer are maidencane (*Panicum hemitomon*), Virginia chain fern (*Woodwardia virginica*), Carolina  
8 redroot (*Lachnanthes caroliana*), blue maidencane (*Amphicarpum muhlenbergianum*), purple  
9 bluestem (*Andropogon glomeratus* var. *glaucoopsis*), broomsedge bluestem (*Andropogon virginicus*),  
10 and low panic grasses (*Dichanthelium* spp.). Spanish moss (*Tillandsia usneoides*) is found occasionally  
11 in the tree canopy.

12 Vines can be abundant and include yellow jessamine (*Gelsemium sempervirens*), Virginia creeper  
13 (*Parthenocissus quinquefolia*), earleaf greenbrier (*Smilax auriculata*), saw greenbrier (*Smilax bona-*  
14 *nox*), cat greenbrier (*Smilax glauca*), laurel greenbrier (*Smilax laurifolia*), and muscadine (*Vitis*  
15 *rotundifolia*).

16 The hooded pitcher plant (*Sarracenia minor*) was documented in wet flatwoods on the CFG, and no  
17 exotic species were documented. Historically, wet prairie occurred within wet flatwoods on Inglis  
18 Island and south of the Inglis Spillway.

19 Disturbances noted within wet flatwoods on the CFG include clearing, fire exclusion, woody  
20 encroachment, forestry operations, fire-breaks, and ditching and canal construction.

## 21 **Fire Regime**

22 The natural fire return interval in wet flatwoods is variable, every two to five years. Without  
23 relatively frequent fires, wet flatwoods succeed into hardwood-dominated forests whose closed  
24 canopy would essentially eliminate the ground cover herbs and shrubs.

## 25 **General Management Measures**

26 For management purposes prescribed fires should be applied on a two-year to four-year cycle to  
27 reduce woody encroachment, sustain herbaceous species, and aid in fuel reduction for the prevention  
28 of catastrophic wildfires. In some areas, mechanical reduction of tall, dense shrubs or the midstory  
29 cover may be needed prior to prescribed burning.

30 Restricting the use of heavy machinery to dry periods when the soil is not saturated will help to avoid  
31 unnecessary vegetation disturbances and soil disturbances such as rutting and erosion.

# 1 Baygall

2 **Global and State Rank: G4/S4**


## 3 **Desired Future Condition**

4 Baygall consist of wet, densely forested, peat-filled  
5 depressions typically near the base of a slope. Seepage  
6 from adjacent uplands will maintain saturated conditions.  
7 Medium to tall trees will mainly consist of sweet bay  
8 (*Magnolia virginiana*), loblolly bay (*Gordonia lasianthus*),  
9 and/or swamp bay (*Persea palustris*); occasionally sparse  
10 pines (*Pinus spp.*) also may exist. A thick understory  
11 consisting of gallberry (*Ilex glabra*), fetterbush (*Lyonia*  
12 *lucida*), dahoon holly (*Ilex cassine*), titi (*Cyrilla*  
13 *racemiflora*), and red maple (*Acer rubrum*) is typical, with  
14 climbing vines such as greenbriar (*Smilax spp.*) and  
15 muscadine grape (*Vitis spp.*) in abundance. The optimal  
16 fire return interval for this community is 25 years to 100  
17 years. Frequent fires from adjacent communities should be  
18 allowed to enter the baygall ecotone.

## 19 **Description and Assessment**

20 In total, as illustrated in Table 8, the Baygall communities  
21 account for 549 acres on the CFG, or 1 percent of CFG  
22 total area. Baygall is documented in the Caravelle Wildlife  
23 Management Area and in the vicinity of Deep Creek; north  
24 and south of the Buckman Lock; and north of the Rodman  
25 Reservoir in the Kenwood and Orange Springs Recreation  
26 Sites. Other areas include the western side border of the  
27 Ocklawaha River from Eureka West to Gores Boat  
28 Landings; the western edge of Marshall Swamp; and in the  
29 Diggings west of SR 200.

30 Within the CFG, baygall generally consists of a dense  
31 overstory and understory and few herbs. Characteristic  
32 canopy and subcanopy trees include loblolly bay,  
33 sweetgum (*Liquidambar styraciflua*), sweet bay, swamp  
34 tupelo (*Nyssa sylvatica* var. *biflora*), swamp bay, slash pine  
35 (*Pinus elliotii*), longleaf pine (*Pinus palustris*), loblolly pine  
36 (*Pinus taeda*), and pond cypress (*Taxodium ascendens*).  
37 Shrubs are diverse. In addition to the same shrubs found in  
38 the upper strata, additional shrub species include Florida  
39 hobblebush (*Agarista populifolia*), dahoon holly (*Ilex*  
40 *cassine*), large gallberry (*Ilex coriacea*), gallberry,  
41 fetterbush, wax myrtle (*Myrica cerifera*), swamp azalea



**Global and State Rank: G4/S4**

**Fire Interval: NA**

### **Listed Species:**

Royal fern, mountain azalea, needle palm

### **Invasive Exotic Species:**

Cogon grass

### **Management Practices:**

- Invasive exotic treatment/control



1 (*Rhododendron viscosum*), cabbage palm (*Sabal palmetto*), saw palmetto (*Serenoa repens*), highbush  
2 blueberry (*Vaccinium corymbosum*), Elliott's blueberry (*Vaccinium elliotii*), blue huckleberry  
3 (*Gaylussacia frondosa* var. *tomentosa*), and dwarf palmetto (*Sabal minor*).

4 Typically, there is little herbaceous cover due to the low light levels under the dense overstory.  
5 Characteristic herbs include Walter's sedge (*Carex striata*), slender woodoats (*Chasmanthium*  
6 *laxum*), witchgrass (*Dichanthelium* sp.), Carolina redroot (*Lachnanthes caroliana*), cinnamon fern  
7 (*Osmunda cinnamomea*), maidencane (*Panicum hemitomon*), bracken fern (*Pteridium aquilinum*),  
8 sphagnum moss (*Sphagnum* spp.), eastern gama grass (*Tripsacum dactyloides*), netted chain fern  
9 (*Woodwardia areolata*), and Virginia chain fern (*Woodwardia virginiana*). The epiphytic and vine  
10 layers of the baygall occur infrequently. Bartram's air-plant (*Tillandsia bartramii*) and Spanish moss  
11 (*Tillandsia usneoides*) are two common epiphytes. Vines include yellow jessamine (*Gelsemium*  
12 *sempervirens*), cat greenbrier (*Smilax glauca*), laurel greenbrier (*Smilax laurifolia*), and muscadine.  
13 The latter often forms thickets around the edges of the baygall and where the canopy trees are sparse.

14 Two state-listed commercially exploited species, the sweet pinxter azalea (*Rhododendron canescens*)  
15 and needle palm (*Rhapidophyllum hystrix*), were found along the north boundary of the Greenway,  
16 north of the Buckman Lock Visitor Center. Only cogon grass was documented. Human disturbances,  
17 such as logging, clearing, fire exclusion, hydrological alteration, roads, and utility corridors are  
18 evident in baygall and likely have changed the vegetation species composition, structure, and  
19 hydrology from the historic condition.

## 20 **Fire Regime**

21 Although the upland ecotones of baygall likely burn historically with the adjacent uplands (typically  
22 every two years to four years), fires in the baygall interior are likely infrequent (every 50 years to  
23 100 years).

## 24 **General Management Measures**

25 The ecotones of the baygall tend to be fire-excluded, with bay trees spreading into the adjacent  
26 natural communities. Fire can be used to control baygall vegetation (primarily loblolly bay and  
27 swamp bay) that has encroached into mesic and wet flatwoods. Fires from the surrounding fire-  
28 dependent communities should be allowed to burn into the baygall during periods of high water and  
29 to extinguish naturally. Management activities also should focus on maintaining and, where needed,  
30 restoring the natural hydrology of these systems.



# 1 Bottomland Forest

## 2 Global and State Rank: NA


### 3 Desired Future Condition

4 Bottomland forest is a low lying, mesic to hydric  
 5 community prone to periodic flooding. Vegetation will  
 6 consist of a mature closed canopy of deciduous and  
 7 evergreen trees. Overstory species may consist of species  
 8 such as sweetgum (*Liquidambar styraciflua*), sweetbay  
 9 (*Magnolia virginiana*), loblolly bay (*Gordonia lasianthus*),  
 10 water oak (*Quercus nigra*), live oak (*Quercus virginiana*),  
 11 swamp chestnut oak (*Quercus michauxii*), loblolly pine  
 12 (*Pinus taeda*), and spruce pine (*Pinus glabra*). Red maple  
 13 (*Acer rubrum*) and bald cypress (*Taxodium distichum*) also  
 14 may be present. The understory may be open or dense.  
 15 Understory species typically will include wax myrtle  
 16 (*Myrica cerifera*), dwarf palmetto (*Sabal minor*), and  
 17 swamp dogwood (*Cornus foemina*). Presence of  
 18 groundcover is variable and may consist of witchgrass  
 19 (*Dicanthelium* sp.) and various sedges (*Carex* spp.).

### 20 Description and Assessment

21 In total, as shown in Table 8, the Bottomland Forest  
 22 represents 1,258 acres of the CFG, or 2 percent of CFG  
 23 total area. Bottomland forest communities in the CFG are  
 24 within the Caravelle Wildlife Management Area in the  
 25 vicinity of the Rodman Reservoir. Other areas include  
 26 different sites along the Ocklawaha River from north of  
 27 Paynes Landing south to CR 314; and north of US 40.

28 The vegetation strata are highly variable within the  
 29 bottomland forests on the CFG. The canopy has a diverse  
 30 array of tree species, which can include red maple (*Acer*  
 31 *rubrum*), American hornbeam (*Carpinus caroliniana*),  
 32 green ash (*Fraxinus pennsylvanica*), sweetgum, sweetbay,  
 33 loblolly pine, diamond leaf oak (*Quercus laurifolia*), swamp  
 34 chestnut oak (*Quercus michauxii*), live oak (*Quercus*  
 35 *virginiana*), cabbage palm (*Sabal palmetto*), and bald  
 36 cypress. Shrubs include Florida hobblebush (*Agarista*  
 37 *populifolia*), wax myrtle, saw palmetto (*Serenoa repens*),  
 38 and small-leaf viburnum (*Viburnum obovatum*).



**Global and State Rank: NA**

**Fire Interval: NA**

### **Listed Species:**

Angle pod, buckthorn, needle palm, pinkroot, Treat’s zephyrlily

### **Invasive Exotic Species:**

Caesar's weed, camphor tree, coral ardisia, hedge bamboo, Japanese climbing Fern, Japanese honey-suckle, nandina, paper mulberry, sword fern, wild taro

### **Management Practices:**

- Hydrologic restoration
- Allow prescribed fire from adjacent communities to burn



1 Herbs typically are sparse and can include switchcane (*Arundinaria gigantea*), longleaf woodoats  
2 (*Chasmanthium laxum* var. *sessiliflorum*), shiny woodoats (*Chasmanthium nitidum*), netted chain fern  
3 (*Woodwardia areolata*), and Virginia chain fern (*Woodwardia virginiana*).

4 Epiphytes and vines are common and include Spanish moss (*Tillandsia usneoides*), eastern poison ivy  
5 (*Toxicodendron radicans*), and muscadine (*Vitis rotundifolia*).

6 Invasive species found within CFG bottomland forests include: Caesar's weed (*Urena lobata*),  
7 camphor tree (*Cinnamomum camphora*), coral ardisia (*Ardisia crenata*), hedge bamboo (*Bambusa*  
8 *multiplex*), Japanese climbing fern (*Lygodium japonicum*), Japanese honeysuckle (*Lonicera japonica*),  
9 nandina (*Nandina domestica*), paper mulberry (*Broussonetia papyrifera*), sword fern (*Nephrolepis*  
10 *cordifolia*), wild taro (*Colocasia esculenta*).

11 Human disturbances—such as clearing, logging, ditching, impoundment, and artificial pond  
12 creation—are evident in some of the bottomland forests. These disturbances likely have changed the  
13 vegetation species composition and structure from the historic condition.

#### 14 **Fire Regime**

15 Fire is infrequent or nonexistent in bottomland forests, possibly occurring only during times of  
16 extreme drought. Prescribed fires from surrounding communities should be allowed to extinguish  
17 naturally at the edges of the bottomland forest to maintain a natural ecotone.

#### 18 **General Management Measures**

19 Management activities in bottomland forest on the CFG could focus on restoring and maintaining the  
20 natural hydrologic patterns of the forest and associated wetlands. Prescribed fires from adjacent  
21 communities should be allowed to burn into the bottomland forest.

# 1 Floodplain Swamp

## 2 Global and State Rank: NA


## 3 Desired Future Condition

4 Floodplain swamps are forested wetlands associated with  
5 rivers or streams, which are inundated for much of the  
6 year. They may be located within the floodplain of any  
7 permanently moving stream or river, and range from  
8 narrow strips of cypress along primary and secondary  
9 streams to expansive stands along large rivers. These  
10 swamps may immediately border the main river or stream  
11 channel or occur farther back within the floodplain in  
12 backswamps, oxbows, overflow channels, and old stream  
13 beds that are only connected to flowing water during flood  
14 conditions. The floodplain swamps intergrade into  
15 bottomland forests, hydric hammock, wet flatwoods, mesic  
16 hammock, and mesic flatwoods.

## 17 Description and Assessment

18 In total, as illustrated in Table 8, floodplain swamps  
19 represent 10,497 acres of the CFG, or 15 percent of CFG  
20 total area. Excellent examples of floodplain swamp occur  
21 within the CFG along the St. Johns River; in the Etoniah area  
22 north of Rodman; along the Ocklawaha River; and within  
23 Marshall Swamp. Other floodplain swamps occur north  
24 and south of the Buckman Lock; north and east of the  
25 Rodman Reservoir; in the vicinity of the Kenwood and  
26 Orange Springs Recreation Areas; in areas along the  
27 Ocklawaha River, including north of Paynes Landing, from  
28 Eureka Bridge south to Gores Landing, from Turkey  
29 Landing south to US 40; east and west of the Ocklawaha  
30 River from CR 314 south; in an area east of US 41 on the  
31 south side of the Withlacoochee River/Lake Rousseau; and  
32 north of the Inglis Spillway.

33 Typically, the CFG floodplain swamps have a semi-closed  
34 to closed canopy and subcanopy dominated by a great  
35 diversity of tree species. These include red maple (*Acer*  
36 *rubrum*), water hickory (*Carya aquatica*), green ash  
37 (*Fraxinus pennsylvanica*), Carolina ash (*Fraxinus*  
38 *caroliniana*), dahoon holly (*Ilex cassine*), sweetgum  
39 (*Liquidambar styraciflua*), sweetbay (*Magnolia virginiana*),  
40 swamp tupelo (*Nyssa sylvatica* var. *biflora*), swamp bay



## Global and State Rank: NA

## Fire Interval: NA

## Listed Species:

Angle pod, cardinal flower, needle palm, cinnamon fern, hairy shadow-witch, mountain azalea, royal fern, toothpetal false rein orchid

## Invasive Exotic Species:

Alligator weed, Caesar's weed, camphor tree, Chinaberry, Chinese brake fern, Chinese privet, Chinese tallow tree, Chinese wisteria, cogon grass, coral ardisia, elephant ear, glossy privet, Japanese climbing fern, Mexican-petunia, nandina, Peruvian primrose-willow, red-tip photinia, skunk vine, small-leaf spiderwort, sword fern, tropical soda apple, Turk's turban, water hyacinth, water lettuce, white mulberry, wild taro, winged yam

## Management Practices:

- Hydrologic restoration
- Allow prescribed fire from adjacent communities to burn
- Invasive exotic treatment/control





1 (*Persea palustris*), slash pine (*Pinus elliottii*), loblolly pine (*Pinus taeda*), diamond leaf oak (*Quercus*  
2 *laurifolia*), water oak (*Quercus nigra*), live oak (*Quercus virginiana*), cabbage palm (*Sabal palmetto*),  
3 coastalplain willow (*Salix caroliniana*), pond cypress (*Taxodium ascendens*), bald cypress (*Taxodium*  
4 *distichum*), and American elm (*Ulmus americana*). The shrub strata cover varies, ranging from  
5 moderate to sparse, and includes many species that occur in the upper layer. Additional species  
6 include groundsel tree (*Baccharis halimifolia*), hackberry (*Celtis laevigata*), common buttonbush  
7 (*Cephalanthus occidentalis*), swamp dogwood (*Cornus foemina*), common persimmon (*Diospyros*  
8 *virginiana*), loblolly bay (*Gordonia lasianthus*), large gallberry (*Ilex coriacea*), Virginia willow (*Itea*  
9 *virginica*), swamp doghobble (*Leucothoe racemosa*), fetterbush (*Lyonia lucida*), wax myrtle (*Myrica*  
10 *cerifera*), eastern hophornbeam (*Ostrya virginiana*), sawtooth blackberry (*Rubus argutus*), dwarf  
11 palmetto (*Sabal minor*), and elderberry (*Sambucus nigra* subsp. *canadensis*).

12 The herbaceous layer varies from sparse to dense. Typical herbs include lizard's tail (*Saururus*  
13 *cernuus*), false nettle (*Boehmeria cylindrica*), hairy bedstraw (*Galium pilosum*), stiff marsh bedstraw  
14 (*Galium tinctorium*), scarlet rosemallow (*Hibiscus coccineus*), large leaf marsh pennywort  
15 (*Hydrocotyle bonariensis*), whorled marsh pennywort (*Hydrocotyle verticillata*), and Virginia iris (*Iris*  
16 *virginica*).

17 A variety of sedges, beakrushes, and grasses are present, such as giant sedge (*Carex gigantea*),  
18 clustered sedge (*Carex glaucescens*), warty sedge (*Carex verrucosa*), spadeleaf (*Centella asiatica*),  
19 narrow fruit horned beaksedge (*Rhynchospora inundata*), millet beaksedge (*Rhynchospora miliacea*)  
20 slender woodoats (*Chasmanthium laxum*), longleaf woodoats (*Chasmanthium laxum* var.  
21 *sessiliflorum*), sawgrass (*Cladium jamaicense*), and eastern gama grass (*Tripsacum dactyloides*).

22 Ferns can be abundant, and the more commonly occurring are netted chain fern (*Woodwardia*  
23 *areolata*), Virginia chain fern (*Woodwardia virginica*), royal fern (*Osmunda regalis*), and marsh fern  
24 (*Thelypteris palustris* var. *pubescens*).

25 Both epiphytes and vines are common. Among the epiphytes are Bartram's air-plant (*Tillandsia*  
26 *bartramii*) and Spanish moss (*Tillandsia usneoides*). Vines are diverse in numbers of species and  
27 include peppervine (*Ampelopsis arborea*), rattan vine (*Berchemia scandens*), crossvine (*Bignonia*  
28 *capreolata*), trumpet creeper (*Campsis radicans*), climbing hydrangea (*Decumaria barbara*), yellow  
29 jessamine (*Gelsemium sempervirens*), climbing hempvine (*Mikania scandens*), Virginia creeper  
30 (*Parthenocissus quinquefolia*), saw greenbrier (*Smilax bona-nox*), cat greenbrier (*Smilax glauca*),  
31 laurel greenbrier (*Smilax laurifolia*), coral greenbrier (*Smilax walteri*), eastern poison ivy  
32 (*Toxicodendron radicans*), muscadine (*Vitis rotundifolia*), and American wisteria (*Wisteria*  
33 *frutescens*).

34 Two state-listed commercially exploited plant species were documented within floodplain swamp  
35 communities: cardinal flower (*Lobelia cardinalis*) and needle palm (*Rhapidophyllum hystrix*).

36 Within the floodplain swamp communities, the following invasive plants were documented,  
37 including alligator weed (*Alternanthera philoxeroides*), wild taro (*Colocasia esculenta*), water  
38 hyacinth (*Eichhornia crassipes*), Japanese climbing fern (*Lygodium japonicum*), skunkvine (*Paederia*  
39 *foetida*), Chinese brake fern (*Pteris vittata*), Chinese tallow tree (*Sapium sebiferum*), tropical soda  
40 apple (*Solanum viarum*), Caesar's weed (*Urena lobata*), Chinaberry (*Melia azedarach*), Chinese privet  
41 (*Ligustrum sinense*), Chinese wisteria (*Wisteria sinensis*), Cogon grass (*Imperata cylindrica*), coral

1 ardisia (*Ardisia crenata*), elephant ear (*Colocasia esculenta*), glossy privet (*Ligustrum lucidum*),  
2 Mexican petunia (*Ruellia simplex*), nandina (*Nandina domestica*), Peruvian primrose-willow  
3 (*Ludwigia peruviana*), red-tip photinia (*Photinia fraseri*), small-leaf spiderwort (*Tradescantia*  
4 *fluminensis*), sword fern (*Nephrolepis cordifolia*), Turk's turban (*Malvaviscus arboreus*), water lettuce  
5 (*Pistia stratiotes*), white mulberry (*Morus alba*), winged yam (*Dioscorea alata*).

6 Feral hogs (*Sus scrofa*) also are present.

7 Previous and current disturbances include clearing, forestry operations, ditching and canal  
8 construction, hydrological alteration, impoundments, roads and off-road vehicle trails, trash  
9 dumping, and introduction of exotic plants and animals.

## 10 **Fire Regime**

11 Floodplain swamps usually are too wet to support fires. However, fires from surrounding uplands  
12 should be allowed to creep into the floodplain swamps to enhance ecotone diversity and to reduce  
13 fuel loads, thereby minimizing the chances of catastrophic fires during drought.

## 14 **General Management Measures**

15 Floodplain swamps provide important wildlife habitat and contribute to the overall water quality of  
16 streams and rivers. Historical construction of ditches, canals, and berms has created serious  
17 hydrological alterations in some locations. Where needed, the natural hydrology should be restored  
18 and maintained. Closure of unnecessary roads and trails to vehicular traffic or redesigns, such as low  
19 water crossings, may be warranted in some areas to prevent erosion and interruption of water flow.  
20 This also may help abate illegal trash dumping.

# 1 Depression Marsh

## 2 Global and State Rank: NA

### 3 Desired Future Condition


4 Dominant vegetation in depression marshes includes  
5 maidencane (*Panicum hemitomon*), panic grasses  
6 (*Panicum spp.*), cutgrass (*Leersia sp.*), pickerelweed  
7 (*Pontederia cordata*), arrowheads (*Sagittaria sp.*),  
8 buttonbush (*Cephalanthus occidentalis*), St. John's wort  
9 (*Hypericum fasciculatum*), and coastalplain willow (*Salix*  
10 *caroliniana*).

### 11 Description and Assessment

12 The formation of depression marshes is associated with  
13 sinkholes and areas where sand has slumped inward.  
14 Rainfall, seepage, and runoff from surrounding uplands fill  
15 in these wetlands. Concentric zones of vegetation may be  
16 present that respond to the hydroperiod and edaphic  
17 conditions within each zone.

18 In total, as illustrated in Table 8, depression marsh  
19 communities represent 735 acres of the CFG, or 1  
20 percent of CFG total area. Numerous depression marshes  
21 occur within the CFG. Some of the best examples occur  
22 north of the Rodman Reservoir and south of CR 310.  
23 Depression marshes also occur in the Etoniah area north of  
24 Rodman; in the vicinity of Deep Creek; north and south of  
25 the Buckman Lock; north of the Rodman Reservoir; in the  
26 Country Club Boulevard area bordering the west side of  
27 the Ocklawaha River; north of US 40; in an area from US  
28 441 west to I-75; from Ross Prairie west to the Pruitt  
29 Trailhead area; and north and south of the Inglis Spillway.

30 Although mostly sparse, trees are sometimes present in  
31 the depression marsh communities on the CFG. Species  
32 include red maple (*Acer rubrum*), common persimmon  
33 (*Diospyros virginiana*), loblolly bay (*Gordonia lasianthus*),  
34 swamp tupelo (*Nyssa sylvatica* var. *biflora*), slash pine  
35 (*Pinus elliottii*), and pond cypress (*Taxodium ascendens*).  
36 The shrub layers of the depression marshes include many  
37 of the same species that occur in the upper strata. Other  
38 shrub species include groundsel tree (*Baccharis*  
39 *halimifolia*), common buttonbush, peelbark St. John's wort,



### Global and State Rank: NA



**Fire Interval: 1-8 years**

### Listed Species:

- Florida Sandhill Crane
- Wood Stork
- Hooded pitcher-plant

### Invasive Exotic Species:

- Camphor tree

### Management Practices:

- Prescribed fire
- Invasive exotic treatment/control



1 fetherbush (*Lyonia lucida*), wax myrtle (*Myrica cerifera*), swamp bay (*Persea palustris*), sand  
2 blackberry (*Rubus cuneifolius*), cabbage palm (*Sabal palmetto*), and coastalplain willow.

3 Herbaceous species usually are the dominant vegetational component of depression marshes, and in  
4 the CFG, the herbaceous layer of the depression marshes is the most diverse in numbers of species  
5 documented. Maidencane and blue maidencane (*Amphicarpum muhlenbergianum*) are typically  
6 dominant. Also common are switchcane (*Arundinaria gigantea*), carpetgrasses (*Axonopus* spp.), false  
7 nettle (*Boehmeria cylindrica*), giant sedge (*Carex gigantea*), clustered sedge (*Carex glaucescens*),  
8 spadeleaf (*Centella asiatica*), sawgrass (*Cladium jamaicense*), Baldwin's spikerush (*Eleocharis*  
9 *baldwinii*), tenangle pipewort (*Eriocaulon decangulare*), rattlesnake master (*Eryngium aquaticum*),  
10 dogfennel (*Eupatorium capillifolium*), falsefennel (*Eupatorium leptophyllum*), southern  
11 umbrellasedge (*Fuirena scirpoidea*), marsh pennywort (*Hydrocotyle* sp.), Carolina redroot  
12 (*Lachnanthes caroliana*), savannah primrosewillow (*Ludwigia virgata*), shade mudflower  
13 (*Micranthemum umbrosum*), rosy camphorweed (*Pluchea rosea*), swamp smartweed (*Polygonum*  
14 *hydropiperoides*), shortbristle horned beaksedge (*Rhynchospora corniculata*), fascicled beaksedge  
15 (*Rhynchospora fascicularis*), narrowfruit horned beaksedge (*Rhynchospora inundata*), sugarcane  
16 plumegrass (*Saccharum giganteum*), grassy arrowhead (*Sagittaria graminea*), water pimpernel  
17 (*Samolus ebracteatus*), sand cordgrass (*Spartina bakeri*), Virginia chain fern (*Woodwardia virginica*),  
18 and yellow-eyed grasses (*Xyris* spp.).

19 Epiphytes are infrequent and include oak mistletoe (*Phoradendron leucarpum*) and Spanish moss  
20 (*Tillandsia usneoides*).

21 Vines include yellow jessamine (*Gelsemium sempervirens*), Japanese climbing fern (*Lygodium*  
22 *japonicum*), and earleaf greenbrier (*Smilax auriculata*).

23 Numerous disturbances, both from historical and current events, have reshaped some of the  
24 depression marshes. Fire exclusion has led to woody encroachment of these normally herb-  
25 dominated systems. Hydrological alteration has occurred due to the woody invasion and land-  
26 clearing practices in the surrounding natural areas. Roads and ORV trails have provided easy access  
27 to the areas and with them has come the introduction of trash dumping, exotic species, and rutting  
28 of delicate terrain.

29 Florida Sandhill Cranes (*Grus canadensis pratensis*), Wood Storks (*Mycteria americana*), and the  
30 hooded pitcher-plant (*Sarracenia minor*) were observed in depression marshes on the CFG during  
31 this survey.

32 In addition, three invasive exotic species were noted. Two plant species—cogon grass (*Imperata*  
33 *cylindrica*) and Japanese climbing fern (*Lygodium japonicum*)—were observed, and one animal  
34 species—feral hogs (*Sus scrofa*)—was documented.

### 35 **Fire Regime**

36 Depression marshes require frequent, light-intensity fires to reduce woody encroachment and  
37 maintain a high herbaceous species component. The natural fire return interval for depression  
38 marshes is every one year to eight years. For depression marshes encroached by woody species,  
39 prescribed burns should be implemented more often (on a one-year to three-year cycle) to reduce

1 the woody species abundance. In addition, fires from surrounding communities should be allowed to  
2 creep into the depression marshes to enhance ecotone diversity and to reduce fuel loads, thereby  
3 minimizing the chances of catastrophic fires during drought.

#### 4 **General Management Measures**

5 Decrease woody species abundance with application of prescribed fires, minimize hydrologic and  
6 soil disturbances, maintain quality of hydrology, and remove and control invasive/exotic species.  
7 Frequent prescribed burns during the early lightning season should aid in decreasing woody species  
8 abundance. Unessential roads and ORV trails that border or are within depression marshes should  
9 be limited in use to vehicular traffic to minimize soil disturbances.



# 1 Dome Swamp

2 **Global and State Rank: NA**


3 **Desired Future Condition**

4 Dome swamps are isolated, forested, depression wetlands  
5 occurring within a fire-maintained matrix, such as mesic  
6 flatwoods. The characteristic dome appearance is created  
7 by smaller trees that grow on the outer edge (shallower  
8 water and less peat) and larger trees that grow in the  
9 interior. Pond cypress (*Taxodium ascendens*) typically will  
10 dominate, but swamp tupelo (*Nyssa sylvatica biflora*) also  
11 may form a pure stand or occur as a co-dominant. Other  
12 subcanopy species can include red maple (*Acer rubrum*),  
13 dahoon holly (*Ilex cassine*), swamp bay (*Persea palustris*),  
14 sweetbay (*Magnolia virginiana*), and loblolly bay (*Gordonia*  
15 *lasianthus*). Shrubs can be absent to moderate (a function  
16 of fire frequency) and can include Virginia willow (*Itea*  
17 *virginica*), fetterbush (*Lyonia lucida*), buttonbush  
18 (*Cephalanthus occidentalis*), wax myrtle (*Myrica cerifera*),  
19 and titi (*Cyrilla racemiflora*). An herbaceous component  
20 can range from absent to dense and includes ferns,  
21 maidencane (*Panicum hemitomon*), sawgrass (*Cladium*  
22 *jamaicense*), sedges, lizard's tail (*Saururus cernuus*), and  
23 sphagnum moss (*Sphagnum spp.*). Vines and epiphytes will  
24 be found commonly.

25 Maintaining the appropriate hydrology and fire frequency  
26 is critical for preserving the structure and species  
27 composition of the community. Dome swamps should be  
28 allowed to burn on the same frequency as the adjacent fire-  
29 type community, allowing fires to naturally burn across  
30 ecotones. Fires should be appropriately planned to avoid  
31 high-severity fuel consumption within the dome swamp.

32 **Description and Assessment**

33 The term “dome” comes from the profile presented, with  
34 the larger trees growing in the interior where water is  
35 deepest and the smaller trees growing around the edges.  
36 Dome swamps may have peat soils, which are thickest  
37 toward the center of the dome and are underlain with  
38 acidic soils. Dome swamps receive water from rainfall,  
39 runoff from adjoining uplands, and near-surface  
40 groundwater. Dome swamps are distinguished from basin



**Global and State Rank: NA**

**Fire Interval: NA**

**Listed Species:**

- Spiny-pod

**Invasive Exotic Species:**

- Skunk vine
- Torpedo grass
- Soldier's orchid
- Tropical soda apple
- Camphor tree

**Management Practices:**

- Restoration of ecotones
- Introduction of prescribed fire
- Maintenance of hydrology
- Exotic minimization





1 swamps primarily by having a more circular shape, smaller size, and shallower depth. During times  
2 of low rainfall, fires may occur more frequently within dome swamps than in the larger, typically  
3 deeper, basin swamps.

4 In total, as presented in Table 8, dome swamps represent 298 acres of the CFG, or 0.3 percent of  
5 CFG total area. Within the CFG, dome swamps occur north and south of the Buckman Lock and on  
6 Inglis Island. Typically, dome swamps on the CFG have an overstory of pond cypress with longleaf  
7 pine (*Pinus palustris*) and loblolly pine (*Pinus taeda*) sometimes present. Species in the subcanopy  
8 layer include red maple, loblolly bay, and swamp tupelo. Shrubs can be denser than the overstory,  
9 and many of the same species that occur in the upper strata also are found in the lower strata. In  
10 addition to the aforementioned trees, species documented within the shrub strata in the dome  
11 swamps include dahoon holly, cabbage palm (*Sabal palmetto*), fetterbush, sweetbay, wax myrtle,  
12 swamp bay, and saw palmetto (*Serenoa repens*). The herbaceous layers in dome swamps vary in  
13 abundance from site to site, depending on available sunlight. Sawgrass, cinnamon fern (*Osmunda*  
14 *cinnamomea*), maidencane, beaksedges (*Rhynchospora* spp.), and Virginia chain fern (*Woodwardia*  
15 *virginiana*) are common. Vines include laurel greenbrier (*Smilax laurifolia*), and muscadine (*Vitis*  
16 *rotundifolia*).

17 Five exotic plant species—skunk vine (*Paederia foetida*), torpedo grass (*Panicum repens*), soldier's  
18 orchid (*Zeuxine strateumatica*), tropical soda apple (*Solanum viarum*), and camphor tree  
19 (*Cinnamomum camphora*)—occur within a disturbed dome swamp on Inglis Island. Historical and  
20 current disturbances that have impacted dome swamps include fire exclusion, woody encroachment,  
21 clearing, forestry operations in surrounding uplands, and exotic plant invasion.

## 22 **Fire Regime**

23 Fire is essential for the maintenance of dome swamps. Without periodic fire, hardwood invasion and  
24 peat accumulation can cause the dome swamp to succeed to a baygall. Fire frequency is greatest at  
25 the periphery of a dome swamp, where a normal fire cycle might be as short as three years to five  
26 years. Fires may occur as infrequently as every 50 years to 150 years in the interior portions.

## 27 **General Management Measures**

28 Management activities in dome swamps on the CFG should include restoration of ecotones,  
29 introduction of prescribed fire, minimization of hydrologic and soil disturbances, maintenance of  
30 hydrology, and eradication of exotic species. Restoration of firebreaks, closure of drainage channels,  
31 and closure of unessential roads and vehicular trails that border the dome swamps will help to  
32 minimize soil disturbances. Burning around dome swamps during years of normal precipitation (as  
33 opposed to drought years) can reduce heavy fuel loads that can facilitate catastrophic fires and  
34 resulting muck fires.

# 1 Floodplain Marsh

## 2 Global and State Rank: NA

### 3 Desired Future Condition

4 Floodplain marshes are herbaceous-dominated wetlands  
5 associated with rivers. They occur in association with  
6 floodplain swamps, basin swamps, and ruderal  
7 communities.


### 8 Description and Assessment

9 In total, as shown in Table 8, floodplain marsh represents  
10 a total of 245 acres of the CFG, or 0.3 percent of total CFG  
11 area. Floodplain marshes within the CFG can be found  
12 along the St. Johns River; along Deep Creek; and north and  
13 south of the Rodman Reservoir.

14 Floodplain marshes are mostly herb dominated. Bald  
15 cypress (*Taxodium distichum*) was the only tree species  
16 observed when there was an overstory present. Shrubs  
17 may be sparse or abundant, depending on the timing of the  
18 last fire. Shrubs include groundsel tree (*Baccharis*  
19 *halimifolia*), common buttonbush (*Cephalanthus*  
20 *occidentalis*), dahoon holly (*Ilex cassine*), wax myrtle  
21 (*Myrica cerifera*), elderberry (*Sambucus nigra* subsp.  
22 *canadensis*), and coastalplain willow (*Salix caroliniana*).

23 The herbaceous layer is the dominant feature, primarily  
24 including yellow pond-lily (*Nuphar advena*), maidencane  
25 (*Panicum hemitomon*), scarlet rosemallow (*Hibiscus*  
26 *coccineus*), swamp dock (*Rumex verticillatus*), marsh fern  
27 (*Thelypteris palustris* var. *pubescens*), bulltongue  
28 arrowhead (*Sagittaria lancifolia*), pickerelweed  
29 (*Pontederia cordata*), giant bulrush (*Scirpus californicus*),  
30 and broadleaf cattail (*Typha latifolia*). Vines are  
31 uncommon and when seen include climbing hempvine  
32 (*Mikania scandens*), and Elliott's aster (*Symphotrichum*  
33 *elliottii*).

34 Previous and current disturbances include clearing,  
35 ditching and canal construction, creation of  
36 impoundments, hydrological alteration, and introduction  
37 of exotic plants.



**Global and State Rank: NA**



**Fire Interval: 2–5 years**

#### **Listed Species:**

- Florida Sandhill Crane
- White Ibis

#### **Invasive Exotic Species:**

- Water lettuce
- Torpedo grass

#### **Management Practices:**

- Restoration of natural hydrology
- Reintroduction of prescribed fire
- Exotic species monitoring and treatment



1 Two species of invasive exotic plants occur within the floodplain marsh, including water lettuce  
2 (*Pistia stratiotes*) and torpedo grass (*Panicum repens*).

3 Two listed bird species were documented within floodplain marsh communities: Florida Sandhill  
4 Crane (*Grus canadensis pratensis*) and White Ibis (*Eudocimus albus*).

#### 5 **Fire Regime**

6 Floodplain marshes are maintained by fire and hydrology. Under natural conditions, fires burn on a  
7 one-year to five-year basis, restricting shrub entry and maintaining the open herbaceous nature of  
8 the marshes. When floodplain marshes burn under drought conditions, the underlying peat also can  
9 burn.

#### 10 **General Management Measures**

11 Management of the floodplain marshes on the CFG includes restoration of natural hydrology (due to  
12 negative impacts stemming from ditching and canal building activities), exotic species monitoring  
13 and treatment, and reintroduction of fire. Natural hydrology is crucial for maintaining species  
14 diversity and water quality. Fires from surrounding uplands should be allowed to creep into the  
15 floodplain marshes to maintain the herb dominance and to keep the woody species from taking over.

# Mesic Hammock


## Global and State Rank: NA

## Desired Future Condition

Mesic hammock is a well-developed evergreen hardwood and/or palm forest that can occur, with variation, through much of peninsular Florida. The often-dense canopy typically will be dominated by live oak (*Quercus virginiana*) with cabbage palm (*Sabal palmetto*) mixed into the understory. Southern magnolia (*Magnolia grandiflora*), sweetgum (*Liquidambar styraciflua*), sugarberry (*Celtis laevigata*), and pignut hickory (*Carya glabra*) can be common components in the subcanopy as well. Slash pines (*Pinus elliotii*) or loblolly pine (*Pinus taeda*) may be sparsely distributed in the canopy. The shrubby understory may be dense or open, tall or short, and is typically composed of saw palmetto (*Serenoa repens*), beautyberry (*Callicarpa Americana*), yaupon (*Ilex vomitoria*), American holly (*Ilex opaca*), gallberry (*Ilex glabra*), common persimmon (*Diospyros virginiana*) and sparkleberry (*Vaccinium arboretum*). The groundcover may be sparse and patchy but generally contains panicgrasses (*Panicum* spp.), switchgrass (*Panicum virgatum*), sedges, and various ferns and forbs. Abundant vines and epiphytes occur on live oaks, cabbage palms, and other subcanopy trees. Mesic hammocks generally will contain sandy soils with organic materials and may have a thick layer of leaf litter at the surface. Mesic hammocks are rarely inundated and not considered to be fire-adapted communities, so they are typically shielded from fire.

## Description and Assessment

In total, as illustrated in Table 8, the mesic hammock communities represent a total of 4,313 acres of the CFG, or 6 percent of total CFG area. High-quality mesic hammocks occur within the CFG south of CR 316 and east of the Ocklawaha River; within the Ross Prairie depression marsh area; and on the western portion of Inglis Island. Mesic hammocks also are located in the Caravelle Wildlife Management Area; in areas such as north of the Buckman Lock; north, south, and east of the Rodman Reservoir; and along Deep Creek. Other areas within mesic hammocks include the vicinity of the Orange Springs Recreation Area;



**Global and State Rank: NA**

**Fire Interval: NA**

### Listed Species:

Coontie, coastal vervain, Florida spiny pod, anglepod, pinewood dainties, green-fly orchid, gopher tortoise, cinnamon fern, mountain azalea, needle palm, spiny-pod, toothpetal false rein orchid

### Invasive Exotic Species:

Mimosa, coral ardisia, paper mulberry, camphor tree, air potato, cogon grass, lantana, Japanese climbing fern, cat's claw fern, Chinaberry, heavenly bamboo, sword fern, skunk vine, golden bamboo, Chinese brake fern, tropical soda apple, Caesar weed, wisteria

### Management Practices:

- Restoration of natural hydrology
- Exotic species monitoring and treatment



1 from north of the Eureka Bridge south to Gores Landing; from Turkey Landing south to US 40; from  
2 Forest Corners east to the Ocklawaha River (north of US 40); Marshall Swamp, from Historic Santos  
3 Recreation Area west to I-75; in the Diggings (from I-75 west to Pruitt Trailhead); from Dunnellon  
4 Baseball Fields and Recreation Complex west to US 41 (south of Withlacoochee River/Lake  
5 Rousseau); Inglis Island; and north and south of the Inglis Spillway.

6 The canopy can be open or closed. The overstory of the mesic hammock is closed to semi-closed, and  
7 typical species include live oak, laurel oak (*Quercus hemisphaerica*), sweet gum, pignut hickory,  
8 Southern magnolia, and occasionally loblolly pine. American hornbeam (*Carpinus caroliniana*),  
9 common persimmon (*Diospyros virginiana*), American holly, Carolina laurelcherry (*Prunus*  
10 *caroliniana*), black cherry (*Prunus serotina*), and cabbage palm are among the species that can  
11 sometimes be found in the subcanopy. The understory is shrubby and may be dense or open, tall or  
12 short. Epiphytes (ferns, orchids, and bromeliads) often are found and may become abundant in  
13 undisturbed stands.

14 Common shrubs include Florida hobblebush (*Agarista populifolia*), indigobush (*Amorpha fruticosa*),  
15 devil's walkingstick (*Aralia spinosa*), coral ardisia (*Ardisia crenata*), woolly pawpaw (*Asimina*  
16 *incana*), smallflower pawpaw (*Asimina parviflora*), eastern redbud (*Cercis canadensis*), hawthorn  
17 (*Crataegus* sp.), upland swampprivet (*Forestiera ligustrina*), blue huckleberry (*Gaylussacia frondosa*  
18 var. *tomentosa*), St. Andrew's cross (*Hypericum hypericoides*), yaupon, wild coffee (*Psychotria*  
19 *nervosa*), myrsine (*Rapanea punctata*), Carolina buckthorn (*Rhamnus caroliniana*), winged sumac  
20 (*Rhus copallinum*), smallflower mock buckthorn (*Sageretia minutiflora*), saw palmetto (*Serenoa*  
21 *repens*), gum bully (*Sideroxylon lanuginosum*), sparkleberry (*Vaccinium arboreum*), deerberry  
22 (*Vaccinium stamineum*), hog plum (*Ximenia americana*), and Hercules' club (*Zanthoxylum clava-*  
23 *herculis*).

24 The herb layer often is sparse or patchy and consists of various grasses. Typical herbs are woods  
25 grass (*Oplismenus hirtellus*), partridgeberry (*Mitchella repens*), lender woodoats (*Chasmanthium*  
26 *laxum*), variable witchgrass (*Dichantheium commutatum*), eggleaf witchgrass (*Dichantheium ovale*),  
27 and bedstraws (*Galium* spp.).

28 Epiphytes are common, and include Florida butterfly orchid (*Encyclia tampensis*), resurrection fern  
29 (*Pleopeltis polypodioides* var. *michauxiana*), Bartram's air-plant (*Tillandsia bartramii*), ballmoss  
30 (*Tillandsia recurvata*), and Spanish moss (*Tillandsia usneoides*).

31 Vines are diverse and abundant, and include rattan vine (*Berchemia scandens*), trumpet creeper  
32 (*Campsis radicans*), air-potato (*Dioscorea bulbifera*), yellow jessamine (*Gelsemium sempervirens*),  
33 cat's claw vine (*Macfadyena unguis-cati*), Virginia creeper (*Parthenocissus quinquefolia*), earleaf  
34 greenbrier (*Smilax auriculata*), saw greenbrier (*Smilax bona-nox*), sarsaparilla vine (*Smilax pumila*),  
35 bristly greenbrier (*Smilax tamnoides*), eastern poison ivy (*Toxicodendron radicans*), muscadine (*Vitis*  
36 *rotundifolia*), and calloose grape (*Vitis shuttleworthii*).

37 Four listed plant species were documented within mesic hammocks on CFG: Florida spiny pod  
38 (*Matelea floridana*), anglepod (*Matelea gonocarpos*), pinewoods dainties (*Phyllanthus liebmannianus*  
39 spp. *platylepis*), and green-fly orchid (*Epidendrum conopseum*). One listed animal species, the gopher  
40 tortoise (*Gopherus polyphemus*), was observed in a few areas, indicating some mesic hammocks may  
41 be historically more open, grassy mesic flatwoods.



1 On the CFG, 18 invasive exotic plant species occur in mesic hammocks: mimosa (*Albizia julibrissin*),  
2 coral ardisia (*Ardisia crenata*), paper mulberry (*Broussonetia papyrifera*), camphor tree  
3 (*Cinnamomum camphora*), air-potato (*Dioscorea bulbifera*), cogon grass (*Imperata cylindrica*),  
4 Lantana (*Lantana camara*), Japanese climbing fern (*Lygodium japonicum*), cat's claw vine, Chinaberry  
5 (*Melia azedarach*), heavenly bamboo (*Nandina domestica*), sword fern (*Nephrolepis cordifolia*), skunk  
6 vine (*Paederia foetida*), golden bamboo (*Phyllostachys aurea*), Chinese brake fern (*Pteris vittata*),  
7 tropical soda apple (*Solanum viarum*), Caesar weed (*Urena lobata*), and wisteria (*Wisteria sinensis*).

8 Mesic hammocks have undergone considerable disturbance from human activities, as these habitats  
9 are desirable home, camp, and recreation sites. Past logging, understory clearing, cattle grazing, and  
10 introduction of feral hogs have altered natural canopies and disturbed soils. Cattle trample  
11 understory plants as they take refuge in shaded oak hammocks, and rooting by hogs causes severe  
12 soil disturbance. These activities leave hammocks vulnerable to invasion by a wide variety of exotic  
13 invasive plants, which compete with native plants and often become the dominant ground or vine  
14 cover.

### 15 **Fire Regime**

16 Although the mesic hammock generally is not considered a fire-adapted community, some small  
17 patches of hammock occurring as islands within marshes or prairies may experience occasional low-  
18 intensity ground fires.

### 19 **General Management Measures**

20 Given that mesic hammocks often are associated with various types of wetlands, either occurring as  
21 a matrix with hydric communities or as a transition to uplands, the hammocks may be sensitive to  
22 hydrologic alteration in the landscape. More frequent and prolonged flooding will kill most  
23 characteristic mesic hammock tree species. Lowered water tables will either shift vegetation to more  
24 xeric species or allow intense fires to burn and destroy the hammock, particularly where surrounding  
25 uplands have been fire excluded in the past. Placement of firebreaks around areas of the hammock  
26 often disrupts the natural ecotone with surrounding pyrogenic communities, leading to invasion of  
27 pine-dominated communities with hardwoods. Protection of mesic hammock habitats must,  
28 therefore, include limitations on development and grazing, restoration of natural fire regime and  
29 hydrology in the overall landscape, and control of invasive species.



# 1 Tidal Marsh

2 **Global and State Rank: NA**

3 **Desired Future Condition**

4 Tidal marshes are characterized as expanses of grasses,  
5 rushes, and sedges along coastlines with low wave energy  
6 and at river mouths. Tidal marsh soils generally are very  
7 poorly drained muck or sandy clay loams. The elevation of  
8 tidal marshes ranges from just below sea level to slightly  
9 above sea level.

10 **Description and Assessment**


11 In total, as presented in Table 8, tidal marshes represent  
12 4,285 acres of the CFG, or 6 percent of total CFG area.  
13 Tidal marshes occur in the western portion of the CFG in  
14 association with the Gulf of Mexico.

15 Typically, tidal marshes on the CFG are dominated by black  
16 needle rush (*Juncus roemerianus*), sawgrass (*Cladium*  
17 *jamaicense*), with saltgrass flats (*Distichlis spicata*) in  
18 shallower areas. Other typical plant species include  
19 Carolina sealavender (*Limonium carolinianum*), big  
20 cordgrass (*Spartina cynosuroides*), starrush whitetop  
21 (*Rhynchospora colorata*), sugarcane plumegrass  
22 (*Saccharum giganteum*), bulltongue arrowhead (*Sagittaria*  
23 *lancifolia*), gulf cordgrass (*Spartina spartinae*), and  
24 broadleaf cattail (*Typha latifolia*). Shrubs are few, mainly  
25 restricted to higher ground, and include silverling  
26 (*Baccharis glomeruliflora*), salt wort (*Batis maritima*), red  
27 cedar (*Juniperus virginiana*), christmasberry (*Lycium*  
28 *carolinianum*), and wax myrtle (*Myrica cerifera*).

29 The Wood Stork (*Mycteria americana*) was observed  
30 within the tidal marsh. Only torpedo grass (*Panicum*  
31 *repens*) was documented within CFG tidal marsh areas.

32 **Fire Regime**

33 It is likely that tidal marshes burn primarily along the  
34 edges near the shoreline, where there are adjacent upland  
35 pyrogenic natural communities, such as flatwoods or wet  
36 prairies. These fires would help reduce woody  
37 encroachment into the marshes from the uplands.



**Global and State Rank: NA**

**Fire Interval: NA**

**Listed Species:**

Wood Stork

**Invasive Exotic Species:**

Torpedo grass

**Management Practices:**

Prescribed fire



1 **General Management Measures**

- 2 Increasing development pressures along Florida’s coasts represent the largest threat to the tidal  
3 marshes. Among the potential problems that could affect the tidal marshes on the CFG are  
4 diminishment of the quality and quantity of fresh water inputs, and pollution from offshore sources.

# 1 Upland Hardwood Forest

## 2 Global and State Rank: NA


### 3 Desired Future Condition

4 Upland hardwood forests are mature, closed-canopy,  
5 hardwood forests typically occurring on slopes and rolling  
6 hills with generally mesic conditions. Overstory tree  
7 species may consist of Southern magnolia (*Magnolia*  
8 *grandiflora*), pignut hickory (*Carya glabra*), sweetgum  
9 (*Liquidambar styraciflua*), live oak (*Quercus virginiana*),  
10 laurel oak (*Quercus laurifolia*), Florida maple (*Acer*  
11 *saccharinum* subsp. *floridanum*), spruce pine (*Pinus*  
12 *glabra*), and swamp chestnut oak (*Quercus michauxii*).  
13 Understory species will include trees and shrubs such as  
14 American holly (*Ilex opaca*), flowering dogwood (*Cornus*  
15 *florida*), eastern hophornbeam (*Ostrya virginiana*),  
16 American hornbeam (*Carpinus caroliniana*), eastern  
17 redbud (*Cercis canadensis*), red bay (*Persea borbonia*),  
18 horse sugar (*Symplocos tinctoria*), and American  
19 beautyberry (*Callicarpa americana*). Ground cover will  
20 comprise shade-tolerant herbaceous species, sedges, and  
21 vines.

### 22 Description and Assessment

23 In total, as shown in Table 8, upland hardwood forests  
24 represent 350 acres of the CFG, or 0.3 percent of total  
25 CFG area. Upland hardwood forests occur on the CFG along  
26 the east side of the Ocklawaha River in the vicinity of  
27 Butterbutt Landing, where they are associated with and  
28 intergrade into mesic hammock, floodplain swamp, and  
29 ruderal communities.

30 The forest is a diverse assemblage of evergreen and  
31 deciduous tree species in the canopy and midstory, shade-  
32 tolerant shrubs, and a sparse groundcover. Characteristic  
33 canopy trees are sand live oak (*Quercus geminata*), live  
34 oak, southern magnolia, laurel oak, pignut hickory, loblolly  
35 pine (*Pinus taeda*), cabbage palm (*Sabal palmetto*), and  
36 sweetgum. The subcanopy includes the aforementioned  
37 species, plus American hornbeam (*Carpinus caroliniana*)  
38 and devil's walking stick (*Aralia spinosa*). Among the  
39 shrubs are American beautyberry, wild olive (*Osmanthus*



**Global and State Rank: NA**

**Fire Interval: NA**

**Listed Species:**

Needle palm

**Invasive Exotic Species:**

Paper mulberry

**Management Practices:**

Invasive exotic treatment/control



1 *americana*), bluestem palmetto (*Sabal minor*), dwarf palmetto (*Sabal minor*), and sparkleberry  
2 (*Vaccinium arboreum*).

3 Few herbs persist in the dense shade, primarily eggleaf witchgrass (*Dichantheium ovale*), woodoats  
4 (*Chasmanthium laxum*), and Caesar weed (*Urena lobata*). Spanish moss (*Tillandsia usneoides*) can be  
5 found in the canopy.

6 Vines include yellow jessamine (*Gelsemium sempervirens*), saw greenbrier (*Smilax bona-nox*), and  
7 muscadine (*Vitis rotundifolia*).

8 The needle palm (*Rhapidophyllum hystrix*) was documented within the CFG upland hardwood forest.

9 Past logging, clearing, and roads and trails have altered natural canopies and disturbed soils, which  
10 likely has encouraged the introduction of invasive species into the hardwood forest at the CFG.  
11 Invasive exotics present within upland hardwood forests include the paper mulberry (*Broussonetia*  
12 *papyrifera*).

### 13 **Fire Regime**

14 Upland hardwood forests are not pyrogenic. The dense canopy and midstory results in low light and  
15 air movement and high relative humidity; thus, fires in adjacent uplands, such as sandhills, extinguish  
16 at the edge of the upland hardwood forest under normal moisture conditions. Localized damage to  
17 upland hardwood forest as a result of low-intensity, naturally occurring fires that creep into the  
18 forest edges from surrounding pyrogenic upland communities appears to be a natural part of the  
19 forest dynamics of this community, but fires should rarely, if ever, burn completely through the  
20 understory.

### 21 **General Management Measures**

22 Damage from invasive exotic plants and animals is a common problem in upland hardwood forest on  
23 the CFG. Control of these pests is the primary management activity needed. For specific control  
24 measures for camphor tree and Caesar weed please refer to the exotic and nuisance species sections.

# 1 Blackwater Stream

## 2 Global and State Rank: NA


### 3 Desired Future Condition

4 Blackwater streams are characterized as perennial or  
5 intermittent watercourses originating in lowlands where  
6 extensive wetlands with organic soils collect rainfall and  
7 runoff, discharging it slowly to the stream. The stained  
8 waters are laden with tannins, particulates, and dissolved  
9 organic matter derived from drainage through adjacent  
10 swamps resulting in sandy bottoms overlain by organic  
11 matter. Emergent and floating vegetation (including  
12 golden club (*Orontium aquaticum*), smartweeds  
13 (*Polygonum spp.*), grasses and sedges) may occur but is  
14 often limited by steep banks and dramatic seasonal  
15 fluctuations in water levels. Desired conditions include  
16 minimizing disturbance and alterations and preserving  
17 adjacent natural communities.

### 18 Description and Assessment

19 In total, as presented in Table 8, blackwater streams  
20 represent 192 acres of the CFG, or 0.2 percent of CFG  
21 total area. There are numerous named and unnamed  
22 blackwater streams on the CFG. Some of the named  
23 waterways include Deep Creek, Sweetwater Creek, Orange  
24 Creek, Cedar Creek, Eaton Creek, Turkey Creek, Ocklawaha  
25 River, St. Johns River, Withlacoochee River, and the Dead  
26 River. Within the CFG, blackwater streams occur within  
27 baygall, hydric hammock, mesic hammock, mesic  
28 flatwoods, and floodplain swamp communities.

29 Primary vegetation associated with blackwater streams on  
30 the CFG originates along the watercourse banks or edges.  
31 The canopy density ranges from open to closed, and can  
32 include red maple (*Acer rubrum*), Carolina ash (*Fraxinus  
33 caroliniana*), green ash (*Fraxinus pennsylvanica*), bald  
34 cypress (*Taxodium distichum*), and cabbage palm (*Sabal  
35 palmetto*). The shrub layer includes several of the same  
36 species that are found in the overstory strata in addition to  
37 common buttonbush (*Cephalanthus occidentalis*), swamp  
38 dogwood (*Cornus foemina*), St. Andrew's cross (*Hypericum  
39 hypericoides*), dahoon holly (*Ilex cassine*), coastalplain



**Global and State Rank: NA**

**Fire Interval: NA**

**Listed Species: NA**

#### **Invasive Exotic Species:**

Torpedo grass, water lettuce, hydrilla, water hyacinth, wild taro, alligator weed, Caesar's weed, Peruvian primrose-willow

#### **Management Practices:**

- Restoration and maintenance of natural hydrologic patterns
- Maintaining existing water quality and quantity
- Invasive and exotic minimization





1 willow (*Salix caroliniana*), and Gulf Sebastian bush (*Sebastiania fruticosa*).

2 Given fluctuating water levels, herbs typically are sparse along these river banks, but more open-  
3 canopied sites can support herbaceous groundcover. Species include false nettle (*Boehmeria*  
4 *cylindrica*), sawgrass (*Cladium jamaicense*), dogfennel (*Eupatorium capillifolium*), yellow pond-lily  
5 (*Nuphar advena*), royal fern (*Osmunda regalis* var. *spectabilis*), maidencane (*Panicum hemitomon*),  
6 green arrow arum (*Peltandra virginica*), narrowfruit horned beaksedge (*Rhynchospora inundata*),  
7 bulltongue arrowhead (*Sagittaria lancifolia*), and southern shield fern (*Thelypteris kunthii*).

8 The epiphytic and vine layers of the blackwater streams are infrequent. Bartram's air-plant  
9 (*Tillandsia bartramii*) is the most common epiphyte and sometimes occurs in great profusion, giving  
10 trees a "hairy" appearance. Vines include climbing hempvine (*Mikania scandens*), saw greenbrier  
11 (*Smilax bona-nox*), and eastern poison ivy (*Toxicodendron radicans*).

12 Four listed animal species were observed in blackwater streams: limpkin (*Aramus guarauna*), Little  
13 Blue Heron (*Egretta caerulea*), White Ibis (*Eudocimus albus*), and American alligator (*Alligator*  
14 *mississippiensis*). Invasive plants documented in and along blackwater streams include torpedo grass  
15 (*Panicum repens*), water lettuce (*Pistia stratiotes*), hydrilla (*Hydrilla verticillata*), water hyacinth  
16 (*Eichhornia crassipes*), wild taro (*Colocasia esculenta*), alligator weed (*Alternanthera philoxeroides*),  
17 Caesar's weed (*Urena lobata*) and Peruvian primrose-willow (*Ludwigia peruviana*).

18 Human disturbances such as clearing and excavation for the Barge Canal, hydrological alteration, and  
19 creation of roads have changed not only the natural course of the larger streams throughout the CFG,  
20 but the vegetation species composition, structure, and hydrology from the historic condition as well.  
21 More recently created disturbances include trash dumping and exotic plant and animal invasions.

## 22 **Fire Regime**

23 Fire is not a requirement for blackwater streams. They may function as natural firebreaks when  
24 adjacent to flatwoods or other communities that do require fire.

## 25 **General Management Measures**

26 Pertinent management objectives for blackwater streams on the CFG include restoration and  
27 maintenance of natural hydrologic patterns, monitoring and control of invasive exotic plants and  
28 animals, and maintaining existing water quality and quantity.

29 Management of water on public conservation lands today includes continued monitoring of and input  
30 into growth management and regulatory processes, with respect to increasing demands for fresh  
31 water from local and regional developments, as well as from neighbors across state lines.



# 1 **Altered Land Cover Types**

2 **Global and State Rank: NA**

**Global and State Rank: NA**

3 **Desired Future Condition**

**Fire Interval: NA**

4 ***Developed Areas***

**Listed Species: NA**

5 The developed areas within the park will be managed to  
6 minimize the effect of the developed areas on adjacent  
7 natural areas. Priority invasive plant species (FLEPPC  
8 Category I and II species) will be controlled from all  
9 developed areas.

**Management Practices:**

- Removal of FLEPPC Category I and II priority invasive exotic species

10 ***Clearcut Pine Plantation***

11 Clearcut pine plantation is an altered community type that  
12 resulted from past commercial management. The  
13 dominant overstory species found in this type is longleaf  
14 pine (*Pinus palustris*), loblolly pine (*Pinus taeda*), or slash  
15 pine (*Pinus elliotti*). As an altered forest type, and  
16 potentially a candidate for restoration, there is currently  
17 no FNAI recommendations on preferred species or  
18 stocking levels for this natural community. Future timber  
19 management activities potentially could transition this  
20 altered type into another natural community type.

21 ***Pine Plantations***

22 These are areas altered by silvicultural activities. These  
23 include lands where either: (1) planted pines are having or  
24 will have an ongoing detrimental effect on native  
25 groundcover, (2) the history of planted pines has damaged  
26 ground cover to the point where further restoration  
27 beyond thinning and burning is required, and/or (3) the  
28 method of planting (e.g., bedding) has severely impacted  
29 groundcover. Pine plantations in Florida often are  
30 dominated by even-aged loblolly (*Pinus taeda*), sand pine  
31 (*Pinus clausa*), or slash pine (*Pinus elliottii*). Dense pine  
32 plantations typically have sparse to absent herbaceous  
33 vegetation as a result of shading or a cover of deep pine  
34 needle duff. These plantations may be very shrubby or  
35 vine-dominated or open at ground level. The groundcover  
36 in most cases has been severely impacted by mechanical  
37 site preparation, such as roller chopping and bedding.  
38 However, while perennial grasses such as wiregrass  
39 (*Aristida stricta* var. *beyrichiana*) may be greatly reduced,

1 many components of the native groundcover persist even though the relative abundance is altered.  
2 Groundcover can be partially restored by thinning and/or frequent burning, although some planting  
3 of perennial grasses such as wiregrass may be required. With activities such as thinning and burning,  
4 plantations with intact native groundcover can be restored to the former natural community.

#### 5 ***Abandoned Field/ Abandoned Pasture***

6 Old fields, fallow pastures, early successional areas formerly grazed or in agriculture without recent  
7 activity to maintain the area as pasture or planted field make up this category. These areas often are  
8 dominated by weedy native species (e.g., *Rubus spp.*, *Myrica cerifera*) and non-native species (e.g.,  
9 *Indigofera hirsuta*). In old pastures, generally designated when weedy cover from woody species  
10 (*Rubus spp.*, *Myrica cerifera*, etc) is greater than 20 percent.

#### 11 **Description and Assessment**

12 The altered areas have been included in the community types in which they occur. These areas  
13 include clearings, impoundment, a utility corridor, ditching and canal construction, and  
14 developments that enhance the greenway visitor experience, including the visitor center,  
15 boardwalks, picnic area with pavilion, main drive, and parking lots.

#### 16 **General Management Measures**

17 The altered areas within the CFG will be managed to remove Florida Exotic Plant Pest Council  
18 (FLEPPC) Category I and II priority invasive exotic plant species. Other management measures may  
19 include limited restoration efforts designed to minimize the effects of the ruderal areas on adjacent  
20 natural areas. Cost-effectiveness and consideration of other higher priority restoration projects  
21 within the park will determine the extent of restoration measures in ruderal areas. The developed  
22 areas within the park will be managed to minimize the effect of the developed areas on adjacent  
23 natural areas.

## CFG Imperiled Species Planning Accomplishments

2007-2016

---

Contracted with FNAI to update surveys of rare plants, including longspurred mint, a federally endangered plant.

---

---

Coordinated with the FWC and the Audubon Society to manage the Spoil Islands for listed shorebird species.

---

---

Continued ongoing documentation of use of the Buckman Lock and Rodman Reservoir by Florida manatees to migrate between the St. Johns River and the Ocklawaha River/Silver River systems.

---

---

Restored 523 acres of Florida Scrub-Jay habitat, for a total of 840 acres out of 1,100 acres of historic habitat. Population increased from 46 birds in 2009 to 111 in 2015.

---

---

Installed wildlife cameras in 2009 to capture wildlife usage at all underpasses on the greenway and on the I-75 land bridge.

---

### 1 **Imperiled Species**

2 For the purposes of this management plan, imperiled  
3 species are species on Florida's Threatened and  
4 Endangered Species List, in accordance with Rules 68A-  
5 27.003 and 68A-27.005, or species protected by  
6 designation under the Federal Endangered Species Act.  
7 Imperiled species also are those species that are tracked  
8 by FNAI as critically imperiled (G1, S1) or imperiled (G2,  
9 S2). Federal agencies that share the authority to list  
10 species as Endangered and Threatened are the National  
11 Oceanic and Atmospheric Administration—National  
12 Marine Fisheries Service (NOAA-NMFS) and the U.S. Fish  
13 and Wildlife Service (USFWS). The NOAA-NMFS is  
14 responsible for the listing of most marine species. The  
15 federal list of animals and plants is administered by the  
16 USFWS and this list is published in 50 Code of Federal  
17 Regulations (CFR) 17 (animals) and 50 CFR 23 (plants).

18 On November 8, 2010, new threatened species rules  
19 approved by the FWC were implemented. All federally  
20 listed species that occur in Florida now will be included on  
21 Florida's list as federally designated endangered species or  
22 federally designated threatened species. In addition, the  
23 state has implemented a listing process to identify species  
24 that are not federally listed, but that may be at risk of  
25 extinction. These species will be called state designated  
26 threatened species. In all, Florida has a total of 57 species  
27 that are considered imperiled, in accordance with the  
28 *Florida's Imperiled Species Management Plan (ISMP) 2016-*  
29 *2026*. For the first time, this plan provides conservation  
30 goals and actions to achieve them for Florida's state-listed  
31 species. Further, the document identifies integrated  
32 conservation strategies and species-specific actions that  
33 could be employed to support populations of threatened  
34 and endangered species.

35 A summary of element and element occurrence status  
36 within the CFG was obtained from the FNAI where  
37 occurrence status was determined solely by the presence  
38 of actual element occurrences within the Biotics database  
39 or by intersections of site boundaries with the Florida  
40 Breeding Bird Atlas (FBBA) polygons. Table 9, below,  
41 provides a list of all known imperiled species within the  
42 CFG and identifies their status as defined by various  
43 entities. It also identifies the types of management actions  
44 that currently are being taken by DRP staff or others and

1 identifies the current level of monitoring effort. The codes used under the column headings for  
 2 management actions and monitoring level are defined following the table. Detailed management  
 3 goals, objectives, and actions for imperiled species in this park are discussed in the Resource  
 4 Management Program section of this component and the Implementation Component of this plan.

5 **Table 9. Imperiled Species on the CFG**

Common and Scientific Name	Imperiled Status				Management Actions	Monitoring Level
	FNAI Global	FNAI State	USFWS	FWC		
Bluenose shiner ( <i>Pteronotropis welaka</i> )	G3/G4	S3/S4	N	SSC	4	Tier 1
Gopher Frog ( <i>Rana capito</i> )	G3	S3	N	SSC	1, 7	Tier 2
American Alligator ( <i>Alligator mississippiensis</i> )	G5	S4	FT (SA)	LS	2, 10	Tier 1
Eastern Indigo Snake ( <i>Drymarchon couperi</i> )	G3/Q	S3	FT	LT	1, 2	Tier 1
Gopher Tortoise ( <i>Gopherus Polyphemus</i> )	G3	S3	C	ST	1, 2	Tier 1
Pine Snake ( <i>Pituophis melanoleucus mugitus</i> )	G4/T3	S3	N	SSC	1, 2	Tier 1
Suwanee Cooter ( <i>Pseudmys concinna suwanniensis</i> )	G5/T3	S3	N	SSC	4	Tier 1
Florida Scrub-Jay ( <i>Aphelocoma coerulescens</i> )	G2	S2	FT	LT	1, 2, 7, 13	Tier 3
Limpkin ( <i>Aramus guarauna</i> )	G5	S3	N	SSC	4	Tier 1
Little Blue Heron ( <i>Egretta caerulea</i> )	G5	S4	N	SSC	4, 10, 13	Tier 2
American Oystercatcher ( <i>Haematopus palliates</i> )	G5	S2	N	SSC	8, 10, 13	Tier 4
Osprey ( <i>Pandicon haliaetus</i> )	G5	S3/S4	N	SSC	5	Tier 2
Least Tern ( <i>Sterna antillarum</i> )	G4	S3	N	ST	8, 10, 13	Tier 3
Florida Mouse ( <i>Podomys floridanus</i> )	G3	S3	N	SSC	1, 2	Tier 1
Sherman's Fox Squirrel ( <i>Sciurus niger shermani</i> )	G5/T3	S3	N	SSC	1	Tier 1
Incised Groove-Bur ( <i>Agrimonia incisa</i> )	G3	S2	N	LE	2	Tier 2

Common and Scientific Name	Imperiled Status				Management Actions	Monitoring Level
	FNAI Global	FNAI State	USFWS	FWC		
Variable-Leaved Indian-Plantain ( <i>Arnoglossum diversifolium</i> )	G2	S2	N	LT	2, 4, 10, 13	Tier 2
Dwarf Spleenwort ( <i>Asplenium pumilum</i> )	G5	S1	N	LE	2, 4	Tier 2
Chapman's Sedge ( <i>Carex chapmanii</i> )	G3	S3	N	LT	2, 4, 10, 13	Tier 2
Longspurred Mint ( <i>Dicerandra cornutissima</i> )	G1	S1	E	FE	1, 2, 7, 13	Tier 3
Coastal Vervain ( <i>Glandularia maritima</i> )	G3	S3	N	E	2	Tier 2
Chapman's Skeletongrass ( <i>Gymnopogon chapmanianus</i> )	G3	S3	N	N	2	Tier 2
Florida Spiny-Pod ( <i>Matelea floridana</i> )	G2	S2	N	LE	2	Tier 2
Garberia ( <i>Garberia heterophylla</i> )	G3/G4	S3/S4	N	LT	2	Tier 2
Sandhill Spiny-Pod ( <i>Matelea pubiflora</i> )	G3/G4	S3/S4	N	LE	2	Tier 2
Pigmy Pipes ( <i>Monotropsis reynoldsiae</i> )	G1	S1	N	E	2, 10	Tier 3
Large-Leaved Grass-of-Parnassus ( <i>Parnassia grandifolia</i> )	G3	S2	N	LE	2, 4, 10, 13	Tier 2
Widespread Polypody ( <i>Pecluma dispersa</i> )	G5	S2	N	E	2	Tier 2
Plume Polypody ( <i>Pecluma plumula</i> )	G5	S2	N	LE	2	Tier 2
Swamp Plume Polypody ( <i>Pecluma ptildodon</i> )	G5?	S2	N	LE	2, 9, 10	Tier 2
Pinewoods Dainties ( <i>Phyllanthus liebmannianus</i> ssp. <i>Platylepis</i> )	G4/T2	S2	N	LE	2, 10	Tier 2
Hooded Pitcherplant ( <i>Sarracenia minor</i> )	G4	S4	N	LT	2, 10	Tier 2
Giant Orchid ( <i>Pteroglossaspis ecristata</i> )	G2/G3	S2	N	LT	1, 2	Tier 2
Florida Willow ( <i>Salix floridana</i> )	G2	S2	N	LE	2, 4, 10, 13	Tier 2

Common and Scientific Name	Imperiled Status				Management Actions	Monitoring Level
	FNAI Global	FNAI State	USFWS	FWC		
Buckthorn ( <i>Sideroxylon lycioides</i> )	G5	S2	N	E	2, 4, 10	Tier 2
Pinkroot ( <i>Spigelia loganioides</i> )	G2/Q	S2	N	LE	2, 4, 10	Tier 2
Scrub Stylisma ( <i>Stylisma abditia</i> )	G3	S3	N	LE	1, 2, 10	Tier 2
Bald Eagle ( <i>Haliaeetus leucocephalus</i> )	G5	S3	N	N	1, 2	Tier 1
Black-Crowned Night-Heron ( <i>Nycticorax nycticorax1</i> )	G5	S3	N	N	10, 13	Tier 2
Eastern Diamondback Rattlesnake ( <i>Crotalus adamanteus</i> )	G4	S3	N	N	1, 2	Tier 1
Florida Black Bear ( <i>Ursus americanus floridanus</i> )	G5/T2	S2	N	N	2, 10, 13	Tier 1
Florida Cebionid Beetle ( <i>Selonodon floridensis</i> )	G2/G4	S2/S4	N	N	2	Tier 1
Florida Long-Tailed Weasel ( <i>Mustela frenata peninsulae</i> )	G5/T3	S3	N	N	2	Tier 1
Florida Olive Hairstreak ( <i>Callophrys gryneus sweadneri</i> )	G5/T2	S2	N	N	2	Tier 1
Florida Sandhill Crane ( <i>Grus Canadensis pratensis</i> )	G5/T2/T <sub>3</sub>	S2/S3	N	ST	2, 4	Tier 1
Florida Scrub Lizard ( <i>Sceloporus woodi</i> )	G2/G3	S2/S3	N	N	1, 2	Tier 1
Hobbs' Cave Amphipod ( <i>Crangonyx hobbsi</i> )	G2/G3	S2/S3	N	N	2, 4, 10	Tier 1
Large-Jawed Cebionid Beetle ( <i>Selonodon mandibularis</i> )	G2/G4	S2/S4	N	N	2	Tier 1



Common and Scientific Name	Imperiled Status				Management Actions	Monitoring Level
	FNAI Global	FNAI State	USFWS	FWC		
Little-Fork Triaenode Caddisfly ( <i>Triaenodes furcellus</i> )	G3	S3	N	N	2	Tier 1
Orange Lake Cave Crayfish ( <i>Procambarus franzi</i> )	G1	S1	N	N	2, 4, 10	Tier 1
Sand Butterfly Pea ( <i>Centrosema arenicola</i> )	G2/Q	S2	N	E	2	Tier 2
Seminole Skipper ( <i>Hesperia attalus slossonae</i> )	G3/G4/T <sub>3</sub>	S3	N	N	2	Tier 1
Short-Tailed Hawk ( <i>Buteo brachyurus</i> )	G4/G5	S1	N	N	1, 2	Tier 1
Snail Bullhead ( <i>Ameiurus brunneus</i> )	G4	S3	N	N	2, 4	Tier 1
Southern Hognose Snake ( <i>Heterodon simus</i> )	G2	S2	N	N	1, 2	Tier 1
Spiked Crested Coralroot ( <i>Hexalectris spicata</i> )	G5	S3	N	LE	2, 4	Tier 2
Tampa Vervain ( <i>Glandularia tampensis</i> )	G3	S3	N	LE	2,4	Tier 2
Umber Shadowfly ( <i>Neurocordulia obsoleta</i> )	G5	S2	N	N	2	Tier 1
White Ibis ( <i>Eudocimus albus</i> )	G5	S4	N	SSC	2, 4	Tier 1
Wilson's Plover ( <i>Charadrius wilsonia</i> )	G5	S2	N	N	2, 10, 13	Tier 2
Wood Stork ( <i>Mycteria americana</i> )	G4	S2	LT	FT	2, 4	Tier 1
Yellow-Crowned Night-Heron ( <i>Nyctanassa violacea</i> )	G5	S3	N	N	2, 8, 10, 13	Tier 2
<p><b>* STATUS/RANK KEY</b></p> <ul style="list-style-type: none"> <li>• <b>Federal Status (USFWS):</b> C = Candidate species for which federal listing agencies have sufficient information on biological vulnerability and threats to support proposing to list the species as Endangered or Threatened, LE = Listed Endangered, LT = Listed Threatened, SAT = Listed Threatened due to similarity of appearance.</li> <li>• <b>State Status (FWC) Plants:</b> LE = Listed Endangered, LT = Listed Threatened, LS = Listed Species of Special Concern, N = Not currently listed, nor currently being considered for listing.</li> <li>• <b>State Status (FWC) Animals:</b> FE=Listed as Endangered Species at the Federal level by the USFWS, FT = Listed as Threatened</li> </ul>						

Common and Scientific Name	Imperiled Status				Management Actions	Monitoring Level
	FNAI Global	FNAI State	USFWS	FWC		
<p>Species at the Federal level by USFWS, FT(S/A) = Federal Threatened due to similarity of appearance, ST = State population listed as Threatened by the FWC. SSC=Listed as Species of Special Concern by the FWC. N= Not currently listed, nor currently being considered for listing.</p> <ul style="list-style-type: none"> <li>• <b>FNAI Global Rank:</b> G1 = Critically Imperiled, G2 = Imperiled, G3 = Very Rare, G4 = Apparently Secure, G5 = Demonstrably Secure, T# = Taxonomic Subgroup; numbers have same definition as G#'s.</li> <li>• <b>FNAI State Rank:</b> S1 = Critically Imperiled, S2 = Imperiled, S3 = Very Rare, S4 = Apparently Secure. ± = Not tracked by FNAI</li> </ul> <p><b>Management Actions:</b></p> <ol style="list-style-type: none"> <li>1. Prescribed Fire</li> <li>2. Exotic Plant Removal</li> <li>3. Population Translocation/Augmentation/Restocking</li> <li>4. Hydrological Maintenance/Restoration</li> <li>5. Nest Boxes/Artificial Cavities</li> <li>6. Hardwood Removal</li> <li>7. Mechanical Treatment</li> <li>8. Predator Control</li> <li>9. Erosion Control</li> <li>10. Protection from visitor impacts (establish buffers)/law enforcement</li> <li>11. Decoys (shorebirds)</li> <li>12. Vegetation planting</li> <li>13. Outreach &amp; Education</li> <li>14. Other</li> </ol> <p><b>Monitoring Level:</b></p> <p><b>Tier 1:</b> Non-Targeted Observation/Documentation: includes documentation of species presence through casual/passive observation during routine park activities (i.e., not conducting species-specific searches). Documentation may be in the form of Wildlife Observation Forms, or other district specific methods used to communicate observations.</p> <p><b>Tier 2:</b> Targeted Presence/Absence: includes monitoring methods/activities that are specifically intended to document presence/absence of a particular species or suite of species.</p> <p><b>Tier 3:</b> Population Estimate/Index: an approximation of the true population size or population index based on a widely accepted method of sampling.</p> <p><b>Tier 4:</b> Population Census: A complete count of an entire population with demographic analysis, including mortality, reproduction, emigration, and immigration.</p> <p><b>Tier 5:</b> Other: may include habitat assessments for a particular species or suite of species or any other specific methods used as indicators to gather information about a particular species.</p>						

1

2 *Listed Species*

3 FNAI conducted surveys on the CFG for about two dozen rare animal species from October 2003 to  
4 May 2004 and located 16 FNAI-tracked species. Among the rarest species located were the southern  
5 hognose snake (*Heterodon simus*; G2/S2; N/N), Florida Scrub-Jay (*Aphelocoma coerulescens*; G2/S2;  
6 LT/LT), gopher tortoise (*Gopherus polyphemus*; G3/S3; N/LS), gopher frog (*Rana capito*; G3G4/S3;  
7 N/LS), Wilson’s Plover (*Charadrius wilsonia*; G5/S2; N/N), American Oystercatcher (*Haematopus*  
8 *palliates*; G5/S2; N/LS), and Wood Stork (*Mycteria Americana*; G5/S2; LE/LE). FNAI also recorded  
9 two wading bird rookeries and a Least Tern rookery. FNAI provided management suggestions for  
10 each species. FNAI also has one or more records for Gulf Hammock dwarf siren (*Pseudobranchius*  
11 *striatus lustricolus*; G5T1/S1; N/N).

12 *Amphibians*

13 The gopher frog is a medium-sized, stocky frog with adults ranging in length from 2.5 inches to 4  
14 inches. Gopher frogs range over most of Florida, with the exception being the Everglades and Florida  
15 Keys. Range-wide, they occur in the Southeastern Gulf and Atlantic Coastal Plain from North Carolina

1 to eastern Louisiana. Six gopher frogs were documented during this inventory and all but one was  
2 caught in the Diggings east of SR 200. The other was caught in a gopher tortoise burrow 1.24 miles  
3 west of SR 200.

#### 4 Mammals

5 Manatees (*Trichechus manatus*) are present on the CFG and frequently wander through Rodman  
6 Reservoir. In response to manatee mortality caused by water control structures operations, manatee  
7 protection grates, acoustic detection devices, and pressure sensors were installed on the lock and  
8 spillway. There is no access for manatees at the western end of the canal, because the lock is not  
9 functional and the spillways are above the water level on the gulf side.

#### 10 Reptiles

11 Gopher tortoises were primarily located on the western segment of the CFG, west of I-75. Some  
12 burrows also were located in the extreme eastern end. As expected, the gopher tortoises were found  
13 in disturbed areas with herbaceous ground cover and sandy substrate, including berms and  
14 powerline cuts, as well as the open natural communities with well-drained sandy substrates, such as  
15 scrub, sandhills, and scrubby flatwoods. Although the population is unknown, it is assumed to be low  
16 due to the lack of an abundance of burrows. However, with continuing restoration of sandhills and  
17 improvement of habitat for gopher tortoises, populations will likely increase.

18 The Southern hognose snake is a relatively small but stocky snake and reaches a maximum body  
19 length of 15 inches to 20 inches. This species is patchily distributed in Panhandle and peninsular  
20 Florida south to the northern edge of Lake Okeechobee. Southern hognose snakes inhabit xeric  
21 uplands, such as sandhills, scrub, and xeric hammocks. One adult Southern hognose snake was caught  
22 on the east side of Ross Prairie.

23 Eastern indigo snakes are the longest snakes in North America, reaching nearly 9 feet in length. These  
24 snakes are denizens of xeric habitats that encompass various wetland communities. Eastern indigo  
25 snakes currently occur patchily throughout Florida and in southern Georgia. Two adult Eastern  
26 indigo snakes were documented on the CFG. The first was a female approximately 4.5 feet in length  
27 crawling across the northern boundary road, south of Highway 484, and into good-quality sandhill.  
28 The second was located on the west side of SR 200 within a dry, open prairie south of the Diggings.

29 Eastern diamondback rattlesnakes are large, heavy-bodied, pit vipers that can reach 6 feet or more  
30 in length. Eastern diamondbacks occur statewide, including in the Keys. Range-wide, they occur in  
31 the Southeastern Coastal Plain from North Carolina to extreme eastern Louisiana. One adult snake  
32 was observed at the edge of a mesic live oak hammock 0.31 miles east-southeast of the Ross Prairie  
33 trailhead.

#### 34 Birds

35 Florida Scrub-Jay is endemic to the low scrub lands of the Florida peninsula, with the largest  
36 populations occurring in Brevard, Highlands, Polk, and Marion counties. Florida Scrub-Jays inhabit  
37 fire-dominated, low oak scrub habitat found on well-drained soils. From the Audobon 2015 Jaywatch  
38 survey, the CFG has approximately 111 Scrub-Jays representing 27 families

39 Limpkin is a medium-sized wading bird with a long, thick, slightly down-curved bill. Apple snails are  
40 an important food item. Range in the United States is chiefly limited to the Florida peninsula. Four

1 Limpkins were documented during this inventory. One was foraging at Rodman Reservoir, while  
2 three others were documented at Sweetwater Creek. One pair was courting and one individual was  
3 foraging.

4 White Ibis is a wading bird with a white body and an orange head. White Ibis nest colonies are located  
5 near water and they feed on invertebrates, fish, and other small vertebrates. A very large feeding  
6 aggregation of more than 100 individuals of multiple age classes was documented on the  
7 northeastern side of Rodman Reservoir. Ibis also were observed nesting within a multi-species  
8 rookery on Lake Rousseau.

9 Wood Stork is a large, heavy, endangered wading bird. Wood Storks are rare to abundant in the  
10 peninsula and Big Bend area of Florida, but generally are rare or lacking in the Panhandle and Florida  
11 Keys. Two Wood Storks were documented foraging on Rodman Reservoir. Also, a Wood Stork was  
12 observed by CFG staff in the Sweetwater Creek area just north of the bridge. No rookeries were  
13 discovered.

14 The rookeries noted by FNAI are at the western end of the CFG. A Black-Crowned Night-Heron  
15 (*Nycticorax nycticorax*) rookery on Spoil Island 7 supports an estimated 50-75 nesting pairs. FNAI  
16 also documented a colony of 75-100 Least Terns (*Sterna antillarum*) on the western end of Spoil  
17 Island 8. No nests were found, but the terns mobbed human observers, possibly indicating a strong  
18 attraction to the area. FNAI also recorded a wading bird rookery with about 100 pairs of nesting birds  
19 on a willow island on the eastern end of Lake Rousseau. White Ibis occupied one half of the island,  
20 and Great Egrets (*Ardea alba*), Cattle Egrets (*Bubulcus ibis*), Little Blue Herons (*Egretta caerulea*), and  
21 Tricolored Herons (*E. tricolor*) occupied the other half. In cooperation with FWC, warning signs were  
22 posted in the rookery areas on the Spoil Islands during critical seasons.

23 Further, the National Audubon Society and Birdlife International have recognized the Citrus County  
24 Spoil Islands at the mouth of the CFG and multiple tracts comprising the Big Bend Ecosystem as  
25 Globally Significant Important Bird Areas.

26 In addition to the above-mentioned birds, the American Oystercatcher (*Haematopus palliatus*) and  
27 Wilson's Plover (*Charadrius wilsonia*) have been documented by the Audubon Society as nesting on  
28 the islands. There also are more than 35 species of shorebirds that use the area during the non-  
29 breeding season. The islands mark the southern end of the area used by the second largest wintering  
30 concentration of American Oystercatchers in their range.

31 The FWC has developed a species action plan for beach-nesting birds, including the American  
32 Oystercatcher. Plan objectives focus on protecting and monitoring nesting sites where  
33 Oystercatchers are particularly susceptible to direct harm and disturbance. Monitoring is crucial for  
34 measuring the effectiveness of conservation efforts and progress toward achieving the plan  
35 objectives.

36 Breeding productivity is monitored semi-annually by Audubon of Florida and the FWC. Table 10  
37 provides nesting data from the Audubon Society counts on the dredge Spoil Islands.

1 **Table 10. Nesting Data on the CFG Dredge Spoil Islands**

Year	Breeding Pairs	Fledglings	Productivity (Fledglings/Pair)
<b>American Oystercatcher</b>			
2001	45	U	U
2011	15	U	U
2012	27	6	0.22
2013	16	4	0.25
2014	14	1	0.07
2015	20	4	0.20
2016	17	5	0.29
<b>Wilson's Plover</b>			
2011	15	Y	U
2012	12	Y	U
2013	6	Y	U
2014	2	Y	U
2015	6	Y	U
2016	7	Y	U
<b>Least Tern</b>			
2011	68	0	0
2012	53	0	0
2013	61	0	0
2014	33	0	0
2015	23	0	0
2016	34	0	0
<b>Notes</b>			
<ol style="list-style-type: none"> <li>1. Part of the statewide AMOY survey</li> <li>2. Monitored via boat by Audubon - no walking the islands</li> <li>3. All fledges lost to Debby</li> <li>4. Infrequent surveys, underestimated nesting pairs?</li> <li>5. Infrequent surveys, underestimated nesting pairs?</li> <li>6. High density of ectoparasites on all chicks, no documented survival</li> <li>7. Infrequent surveys, underestimated nesting pairs?</li> </ol>			

2

3 Bald Eagles (*Haliaeetus leucocephalus*) and Osprey (*Pandion haliaetus*) are large birds of prey that  
 4 feed on fish and waterfowl. The Bald Eagle was removed from the USFWS endangered species list on  
 5 June 28, 2007, and is no longer protected under the Endangered Species Act, but remains protected  
 6 under the Bald and Golden Eagle Protection Act and the Migratory Bird Treaty Act (U.S. Department  
 7 of the Interior, 2007). The Bald Eagle was delisted by the FWC through adoption of the Bald Eagle  
 8 Management Plan on April 9, 2008 (FWC, 2008). The FWC Bald Eagle Management Plan recommends

1 maintaining a 660-foot buffer zone, with certain activities allowable between 330 feet and 660 feet  
2 of an active nest outside of the nesting season.

3 Nesting surveys for Ospreys and Bald Eagles were conducted on Rodman Reservoir, Deep Creek, and  
4 Sweetwater Creek. Rookeries are visited at least every two years. Osprey platforms in Rodman  
5 Reservoir and Lake Rousseau are visited several times during the breeding season, at least biennially.  
6 Breeding pairs are gps'd and recorded as well as nest productivity.

7 The largest concentration of nesting Ospreys was observed at Rodman Reservoir, where 34 active  
8 nests were documented. During this time, no nests were observed at Lake Rousseau, although an  
9 active Osprey nest was present on a very tall platform approximately 100 meters downstream of the  
10 Lake Rousseau Dam. Two nests were observed at Deep Creek, and one on Sweetwater Creek; all three  
11 were active.

12 Two Bald Eagle nests were documented during spring/winter 2004. One was in Putnam County 0.31  
13 miles northeast of the Buckman Lock and the other in Marion County about 1.86 miles east of the  
14 town of Dunnellon. Both nests were in pine trees.

#### 15 Plants

16 The CFG contains 70,817 acres and spans the state from the Gulf of Mexico to the St. Johns River. In  
17 2008, FNAI mapped the natural communities on the CFG. The CFG harbors excellent examples of  
18 Florida's rapidly disappearing natural communities, including bottomland forest, depression marsh,  
19 floodplain swamp, four types of hammocks (hydric, maritime, mesic, and xeric), mesic and wet  
20 flatwoods, sandhill, scrub, and upland hardwood forest. These communities provide habitat for a  
21 number of FNAI, state, and federally listed plant species.

22 FNAI completed surveys in July, August, and October 2016 and January, February, March, April, and  
23 May 2017. At the completion of the surveys, FNAI prepared the *2017 Assessment of Listed and Rare*  
24 *Plant Species at Marjorie Harris Carr Cross Florida Greenway Levy, Citrus, Marion and Putnam Counties,*  
25 *Florida*, which was used to develop this section of the CFG UMP. Results of these surveys include the  
26 occurrence of 23 rare plant species listed by the state of Florida, of which 12 are state listed as  
27 endangered, 10 are state listed threatened, and the federally listed endangered rare longspurred  
28 mint (*Dicerandra cornutissima*) also was found. Several locations were recorded for five species listed  
29 by the state as commercially exploited: Green fly orchid (*Epidendrum conopseum*), cinnamon fern  
30 (*Osmunda cinnamomea*), royal fern (*Osmunda regalis var. spectabilis*), needle palm (*Rhapidophyllum*  
31 *hystrix*), and coontie (*Zamia integrifolia* [formerly called *Z. pumila*]).

32 Populations of dwarf spleenwort (*Asplenium pumilum*) and Tampa vervain (*Glandularia tampensis*),  
33 two species new to the CFG, were found during the current survey. A new population of Chapman's  
34 skeletongrass (*Gymnopogon chapmanianus*), tracked by FNAI but not listed, also was documented. In  
35 addition, new populations were recorded of angle pod (*Matelea carolinensis*), Florida spiny-pod  
36 (*Matelea floridana*), sandhill spiny-pod (*Matelea pubiflora*), blueflower butterwort (*Pinguicula*  
37 *caerulea*), cardinal flower (*Lobelia cardinalis*), cinnamon fern, royal fern, green fly orchid, coontie,  
38 giant orchid (*Eulophia ecristata*), plume polypody (*Pecluma plumula*), and swamp plume poypody  
39 (*Pecluma ptildodon*). The known population of scrub stylisma (*Stylisma abdita*) was determined to  
40 be much larger than originally documented. One 15-foot-tall star anise (*Illicium parviflorum*; state  
41 listed endangered) was found during an invasive exotic plant survey in January 2016; this small tree



1 was near a north boundary and we suspect it may be an introduction from a neighboring yard. The  
2 anise was not re-verified during the current survey.

3 Further, 11 significant botanical sites were identified on the CFG, based on current and earlier  
4 surveys (Herring and Schultz 2003, Herring 2005, FNAI 2008, FNAI 2015). These botanically  
5 significant sites were determined based on: rarity of a given species, numbers of listed species,  
6 numbers of species in general, lack of disturbance, or with lots of disturbance that warrants attention.  
7 The 11 sites include: the vicinity of Inglis Canal and west of US 19, Inglis Island, the Diggings scrub &  
8 sandhill (from approximately two miles west of SR 200 to just east of I-75), SE 25th Street (west of  
9 Santos), Marshall Swamp Trail, Ocklawaha River floodplain, Eureka Dam, Deep Creek, select areas  
10 surrounding the Rodman Reservoir, Caravelle Ranch West, and select areas surrounding the  
11 Buckman Lock (Herring and Schultz 2003, Herring 2005).

12 From the western end of the CFG, in Citrus and Levy Counties, bordering the Gulf Coast, maritime  
13 hammock and mesic hammock that have exposed limestone provide habitat to several rare plant  
14 species. Three listed plants—Tampa vervain, spiked crested coralroot (*Hexalectris spicata*), and  
15 angle pod—were documented in maritime hammock in the vicinity of the Inglis Canal. Angle pod and  
16 pinewoods dainties (*Phyllanthus liebmannianus*) were observed in mesic hammock on Inglis Island.  
17 Most plants were located immediately adjacent to a jeep trail that runs along the southern end of  
18 mesic hammock. A few plants were in the trail; we recommend the trail be gated or closed to prevent  
19 excessive foot and vehicular traffic.

20 Further east in Marion County, within a linear east-west strip of the CFG (approximately beginning  
21 with the Diggings west of SR 200 east to I-75), seven listed plants occur in sandhill, scrub, and  
22 successional hardwoods that surround and include the Diggings, the site of the slated Cross Florida  
23 Barge Canal. Scrub and, to a lesser extent, sandhill harbor the population stronghold of the federal  
24 and state listed endangered longspurred mint (*Dicerandra cornutissima*). State listed commercially  
25 exploited garberia (*Garberia heterophylla*) flourish within this region. The highest quality sandhills  
26 are found in this region of the CFG and contain the following rare species: giant orchid, longspurred  
27 mint, garberia, scrub stylisma, and sandhill spiny-pod. In the Diggings proper within successional  
28 hardwoods, numerous large limestone boulders dot the landscape and are covered with a multitude  
29 of fern species forming a fern grotto. State listed plume polypody and swamp plume polypody were  
30 found on the limestone boulders that provide an ideal substrate. Sword fern (*Nephrolepis cordifolia*)  
31 poses perhaps the biggest threat to the ferns. A new population of Florida spiny-pod was found in  
32 this area on top of the old canal berm.

33 East of I-75, another fern grotto-like habitat occurs on the east side of SE 25th Street (west of Santos)  
34 in Marion County within successional hardwood forest. A series of boulders run parallel, north-south,  
35 to SE 25th Street. Plume polypody, swamp plume polypody, and dwarf spleenwort (new species for  
36 the CFG) were occasional, common, and rare, respectively, on the limestone boulders. The sword fern  
37 is of concern here also.

38 Continuing northeast, Marshall Swamp Trail and Ocklawaha River floodplain in the Sharpes Ferry  
39 area offer high-quality hydric hammock and bottomland forest in which several rare plants are  
40 known, such as angle pod, needle palm (*Rhapidophyllum hystrix*), pinkroot (*Spigelia marilandica*),  
41 and Treat's zephyrlily (*Zephyranthes rosea*). In areas where limestone is exposed, such as Butterbutt  
42 Landing, swamp plume polypody occurs.

1 Eureka Dam, east of the Ocklawaha River, was known to have populations of the rare pigmy pipes  
2 (*Monotropsis odorata*) and giant orchid. Despite repeated searches, neither species has been found  
3 in recent years.

4 In Putnam County, within the Deep Creek site, seven listed and one commercially exploited species  
5 were documented within excellent hydric hammock: angle-pod, cardinal flower, Chapman's sedge  
6 (*Carex chapmanii*), Florida spiny pod, Florida willow (*Salix floridana*), large-leaved grass-of-  
7 parnassus (*Parnassia grandifolia*), needle palm, and variable leaf Indian plantain (*Arnoglossum*  
8 *diversifolium*).

9 Several areas surrounding the Rodman Reservoir in Putnam County have listed or rare plants. A  
10 population of Chapman's skeletongrass (*Gymnopogon chapmanianus*) and garberia occurs on the  
11 scrub island north of the Rodman Reservoir. Populations of hooded pitcher plants (*Sarracenia minor*)  
12 also grow in the vicinity of the Rodman Reservoir and north and south of Rodman Road in mesic  
13 flatwoods.

14 In the Caravelle Ranch West portion of the CFG in Putnam County, five listed species were  
15 documented: blue butterwort, Chapman's skeletongrass, garberia, yellow-flowered butterwort  
16 (*Pinguicula lutea*), and hooded pitcher plant in mesic flatwoods, wet prairie, and scrubby flatwoods.

17 Several areas surrounding the Buckman Lock had listed plants. Along Rodeheaver Boys Ranch Road  
18 and west of the visitor's center, populations of hooded pitcher plants occur in mesic flatwoods.  
19 Garberia occurs north and south of the Buckman Lock canal in sandhill and scrub.

# 1 FLORIDA SCRUB-JAY

2 *Aphelocoma coerulescens*

## 3 Description

4 The Florida Scrub-Jay is Florida's only endemic bird  
5 species, found nowhere else in the world. It was listed as  
6 federally Threatened by the USFWS under the Endangered  
7 Species Act in 1987, largely due to loss of its native scrub  
8 habitat and decades of fire suppression that allowed the  
9 scrub to become overgrown and unsuitable for Scrub-Jays.

10 The Florida Scrub-Jay's appearance is similar to the far  
11 more common Blue Jay. Both are the same size,  
12 approximately 12 inches in length, but the Scrub-Jay is  
13 paler in color and lacks a crest. The Scrub-Jay also lacks the  
14 white wing spots and white tail feather tips typical of the  
15 Blue Jay. Rather, the Scrub-Jay wears a collar of blue  
16 feathers that separates its white throat from its gray  
17 underparts, and it has a white line over the eye that blends  
18 into a whitish forehead. The white forehead and eyebrows  
19 distinguish the Florida Scrub-Jay from those of western  
20 states.

21 The Scrub-Jay's diet is quite varied. Acorns are consumed  
22 year-round and are a main staple during the fall and  
23 winter. In spring and summer, insects become the main  
24 food source, supplemented by frogs, mice, toads, lizards,  
25 snakes, and birds' eggs. Saw palmetto drupes, and  
26 greenbrier berries also are eaten when available.

## 27 Habitat

28 The Florida Scrub-Jay has extremely specific habitat  
29 requirements that have been degraded throughout Florida  
30 due to fire exclusion and habitat destruction. In ideal  
31 Scrub-Jay habitat, oaks between three feet and eight feet  
32 tall blanket between 50 percent and 90 percent of an area,  
33 while sparse vegetation no higher than six inches (or  
34 perhaps only bare ground) covers the remaining region.  
35 Fire is essential for maintaining Scrub-Jay habitat. An area  
36 needs to be burned every five years to 20 years to keep  
37 scrub vegetation at the proper height.

38 Based on their field surveys, FNAI only reported Scrub-  
39 Jays from CFG-managed scrub immediately west of I-75.  
40 Scrub-Jays also are reported very near the scrub formerly



## Species Status:

**Federal Listing:** Federally Threatened (FT)

**State Listing:** Listed Threatened (LT)

**FNAI Global and State Rank:** G2, S2

## Habitat:

- Oaks between 3 feet and 8 feet tall
- Sparse vegetation not higher than 6 inches tall or bare ground
- Periodic fire every 5 to 20 years

## Nesting:

- Family territory between 5 acres and 50 acres
- Mate for life
- Mating season from early March to late May, sometimes into June

## Management Practices:

- Prescribed fire
- Exotic minimization
- Scrub restoration
- Monitoring and tracking

1 managed by the FDEP's Office of Coastal and Aquatic Managed Areas (CAMA) on the western end of  
2 the greenway near Yankeetown.

### 3 **Reproduction**

4 The family life of Scrub-Jays is unusually complex. A family, which consists of a breeding pair and  
5 some of their offspring, establishes its own territory and strongly protects it from other Scrub-Jays.  
6 A family's territory may average between five acres and 50 acres, though 25 acres is most common.

7 The breeding pair, which mate for life, usually are around three or four years of age. Their mating  
8 season is short, from early March to late May and sometimes into June.

9 A nest is built between three feet and 10 feet above ground in one of the scrub oaks. Nests, made of  
10 twigs and lined with finer material, are used only once.

11 The average clutch is three greenish, brown-spotted eggs, which hatch after about 17 days. It also  
12 takes about 17 days, on average, for the nestlings to fledge from the time they hatch. The juveniles  
13 are distinctive, with a dusky brown head and neck that lasts until their first molt, following the first  
14 summer.

15 Unlike any of Florida's other songbirds, both non-nesting females but particularly males remain part  
16 of the family for several years. These hangers-on serve as valuable helpers by defending the family  
17 territory and feeding the nestlings and fledglings. However, helpers do not assist in nest building or  
18 incubating. Not surprisingly, studies have shown that breeding pairs with helpers raise their young  
19 more successfully than do birds without helpers.

### 20 **Management**

21 The CFG has worked with the FWC to identify and restore scrub communities on the greenway.  
22 Restoration of these areas includes the mechanical removal of sand pine and large oaks to allow the  
23 return of fire to these communities. Specifically, CFG has restored a total of 840 acres out of 1,100  
24 acres of historic habitat on the CFG. Increase in burning and restoration activities resulted in an  
25 increase of Scrub-Jays from 46 birds in 2009 to a total of 111 birds in 2015. Current management  
26 activities include the continued coordination with Florida Audubon Society for monitoring and  
27 includes contracting with wildlife consultants for trapping and banding of the populations on the  
28 Triangle. In fact, the "Triangle" area of the CFG—within Marion County—is designated through the  
29 Audubon Society as Jay Watch Site. Additionally, the CFG will continue to coordinate with the  
30 Audubon Society, FWC, and private wildlife consultants to complete necessary management and  
31 restoration activities on CFG scrub habitat, specifically within the Triangle area.



# 1 **LONGSPURRED MINT**

2 *Dicerandra cornutissima*

## 3 **Description**

4 A strongly aromatic plant, longspurred mint grows up to  
5 1.6 feet tall, with erect, non-woody flowering shoots  
6 growing from a woody base. The leaves are just over 0.5-  
7 inch long, linear, with entire margins, and covered with  
8 conspicuous sunken glands. The leaves are borne opposite  
9 one another on the stems, often with two smaller leaves at  
10 each node. Flowers are borne in groups in the axils of the  
11 leaves on the upper parts of the stems. The petals are 7  
12 millimeters (0.3 inch) long, forming a tube with two lips,  
13 bent at a 90-degree angle in the middle, and colored  
14 purplish-rose with deep purple markings and a whitish  
15 throat. The anthers are tipped by a spur 1.2 millimeters  
16 long, for which the plant is named.

## 17 **Reproduction**

18 Longspurred mint flowers in September and produces  
19 fruits in the form of four small nutlets. This is a short-lived  
20 perennial that grows from seed; the species does not  
21 spread vegetatively.

## 22 **Range and Population**

23 There are 15 occurrences of Longspurred mint in Marion  
24 and Sumter Counties, of which six of these populations are  
25 on the CFG. The plant has been extirpated from several  
26 sites in these counties. The CFG is the only conservation  
27 land that supports a mint population.

## 28 **Habitat**

29 Longspurred mint is found only in open areas in sand pine  
30 scrub or oak scrub, and in the ecotones between these and  
31 turkey oak communities. It can colonize the edges of road  
32 rights-of-way, and has spread vigorously along streets.

33 Within the CFG, populations are found in mostly sand  
34 pine-dominated scrub with sandhills interspersed  
35 within. As with the longspurred mint locations farther  
36 west on the CFG, roadside edges are the preferred habitat.  
37 Care should be taken when moving dirt along the jeep  
38 trails, in driving any heavy equipment off-road, or when  
39 establishing/ maintaining fire plow lines.



## **Species Status:**

**Federal Listing:** Federally  
Endangered (FE)

**State Listing:** Listed Endangered  
(LE)

**FNAI Global and State Rank:** G1,  
S1

## **Habitat:**

- Open areas in sand pine scrub or oak scrub
- In the ecotones between scrub and turkey oak communities

## **Management Practices:**

- Preserve the species in the extant sites
- Evaluate establishing additional populations within historic range
- Control exotic plants (particularly natal grass and cogon grass).
- Limit off-road activity such as foot, horse, or vehicular traffic.

1 Occurrences have been observed on the CFG, in a small area between I-75 and SR 200 in scrub and  
2 sandhill. The CFG provides critical habitat for this species and contains a significant portion of the  
3 known plants. A few known associates observed with longspurred mint include longleaf pine (*Pinus*  
4 *palustris*), sand pine (*Pinus clausa*), turkey oak (*Quercus laevis*), sand live oak (*Quercus geminata*),  
5 myrtle oak (*Quercus myrtifolia*), Chapman's oak (*Quercus chapmanii*), saw palmetto (*Serenoa repens*),  
6 scrub palmetto (*Sabal etonia*), and a lichen (*Cladina evansii*). Longspurred mint was not a high-  
7 priority target species during the current survey; a more thorough survey, concentrating exclusively  
8 on the mint, is scheduled for summer and fall of 2017.

### 9 **Management and Protection**

10 The species should be preserved in the extant sites, and, to provide greater security, the possibility  
11 of establishing additional populations within the historic range should be evaluated. Mild  
12 disturbances appear to have little effect and probably stimulate the species by reducing competition.

13 While it is not known how the mint responds to fire, it does appear to favor open areas as evident  
14 from its proliferation along roadsides. Other management needs are to control exotic plants  
15 (especially natal grass and cogon grass) and limit off-road activity, such as foot, horse, or vehicular  
16 traffic.

## CFG Exotic Invasive Species Planning Accomplishments

2007-2016

According to the FDEP, DRP database, the CFG has a total of 70,531 acres, of which 1,716 acres are considered to be infested with exotic invasive plant species. In all, the infested acres are made up of 4,410 individual points.

All infestations have been treated at least biannually since 2008/2009 and are considered to be in maintenance condition.

Ongoing steps in this successful management process include:

- Continue to manage a systematic upland invasive plant maintenance program.
- Use a combination of Florida Fish and Wildlife Conservation Commission (FWC) and in-house funding to hire contractors and staff to perform invasive species control work.

### 1 **Exotic and Nuisance Species**

2 Non-native invasive plant species pose a significant threat  
3 to Florida’s natural areas. Florida is particularly vulnerable  
4 to non-native invasive species because of its peninsular  
5 geography, tropical/subtropical climate, and diverse  
6 ecosystems. More than 1.5 million acres of Florida’s public  
7 conservation lands have been invaded by alien (exotic,  
8 non-native, non-indigenous) plants such as melaleuca  
9 (*Melaleuca quinquenervia*), Brazilian pepper (*Schinus*  
10 *terebinthifolius*), cogon grass (*Imperata cylindrica*), and  
11 climbing ferns. It is estimated that approximately 1,400  
12 non-native plant species are present in the state, with 124  
13 species currently present in state parks (Adams et al.,  
14 2011; FLEPPC, 2011).

15 In an ecological context, an invasive species is one that is  
16 aggressive in growth and expansion of range and tends to  
17 dominate other species; its establishment and dominance  
18 can cause widespread harm to an ecological system by  
19 altering the species composition, susceptibility to fire, and  
20 hydrology of an area. The characteristics of some of these  
21 species (high rate of growth/reproduction, no natural  
22 predators, easily dispersed, able to out-compete native  
23 species) make them invasive. Invasions by native and non-  
24 native species often follow an alteration to ecosystem  
25 function, disruption of the food web, large-scale  
26 fragmentation of an ecosystem, and/or disturbance (e.g.,  
27 clearing, fire, drought, etc.) of an area.

28 The FWC’s Invasive Plant Management Section (IPMS) is  
29 the designated lead entity in Florida responsible for  
30 coordinating and funding the statewide control of invasive  
31 aquatic and upland plants in public waterways and on  
32 public conservation land. The Upland Invasive Exotic Plant  
33 Management Program was established in 1997 to address  
34 the need for a statewide coordinated approach to the  
35 terrestrial (vs. aquatic) invasive exotic plant problem. The  
36 Uplands Program (a subsection of IPMS) funds individual  
37 exotic plant removal projects statewide on public  
38 conservation land. Projects are considered for funding  
39 based on recommendations from 11 Regional Invasive  
40 Plant Working Groups.

### 41 *Invasive Exotic Species and Distribution on CFG*

42 Given the disturbed nature of a significant portion of the  
43 lands and waters of the CFG, the large proportion of

1 boundary edge, numerous intersecting transportation routes, and large amount of disturbance on  
 2 adjoining properties, the CFG has serious invasive exotic species problems. In fact, according to the  
 3 DRP database, of the 70, 531 acres on the CFG, 1,716 acres are considered to be infested with non-  
 4 native invasive species. These infested areas are comprised of 4,410 individual points. Surveys  
 5 concluded that of the 4,410 point occurrences on the CFG, 54 percent, or 2,381 points, are either  
 6 cogon grass (*Imperata cylindrica*), camphor tree (*Cinnamomum camphora*), mimosa (*Albizia*  
 7 *julibrissin*), or Caesar's weed (*Urena lobata*). Aquatic invasive species also exist in the water resources  
 8 (8,270 acres, or 12 percent of total CFG area). The most pervasive water invasive species on the CFG  
 9 include the water hyacinth (*Eichhornia crassipes*) and water lettuce (*Pistia stratiotes*), which is  
 10 present within the Rodman Reservoir.

11 Additionally, because of the overall size of the greenway and the amount of invasive exotic species  
 12 present, the CFG staff subdivided the greenway into large management zones (from west to east 1  
 13 through 6). Table 11 demonstrates that the central portions of the greenway are more heavily  
 14 infested with invasive and exotic plant species, while the eastern portions of the greenway appear to  
 15 be least infested. In general, the worst problems were identified in disturbed areas, including areas  
 16 with numerous trails.

17 **Table 11. CFG Large Management Zones and Degree of Infestation**

Large Management Zone	Infestation Degree*	Acres	Area (sq. miles)	Perimeter (miles)	Most Prevalent Invasive Exotic Species by Large Zone
1	4	16,089	25.14	145.62	cogon grass, skunk vine, Camphor tree, lantana, Caesar's weed
2	2	5,478	8.56	24.86	cogon grass, mimosa, ravenegrass, Natal grass, sword fern
3	1	4,295	6.71	24.04	mimosa, cogon grass, Camphor tree, air potato, paper mulberry
4	3	8,732	13.64	60.65	Cogon grass, mimosa, Camphor tree, Caesar's weed, paper mulberry
5	5	9,002	14.07	107.05	Camphor tree, coral ardisia, Caesar's weed, cogon grass, mimosa
6	6	26,969	42.14	178.08	Japanese climbing fern, camphor tree, torpedograss, Peruvian primrosewillow, cogon grass

\*Infestation Degree = 1-6 Highest to Lowest

18 Table 12, below, contains a list of the FLEPPC Category I and II invasive exotic plant species found  
 19 within the park (FLEPPC, 2013). The table also identifies relative distribution for each species and  
 20 the management zones in which they are known to occur. An explanation of the codes is provided  
 21 following the table.



1 **Table 12. Inventory of FLEPPC Category I and II Invasive Exotic Plant Species**

Common Name/Scientific Name	FLEPPC Cat <sup>a</sup>	Large Zone(s)	Florida Noxious Weed List	Management Zone
Lantana ( <i>Lantana camara</i> )	I	1, 3	N	<b>West:</b> W021, W025, W031, W032, W040, W041, W043, W044, W055, W056, W093, W107, W114, W116, W118, W119, W120, W123, W124 <b>Central:</b> C022, C027, C029, C139, C140, C143, C162, C163, C165, C166, C168, C175, C179, C180, C184, C217, C264, C265, C266, C275
Mimosa ( <i>Albizia julibrissin</i> )	I	1, 2, 3, 4, 5, 6	N	<b>West:</b> W006, W016, W017, W018, W019, W021, W026, W033, W036, W041, W043, W044, W047, W056, W060, W107, W114, W116, W118, W123, <b>Central:</b> C005, C009, C013, C022, C025, C026, C027, C029, C048, C079, C081, C084, C122, C137, C138, C139, C140, C141, C142, C143, C144, C145, C146, C147, C148, C149, C151, C152, C153, C156, C157, C158, C160, C162, C163, C166, C167, C168, C169, C170, C171, C172, C173, C174, C175, C176, C177, C178, C179, C180, C181, C182, C184, C185, C188, C189, C190, C192, C193, C194, C195, C197, C198, C199, C200, C203, C204, C205, C206, C207, C209, C211, C212, C214, C216, C217, C218, C220, C223, C224, C242, C244, C255, C256, C257, C258, C259, C260, C261, C263, C265, C266, C267, C269, C272, C274 <b>East:</b> E021, E141, E165, E166, E167, E190, E233, E242, E243, E275, E277, E278, E294, E295
Glossy Privet ( <i>Ligustrum lucidum</i> )	I	1, 3, 4, 5	N	<b>West:</b> W034, W056, W058, W060 <b>Central:</b> C023, C027, C031, C104, C122, C123, C126, C127, C139, C141, C142, C146, C149, C162, C168, C169, C170, C172, C175, C184, C190, C207, C217 <b>East:</b> E277
Natal Grass ( <i>Melinis repens</i> )	II	2	N	<b>West:</b> W033, W116, W118, W120, W123 <b>Central:</b> C079, C110, C112, C235, C236, C255, C256, C259, C261, C263, C264, C266, C267, E260, E261
Tung oil tree ( <i>Aleurites fordii</i> )	II	1, 3	N	<b>West:</b> W021, W043 <b>Central:</b> C027, C168
Chinese privet ( <i>Ligustrum sinense</i> )	I	1, 3, 4, 5	N	<b>West:</b> W034 <b>Central:</b> C025, C027, C149, C168, C170, C207 <b>East:</b> E277
Mexican petunia ( <i>Ruellia simplex</i> )	I	5	N	<b>Central:</b> C026 <b>East:</b> E272
Coral ardisia ( <i>Ardisia crenata</i> )	I	1, 5	N	<b>West:</b> W034, W043, W044, W055, W058, W060 <b>Central:</b> C005, C012, C013, C014, C021, C022, C023, C025, C026, C027, C031, C032, C079, C094, C146, C166, C217 <b>East:</b> E206, E250

Common Name/Scientific Name	FLEPPC Cat <sup>a</sup>	Large Zone(s)	Florida Noxious Weed List	Management Zone
Japanese honeysuckle ( <i>Lonicera japonica</i> )	I	5	N	<b>West:</b> W034, W044, W055, W056, W058, W060 <b>Central:</b> C022, C023, C025, C027, C084, C127, C168, C170, C172, C175, C180, C203, C264, C275
Flamegold tree ( <i>Koelreuteria elegans</i> ssp. <i>Formosana</i> )	II	4	N	<b>Central:</b> C205
Chinese tallow tree ( <i>Sapium sebiferum</i> )	I	4	Y	<b>West:</b> W005, W026, W028, W031, W041 <b>Central:</b> C025, C104, C105, C110, C112, C118, C119, C121, C122, C231, C265, C272 <b>East:</b> E234, E301
Durban crowfoot grass ( <i>Dactyloctenium aegyptium</i> )	II	6	N	<b>East:</b> E014, E021
Chinese wisteria ( <i>Wisteria sinensis</i> )	II	5	N	<b>Central:</b> C027, C029, C084, C142, C163, C205, C207, C264, C265 <b>East:</b> E275, E277
Asparagus fern ( <i>Asparagus Aethiopicus</i> )	I	5	N	<b>Central:</b> C026, C079, C142
Japanese climbing fern ( <i>Lygodium japonicum</i> )	I	1, 2, 3, 4, 5, 6	Y	<b>West:</b> W008, W017, W027, W036, W043, W044, W055, W056, W058, W060, W088, W096, W124 <b>Central:</b> C021, C022, C023, C026, C046, C079, C098, C099, C101, C102, C103, C104, C105, C115, C168, C171, C181, C184, C190, C206, C221, C254, C255, C265, C266, C275 <b>East:</b> E017, E018, E022, E023, E154, E167, E171, E195, E206, E215, E216, E219, E222, E233, E234, E235, E243, E270
Japanese jasmine			N	<b>Central:</b> C026, C027
Brazilian pepper ( <i>Schinus terebinthifolius</i> )	I	1	Y	<b>West:</b> W123, W125, W126, W139
Nandina ( <i>Nandina domestica</i> )	I	1, 2, 3, 5	N	<b>West:</b> W005, W043 <b>Central:</b> C022, C025, C027, C168, C184, C265 <b>East:</b> E277
Napier grass ( <i>Pennisetum purpureum</i> )	I	6	N	<b>East:</b> E233

Common Name/Scientific Name	FLEPPC Cat <sup>a</sup>	Large Zone(s)	Florida Noxious Weed List	Management Zone
Paper mulberry ( <i>Broussonetia papyrifera</i> )	II	3, 4, 5	N	<b>West:</b> W041, W042 <b>Central:</b> C025, C027, C029, C031, C073, C124, C126, C136, C138, C139, C140, C142, C143, C144, C145, C146, C147, C148, C149, C151, C152, C153, C162, C163, C168, C169, C172, C173, C178, C179, C180, C184, C188, C190, C205, C209, C211, C217 <b>East:</b> E243, E244, E252, E294
Cat's claw vine ( <i>Dolichandra unguis-cati</i> )	I	5	N	<b>Central:</b> C027, C031, C032, C033, C205
Rattlebox ( <i>Sesbania punicea</i> )	II	5, 6	N	<b>Central:</b> C022 <b>East:</b> E170, E171, E175, E176, E206, E214, E237, E307
Australian pine ( <i>Casuarina equisetifolia</i> )	I	1	Y	<b>West:</b> W139
Chinaberry ( <i>Meia azedarach</i> )	II	1, 3, 4	N	<b>West:</b> W031, W032, W034, W043, W056, W058, W090, W116, W117, W118 <b>Central:</b> C023, C026, C027, C099, C103, C104, C110, C121, C122, C126, C127, C136, C140, C141, C142, C143, C144, C145, C148, C149, C154, C157, C159, C162, C163, C164, C168, C200, C204, C205, C207, C211 <b>East:</b> E235
Tropical soda apple ( <i>Solanum viarum</i> )	I	1, 5	Y	<b>West:</b> W021, W027, W028, W030, W031, W032, W033, W034, W043, W055, W056 <b>Central:</b> C021, C022, C025, C026, C027, C031, C046, C098, C110, C115, C217 <b>East:</b> E162, E167, E171, E176, E242, E243
Camphor tree ( <i>Cinnamomum camphora</i> )	I	1, 3, 4, 5, 6	N	<b>West:</b> W005, W006, W008, W014, W015, W016, W017, W018, W020, W021, W024, W025, W026, W027, W030, W031, W032, W033, W034, W035, W036, W037, W039, W041, W042, W043, W044, W045, W047, W055, W056, W058, W060, W061, W093, W096, W132 <b>Central:</b> C013, C014, C021, C022, C023, C025, C026, C027, C029, C031, C032, C033, C038, C044, C046, C048, C052, C054, C079, C084, C098, C102, C103, C104, C110, C118, C121, C122, C123, C125, C126, C127, C137, C138, C139, C140, C141, C142, C143, C144, C145, C146, C149, C152, C153, C154, C156, C157, C162, C163, C168, C169, C170, C171, C172, C174, C175, C176, C178, C181, C183, C184, C185, C188, C189, C190, C191, C193, C200, C204, C205, C214, C217, C220, C224, C255, C256, C266 <b>East:</b> E039, E147, E154, E155, E162, E163, E166, E167, E169, E176, E177, E199, E205, E206, E208, E210, E215, E221, E222, E243, E250, E285, E308

Common Name/Scientific Name	FLEPPC Cat <sup>a</sup>	Large Zone(s)	Florida Noxious Weed List	Management Zone
Golden bamboo	II	2	N	<b>Central:</b> C264
Air potato ( <i>Dioscoria bulbifera</i> )	I	3, 4, 5	Y	<b>West:</b> W021, W041, W043, W056 <b>Central:</b> C027, C029, C079, C081, C142, C143, C146, C149, C151, C152, C153, C156, C157, C162, C163, C168, C169, C172, C175, C176, C178, C179, C180, C181, C184, C205, C207, C214, C265 <b>East:</b> E161, E235, E237, E242, E243, E259, E277
Alligator weed ( <i>Alternanthera philoxeroides</i> )	II	4	Y	<b>Central:</b> C086, C089, C094, C095, C098, C101, C102, C103
Arrowhead vine ( <i>Synгонium podophyllum</i> )	I	5	N	<b>Central:</b> C025, C026
Sword fern ( <i>Nephrolepis cordifolia</i> )	I	2, 3, 4, 5	N	<b>West:</b> W021, W043, W058, W060 <b>Central:</b> C021, C022, C023, C025, C026, C027, C058, C079, C098, C099, C103, C104, C119, C122, C140, C149, C151, C156, C163, C164, C168, C172, C175, C176, C178, C180, C181, C205, C211, C266, C275 <b>East:</b> E206, E236, E250
Small-leaf spiderwort ( <i>Tradescantia fluminensis</i> )	I	5	N	<b>West:</b> W034 <b>Central:</b> C023, C025, C026, C027, C084 <b>East:</b> E277
Caesar weed ( <i>Urena lobate</i> )	I	1, 3, 4, 5, 6	N	<b>West:</b> W025, W026, W027, W030, W031, W032, W033, W034, W043, W044, W047, W055, W081, W093, W095, W096 <b>Central:</b> C007, C013, C014, C021, C022, C023, C025, C026, C046, C048, C052, C079, C084, C086, C089, C094, C095, C098, C099, C101, C102, C103, C104, C105, C118, C119, C142, C143, C163, C168, C184, C188, C190, C203, C205, C217, C244, C266, C275 <b>East:</b> E167, E206, E215, E233, E234, E240, E242, E243, E245
Silverthorn ( <i>Elaeagnus pungens</i> )	II	1, 2, 4, 5	N	<b>West:</b> W037 <b>Central:</b> C027, C138, C139, C264
Senegal date palm ( <i>Phoenix reclinata</i> )	II	4	N	<b>Central:</b> C146, C156, C162
Skunk vine ( <i>Paederia foetida</i> )	I	1	Y	<b>West:</b> W008, W014, W017, W023, W025, W026, W027, W028, W030, W031, W032, W033, W034, W035, W036, W043, W044, W047, W055, W056, W058, W060, W088, W090, W091, W095, W096, W099, W107 <b>Central:</b> C266, C275, C027
Para grass ( <i>Urochloa mutica</i> )	I	6	N	<b>East:</b> E220



Common Name/Scientific Name	FLEPPC Cat <sup>a</sup>	Large Zone(s)	Florida Noxious Weed List	Management Zone
Cogon grass ( <i>Imperata cylindrical</i> )	I	1, 2, 3, 4, 5, 6	Y	<p><b>West:</b> W005, W008, W015, W017, W018, W021, W024, W025, W026, W030, W036, W047, W084, W088, W094, W095, W105, W107, W108, W109, W111, W112, W114, W116, W118, W119, W120, W123, W124, W125, W126, W128, W129, W130, W131, W135</p> <p><b>Central:</b> C008, C013, C014, C021, C022, C023, C025, C026, C027, C028, C029, C035, C046, C048, C061, C079, C084, C099, C103, C104, C106, C107, C108, C109, C110, C111, C112, C116, C117, C118, C120, C121, C122, C123, C125, C126, C136, C137, C138, C139, C140, C141, C142, C144, C146, C151, C152, C156, C157, C162, C163, C165, C166, C167, C168, C173, C175, C176, C178, C179, C180, C181, C184, C185, C188, C189, C190, C192, C193, C194, C205, C206, C207, C212, C214, C217, C222, C223, C224, C226, C244, C254, C255, C256, C257, C258, C259, C261, C262, C263, C264, C266, C267, C268, C269, C274, C275, C276</p> <p><b>East:</b> E025, E062, E098, E141, E142, E147, E165, E186, E205, E211, E214, E216, E222, E232, E233, E234, E243, E249, E259, E260, E266, E267, E275, E277</p>
Torpedo grass ( <i>Panicum repens</i> )	I	1, 4, 6	N	<p><b>West:</b> W028, W108, W124, W125, W126</p> <p><b>Central:</b> C089, C098, C099, C102, C104, C110, C163</p> <p><b>East:</b> E012, E014, E020, E022, E023, E036, E056, E062, E064, E065, E068, E070, E074, E075, E077, E092, E099, E120, E136, E138, E166, E167, E182, E195, E204, E205, E206, E212, E215, E216, E217, E219, E221, E222, E226, E234, E235, E236, E242, E308</p>
Winged yam ( <i>Dioscorea alata</i> )	I	4, 5	N	<p><b>Central:</b> C143</p> <p><b>East:</b> E277</p>
Wild taro ( <i>Colocasia esculenta</i> )	I	4, 5, 6	N	<p><b>West:</b> W106, W109</p> <p><b>Central:</b> C021, C022, C025, C026, C058, C079, C084, C086, C094, C095, C098, C099, C101, C102, C103, C168</p> <p><b>East:</b> E063, E099, E154, E161, E167, E205, E206, E207, E241, E245, E250, E272, E277, E293, E296</p>
Chinese brake fern ( <i>Pteris vittata</i> )	II	2, 5	N	<p><b>West:</b> W008, W017, W018, W043, W056, W060, W087, W093, W096, W105, W106, W109, W114, W116, W118, W123, W124, W125, W126</p> <p><b>Central:</b> C021, C022, C026, C027, C032, C046, C058, C060, C061, C074, C103, C104, C118, C168, C172, C173, C178, C180, C259, C266, C274, C275</p> <p><b>East:</b> E062, E219, E223, E236</p>
Elephant ear ( <i>Xanthosoma sagittifolium</i> )	II	4, 5	N	<p><b>Central:</b> C026, C084, C122, C142, C168</p> <p><b>East:</b> E277</p>

Common Name/Scientific Name	FLEPPC Cat <sup>a</sup>	Large Zone(s)	Florida Noxious Weed List	Management Zone
Peruvian primrosewillow ( <i>Ludwigia peruviana</i> )	I	6	N	<b>West:</b> W096 <b>Central:</b> C025, C058, C060, C094, C095, C098, C101, C102, C156 <b>East:</b> E006, E017, E018, E023, E154, E167, E171, E172, E175, E176, E205, E206, E216, E217, E219, E221, E223, E224, E234, E235, E236, E237, E241, E242, E281, E293
Water hyacinth ( <i>Eichhornia crassipes</i> )	I	6	N	<b>Central:</b> C073, C089, C101, C102, C104, C115, C119 <b>East:</b> E066, E161, E167, E175, E205, E206, E220, E241, E279, E309
Water lettuce ( <i>Pistia stratiotes</i> )	I	5, 6	N	<b>West:</b> W046 <b>Central:</b> C007, C022, C025, C026, C058, C102 <b>East:</b> E099, E166, E167, E194, E205, E241, E249, E250, E252, E257, E286
<b>Florida Exotic Pest Plant Council (FLEPPC) 2007 designations:</b> I: Invasive exotics that are altering native plant communities by displacing native species, changing community structures or ecological functions, or hybridizing with natives. II: Invasive exotics that have increased in abundance or frequency but have not yet altered Florida plant communities to the extent shown by Category I species.				

1

2 ***Invasive Exotic Plant Species Management on the CFG***

3 FNAI conducted upland surveys for invasive exotic plant species. Numerous species of upland exotic  
4 pest plant species were documented on CFG lands. Table 12, provided above, indicates the species  
5 that are considered Category I invasive non-native plant species. The CFG will continue efforts to  
6 control the establishment and spread of Florida Exotic Pest Plant Council (FLEPPC) Category I or II  
7 plants as provided in Table 12. Control techniques are ongoing and include mechanical, chemical,  
8 biological, and other appropriate treatments. Treatments utilizing herbicides will comply with  
9 instructions found on the herbicide label and employ the Best Management Practices for their  
10 application.

11 Since the approval of the last unit management plan, the greenway has been subdivided into six zones  
12 to facilitate management. Over the last few years, staff and contractors have treated the population  
13 of exotic plant species within the park at least biannually since 2008-2009, resulting in all  
14 populations being in a maintenance condition.

15 CFG has an aggressive program to control invasive non-native plant species, which incorporates  
16 three basic guidelines. First, the implementation of a systematic approach ensures that infestations  
17 are not overlooked or missed, and that all required follow-up treatments are easier. Use of this  
18 approach has proven to be the most cost-effective way to utilize private contractors. For private  
19 contractors, mobilization to a work site normally is the largest expense. Therefore, it is advantageous  
20 to minimize the number of mobilizations by requesting that the contractors treat all infestations in a  
21 given area, regardless of species type.

1 Second, the CFG deploys contractors to a site if new large infestations are discovered. With this in  
2 mind, the goal is to reduce the overall size of the infestation to be manageable through the use of in-  
3 house staff.

4 The third guideline is in place to effectively and efficiently use in-house staff resources, which are  
5 ultimately used in several capacities. In-house staff primarily focuses on small maintenance retreats  
6 and additional applications necessary for the control of grasses and vines on the CFG. To accomplish  
7 this, in-house staff apply the principles of Early Detection and Rapid Response (EDRR) to minimize  
8 the overall impact of these small infestations by ensuring that they remain in check and do not  
9 become something that requires outside assistance.

10 Finally, in addition to responding to small infestations, in-house crews are able to investigate the  
11 effectiveness of new recommended herbicide and treatment methodologies through the use of  
12 informal practical test plots. In addition to funding FNAI's invasive exotic plants survey, the Office of  
13 Greenways and Trails (OGT) contracted with the University of Florida Weed Science Department to  
14 develop individual weed control plans. These plans recommended the latest and most efficient  
15 eradication methodologies for those invasives documented on the CFG.

#### 16 ***Aquatic Invasive Plant Species Management on the CFG***

17 Invasive aquatic plant species, such as water lettuce and water hyacinth, exist in the water resources  
18 on CFG, particularly within Lake Rousseau and Rodman Reservoir. Invasive aquatic plant  
19 management strategy requires the control of these species on Lake Rousseau and Rodman Reservoir.  
20 On Rodman Reservoir, aquatic invasive plants are controlled to a degree by fluctuating water levels  
21 annually and conducting temporary drawdowns every three to four years. Lowering the  
22 reservoir's water level dries the shallow areas, killing the aquatic invasive plants, in addition to  
23 numerous other benefits. FWC budgets approximately \$30,000 per year for aquatic plant control  
24 in Rodman Reservoir. The FWC budgets up to \$1 million per year to control aquatic non-native  
25 invasive plant species in Lake Rousseau by using contractors.

26 Lake Rousseau, also known as the "Backwaters," is a 100-year-old, 4,000-acre impoundment and is  
27 bordered to the north by Levy and Marion Counties and on the south by Citrus County. The reservoir  
28 is supplied by water from two main sources, the Withlacoochee River and the Rainbow River.

29 In 2014, FWC and FDEP prepared an *Aquatic Plant Management Plan* for Lake Rousseau. The Florida  
30 Aquatic Weed Control Act, Section 369.20(2) Florida Statutes states, "The FWC shall direct the  
31 control, eradication, and regulation of noxious aquatic weeds and direct the research and planning  
32 related to these activities, as provided in this section, so as to protect human health, safety, and  
33 recreation and, to the greatest degree practicable, prevent injury to plant and animal life and  
34 property."

35 The first priority will be to keep the invasive non-native floating plants (water hyacinth/water  
36 lettuce) under maintenance control. The next priority will be to keep established navigation channels  
37 and boat trails open and to control any plants blocking access and navigation from public boat ramps.  
38 The third priority will be to keep open areas for fishing in dense hydrilla mats, as technology, current  
39 conditions, and funding will allow. Aquatic plant control conducted on Lake Rousseau is  
40 accomplished through use of herbicides that are registered with both the U.S. Environmental

1 Protection Agency and FDACS for use specifically in water. Prior to all herbicide applications, the lake  
2 is surveyed by the FWC aquatic plant biologist.

3 A maintenance control strategy is used for floating plants at Lake Rousseau. This strategy is one in  
4 which plants are controlled on a routine basis to prevent small populations of invasive aquatic plants  
5 from expanding into large problems. Maintenance control also prevents aquatic weeds from taking  
6 over large areas of the lake, which maintains more of the lake in a condition that will support native  
7 plants for fish and wildlife habitat, afford the public better access for recreation, and protect the flood  
8 control functions of the system. Frequent small-scale herbicide applications also reduce management  
9 costs and herbicide use.

10 Lake Rousseau also has a well-established bird population, so maintaining a healthy environment for  
11 birds on the lake is an important part of FWC's invasive plant management program. For the past 30-  
12 plus years, Lake Rousseau's bird population and FWC's Invasive Plant Management program that is  
13 designed to conserve or enhance native plant diversity and habitat have coexisted without problem.  
14 Species of concern observed on Lake Rousseau include: Brown Pelicans, Florida Sandhill Cranes,  
15 Limpkins, Little Blue Herons, Snowy Egrets, Tricolored Herons, White Ibis, and Wood Storks.

16 An additional challenge in managing aquatic invasives on Lake Rousseau is that bird nesting and the  
17 invasive plant growing season overlap. To maintain invasive plants at a reasonable level, some  
18 herbicide control will be required near nesting birds, which are found all around Lake Rousseau. As  
19 in the past, herbicide applicators will continue to use caution and good judgment while operating  
20 near these areas. Lake Rousseau's herbicide applicators are instructed to: (1) reduce noise by idling  
21 boats as much as possible and trying to avoid revving them up to a high rpm around rookery areas,  
22 (2) carefully observe bird behavior while applying herbicides, and (3) exit areas adjacent to rookeries  
23 immediately if birds leave their nests or an upflight (large numbers of birds exiting a rookery) occurs.  
24 If such disturbance occurs, applicators are instructed to return later in the day or during another  
25 application time- period when nesting is completed if possible.

26 The following sections describe the main invasive plant species that comprise up to 54 percent  
27 of the infestation on the CFG. These species are described, impacts are noted, then management  
28 methodologies are discussed, including treatment options.



# 1 Cogon Grass

2 *Imperita cylindrical*

## 3 Introduction

4 Cogon grass is an aggressive, rhizomatous, perennial grass  
5 that is distributed throughout the tropical and subtropical  
6 regions of the world. It has become established in the  
7 southeastern United States within the last 50 years, and  
8 Alabama, Mississippi, and Florida have extensive acreage of  
9 roadways and pastures infested with cogon grass. Cogon  
10 grass first appeared in the area around Grand Bay, Alabama,  
11 in 1912, escaping from Satsuma orange crate packing.  
12 Cogon grass was introduced into Florida in the 1930s and  
13 1940s as a potential forage and for soil stabilization  
14 purposes.

## 15 Description

16 Cogon grass is a perennial grass that varies greatly in  
17 appearance. The leaves appear light green, with older  
18 leaves becoming orange-brown in color. In areas with  
19 killing frosts, the leaves will turn light brown during winter  
20 months and present a substantial fire hazard. Cogon grass  
21 grows in loose to compact bunches, each “bunch”  
22 containing several leaves arising from a central area along  
23 a rhizome. The leaves originate directly from ground level  
24 and range from one to four feet in length. Each leaf is 0.5-  
25 inch to 0.75-inch wide with a prominent, off-center, white  
26 mid-rib. The leaf margins are finely serrated; contributing  
27 to the undesirable forage qualities of this grass.

28 Seed production predominantly occurs in the spring, with  
29 long, fluffy-white seedheads. Mowing, burning, or  
30 fertilization also can induce sporadic seedhead formation.  
31 Seeds are extremely small and attached to a plume of long  
32 hairs. Although the seeds can be carried long distances by  
33 wind and animals, the spread of cogon grass by seed is  
34 questionable and still under investigation.

## 35 Management

36 Extensive research has been conducted in Africa, southeast  
37 Asia, and the United States to evaluate the best methods to  
38 control cogon grass. Burning, cultivation, cover crops, and

CFG Distribution: 1, 2, 3, 4, 5, 6



### Status:

- Federal noxious weed
- State listed in Florida
- Category I FLEPPC

### Identification:

- Cylindrical in shape
- 2 inches to 8 inches in length
- Silvery white in color
- Light, fluffy, dandelion-like seeds
- Blooms from late March to mid-June

### Treatment:

- Prescribed fire
- Herbicide application
- Seeding with mix of native species
- Control



1 herbicides have been used with varying degrees of effectiveness. To eliminate cogon grass, the  
2 rhizomes must be destroyed to avoid regrowth. Cultivation and herbicides have been the two most  
3 frequently utilized control strategies. One of the oldest and most successful methods is to deep plow  
4 or disk several times during the dry season to desiccate the rhizomes and exhaust the food reserves.  
5 It is essential to cut to a depth of at least six inches to ensure that most, if not all, of the rhizomes have  
6 been cut. Results from these practices are evident when observing cogon grass growing up to the  
7 edge of a cultivated field with no evidence of spread into the field itself.

8 Although tillage and herbicides will provide some control and suppression of cogon grass, long-term  
9 eradication is seldom achieved. It has been shown that an integrated approach that combines  
10 burning, tillage (mechanical disturbance), and chemical applications provides the best solution for  
11 cogon grass management. Initially, cogon grass should be burned or mowed to remove excess thatch  
12 and older leaves. This initiates regrowth from the rhizomes, thereby reducing rhizome biomass. It  
13 also allows herbicides to be applied to only actively growing leaves, maximizing herbicide absorption  
14 into the plant. Ideally, burning should take place in the summer. A one- to four-month regrowth  
15 period has been shown to provide a sufficient level of leaf biomass for herbicide treatment. This  
16 targets herbicide applications to be made in the late summer/early fall—approximately one month  
17 prior to the average killing frost, depending on the area. If tillage can be incorporated, then a discing  
18 treatment directly following a burn is the best approach. This will further deplete the rhizome  
19 reserve through dessication and increase the number of shoots per given area. A one- to four-month  
20 regrowth period before herbicide treatment also is needed with this approach as well.

21 When good control of cogon grass has been achieved, it is essential to introduce desirable vegetation  
22 as quickly as possible to prevent cogon grass from re-infesting the area. Several species have been  
23 shown to colonize rapidly and tolerate the residual effects of imazapyr. A wider range of plant species  
24 can be used with glyphosate due to the lack of soil activity. However, cogon grass will eventually  
25 begin to re-infest, regardless of control. Therefore, diligence and persistence are essential to  
26 remove/treat re-infested areas before this grass regains a foothold.

## 27 **Education and Outreach**

28 Heightening awareness among land managers will be necessary to reduce the potential movement of  
29 cogon grass. In an effort to reduce the introduction of new infestations, preventative measures—  
30 including equipment sanitation and off-site material quarantines—should be highlighted through  
31 educational programs.

32 Encouraging proper equipment sanitation practices when operating on infested sites and moving  
33 equipment to other locations to prevent spread include:

- 34 • Cleaning of radiators, screens, and equipment parts that collect seed or come into contact  
35 with the soil and rhizomes
- 36 • Inspecting sources of off-site material, including soil, gravel, and mulch, for invasive species
- 37 • Establishing central staging areas on a property when equipment and material from off-site  
38 are stored or staged to allow easy inspection and monitoring for the introduction of  
39 invasives

# 1 **Mimosa**

2 *Albizia julibrissin*

## 3 **Introduction**

4 Originally from China, Mimosa, or Silk tree, was introduced  
5 to the United States in 1745 and has been cultivated since  
6 the 18th century primarily for use as an ornamental plant.  
7 Mimosa remains a popular ornamental plant because of its  
8 fragrant and showy flowers. Due to its ability to grow and  
9 reproduce along roadways and disturbed areas, and its  
10 tendency to readily establish after escaping from  
11 cultivation, mimosa is considered a Category II invasive by  
12 FLEPPC.

## 13 **Description**

14 Mimosa is a deciduous, small- to medium-sized tree that  
15 can grow 20 feet to 40 feet tall. It is a member of the legume  
16 (*Fabaceae*) plant family and is capable of fixing nitrogen.  
17 The bark is light brown and smooth, while young stems are  
18 lime green in color, turning light brown and covered with  
19 lenticels. Leaves are alternately arranged and bipinnately  
20 compound (six to 20 inches long), having 20 to 60 leaflets  
21 per branch. The leaf arrangement gives mimosa a fern-like  
22 or feathery appearance. Mimosa flowering occurs from  
23 May through July. Pompom-like flowers are borne in  
24 terminal clusters at the base of the current year's twigs.  
25 The flowers are fragrant and pink in color, about 1.5 inches  
26 long. Fruits are flat and in pods, a characteristic of many  
27 legumes. Pods are straw-colored and six inches long,  
28 containing five to 10 light brown oval-shaped seeds about  
29 0.5-inch in length. Pods typically persist on the plant  
30 through the winter months.

31 Mimosa reproduces both vegetatively and by seed. This  
32 characteristic allows the seed to remain dormant for many  
33 years. Normally, seeds are dispersed in close proximity to  
34 the parent plant; however, seeds also can be dispersed by  
35 water. Wildlife also may contribute to the spread of  
36 mimosa through the ingestion and excretion of the seeds.  
37 Vegetative reproduction occurs when trees are cut back,  
38 causing quick resprouting and regrowth.

**CFG Distribution:** 1, 2, 3, 4, 5, 6



### **Status:**

- Category II FLEPPC

### **Identification:**

- Fern-like leaves
- 20 feet to 40 feet tall
- 20 to 60 leaflets per branch
- Seed pods
- Blooms—May through July

### **Treatment:**

- Cutting
- Herbicide application
- Cut stump, basal bark, foliar



1 **Impacts**

2 Mimosa is a strong competitor in open areas or forest edges due to its ability to grow in various soil  
3 types, to produce large numbers of seed, and to resprout when cut back or damaged. An opportunist,  
4 mimosa will take advantage of disturbed areas, either spreading by seed or germinating in  
5 contaminated soil. Mimosa often is seen along roadsides and open vacant lots in urban/suburban  
6 areas and can become a problem along banks of waterways, where its seeds are easily transported  
7 in water.

8 **Management**

9 The first step in preventive control of mimosa is to limit planting and removal of existing plants  
10 within the landscape.

11 There are many native or non-invasive plants that make excellent alternatives to mimosa. These  
12 include serviceberry (*Amelanchier arborea*), redbud (*Cercis canadensis*), flowering dogwood (*Cornus*  
13 *florida*), river birch (*Betula nigra*), fringe tree (*Chionanthus virginicus*), American holly (*Ilex opaca*),  
14 and sweetgum (*Liquidambar styraciflua*).

15 Mimosa can be controlled using a variety of mechanical controls. Power or manual saws can be used  
16 to cut trees at ground level. Control is best achieved by cutting at flowering time before seed  
17 production. Cutting is an initial control measure and will require either an herbicidal control or  
18 repeated cutting for resprouts. In cases where herbicide use is impractical, girdling can be effective  
19 on larger trees. Make a cut through the bark, encircling the base of the tree approximately six inches  
20 above the ground, ensuring the cut goes well below the bark. This will kill the top of the tree but the  
21 tree may resprout and require a follow-up treatment with an herbicide. Hand pulling will effectively  
22 control young seedlings. Plants should be pulled as soon as possible to prevent maturation. The entire  
23 root must be removed since broken fragments may resprout.

24 There are no known biological agents for the control of mimosa.

25 The cut-stump and basal bark herbicidal methods should be considered when treating individual  
26 trees or where the presence of desirable species preclude foliar application. Stump treatments can  
27 be used as long as the ground is not frozen. Horizontally cut stems at or near ground level. Basal bark  
28 applications are effective throughout the year as long as water is not standing at the time of  
29 application.



# 1 Camphor Tree

2 *Cinnamomum camphora*

## 3 Introduction

4 Camphor tree grows natively in China and Japan, where it  
5 is used for oils and timber. In 1875, camphor tree was  
6 introduced into Florida and established in plantations for  
7 camphor production, although it was not profitable for  
8 growers.

9 In Florida, camphor tree is able to rapidly displace native  
10 trees and infest forests and other natural areas. This  
11 invasive species displaces native plants due to its fast  
12 growth habit and the ability to produce large numbers of  
13 seed. This seed is readily eaten and spread by birds.  
14 Nurseries and garden centers sell camphor tree as a  
15 popular ornamental plant, which aids in its dispersal in  
16 landscaped areas.

## 17 Description

18 A quick and easy method of identifying camphor is by  
19 crushing the leaves or peeling a twig or bark. This will  
20 release oils and the scent of camphor. Camphor is an  
21 evergreen tree with oval to elliptical leaves, arranged  
22 alternately on the stem. Slender twigs are initially green  
23 but change to reddish brown. Buds are sharply pointed,  
24 roughly 0.5 inch in length. Camphor tree bark is variable,  
25 from scaly to irregularly furrowed with flat-topped ridges.

26 The camphor tree habit ranges from small to medium (25  
27 feet to 40 feet tall), but some specimens have grown to  
28 more than 100 feet. Leaf margins are entire, but can be  
29 wavy with a shiny, dark green color. Fragrant flowers are  
30 greenish white to pale yellow, borne on panicles about  
31 three inches long. The fruit is dark blue to black, fleshy, and  
32 approximately 1.0 to 1.5 cm in diameter. These are  
33 produced in large quantities during the winter and spring  
34 months in central and north Florida.

## 35 Impacts

36 Camphor tree can be found throughout Florida, Georgia,  
37 and western Texas. Habitats conducive for camphor tree

CFG Distribution: 1, 2, 3, 4, 5, 6



### Status:

- Category I FLEPPC

### Identification:

- Fern-like leaves
- 20 feet to 40 feet tall
- 20 to 60 leaflets per branch
- Seed pods
- Blooms from May through July

### Treatment:

- Cutting
- Herbicide application
- Cut stump, basal bark, foliar



1 establishment are dry, disturbed areas, such as roadsides. Camphor tree also will invade natural  
2 areas. The Florida jujube (*Ziziphus celatus*) is an endangered native species in Polk County that is  
3 being pushed out by camphor tree. Because camphor tree is available in garden centers and  
4 nurseries, homeowners are able to purchase plants, ensuring its survival and spread. This species  
5 also is spread by wildlife, such as birds and other animals that eat the fruit, spreading the seed to  
6 different areas.

## 7 **Management**

8 Preventing the spread and establishment of camphor tree is the first step in a successful management  
9 plan. Since the fruit is the primary means of spread, controlling trees before maturation and fruit  
10 development is critical. With this caution in mind, large trees with heavy fruit potential should be  
11 eliminated first. However, since birds vector the seeds, constant monitoring will be necessary to keep  
12 this species in check.

13 Weeds such as camphor tree generally invade open or disturbed areas—following a burn, clearing,  
14 mowing, etc.—so these areas are particularly vulnerable to invasion. Therefore, a healthy ecosystem  
15 with good species diversity will help to deter infestation. Given this, disturbed areas should be  
16 monitored more frequently and extensively.

17 Mechanical control is particularly effective on seedling trees when smaller equipment can be used to  
18 remove/destroy the plants. Mowing will kill seedling trees and continuous mowing will eventually  
19 kill resprouting shoots from a cut-stump treatment. Discing or other mechanical tillage will kill small  
20 plants, but may encourage subsequent re-infestation due to disturbance. Burning also may provide  
21 good control of camphor tree, but resprouting will likely occur on larger trees. Physical removal of  
22 seedlings and young trees is another tactic, although this may be labor intensive. Take care when  
23 removing small trees.

24 There is limited research and data on biological control of camphor tree.

25 Chemical control can be separated into cut-stump, basal bark, and foliar treatments. Foliar  
26 treatments will work well on young trees less than 10 feet tall.

27 Basal bark treatments are effective for trees up to six inches in diameter with smooth bark. Be sure  
28 to spray around the entire tree, up 12 inches from the base. For larger trees with thick bark, a frill  
29 treatment is recommended. For this application, cut into the bark and peel it back to form a cup.  
30 Herbicide then can be poured into the pocket created by the frill. The number of frills per tree  
31 depends on tree diameter and herbicide used. Cut-stump treatments are effective on trees of all sizes.

# 1 Caesar's Weed

2 *Urena lobata*

## 3 Introduction

4 There are many plants in the family *Malvaceae* that are  
5 grown for ornamental purposes, including hibiscus,  
6 abutilon, and alcea. Cotton (*Gossypium hirsutum*) also is in  
7 this family. Not only does this plant family contain many  
8 ornamentals, but there are also many weedy species, such  
9 as malva, malachra, and urena.

## 10 Description

11 Caesar's weed is an erect shrub that grows up to 10 feet in  
12 height. The plant is single stalked, with free-branching  
13 stems that comprise a bushy appearance. The leaves are  
14 palmately lobed, pubescent with stellate hairs, and four cm  
15 to eight cm long. Flowers are borne in axillary clusters,  
16 pinkish-violet, about one cm across. Fruit is pubescent  
17 with hooked bristles or barbs that cling to clothing or fur.  
18 It grows as an annual species in many areas of Florida, but  
19 may perennate in south Florida.

## 20 Impacts

21 Caesar's weed invades disturbed areas, pastures, eroded  
22 areas, and perennial crop plantations. The species does not  
23 compete well in tall grass and brush lands and does not  
24 grow under forest canopies. Caesar's weed tolerates salt  
25 spray, but does not grow in saturated soils. Caesar's weed  
26 grows rapidly and can reach two feet to seven feet tall by  
27 the end of the first year.

## 28 Management

29 Take care to prevent seed spread into "clean areas." The  
30 seed of Caesar's weed clings to clothing; therefore, treat  
31 plants before seed set. Avoid treating and then traveling to  
32 other areas. Also, avoid driving vehicles through areas of  
33 Caesar's weed. Shade will help to deter growth and limit  
34 seedling establishment. Mulches or other ground cover  
35 also will prevent seed germination. There are no known  
36 biological controls for this species. There is only limited  
37 research in the area of chemical control.

CFG Distribution: 1, 2, 3, 4, 5, 6



### Status:


- Florida Noxious Weed List
- Category I FLEPPC

### Identification:

- Single-stalked with free branching stems
- Palmately lobed leaves
- Up to 10 feet tall
- 20 to 60 leaflets per branch
- Axillary cluster flowers

### Treatment:

- Herbicide application



# 1 Skunk vine

2 *Paederia foetida*

## 3 Introduction

4 Sometime before 1897 at a USDA Field Station, *Paederia*  
5 *foetida*, or skunk vine, was introduced from Asia to  
6 Hernando County, Florida, as a potential fiber crop. Skunk  
7 vine was reported as a troublesome weed very early in its  
8 introduction, escaping into native areas throughout  
9 Florida. It was soon recognized as an economically  
10 important invasive weed. Currently, skunk vine is found in  
11 at least 17 counties in central and north central Florida.

## 12 Description

13 Skunk vine is a woody vine that does not have thorns. Its  
14 vines are able to grow 30 feet in length, climbing up into  
15 tree canopies or crawling along the ground. For some  
16 unknown reason, the vines constantly twine to the right.  
17 The smelly, foul odor released when skunk vine is crushed  
18 is a useful characteristic that can aid in identification.  
19 Skunk vine leaves vary in size and shape. Generally, skunk  
20 vine leaf blades have rounded to cordate (heart) shaped  
21 bases and acuminate (pointed) tips, with entire (smooth)  
22 margins. Leaves may be opposite on the stem. In rare  
23 instances, leaves also have been found in whorls of three.  
24 Leaves and flowers are on petioles about 2.5 inches long.  
25 Skunk vine flowers are small, light grayish pink or lilac,  
26 with red centers. The fruit is small, spherical, and shiny  
27 brown, having two black, non-winged seeds. Skunk vine is  
28 able to reproduce vegetatively and via seed. Its stems are  
29 able to root readily in soil. It is thought that seeds are eaten  
30 by frugivorous birds and spread, but this has not yet been  
31 verified.

## 32 Impacts

33 Skunk vine is able to survive in a variety of Florida habitats,  
34 including hardwood, mixed forests, and pine forests,  
35 sandhills, and floodplain forest and marsh. A serious  
36 invasive weed, skunk vine is able to displace native  
37 vegetation. The dense layer of vegetation created by skunk  
38 vine can both damage and kill native vegetation. Climbing  
39 vines can engulf and cover trees and shrubs. The weight of

### CFG Distribution: 1



### Status:

- Florida Noxious Weed List
- Category I FLEPPC

### Identification:

- Woody vine
- Leaf blades have rounded to cordate-shaped bases and acuminate tips with smooth margins
- Up to 30 feet in length
- Leaves and flowers on petioles about 2.5 inches long

### Treatment:

- Mowing and tillage
- Triclopyr





1 the vine mass climbing over vegetation can cause branches or entire trees to break or collapse.  
2 Crawling vines can form a dense layer of vegetation, smothering many shrubs and other plants  
3 growing in the understory.

#### 4 **Management**

5 Take care when disposing of skunk vine, since it is able to regrow when cut back and produce new  
6 plants from stem fragments. Seeds also are able to germinate in brush piles. Prevent the transport of  
7 stem fragments and seeds to other locations by ensuring machinery is free of seed and stem  
8 fragments. Flooded conditions can decrease the vigor of skunk vine; however, skunk vine can live in  
9 marsh-like conditions, able to survive for approximately 190 days under water.

10 Weeds such as skunk vine generally invade open or disturbed areas—following a burn, clearing  
11 mowing, etc.—so these areas are particularly vulnerable to invasion. Therefore, a healthy ecosystem  
12 with good species diversity will help to deter infestation.

13 Mowing and tillage will provide some measure of control, but are impractical in most situations.

14 There is much hope in biological control agents collected in Japan and Nepal by Agricultural Research  
15 Service Entomologists Robert Pemberton and Paul Pratt. Chrysomelid leaf beetles and two sawfly  
16 species were found feeding on the leaves of skunk vine. A stem gallmaker and a moth in the *Sessiidae*  
17 family also were found to attack skunk vine by causing the formation of galls on vine stems. The flea  
18 beetle (*Trachyaphthona sordida*) has the greatest potential of all of the biological control agents listed  
19 to potentially control skunk vine. Flea beetles damage the host root system by feeding on roots and  
20 root hairs, leading to reduced uptake of nutrients and water by the host plant.

21 Chemical control is one of the most effective means of control for skunk vine, but single applications  
22 generally will not provide complete control. This is due to resprouting from rootstocks or root  
23 crowns. If skunk vine is growing up into trees or other desirable species, vines should be cut or pulled  
24 down to minimize damage to the desirable vegetation. Pulling the vines down without severing them  
25 from the root crown will allow the herbicide to move into the root and provide better control. The  
26 best time to apply an herbicide is in the spring and summer when skunk vine is actively growing. Be  
27 sure to allow adequate time for the plant to regrow from the winter to ensure movement of the  
28 herbicide back into the roots. (As plants grow and mature, they begin to move sugars back into the  
29 roots.)

# 1 Natal Grass

2 *Melinis repens*

## 3 Introduction

4 Natal grass, is a short-lived perennial grass native to South  
5 Africa, the Arabian Peninsula, India, the Seychelle Islands,  
6 and the Canary Islands. It is already widely distributed in  
7 tropical and subtropical regions due to its long use as a  
8 pasture grass and ornamental plant. Although considered  
9 a weed in many countries, it is not currently regulated.

10 It mainly occurs in disturbed areas, such as along roadsides  
11 and railway lines, but it can spread into natural areas  
12 interfering with early successional processes. In central  
13 and subtropical Florida, natal grass occurs in disturbed  
14 uplands, drained hydric pine and cypress prairies, pine  
15 rocklands, scrub habitats, flatwoods, firebreaks, and  
16 sandhill restoration areas. As of 1999, natal grass occurred  
17 in 49 percent of nature preserves in southern Florida.  
18 Natal grass also occurs in sand pine scrub; experiments are  
19 ongoing in the Ocala National Forest, central Florida. Natal  
20 grass does not persist on soils that retain moisture and is  
21 inhibited by flooding (as little as one month of inundation).  
22 It also is inhibited by shade and cold, although seeds can  
23 germinate after freezing.

## 24 Description

25 An annual or short-lived perennial grass growing 20 cm to  
26 150 cm in height. Culms (stems) root from the lower nodes,  
27 but stems are held upright. The leaf blades are flat, five cm  
28 to 30 cm long; two mm to 10 mm wide. The flowers are  
29 clustered in a fluffy oblong or ovate panicle, five cm to 20  
30 cm long. Spikelets are two mm to 10 mm long, two-  
31 flowered, with the lower floret male, the upper floret a  
32 hermaphrodite. They are densely villous with hairs up to  
33 eight mm long, on very fine pedicels with sparse long hairs.  
34 Panicles often have a rosy color from the long silky hairs  
35 attached to the triangular fruits. The color fades to silvery-  
36 white with age.

**CFG Distribution:** 1 and 2



### Status:

- Florida Noxious Weed List
- Category I FLEPPC

### Identification:

- Woody vine
- Leaf blades flat 5-30 cm long; 2-10 mm wide.
- Up to 30 feet in length
- Flowers are clustered

### Treatment:

- Remove seed heads  
Herbicides containing  
fluazifop or glyphosate



1 **Impacts**

2 Natal grass establishes and regenerates most successfully on sites that have been mechanically  
3 disturbed, including road edges and firelines, but it also can spread from the edges into the interior  
4 of undisturbed areas. It can thrive in drained sites, mowed firebreaks, fire-created openings, and  
5 mined areas. It can form a monoculture in disturbed areas, reducing native species, particularly  
6 graminoids.

7 **Management**

8 Public awareness is important in potentially reducing the purchase of natal grass as a landscape  
9 plant. Natal grass is currently controlled mainly through physical/ mechanical and chemical methods  
10 both in agricultural systems and natural areas. Preventing seed set reduces local dispersal. It is also  
11 important to clean agricultural and landscape equipment when used in infested areas.

12 Prevent seed-set by removing or cutting grass prior to flowering. Cut out small populations. Fire kills  
13 seeds and adult plants. Fire may be ineffective in reducing natal grass if rainfall is abundant after the  
14 fire. Cattle and sheep eat natal grass but have not been used to control it. Since seeds are wind-  
15 dispersed, removal of seed heads could significantly reduce spread. No biological control is currently  
16 available. Herbicides may be used to control natal grass. However, the best control is achieved when  
17 plants are sprayed before flowering and seed-set or after germination following a fire.

1 *Nuisance and Exotic Animals*

2 Exotic animal species include non-native wildlife species, free-ranging domesticated pets or livestock, and  
3 feral animals. Because of the negative impacts to natural systems attributed to exotic animals, the DRP  
4 actively removes exotic animals from state parks, with priority being given to those species causing the  
5 most ecological damage. A nuisance animal is an individual native animal whose presence or activities  
6 create special management problems.

7 Invasive non-native animal species are not as large an issue on the CFG as invasive plants. An exotic animal  
8 species of concern found on the CFG is the feral hog. Having an inherently high reproductive rate, when  
9 populations of wild hogs are left unchecked, these voracious omnivores are known to significantly degrade  
10 natural communities through foraging activity (rooting). Feral hogs forage in the park's wetter areas and  
11 seasonally in scrub when acorn crops are plentiful. In the eastern region of the CFG, hogs are having  
12 negative impacts on wetlands, depression marshes, and listed plants. The greenway has an active program  
13 of feral hog removal. Park staff and contractors both participate in hog removal as conditions warrant. CFG  
14 issues permits for hog trappers in areas where hogs are known to be a problem, such as Putnam County  
15 CFG lands, Marshall Swamp, and Inglis Island. The CFG staff also will consult with other regional natural  
16 resource managing agencies and private land owners to coordinate wild hog control measures as  
17 necessary.

18 Immigration of feral cats (*Felis catus*) and dogs (*Canis lupus familiaris*) from the surrounding residential  
19 areas to park lands is an ongoing concern for the protection of natural and cultural resources. Local animal  
20 control agencies are contacted for control of these species when needed. Some monkeys from Silver  
21 Springs are on the CFG. Control of these non-native animals is under the authority of the FWC. Coyotes are  
22 reportedly common in the eastern region of the CFG; no control methods are being used for this species.  
23 Management goals, objectives, and actions for management of invasive exotic plants and exotic and  
24 nuisance animals are discussed in the Resource Management Program section of this component.

25 **Cultural Resources**

26 The Cross Florida Barge Canal project was a canal project to connect the Gulf of Mexico and the Atlantic  
27 Ocean across Florida for barge traffic. The idea of such a canal was first proposed by Philip II of Spain in  
28 1567. It was repeatedly considered over the years, but found to be economically unviable. Secretary of  
29 War John C. Calhoun once again proposed a canal in 1818 in an attempt to solve the losses due to  
30 shipwrecks and piracy.

31 In the 1930s, regional politicians lobbied the federal government to fund canal construction as an  
32 economic recovery program, and President Franklin D. Roosevelt allocated emergency funds in 1935.  
33 Local opponents of the canal protested that the canal would deplete Florida's aquifers, and work was  
34 stopped one year later.

35 Work was reauthorized in 1942 as a national defense project, with dams and locks to protect the  
36 underground water supply. Support for the project from Washington was sporadic, and funds were never  
37 allocated to the USACE for construction. Planning was once again initiated in 1963 with support from  
38 President John F. Kennedy, who allocated \$1 million dollars for the project. The next year, President  
39 Lyndon B. Johnson set off the explosives that initiated construction. It was hoped, that the canal along with  
40 the St. Johns-Indian River Barge Canal, would provide a quicker safer route across Florida by 1971.



1 Opponents subsequently campaigned against the canal based on environmental concerns and the project  
2 stopped again in January 1971. The project was officially cancelled in 1991. The right of way was turned  
3 over to the state and officially became the Marjorie Harris Carr Cross Florida Greenway, named in honor  
4 of the woman who led opposition to the canal.

5 This section addresses the cultural resources present in the CFG that may include archaeological sites,  
6 historic buildings and structures, cultural landscapes, and collections. The Florida Department of State  
7 (FDOS) maintains the master inventory of these resources through the Florida Master Site File (FMSF).  
8 State law requires that all state agencies locate, inventory, and evaluate cultural resources that appear to  
9 be eligible for listing in the National Register of Historic Places (NRHP). For the purposes of this plan,  
10 significant archaeological sites, significant structures, and landscapes mean those cultural resources listed  
11 or eligible for listing in the NRHP. The terms “archaeological site,” “historic structure,” or “historic  
12 landscape” refer to all resources that are or will become 50 years old during the term of this plan.

### 13 **Cultural Resources in the CFG**

14 The archaeological sites and other historical resources within the CFG represent the range of Florida’s  
15 cultural periods from the Paleoindian period (10,000 B.C.–8,000 B.C.) to the twentieth century and a range  
16 of site types as well. Prehistoric artifact scatters to 19th to 20th century railroad corridors to historic  
17 standing structures have been identified within the park. However, not all of the resources have been  
18 considered by the State Historic Preservation Office (SHPO) to be significant. At the same time, it should  
19 be noted that the majority of the archaeological sites in the CFG have not been evaluated by SHPO and the  
20 majority of the park has not been subjected to Phase I level cultural resource assessments.

21 According to the information in the FMSF, seven archaeological sites (8MR1082, 8MR1878, 8MR2357,  
22 8MR3526, 8MR3863, 8MR3865 and 8MR3866), three resource groups (8MR3410, 8MR3563 and  
23 8MR3762)), and a historic bridge (8MR3858) have been determined to be eligible for listing in the National  
24 Register of Historic Places. SHPO has determined that insufficient information exists to make a significance  
25 determination for three archaeological sites (8MR2549, 8MR2556, and 8MR3925), a linear resource  
26 (8CI1223), and a historic structure (8MR1515). All archaeological sites and historic resources in the park  
27 are protected; however, the sites listed or eligible for listing in the National Register of Historic Places  
28 warrant higher profile monitoring and measures to stabilize and mitigate deterioration and disturbance.  
29 Additional information or data relative to any of the sites will be submitted to DHR and the FMSF.

30 There is a cemetery (8MR3923) on the CFG, which is indicated as abandoned in the FMSF records. SHPO  
31 has determined the cemetery to be ineligible for listing in the National Register of Historic Places. The  
32 condition of the cemetery will be monitored and access will be maintained for family members of those  
33 interred at the cemetery. Finally, there are several archaeological sites recorded as middens, mounds, or  
34 burial mounds on the CFG. These sites will be monitored due to potential for human burials/remains at  
35 these locations.

36 Table 13 provides a complete listing by county of the 265 CFG cultural sites listed in the FMSF.

### 37 **Condition Assessment**

38 The CFG contains 265 sites listed in the FMSF, which includes 85 sites within Citrus County; five sites  
39 within Levy County; 162 sites within Marion County; and 13 sites within Putnam County. Given this large  
40 number of sites and the limited staff at the CFG, evaluation of the condition of all of these resources has

1 not been completed nor is it anticipated to be completed by the time this plan is adopted. Additional  
2 resources through contractors and university researchers should be acquired to assist with this effort.

3 Condition assessments of these cultural resources must be completed by the authorized CFG staff (who  
4 have taken the Archeological Resources Monitoring training). Meetings with the Bureau of Natural and  
5 Cultural Resources and Office of Park Planning concluded that prioritization of these sites would be  
6 acceptable for the completion of condition assessments. Recommendations were made that the sites be  
7 prioritized and condition assessments should only be conducted for the UMP based on the following  
8 priority levels:

- 9 1. The 11 sites that are listed or eligible for listing on the NRHP
- 10 2. Sites in/around areas proposed for development (e.g., new trailheads, trails, etc.)
- 11 3. Sites in/around areas immediately accessible by visitors to the park, primarily involving  
12 trailheads

13 Twenty-three sites that have already been determined “Not Significant” do not require an assessment. The  
14 remaining sites (and any sites not assessed from the three priority levels stated above) are listed in the  
15 UMP as “Not Assessed” and will be assessed during the 2017-2027 planning period.

16 Condition assessments should be accomplished using a three-part evaluation scale, expressed as good,  
17 fair, and poor. These terms describe the present condition, rather than comparing what exists to the ideal  
18 condition. Good describes a condition of structural stability and physical wholeness, where no obvious  
19 deterioration other than normal decline occurs. Fair describes a condition in which there is a discernible  
20 decline in condition between inspections, and the wholeness or physical integrity is and continues to be  
21 threatened by factors other than normal wear. A fair assessment usually is a cause for concern. Poor  
22 describes an unstable condition where there is palpable, accelerating decline and physical integrity is  
23 being compromised quickly. A resource in poor condition suffers obvious deterioration in physical  
24 integrity from year to year. A poor condition suggests immediate action is needed to re-establish physical  
25 stability.

## 26 **Level of Significance**

27 Applying the criteria for listing in the NRHP involves the use of contexts as well as an evaluation of integrity  
28 of the site. A cultural resource’s significance derives from its historical, architectural, ethnographic, or  
29 archaeological context. Evaluation of cultural resources will result in a designation of NRL (National  
30 Register or National Landmark Listed or located in an NR district), NR (National Register eligible), NE (not  
31 evaluated), or NS (not significant), as indicated in Table 13.

32 There are no criteria for use in determining the significance of collections or archival material. Usually,  
33 significance of a collection is based on what or whom it may represent. For instance, a collection of  
34 furniture from a single family and a particular era in connection with a significant historic site would be  
35 considered highly significant. In the same way, a high-quality collection of artifacts from a significant  
36 archaeological site would be of important significance. A large herbarium collected from a specific park  
37 over many decades could be valuable to resource management efforts. Archival records are most  
38 significant as a research source. Any records depicting critical events in the park’s history, including  
39 construction and resource management efforts, would all be significant.

1 **Table 13. Cultural Sites Listed in the Florida Master Site File**

Site Name	Site Number	Culture/Period	Description	Significance	Condition	Treatment
<b>Citrus County Sites</b>						
Burtine's	8CI0058	Deptford	Archaeological Site	NE	NE	TBD
Captain Joe Island 1	8CI0062	Prehistoric (redeposited)	Archaeological Site	NE	NE	TBD
Captain Joe Island 2	8CI0063	Prehistoric	Archaeological Site	NE	NE	TBD
Everett Island	8CI0064	Prehistoric	Archaeological Site	NE	NE	TBD
Richardson Creek	8CI0065	Prehistoric	Archaeological Site	NE	NE	TBD
Florida Barge Canal 1	8CI0066	Prehistoric	Archaeological Site	NE	NE	TBD
Florida Barge Canal 12	8CI0070	Prehistoric	Archaeological Site	NE	NE	TBD
Florida Barge Canal 14	8CI0072	Prehistoric	Archaeological Site	NE	NE	TBD
Florida Barge Canal 16	8CI0074	Prehistoric	Archaeological Site	NE	NE	TBD
Florida Barge Canal 18	8CI0075	Prehistoric	Archaeological Site	NE	NE	TBD
Bennett's Creek 2	8CI0090	Deptford, Weeden Island	Archaeological Site	NE	NE	TBD
Withlacochee 13 Midden	8CI0096	Archaic, St. Johns, Swift Creek	Archaeological Site	NE	NE	TBD
Withlacochee 10 Midden	8CI0100	Prehistoric	Archaeological Site	NE	NE	TBD
Pat's Elbow, John Gibson Homestead	8CI0101	Prehistoric, Deptford, Weeden Island, Historic	Archaeological Site	NE	NE	TBD
Withlacochee 5 Midden	8CI0102	Weeden Island	Archaeological Site	NE	NE	TBD
FPC 18	8CI0106	Prehistoric	Archaeological Site	NE	NE	TBD
FPC 17	8CI0107	Prehistoric	Archaeological Site	NE	NE	TBD
Steamship LT Izard Wreck	8CI0122	Historic 1821-1845	Underwater Site	NE	NE	TBD
Ouithla 3	8CI0330	Weeden Island	Archaeological Site	NE	NE	TBD

Site Name	Site Number	Culture/Period	Description	Significance	Condition	Treatment
High Knoll	8CI0893	Prehistoric	Archaeological Site	NE	NE	TBD
Withlacoochee Bend	8CI1076	Orange, Deptford, Weeden Island	Archaeological Site	NE	NE	TBD
Bennett's Creek	8CI1089	Paleoindian, Orange, Deptford, Weeden Island	Archaeological Site	NE	NE	TBD
Buckford 1	8CI1134	Prehistoric	Archaeological Site	NE	NE	TBD
Buckford 2	8CI1135	Prehistoric	Archaeological Site	NE	NE	TBD
Wekiwa	8CI1136	Deptford, Weeden Island	Archaeological Site	NE	NE	TBD
Sinte 1	8CI1137	Prehistoric Weeden Island	Archaeological Site	NE	NE	TBD
Sinte 2	8CI1138	Prehistoric, Weeden Island	Archaeological Site	NE	NE	TBD
Beaten Face	8CI1139	Prehistoric	Archaeological Site	NE	NE	TBD
Thla Rakke 1	8CI1140	Prehistoric	Archaeological Site	NE	NE	TBD
Little Peak	8CI1141	Deptford, Weeden Island	Archaeological Site	NE	NE	TBD
Thla Rakke 2	8CI1142	Prehistoric	Archaeological Site	NE	NE	TBD
Stuck Nowhere	8CI1143	Deptford	Archaeological Site	NE	NE	TBD
Blowing Willows	8CI1144	Deptford, Weeden Island	Archaeological Site	NE	NE	TBD
Falling off the Rock	8CI1145	Prehistoric	Archaeological Site	NE	NE	TBD
Many Places	8CI1146	Prehistoric	Archaeological Site	NE	NE	TBD
Crunchy Ground	8CI1147	Prehistoric	Archaeological Site	NE	NE	TBD
Old Snake's Path	8CI1148	Prehistoric	Archaeological Site	NE	NE	TBD



Site Name	Site Number	Culture/Period	Description	Significance	Condition	Treatment
Berry Cakes	8CI1149	Prehistoric	Archaeological Site	NE	NE	TBD
Grandfather Egret's Pool	8CI1150	Prehistoric	Archaeological Site	NE	NE	TBD
Mother's Dimple	8CI1151	Prehistoric	Archaeological Site	NE	NE	TBD
Moving Dirt	8CI1152	Prehistoric	Archaeological Site	NE	NE	TBD
Two Rock Houses	8CI1153	Prehistoric	Archaeological Site	NE	NE	TBD
Crab Warriors	8CI1154	Prehistoric	Archaeological Site	NE	NE	TBD
Gossiping Palms	8CI1155	Prehistoric	Archaeological Site	NE	NE	TBD
Limus	8CI1156	Prehistoric	Archaeological Site	NE	NE	TBD
Terrapin Wipes His Nose	8CI1157	Prehistoric	Archaeological Site	NE	NE	TBD
Weeping Rock	8CI1158	Prehistoric	Archaeological Site	NE	NE	TBD
Dying Cedars	8CI1161	Prehistoric	Archaeological Site	NE	NE	TBD
Grandma Sits Alone	8CI1162	Prehistoric	Archaeological Site	NE	NE	TBD
Fish Splashers	8CI1163	Prehistoric	Archaeological Site	NE	NE	TBD
Fiery Palms	8CI1164	Prehistoric, 19 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Rocky Place	8CI1165	Prehistoric	Archaeological Site	NE	NE	TBD
Broken Cups	8CI1166	Prehistoric	Archaeological Site	NE	NE	TBD
Bird Town	8CI1167	Prehistoric	Archaeological Site	NE	NE	TBD
Pond in the Rock	8CI1168	Prehistoric	Archaeological Site	NE	NE	TBD
JD's Site	8CI1169	Weeden Island	Archaeological Site	NE	NE	TBD
Kathy's	8CI1170	Deptford, Weeden Island	Archaeological Site	NE	NE	TBD
Melissa's Site	8CI1178	Prehistoric	Archaeological Site	NE	NE	TBD
Telling Secrets	8CI1182	Prehistoric	Archaeological Site	NE	NE	TBD

Site Name	Site Number	Culture/Period	Description	Significance	Condition	Treatment
Feet Getting Wet	8CI1183	Prehistoric	Archaeological Site	NE	NE	TBD
Everett Island 1	8CI1184	Prehistoric	Archaeological Site	NE	NE	TBD
Everett Island 6	8CI1185	Prehistoric, Deptford, Weeden Island	Archaeological Site	NE	NE	TBD
Turtle Left His Shell	8CI1186	Prehistoric	Archaeological Site	NE	NE	TBD
Scorpion Palace	8CI1187	Prehistoric	Archaeological Site	NE	NE	TBD
Toppled Cedars	8CI1188	Prehistoric, Weeden Island	Archaeological Site	NE	NE	TBD
Everett Island 2	8CI1189	Prehistoric	Archaeological Site	NE	NE	TBD
Everett 3	8CI1190	Prehistoric, Weeden Island	Archaeological Site	NE	NE	TBD
Everett 4	8CI1191	Prehistoric, Weeden Island	Archaeological Site	NE	NE	TBD
Everett 5	8CI1192	Deptford, Weeden Island	Archaeological Site	NE	NE	TBD
Dunnellon Abandoned Railroad Line	8CI1223	Historic	Linear Resource	INSF	NE	TBD
Trout Creek 1	8CI1314	Deptford, Weeden Island	Archaeological Site	NE	NE	TBD
Trout Creek 2	8CI1315	Deptford, Weeden Island	Archaeological Site	NE	NE	TBD
Trout Creek 3	8CI1316	Prehistoric	Archaeological Site	NE	NE	TBD
Trout Creek 4	8CI1317	Weeden Island	Archaeological Site	NE	NE	TBD
Trout Creek 5	8CI1318	Deptford, Weeden Island	Archaeological Site	NE	NE	TBD
John's Creek	8CI1319	Deptford, Weeden Island	Archaeological Site	NE	NE	TBD
Trout Creek 6	8CI1320	Prehistoric	Archaeological Site	NE	NE	TBD

Site Name	Site Number	Culture/Period	Description	Significance	Condition	Treatment
Bennett's Creek 1	8CI1321	Prehistoric	Archaeological Site	NE	NE	TBD
Bennett's Creek 4	8CI1324	Prehistoric	Archaeological Site	NE	NE	TBD
Bennett's Creek	8CI1324	Prehistoric	Archaeological Site	NE	NE	TBD
John's Creek 2	8CI1325	Deptford, Weeden Island	Archaeological Site	NE	NE	TBD
John's Creek 3	8CI1326	Deptford	Archaeological Site	NE	NE	TBD
John's Creek 4	8CI1327	Weeden Island	Archaeological Site	NE	NE	TBD
Ouithla 1	8CI1328	Prehistoric	Archaeological Site	NE	NE	TBD
Ouithla 5	8CI1332	Prehistoric	Archaeological Site	NE	NE	TBD
<b>Levy County Sites</b>						
Florida Barge Canal 1	8LV0089	Prehistoric	Archaeological Site	NE	NE	TBD
Florida Barge Canal 8	8LV0090	Archaic	Archaeological Site	NE	NE	TBD
Florida Barge Canal 9	8LV0091	Prehistoric	Archaeological Site	NE	NE	TBD
Florida Barge Canal 10	8LV0092	Prehistoric	Archaeological Site	NE	NE	TBD
Florida Barge Canal 11	8LV0093	Prehistoric	Archaeological Site	NE	NE	TBD
<b>Marion County Sites</b>						
Cedar Landing 2	8MR0005	Prehistoric	Archaeological Site	NE	NE	TBD
Cedar Landing 3	8MR0007	Orange, St. Johns	Archaeological Site	NE	NE	TBD
Watkin's Camp Mound	8MR0008	Prehistoric	Archaeological Site	NE	NE	TBD
Tobacco Patch Landing	8MR0010	St. Johns	Archaeological Site	NE	NE	TBD
Eureka Log Landing	8MR0012	St. Johns	Archaeological Site	NE	NE	TBD

Site Name	Site Number	Culture/Period	Description	Significance	Condition	Treatment
Sunday Bluff	8MR0013	Orange, Deptford, Middle and Late Archaic, St. Johns, 19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Old Site Eaton Creek	8MR0014	Prehistoric, Orange, St. Johns	Archaeological Site	NE	NE	TBD
Eaton Creek	8MR0015	Indeterminate	Archaeological Site	NE	NE	TBD
Eaton Creek Midden	8MR0016	Orange	Archaeological Site	NE	NE	TBD
Palmetto Landing Mound 6	8MR0024	Orange, St. Johns	Archaeological Site	NE	NE	TBD
Gore's Landing Midden	8MR0030	Prehistoric	Archaeological Site	NE	NE	TBD
Gore's Landing Mound	8MR0031	Prehistoric, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Dek's Landing Mound	8MR0032	Prehistoric	Archaeological Site	NE	NE	TBD
No Name	8MR0044	St. Johns	Archaeological Site	NE	NE	TBD
Colby Landing (Florida Barge Canal #28)	8MR0057	Late Archaic, Transitional, St. Johns	Archaeological Site	NE	NE	TBD
No Name	8MR0073	Prehistoric, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Shell Knoll Mound	8MR0075	Prehistoric, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Shell Knoll Landing	8MR0076	Prehistoric, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Payne's Landing	8MR0077	Indeterminate	Archaeological Site	NE	NE	TBD
No Name	8MR0079	20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Gore's Landing Borrow Pit	8MR0080	Archaic	Archaeological Site	NE	NE	TBD
No Name	8MR0091	Indeterminate	Archaeological Site	NE	NE	TBD
Eureka Bluff	8MR0096	Prehistoric	Archaeological Site	NE	NE	TBD
Florida Barge Canal 29	8MR0097	Indeterminate	Archaeological Site	NE	NE	TBD



Site Name	Site Number	Culture/Period	Description	Significance	Condition	Treatment
Ross Prairie 1	8MR0100	Archaic, Deptford, Weeden Island	Archaeological Site	NE	NE	TBD
Ross Prairie 2	8MR0101	Archaic	Archaeological Site	NE	NE	TBD
Ross Prairie 3	8MR0102	Archaic	Archaeological Site	NE	NE	TBD
Marion County Farm	8MR0103	Archaic	Archaeological Site	NE	NE	TBD
175 A	8MR0104	Archaic, 19 <sup>th</sup> and 29 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
175 B	8MR0105	Archaic	Archaeological Site	NE	NE	TBD
Road 275	8MR0106	Archaic, St. Johns	Archaeological Site	NE	NE	TBD
Near Blue Springs	8MR0107	St. Johns	Archaeological Site	NE	NE	TBD
Orange Springs Ferry Road Mound	8MR0127	Prehistoric, St. Johns, 19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
MacDonald Tobacco Shed	8MR0133	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
New Yarbrough Place	8MR0134	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Double Bridge Mound A	8MR0148	Prehistoric, Orange, Deptford, St. Johns	Archaeological Site	NE	NE	TBD
Double Bridge Mound B	8MR0149	Prehistoric, Archaic, Deptford	Archaeological Site	NE	NE	TBD
Turtle Egg	8MR0164	St. Johns	Archaeological Site	NE	NE	TBD
Oklawaha River Shell Mound II	8MR0224	Orange, St. Johns	Archaeological Site	NE	NE	TBD
No Name	8MR0231	Prehistoric, Archaic	Archaeological Site	NE	NE	TBD
No Name	8MR0232	Prehistoric, Archaic	Archaeological Site	NE	NE	TBD
USFS 81-60	8MR0247	Prehistoric, Archaic	Archaeological Site	NE	NE	TBD

Site Name	Site Number	Culture/Period	Description	Significance	Condition	Treatment
TJ's Midden (Piney Island Midden)	8MR0255	Mt. Taylor, St. Johns	Archaeological Site	NE	NE	TBD
Nina's Dream	8MR0262	Prehistoric, Late Archaic, Transitional, St. Johns, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Cedar Landing South	8MR0390	Prehistoric, St. Johns, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
North Barge Canal	8MR0475	Archaic, Alachua	Archaeological Site	NE	NE	TBD
I-75 Pond	8MR0476	Archaic, Deptford, St. Johns	Archaeological Site	NE	NE	TBD
South Barge Canal	8MR0477	Archaic	Archaeological Site	NE	NE	TBD
Barge Canal Historic	8MR0478	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Barge Canal Farm 2	8MR0479	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Crepe Myrtle Farm 3	8MR0480	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Turpentine	8MR0481	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Barge Canal Farm 5	8MR0482	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Barge Canal Farm 8	8MR0483	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Barge Canal Farm 9	8MR0484	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Marshall Swamp	8MR0798	Orange, Transitional	Archaeological Site	NE	NE	TBD
Cedar Creek Still	8MR0825	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Piney Island	8MR0848	Paleoindian, Early and Late Archaic, Alachua, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Ross Prairie 4	8MR0932	Prehistoric	Archaeological Site	NS	NE	TBD
Shady Sink	8MR0998	Prehistoric	Archaeological Site	NE	NE	TBD

Site Name	Site Number	Culture/Period	Description	Significance	Condition	Treatment
Canal	8MR1007	Paleoindian, Early and Late Archaic, St. Johns	Archaeological Site	NE	NE	TBD
Franklin 3	8MR1071	Indeterminate	Archaeological Site	NE	NE	TBD
Franklin 8	8MR1073	Indeterminate	Archaeological Site	NE	NE	TBD
Franklin 15	8MR1082	Prehistoric, Middle and Late Archaic, Orange, 19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NR	NE	TBD
F 84	8MR1103	Indeterminate	Archaeological Site	NE	NE	TBD
*Ross Prairie A	8MR1119	Unknown	Archaeological Site	NE	NE	TBD
*Ross Prairie B	8MR1120	Unknown	Archaeological Site	NE	NE	TBD
*Ross Prairie E	8MR1122	Unknown	Archaeological Site	NE	NE	TBD
*Ross Prairie F	8MR1123	Unknown	Archaeological Site	NE	NE	TBD
*Ross Prairie G	8MR1124	Unknown	Archaeological Site	NE	NE	TBD
*Ross Prairie H	8MR1125	Unknown	Archaeological Site	NE	NE	TBD
Holy Band of Inspiration #1	8MR1515	Circa 1890	Historic Structure	INSF	NE	TBD
Butterbutt Landing	8MR1869	Prehistoric	Archaeological Site	NE	NE	TBD
Cactus Flower	8MR1878	Prehistoric, Paleoindian, Middle Archaic, Achaic, Orange, Alachua, St. Johns	Archaeological Site	NR	NE	TBD
Turkey Landing	8MR1926	Prehistoric	Archaeological Site	NE	NE	TBD
Tuten Creek Mounds	8MR1972	Paleoindian, Archaic, Late Archaic, Orange, St. Johns	Archaeological Site	NE	NE	TBD

Site Name	Site Number	Culture/Period	Description	Significance	Condition	Treatment
DiCarlo	8MR2060	Early Archaic	Archaeological Site	NE	NE	TBD
Backcurrent	8MR2062	Early Archaic, Historic	Archaeological Site	NE	NE	TBD
Turkey Landing	8MR2063	Late Archaic, St. Johns, First Spanish, 19 <sup>th</sup> and 29 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Conner Landing	8MR2064	Late Archaic, St. Johns, Historic, 19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Stallings	8MR2065	Early Archaic	Archaeological Site	NE	NE	TBD
Gore's Landing	8MR2066	St. Johns, Historic	Archaeological Site	NE	NE	TBD
Olsen	8MR2067	Historic	Archaeological Site	NE	NE	TBD
Osceola	8MR2076	Prehistoric, Historic	Archaeological Site	NE	NE	TBD
Llama	8MR2117	Prehistoric	Archaeological Site	NS	NE	TBD
Heather Island Preserve	8MR2223	Prehistoric, Paleoindian, Archaic, Weeden Island, St. Johns, First Spanish Period, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Pond D	8MR2355	Prehistoric	Archaeological Site	NS	NE	TBD
Surface	8MR2356	Prehistoric	Archaeological Site	NS	NE	TBD
Ross Prairie	8MR2357	Prehistoric, Weeden Island, Safety Harbor	Archaeological Site	NR	NE	TBD
Sharps Ferry Office	8MR2402	Prehistoric, Archaic	Archaeological Site	NE	NE	TBD
Sharps Ferry Field	8MR2403	Prehistoric, St. Johns	Archaeological Site	NE	NE	TBD
The Iron Bridge	8MR2548	1929	Historic Bridge	NE	NE	TBD



Site Name	Site Number	Culture/Period	Description	Significance	Condition	Treatment
98-3, Ocala	8MR2549	Middle Archaic, St. Johns, 20 <sup>th</sup> Century	Archaeological Site	INSF	NE	TBD
98-5, Ocala	8MR2551	Prehistoric	Archaeological Site	NS	NE	TBD
98-9, Ocala	8MR2555	Prehistoric	Archaeological Site	NE	NE	TBD
98-10, Ocala	8MR2556	Prehistoric	Archaeological Site	INSF	NE	TBD
Shangri-La	8MR2662	19 <sup>th</sup> and 29 <sup>th</sup> Century	Archaeological Site	NS	NE	TBD
Oklawaha River Canoe #4	8MR3167	Prehistoric	Underwater site	NE	NE	TBD
Sunday Bluff Canoe	8MR3169	Prehistoric	Archaeological Site	NE	NE	TBD
Newman Point	8MR3177	indeterminate	Archaeological Site	NE	NE	TBD
Abandoned Railroad Grade	8MR3270	Historic	Linear Resource	NE	NE	TBD
Voorhees Bluff	8MR3351	Prehistoric	Archaeological Site	NS	NE	TBD
Lithic Scatter 2, Piney Island South	8MR3358	Prehistoric, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Lithic Scatter 3, Piney Island South	8MR3359	Prehistoric, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Lithic Scatter 4, Piney Island South	8MR3360	Prehistoric, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Eaton Creek Island Pilings	8MR3361	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Eaton Creek Railroad Spike	8MR3362	19 <sup>th</sup> and 30 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
McCarthy's Midden	8MR3363	Prehistoric, 19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Eaton Creek Lithic Scatter #2	8MR3364	Prehistoric, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Homesteader's Site, Eaton Creek Road	8MR3365	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Mason Bay West Bridge	8MR3366	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD

Site Name	Site Number	Culture/Period	Description	Significance	Condition	Treatment
Eaton Creek Bridge	8MR3367	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Tuten Creek Borrow Pits	8MR3368	Prehistoric, Lake Archaic, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Kelly and DJ's Camp	8MR3369	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Cedar Creek North Midden	8MR3370	Prehistoric, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Conner Homestead	8MR3372	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Charley Perry's Mound 1	8MR3373	Prehistoric 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Charley Perry's Mound 2	8MR3374	Prehistoric, Late Archaic, Archaic, St. Johns, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Charley Perry's Village	8MR3375	Prehistoric, St. Johns, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Charley Perry's Midden	8MR3376	Prehistoric, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Dek's Bluff Midden	8MR3377	Prehistoric, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Tuten Creek Midden	8MR3378	Prehistoric, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Cedar Creek East Midden	8MR3379	Prehistoric, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Eaton Creek Lithic Scatter #3	8MR3381	Prehistoric, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Eaton Creek Lithic Scatter 31	8MR3382	Prehistoric, 20 <sup>th</sup> Century	Archaeological Site	NE	NE	TBD
Cross Florida Greenway	8MR3410	American, Depression/ New Deal, 19 <sup>th</sup> and 20 <sup>th</sup> Century	Linear Resource	NR	NE	TBD
Cedar Creek Bell Site	8MR3446	First Spanish Period	Archaeological Site	NE	NE	TBD
SR 40/CR 326 Site	8MR3477	Prehistoric	Archaeological Site	NS	NE	TBD
Quonset Hut #1	8MR3521	Circa 1940	Historic Structure	NS	NE	TBD
Quonset Hut #2	8MR3522	Circa 1940	Historic Structure	NS	NE	TBD

Site Name	Site Number	Culture/Period	Description	Significance	Condition	Treatment
Rat House	8MR3523	Circa 1950	Historic Structure	NS	NE	TBD
Lung Building	8MR3524	Circa 1950	Historic Structure	NS	NE	TBD
Tractor Shed	8MR3525	Circa 1950	Historic Structure	NS	NE	TBD
MR3526	8MR3526	Archaic, Orange, Woodland, St. Johns, St. Johns, 20 <sup>th</sup> Century	Archaeological Site	NR	NE	TBD
MR3527	8MR3527	Prehistoric	Archaeological Site	NE	NE	TBD
MR3529	8MR3529	Prehistoric	Archaeological Site	NE	NE	TBD
MR3530	8MR3530	Prehistoric	Archaeological Site	NE	NE	TBD
MR3531	8MR3531	Prehistoric	Archaeological Site	NE	NE	TBD
Eureka Lock and Dam Complex	8MR3563	20 <sup>th</sup> Century	Resource Group	NR	NE	TBD
Eureka Lock	8MR3564	20 <sup>th</sup> Century	Historic Structure	NE	NE	TBD
Eureka Lock Building 1	8MR3565	20 <sup>th</sup> Century	Historic Structure	NE	NE	TBD
Eureka Lock Building 2	8MR3566	20 <sup>th</sup> Century	Historic Structure	NE	NE	TBD
Eureka Lock Building 3	8MR3567	20 <sup>th</sup> Century	Historic Structure	NE	NE	TBD
Eureka Lock Building 4	8MR3568	20 <sup>th</sup> Century	Historic Structure	NE	NE	TBD
Horse Park Historic Scatter	8MR3569	20 <sup>th</sup> Century	Archaeological Site	NS	NE	TBD
Horse Park Lithic Scatter	8MR3570	Prehistoric	Archaeological Site	NS	NE	TBD
CR-316 / Proposed Cross FL Canal Bridge	8MR3585	1969	Historic Bridge	NR	NE	TBD
Eureka Dam Dugout	8MR3675	Prehistoric or Historic	Archaeological Site	NE	NE	TBD
Original Little Chapel UMC site	8MR3679	Prehistoric, 19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NS	NE	TBD
FPC-R1C Two	8MR3748	Prehistoric	Archaeological Site	NS	NE	TBD

Site Name	Site Number	Culture/Period	Description	Significance	Condition	Treatment
No Name	8MR3750	Historic	Historic Bridge	NE	NE	TBD
Silver Springs Head Springs Site Complex	8MR3762	Archaic, Historic	Resource Group	NE	NE	TBD
Shed	8MR3791	Historic, 1964	Historic Structure	NS	NE	TBD
CR 316/Proposed Cross FL Canal Bridge	8MR3858	Historic	Bridge	NR	NE	TBD
FL-817	8MR3863	Prehistoric, 19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NR	NE	TBD
FL-817B	8MR3865	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NR	NE	TBD
FL-817C	8MR3866	Prehistoric, 19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NR	NE	TBD
Greenway Trail 1	8MR3921	Prehistoric, 19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site	NS	NE	TBD
Greenway Trail 2	8MR3922	Prehistoric	Archaeological Site	NS	NE	TBD
Fisher Cemetery	8MR3923	Started 1940 (abandoned)	Historic Cemetery	NS	NE	TBD
Heidtville	8MR3925	19 <sup>th</sup> and 20 <sup>th</sup> Century	Archaeological Site/Historic Town	INSF	NE	TBD
<b>Putnam County Sites</b>						
½ Mile North of Horse Landing	8PU0026	Orange, St. Johns	Archaeological Site	NE	NE	TBD
No Name	8PU0052	Prehistoric	Archaeological Site	NE	NE	TBD
Mound A, Ditch Creek	8PU0053	St. Johns	Archaeological Site	NE	NE	TBD
Cedar Hammock Midden	8PU0058	Prehistoric	Archaeological Site	NE	NE	TBD
No Name	8PU0078	Archaic	Archaeological Site	NE	NE	TBD
No Name	8PU0078	St. Johns	Archaeological Site	NE	NE	TBD
EH & A Putnam 28	8PU0113	Prehistoric	Archaeological Site	NE	NE	TBD
Structure 98	8PU0118	St. Johns	Archaeological Site	NE	NE	TBD



Site Name	Site Number	Culture/Period	Description	Significance	Condition	Treatment
No Name	8PU0674	Prehistoric	Archaeological Site	NE	NE	TBD
Abandoned Seaboard Coastline	8PU0800	19 <sup>th</sup> and 20 <sup>th</sup> Century	Linear Resource	NS	NE	TBD
PU1568	8PU1568	St. Johns	Archaeological Site	NS	NE	TBD
PU1569	8PU1569	Prehistoric	Archaeological Site	NE	NE	TBD
Ocala Northern Railroad	8PU1633	20 <sup>th</sup> Century	Linear Resource	NE	NE	TBD
<p><b>Key:</b></p> <p><b>Significance:</b> NRL = National Register listed; NR = National Register eligible; NE = not evaluated; NS = not significant; INSF = insufficient information</p> <p><b>Condition:</b> G = Good; F = Fair; P = Poor; NA = Not accessible; NE = Not evaluated</p> <p><b>Treatment:</b> RS = Restoration; RH = Rehabilitation; ST = Stabilization; P = Preservation; R = Removal; TBD = To Be Determined; N/A = Not applicable</p>						

1

# 1 RESOURCE MANAGEMENT PROGRAM

## 2 Management Goals, Objectives, and Actions

3 Measurable objectives and actions have been identified for each of DRP's management goals for the  
4 CFG. Please refer to the Implementation Schedule and Cost Estimates in the Implementation  
5 Component of this plan for a consolidated spreadsheet of the recommended actions, measures of  
6 progress, target year for completion, and estimated costs to fulfill the management goals and  
7 objectives of this park.

8 While DRP utilizes the 10-year management plan to serve as the basic statement of policy and future  
9 direction for each park, a number of annual work plans provide more specific guidance for DRP staff  
10 to accomplish many of the resource management goals and objectives of the park. Where such  
11 detailed planning is appropriate to the character and scale of the park's natural resources, annual  
12 work plans are developed for prescribed fire management, exotic plant management, and imperiled  
13 species management. Annual or longer-term work plans are developed for natural community  
14 restoration and hydrological restoration. The work plans provide DRP with crucial flexibility in its  
15 efforts to generate and implement adaptive resource management practices in the state park system.

16 The work plans are reviewed and updated annually. Through this process, DRP's resource  
17 management strategies are systematically evaluated to determine their effectiveness. The process  
18 and the information collected is used to refine techniques, methodologies, and strategies, and ensures  
19 that each park's prescribed management actions are monitored and reported as required by Chapters  
20 253.034 and 259.037, F.S.

21 The goals, objectives, and actions identified in this management plan will serve as the basis for  
22 developing annual work plans for the park. The 10-year management plan is based on conditions  
23 that exist at the time the plan is developed, and the annual work plans provide the flexibility needed  
24 to adapt to future conditions as they change during the 10-year management planning cycle. As the  
25 park's annual work plans are implemented through the 10-year cycle, it may become necessary to  
26 adjust the management plan's priority schedules and cost estimates to reflect these changing  
27 conditions.

## 28 Natural Resource Management

### 29 Hydrological Management

30 **Goal: Protect water quality and quantity in the park, restore hydrology to the greatest extent**  
31 **feasible, and maintain the restored condition.**

32 The natural hydrology of most state parks has been impaired prior to acquisition to one degree or  
33 another. Florida's native habitats are precisely adapted to natural drainage patterns and seasonal  
34 water level fluctuations. Variations in these factors frequently determine the types of natural  
35 communities that occur on a particular site. Even minor changes to natural hydrology can result in  
36 the loss of plant and animal species from a landscape. Restoring state park lands to original, natural  
37 conditions often depends on returning natural hydrological processes and conditions to the park.  
38 This is done primarily by filling or plugging ditches, removing obstructions to surface water "sheet

1 flow,” installing culverts or low-water crossings on roads, and installing water control structures to  
2 manage water levels.

3 *Objective A: Conduct periodic inspections, repairs and maintenance at former Cross Florida Barge Canal*  
4 *Water Control Structures per Federal Emergency Management Agency guidelines and professional*  
5 *engineer recommendations.*

6 *Objective B: Repair hydrological conditions and function to approximately 16,700 acres – Etoniah,*  
7 *Marshall Swamp and Gore’s Landing.*

## 8 **Natural Communities Management**

9 **Goal: Restore and maintain the natural communities/ habitats of the park.**

10 As discussed above, DRP practices natural systems management. It is most important to restore and  
11 maintain native plant communities, to the greatest extent practical. In most cases, this entails  
12 returning fire to its natural role in fire-dependent natural communities. Other methods to implement  
13 this goal include large-scale restoration projects, as well as smaller-scale improvements to natural  
14 communities. Following are the natural community management objectives and actions  
15 recommended for the state park.

16 To ensure restoration priorities on the CFG, the evaluation should be completed of the overall quality  
17 of vegetative communities throughout the greenway for potential restoration. Assessments were  
18 based on field evaluations and ranged from poor, fair, good to excellent. The field evaluations also  
19 provided prescriptions for future management actions. CFG should coordinate with surrounding  
20 landholders, such as the FWC, to assist with both the assessment and identification of restoration  
21 priorities on the greenway. Additionally, CFG should work with universities, federal agencies, and  
22 non-governmental organizations to gather basic data on natural resources and to help develop  
23 planning and evaluation tools.

### 24 *Prescribed Fire Management*

25 Prescribed fire is used to mimic natural lightning-set fires, which are one of the primary natural  
26 forces that shaped Florida’s ecosystem. Prescribed burning increases the abundance and health of  
27 many wildlife species. Many of Florida’s imperiled species of plants and animals are dependent on  
28 periodic fire for their continued existence. Fire-dependent natural communities gradually  
29 accumulate flammable vegetation; therefore, prescribed fire reduces wildfire hazards by reducing  
30 these wild land fuels.

31 All prescribed burns in the Florida state park system are conducted with authorization from the FFS.  
32 Wildfire suppression activities in the park are coordinated with the FFS.

33 *Objective A: Within 10 years have 25,865 acres of the park maintained within optimal fire return*  
34 *interval.*

35 *Action 1: Develop/update annual burn plan.*

36 *Action 2: Manage fire dependent communities for ecosystem function, structure and process by*  
37 *burning between 7,500 – 8,000 acres annually, as identified by the annual burn plan.*

1 Funding will be required to secure contract burning assistance to meet this goal of burning 7,500  
 2 acres to 8,000 acres annually.

3 *Action 3: Establish and maintain 500 miles of fire breaks.*

4 Table 14 contains a list of all fire-dependent communities, including ruderal fire type communities  
 5 located within the park, their associated acreage and optimal fire return interval, and the annual  
 6 average target for acres to be burned.

7 The CFG is partitioned into management zones including those designated as burn zones (see  
 8 Management Zones Table and Map). Prescribed fire is planned for each burn zone on the appropriate  
 9 interval. The park's burn plan is updated annually because fire management is a dynamic process. To  
 10 provide adaptive responses to changing conditions, fire management requires careful planning based  
 11 on annual and very specific burn objectives. Each annual burn plan is developed to support and  
 12 implement the broader objectives and actions outlined in this ten-year management plan.

13 **Table 14. Prescribed Fire Management**

Natural Community	Acres	Optimal Fire Return Interval (Years)	*Average Number of Acres Needed to be Burned per Year
<b>Fire-Type Acres</b>			
Basin Marsh	14	2-10 years (5)	3
Floodplain Marsh	245	2-5 years	82
Mesic Flatwoods	5,041	1-4 years (3)	1,680
Sandhill	6,408	1-3 years (3)	2,136
Scrub	1,630	5-20 years (5)	326
Scrubby Flatwoods	683	3-14 years (10)	68
Upland Pine Forest	6	1-3 years (3)	2
Wet Flatwoods	1,773	5-10 years (7)	253
Subtotal Fire Type Acres	15,800		4,550
<b>Ruderal Fire Type Acres</b>			
Clear Cut Pine Plantations	161	3 years	54
Pine Plantation	9,071	3 years	3,024
Subtotal Ruderal Fire Type Acres	9,232		3,078
Total Fire Type	25,032		7,628
* Average Number of Acres Needed to be Burned per Year is based on the fire return interval assigned to each burn zone. Each burn zone may include multiple natural communities.			

14

15 Fire has historically been a significant force in shaping the natural Florida landscape. The fire  
 16 management program on the CFG is intended to restore the natural process of fire to the landscape.

1 Upland communities normally are burned in the lightning season during the late spring and summer.  
 2 However, natural lightning-caused ignitions may occur in any month of the year. In some cases, areas  
 3 will be burned during the winter season to reduce fuel loads before switching to lightning season  
 4 burning. Fuel loads, restoration goals, and natural community type will be considered when  
 5 scheduling prescribed fires.

6 Table 15 contains a history of acres burned across the greenway and a projection for the year 2016-  
 7 2017.

8 **Table 15. CFG Prescribed Fire History**

FY	Acres
1998-99	1,540
1999-00	1474
2000-01	0
2001-02	83
2002-03	2,438 (1,002 contracted + 1,436 in-house)
2003-04	1,297
2004-05	1,208
2005-06	1,763
2006-07	1,559
2007-08	1,638
2008-09	1,152
2009-10	2,639 (584 contracted + 2,055 in-house)
2010-11	1,368
2011-12	0
2012-13	981
2013-14	2,500 (867 contracted + 1,633 in-house)
2014-15	1,640 (651 contracted + 989 in-house)
2015-16	4,059 (3,574 contracted + 485 in-house)
2016-17	6,761 (5,161 contracted + 1,600 in-house)

9

10 To track fire management activities, the DRP maintains a statewide burn database. The database  
 11 allows staff to track various aspects of each park's fire management program, including individual  
 12 burn zone histories and fire return intervals, staff training/experience, backlog, if burn objectives  
 13 have been met, etc. The database also is used for annual burn planning, which allows DRP to  
 14 document fire management goals and objectives on an annual basis. Each quarter, the database is  
 15 updated and reports are produced that track progress toward meeting annual burn objectives.

16 *Natural Communities Restoration*

17 In some cases, the reintroduction and maintenance of natural processes is not enough to reach the  
 18 natural community desired future conditions in the park, and active restoration programs are  
 19 required. Restoration of altered natural communities to healthy, fully functioning, natural landscapes  
 20 often requires substantial efforts that may include mechanical treatment of vegetation or soils and



1 reintroduction or augmentation of native plants and animals. For the purposes of this management  
2 plan, restoration is defined as the process of assisting the recovery and natural functioning of  
3 degraded natural communities to desired future condition, including the re-establishment of  
4 biodiversity, ecological processes, vegetation structure, and physical characters.

5 Examples that would qualify as natural communities' restoration, requiring annual restoration plans,  
6 include large mitigation projects, large-scale hardwood removal and timbering activities, and roller-  
7 chopping and other large-scale vegetative modifications. The key concept is that restoration projects  
8 will go beyond management activities routinely done as standard operating procedures, such as  
9 routine mowing, the reintroduction of fire as a natural process, spot treatments of exotic plants, and  
10 small-scale vegetation management.

#### 11 *Natural Communities Improvement*

12 Improvements are similar to restoration but on a smaller, less-intense scale. Improvements typically  
13 include small-scale vegetative management activities or minor habitat manipulation. Following are  
14 the natural community/habitat improvement actions recommended at the park.

15 *Objective: Conduct habitat/natural community restoration activities on 2,555 acres of ruderal*  
16 *communities.*

17 *Action 1: Plant 50-100 acres of wiregrass annually in 2,000 acres of old pastures planted in*  
18 *longleaf pine.*

19 *Action 2: Replant 555 acres of slash pine in the Etoniah properties.*

#### 20 **Imperiled Species Management**

21 **Goal: Maintain, improve, or restore imperiled species populations and habitats in the park.**

22 DRP strives to maintain and restore viable populations of imperiled plant and animal species  
23 primarily by implementing effective management of natural systems. Single-species management is  
24 appropriate in state parks when the maintenance, recovery, or restoration of a species or population  
25 is complicated due to constraints associated with long-term restoration efforts, unnaturally high  
26 mortality, or insufficient habitat. Single-species management should be compatible with the  
27 maintenance and restoration of natural processes, and should not imperil other native species or  
28 seriously compromise park values.

29 In the preparation of this management plan, DRP staff consulted with staff of the FWC's Imperiled  
30 Species Management or that agency's Regional Biologist and other appropriate federal, state, and  
31 local agencies for assistance in developing imperiled animal species management objectives and  
32 actions. Data collected by the USFWS, FWC, and FNAI as part of their ongoing research and  
33 monitoring programs will be reviewed by park staff periodically to inform management of decisions  
34 that may have an impact on imperiled species at the park.

35 Ongoing inventory and monitoring of imperiled species in the state park system is necessary to meet  
36 DRP's mission. Long-term monitoring also is essential to ensure the effectiveness of resource  
37 management programs. Monitoring efforts must be prioritized so that the data collected provide  
38 information that can be used to improve or confirm the effectiveness of management actions on

1 conservation priorities. Monitoring intensity must at least be at a level that provides the minimum  
2 data needed to make informed decisions to meet conservation goals. Not all imperiled species require  
3 intensive monitoring efforts on a regular interval. Priority must be given to those species that can  
4 provide valuable data to guide adaptive management practices. Those species selected for specific  
5 management action and those that will provide management guidance through regular monitoring  
6 are addressed in the objectives below.

7 Given the significant effort to restore scrub habitat, efforts should include monitoring and  
8 documenting of imperiled species that often inhabit scrub communities. Also, efforts should be  
9 focused on coordination with the appropriate federal, state, and local agencies to monitor, document,  
10 and research bird species on the Spoil Islands in the western part of the CFG.

11 *Objective A: Update baseline imperiled species occurrence inventory lists for plants and animals with*  
12 *FNAI.*

13 Depending on funding, a full plant survey needs to be conducted at the park to determine presence  
14 and location of other listed plant species. In addition, the park has never been fully surveyed for  
15 herptofauna, insects, bats, or birds. If funding is available, surveys for these species should be  
16 conducted and the species list updated. Surveys for these species are particularly important around  
17 wet community types, such as depression marsh, basin swamp, or dome swamp, that are critical for  
18 breeding amphibians, but where arthropod control may occur to limit the presence of mosquitoes.  
19 District biologists in partnership with FWC may survey for herptofauna. The park will work with  
20 district biologists to conduct limited surveys, update the imperiled species lists, and utilize  
21 observations to update the arthropod control plan to minimize the impacts of spraying to potentially  
22 sensitive species.

23 *Objective B: Monitor and document 3 selected imperiled animal species in the park.*

24 *Objective C: Monitor and document 1 selected imperiled plant species in the park.*

25 *Objective D: Maintain/improve scrub jay habitat on the CFG.*

26 *Action 1: Apply mechanical treatments to 10 percent (100 acres) of our scrub jay habitat*  
27 *annually.*

28 *Action 1: Band scrub jay population for three to five days annually.*

## 29 **Exotic Species Management**

30 **Goal: Remove invasive exotic plants and animals from the park and conduct needed**  
31 **maintenance control.**

32 DRP actively removes invasive exotic plant species from state parks, with priority being given to  
33 those causing ecological damage. Removal techniques may include mechanical treatment, herbicides,  
34 or biocontrol agents. Over the past few years, CFG has implemented an extremely aggressive program  
35 to remove invasive exotic plants from the park. This program has involved treatment of the entire  
36 greenway with biannual treating of at least two large management zones. As a result of this  
37 aggressive treatment program, the greenway is in maintenance for invasive exotic plant species.

1 Maintenance condition describes a formerly active infestation that has been treated to the extent that  
2 any plants remaining are manageable with existing staff and resources, total area is stable or  
3 declining, mature reproducing individuals are absent, and the species poses no significant threat to  
4 listed plants or animals.

5 *Objective A: Annually treat approximately 1,716 infested acres of exotic upland plant species in the park.*

6 *Action 1: Annually develop/update exotic plant management work plan. Action 2: Implement*  
7 *annual work plan by treating 1,716 acres in park annually and continuing maintenance and*  
8 *follow-up treatments, as needed.*

9 Therefore, the focus for invasive exotic species management on the CFG should move to individual  
10 species management. The four most persistent species are cogon grass, Caesar’s weed, mimosa, and  
11 camphor tree. In addition, specific efforts should be focused on skunk vine, which is becoming more  
12 pervasive in the western portion of the CFG (Large Management Zone 1). Also, efforts should be given  
13 to managing natal grass, which is infringing on the scrub areas throughout the greenway. Natal grass  
14 is in the early infestation stage and will be easier to develop an eradication program.

15 *Objective B: Implement control measures on feral hogs in the park.*

## 16 **Special Management Considerations**

### 17 **Timber Management Analysis**

18 On all parcels larger than 1,000 acres, if the DRP determines that timber management does not  
19 conflict with the primary management objectives of the land, Florida Statutes Chapters 253 and 259  
20 require:

- 1) An analysis of the multiple-use potential of the parcel. Such analysis shall include the potential of the parcel to generate revenues to enhance the management of the parcel.
- 2) An assessment of the feasibility of managing timber resources for conservation and revenue generation purposes through a stewardship ethic that embraces sustainable forest management practices in land management plans.

21 The CFG spans 110 miles from Yankeetown on Florida’s west coast to south of Palatka on the St. Johns  
22 River, near the east coast of Florida. The CFG ranges from 300 yards wide to one mile wide and  
23 includes portions of four counties in the upper Florida peninsula: Citrus, Levy, Marion, and Putnam.  
24 Over the 110 miles, the CFG traverses numerous natural, physiographic, and developed areas. The  
25 CFG is designated as a multiple-use park. The feasibility of harvesting timber on the CFG during the  
26 period covered by this UMP was considered pursuant to the DRP statutory responsibilities to analyze  
27 the park’s resource needs and values.

28 The long-term management goal for forest communities in the state park system is to maintain or re-  
29 establish old-growth characteristics to the greatest degree practicable, except in those forest  
30 communities specifically managed as early successional. Timber management is utilized for the  
31 specific purpose of helping restore or improve current habitat conditions and enhance the overall  
32 integrity of the natural community. Revenue generation from timber management is not the goal, but  
33 rather a by-product of taking such actions to help restore/improve target conditions of specific  
34 natural communities. In all situations, forest/stand/timber management activities undertaken will

1 adhere to the current Florida Silvicultural Best Management Practices and Florida Forestry Wildlife  
2 Best Management Practices for State Imperiled Species.

3 Many of the natural communities evaluated on the CFG had overstory stocking levels at, or above, the  
4 upper limits for corresponding FNAI Reference Sites. A subset of these stands has overstocked  
5 conditions in the preferred pine component, while the remainder has overstocked conditions in the  
6 non-preferred pine or hardwood components. This overstocked condition makes overstory thinning  
7 a potential management tool that should be considered. Activities related to stand improvement,  
8 including palmetto and midstory reduction, are needed in many areas.

9 The Timber Management Analysis found in Addendum 4 provides additional details. This analysis  
10 has been evaluated and found to be consistent with the recommendations found in the subject RMC.

## 11 **Additional Considerations**

### 12 *Spoil Island Management*

13 Nesting shore birds have been identified on Spoil Islands 4 through 7. These include the  
14 oystercatcher, which is listed as a species of high conservation concern in the U.S. Shorebird  
15 Conservation Plan (Brown et al., 2001). There are approximately 1,500 nesting pairs along the  
16 Atlantic and Gulf Coasts of the U.S. (Brown et al., 2005) and the species appears to be declining in the  
17 southeastern portion of the range (Davis et al., 2001). In Florida, a statewide survey of Oystercatcher  
18 nesting conducted in 2001 documented a total of 391 probable breeding pairs (Douglass and Clayton,  
19 2004). A 2010 survey estimated 170 breeding pairs (Brush, 2010), documenting a 56-percent loss.  
20 Due to the current population size and trend, Oystercatchers have been proposed to be state listed  
21 as threatened (FWC, 2011). While recent research has been focused on winter habitat limitations and  
22 restoration (Brush et al., 2015), little research has been devoted to understanding or improving  
23 breeding productivity and habitat for the Oystercatcher population breeding along the Nature Coast.

24 Within Florida, Oystercatchers are breeding at about 50 areas (managed land boundaries, DOD  
25 properties, etc.), with greater than 90 percent of the population concentrated on the Gulf Coast.  
26 Nesting habitat within the Nature Coast is limited to a few small offshore islands around Cedar Key,  
27 the Horseshoe Beach jetties, and on Spoil Islands in Citrus County. The CFG Spoil Islands in Citrus  
28 County support the largest concentration of nesting Oystercatchers along the Nature Coast Region  
29 and the fourth largest concentration in the state. FWC has been intensively monitoring nesting  
30 Oystercatchers on these islands since 2012, and nesting has been documented on all the Spoil Islands.  
31 Reproductive effort is high as the number of nesting pairs at the site has increased from 14 to 22, and  
32 most pairs that fail early in the season re-nest. However, annual site productivity is low (average of  
33 0.05 chicks per pair) compared to annual statewide productivity during the same time period  
34 (average greater than 0.20 chicks per pair).

35 Predation of eggs and young has been shown to be the greatest limitation in breeding success of  
36 Oystercatchers (Nol, 1989; Morse et al., 2006; Tessler et al., 2007). Semiprecocial young are  
37 particularly vulnerable to predation, starvation, and weather events within two weeks of hatching  
38 (Colwell et al., 2007; American Oystercatcher Working Group et al., 2012; Schulte, 2012). A high  
39 reproductive effort in an area with low success warrants further investigation to determine what is  
40 limiting the reproductive performance of this breeding population of Oystercatchers. This  
41 information is critical to the management of these sites as Oystercatcher breeding areas.

1 In addition, the Spoil Islands on the CFG are a popular destination for boating and kayaking  
2 recreationists. Most of the habitat near the CFG is intertidal oyster rakes that lack beach areas and  
3 shade. The Spoil Islands offer inviting areas for boaters and kayakers to land and recreate for long  
4 periods of time. During the summer months, human presence on the islands is highest, coinciding  
5 with the Oystercatcher breeding season; all of the islands have been documented as Oystercatcher  
6 nesting sites. While some of the islands are managed to minimize disturbance using FWC signs, boats  
7 are still able to land on the islands and often remain on the island. Oystercatchers on the CFG are  
8 actively incubating or caring for young in a habitat that already has a large number of avian predators  
9 and affords little protection from human disturbances. Additionally, unique to the CFG islands is the  
10 fact that Oystercatchers and their young tend to stay close to the nest location rather than roving or  
11 foraging at adjacent areas at low tide, which could lead to increased vulnerability if disturbed.

12 A study is currently underway to assist with determining the effects of disturbance from humans and  
13 predators on Oystercatcher productivity on the CFG Spoil Islands. Information from this study will  
14 better focus future conservation efforts in the area. The objectives of this study are to use existing  
15 data and collect additional field data (2016-2018) to: (1) determine causes of mortality and  
16 disturbance (using existing data as well as conducting direct observations and use of remote  
17 cameras); (2) estimate annual productivity (2012-2018) with respect to covariates, such as potential  
18 sources of disturbance, presence of predators, and food supply; (3) propose management actions to  
19 increase annual productivity of Oystercatchers nesting on Spoil Islands on the CFG.

#### 20 *Canal Landfill Site Restoration*

21 The canal landfill site or “Pedro Landfill” is located at the junction of SR 475B and SR 475 (see Base  
22 Map). This landfill was operated as a Class I landfill from the mid-1960s until 1984 and received solid  
23 waste generated from surrounding unincorporated areas. During this period, local residents and  
24 commercial garbage haulers brought garbage that was disposed of in trenches that were 12 feet to  
25 15 feet deep and 30 feet wide. Initially, this site was operated as a landfill for local residents. After  
26 1974, this landfill had a full-time County staff that operated the landfill continuously until operations  
27 ceased in 1984. Although it stated it was closed in 1984, based on information received from the  
28 Marion County SWD, it is believed that the site continued to be used illegally by residents and  
29 commercial vendors for solid waste disposal from 1984 until 1989. The site then was converted into  
30 the Old Canal Greenbox Recycling Center in approximately 1989, and continued to be used for this  
31 purpose until approximately 2007.

32 A closure plan was submitted to the Florida Department of Regulation (DER) in 1985; however, this  
33 plan has yet to be implemented. Low levels of synthetic organic compounds have been found in  
34 monitoring wells on the site. Site recommendations include continuing the monitoring program. CFG  
35 staff would like to reclaim or restore the site to render it safe to be used by wildlife and the public.

#### 36 *Lower Ocklawaha River Shoreline Management*

37 The CFG needs shoreline and water management plans for the Lower Ocklawaha River and Rodman  
38 Reservoir. The CFG possesses and manages 60+ miles of littoral shoreline ownership along both these  
39 waterbodies in the eastern half of the CFG. Both of these areas are bordered by private ownership up  
40 the hill from the water’s edge. Adjacent private landowners have established and maintained  
41 numerous encroachments on state-owned lands in these areas, including, but not limited to:  
42 vegetation/under brushing/ mowing and even large tree removal in numerous instances. There also



1 are more than 100 illegal docks placed on these state-owned lands without appropriate permits,  
2 based on an inventory of these docks by the FDEP Division of State Lands in 2007. Due to the  
3 politically charged nature of the Rodman/ Ocklawaha issue, staff need clear and legal guidance in  
4 dealing with these issues and protecting the state’s interest and the environment.

5 The Rodman Reservoir is an approximately 9,000-acre man-made impoundment of the Ocklawaha  
6 River that was constructed as part of the former Cross Florida Barge Canal Project. Due to the 48+  
7 year-old effort by environmental groups to have the dam and reservoir removed and the Ocklawaha  
8 River restored, there has never been a formal water management plan developed for this reservoir.  
9 DRP/CFG management should have and has recurrently asked for the development and  
10 implementation of such a plan for the protection of personal property and life downstream from the  
11 Kirkpatrick Dam along the St. Johns River. Approximately 400+ properties were shown to be in  
12 potential harm’s way if the Kirkpatrick Dam failed and the impounded water in the reservoir flowed  
13 downstream in an uncontrolled discharge.

#### 14 *Impoundments and Water Control Structures*

15 CFBC construction/development resulted in the construction of towering bridges, locks, and dams.  
16 Remnants of the sea-level 1930s ship canal include the deep canal digs and the monolithic concrete  
17 bridge stanchions located in the median of US 441. Three large lock structures, H.H. Buckman Lock,  
18 Eureka Lock and Dam, and Inglis Lock and Dam, are the most prominent. Although still in place, the  
19 Eureka Lock and Dam have never been functional.

20 CFG staff maintain and operate the Buckman Lock and Kirkpatrick Dam and Spillway. The Buckman  
21 Lock controls access to Rodman Reservoir from the St. Johns River through the east barge canal. The  
22 Kirkpatrick Dam spillway controls the level of Rodman Reservoir—a 9,000-acre reservoir. In general,  
23 the water level of the reservoir is maintained at the 18 feet to 20 feet national geodetic vertical datum  
24 (NGVD) level. The water level is drawn down about every three years to about 11 feet to consolidate  
25 bottom sediments, enhance the fishery and wildlife habitats, and assist in control of aquatic plants.

26 Since the 1970s, numerous groups have urged the removal of the Kirkpatrick (formerly Rodman)  
27 Dam and restoration of Rodman Reservoir to the Ocklawaha River floodplain because of the impact  
28 of the reservoir on the Ocklawaha River floodplain and associated ecosystems. There is resistance to  
29 this from other groups, such as sports-fishing related organizations and businesses. The Governor  
30 and Cabinet, sitting as the Board of Trustees of the Internal Improvement Trust Fund, have an  
31 established policy that the Ocklawaha River should be “partially restored” (see explanation below),  
32 with the FDEP as the lead agency. However, the Legislature has not appropriated funds for this  
33 purpose. If funds are made available and permits are issued, it is the intent of the FDEP to undertake  
34 this restoration. The SJRWMD is investigating the potential impact on the lower St. Johns River from  
35 restoration of the Ocklawaha River.

36 Kirkpatrick Dam caused the flooding of a portion of Ocala National Forest lands. The flooding and  
37 occupying of these lands was allowed under a special permit from the U.S. Department of Agriculture,  
38 Forest Service, first issued in 1994. This special permit expired in 2002. FDEP applied for a new  
39 permit, but the Forest Service included conditions about the issuance of the new permit, including a  
40 schedule for the reservoir to be drawn down and the dam to be breached. The Secretary of the FDEP  
41 did not sign the Forest Service permit because the permit(s) from SJRWMD necessary to draw down  
42 the reservoir had not been approved.

1 “Partial restoration,” the restoration alternative selected by the Governor and Cabinet in 1995, is  
2 intended to restore river hydrology and floodplain function to near preconstruction conditions  
3 through breaching of the dam, with limited removal and/or alteration of structures and alteration of  
4 topography. This alternative will retrieve National Forest System lands at the lowest cost while  
5 restoring river and floodplain hydrology. The major components of partial restoration are:

- 6 1. Drawdown of the reservoir to be accomplished in three phases (three years to drop from 18  
7 feet to four feet NGVD)
- 8 2. Limited construction of channel stabilization and erosion control structures in the Ocklawaha  
9 River
- 10 3. Limited planting of native plant species to provide for erosion control
- 11 4. Partial leveling of the exposed barge canal side-cast spoil berms
- 12 5. Restoration of the historic Ocklawaha River channel flow by filling the barge canal where it  
13 intersects the river channel
- 14 6. Restoration of the historic Deep Creek channel flow by filling the barge canal where it  
15 intersects the creek channel
- 16 7. Restoration of the historic Camp Branch floodplain and channel flow by filling the barge canal  
17 where it intersects the creek channel
- 18 8. Closure and securing of the Buckman Lock
- 19 9. Removal of 2,000 feet of the Kirkpatrick Dam (earthen portion)
- 20 10. Partial filling and restoration of the spillway tailrace to natural grade
- 21 11. Development and implementation of a cultural resources operating plan

22 The projected cost for repairs and to bring the Buckman Lock, Kirkpatrick Dam and Spillway, and the  
23 Eureka Lock and Dam up to latest standards for operation is \$4 million. This estimate is based on the  
24 July 2015 inspection report and years of historical data from the Inglis Lock and Dam repair projects  
25 and assessment reports. The estimated cost for restoration of the Ocklawaha River is \$25.8 million.  
26 The yearly operating costs for the dam components and activities and the operating costs for the area  
27 after restoration are approximately the same: \$198,000 and \$234,000, respectively. Snagging for  
28 navigation maintenance is expected to increase after river restoration, contributing to almost half of  
29 the annual operating cost after restoration.

30 The Withlacoochee River is the main river system on the western end of the CFG. A small portion of  
31 the CFG fronts the Withlacoochee River near Dunnellon, where the Rainbow River, arising from a  
32 first-magnitude spring, flows into the river. Downstream of the confluence with the Rainbow River,  
33 the Withlacoochee River becomes Lake Rousseau. The Withlacoochee River was dammed in the early  
34 1900s to generate electric power, creating Lake Rousseau. The reservoir is no longer used for power  
35 generation. Below Lake Rousseau, the CFG forms the south bank of the Withlacoochee in several  
36 places as it flows to the Gulf.

37 Current water control structures include the Inglis Dam and Spillway at the western end of Lake  
38 Rousseau, the Inglis Lock as part of the western barge canal, and the Inglis Bypass Canal and Spillway.  
39 The bypass canal funnels water from Lake Rousseau just east of the lock to a spillway that provides  
40 water to the lower reaches of the Withlacoochee River. The Inglis Lock is no longer operational due  
41 to its deteriorated condition. The Governor and Legislature will decide whether the lock will be made  
42 operational or permanently closed. Permanent closure may require deauthorization by Congress.  
43 The SWFWMD operates the western barge canal water control dams and spillways under a contract  
44 with and funding through DEP; DEP is still currently responsible for the lock. The Lake Rousseau

1 water level generally is at a fixed elevation of 27.5 NGVD. In times of heavy rainfall, additional water  
2 can be released to prevent or minimize flooding around Lake Rousseau.

3 The western barge canal is about nine miles long. It extends from near the western end of Lake  
4 Rousseau into the Gulf, where it extends for approximately 10.5 miles. It cuts through the lower  
5 reaches of the Withlacoochee River between the Inglis Bypass Spillway and the western end of Lake  
6 Rousseau. The coastal wetlands and wet flatwoods were disrupted by construction of the canal. Inglis  
7 Island, formerly land bordering the north side of Lake Rousseau and the Withlacoochee River, was  
8 surrounded by water by the canal being cut through on the north side of the land mass.

9 Discussions will continue regarding the impoundments and water control structures, ownership, and  
10 future management.

### 11 *Highway Corridor Management*

12 The CFG is surrounded and split by several miles of roads and highways ranging from two-lane  
13 state roads to a six-lane high-speed interstate. The effects of these roads extend far beyond the rights  
14 of way. Roads not only have direct mortality effects on wildlife, but they can alter hydrology, hamper  
15 prescribed burning efforts, act as corridors for invasive plants and animals, and serve as isolation  
16 mechanisms for some wildlife species. Fortunately, these effects are recognized and steps have been  
17 taken to mitigate them wherever possible.

18 To mitigate these effects, the iconic Cross Florida Greenway Land Bridge was erected in 1999-2000.  
19 This was the first true land bridge in the United States, modeled after a design used in the  
20 Netherlands. Although this has been extremely effective from both a recreational as well as wildlife  
21 management perspective, recent discussions have included either the expansion of this land bridge  
22 or even the construction of an additional land bridge over I-75. Particularly, given the development  
23 of the new paved trail funded by FDOT across the CFG from Ocala to Dunnellon expansion of existing  
24 or addition of a second bridge to further enhance ecological and recreational connectivity is needed.

25 In addition to these land bridge discussions, other discussions have been centered around improved  
26 wildlife crossings around other adjacent roadways. This includes an off-grade crossing on US 441, as  
27 well as expanded underpasses when two-lane roads are expanded. This will be discussed and  
28 coordinated with both state and federal DOT.

### 29 **Cultural Resources Management**

30 Cultural resources are individually unique, and collectively, very challenging for the public land  
31 manager whose goal is to preserve and protect them in perpetuity. DRP is implementing the following  
32 goals, objectives, and actions, as funding becomes available, to preserve the cultural resources found  
33 on the CFG.

#### 34 **Goal: Protect, preserve, and maintain the cultural resources of the park.**

35 The management of cultural resources often is complicated because these resources are  
36 irreplaceable and extremely vulnerable to disturbances. The advice of historical and archaeological  
37 experts is required in this effort. All activities related to land clearing, ground-disturbing activities,  
38 major repairs or additions to historic structures 50 years old or older, or those structures listed or

1 eligible for listing in the NRHP will be submitted to the DHR for review and comment prior to  
2 undertaking the proposed project.

3 Recommendations from DHR may include, but are not limited to, concurrence of no effect to  
4 significant cultural resources for the submitted project, monitoring of project activities by a certified  
5 archaeological monitor or qualified professional archaeologist, cultural resource assessment survey  
6 by a qualified professional archaeologist, and modifications to the proposed project to avoid or  
7 mitigate potential adverse effect. In addition, any demolition or substantial alteration to any historic  
8 structure or resource 50 years old or older must be submitted to DHR for consultation and DRP must  
9 demonstrate that there is no feasible alternative to removal and must provide a strategy for  
10 documentation of the resource. Section 267.016(2)(b), F.S., further requires that DRP consider the  
11 reuse of historic buildings in the park in lieu of new construction and must undertake a cost  
12 comparison of new development versus rehabilitation of a building before electing to construct a  
13 new or replacement building. This comparison must be accomplished with the assistance of DHR.

14 *Objective: Assess and evaluate 30 of 300 recorded cultural resources in the park annually.*

15 *Action 1: Complete 300 assessments/evaluations of archaeological sites. Prioritize preservation*  
16 *and stabilization projects.*

17 *Action 2: Complete Historic Structures Reports (HSRs) for historic buildings and cultural*  
18 *landscapes. Prioritize stabilization, restoration, and rehabilitation projects.*

19 All recorded cultural sites will be assessed and evaluated within the 10-year period of this  
20 management plan. The assessments will include an examination of each site with attention being paid  
21 to any threats to the site's condition, such as natural erosion, damage, looting, construction, animal  
22 damage, plant or root damage, or other factors that might cause deterioration of the site. Any  
23 preservation and stabilization identified by the assessments/ evaluations will need to be prioritized.  
24 Due to the numerous cultural resource sites on the CFG and current staffing limitations, DRP may  
25 consider partnering with an archaeology program at a state university to utilize student participation  
26 with the assessments.

27 *Objective: Compile reliable documentation for all recorded historic and archaeological sites.*

28 *Action 1: Ensure all known sites are recorded or updated in the Florida Master Site File. Would*  
29 *be part of Objective B of having professional archaeologist inventory and assess all known and*  
30 *listed sites.*

31 The potential exists for other unrecorded sites; staff will continue to monitor the park for this  
32 possibility and consult with the Bureau of Natural and Cultural Resources and Division of Historic  
33 Resources staff. An archaeological resource predictive model was completed for the park in 2010;  
34 while areas of high, medium, and low sensitivity were identified, there weren't any priority areas  
35 noted where a Phase I cultural resources assessment survey should be conducted. A Scope of  
36 Collections Statement has been developed and adopted and is available at the park.

37 *Objective: Maintain 263 of 263 cultural resource sites in good condition.*

1 This will be achieved by regular monitoring, site stabilization, and protection from disturbance. The  
2 specific sites will be determined after further condition assessments have been conducted.

3 *Objective: Interpret cultural and historical resources on the CFG for the public.*

4 Interpretation on the CFG will be accomplished through a multi-faceted approach with kiosk-based  
5 descriptions of the pre-historic to modern history information related to the natural and cultural  
6 resources that occur on the CFG. There will be a specific emphasis on the story of the 1930s Great  
7 Depression-era sea level Cross Florida Ship Canal that created the numerous large-scale ship canal  
8 “diggings” and the later 1960s to 1970s Cross Florida Barge Canal.

9 Tying both these projects back to the Spanish explorers and the first written record of the desire to  
10 find a water route across the Florida peninsula in the 1500s will be critical, as well as well as the  
11 political machinations over the following several hundred years. We obtained written permission  
12 from the Florida Press to utilize excerpts from the *Ditch of Dreams* book that chronicles the history  
13 of the canal projects and their ultimate transition into the CFG.

14 Eventual interpretive opportunities will be sought using recent technology with qwerty codes where  
15 people can look up various interpretive stories and information with their mobile devices to help  
16 expand the long history of this significant piece of Florida and our nation’s history.

### 17 **Resource Management Schedule**

18 To enhance the resource values, a priority schedule for conducting all management activities, which  
19 is based on the purposes for which these lands were acquired, is located in the Implementation  
20 Component of this management plan.

### 21 **Land Management Review**

22 Section 259.036, F.S., established land management review teams to determine whether  
23 conservation, preservation, and recreation lands titled in the name of the Board of Trustees are being  
24 managed for the purposes for which they were acquired and in accordance with their approved land  
25 management plans.

26 The CFG was subject to a land management review of the entire greenway on January 26, 2010. The  
27 review team made the following determinations:

- 28           1. The land is being managed for the purpose for which it was acquired.
- 29           2. The actual management practices, including public access, complied with the management  
30           plan for this site.

31 On May 26, 2015, a land management review was performed on the eastern portion of the CFG. The  
32 review team made the following recommendations:

- 33           1. The land is being managed for the purpose for which it was acquired.
- 34           2. The actual management practices, including public access, complied with the management  
35           plan for this site.



# LAND USE COMPONENT

## *CFG Planning and Recreational Accomplishments, 2007 to 2016*

### INTRODUCTION

Land use planning and park development decisions for the state park system are based on the dual responsibilities of the FDEP, DRP. These responsibilities are to preserve representative examples of original natural Florida and its cultural resources, and to provide outdoor recreation opportunities for Florida's citizens and visitors.

The general planning and design process begins with an analysis of the natural and cultural resources of the unit, and then proceeds through the creation of a conceptual land use plan to guide the location and extent of future park development. Input to the plan is provided by experts in environmental sciences, cultural resources, park operation and management, and through public workshops and user groups. With this approach, DRP's objective is to provide quality development for resource-based recreation with a high level of sensitivity to the natural and cultural resources at each park throughout the state.

This component of the unit management plan includes a brief inventory of the external conditions and the recreational potential of the unit. Existing uses, facilities, special conditions of use, and specific areas within the park that will be given special protection are identified. The land use component then summarizes the current conceptual land use plan for the park, identifying the existing or proposed activities suited to the resource base of the park. Any new facilities needed to support the proposed activities are described and located in general terms.

Given the size of the CFG and the fact that it traverses four counties (Citrus, Levy, Marion, and Putnam), this section is subdivided into county discussions. Each discussion takes into consideration the external conditions, such as existing and planned use of lands adjacent to the greenway within each county, existing and future land use, and population/demographics. We also analyze each portion of the greenway—its land, water, natural features, and existing recreational opportunities—to determine any additional recreational potential for each area.

- New floating boat dock at Kenwood Boat Ramp
- New floating dock and ADA ramp at the US 19 boat ramp
- New picnic pavilions with ADA access at Eureka Rec Area West, Orange Springs, Kenwood, Rodman East Rec Area, Buckman Rec Area
- Phase 2 Rodman Campground
- Logging Miller Tract
- Road repair and culvert replacement Miller Tract
- St. Johns Trail Loop South Primitive Equestrian Camp Ground
- Hunter Road Trailhead
- Updated Inglis Bypass Recreation area with ADA sidewalks
- Bulkheads at Inglis Lock and equipment bridge
- New section of Withlacoochee Bay Trail going east to Inglis Island
- Dunnellon Trail and Bridge
- Coordinating with Marion County on the new proposed paved trail in between SR 200 and Dunnellon Trail
- Developed and opened Shangri La Campground and Trailhead.
- Developed and opened Vortex Trailhead.
- Developed and opened Ned Folks Pavilion.
- Repair gates on Kilpatrick Dam
- Repair and improve accessibility on fishing dock below Kilpatrick Dam.

1 As indicated previously, for management purposes, the CFG is subdivided into three regions. These  
2 include the following:

- 3 • Gulf of Mexico to SR 200
- 4 • SR 200 to CR 316
- 5 • CR 316 to St. Johns River

6 These are recognized divisions of the CFG, each with individual managers to assist with the overall  
7 management of the CFG, which is complex. CFG management provides challenges that are different  
8 from any other piece of the Florida State Parks system. For instance, the CFG over the past few years  
9 has increasingly utilized contractor labor for burning, timber management, improving wildlife  
10 habitat and invasive species removal. By doing this, the CFG has been successful at meeting specified  
11 resource management goals and objectives. The CFG would like to continue to utilize contractors in  
12 this capacity, and perhaps look at additional uses for this resource.

13 Additionally, the CFG is unique in the number of special use permitted activities that occur on the  
14 greenway. For ease of management of these activities, which totaled 83 last year, as detailed later in  
15 this document, the CFG would like to create an on-line permit tracking system. This would further  
16 assist with the tracking and management of these types of activities.

### 17 **CFG Acquisition History**

18 The Board of Trustees of the Internal Improvement Trust Fund of the State of Florida (Trustees)  
19 acquired the CFG to create a cross Florida greenway corridor within the right of way of the former  
20 Cross Florida Barge Canal. The purpose of this acquisition was so the Trustees could effectively and  
21 efficiently protect, conserve, and preserve the natural resources and scenic beauty of Florida, while  
22 providing and enhancing compatible public recreational opportunities such as fishing, camping,  
23 boating, bicycling, nature study, horseback riding, hiking, hunting, paddling, and other outdoor  
24 interests.

25 On July 26, 1993, the Trustees obtained title to approximately 107 miles of former Cross Florida  
26 Barge Canal project right of way, constituting the initial area of the CFG. The Trustees acquired this  
27 property through a Quitclaim Deed from the United States of America, acting through then Secretary  
28 of the Army, John W. Shannon.

29 Since the initial 1993 acquisition, the Trustees have purchased more parcels using different land  
30 acquisition programs—mainly Preservation 2000 and Florida Forever—and added them to the CFG.  
31 Between 1995 and 2011, the Trustees received six donations of parcels from private individuals, local  
32 governments, private Trustees, and/or private corporations and added them to the CFG. For more  
33 details on the acquisition, see Addendum 1.

34 Between 1996 and 2011, the OGT entered into multiple management lease agreements with  
35 SJRWMD, SWFWMD, and the Felburn Foundation, a private nonprofit corporation. These entities  
36 own title to certain lands now managed as a portion of the CFG.

1 **Subleases, Agreements, and Easements**

2 DRP subleases 29 parcels to other entities for a variety of purposes. All are on former barge canal  
 3 lands. Table 16 lists the major subleases on the CFG. For those subleases ending during the period  
 4 covered by this management plan period, the intended future of each lease is indicated. In addition  
 5 to the subleases in Table 16, DRP operates seven buildings as residences to staff and to law  
 6 enforcement personnel who provide security for the CFG by virtue of their presence.

7 In addition, the Baseline Road to Marshall Swamp Trail is leased to Marion County for operation and  
 8 maintenance.

9 **Table 16. Subleases on the CFG**

State Lands No.	OGT NO.	County	Lessee	Name	Acres	Beginning and Ending Date	Comments
4013-101	PF-69	Marion	Marion County	Orange Springs Boat Ramp (f/k/a 4013-80)	15.5	7/1/95-10/26/16	Currently being renewed
4013-124		Marion	Marion County	Sheriff's Work Farm Project **Amended	58.73	9/30/16 - 9/29/20	
4013-125	PF-57	Marion	City of Dunnellon	Dunnellon Ball Fields	19	9/14/16 - 9/13/21	
4013-102	PF-76	Marion	Marion County	Gore's Landing	106	10/4/06 - 10/3/16	Currently being renewed
4013-117	PF-66	Marion	Marion County	Median of 441 South of 80th Street (S.O. station)	17.75	11/20/06-11/19/16	Currently being renewed
DACW-17-5-14-0002	77	Putnam	USACE	Chemical Storage—Buckman Lock	1.8	7/21/09 - 4/30/19	
4013-115		Marion	Marion County	Recycling Center	3.01	11/16/06-12/31/17	
4013-92		Marion	Marion Therapeutic Riding Association	Therapeutic Equestrian Facility	30.2	7/1/03-6/30/23	
4013-107(?)	83	Marion	Marion County	Baseline Road to Marshall Swamp Trail	830	3/1/99-2/28/24	
	79	Levy	Southern Hy-Power Corp.	Hydroelectric Facility Inglis Spillway	0.61	10/17/95-10/16/25	

State Lands No.	OGT NO.	County	Lessee	Name	Acres	Beginning and Ending Date	Comments
	82	Marion	Marion County	4-H Club Facility SW Corner Highway 464 & CR 35	45	2/1/99- 1/31/43	
	PF-77	Marion	Marion County	Sheriff's Substation	3.43	7/1/95- 6/30/45	
	PF-78	Citrus	Florida Marine Patrol	Marine Patrol Station—Inglis	8.6	1/16/96 - 6/30/45	
4013- 108	PF-79	Putnam	FWC	Caravelle Wildlife Management Area	3,000	7/1/95- 6/30/45	
4013- 90	PF- 97-22	Marion	DACS/ Horsepark Authority	DACS Horsepark— Highway 475 (Subleased to Horsepark Authority) **Amended on 02/11/02	500	4/9/13- 10/26/7 2	
	PF-80	Marion	Marion County	Rotary Sports Complex Sublease	78	9/1/97- 9/1/47	
Totals					4,639.9		

- 1
- 2 The FWC leases 3,000 acres of former barge canal land that is part of the Caravelle Ranch Wildlife  
3 Management Area. Four boat ramps are leased to Marion County: Gore's Landing, the Orange Springs  
4 boat ramp, and two ramps at Eureka, all of which provide access to the Ocklawaha River.
- 5 Approximately 30 acres are leased to the Marion Therapeutic Riding Association, Inc. This non-profit  
6 group provides the opportunity for individuals challenged by physical, mental, and emotional  
7 disabilities to take advantage of the extraordinary physical and psychological benefits of horseback  
8 riding and driving (<http://www.mtraocala.org/>). Marion County also leases land for a 4-H club  
9 facility.
- 10 The FDACS leases 500 acres from the CFG, which it then leases to the Florida Agriculture Center and  
11 Horse Park Authority for the Florida Horse Park, as authorized in Florida Statutes (Ch. 253.7825).  
12 Equestrian facilities are located at the park, and special events are held there.
- 13 CFG leases 19 acres of land to the City of Dunnellon for the T.K. Egan Sports Complex; approximately  
14 10 of these acres are developed. Marion County leases 78 acres for the Ocala Rotary Sports Complex.  
15 These ballfields are widely used by the communities.
- 16 CFG also leases land to the Marion County Sheriff and the FWC for law enforcement stations. An area  
17 formerly heavily infested with cogon grass is leased to the Marion County Sheriff for an inmate work

1 farm project. The cogon grass was eliminated by their operations. In exchange for the land use,  
2 inmate labor is provided to the CFG. The work farm, sports complex, and 4H leases are all located at  
3 the intersection of CR 464 and SR 35.

4 About two acres are leased to the USACE for chemical storage near the Buckman Lock. Less than one  
5 acre is leased to Southern Hy-Power Corporation for a hydroelectric facility, which has not yet been  
6 permitted or constructed at the Inglis Bypass Spillway.

7 Three acres are leased to Marion County for use as a recycling staging area. The City of Dunnellon has  
8 an easement on CFG land used for a water/wastewater plant. Once a better location is identified for  
9 the water/wastewater plant, the easement will be phased out and the lands will be returned to the  
10 CFG.

11 Six parcels managed by the CFG are either leased from other entities or are covered by management  
12 agreements. The SWFWMD has an agreement with CFG/DRP that gives the CFG management of 301  
13 acres that help bridge the gap between the City of Dunnellon and SR 200. The Division of Forestry  
14 has an agreement with CFG/DRP that provides access to the Ross Prairie Trailhead. In exchange,  
15 CFG/DRP includes information at the trailhead about Ross Prairie State Forest. The Felburn  
16 Foundation leases 135 acres to CFG/DRP on the western end of the Greenway for use as a trailhead.  
17 Felburn Park is a former mining area with water features adjacent to the western barge canal.  
18 CFG/DRP also leases space in a chemical storage room at Buckman Lock from the U.S. Army Corps of  
19 Engineers. The land the chemical storage room is on is leased by CFG/DRP to the Corps.

## 20 **EXTERNAL CONDITIONS**

21 An assessment of the conditions that exist beyond the boundaries of the unit is important in  
22 identifying any special development problems or opportunities that exist because of the unit's unique  
23 setting or environment. This also provides an opportunity to deal systematically with various  
24 planning issues, such as location, regional demographics, adjacent land uses, and park interaction  
25 with other facilities.

26 Given that the CFG is more than 110 miles long and traverses four counties (Citrus, Levy, Marion and  
27 Putnam), many opportunities exist to analyze the existing and future development activities within  
28 each county. Further, this plan briefly analyzes the population and demographic characteristics of  
29 the state of Florida, which is highly urbanized. More than 94 percent of Florida's population lived in  
30 metropolitan areas in 2010, and many others lived in small- and medium-sized towns in non-  
31 metropolitan counties. Only a small fraction of Florida's population may be truly classified as rural.

32 Overall, from 2000 to 2010, Florida has experienced a 17.6-percent population increase to 18.8  
33 million people. Today, Florida is the third most populated state in the nation and its population is  
34 projected to reach nearly 24.5 million by 2025 (SCORP, 2013). In addition, Florida's population is  
35 getting older. In 2010, 17.3 percent of Floridians were aged 65 or older, in comparison to the national  
36 average of 13 percent. Florida's mean age of 40.7 was fifth highest in the nation (the national average  
37 is 37.2 years), an upward shift compared to the mean age of 38.7 in 2000.



1 If these current trends continue, the majority of the population increase will concentrate in relatively  
 2 unpopulated counties adjacent to highly developed metropolitan areas, particularly along the coast.  
 3 As these areas become more crowded, problems such as loss of open space and natural areas,  
 4 crowding, and a higher cost of living will provide an impetus to expand farther into rural areas. This  
 5 sprawling pattern of growth reduces the availability of outdoor recreation land and facilities unless  
 6 more land is acquired and more facilities are developed to keep up with the expansion.

7 In the following sections, this plan evaluates the population and demographic characteristics of each  
 8 county the CFG traverses to better understand and detail the opportunities, constraints, and  
 9 interactions with surrounding land uses. Additionally, the Land Use Component evaluates the  
 10 recreational opportunities provided by the CFG within each county. As indicated previously, each  
 11 piece of the CFG is unique in its resources, recreational opportunities, and regional setting.

12 Table 17 provides a listing of the many resource-based recreation areas within a 15-mile radius  
 13 of the CFG. These lands and waters support an array of resource-based outdoor activities, including  
 14 swimming, fishing, canoeing/kayaking, boating, camping, picnicking, hiking, biking, horseback riding,  
 15 wildlife viewing, nature study, and visiting historical sites.

16 **Table 17. Conservation Lands and Activities within 15 Miles of the CFG**

Conservation Area	Managing Agency	Activities
Ocala National Forest	U.S. Department of Agriculture, Forest Service	Fishing, hunting, canoeing, kayaking, hiking, camping, boating, picnicking, wildlife viewing, and bicycling
Withlacoochee State Forest	Florida Department of Agriculture and Consumer Services, Florida Forestry Service	Hiking, bicycling, horseback riding, canoeing, seasonal hunting, fishing, camping, picnicking, and wildlife viewing
Big Bend Seagrasses Aquatic Preserve	Florida Department of Environmental Protection, Office of Coastal & Aquatic Managed Areas	Swimming, wildlife viewing, boating, canoeing, kayaking, scalloping, and fishing
Goethe State Forest	Florida Department of Agriculture and Consumer Services, Florida Forestry Service	Picnicking, hiking, bicycling, fishing, wildlife viewing, and horseback riding
Waccasassa Bay Preserve State Park	Florida Department of Environmental Protection, Division of Recreation and Parks	Birding, boating, canoeing, kayaking, fishing, wildlife viewing, and camping
St. Martins Marsh Aquatic Preserve	Florida Department of Environmental Protection, Office of Coastal & Aquatic Managed Areas	Boating, kayaking, canoeing, hiking, fishing, snorkeling, scuba diving, birding, and wildlife viewing
Crystal River Preserve State Park	Florida Department of Environmental Protection, Division of Recreation and Parks	Biking, hiking, wildlife viewing, fishing, canoeing, and kayaking
Caravelle Ranch Wildlife Management Area	Florida Fish and Wildlife Conservation Commission	Hunting, fishing, hiking, birding, and wildlife viewing
Potts Preserve	Southwest Florida Water Management District	Birding, hiking, biking, horseback riding, boating, fishing, canoeing, and hunting

<b>Conservation Area</b>	<b>Managing Agency</b>	<b>Activities</b>
Halpata Tastanaki Preserve	Southwest Florida Water Management District	Hiking, biking, horseback riding, and fishing
Ocklawaha River Aquatic Preserve	Florida Department of Environmental Protection, Office of Coastal & Aquatic Managed Areas	Swimming, wildlife viewing, boating, canoeing, kayaking, and fishing
Dunns Creek	Florida Department of Environmental Protection, Division of Recreation and Parks	Biking, hiking, horseback riding, picnicking, and wildlife viewing
Ocklawaha Prairie Restoration Area	St. Johns River Water Management District	Wildlife viewing, hiking, horseback riding, bicycling, fishing, seasonal waterfowl hunting, nature study, camping, boating, and canoeing
Gum Slough SWFWMD Conservation Easement	Southwest Florida Water Management District	Hiking, biking, and wildlife viewing
Silver River State Park	Florida Department of Environmental Protection, Division of Recreation and Parks	Biking, birding, camping, canoeing, kayaking, horseback riding, and wildlife viewing
Sunnyhill Restoration Area	St. Johns River Water Management District	Hiking, horseback riding, wildlife viewing, bicycling, picnicking, and primitive camping
Rice Creek Conservation Area	St. Johns River Water Management District	Hiking, bicycling, horseback riding, and primitive camping
Ross Prairie State Forest	Florida Department of Agriculture and Consumer Services, Florida Forestry Service	Hiking, horseback riding, birding, seasonal hunting, and picnicking
Dunns Creek Conservation Area	St. Johns River Water Management District	Seasonal hunting, hiking, bicycling, horseback riding, fishing, and wildlife viewing
Murphy Creek Conservation Area	St. Johns River Water Management District	Hiking, bicycling, horseback riding, wildlife viewing, and nature study; primitive camping is allowed only at designated sites
Rainbow Springs State Park	Florida Department of Environmental Protection, Division of Recreation and Parks	Birding, boating, canoeing, kayaking, fishing, wildlife viewing, and camping
Withlacoochee State Trail	Florida Department of Environmental Protection, Division of Recreation and Parks	Hiking, bicycling, horseback riding, and birding
Palatka-Lake Butler State Trail	Florida Department of Environmental Protection, Division of Recreation and Parks	Hiking, bicycling, horseback riding, and birding
Carl Duval Moore State Forest and Park	Florida Department of Agriculture and Consumer Services, Florida Forestry Service	Hiking, birding, and fishing
Seven Sisters Islands	St. Johns River Water Management District	Birding, boating, canoeing, kayaking, fishing, wildlife viewing, and camping

Conservation Area	Managing Agency	Activities
Rainbow Springs Aquatic Preserve	Florida Department of Environmental Protection, Office of Coastal & Aquatic Managed Areas	Tubing, boating, fishing, snorkeling, kayaking, canoeing, and swimming
Ravine Gardens State Park	Florida Department of Environmental Protection, Division of Recreation and Parks	Wildlife viewing, hiking, bicycling, and picnicking
Crystal River Archaeological State Park	Florida Department of Environmental Protection, Division of Recreation and Parks	Picnicking, fishing, wildlife viewing, and an interpretive exhibit

1

2 **Past Uses**

3 The lands of the Marjorie Harris Carr Cross Florida Greenway have a long history of human use and  
 4 construction. In addition to the Native Americans who settled the area nearly 10,000 years ago,  
 5 people of European descent have been active in the area since the 16th century. Subsistence farming,  
 6 plantations, and timbering were common. Around 1870, the Ocklawaha River saw increased  
 7 steamboat trade and tourism.

8 Along with the more obvious manipulations related to the barge canal, the Cross Florida Greenway  
 9 has seen a variety of uses since its acquisition for the barge canal and prior to establishment of the  
 10 Greenway. Land was acquired for the sea level ship canal in the 1930s and for the barge canal in the  
 11 1960s. When canal work was not taking place on these lands, they were often leased out for caretaker  
 12 purposes, and timbering and grazing were common. The state managed most of the barge canal lands  
 13 from the 1960s to 1990 and the state took over management of all former barge canal lands in 1991.

14 **Future Land Use and Zoning**

15 The DRP works with local governments to establish designations that provide both consistency  
 16 between comprehensive plans and zoning codes that permit typical state park uses and facilities  
 17 necessary for the provision of resource-based recreation opportunities.

18 Each county works to designate park lands with “conservation” zoning and “conservation” future  
 19 land use. Typical state park development is permitted in these categories with review by the county.

20 **Current Recreation Use and Visitor Programs**

21 Given the significant amount of natural resources in Florida, resource-based outdoor recreation is  
 22 very popular. Resource-based outdoor recreation differs from user-oriented recreation in that it  
 23 cannot be provided just anywhere, but is dependent upon some element or combination of elements  
 24 in the natural or cultural environments that cannot be easily duplicated by man. Examples of  
 25 activities include fishing, hiking, biking, horseback riding, hunting, camping, boating, surfing, nature  
 26 study, and visiting historical sites. Throughout the CFG, amenities are offered to all visitors to  
 27 enhance individual recreational experiences. These amenities include, but are not limited to: picnic  
 28 pavilions and benches, grills, playgrounds, boat launches, paved multi-use trails, ADA accessible  
 29 natural surface trails, fishing piers, and accessible campsites.

30 The growth of Florida’s resident and tourist populations brings increasing pressure for more  
 31 widespread access for denser levels of public use in the natural areas available to the public.

1 Consequently, one of the greatest challenges for public land management today is to balance  
 2 reasonable levels of public access with the need to preserve and enhance the natural and cultural  
 3 resources of the protected landscapes.

4 Table 18 provides visitor estimates from the last eight years at the CFG.

5 **Table 18. CFG Visitor Estimates (FY 09 to FY 17)**

Fiscal Year	Traffic Count	Trail Count	Total Visitation	Comments:
FY 09-10	303,653	861,987	1,165,640	
FY 10-11	928,164	728,010	1,656,174	
FY 11-12	965,686	404,093	1,369,779	Removed 12 trail counters in April 2012
FY 12-13	847,237	159,627	1,006,864	
FY 13-14	747,634	135,139	882,773	
FY 14-15	757,234	161,257	918,491	
FY 15-16	1,042,420	159,351	1,201,771	
FY 16-17 (thru 1/31/17)	571,412	89,399	660,811	
<b>Total</b>			<b>8,862,303</b>	

6 Source: Cross Florida Greenway

7 Key: These numbers are the raw traffic and trail counter numbers prior to using the 2.5 x multiplier  
 8 per vehicle counted until Oct 2014, then the DRP changed the standard vehicle counted multiplier  
 9 to 3x statewide. This explains why these are qualified as estimated visitation. Also, the note about  
 10 elimination of 12 trail counters was after merger into DRP in July 2011 and District leadership  
 11 wanted to reduce those to minimize double counts for conservative estimated visitation.

# 1 PROPERTY ANALYSIS BY COUNTY

## 2 Citrus County

### 3 History/Setting

4 Citrus County was created in 1887 and was named for its  
5 primary industry: citrus growing. Citrus production  
6 declined significantly following the “Big Freeze of 1894-  
7 1895.” With the decline of the citrus industry, phosphate  
8 mining became the largest industry, which continued until  
9 World War I. Planned industrial development surrounding  
10 the construction of the 1930s ship canal never came to  
11 fruition when the partially built canal was terminated after  
12 economic and environmental opposition.

13 Within Citrus County, the CFG extends briefly along the  
14 northern county boundary, beginning at the Gulf of Mexico  
15 (see map at top of column to the right). Eleven dredge Spoil  
16 Islands exist along the westernmost portion of the CFG out  
17 into the Gulf of Mexico. These Spoil Islands are remnants  
18 from the original failed public works project. Also within  
19 Citrus County is the Inglis Dam, Felburn Park/Bay Trail  
20 and a portion of the Barge Canal.

### 21 Population/Demographics

22 Although Citrus County is within an hour’s drive from the  
23 city of Tampa, the county remains relatively rural in nature  
24 with a population density of 182.64 persons per square  
25 mile. This number is much lower than the state average  
26 population density of 375.7 people per square mile, but is  
27 much higher than the national average population density  
28 of 82.73 people per square mile. According to the U.S.  
29 Census Bureau, in 2010 the population of Citrus County  
30 was 141,236. Estimates from the University of Florida,  
31 Bureau of Economic and Business Research (BEBR)  
32 indicate that the 2016 population of Citrus County was  
33 143,054.

34 Between 2010 and 2016, the population of Citrus County  
35 increased at an average annual rate of 1.3 percent, which  
36 was lower than the rate of growth recorded throughout  
37 Florida (7.2 percent per year), but higher than the national  
38 average of 0.9 percent per year. According to the BEBR,  
39 over the next 25 years, Citrus County’s population is  
40 projected to grow steadily, reaching 156,200 people by the  
41 year 2025.



#### **Land Area:**

Total: 773 square miles (2,002 km<sup>2</sup>)  
Land: 582 square miles (1,507 km<sup>2</sup>)  
Water: 192 square miles (497 km<sup>2</sup>)


#### **Population:**

143,054 (2016)

**Median Age:** 54 years old

#### **Population Density:**

182.64 people per square mile





1 Table 19, provided below, indicates the low, medium, and high population projections through 2045.

2 **Table 19. Historic, Current and Projected Population through 2045—Citrus County**

	2015	2020	2025	2030	2035	2040	2045
Low	141,501	141,800	143,300	144,700	145,400	145,100	144,200
Medium		149,300	156,200	162,100	167,500	171,700	175,500
High		155,900	167,500	178,900	190,100	200,700	211,000
State of Florida*	19,815,183	21,372,200	22,799,500	24,071,000	25,212,400	26,252,100	27,217,600

3

4 Source: University of Florida, Bureau of Economic and Business Research, 2015.

5 \*Used medium population projections.

6 Table 20, provided below, indicates the historic, current, and future population change in Citrus  
7 County.

8 **Table 20. Population Change—Citrus County**

	1980-1990 Percent Change	1990-2000 Percent Change	2000-2010 Percent Change	2010-2015 Percent Change
Citrus County	70.9	26.3	19.6	-0.1
State of Florida	32.7	23.5	17.6	8.0

9

Source: U.S. Census Bureau, 2010.

10 In 2015, 50.8 percent of residents of the County were age 55 and older compared to 30.5 percent of  
11 residents of Florida, and 21.1 percent of the resident population of the U.S. Further, in 2015, the  
12 median age of residents of Citrus County was 55.4 years of age, which was significantly higher than  
13 the median age of residents of Florida (41.6 years), and particularly the U.S. (37.4 years). Table 21  
14 provides a breakdown of the population in Citrus County by age from 2010 to 2045.

1 **Table 21. Population by Age—Citrus County**

Age	Census	Estimates	Projections					
	2010	2015	2020	2025	2030	2035	2040	2045
0-4	5,537	5,280	5,390	5,717	5,901	6,050	6,055	6,202
5-17	16,857	15,875	16,480	16,997	17,474	18,146	18,726	18,987
18-24	8,021	8,292	7,630	7,652	8,201	8,264	8,462	8,845
25-54	42,279	40,147	40,505	41,399	42,712	45,594	47,231	48,231
55-64	23,501	24,175	27,185	26,649	24,234	22,685	23,309	26,207
65-79	33,463	35,221	37,843	41,290	44,948	46,367	44,634	40,978
80+	11,578	12,511	14,242	16,481	18,660	20,349	23,283	26,051
<b>Total</b>	141,236	141,501	149,275	156,185	162,130	167,455	171,700	175,501

2 Source: University of Florida, Bureau of Economic and Business Research, 2015.

3  
 4 Over the long term, projections for Florida indicate that the state’s population will increase at an  
 5 average annual rate of 2.1 percent between 2020 and 2030, reaching 28.7 million people by 2030.  
 6 This rate of growth is significantly higher than the growth rate forecast throughout the U.S. (0.8  
 7 percent per annum), reflecting long-term growth rates in the state.

8 **Existing Use of Adjacent Lands**

9 Citrus County, with an abundance of natural resources, is located just west of I-75, and is accessible  
 10 by several interstate highways. Florida State Highways 98, 44, and 41 all provide access to the County  
 11 and link residents with several major cities, including Orlando, Tampa, and Gainesville. The provision  
 12 of four lanes to accommodate traffic on U.S. Highway 19 (US 19) across the Cross Florida Barge Canal  
 13 also supports the area’s mobility.

14 Within Citrus County, the CFG is not within proximity of heavily populated areas, but it is located  
 15 adjacent to Yankeetown and Inglis in Levy County, as well as Dunnellon in Marion County. The CFG  
 16 is situated in the northwestern portion of Citrus County and runs along the northern border of the  
 17 county prior to leading into Lake Rousseau. At its closest, the CFG is within 0.2 miles north of the  
 18 Crystal River Energy Complex. In all, approximately 141,236 people live in Citrus County (U.S. Census  
 19 2010) and are within 30 miles of the CFG.

20 Inside the one-mile buffer area of the CFG in Citrus County lies a variety of existing land uses. The  
 21 most common includes Agriculture (42 percent), Residential (40 percent), Public/ Institutional (10  
 22 percent), Non-Agricultural Acreage (3 percent), and Other (2 percent). Agriculture and Public/  
 23 Institutional land uses, although scattered throughout the study area, are primarily located in the  
 24 northwestern portion of Citrus County. Residential land uses are prominent along Lake Rousseau  
 25 and across the study area. Non-Agricultural Acreage and Other, although not as prominent, also are  
 26 scattered across the study area.

27 Figure 7, below, displays the generalized existing land use for Citrus County within one mile of the  
 28 CFG.

1 **Planned Use of Adjacent Lands**

2 The northwest portion of Citrus County has many attractive qualities. As indicated earlier, travel to  
3 and from this area is easy given the abundant roadway access, as well as waterfront availability. In  
4 addition, the relatively pristine land with proximity to the coast makes it attractive for future  
5 development.

6 Although the area adjacent to the CFG has significant limitations for future development, there are  
7 some plans for future development within the area. The current comprehensive plan directs growth  
8 toward the central portion of the county, from Citrus Springs, south to Homosassa and northwest of  
9 Inverness, rather than its outer reaches. Reasons for inland development include the environmental  
10 sensitivity of the west coast and its susceptibility to flooding in low-lying areas as experienced during  
11 Hermine (2016). Additionally, the infrastructure necessary to support growth and development in  
12 this area of the county is not present.

13 Hollinswood Harbor is an approved development with a subarea plan that contains commercial,  
14 industrial, and water-dependent uses. It is planned as a working waterfront with a marina, resort,  
15 recreational, residential, industrial, and support education/institution uses incorporated in the  
16 Master Plan. Both applications were approved and are reflected within the Citrus County  
17 Comprehensive Plan and the Future Land Use Map. Including canal bottom, this project will contain  
18 545 acres and border the north shore of the Barge Canal west of the US 19 bridges. Another possible  
19 development includes a boat ramp with a proposed location on the Cross Florida Barge Canal west  
20 of US 19. At this time, the infrastructure necessary to support growth and development in this area  
21 of the county is not present. Citrus County is a member of the Hernando/Citrus Metropolitan  
22 Planning Organization (MPO). Based on a review of the *Hernando/Citrus MPO 2040 Long Range*  
23 *Transportation Plan*, Citrus County did not have any proposed widening or new alignment roadway  
24 projects within the vicinity of the CFG.

25 Inside the one-mile buffer area of the CFG in Citrus County, some changes can be noticed from existing  
26 land use to future land use. These changes include a substantial increase in Residential (57 percent)  
27 across most of the CFG area, Industrial (19 percent) in the northwestern portion, Infrastructure (7  
28 percent) primarily focused in the southwestern portion, Mixed Use (7 percent) primarily in the  
29 northeastern portion, and Commercial (4 percent) along US 98 in the western portion as well as along  
30 CR 488 and US 41. The increase in the Infrastructure future land use is attributed mostly to the  
31 County's land use re-classification of the Crystal River Energy Complex, which was considered as  
32 Public/Institutional under existing land use. The most substantial decrease occurs in the Agriculture  
33 (4 percent) land use within the CFG study area.

34 Figure 8, below, displays the generalized future land use for Citrus County within one mile of the CFG.

35 **Property Analysis**

36 As indicated previously, in an attempt to better understand and detail the specific recreation  
37 resource elements, this plan describes these resources by county. Because effective planning requires  
38 a thorough understanding of the unit's natural and cultural resources, this section describes the  
39 resource characteristics and existing uses of the property. The unit's recreation resource elements  
40 are examined to identify the opportunities and constraints they present for recreational

1 development. Past and present uses are assessed for their effects on the property, compatibility with  
2 the site, and relation to the unit's classification.

3 *Recreation Resource Elements*

4 This section of the Land Use Component assesses the park's recreational resource elements—those  
5 physical qualities that, either singly or in certain combinations, can support various resource-based  
6 recreation activities. Breaking down the property into such elements provides a means for measuring  
7 the property's capability to support potential recreational activities. This process also analyzes the  
8 existing spatial factors that either favor or limit the provision of each activity.

9 **Land Area**

10 Overall, the CFG encompasses a total of 6,729.1 acres within Citrus County. Table 22 shows that there  
11 are 18.7 miles of trails in Citrus County. As demonstrated in Table 23, 1,884.1 acres within the Citrus  
12 County portion of the CFG are considered salt marsh extending from the Gulf of Mexico. As salt marsh,  
13 these lands are subject to extreme tidal events—particularly during storm events—so they are  
14 limited regarding recreational opportunities. However, opportunities may exist for walking trails and  
15 wildlife viewing, particularly since portions of the CFG through the western portion of the greenway  
16 are within the Great Florida Bird Watching Trail (GFBWT).

17 **Table 22. Trails and Mileage on the CFG—Citrus County**

Trail Name	Length in County (Feet)	Length in County (Miles)
Dixon Hammock	39,084.16	7.4
Dunnellon Trail	5,083.68	1.0
Felburn	12,928.37	2.4
Felburn Trail	8,107.52	1.5
North Canal	8,111.47	1.5
Withlacoochee Bay Trail	25,981.54	4.9
<b>Total</b>		<b>18.7</b>

18

19 Additionally, this portion of the CFG contains 2,336.2 acres, or 35 percent, of altered lands, which  
20 include the Barge Canal and the Inglis Dam. Many of the recreational opportunities within this area  
21 of the CFG are focused along and within developed areas, as demonstrated by the 2.5 miles of the  
22 Withlacoochee Bay Trail—a 12-foot-wide multi-use trail that follows the south side of the barge  
23 canal. The remaining 2.5 miles of the 5-mile-long Withlacoochee Bay Trail traverses tidal marsh and  
24 hydric hammock natural communities.

25 **Table 23. Natural Communities and Acreage on the CFG—Citrus County**

Community	Acreage	Percent Total
Salt Marsh	1,884.1	28
Blackwater Stream	15.6	.002
Altered Lands	2,336.2	35
Depression Marsh	137.4	2
Dome Swamp	6.0	.0009

Community	Acreage	Percent Total
Floodplain Swamp	251.1	4
Hydric Hammock	877.7	13
Mesic Flatwoods	100.9	1
Mesic Hammock	585.8	9
Sandhill	47.9	1
Scrubby Flatwoods	301.7	4
Wet Flatwoods	167.5	2
Xeric Hammock	17.2	1
<b>Total</b>	<b>6,729.1</b>	<b>100</b>

1 **Key:** Altered land use includes abandoned fields, canals/ditches, clearing, developed areas, impoundment/artificial pond, pine plantation,  
2 road, spoil areas, and utility corridors.

### 3 Water Area

4 Water resources on the CFG within Citrus County are a mix of manmade and natural resources. The  
5 CFG does not include the submerged boundary beyond the mean high waterline; however, the  
6 shoreline that is present within Citrus County offers boating, fishing, paddling, swimming, wildlife  
7 viewing and photography opportunities. The Gulf of Mexico area at the westernmost portion of the  
8 CFG also offers paddling and opportunities to view communities of nesting shorebirds, as well as  
9 access to other salt marsh areas and barrier islands south and west of the paved trail and barge canal.  
10 These areas contain some of the largest populations of nesting American Oystercatchers.  
11 Additionally, this area provides the starting point for the Segment 7 (Nature Coast) portion of the  
12 Florida Circumnavigation Saltwater Paddling Trail.

13 By its very nature, the CFG includes structures and impoundments that remain from the original  
14 construction of the Cross Florida Barge Canal. The CFG also includes the 3,400-acre Lake Rousseau  
15 impoundment, which was formed by the construction of Inglis Dam in 1909 by Florida Power  
16 Corporation to create hydropower. Hydropower operations ceased in 1965. In the 1960s, the U.S.  
17 Army Corps of Engineers built the Citrus County portion of the Cross Florida Barge Canal between  
18 Lake Rousseau and the Gulf of Mexico. The construction of the barge canal interrupted the natural  
19 flow of water from the upper segment into the lower segment of the river. The Corps constructed the  
20 8,900-foot-long Inglis Lock bypass channel and bypass spillway to discharge fresh water from Lake  
21 Rousseau into the lower Withlacoochee River segment.

### 22 Natural Scenery

23 This portion of the CFG offers expansive views of the Gulf of Mexico from the pavilion at the end of  
24 the greenway. The Withlacoochee Bay Trail offers scenic views. Also, this portion of the CFG includes  
25 Lake Rousseau and the Withlacoochee River which offer excellent opportunities to view wildlife. This  
26 scenic setting is conducive to nature study, wildlife viewing, and photography.

### 27 Significant Habitat

28 The shorebird rookery on the large island is one of the park's important habitats. During the nesting  
29 season, the activities of parents and young can be observed from the dock and the scenic overlook.  
30 The park's maritime hammock is an important habitat for migrating songbirds and provides visitors  
31 with good opportunities for wildlife watching. The salt marsh is another significant habitat in the  
32 park, which provides excellent opportunities to observe numerous species of wading birds and other



1 avian species, including Pelicans, Osprey, and Bald Eagles. The dock and boardwalk provide access  
2 to this community and should have interpretive signage placed to inform visitors about the important  
3 role that marshes play in marine ecology.

4 **Natural Features**

5 The salt marsh and hydric hammock are the most significant natural features in this portion of the  
6 CFG. They provide a setting for a variety of recreational activities, including hiking, nature study,  
7 wildlife viewing, photography, picnicking, and the interpretation of natural and cultural resources.

8 **Archaeological and Historical Features**

9 The previously recorded archaeological sites and historic features in the Citrus County section of the  
10 CFG consist exclusively of a variety of prehistoric archaeological sites. There is a recorded steamship  
11 wreck just off the coast that dates to the mid-19th century. The nature of a number of the  
12 archaeological sites is unclear, but each in its own way offers a good cross section of Central Florida's  
13 prehistoric past.

14 **Assessment of Use**

15 All legal boundaries, significant natural features, structures, facilities, roads, and trails existing in the  
16 unit are delineated on the base map (see Base Map). Specific uses made of the unit are briefly  
17 described in the following sections.

18 **Current Recreation Use and Visitor Programs**

19 The following recreational activities occur along the CFG within Citrus County.

20 The *Felburn Park Trailhead*, *Withlacoochee Bay Trail*, and *Inglis Dam Recreation Area* are recreational  
21 resources that are located within the Citrus County portion of the CFG. Also, the dredge Spoil Islands  
22 off the shoreline provide excellent bird-watching opportunities.



The *Withlacoochee Bay Trail* traverses five miles west from the Felburn Park Trailhead to the Gulf of Mexico, along the south shore of the Cross Florida Barge Canal. This multi-purpose trail is 12 feet wide and runs adjacent to the barge canal for the first 2.5 miles, then switching its path to the south side of the "berm" that was created by the canal's excavation. The westernmost 2.5 miles of the trail run through scenic maritime hammock and salt marsh habitats. Bicycling, walking, or inline skating on the paved trail

32 provide numerous opportunities to observe wildlife along the route. The trail also is one of the two  
33 GFBWT locations on the CFG.

34 *Felburn Park* is an approximately 140-acre property that is owned by the private non-profit Felburn  
35 Foundation and is leased to the state of Florida as part of the CFG. Felburn Park, formerly a limerock  
36 mine of approximately 140 acres, features the 40-acre "Phil's Lake," named after Phil Felburn,  
37 founder of the Felburn Foundation. Located just east of US 19 on the south side of the barge canal,  
38 the park has paved parking, potable water, several picnic pavilions, a small playground, and it  
39 provides access to paved trails running east and west along the barge canal. Traveling to the west is  
40 the Withlacoochee Bay Trail, which contains multi-use platforms available for picnicking, taking a

1 break, or fishing in the barge canal before terminating at a scenic overlook pavilion adjacent to the  
2 entrance of the barge canal channel to the Gulf of Mexico.

3 To the east from Felburn Park, the trail travels between the approximately 40-acre freshwater Phil's  
4 Lake and the brackish water barge canal. The trail continues eastward and rises onto the top of the  
5 berm created by the construction of the barge canal. The trail ventures 1.25 miles east of Felburn  
6 Park before ending near the old Withlacoochee River channel, where it was bifurcated by the barge  
7 canal.

8 In addition to the paved trails at Felburn Park, there are a few miles of mowed grass and natural  
9 surface trails that circle Phil's Lake. When the mine was operational, it eventually hit the local  
10 groundwater table and freshwater began to seep into the pit. Eventually, the pumping of the water  
11 became too much to continue and the mining ceased. Phil's Lake was the result, which is home to  
12 numerous species of freshwater fish. Recreational enthusiasts can use canoes, kayaks, and other non-  
13 motorized craft to paddle and fish the lake. Bank fishing also is possible in some locations to catch  
largemouth bass, bluegill, and catfish.



*Inglis Dam Recreation Area* is located off West Riverwood Drive approximately two miles east of US 19. Paved parking is located south of the Main Dam, along with a boat ramp to access the upstream Lake Rousseau side of the dam. Multi-use platforms run along the lakeshore for freshwater fishing, bird watching, or picnicking.

The Inglis Main Dam, the larger of two spillway structures for Lake Rousseau, is located at this site and provides an  
23 access bridge onto Inglis Island's south side. At the dam, as a part of the GFBWT, wintering waterfowl  
24 may be observed, such as the Ring-Necked Duck, Greater Scaup, Ruddy Duck, and Common Loon.

25 On the lower downstream side of the dam, there is another large paved parking area with a boat ramp  
26 that also serves as a canoe/kayak launch. This ramp provides access to the Withlacoochee River  
27 segment that runs 1.5 miles from below the Main Dam and Spillway to the barge canal channel. This  
28 segment of the old river can have high freshwater flows if significant discharges from the lake are  
29 needed, but often there is little or no flow and the river can be tidally influenced from the Gulf of  
30 Mexico. There are both fresh and saltwater fish species to catch on the downstream side of the dam,  
31 and occasionally manatees are seen in these waters during the warmer months.

### 32 Other Uses

33 The FWC has a law enforcement field office located on the CFG. FWC also has a dedicated office space  
34 at the park.

### 35 Protected Zones


36 A protected zone is an area of high sensitivity or outstanding character from which most types of  
37 development are excluded as a protective measure. Generally, facilities requiring extensive land  
38 alteration or resulting in intensive resource use—parking lots, camping areas, shops, or maintenance  
39 areas—are not permitted in protected zones. Facilities with minimal resource impacts—trails,

1 interpretive signs, and boardwalks—generally are allowed. All decisions involving the use of  
2 protected zones are made on a case-by-case basis after careful site planning and analysis.

3 Although, by definition, the dredge Spoil Islands are not considered to be high-quality natural  
4 communities, these areas are used extensively by nesting shorebirds and are being studied through  
5 a multi-partnered research operation. The dredge Spoil Islands should be considered as potential  
6 protected zones and not incur extensive recreational opportunities.




1 **Figure 7. Citrus County Generalized Existing Land Use**



2



1 **Figure 8. Citrus County Generalized Future Land Use**



2



# 1 Levy County

## 2 History/Setting

3 Levy County was created in 1845, after the Seminole  
4 Wars. The county was named for David Levy, a planter  
5 elected in 1841 as the state's territorial delegate to the  
6 United States House of Representatives, where he  
7 served two terms. When Florida was admitted as a  
8 state, Levy was elected by the new state legislature as  
9 one of Florida's first two U.S. senators; he served from  
10 1845 to 1851, and again from 1855 to 1861.


11 Within Levy County, the CFG extends briefly along the  
12 southeastern corner of the county, primarily along  
13 Lake Rousseau. Similar to Citrus County, this portion is  
14 primarily represented by disturbed lands associated  
15 with the failed public works project. However, there  
16 are a few areas of nice natural habitat remaining on  
17 Inglis Island. This area also includes the bypass canal  
18 and lock.

## 19 Population/Demographics

20 Levy County is comprised of eight incorporated cities  
21 and towns: Bronson, Cedar Key, Chiefland, Fanning  
22 Springs, Inglis, Otter Creek, Williston, and Yankeetown.  
23 Levy County is located along the western coast of  
24 North-Central Florida just north of Citrus County and  
25 is ranked ninth out of 67 counties in Florida in overall  
26 size.

27 In total, Levy County encompasses 1,412 square miles  
28 (903,680 acres), with a population density of 36.5  
29 people per square mile, substantially lower than the  
30 state average of 369.5. Contributing to this is the fact  
31 that the county contains a total of 353,400 acres of  
32 private forest land.

33 According to the University of Florida, BEBR (2015),  
34 Levy County's population is projected to grow steadily  
35 for the next 30 years, reaching 45,900 people by the  
36 year 2030, as indicated below in Table 24. Further, as  
37 illustrated in Table 25, between 1980 and 2010, the  
38 population of Levy County increased at an average rate



### **Land Area:**

Total: 1,412 square miles (3,660 km<sup>2</sup>)  
Land: 1,118 square miles (2,900 km<sup>2</sup>)  
Water: 295 square miles (760 km<sup>2</sup>)

### **Population:**

40,801 (2016)

**Median Age:** 45 years old

### **Population Density:**

36.5 people per square mile

1 of 27.2 percent. However, from 2010 to 2016, the population of Levy County decreased by  
 2 0.6 percent.

3 **Table 24. Historic, Current and Projected Population through 2045—Levy County**

	2015	2020	2025	2030	2035	2040	2045
Low	40,448	40,400	40,700	41,000	41,000	41,000	40,700
Medium		42,500	44,300	45,900	47,200	48,500	49,600
High		44,400	47,600	50,600	53,700	56,700	59,600
State of Florida*	19,815,183	21,372,200	22,799,500	24,071,000	25,212,400	26,252,100	27,217,600

4 Source: University of Florida, Bureau of Economic and Business Research, 2015.

5 \*Used medium population projections.

6  
 7 **Table 25. Population Change—Levy County**

	1980-1990 Percent Change	1990-2000 Percent Change	2000-2010 Percent Change	2010-2016 Percent Change	2015-2020 Percent Change	2020-2025 Percent Change
Levy County	30.4	32.9	18.4	-0.6	4.8	4.3
State of Florida	32.7	23.5	17.6	7.2	6.1	6.7

8  
 9 In 2015, 38 percent of residents of the county were age 55 and older, compared to 30.5  
 10 percent of residents of Florida and 21.1 percent of the resident population of the U.S. Further,  
 11 in 2015, the median age of residents of Levy County was 45 years of age, which was slightly  
 12 higher than the median age of residents of Florida (41.6 years), and considerably higher than  
 13 the median age of the U.S. population (37.4 years). Table 26 provides a breakdown of the  
 14 population in Levy County by age from 2010 through 2045.

15 **Table 26. Population by Age—Levy County**

Age	Census	Estimates	Projections					
	2010	2015	2020	2025	2030	2035	2040	2045
0-4	2,299	2,185	2,234	2,304	2,336	2,376	2,387	2,438
5-17	6,330	5,889	5,989	6,096	6,227	6,372	6,500	6,564
18-24	3,073	3,118	2,956	2,984	3,065	3,096	3,186	3,272
25-54	14,881	14,021	14,163	14,670	15,142	15,822	16,243	16,595
55-64	6,304	6,481	7,045	6,740	6,196	6,194	6,447	6,941

Age	Census	Estimates	Projections					
	2010	2015	2020	2025	2030	2035	2040	2045
65-79	6,223	6,990	7,998	8,944	9,771	9,736	9,512	9,014
80+	1,691	1,764	2,120	2,581	3,129	3,645	4,205	4,738
<b>Total</b>	40,801	40,448	42,505	44,319	45,866	47,241	48,480	49,562

Source: University of Florida, Bureau of Economic and Business Research, 2015.

By comparison, over the long term, projections for Florida indicate that the state’s population will increase at an average annual rate of 2.1 percent between 2020 and 2030, reaching 28.7 million people in 2030. This rate of growth is significantly higher than the growth rate forecast for the U.S. (0.8 percent per annum), reflecting long-term growth rates in the state.

**Existing Use of Adjacent Lands**

Levy County, a coastal county situated along Florida’s Gulf Coast, also is easily accessible. Located west of I-75, Florida State Highways 27, 41, and 98 also provide access to the county and link residents with several major cities, including Gainesville, Ocala, Orlando, and Tampa.

Within Levy County, the CFG is within proximity of some populated areas, including Inglis and Yankeetown. The CFG crosses the southern portion of Levy County. In all, approximately 40,801 people live in Levy County (U.S. Census 2010) and are within 40 miles of the CFG.

Inside the one-mile buffer area of the CFG in Levy County, there are a variety of existing land uses. The most common land uses include Agriculture (49 percent), Residential (30 percent), Public/Institutional (12 percent), Non-Agricultural Acreage (4 percent), and Recreation (2 percent). Agriculture land uses are predominantly west of Inglis in the one-mile buffer area of the CFG. Residential land uses are scattered throughout, with some concentration in the Yankeetown and Inglis areas. Public/Institutional and Non-Agricultural Acreage land uses are predominately within the eastern portion of Yankeetown, with some scattered throughout the CFG area. Recreation land use is limited to the west in Yankeetown.

Figure 9, below, displays the generalized existing land use for Levy County within one mile of the CFG.

**Planned Use of Adjacent Lands**

Southern Levy County has its fair share of attractive qualities. As indicated earlier, travel to and from this area is easy given the abundant roadway access. Its location along the Gulf of Mexico and Lake Rousseau—as well as its proximity to larger cities—make it attractive for future development.

Overall, Levy County expects a sustained annual population growth rate of at least 2 percent. Most of this growth will be within the unincorporated area, converting approximately 300

1 acres county-wide per year to residential use. Commercial, industrial, public/quasi-public,  
2 recreational, and agricultural uses are expected to change at a rate dependent upon  
3 population growth, the overall economy, and government decisions. A review of the 2018-  
4 2022 Florida Department of Transportation (FDOT) Five-Year Work Program and Levy  
5 County’s website did not identify any proposed widening or new alignment roadway projects  
6 within the vicinity of the CFG in Levy County.

7 Inside the one-mile buffer area of the CFG in Levy County, some changes can be noticed from  
8 existing land use to future land use. These changes include a substantial increase in  
9 Residential (45 percent) and a decrease of Agriculture (22 percent) land uses. The other most  
10 common types of land use include Municipality (21 percent), Mixed Use (4 percent), and  
11 Conservation (3 percent). The area experiencing perhaps the most change is the area  
12 between the towns of Inglis and Dunnellon north of Lake Rousseau, which has experienced a  
13 significant increase in Residential land use and a decrease in Agriculture land use.  
14 Municipality land use includes the coastal and some inland portions of Yankeetown around  
15 CR 40A. Mixed-Use land use is planned for the area between CR 40 and CR 40A west of  
16 Yankeetown School. Conservation areas are primarily in the Inglis area around US 98.

17 Figure 10, below, displays the generalized future land use for Levy County within one mile of  
18 the CFG.

19 **Property Analysis**

20 As indicated previously, in an attempt to better understand and detail the specific recreation  
21 resource elements, this plan describes these resources by county. Effective planning requires  
22 a thorough understanding of the unit's natural and cultural resources. This section describes  
23 the resource characteristics and existing uses of the property. The unit's recreation resource  
24 elements are examined to identify the opportunities and constraints they present for  
25 recreational development. Past and present uses are assessed for their effects on the  
26 property, compatibility with the site, and relation to the unit's classification.

27 *Recreation Resource Elements*

28 This section of the Land Use Component assesses the park’s recreational resource elements,  
29 those physical qualities that, either singly or in certain combinations, can support various  
30 resource-based recreation activities. Breaking down the property into such elements  
31 provides a means for measuring the property's capability to support potential recreational  
32 activities. This process also analyzes the existing spatial factors that either favor or limit the  
33 provision of each activity.

34 **Land Area**

35 Overall, Levy County contains approximately 2,262.4 acres—the smallest county traversed  
36 by the CFG. Table 27 shows that there are only 12.1 miles of trails in Levy County, in contrast  
37 with the more than 200 miles of trails in Marion County. As shown in Table 28, a total of  
38 1,539.4 acres, or 68 percent of land within the CFG boundary in Levy County, is considered  
39 altered land use, which includes the Bypass Canal and Lock. The primary land feature within  
40 Levy County is Inglis Island, which is approximately 1,200 acres and is located between Inglis

1 Lock and Dam. Inglis Island was developed as part of the construction of the Inglis Lock and  
 2 Barge Canal during the 1960s.

3 **Table 27. Trails and Mileage on the CFG—Levy County**

Trail Name	Length in County (Feet)	Length in County (Miles)
Inglis Bypass	4,565.43	0.9
Inglis Island	52,432.90	9.9
North Canal	6,965.09	1.3
<b>Total</b>		<b>12.1</b>

4

5 Inglis Island contains 10 miles of trails (2.1 miles of paved trails and 7.9 miles of unpaved  
 6 trails) used for hiking, mountain biking, and equestrian uses. These trails are featured as  
 7 locations along the GFBWT and offers excellent views of the island’s natural communities,  
 8 which include primarily mesic flatwoods and mesic hammocks.

9 **Table 28. Natural Communities and Acreage on the CFG—Levy County**

Community	Acreage	Percent Total
Salt Marsh	4.5	.002
Blackwater Stream	3.1	.001
Altered Lands	1,539.4	68
Dome Swamp	112.6	5
Floodplain Swamp	123.1	5
Hydric Hammock	0.2	0
Mesic Flatwoods	185.3	8
Mesic Hammock	179.1	8
Sandhill	12.8	.006
Wet Flatwoods	102.3	5
<b>Total</b>	<b>2,262.1</b>	<b>100</b>

10 Key: Altered land use includes abandoned fields, canals/ditches, clearing, developed areas, impoundment/  
 11 artificial pond, pine plantation, roads, spoil areas, and utility corridors.

12 **Water Area**

13 Similar to Citrus County, the water resources on the CFG within Levy County include a mix of  
 14 manmade and natural resources. These water resources primarily include the Bypass Canal  
 15 and Lake Rousseau—totaling 1,077.9 acres of water resources within Levy County. Lake  
 16 Rousseau provides excellent fishing opportunities for certain species, including bluegill,  
 17 redear sunfish, catfish, black crappie, and largemouth bass. Duck-hunting opportunities occur  
 18 along Lake Rousseau.

19 The CFG, by its very nature, includes structures and impoundments that remain from the  
 20 original construction of the Cross Florida Barge Canal. Included is the 3,400-acre Lake  
 21 Rousseau impoundment, which was formed by the construction of Inglis Dam in 1909 by



1 Florida Power Corporation to create hydropower. Hydropower operations ceased in 1965. In  
2 the 1960s, the U.S. Army Corps of Engineers built the portion of the Cross Florida Barge Canal  
3 between Lake Rousseau and the Gulf of Mexico. The construction of the barge canal  
4 interrupted the natural flow of water from the upper segment of the Withlacoochee River  
5 into the lower segment of the river. The Corps constructed the 8,900-foot long Inglis Bypass  
6 channel to discharge fresh water from Lake Rousseau into the lower Withlacoochee River  
7 segment.

#### 8 Natural Scenery

9 This portion in certain areas of the CFG offers scenic views of Lake Rousseau. This scenic  
10 setting is conducive to nature study, wildlife viewing, and photography. Also, the area within  
11 Inglis Island is considered nice mesic flatwood and mesic hammock for hiking and potential  
12 wildlife viewing.

#### 13 Significant Habitat

14 Inglis Island is considered a significant botanical site located within the Levy County portion  
15 of the CFG. According to FNAI, a population of pinewoods dainties (*Phyllanthus liebmannianus*  
16 *ssp. Platylepis*) was seen both in flower and fruit in openings (road edges) of mesic hammock  
17 on Inglis Island on the north side of the Withlacoochee River in Levy County.

#### 18 Natural Features

19 The mesic hammock and mesic flatwood communities on Inglis Island likely are the most  
20 natural features provided in this portion of the CFG. They provide a setting for a variety of  
21 recreational activities, including hiking, nature study, wildlife viewing, photography,  
22 picnicking, and the interpretation of natural and cultural resources.

#### 23 Archaeological and Historical Features

24 Only a handful of previously recorded archaeological sites exist on the CFG within Levy  
25 County. The nature of these resources is unknown and SHPO has not determined the  
26 significance of these resources.

#### 27 *Assessment of Use*

28 All legal boundaries, significant natural features, structures, facilities, roads, and trails  
29 existing in the unit are delineated on the base map (see Base Map). Specific uses made of the  
30 unit are briefly described in the following sections.


#### 31 Current Recreation Use and Visitor Programs

32 The following recreational activities occur along the CFG within Levy County.

33 The *Inglis Island Trails* are located in the CFG's west region. Inglis Island was developed as  
34 part of the construction of the Inglis Lock and Barge Canal during the 1960s. This island,  
35 approximately 1,200 acres, is situated between Inglis Lock and Dam. The recreation area at  
36 the dam provides access to the trails. The trails offer views of the island's natural  
37 communities, which include cypress swamp, pine flatwoods, and mixed hardwood  
38 hammocks.

- 1 Other Uses
- 2 FWC facility and boat launch provide access to Lake Rousseau.
- 3 Protected Zones
- 4 There are no known protected zones within the Levy County portion of the CFG.


1 **Figure 9. Levy County Generalized Existing Land Use**



2



1 **Figure 10. Levy County Generalized Future Land Use**



2



# 1 Marion County

## 2 History/Setting


3 Marion County is considered the southernmost county  
4 in North Central Florida, and the northernmost county  
5 in Central Florida between the Atlantic Ocean and the  
6 Gulf of Mexico. Marion County generally is comprised  
7 of rolling hills, some high and some low. Contributing  
8 to Marion County’s appeal is that it is within a two-  
9 hour drive from many of Florida’s major cities. Orlando  
10 is 75 minutes to the southeast, while Daytona Beach is  
11 about 90 minutes to the east. Tampa is about 75  
12 minutes to the southwest. Jacksonville is roughly a  
13 two-hour drive northeast.

14 The county seat is Ocala, Florida. Marion County  
15 occupies 1,057,280 acres of land with an additional  
16 53,120 acres of water. Marion County ranks as the fifth  
17 largest in size of Florida counties. Approximately  
18 276,000 acres of land within Marion County is  
19 dedicated to the Ocala National Forest. Two additional  
20 parks—Silver Springs and Rainbow Springs—  
21 comprise 5,686 acres of open space.

22 Numerous lakes and rivers—including the Ocklawaha  
23 River, Rainbow River, Silver River, and Withlacoochee  
24 River, Lake Weir, Lake Kerr, and other significant  
25 waterbodies—provide opportunities for fishing,  
26 boating, swimming, tubing, and snorkeling. Major  
27 roadways that provide routes to Marion County  
28 include I-75, United States Highways 27, 41, 301, and  
29 441, and State Roads 40, 200, and 326. At some point,  
30 the roads converge in Ocala, which is centrally located  
31 within the county.

## 32 Population/Demographics

33 Marion County is Florida’s 17th most populous county.  
34 According to the 2010 U.S. Census, Marion County had  
35 a total population of 331,303. This is a 28-percent  
36 increase from the 2000 U.S. Census. Using the BEBR  
37 data, by the year 2025, the estimated population of  
38 Marion County is expected to reach 401,110. Table 29,  
39 below, provides estimates until the year 2045.



### **Land Area:**

Total: 1,663 square miles (4,307 km<sup>2</sup>)  
Land: 1,585 square miles (4,105 km<sup>2</sup>)  
Water: 78 square miles (202 km<sup>2</sup>)


### **Population:**

40,801 (2016)

**Median Age:** 47 years old

### **Population Density:**

218 people per square mile





1 Overall, Marion County is 69 percent urban and 31 percent rural. As such, Marion County  
 2 represents the most urban county that the CFG traverses. This is demonstrated by a higher  
 3 population density of 218.2 persons per square mile, which is much lower than the state  
 4 average population density of 375.7 people per square mile, but is much higher than the  
 5 national average density of 82.73 people per square mile. As shown in Table 30, between  
 6 2010 and 2016, the population of Marion County increased at an average annual rate of 4.4  
 7 percent, which was lower than the rate of growth recorded throughout Florida (7.2 percent  
 8 per year), but higher than the national average of 0.9 percent per year. According to the  
 9 University of Florida, BEBR (2015), over the next 30 years, Marion County’s population is  
 10 projected to grow steadily, reaching 495,600 people by the year 2045.

11 **Table 29. Historic, Current, and Projected Population through 2045—Marion County**

	2015	2020	2025	2030	2035	2040	2045
Low	341,205	352,600	365,600	378,000	388,300	396,800	403,000
Medium		372,300	401,100	427,100	451,400	474,400	495,600
High		387,700	427,600	468,000	509,100	551,200	593,300
State of Florida*	19,815,183	21,372,200	22,799,500	24,071,000	25,212,400	26,252,100	27,217,600

12 Source: University of Florida, Bureau of Economic and Business Research, 2015.

13 \*Used medium population projections.

14  
 15 **Table 30. Population Change—Marion County**

	1980-1990 Percent Change	1990-2000 Percent Change	2000-2010 Percent Change	2010-2016 Percent Change	2015-2020 Percent Change	2020-2025 Percent Change
Marion County	59.1	32.9	28.0	4.4	5.2	5.4
State of Florida	32.7	23.5	17.6	7.2	6.1	6.7

16

17 In 2015, 42 percent of residents of Marion County were age 55 and older compared to 30.5  
 18 percent of residents of Florida, and 21.1 percent of the resident population of the U.S. Further,  
 19 in 2015, the median age of residents of Marion County was 47 years old, which was higher  
 20 than the median age of residents of Florida (41.6 years old), and considerably higher than the  
 21 median age of residents of the U.S. (37.4 years old). Table 31, below, provides a breakdown  
 22 of the population in Marion County by age from 2010 to 2045.

1 **Table 31. Population by Age—Marion County**

Age	Census	Estimates	Projections					
	2010	2015	2020	2025	2030	2035	2040	2045
0-4	17,112	17,433	18,559	19,850	20,644	21,508	22,269	23,362
5-17	47,070	45,932	49,228	52,236	55,123	57,868	60,399	62,467
18-24	23,743	24,134	23,624	24,710	26,264	27,631	29,053	30,554
25-54	112,059	109,538	114,439	119,439	125,591	133,433	138,607	143,852
55-64	46,001	49,602	57,368	57,496	54,109	52,882	57,444	64,114
65-79	63,856	70,897	80,813	92,931	104,093	110,706	109,249	105,113
80+	21,462	23,669	28,227	34,448	41,305	47,339	57,341	66,148
<b>Total</b>	<b>331,303</b>	<b>341,205</b>	<b>372,258</b>	<b>401,110</b>	<b>427,129</b>	<b>451,427</b>	<b>474,362</b>	<b>495,610</b>

2 Source: University of Florida, Bureau of Economic and Business Research, 2015.

3

4 Over the long-term, projections for Florida indicate that the state’s population will increase  
 5 at an average annual rate of 2.1 percent between 2020 and 2030, reaching 28.7 million people  
 6 in 2030. This rate of growth is significantly higher than the rate of increase forecast  
 7 throughout the U.S. (0.8 percent per year), reflecting long-term growth rates in the state.

8 **Existing Use of Adjacent Lands**

9 Marion County, an inland county centered around Ocala, is easily accessible by several  
 10 interstate highways. Intersected by I-75, Florida State Highways 19, 25, 35, 40, 200, 326, and  
 11 464 also provide access to the county and link residents with several major cities, including  
 12 Orlando, Tampa, and Gainesville.

13 Within Marion County, the CFG is within proximity of some populated areas, including  
 14 Dunnellon, Chatmire, Huntington, Silver Springs, and Orange Springs. The CFG intersects the  
 15 south/southwestern portion of Marion County beginning at Lake Rousseau and the  
 16 Withlacoochee River before proceeding on an east/northeast direction toward Orange  
 17 Springs and Putnam County. In all, approximately 331,298 people live in Marion County (U.S.  
 18 Census 2010) and are within 30 miles of the CFG.

19 Inside the one-mile buffer area of the CFG in Marion County, there are a variety of existing  
 20 land uses. The most common include Agriculture (36 percent), Public/Institutional (29  
 21 percent), Residential (20 percent), Recreation (5 percent), and Other (4 percent). Agriculture,  
 22 Residential, Recreation, and Other land uses are scattered throughout the study area.  
 23 Public/Institutional land use includes the Ocala National Forest, Halpata Tastanaki Preserve,  
 24 and Silver Springs State Park, among other areas.

25 Figure 11 and Figure 12, below, display the generalized existing land use for Marion County  
 26 within one mile of the CFG.

1 **Planned Use of Adjacent Lands**

2 Marion County, composed of rolling hills with an abundance of farmland and forests, has  
3 many attractive qualities. As indicated earlier, travel to and from this area is easy given the  
4 abundant roadway access. The proximity to both the Gulf of Mexico and the Atlantic coast  
5 make it attractive for future development.

6 The area surrounding the CFG also has some future planned development. West of Dunnellon,  
7 just north and south of the CFG along SR 200, there are several planned unit developments  
8 (PUDs), including Mixed Use. Per the Marion County Planning Department, to date, this is the  
9 only area near the CFG for which new development has been planned. Marion County is a  
10 member of the Ocala/Marion County Transportation Planning Organization (TPO). The  
11 *Ocala/Marion County 2040 Long Range Transportation Plan* indicated there are four roadway  
12 projects within the vicinity of the CFG in Marion County. These projects are in various phases  
13 of project development and details are provided below in Table 32.

14 **Table 32. Roadway Projects—Marion County**

Project Name	Project Description	Funding Timeframe
SR 200 from Citrus County Line to CR 484	Add 2 lanes	2021-2030
SW 49 <sup>th</sup> Avenue from SW 95 <sup>th</sup> Street to Marion Oaks Trail	Add 2 lanes	2026-2040
SR 40 from NE 60 <sup>th</sup> Court to CR 314	Add 2 lanes	2016-2020
SR 35 from SE 92 <sup>nd</sup> Place to CR 464	Add 2 lanes	2016-2018

15 Source: Ocala/Marion County 2040 Long Range Transportation Plan

16 It should be noted that CFG staff will coordinate with all entitifes to include, at a minimum,  
17 double off-grade road crossings for wildlife and recreational connectivity when roads are  
18 expanded from two lanes to four lanes or more. This also would include a more  
19 comprehensive consideration of impacts and how to best mitigate and/or compensate for  
20 those potential impacts during roadway planning, design, and construction.

21 Inside the one-mile buffer area of the CFG in Marion County, several changes can be noticed  
22 from existing land use to future land use. These changes include a substantial increase in  
23 Agriculture (44 percent) and Conservation (31 percent). The increase in Conservation is  
24 mostly due to the County’s land use re-classification. Areas such as the CFG, Ocala National  
25 Forest, Halpata Tastanaki Preserve, and Silver Springs State Park were previously classified  
26 as Public/Institutional or Recreation under existing land use and are now classified as  
27 Conservation under future land use. Despite PUDs, Residential land use is expected to  
28 decrease (16 percent) across most of the CFG area. Commercial land use increases slightly,  
29 including in the areas around SR 200 and US 301.

1 Figure 13 and Figure 14, below, display the generalized future land use for Marion County  
2 within one mile of the CFG.

3 **Property Analysis**

4 As indicated previously, in an attempt to better understand and detail the specific recreation  
5 resource elements, this plan describes these resources by county. Effective planning requires  
6 a thorough understanding of the unit's natural and cultural resources. This section describes  
7 the resource characteristics and existing uses of the property. The unit's recreation resource  
8 elements are examined to identify the opportunities and constraints they present for  
9 recreational development. Past and present uses are assessed for their effects on the  
10 property, compatibility with the site, and relation to the unit's classification.

11 *Recreation Resource Elements*

12 This section of the Land Use Component assesses the park's recreational resource elements,  
13 those physical qualities that—either singly or in certain combinations—can support various  
14 resource-based recreation activities. Breaking down the property into such elements  
15 provides a means for measuring the property's capability to support potential recreational  
16 activities. This process also analyzes the existing spatial factors that either favor or limit the  
17 provision of each activity.

18 **Land Area**

19 Overall, Marion County contains 36,594 acres—the largest county traversed by the CFG. In  
20 total, it contains more than 200 miles of trails, as shown in Table 33, vastly out-distancing the  
21 trail mileage in the other three counties combined. As illustrated in Table 34, a total of  
22 10,614.7 acres, or 29 percent of land within the CFG boundary in Marion County, is  
23 considered altered land use. Land within the Marion County portion of the CFG is dominated  
24 by basin swamp, floodplain swamp, hydric hammock, mesic flatwoods, mesic hammock,  
25 sandhills, and scrub.

26 ***Table 33. Trails and Mileage on CFG—Marion County***

<b>Trail Name</b>	<b>Length in County (Feet)</b>	<b>Length in County (Miles)</b>
110th Street	2,004.70	0.4
25th Crossing	172.31	0.0
25th Street ER Access	856.87	0.2
49th Avenue	22,188.41	4.2
Anthill	1,212.66	0.2
Backdoor	514.78	0.1
Baseline Park Trail	25,103.05	4.8
Beyond	15,781.85	3.0
Blue Highway	1,960.75	0.4
Bridge Station Trail	2,533.59	0.5
Bunny	6,649.22	1.3

<b>Trail Name</b>	<b>Length in County (Feet)</b>	<b>Length in County (Miles)</b>
Cactus Jack Trail	1,493.60	0.3
Canopy	2,093.17	0.4
Cow Bone	6,202.92	1.2
Coyote Corner	684.76	0.1
Coyote Corner North	4,167.21	0.8
Coyote Corner South	3,070.30	0.6
Dog Bone	11,065.20	2.1
Dog Tail	2,757.83	0.5
Dr Ruth	5,535.31	1.0
Dunnellon Trail	8,069.36	1.5
Equestrian Access Trail	9,966.17	1.9
Ern N Burn	14,184.76	2.7
Florida Trail	191,929.80	36.4
Florida Trail Connector	2,321.20	0.4
Hiking/Biking	6,345.03	1.2
Hiking/Biking	6,345.03	1.2
Horse Park Connector	1,557.79	0.3
Horse Park Trail	11,940.98	2.3
John Brown	2,641.92	0.5
John Frank Access Trail	4,763.06	0.9
LandBridge	1,527.94	0.3
LandBridge North	15,710.02	3.0
LandBridge South	6,350.42	1.2
Limerock Service Road	81,651.77	15.5
Magic Mountain	472.58	0.1
Marshmallow	2,057.25	0.4
Nayls North	28,782.97	5.5
Nayls South	18,266.46	3.5
Ocklawaha Visitor Center	7,784.29	1.5
Pine Tree	16,558.96	3.1
Puppy Loop	2,486.80	0.5
Rattlesnake Ridge	1,767.03	0.3
Ross Prairie	5,707.69	1.1



Trail Name	Length in County (Feet)	Length in County (Miles)
Shangi-La	4,689.75	0.9
Shangri-La Day Use	7,866.76	1.5
Shorty	1,063.95	0.2
Sinkhole	4,622.23	0.9
Speedway	2,848.42	0.5
Spider Kingdom North	16,841.06	3.2
Spider Kingdom South	13,086.28	2.5
Trail #1	108,876.57	20.6
Trail #2	95,437.65	18.1
Trail #3	138,440.56	26.2
Trail #4	38,758.66	7.3
Tricycle	37,981.30	7.2
Twister	13,719.15	2.6
Vortex	2,756.79	0.5
Vortex Trail	7,256.54	1.4
<b>Total</b>		<b>201.7</b>

1

2 **Table 34. Natural Communities and Acreage on CGF—Marion County**

Community	Acreage	Percent Total
Basin Marsh	13.6	0
Basin Swamp	3,376.2	9
Baygall	304.7	0.008
Blackwater Stream	145.9	0.004
Bottomland Forest	163.2	0.005
Altered Lands	10,614.7	29
Depression Marsh	495.4	1
Dome Swamp	5.2	0
Floodplain Swamp	6,540.2	18
Hydric Hammock	1,367.0	4
Mesic Flatwoods	1,472.8	4
Mesic Hammock	3,365.1	9
Sandhill	6,203.4	17
Scrub	1,244.0	3
Scrubby Flatwoods	226.7	0.006
Swamp Lake	28.3	0
Upland Hardwood Forest	243.9	0.007

Community	Acreage	Percent Total
Upland Pine Forest	6.4	0
Wet Flatwoods	588.0	2
Xeric Hammock	189.3	0.005
<b>Total</b>	<b>36,594</b>	<b>100</b>

**Key:** Altered land use includes abandoned fields, canals/ditches, clearings, developed areas, impoundment/artificial pond, pine plantation, roads, spoil areas, and utility corridors.

1  
2  
3

4 The westernmost portion of the CFG within Marion County is dominated by wet flatwoods  
5 and floodplain swamp associated with the easternmost portion of Lake Rousseau and  
6 impoundment. Moving east across US 41, near Dunnellon, the CFG is characterized by a  
7 predominantly altered landscape, with higher-intensity uses such as the Dunnellon Ballfields  
8 and the Wastewater Treatment Plant. Higher-intensity land uses should be shifted out of the  
9 main CFG corridor.

10 Further, this area includes the highest number of trails given that there are no major  
11 waterbodies. Trails include 1.5 miles of the Dunnellon Trail and the Florida Trail (37 miles  
12 within Marion County). The Dunnellon multi-use trail is primarily located along a historic  
13 railroad bed that runs along the Rainbow River south of CR 484 and the Blue Run Park and  
14 tubers take out. The trailhead is located along Bridges Road on the Marion County side. The  
15 trailhead provides access to this popular trail, which provides scenic views of the  
16 Withlacoochee River from the trail bridge, which connects Marion and Citrus Counties  
17 together. The trail bridge was designed to capture the look of the former historic railroad  
18 bridges that were built in the late 1800s when phosphate was discovered in Dunnellon and it  
19 enjoyed its “Boomtown” days that are still celebrated annually. Throughout this portion of  
20 the CFG, consideration should be given to possible interconnectivity with existing  
21 surrounding trails.

22 Continuing east, the CFG begins to traverse higher, drier communities, mostly consisting of  
23 sandhills. Of note within this section of the CFG is the historic “Diggings,” which total 2,946.5  
24 acres. This region is dominated by longleaf pine sandhill and sand pine scrub natural  
25 communities. These 1930s sea-level ship-to-canal project canal diggings serve as remnants  
26 of the former canal project. These features are wide linear areas that were dug out of the  
27 landscape to create the canal. Today, these areas exist as reforested, small-scale valleys.  
28 However, because the bottoms of “diggings” are comprised of clayey soils that hold moisture,  
29 they have become dominated by loblolly pines that prefer wetter areas than longleaf or sand  
30 pines. In many areas, they also offer challenging terrain for trail goers.

31 In addition, this portion of the CFG offers excellent opportunities for wildlife viewing as well  
32 as cultural interpretation. Within this section, located along the southern boundary of the CFG  
33 along the east side of SR 200 and adjacent to the entrance of Ross Prairie State Forest is the  
34 Ross Prairie Trailhead and Campground. Ross Prairie is a unique ecological feature due to the  
35 adjacent placement of the high, dry sandhill community (where the trailhead and  
36 campground are located) and the ephemeral wetland Ross Prairie (located just north of the

1 public use facilities). Specifically, this area contains 239 acres of depression marsh that serves  
2 as a prairie-type landscape within a predominantly longleaf pine forested area of the CFG.  
3 Moving west to east, the elevation continues to increase and contains outstanding hardwood  
4 live oak hammocks in between Pruitt and SR 200 and oak islands in Ross Prairie. Additionally,  
5 this area contains approximately 1,000 acres of scrub immediately west of I-75.

6 This area has undergone significant restoration activities and contains a number of breeding  
7 pairs of Scrub-Jays. CFG has partnered with the Florida Audobon Society to monitor and track  
8 Scrub-Jays. This area is considered to be a protected area of the CFG. This area also contains  
9 Shangri-La and Pruitt Trailheads. As indicated in Table 33, this section of the CFG contains a  
10 significant trail network, including equestrian, hiking, and biking trails. Equestrian trails are  
11 predominantly located in the southernmost portion of the CFG, while hiking and biking trails  
12 are to the north.

13 Moving across I-75, equestrian, hiking, and biking trails continue up to the Santos Trailhead  
14 and camping area. The Santos Trail is widely recognized for its mountain biking trails and  
15 other trail networks—in particular, the winding trails through the rock quarry country,  
16 which are a favorite of the mountain bike community. This portion of the CFG begins to move  
17 through more populated areas of Marion County and, as such, begins to have more  
18 altered/developed types of land uses, with a higher number of road crossings. Baseline Road  
19 Trailhead is located within this area, which is a community park with trails and is sub-leased  
20 to Marion County for management.

21 Marshall Swamp is the next portion of the CFG and contains primarily mesic hammock, hydric  
22 hammock, and floodplain swamp. Moving from developed to undeveloped land, this area is  
23 adjacent to the Ocala National Forest and Silver Springs. Following Marshall Swamp, this area  
24 continues along the Ocklawaha River, incorporating floodplain swamp, mesic hammock, and  
25 other wet communities until the Marion County boundary. This area along the Ocklawaha  
26 River also provides paddling, fishing, and boating opportunities, scenic views, wildlife  
27 watching, and interspersed boat landings.

## 28 Water Area

29 As indicated previously, the westernmost portion of the CFG within Marion County contains  
30 the eastern extent of Lake Rousseau. From Marshall Swamp to the Marion County boundary,  
31 the CFG contains the Ocklawaha River and its floodplain.

## 32 Natural Scenery

33 In general, the highest quality sandhills are located within the Marion County portion of the  
34 CFG. In the western portion of the CFG through Marion County, wildlife viewing opportunities  
35 and scenic views are offered along the Dunnellon multi-use trail. Farther to the east within  
36 Marion County, the Ross Prairie area provides excellent examples of unique prairie-type  
37 communities within the CFG. This habitat is conducive to wildlife viewing and hiking.

38 Continuing east within Marion County, the “Diggings” sites—although significantly impacted  
39 by previous canal digging activities—continue to provide wildlife viewing opportunities.

1 Additionally, given the historic nature of these sites, they also provide significant cultural  
2 interpretation opportunities.

3 A large portion of the CFG within Marion County contains the Ocklawaha River and its  
4 floodplain communities. Within this portion of the CFG, there is little to no development, with  
5 only a few adjacent publicly owned lands. As such, this area also provides significant  
6 opportunities to view natural scenery and wildlife. Further, Marshall Swamp Trail and  
7 Ocklawaha River floodplain in the Sharpes Ferry area offer high-quality hydric hammock and  
8 bottomland forest in which several rare plants are known to occur.

#### 9 Significant Habitat

10 There is a total of 1,244 acres of scrub habitat within this portion of the CFG. This community  
11 is a significant habitat given that it is home to the endangered Florida Scrub-Jay. This habitat  
12 is a primary restoration activity on the CFG. Further, the CFG works closely with the Florida  
13 Audobon Society to monitor and count nesting Scrub-Jays. These areas also are considered  
14 protected throughout the CFG and offer passive recreational opportunities.

15 FNAI identified several areas throughout the CFG as significant botanical sites. This includes  
16 a linear east/west strip beginning with the Diggings west of SR 200 east to I-75, in which  
17 seven listed plants were identified in sandhill, scrub, and successional hardwoods that  
18 surround and include the Diggings. Scrub and sandhill (less so) also harbor the population  
19 stronghold of the federal and state listed endangered long-spurred mint.

#### 20 Natural Features

21 Ross Prairie is the most unique natural feature within this section of the CFG. Having the  
22 sandhill and wetland prairie systems adjacent to each other helps to provide habitat to a wide  
23 range of species. The habitat changes with the seasons and water volume of the prairie; when  
24 wet, dozens of waterfowl avian species, wading birds, reptiles, amphibians and mammals can  
25 be spotted.

#### 26 Archaeological and Historical Features

27 The previously recorded archaeological sites and historic features in the Marion County  
28 section of the CFG consist of a variety of prehistoric and historic period archaeological sites.  
29 These represent an excellent cross section of Central Florida's prehistoric past. The exact  
30 nature of the historic sites are unclear. They represent historic artifact scatters, historic  
31 dumps, or possible historic home sites. There are also several historic railway corridors,  
32 bridges, and the Eureka Lock and Dam Complex in this section of the park. Florida SHPO has  
33 not determined the significance of the vast majority of these sites. Six archaeological sites  
34 (8MR1878, 8MR2351, 8MR3863, 8MR3865, and 8MR3866), the Eureka Lock and Dam  
35 Complex (8MR3563), a linear resource (8MR3410), and a historic bridge (8MR3585) have  
36 been listed or determined eligible of listing on the NRHP.

#### 37 *Assessment of Use*

38 All legal boundaries, significant natural features, structures, facilities, roads, and trails  
39 existing in the unit are delineated on the base map (see Base Map). Specific uses made of the  
40 unit are briefly described in the following sections.

1 Current Recreation Use and Visitor Programs

2 As mentioned above, Marion County is home to a large portion of the CFG. In fact, Marion  
3 County works closely with the CFG staff and operates several trailheads along the greenway,  
4 providing unlimited recreational access. The Ocala area boasts numerous multi-use trails.  
5 The Santos Bike Trails, winding through rock quarry country, are a favorite of mountain  
6 bikers. Below is a discussion of the recreational facilities located along the CFG within Marion  
7 County.

8 *Dunn-Bridges Road Trailhead* features a 2.4 mile, paved multi-use trail that is located along  
9 former pastures that are being restored to the original longleaf pine sandhill community. This  
10 trailhead, managed by Marion County Parks and Recreation, offers parking and amenities and  
11 a barrier-free Boundless Playground.



*Ross Prairie Trailhead and Campground* is located along the southern boundary of the CFG along the east side of SR 200 and adjacent to the entrance to Ross Prairie State Forest. This area is ecologically unique due to the adjacent placement of the high, dry sandhill community (where the trailhead and campground are located) and the ephemeral wetland, Ross Prairie (located just north of the public use facilities).

22 The day-use trailhead facilities include an equestrian staging area and bike and hike  
23 trailheads for the natural surface mountain bike and Florida Trail hiking trails. There are  
24 restrooms with potable water and a picnic pavilion as well. The Ross Prairie Campground has  
25 14 public-use campsites with water and 30/50 amp hookups available. An on-site  
26 campground host is available 365 days a year.

27 The trails in this area meander through beautiful oak hammocks and islands interspersed  
28 around the edge of Ross Prairie. Wildlife in this area is abundant, with whitetail deer, wild  
29 turkey, bobcat, alligators in the prairie, gopher tortoises in the sandhills, and numerous other  
species to observe. This part of the Greenway also is conveniently located within two miles of major shopping and dining opportunities just north of the CR 484 and SR 200 intersection.



*Shangri-La Trailhead and Campground* is located along the southern boundary of the CFG in the sandhills community five miles west of I-75 off the CR 484 exit. Shangri-La is just south of one of the historic sea-level Cross Florida Ship Canal diggings. Shangri-La provides optimal access and overnight opportunities to explore the myriad network of well-marked trails through



1 xeric hammock. Imagine what it must have been like during the Great Depression when  
2 thousands of workers armed with picks and shovels worked incessantly to dig toward the  
3 bedrock and water table below to facilitate a canal below sea-level. Now, 80 years later, the  
4 scars have healed, but the excavated berms still stand as reminders of hardships and  
5 endeavors long gone. Towering pines, oaks and mixed hardwoods now cover the lands and  
6 provide habitat for whitetail deer, fox, squirrels, wild turkey, bobcat, and more.

7 Shangri-La has an ample day-use equestrian and bike trailhead with a picnic pavilion and  
8 restrooms with potable water. The campground has 24 campsites available. The compacted  
9 shell sites have communal water and a full bathhouse facility. The sites do not have power;  
10 however, generators are acceptable to use except during quiet hours. There are also two  
11 covered picnic pavilions in the campground. Campground hosts are on-site 365 days a year.  
12 The campground also has nearby supply shops as well as numerous restaurants within only  
13 a couple of miles.

14 *49th Avenue Trailhead* is located along SW 49th Avenue within the CFG. The trailhead  
15 contains only dirt parking, has no potable water, and only has portable toilets, but there are  
16 covered picnic tables at this facility. It is also shared with equestrian users and is at the east  
17 end of the Nays Trail.



*The LandBridge Trailhead* is the primary trailhead developed and located to provide access to the “First of Its Kind in the U.S.” I-75 Landbridge. The structure, which is 52 feet wide by 200 feet long, follows a natural ridge over 100 feet in elevation to minimize ecological damage. The planters on both sides of the bridge are vegetated with plants native to the surrounding area. The walls were built from local fieldstone.

28 Located on the bridge, wildlife cameras—monitored  
29 by Greenway staff—have captured photos of numerous crossings of the bridge by bobcats,  
coyotes, wild turkeys, and even Florida black bears.

30 The trailhead is located approximately 1.4 miles east of the LandBridge off County Road 475A  
31 and is accessible from I-75 via the CR 484 exit. Facilities include restrooms with potable  
32 water, a beautiful shaded picnic area along with equine, hiking, natural surface bike trails,  
33 and paved trail (under design/construction). In addition, a concessionaire offers guided  
34 equestrian horseback rides along the Greenway’s shaded trails, including rides over the  
35 LandBridge.



*Santos Trailhead* often is referred to as a “Mecca” of mountain biking throughout Florida, the southeastern U.S., and even internationally. In fact, Santos has been dedicated as a bronze-level Ride Center by the International Mountain Bike Association (IMBA). The Santos Trails includes 80+ miles of single-track bike trails and includes trails for beginners as well as expert-level riders that will challenge even the most elite riders. Santos is perennially rated as one of the best mountain bike facilities in existence. Mountain biking trails are developed

13 and maintained through a partnership with the OMBA volunteer organization. Numerous  
14 wooden “features” and the Vortex Freeride area located in a former limerock quarry provide  
15 extreme terrain in the flatlands of Florida.

16 The Santos Trailhead has restrooms, potable water, a bike wash rack, and several picnic  
17 pavilions. The trailhead and campground are located just west off US 441 on SE 80th Street,  
18 between Ocala and Belleview, across from the Marion County Sheriff’s Santos Station. Also  
19 available at the Santos Trailhead is access to the CFG’s extensive equestrian trail network—  
20 with 90+ miles of horse trails, including varying levels of difficulty, and wagon/carriage  
21 opportunities. Santos Trailhead serves as the eastern end of equestrian trails on the  
22 Greenway and is adjacent to the Santos Campground. Santos campground contains six  
23 primitive campsites for horse camping; four have poles for overhead picket lines. The first  
24 campsite has a pen that will hold up to two horses. Trails are marked for equestrian, wagon,  
25 or carriage, and some multi-use trails run west to the Pruitt Trailhead. The Santos Trailhead  
26 campground area offers equestrian parking and a fenced equestrian staging area. Trails in  
27 this area are generally flat and sandy, and much of the trail runs through the woods. There  
28 are approximately 1.5 miles of fireline roads accessible to riding and driving and two miles  
29 of shaded singletrack for horseback riders only.

30 The Florida National Scenic Trail (FNST) hiking trail also traverses through the Santos  
31 Trailhead and runs westward to Dunnellon from Santos for approximately 24 miles and  
32 eastward 10+ miles before entering the Ocala National Forest. As one of only 11 National  
33 Scenic Trails nationwide (including the Pacific Crest Trail and Appalachian Trail), this  
34 congressionally designated trail runs for approximately 1,300 miles through diverse  
35 ecological communities throughout Florida, from the Gulf National Seashore near Pensacola  
36 Beach to the Fakahatchee Strand in southwest Florida. The segment from Santos west to  
37 Dunnellon goes through and along several of the Historic Sea-Level Cross Florida Ship Canal  
38 project diggings with 40+ foot berms measuring from one-quarter mile to two miles long.  
39 Adjacent to the Santos Trailhead and its diverse trail offerings is the Santos Campground. It  
40 features 23 full hookup sites available year-round for overnight trips.

1 The *Baseline Road to Marshall Swamp* area of the CFG offers two trails, providing  
2 opportunities to experience a paved urban trail or escape to a secluded multi-use trail that



weaves through the cypress swamp and hardwood hammock communities of Marshall Swamp. Access to both is just minutes from downtown Ocala and Silver Springs. The Greenway's SE 64th Avenue Trailhead provides parking and access to the paved and unpaved trails where they connect, offering an easy opportunity to experience two very different types of trails.

13 The Baseline Road Trailhead features a five-mile,  
14 paved, multi-use trail, which is situated along former pastures that are being restored to the  
15 original longleaf pine sandhill community. This trailhead, managed by Marion County Parks  
16 and Recreation, offers parking and amenities and, thanks to the generosity of the Felburn  
17 Foundation, a barrier-free, Boundless Playground®, designed to enable children of all abilities  
to learn and play freely together.

18 The Marshall Swamp Trail can be accessed from the Marshall Swamp Trailhead, which is  
19 adjacent to the CFG's Sharpe's Ferry Office. This unpaved trail provides access to a segment  
20 of the FNST as it runs nearly three miles through Marshall Swamp, a jungle-like, hydric  
21 hammock floodplain of the Ocklawaha and Silver rivers.

22 *Gores Landing* is in northeast Marion County and offers primitive camping, a boat ramp,  
23 restrooms, and picnic tables on the Ocklawaha River and provides southern access to the  
24 Gores Landing WMA. Camping is available on a first-come, first-served basis. This state-  
25 owned property is leased to the Marion County Board of County Commissioners, who have  
26 managed the facility since 1966. In addition to fishing, the Gore's Landing area offers 52 days  
27 of hunting each year, in accordance with WMA regulations and rules per FWC, with the same  
28 species sought as at Caravelle.

29 *Eureka East and West Boat Ramps* are in northeast Marion County on the Ocklawaha River  
30 and both include a boat ramp and picnic pavillion, while Eureka West also contains a fishing  
31 dock. This state-owned property is leased to the Marion County Board of County  
32 Commissioners, who have managed the facility since 1977.

33 The *Orange Springs Boat Ramp* provides paddle and watercraft access to the Ocklawaha  
34 River, as well as a picnic pavilion. Marion County has leased the boat ramp from the state  
35 since 1993.


### 36 Protected Zones

37 There are 15 occurrences of longspurred mint in Marion and Sumter Counties, of which six  
38 of these populations are on the CFG. The plant has been extirpated from several sites in these  
39 counties. The CFG is the only conservation land that supports a mint population. These areas  
40 will be considered protected zones. The primary protected area within the Marion County

1 portion of the CFG is the scrub habitat immediately adjacent to I-75. CFG staff have  
2 contributed significant resources to the restoration of this area. The total area is 1,088 acres  
3 and there are numerous nesting Scrub-Jays within the area.



1 **Figure 11. Marion County Generalized Existing Land Use, Page 1**



### Marion County Generalized Existing Land Use

**Legend**

- 1-mile Buffer Area
- Greenway Area
- County Boundary

Generalized Existing Land Use (FDOR et al.) 112,060.59 acres

- Agriculture (35.68%)
- Public/Institutional (28.51%)
- Residential (20.19%)
- Recreation (5.28%)
- Other (4.27%)
- Non-agricultural Acreage (3.35%)


- Commercial (1.4%)
- Infrastructure (1.02%)
- Industrial (0.31%)

**Generalized Existing Land Use**

- Agriculture
- Commercial
- Industrial
- Infrastructure
- Public/Institutional
- Recreation
- Residential
- Non-agricultural Acreage
- Other

**Marion County Existing Land Use**


- Agricultural
- Retail/Office
- Industrial, Mining
- ROW
- Public/Semi-Public, Institutional
- Recreation
- Residential, Vacant Residential
- Acreage Not Zoned For Agriculture
- Other, Vacant/Nonresidential, Parcels With No Values, No Data Available, Centrally Assessed



2




1 **Figure 12. Marion County Generalized Existing Land Use, Page 2**



2




1 **Figure 13. Marion County Generalized Future Land Use, Page 1**



2



1 **Figure 14. Marion County Generalized Future Land Use, Page 2**



2

# 1 Putnam County

## 2 History/Setting

3 Putnam County was created in 1849. It was Florida's  
4 28th county created from parts of St. Johns, Alachua,  
5 Orange, Duval, and Marion. The county was named for  
6 Benjamin A. Putnam, who was a soldier in the First  
7 Seminole War, a lawyer, a Florida legislator, and the  
8 first president of the Florida Historical Society. The  
9 Putnam County seat is Palatka.

10 Putnam County is centrally located between  
11 Jacksonville, Gainesville, St. Augustine, and Daytona  
12 Beach. According to the U.S. Census Bureau, the county  
13 has a total area of 827 square miles (2,140 km<sup>2</sup>), of  
14 which 728 square miles (1,890 km<sup>2</sup>) is land and 99  
15 square miles (260 km<sup>2</sup>) is water.

16 Putnam County contains a wealth of important natural  
17 resources, including lakes, creeks, rivers, wetlands,  
18 mineral resources, aquifer recharge areas, and fish and  
19 wildlife. The dominant natural feature in the county is  
20 the St. Johns River, which flows through the eastern  
21 portion of the county and forms the eastern city limits  
22 of Palatka. Putnam County contains approximately 260  
23 lakes of 10 acres or more, plus numerous lakes smaller  
24 in size scattered throughout the county. These lakes  
25 cover an area of approximately 47,220 acres (74  
26 square miles) and are especially numerous in the  
27 western and southeastern portions of the County.

28 The CFG enters Putnam County along the south-central  
29 county line. Kirkpatrick Dam and Rodman Reservoir  
30 form the south-central boundary line of Putnam  
31 County.

## 32 Population/Demographics

33 Putnam County is Florida's 39th most populous  
34 county. According to the 2010 U.S. Census, Putnam  
35 County had a total population of 74,364. This is a 5.6  
36 percent increase from the 2000 U.S. Census. Using  
37 medium population projections prepared by the BEBR,  
38 by the year 2025, the estimated population of Putnam



### **Land Area:**

Total: 827 square miles (2,140 km<sup>2</sup>)  
Land: 728 square miles (1,890 km<sup>2</sup>)  
Water: 99 square miles (260 km<sup>2</sup>)


### **Population:**

72,023 (2016)

**Median Age:** 44 years old

### **Population Density:**

100 people per square mile





1 County is expected to decrease slightly, totaling 73,700. Table 35 below, provides estimates  
 2 through the year 2045.

3 Although, as indicated previously, Putnam County is centrally located between several large  
 4 cities, the county remains very rural in nature with an average population density of 100.3  
 5 persons per square mile, which is significantly lower than the state average population  
 6 density of 375.7 people per square mile, but is slightly higher than the national average  
 7 density of 82.73 people per square mile. Table 36 shows that, between 2010 and 2016, the  
 8 population of Putnam County actually decreased by 1.9 percent, which was lower than the  
 9 rate of growth recorded throughout Florida (7.2 percent per year) and the national average  
 10 of 0.9 percent per year. According to the University of Florida, BRER (2015), over the next 30  
 11 years, Putnam County’s population is projected to grow at a very small rate, reaching 75,500  
 12 people by the year 2045.

13 **Table 35. Historic, Current, and Projected Population through 2045—Putnam County**

	2015	2020	2025	2030	2035	2040	2045
Low		62,000	63,500	65,000	66,500	68,000	69,900
Medium		73,200	73,700	74,200	74,600	75,100	75,500
High		76,900	79,500	82,300	85,000	87,800	90,700
State of Florida*	19,815,183	21,372,200	22,799,500	24,071,000	25,212,400	26,252,100	27,217,600

14 Source: University of Florida, Bureau of Economic and Business Research, 2015.  
 15 \*Used medium population projections.

17 **Table 36. Population Change—Putnam County**

	1980-1990 Percent Change	1990-2000 Percent Change	2000-2010 Percent Change	2010-2016 Percent Change	2015-2020 Percent Change	2020-2025 Percent Change
Putnam County	28.7	8.2	5.6	-1.9	0.4	0.7
State of Florida	32.7	23.5	17.6	7.2	6.1	6.7

18  
 19 In 2015, 36 percent of residents of the county were age 55 and older, compared to 30.5  
 20 percent of residents of Florida, and 21.1 percent of the resident population of the U.S. Further,  
 21 as shown in Table 37 in 2015, the median age of residents of Putnam County was 44.2 years  
 22 old, which was higher than the median age of residents of Florida (41.6 years old), and  
 23 considerably higher than the median age of residents of the U.S. (37.4 years old).



1 **Table 37. Population by Age—Putnam County**

Age	Census	Estimates	Projections					
	2010	2015	2020	2025	2030	2035	2040	2045
0-4	4,689	4,441	4,359	4,366	4,300	4,299	4,298	4,346
5-17	12,096	11,430	11,372	11,318	11,274	11,281	11,301	11,312
18-24	6,061	5,920	5,345	5,289	5,451	5,391	5,426	5,486
25-54	26,748	24,781	24,028	23,869	23,879	24,188	24,123	24,212
55-64	10,700	11,145	11,555	10,434	9,323	9,268	9,724	10,087
65-79	10,670	11,452	12,690	14,057	14,829	14,358	13,469	12,733
80+	3,400	3,587	3,893	4,388	5,126	5,850	6,727	7,342
<b>Total</b>	<b>74,364</b>	<b>72,756</b>	<b>73,242</b>	<b>73,721</b>	<b>74,182</b>	<b>74,635</b>	<b>75,078</b>	<b>75,518</b>

2 Source: University of Florida, Bureau of Economic and Business Research, 2015.

3

4 Over the long-term, projections for Florida indicate that the state’s population will increase  
 5 at an average annual rate of 2.1 percent between 2020 and 2030, reaching 28.7 million people  
 6 in 2030. This rate of growth is significantly higher than the rate of increase forecast  
 7 throughout the U.S. (0.8 percent per annum), reflecting long-term growth rates in the state.

8 **Existing Use of Adjacent Lands**

9 Putnam County, an inland county along the St. Johns River, is easily accessible by several  
 10 interstate highways. Located between I-75 to the west and I-95 to the east, Florida State  
 11 Highways 17, 19, 20, 100, and 207 also provide access to the county and link residents with  
 12 several major cities, including Jacksonville, Gainesville, and Orlando.

13 The CFG within Putnam County is close to some populated areas, including Palatka, East  
 14 Palatka, San Mateo, and Satsuma. The CFG crosses Putnam County from Lake Ocklawaha to  
 15 the St. Johns River. In all, approximately 74,364 people live in Putnam County (U.S. Census  
 16 2010) and are within 20 miles of the CFG.

17 Inside the one-mile buffer area of the CFG in Putnam County, there are a variety of existing  
 18 land uses. The most common include Agriculture (45 percent), Residential (20 percent),  
 19 Recreation (13 percent), Public/Institutional (11 percent), and Non-Agricultural Acreage (9  
 20 percent). Agriculture land uses are predominantly around the northern portion of the CFG,  
 21 Residential land uses are predominantly located near SR 19 and CR 309, while Recreation  
 22 and Public/Institutional land uses are predominantly south of the CFG. Public/Institutional  
 23 land uses include the Ocala National Forest, Murphy Creek Conservation Area, and Seven  
 24 Sisters Islands, among other areas. Non-Agricultural Acreage is predominantly located to the  
 25 north of the CFG around SR 20.

1 Figure 15, below, displays the generalized existing land use for Putnam County within one  
2 mile of the CFG.

### 3 **Planned Use of Adjacent Lands**

4 Putnam County, with an abundance of natural features, also has a variety of appealing  
5 qualities. As indicated earlier, travel to and from this area is easy given the abundant roadway  
6 access. The convenience of the St. Johns River and its proximity to the Gulf of Mexico make it  
7 attractive for future development while being mindful of the threat of inland flooding. A  
8 review of the 2018-2022 Florida Department of Transportation Five-Year Work Program and  
9 Putnam County's website did not identify any proposed widening or new alignment roadway  
10 projects within the vicinity of the CFG in Putnam County

11 Inside the one-mile buffer area of the CFG in Putnam County, some changes can be noticed  
12 from existing land use to future land use. These changes include a substantial increase in  
13 Agriculture (60 percent) and Conservation (33 percent), addition of Mixed Use (4 percent),  
14 and a decrease in Residential (2 percent) and Public/Institutional (1 percent). The increase  
15 in Conservation is mostly due to the County's land use re-classification. Areas such as the CFG,  
16 Ocala National Forest, Seven Sisters Islands, and Murphy Creek Conservation Area were  
17 classified previously as Public/Institutional or Recreation under then-existing land use and  
18 are now classified as Conservation under future land use. Given that most of the land  
19 surrounding the CFG is Conservation, currently the county does not foresee any significant  
20 development occurring within the one-mile buffer of the CFG.

21 Figure 16, below, displays the generalized future land use for Putnam County within one mile  
22 of the CFG.

### 23 **Property Analysis**

24 As indicated previously, in an attempt to better understand and detail the specific recreation  
25 resource elements, this plan details these resources by county. Effective planning requires a  
26 thorough understanding of the unit's natural and cultural resources. This section describes  
27 the resource characteristics and existing uses of the property. The unit's recreation resource  
28 elements are examined to identify the opportunities and constraints they present for  
29 recreational development. Past and present uses are assessed for their effects on the  
30 property, compatibility with the site, and relation to the unit's classification.

#### 31 *Recreation Resource Elements*

32 This section of the Land Use Component assesses the park's recreational resource elements,  
33 those physical qualities that, either singly or in certain combinations, can support various  
34 resource-based recreation activities. Breaking down the property into such elements  
35 provides a means for measuring the property's capability to support potential recreational  
36 activities. This process also analyzes the existing spatial factors that either favor or limit the  
37 provision of each activity.

1 Land Area  
 2 After Marion County, Putnam County contains the 2nd largest amount of acreage on the CFG,  
 3 though it compares in trail miles to Citrus and Levy counties, as shown in Table 38. As  
 4 illustrated in Table 39, a total of 11,638.2 acres, or 52 percent of land within the CFG  
 5 boundary in Putnam County, is considered altered land use, which includes the Rodman  
 6 Reservoir and large amounts of pine plantations. Bottomland forest, floodplain swamp, mesic  
 7 flatwoods, and hydric hammock comprise a total of 36 percent of the CFG through Putnam  
 8 County.

9 **Table 38. Trails and Mileage on CFG—Putnam County**

Trail Name	Length in County (Feet)	Length in County (Miles)
Buckman Interpretive Trail	16,022.39	3.0
Florida Trail	49,257.00	9.3
Park Road Unstabilized	413,464.90	78.31
<b>Total Trails</b>		<b>90.61</b>

10

11 **Table 39. Natural Communities and Acreage on CFG—Putnam County**

Community	Acreage	Percent Total
Basin Swamp	11.1	0
Baygall	239.9	1
Bottomland Forest	1,092.1	5
Altered Lands	11,638.2	52
Depression Marsh	266.1	1
Dome Swamp	370.1	2
Floodplain Marsh	243.5	1
Floodplain Swamp	2,834.4	13
Hydric Hammock	999.6	4
Mesic Flatwoods	3,099.6	14
Mesic Hammock	197.0	1
Sandhill	145.8	0
Scrub	341.7	2
Scrubby Flatwoods	198.8	0
Upland Hardwood Forest	1.7	0
Wet Flatwoods	744.0	3
Xeric Hammock	117.0	0
<b>Total</b>	<b>22,560.6</b>	<b>100</b>

**Key:** Altered land use includes abandoned fields, canals/ditches, clearings, developed areas, impoundment/artificial pond, pine plantation, roads, spoil areas, and utility corridors.

12  
 13

1 Outstanding examples of hydric hammock within the CFG occur along Deep Creek, north of  
2 Hunter Road, and several areas that join the Ocklawaha River. Other areas include north and  
3 south of the Buckman Lock and north and east of the Rodman Reservoir, near the Kenwood  
4 Recreation Area.

5 As stated previously, restoration of scrub habitat is a priority. This portion of the CFG contains  
6 341.7 acres of scrub, of which 157.3 acres are located within the Deep Creek area. This area  
7 also contains 106.5 acres of sandhill and approximately 13.0 acres of scrubby flatwoods. One  
8 of the most significant examples of quality hydric hammock on the CFG is in the Deep Creek  
9 area.

#### 10 Water Area

11 As the reservoir makes its turn to the east, Orange Creek flows in from the northwest. Farther  
12 east, the Deep Creek and Sweetwater Creek complexes flow into the reservoir. Much of the  
13 Sweetwater Creek complex is on the CFG. East of the reservoir, the east barge canal extends  
14 about nine miles to the east-northeast, where it joins with the St. Johns River. This part of the  
15 canal bisected the Camp Branch Creek system and disrupted the natural surficial flows. The  
16 Camp Branch Creek system originally flowed south-southeast, connecting the Cow Heaven  
17 Bay Swamp to the St. Johns River. Buckman Lock, still operational, is located in the eastern  
18 canal. CFG lands continue about five miles northeast along the St. Johns River.

19 CFG staff maintain and operate the Buckman Lock and Kirkpatrick Dam and Spillway. The  
20 Buckman Lock controls water access to Rodman Reservoir from the St. Johns River through  
21 the east barge canal. It should also be noted that during storm events, Department of Defense  
22 vessels (USCG, COE and USN) use Buckman Lock as a safe harbor. The Kirkpatrick Dam  
23 spillway controls the level of Rodman Reservoir. Generally, the water level of the reservoir is  
24 kept at the 18 feet to 20 feet national geodetic vertical datum (NGVD) level which is equal to  
25 MSL at the dam. The water level is drawn down every three to four years to 11 feet based  
26 upon mutual assessment with FWC fisheries and aquatic plant staff. This is completed to  
27 consolidate bottom sediments, enhance the fishery and wildlife habitats, and to assist in  
28 control of aquatic plants.

#### 29 Natural Scenery

30 Deep Creek is scenic and provides significant habitat to numerous rare species with its  
31 excellent-quality hydric hammock.

#### 32 Significant Habitat

33 Several areas of significant botanical habitat were identified by FNAI throughout the Putnam  
34 County portion of the CFG. These include: Ocklawaha River floodplain, Eureka Dam, Deep  
35 Creek, select areas surrounding the Rodman Reservoir, Caravelle Ranch West, and select  
36 areas surrounding the Buckman Lock.

#### 37 Archaeological and Historical Features

38 With the exception of two abandoned rail lines, the previously recorded cultural sites in the  
39 Putnam County section of the CFG consist exclusively of a variety of prehistoric  
40 archaeological sites. The nature of a number of the archaeological sites is unclear, but each in



1 its own way offers a good cross section of Central Florida’s prehistoric past. Florida SHPO has  
2 determined that 8PU800 and 8PU1568 are not eligible for listing in the NRHP.

3 *Assessment of Use*

4 All legal boundaries, significant natural features, structures, facilities, roads, and trails  
5 existing in the unit are delineated on the base map (see Base Map). Specific uses made of the  
6 unit are briefly described in the following sections.

7 *Current Recreation Use and Visitor Programs*

8 Recreational facilities in Putnam County are primarily associated with fishing and boating  
9 activities and some hunting, as discussed below. Ramps at Kenwood and Rodman  
10 Campground provide water access to the Rodman area, which is nationally known for its  
11 fishing.



The *Kenwood Recreation Area* is located on the north side of Rodman Reservoir off of CR 315, approximately six miles south of the town of Interlachen and five miles northeast of the small community of Orange Springs. There is a two-lane boat ramp, two picnic pavilions, and ample parking located at Kenwood.

23 events (150+ boats). These events make Kenwood the most popular fishing excursion  
24 launching spot on Rodman Reservoir.

25 During the temporary drawdowns conducted at Rodman every three to four years, when the  
26 normal reservoir level of 18 feet to 20 feet is lowered to 11 feet, a temporary ramp is available  
27 at the end of Kenwood Road into the former barge canal channel.



The *Rodman Recreation Area* is comprised of the facilities and recreational opportunities surrounding and located upon the former Rodman Dam, designated by the Florida Legislature as the Kirkpatrick Dam in 1998 (named after former State Senator George Kirkpatrick, who was an ardent defender of keeping the dam and reservoir intact). The 7,200-foot-long earthen dam has a four-gate spillway designed to discharge up to 36,000 cubic feet of water per second from the Rodman

40 Reservoir, which is located on the upstream side of the dam and spillway. The approximately

1 9,500-acre reservoir has a drainage basin of almost 2,800 square miles, and its headwaters  
2 start in the Green Swamp and Lake Apopka. The recreation area is located approximately  
3 three-quarters of a mile west of the Rodman Campground on Rodman Dam Road.

4 Freshwater fishing is the primary recreational pursuit, with bank fishing opportunities on the  
5 downstream discharge side of the spillway; two accessible recreational fishing piers are  
6 located there. There is one wooden pier on the eastern side of the spillway discharge channel  
7 and an aluminum pier on the western side. The moving water flowing through the spillway  
8 and down past the fishing piers provides excellent freshwater fishing opportunities, which  
9 include species such as bass, bream, catfish, and more. There are also hardened areas along  
10 the spillway wing walls where people line up to fish.

11 The upstream reservoir side of the earthen dam, adjacent to the spillway, provides excellent  
12 additional bank fishing opportunities dependent upon the season, weather patterns, and  
13 vegetation. The reservoir side is known for producing bass, bream, catfish, and mullet.

14 For boat fishing or paddling enthusiasts, there is a two-lane boat ramp on the lower east side  
15 of the spillway with paved parking, potable water, picnic pavilions, and restrooms. These  
16 ramps access the lower Ocklawaha River below the dam. The Ocklawaha River flows  
17 approximately nine miles downstream from the dam into the St. Johns River. The Ocklawaha  
18 River is the largest tributary of the St. Johns.

19 People also launch canoes and kayaks here or watch the numerous bird species that frequent  
20 the dam and spillway area. It is common to see Anhinga, Cormorant, Bald Eagles, Osprey, and  
21 numerous types of Herons and Egrets looking around for an easy meal when available.



The 67-site *Rodman Campground* is located approximately nine miles southwest of Palatka off SR 19 on Rodman Dam Road. This campground provides optimal access via its two-lane boat ramp to some of Florida's finest freshwater fishing in the 9,000-acre Rodman Reservoir, which is perennially rated in the Top 10 Trophy Bass Lakes in Florida by the FWC.

32 and extensive habitat not only for trophy fish, but for numerous avian species as well. Many  
33 endangered and threatened species of wading birds, waterfowl, Bald Eagles, and others use  
34 the reservoir, particularly during the cooler months. There are also numerous alligators,  
35 turtles, and even manatees that travel through the reservoir seasonally.

36 The campground and reservoir were created during the 1960s era Cross Florida Barge Canal  
37 project and serve as reminders today of this mammoth public works project, which was  
38 started in 1964 when then President Lyndon B. Johnson flew into Palatka and started the  
39 project with a ground-breaking explosion at the nearby Rodeheaver's Boys Ranch.



The *Buckman Lock* provides navigational connectivity between the St. Johns River and the Rodman Reservoir. The Lock was constructed during the mid to late 1960s as part of the former Cross Florida Barge Canal project, which was de-authorized in 1992 by the U.S. government and the state of Florida when it was approximately one-third complete.

11 The St. Johns River level averages  
12 approximately two feet above mean sea level, and the Rodman Reservoir is normally  
13 operated at a level between 18 feet and 20 feet msl. The interior of the lock chamber is 600  
14 feet long and 84 feet wide with a designed minimum draft of 12 feet with the concrete sill on  
15 the upstream (reservoir) side at 6 feet msl. In addition to boats and watercraft utilizing the  
16 lock, it functions as a fish ladder allowing aquatic species such as manatees, eels, and various  
17 types of fish to traverse the lock to migrate between the river and the reservoir. The St. Johns  
18 Loop North Trailhead provides access to non-motorized multi-use trails to the north and east  
19 of the lock. The trails pass through predominantly pine flatwoods habitat with gently sloping  
topography and moderately drained soils.


20 In addition to the boating opportunities identified above, the FWC manages all hunting on  
21 CFG lands. The FWC leases 16,027 acres of former barge canal lands at the eastern end of the  
22 CFG for management as part of the Caravelle Ranch Wildlife Management Area (WMA). As  
23 such, the Caravelle WMA has 65 days of hunting each year, including separate seasons for  
24 archery, muzzleloaders, and general gun. Small game hunting also is allowed. Popular species  
25 to hunt include deer, turkey, quail, hogs, and squirrel. Hunting seasons generally are of short  
26 duration, with a limited number of hunters, and hunter satisfaction is good.

#### 27 Protected Zones

28 Given the limited amount of disturbance and the number of rare plant species present, the  
29 Deep Creek area is considered a protected area on the CFG.




1 **Figure 15. Putnam County Generalized Existing Land Use**



2



1 **Figure 16. Putnam County Generalized Future Land Use**



2

# 1 CONCEPTUAL LAND USE PLAN

2 The following narrative and Figures 17 through 19, presented below, represent the current  
3 conceptual land use proposal for this park. The conceptual land use plan is the long-term,  
4 optimal development plan for the park, based on current conditions and knowledge of the  
5 park's resources, landscape, and social setting (see Conceptual Land Use Plan). The  
6 conceptual land use plan is modified or amended as new information becomes available  
7 regarding the park's natural and cultural resources or trends in recreational uses to adapt to  
8 changing conditions. Additionally, the acquisition of new parkland may provide  
9 opportunities for alternative or expanded land uses. The DRP develops a detailed  
10 development plan for the park and a site plan for specific facilities based on this conceptual  
11 land use plan, as funding becomes available.

12 During the development of the conceptual land use plan, the DRP assessed the potential  
13 impact of proposed uses or development on the park resources and applied that analysis to  
14 determine the future physical plan of the park, as well as the scale and character of proposed  
15 development. Potential resource impacts also are identified and assessed as part of the site  
16 planning process when funding is available for facility development. At that stage, design  
17 elements (such as existing topography and vegetation, sewage disposal, and stormwater  
18 management) and design constraints (such as imperiled species or cultural site locations) are  
19 investigated in greater detail. Municipal sewer connections, advanced wastewater treatment,  
20 or best available technology systems are applied for on-site sewage disposal. Creation of  
21 impervious surfaces is minimized to the greatest extent feasible to limit the need for  
22 stormwater management systems, and all facilities are designed and constructed using best  
23 management practices to limit and avoid resource impacts. Federal, state, and local permit  
24 and regulatory requirements are addressed during facility development. This includes the  
25 design of all new park facilities consistent with the universal access requirements of the  
26 Americans with Disabilities Act (ADA). After new facilities are constructed, park staff  
27 monitors conditions to ensure that impacts remain within acceptable levels.

## 28 Potential Uses

### 29 **Public Access and Recreational Opportunities**

30 **Goal:** Provide public access and recreational opportunities in the park.

#### 31 **A. Public Accessibility**

32 The CFG is a long linear corridor with a permeable boundary. Access points exist along the  
33 extent of the 110-mile long CFG. It is not feasible to control every ingress/egress point along  
34 the CFG, resulting in unmonitored access. CFG staff work with adjacent property owners to  
35 provide communal access locations. This encourages a "stewardship" philosophy among  
36 adjacent landowners towards the CFG. While public access is considered good, unrestrained  
37 access can cause management problems and unnecessarily impact the natural resources of  
38 the CFG. Given the size and access along the CFG, staff has and will continue to work with  
39 emergency responders to develop maps and other locators to increase safety of CFG



1 visitors. In addition, CFG will continue to work with area developers on public access  
2 opportunities.

3 Due to the extensive management actions required to maintain and restore existing natural  
4 communities, the CFG reserves the right to change access points and trails, both existing and  
5 planned as they deem appropriate. CFG will continue to engage private businesses along the  
6 CFG to encourage the provision of recreation and tourism services. Some objectives related  
7 to access for a specified user group are addressed in other sections, such as biking and  
8 equestrian.

9 *Objective A1: Maintain the park's current recreational carrying capacity of 10,000 users*  
10 *per day.*

11 *Objective A2: Develop and implement an Interpretive Master Plan.*

12 *Objective A3: Update and formalize a policy addressing adjacent landowner access to the*  
13 *CFG.*

14 *Objective A4: Review the existing trail network to determine if any need to be closed or realigned*  
15 *for safety, protection of natural resources, or navigability.*

## 16 **B. Education Facilities**

17 The CFG offers opportunities to view extensive natural and cultural resources. Opportunities  
18 exist to improve the signage and interpretation for these resources throughout the CFG. CFG  
19 will work to provide natural, recreational, cultural and historical resources interpretation  
20 materials and make them available to visitors. New digital technologies are available to  
21 improve the interpretive experience and will be incorporated where appropriate.

22 *Objective B1: Continue to provide natural, recreational, cultural and historical resources*  
23 *interpretation (e.g., signage, checklists, kiosks) where appropriate along the CFG.*

24 *Objective B2: Evaluate interpretive opportunities at the lock system and more*  
25 *specifically, the Buckman Lock.*

26 *Objective B3: Coordinate with OTIS to update the park's website to be more user friendly,*  
27 *educational, and interpretive.*

28 *Objective B4: Develop interpretive brochures and identify locations for distribution, both within*  
29 *and outside the CFG.*

30 *Objective B5: Evaluate opportunities to partner with developers and surrounding land*  
31 *owners to enhance interpretive opportunities through the CFG.*

32 *Objective B6: Develop and implement an Interpretive Cultural Master Plan. Consider*  
33 *securing grant funds for this effort in conjunction with surrounding municipalities.*

1 **C. Multi-Use Trails**

2 Given the boundaries and the limited land availability, CFG staff will seek to utilize multi-  
3 use trails where appropriate as opposed to individual use trails. Additional multi-use  
4 trails, including crossings, are planned for the CFG. Crossings are necessary for visitor  
5 safety and wildlife. Not all land necessary for some of the listed trails is in public  
6 ownership. A combination of land acquisition and private landowner agreements may be  
7 necessary to establish future trails.

8 *Objective C1: Working with FDOT and other partners, establish off-grade road crossings*  
9 *where appropriate, including SR 200, US 441 and future expansion of I-75.*

10 *Objective C2: Work with the City of Palatka, Putnam County, other state agencies, and public*  
11 *interest groups to improve connectivity to the Water Works Environmental Education Center in*  
12 *Palatka.*

13 *Objective C3: Evaluate other opportunities to establish linkages to other publicly-accessible*  
14 *multi-use trails.*

15 *Objective C4: Add mile marker signage to multi-use trails in the CFG.*

16 Continue coordination with EMS/LE on-trail reference points. Location reference markers  
17 are better when latitude/longitude is incorporated for emergency management services and  
18 law enforcement use.

19 **D. Hiking**

20 Trails for hiking-only are maintained by the Florida Trail Association (FTA). The CFG contains  
21 43 miles of the FNST. Additional incorporation of the FNST within the CFG also should be  
22 considered and encouraged. Linkages to other publicly accessible hiking trails also would be  
23 beneficial. Any proposed new hiking trails will be carefully evaluated on a case by case basis.  
24 Currently, an extensive network of hiking trails exists throughout the CFG. Given this, the CFG  
25 has a limitation placed on the number of trails that can be developed while balancing the  
26 maintenance of quality wildlife habitat and ecological function. CFG staff remain committed  
27 to working in concert with FTA to maintain the existing hiking trail network. The inclusion of  
28 more interpretive materials along the trails is warranted.

29 *Objective D1: Evaluate and update the existing maintenance agreement with FTA, which*  
30 *encourages the responsible use and maintenance of hiking trails.*

31 *Objective D2: Encourage relationships with other local groups focused on responsible use and*  
32 *maintenance of hiking, walking, running and interpretive trails.*

33 *Objective D3: Evaluate other opportunities to establish linkages to other publicly-accessible*  
34 *hiking trails.*



1 **E. Biking**

2 Responsible biking on the CFG is necessary to protect the natural resources along the  
3 Greenway. The Ocala Mountain Bike Association (OMBA) is active in promoting proper trail  
4 usage. Similar efforts by other bike groups should be encouraged where appropriate. Any  
5 proposed new biking trails will be carefully evaluated on a case by case basis. Currently, there  
6 is an extensive network of biking trails throughout the CFG. Given this, the CFG has a  
7 limitation placed on the number of trails that can be developed while balancing the quality  
8 wildlife habitat and ecological function. CFG staff remain committed to working in concert  
9 with OMBA to maintain the existing biking trail network.

10 *Objective E1: Continue to work with the OMBA to promote responsible use and maintenance of*  
11 *bicycle trails*

12 *Objective E2: Evaluate and update as necessary existing agreements with biking associations to*  
13 *formalize the planning and maintenance process for mountain biking facilities along the CFG.*

14 *Objective E3: Evaluate other opportunities to establish linkages to other publicly-accessible*  
15 *biking trails.*

16 **F. Equestrian**

17 As with biking, responsible equestrian usage of trails is essential on the Greenway. CFG staff  
18 will work with the Greenway Equestrians group and others to encourage responsible use and  
19 maintenance of equestrian trails. CFG staff will continuously evaluate and implement  
20 adaptations to existing equestrian trails and equestrian facilities as warranted by changing  
21 conditions.

22 *Objective F1: Encourage the further development of the Greenway Equestrians group and other*  
23 *equestrian groups along the CFG to promote responsible use and maintenance of equestrian*  
24 *trails.*

25 *Objective F2: Relocate equestrian facilities from Santos Trailhead farther south along US*  
26 *441 and evaluate potential expansion and improvement opportunities (e.g., new trailhead,*  
27 *campground, and concessionaire).*

28 *Objective F3: Evaluate other opportunities to establish linkages to other publicly-accessible*  
29 *equestrian trails.*

30 *Objective F4: Continue to provide assistance to those entities that offer equestrian interaction*  
31 *to physically and emotionally challenged individuals where possible.*

32 **G. Paddling**

33 Although paddling is a popular sport on the Greenway, no trails are currently designated.  
34 CFG will work with paddling groups to establish paddling trails and primitive  
35 campgrounds along the paddling routes, as appropriate. It is the Army Corps of  
36 Engineers' responsibility to maintain the navigability of the Ocklawaha River. CFG is

1 already working with Marion County and other agencies to remove snags from the river  
2 to enhance navigability and will continue these cooperative efforts.

3 *Objective G1: Designate and consider marking a paddling trail on the Ocklawaha River with*  
4 *designated official primitive campsites.*

5 *Objective G2: Promote the designation of the Ocklawaha River from SR 40 to Eureka*  
6 *(approximately 15 miles) as a Wild and Scenic River.*

7 *Objective G3: Evaluate other opportunities to establish linkages to other publicly accessible*  
8 *paddling trails.*

## 9 **H. Boating**

10 Motorized boating is popular on the CFG, especially for fishing.

11 *Objective H1: Pave Kenwood Recreation Area road and parking area, provide restrooms,*  
12 *and provide one or more staff/security residences.*

13 *Objective H2: In cooperation with FWC and the Coast Guard, provide channel markers for the*  
14 *original river channel (which serves as the navigation channel in many locations) within*  
15 *Rodman Reservoir, if funds are available. Additional funds are needed to implement and be*  
16 *completed in priority order.*

17 *Objective H3: Continue partnerships with local governments in the operation and maintenance*  
18 *of boat launches on the CFG.*

19 *Objective H4: Evaluate the opportunity to develop a Guided Historical Interpretive Boat Tour*  
20 *along the greenway canal and lock system.*

## 21 **I. Camping**

22 Camping is available in a limited number of areas along the CFG. CFG staff plan to  
23 improve and provide additional camping facilities, some in association with other  
24 activities, such as paddling along the Ocklawaha River.

25 *Objective I1: Evaluate the need for and resource impact of expansion of current*  
26 *campgrounds (i.e., Santos Campground Loop 2 and Ross Prairie).*

27 *Objective I2: Evaluate the need for and resource impact of improvements at Kenwood*  
28 *Recreation Area.*

29 *Objective I3: Identify and designate current primitive campground facilities (no vehicle access,*  
30 *minimal facilities) in several locations along the Ocklawaha River (boater friendly,*  
31 *improvement of current facilities), along the proposed natural surface trail corridor connecting*  
32 *Marshall Swamp and Rodman (possibly the same as the boater campsites) and between Pruitt*  
33 *Trailhead and Felburn Park. Lack of ability to have natural surface trails between Silver*  
34 *Springs State Park and Rodman area due to low lying lands. Better to use existing aquatic*  
35 *corridor.*

1 **J. Fishing**

2 Fishing, especially from boats, is popular along the CFG. The opportunity to provide  
3 more land-based fishing access points will be evaluated and implemented, if feasible.

4 *Objective J1: Assess the impacts, desirability, demand for, and cost of installing multi-*  
5 *purpose boardwalks/docks that would allow for non-boat fishing on the CFG.*

6 *Objective J2: Establish additional fishing access points at appropriate locations.*

7 **K. Hunting**

8 The feasibility of providing additional hunting opportunities will be discussed with  
9 FWC, which manages all hunts on CFG lands. Care will be taken to ensure that hunting  
10 does not unduly detract from other user groups use of the CFG.

11 *Objective K1: Continue cooperation with FWC managing hunts in designated hunting areas on*  
12 *the CFG.*

13 *Objective K2: Identify and map designated hunting areas within the CFG.*

14 **Proposed Facilities**

15 **Capital Facilities and Infrastructure**

16 **Goal:** Develop and maintain the capital facilities and infrastructure necessary to implement  
17 the recommendations of the management plan.

18 The existing facilities of this state park are appropriate to the natural and cultural resources  
19 contained in the park and should be maintained. New construction, as discussed further  
20 below, is recommended to improve the quality and safety of the recreational opportunities,  
21 to improve the protection of park resources, and to streamline the efficiency of park  
22 operations. The following is a summary of improved facilities needed to implement the  
23 conceptual land use plan for the CFG.

24 *Objective A: Maintain all public and support facilities in the park.*

25 All capital facilities, trails and roads within the park will be kept in proper condition through  
26 the daily or regular work of park staff and/or contracted help.

27 *Objective B: Continue to implement the park's transition plan to ensure facilities are accessible*  
28 *in accordance with the Americans with Disabilities Act of 1990.*

29 *Objective C: Expand maintenance activities as existing facilities are improved and new facilities*  
30 *are developed.*

31 *Objective D: Develop a prioritization process for necessary capital improvements on the CFG.*

32 As previously indicated, the CFG is 110 miles long and contains numerous recreational  
33 opportunities and facilities and limited resources. As such, it is extremely important that the

1 resources are utilized in an efficient manner. By prioritizing these capital improvement  
2 activities, it reduces/minimizes maintenance and repair needs.

3 *Objective E: Evaluate and consider the transfer of operation and management responsibilities*  
4 *for the Kirkpatrick Dam and Buckman Lock to the SJRWMD.*

## 5 **Facilities Development**

6 Preliminary cost estimates for these recommended facilities and improvements are provided  
7 in the Implementation Component of this plan. These cost estimates are based on the most  
8 cost-effective construction standards available at this time. The preliminary estimates are  
9 provided to assist DRP in budgeting future park improvements, and may be revised as more  
10 information is collected through the planning and design processes.

11 New facilities and improvements to existing facilities recommended by the plan include:

### 12 **Eureka Lock Law Enforcement**

13 The Eureka Lock is a heavily disturbed borrow pit with extensive unsanctioned OHV activity.  
14 CFG would like to place (up to two) law enforcement residences. Sewer and water hookups  
15 are present at the site.

### 16 **Withlacochee Bay Trail Road**

17 The CFG strongly desires to pave the Withlacochee Bay Trail maintenance road. By paving  
18 this road, the CFG will significantly minimize its maintenance obligations to be able to more  
19 efficiently utilize existing resources.

### 20 **Relocation of Santos Equestrian Facilities**

21 Relocate equestrian facilities from Santos Trailhead farther south and west along the  
22 south side of the CFG corridor to minimize the need for accel/decal lanes. Most of CFG  
23 equestrian trails are located along the southern boundary of the greenway. By  
24 relocating these facilities, it will ensure the appropriate collocation of like uses. In  
25 addition to relocation of the facilities, consideration will be given to expansion and  
26 improvement opportunities (e.g., new trailhead, campground, and concessionaire).

### 27 **Withlacochee Bay Trail Phase III**

28 Phase III of the Withlacochee Bay Trail has been completed, extending the existing trail from  
29 Felburn Park along the canal berm to where the Withlacochee River was bisected by the  
30 canal to Inglis Island. The only remaining portion of this project is the construction of the  
31 eastward bridge, which will facilitate the connection of the trails on the western portion of  
32 the CFG.

## 33 **Other Potential Future Trail/Facilities Developments**

34 In addition to the abovementioned projects, there are several future projects located either  
35 in proximity to or on the CFG that will require DRP/CFG involvement during the planning  
36 process. These projects are not listed or included in the Implementation Component.



1 **Dunnellon to Inglis Lock Multi-Use Trail Corridor**

2 CFG has very little land going east-west along the north and south shores of Lake Rousseau,  
3 which creates a gap in the CFG trail system. CFG would like to establish a paved connector  
4 trail north of Lake Rousseau from Dunnellon to the Inglis Lock, primarily using existing right  
5 of way of CR 40. Environmental impact is expected to be minimal, given the trail's location in  
6 a right of way. A trail separated from the road is planned. Anticipated partners include Marion  
7 County, Levy County, FDOT, and the Cities of Dunnellon and Inglis.

8 **CFG to Nature Coast Multi-Use Connector Trail Corridor**

9 Duke Energy has conveyed a perpetual easement on abandoned railroad right-of-way to the  
10 state for the establishment of a 42-mile paved multi-use trail from the north side of Dunnellon  
11 to Chiefland through the Goethe State Forest connecting to the Nature Coast State Trail. The  
12 paved connector from the CFG to Goethe State Forest to Chiefland and the Nature Coast trail  
13 system would greatly expand the opportunities for CFG trail users. In addition to the 53,000-  
14 acre Goethe State Forest, visitors could access the Nature Coast State Trail. The state does not  
15 own most of the land necessary to make the connection to Goethe; the cooperation of private  
16 landowners likely will be necessary. Because of the potential to link two major equestrian  
17 trail systems, accommodations for equestrian use will be made if feasible. Anticipated  
18 partners are the City of Dunnellon, FDOT, FFS, and private landowners.

19 **Dunnellon to Pruitt Trailhead Multi-Use Trail Corridor**

20 A multi-use trail corridor is planned to link the Pruitt Trailhead to the ballfields in Dunnellon.  
21 Additional land acquisition will be necessary for this trail to be completed. Project may use  
22 existing road right-of-way where needed to make connection if other lands are not feasible  
23 to acquire.

24 **Ocklawaha Paddling Trail with Camping**

25 The feasibility of establishing a paddling trail on the Ocklawaha River and into Rodman  
26 Reservoir will be evaluated. The trail may be established with GPS points and maps, rather  
27 than posted signage. If feasible, a system of primitive campsites may be established and  
28 designated.

29 **Kenwood Recreation Area Improvements**

30 The existing one-mile plus dirt road at the Kenwood Recreation Area will be paved to the boat  
31 ramp. This ramp is heavily used by recreational and tournament fishermen. If feasible, the  
32 campground will be reopened if funds and staffing are available for development, staffing  
33 O&M and security.

34 **Ocklawaha River Visitor Center and Interpretive Trail Improvements**

35 The Felburn Foundation leases the Ocklawaha Visitor Center located on 2.7 acres on the  
36 corner of State Hwy 40 and County Rd 315. The Felburn Foundation will fund all necessary  
37 improvements to the interior and exterior of the structure. In addition to these  
38 improvements, the Felburn Foundation would erect a kiosk at the trail head with interpretive

1 information regarding the trail to the Ocklawaha River with shared information regarding  
2 the public lands trails systems that are found throughout this area. They would also develop  
3 and install interpretive trail signs approved by DEP's biologists on the trail to the Ocklawaha  
4 River.

5 **US 19 Boat Ramp**

6 CFG has been approached by Citrus County to consider leasing 16 acres of land on the north  
7 side of the barge canal just west of US 19 for a boat ramp. This property is isolated from other  
8 DEP lands. This parcel is somewhat disturbed. A 30-year lease is planned that will give the  
9 county five years to fund, design, permit, and construct the boat ramp, or the lease will be  
10 canceled if this project is not pursued further by the county. Anticipated partners include  
11 Citrus County and FWC.



1 **Figure 17. Cross Florida Greenway Conceptual Land Use Map—West Section**



2




1 **Figure 18. Cross Florida Greenway Conceptual Land Use Map—Central Section**



2



1 Figure 19. Cross Florida Greenway Conceptual Land Use Map—East Section



2



1 **Existing Use and Recreational Carrying Capacity**

2 Carrying capacity is an estimate of the number of users a recreation resource or facility can  
3 accommodate and still provide a high quality recreational experience and preserve the  
4 natural values of the site. The carrying capacity of a unit is determined by identifying the land  
5 and water requirements for each recreation activity at the unit, and then applying these  
6 requirements to the unit's land and water base. Next, guidelines are applied which estimate  
7 the physical capacity of the unit's natural communities to withstand recreational uses  
8 without significant degradation. This analysis identifies a range within which the carrying  
9 capacity most appropriate to the specific activity, the activity site, and the unit's classification  
10 is selected (see Table 40).

11 The recreational carrying capacity for this park is a preliminary estimate of the number of  
12 users the unit could accommodate after the current conceptual development program has  
13 been implemented. However, a recreational carrying capacity study should not be used as the  
14 sole determining factor for limiting recreational use or access. Rather, a recreational carrying  
15 capacity analysis should be used as a tool to evaluate the range of options that are available  
16 to help minimize multi-use conflicts, environmental concerns, and other problems associated  
17 with overuse and overcrowding. A recreational carrying capacity study can establish a  
18 framework for decision making and provide a basis for regulatory action. When developed,  
19 the proposed new facilities would increase the unit's carrying capacity, as shown in Table 40.

1 **Table 40. Recreational Carrying Capacity**

Recreation Activity	Type	Existing Capacity		Proposed Additional Capacity		Estimated Recreational Capacity	
		One Time	Daily	One Time	Daily	One Time	Daily
<b>Camping</b>							
	Primitive	104	104				
	Short-walk, Tent	208	208				
	Standard	944	944	96	96	1,040	1,040
<b>Trails</b>							
	<b>Biking</b>						
	Maximum	605	2,421			605	2,421
	Minimum		2,421			-	2,421
	Average		2,421			-	2,421
	<b>Equestrian</b>					-	-
	Maximum	2,382	4,764			2,382	4,764
	Minimum		595			-	595
	Average		1,489			-	1,489
	<b>Hiking</b>					-	-
	Maximum	2,105	8,422			2,105	8,422
	Minimum		2,105			-	2,105
	Average		5,053			-	5,053
	<b>Hiking/Biking</b>					-	-
	Maximum	1,176	4,702			1,176	4,702
	Minimum		2,351			-	2,351
	Average		3,527			-	3,527
	<b>Hiking/Biking/ Equestrian</b>					-	-
	Maximum	1,124	4,495			1,124	4,495
	Minimum		2,248			-	2,248
	Average		3,371			-	3,371
<b>Other</b>							
	Campfire Circle	628	628			628	628
	Picnicking	784	784			784	784
	Boat Ramps	26	20,800	2	800	28	21,600
	<b>TOTAL</b>	<b>10,086</b>	<b>37,856</b>			<b>9,872</b>	<b>38,440</b>

2

3

## 1 **Optimum Boundary**

2 The Optimum Boundary Map reflects lands considered desirable for direct management by  
3 the DRP as part of the state park. These parcels may include publicly owned or privately  
4 owned land that would improve the continuity of existing parklands, provide the most  
5 efficient boundary configuration, improve access to the park, provide additional natural and  
6 cultural resource protection or allow for future expansion of recreational activities.  
7 Parklands that are potentially surplus to the management needs of DRP also are identified.  
8 As additional needs are identified through park use, development, and research, and as land  
9 use changes on adjacent property, modification of the park’s optimum boundary may be  
10 necessary.

11 Identification of parcels on the optimum boundary map is intended solely for planning  
12 purposes. It is not to be used in conjunction with any regulatory purposes. Any party or  
13 governmental entity should not use a property’s identification on the optimum boundary  
14 map to reduce or restrict the lawful rights of private landowners. Identification on the map  
15 does not empower or suggest that any government entity should impose additional or more  
16 restrictive environmental land use or zoning regulations. Identification should not be used as  
17 the basis for permit denial or the imposition of permit conditions.

18 Several parcels have been discussed during the planning process as being necessary to  
19 optimize the management of the CFG, as well as enhance connectivity within the CFG.  
20 Outlined below are the parcels in priority order that are identified as part of the optimum  
21 boundary. CFG staff have identified parcels of property to be obtained by acquisition, also  
22 parcels that have been identified as surplus property. It should be noted that even if the  
23 parcels recommended as surplus cannot be surplus, consideration should be given to  
24 removing these parcels from managed lands. Property acquisition should be considered the  
25 most ideal option moving forward. However, parcels designated as acquisition may be  
26 considered for exchange upon approval by all parties and would be subject to review and  
27 approval by the Board of Trustees.

## 28 **Property Acquisition**

- 29 1) Santos Gap
- 30 2) FDOT Scrub Triangle keyhole parcel
- 31 3) Florida Power and Light Ocklawaha River parcels
- 32 4) DECCA—adjacent to west side of I-75 along CFG south boundary
- 33 5) Greenberg Properties—north side of Dunnellon Rainbow River area
- 34 6) Cannon/Folks
- 35 7) USFS parcels along Ocklawaha River between SR 40 and gas pipeline to consolidate  
36 and clean up management boundary lines.
- 37 8) 40-acre parcel owned by Marion County (Dinkins Parcel)
- 38 9) Inholdings along North side of SR 40 between SCR 326 and SR 35



1 **Surplus**

- 2 1) River Gardens—south side of Lake Rousseau west of Dunnellon
- 3 2) Citrus County US 19 boat ramp parcel—surplus in exchange for county paving the
- 4 Withlacoochee Bay Trail Road
- 5 3) USFS—approximately a dozen inholdings each agency has within CFG/ONF to
- 6 improve respective management boundaries
- 7 4) SJRWMD—St. Johns River parcels to water management district
- 8 5) Ernie Cremer—consolidated perimeter parcel of current state-owned lands to clean
- 9 up management boundaries and reduce easements to others
- 10 6) Marion County Parks—Independence Park request—east side of SW 49th Avenue
- 11 along CFG south boundary
- 12 7) 10-acre rectangle and angular parcel within residential area west of SR 35 and north
- 13 of CR 464; DSL # CF-714-4116
- 14 8) 464 frontage and mobile home park parcels west of SR 35 and north of Rotary
- 15 Sportsplex; DSL Surplus # FMLA\_177 and CF-714-4111
- 16 9) Rainey Inholdings—Rainey Pasture east side of CR 315 and north of SR 40
- 17 10) DSL FLMA\_184
- 18 11) DSL FLMA\_171
- 19 12) Parcel at the intersection of NE 35th Street and NE 60th Court.




1 **Figure 20. Cross Florida Greenway Optimum Boundary Map, Page 1 of 7**



2




1 **Figure 21. Cross Florida Greenway Optimum Boundary Map, Page 2 of 7**



2




1 **Figure 22. Cross Florida Greenway Optimum Boundary Map, Page 3 of 7**



2




1 **Figure 23. Cross Florida Greenway Optimum Boundary Map, Page 4 of 7**



2




1 **Figure 24. Cross Florida Greenway Optimum Boundary Map, Page 5 of 7**



2




1 **Figure 25. Cross Florida Greenway Optimum Boundary Map, Page 6 of 7**



2



1 **Figure 26. Cross Florida Greenway Optimum Boundary Map, Page 7 of 7**



2



# 1      **IMPLEMENTATION COMPONENT**

2      The resource management and land use components of  
3      this management plan provide a thorough inventory of  
4      the park’s natural, cultural, and recreational resources.  
5      They outline the park’s management needs and  
6      problems, and recommend both short- and long-term  
7      objectives and actions to meet those needs. The  
8      implementation component addresses the  
9      administrative goals for the park and reports on the  
10     DRP’s progress toward achieving resource  
11     management, operational, and capital improvement  
12     goals and objectives since approval of the previous  
13     management plan for this park. This component also  
14     compiles the management goals, objectives, and  
15     actions expressed in the separate parts of this  
16     management plan for easy review. Estimated costs for  
17     the 10-year period of this plan are provided for each  
18     action and objective, and the costs are summarized  
19     under standard categories of land management  
20     activities.

## 21     **MANAGEMENT PROGRESS**

22     Since the approval of the last management plan for the  
23     CFG in 2007, significant work has been accomplished  
24     and progress has been made toward meeting the DRP’s  
25     management objectives for the park. These  
26     accomplishments fall within three of the five general  
27     categories that encompass the mission of the park and  
28     the DRP.

### 29     **Park Administration and Operations**

30     Continued to fulfill the goals of the DRP mission,  
31     maintaining infrastructure and meeting visitors’  
32     expectations

### 33     **Resource Management**

#### 34     **Fire**

35     15,977 acres burned on the CFG

- 36         • 66 burn zones in rotation (7,325 acres)
- 37         • 29 new burn zones since 2007

- 1 • All the CFG fire type acreage is now divided into burn zones
- 2 • Utilization of prescribed burn contractors to increase annual acreage

### 3 **Natural Community Restoration**

- 4 • 3,925 acres in timber harvests for restoration of natural communities with the side
- 5 benefit of earning \$1.065 million in revenue
- 6 • 498 acres of trees planted
- 7 • 157 acres of groundcover planted
- 8 • 540 acres of mechanical treatments
- 9 • A timber inventory and management plan for the 9,000 acre Etoniah addition is
- 10 planned

### 11 **Endangered Species**

- 12 • In 2008, coordinated with Audubon Florida to join their annual Florida Scrub-Jay
- 13 Watch program to monitor the Florida Scrub-Jay population on the CFG Triangle
- 14 Scrub tract
- 15 • Restored 523 additional acres of scrub, for a total of 840 acres out of 1,100 acres of
- 16 historic scrub restored
- 17 • Contracted to band Jays for better data collection methods on population and family
- 18 responses to management actions
- 19 • Mechanically treated 538 acres
- 20 • Increased the Florida Scrub-Jay population from 46 birds in 2009 to 111 in 2015
- 21 • Coordinated with FWC to set up boxes for kestrel recruitment
- 22 • Set up wildlife cameras in 2009 at all the underpasses on the CFG and the Landbridge
- 23 to capture wildlife usage

### 24 **Recreational Facilities and Visitor Services**

- 25 • Added a new floating boat dock at Kenwood Boat Ramp
- 26 • Added a new floating boat dock and ADA ramp at the US 19 Boat Ramp
- 27 • Added new picnic pavilions with ADA access at Eureka Recreation Area West, Orange
- 28 Springs, Kenwood, Rodman East Recreation Area, Buckman Recreation Area
- 29 • Phase 2 Rodman Campground
- 30 • Logged Miller Tract
- 31 • Initiated road repair and culvert replacement on the Miller Tract
- 32 • St. Johns Trail Loop South Primitive Equestrian Campground
- 33 • Hunter Road Trailhead
- 34 • Updated Inglis Bypass Recreation Area with ADA sidewalks
- 35 • Bulkheads at Inglis Lock and equipment bridge
- 36 • New section of Withlacoochee Bay Trail going east to Inglis Island
- 37 • Dunnellon Trail and Bridge
- 38 • Coordinating with Marion County on the proposed new paved trail between SR 200 and
- 39 Dunnellon Trail
- 40 • Developed and opened Shangri La Campground and Trailhead
- 41 • Developed and opened Vortex Trailhead
- 42 • Developed and opened Ned Folks Pavilion

## 1 **MANAGEMENT PLAN IMPLEMENTATION**

2 This management plan is written for a timeframe of 10 years, as required by Section 253.034,  
3 Florida Statutes. The 10-Year Implementation Schedule and Cost Estimates (Table 41)  
4 summarizes the management goals, objectives, and actions that are recommended for  
5 implementation during this period and beyond. Measures are identified for assessing  
6 progress toward completing each objective and action. A time frame for completing each  
7 objective and action also is provided. Preliminary cost estimates for each action are provided  
8 and the estimated total costs to complete each objective are computed. Finally, all costs are  
9 consolidated under the following five standard land management categories: Resource  
10 Management, Administration and Support, Capital Improvements, Recreation Visitor  
11 Services, and Law Enforcement.

12 Some of the actions identified in the plan can be implemented using existing staff and funding  
13 sources based on grants, partnerships, and legislative appropriations. However, as the plan  
14 guides long-term management over a period of 10 years, some actions have been identified  
15 that may require additional resources. The 10-year Implementation Schedule and Cost  
16 Estimates table, therefore, includes both “funded” and “unfunded” needs. It should be noted  
17 that the costs associated with each of the five standard land management categories are  
18 expected to increase over the 10-year period covered by this plan. The estimate of costs  
19 provided herewith is based on the best information available at the time this plan was  
20 completed and cannot be considered a final determination of actual costs over the 10-year  
21 life of the plan.

22 The administration of the state park is an ongoing cost that will increase in the future as  
23 additional staff, programs, and responsibilities are assigned. These administrative costs  
24 include a variety of activities, such as the administration of personnel, the management of  
25 vendors and contractors for all the park’s supply and service needs, and the coordination of  
26 the park’s Citizen Support Organization, to name a few. A high degree of adaptability and  
27 flexibility is necessary for implementation of this management plan to ensure that the  
28 Division can adjust to changes in the availability of funds, create improved understanding of  
29 the park’s natural and cultural resources, and remain current with changes in statewide land  
30 management issues, priorities, and policies.

31 Statewide priorities for all aspects of land management are evaluated each year as part of the  
32 process for developing the Division’s annual legislative budget requests. When preparing  
33 these annual requests, the Division considers the needs and priorities of the entire state park  
34 system and the projected availability of funding from all sources during the upcoming fiscal  
35 year. In addition to annual legislative appropriations, the Division pursues supplemental  
36 sources of funds and staff resources wherever possible, including grants, volunteers, and  
37 partnerships with other entities. The Division’s ability to accomplish the specific actions  
38 identified in the plan will be determined largely by the availability of funds and staff for these  
39 purposes, which may vary from year to year. Consequently, the target schedules and  
40 estimated costs identified in Table 41 may need to be adjusted during the 10-year  
41 management planning cycle.

1 **Table 41. Marjorie Harris Carr Cross Florida Greenway State Recreation and Conservation Area 10-Year Implementation Schedule and Cost**  
 2 **Estimates Sheet**

<b>NOTE: THE DIVISION'S ABILITY TO COMPLETE THE OBJECTIVES OUTLINED BY THE MANAGEMENT PLAN IS CONTINGENT ON THE AVAILABILITY OF FUNDING AND OTHER RESOURCES FOR THESE PURPOSES.</b>						
<b>Goal I: Provide administrative support for all park functions.</b>		<b>Measure</b>	<b>Planning Period</b>	<b>Estimated Manpower and Expense Cost (10 Years)</b>	<b>Estimated Manpower Cost (10 Years)</b>	<b>Estimated Expense Cost (10 Years)</b>
<b>Objective A</b>	<b>Continue day-to-day administrative support at current levels.</b>	Administrative support ongoing	C	\$954,070	\$942,070	\$12,000
<b>Objective B</b>	<b>Expand administrative support as new lands are acquired, new facilities are developed, or as other needs arise.</b>	Administrative support expanded	C	\$433,536	\$419,536	\$14,000
<b>Goal II: Protect water quality and quantity in the park, restore hydrology to the extent feasible, and maintain the restored condition.</b>		<b>Measure</b>	<b>Planning Period</b>	<b>Estimated Manpower and Expense Cost (10-years)</b>	<b>Estimated Manpower Cost (10 Years)</b>	<b>Estimated Expense Cost (10 Years)</b>
<b>Objective A</b>	<b>Conduct periodic former Cross Florida Barge Canal Water Control Structures inspections, repairs and maintenance per FEMA guidelines and professional engineers' recommendations.</b>	# of structures maintained according to accepted guidelines	C	\$5,750,000	\$1,750,000	\$4,000,000.00
<b>Objective B</b>	<b>Repair hydrological conditions and function to approximately 16,700 acres—Etoniah, Marshall Swamp, and Gore's Landing.</b>	# of acres restored or with restoration underway	C	\$1,900,000	\$950,000	\$950,000



<b>Goal III: Maintain and restore the natural communities/habitats in the park</b>		<b>Measure</b>	<b>Planning Period</b>	<b>Estimated Manpower and Expense Cost (10-years)</b>	<b>Estimated Manpower Cost (10 Years)</b>	<b>Estimated Expense Cost (10 Years)</b>
<b>Objective A</b>	<b>Within 10 years, have 25,865 acres of the park maintained within optimal fire return interval.</b>	# of acres within fire return interval target	LT	<b>\$10,570,300</b>	<b>\$6,081,800</b>	<b>\$4,488,200</b>
Action 1	Develop/update annual burn plan.	Plan updated	C	\$30,000	\$21,800	\$8,200
Action 2	Manage fire-dependent communities for ecosystem function, structure, and processes by burning between 7,500-8,000 acres annually, as identified by the annual burn plan.	Average # of acres burned annually	C	\$7,900,300	\$4,740,000	\$3,160,000
Action 3	Establish and maintain 500 miles of fire breaks.	# of miles established	LT	\$2,640,000	\$1,320,000	\$1,320,000
<b>Objective B</b>	<b>Conduct habitat/natural community restoration activities on 2,555 acres of ruderal community(ies).</b>	# of acres restored or with restoration underway	LT	<b>\$1,376,550</b>	<b>\$521,565</b>	<b>\$854,985</b>
Action 1	Plant 50-100 acres of wiregrass annually in 2,000 acres of old pastures planted in longleaf pine.	# of acres planted w/ wiregrass	LT	\$1,260,000	\$420,000	\$840,000
Action 2	Replant 555 acres of slash pine in Etoniah properties.	# of acres replanted in Etoniah	LT	\$116,550	\$101,565	\$14,985
<b>Goal IV: Maintain, improve or restore imperiled species populations and habitats in the park.</b>		<b>Measure</b>	<b>Planning Period</b>	<b>Estimated Manpower and Expense Cost (10-years)</b>	<b>Estimated Manpower Cost (10 Years)</b>	<b>Estimated Expense Cost (10 Years)</b>
<b>Objective A</b>	<b>Update baseline imperiled species occurrence inventory lists for plants and animals w/ FNAI.</b>	Updated	C	<b>\$40,000</b>	\$30,000	<b>\$10,000</b>
<b>Objective B</b>	<b>Monitor and document three selected imperiled animal species in the park.</b>	Population trends of species monitored	C	<b>\$150,000</b>	\$125,050	\$24,950
<b>Objective C</b>	<b>Monitor and document one selected imperiled plant species in the park.</b>	Population and dispersal trend/s	C	<b>\$25,000</b>	\$20,410	<b>\$4,590</b>
<b>Objective D</b>	<b>Maintain/improve scrub jay habitat on the CFG.</b>	# of acres maintained/imp roved of scrub jay habitat	C	<b>\$285,200</b>	\$35,200	<b>\$250,000</b>

Action 1	Mechanically treat approximately 100 acres of scrub jay habitat per year to maintain good habitat conditions.	# of acres of scrub jay habitat mechanically treated	C	\$250,000	\$0	\$250,000
Action 2	Conduct seasonal scrub jay banding to track the species population within the park.	# of banded scrub jays	C	\$35,200	\$35,200	\$0
<b>Goal V: Remove exotic and invasive plants and animals from the park and conduct needed maintenance-control.</b>		<b>Measure</b>	<b>Planning Period</b>	<b>Estimated Manpower and Expense Cost (10-years)</b>	<b>Estimated Manpower Cost (10 Years)</b>	<b>Estimated Expense Cost (10 Years)</b>
<b>Objective A</b>	<b>Annually treat approximately 7,500 infested acres of exotic upland plant species in the park.</b>	# of acres treated	C	\$4,144,000	\$3,700,000	\$517,000
Action 1	Annually develop/update exotic plant management work plan.	Plan developed/updated	C	\$25,524	\$25,524	\$0
Action 2	Implement annual work plan by treating 7,500 acres in park, annually, and continuing maintenance and follow-up treatments, as needed.	Plan implemented	C	\$4,118,476	\$3,674,476	\$517,000
<b>Objective B</b>	<b>Implement control measures for feral hogs on the CFG.</b>	# of hogs controlled	C	\$7,000	\$6,000	\$1,000
<b>Goal VI: Protect, preserve and maintain the cultural resources of the park.</b>		<b>Measure</b>	<b>Planning Period</b>	<b>Estimated Manpower and Expense Cost (10-years)</b>	<b>Estimated Manpower Cost (10 Years)</b>	<b>Estimated Expense Cost (10 Years)</b>
<b>Objective A</b>	<b>Assess and evaluate 30 of 300 recorded cultural resources in the park annually.</b>	Ongoing	C	\$420,000	\$327,600	\$92,400
Action 1	Complete 300 assessments/evaluations of archaeological sites. Prioritize preservation and stabilization projects.	Assessments complete	LT	\$400,000	\$312,000	\$88,000
Action 2	Complete Historic Structures Reports (HSRs) for historic buildings and cultural landscape. Prioritize stabilization, restoration and rehabilitation projects.	Reports and priority lists completed	LT	\$20,000	\$15,600	\$4,400
<b>Objective B</b>	<b>Compile reliable documentation for all recorded historic and archaeological sites.</b>	Documentation complete	LT	\$600,000	\$450,000	\$150,000

Action 1	Ensure all known sites are recorded or updated in the Florida Master Site File. Would be part of Objective B of having professional archaeologist inventory and assess all know and listed sites.	# of sites recorded or updated	ST	See Objective B	See Objective B	See Objective B
<b>Goal VII: Provide public access and recreational opportunities in the park.</b>		<b>Measure</b>	<b>Planning Period</b>	<b>Estimated Manpower and Expense Cost (10-years)</b>	<b>Estimated Manpower Cost (10 Years)</b>	<b>Estimated Expense Cost (10 Years)</b>
<b>Objective A</b>	<b>Maintain the park's current recreational carrying capacity of 10,000 users per day.</b>	# of recreation/visitor opportunities per day	C	\$24,000,000	\$12,300,000	\$11,700,000
<b>Objective B</b>	<b>Develop and implement Interpretive Master Plan.</b>	Plan implemented	LT	\$550,000	\$50,000	\$500,000
<b>Goal VIII: Develop and maintain the capital facilities and infrastructure necessary to meet the goals and objectives of this management plan.</b>		<b>Measure</b>	<b>Planning Period</b>	<b>Estimated Manpower and Expense Cost (10-years)</b>	<b>Estimated Manpower Cost (10 Years)</b>	<b>Estimated Expense Cost (10 Years)</b>
<b>Objective A</b>	<b>Maintain all public and support facilities in the park.</b>	Facilities maintained	C	\$24,000,000	\$12,300,000	\$11,700,000
<b>Objective B</b>	<b>Continue to implement the park's transition plan to ensure facilities are accessible in accordance with the Americans with Disabilities Act of 1990.</b>	Plan implemented	LT	\$175,000	\$50,000	\$125,000
<b>Objective C</b>	<b>Improve and/or repair as identified in the Land Use Component.</b>	# of facilities/miles of trail/miles of road	LT	\$4,000,000	\$0	\$4,000,000
Action 1	Pave 6+ miles of existing graded roads—Withlacoochee Bay Trail Road and Kenwood Road.	# of miles paved at these facilities	LT	\$4,000,000	\$0	\$4,000,000
<b>Objective D</b>	<b>Construct facilities needed for operational improvements and increased public visitation.</b>	# of facilities/miles of trail/miles of road	LT	\$4,475,000	\$435,000	\$4,040,000
Action 1	Construct a new bathhouse at Santos Campground and open second camping loop there.	Campground capacity increased	LT	\$350,000	\$20,000	\$330,000
Action 2	Construct Ranger Entrance station at Ross Prairie Campground.	Construction of facility listed	LT	\$125,000	\$15,000	\$110,000

Action 3	Construct Inglis Island to Mullet Point Trail Bridge.	Construction of facility listed	LT	\$4,000,000	\$400,000	\$3,600,000
<b>Objective E</b>	<b>Expand maintenance activities as existing facilities are improved and new facilities are developed.</b>	Facilities maintained	C	<b>\$800,000</b>	<b>\$100,000</b>	<b>\$700,000</b>
<b>Summary of Estimated Costs</b>						
<b>Management Categories</b>				<b>Total Estimated Manpower and Expense Cost* (10-years)</b>		
	Resource Management			\$25,052,750	\$13,997,625	\$11,343,125
	Administration and Support			<b>\$1,387,606</b>	\$1,361,606	\$26,000
	Capital Improvements			<b>\$33,450,000</b>	\$12,885,000	\$20,565,000
	Recreation Visitor Services			<b>\$24,550,000</b>	\$12,350,000	\$12,200,000
	Law Enforcement Activities	Note: Law enforcement activities in Florida State Parks are conducted by the FWC Division of Law Enforcement and by local law enforcement agencies.				

1



**ADDENDUM 1:**

**ACQUISITION HISTORY**



# **Marjorie Harris Carr Cross Florida Greenway State Recreation and Conservation Area**

## **Acquisition History**

(10/10/17)

### **Purpose of Acquisition:**

The Board of Trustees of the Internal Improvement Trust Fund of the State of Florida (Trustees) acquired the Marjorie Harris Carr Cross Florida Greenway State Recreation and Conservation Area (Cross Florida Greenway) to create a cross Florida greenway corridor within the right-of-way of the former Cross Florida Barge Canal, so that it can effectively and efficiently protect, conserve and preserve the natural resources and scenic beauty of Florida, while providing and enhancing compatible public recreational opportunities such as fishing, camping, boating, bicycling, nature study, horseback riding, hiking, hunting, paddling, and other outdoor interests.

### **Sequence of Acquisition:**

On July 26, 1993, Trustees obtained title to approximately 107 miles of former Cross Florida Barge Canal project right-of-way, constituting the initial area of the Cross Florida Greenway. The Trustees acquired this property through a Quitclaim Deed from the United States of America, acting through then Secretary of the Army, John W. Shannon.

Since the initial 1993 acquisition, the Trustees purchased more parcels using different land acquisition programs, mainly Preservation 2000 and Florida Forever, and added them to the Cross Florida Greenway. Between 1995 and 2011, the Trustees received about six (6) donations from private individuals, local governments, private Trustees, and/or private corporations and added them to the Cross Florida Greenway.

Between 1996 and 2011, the State of Florida Department of Environmental Protection, Office of Greenways and Trails (OGT), entered into multiple management lease agreements with several landowners/entities: St. Johns River Water Management District, Southwest Florida Water Management District, and the Felburn Foundation, a private nonprofit corporation. These entities own a portion of the Cross Florida Greenway.

As outlined above, the land that constitutes the present area of the Cross Florida Greenway—about 70,833 acres—primarily came from the right-of-way of the former Cross Florida Barge Canal, new acquisitions by the Trustees, new donations to the Trustees, and leases between OGT and different landowners/entities.

### **Brief History of Cross Florida Barge Canal:**

The following paragraphs are devoted to the Cross Florida Barge Canal. The idea behind this effort is to draw a mental sketch of how the canal project itself started in the first place, without writing a comprehensive history of the canal; rather explaining how the right-of-way of the former Cross Florida Barge Canal ended up being a state-owned greenway corridor, which we now call the Marjorie Harris Carr Greenways State Recreational and Conservation Area.

Many historians believe that the idea of creating an inland waterway across Florida dates to 1567 when Pedro Menendez de Aviles received instruction from his boss, King Philip the 2<sup>nd</sup> of Spain, to explore the Florida peninsula and to determine a suitable water route for crossing the isthmus. The route Pedro Menendez de Aviles recommended to the king was by and large the same as the route that was recommended by the United States Congress for a canal project more than 300 years later.

The history of the construction of the Cross Florida Barge Canal by the United States involves a succession of political controversies, heated opposition from various parties, and two abortive attempts by the federal government to start and complete the construction.

In 1818, Secretary of War, John C. Calhoun, proposed building a canal across Florida to solve the losses due to shipwrecks and piracy associated with open-sea voyage around the tip of the Florida peninsula. Around 1824, Florida's elected territorial representative, Joseph White, asked the federal government for funds to survey the best route and to construct a canal across the state contending that such canal would shorten the distance of going around the south of the state by 1,000 miles. The canal would also benefit the territory by providing the facilities and defenses it needed. After several calls for canal construction by different individuals and groups, the federal government decided, in 1930, that the St. Johns River to Withlacoochee River route was the most desirable, practical and economic route. This route would follow the St. Johns River from its mouth at the Atlantic Ocean to the City of Palatka. Then the route would go along the Ocklawaha River to a point near Silver Springs. From there, the route would continue



westward across uplands below the City of Ocala to the City of Dunnellon, and then the route would follow the course of the Withlacoochee River until it would finally enter the Gulf of Mexico near the small towns of Inglis and Yankeetown. In 1931, the Florida legislature established the Florida State Canal Commission, which was given the power to acquire lands for construction of the proposed canal that was estimated to be approximately two hundred (200) miles long. The construction started in earnest on September 6, 1935, with five million dollars (\$5M) that President Roosevelt had allocated as part of his federal funds to combat unemployment during the Great Depression.

Not long after the canal work started, different entities opposed the construction of the canal both at the local and federal levels. Locally, Central Florida Citrus and Vegetable Growers opposed the canal construction project stating that they would lose labor due to the canal project. Also, port officials from Miami and Tampa feared competition from new ports. Nationally, conservative members of the United States Congress saw the canal project as a symbol of Franklin Roosevelt's profligate spending and unprecedented federal power. Because of these oppositions, within a year of its start, the work of canal construction was effectively and completely stopped, and the cranes and bulldozers fell silent for the following 30 years (from the 1930s to 1960s).

In 1963, President John F. Kennedy (JFK) allocated one million dollars (\$1M) for the canal construction. The following year, after the assassination of JFK, President Lyndon Johnson came to Florida giving a speech outlining the many benefits which would result from constructing the Cross Florida Canal project, and explosives were set off that started construction. However, the following administration undermined the canal project, set in motion by President Johnson. Both President Richard Nixon and Florida Governor Claude Kirk were anxious to consolidate their gains, and they saw a chance to attract more votes by embracing an emerging environmental movement.

At the time, the environmental movement strongly opposed the construction of the Cross Florida Barge Canal and asked a Florida federal court to stop the construction. The court ordered a preliminary halting of further work on the canal construction, and President Nixon supported the court's decision. In 1971, President Nixon signed an executive order causing the canal project to be stopped. In 1976, the Florida governor and cabinet voted to withdraw its support for the canal project. In 1986, the United States Congress de-authorized the canal project, and the

Florida legislature approved the de-authorization four years later. In November of 1990, President George W. Bush de-authorized the federal project, and the 110 miles of canal lands were reverted to the State of Florida. This property was transferred to the State of Florida through a Quitclaim Deed executed by the Secretary of the Army on July 26, 1993. This transfer is subject to certain terms and conditions.

Many major political figures, including several Florida governors and senators, played significant roles in the development and eventual demise of the Cross Florida Barge Canal. However, the work of one Floridian stands out. Marjorie Harris Carr, who was a trained field biologist, housewife and environmental activist/leader, managed to galvanize the efforts to halt the construction of the Cross Florida Barge Canal for good. In recognition of her contribution in this effort, the Florida Legislature in 1998 named the 110-mile corridor of canal lands the “Marjorie Harris Carr Cross Florida Greenways State Recreation and Conservation Area.”

### **Management Lease:**

On October 27, 1993, the Trustees leased the Cross Florida Greenway to the State of Florida Department of Environmental Protection, for the use and benefit of the public, to the Office of Greenways Management under a 50-year term lease, Lease No. 4013. Lease No. 4013 is scheduled to expire on October 27, 2043. The Trustees has amended this lease twenty-seven (27) times mostly to add new parcels it has acquired through purchases or donations.

Per the terms and conditions of Lease No. 4013, the Office of Greenways Management (now commonly known as the Office of Greenways and Trails, or OGT) manages the property as a multiple-use area and for such other purposes as authorized by the provisions of the Greenways Bill adopted by the Florida legislature.

OGT was merged into the State of Florida Department of Environmental Protection, Division of Recreation and Parks, in July 2011. Thus, the Division of Recreation and Park (which is commonly known as “the Florida Park Service”) manages the parcels owned by the Trustees as well as some parcels owned by the St. Johns River Water Management District, the Southwest Florida Water Management District, and the private non-profit Felburn Foundation, as part the Cross Florida Greenway.

**Title Interest:**

The Trustees holds fee simple title to a majority of the lands within Cross Florida Greenway. Additionally, some areas are owned by the St. Johns River Water Management District, the Southwest Florida Water Management District, and the Felburn Foundation.

**Special Conditions on Use:**

The Cross Florida Greenway is designated as a multiple-use property to provide resource-based public outdoor recreation and other park related uses. Uses such as water resource development projects, water supply projects, storm-water management projects, and linear facilities and sustainable agriculture and forestry are not consistent with the purpose for which the State of Florida Department of Environmental Protection, Division of Recreation and Parks, manages the property.

**Outstanding Issues:**

There is no known deed-related restriction or reservation on the use of the Cross Florida Greenway. However, if any entity makes a claim for or against use of any portion of the Cross Florida Greenway, the Office of Park Planning would assist that entity and would validate the existence of such restriction or reservation. The Office of Park Planning can be reached at (850) 245-3051.





**LAND ACQUISITION HISTORY REPORT**

<b>Park Name</b>	Marjorie Harris Carr Cross Florida Greenways State Recreation and Conservation Area				
<b>Date Updated</b>	12/1/2016				
<b>County</b>	Citrus, Levy, Marion and Putnam Counties				
<b>Trustees Lease Number</b>	Trustees Lease No. 4013				
<b>Current Park Size</b>	70,833.51 acres				
<b>Purpose of Acquisition</b>	The Board of Trustees of the Internal Improvement Trust Fund of the State of Florida acquired Marjorie Harris Carr Cross Florida Greenways State Recreation and Conservation Area to create a Cross Florida Greenway corridor within the right-of-way of the former Cross Florida Barge Canal.				
<b>ACQUISITION HISTORY</b>					
<b>Parcel Name or Parcel DM-ID</b>	<b>Date Acquired</b>	<b>Initial Seller</b>	<b>Initial Purchaser</b>	<b>Size in acres</b>	<b>Instrument Type</b>
DMID 310713	10/18/2000 & 10/28/2000	Peter Thralls Miller and his wife Linda C. Miller Ardrenn M. Suttlemyre	The Board of Trustees of the Internal Improvement Trust Fund of the State of Florida (Trustees)	3,864.20	Warranty Deed

Parcel Name or Parcel DM-ID	Date Acquired	Initial Seller	Initial Purchaser	Size in acres	Instrument Type
DMID 310712	8/2/2000	Earl M. Miller Jr. Margarete Miller Carlson	Trustees	3,323.54	Warranty Deed
DMID 310710	6/30/2000	Earl M. Miller Jr. Joe C. Miller and his wife Judith G. Shine Douglas M. Miller and his wife Barbara I. Miller	Trustees	2,877.00	Warranty Deed
DMID 310786	19/09/2000	Steve W. Conner	Trustees	2,292.15	Warranty Deed
DMID 347650	NA (no date)	NA (the amendment has only legal des.)	NA	1,940.00	NA
DMID 310784	9/23/2003	P.N. Philips and his wife Elizabeth Ann Philips	Trustees	1,541.87	Warranty Deed
DMID 310711	1/9/2001	Joe C. Miller Jr . and his wife Judith G. Sine	Trustees	1,330.02	Warranty Deed
DMID 310714	8/31/1999	Douglas M. Miller and his wife Barbara L. Miller	Trustees	702.99	Quitclaim Deed
DMID 310709	8/23/1999	Silver Run Properties, Inc.	Trustees	630.8	Warranty Deed
DMID 383104	2/28/2003	James Theodore Miller John Robert Miller Peter Thralls Miller Susan Eleanor Miller Thomas	Trustees	578	Warranty Deed
DMID 310716	No date given	Huntington National Bank of Florida	Trustees	446.18	S. W. Deed

<b>Parcel Name or Parcel DM-ID</b>	<b>Date Acquired</b>	<b>Initial Seller</b>	<b>Initial Purchaser</b>	<b>Size in acres</b>	<b>Instrument Type</b>
DMID 310717	6/15/2001	Herb Davis Plumbing, Inc.	Trustees	398.8	Warranty Deed
DMID 335205	5/11/2004	James Douglas Mcdowall and his wife Linda Anne Mcdowall	Trustees	255.27	Warranty Deed
DMID 327029	6/13/2003	Silver Springs Shore Land Trust LTD	Trustees	203.45	Warranty Deed
DMID 353161	1/2/2008	Robert P. Drake, Individually and as Trustees	Trustees	107.63	Warranty Deed
DMID 355888	12/11/2006	Felburn Foundation	Trustees	44.33	Warranty Deed
DMID 365899	12/28/2010	Thomas Rex Dull	Trustees	44.33	Warranty Deed
DMID 367904	12/30/2010	Citrus Mining & Timber Inc.	Trustees	18.54	Warranty Deed
<b>MANAGEMENT LEASE</b>					
<b>Parcel Name or Lease Number</b>	<b>Date Leased</b>	<b>Initial Lessor</b>	<b>Initial Lessee</b>	<b>Current Term</b>	<b>Expiration Date</b>
Trustee Lease No. 4013	10/27/1993	The Board of Trustees of the Internal Improvement Trust Fund of the State of Florida	State of Florida Department of Environmental Protection, for the use and benefit of Office of Greenways Management	50 years	10/26/2043

<b>Outstanding Issue</b>	<b>Type of Instrument</b>	<b>Brief Description of the Outstanding Issue</b>	<b>Term of the Outstanding Issue</b>
<p>There is no known deed-related restriction or reservation that applies to this property. However, if any entity makes a claim related to use of the property, please call Office Park Planning (850-245-3051) for assistance</p>			



**ADDENDUM 2:**  
**REFERENCES**



## REFERENCES

- Brooks, H.K. 1981. *Guide to the Physiographic Divisions of Florida*. Institute of Food and Agricultural Sciences. Gainesville, Fla.: University of Florida.
- Brush, Janell. Avian Research Scientist, Florida Fish and Wildlife Conservation Commission. Personal communication shorebird data for Cross Florida Greenway spoil islands.
- Bureau of Economic and Business Research (BEBR). 2011. *Florida Statistical Abstract*. Warrington College of Business Administration, University of Florida.
- Citrus County. 2015. *Citrus County Comprehensive Plan*. Citrus County, Florida.
- City of Ocala. 2010. *City of Ocala Comprehensive Plan*. Ocala, Florida.
- Davis, A., and D.R. Jackson. 2004. *Marjorie Harris Carr Cross Florida Greenway Rare Animal Survey, Final Report*. Submitted to Florida Department of Environmental Protection, Office of Greenways and Trails. Florida Natural Areas Inventory, Florida State University.
- Faulkner, Glen L. 1973. *Geohydrology of the Cross Florida Barge Canal Area with Special Reference to the Ocala Vicinity*. U.S. Geological Survey Water Resources Investigations Report I-73. Prepared in cooperation with the U.S. Army Corps of Engineers.
- Fernald E.A., and E.D. Purdum. 1998. *Water Resources Atlas of Florida*. Florida State University, Institute of Science and Public Affairs.
- Florida Department of Environmental Protection (FDEP). 2001. *Group 1 Basin Status Report: Ocklawaha*. FDEP, Division of Water Resource Management, November 2001. 315pp.
- FDEP. 2010. *Florida State Park System Economic Impact Assessment for Fiscal Year 2010/2011*. Tallahassee, Florida.
- FDEP. 2005. *Elements of Florida's Water Monitoring and Assessment Program*. Florida Department of Environmental Protection, Tallahassee, Florida Technical Report 113pp.
- FDEP. 2008a. Florida Department of Environmental Protection STORET database. Web-based water quality data database. <http://storet.dep.state.fl.us/WrmSpa/>
- FDEP. 2008b. Florida Department of Environmental Protection Bureau of Laboratories. Ecosummary Reports. <http://www.floridadep.org/labs/cgibin/reports/search.asp>
- Florida Exotic Pest Plant Council (FLEPPC). 2009. *Florida Exotic Pest Plant Council's 2009 List of Invasive Plant Species*. Internet: <http://www.fleppc.org/list/list.htm> or Wildland Weeds Vol. 12(4):13-16. Fall 2009.
- Florida Memory State Library and Archives of Florida. Florida archived photos of Marjorie Harris Carr Cross Florida Greenway. Florida Department of State, Division of Library and Information Services. <https://www.floridamemory.com/solr-search/results/?q=%28barge%20canal%20OR%20tt%3Abarge%20canal%5E10%29%20AND%20collection%3A%22Florida%20Photographic%20Collection%22&searchbox=1&query=barge%20canal&year=&gallery=0&search-type>

Florida's Wildlife Legacy Initiative: Florida's State Wildlife Action Plan. Tallahassee, Florida, USA.

Florida Natural Areas Inventory (FNAI). 2010. *Guide to the Natural Communities of Florida: 2010 edition*. Florida Natural Areas Inventory, Tallahassee, FL.

Levy County. 2015. *Levy County Comprehensive Plan*. Levy County, Florida.

Marion County. 2015. *Marion County Comprehensive Plan*. Marion County, Florida.

Putnam County. 2015. *Putnam County Comprehensive Plan*. Putnam County, Florida.

Scott, 2001. Text to Accompany the Geologic Map of Florida. Florida Geological Survey, Tallahassee. Open File Report No. 80.

Scott, T.M., G.H. Means, R.P. Meegan, R.C. Means, S.B. Upchurch, R.E. Copeland, J. Jones, T. Roberts, and A. Willet. 2004. *Springs of Florida*. Florida Geological Survey Bulletin No. 66. 377pp.

SCS Engineers. 2010. *Site Investigation Report for the Old Canal Green Box Recycling Center, South Highway 475, Ocala, Marion County, Florida*. Report to Marion County Solid Waste Department, Ocala, Florida. 30pp.

([https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/use/hydric/?cid=nrcs142p2\\_053961hydric](https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/use/hydric/?cid=nrcs142p2_053961hydric))

St. Johns River Water Management District. 2011. Hydrologic data. Online GIS data retrieval system. <http://www.sjrwmd.com/toolsGISdata/index.html>

United States Department of Commerce, Bureau of the Census. 2010. U.S. Census 2010.

(<http://www.floridacaving.com/pages/misc/geoflorida.htm>)



**ADDENDUM 3:  
PRELIMINARY PUBLIC WORKSHOP  
SUMMARY**



# Marjorie Harris Carr Cross Florida Greenways State Recreation and Conservation Area

## *SUMMARY OF PRELIMINARY PUBLIC WORKSHOP*

*December 2016*

---

As part of the planning process for the 2017 Unit Management Plan for the Marjorie Harris Carr Cross Florida Greenways State Recreation and Conservation Area, three (3) preliminary public workshops were held in three locations along the Greenway. On November 25, 2016, public advertisements were published in the *Florida Administrative Register* and the following local newspapers: the *Palatka Daily News*, the *Ocala Star Banner*, and the *Citrus County Chronicle*. Affidavits of the newspaper advertisements are provided in **Attachment A**.

The preliminary public workshops were held as follow and scanned copies of the sign-in sheets from each meeting are provided in **Attachment B**.

- **December 6, 2016 (5:30-7:30 PM)**  
St. Johns Water Management District Governing Board Room  
4049 Reid Street  
Palatka, FL 32177
  - 26 attendees – Five (5) written comments received at this workshop.
  
- **December 7, 2016 (5:30-7:30 PM)**  
Marion County Growth Services Training Room  
2710 East Silver Springs Boulevard  
Ocala, FL 34470
  - 31 attendees – Two (2) written comments received at this workshop.
  
- **December 8, 2016 (5:30-7:30 PM)**  
Inglis Community Center  
137 Highway 40 West  
Inglis, FL 34449
  - 12 attendees – Four (4) written comments received at this workshop.

An online survey was also employed to obtain feedback from the public as a result of the preliminary public workshops. The survey will remain open and public comments accepted until December 21, 2016, with results included in the public comment documentation.

**Marjorie Harris Carr Cross Florida Greenways  
State Recreation and Conservation Area**

*SUMMARY OF PRELIMINARY PUBLIC WORKSHOPS  
December 2016*

---

**ATTACHMENT A**

**AFFIDAVITS FOR PUBLICATION  
OF NEWSPAPER ADVERTISEMENTS**



# AFFIDAVIT OF PUBLICATION

**Star-Banner**  
Published – Daily  
Ocala, Marion County, Florida

STATE OF FLORIDA  
COUNTY OF MARION

Before the undersigned, a Notary Public of Said County and State, [Signature] who on oath says that they are an authorized employee of the Star-Banner, a daily newspaper published at Ocala, in Marion County, Florida; that the attached copy of advertisement, being a notice in the matter of

*The Florida Department of Environmental Protection, Division of Recreation and Parks, announces a public workshop to which all persons are invited. DATE AND TIME: Wednesday, December 7, 2016, 5:30 p.m. EST. PLACE: Marion County Growth Services Training Ro*

was published in said newspaper in the issues of:

11/25 1x

Affiant further says that the said STAR-BANNER is a daily newspaper published at Ocala, in said Marion County, Florida, and that the said newspaper has heretofore been continuously published in said Marion County, Florida, daily, and has been entered as second class mail matter at the post office in Ocala in said Marion County, Florida, for a period of one year next preceding the first publication of the attached copy of advertisement; and affiant further says that he has neither paid nor promised any person, firm or corporation any discount, rebate, commission or refund for the person of securing this advertisement for publication in the said newspaper.

Sworn to and subscribed before me this 25 day of November, A.D., 2016



[Signature]  
Notary Public  
**HARMONY STALTER**  
(Print, Type or Stamp Name of Notary Public)

The Florida Department of Environmental Protection, Division of Recreation and Parks, announces a public workshop to which all persons are invited.  
**DATE AND TIME:** Wednesday, December 7, 2016, 5:30 p.m. (EST).  
**PLACE:** Marion County Growth Services Training Room, 2710 East Silver Springs Boulevard, Ocala, Florida 34470.  
**GENERAL SUBJECT MATTER TO BE CONSIDERED:** The purpose of this workshop is to gather input for the update to the Marjorie Harris Carr Cross Florida Greenways State Recreation and Conservation Area Unit Management Plan at the beginning of the planning process.  
**A COPY OF THE AGENDA MAY BE OBTAINED BY CONTACTING:** Mickey Thomason, Park Manager, 8262 SE Highway 314, Ocala, Florida 33470, PH # (352) 236-7143 or email [mickey.thomason@dep.state.fl.us](mailto:mickey.thomason@dep.state.fl.us). A copy of the agenda is available before the date of the public workshop online at <https://www.fldepnet.org/public-notices>. Pursuant to the provisions of the Americans with Disabilities Act, any person requiring special accommodations to participate in this workshop is asked to advise the agency at least 48 hours before the workshop by contacting: Mickey Thomason as listed above.  
If you are hearing or speech impaired, please contact the agency using the Florida Relay Service, 1(800)955-8771 (TDD) or 1(800)955-8770 (Voice)

November 25, 2016  
#A000900598

Ad #: A000900598

STATE OF FLORIDA

County of Putnam

The undersigned personally appeared before me, a Notary Public for the State of Florida, and deposes that the Palatka Daily News is a daily newspaper of general circulation, printed in the English language and published in the City of Palatka in said County and State; and that the attached order, notice, publication and/or advertisement:

The Florida Department of Envi

Was published in said newspaper 1 time with said being made on the following dates:

11/25/2016

The Palatka Daily News has been continuously published as a daily newspaper, and has been entered as second class matter at the post office at the City of Palatka, Putnam County, Florida, each for a period of more than one year next preceding the date of the first publication of the above described order, notice and/or advertisement.

*Jeannette Eveland*

Sworn to and subscribed to before me this 25th day of November, 2016 by Jeannette Eveland, Administrative Assistant, of the Palatka Daily News, a Florida corporation, on behalf of the corporation.

*Mary Kaye Wells*

Mary Kaye Wells, Notary Public  
My commission expires: July 22, 2019

Notary Seal  
Seal of Office:

- Personally known to me, or
- Produced identification:
- Did take an oath



**PUBLIC NOTICE**

The Florida Department of Environmental Protection, Division of Recreation and Parks, announces a public workshop to which all persons are invited.

**DATE AND TIME:** Tuesday, December 6, 2016, 5:30 p.m. (EST).

**PLACE:** St. Johns River Water Management District Governing Board Room, 4049 Reid Street, Palatka, Florida 32177.

**GENERAL SUBJECT MATTER TO BE CONSIDERED:** The purpose of this workshop is to gather input for the update to the Marjorie Harris Carr Cross Florida Greenways State Recreation and Conservation Area Unit Management Plan at the beginning of the planning process.

**A COPY OF THE AGENDA MAY BE OBTAINED BY CONTACTING:** Mickey Thomason, Park Manager, 8282 SE Highway 314, Ocala, Florida 33470, PH # (352) 236-7143 or email [mickey.thomason@dep.state.fl.us](mailto:mickey.thomason@dep.state.fl.us). A copy of the agenda is available before the date of the public workshop online at <https://www.fldepnet.org/public-notices>.

Pursuant to the provisions of the Americans with Disabilities Act, any person requiring special accommodations to participate in this workshop is asked to advise the agency at least 48 hours before the workshop by contacting: Mickey Thomason as listed above.

If you are hearing or speech impaired, please contact the agency using the Florida Relay Service, 1(800)955-8771 (TDD) or 1(800)955-8770 (Voice).

11/25/16  
Legal No. 00047068

# Proof of Publication

from the  
**CITRUS COUNTY CHRONICLE**  
Crystal River, Citrus County, Florida  
**PUBLISHED DAILY**

STATE OF FLORIDA  
COUNTY OF CITRUS

Before the undersigned authority personally appeared


John Murphy and/or Mary Ann Naczi and/or Mishayla Coffas

Of the Citrus County Chronicle, a newspaper published daily at Crystal River, in Citrus County, Florida, that the attached copy of advertisement being a public notice in the matter of the

8066-1125 FCRN The Florida Department of Environmental Protection, Division of Recreation and Parks, announces a public workshop to which all persons are invited. DATE AND TIME: Thursday, December 8, 2016, 5:30 p.m. (EST). PLACE: Inglis Commu

Court, was published in said newspaper in the issues of November 25th, 2016,


Affiant further says that the Citrus County Chronicle is a Newspaper published at Crystal River in said Citrus County, Florida, and that the said newspaper has heretofore been continuously published in Citrus County, Marion County and Levy County, Florida, each week and has been entered as second class mail matter at the post office in Inverness in said Citrus County, Florida, for a period of one year next preceding the first publication of the attached copy of advertisement; and affiant further says that he/she has neither paid nor promised any person, firm or corporation any discount, rebate, commission or refund for the purpose of securing this advertisement for publication in the said newspaper.




The foregoing instrument was acknowledged before me

This 25<sup>th</sup> day of Nov., 2016  
By: John Murphy and/or Mary Ann Naczi and/or Mishayla Coffas

who is personally known to me and who did take an oath.

  
Notary Public



**Marjorie Harris Carr Cross Florida Greenways  
State Recreation and Conservation Area**

*SUMMARY OF PRELIMINARY PUBLIC WORKSHOPS*

*December 2016*

---

**ATTACHMENT B**

**SIGN-IN SHEETS FROM  
PRELIMINARY PUBLIC WORKSHOPS**

































**ADDENDUM 4:  
TIMBER MANAGEMENT ANALYSIS FROM  
TIMBER ASSESSMENT**



Addendum 4  
Timber Management Analysis

1. *Management Context and Best Management Practices*

Timber management prescriptions and actions on the Cross Florida Greenway (CFG) are based on the desired future condition (DFC) of a stand or natural community (NatCom) as determined by guidelines determined by the Division of Recreation and Parks (DRP). In most cases, the DFC will be closely related to the historic NatCom. However, where the historic community has been severely altered by past land use practices, the DFC may not always be the same as the historic NatCom. All forest/stand/timber management activities undertaken will adhere to the current Florida Silvicultural Best Management Practices and Florida Forestry Wildlife Best Management Practices for State Imperiled Species. DRP is responsible for managing timber resources within corresponding management zones. This timber assessment was conducted by F4 Tech on behalf of DRP.

2. *Purpose of Timber Management Activities*

Timber management activities will be conducted to help restore and/or improve current conditions so that the associated DFC (typically an historic condition) can be achieved or maintained. Timber management will primarily be conducted in pine-dominated natural communities. Upland communities typically include mesic flatwoods, sandhill, upland pine, upland mixed woodland, and altered landcover areas such as successional hardwood forest and pine plantations. Other historically hardwood-dominated natural communities will likely have little to no scheduled timber management activities. In some circumstances, actions may be conducted to remove overstory invasive/exotic trees, e.g. Melaleuca, Chinese tallow, Brazilian pepper, occupying contiguous areas of land to help restore or maintain natural communities.

3. *Potential Silvicultural Treatments*

Several silvicultural treatments may be considered and utilized over the next ten years to achieve the long-term DFC for candidate natural communities at the CFG. These treatments include timber harvests, timber stand improvement, and reforestation. The various types of timber harvests may include pine thinning, targeted hardwood removal, and clearcutting. Silvicultural treatments should be implemented to minimize disturbance to non-target vegetation, soil, and wildlife.

Thinning is conducted to reduce the basal area (BA) or density of stems in a stand to **improve forest health and growth conditions for residual trees.** The “opening up” of high density forest stands increases tree and stand vigor, which helps mitigate the potential for damaging insect outbreaks. Thinning also increases sunlight reaching the forest floor, which when combined with routine prescribed fire, can increase groundcover vegetation abundance, species richness, and overall ecological diversity. The disruption of a historic natural fire regime and/or fire return interval can often result in the need to remove undesirable or overstocked hardwood stems that currently occupy growing space in the canopy and sub-canopy. Tree removal/harvest also increases groundcover vegetation, ecological diversity, and fine fuels that facilitate consistent fire return intervals and responses.

## Florida State Parks Timber Management Analyses

Clearcutting supports restoration goals by removing offsite pine or hardwood species and is a precursor to establishing site-appropriate species. It is also used to control insect infestations that are damaging or threatening forest resources and ecosystem conditions on or off site.

A tangible by-product of conducting timber harvests for restoring or improving forested communities is the generation of revenue.

Stand or natural community improvement activities are often conducted to reduce unwanted hardwood or palmetto competition. Stand improvement treatments reduce fuel or fuel height, which can improve groundcover conditions and aid in maintaining proper prescribed burning return intervals. The two main stand improvement activities used on park property are herbicide treatments and mechanically cutting vegetation. Herbicide may be applied aerially, by mechanized ground-based equipment, or via backpack sprayers. Herbicides are used to reduce the amount of hardwood competition in areas that are unable to carry sufficient prescribed fire due to shading and lack of adequate groundcover fuels. Mechanical cutting is used to reduce the height of smaller shrub and hardwood competition, allowing for the establishment of fire-dependent herbs and grasses. Decreasing fuel loadings and enhancing groundcover allows prescribed fire to be reintroduced safely into a stand that has been unable to carry fire adequately.

Reforestation is used to establish the appropriate southern pine species in areas that have been harvested and lack sufficient natural regeneration in terms of abundance (seedlings/acre) and/or species composition. Reforestation candidate areas can also include those that are fire suppressed or have been recently impacted by natural events such as windthrow, bark beetle attack, or wildfire. The two methods used to reestablish the overstory will be natural and artificial regeneration. Both methods may require site preparation to facilitate survival of the desired species. Site preparation activities may include the use of prescribed fire, herbicides, and/or mechanical treatments such as roller chopping. Site preparation technique(s) will be selected that address the current vegetative cover type and condition, and the need to minimize seedling competition while avoiding/minimizing any long-term impacts to native groundcover species and native wildlife. Where artificial regeneration is not needed, natural regeneration may be used, specifically in areas that have an adequate seed source of the desired tree species located on site or in the immediate vicinity. Artificial regeneration may include machine or hand planting. Hand planting is preferred on wetter sites, rougher sites, and/or sites where groundcover protection is a concern and a more natural appearance of randomly spaced trees is desired. Machine planting generally allows for more consistent planting and often allows higher survival rates if the site is properly prepared.



## Florida State Parks Timber Management Analyses

### *4. Inventory Data and Potential Actions per Area of Interest or Management Zone*

CFG comprises 70,531 acres across Citrus, Levy, Marion and Putnam Counties. A total of 28,005 acres are associated with several upland natural communities that are potential candidates for timber management. For this region, upland natural communities include mesic flatwoods, mesic hammock, sandhill, scrub, scrubby flatwoods, upland hardwood forests, upland pine forest, xeric hammock, and altered communities/landcovers such as clearcut pine plantation, pine plantation, and invasive-exotic monoculture. From January 2016 through April 2016, a plot-based forest/vegetation inventory was conducted across and within these areas to quantify overstory, midstory and understory conditions. Table 1 below provides general statistics generated by this inventory of the CFG. Table 2 below provides current stocking levels and potential management activities of candidate management zones and natural communities.

This timber assessment was based on GIS data (management zone and NatCom boundary data) provided by DRP in June 2016. This assessment identifies opportunities for potential actions over the next 10-year UMP planning horizon (2017-2027) based on current conditions compared against desired future conditions. It is not intended to be prescriptive. State park staff responsible for developing operational plans should view this timber assessment and all supporting data as a guide for potential actions to consider. Given the dynamic nature of property ownership and land management activities at the CFG, together with the timeframe required to create or update a UMP (12-18 months), it is possible that some tabular data may be dated. Therefore, NatCom acreages and recent treatments that occurred after the June 2016 period are not reflected in the tables herein.

A review and analysis of this data suggests that current ecological conditions for multiple management zones and associated forested communities could benefit from vegetation treatments (non-revenue generating). This assessment was based on a comparison of current conditions and the corresponding natural community analog or target conditions as defined per Florida Natural Areas Inventory (FNAI) Reference Site descriptions. In general, inventory data indicates that upland habitats in several management zones have an average pine BA that is outside the acceptable range for the DFC of the natural community types. Some natural communities considered may need midstory and overstory control to become, or remain, in compliance with FNAI defined ranges for palmetto and non-pine midstory. Stands with low stocking levels or a complete lack of preferred tree species would likely benefit from midstory control and artificial regeneration. In areas where planting is deemed necessary, the site should be assessed for site preparation needs including midstory/understory reduction.

The following contains a general description of each management zone within the CFG that contains upland natural communities as well as their general condition and need for restoration and/or improvement actions via timber management.

Florida State Parks  
Timber Management Analyses

Table 1. General summary statistics for CFG

Number of Management Zones within the Park	734
Number of Management Zones needing timber management	438
Number of unique upland Natural Community Polygons (split by management zone)	571
Number of unique upland Natural Community Polygons potentially needing timber management	571
Upland Natural Community acres	28,005
Acres potentially needing timber management	28,005

Clearcut Pine Plantation (161 acres)

Clearcut pine plantation is an altered community type which resulted from past commercial forest management. The dominant overstory species found in this type is longleaf pine (*Pinus palustris*), loblolly pine (*P. taeda*), or slash pine (*P. elliottii*). As an altered forest/community type, and potentially a candidate for restoration, there are currently no FNAI recommendations on preferred species or stocking levels. Future timber management activities could potentially transition this altered type into another natural community type per on-site and/or nearby land-use conditions.

Management Zones	Clearcut Pine Plantation (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-E007*	2	--	--	--	--
CFG-E008	77	114	85	28	4.6
CFG-E032*	2	--	--	--	--
CFG-E074*	2	--	--	--	--
CFG-E088*	4	--	--	--	--
CFG-E105*	<1	--	--	--	--
CFG-E106*	1	--	--	--	--
CFG-E107*	<1	--	--	--	--

Florida State Parks  
Timber Management Analyses

Management Zones	Clearcut Pine Plantation (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-E108*	<1	--	--	--	--
CFG-E110*	4	--	--	--	--
CFG-E111	14	58	20	38	3.7
CFG-E142*	<1	--	--	--	--
CFG-E143	14	67	56	10	4
CFG-E144*	2	--	--	--	--
CFG-E145	18	28	9	19	3.5
CFG-E199	9	32	32	0	4.6
CFG-E200*	5	--	--	--	--

\*Unsampled upland areas are present in this analysis and could require vegetation management in the future.

### Invasive Exotic Monoculture (35 acres)

Invasive exotic monoculture is an altered community type which resulted from infestation of one or more species of invasive exotic. There is no preferred species in the overstory. As an altered forest/community type, and potentially a candidate for restoration, there are currently no FNAI recommendations on preferred species or stocking levels. Future timber management activities could potentially transition this altered type into another natural community type per on-site and/or nearby land-use conditions.

Management Zones	Invasive Exotic Monoculture (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C167	34	110	--	110	18.6
CFG-C168*	<1	--	--	--	--

\*Unsampled upland areas are present in this analysis and could require vegetation management in the future.

### Mesic Flatwoods (5,041 acres)

Dominant pine in mesic flatwoods in the region is usually longleaf pine (*P. palustris*) with occasional stands of slash pine (*P. elliotii*) in coastal situations adjacent to tidal marsh. Native herbaceous groundcover will cover at least 50% of the area at a height of less than three feet. Saw palmetto (*Serenoa repens*) will comprise less than 50% of the total shrub cover, also at a height of less than three feet. Other common shrub species may include gallberry (*Ilex glabra*), winged sumac (*Rhus copallinum*),

Florida State Parks  
Timber Management Analyses

fetterbush (*Lyonia lucida*), wax myrtle (*Myrica cerifera*), yaupon holly (*Ilex vomitoria*), running oak (*Quercus pumila*), pawpaw (*Asimina* spp.), dwarf live oak (*Quercus minima*), shiny blueberry (*Vaccinium myrsinites*), and coontie (*Zamia pumila*). The optimal fire return interval for this community is two to three years. The preferred pine species, as determined by FNAI reference sites, is longleaf pine and should be stocked at a level of 10 to 50 square feet per acre BA while non-pine species should remain between 0 and 26.2 stems per acre. The following management zone(s) contain mesic flatwoods which could be considered for some form of timber management including overstory removal, midstory mitigation, site preparation, and planting of preferred pine species.

Management Zones	Mesic Flatwoods (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C012	35	23	3	20	7.7
CFG-C013*	<1	--	--	--	--
CFG-C018*	2	--	--	--	--
CFG-C020*	33	--	--	--	--
CFG-C021*	5	--	--	--	--
CFG-C036*	<1	--	--	--	--
CFG-C038*	2	--	--	--	--
CFG-C039	5	60	--	60	14.1
CFG-C040	7	100	--	100	13.7
CFG-C041*	2	--	--	--	--
CFG-C042*	<1	--	--	--	--
CFG-C043*	1	--	--	--	--
CFG-C044	14	118	20	98	8.2
CFG-C045	10	99	1	97	8.2
CFG-C046	42	83	11	72	8.5
CFG-C047*	<1	--	--	--	--
CFG-C048	302	112	13	98	9.4
CFG-C049*	7	--	--	--	--
CFG-C054*	1	--	--	--	--
CFG-C058	21	85	12	73	6.2
CFG-C060	38	157	17	139	9.7
CFG-C061*	3	--	--	--	--
CFG-C062	67	144	4	139	7.6
CFG-C063	26	125	60	65	6
CFG-C064	34	180	6	173	8
CFG-C065	10	92	--	92	7.3



Florida State Parks  
Timber Management Analyses

Management Zones	Mesic Flatwoods (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C066	15	95	--	95	8.8
CFG-C067	33	133	--	133	6.2
CFG-C068	63	128	--	128	5.7
CFG-C069	15	105	62	43	6.7
CFG-C070	49	94	--	94	8.8
CFG-C071	26	118	--	118	7.1
CFG-C072	52	117	10	106	7.7
CFG-C074	135	98	--	98	5.7
CFG-C075	22	66	28	38	6.4
CFG-C076	10	2	--	2	0.7
CFG-C077	53	118	22	96	6.7
CFG-C078*	37	--	--	--	--
CFG-C079	60	110	24	85	6.3
CFG-C080*	1	--	--	--	--
CFG-C081	3	100	--	100	9.6
CFG-C097*	4	--	--	--	--
CFG-C098*	<1	--	--	--	--
CFG-C103	5	188	10	178	15.7
CFG-C104	30	124	20	104	11.8
CFG-C114	6	105	--	105	9.5
CFG-C115*	<1	--	--	--	--
CFG-C116*	<1	--	--	--	--
CFG-C254	3	163	--	163	13.5
CFG-C255	49	154	1	153	7.8
CFG-E015*	<1	--	--	--	--
CFG-E016	8	60	--	60	6
CFG-E017*	<1	--	--	--	--
CFG-E020	5	105	5	100	11.7
CFG-E021*	4	--	--	--	--
CFG-E044	18	45	--	45	4.5
CFG-E045*	3	--	--	--	--
CFG-E046	3	19	--	19	2.1
CFG-E051*	4	--	--	--	--
CFG-E052*	<1	--	--	--	--
CFG-E053*	42	--	--	--	--
CFG-E054*	<1	--	--	--	--
CFG-E059*	<1	--	--	--	--

Florida State Parks  
Timber Management Analyses

Management Zones	Mesic Flatwoods (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-E060	9	52	--	52	5.8
CFG-E062*	1	--	--	--	--
CFG-E065	8	96	5	91	6.9
CFG-E068*	4	--	--	--	--
CFG-E070*	13	--	--	--	--
CFG-E071*	<1	--	--	--	--
CFG-E072*	34	--	--	--	--
CFG-E073*	1	--	--	--	--
CFG-E079	<1	51	--	51	4.1
CFG-E085*	<1	--	--	--	--
CFG-E092*	<1	--	--	--	--
CFG-E093*	<1	--	--	--	--
CFG-E094*	19	--	--	--	--
CFG-E097	133	48	1	47	5.2
CFG-E099	7	125	--	125	6.5
CFG-E100*	12	--	--	--	--
CFG-E103	17	24	--	24	3.3
CFG-E124*	<1	--	--	--	--
CFG-E125*	<1	--	--	--	--
CFG-E126*	<1	--	--	--	--
CFG-E135*	<1	--	--	--	--
CFG-E136*	4	--	--	--	--
CFG-E138*	1	--	--	--	--
CFG-E139*	1	--	--	--	--
CFG-E140	33	62	20	42	8.9
CFG-E142	62	59	--	59	7.2
CFG-E145*	<1	--	--	--	--
CFG-E154*	2	--	--	--	--
CFG-E155*	11	--	--	--	--
CFG-E156*	22	--	--	--	--
CFG-E157*	1	--	--	--	--
CFG-E158*	<1	--	--	--	--
CFG-E159*	1	--	--	--	--
CFG-E160*	1	--	--	--	--
CFG-E162	33	41	12	29	8.7
CFG-E163	57	39	22	17	12.8
CFG-E164*	<1	--	--	--	--

Florida State Parks  
Timber Management Analyses

Management Zones	Mesic Flatwoods (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-E165*	<1	--	--	--	--
CFG-E166*	<1	--	--	--	--
CFG-E167*	<1	--	--	--	--
CFG-E168	126	94	10	83	11.2
CFG-E169	97	95	--	95	10.9
CFG-E170	50	118	--	118	9.7
CFG-E171*	27	--	--	--	--
CFG-E172	22	136	--	136	10.7
CFG-E174*	<1	--	--	--	--
CFG-E175	71	100	16	83	11.7
CFG-E178	114	93	1	91	9.3
CFG-E179	131	81	13	67	10.6
CFG-E180	143	73	20	53	11.5
CFG-E181	144	83	4	79	9.9
CFG-E182	60	112	8	104	9.9
CFG-E185*	<1	--	--	--	--
CFG-E186	39	61	13	47	6.1
CFG-E187*	<1	--	--	--	--
CFG-E188	24	102	20	82	7.1
CFG-E204*	1	--	--	--	--
CFG-E205*	<1	--	--	--	--
CFG-E206*	2	--	--	--	--
CFG-E208	6	79	--	79	7.2
CFG-E209	29	2	2	0	2.2
CFG-E210	76	95	0	94	9.2
CFG-E211	64	106	12	94	11.9
CFG-E213*	<1	--	--	--	--
CFG-E214*	<1	--	--	--	--
CFG-E215	168	99	3	95	11.5
CFG-E216	129	94	2	92	11.4
CFG-E217*	4	--	--	--	--
CFG-E218	32	76	2	74	8.8
CFG-E219	82	68	0	67	10.5
CFG-E220*	<1	--	--	--	--
CFG-E221*	2	--	--	--	--
CFG-E222*	3	--	--	--	--
CFG-E223	176	100	6	93	12

Florida State Parks  
Timber Management Analyses

Management Zones	Mesic Flatwoods (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-E224	74	76	6	70	10.8
CFG-E225	112	75	7	68	11.7
CFG-E226*	<1	--	--	--	--
CFG-E227*	2	--	--	--	--
CFG-E229*	<1	--	--	--	--
CFG-E230	2	62	--	62	12.5
CFG-E231	29	54	1	53	8.3
CFG-E232*	1	--	--	--	--
CFG-E233*	<1	--	--	--	--
CFG-E234	56	83	12	70	9.6
CFG-E235	44	146	--	146	11.8
CFG-E236	21	120	--	120	11.7
CFG-E237*	7	--	--	--	--
CFG-E238	117	98	1	96	10.3
CFG-E239	3	90	20	70	13.8
CFG-E240	20	126	--	126	11.9
CFG-E241	7	82	10	72	10.5
CFG-E242*	<1	--	--	--	--
CFG-E279*	35	--	--	--	--
CFG-E283*	<1	--	--	--	--
CFG-E285*	12	--	--	--	--
CFG-E286*	4	--	--	--	--
CFG-E288*	39	--	--	--	--
CFG-E295*	14	--	--	--	--
CFG-E296*	6	--	--	--	--
CFG-E297*	1	--	--	--	--
CFG-E298*	<1	--	--	--	--
CFG-E300	27	59	--	59	5.1
CFG-E301*	<1	--	--	--	--
CFG-E308*	15	--	--	--	--
CFG-E309*	<1	--	--	--	--
CFG-W087	49	55	35	20	10.4
CFG-W089*	11	--	--	--	--
CFG-W090*	2	--	--	--	--
CFG-W092	30	96	32	63	11.2
CFG-W093	6	70	13	57	13.7
CFG-W094	73	61	38	23	10.7



Florida State Parks  
Timber Management Analyses

Management Zones	Mesic Flatwoods (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-W095	10	57	30	27	11.7
CFG-W108	24	76	62	14	8.6
CFG-W109*	<1	--	--	--	--
CFG-W123	8	81	--	81	8
CFG-W124*	2	--	--	--	--
CFG-W126	48	99	20	79	8.4
CFG-W129*	<1	--	--	--	--
CFG-W130*	7	--	--	--	--
CFG-W132	7	102	10	92	13.2
CFG-W135*	<1	--	--	--	--

\*Unsampled upland areas are present in this analysis and could require vegetation management in the future.

Mesic Hammock (4,313 acres)

Mesic hammocks are characterized by a well-developed evergreen hardwood and/or palm forest which can occur through much of peninsular Florida. The canopy, often dense, will typically be dominated by live oak (*Q. virginiana*) with cabbage palm (*Sabal palmetto*) mixed into the understory. Southern magnolia (*Magnolia grandiflora*) and pignut hickory (*Carya glabra*) can be common components in the subcanopy. Pine trees, particularly slash pine or loblolly pine (*P. taeda*), may form a sparse emergent layer. Mesic hammocks can arise in naturally pine-dominated areas when shielded from fire because of human activities and timber management activities can support restoration goals. Preferred species for this NatCom include those specified above. The following management zone(s) contain mesic hammock which could be considered for some form of timber management including overstory removal, midstory mitigation, site preparation, and planting of preferred pine species.

Management Zones	Mesic Hammock (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C002*	3	--	--	--	--
CFG-C003*	1	--	--	--	--
CFG-C004*	1	--	--	--	--
CFG-C005	19	147	65	82	11.4
CFG-C006	22	133	44	89	12.5
CFG-C007*	4	--	--	--	--

Florida State Parks  
Timber Management Analyses

CFG-C008	38	133	60	73	8.6
CFG-C009	24	46	20	26	9.6
CFG-C010	79	63	38	25	9.2
CFG-C012*	20	--	--	--	--
CFG-C013	93	99	51	48	11
CFG-C014*	74	--	--	--	--
CFG-C015*	3	--	--	--	--
CFG-C018*	4	--	--	--	--
CFG-C025*	20	--	--	--	--
CFG-C026*	29	--	--	--	--
CFG-C027	41	123	41	81	15.6
CFG-C028*	<1	--	--	--	--
CFG-C029	21	116	52	64	12.3
CFG-C030*	2	--	--	--	--
CFG-C031*	2	--	--	--	--
CFG-C032	7	77	25	52	10
CFG-C033	3	104	90	14	12.9
CFG-C034	<1	108	60	48	8.5
CFG-C035	5	121	100	21	11.1
CFG-C036	3	80	65	15	9.8
CFG-C037	<1	179	160	19	15.1
CFG-C038*	<1	--	--	--	--
CFG-C039*	<1	--	--	--	--
CFG-C040*	<1	--	--	--	--
CFG-C041*	<1	--	--	--	--
CFG-C046	101	91	43	47	11.1
CFG-C058*	12	--	--	--	--
CFG-C059*	3	--	--	--	--
CFG-C060	89	119	50	68	9.4
CFG-C061	24	150	63	86	10.4
CFG-C072	15	124	66	57	10.5
CFG-C074	9	122	100	22	6.6
CFG-C077	15	155	120	35	8.3
CFG-C079*	24	--	--	--	--
CFG-C085*	14	--	--	--	--
CFG-C086*	<1	--	--	--	--
CFG-C087*	16	--	--	--	--

Florida State Parks  
Timber Management Analyses

CFG-C088*	9	--	--	--	--
CFG-C090*	<1	--	--	--	--
CFG-C091*	6	--	--	--	--
CFG-C092*	5	--	--	--	--
CFG-C095*	<1	--	--	--	--
CFG-C096*	4	--	--	--	--
CFG-C097*	<1	--	--	--	--
CFG-C104*	1	--	--	--	--
CFG-C106	51	99	37	62	8.8
CFG-C107	3	73	10	63	5.4
CFG-C112*	<1	--	--	--	--
CFG-C113*	3	--	--	--	--
CFG-C114	8	80	--	80	16.3
CFG-C115*	<1	--	--	--	--
CFG-C116	24	96	56	39	8.7
CFG-C117*	<1	--	--	--	--
CFG-C118	34	113	103	9	14.5
CFG-C119*	<1	--	--	--	--
CFG-C130*	5	--	--	--	--
CFG-C132	13	128	32	95	19.1
CFG-C135*	5	--	--	--	--
CFG-C138*	1	--	--	--	--
CFG-C140	5	140	5	135	15.2
CFG-C163	21	40	15	25	30.9
CFG-C164*	<1	--	--	--	--
CFG-C168*	<1	--	--	--	--
CFG-C205	117	97	24	73	14
CFG-C209*	<1	--	--	--	--
CFG-C210	7	77	60	17	17.5
CFG-C214	59	111	10	101	12.5
CFG-C218*	<1	--	--	--	--
CFG-C220	7	155	150	5	35
CFG-C222*	<1	--	--	--	--
CFG-C223	255	93	27	65	15
CFG-C224	15	109	30	79	7.2
CFG-C255	35	131	43	87	14.6
CFG-C261	7	112	70	42	10.1

Florida State Parks  
Timber Management Analyses

CFG-C263	10	101	20	81	9.1
CFG-C275*	4	--	--	--	--
CFG-C276B	30	98	--	98	17.3
CFG-C277	47	124	--	124	18.6
CFG-C278	70	135	--	135	16.6
CFG-C279	2	82	70	12	10.2
CFG-C280	17	59	55	4	10.2
CFG-C281	10	73	60	13	8.8
CFG-E085*	3	--	--	--	--
CFG-E088*	3	--	--	--	--
CFG-E099*	8	--	--	--	--
CFG-E103	14	151	--	151	5.4
CFG-E141	62	59	24	35	8
CFG-E154*	<1	--	--	--	--
CFG-E159*	1	--	--	--	--
CFG-E160*	1	--	--	--	--
CFG-E167	<1	126	100	26	12.6
CFG-E170	27	120	81	39	10.9
CFG-E171	9	64	45	19	10.6
CFG-E172	16	164	105	59	12.6
CFG-E175*	3	--	--	--	--
CFG-E178*	4	--	--	--	--
CFG-E179	13	56	40	16	10.3
CFG-E182*	4	--	--	--	--
CFG-E183	19	83	62	21	11.1
CFG-E184*	<1	--	--	--	--
CFG-E185*	<1	--	--	--	--
CFG-E186*	2	--	--	--	--
CFG-E187*	<1	--	--	--	--
CFG-E220*	<1	--	--	--	--
CFG-E221*	<1	--	--	--	--
CFG-E241*	<1	--	--	--	--
CFG-E242*	<1	--	--	--	--
CFG-E259	13	107	42	64	9.2
CFG-E260*	<1	--	--	--	--
CFG-E262	10	78	25	53	8.5
CFG-E275	10	129	53	75	10.2



Florida State Parks  
Timber Management Analyses

CFG-E276*	2	--	--	--	--
CFG-E277	9	101	20	81	6.6
CFG-E278	12	123	60	63	9
CFG-E285*	5	--	--	--	--
CFG-E291*	2	--	--	--	--
CFG-E292*	<1	--	--	--	--
CFG-E293*	8	--	--	--	--
CFG-W001	19	83	26	57	11.4
CFG-W002	8	82	25	57	17.6
CFG-W003	42	133	46	86	13
CFG-W004	87	104	36	68	14.1
CFG-W005	8	124	20	104	10.6
CFG-W006	17	100	30	70	10.2
CFG-W007	157	126	86	40	13.6
CFG-W008	227	97	69	27	14.6
CFG-W009	7	110	90	20	21.4
CFG-W010*	1	--	--	--	--
CFG-W011	72	104	80	24	13
CFG-W012*	<1	--	--	--	--
CFG-W013*	<1	--	--	--	--
CFG-W014	13	113	100	13	22.3
CFG-W015*	1	--	--	--	--
CFG-W016	58	59	28	30	16.7
CFG-W017	76	119	87	31	15.5
CFG-W018	124	78	49	29	17.9
CFG-W019*	<1	--	--	--	--
CFG-W020*	4	--	--	--	--
CFG-W021	82	78	28	50	13.1
CFG-W022*	2	--	--	--	--
CFG-W023	2	20	--	20	1.7
CFG-W024	106	112	73	39	14.9
CFG-W025	28	110	65	45	14.3
CFG-W026	62	147	82	64	17.1
CFG-W027	20	105	105	--	26.5
CFG-W028*	<1	--	--	--	--
CFG-W030*	3	--	--	--	--
CFG-W031*	<1	--	--	--	--

Florida State Parks  
Timber Management Analyses

CFG-W032	48	99	52	46	18
CFG-W033*	<1	--	--	--	--
CFG-W034	17	66	20	46	13.3
CFG-W038*	<1	--	--	--	--
CFG-W040	1	118	40	78	11.2
CFG-W041	27	62	21	41	10
CFG-W042*	<1	--	--	--	--
CFG-W043	50	142	74	68	13.1
CFG-W044	38	115	51	64	12.9
CFG-W045*	2	--	--	--	--
CFG-W046*	<1	--	--	--	--
CFG-W047*	6	--	--	--	--
CFG-W055	61	123	56	67	18.6
CFG-W056	17	118	30	88	15.6
CFG-W060	15	83	61	21	11
CFG-W061*	1	--	--	--	--
CFG-W081*	<1	--	--	--	--
CFG-W082*	<1	--	--	--	--
CFG-W083	57	105	90	15	12.7
CFG-W084*	12	--	--	--	--
CFG-W085	7	90	70	20	11.7
CFG-W086*	3	--	--	--	--
CFG-W087	86	101	67	34	13.3
CFG-W088*	<1	--	--	--	--
CFG-W089	6	200	115	85	15.3
CFG-W090	1	120	20	100	6.5
CFG-W093	24	140	73	67	14.3
CFG-W095	53	97	42	54	14.5
CFG-W096	4	154	130	24	16
CFG-W106*	<1	--	--	--	--
CFG-W107	49	47	18	28	9.2
CFG-W116	12	16	10	6	10.1
CFG-W121*	33	--	--	--	--
CFG-W122*	<1	--	--	--	--
CFG-W124	11	64	18	46	4.9
CFG-W126	275	69	28	41	6.7
CFG-W133	6	182	30	152	19.4

Florida State Parks  
Timber Management Analyses

CFG-W135*	28	--	--	--	--
CFG-W138*	4	--	--	--	--
CFG-W139*	<1	--	--	--	--

\*Unsampled upland areas are present in this analysis and could require vegetation management in the future.

Pine Plantation (9,071 acres)

Pine plantation is an altered community type which resulted from past commercial management. The dominant overstory species found in this type is longleaf pine (*P. palustris*), loblolly pine (*P. taeda*), or slash pine (*P. elliottii*). As an altered forest/community type, and potentially a candidate for restoration, there are currently no FNAI recommendations on preferred species or stocking levels for this natural community. For the purpose of this timber assessment, any southern yellow pine species is considered preferred and all other overstory species are considered non-preferred. Future timber management activities could potentially transition this altered type into another natural community type per on-site and/or nearby land-use conditions.

Management Zones	Pine Plantation (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C021*	4	--	--	--	--
CFG-C027	13	134	53	80	11.1
CFG-C032*	<1	--	--	--	--
CFG-C035*	<1	--	--	--	--
CFG-C036*	<1	--	--	--	--
CFG-C037*	<1	--	--	--	--
CFG-C058	6	150	147	3	7.1
CFG-C060	4	171	50	121	9.3
CFG-C061	6	120	60	60	8.1
CFG-C079*	<1	--	--	--	--
CFG-C098*	11	--	--	--	--
CFG-C099*	4	--	--	--	--
CFG-C100*	4	--	--	--	--
CFG-C122*	3	--	--	--	--
CFG-C123	70	38	35	3	8.7
CFG-C124	51	81	81	--	6
CFG-C125	66	44	39	4	6.2

Florida State Parks  
Timber Management Analyses

CFG-C126*	<1	--	--	--	--
CFG-C127*	<1	--	--	--	--
CFG-C128	56	56	51	5	5.9
CFG-C129*	2	--	--	--	--
CFG-C141*	<1	--	--	--	--
CFG-C142	271	35	31	3	6.7
CFG-C143	18	15	7	7	20
CFG-C144	34	38	22	15	9.3
CFG-C145	10	25	24	0	2.7
CFG-C146*	6	--	--	--	--
CFG-C147*	2	--	--	--	--
CFG-C148	24	40	40	0	7.4
CFG-C149*	<1	--	--	--	--
CFG-C157*	1	--	--	--	--
CFG-C158*	16	--	--	--	--
CFG-C160*	<1	--	--	--	--
CFG-C168	18	50	20	30	8.3
CFG-C169	13	143	110	33	8.6
CFG-C170	124	58	35	23	7.4
CFG-C171	17	119	90	29	12.1
CFG-C176*	<1	--	--	--	--
CFG-C177	22	107	46	61	12.2
CFG-C178*	<1	--	--	--	--
CFG-C179	16	123	103	19	10
CFG-C181*	<1	--	--	--	--
CFG-C183	14	69	35	34	7.9
CFG-C184	34	60	8	52	11.4
CFG-C185	25	37	27	10	3.2
CFG-C186	11	11	8	2	2.1
CFG-C187*	3	--	--	--	--
CFG-C188*	7	--	--	--	--
CFG-C189	32	41	30	10	4.8
CFG-C190	42	67	43	24	9.9
CFG-C191	6	15	7	7	2.3
CFG-C193	31	56	51	5	4.4
CFG-C194	42	57	57	0	6.9
CFG-C195	48	58	42	15	3.3



Florida State Parks  
Timber Management Analyses

CFG-C196	18	8	8	0	2
CFG-C197	29	54	50	4	6.7
CFG-C198	15	73	66	7	3.3
CFG-C199	16	80	80	0	7
CFG-C200	15	35	0	34	16.4
CFG-C201	15	116	113	2	6
CFG-C202	15	70	67	3	4.2
CFG-C203	9	54	38	16	8.2
CFG-C204	33	89	62	26	12.8
CFG-C205	46	79	44	34	10.9
CFG-C206	130	28	24	3	3.9
CFG-C207*	5	--	--	--	--
CFG-C208*	<1	--	--	--	--
CFG-C213*	19	--	--	--	--
CFG-C218	60	64	48	16	8
CFG-C220	17	60	60	0	5.7
CFG-C221	75	65	60	5	6.8
CFG-C222	118	72	51	21	9
CFG-C223	18	71	5	66	15.2
CFG-C224*	7	--	--	--	--
CFG-C266*	<1	--	--	--	--
CFG-C267*	<1	--	--	--	--
CFG-C268	46	57	45	11	4.5
CFG-C269	12	18	16	2	3.3
CFG-C270*	3	--	--	--	--
CFG-C271*	<1	--	--	--	--
CFG-E005*	5	--	--	--	--
CFG-E006	12	87	70	16	4.6
CFG-E008	40	110	78	31	5.3
CFG-E009*	1	--	--	--	--
CFG-E010*	40	--	--	--	--
CFG-E011*	<1	--	--	--	--
CFG-E012	116	123	95	28	7.7
CFG-E013*	44	--	--	--	--
CFG-E014	136	92	64	27	7.4
CFG-E015*	1	--	--	--	--
CFG-E016*	3	--	--	--	--

Florida State Parks  
Timber Management Analyses

CFG-E017*	1	--	--	--	--
CFG-E018	60	154	154	0	6.1
CFG-E019*	<1	--	--	--	--
CFG-E021	7	132	132	--	6.6
CFG-E022*	<1	--	--	--	--
CFG-E023*	<1	--	--	--	--
CFG-E024*	<1	--	--	--	--
CFG-E025	5	34	34	0	4.4
CFG-E026	9	146	145	1	6.8
CFG-E027	6	140	134	5	6.5
CFG-E028*	12	--	--	--	--
CFG-E029	44	64	54	9	6.3
CFG-E030	52	67	58	9	6
CFG-E031*	2	--	--	--	--
CFG-E032	50	85	84	0	7.5
CFG-E033	5	63	63	0	7.6
CFG-E034	143	116	115	0	6.3
CFG-E035	51	53	52	0	6.6
CFG-E036*	41	--	--	--	--
CFG-E037	85	109	96	13	4.6
CFG-E038	28	19	7	11	3.2
CFG-E039	125	51	50	0	7.4
CFG-E040	44	118	117	0	5.8
CFG-E041	23	81	77	4	7
CFG-E042	7	160	160	0	6.5
CFG-E043	26	51	47	4	7.6
CFG-E044	154	40	24	16	4.5
CFG-E045	89	52	51	0	5.4
CFG-E046	62	25	25	0	2.5
CFG-E047	15	75	75	0	7
CFG-E048	10	9	9	0	2.6
CFG-E049	104	46	46	0	5.3
CFG-E050	53	21	10	10	2.4
CFG-E051*	<1	--	--	--	--
CFG-E052	11	57	57	0	5.5
CFG-E053*	<1	--	--	--	--
CFG-E054	39	68	58	9	6.1

Florida State Parks  
Timber Management Analyses

CFG-E055	21	130	86	44	6.2
CFG-E056	72	77	74	2	7.3
CFG-E057	14	55	55	0	6.2
CFG-E058	58	69	64	5	6.1
CFG-E059	42	53	50	2	5.7
CFG-E061	47	106	86	20	5.9
CFG-E062	102	124	120	4	6.5
CFG-E063	29	80	77	3	8.2
CFG-E064	88	96	86	10	6.4
CFG-E065	403	40	21	19	5.4
CFG-E066*	<1	--	--	--	--
CFG-E067	8	111	87	24	7.9
CFG-E068	49	73	66	6	6.1
CFG-E069	11	107	100	7	6.4
CFG-E070	22	58	52	5	3.8
CFG-E072*	<1	--	--	--	--
CFG-E073	113	59	56	2	6.8
CFG-E074	161	24	24	0	3.2
CFG-E075*	<1	--	--	--	--
CFG-E076	17	78	77	0	6.7
CFG-E077	56	60	60	0	5.9
CFG-E078*	3	--	--	--	--
CFG-E079	133	71	38	32	5.3
CFG-E080	4	109	55	53	4.8
CFG-E081*	35	--	--	--	--
CFG-E082	25	103	81	21	6.1
CFG-E083	22	99	94	4	4.6
CFG-E084	10	157	146	10	6
CFG-E085	154	86	75	11	7.2
CFG-E086*	1	--	--	--	--
CFG-E087	5	114	93	20	5.1
CFG-E088	211	23	14	8	3.2
CFG-E089	16	50	46	4	6.9
CFG-E090	33	61	50	11	7.7
CFG-E091	13	7	0	7	2.8
CFG-E092	78	84	84	0	8.4
CFG-E093	43	89	88	0	7.8

Florida State Parks  
Timber Management Analyses

CFG-E094*	35	--	--	--	--
CFG-E095	18	66	62	4	7.5
CFG-E096	6	71	53	18	5.4
CFG-E097*	<1	--	--	--	--
CFG-E099*	25	--	--	--	--
CFG-E101*	<1	--	--	--	--
CFG-E102*	3	--	--	--	--
CFG-E103	90	59	38	21	5
CFG-E104	26	65	62	2	6.7
CFG-E105	55	92	79	12	6.4
CFG-E106*	<1	--	--	--	--
CFG-E107	12	165	165	0	6.4
CFG-E108	30	43	34	9	4.7
CFG-E109*	<1	--	--	--	--
CFG-E110*	<1	--	--	--	--
CFG-E111*	<1	--	--	--	--
CFG-E112	17	35	35	0	6.5
CFG-E113	22	52	46	6	4.9
CFG-E114	38	70	58	11	5.4
CFG-E115	10	67	60	7	4.2
CFG-E116*	4	--	--	--	--
CFG-E117	41	59	50	9	4.1
CFG-E118	15	55	50	5	5.7
CFG-E119	168	75	48	26	6.5
CFG-E120	137	71	66	4	6.6
CFG-E121	12	17	7	10	2.8
CFG-E122	37	117	92	24	6
CFG-E123	11	73	66	6	6
CFG-E124	28	45	38	7	4.9
CFG-E125	4	118	92	25	5.1
CFG-E126	196	59	52	7	4.9
CFG-E127*	3	--	--	--	--
CFG-E128	33	58	48	10	4.4
CFG-E129	52	51	42	8	5.2
CFG-E130	32	88	54	33	5.2
CFG-E131*	9	--	--	--	--
CFG-E132	43	100	90	9	6.8



Florida State Parks  
Timber Management Analyses

CFG-E133	49	33	1	32	3.3
CFG-E134	20	28	27	0	4.5
CFG-E136	50	80	67	13	5.6
CFG-E137	33	43	38	5	5.8
CFG-E138	100	57	52	5	5
CFG-E139	24	68	63	5	5
CFG-E140	27	49	20	29	6.8
CFG-E142	21	232	60	172	10.7
CFG-E146	9	150	150	0	4.3
CFG-E147	25	148	98	50	6.9
CFG-E148	15	180	170	10	6.3
CFG-E149	24	188	66	122	9.3
CFG-E150	4	50	50	0	4.2
CFG-E151*	3	--	--	--	--
CFG-E152*	10	--	--	--	--
CFG-E153	11	73	56	16	5.5
CFG-E154	106	260	20	240	10.5
CFG-E155	56	71	69	1	7.7
CFG-E156*	<1	--	--	--	--
CFG-E157*	15	--	--	--	--
CFG-E158*	<1	--	--	--	--
CFG-E161*	3	--	--	--	--
CFG-E162*	2	--	--	--	--
CFG-E163	35	68	42	25	8.5
CFG-E164*	<1	--	--	--	--
CFG-E165*	<1	--	--	--	--
CFG-E166	21	129	70	59	12.6
CFG-E174	6	170	160	10	9.4
CFG-E175	11	164	130	34	10.2
CFG-E195*	<1	--	--	--	--
CFG-E196*	8	--	--	--	--
CFG-E197	48	65	55	10	5.7
CFG-E198	12	111	87	23	6
CFG-E199	191	34	30	4	4.6
CFG-E200	14	20	20	0	3.7
CFG-E201	43	97	78	18	4.8
CFG-E202	93	70	60	9	4.4

Florida State Parks  
Timber Management Analyses

CFG-E203	19	23	5	18	3.8
CFG-E204	87	73	62	11	5.1
CFG-E206*	3	--	--	--	--
CFG-E211	5	85	25	60	8.8
CFG-E212	13	84	70	14	12.2
CFG-E213*	<1	--	--	--	--
CFG-E214*	13	--	--	--	--
CFG-E215*	<1	--	--	--	--
CFG-E234	12	120	120	0	8.1
CFG-E235	16	126	126	0	8.7
CFG-E279	28	132	89	43	6.2
CFG-E284*	<1	--	--	--	--
CFG-E300*	2	--	--	--	--
CFG-E301	45	60	38	22	6.3
CFG-E302*	2	--	--	--	--
CFG-E303*	<1	--	--	--	--
CFG-E304*	1	--	--	--	--
CFG-E305*	4	--	--	--	--
CFG-E306	11	155	126	29	7.9
CFG-E307	12	172	110	62	9.5
CFG-E308	2	209	200	9	7
CFG-W008*	<1	--	--	--	--
CFG-W011*	<1	--	--	--	--
CFG-W012	55	52	49	2	6.2
CFG-W013	90	52	51	0	5.5
CFG-W014*	1	--	--	--	--
CFG-W015	61	54	37	17	5.4
CFG-W016*	1	--	--	--	--
CFG-W017*	1	--	--	--	--
CFG-W018*	<1	--	--	--	--
CFG-W019	12	6	--	6	3.5
CFG-W020	62	38	35	3	5.6
CFG-W021	2	72	--	72	12.7
CFG-W023	37	59	55	3	5.2
CFG-W024*	1	--	--	--	--
CFG-W025*	<1	--	--	--	--
CFG-W026*	<1	--	--	--	--

Florida State Parks  
Timber Management Analyses

CFG-W031*	1	--	--	--	--
CFG-W032*	2	--	--	--	--
CFG-W033	153	76	76	0	6.5
CFG-W034*	1	--	--	--	--
CFG-W035	89	45	44	0	6.8
CFG-W036*	1	--	--	--	--
CFG-W037*	24	--	--	--	--
CFG-W039	91	31	29	2	6.9
CFG-W040*	<1	--	--	--	--
CFG-W044	13	41	1	40	12.3
CFG-W123*	<1	--	--	--	--
CFG-W124	98	52	12	39	6.9
CFG-W125	58	26	14	12	5.8
CFG-W126	70	147	66	80	6.4

\*Unsampled upland areas are present in this analysis and could require vegetation management in the future.

Sandhill (6,408 acres)

The dominant pine tree in sandhill communities in the region is longleaf pine (*P. palustris*). Herbaceous cover, dominated by wiregrass (*Aristida stricta*), should be 80% or greater and reach a height of less than three feet. Sandhill communities in the region will contain scattered individual trees, clumps, or ridges of onsite oak species such as turkey oak (*Q. laevis*), sand post oak (*Q. margaretta*), and bluejack oak (*Q. incana*). In old growth conditions, sand post oaks will commonly be 150-200 years old, and some turkey oaks will be over 100 years old. The optimal fire return interval for this community is two to three years. In this region, the preferred species, as determined by FNAI reference sites, is longleaf pine and should be stocked at a level of 20 to 60 square feet per acre BA while non-pine species should remain between 0 and 78.8 stems per acre. The following management zone(s) contain sandhill which could be considered for some form of timber management including midstory mitigation, site preparation, and planting of preferred pine species.

Management Zones	Sandhill (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C077	8	102	80	22	3.6
CFG-C079	23	39	5	34	7.7
CFG-C080	23	125	--	125	7.5
CFG-C081	33	102	24	77	8.3

Florida State Parks  
Timber Management Analyses

Management Zones	Sandhill (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C106	6	66	--	66	8.4
CFG-C111	17	66	15	51	5.8
CFG-C116	29	79	14	65	7.9
CFG-C120*	9	--	--	--	--
CFG-C121	30	62	4	58	6.7
CFG-C122	147	76	7	69	7.9
CFG-C123*	<1	--	--	--	--
CFG-C136*	<1	--	--	--	--
CFG-C137	24	88	30	58	6.9
CFG-C138*	<1	--	--	--	--
CFG-C139	71	88	22	66	7.8
CFG-C141	36	120	--	120	9.2
CFG-C142*	<1	--	--	--	--
CFG-C145*	<1	--	--	--	--
CFG-C146	3	114	--	114	13.9
CFG-C149	44	100	8	91	11.1
CFG-C150*	1	--	--	--	--
CFG-C151	3	100	20	80	14
CFG-C152	9	101	30	71	15.8
CFG-C153	12	90	--	90	15.4
CFG-C156	30	99	7	92	13.7
CFG-C157	35	124	10	114	14.2
CFG-C158	<1	92	20	72	15.2
CFG-C159*	<1	--	--	--	--
CFG-C160	11	86	--	86	12.7
CFG-C161*	<1	--	--	--	--
CFG-C162	15	120	17	102	14.6
CFG-C163*	<1	--	--	--	--
CFG-C164*	<1	--	--	--	--
CFG-C165*	<1	--	--	--	--
CFG-C166*	<1	--	--	--	--
CFG-C167*	<1	--	--	--	--
CFG-C168	432	112	8	103	15.7



Florida State Parks  
Timber Management Analyses

Management Zones	Sandhill (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C169*	<1	--	--	--	--
CFG-C170	13	147	--	147	17.2
CFG-C171*	<1	--	--	--	--
CFG-C172*	<1	--	--	--	--
CFG-C173*	<1	--	--	--	--
CFG-C174	8	116	10	106	9.2
CFG-C175	17	150	12	138	14.2
CFG-C176	28	120	--	120	15.6
CFG-C177	5	84	--	84	11.9
CFG-C178	76	87	15	72	10.5
CFG-C179	8	101	--	101	14.1
CFG-C180*	<1	--	--	--	--
CFG-C181	152	108	18	90	12.2
CFG-C182	9	100	20	80	8.9
CFG-C183	13	71	--	71	17.5
CFG-C184	92	93	1	92	14.6
CFG-C185*	<1	--	--	--	--
CFG-C186	<1	13	--	13	2.1
CFG-C187*	<1	--	--	--	--
CFG-C188*	<1	--	--	--	--
CFG-C189*	<1	--	--	--	--
CFG-C190	112	117	8	108	12.4
CFG-C191*	<1	--	--	--	--
CFG-C192	10	47	--	47	2.9
CFG-C193*	<1	--	--	--	--
CFG-C194*	<1	--	--	--	--
CFG-C195*	1	--	--	--	--
CFG-C197*	<1	--	--	--	--
CFG-C198*	<1	--	--	--	--
CFG-C199*	<1	--	--	--	--
CFG-C200	29	149	1	147	16.2
CFG-C201*	<1	--	--	--	--
CFG-C202*	<1	--	--	--	--

Florida State Parks  
Timber Management Analyses

Management Zones	Sandhill (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C203	12	103	--	103	21
CFG-C204*	<1	--	--	--	--
CFG-C205	117	96	21	74	13.7
CFG-C206*	1	--	--	--	--
CFG-C207	107	80	5	75	10
CFG-C208*	<1	--	--	--	--
CFG-C210	8	55	10	45	8.4
CFG-C211	54	40	1	39	15
CFG-C212	1	180	--	180	17.1
CFG-C214*	1	--	--	--	--
CFG-C215	26	9	9	0	3.1
CFG-C216	28	13	6	6	2.3
CFG-C223*	3	--	--	--	--
CFG-C224	91	93	19	74	9.7
CFG-C225*	<1	--	--	--	--
CFG-C237*	1	--	--	--	--
CFG-C238*	33	--	--	--	--
CFG-C239*	<1	--	--	--	--
CFG-C240*	<1	--	--	--	--
CFG-C242*	<1	--	--	--	--
CFG-C243	24	51	48	3	6.2
CFG-C244	50	75	65	10	7.7
CFG-C245*	2	--	--	--	--
CFG-C252*	<1	--	--	--	--
CFG-C253*	<1	--	--	--	--
CFG-C254*	<1	--	--	--	--
CFG-C255	77	72	39	33	6.8
CFG-C256	154	46	37	9	5.5
CFG-C257	137	35	29	6	6.5
CFG-C258	144	81	42	38	7.9
CFG-C259	139	79	52	26	7.5
CFG-C260	9	127	64	62	6.8
CFG-C261	289	38	16	21	5.7

Florida State Parks  
Timber Management Analyses

Management Zones	Sandhill (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C262	21	93	19	73	6.7
CFG-C263	284	58	18	40	5.8
CFG-C264	200	60	30	30	7.7
CFG-C265	212	71	40	30	7.1
CFG-C266	41	109	8	101	8.5
CFG-C267	185	33	21	11	5.8
CFG-C270*	<1	--	--	--	--
CFG-C271	19	88	20	68	7.1
CFG-C272	180	30	13	17	6.4
CFG-C273	160	25	17	8	6.5
CFG-C274	229	52	30	21	6.5
CFG-C275	70	23	13	10	6.9
CFG-C276	124	35	21	13	6.4
CFG-C276A	82	34	20	14	5.4
CFG-C276B	120	42	23	19	6.3
CFG-C278	10	57	--	57	6.6
CFG-E066*	4	--	--	--	--
CFG-E124*	<1	--	--	--	--
CFG-E126	8	67	--	67	5.6
CFG-E127	43	65	--	65	7.2
CFG-E128*	1	--	--	--	--
CFG-E161	21	84	4	80	6.3
CFG-E162*	5	--	--	--	--
CFG-E207	22	71	--	71	7.3
CFG-E208	3	104	34	70	9.1
CFG-E209	34	34	7	27	6.2
CFG-W001	30	61	30	31	6.7
CFG-W003	53	60	20	39	8.9
CFG-W004	46	86	13	72	7.7
CFG-W005	91	100	45	55	9.9
CFG-W006	172	98	28	69	11.4
CFG-W007	129	63	5	58	8.4
CFG-W008	39	39	5	33	9.8

Florida State Parks  
Timber Management Analyses

Management Zones	Sandhill (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-W009	57	55	11	44	7.8
CFG-W010*	<1	--	--	--	--
CFG-W011	13	78	27	50	9.6
CFG-W016	11	20	15	5	9.8
CFG-W017*	<1	--	--	--	--
CFG-W018	5	75	10	65	10.7
CFG-W021	66	72	14	57	11.2
CFG-W035*	<1	--	--	--	--
CFG-W036	44	44	5	39	12.1
CFG-W059*	3	--	--	--	--
CFG-W060	66	114	50	64	9.8
CFG-W061	18	63	6	57	7.2
CFG-W083*	1	--	--	--	--
CFG-W084	9	74	13	61	8
CFG-W087	12	120	5	115	12.3
CFG-W089*	<1	--	--	--	--
CFG-W109	3	81	10	71	9.7
CFG-W110	34	61	--	61	10

\*Unsampled upland areas are present in this analysis and could require vegetation management in the future.

### Scrub (1,630 acres)

Dominant plant species in scrub include rusty staggerbush (*Lyonia ferruginea*), sand live oak (*Q. geminata*), myrtle oak (*Q. myrtifolia*), **Chapman's oak** (*Q. chapmanii*), fetterbush (*Lyonia lucida*), shiny blueberry (*Vaccinium myrsinites*), and saw palmetto (*Serenoa repens*). In this region, preferred or likely pine species, as determined by FNAI reference sites, are longleaf (*P. palustris*) and slash (*P. elliotii*) and should be stocked at a level of 0 to 20 square feet per acre BA while non-pine species should remain between 0 and 13.1 stems per acre. The following management zone(s) contain scrub which could be considered for some form of timber management including overstory removal, midstory mitigation, site preparation, and planting of preferred pine species.



Florida State Parks  
Timber Management Analyses

Management Zones	Scrub (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C008*	2	--	--	--	--
CFG-C009	36	102	10	92	5.6
CFG-C010	75	53	9	43	4.6
CFG-C013*	4	--	--	--	--
CFG-C226*	4	--	--	--	--
CFG-C227*	9	--	--	--	--
CFG-C228*	30	--	--	--	--
CFG-C229*	22	--	--	--	--
CFG-C230*	26	--	--	--	--
CFG-C231*	15	--	--	--	--
CFG-C232*	16	--	--	--	--
CFG-C233*	28	--	--	--	--
CFG-C234*	17	--	--	--	--
CFG-C235*	28	--	--	--	--
CFG-C236*	26	--	--	--	--
CFG-C237*	23	--	--	--	--
CFG-C238*	2	--	--	--	--
CFG-C239*	44	--	--	--	--
CFG-C240*	23	--	--	--	--
CFG-C241*	13	--	--	--	--
CFG-C242*	68	--	--	--	--
CFG-C244*	4	--	--	--	--
CFG-C245*	17	--	--	--	--
CFG-C245A*	27	--	--	--	--
CFG-C246*	13	--	--	--	--
CFG-C247*	12	--	--	--	--
CFG-C248*	13	--	--	--	--
CFG-C249*	14	--	--	--	--
CFG-C250*	17	--	--	--	--
CFG-C251*	4	--	--	--	--
CFG-C252*	31	--	--	--	--
CFG-C253*	5	--	--	--	--
CFG-C254	114	71	10	60	6.7

Florida State Parks  
Timber Management Analyses

Management Zones	Scrub (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C255	393	25	5	20	6.1
CFG-C256*	4	--	--	--	--
CFG-C257*	11	--	--	--	--
CFG-C273	10	6	3	2	3.5
CFG-C274*	1	--	--	--	--
CFG-C275	21	36	24	12	7.4
CFG-E100	59	99	54	44	5.1
CFG-E101	11	141	20	121	6.3
CFG-E135	6	36	10	26	3.1
CFG-E141	25	63	--	63	5.7
CFG-E149	4	7	--	7	0.3
CFG-E150	3	0	--	0	0
CFG-E151	2	10	--	10	2.8
CFG-E152	4	85	--	85	1.5
CFG-E154*	<1	--	--	--	--
CFG-E161	9	24	--	24	10.6
CFG-E162	27	63	6	56	8
CFG-E166*	<1	--	--	--	--
CFG-E182*	<1	--	--	--	--
CFG-E183*	1	--	--	--	--
CFG-E184	16	50	17	32	3.6
CFG-E185	42	60	24	36	5.4
CFG-E186*	<1	--	--	--	--
CFG-E187	28	74	53	20	4.6
CFG-E188*	<1	--	--	--	--
CFG-E189	49	65	38	27	6
CFG-E190	45	68	25	42	6.4
CFG-E191*	<1	--	--	--	--
CFG-E264*	2	--	--	--	--

\*Unsampled upland areas are present in this analysis and could require vegetation management in the future.

### Scrubby Flatwoods (683 acres)

The dominant tree in the scrubby flatwoods of north Florida will usually be longleaf

Florida State Parks  
Timber Management Analyses

pine (*P. palustris*). Mature sand pines (*P. clausa*) will typically be absent. A diverse shrub understory will be characteristic, with up to 25% bare sand coverage. Dominant shrubs include sand live oak (*Q. geminata*), myrtle oak (*Q. myrtifolia*), **Chapman's** oak (*Q. chapmanii*), saw palmetto (*Serenoa repens*), rusty staggerbush (*Lyonia ferruginea*), and tarflower (*Bejaria racemosa*). The optimal fire return interval for this community is regionally variable, but coastal scrub has shown an ability to reach fuel height and fire carrying potential faster than interior examples. Areas may be burned as frequently as every three to eight years when burn prescriptions are designed to achieve a mosaic of burned and unburned areas. In this region, the preferred species, as determined by FNAI reference sites, is longleaf pine and should be stocked at a level of 10 to 60 square feet per acre BA while non-pine species should remain between 0 and 26.2 stems per acre. The following management zone(s) contain scrubby flatwoods which could be considered for some form of timber management including overstory removal, midstory mitigation, site preparation, and planting of preferred pine species.

Management Zones	Scrubby Flatwoods (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C012	34	55	34	21	5.3
CFG-C013	2	60	50	10	9.9
CFG-C058	5	65	62	3	6.1
CFG-C061	14	139	70	69	7
CFG-C064*	3	--	--	--	--
CFG-C065*	2	--	--	--	--
CFG-C068*	2	--	--	--	--
CFG-C069	8	101	90	11	9.7
CFG-C070*	<1	--	--	--	--
CFG-C072*	2	--	--	--	--
CFG-C074*	4	--	--	--	--
CFG-C077*	1	--	--	--	--
CFG-C078*	2	--	--	--	--
CFG-C079	20	69	53	16	6.4
CFG-C080	3	13	--	13	0.3
CFG-C081	16	162	140	22	4.5
CFG-C116	26	63	52	11	4.4
CFG-C118	57	123	45	78	10.7
CFG-C119*	<1	--	--	--	--
CFG-E097	40	96	14	82	5.9

Florida State Parks  
Timber Management Analyses

Management Zones	Scrubby Flatwoods (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-E098	28	54	30	24	5.3
CFG-E099*	2	--	--	--	--
CFG-E100*	2	--	--	--	--
CFG-E103	19	20	--	20	1.6
CFG-E135	6	45	42	3	3.5
CFG-E141	17	19	10	9	3.1
CFG-E157*	1	--	--	--	--
CFG-E158*	1	--	--	--	--
CFG-E159*	2	--	--	--	--
CFG-E161*	<1	--	--	--	--
CFG-E162	38	49	44	4	8.3
CFG-E163*	4	--	--	--	--
CFG-E191*	<1	--	--	--	--
CFG-E192	8	60	15	45	7
CFG-E209	24	19	5	14	3.8
CFG-E279	5	20	3	16	9.3
CFG-E285*	2	--	--	--	--
CFG-E286*	4	--	--	--	--
CFG-E287*	<1	--	--	--	--
CFG-E288*	3	--	--	--	--
CFG-W109*	<1	--	--	--	--
CFG-W111	12	29	13	16	9.1
CFG-W127	42	52	25	27	6.2
CFG-W128	46	19	13	6	7.8
CFG-W129	12	54	40	14	12.6
CFG-W130	50	42	17	25	4
CFG-W131*	24	--	--	--	--
CFG-W132	48	41	14	27	5.9
CFG-W133	31	42	12	29	5.4
CFG-W134*	10	--	--	--	--
CFG-W135	24	42	40	2	7.8

\*Unsampled upland areas are present in this analysis and could require vegetation management in the future.



Florida State Parks  
Timber Management Analyses

Upland Hardwood Forest (350 acres)

Upland hardwood forest is a well-developed, closed-canopy forest dominated by deciduous hardwood trees on mesic soils in areas sheltered from fire. It typically has a diverse assemblage of deciduous and evergreen tree species in the canopy and midstory, shade-tolerant shrubs, and a sparse groundcover. Characteristic canopy trees include southern magnolia (*Magnolia grandiflora*), pignut hickory (*Carya glabra*), sweetgum (*Liquidambar styraciflua*), Florida maple (*Acer saccharum* ssp. *floridanum*), live oak (*Quercus virginiana*), laurel oak (*Q. hemisphaerica*), swamp chestnut oak (*Q. michauxii*), southern hackberry (*Celtis occidentalis*), white ash (*Fraxinus americana*), and loblolly pine (*P. taeda*). Species commonly found in Florida Panhandle and northern peninsula but not farther south include American beech (*Fagus grandifolia*), white oak (*Q. alba*), and spruce pine (*P. glabra*). There are currently no FNAI recommendations on preferred species or stocking levels for this natural community.

Management Zones	Upland Hardwood Forest (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C008	9	209	55	154	9.7

\*Unsampled upland areas are present in this analysis and could require vegetation management in the future.

Upland Pine Forest (6 acres)

Upland pine is a woodland of widely spaced pines with a sparse to moderate shrub layer and a dense, species-rich groundcover of grasses and herbs, occurring on gently rolling terrain. The canopy is dominated by longleaf pine (*P. palustris*); shortleaf pine (*P. echinata*) also may be present. There is an intermittent subcanopy layer of smaller pines, and hardwoods including southern red oak (*Q. falcata*), blackjack oak (*Q. marilandica*), flowering dogwood (*Cornus florida*), bluejack oak (*Q. incana*), post oak (*Q. stellata*), sassafras (*Sassafras albidum*), laurel oak (*Q. hemisphaerica*), winged sumac (*Rhus copallinum*), common persimmon (*Diospyros virginiana*), sand post oak (*Q. margaretta*), mockernut hickory (*Carya alba*), and sourgum (*Nyssa sylvatica*). There are currently no FNAI recommendations on preferred species or stocking levels for this natural community.

Management Zones	Upland Pine Forest (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C142*	6	--	--	--	--

\*Unsampled upland areas are present in this analysis and could require vegetation management in the future.

Florida State Parks  
Timber Management Analyses

Wet Flatwoods (1,773 acres)

Within wet flatwoods the dominant pines species will usually be longleaf pine (*P. palustris*), slash pine (*P. elliottii*), pond pine (*P. serotina*), and/or loblolly pine (*P. taeda*). The species composition within a location will be determined by drainage and periods of higher moisture content. Pond cypress (*Taxodium ascendens*) may reach canopy in some locations. The canopy will be open, with pines being widely scattered and of variable age classes. In this region, the preferred species, as determined by FNAI reference sites, is slash pine and should be stocked at a level of 10 to 50 square feet per acre BA while non-pine species should remain at 0 stems per acre. The following management zone(s) contain wet flatwoods which could be considered for some form of timber management including overstory removal, midstory mitigation, site preparation, and planting of preferred pine species.

Management Zones	Wet Flatwoods (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C011*	<1	--	--	--	--
CFG-C012	134	99	37	62	7.2
CFG-C013*	2	--	--	--	--
CFG-C017*	18	--	--	--	--
CFG-C054*	4	--	--	--	--
CFG-C058	123	153	115	38	8.8
CFG-C061	22	132	100	32	7.7
CFG-C062	8	135	80	55	6.6
CFG-C063*	8	--	--	--	--
CFG-C064	24	232	140	92	9.8
CFG-C065	1	223	80	143	11.8
CFG-C075	10	128	95	32	5.7
CFG-C076*	1	--	--	--	--
CFG-C077	10	112	50	62	5.3
CFG-C104*	<1	--	--	--	--
CFG-C106*	5	--	--	--	--
CFG-C114*	10	--	--	--	--
CFG-C115*	<1	--	--	--	--
CFG-C116*	1	--	--	--	--
CFG-C120*	2	--	--	--	--
CFG-E003*	<1	--	--	--	--
CFG-E036*	1	--	--	--	--

Florida State Parks  
Timber Management Analyses

Management Zones	Wet Flatwoods (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-E044*	8	--	--	--	--
CFG-E049*	<1	--	--	--	--
CFG-E050*	<1	--	--	--	--
CFG-E074	22	74	74	0	5
CFG-E075	32	64	59	4	6.4
CFG-E077*	1	--	--	--	--
CFG-E078*	24	--	--	--	--
CFG-E088*	25	--	--	--	--
CFG-E096*	<1	--	--	--	--
CFG-E097*	2	--	--	--	--
CFG-E099*	16	--	--	--	--
CFG-E100	37	84	30	54	4.4
CFG-E101	21	79	20	59	4.3
CFG-E102*	<1	--	--	--	--
CFG-E103	118	87	22	64	7.2
CFG-E105	<1	76	10	66	5.9
CFG-E106*	<1	--	--	--	--
CFG-E112*	<1	--	--	--	--
CFG-E113*	1	--	--	--	--
CFG-E124*	1	--	--	--	--
CFG-E126*	1	--	--	--	--
CFG-E128*	<1	--	--	--	--
CFG-E129	6	10	--	10	4
CFG-E131*	16	--	--	--	--
CFG-E141	55	65	13	51	6
CFG-E161	12	24	--	24	7.3
CFG-E180*	2	--	--	--	--
CFG-E182	45	112	68	44	10.4
CFG-E183	37	110	52	57	10.7
CFG-E184*	<1	--	--	--	--
CFG-E191	2	132	70	62	9.3
CFG-E192	42	145	54	91	12.8
CFG-E193	19	33	--	33	4.2

Florida State Parks  
Timber Management Analyses

Management Zones	Wet Flatwoods (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-E194*	<1	--	--	--	--
CFG-E195	90	97	74	23	10
CFG-E197*	<1	--	--	--	--
CFG-E199*	<1	--	--	--	--
CFG-E221*	<1	--	--	--	--
CFG-E223*	<1	--	--	--	--
CFG-E224	45	101	70	31	11.2
CFG-E227	4	130	120	10	15.7
CFG-E229	4	100	80	20	13
CFG-E241*	<1	--	--	--	--
CFG-E279	60	8	--	8	8
CFG-E280*	<1	--	--	--	--
CFG-E285*	<1	--	--	--	--
CFG-E287*	<1	--	--	--	--
CFG-E288*	2	--	--	--	--
CFG-E290*	7	--	--	--	--
CFG-E291*	<1	--	--	--	--
CFG-E298*	<1	--	--	--	--
CFG-E307	37	113	41	72	5.3
CFG-E308*	<1	--	--	--	--
CFG-W030*	3	--	--	--	--
CFG-W077	55	78	20	58	11.7
CFG-W078	65	96	51	44	10.6
CFG-W079	7	108	80	27	13.6
CFG-W080	28	117	26	91	11
CFG-W094*	3	--	--	--	--
CFG-W095	17	45	22	22	15.7
CFG-W096	49	70	21	48	14.7
CFG-W107	2	0	--	0	5.5
CFG-W123*	<1	--	--	--	--
CFG-W124	152	17	1	16	4.2
CFG-W135	8	62	--	62	12.4

\*Unsampled upland areas are present in this analysis and could require vegetation management in the future.



Florida State Parks  
Timber Management Analyses

Xeric Hammock (307 acres)

Typically considered a late successional stage of scrub or sandhill that generally occurs in small isolated patches on excessively well drained soils. Vegetation will consist of a low closed canopy dominated by live oak (*Q. virginiana*) which provides **shady conditions**. Typical plant species may also include Chapman's oak (*Q. chapmanii*), and laurel oak (*Q. laurifolia*). Sand pine (*P. clausa*), slash pine (*P. elliottii*), or longleaf pine (*P. palustris*) may also be a minor component. Areas that have been determined to be severely degraded sandhill instead of true xeric hammock should be considered for restoration efforts to return the community to historic conditions. There is currently no FNAI recommendations or preferred species or stocking levels for this natural community but in areas where restoration is considered, longleaf pine will be considered the preferred species. The following management zones contain xeric hammock which could be considered for some form of timber management including overstory removals, and midstory mitigation.

Management Zones	Xeric Hammock (Acres)	Basal Area (ft <sup>2</sup> /acre)	Basal Area Preferred Species	Basal Area Non-Preferred Species	Average Diameter at breast height (inches)
CFG-C015*	34	--	--	--	--
CFG-C016*	16	--	--	--	--
CFG-C106	14	75	65	10	7
CFG-C107	10	68	23	45	8.7
CFG-C108*	10	--	--	--	--
CFG-C109*	<1	--	--	--	--
CFG-C111	9	95	80	15	7.9
CFG-C112*	<1	--	--	--	--
CFG-C114	15	100	12	87	8.4
CFG-C115*	1	--	--	--	--
CFG-C116	7	66	50	16	4.8
CFG-C120	17	73	40	33	7.1
CFG-C154*	3	--	--	--	--
CFG-C275	11	84	62	22	10.1
CFG-C277*	<1	--	--	--	--
CFG-E182	51	58	26	32	7.1
CFG-E184*	<1	--	--	--	--
CFG-E185*	<1	--	--	--	--
CFG-E186	23	90	32	58	7.2
CFG-E187*	<1	--	--	--	--
CFG-E242*	2	--	--	--	--
CFG-E279*	1	--	--	--	--

Florida State Parks  
Timber Management Analyses

CFG-E281*	<1	--	--	--	--
CFG-E285*	13	--	--	--	--
CFG-E286*	11	--	--	--	--
CFG-E290*	<1	--	--	--	--
CFG-E291*	9	--	--	--	--
CFG-E292*	<1	--	--	--	--
CFG-E303*	1	--	--	--	--
CFG-E306	34	103	66	36	8.8
CFG-E307	1	144	140	4	4.8
CFG-E308	1	191	140	51	5.2
CFG-W069	4	70	70	--	13.8
CFG-W070	1	167	--	167	10.6
CFG-W071	9	110	--	110	10.5
CFG-W072*	<1	--	--	--	--
CFG-W073*	<1	--	--	--	--

\*Unsampled upland areas are present in this analysis and could require vegetation management in the future.

Florida State Parks  
Timber Management Analyses

Table 2. Summary of potential timber management actions for upland natural community (NatCom) types to help restore or improve ecosystem conditions.

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C002**	12	Mesic Hammock	3	--	--	--	--	Y	Y	Y	Y
CFG-C003**	5	Mesic Hammock	1	--	--	--	--	Y	Y	Y	Y
CFG-C004**	17	Mesic Hammock	1	--	--	--	--	Y	Y	Y	Y
CFG-C005	26	Mesic Hammock	19	15	10 - 30	733	0 - 79	Y	Y	N	N
CFG-C006	22	Mesic Hammock	22	36	10 - 30	774	0 - 79	Y	Y	N	N
CFG-C007**	225	Mesic Hammock	4	--	--	--	--	Y	Y	Y	Y
CFG-C008	65	Upland Hardwood Forest	9	20	30 - 80	837	0 - 263	Y	Y	Y	Y
CFG-C008	65	Mesic Hammock	38	20	10 - 30	837	0 - 79	Y	Y	N	N
CFG-C008**	65	Scrub	2	--	--	--	--	Y	Y	Y	Y
CFG-C009	71	Scrub	36	40	0 - 20	1124	0 - 13	Y	Y	N	N
CFG-C009	71	Mesic Hammock	24	40	10 - 30	1124	0 - 79	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C010	181	Mesic Hammock	79	5	10 - 30	1572	0 - 79	Y	Y	Y	Y
CFG-C010	181	Scrub	75	5	0 - 20	1572	0 - 13	Y	Y	N	N
CFG-C011**	112	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-C012	431	Wet Flatwoods	134	30	10 - 50	1707	0 - 0	Y	Y	N	N
CFG-C012	431	Scrubby Flatwoods	34	30	20 - 60	1707	0 - 26	Y	Y	N	N
CFG-C012	431	Mesic Flatwoods	35	30	10 - 50	1707	0 - 0	Y	Y	N	N
CFG-C012**	431	Mesic Hammock	20	--	--	--	--	Y	Y	Y	Y
CFG-C013	1481	Scrub	4	26	0 - 20	608	0 - 13	Y	Y	N	N
CFG-C013	1481	Scrubby Flatwoods	2	26	20 - 60	608	0 - 26	Y	Y	N	N
CFG-C013	1481	Mesic Hammock	93	26	10 - 30	608	0 - 79	Y	Y	N	N
CFG-C013**	1481	Wet Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-C013**	1481	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-C014**	168	Mesic Hammock	74	--	--	--	--	Y	Y	Y	Y



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C015**	110	Mesic Hammock	3	--	--	--	--	Y	Y	Y	Y
CFG-C015**	110	Xeric Hammock	34	--	--	--	--	Y	Y	Y	Y
CFG-C016**	33	Xeric Hammock	16	--	--	--	--	Y	Y	Y	Y
CFG-C017**	27	Wet Flatwoods	18	--	--	--	--	Y	Y	Y	Y
CFG-C018**	15	Mesic Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-C018**	15	Mesic Hammock	4	--	--	--	--	Y	Y	Y	Y
CFG-C020**	140	Mesic Flatwoods	33	--	--	--	--	Y	Y	Y	Y
CFG-C020**	140	Upland Hardwood Forest	64	--	--	--	--	Y	Y	Y	Y
CFG-C021**	39	Pine Plantation	4	--	--	--	--	Y	Y	Y	Y
CFG-C021**	39	Mesic Flatwoods	5	--	--	--	--	Y	Y	Y	Y
CFG-C021**	39	Upland Hardwood Forest	18	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C022**	42	Upland Hardwood Forest	13	--	--	--	--	Y	Y	Y	Y
CFG-C023**	8	Upland Hardwood Forest	8	--	--	--	--	Y	Y	Y	Y
CFG-C025**	96	Mesic Hammock	20	--	--	--	--	Y	Y	Y	Y
CFG-C026**	61	Mesic Hammock	29	--	--	--	--	Y	Y	Y	Y
CFG-C027	60	Pine Plantation	13	43	30 - 80	313	0 - 26	Y	Y	Y	Y
CFG-C027	60	Mesic Hammock	41	43	10 - 30	313	0 - 79	Y	Y	N	N
CFG-C028**	0	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C029	21	Mesic Hammock	21	14	10 - 30	517	0 - 79	Y	Y	N	N
CFG-C030**	2	Mesic Hammock	2	--	--	--	--	Y	Y	Y	Y
CFG-C031**	17	Mesic Hammock	2	--	--	--	--	Y	Y	Y	Y
CFG-C032**	8	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C032**	8	Mesic Hammock	7	--	--	--	--	Y	Y	Y	Y
CFG-C033**	3	Mesic Hammock	3	--	--	--	--	Y	Y	Y	Y
CFG-C034**	0	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C035	5	Mesic Hammock	5	5	10 - 30	1003	0 - 79	Y	Y	Y	Y
CFG-C035**	5	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-C036	4	Mesic Hammock	3	40	10 - 30	516	0 - 79	Y	Y	N	N
CFG-C036**	4	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-C036**	4	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-C037	1	Mesic Hammock	<1	130	10 - 30	281	0 - 79	Y	Y	N	N
CFG-C037**	1	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-C038**	3	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C038**	3	Mesic Flatwoods	2	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C039	5	Mesic Flatwoods	5	50	10 - 50	209	0 - 0	Y	Y	N	N
CFG-C039**	5	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C040	7	Mesic Flatwoods	7	100	10 - 50	500	0 - 0	Y	Y	N	N
CFG-C040**	7	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C041**	2	Mesic Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-C041**	2	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C042**	0	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-C043**	1	Mesic Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-C044	14	Mesic Flatwoods	14	87	10 - 50	1263	0 - 0	Y	Y	N	N
CFG-C045	15	Mesic Flatwoods	10	80	10 - 50	851	0 - 0	Y	Y	N	N
CFG-C046	145	Mesic Hammock	101	44	10 - 30	343	0 - 79	Y	Y	N	N
CFG-C046	145	Mesic Flatwoods	42	44	10 - 50	343	0 - 0	Y	Y	N	N



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C047**	3	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-C048	400	Mesic Flatwoods	302	50	10 - 50	715	0 - 0	Y	Y	N	N
CFG-C049**	44	Mesic Flatwoods	7	--	--	--	--	Y	Y	Y	Y
CFG-C054**	6	Wet Flatwoods	4	--	--	--	--	Y	Y	Y	Y
CFG-C054**	6	Mesic Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-C058	852	Wet Flatwoods	123	62	10 - 50	296	0 - 0	Y	Y	N	N
CFG-C058	852	Pine Plantation	6	62	30 - 80	296	0 - 26	Y	Y	Y	Y
CFG-C058	852	Mesic Flatwoods	21	62	10 - 50	296	0 - 0	Y	Y	N	N
CFG-C058	852	Scrubby Flatwoods	5	62	20 - 60	296	0 - 26	Y	Y	N	N
CFG-C058**	852	Mesic Hammock	12	--	--	--	--	Y	Y	Y	Y
CFG-C059**	22	Mesic Hammock	3	--	--	--	--	Y	Y	Y	Y
CFG-C060	137	Mesic Flatwoods	38	130	10 - 50	799	0 - 0	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C060	137	Mesic Hammock	89	130	10 - 30	799	0 - 79	Y	Y	N	N
CFG-C060	137	Pine Plantation	4	130	30 - 80	799	0 - 26	Y	Y	Y	Y
CFG-C061	167	Pine Plantation	6	21	30 - 80	362	0 - 26	Y	Y	Y	Y
CFG-C061	167	Scrubby Flatwoods	14	21	20 - 60	362	0 - 26	Y	Y	N	N
CFG-C061	167	Wet Flatwoods	22	21	10 - 50	362	0 - 0	Y	Y	N	N
CFG-C061	167	Mesic Hammock	24	21	10 - 30	362	0 - 79	Y	Y	N	N
CFG-C061**	167	Mesic Flatwoods	3	--	--	--	--	Y	Y	Y	Y
CFG-C062	154	Wet Flatwoods	8	80	10 - 50	1786	0 - 0	Y	Y	N	N
CFG-C062	154	Mesic Flatwoods	67	80	10 - 50	1786	0 - 0	Y	Y	N	N
CFG-C063	131	Mesic Flatwoods	26	60	10 - 50	1653	0 - 0	Y	Y	N	N
CFG-C063**	131	Wet Flatwoods	8	--	--	--	--	Y	Y	Y	Y
CFG-C064	65	Mesic Flatwoods	34	131	10 - 50	1338	0 - 0	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C064	65	Wet Flatwoods	24	131	10 - 50	1338	0 - 0	Y	Y	N	N
CFG-C064**	65	Scrubby Flatwoods	3	--	--	--	--	Y	Y	Y	Y
CFG-C065	24	Mesic Flatwoods	10	3	10 - 50	154	0 - 0	Y	Y	Y	Y
CFG-C065	24	Wet Flatwoods	1	3	10 - 50	154	0 - 0	Y	Y	Y	Y
CFG-C065**	24	Scrubby Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-C066	38	Mesic Flatwoods	15	80	10 - 50	287	0 - 0	Y	Y	N	N
CFG-C067	59	Mesic Flatwoods	33	110	10 - 50	1796	0 - 0	Y	Y	N	N
CFG-C068	92	Mesic Flatwoods	63	80	10 - 50	3833	0 - 0	Y	Y	N	N
CFG-C068**	92	Scrubby Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-C069	24	Scrubby Flatwoods	8	40	20 - 60	622	0 - 26	Y	Y	N	N
CFG-C069	24	Mesic Flatwoods	15	40	10 - 50	622	0 - 0	Y	Y	N	N
CFG-C070	57	Mesic Flatwoods	49	90	10 - 50	2634	0 - 0	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C070**	57	Scrubby Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-C071	45	Mesic Flatwoods	26	100	10 - 50	866	0 - 0	Y	Y	N	N
CFG-C072	96	Mesic Flatwoods	52	10	10 - 50	1568	0 - 0	Y	Y	Y	Y
CFG-C072	96	Mesic Hammock	15	10	10 - 30	1568	0 - 79	Y	Y	Y	Y
CFG-C072**	96	Scrubby Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-C074	170	Mesic Hammock	9	80	10 - 30	907	0 - 79	Y	Y	N	N
CFG-C074	170	Mesic Flatwoods	135	80	10 - 50	907	0 - 0	Y	Y	N	N
CFG-C074**	170	Scrubby Flatwoods	4	--	--	--	--	Y	Y	Y	Y
CFG-C075	45	Wet Flatwoods	10	61	10 - 50	917	0 - 0	Y	Y	N	N
CFG-C075	45	Mesic Flatwoods	22	61	10 - 50	917	0 - 0	Y	Y	N	N
CFG-C076	11	Mesic Flatwoods	10	2	10 - 50	2500	0 - 0	Y	Y	Y	Y
CFG-C076**	11	Wet Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-C077	127	Sandhill	8	20	20 - 60	1244	0 - 79	Y	Y	Y	Y



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C077	127	Mesic Hammock	15	20	10 - 30	1244	0 - 79	Y	Y	N	N
CFG-C077	127	Wet Flatwoods	10	20	10 - 50	1244	0 - 0	Y	Y	N	N
CFG-C077	127	Mesic Flatwoods	53	20	10 - 50	1244	0 - 0	Y	Y	N	N
CFG-C077**	127	Scrubby Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-C078**	40	Scrubby Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-C078**	40	Mesic Flatwoods	37	--	--	--	--	Y	Y	Y	Y
CFG-C079	236	Scrubby Flatwoods	20	37	20 - 60	343	0 - 26	Y	Y	N	N
CFG-C079	236	Mesic Hammock	24	37	10 - 30	343	0 - 79	Y	Y	N	N
CFG-C079	236	Mesic Flatwoods	60	37	10 - 50	343	0 - 0	Y	Y	N	N
CFG-C079	236	Sandhill	23	37	20 - 60	343	0 - 79	Y	Y	N	N
CFG-C079**	236	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-C080	30	Scrubby Flatwoods	3	75	20 - 60	2900	0 - 26	Y	Y	N	N
CFG-C080	30	Sandhill	23	75	20 - 60	2900	0 - 79	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C080**	30	Mesic Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-C081	54	Mesic Flatwoods	3	100	10 - 50	100	0 - 0	Y	Y	N	N
CFG-C081	54	Scrubby Flatwoods	16	100	20 - 60	100	0 - 26	Y	Y	N	N
CFG-C081	54	Sandhill	33	100	20 - 60	100	0 - 79	Y	Y	N	N
CFG-C085**	23	Mesic Hammock	14	--	--	--	--	Y	Y	Y	Y
CFG-C086**	61	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C087**	25	Mesic Hammock	16	--	--	--	--	Y	Y	Y	Y
CFG-C088**	11	Mesic Hammock	9	--	--	--	--	Y	Y	Y	Y
CFG-C090**	4	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C091**	7	Mesic Hammock	6	--	--	--	--	Y	Y	Y	Y
CFG-C092**	8	Mesic Hammock	5	--	--	--	--	Y	Y	Y	Y
CFG-C095**	23	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C096**	7	Mesic Hammock	4	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C097**	5	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C097**	5	Mesic Flatwoods	4	--	--	--	--	Y	Y	Y	Y
CFG-C098**	309	Pine Plantation	11	--	--	--	--	Y	Y	Y	Y
CFG-C098**	309	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-C099**	337	Pine Plantation	4	--	--	--	--	Y	Y	Y	Y
CFG-C100**	4	Pine Plantation	4	--	--	--	--	Y	Y	Y	Y
CFG-C103	363	Mesic Flatwoods	5	20	10 - 50	1128	0 - 0	Y	Y	N	N
CFG-C104	2262	Mesic Flatwoods	30	20	10 - 50	334	0 - 0	Y	Y	N	N
CFG-C104**	2262	Mesic Hammock	1	--	--	--	--	Y	Y	Y	Y
CFG-C104**	2262	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-C106	93	Sandhill	6	5	20 - 60	195	0 - 79	Y	Y	Y	Y
CFG-C106	93	Xeric Hammock	14	5	10 - 30	195	0 - 79	Y	Y	Y	Y
CFG-C106	93	Mesic Hammock	51	5	10 - 30	195	0 - 79	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C106**	93	Wet Flatwoods	5	--	--	--	--	Y	Y	Y	Y
CFG-C107	15	Xeric Hammock	10	15	10 - 30	1128	0 - 79	Y	Y	N	N
CFG-C107	15	Mesic Hammock	3	15	10 - 30	1128	0 - 79	Y	Y	N	N
CFG-C108**	11	Xeric Hammock	10	--	--	--	--	Y	Y	Y	Y
CFG-C109**	7	Xeric Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C111	31	Sandhill	17	4	20 - 60	495	0 - 79	Y	Y	Y	Y
CFG-C111	31	Xeric Hammock	9	4	10 - 30	495	0 - 79	Y	Y	Y	Y
CFG-C112**	22	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C112**	22	Xeric Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C113**	5	Mesic Hammock	3	--	--	--	--	Y	Y	Y	Y
CFG-C114	74	Mesic Flatwoods	6	50	10 - 50	78	0 - 0	Y	Y	N	N
CFG-C114	74	Xeric Hammock	15	50	10 - 30	78	0 - 79	Y	N	N	N
CFG-C114	74	Mesic Hammock	8	50	10 - 30	78	0 - 79	Y	N	N	N



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C114**	74	Wet Flatwoods	10	--	--	--	--	Y	Y	Y	Y
CFG-C115**	273	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-C115**	273	Xeric Hammock	1	--	--	--	--	Y	Y	Y	Y
CFG-C115**	273	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C115**	273	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-C116	165	Xeric Hammock	7	50	10 - 30	344	0 - 79	Y	Y	N	N
CFG-C116	165	Scrubby Flatwoods	26	50	20 - 60	344	0 - 26	Y	Y	N	N
CFG-C116	165	Mesic Hammock	24	50	10 - 30	344	0 - 79	Y	Y	N	N
CFG-C116	165	Sandhill	29	50	20 - 60	344	0 - 79	Y	Y	N	N
CFG-C116**	165	Wet Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-C116**	165	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-C117**	9	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C118	104	Scrubby Flatwoods	57	40	20 - 60	1642	0 - 26	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C118	104	Mesic Hammock	34	40	10 - 30	1642	0 - 79	Y	Y	N	N
CFG-C119**	468	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C119**	468	Scrubby Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-C120	47	Xeric Hammock	17	30	10 - 30	994	0 - 79	Y	Y	N	N
CFG-C120**	47	Sandhill	9	--	--	--	--	Y	Y	Y	Y
CFG-C120**	47	Wet Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-C121	30	Sandhill	30	13	20 - 60	270	0 - 79	Y	Y	Y	Y
CFG-C122	152	Sandhill	147	21	20 - 60	509	0 - 79	Y	Y	N	N
CFG-C122**	152	Pine Plantation	3	--	--	--	--	Y	Y	Y	Y
CFG-C123	70	Pine Plantation	70	35	30 - 80	277	0 - 26	Y	Y	Y	Y
CFG-C123**	70	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C124	51	Pine Plantation	51	81	30 - 80	--	0 - 26	Y	N	Y	Y
CFG-C125	67	Pine Plantation	66	39	30 - 80	236	0 - 26	Y	Y	Y	Y
CFG-C126**	77	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C127**	20	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-C128	57	Pine Plantation	56	51	30 - 80	189	0 - 26	Y	Y	Y	Y
CFG-C129**	2	Pine Plantation	2	--	--	--	--	Y	Y	Y	Y
CFG-C130**	15	Mesic Hammock	5	--	--	--	--	Y	Y	Y	Y
CFG-C132**	59	Mesic Hammock	13	--	--	--	--	Y	Y	Y	Y
CFG-C135**	5	Mesic Hammock	5	--	--	--	--	Y	Y	Y	Y
CFG-C136**	77	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C137	24	Sandhill	24	30	20 - 60	551	0 - 79	Y	Y	N	N
CFG-C138**	61	Mesic Hammock	1	--	--	--	--	Y	Y	Y	Y
CFG-C138**	61	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C139	74	Sandhill	71	24	20 - 60	758	0 - 79	Y	Y	N	N
CFG-C140	54	Mesic Hammock	5	5	10 - 30	108	0 - 79	Y	Y	Y	Y
CFG-C141	36	Sandhill	36	2	20 - 60	625	0 - 79	Y	Y	Y	Y
CFG-C141**	36	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C142	278	Upland Pine Forest	6	32	--	1626	--	N	N	N	N
CFG-C142	278	Pine Plantation	271	32	30 - 80	1626	0 - 26	Y	Y	Y	Y
CFG-C142**	278	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C143	19	Pine Plantation	18	7	30 - 80	76	0 - 26	Y	Y	Y	Y
CFG-C144	34	Pine Plantation	34	22	30 - 80	890	0 - 26	Y	Y	Y	Y
CFG-C145	10	Pine Plantation	10	24	30 - 80	100	0 - 26	Y	Y	Y	Y
CFG-C145**	10	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C146**	10	Pine Plantation	6	--	--	--	--	Y	Y	Y	Y
CFG-C146**	10	Sandhill	3	--	--	--	--	Y	Y	Y	Y
CFG-C147**	34	Pine Plantation	2	--	--	--	--	Y	Y	Y	Y
CFG-C148	24	Pine Plantation	24	40	30 - 80	140	0 - 26	Y	Y	Y	Y
CFG-C149	45	Sandhill	44	8	20 - 60	432	0 - 79	Y	Y	Y	Y
CFG-C149**	45	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-C150**	1	Sandhill	1	--	--	--	--	Y	Y	Y	Y



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C151	3	Sandhill	3	20	20 - 60	120	0 - 79	Y	Y	Y	Y
CFG-C152	9	Sandhill	9	30	20 - 60	337	0 - 79	Y	Y	N	N
CFG-C153**	13	Sandhill	12	--	--	--	--	Y	Y	Y	Y
CFG-C154**	3	Xeric Hammock	3	--	--	--	--	Y	Y	Y	Y
CFG-C156	30	Sandhill	30	12	20 - 60	430	0 - 79	Y	Y	Y	Y
CFG-C157	37	Sandhill	35	10	20 - 60	351	0 - 79	Y	Y	Y	Y
CFG-C157**	37	Pine Plantation	1	--	--	--	--	Y	Y	Y	Y
CFG-C158	17	Sandhill	<1	30	20 - 60	191	0 - 79	Y	Y	N	N
CFG-C158**	17	Pine Plantation	16	--	--	--	--	Y	Y	Y	Y
CFG-C159**	3	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C160	12	Sandhill	11	15	20 - 60	257	0 - 79	Y	Y	Y	Y
CFG-C160**	12	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-C161**	1	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C162	15	Sandhill	15	20	20 - 60	334	0 - 79	Y	Y	Y	Y
CFG-C163	21	Mesic Hammock	21	10	10 - 30	12	0 - 79	N	N	Y	Y
CFG-C163**	21	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C164**	17	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C164**	17	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C165**	7	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C166**	8	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C167**	34	Invasive Exotic Monoculture	34	--	--	--	--	Y	Y	Y	Y
CFG-C167**	34	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C168	465	Sandhill	432	20	20 - 60	382	0 - 79	Y	Y	Y	Y
CFG-C168	465	Pine Plantation	18	20	30 - 80	382	0 - 26	Y	Y	Y	Y
CFG-C168**	465	Invasive Exotic Monoculture	<1	--	--	--	--	Y	Y	Y	Y
CFG-C168**	465	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C169	13	Pine Plantation	13	110	30 - 80	339	0 - 26	Y	Y	Y	Y
CFG-C169**	13	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C170	137	Sandhill	13	13	20 - 60	1155	0 - 79	Y	Y	Y	Y
CFG-C170	137	Pine Plantation	124	13	30 - 80	1155	0 - 26	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C171	17	Pine Plantation	17	90	30 - 80	292	0 - 26	Y	Y	Y	Y
CFG-C171**	17	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C172**	12	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C173**	2	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C174	9	Sandhill	8	70	20 - 60	110	0 - 79	Y	Y	N	N
CFG-C175	59	Sandhill	17	5	20 - 60	252	0 - 79	Y	Y	Y	Y
CFG-C176	30	Sandhill	28	0	20 - 60	394	0 - 79	Y	Y	Y	Y
CFG-C176**	30	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-C177	28	Pine Plantation	22	46	30 - 80	873	0 - 26	Y	Y	Y	Y
CFG-C177	28	Sandhill	5	46	20 - 60	873	0 - 79	Y	Y	N	N
CFG-C178	78	Sandhill	76	26	20 - 60	299	0 - 79	Y	Y	N	N
CFG-C178**	78	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-C179	25	Pine Plantation	16	23	30 - 80	479	0 - 26	Y	Y	Y	Y
CFG-C179	25	Sandhill	8	23	20 - 60	479	0 - 79	Y	Y	N	N
CFG-C180**	21	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C181	155	Sandhill	152	23	20 - 60	456	0 - 79	Y	Y	N	N
CFG-C181**	155	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C182	9	Sandhill	9	26	20 - 60	252	0 - 79	Y	Y	N	N
CFG-C183	27	Pine Plantation	14	35	30 - 80	465	0 - 26	Y	Y	Y	Y
CFG-C183	27	Sandhill	13	35	20 - 60	465	0 - 79	Y	Y	N	N
CFG-C184	130	Sandhill	92	8	20 - 60	649	0 - 79	Y	Y	Y	Y
CFG-C184	130	Pine Plantation	34	8	30 - 80	649	0 - 26	Y	Y	Y	Y
CFG-C185	25	Pine Plantation	25	27	30 - 80	560	0 - 26	Y	Y	Y	Y
CFG-C185**	25	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C186	11	Pine Plantation	11	8	30 - 80	120	0 - 26	Y	Y	Y	Y
CFG-C186	11	Sandhill	<1	8	20 - 60	120	0 - 79	Y	Y	Y	Y
CFG-C187**	4	Pine Plantation	3	--	--	--	--	Y	Y	Y	Y
CFG-C187**	4	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C188**	8	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C188**	8	Pine Plantation	7	--	--	--	--	Y	Y	Y	Y
CFG-C189	33	Pine Plantation	32	30	30 - 80	250	0 - 26	Y	Y	Y	Y
CFG-C189**	33	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C190	159	Pine Plantation	42	39	30 - 80	383	0 - 26	Y	Y	Y	Y



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C190	159	Sandhill	112	39	20 - 60	383	0 - 79	Y	Y	N	N
CFG-C191	6	Pine Plantation	6	7	30 - 80	333	0 - 26	Y	Y	Y	Y
CFG-C191**	6	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C192**	11	Sandhill	10	--	--	--	--	Y	Y	Y	Y
CFG-C193	31	Pine Plantation	31	51	30 - 80	110	0 - 26	Y	Y	Y	Y
CFG-C193**	31	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C194	42	Pine Plantation	42	57	30 - 80	14	0 - 26	N	N	Y	Y
CFG-C194**	42	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C195	50	Pine Plantation	48	42	30 - 80	600	0 - 26	Y	Y	Y	Y
CFG-C195**	50	Sandhill	1	--	--	--	--	Y	Y	Y	Y
CFG-C196	18	Pine Plantation	18	8	30 - 80	300	0 - 26	Y	Y	Y	Y
CFG-C197	29	Pine Plantation	29	50	30 - 80	188	0 - 26	Y	Y	Y	Y
CFG-C197**	29	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C198	15	Pine Plantation	15	66	30 - 80	480	0 - 26	Y	Y	Y	Y
CFG-C198**	15	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C199	17	Pine Plantation	16	80	30 - 80	100	0 - 26	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C199**	17	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C200	44	Pine Plantation	15	0	30 - 80	674	0 - 26	Y	Y	Y	Y
CFG-C200	44	Sandhill	29	0	20 - 60	674	0 - 79	Y	Y	Y	Y
CFG-C201	15	Pine Plantation	15	113	30 - 80	500	0 - 26	Y	Y	Y	Y
CFG-C201**	15	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C202	15	Pine Plantation	15	67	30 - 80	225	0 - 26	Y	Y	Y	Y
CFG-C202**	15	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C203	29	Pine Plantation	9	38	30 - 80	638	0 - 26	Y	Y	Y	Y
CFG-C203	29	Sandhill	12	38	20 - 60	638	0 - 79	Y	Y	N	N
CFG-C204	34	Pine Plantation	33	62	30 - 80	107	0 - 26	Y	Y	Y	Y
CFG-C204**	34	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C205	287	Sandhill	117	35	20 - 60	381	0 - 79	Y	Y	N	N
CFG-C205	287	Mesic Hammock	117	35	10 - 30	381	0 - 79	Y	Y	N	N
CFG-C205	287	Pine Plantation	46	35	30 - 80	381	0 - 26	Y	Y	Y	Y
CFG-C206	132	Pine Plantation	130	24	30 - 80	218	0 - 26	Y	Y	Y	Y
CFG-C206**	132	Sandhill	1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C207	135	Sandhill	107	18	20 - 60	87	0 - 79	Y	Y	Y	Y
CFG-C207**	135	Pine Plantation	5	--	--	--	--	Y	Y	Y	Y
CFG-C208**	3	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-C208**	3	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C209**	6	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C210	18	Mesic Hammock	7	10	10 - 30	616	0 - 79	Y	Y	Y	Y
CFG-C210	18	Sandhill	8	10	20 - 60	616	0 - 79	Y	Y	Y	Y
CFG-C211	75	Sandhill	54	1	20 - 60	438	0 - 79	Y	Y	Y	Y
CFG-C212**	68	Sandhill	1	--	--	--	--	Y	Y	Y	Y
CFG-C213**	20	Pine Plantation	19	--	--	--	--	Y	Y	Y	Y
CFG-C214	64	Mesic Hammock	59	12	10 - 30	360	0 - 79	Y	Y	N	N
CFG-C214**	64	Sandhill	1	--	--	--	--	Y	Y	Y	Y
CFG-C215	26	Sandhill	26	9	20 - 60	1580	0 - 79	Y	Y	N	N
CFG-C216	28	Sandhill	28	6	20 - 60	786	0 - 79	Y	Y	N	N
CFG-C218	61	Pine Plantation	60	48	30 - 80	378	0 - 26	Y	Y	Y	Y
CFG-C218**	61	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C220	40	Mesic Hammock	7	60	10 - 30	100	0 - 79	Y	Y	N	N
CFG-C220	40	Pine Plantation	17	60	30 - 80	100	0 - 26	Y	Y	Y	Y
CFG-C221	75	Pine Plantation	75	60	30 - 80	77	0 - 26	Y	Y	Y	Y
CFG-C222	119	Pine Plantation	118	51	30 - 80	167	0 - 26	Y	Y	Y	Y
CFG-C222**	119	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C223	286	Mesic Hammock	255	5	10 - 30	352	0 - 79	Y	Y	Y	Y
CFG-C223	286	Pine Plantation	18	5	30 - 80	352	0 - 26	Y	Y	Y	Y
CFG-C223**	286	Sandhill	3	--	--	--	--	Y	Y	Y	Y
CFG-C224	115	Mesic Hammock	15	20	10 - 30	1567	0 - 79	Y	Y	N	N
CFG-C224	115	Sandhill	91	20	20 - 60	1567	0 - 79	Y	Y	N	N
CFG-C224**	115	Pine Plantation	7	--	--	--	--	Y	Y	Y	Y
CFG-C225**	0	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C226**	4	Scrub	4	--	--	--	--	Y	Y	Y	Y
CFG-C227**	9	Scrub	9	--	--	--	--	Y	Y	Y	Y
CFG-C228**	30	Scrub	30	--	--	--	--	Y	Y	Y	Y



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C229**	22	Scrub	22	--	--	--	--	Y	Y	Y	Y
CFG-C230**	26	Scrub	26	--	--	--	--	Y	Y	Y	Y
CFG-C231**	15	Scrub	15	--	--	--	--	Y	Y	Y	Y
CFG-C232**	16	Scrub	16	--	--	--	--	Y	Y	Y	Y
CFG-C233**	28	Scrub	28	--	--	--	--	Y	Y	Y	Y
CFG-C234**	17	Scrub	17	--	--	--	--	Y	Y	Y	Y
CFG-C235**	28	Scrub	28	--	--	--	--	Y	Y	Y	Y
CFG-C236**	26	Scrub	26	--	--	--	--	Y	Y	Y	Y
CFG-C237**	24	Sandhill	1	--	--	--	--	Y	Y	Y	Y
CFG-C237**	24	Scrub	23	--	--	--	--	Y	Y	Y	Y
CFG-C238**	36	Sandhill	33	--	--	--	--	Y	Y	Y	Y
CFG-C238**	36	Scrub	2	--	--	--	--	Y	Y	Y	Y
CFG-C239**	44	Scrub	44	--	--	--	--	Y	Y	Y	Y
CFG-C239**	44	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C240**	24	Scrub	23	--	--	--	--	Y	Y	Y	Y
CFG-C240**	24	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C241**	13	Scrub	13	--	--	--	--	Y	Y	Y	Y
CFG-C242**	69	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C242**	69	Scrub	68	--	--	--	--	Y	Y	Y	Y
CFG-C243	25	Sandhill	24	48	20 - 60	275	0 - 79	Y	Y	N	N
CFG-C244	71	Sandhill	50	65	20 - 60	415	0 - 79	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C244**	71	Scrub	4	--	--	--	--	Y	Y	Y	Y
CFG-C245**	20	Sandhill	2	--	--	--	--	Y	Y	Y	Y
CFG-C245**	20	Scrub	17	--	--	--	--	Y	Y	Y	Y
CFG-C245A**	27	Scrub	27	--	--	--	--	Y	Y	Y	Y
CFG-C246**	13	Scrub	13	--	--	--	--	Y	Y	Y	Y
CFG-C247**	12	Scrub	12	--	--	--	--	Y	Y	Y	Y
CFG-C248**	13	Scrub	13	--	--	--	--	Y	Y	Y	Y
CFG-C249**	14	Scrub	14	--	--	--	--	Y	Y	Y	Y
CFG-C250**	17	Scrub	17	--	--	--	--	Y	Y	Y	Y
CFG-C251**	4	Scrub	4	--	--	--	--	Y	Y	Y	Y
CFG-C252**	31	Scrub	31	--	--	--	--	Y	Y	Y	Y
CFG-C252**	31	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C253**	5	Scrub	5	--	--	--	--	Y	Y	Y	Y
CFG-C253**	5	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C254	118	Scrub	114	160	0 - 20	1100	0 - 13	Y	Y	N	N
CFG-C254	118	Mesic Flatwoods	3	160	10 - 50	1100	0 - 0	Y	Y	N	N
CFG-C254**	118	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C255	561	Mesic Flatwoods	49	10	10 - 50	300	0 - 0	Y	Y	Y	Y
CFG-C255	561	Mesic Hammock	35	10	10 - 30	300	0 - 79	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C255	561	Sandhill	77	10	20 - 60	300	0 - 79	Y	Y	Y	Y
CFG-C255	561	Scrub	393	10	0 - 20	300	0 - 13	Y	Y	N	N
CFG-C256	159	Sandhill	154	37	20 - 60	790	0 - 79	Y	Y	N	N
CFG-C256**	159	Scrub	4	--	--	--	--	Y	Y	Y	Y
CFG-C257	149	Sandhill	137	30	20 - 60	330	0 - 79	Y	Y	N	N
CFG-C257**	149	Scrub	11	--	--	--	--	Y	Y	Y	Y
CFG-C258	146	Sandhill	144	57	20 - 60	319	0 - 79	Y	Y	N	N
CFG-C259	140	Sandhill	139	55	20 - 60	220	0 - 79	Y	Y	N	N
CFG-C260	12	Sandhill	9	64	20 - 60	208	0 - 79	Y	Y	N	N
CFG-C261	301	Mesic Hammock	7	5	10 - 30	1509	0 - 79	Y	Y	Y	Y
CFG-C261	301	Sandhill	289	5	20 - 60	1509	0 - 79	Y	Y	Y	Y
CFG-C262	58	Sandhill	21	36	20 - 60	689	0 - 79	Y	Y	N	N
CFG-C263	299	Sandhill	284	20	20 - 60	1246	0 - 79	Y	Y	N	N
CFG-C263	299	Mesic Hammock	10	20	10 - 30	1246	0 - 79	Y	Y	N	N
CFG-C264	201	Sandhill	200	34	20 - 60	426	0 - 79	Y	Y	N	N
CFG-C265	212	Sandhill	212	42	20 - 60	651	0 - 79	Y	Y	N	N
CFG-C266	394	Sandhill	41	62	20 - 60	585	0 - 79	Y	Y	N	N
CFG-C266**	394	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-C267	186	Sandhill	185	22	20 - 60	648	0 - 79	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C267**	186	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-C268	46	Pine Plantation	46	45	30 - 80	1880	0 - 26	Y	Y	Y	Y
CFG-C269	12	Pine Plantation	12	16	30 - 80	480	0 - 26	Y	Y	Y	Y
CFG-C270**	14	Pine Plantation	3	--	--	--	--	Y	Y	Y	Y
CFG-C270**	14	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-C271	20	Sandhill	19	26	20 - 60	1431	0 - 79	Y	Y	N	N
CFG-C271**	20	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-C272	181	Sandhill	180	17	20 - 60	523	0 - 79	Y	Y	Y	Y
CFG-C273	171	Sandhill	160	6	20 - 60	87	0 - 79	Y	Y	Y	Y
CFG-C273	171	Scrub	10	6	0 - 20	87	0 - 13	Y	Y	N	N
CFG-C274	230	Scrub	1	30	0 - 20	993	0 - 13	Y	Y	N	N
CFG-C274	230	Sandhill	229	30	20 - 60	993	0 - 79	Y	Y	N	N
CFG-C275	475	Xeric Hammock	11	5	10 - 30	635	0 - 79	Y	Y	Y	Y
CFG-C275	475	Sandhill	70	5	20 - 60	635	0 - 79	Y	Y	Y	Y
CFG-C275	475	Scrub	21	5	0 - 20	635	0 - 13	Y	Y	N	N
CFG-C275**	475	Mesic Hammock	4	--	--	--	--	Y	Y	Y	Y



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-C276	126	Sandhill	124	22	20 - 60	635	0 - 79	Y	Y	N	N
CFG-C276A	82	Sandhill	82	20	20 - 60	913	0 - 79	Y	Y	N	N
CFG-C276B	157	Sandhill	120	6	20 - 60	189	0 - 79	Y	Y	Y	Y
CFG-C276B	157	Mesic Hammock	30	6	10 - 30	189	0 - 79	Y	Y	Y	Y
CFG-C277	182	Mesic Hammock	47	0	10 - 30	414	0 - 79	Y	Y	Y	Y
CFG-C277**	182	Xeric Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-C278	107	Sandhill	10	1	20 - 60	674	0 - 79	Y	Y	Y	Y
CFG-C278	107	Mesic Hammock	70	1	10 - 30	674	0 - 79	Y	Y	Y	Y
CFG-C279**	28	Mesic Hammock	2	--	--	--	--	Y	Y	Y	Y
CFG-C280**	17	Mesic Hammock	17	--	--	--	--	Y	Y	Y	Y
CFG-C281	11	Mesic Hammock	10	3	10 - 30	1172	0 - 79	Y	Y	Y	Y
CFG-E003**	108	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E005**	5	Pine Plantation	5	--	--	--	--	Y	Y	Y	Y
CFG-E006	12	Pine Plantation	12	70	30 - 80	432	0 - 26	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E007**	2	Clearcut Pine Plantation	2	--	--	--	--	Y	Y	Y	Y
CFG-E008	125	Pine Plantation	40	80	30 - 80	964	0 - 26	Y	Y	Y	Y
CFG-E008	125	Clearcut Pine Plantation	77	80	30 - 80	964	0 - 26	Y	Y	Y	Y
CFG-E009**	21	Pine Plantation	1	--	--	--	--	Y	Y	Y	Y
CFG-E010**	42	Pine Plantation	40	--	--	--	--	Y	Y	Y	Y
CFG-E011**	81	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E012	135	Pine Plantation	116	95	30 - 80	525	0 - 26	Y	Y	Y	Y
CFG-E013**	46	Pine Plantation	44	--	--	--	--	Y	Y	Y	Y
CFG-E014	155	Pine Plantation	136	64	30 - 80	657	0 - 26	Y	Y	Y	Y
CFG-E015**	31	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E015**	31	Pine Plantation	1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E016	16	Mesic Flatwoods	8	60	10 - 50	400	0 - 0	Y	Y	N	N
CFG-E016**	16	Pine Plantation	3	--	--	--	--	Y	Y	Y	Y
CFG-E017**	31	Pine Plantation	1	--	--	--	--	Y	Y	Y	Y
CFG-E017**	31	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E018	80	Pine Plantation	60	154	30 - 80	680	0 - 26	Y	Y	Y	Y
CFG-E019**	32	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E020	7	Mesic Flatwoods	5	20	10 - 50	256	0 - 0	Y	Y	N	N
CFG-E021	25	Pine Plantation	7	132	30 - 80	--	0 - 26	Y	N	Y	Y
CFG-E021**	25	Mesic Flatwoods	4	--	--	--	--	Y	Y	Y	Y
CFG-E022**	1	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E023**	2	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E024**	0	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E025	6	Pine Plantation	5	34	30 - 80	650	0 - 26	Y	Y	Y	Y
CFG-E026	9	Pine Plantation	9	145	30 - 80	1725	0 - 26	Y	Y	Y	Y
CFG-E027	7	Pine Plantation	6	134	30 - 80	800	0 - 26	Y	Y	Y	Y
CFG-E028**	16	Pine Plantation	12	--	--	--	--	Y	Y	Y	Y
CFG-E029	44	Pine Plantation	44	54	30 - 80	868	0 - 26	Y	Y	Y	Y
CFG-E030	53	Pine Plantation	52	58	30 - 80	683	0 - 26	Y	Y	Y	Y
CFG-E031**	13	Pine Plantation	2	--	--	--	--	Y	Y	Y	Y
CFG-E032	53	Pine Plantation	50	84	30 - 80	1662	0 - 26	Y	Y	Y	Y
CFG-E032**	53	Clearcut Pine Plantation	2	--	--	--	--	Y	Y	Y	Y
CFG-E033	5	Pine Plantation	5	63	30 - 80	966	0 - 26	Y	Y	Y	Y
CFG-E034	155	Pine Plantation	143	115	30 - 80	1169	0 - 26	Y	Y	Y	Y
CFG-E035	53	Pine Plantation	51	87	30 - 80	1471	0 - 26	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E036**	200	Wet Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E036**	200	Pine Plantation	41	--	--	--	--	Y	Y	Y	Y
CFG-E037	85	Pine Plantation	85	96	30 - 80	2316	0 - 26	Y	Y	Y	Y
CFG-E038	32	Pine Plantation	28	7	30 - 80	1249	0 - 26	Y	Y	Y	Y
CFG-E039	134	Pine Plantation	125	50	30 - 80	1711	0 - 26	Y	Y	Y	Y
CFG-E040	47	Pine Plantation	44	117	30 - 80	1650	0 - 26	Y	Y	Y	Y
CFG-E041	23	Pine Plantation	23	77	30 - 80	425	0 - 26	Y	Y	Y	Y
CFG-E042	7	Pine Plantation	7	160	30 - 80	100	0 - 26	Y	Y	Y	Y
CFG-E043	28	Pine Plantation	26	47	30 - 80	1738	0 - 26	Y	Y	Y	Y
CFG-E044	448	Pine Plantation	154	45	30 - 80	1600	0 - 26	Y	Y	Y	Y
CFG-E044	448	Mesic Flatwoods	18	45	10 - 50	1600	0 - 0	Y	Y	N	N
CFG-E044**	448	Wet Flatwoods	8	--	--	--	--	Y	Y	Y	Y



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E045	97	Pine Plantation	89	51	30 - 80	955	0 - 26	Y	Y	Y	Y
CFG-E045**	97	Mesic Flatwoods	3	--	--	--	--	Y	Y	Y	Y
CFG-E046	67	Pine Plantation	62	4	30 - 80	1860	0 - 26	Y	Y	Y	Y
CFG-E046	67	Mesic Flatwoods	3	4	10 - 50	1860	0 - 0	Y	Y	Y	Y
CFG-E047	15	Pine Plantation	15	75	30 - 80	1213	0 - 26	Y	Y	Y	Y
CFG-E048	10	Pine Plantation	10	9	30 - 80	666	0 - 26	Y	Y	Y	Y
CFG-E049	116	Pine Plantation	104	46	30 - 80	941	0 - 26	Y	Y	Y	Y
CFG-E049**	116	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E050	64	Pine Plantation	53	21	30 - 80	1700	0 - 26	Y	Y	Y	Y
CFG-E050**	64	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E051**	4	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E051**	4	Mesic Flatwoods	4	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E052	11	Pine Plantation	11	57	30 - 80	1925	0 - 26	Y	Y	Y	Y
CFG-E052**	11	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E053**	49	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E053**	49	Mesic Flatwoods	42	--	--	--	--	Y	Y	Y	Y
CFG-E054	50	Pine Plantation	39	58	30 - 80	2084	0 - 26	Y	Y	Y	Y
CFG-E054**	50	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E055	22	Pine Plantation	21	86	30 - 80	6347	0 - 26	Y	Y	Y	Y
CFG-E056	73	Pine Plantation	72	74	30 - 80	2249	0 - 26	Y	Y	Y	Y
CFG-E057	14	Pine Plantation	14	55	30 - 80	650	0 - 26	Y	Y	Y	Y
CFG-E058	61	Pine Plantation	58	64	30 - 80	4673	0 - 26	Y	Y	Y	Y
CFG-E059	46	Pine Plantation	42	50	30 - 80	3102	0 - 26	Y	Y	Y	Y
CFG-E059**	46	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E060	12	Mesic Flatwoods	9	50	10 - 50	4125	0 - 0	Y	Y	N	N
CFG-E061	55	Pine Plantation	47	86	30 - 80	1077	0 - 26	Y	Y	Y	Y
CFG-E062	111	Pine Plantation	102	120	30 - 80	2492	0 - 26	Y	Y	Y	Y
CFG-E062**	111	Mesic Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E063	35	Pine Plantation	29	77	30 - 80	4801	0 - 26	Y	Y	Y	Y
CFG-E064	91	Pine Plantation	88	86	30 - 80	1137	0 - 26	Y	Y	Y	Y
CFG-E065	474	Pine Plantation	403	10	30 - 80	534	0 - 26	Y	Y	Y	Y
CFG-E065	474	Mesic Flatwoods	8	10	10 - 50	534	0 - 0	Y	Y	Y	Y
CFG-E066**	105	Sandhill	4	--	--	--	--	Y	Y	Y	Y
CFG-E066**	105	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E067	8	Pine Plantation	8	87	30 - 80	429	0 - 26	Y	Y	Y	Y
CFG-E068	57	Pine Plantation	49	80	30 - 80	2000	0 - 26	Y	Y	Y	Y
CFG-E068**	57	Mesic Flatwoods	4	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E069	12	Pine Plantation	11	100	30 - 80	1675	0 - 26	Y	Y	Y	Y
CFG-E070	745	Pine Plantation	22	40	30 - 80	1273	0 - 26	Y	Y	Y	Y
CFG-E070**	745	Mesic Flatwoods	13	--	--	--	--	Y	Y	Y	Y
CFG-E071**	10	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E072**	35	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E072**	35	Mesic Flatwoods	34	--	--	--	--	Y	Y	Y	Y
CFG-E073	115	Pine Plantation	113	56	30 - 80	823	0 - 26	Y	Y	Y	Y
CFG-E073**	115	Mesic Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E074	190	Pine Plantation	161	20	30 - 80	1000	0 - 26	Y	Y	Y	Y
CFG-E074	190	Wet Flatwoods	22	20	10 - 50	1000	0 - 0	Y	Y	N	N
CFG-E074**	190	Clearcut Pine Plantation	2	--	--	--	--	Y	Y	Y	Y
CFG-E075	36	Wet Flatwoods	32	59	10 - 50	3184	0 - 0	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E075**	36	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E076	21	Pine Plantation	17	77	30 - 80	1480	0 - 26	Y	Y	Y	Y
CFG-E077	58	Pine Plantation	56	60	30 - 80	1850	0 - 26	Y	Y	Y	Y
CFG-E077**	58	Wet Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E078**	160	Wet Flatwoods	24	--	--	--	--	Y	Y	Y	Y
CFG-E078**	160	Pine Plantation	3	--	--	--	--	Y	Y	Y	Y
CFG-E079	135	Pine Plantation	133	10	30 - 80	514	0 - 26	Y	Y	Y	Y
CFG-E079	135	Mesic Flatwoods	<1	10	10 - 50	514	0 - 0	Y	Y	Y	Y
CFG-E080	41	Pine Plantation	4	55	30 - 80	662	0 - 26	Y	Y	Y	Y
CFG-E081**	35	Pine Plantation	35	--	--	--	--	Y	Y	Y	Y
CFG-E082	25	Pine Plantation	25	81	30 - 80	1485	0 - 26	Y	Y	Y	Y
CFG-E083	23	Pine Plantation	22	94	30 - 80	883	0 - 26	Y	Y	Y	Y



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E084	10	Pine Plantation	10	146	30 - 80	1433	0 - 26	Y	Y	Y	Y
CFG-E085	163	Pine Plantation	154	75	30 - 80	523	0 - 26	Y	Y	Y	Y
CFG-E085**	163	Mesic Hammock	3	--	--	--	--	Y	Y	Y	Y
CFG-E085**	163	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E086**	41	Pine Plantation	1	--	--	--	--	Y	Y	Y	Y
CFG-E087	7	Pine Plantation	5	20	30 - 80	1584	0 - 26	Y	Y	Y	Y
CFG-E088	260	Pine Plantation	211	14	30 - 80	2180	0 - 26	Y	Y	Y	Y
CFG-E088**	260	Clearcut Pine Plantation	4	--	--	--	--	Y	Y	Y	Y
CFG-E088**	260	Mesic Hammock	3	--	--	--	--	Y	Y	Y	Y
CFG-E088**	260	Wet Flatwoods	25	--	--	--	--	Y	Y	Y	Y
CFG-E089	18	Pine Plantation	16	46	30 - 80	608	0 - 26	Y	Y	Y	Y
CFG-E090	35	Pine Plantation	33	50	30 - 80	189	0 - 26	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E091	13	Pine Plantation	13	0	30 - 80	2173	0 - 26	Y	Y	Y	Y
CFG-E092	78	Pine Plantation	78	84	30 - 80	33	0 - 26	Y	Y	Y	Y
CFG-E092**	78	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E093	45	Pine Plantation	43	88	30 - 80	57	0 - 26	Y	Y	Y	Y
CFG-E093**	45	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E094**	55	Pine Plantation	35	--	--	--	--	Y	Y	Y	Y
CFG-E094**	55	Mesic Flatwoods	19	--	--	--	--	Y	Y	Y	Y
CFG-E095	18	Pine Plantation	18	62	30 - 80	273	0 - 26	Y	Y	Y	Y
CFG-E096	7	Pine Plantation	6	53	30 - 80	1176	0 - 26	Y	Y	Y	Y
CFG-E096**	7	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E097	232	Scrubby Flatwoods	40	18	20 - 60	2237	0 - 26	Y	Y	Y	Y
CFG-E097	232	Mesic Flatwoods	133	18	10 - 50	2237	0 - 0	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E097**	232	Wet Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-E097**	232	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E098	28	Scrubby Flatwoods	28	15	20 - 60	1442	0 - 26	Y	Y	Y	Y
CFG-E099	1018	Mesic Flatwoods	7	20	10 - 50	1066	0 - 0	Y	Y	N	N
CFG-E099**	1018	Scrubby Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-E099**	1018	Mesic Hammock	8	--	--	--	--	Y	Y	Y	Y
CFG-E099**	1018	Wet Flatwoods	16	--	--	--	--	Y	Y	Y	Y
CFG-E099**	1018	Pine Plantation	25	--	--	--	--	Y	Y	Y	Y
CFG-E100	113	Wet Flatwoods	37	30	10 - 50	3444	0 - 0	Y	Y	N	N
CFG-E100	113	Scrub	59	30	0 - 20	3444	0 - 13	Y	Y	N	N
CFG-E100**	113	Scrubby Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-E100**	113	Mesic Flatwoods	12	--	--	--	--	Y	Y	Y	Y
CFG-E101	90	Wet Flatwoods	21	20	10 - 50	1552	0 - 0	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E101	90	Scrub	11	20	0 - 20	1552	0 - 13	Y	Y	N	N
CFG-E101**	90	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E102**	3	Pine Plantation	3	--	--	--	--	Y	Y	Y	Y
CFG-E102**	3	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E103	1139	Scrubby Flatwoods	19	19	20 - 60	3300	0 - 26	Y	Y	Y	Y
CFG-E103	1139	Pine Plantation	90	19	30 - 80	3300	0 - 26	Y	Y	Y	Y
CFG-E103	1139	Mesic Flatwoods	17	19	10 - 50	3300	0 - 0	Y	Y	N	N
CFG-E103	1139	Wet Flatwoods	118	19	10 - 50	3300	0 - 0	Y	Y	N	N
CFG-E103	1139	Mesic Hammock	14	19	10 - 30	3300	0 - 79	Y	Y	N	N
CFG-E104	26	Pine Plantation	26	62	30 - 80	183	0 - 26	Y	Y	Y	Y
CFG-E105	60	Pine Plantation	55	10	30 - 80	296	0 - 26	Y	Y	Y	Y
CFG-E105	60	Wet Flatwoods	<1	10	10 - 50	296	0 - 0	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E105**	60	Clearcut Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E106**	24	Clearcut Pine Plantation	1	--	--	--	--	Y	Y	Y	Y
CFG-E106**	24	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E106**	24	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E107	12	Pine Plantation	12	165	30 - 80	1500	0 - 26	Y	Y	Y	Y
CFG-E107**	12	Clearcut Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E108	31	Pine Plantation	30	34	30 - 80	2202	0 - 26	Y	Y	Y	Y
CFG-E108**	31	Clearcut Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E109**	4	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E110**	4	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E110**	4	Clearcut Pine Plantation	4	--	--	--	--	Y	Y	Y	Y
CFG-E111	16	Clearcut Pine Plantation	14	20	30 - 80	2480	0 - 26	Y	Y	Y	Y
CFG-E111**	16	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E112	18	Pine Plantation	17	35	30 - 80	250	0 - 26	Y	Y	Y	Y
CFG-E112**	18	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E113	23	Pine Plantation	22	46	30 - 80	778	0 - 26	Y	Y	Y	Y
CFG-E113**	23	Wet Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E114	41	Pine Plantation	38	58	30 - 80	2968	0 - 26	Y	Y	Y	Y
CFG-E115	10	Pine Plantation	10	60	30 - 80	1491	0 - 26	Y	Y	Y	Y
CFG-E116**	5	Pine Plantation	4	--	--	--	--	Y	Y	Y	Y
CFG-E117	42	Pine Plantation	41	50	30 - 80	2681	0 - 26	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E118	16	Pine Plantation	15	50	30 - 80	1672	0 - 26	Y	Y	Y	Y
CFG-E119	188	Pine Plantation	168	45	30 - 80	787	0 - 26	Y	Y	Y	Y
CFG-E120	142	Pine Plantation	137	66	30 - 80	803	0 - 26	Y	Y	Y	Y
CFG-E121	12	Pine Plantation	12	7	30 - 80	786	0 - 26	Y	Y	Y	Y
CFG-E122	39	Pine Plantation	37	92	30 - 80	963	0 - 26	Y	Y	Y	Y
CFG-E123	11	Pine Plantation	11	66	30 - 80	1814	0 - 26	Y	Y	Y	Y
CFG-E124	30	Pine Plantation	28	40	30 - 80	2549	0 - 26	Y	Y	Y	Y
CFG-E124**	30	Wet Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E124**	30	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E124**	30	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-E125	4	Pine Plantation	4	92	30 - 80	1990	0 - 26	Y	Y	Y	Y
CFG-E125**	4	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E126	230	Sandhill	8	7	20 - 60	768	0 - 79	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E126	230	Pine Plantation	196	7	30 - 80	768	0 - 26	Y	Y	Y	Y
CFG-E126	230	Upland Hardwood Forest	1	7	30 - 80	768	0 - 263	Y	Y	Y	Y
CFG-E126**	230	Wet Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E126**	230	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E127	47	Sandhill	43	4	20 - 60	767	0 - 79	Y	Y	Y	Y
CFG-E127**	47	Pine Plantation	3	--	--	--	--	Y	Y	Y	Y
CFG-E128	34	Pine Plantation	33	48	30 - 80	1453	0 - 26	Y	Y	Y	Y
CFG-E128**	34	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E128**	34	Sandhill	1	--	--	--	--	Y	Y	Y	Y
CFG-E129	59	Wet Flatwoods	6	42	10 - 50	8424	0 - 0	Y	Y	N	N
CFG-E129	59	Pine Plantation	52	42	30 - 80	8424	0 - 26	Y	Y	Y	Y
CFG-E130	39	Pine Plantation	32	20	30 - 80	1275	0 - 26	Y	Y	Y	Y
CFG-E131**	78	Pine Plantation	9	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E131**	78	Wet Flatwoods	16	--	--	--	--	Y	Y	Y	Y
CFG-E132	45	Pine Plantation	43	90	30 - 80	408	0 - 26	Y	Y	Y	Y
CFG-E133	53	Pine Plantation	49	1	30 - 80	4679	0 - 26	Y	Y	Y	Y
CFG-E134	21	Pine Plantation	20	27	30 - 80	2600	0 - 26	Y	Y	Y	Y
CFG-E135	13	Scrubby Flatwoods	6	10	20 - 60	2300	0 - 26	Y	Y	Y	Y
CFG-E135	13	Scrub	6	10	0 - 20	2300	0 - 13	Y	Y	N	N
CFG-E135**	13	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E136	99	Pine Plantation	50	80	30 - 80	914	0 - 26	Y	Y	Y	Y
CFG-E136**	99	Mesic Flatwoods	4	--	--	--	--	Y	Y	Y	Y
CFG-E137	35	Pine Plantation	33	100	30 - 80	887	0 - 26	Y	Y	Y	Y
CFG-E138	102	Pine Plantation	100	52	30 - 80	4032	0 - 26	Y	Y	Y	Y
CFG-E138**	102	Mesic Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E139	26	Pine Plantation	24	63	30 - 80	4840	0 - 26	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E139**	26	Mesic Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E140	62	Pine Plantation	27	20	30 - 80	935	0 - 26	Y	Y	Y	Y
CFG-E140	62	Mesic Flatwoods	33	20	10 - 50	935	0 - 0	Y	Y	N	N
CFG-E141	168	Wet Flatwoods	55	5	10 - 50	2281	0 - 0	Y	Y	Y	Y
CFG-E141	168	Scrubby Flatwoods	17	5	20 - 60	2281	0 - 26	Y	Y	Y	Y
CFG-E141	168	Mesic Hammock	62	5	10 - 30	2281	0 - 79	Y	Y	Y	Y
CFG-E141	168	Scrub	25	5	0 - 20	2281	0 - 13	Y	Y	N	N
CFG-E142	87	Pine Plantation	21	60	30 - 80	729	0 - 26	Y	Y	Y	Y
CFG-E142	87	Mesic Flatwoods	62	60	10 - 50	729	0 - 0	Y	Y	N	N
CFG-E142**	87	Clearcut Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E143	14	Clearcut Pine Plantation	14	56	30 - 80	2600	0 - 26	Y	Y	Y	Y



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E144**	23	Clearcut Pine Plantation	2	--	--	--	--	Y	Y	Y	Y
CFG-E145	24	Clearcut Pine Plantation	18	9	30 - 80	3687	0 - 26	Y	Y	Y	Y
CFG-E145**	24	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E146	9	Pine Plantation	9	150	30 - 80	1000	0 - 26	Y	Y	Y	Y
CFG-E147	27	Pine Plantation	25	98	30 - 80	621	0 - 26	Y	Y	Y	Y
CFG-E148	15	Pine Plantation	15	170	30 - 80	2403	0 - 26	Y	Y	Y	Y
CFG-E149	29	Pine Plantation	24	66	30 - 80	1300	0 - 26	Y	Y	Y	Y
CFG-E149	29	Scrub	4	66	0 - 20	1300	0 - 13	Y	Y	N	N
CFG-E150	7	Pine Plantation	4	50	30 - 80	300	0 - 26	Y	Y	Y	Y
CFG-E150	7	Scrub	3	50	0 - 20	300	0 - 13	Y	Y	N	N
CFG-E151	6	Scrub	2	10	0 - 20	1000	0 - 13	Y	Y	N	N
CFG-E151**	6	Pine Plantation	3	--	--	--	--	Y	Y	Y	Y
CFG-E152**	15	Pine Plantation	10	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E152**	15	Scrub	4	--	--	--	--	Y	Y	Y	Y
CFG-E153	11	Pine Plantation	11	56	30 - 80	1487	0 - 26	Y	Y	Y	Y
CFG-E154	114	Pine Plantation	106	40	30 - 80	1500	0 - 26	Y	Y	Y	Y
CFG-E154**	114	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-E154**	114	Mesic Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-E154**	114	Scrub	<1	--	--	--	--	Y	Y	Y	Y
CFG-E155	69	Pine Plantation	56	69	30 - 80	5657	0 - 26	Y	Y	Y	Y
CFG-E155**	69	Mesic Flatwoods	11	--	--	--	--	Y	Y	Y	Y
CFG-E156**	23	Mesic Flatwoods	22	--	--	--	--	Y	Y	Y	Y
CFG-E156**	23	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E157**	19	Mesic Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E157**	19	Scrubby Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E157**	19	Pine Plantation	15	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E158**	11	Scrubby Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E158**	11	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E158**	11	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E159**	22	Scrubby Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-E159**	22	Mesic Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E159**	22	Mesic Hammock	1	--	--	--	--	Y	Y	Y	Y
CFG-E160**	5	Mesic Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E160**	5	Mesic Hammock	1	--	--	--	--	Y	Y	Y	Y
CFG-E161	420	Wet Flatwoods	12	22	10 - 50	269	0 - 0	Y	Y	N	N
CFG-E161	420	Sandhill	21	22	20 - 60	269	0 - 79	Y	Y	N	N
CFG-E161	420	Scrub	9	22	0 - 20	269	0 - 13	Y	Y	N	N
CFG-E161**	420	Pine Plantation	3	--	--	--	--	Y	Y	Y	Y
CFG-E161**	420	Scrubby Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E162	107	Scrub	27	25	0 - 20	165	0 - 13	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E162	107	Scrubby Flatwoods	38	25	20 - 60	165	0 - 26	Y	Y	N	N
CFG-E162	107	Mesic Flatwoods	33	25	10 - 50	165	0 - 0	Y	Y	N	N
CFG-E162**	107	Sandhill	5	--	--	--	--	Y	Y	Y	Y
CFG-E162**	107	Pine Plantation	2	--	--	--	--	Y	Y	Y	Y
CFG-E163	105	Pine Plantation	35	40	30 - 80	2449	0 - 26	Y	Y	Y	Y
CFG-E163	105	Mesic Flatwoods	57	40	10 - 50	2449	0 - 0	Y	Y	N	N
CFG-E163**	105	Scrubby Flatwoods	4	--	--	--	--	Y	Y	Y	Y
CFG-E164**	1	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E164**	1	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E165**	32	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E165**	32	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E166	50	Pine Plantation	21	70	30 - 80	100	0 - 26	Y	Y	Y	Y
CFG-E166**	50	Scrub	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E166**	50	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E167	68	Mesic Hammock	<1	80	10 - 30	1022	0 - 79	Y	Y	N	N
CFG-E167**	68	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E168	151	Mesic Flatwoods	126	37	10 - 50	228	0 - 0	Y	Y	N	N
CFG-E169	108	Mesic Flatwoods	97	22	10 - 50	3200	0 - 0	Y	Y	N	N
CFG-E170	86	Mesic Flatwoods	50	101	10 - 50	750	0 - 0	Y	Y	N	N
CFG-E170	86	Mesic Hammock	27	101	10 - 30	750	0 - 79	Y	Y	N	N
CFG-E171	74	Mesic Hammock	9	25	10 - 30	897	0 - 79	Y	Y	N	N
CFG-E171**	74	Mesic Flatwoods	27	--	--	--	--	Y	Y	Y	Y
CFG-E172	39	Mesic Flatwoods	22	70	10 - 50	670	0 - 0	Y	Y	N	N
CFG-E172	39	Mesic Hammock	16	70	10 - 30	670	0 - 79	Y	Y	N	N
CFG-E174	7	Pine Plantation	6	161	30 - 80	600	0 - 26	Y	Y	Y	Y



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E174**	7	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E175	102	Pine Plantation	11	10	30 - 80	1737	0 - 26	Y	Y	Y	Y
CFG-E175	102	Mesic Flatwoods	71	10	10 - 50	1737	0 - 0	Y	Y	Y	Y
CFG-E175**	102	Mesic Hammock	3	--	--	--	--	Y	Y	Y	Y
CFG-E178	122	Mesic Flatwoods	114	80	10 - 50	817	0 - 0	Y	Y	N	N
CFG-E178**	122	Mesic Hammock	4	--	--	--	--	Y	Y	Y	Y
CFG-E179	151	Mesic Flatwoods	131	25	10 - 50	528	0 - 0	Y	Y	N	N
CFG-E179	151	Mesic Hammock	13	25	10 - 30	528	0 - 79	Y	Y	N	N
CFG-E180	153	Mesic Flatwoods	143	58	10 - 50	302	0 - 0	Y	Y	N	N
CFG-E180**	153	Wet Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-E181	145	Mesic Flatwoods	144	69	10 - 50	369	0 - 0	Y	Y	N	N
CFG-E182	299	Wet Flatwoods	45	16	10 - 50	1091	0 - 0	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E182	299	Mesic Flatwoods	60	16	10 - 50	1091	0 - 0	Y	Y	N	N
CFG-E182	299	Xeric Hammock	51	16	10 - 30	1091	0 - 79	Y	Y	N	N
CFG-E182**	299	Mesic Hammock	4	--	--	--	--	Y	Y	Y	Y
CFG-E182**	299	Scrub	<1	--	--	--	--	Y	Y	Y	Y
CFG-E183	61	Mesic Hammock	19	24	10 - 30	678	0 - 79	Y	Y	N	N
CFG-E183	61	Wet Flatwoods	37	24	10 - 50	678	0 - 0	Y	Y	N	N
CFG-E183**	61	Scrub	1	--	--	--	--	Y	Y	Y	Y
CFG-E184	16	Scrub	16	17	0 - 20	966	0 - 13	Y	Y	N	N
CFG-E184**	16	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-E184**	16	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E184**	16	Xeric Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-E185	43	Scrub	42	27	0 - 20	1125	0 - 13	Y	Y	N	N
CFG-E185**	43	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-E185**	43	Xeric Hammock	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E185**	43	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E186	79	Mesic Flatwoods	39	60	10 - 50	1755	0 - 0	Y	Y	N	N
CFG-E186	79	Xeric Hammock	23	60	10 - 30	1755	0 - 79	Y	Y	N	N
CFG-E186**	79	Scrub	<1	--	--	--	--	Y	Y	Y	Y
CFG-E186**	79	Mesic Hammock	2	--	--	--	--	Y	Y	Y	Y
CFG-E187	29	Scrub	28	53	0 - 20	590	0 - 13	Y	Y	N	N
CFG-E187**	29	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E187**	29	Xeric Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-E187**	29	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-E188	164	Mesic Flatwoods	24	73	10 - 50	1696	0 - 0	Y	Y	N	N
CFG-E188**	164	Scrub	<1	--	--	--	--	Y	Y	Y	Y
CFG-E189	50	Scrub	49	43	0 - 20	704	0 - 13	Y	Y	N	N
CFG-E190	53	Scrub	45	20	0 - 20	5100	0 - 13	Y	Y	N	N
CFG-E191	37	Wet Flatwoods	2	70	10 - 50	1363	0 - 0	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E191**	37	Scrubby Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E191**	37	Scrub	<1	--	--	--	--	Y	Y	Y	Y
CFG-E192	75	Wet Flatwoods	42	55	10 - 50	500	0 - 0	Y	Y	N	N
CFG-E192	75	Scrubby Flatwoods	8	55	20 - 60	500	0 - 26	Y	Y	N	N
CFG-E193**	27	Wet Flatwoods	19	--	--	--	--	Y	Y	Y	Y
CFG-E194**	74	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E195	110	Wet Flatwoods	90	90	10 - 50	636	0 - 0	Y	Y	N	N
CFG-E195**	110	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E196**	55	Pine Plantation	8	--	--	--	--	Y	Y	Y	Y
CFG-E197	51	Pine Plantation	48	55	30 - 80	810	0 - 26	Y	Y	Y	Y
CFG-E197**	51	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E198	14	Pine Plantation	12	87	30 - 80	1227	0 - 26	Y	Y	Y	Y
CFG-E199	210	Pine Plantation	191	15	30 - 80	500	0 - 26	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E199	210	Clearcut Pine Plantation	9	15	30 - 80	500	0 - 26	Y	Y	Y	Y
CFG-E199**	210	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E200	28	Pine Plantation	14	20	30 - 80	3400	0 - 26	Y	Y	Y	Y
CFG-E200**	28	Clearcut Pine Plantation	5	--	--	--	--	Y	Y	Y	Y
CFG-E201	51	Pine Plantation	43	80	30 - 80	4570	0 - 26	Y	Y	Y	Y
CFG-E202	181	Pine Plantation	93	7	30 - 80	3853	0 - 26	Y	Y	Y	Y
CFG-E203	19	Pine Plantation	19	5	30 - 80	1406	0 - 26	Y	Y	Y	Y
CFG-E204	128	Pine Plantation	87	60	30 - 80	1089	0 - 26	Y	Y	Y	Y
CFG-E204**	128	Mesic Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E205**	126	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E206**	88	Pine Plantation	3	--	--	--	--	Y	Y	Y	Y



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E206**	88	Mesic Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-E207**	150	Sandhill	22	--	--	--	--	Y	Y	Y	Y
CFG-E208	12	Sandhill	3	25	20 - 60	94	0 - 79	Y	Y	N	N
CFG-E208	12	Mesic Flatwoods	6	25	10 - 50	94	0 - 0	Y	Y	N	N
CFG-E209	88	Mesic Flatwoods	29	2	10 - 50	75	0 - 0	Y	Y	Y	Y
CFG-E209	88	Scrubby Flatwoods	24	2	20 - 60	75	0 - 26	Y	Y	Y	Y
CFG-E209	88	Sandhill	34	2	20 - 60	75	0 - 79	N	N	Y	Y
CFG-E210	96	Mesic Flatwoods	76	70	10 - 50	337	0 - 0	Y	Y	N	N
CFG-E211	75	Pine Plantation	5	25	30 - 80	811	0 - 26	Y	Y	Y	Y
CFG-E211	75	Mesic Flatwoods	64	25	10 - 50	811	0 - 0	Y	Y	N	N
CFG-E212	13	Pine Plantation	13	70	30 - 80	354	0 - 26	Y	Y	Y	Y
CFG-E213**	3	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E213**	3	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E214**	26	Pine Plantation	13	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E214**	26	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E215	173	Mesic Flatwoods	168	84	10 - 50	503	0 - 0	Y	Y	N	N
CFG-E215**	173	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E216	136	Mesic Flatwoods	129	89	10 - 50	380	0 - 0	Y	Y	N	N
CFG-E217**	129	Mesic Flatwoods	4	--	--	--	--	Y	Y	Y	Y
CFG-E218	35	Mesic Flatwoods	32	69	10 - 50	2072	0 - 0	Y	Y	N	N
CFG-E219	84	Mesic Flatwoods	82	63	10 - 50	1234	0 - 0	Y	Y	N	N
CFG-E220**	96	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E220**	96	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-E221**	271	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E221**	271	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-E221**	271	Mesic Flatwoods	2	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E222**	65	Mesic Flatwoods	3	--	--	--	--	Y	Y	Y	Y
CFG-E223	200	Mesic Flatwoods	176	138	10 - 50	543	0 - 0	Y	Y	N	N
CFG-E223**	200	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E224	157	Mesic Flatwoods	74	40	10 - 50	289	0 - 0	Y	Y	N	N
CFG-E224	157	Wet Flatwoods	45	40	10 - 50	289	0 - 0	Y	Y	N	N
CFG-E225	117	Mesic Flatwoods	112	61	10 - 50	281	0 - 0	Y	Y	N	N
CFG-E226**	30	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E227	61	Wet Flatwoods	4	120	10 - 50	6	0 - 0	Y	Y	N	N
CFG-E227**	61	Mesic Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-E229	40	Wet Flatwoods	4	80	10 - 50	74	0 - 0	Y	Y	N	N
CFG-E229**	40	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E230	52	Mesic Flatwoods	2	62	10 - 50	384	0 - 0	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E231	31	Mesic Flatwoods	29	52	10 - 50	1072	0 - 0	Y	Y	N	N
CFG-E232**	15	Mesic Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E233**	19	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E234	69	Mesic Flatwoods	56	120	10 - 50	900	0 - 0	Y	Y	N	N
CFG-E234	69	Pine Plantation	12	120	30 - 80	900	0 - 26	Y	Y	Y	Y
CFG-E235	183	Mesic Flatwoods	44	126	10 - 50	900	0 - 0	Y	Y	N	N
CFG-E235	183	Pine Plantation	16	126	30 - 80	900	0 - 26	Y	Y	Y	Y
CFG-E236	155	Mesic Flatwoods	21	80	10 - 50	1120	0 - 0	Y	Y	N	N
CFG-E237**	30	Mesic Flatwoods	7	--	--	--	--	Y	Y	Y	Y
CFG-E238	126	Mesic Flatwoods	117	87	10 - 50	1550	0 - 0	Y	Y	N	N
CFG-E239	10	Mesic Flatwoods	3	50	10 - 50	60	0 - 0	Y	Y	N	N
CFG-E240	36	Mesic Flatwoods	20	93	10 - 50	1131	0 - 0	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E241	2765	Mesic Flatwoods	7	20	10 - 50	307	0 - 0	Y	Y	N	N
CFG-E241**	2765	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E241**	2765	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-E242**	110	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-E242**	110	Xeric Hammock	2	--	--	--	--	Y	Y	Y	Y
CFG-E242**	110	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E243**	63	Upland Hardwood Forest	15	--	--	--	--	Y	Y	Y	Y
CFG-E244**	147	Upland Hardwood Forest	19	--	--	--	--	Y	Y	Y	Y
CFG-E245**	175	Upland Hardwood Forest	4	--	--	--	--	Y	Y	Y	Y
CFG-E246**	14	Upland Hardwood Forest	14	--	--	--	--	Y	Y	Y	Y



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E247**	37	Upland Hardwood Forest	<1	--	--	--	--	Y	Y	Y	Y
CFG-E248**	4	Upland Hardwood Forest	<1	--	--	--	--	Y	Y	Y	Y
CFG-E249**	84	Upland Hardwood Forest	11	--	--	--	--	Y	Y	Y	Y
CFG-E250**	59	Upland Hardwood Forest	8	--	--	--	--	Y	Y	Y	Y
CFG-E251**	64	Upland Hardwood Forest	<1	--	--	--	--	Y	Y	Y	Y
CFG-E252**	170	Upland Hardwood Forest	13	--	--	--	--	Y	Y	Y	Y
CFG-E255**	152	Upland Hardwood Forest	4	--	--	--	--	Y	Y	Y	Y
CFG-E257**	194	Upland Hardwood Forest	19	--	--	--	--	Y	Y	Y	Y
CFG-E259	18	Mesic Hammock	13	31	10 - 30	860	0 - 79	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E260**	29	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-E262**	13	Mesic Hammock	10	--	--	--	--	Y	Y	Y	Y
CFG-E264**	2	Scrub	2	--	--	--	--	Y	Y	Y	Y
CFG-E275	25	Mesic Hammock	10	40	10 - 30	294	0 - 79	Y	Y	N	N
CFG-E276**	6	Mesic Hammock	2	--	--	--	--	Y	Y	Y	Y
CFG-E277	14	Mesic Hammock	9	20	10 - 30	390	0 - 79	Y	Y	N	N
CFG-E278	28	Mesic Hammock	12	40	10 - 30	877	0 - 79	Y	Y	N	N
CFG-E279	340	Mesic Flatwoods	35	10	10 - 50	179	0 - 0	Y	Y	Y	Y
CFG-E279	340	Pine Plantation	28	10	30 - 80	179	0 - 26	Y	Y	Y	Y
CFG-E279	340	Scrubby Flatwoods	5	10	20 - 60	179	0 - 26	Y	Y	Y	Y
CFG-E279	340	Wet Flatwoods	60	10	10 - 50	179	0 - 0	Y	Y	Y	Y
CFG-E279**	340	Xeric Hammock	1	--	--	--	--	Y	Y	Y	Y
CFG-E280**	46	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E281**	630	Xeric Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-E283**	57	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E284**	2	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E285**	203	Xeric Hammock	13	--	--	--	--	Y	Y	Y	Y
CFG-E285**	203	Scrubby Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-E285**	203	Mesic Flatwoods	12	--	--	--	--	Y	Y	Y	Y
CFG-E285**	203	Mesic Hammock	5	--	--	--	--	Y	Y	Y	Y
CFG-E285**	203	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E286**	68	Mesic Flatwoods	4	--	--	--	--	Y	Y	Y	Y
CFG-E286**	68	Xeric Hammock	11	--	--	--	--	Y	Y	Y	Y
CFG-E286**	68	Scrubby Flatwoods	4	--	--	--	--	Y	Y	Y	Y
CFG-E287**	325	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E287**	325	Scrubby Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E288**	55	Mesic Flatwoods	39	--	--	--	--	Y	Y	Y	Y
CFG-E288**	55	Scrubby Flatwoods	3	--	--	--	--	Y	Y	Y	Y
CFG-E288**	55	Wet Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-E290**	8	Wet Flatwoods	7	--	--	--	--	Y	Y	Y	Y
CFG-E290**	8	Xeric Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-E291**	38	Upland Hardwood Forest	18	--	--	--	--	Y	Y	Y	Y
CFG-E291**	38	Mesic Hammock	2	--	--	--	--	Y	Y	Y	Y
CFG-E291**	38	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E291**	38	Xeric Hammock	9	--	--	--	--	Y	Y	Y	Y
CFG-E292**	579	Xeric Hammock	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E292**	579	Upland Hardwood Forest	<1	--	--	--	--	Y	Y	Y	Y
CFG-E292**	579	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-E293**	32	Mesic Hammock	8	--	--	--	--	Y	Y	Y	Y
CFG-E295**	57	Mesic Flatwoods	14	--	--	--	--	Y	Y	Y	Y
CFG-E296**	24	Mesic Flatwoods	6	--	--	--	--	Y	Y	Y	Y
CFG-E297**	13	Mesic Flatwoods	1	--	--	--	--	Y	Y	Y	Y
CFG-E298**	534	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E298**	534	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E300	54	Mesic Flatwoods	27	37	10 - 50	2276	0 - 0	Y	Y	N	N
CFG-E300**	54	Pine Plantation	2	--	--	--	--	Y	Y	Y	Y
CFG-E301	55	Pine Plantation	45	40	30 - 80	1850	0 - 26	Y	Y	Y	Y
CFG-E301**	55	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E302**	47	Pine Plantation	2	--	--	--	--	Y	Y	Y	Y
CFG-E303**	69	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-E303**	69	Xeric Hammock	1	--	--	--	--	Y	Y	Y	Y
CFG-E304**	1	Pine Plantation	1	--	--	--	--	Y	Y	Y	Y
CFG-E305**	4	Pine Plantation	4	--	--	--	--	Y	Y	Y	Y
CFG-E306	50	Pine Plantation	11	22	30 - 80	402	0 - 26	Y	Y	Y	Y
CFG-E306	50	Xeric Hammock	34	22	10 - 30	402	0 - 79	Y	Y	N	N
CFG-E307	109	Pine Plantation	12	80	30 - 80	2519	0 - 26	Y	Y	Y	Y
CFG-E307	109	Xeric Hammock	1	80	10 - 30	2519	0 - 79	Y	Y	N	N
CFG-E307	109	Wet Flatwoods	37	80	10 - 50	2519	0 - 0	Y	Y	N	N
CFG-E308	143	Pine Plantation	2	20	30 - 80	157	0 - 26	Y	Y	Y	Y
CFG-E308	143	Xeric Hammock	1	20	10 - 30	157	0 - 79	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-E308**	143	Mesic Flatwoods	15	--	--	--	--	Y	Y	Y	Y
CFG-E308**	143	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-E309**	141	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-W001	51	Mesic Hammock	19	30	10 - 30	303	0 - 79	Y	Y	N	N
CFG-W001	51	Sandhill	30	30	20 - 60	303	0 - 79	Y	Y	N	N
CFG-W002**	15	Mesic Hammock	8	--	--	--	--	Y	Y	Y	Y
CFG-W003	120	Mesic Hammock	42	20	10 - 30	27	0 - 79	N	N	N	N
CFG-W003	120	Sandhill	53	20	20 - 60	27	0 - 79	N	N	N	N
CFG-W004	202	Mesic Hammock	87	5	10 - 30	548	0 - 79	Y	Y	Y	Y
CFG-W004	202	Sandhill	46	5	20 - 60	548	0 - 79	Y	Y	Y	Y
CFG-W005	99	Mesic Hammock	8	10	10 - 30	376	0 - 79	Y	Y	Y	Y
CFG-W005	99	Sandhill	91	10	20 - 60	376	0 - 79	Y	Y	Y	Y
CFG-W006	190	Mesic Hammock	17	28	10 - 30	169	0 - 79	Y	Y	N	N
CFG-W006	190	Sandhill	172	28	20 - 60	169	0 - 79	Y	Y	N	N

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-W007	293	Mesic Hammock	157	5	10 - 30	391	0 - 79	Y	Y	Y	Y
CFG-W007	293	Sandhill	129	5	20 - 60	391	0 - 79	Y	Y	Y	Y
CFG-W008	299	Mesic Hammock	227	8	10 - 30	173	0 - 79	Y	Y	Y	Y
CFG-W008	299	Sandhill	39	8	20 - 60	173	0 - 79	Y	Y	Y	Y
CFG-W008**	299	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-W009	65	Sandhill	57	14	20 - 60	51	0 - 79	N	N	Y	Y
CFG-W009	65	Mesic Hammock	7	14	10 - 30	51	0 - 79	N	N	N	N
CFG-W010**	24	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-W010**	24	Mesic Hammock	1	--	--	--	--	Y	Y	Y	Y
CFG-W011	90	Sandhill	13	5	20 - 60	308	0 - 79	Y	Y	Y	Y
CFG-W011	90	Mesic Hammock	72	5	10 - 30	308	0 - 79	Y	Y	Y	Y
CFG-W011**	90	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-W012	55	Pine Plantation	55	49	30 - 80	10	0 - 26	N	N	Y	Y
CFG-W012**	55	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-W013	90	Pine Plantation	90	51	30 - 80	17	0 - 26	N	N	Y	Y
CFG-W013**	90	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-W014**	15	Pine Plantation	1	--	--	--	--	Y	Y	Y	Y
CFG-W014**	15	Mesic Hammock	13	--	--	--	--	Y	Y	Y	Y
CFG-W015	63	Pine Plantation	61	37	30 - 80	54	0 - 26	Y	Y	Y	Y
CFG-W015**	63	Mesic Hammock	1	--	--	--	--	Y	Y	Y	Y
CFG-W016	71	Sandhill	11	15	20 - 60	2	0 - 79	N	N	Y	Y
CFG-W016	71	Mesic Hammock	58	15	10 - 30	2	0 - 79	N	N	N	N
CFG-W016**	71	Pine Plantation	1	--	--	--	--	Y	Y	Y	Y
CFG-W017	95	Mesic Hammock	76	10	10 - 30	556	0 - 79	Y	Y	N	N
CFG-W017**	95	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-W017**	95	Pine Plantation	1	--	--	--	--	Y	Y	Y	Y
CFG-W018	147	Mesic Hammock	124	10	10 - 30	172	0 - 79	Y	Y	Y	Y
CFG-W018	147	Sandhill	5	10	20 - 60	172	0 - 79	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-W018**	147	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-W019**	12	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-W019**	12	Pine Plantation	12	--	--	--	--	Y	Y	Y	Y
CFG-W020	67	Pine Plantation	62	35	30 - 80	40	0 - 26	Y	Y	Y	Y
CFG-W020**	67	Mesic Hammock	4	--	--	--	--	Y	Y	Y	Y
CFG-W021	151	Pine Plantation	2	27	30 - 80	228	0 - 26	Y	Y	Y	Y
CFG-W021	151	Mesic Hammock	82	27	10 - 30	228	0 - 79	Y	Y	N	N
CFG-W021	151	Sandhill	66	27	20 - 60	228	0 - 79	Y	Y	N	N
CFG-W022**	10	Mesic Hammock	2	--	--	--	--	Y	Y	Y	Y
CFG-W023	39	Mesic Hammock	2	55	10 - 30	360	0 - 79	Y	Y	N	N
CFG-W023	39	Pine Plantation	37	55	30 - 80	360	0 - 26	Y	Y	Y	Y
CFG-W024	147	Mesic Hammock	106	0	10 - 30	2	0 - 79	N	N	Y	Y
CFG-W024**	147	Pine Plantation	1	--	--	--	--	Y	Y	Y	Y



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-W025	48	Mesic Hammock	28	15	10 - 30	1154	0 - 79	Y	Y	N	N
CFG-W025**	48	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-W026	67	Mesic Hammock	62	1	10 - 30	1020	0 - 79	Y	Y	Y	Y
CFG-W026**	67	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-W027**	87	Mesic Hammock	20	--	--	--	--	Y	Y	Y	Y
CFG-W028**	107	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-W030**	262	Wet Flatwoods	3	--	--	--	--	Y	Y	Y	Y
CFG-W030**	262	Mesic Hammock	3	--	--	--	--	Y	Y	Y	Y
CFG-W031**	68	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-W031**	68	Pine Plantation	1	--	--	--	--	Y	Y	Y	Y
CFG-W032	56	Mesic Hammock	48	6	10 - 30	317	0 - 79	Y	Y	Y	Y
CFG-W032**	56	Pine Plantation	2	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-W033	153	Pine Plantation	153	76	30 - 80	8	0 - 26	N	N	Y	Y
CFG-W033**	153	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-W034	19	Mesic Hammock	17	2	10 - 30	773	0 - 79	Y	Y	Y	Y
CFG-W034**	19	Pine Plantation	1	--	--	--	--	Y	Y	Y	Y
CFG-W035	89	Pine Plantation	89	44	30 - 80	516	0 - 26	Y	Y	Y	Y
CFG-W035**	89	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-W036	45	Sandhill	44	5	20 - 60	649	0 - 79	Y	Y	Y	Y
CFG-W036**	45	Pine Plantation	1	--	--	--	--	Y	Y	Y	Y
CFG-W037**	24	Pine Plantation	24	--	--	--	--	Y	Y	Y	Y
CFG-W038**	0	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-W039	91	Pine Plantation	91	29	30 - 80	270	0 - 26	Y	Y	Y	Y
CFG-W040**	20	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y
CFG-W040**	20	Mesic Hammock	1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-W041	28	Mesic Hammock	27	0	10 - 30	633	0 - 79	Y	Y	Y	Y
CFG-W042**	9	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-W043	66	Mesic Hammock	50	32	10 - 30	817	0 - 79	Y	Y	N	N
CFG-W044	67	Mesic Hammock	38	3	10 - 30	312	0 - 79	Y	Y	Y	Y
CFG-W044	67	Pine Plantation	13	3	30 - 80	312	0 - 26	Y	Y	Y	Y
CFG-W045**	8	Mesic Hammock	2	--	--	--	--	Y	Y	Y	Y
CFG-W046**	6	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-W047**	28	Mesic Hammock	6	--	--	--	--	Y	Y	Y	Y
CFG-W055	68	Mesic Hammock	61	10	10 - 30	937	0 - 79	Y	Y	Y	Y
CFG-W056	55	Mesic Hammock	17	3	10 - 30	309	0 - 79	Y	Y	Y	Y
CFG-W059**	3	Sandhill	3	--	--	--	--	Y	Y	Y	Y
CFG-W060	92	Sandhill	66	20	20 - 60	981	0 - 79	Y	Y	N	N
CFG-W060	92	Mesic Hammock	15	20	10 - 30	981	0 - 79	Y	Y	N	N
CFG-W061	19	Sandhill	18	6	20 - 60	1919	0 - 79	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-W061**	19	Mesic Hammock	1	--	--	--	--	Y	Y	Y	Y
CFG-W069	4	Xeric Hammock	4	10	10 - 30	45	0 - 79	N	N	Y	Y
CFG-W070**	1	Xeric Hammock	1	--	--	--	--	Y	Y	Y	Y
CFG-W071**	9	Xeric Hammock	9	--	--	--	--	Y	Y	Y	Y
CFG-W072**	0	Xeric Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-W073**	0	Xeric Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-W077	59	Wet Flatwoods	55	20	10 - 50	976	0 - 0	Y	Y	N	N
CFG-W078	70	Wet Flatwoods	65	51	10 - 50	671	0 - 0	Y	Y	N	N
CFG-W079	3155	Wet Flatwoods	7	80	10 - 50	763	0 - 0	Y	Y	N	N
CFG-W080	60	Wet Flatwoods	28	26	10 - 50	739	0 - 0	Y	Y	N	N
CFG-W081**	15	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-W082**	12	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-W083	62	Mesic Hammock	57	44	10 - 30	1200	0 - 79	Y	Y	N	N
CFG-W083**	62	Sandhill	1	--	--	--	--	Y	Y	Y	Y
CFG-W084	27	Sandhill	9	36	20 - 60	579	0 - 79	Y	Y	N	N
CFG-W084**	27	Mesic Hammock	12	--	--	--	--	Y	Y	Y	Y
CFG-W085	40	Mesic Hammock	7	3	10 - 30	659	0 - 79	Y	Y	Y	Y
CFG-W086**	12	Mesic Hammock	3	--	--	--	--	Y	Y	Y	Y
CFG-W087	155	Sandhill	12	10	20 - 60	490	0 - 79	Y	Y	Y	Y
CFG-W087	155	Mesic Hammock	86	10	10 - 30	490	0 - 79	Y	Y	Y	Y
CFG-W087	155	Mesic Flatwoods	49	10	10 - 50	490	0 - 0	Y	Y	Y	Y
CFG-W088**	39	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-W089	20	Mesic Hammock	6	60	10 - 30	562	0 - 79	Y	Y	N	N
CFG-W089**	20	Mesic Flatwoods	11	--	--	--	--	Y	Y	Y	Y
CFG-W089**	20	Sandhill	<1	--	--	--	--	Y	Y	Y	Y
CFG-W090	16	Mesic Hammock	1	10	10 - 30	1366	0 - 79	Y	Y	Y	Y



Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-W090**	16	Mesic Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-W092	71	Mesic Flatwoods	30	10	10 - 50	229	0 - 0	Y	Y	Y	Y
CFG-W093	33	Mesic Flatwoods	6	40	10 - 50	1	0 - 0	Y	Y	N	N
CFG-W093	33	Mesic Hammock	24	40	10 - 30	1	0 - 79	Y	N	N	N
CFG-W094	135	Mesic Flatwoods	73	2	10 - 50	825	0 - 0	Y	Y	Y	Y
CFG-W094**	135	Wet Flatwoods	3	--	--	--	--	Y	Y	Y	Y
CFG-W095	100	Wet Flatwoods	17	15	10 - 50	293	0 - 0	Y	Y	N	N
CFG-W095	100	Mesic Flatwoods	10	15	10 - 50	293	0 - 0	Y	Y	N	N
CFG-W095	100	Mesic Hammock	53	15	10 - 30	293	0 - 79	Y	Y	N	N
CFG-W096	158	Wet Flatwoods	49	100	10 - 50	394	0 - 0	Y	Y	N	N
CFG-W096	158	Mesic Hammock	4	100	10 - 30	394	0 - 79	Y	Y	N	N
CFG-W106**	87	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-W107	104	Mesic Hammock	49	2	10 - 30	700	0 - 79	Y	Y	Y	Y
CFG-W107	104	Wet Flatwoods	2	2	10 - 50	700	0 - 0	Y	Y	Y	Y
CFG-W108	107	Mesic Flatwoods	24	2	10 - 50	1104	0 - 0	Y	Y	Y	Y
CFG-W109	68	Sandhill	3	10	20 - 60	758	0 - 79	Y	Y	Y	Y
CFG-W109**	68	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-W109**	68	Scrubby Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-W110	43	Sandhill	34	4	20 - 60	583	0 - 79	Y	Y	Y	Y
CFG-W111	27	Scrubby Flatwoods	12	6	20 - 60	1097	0 - 26	Y	Y	Y	Y
CFG-W116**	19	Mesic Hammock	12	--	--	--	--	Y	Y	Y	Y
CFG-W121**	112	Mesic Hammock	33	--	--	--	--	Y	Y	Y	Y
CFG-W122**	97	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y
CFG-W123	133	Mesic Flatwoods	8	10	10 - 50	860	0 - 0	Y	Y	Y	Y
CFG-W123**	133	Pine Plantation	<1	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-W123**	133	Wet Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-W124	553	Mesic Hammock	11	6	10 - 30	1041	0 - 79	Y	Y	Y	Y
CFG-W124	553	Wet Flatwoods	152	6	10 - 50	1041	0 - 0	Y	Y	Y	Y
CFG-W124	553	Pine Plantation	98	6	30 - 80	1041	0 - 26	Y	Y	Y	Y
CFG-W124**	553	Mesic Flatwoods	2	--	--	--	--	Y	Y	Y	Y
CFG-W125	126	Pine Plantation	58	5	30 - 80	2300	0 - 26	Y	Y	Y	Y
CFG-W126	1831	Mesic Flatwoods	48	4	10 - 50	1583	0 - 0	Y	Y	Y	Y
CFG-W126	1831	Mesic Hammock	275	4	10 - 30	1583	0 - 79	Y	Y	Y	Y
CFG-W126	1831	Pine Plantation	70	4	30 - 80	1583	0 - 26	Y	Y	Y	Y
CFG-W127	42	Scrubby Flatwoods	42	4	20 - 60	1071	0 - 26	Y	Y	Y	Y
CFG-W128	46	Scrubby Flatwoods	46	8	20 - 60	760	0 - 26	Y	Y	Y	Y
CFG-W129	13	Scrubby Flatwoods	12	10	20 - 60	379	0 - 26	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-W129**	13	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-W130	83	Scrubby Flatwoods	50	2	20 - 60	394	0 - 26	Y	Y	Y	Y
CFG-W130**	83	Mesic Flatwoods	7	--	--	--	--	Y	Y	Y	Y
CFG-W131**	31	Scrubby Flatwoods	24	--	--	--	--	Y	Y	Y	Y
CFG-W132	84	Scrubby Flatwoods	48	20	20 - 60	500	0 - 26	Y	Y	Y	Y
CFG-W132	84	Mesic Flatwoods	7	20	10 - 50	500	0 - 0	Y	Y	N	N
CFG-W133	75	Mesic Hammock	6	5	10 - 30	381	0 - 79	Y	Y	Y	Y
CFG-W133	75	Scrubby Flatwoods	31	5	20 - 60	381	0 - 26	Y	Y	Y	Y
CFG-W134**	15	Scrubby Flatwoods	10	--	--	--	--	Y	Y	Y	Y
CFG-W135	1161	Scrubby Flatwoods	24	17	20 - 60	196	0 - 26	Y	Y	Y	Y
CFG-W135	1161	Wet Flatwoods	8	17	10 - 50	196	0 - 0	Y	Y	N	N
CFG-W135**	1161	Mesic Hammock	28	--	--	--	--	Y	Y	Y	Y

Florida State Parks  
Timber Management Analyses

Management Zones (MZ)	MZ (acres)	Candidate NatCom Type	Candidate NatCom (acres)	Current Average Overstory Pine BA (ft <sup>2</sup> /AC)	Target Overstory Pine BA (ft <sup>2</sup> /AC)	Current Non-Pine Overstory TPA	Target Non-Pine Overstory TPA	Potential Actions/Treatments			
								Harvest or Thin	Stand Improvement*	Site Prep	Plant
CFG-W135**	1161	Mesic Flatwoods	<1	--	--	--	--	Y	Y	Y	Y
CFG-W138**	195	Mesic Hammock	4	--	--	--	--	Y	Y	Y	Y
CFG-W139**	2272	Mesic Hammock	<1	--	--	--	--	Y	Y	Y	Y

\*Stand improvement, per Section 3 above, includes palmetto/midstory reduction. While inventory data was not used to estimate this metric, remotely sensed images and on-site observations have indicated that the selected areas could benefit from such treatments.

\*\*Unsampled upland areas are present in this analysis and could require vegetation management in the future.





**ADDENDUM 5:**  
**ADVISORY GROUP MEMBERS AND SUMMARY REPORT**  
**(Placeholder pending report completion)**



**ADDENDUM 6:  
MANATEE MANAGEMENT**





# Buckman Lock

## Lock Operation Procedures for Manatee Protection

Updated 3/27/12


**General:** All Lock Operators must be thoroughly familiar with the Manatee Protection System (MPS) and demonstrate the ability to safely perform locking operations during a training period before being allowed to operate Buckman Lock independent of supervision.

Although manatees frequent the lock area more often during the warmer months, they may be in the area at any time of the year. Constant vigilance and awareness of manatee movements should be maintained while on duty.

### Manatee Protection System Operation

- (1) Prior to entering the machinery house of the gates to be operated, observe the water upstream and downstream for manatee presence for several minutes. Advise boaters of their location if possible.
- (2) Before operating any gate the operator must accomplish a **system self test** of the MPS. This tests all circuits and lets the operator know that the MPS is functioning correctly.
- (3) If the system indicates a Passed condition on the PEC's (2) and Pecop (1) the operator may continue and open the gates. The gates are electronically preset to a 45% opening and the operator should visually insure that the gates do not exceed the 45% opening. Gates may be opened at a normal speed.
- (4) If the MPS indicates a failed system test the Lock operation will be halted until the problem is solved and a Pass system test is indicated.
- (5) Notify boaters of the presence of manatees and the need to maintain idle speed and steer clear.
- (6) **Gate Closing:** Closing the miter gates is **always** accomplished at **slow speed**. While gates are closing observe area around and to each side of gate for manatees.
- (7) **Rejected Gate Closing:** If the MPS alerts the operator to a rejected condition the operator must see that the gate moves back toward the open position and note the percent gate open at which the system rejected it.

- a. Most of the alerts will come from the lower sensors on the bottom of the gate (PEC's) which are pressure sensors and can be activated by debris, heavy vegetation, ect.
  
- (8)** After the gate has returned to the open position the operator will go out on the gate and visually check the immediate area for evidence of a manatee. If nothing is sighted the operator may try to close the gate again. If the gate is rejected at the same % of the gate opening the above procedure should be repeated. On the third try a similar rejection indicates that the trouble is debris or possibly a bad sensor. The operator can then manually bypass that specific opening and reactivate the PEC immediately after, to continue protection as the gate closes.
  - a. **For example:** If the gate rejects at 39% closed for the third time, go to bypass and start closing. At 35% closed reactivate the PEC and continue to close with the protection. This will usually solve the problem.
  
- (9)** When manatees are in the Lock with boaters, advise the boaters and allow manatees to exit the lock first. Then advise the boaters of the manatee's location and direct them to pass at a safe distance.
  
- (10)** If the Lock tender must leave the Lock and there is a manatee inside the Lock chamber, (such as the end of the shift) the upstream gates must be left open at the 45% opening to allow the manatee to exit safely.



**PECOP MANATEE PROTECTION SYSTEM FOR NAVIGATION LOCKS**  
**Manatee Protection Systems Development Team**



**US Army Corps  
of Engineers®**



**Government and Industry Working Together**

**Larry Taylor**  
Electrical/Electronic Engineering Production Manager  
**Harbor Branch Oceanographic Institution, Inc**  
5600 U.S. 1 North  
Fort Pierce, Florida 24946  
561-465-2400



## **PECOP MANATEE PROTECTION SYSTEM FOR NAVIGATION LOCKS**

### **HBOI Manatee Protection System Development Timeline**



#### **PED (Piezo-Electric-Detector) for Vertical Lift Gates**

1996 - S-26 in Miami - First Prototype - SFWMD

1997 - S-29 Miami - SFWMD

1998 - S-27 Miami, S20F Homestead - SFWMD  
S-25B Miami, USACE

1999 - IDT Contract Bid for 13 Structures First Advertisement

2000 - S-28 Miami, G-36 Okeechobee - SFWMD

#### **PECOP (Piezo-Electric-CO-Polymer) Non-Contact Sensor for Navigation Locks**

1997 - Proof of Application - October

Critical Design Review - December

1998 - St. Lucie Lock- Upper Gates


1999 - St. Lucie - Lower Gates

2000 - Port Canaveral - On-Line March 2000.



## PECOP MANATEE PROTECTION SYSTEM FOR NAVIGATION LOCKS

### Vertical Lift Gate Hazard to Manatees




**Gates move at 6" per Minute, Gates Remotely Controlled, Gate openings vary**  
**Manatees approach from both directions, Flow high or low**



PECOP MANATEE PROTECTION SYSTEM FOR NAVIGATION LOCKS

**PED Bumper Design Features**



- **Ruggedness**
  - Solid monolithic block of tough, polyurethane
  - Sensor has no moving parts
- **Engineered Shape**
  - Sensor Protected from debris impact
  - Highly Sensitive to “normal” axis
  - Low sensitivity off-axis
- **Mechanical Distortion**
  - Sensor nulls-out stresses
    - Sensor Adapts to distortion without recalibration or adjustment
    - Depth, biofouling do not affect sensitivity
- Each Sensor has individual twisted pair conductors
- **Self-Test Feature**



## PECOP MANATEE PROTECTION SYSTEM FOR NAVIGATION LOCKS


### PECOP Sensor System Design Goals



- Detect the presence of a submerged manatee that is in the path of the closing gates.
- Intervene in the gate control system to automatically stop the gate and then partially open the gate to allow free passage of the manatee.
- System should require little or no maintenance
- System must be protected from impact
- System must be resistant to corrosion and biofouling
- System must be adaptable to eight locks with minimal changes :  
Saint Lucie, Port Canaveral, Port Mayaca, W.P. Franklin, Ortona, Moore Haven, Okeechobee, Clewiston.
- System must have a "BYPASS" Mode

## PECOP MANATEE PROTECTION SYSTEM FOR NAVIGATION LOCKS

### Non-Contact Blocked Beam Sensor



**Low Power, High Frequency, Highly Directional Array of Acoustic Beams.  
Sensor Activated by blocking path between transmitter and receiver  
Gates will Automatically Stop and Allow Manatee to Pass Unharmed**



## PECOP MANATEE PROTECTION SYSTEM FOR NAVIGATION LOCKS

### Sensor Technology



PECOP is an acronym for Piezo Electric COPolymer.

A piezo-electric material changes dimensions when an electric potential is applied, and it generates electric charge when it is mechanically deformed.

The PECOP Sensor elements are 3” long copolymer (plastic) cylinders.

One set serves as the transmitting “loudspeakers” and another set serves as the receivers or “ear pieces” .

Transmit Frequency: 700,000 Hz to 1,000,000 Hz

- 1) This frequency is above the hearing range of manatees.
- 2) The high frequency waves are rapidly attenuated in the water  
( echoes dampen-out quickly).




# PECOP MANATEE PROTECTION SYSTEM FOR NAVIGATION LOCKS

## Highly Directional Sound Waves




Sensor Element Vertical Spacing - 6" Center to Center.  
Minimum Height - 12" Above Highest Recorded Water Level  
Maximum Sensor Height - 20'  
Standardized - 5' Cartridge (10' and 5' St. Lucie)








# PECOP MANATEE PROTECTION SYSTEM FOR NAVIGATION LOCKS Port Canaveral PECOP Sensor Mounting Locations





# PECOP MANATEE PROTECTION SYSTEM FOR NAVIGATION LOCKS

## Dual Gate Operation




BOTH TRANSMITTERS COUPLE  
TO BOTH RECEIVERS UNTIL ALMOST CLOSED,  
THEN DIRECT SKIN-TO-SKIN AND  
TIMBER-TO-TIMBER COUPLING OCCURS



# PECOP MANATEE PROTECTION SYSTEM FOR NAVIGATION LOCKS


## Manatee Protection Zone Geometry





## PECOP MANATEE PROTECTION SYSTEM FOR NAVIGATION LOCKS

### Detection Criteria



#### **Criteria #1: Amplitude**

Alarm Determined by Threshold Setting in Receiver Module.

Manatee Blocks Beam - Received Signal drops to zero.

#### **Criteria #2: Time Of Blockage**


Programmable Time Response to eliminate triggering on fast moving targets

#### **Criteria #3: Number of Blocked Beams**

Adaptive dual/single beam detection



# PECOP MANATEE PROTECTION SYSTEM FOR NAVIGATION LOCKS OPERATOR'S DISPLAY - "BEAMS" SCREEN




Record of Detection and Intervention Cycle - St. Lucie Lock Feb. 2, 1999






# PECOP MANATEE PROTECTION SYSTEM FOR NAVIGATION LOCKS

## St. Lucie Lock MPS System Configuration





**PECOP MANATEE PROTECTION SYSTEM FOR NAVIGATION LOCKS**  
**Port Canaveral Lock MPS Configuration**





## **PECOP MANATEE PROTECTION SYSTEM FOR NAVIGATION LOCKS**

**Target Design for St. Johns Lock**



- **Keep Cost Down**
- **No Data Logger Required (Can be added Later)**
- **Automatic Intervention Optional**

**ADDENDUM 7:  
LAND MANAGEMENT REVIEW**







# FLORIDA DEPARTMENT OF ENVIRONMENTAL PROTECTION

MARJORY STONEMAN DOUGLAS BUILDING  
3900 COMMONWEALTH BOULEVARD  
TALLAHASSEE, FLORIDA 32399-3000

RICK SCOTT  
GOVERNOR

CARLOS LOPEZ-CANTERA  
LT. GOVERNOR

JONATHAN P. STEVERSON  
INTERIM SECRETARY

## MEMORANDUM

TO: Aric Larson, Governmental Operations Consultant III  
Division of State Lands

FROM: Parks Small, Chief, Bureau of Natural and Cultural Resources  
Division of Recreation and Parks

Lew Scruggs, Chief, Office of Park Planning *LPS*  
Division of Recreation and Parks

SUBJECT: Response to Draft Land Management Review (LMR)  
Marjorie Harris Carr Cross Florida Greenway (Eastern Portion)

DATE: May 26, 2015

---

The Land Management Review draft report provided to DRP determined that management of Marjorie Harris Carr Cross Florida Greenway (Eastern Portion) by the Division of Recreation and Parks met the two tests prescribed by law. Namely, the review team concluded that the land is being managed for the purposes for which it was acquired and in accordance with the land management plan.

Below are Additional Recommendations and Checklist Findings (items the LMR determined should be further addressed in the Management Plan update) of the draft LMR report, with our Manager's Response to each. The responses were prepared via a coordinated effort of the park, district office, and our offices.

### CONSENSUS RECOMMENDATIONS

- The team recommends that DRP seek additional FTE positions and additional funding commensurate with the unique management needs associated with this property (7+, 0-)**  
*Managing Agency Response: Agree. If it is determined that additional staff are needed at the time of the next unit management plan revision, it will be included in the plan. However, no new staff can be assigned to this or any other park unit unless they are appropriated by the Legislature or reassigned from other units. Funding is determined annually by the Florida Legislature. The updated unit management plan will address land management funding needs. However, Division funding is determined annually by the Florida Legislature and funds are allocated to the 171 state parks and trails according to priority needs.*
- The team recommends that DRP re-visit efforts to transfer management of water control structures to a more appropriate agency such as the St. Johns River Water Management District. (7+, 0-)**  
*Managing Agency Response: Agree. Ultimate responsibility for the assignment of management and operations of the former Cross Florida Barge Canal (CFBC) project works is determined by the*

*department working with the legislative, executive branches and respective water management districts.*

- 3. The team recommends that DRP and Greenway staff continue efforts to seek funding for contract burning and staff enhancements necessary to achieve increased fire frequency and acreage burned. (7+, 0-)**

*Managing Agency Response: Agree. DRP continues to seek funds for contract funding for prescribed fire. Currently, DRP is working on securing funding for a 1,200 acre contract for FY 2015-2016 with our existing fire contractor that we have used for the past two years. We are also involved in a pilot timber sale project that would provide contract burning services in lieu of timber revenue. We will continue to pursue viable opportunities to increase annual prescribed fire acreages to fullest extent possible.*

- 4. The team recommends that DRP seek funding or additional release of timber sale revenues necessary to implement reforestation goals on extensive areas of cutover flatwoods. (7+, 0-)**

*Managing Agency Response: Agree. The pilot timber sale project mentioned above would also call for planting services in lieu of timber revenue.*

- 5. The team recommends that DRP/Greenways staff continue efforts to coordinate with DHR to prioritize archaeological/cultural sites for annual monitoring. (7+, 0-)**

*Managing Agency Response: Agree. DRP staff will increase coordination and training with the DHR to monitor priority sites and increase the number of staff who attend the revised Archaeological Resource Management (ARM) trainings provided by DHR.*

#### FIELD REVIEW

- 1. The maintenance condition of the Natural Communities, specifically sandhill, scrub and wet flatwoods, received below average scores. The review team is asked to evaluate, based on their perspective, what percent of the natural community is in maintenance condition. The scores range from 1 to 5, with 1 being 0-20% in maintenance condition, 2 being 21-40%, 3 being 41-60%, 4 being 61-80% and 5 being 81-100%.**

*Managing Agency Response: Agree. DRP is in the process of restoring many areas of the CFG. On the east end of the CFG, the quality of the sandhill is not as high as that on the west end because much of the sandhill in the Marion County area has been disturbed and will require significant funding efforts to restore. Many areas have been replanted with longleaf pine and in some areas with wiregrass plugs. The sandhill in Putnam County occurs in small parcels and is usually included in the larger natural communities which surround it. Sandhill restoration is in process in these areas via removing oaks and prescribed fire.*

*DRP has developed a scrub restoration plan for the scrub habitat within the CFG. A significant portion of the scrub is already planned to be timbered by a Florida Forest Service contractor. The wet flatwoods on the east end has good structure for the most part. Many of the management zones are also in need of a second thinning treatment in order to decrease the basal area and encourage greater understory diversity. The pilot timber sale project is comprised of second timber thinnings in much of this natural community.*

- 2. Cultural Resources (Archeological & Historic sites), specifically protection and preservation, received a below average score. The review team is asked to evaluate, based information provided by the managing agency, whether the protection and preservation of cultural resources is sufficient.**

**Managing Agency Response:** *Agree. DRP will continue coordination and training with the DHR to monitor priority sites and increase the number of CFG staff who attend the revised ARM trainings provided by DHR.*

3. **Restoration, specifically deep creek scrub, received a below average score. The review team is asked to evaluate, based on their perspective, whether restoration efforts are adequate.**  
**Managing Agency Response:** *Agree. As mentioned in a previous response, the restoration of the Deep Creek scrub area will benefit a large suite of species. This scrub restoration project was initiated through a contract with FNAI to perform a rare plant inventory which was completed in October 2014. The next stage is a timber sale which will remove sand pine and large oaks in this area. Once the site has a chance for the scrub community to regrow, DRP will manage this area through a combination of mechanical treatments and prescribed fire. Other restoration projects on the east end include pine thinning in natural flatwoods to aid in prescribed fire, to provide the structure and density recommended for flatwoods, and to reduce fuel buildup.*
4. **Resource Management, Prescribed Fire, specifically area being burned and frequency received below average scores. The review team is asked to evaluate, based on information provided by the managing agency, to what degree prescribed fire is accomplished according to the objectives for prescribed fire management. The scores range from 1 to 5, with 1 being 0-20% accomplished, 2 being 21-40%, 3 being 41-60%, 4 being 61-80% and 5 being 81-100%.**  
**Managing Agency Response:** *Agree. DRP's ecological prescribed fire goal for the east end of the Greenway is approximately 5,000 acres per year. The largest acreage in a given year burned was approximately 2,500 acres with the assistance from contract burning. Current staffing and budget does not allow for the recommended ecological goal to be met. The timber sale in-kind services pilot project for the Greenway could potentially result in a great deal more funding for prescribed fire contracting. Many flatwoods zones have a very tight burn window, making it more difficult to burn them when the weather is right. DRP strives to keep zones in rotation burned at the recommended frequency while adding backlogged zones to the burn plan to the extent possible.*
5. **Non-Native, Invasive & Problem Species, specifically prevention of pest/pathogens, received a below average score. The review team is asked to evaluate, based on information provided by the managing agency, as well as overall management actions, whether prevention and control are sufficient.**  
**Managing Agency Response:** *Agree. DRP will continue to work with local, state, and federal agencies in order to prevent the spread of pest and pathogens to best degree practical.*
6. **Hydrologic/Geologic function Hydro-Alteration, specifically roads/culverts, received a below average score. The review team is asked to evaluate, based on information provided by the managing agency, whether consideration of past and present hydrologic and geologic functions are sufficient.**  
**Managing Agency Response:** *Agree. The CFG's history as a sea-level Cross Florida Ship Canal and later Barge Canal are the primary reasons that the CFG has the extensive large scale hydrologic/geologic disturbances and disruptions. The ongoing contentious debate over restoration of the Ocklawaha River vs. retention of the Rodman Reservoir and the ultimate determination of other remnant CFBC project works will require legislative and executive actions to be addressed substantively.*

7. **Resources Protection, specifically signage, received a below average score. The review team is asked to evaluate, based on information provided by the managing agency, whether resources are sufficient to protect the property.**  
*Managing Agency Response: Agree. DRP staff will continue to assess signage needs regarding protection and increase as needed. This is an ongoing process.*
8. **Adjacent Property Concerns, specifically Rodman docks and clearing of uplands adjacent to Rodman, received below average scores. The review team is asked to evaluate, based on information provided by the managing agency, whether adjacent property concerns are sufficiently addressed.**  
*Managing Agency Response: Agree. Because of the extremely sensitive nature of the political turmoil that surrounds the restoration of the Ocklawaha River vs. retention of Rodman Reservoir, garnering support/desire to increase/improve protection and management of CFG shoreline resources along the lower Ocklawaha River and Rodman Reservoir has not been forthcoming.*
9. **Environmental Education & Outreach, specifically wildlife, invasive species, habitat management activities and interpretive facilities and signs, received below average scores. The review team is asked to evaluate, based on information provided by the managing agency, whether environmental education & outreach are sufficient.**  
*Managing Agency Response: Agree. Due to budget and staffing reductions over the past eight years the CFG has closed two former public visitor and information centers that provided CFG history and interpretation information as well as environmental education opportunities for school groups and others. CFG management is now looking at increasing interpretive signage opportunities across the CFG to focus on the unique history and story of this former massive public works canal project corridor.*
10. **Management Resources, staff and funding, received below average scores. The review team is asked to evaluate, based on information provided by the managing agency, whether management resources are sufficient.**  
*Managing Agency Response: Agree. If it is determined that additional staff are needed at the time of the next unit management plan revision, it will be included in the plan. However, no new staff can be assigned to this or any other park unit unless they are appropriated by the Legislature or reassigned from other units. Funding is determined annually by the Florida Legislature. The updated unit management plan will address land management funding needs. However, Division funding is determined annually by the Florida Legislature and funds are allocated to the 171 state parks and trails according to priority needs.*

#### PLAN REVIEW

1. **Natural Communities, specifically basin swamp, blackwater stream and hydric hammock, received below average scores. This is an indication that the management plan does not sufficiently address current or desired condition and/or future management actions to protect or restore.**  
*Managing Agency Response: Agree. DRP staff will add a description and assessment of these communities in the upcoming management plan update. The current management plan was reviewed by the relevant agencies and was in full compliance with Chapters 253 and 259, F.S. and Chapter 18-2 F.A.C. when it was approved by ARC. The next update of this plan will be in full compliance with changes made to the statutes noted above by the Florida Legislature in 2008.*

- 2. Restoration, specifically Deep Creek scrub, received a below average score. This is an indication that the management plan does not sufficiently address restoration.**  
*Managing Agency Response: Agree. DRP staff will add specifics on how to restore the Deep Creek scrub area in the upcoming management plan update. The current management plan was reviewed by the relevant agencies and was in full compliance with Chapters 253 and 259, F.S. and Chapter 18-2 F.A.C. when it was approved by ARC. The next update of this plan will be in full compliance with changes made to the statutes noted above by the Florida Legislature in 2008.*
- 3. Non-native, Invasive & Problem Species, specifically prevention of animals and pests/pathogens and control of pest/pathogens, received below average scores. This is an indication that the management plan does not sufficiently address prevention of invasive species.**  
*Managing Agency Response: Agree. The plan does not specifically address these subjects within the current management plan for the park. DRP staff will add prevention and control methods for pest and pathogens along with exotic animal species in the upcoming management plan update. The current management plan was reviewed by the relevant agencies and was in full compliance with Chapters 253 and 259, F.S. and Chapter 18-2 F.A.C. when it was approved by ARC. The next update of this plan will be in full compliance with changes made to the statutes noted above by the Florida Legislature in 2008.*
- 4. Hydrologic/Geologic function, Hydro-Alteration, specifically roads/culverts and ditches, received below average scores. This is an indication that the management plan does not sufficiently address hydrologic and geologic function.**  
*Managing Agency Response: Agree. On page 55 of the current unit management plan, the plan describes some of the overall problems and issues that the CFG faces in a very wide scale. The plan calls for an assessment of hydrological impacts which is to be conducted in the future. However, a great deal of detail cannot be found in the plan related specifically to the road, culverts and ditches and will be included in the upcoming management plan update when the assessment and inventory is completed. The current management plan was reviewed by the relevant agencies and was in full compliance with Chapters 253 and 259, F.S. and Chapter 18-2 F.A.C. when it was approved by ARC. The next update of this plan will be in full compliance with changes made to the statutes noted above by the Florida Legislature in 2008.*
- 5. Ground Water Monitoring, specifically ground water quality and quantity, received a below average score. This is an indication that the management plan does not sufficiently address ground water quality and quantity.**  
*Managing Agency Response: Agree. The current unit management plan on page 66 and 67 describes goals and objectives as well as research needs for the CFG related to ground water quality and quantity. However, it does not state specific data collected by local water management districts. DRP staff will work with the water management districts in order to include general water quality and quantity data in the next management plan for the park. The current management plan was reviewed by the relevant agencies and was in full compliance with Chapters 253 and 259, F.S. and Chapter 18-2 F.A.C. when it was approved by ARC. The next update of this plan will be in full compliance with changes made to the statutes noted above by the Florida Legislature in 2008.*
- 6. Adjacent Property Concerns, specifically Rodman docks, clearing of uplands adjacent to Rodman and discussion of potential surplus land determination, received below average scores. This is an indication that the management plan does not sufficiently address surplus lands.**



***Managing Agency Response:** Agree. DRP has evaluated the areas in question for possible surplus because more support to enforce existing rules and statutes on state-owned riparian areas is needed. Adjacent property concerns and the determination of surplus lands will be addressed more thoroughly in the next management plan update. The current management plan was reviewed by the relevant agencies and was in full compliance with Chapters 253 and 259, F.S. and Chapter 18-2 F.A.C. when it was approved by ARC. The next update of this plan will be in full compliance with changes made to the statutes noted above by the Florida Legislature in 2008.*

7. **Environmental Education & Outreach, specifically wildlife, invasive species, habitat management activities and interpretive facilities and signs, received below average scores. This is an indication that the management plan does not sufficiently address environmental education and outreach.**

***Managing Agency Response:** Disagree. On pages 66-67 of the current unit management plan it described many environmental education and outreach goals for the Greenway and what facilities currently exist. DRP staff will further expand this section of the plan in the next unit management plan rewrite to further explain this topic. The current management plan was reviewed by the relevant agencies and was in full compliance with Chapters 253 and 259, F.S. and Chapter 18-2 F.A.C. when it was approved by ARC. The next update of this plan will be in full compliance with changes made to the statutes noted above by the Florida Legislature in 2008.*

Thank you for your attention.

/ca

cc: Larry Fooks, Chief, Bureau of Parks District 3  
Robert Yero, Assistant Chief, Bureau of Parks District 3  
Mickey Thomason, Park Manager, Marjorie Harris Carr Cross Florida Greenway  
Jason DePue, Environmental Specialist, Bureau of Parks District 3