


ARROW SQUID (SQU)

(*Nototodarus gouldi*, *N. sloanii*)
 Wheketere



1. FISHERY SUMMARY

1.1 Commercial fisheries

The New Zealand arrow squid fishery is based on two related species. *Nototodarus gouldi* is found around mainland New Zealand north of the Subtropical Convergence, whereas *N. sloanii* is found in and to the south of the convergence zone.

Except for the Southern Islands fishery, for which a separate TACC is set, the two species are managed as a single fishery within an overall TACC. The Southern Islands fishery (SQU 6T) is almost entirely a trawl fishery. Although the species (*N. sloanii*) is the same as that found around the south of the South Island, there is evidence to suggest that the Auckland Island shelf stock is different from the mainland stocks. Because the Auckland Island shelf squid are readily accessible to trawlers, and because they can be caught with little finfish bycatch and are therefore an attractive resource for trawlers, a quota has been set separately for the Southern Islands. Total reported landings and TACCs for each stock are shown in Table 1, while historical landings and TACC are depicted in Figure 1.

The New Zealand squid fishery began in the late 1970s and reached a peak in the early 1980s when over 200 squid jigging vessels came to fish in the New Zealand EEZ. The discovery and exploitation of the large squid stocks in the southwest Atlantic substantially increased the supply of squid to the Asian markets causing the price to fall. In the early 1980s, Japanese squid jiggers would fish in New Zealand for a short time before continuing on to the southwest Atlantic. In the late 1980s, the jiggers stopped transit fishing in New Zealand and the number of jiggers fishing declined from over 200 in 1983 to around 15 in 1994. The jig catch in SQU 1J declined from 53 872 t in 1988–89 to 4865 t in 1992–93 but increased significantly to over 30 000 t in 1994–95, before declining to just over 9000 t in 1997–98. The jig catch declined to low levels for the next four years but then increased back up to almost 9000 t in 2004–05, before declining again to 891 t in 2009–10. The 2010–11 and 2011–12 fishing years have seen an increase from this eight year low to 1811 t.

From 1987 to 1998 the trawl catch fluctuated between about 30 000–70 000 t, but in SQU 6T the impact of management measures to protect the Hooker's sea lion (*Phocarctos hookeri*) restricted the total catch in some years between 1999 and 2005.

Catch and effort data from the SQU 1T fishery show that the catch occurs between December and May, with peak harvest from January to April. The catch has been taken from the Snares shelf on the south

ARROW SQUID (SQU)

coast of the South Island right through to the Mernoo Bank (east coast), but statistical area 028 (Snares shelf and Snares Island region) has accounted for over 77% of the total in recent years. Based on Observer data, squid accounts for 67% of the total catch in the target trawl fishery, with bycatch principally of barracouta, jack mackerel, silver warehou and spiny dogfish.

For 2005–06 a 10% in-season increase to the SQU 1T TACC was approved by the Minister of Fisheries. The catch for December–March was 40% higher than the average over the previous eight years and catch rates were double the average, indicating an increased abundance of squid. Previously, in 2003–04, a 30% in-season increase to the TACC was agreed, but catches did not reach the higher limit. Note that the TACC automatically reverts to the original value at the end of the fishing year.

Table 1: Reported catches (t) and TACCs (t) of arrow squid from 1986–87 to 2013–14. Source - QMS.

Fishstock	SQU 1J*		SQU 1T*		SQU 6T†		SQU 10T‡		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1986–87	32 394	57 705	25 621	30 962	16 025	32 333	0	10	74 040	121 010
1987–88	40 312	57 705	21 983	30 962	7 021	32 333	0	10	69 316	121 010
1988–89	53 872	62 996	26 825	36 081	33 462	35 933	0	10	114 160	135 080
1989–90	13 895	76 136	13 161	47 986	19 859	42 118	0	10	46 915	166 250
1990–91	11 562	46 087	18 680	42 284	10 658	30 190	0	10	40 900	118 571
1991–92	12 985	45 766	36 653	42 284	10 861	30 190	0	10	60 509	118 571
1992–93	4 865	49 891	30 862	42 615	1 551	30 369	0	10	37 278	122 875
1993–94	6 524	49 891	33 434	42 615	34 534	30 369	0	10	74 492	122 875
1994–95	33 615	49 891	35 017	42 741	30 683	30 369	0	10	99 315	123 011
1995–96	30 805	49 891	17 823	42 741	14 041	30 369	0	10	62 668	123 011
1996–97	20 792	50 212	24 769	42 741	19 843	30 369	0	10	65 403	123 332
1997–98	9 329	50 212	28 687	44 741	7 344	32 369	0	10	45 362	127 332
1998–99	3 240	50 212	23 362	44 741	950	32 369	0	10	27 553	127 332
1999–00	1457	50 212	13 049	44 741	6 241	32 369	0	10	20 747	127 332
2000–01	521	50 212	31 297	44 741	3 254	32 369	< 1	10	35 071	127 332
2001–02	799	50 212	35 872	44 741	11 502	32 369	0	10	48 173	127 332
2002–03	2 896	50 212	33 936	44 741	6 887	32 369	0	10	43 720	127 332
2003–04	2 267	50 212	48 060	*58 163	34 635	32 369	0	10	84 962	127 332
2004–05	8 981	50 212	49 780	44 741	27 314	32 369	0	10	86 075	127 332
2005–06	5 844	50 212	49 149	*49 215	17 425	32 369	0	10	72 418	127 332
2006–07	2 278	50 212	49 495	44 741	18 479	32 369	0	10	70 253	127 332
2007–08	1 371	50 212	36 171	44 741	18 493	32 369	0	10	56 035	127 332
2008–09	1 032	50 212	16 407	44 741	28 872	32 369	0	10	46 311	127 332
2009–10	891	50 212	16 759	44 741	14 786	32 369	0	10	32 436	127 332
2010–11	1 414	50 212	14 957	44 741	20 934	32 369	0	10	37 304	127 332
2011–12	1 811	50 212	18 969	44 741	14 427	32 369	0	10	35 207	127 332
2012–13	741	50 212	13 951	44 741	9 944	32 369	0	10	24 637	127 332
2013–14	167	50 212	7483	44 741	7 403	32 369	0	10	15 053	127 332

* All areas except Southern Islands and Kermadec.

† Southern Islands.

‡ Kermadec.

In season increase of 30% for 2003–04 and 10% for 2005–06

1.2 Recreational fisheries

The amount of arrow squid caught by recreational fishers is not known.

1.3 Customary non-commercial fisheries

No quantitative information is available on the current level of customary non-commercial take.

1.4 Illegal catch

There is no quantitative information available on the level of illegal catch.

1.5 Other sources of mortality

No information is available on other sources of mortality.

2. BIOLOGY

Two species of arrow squid are caught in the New Zealand fishery. Both species are found over the continental shelf in water up to 500 m depth, though they are most prevalent in water less than 300 m depth. Both species are sexually dimorphic, though similar in biology and appearance. Individuals can be identified to species level based on sucker counts on Arm I and differences in the hectocotylized arm of males.




Figure 1: Reported commercial landings and TACC for the three main SQU stocks. Top to bottom: SQU 1J (All Waters Except 10T and 6T, Jigging), SQU 1T (All Waters Except 10T and 6T, All Other Methods) and SQU 6T (Southern Islands, All Methods). Note that these figures do not show data prior to entry into the QMS.

ARROW SQUID (SQU)

Recent work on the banding of statoliths from *N. sloanii* suggests that the animals live for around one year. Growth is rapid. Modal analysis of research data has shown increases of 3.0–4.5 cm per month for Gould's arrow squid measuring between 10 and 34 cm Dorsal Mantle Length (DML).

Estimated ages suggest that *N. sloanii* hatches in July and August, with spawning occurring in June and July. It also appears that *N. gouldi* may spawn one to two months before *N. sloanii*, although there are some indications that *N. sloanii* spawns at other times of the year. The squid taken by the fishery do not appear to have spawned.

Tagging experiments indicate that arrow squid can travel on average about 1.1 km per day with a range of 0.14–5.6 km per day.

Biological parameters relevant to stock assessment are shown in Table 2.

Table 2: Estimates of biological parameters.

Fishstock		Estimate		Source
<u>1. Weight = a (length)^b (Weight in g, length in cm dorsal length)</u>				
		a	b	
<i>N. gouldi</i>	≤ 12 cm DML	0.0738	2.63	Mattlin et al (1985)
<i>N. sloanii</i>	≥ 12 cm DML	0.029	3	
<u>2. von Bertalanffy growth parameters</u>				
	<i>K</i>	<i>t₀</i>	<i>L_∞</i>	
<i>N. gouldi</i>	2.1–3.6	0	35	Gibson & Jones (1993)
<i>N. sloanii</i>	2.0–2.8	0	35	

3. STOCKS AND AREAS

There are no new data which would alter the stock boundaries given in previous assessment documents. It is assumed that the stock of *N. gouldi* (the northern species) is a single stock, and that *N. sloanii* around the mainland comprises a unit stock for management purposes, though the detailed structure of these stocks is not fully understood. The distribution of the two species is largely geographically separate but those occurring around the mainland are combined for management purposes. The Auckland Islands Shelf stock of *N. sloanii* appears to be different from the mainland stock and is managed separately.

4. ENVIRONMENTAL AND ECOSYSTEM CONSIDERATIONS

This section was last reviewed by the Aquatic Environment Working Group for the May 2012 Fishery Assessment Plenary. Tables were updated and minor corrections to the text were made for the May 2013 Fishery Assessment Plenary. This summary is from the perspective of the squid trawl fishery; a more detailed summary from an issue by issue perspective is available in the Aquatic Environment & Biodiversity Annual Review (www.mpi.govt.nz/Default.aspx?TabId=126&id=1644).

4.1 Role in the ecosystem

Arrow squid are short-lived and highly variable between years (see Biology section). Hurst et al (2012) reviewed the literature and noted that arrow squid are an important part of the diet for many species. Stevens et al (2012) reported that, between 1960 and 2000, squids (including arrow squid) were important in the diet of banded stargazer (59% of non-empty stomachs), bluenose (26%), giant stargazer (34%), gemfish (43%), and hapuku (21%), and arrow squid were specifically recorded in the diets of alfonsino, barracouta, hake, hoki, ling, red cod, red gurnard, sea perch, and southern blue whiting. In a detailed study on the Chatham Rise (Dunn et al 2009), cephalopods were identified as prey of almost all demersal fish species, and arrow squid were identified in the diet of hake, hoki, ling, Ray's bream, shovelnose spiny dogfish, sea perch, smooth skate, giant stargazer and silver warehou, and was a significant component (over 10% prey weight) of the diet of barracouta and spiny dogfish.

Arrow squid have been recorded as important in the diet of marine mammals such as NZ fur seals and NZ sea lions, particularly during summer and autumn (Fea et al 1999, Harcourt et al 2002, Chilvers 2008, Boren 2008) and in the diet of common dolphins (Meynier et al 2008, Stockin 2008). They are also important in the diet of seabirds such as shy albatross in Australia (Hedd & Gales 2001) and Buller's albatross at the Snares and Solander Islands (James & Stahl 2000). Cephalopods in general are important in the diet of a wide range of Australasian albatrosses, petrels and penguins (Marchant & Higgins 2004).

Arrow squid in New Zealand waters have been reported to feed on myctophids, sprats, pilchards, barracouta, euphausiids, mysids, isopods and squid, probably other arrow squid (Yatsu 1986, Uozumi 1998). Uozumi found that the importance of various food items changed between years, and the percentage of empty stomachs was influenced by area, season, size, maturation, and time of day. In Australia, *N. gouldi* was found to feed mostly on pilchard, barracouta, and crustaceans (O'Sullivan & Cullen 1983). Cannibalism was also recorded.

4.2 Incidental catch (fish and invertebrates)

Based on models using observer and fisher-reported data, total bycatch in the arrow squid trawl fishery ranged from 4500 to 25 000 t per year between 1991 and 2010–11 (Anderson 2013). Over that time period arrow squid comprised about 80% of the total estimated catch recorded by observers in this fishery (Figure 2). The remainder of the observed catch comprised mainly the commercial fish species barracouta (8.5%), spiny dogfish (1.7%), and jack mackerel (1.1%). Invertebrate species made up a much smaller fraction of the bycatch overall (about 1%), but crabs (0.8%), especially the smooth red swimming crab (*Nectocarcinus bennetti*, 0.5%), were frequently caught.

Estimated total annual discards ranged from just over 200 t in 1995–96 to about 5500 in 2001–02 and, like bycatch, peaked in the early 1990s and were at relatively low levels after 2006–07 (Anderson 2013). Most discards were QMS species (about 62% over all years), followed by non-QMS species (19%), invertebrate species (11%), and arrow squid (7%). Absolute levels of discards increased in all categories over the 21-year period; this increase was strongly significant for non-QMS species and total discards, and also marginally significant for QMS species and invertebrates. The species discarded in the greatest amounts were spiny dogfish, redbait, rattails, and silver dory. Discards peaked at 0.13 kg of discarded fish for every 1 kg of arrow squid caught in the early 1990s and declined to 0.02–0.07 kg after 2002–03.

ARROW SQUID (SQU)




Figure 2: Percentage of the total catch contributed by the main bycatch species (those representing 0.05% or more of the total catch) in the observed portion of the arrow squid fishery, and the percentage discarded. The Other category is the sum of all bycatch species representing less than 0.05% of the total catch (Anderson 2013).

4.3 Incidental catch (seabirds, mammals, and protected fish)

For protected species, capture estimates presented here include all animals recovered to the deck (alive, injured or dead) of fishing vessels but do not include any cryptic mortality (e.g., seabirds struck by a warp but not brought onboard the vessel, Middleton & Abraham 2007).

4.3.1 NZ sea lion interactions

The New Zealand (or Hooker's) sea lion was classified in 2008 as "Vulnerable" by the International Union for Conservation of Nature (IUCN) and in 2010 as "Nationally Critical" under the NZ Threat Classification System (Baker et al 2010). Pup production at the main Auckland Island rookeries shows a steady decline since the late 1990s.

NZ sea lions are sometimes caught by vessels trawling for arrow squid (Smith & Baird 2005a, 2007a & b, Thompson & Abraham 2010a, Abraham & Thompson 2011). The trend in observed and estimated captures is downwards. Until recently, captures occurred most frequently in the SQU 6T fishery around the Auckland Islands, and a limit on the number of fishery-related mortalities in this fishery has been set since 1992 (Table 3). These limits have been determined using various approaches, but the current approach is to limit the number of sea lions estimated to have been captured using control rules calculated using the number of pups born in the previous two years. Estimated captures for a year are calculated from the estimated strike rate per tow and the number of tows. The average length of tows has increased substantially over the past decade, but this should be incorporated in the estimated strike rate per tow, albeit with high uncertainty. The likely performance of candidate control rules has been tested using an integrated population and fishery model (Breen et al 2010). Candidate rules are assessed against management criteria developed and agreed in 2003 by a Technical Working Group comprising

Ministry of Fisheries, DOC, NIWA, squid industry representatives, and environmental groups (details can be found in the Aquatic Environment and Biodiversity Annual Review 2012).

Sea Lion Exclusion Devices (SLEDs) were introduced into the SQU 6T fishery in 2001–02 and were in widespread use by 2004–05 leading to a sharp drop in observed incidental captures (Table 4). SLEDs are designed to allow sea lions to escape from a trawl and consist of a grid of steel bars that prevents sea lions entering the codend and an escape hole. From their introduction, SLEDs were subject to continuous design improvements for 10–15 years and, since 2007, a standard Mark 3/13 version has been used by all vessels in the SQU 6T fishery. Tows undertaken using an approved SLED receive a discount on the pre-determined sea lion strike rate, based on the assumption that some sea lions that encounter a trawl equipped with a SLED that would have drowned in the absence of a SLED will survive. This discount was originally set at 20%, was increased to 35% in 2007–08, and further increased to 82% in August 2012. The recent increase in discount rate was made to acknowledge recent research indicating that a high proportion of sea lions encountering a SLED are likely to survive the encounter (summarised in Abraham 2011). There is some remaining uncertainty, including the unknown probability that a sea lion that enters a net but is not subsequently captured will exceed its breath holding limit and die after exiting the trawl via the SLED or the front of the net. This uncertainty is discussed in the 2012 Aquatic Environment and Biodiversity Annual Review (MPI 2012).

Smaller numbers of NZ sea lions are captured in the squid trawl fishery on the Stewart-Snares shelf (SQU 1T, Table 5). Formal estimates of total captures in this fishery have not been calculated but captures across all trawl fisheries on the Stewart-Snares shelf were estimated by Thompson & Abraham (2010a) to vary from 3 to 9 sea lions each year.

Table 3: Fisheries-related mortality limit (FRML) from 1991 to 2012 (♀ = females; numbers in parentheses are FRMLs modified in-season). Direct comparisons among years are not useful because the assumptions underlying the FRML changed over time.

Year	FRML	Discount rate	Management actions
1991–92	16 (♀)		
1992–93	63		
1993–94	63		
1994–95	69		
1995–96	73		Fishery closed by MFish (4 May)
1996–97	79		Fishery closed by MFish (28 Mar)
1997–98	63		Fishery closed by MFish (27 Mar)
1998–99	64		
1999–00	65		Fishery closed by MFish (8 Mar)
2000–01	75		Voluntary withdrawal by industry
2001–02	79		Fishery closed by MFish (13Apr)
2002–03	70		Fishery closed by MFish (29 Mar), overturned by High Court
2003–04	62 (124)	20%	Fishery closed by MFish (22 Mar), overturned by High Court
2004–05	115	20%	Voluntary withdrawal by industry on reaching the FRML
2005–06	97 (150)	20%	FRML increased in mid-March due to abundance of squid
2006–07	93	20%	
2007–08	81	35%	
2008–09	113 (95)	35%	Lower interim limit agreed following decrease in pup numbers
2009–10	76	35%	
2010–11	68	35%	
2011–12	68	35%	
2012–13	68	82%	

ARROW SQUID (SQU)

Table 4: Annual trawl effort, observer coverage, observed numbers of sea lions captured, observed capture rate (sea lions per 100 trawls), estimated sea lion captures, interactions, and the estimated strike or capture rate (with 95% confidence intervals) for the squid trawl fisheries operating in SQU 6T (Auckland Islands). Estimates are based on methods described in Thompson et al (2013) and available via <http://www.fish.govt.nz/en-nz/Environmental/Seabirds/>. Data for 1995–96 to 2001–02 and all estimated strike rates data are from Thompson et al (2013). Data for 2002–03 to 2011–12 are based on data version 20130304 and provisional data for 2012–13 are based on data version 20140131.

Year	Tows	Obs. captures			Est. captures		Est. interactions		Est. strike rate	
		% obs.	No.	Rate	Mean	95% c.i.	Mean	95% c.i.	Mean	95% c.i.
1995–96	4 467	12	13	2.4	131	69–226	131	67–244	2.9	1.6–5.0
1996–97	3 716	19	28	3.9	142	91–208	142	89–210	3.8	2.6–5.5
1997–98	1 441	22	13	4.2	60	33–102	60	31–104	4.2	2.5–6.9
1998–99	402	38	5	3.2	14	7–27	15	5–29	3.6	2.1–5.9
1999–00	1 206	36	25	5.7	69	45–107	69	41–108	5.8	4.0–8.6
2000–01	583	99	39	6.7	39	39–40	61	39–87	10.4	8.6–13.1
2001–02*	1 648	34	21	3.7	43	30–64	73	43–116	4.4	3.0–6.6
2002–03	1 466	28	11	2.6	19	13–28	46	24–77	3.2	2.0–5.1
2003–04	2 594	31	16	2.0	40	26–60	200	98–370	7.5	4.0–13.5
2004–05^	2 693	30	9	1.1	31	17–53	165	73–320	5.9	2.7–11.1
2005–06	2 459	22	10	1.8	27	15–45	149	63–309	6.0	2.7–12.5
2006–07	1 317	41	7	1.3	16	9–26	89	28–200	6.6	2.3–14.8
2007–08	1 265	47	5	0.9	12	6–21	116	21–489	8.0	1.6–30.9
2008–09	1 925	40	2	0.3	7	3–16	97	12–441	4.6	0.7–18.4
2009–10	1 188	26	3	1.0	13	5–26	124	19–508	9.0	1.7–33.6
2010–11	1 583	34	0	0.0	4	0–11	60	4–278	3.5	0.4–14.9
2011–12	1 281	45	0	0.0	2	0–7	43	2–206	-	-
2012–13†	1 027	86	3	0.3	-	-	-	-	-	-

* SLEDs were introduced. ^ SLEDs were standardised and in widespread use. † Provisional data, no model estimates available.

Table 5: Number of tows by fishing year and observed NZ sea lion captures in squid trawl fisheries on the Stewart-Snares shelf, 2002–03 to 2011–12. No. obs, number of observed tows; % obs, percentage of tows observed; Rate, number of captures per 100 observed tows. Estimates are based on methods described in Thompson et al (2013) and available via <http://www.fish.govt.nz/en-nz/Environmental/Seabirds/>. Data for 2002–03 to 2011–12 are based on data version 20130304 and provisional data for 2012–13 are based on data version 20140131.

	Tows	Fishing effort		Observed captures		Estimated interactions		
		No. obs	% obs	Captures	Rate	Mean	95% c.i.	% included
2002–03	3 281	506	15.4	0	0.00	2	0–5	100.0
2003–04	4 534	957	21.1	1	0.10	3	1–7	100.0
2004–05	5 861	1 581	27.0	3	0.19	6	3–11	100.0
2005–06	4 481	537	12.0	1	0.19	4	1–8	100.0
2006–07	2 925	706	24.1	1	0.14	2	1–5	100.0
2007–08	2 412	864	35.8	0	0.00	1	0–3	100.0
2008–09	1 809	531	29.4	0	0.00	1	0–3	100.0
2009–10	2 259	764	33.8	1	0.13	2	1–4	100.0
2010–11	2 177	685	31.5	0	0.00	1	0–3	100.0
2011–12	1 983	800	40.3	0	0.00	1	0–2	100.0
2012–13†	1 529	1 343	87.8	0	0.00	0	0–1	100.0

† Provisional data, no model estimates available.

4.3.2 NZ fur seal interactions

The New Zealand fur seal was classified in 2008 as “Least Concern” by IUCN and in 2010 as “Not Threatened” under the NZ Threat Classification System.

Vessels targeting arrow squid incidentally catch fur seals (Baird & Smith 2007a, Smith & Baird 2009, Thompson & Abraham 2010b, Baird 2011), mostly off the east coast South Island, on the Stewart-

Snares shelf, and close to the Auckland Islands. In the 2012–13 fishing year there were six observed captures of New Zealand fur seal in squid trawl fisheries. In the 2011–12 fishing year, there were 25 (95% c.i.: 12–53) estimated captures, with the estimates made using a statistical model (Thompson et al 2013, Table 6). Total estimated captures in squid trawl fisheries varied from 21 to 168 between 2002–03 and 2011–12, representing about 9% of the total estimated captures in trawl fisheries over those years (noting that less than 50% of all trawl effort is included in the estimates, except for the most recent year). The rate of capture over this period varied from 0.08 to 0.96 captures per hundred tows without obvious trend (Table 6), a rate that is about 40% of the rate for all trawl

Table 6: Number of tows by fishing year and observed and model-estimated total NZ fur seal captures in squid trawl fisheries, 2002–03 to 2012–13. No. obs, number of observed tows; % obs, percentage of tows observed; Rate, number of captures per 100 observed tows, % inc, percentage of total effort included in the statistical model. Estimates are based on methods described in Thompson et al (2013) and available via <http://www.fish.govt.nz/en/nz/Environmental/Seabirds/>. Data for 2002–03 to 2011–12 are based on data version 20130304 and provisional data for 2012–13 are based on data version 20140131

	Tows	Observed				Estimated		
		No. obs	% obs	Captures	Rate	Captures	95% c.i.	% inc.
2002–03	8 410	1 308	15.6	8	0.61	56	27 – 108	100.0
2003–04	8 336	1 771	21.2	17	0.96	89	46 – 166	100.0
2004–05	10 486	2 511	23.9	16	0.64	155	82 – 284	100.0
2005–06	8 575	1 103	12.9	4	0.36	96	43 – 192	100.0
2006–07	5 906	1 289	21.8	8	0.62	42	20 – 83	100.0
2007–08	4 236	1 457	34.4	6	0.41	33	14 – 67	100.0
2008–09	3 867	1 298	33.6	1	0.08	19	6 – 45	100.0
2009–10	3 789	1 070	28.2	8	0.75	33	15– 66	100.0
	Tows	Observed				Estimated		
		No. obs	% obs	Captures	Rate	Captures	95% c.i.	% inc.
2010–11	4 214	1.261	29.9	8	0.63	23	12 – 42	100.0
2011–12	3 505	1 380	39.4	8	0.58	23	11 – 48	100.0
2012–13	2 646	2 273	85.9	6	0.26	8	6– 17	-

4.3.3 Seabird interactions

Vessels targeting arrow squid incidentally catch seabirds. Baird (2005a) summarised observed seabird captures in the arrow squid target fishery for the fishing years 1998–99 to 2002–03 and calculated total seabird captures for the areas with adequate observer coverage using ratio based estimations. Baird & Smith (2007b, 2008) summarised observed seabird captures and used both ratio-based and model-based predictions to estimate the total seabird captures for 2003–04, 2004–05 and 2005–06. Abraham & Thompson (2011) summarised captures of protected species and used model and ratio-based predictions of the total seabird captures for 1989–90 and 2008–09.

In the 2012–13 fishing year there were 450 observed captures of birds in squid trawl fisheries. In the 2011–12 fishing year, there were 327 (95% c.i.: 261–422) estimated captures, with the estimates made using a statistical model (Table 7).

ARROW SQUID (SQU)

Table 7: Number of tows by fishing year and observed and model-estimated total bird captures in squid trawl fisheries, 2002–03 to 2011–12. No. obs, number of observed tows; % obs, percentage of tows observed; Rate, number of captures per 100 observed tows, % inc, percentage of total effort included in the statistical model. Estimates are based on methods described in Abraham et al (2013) and are available via <http://www.fish.govt.nz/en/nz/Environmental/Seabirds/>. Estimates from 2002–03 to 2010–11 are based on data version 20120531 and preliminary estimates for 2011–12 are based on data version 20130304.

	Tows	Observed				Estimated		
		No. obs	% obs	Captures	Rate	Captures	95% c.i.	% inc.
2002–03	8 410	1 308	15.6	159	12.16	929	761 – 1 227	100.0
2003–04	8 336	1 771	21.2	204	11.52	857	717 – 1027	100.0
2004–05	10 486	2 511	23.9	384	15.29	1 442	1 236 – 1 710	100.0
2005–06	8 575	1 103	12.9	200	18.13	1 203	947 – 1 568	100.0
2006–07	5 906	1 289	21.8	127	9.85	573	440 – 767	100.0
2007–08	4 236	1 457	34.4	162	11.12	540	425 – 705	100.0
2008–09	3 867	1 298	33.6	259	19.95	624	523 – 764	100.0
2009–10	3 789	1 070	28.2	92	8.60	401	311 – 521	100.0
2010–11	4 214	1 261	29.9	141	11.18	600	468 – 779	100.0
2011–12	3 505	1 380	39.4	106	7.68	344	277 – 438	100.0
2012–13†	2 646	2 273	85.9	450	19.78	505	477 – 553	100.0

† Provisional data, no model estimates available

Total estimated seabird captures in squid trawl fisheries varied from 327 to 1397 between 2002–03 and 2011–12 at a rate of 8.6 to 20.0 captures per hundred tows without obvious trend (Table 7). These estimates include all bird species and should be interpreted with caution because trends by species can be masked. The average capture rate in squid trawl fisheries over the last ten years is about 13.79 birds per 100 tows, a high rate relative to trawl fisheries for scampi (5.57 birds per 100 tows) and hoki (2.16 birds per 100 tows) over the same years. The squid fishery accounted for about 58% of seabird captures in the trawl fisheries modelled by Abraham et al (2013).

Observed seabird captures since 2002–03 have been dominated by four species: white-capped and southern Buller’s albatrosses make up 85% and 9% of the albatrosses captured, respectively; and white-chinned petrels and sooty shearwaters make up 48% and 45% of other birds, respectively, the total and fishery risk ratios presented in Table 8. Most captures occur on the Stewart-Snares shelf (60%) or close to the Auckland Islands (37%). These numbers should be regarded as only a general guide on the distribution of captures because observer coverage is not uniform across areas and may not be representative.

Table 8: Risk ratio of seabirds predicted by the level two risk assessment for SLL fisheries (grouped by vessel size) and all fisheries included in the level two risk assessment, 2002–03 to 2011–12, showing seabird species with a risk ratio of at least 0.001 of PBR₁. The risk ratio is an estimate of aggregate potential fatalities across trawl and longline fisheries relative to the Potential Biological Removals, PBR₁ (from Richard and Abraham 2013 where full details of the risk assessment approach can be found). PBR₁ applies a recovery factor of 1.0. Typically a recovery factor of 0.1 to 0.5 is applied (based on the state of the population) to allow for recovery from low population sizes as quickly as possible. This should be considered when interpreting these results. The DOC threat classifications are shown (Robertson et al 2013 at <http://www.doc.govt.nz/documents/science-and-technical/nztc4entire.pdf>).

Species name	PBR ₁ (mean)	Risk ratio		Risk category	DoC Threat Classification
		Squid target trawl	TOTAL		
Salvin's albatross	975	0.035	2.756	Very high	Threatened: Nationally Critical
Flesh-footed shearwater	590	0.001	1.321	Very high	Threatened: Nationally Vulnerable
Southern Buller's albatross	513	0.292	1.292	Very high	At Risk: Naturally Uncommon
Chatham Island albatross	159	0.004	1.291	Very high	At Risk: Naturally Uncommon
NZ white-capped albatross	4 044	0.229	0.700	Very high	At Risk: Naturally Uncommon
Northern Buller's albatross	617	0.002	0.678	High	At Risk: Naturally Uncommon

Table 8 [Continued]

Species name	PBR _i (mean)	Risk ratio		Risk category	DoC Threat Classification
		Squid target trawl	TOTAL		
Gibson's albatross	260	0.018	0.467	High	Threatened: Nationally Critical
Cape petrel	840	0.007	0.303	High	At Risk: Naturally Uncommon
Antipodean albatross	295	0.005	0.301	High	Threatened: Nationally Critical
Northern royal albatross	396	0.011	0.271	Medium	At Risk: Naturally Uncommon
Southern royal albatross	441	0.012	0.264	Medium	At Risk: Naturally Uncommon
Westland petrel	241	0.003	0.263	Medium	At Risk: Naturally Uncommon
Northern giant petrel	217	0.008	0.215	Medium	At Risk: Naturally Uncommon
White-chinned petrel	7 925	0.066	0.211	Medium	At Risk: Declining
Campbell black-browed albatross	1 017	0.002	0.189	Medium	At Risk: Naturally Uncommon
Grey petrel	2 172	0.001	0.114	Medium	At Risk: Naturally Uncommon
Light-mantled sooty albatross	237	0.002	0.028	Low	At Risk: Declining
Grey-headed albatross	333	0.001	0.018	Low	Threatened: Nationally Vulnerable
Sooty shearwater	348 096	0.002	0.005	Low	At Risk: Declining

Mitigation methods such as streamer (tori) lines, Brady bird bafflers, warp deflectors, and offal management are used in the squid trawl fishery. Warp mitigation was voluntarily introduced from about 2004 and made mandatory in April 2006 (Ministry of Fisheries 2006). The 2006 notice mandated that all trawlers over 28 m in length use a seabird scaring device while trawling (being “paired streamer lines”, “bird baffle” or “warp deflector” as defined in the notice). During the 2005–06 fishing year a large trial of mitigation devices was conducted in the squid fishery (Middleton & Abraham 2007). Eighteen vessels were involved in the trial which used observations of seabird heavily contacting the trawl warps (‘warp strikes’) to quantify the effect of using three mitigation devices; paired streamer/tori lines, four boom bird bafflers and warp scarers. Few warp strikes occurred in the absence of offal discharge. When offal was present the tori lines were most effective at reducing warp strikes. All mitigation devices were more effective for reducing large bird warp strikes than small bird. There were, however, about as many bird strikes on the tori lines as the number of strikes on unmitigated warps. The effect of these strikes has not been assessed (Middleton & Abraham 2007).

In the four complete fishing years after mitigation was made mandatory, the average rate of capture for white-capped albatross (90% of albatross captures in this fishery) was 3.2 birds per 100 tows compared with 7.9 per 100 tows in the three complete years before mitigation was made mandatory. This trend is masked in Table 7 by continued captures of smaller birds, mostly in trawl nets as opposed to on trawl warps (where mitigation is focused).

4.4 Benthic interactions

Between 1989–90 and 2004–05, 131 973 trawl tows for squid on or within 1 m of the seabed were reported, comprising 13.7% of all trawl tows on or within 1 m of the seabed reported on TCEPR forms in those years (range 8–23% by year, Baird et al 2011). Black et al (2013) estimated that hoki arrow squid has accounted for 13.5% of all tows reported on TCEPR forms since 1989–90. Between 2006–07 and 2010–11, 95% of arrow squid catch was reported on TCEPR forms. The great majority of tows are conducted on the Stewart-Snares shelf or north and east of the Auckland Islands, with smaller numbers off the east coast of the South Island and the Chatham Rise. Tows were located in Benthic Optimised Marine Environment Classification (BOMECE, Leathwick et al 2009) classes E (outer shelf), F, H (upper slope), I, J, L, and M (mid-slope) (Baird & Wood 2012), and 92% were between 100 and 300 m depth (Baird et al 2011). Tables 4–7 show that the number of trawl tows for squid varies between years, largely without trend and presumably in response to variations in the abundance of squid and management measures to limit the number of sea lions caught. The average duration of trawls has increased over this time so the trend in aggregate swept area will not be the same.

ARROW SQUID (SQU)

Bottom trawling for squid, like trawling for other species, is likely to have effects on benthic community structure and function (e.g., see Rice 2006 for an international review) and there may be consequences for benthic productivity (e.g., Jennings 2001, Hermsen et al 2003, Hiddink et al 2006, Reiss et al 2009). These are not considered in detail here but are discussed in the 2012 Aquatic Environment and Biodiversity Annual Review.

4.5 Other considerations

A substantial decline in the west coast jig fishery for squid will have reduced any trophic implications of that fishery.

5. STOCK ASSESSMENT

Arrow squid live for one year, spawn once then die. Every squid fishing season is therefore based on what amounts to a new stock. It is not possible to calculate reliable yield estimates from historical catch and effort data for a resource which has not yet hatched, even when including data which are just one year old. Furthermore, because of the short life span and rapid growth of arrow squid, it is not possible to estimate the biomass prior to the fishing season. Moreover, the biomass increases rapidly during the season and then decreases to low levels as the animals spawn and die.

5.1 Estimates of fishery parameters and abundance

No estimates are available.

5.2 Biomass estimates

Biomass estimates are not available for squid.

5.3 Yield estimates and projections

It is not possible to estimate *MCY*.

It is not possible to estimate *CAY*.

5.4 Other yield estimates and stock assessment results

There are no other yield estimates of stock assessment results available for arrow squid.

5.5 Other factors

N. gouldi spawns one to two months before *N. sloanii*. This means that at any given time *N. gouldi* is older and larger than *N. sloanii*. The annual squid jigging fishery begins on *N. gouldii* and at some time during the season the biomass of *N. sloanii* will exceed that of *N. gouldi* and the fleet will move south. If *N. sloanii* are abundant the fleet will remain in the south fishing for *N. sloanii*. If *N. sloanii* are less abundant the fleet will return north and resume fishing *N. gouldi*.

6. STATUS OF THE STOCKS

No estimates of current and reference biomass are available. There is also no proven method at this time to estimate yields from the squid fishery before a fishing season begins based on biomass estimates or CPUE data.

Because squid live for about one year, spawn and then die, and because the fishery is so variable, it is not practical to predict future stock size in advance of the fishing season. As a consequence, it is not possible to estimate a long-term sustainable yield for squid, nor determine if recent catch levels or the current TACC will allow the stock to move towards a size that will support the *MSY*. There will be some years in which economic or other factors will prevent the TACC from being fully taken, while in other years the TACC may be lower than the potential yield. It is not known whether New Zealand squid stocks have ever been stressed through fishing mortality.

TACCs and reported landings for the 2013–14 fishing year are summarised in Table 9.

Table 9: Summary of TACCs (t) and reported landings (t) of arrow squid for the most recent fishing year.

Fishstock	2013–14	2013–14
	Actual TACC	Reported landings
SQU 1J	50 212	167
SQU 1T	44 741	7 483
SQU 6T	32 369	7 403
SQU 10T	10	0
Total	127 332	15 503

7. FOR FURTHER INFORMATION

- Abraham, E R (2011) Probability of Mild Traumatic Brain Injury for sea lions interacting with SLEDs. Final Research Report for project SRP2011-03 (Unpublished report held by the Ministry for Primary Industries, Wellington). 21 p.
- Abraham, E R; Thompson, F N (2011) Summary of the capture of seabirds, marine mammals, and turtles in New Zealand commercial fisheries, 1998–99 to 2008–09. *New Zealand Aquatic Environment and Biodiversity Report No. 80* 172 p.
- Abraham, E R; Thompson, F N; Berkenbusch, K (2013) Estimated capture of seabirds in New Zealand trawl and longline fisheries, 2002–03 to 2010–11. Final Research Report for Ministry for Primary Industries project PRO2010-01 (Unpublished report held by Ministry for Primary Industries, Wellington).
- Abraham, E R; Thompson, F N; Oliver, M D (2010) Summary of the capture of seabirds, mammals, and turtles in New Zealand commercial fisheries, 1998–99 to 2007–08. *New Zealand Aquatic Environment and Biodiversity Report No. 45*. 149p.
- Anderson, O F (2004) Fish discards and non-target fish catch in the trawl fisheries for arrow squid, jack mackerel, and scampi in New Zealand Waters. *New Zealand Fisheries Assessment Report. 2004/10*. 61 p.
- Anderson, O F (2013) Fish and invertebrate bycatch and discards in New Zealand arrow squid fisheries from 1990–91 until 2010–11. *New Zealand Aquatic Environment and Biodiversity Report No. 112*. 62 p.
- Baird, S J (2005a) Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2002–03. *New Zealand Fisheries Assessment Report 2005/2*. 50 p.
- Baird, S J (2005b) Incidental capture of *Phocarctos hookeri* (New Zealand sea lions) in New Zealand commercial fisheries, 2001–02. *New Zealand Fisheries Assessment Report. 2005/08*.
- Baird, S J (2005c) Incidental capture of *Phocarctos hookeri* (New Zealand sea lions) in New Zealand commercial fisheries, 2002–03. *New Zealand Fisheries Assessment Report. 2005/09*.
- Baird, S J (2011) New Zealand fur seals – summary of current knowledge. *New Zealand Aquatic Environment and Biodiversity Report No. 72*. 50 p.
- Baird, S J; Doonan, I J (2005) *Phocarctos hookeri* (New Zealand sea lions): incidental captures in New Zealand commercial fisheries during 2000–01 and in-season estimates of captures during squid trawling in SQU 6T in 2002. *New Zealand Fisheries Assessment Report, 2005/17*.
- Baird, S J; Smith, M H (2007a) Incidental capture of New Zealand fur seals (*Arctocephalus forsteri*) in commercial fisheries in New Zealand waters, 2003–04 to 2004–05. *New Zealand Aquatic Environment and Biodiversity Report No. 14*. 98 p.
- Baird, S J; Smith, M H (2007b) Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2003–04 and 2004–05. *New Zealand Aquatic Environment and Biodiversity Report No. 9*. 108 p.
- Baird, S J; Smith, M H (2008) Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2005–06. *New Zealand Aquatic Environment and Biodiversity Report No. 18*. 124 p.
- Baird, S J; Wood, B A (2012) Extent of coverage of 15 environmental classes within the New Zealand EEZ by commercial trawling with seafloor contact. *New Zealand Aquatic Environment and Biodiversity Report No. 89*.
- Baird, S J; Wood, B A; Bagley, N W (2011) Nature and extent of commercial fishing effort on or near the seafloor within the New Zealand 200 n. mile Exclusive Economic Zone, 1989–90 to 2004–05. *New Zealand Aquatic Environment and Biodiversity Report No. 73*.
- Baker, C S; Chilvers, B L; Constantine, R; DuFresne, S; Mattlin, R H; van Helden, A; Hitchmough, R (2010) Conservation status of New Zealand marine mammals (suborders Cetacea and Pinnipedia), 2009. *New Zealand Journal of Marine and Freshwater Research 44*: 101–115.
- Ballara, S L; Anderson, O F (2009) Fish discards and non-target fish catch in the trawl fisheries for arrow squid and scampi in New Zealand waters. *New Zealand Aquatic Environment and Biodiversity Report No. 38*. 102 p.
- Black, J; Wood, R; Berthelsen, T; Tilney, R (2013) Monitoring New Zealand's trawl footprint for deepwater fisheries: 1989–1990 to 2009–2010. *New Zealand Aquatic Environment and Biodiversity Report No. 110*. 57 p.
- Boren, L (2008) Diet of New Zealand fur seals (*Arctocephalus forsteri*): a summary. *DOC Research & Development Series 319*. 19 p.
- Bradford, E (2002) Estimation of the variance of mean catch rates and total catches of non-target species in New Zealand fisheries. *New Zealand Fisheries Assessment Report 2002/54*. 60 p.
- Breen, P A; Fu, D; Gilbert, D J (2010) Sea lion population modelling and management procedure evaluations: Report for Project SAP2008/14, Objective 2. Final Research Report for Ministry of Fisheries project SAP2008/14, Objective 2. (Unpublished report held by Ministry for Primary Industries, Wellington.)
- Brothers, N; Duckworth, A R; Safina, C; Gilman, E L (2010) Seabird bycatch in pelagic longline fisheries is grossly underestimated when using only haul data. *PLoS One 5*: e12491. doi: 10.1371/journal.pone.001249
- Chilvers, B L (2008) New Zealand sea lions *Phocarctos hookeri* and squid trawl fisheries: bycatch problems and management options. *Endangered Species Research 5*: 193–204.
- Doonan, I J (1995) Incidental catch of Hooker's sea lion in the southern trawl fishery for squid, summer 1994. Fisheries Assessment Research Document. 1995/22. (Unpublished document held by NIWA library, Wellington.)
- Doonan, I J (2000) Estimation of Hooker's sea lion *Phocarctos hookeri* captures in the southern squid trawl fisheries, 2000. *New Zealand Fisheries Assessment Report. 2000/41*.
- Doonan, I J (2001) Estimation of Hooker's sea lion *Phocarctos hookeri* captures in the southern squid trawl fisheries, 2001. *New Zealand Fisheries Assessment Report. 2001/67*.
- Dunn, M (2009) Feeding habits of the ommastrephid squid *Nototodarus sloanii* on the Chatham Rise, New Zealand. *New Zealand Journal of Marine and Freshwater Research, 43(5)*: 1103–1113.
- Dunn, M., Horn, P, Connell, A, Stevens, D, Forman, J, Pinkerton, M, Griggs, L, Notman, P, Wood, B. (2009) Ecosystem-scale trophic relationships: diet composition and guild structure of middle-depth fish on the Chatham Rise. Final Research Report for Ministry of Fisheries Project, ZBD2004-02. (Unpublished report held by Ministry for Primary Industries, Wellington.)

ARROW SQUID (SQU)

- Fea, N I; Harcourt, R; Lalas, C (1999) Seasonal variation in the diet of New Zealand fur seals (*Arctocephalus forsteri*) at Otago Peninsula, New Zealand. *Wildlife Research* 26(2): 147–160.
- Förch, E C (1983) Squid - current research. In: Taylor, J L; Baird, G G (eds.) New Zealand finfish fisheries: the resources and their management, pp. 33–34. Trade Publications Ltd., Auckland.
- Gibson, D; Jones, J B (1993) Fed up with parasites? — old fish are. *Marine Biology* 117: 495–500.
- Gibson, D J M (1995) The New Zealand Squid Fishery, 1979–93. *MAF Fisheries Technical Report No 42*. 43 p.
- Harcourt, R G; Bradshaw, C J A; Dickson, K; Davis, L S (2002) Foraging ecology of a generalist predator, the female New Zealand fur seal. *Marine Ecology Progress Series* 227: 11–24.
- Hedd, H; Gales, R (2001) The diet of shy albatrosses (*Thalassarche cauta*) at Albatross Island, Tasmania. *Journal of Zoology* 253(1): 69–90.
- Hermesen, J M; Collie, J S; Valentine, P C (2003) Mobile fishing gear reduces benthic megafaunal production on Georges Bank. *Marine Ecology Progress Series* 260: 97–108.
- Hiddink, J G; Jennings, S; Kaiser, M J; Queiros, A M; Duplisea, D E; Piet, G J (2006) Cumulative impacts of seabed trawl disturbance on benthic biomass, production, and species richness in different habitats. *Canadian Journal of Fisheries and Aquatic Sciences* 63:721–36.
- Hurst, R J; Ballara, S L; MacGibbon, D; Triantafillos, L (2012) Fishery characterisation and standardised CPUE analyses for arrow squid (*Nototodarus gouldi* and *N. sloanii*), 1989–90 to 2007–08, and potential management approaches for southern fisheries. *New Zealand Fisheries Assessment Report 2012/47*. 303 p.
- James, G D; Stahl, J C (2000) Diet of the Buller's albatross (*Diomedea bulleri bulleri*) and the importance of fishery discards during chick rearing. *New Zealand Journal of Marine and Freshwater Research* 34: 435–454.
- Jennings, S; Dinmore, T A; Duplisea, D E; Warr, K J; Lancaster, J E (2001) Trawling disturbance can modify benthic production processes. *Journal of Animal Ecology* 70: 459–475.
- Jones, M R L (2007) Historic trawl data and recent information infers temporal change in the occurrence of squid in the diet of orange roughy (*Hoplostethus atlanticus* Collett) in New Zealand. *Reviews in Fish Biology and Fisheries* 17(2–3): 493–499.
- Langley, A D (2001) Summary of catch and effort data from the SQU 1J, SQU 1T, and SQU 6T fisheries for 1989–90 to 1999–2000. *New Zealand Fisheries Assessment Report*. 2001/51.
- Leathwick, J R; Rowden, A; Nodder, S; Gorman, R; Bardsley, S; Pinkerton, M; Baird, S J; Hadfield, M; Currie, K; Goh, A (2009) Benthic-optimised marine environment classification for New Zealand waters. Final Research Report project for Ministry of Fisheries project BEN2006/01. 52 p. (Unpublished report held by Ministry for Primary Industries, Wellington.)
- McKenzie, D; Fletcher, D (2006) Characterisation of seabird captures in commercial trawl and longline fisheries in New Zealand 1997/98 to 2003/04. Final Research Report for Ministry of Fisheries project ENV2004/04. (Unpublished report held by Ministry for Primary Industries, Wellington.) 102 p.
- Marchant, S; Higgins, P J (2004) (Co-ordinators). Handbook of Australian, New Zealand and Antarctic birds. Volume 1 Ratites to Ducks. Oxford University Press, Melbourne, Australia. 735 p.
- Mattlin, R H (1983) Squid. In: Taylor, J. L.; Baird, G. G. (eds.) New Zealand finfish fisheries: the resources and their management, pp. 30–32. Trade Publications Ltd., Auckland.
- Mattlin, R H; Colman, J A (1988) Arrow squid. New Zealand Fisheries Assessment Research Document 88/34. 16 p. (Unpublished document held in NIWA library, Wellington).
- Mattlin, R H; Scheibling, R E; Förch, E C (1985) Distribution, abundance and size structure of arrow squid (*Nototodarus* sp.) off New Zealand. *Northwest Atlantic Fisheries Organisation Scientific Council Studies* 9: 39–45.
- Meynier, L; Mackenzie, D D S; Duignan, P J; Chilvers, B L; Morel, P C H (2009) Variability in the diet of New Zealand sea lion (*Phocarcos hookeri*) at the Auckland Islands, New Zealand. *Marine Mammal Science* 25(2): 302–326.
- Meynier, L; Stockin, K A; Bando, M K H; Duignan, P J (2008) Stomach contents of common dolphin (*Delphinus* sp.) from New Zealand waters. *New Zealand Journal of Marine and Freshwater Research* 42: 257–268.
- Middleton, D A J; Abraham, E R (2007) The efficacy of warp strike mitigation devices: Trials in the 2006 squid fishery. Final Research Report for Ministry of Fisheries project IPA2006/02. (Unpublished report held by Ministry for Primary Industries, Wellington).
- Ministry of Fisheries (2006) Fisheries (Commercial Fishing) Amendment Regulations 2006. Seabird Scaring Devices – Circular Issued under Authority of the Fisheries (Commercial Fishing) Amendment Regulations 2006 (No. F361). *New Zealand Gazette No. 33*, 6 April 2006. 1 p.
- MPI (2012) Aquatic Environment and Biodiversity Annual Review 2012. Compiled by the Fisheries Management Science. Ministry for Primary Industries, Wellington, New Zealand. 388 p.
- O'Sullivan, D; Cullen, J M (1983) Food of the squid *Nototodarus gouldi* in Bass Strait. *Australian Journal of Marine and Freshwater Research* 34: 261–285.
- Reiss, H; Greenstreet, S P R; Sieben, K; Ehrlich, S; Piet, G J; Quirijns, F; Robinson, L; Wolff, W J; Kröncke, I (2009) Effects of fishing disturbance on benthic communities and secondary production within an intensively fished area. *Marine Ecology Progress Series* 394: 201–213.
- Rice, J (2006) Impacts of Mobile Bottom Gears on Seafloor Habitats, Species, and Communities: A Review and Synthesis of Selected International Reviews. *Canadian Science Advisory Secretariat Research Document 2006/057*. 35 p. (available from http://www.dfo-mpo.gc.ca/CSAS/Csas/DocREC/2006/RES2006_057_e.pdf).
- Richard, Y; Abraham, E R (2013) Risk of commercial fisheries to New Zealand seabird populations. *New Zealand Aquatic Environment and Biodiversity Report No. 109*. 58 p.
- Smith, M H; Baird, S J (2005) Representativeness of past observer coverage, and future coverage required for estimation of New Zealand sea lion (*Phocarcos hookeri*) captures in the SQU 6T fishery. *New Zealand Fisheries Assessment Report*. 2005/05.
- Smith, M H; Baird, S J (2005a) Factors that may influence the level of incidental mortality of New Zealand sea lions (*Phocarcos hookeri*) in the squid (*Nototodarus* spp.) trawl fishery in SQU 6T. *New Zealand Fisheries Assessment Report* 2005/20.
- Smith, M H; Baird, S J (2007a) Estimation of incidental captures of New Zealand sea lions (*Phocarcos hookeri*) in New Zealand fisheries in 2003–04, with particular reference to the SQU 6T squid trawl fishery. *New Zealand Fisheries Assessment Report* 2007/7. 32 p.
- Smith, M H; Baird, S J (2007b) Estimation of incidental captures of New Zealand sea lions (*Phocarcos hookeri*) in New Zealand fisheries in 2004–05, with particular reference to the SQU 6T squid trawl fishery. *New Zealand Aquatic Environment and Biodiversity Report No. 12*. 31 p.
- Smith, M H; Baird, S J (2009) Model-based estimation of New Zealand fur seal (*Arctocephalus forsteri*) incidental captures and strike rates for trawl fishing in New Zealand waters for the years 1994–95 to 2005–06. *New Zealand Aquatic Environment and Biodiversity Report No. 40*. 92 p.
- Smith, P J; Mattlin, R H; Roeleveld, M A; Okutani, T (1987) Arrow squids of the genus *Nototodarus* in New Zealand waters: systematics, biology and fisheries. *New Zealand Journal of Marine and Freshwater Research* 21: 315–326.
- Stevens, D W; Hurst, R J; Bagley, N W (2012) Feeding habits of New Zealand fishes; a literature review and summary of research trawl database records 1960 to 2000. *New Zealand Aquatic Environment and Biodiversity Report No. 85*.
- Stockin, K A (2008) The New Zealand common dolphin (*Delphinus* sp.): identity, ecology and conservation : a thesis submitted in partial fulfilment of the requirements for the degree of Doctor of Philosophy in Zoology, Massey University, Auckland, New Zealand.

- Thompson, D R (2008) Autopsy report for seabirds killed and returned from New Zealand fisheries, 1 October 2006 to 30 September 2007. Report prepared for the Conservation Services Programme, Department of Conservation: Contract INT2006/02. Available on www.doc.govt.nz
- Thompson, D R (2009) Seabird Autopsy Project: Summary Report for the 2007–08 Fishing Year. Report prepared for the Conservation Services Programme, Department of Conservation: Contract INT2007/02. Available on www.doc.govt.nz
- Thompson, F N; Abraham, E R (2010a) Estimation of the capture of New Zealand sea lions (*Phocarctos hookeri*) in trawl fisheries, from 1995–96 to 2008–09. *New Zealand Aquatic Environment and Biodiversity Report No. 66*.
- Thompson, F N; Abraham, E R (2010b) Estimation of fur seal (*Arctocephalus forsteri*) bycatch in New Zealand trawl fisheries, 2002–03 to 2008–09. *New Zealand Aquatic Environment and Biodiversity Report No. 61*. 37 p.
- Thompson, F N; Abraham, E R (2012) Captures of New Zealand sea lion in squid trawl fisheries, during the 2008/2009 fishing year. Retrieved from <http://bycatch.dragonfly.co.nz/v20120315/new-zealand-sea-lion/squid-trawl/all-vessels/eez/2009/>, Apr 27, 2012
- Thompson, F N; Abraham, E R; Berkenbusch, K (2011) Marine mammal bycatch in New Zealand trawl fisheries, 1995–96 to 2009–10. Final Research Report for Ministry of Fisheries project PRO2010-01 (Unpublished report held by the Ministry for Primary Industries, Wellington). 80 p.
- Thompson, F N; Abraham, E R; Oliver, M D (2010a) Estimation of fur seal bycatch in New Zealand sea lions trawl fisheries, 2002–03 to 2007–08. *New Zealand Aquatic Environment and Biodiversity Report No. 56*.
- Thompson, F N; Berkenbusch, K; Abraham, E R (2013) Marine mammal bycatch in New Zealand trawl fisheries, 1995–96 to 2010–11. *New Zealand Aquatic Environment and Biodiversity Report No. 105*. 73 p.
- Thompson, F N; Oliver, M D; Abraham, E R (2010b) Estimation of the capture of New Zealand sea lions (*Phocarctos hookeri*) in trawl fisheries, from 1995–96 to 2007–08. *New Zealand Aquatic Environment and Biodiversity Report No. 52*.
- Uozumi, Y (1998) Fishery biology of arrow squids, *Nototodarus gouldi* and *sloanii*, in New Zealand waters. *Bulletin of the National Institute of Far Seas Fisheries* 35. 111 p.
- Uozumi, Y; Ohara, H (1992) Age and growth of *Nototodarus sloanii* (Cephalopoda: Oegopsida) based on daily increment counts in statoliths. *Nippon Suisan Gakkaishi* 59: 1469–1477.
- Yatsu, A (1986) Feeding habit of *N. sloanii* caught by bottom trawl. *Japanese Marine Fishery Resource research Centre* 30: 45–52.