

Caridea (Crustacea, Decapoda) collected on the Brazilian (13°/22°S) continental shelf and slope

IRENE CARDOSO

Museu Nacional / UFRJ, Quinta da Boa Vista, 20940-040 - Rio de Janeiro - RJ, Brazil.

Table of contents

Abstract	2
Introduction	2
Material and methods	3
Systematics	3
Family Anchistioitidae Borradaile, 1915	3
<i>Anchistiooides</i> Paulson, 1875	3
<i>Anchistiooides antiquensis</i> (Schmitt, 1924)	6
Family Bresiliidae Calman, 1896	10
<i>Pseudocheles</i> Chace & Brown, 1978	10
<i>Pseudocheles chacei</i> Kensley, 1983	10
Family Hippolytidae Dana, 1852	15
<i>Latreutes</i> Stimpson, 1860	15
<i>Latreutes fucorum</i> (Fabricius, 1798)	16
<i>Trachycaris</i> Calman, 1906	21
<i>Trachycaris restricta</i> (A. Milne Edwards, 1878)	21
Family Palaemonidae Rafinesque, 1815	26
Subfamily Palaemoninae Rafinesque, 1815	26
<i>Brachycarpus</i> Bate, 1888	26
<i>Brachycarpus biunguiculatus</i> (Lucas, 1846)	27
<i>Leander</i> Desmarest, 1849	27
<i>Leander tenuicornis</i> (Say, 1818)	27
Subfamily Pontoniinae Kingsley, 1878	28
<i>Pontonia</i> Latreille, 1829	28
<i>Pontonia manningi</i> Fransen, 2000	28
Pasiphaeidae Dana, 1852	34
<i>Leptochela</i> Stimpson, 1860	34
<i>Leptochela serratorbita</i> Bate, 1888	34

Family Processidae Ortmann, 1890	40
<i>Processa</i> Leach, 1815	40
<i>Processa brasiliensis</i> Christoffersen, 1979	40
References	40

Abstract

The REVIZEE (Survey of the live resources from the Economic Exclusive Zone) Program collected 158 benthic samples, in Brazilian continental shelf and slope, in latitudes ranging from 13°00'S to 22°30'S, and depths ranging from 50 to 500m. In these samples, ten caridean species, distributed in six families were identified: *Latreutes fucorum* (Fabricius, 1798), *Trachycaris restricta* (A. Milne Edwards, 1878) (Hippolytidae); *Anchistiooides antiguensis* (Schmitt, 1924) (Anchistioitidae); *Brachycarpus biunguiculatus* (Lucas, 1849), *Leander tenuicornis* (Say, 1818) (Palaemonidae, subfamily Palaemoninae); *Periclimenaeus bermudensis* (Armstrong, 1940), *Pontonia manningi* Fransen, 2000 (Palaemonidae, subfamily Pontoniinae); *Leptochela serratorbita* Bate, 1888 (Pasiphaeidae); *Processa brasiliensis* Christoffersen, 1979 (Processidae) and *Pseudocheles chacei* Kensley, 1983 (Bresiliidae). Nine of these species are herein redescribed and compared with data from the literature. Of these, *Pseudocheles chacei* is a new record for the Brazilian coast. *Anchistiooides antiguensis*, *P. manningi*, *L. serratorbita* and *P. brasiliensis* had their latitudinal distribution expanded to the study area.

Key words: Caridea, Brazil, continental shelf, slope

Introduction

The Infraorder Caridea Dana, 1852, comprises 36 families, 16 of which occur in Brazilian waters (Ramos-Porto & Coelho, 1998; Martin & Davis, 2001). In the present study, nine species from six families were included: *Anchistiooides antiguensis* (Schmitt, 1924), Anchistioitidae Borradaile, 1915; *Pseudocheles chacei* Kensley, 1983, Bresiliidae Calman, 1896; *Latreutes fucorum* (Fabricius, 1798) and *Trachycaris restricta* (A. Milne Edwards, 1878), Hippolytidae Bate, 1888; *Brachycarpus biunguiculatus* (Lucas, 1849) and *Leander tenuicornis* (Say, 1818), Palaemonidae, subfamily Palaemoninae Rafinesque, 1815; *Pontonia manningi* Fransen, 2000, Palaemonidae, subfamily Pontoniinae Kingsley, 1878; and *Processa brasiliensis* Christoffersen, 1979, Processidae Ortmann, 1890 (Holthuis, 1993; Ramos-Porto & Coelho, 1998).

This study describes the caridean shrimps (except Alpheidae) collected by the Revizee Benthos Program. Three species collected by the Revizee Benthos Program were collected at Rocas Atoll: *B. biunguiculatus* and *L. tenuicornis* and *P. brasiliensis*. These species were redescribed recently by Cardoso & Young (*in press*), so, they were only cited with a brief diagnosis given.

The six species redescribed here were previously recorded in Brazilian waters, but