
Micranthes hitchcockiana

Saxifragaceae Saddle Mountain saxifrage, Hitchcock's saxifrage









Perennial herb occasionally from thick, ascending rhizomes at least 5 mm long. **Stems** thick and fleshy, unbranched. **Leaves** 4-12(15) cm long, densely covered with white, multicellular hairs, elliptic to obovate, tapering gradually to a broad, indistinct petiole, the margin usually strongly serrate to dentate, ciliate; leafless flowering stems 10-25 cm tall, densely pubescent. **Inflorescence** an open, broadly cymose panicle, 5-10 cm tall, densely covered with wavy multicellular glandular hairs ca. 2 mm long; main axis with 4-8 lateral branches each 1-4 cm long; pedicels 1-2 mm long, strongly reflexed; bracts elliptic to ovate, green. **Flowers** with petals 2-5 mm long, more than 1.5 times the length of the sepals, elliptic, white; stamens ~equal to petal length, filaments more subulate than clavate. **Fruits** 2.5-6mm long, greenish to reddish-purple.

Lookalikes Saxifraga cespitosa Micranthes oregana

differs from featured plant by smaller, lobed leaves


inflorescence pyramidal and open with spreading lower branches

Micranthes hitchcockiana (Elvander) Brouillet & Gornall

Saddle Mountain saxifrage, Hitchcock's saxifrage

PLANTS symbol: SAHI7

August 2019 status Federal:SOC; Oregon:C; ORBIC: List 1



Distribution: Endemic to higher peaks in northwest Oregon Coast Range.

Habitat: Grassy balds, thin, rocky soils and rock crevices.

Elevation: 670-1020 m

Best survey time(in flower): late May-July

Associated species:

Abies amabilis (Pacific silver fir) Tsuga heterophylla (Western hemlock) Iris tenax (Oregon iris) Lomatium martindalei (Few fruited Iomatium) Saxifraga cespitosa (Tufted alpine saxifrage)







1 cm





