



Taxonomic Paper

Snake richness in urban forest fragments from Niterói and surroundings, state of Rio de Janeiro, southeastern Brazil

Nathalie Citeli[†], Breno Hamdan^{§,‡}, Thais Guedes[|]

[‡] Laboratório de Coleções Biológicas e Biodiversidade, Instituto Vital Brazil, Rua Maestro José Botelho, 64, 24230-410, Niterói, Brazil

[§] Laboratório de Biologia Evolutiva Teórica e Aplicada, Departamento de Genética, Instituto de Biologia, UFRJ, Ilha do Fundão, Rio de Janeiro, Brazil

[|] University of Gothenburg. Department of Biological and Environmental Sciences. Carl Skottsberg gata 22B, Se 405 30, Göteborg, Sweden

Corresponding author: Thais Guedes (thaisbguedes@yahoo.com.br)

Academic editor: Truong Nguyen

Received: 10 Nov 2015 | Accepted: 27 Jan 2016 | Published: 04 Feb 2016

Citation: Citeli N, Hamdan B, Guedes T (2016) Snake richness in urban forest fragments from Niterói and surroundings, state of Rio de Janeiro, southeastern Brazil. Biodiversity Data Journal 4: e7145. doi: [10.3897/BDJ.4.e7145](https://doi.org/10.3897/BDJ.4.e7145)

Abstract

Background

The Atlantic Forest is a hotspot for biodiversity, an area which houses high species richness and endemism, but with high level of threat. With reference to the herpetofauna, until recently there has been no detailed information regarding diversity of snakes recorded in the Atlantic Forest, the number of endemic species and their distribution ranges. While these basic data were missing, large areas of original forest have continued to be lost to increased urbanization and human population, representing a severe threat to the biodiversity.

New information

We recorded 28 snake species in our study area. Dipsadidae was the richest family with 14 species, followed by Colubridae (eight species), Boidae (two species), Viperidae (two species), and Anomalepididae, Elapidae and Typhlopidae (one species each). Most species were forest inhabitants (61%), of which 13 are endemic to the Atlantic Forest. There were no clearly defined species clusters regarding species composition. None of the species recorded in Niterói are listed as threatened in the Brazilian Redlist. However, most of them are strongly associated with forested areas and, perhaps, are not adapted to live in small fragments. Thus, more initiatives should be implemented to evaluate the true conservation status of these species in order to better protect them.

Keywords

Urban Inventory, Serpentes, Species Richness, Forested Areas, Endemism, Conservation

Introduction

The Brazilian Atlantic Forest is one of the largest Neotropical rainforests originally covering around 150 million ha (Tabarelli et al. 2005, Ribeiro et al. 2009). This forest has one of the highest species richness and endemism levels on Earth (Myers et al. 2000). However, it is highly threatened (Ribeiro et al. 2009) which, along with its high endemism, makes it a biodiversity hotspot for conservation (Myers et al. 2000). Recent data suggested that the Atlantic Forest has already lost more than 84 to 89% of its area, while the remaining 100,000 km² comprises only small isolated fragments (Tabarelli et al. 2005, Ribeiro et al. 2009). These fragments may still house many unknown species, with several new snake species having been described recently (Lewinsohn and Prado 2005, Passos et al. 2010, Pires et al. 2014).

The future of the biodiversity in the Atlantic forest is under threat, since deforestation and habitat fragmentation impose barriers for individual migration, leading to the decrease and isolation of population and extinction in the long term (Tabarelli et al. 2005, Becher et al. 2007). The increase in urbanization and a growing human population, represent a considerable pressure to this area's biodiversity (Nations 2008). Niterói is a city on the coast of Rio de Janeiro, Brazil. About 23% of the state of Rio de Janeiro is covered by Atlantic Forest (SOS Mata Atlântica and INPE 2014). Recent studies have surveyed the herpetofauna of particular areas of the state and provided lists of snake species (e.g. Rocha et al. 2000, Rocha et al. 2004, Izecksohn and Carvalho-e-Silva 2001, Rocha and Van Sluys 2006, Almeida-Gomes et al. 2008, Almeida-Gomes et al. 2010, Pontes et al. 2009, Pontes and Rocha 2008, Vrcibradic et al. 2011, Martins et al. 2012), but urban areas remain poorly studied (Rocha et al. 2004). Certainly, no previous snake survey has been conducted in the municipality of Niterói.

Here we provide the first account of snake species richness in the urban area of Niterói. We compare our results with other surveys carried out in nearby areas in Rio de Janeiro and the Atlantic Forest. This study provides basic knowledge about diversity, geographic distribution, endemism, and ecology of species that could help in future conservation planning (Whittaker et al. 2005).

Materials and methods

Study area

The city of Niterói ($43^{\circ}06'13''\text{W}$, $22^{\circ}53'00''\text{S}$; Fig. 1) is currently the fifth most populous urban area of the state of Rio de Janeiro, housing about 500,000 inhabitants in an area of $133,000 \text{ km}^2$ (IBGE 2014). The city is located in the Atlantic Forest Domain, with altitude varying from sea level up to 412 m in the Elefante Mountain Range (CEPERJ 2013, Pontes 1987). About 99% of the city's original area was originally covered by the Atlantic Rain Forest, but vegetation cover today is reduced to about 23% (Ab'Saber 1977, Rizzini 1979, IBGE 1992, SOS Mata Atlantica and INPE 2014).




Figure 1.

Map showing the localization of the municipality of Niterói in the Atlantic Forest Domain and in the state of Rio de Janeiro (featured area).

The climate in Niterói is of the Köppen's type Aw Tropical, with high temperatures, rainy summers, and dry winters (Köppen 1936). Relative humidity is 79.1%, mean annual maximum and minimum temperatures are 27.3°C and 21°C , respectively, with extremes recorded of 10.1°C and 42°C (INMET 1992). The mean annual rainfall is 98 mm (Sentelhas et al. 2003) and the main sources of fresh water are the Macacu and Guapiaçu rivers (Águas 2010).

Data collection

Species identification was based on the specimens examined by direct analysis of voucher specimens (Suppl. material 1) collected between 1956 and 2014 and housed in the following scientific collections: Museu Nacional da Universidade Federal do Rio de Janeiro (MNRJ), Coleção Científica de Serpentes do Instituto Vital Brazil (IVB), Coleção de Répteis da Universidade Federal do Rio de Janeiro (ZUFRJ), and Museu de Zoologia da Universidade de Campinas (ZUEC-REP). Museum acronyms are taken from Sabaj Pérez 2014, except IVB.

Specimens were examined by the authors and identified based on current and reliable taxonomic literature (e.g., Peters and Orejas-Miranda 1970, Peters 1960, Dixon and Kofron 1984, Dixon 1989, Dixon and Markezich 1992, Dixon et al. 1993, Franco and Ferreira 2002, Campbell and Lamar 2004, Hedges et al. 2014). The systematic arrangement followed Zaher et al. 2009 and nomenclature follows Costa and Bérnls 2014. We also provide additional information about habitat use and diet for each species based on data from the literature (Almeida-Gomes et al. 2008, Argôlo 2004, Gaiarsa et al. 2013, Hartmann et al. 2009, Marques et al. 2004, Vrcibradic et al. 2011, Pontes and Rocha 2008).

Analysis

We performed a dissimilarity analysis with Jaccard's coefficient and centroid clustering method of the species presence/absence matrix to compare our species list with others from the Atlantic Forest of the state of Rio de Janeiro. The analysis was implemented in the package *vegan* (Oksanen et al. 2015) of the R software (R Core Team 2014). The raw data used to perform the similarity analysis is provided in Suppl. material 2.

Data resources

We recorded 28 snake species from 18 genera and seven families (see Checklist, Suppl. material 1). The most speciose family was Dipsadidae (14 species), followed by Colubridae (eight species), Boidae (two species), Viperidae (two species), Anomalepididae and Typhlopidae (one species each). The most common species found in the collections were *Bothrops jararacussu* (103 specimens), followed by *Bothrops jararaca* and *Micrurus corallinus* (78 and 32 specimens, respectively). The least common were *Chironius exoletus*, *Erythrolamprus aesculapii*, *Leptophis ahaetulla*, and *Liophidium wilderi* (one specimen each). We also established a first record from Niterói for *Clelia plumbea*, though it also seems to be very rare.

Most species are forest inhabitants (17 species, 61%), with a smaller percentage (39%) inhabiting both forested and open areas. All of them are typical of the Altantic Forest Domain. However, none of them exclusively inhabit open areas. The most species rich locality is the Serra do Mar with 60 species, followed by Duque de Caxias with 33 species

(Fig. 2). We also recorded thirteen endemic Atlantic Forest species. The species composition is most similar to those of Duque de Caxias, Serra do Medanha and Estação Ecológica do Paraíso. However, the analysis did not recover distinct species clusters for the snakes in the Atlantic Forest of the state of Rio de Janeiro (see Fig. 3). All species recorded were listed as Least Concern (LC) in the Brazilian National Redlist (ICMBio 2014) since they are common in the Atlantic Forest and have a wide distribution range (see Checklist).




Figure 2.

Richness of snake species in the Atlantic Forest of Niterói and other regions of the state of Rio de Janeiro (Raw data available in Suppl. material 2).




Figure 3.

Dendrogram representing a cluster analysis, with Jaccard's coefficient and centroid clustering method showing the taxonomic similarities between the Atlantic Forest of Niterói and Atlantic Forest in other municipalities of Rio de Janeiro (Raw data available in Suppl. material 2).

List of species of snakes recorded in the municipality of Niterói, state of Rio de Janeiro, Brazil in this study

Liophidium wilderi (Garman, 1883)

Conservation status: Least Concern

Distribution: Recorded in forested and open areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: Endemic of the Atlantic Forest (Fig. 4). It is fossorial, nocturnal and diurnal, and feeds on arthropods.



Figure 4.

Liophidium wilderi (Anomalepididae) from Niterói, state of Rio de Janeiro, Brazil.

Amerotyphlops brongersmianus* (Vanzolini, 1976)*Conservation status:** Least Concern**Distribution:** Recorded in forested areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil**Notes:** It is fossorial, nocturnal and diurnal, and feeds on arthropods (Fig. 5).

Figure 5.

Amerotyphlops brongersmianus (Typhlopidae).

***Boa constrictor* Linnaeus, 1758**

Conservation status: Least Concern

Distribution: Recorded in forested and open areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: It is semi-arboreal, diurnal and nocturnal and feeds mostly on vertebrates (birds and mammals) (Fig. 6).



Figure 6.

Boa constrictor (Boidae) from Niterói, state of Rio de Janeiro, Brazil.

Corallus hortulanus* (Linnaeus, 1758)*Conservation status:** Least Concern**Distribution:** Recorded in forested areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil**Notes:** It is arboreal, nocturnal and feeds mostly on vertebrates (birds and mammals) (Fig. 7).

Figure 7.

Corallus hortulanus (Boidae). Photo: G. Jones.

***Chironius bicarinatus* (Wied, 1820)**

Conservation status: Least Concern

Distribution: Recorded in forested areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: Endemic of the Atlantic Forest (Fig. 8). It is semi-arboreal, diurnal and feeds mostly on frogs.



Figure 8.

Chironius bicarinatus (Colubridae). Photo: D. Loebmann.

Chironius exoletus* (Linnaeus, 1758)*Conservation status:** Least Concern**Distribution:** Recorded in forested areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil**Notes:** It is semi-arboreal, diurnal and feeds mostly on frogs (Fig. 9).

Figure 9.

Chironius exoletus (Colubridae).

***Chironius laevicollis* (Wied, 1824)**

Conservation status: Least Concern

Distribution: Recorded in forested areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: Endemic of the Atlantic Forest (Fig. 10). It is semi-arboreal, diurnal and feeds mostly on frogs.



Figure 10.

Chironius laevicollis (Colubridae) from Niterói, state of Rio de Janeiro, Brazil.

Leptophis ahaetulla* subsp. *ahaetulla* (Wied, 1824)*Conservation status:** Least Concern**Distribution:** Recorded in forested and open areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil**Notes:** It is semi-arboreal, diurnal and feeds on vertebrates (frogs, lizards and birds) (Fig. 11).

Figure 11.

Leptophis ahaetulla (Colubridae). Photo: D. Loebmann.

***Mastigodryas bifossatus* (Raddi, 1820)**

Conservation status: Least Concern

Distribution: Recorded in forested and open areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: It is terrestrial, diurnal and feeds on vertebrates (frogs and mammals) (Fig. 12).



Figure 12.

Mastigodryas bifossatus (Colubridae).

***Spilotes pullatus* subsp. *anomalepis* (Linnaeus, 1758)**

Conservation status: Least Concern

Distribution: Recorded in forested and open areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: Endemic of the Atlantic Forest (Fig. 13). It is arboreal, diurnal and feeds on vertebrates (birds and mammals).



Figure 13.

Spilotes pullatus anomalepis (Colubridae). Photo: W. Pessoa.

***Spilotes sulphureus* subsp. *poecilostoma* (Wied, 1824)**

Conservation status: Least Concern

Distribution: Recorded in forested areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: It is semi-arboreal, diurnal and feeds on vertebrates (frogs, lizards and mammals) (Fig. 14).



Figure 14.

Spilotes sulphureus poecilostoma (Colubridae) from Niterói, state of Rio de Janeiro, Brazil.
Photo: J. L. Pontes.

Oxybelis aeneus* (Wagler in Spix, 1824)*Conservation status:** Least Concern**Distribution:** Recorded in forested and open areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil**Notes:** It is arboreal, diurnal and feeds on vertebrates (lizards) (Fig. 15).

Figure 15.

Oxybelis aeneus (Colubridae) from Niterói, state of Rio de Janeiro, Brazil.

***Clelia plumbea* (Wied, 1820)**

Conservation status: Least Concern

Distribution: Recorded in forested areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: It is terrestrial, nocturnal and feeds on vertebrates (snakes and lizards) (Fig. 16).



Figure 16.

Clelia plumbea (Dipsadidae). Photo: M. A. Freitas.

Elapomorphus quinquelineatus* (Raddi, 1820)*Conservation status:** Least Concern**Distribution:** Recorded in forested areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil**Notes:** Endemic of the Atlantic Forest (Fig. 17). It is fossorial, diurnal and feeds on vertebrates (amphisbaena and snakes).

Figure 17.

Elapomorphus quinquelineatus (Dipsadidae) from Niterói, state of Rio de Janeiro, Brazil.
Photo: J. L. Pontes.

***Erythrolamprus aesculapii* (Linnaeus, 1766)**

Conservation status: Least Concern

Distribution: Recorded in forested areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: It is terrestrial, diurnal and feeds on vertebrates (amphisbaena, snakes and eventually lizards) (Fig. 18)



Figure 18.

Erythrolamprus aesculapii (Dipsadidae). Photo: M. A. Freitas.

***Erythrolamprus miliaris* subsp. *orinus* (Linnaeus, 1758)**

Conservation status: Least Concern

Distribution: Recorded in forested areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: Endemic of the Atlantic Forest (Fig. 19). It is semi-aquatic, diurnal and feeds on vertebrates (frogs and fishes).



Figure 19.

Erythrolamprus miliaris orinus (Dipsadidae) from Niterói, state of Rio de Janeiro, Brazil.

***Erythrolamprus poecilogyrus* subsp. *poecilogyrus* (Wied, 1825)**

Conservation status: Least Concern

Distribution: Recorded in forested and open areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: Endemic of the Atlantic Forest (Fig. 20). It is semi-aquatic, diurnal and feeds on vertebrates (frogs and fishes).



Figure 20.

Erythrolamprus poecilogyrus poecilogyrus (Dipsadidae) from Niterói, state of Rio de Janeiro, Brazil.

Erythrolamprus poecilogyrus* subsp. *schotti* (Wied, 1825)*Conservation status:** Least Concern**Distribution:** Recorded in forested and open areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil**Notes:** It is terrestrial, diurnal and feeds on vertebrates (frogs). (Fig. 21)

Figure 21.

Erythrolamprus poecilogyrus schotti (Dipsadidae) from Niterói, state of Rio de Janeiro, Brazil.

***Helicops carinicaudus* (Wied, 1825)**

Conservation status: Least Concern

Distribution: Recorded in forested areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: It is aquatical, diurnal and feeds on vertebrates (fishes and frogs) (Fig. 22)



Figure 22.

Helicops carinicaudus (Dipsadidae) from Niterói, state of Rio de Janeiro, Brazil.

Oxyrhopus petolarius* subsp. *digitalis* (Reuss, 1834)*Conservation status:** Least Concern**Distribution:** Recorded in forested and open areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil**Notes:** It is terrestrial, nocturnal and feeds on vertebrates (lizards and mammals) (Fig. 23)**Figure 23.***Oxyrhopus petolarius digitalis* (Dipsadidae). Photo: M. A. Freitas

***Oxyrhopus clathratus* Duméril, Bibron & Duméril, 1854**

Conservation status: Least Concern

Distribution: Recorded in forested and open areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: Endemic of the Atlantic Forest (Fig. 24). It is terrestrial, nocturnal and feeds on vertebrates (lizards and mammals).



Figure 24.

Oxyrhopus clathratus (Dipsadidae) from Niterói, state of Rio de Janeiro, Brazil.

***Philodryas olfersii* subsp. *olfersii* (Liechtenstein, 1823)**

Conservation status: Least Concern

Distribution: Recorded in forested and open areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: It is semi-arboreal, diurnal and feeds on vertebrates (lizards, mammals and birds) (Fig. 25)



Figure 25.

Philodryas olfersii olfersii (Dipsadidae). Photo: J. L. Pontes.

***Philodryas patagoniensis* (Girard, 1858)**

Conservation status: Least Concern

Distribution: Recorded in forested and open areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: It is terrestrial, diurnal and feeds on vertebrates (lizards and mammals) (Fig. 26)



Figure 26.

Philodryas patagoniensis (Dipsadidae) from Niterói, state of Rio de Janeiro, Brazil.

Sibynomorphus neuwiedi* (Ihering, 1911)*Conservation status:** Least Concern**Distribution:** Recorded in forested areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil**Notes:** Endemic of the Atlantic Forest (Fig. 27). It is terrestrial, nocturnal and feeds on molluscs.

Figure 27.

Sibynomorphus neuwiedi (Dipsadidae) from Niterói, state of Rio de Janeiro, Brazil.

***Thamnodynastes cf. nattereri* (Mikan, 1828)**

Conservation status: Least Concern

Distribution: Recorded in forested areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: It is semi-arboreal, nocturnal and feeds on vertebrates (frogs) (Fig. 28). See taxonomic notes for the species taxonomic status.



Figure 28.

Thamnodynastes cf. nattereri (Dipsadidae) from Niterói, state of Rio de Janeiro, Brazil.

Xenodon neuwiedii* Günther, 1863*Conservation status:** Least Concern**Distribution:** Recorded in forested and open areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil**Notes:** It is terrestrial, diurnal and feeds on vertebrates (frogs) (Fig. 29)

Figure 29.

Xenodon neuwiedii (Dipsadidae) from Niterói, state of Rio de Janeiro, Brazil.

***Micrurus corallinus* (Merrem, 1820)**

Conservation status: Least Concern

Distribution: Recorded in forested areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: Endemic of the Atlantic Forest (Fig. 30). It is fossorial, diurnal and nocturnal and feeds on vertebrates (lizards and snakes).



Figure 30.

Micrurus corallinus (Elapidae) from Niterói, state of Rio de Janeiro, Brazil.

Bothrops jararaca* (Wied, 1824)*Conservation status:** Least Concern**Distribution:** Recorded in forested areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil**Notes:** Endemic of the Atlantic Forest (Fig. 31). It is terrestrial, nocturnal and feeds on vertebrates (mammals, frogs and lizards).

Figure 31.

Bothrops jararaca (Viperidae) from Niterói, state of Rio de Janeiro, Brazil.

***Bothrops jararacussu* Lacerda, 1884**

Conservation status: Least Concern

Distribution: Recorded in forested areas of the Atlantic Forest. Municipality of Niterói. State of Rio de Janeiro. Brazil

Notes: It is terrestrial, nocturnal and feeds on vertebrates (mammals and frogs) (Fig. 32).



Figure 32.

Bothrops jararacussu (Viperidae) from Niterói, state of Rio de Janeiro, Brazil.

Discussion

The species richness found in Niterói represents 34% of the snake species known for the state of Rio de Janeiro (Rocha et al. 2004) and 7.3% of the species known for Brazil (Costa and Bérnails 2014). Additionally, the richness recorded corresponds to 47% of the snake species occurring in the southern portion of the Atlantic Forest (Marques et al. 2004). The richness found in Niterói is similar to that reported for other localities in the Atlantic Forest, such as Viçosa (Minas Gerais; 27 species; Costa et al. 2010), Duque de Caxias (Rio de Janeiro, 33 spp., Salles and Silva-Soares 2010), Serra do Medanha (Rio de Janeiro, 27 spp., Pontes and Rocha 2008) and Vitória (Espírito Santo, 27 spp., Silva-Soares et al. 2011). The surveys in protected areas had lower richness (e.g., Martins et al. 2012) when compared with our study. This is probably an artifact of the size of the area or might be a consequence of the environmental education program developed by the Vital Brazil Institute that receive venomous snakes species from the population for snakebite serum production (it also justify the higher abundance of *Bothrops jararaca* and *Micrurus corallinus* in our samples).

Information about diversity, taxonomy, and geographical distribution of species from scientific collections can be used for a broad range of purposes (Graham et al. 2004), including conservation. Thus, Brazilian scientific collections play a key role in recording part of the country's biodiversity. For example, by analyzing specimens housed in scientific collections we were able to recover the snake richness of an area of Atlantic Forest that is today primarily covered by a large city.

Currently, no complete snake species list for the entire Atlantic Forest, let alone a list of the potentially threatened endemic species but, based on the literature (e.g., Cunha and Nascimento 1993, Cunha and Nascimento 1978, Marques et al. 2004, Strussmann and Sazima 1993), it seems that 13 species are endemic to this biome (see checklist). Most of the species (17 species, 61%) are strictly forest inhabitants (e.g., species of *Chironius* and *Bothrops*), and eleven (39%) inhabit both open and forested areas (e.g., genus *Philodryas*; Argôlo 2004, Hartmann et al. 2009, Marques et al. 2009, Costa et al. 2010, Gaiarsa et al. 2013). Our results suggest that the Atlantic Forest remnants in Niterói are important for maintaining the high snake diversity, since most species are not adapted to live in urban environments (Marques et al. 2009).

The snake fauna of Niterói is similar to the nearby localities of Duque de Caxias, Serra do Medanha, and Estação Ecológica do Paraíso. This suggests that there is little turnover in snake species composition between lowland Atlantic Forest areas of the state of Rio de Janeiro (Salles and Silva-Soares 2010). It seems likely that the areas surrounding the Bay of Guanabara share the same species, and perhaps the same evolutionary history.

None of the species recorded were listed as threatened (ICMBio 2014). However, most of them are strongly associated with forested areas and are not adapted to live in small fragments, forest edges, or urban areas (Argôlo 2004, Marques et al. 2009). The Brazilian national redlist takes into account the whole species range and the occurrence of each

species in protected areas. But the conservation status in each region can be different (e.g., Bressan et al. 2009).

Most of the protected areas in Brazil have less than 500 ha, an inadequate size to maintain their biota (Viana and Pinheiro 1998). Thus, strategies for biodiversity conservation should expand the boundaries of protected areas as part of the large potential area to be protected (Viana and Pinheiro 1998). The municipal and state protected areas are apparently effective in preserving this high snake richness in the study region (Viana and Pinheiro 1998). However, increased urbanization rate and population put the survival of species at risk and may considerably alter patterns of species richness and composition (e.g., Viviani et al. 2010). There is an increasing trend towards deforestation in Niterói due to the construction of ports and houses around the forest fragments in the Bay of Guanabara. This removal of vegetation cover could exert considerable pressure on these Atlantic Forest remnants over time. Thus, unprotected forested areas in and around Niterói (e.g., Maricá, São Gonçalo Saquarema, Araruama, Rio Bonito, and Itaboraí) should be protected. Our study reinforces the importance of forested areas in Niterói for protecting species diversity. We suggest that public policies related to environmental education should be implemented, as well as the expansion of protected areas and the creation of new ones.

Taxonomic notes

Erythrolamprus poecilogyrus schotti (IVB 1399, 2010) is widely distributed in the Caatinga and Cerrado, with dubious records in the Atlantic Forest of Rio de Janeiro (Dixon and Markezich 1992). The specimens analyzed here have diagnostic characteristics of this subspecies, namely: 19-19-15 of dorsal rows, 154-161 ventral scales, eight supralabials, nine infralabials, and 53-56 subcaudals. Therefore, we confirm that this subspecies occurs in Niterói.

The specimens of *Thamnodynastes* from Niterói have dorsal scales weakly keeled, ventral portion of the head immaculate, longitudinal stripes little conspicuous, posterior region of the vent darker, 3rd and 4th supralabials in contact with the orbits, 19 dorsal rows, loreal present, ventrals ranging from 141 to 165 in males and from 147 to 162 in females. These characteristics correspond to *Thamnodynastes* cf. *nattereri* (Franco and Ferreira 2002). The taxonomy of *Thamnodynastes* cf. *nattereri* is complicated, since the type material is lost (F. Franco pers. comm.). This species is morphologically similar to *T. hypoconia* that also occurs in the Atlantic Forest of Rio de Janeiro. However, it cannot be treated as *T. hypoconia* because the later has dorsal scales strongly keeled, anterior and posterior region of the vent homogeneous, with longitudinal stripes very conspicuous, and dark gular region (Franco and Ferreira 2002).

Acknowledgements

The authors are grateful to all curators of the institutions visited: Paulo Passos (MNRJ), Daniel Fernandes da Silva (UFRJ) and Paulo Manzani (ZUEC-REP). We thanks to Daniel Fernandes (UFRJ) and Barbara Francisco (MNRJ) for confirm the identification of some snakes species. We are grateful to Mark O'Shea (Univ. Wolverhampton) for valuable comments on the manuscript. This research was supported by Vital Brazil Institute. Diogo B. Provete and Anne d'Heursel-Baldisseri reviewed the English language. T. B. Guedes received post-doctoral fellowships (Grants Nº 2013/04170-8 and 2014/18837-7) from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP).

References

- Ab'Saber AN (1977) Os domínios morfoclimáticos da América do Sul. *Geomorfologia* 52: 1-21.
- Águas AND (2010) Atlas Brasil Abastecimento Urbano de Água. <http://atlas.ana.gov.br/atlas/forms/analise/RegiaoMetropolitana.aspx?rme=18>. Accession date: 2015 6 02.
- Almeida-Gomes M, Almeida-Santos M, Goyannes-Araújo P, Borges-Júnior VNT, Vrcibradic D, Siqueira CC, Ariani CV, Dias AS, Souza VV, Pinto RR, Van Sluys M, Rocha CFD (2010) Anurofauna of an Atlantic Rainforest fragment and its surroundings in Northern Rio de Janeiro State, Brazil. *Brazilian Journal of Biology* 70: 871-877. DOI: [10.1590/s1519-69842010000400018](https://doi.org/10.1590/s1519-69842010000400018)
- Almeida-Gomes M, Vrcibradic D, Siqueira CC, Kiefer MC, Klaion T, Almeida-Santos P, Nascimento D, Ariani CV, Borges-Junior VNT, Freitas-Filho RF, Van Sluys M, Rocha CFD (2008) Herpetofauna of an Atlantic rainforest area (Morro São João) in Rio de Janeiro State, Brazil. *Anais da Academia Brasileira de Ciências* 80: 291-300. DOI: [10.1590/s0001-37652008000200007](https://doi.org/10.1590/s0001-37652008000200007)
- Argôlo AJS (2004) As serpentes dos cacauais do Sudeste da Bahia. 1. Editora da UESC, 260 pp.
- Becher CG, Fonseca CR, Haddad CFB, Bastista RF, Prado PI (2007) Habitat Split and the global decline of amphibians. *Science* 318: 1775-1777. DOI: [10.1126/science.1149374](https://doi.org/10.1126/science.1149374)
- Bressan PM, Kierulf MCM, Sugieda AM (2009) Fauna ameaçada de extinção do estado de São Paulo: Vertebrados. Fundação Parque Zoológico de São Paulo, São Paulo, 645 pp.
- Campbell JA, Lamar WW (2004) *The Venomous Reptiles of Latin America*. Comstock Publishing Associates, 976 pp.
- CEPERJ F (2013) Anuário Estatístico do Estado do Rio de Janeiro 2013. Centro Estadual de Estatísticas Pesquisas e Formação de Servidores Públicos do Rio de Janeiro. Rio de Janeiro, RJ, Brasil. http://www.ceperj.rj.gov.br/ceep/ent/anu_online.html. Accession date: 2015 6 10.
- Costa HC, Bérnails RS (2014) Brazilian reptiles – List of species. Sociedade Brasileira de Herpetologia. <http://www.sbherpetologia.org.br/checklist/repteis.htm>. Accession date: 2015 6 21.

- Costa HC, Pantoja DL, Pontes JL, Feio RN (2010) Serpentes do Município de Viçosa, Mata Atlântica do Sudeste do Brasil. *Biota Neotropica* 10: 353-377. DOI: [10.1590/s1676-06032010000300033](https://doi.org/10.1590/s1676-06032010000300033)
- Cunha OR, Nascimento FP (1978) Ofídios da Amazônia. X – As cobras da região Leste do Pará. *Bol. Boletim do Museu Paraense Emilio Goeldi* 31: 1-218.
- Cunha OR, Nascimento FP (1993) Ofídios da Amazônia. As cobras da região leste do Pará. *Bol. Mus. Boletim do Museu Paraense Emilio Goeldi* 9: 1-191.
- Dixon JR (1989) A key and checklist to the Neotropical snake genus *Liophis* with country lists and maps. *Smithsonian Herpetol. Inf. Serv.* 79: 1-40. DOI: [10.5479/si.23317515.79.1](https://doi.org/10.5479/si.23317515.79.1)
- Dixon JR, Kofron CP (1984) The Central and South American anomalepid snakes of the genus *Liophlops*. *Amphibia-Reptilia* 4 (2): 241-264. DOI: [10.1163/156853883X00120](https://doi.org/10.1163/156853883X00120)
- Dixon JR, Markezich AL (1992) Taxonomy and geographic variation of *Liophis poecilogyrus* (Wied) from South America (Serpentes: Colubridae). *The Texas Journal of Science* 44: 131-166.
- Dixon JR, Wiest JA, Cei JM (1993) Revision of the tropical snake *Chironius Fitzinger* (Serpentes, Colubridae). *Monogr. Mus. Reg. Sci. Natur.* 13: 1-279.
- Franco FL, Ferreira TG (2002) Descrição de uma nova espécie de *Thamnodynastes Wagler, 1830* (Serpentes, Colubridae) do nordeste brasileiro, com comentários sobre o gênero. *Phyllomedusa* 1: 57-74. DOI: [10.11606/issn.2316-9079.v1i2p57-74](https://doi.org/10.11606/issn.2316-9079.v1i2p57-74)
- Gaiarsa MP, Alencar LRV, Martins M (2013) Natural history of Pseudoboine snakes. *Papéis Avulsos de Zoologia* 53 (19): 261-283. DOI: [10.1590/s0031-10492013001900001](https://doi.org/10.1590/s0031-10492013001900001)
- Graham CH, Ferrier S, Huettman F, Moritz C, Peterson AT (2004) New developments in museum-based informatics and applications in biodiversity analysis. *Trends in ecology and evolution* 19: 497-503. DOI: [10.1016/j.tree.2004.07.006](https://doi.org/10.1016/j.tree.2004.07.006)
- Hartmann PA, Hartmann MT, Martins M (2009) Ecologia e história natural de uma taxocenose de serpentes no Núcleo Santa Virgínia do Parque Estadual da Serra do Mar, no sudeste do Brasil. *Biota Neotropica* 9 (3): 173-184. DOI: [10.1590/s1676-06032009000300018](https://doi.org/10.1590/s1676-06032009000300018)
- Hedges SB, Marion AB, Lipp KM, Marin J, Vidal N (2014) A taxonomic framework for Typhlopidae snakes from the Caribbean and other regions (Reptilia, Squamata). *Caribbean Herpetology* 49: 1-61.
- IBGE (1992) Manual técnico da vegetação brasileira, Manuais técnicos em geociências. IBGE, Rio de Janeiro, 166 pp.
- IBGE (2014) Informações Estatísticas 2014. Instituto Brasileiro de Geografia e Estatística. <http://cod.ibge.gov.br/233F4>. Accession date: 2015 6 03.
- ICMBio (2014) Lista das espécies ameaçadas de extinção do Brasil. <http://www.icmbio.gov.br/portal/comunicacao/noticias/4-destaques/6658-mma-e-icmbio-divulga-novas-listas-de-especies-ameacadas-de-extincao.html>. Accession date: 2015 10 07.
- INMET (1992) Normais Climatológicas 1961 – 1990. Instituto Nacional de Meteorologia, Brasília, 84 pp.
- Izecksohn E, Carvalho-e-Silva SP (2001) Anfíbios do Município do Rio de Janeiro. Editora UFRJ, Rio de Janeiro, 148 pp.
- Köppen W (1936) Das geographische System der Klimate. In: Köppen W, Geiger R (Eds) *Handbuch der Klimatologie*. Gebrüder Borntraeger, Berlin.

- Lewinsohn T, Prado PI (2005) How many species are there in Brazil. *Conservation Biology* 19 (3): 619-624. DOI: [10.1111/j.1523-1739.2005.00680.x](https://doi.org/10.1111/j.1523-1739.2005.00680.x)
- Marques OAV, Eterovic A, Sazima I (2004) Snakes of the Brazilian Atlantic Forest: an illustrated field guide for the Serra do Mar range. 1. Holos Editora, Ribeirão Preto, 204 pp.
- Marques OAV, Pereira DN, Barbo FE, Germano VJ, Sawaya JR (2009) Os Répteis do Município de São Paulo: diversidade e ecologia da fauna pretérita e atual. *Biota Neotropica* 9 (2): 139-150. DOI: [10.1590/s1676-06032009000200014](https://doi.org/10.1590/s1676-06032009000200014)
- Martins AR, Bruno SF, Navegantes AQ (2012) Herpetofauna of Núcleo Experimental de Iguaba Grande, Rio de Janeiro state, Brazil. *Brazilian Journal of Biology* 72: 553-562. DOI: [10.1590/S1519-69842012000300018](https://doi.org/10.1590/S1519-69842012000300018)
- Myers N, Mittermeyer RA, Mittemeyer CG, Fonseca GAB, Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature* 403: 853-858. DOI: [10.1038/35002501](https://doi.org/10.1038/35002501)
- Nations U (2008) The Millennium Development Goals Report. United Nations Publications, 72 pp.
- Oksanen JF, Blanchet G, Kindt R, Legendre P, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Wagner H (2015) Vegan: Community Ecology Package. R package version 2.2-1. <http://CRAN.R-project.org/package=vegan>. Accession date: 2015 6 08.
- Passos P, Fernandes R, Bérnuls R, Moura-Leite Jd (2010) Taxonomic revision of the Brazilian Atlantic Forest *Atractus* (Reptilia: Serpentes: Dipsadidae). *Zootaxa* 2364: 1-63.
- Peters JA (1960) The snakes of the subfamily Dipsadinae . *Misc. publ. - Mus. Zool., Univ. Mich* 114: 1-224.
- Peters JA, Orejas-Miranda B (1970) Catalogue of the neotropical squamata. Part I. Snakes. *Bull. U. S. Natl. Mus* 297: 1-347. DOI: [10.5962/bhl.title.46653](https://doi.org/10.5962/bhl.title.46653)
- Pires MG, Jr. NJdS, Feitosa DT, Prudente ALdC, Pereira GA, Zaher H (2014) A new species of triadal coral snake of the genus *Micrurus* Wagler, 1824 (Serpentes: Elapidae) from northeastern Brazil. *Zootaxa* 3811 (4): 569-584. DOI: [10.11646/zootaxa.3811.4.8](https://doi.org/10.11646/zootaxa.3811.4.8)
- Pontes JAL (1987) Serra da Tiririca, RJ: necessidade de conservação (Primeira Contribuição). *Bol. FBCN* 22: 89-94.
- Pontes JAL, Rocha CFD (2008) Serpentes da Serra do Mendenha, Rio de Janeiro, RJ: ecologia e conservação. Technical Books Editora Ltda, Rio de Janeiro, 147 pp.
- Pontes JAL, Pontes RC, Rocha CFD (2009) The snake community of Serra do Mendenha, in Rio de Janeiro State, southeastern Brazil: composition, abundance, richness and diversity in areas with different conservation degrees. *Brazilian Journal of Biology* 69 (3): 795-804. DOI: [10.1590/s1519-69842009000400006](https://doi.org/10.1590/s1519-69842009000400006)
- R Core Team (2014) A language and environment for statistical computing. <http://www.R-project.org>. Accession date: 2015 6 08.
- Ribeiro MC, Metzger JP, Martensen AC, Ponsoni F, Hirota MM (2009) Brazilian Atlantic forest: how much is left and how is the remaining forest distributed? *Biological Conservation* 142: 1141-1153. DOI: [10.1016/j.biocon.2009.02.021](https://doi.org/10.1016/j.biocon.2009.02.021)
- Rizzini CT (1979) Tratado de Fitogeografia do Brasil. Aspectos Sociológicos e Florísticos. HUCITEC, São Paulo, 374 pp.
- Rocha CFD, Van Sluys M (2006) New records of reptiles from Ilha Grande Island in Rio de Janeiro state, Brazil. *Herpetological review* 37 (1): 112-114.

- Rocha CFD, Araújo AFB, Vrcibradic D, Costa EM (2000) New *Cnemidophorus* (Squamata: Teiidae) from coastal Rio de Janeiro State, Southeastern Brazil. Copeia 2: 501-509. DOI: [10.1643/0045-8511\(2000\)000\[0501:NCSTFC\]2.0.CO;2](https://doi.org/10.1643/0045-8511(2000)000[0501:NCSTFC]2.0.CO;2)
- Rocha DFD, Bergallo HG, Pombal Júnior JP, Lena G, Van Sluys MV, Fernandes R, Caramaschi (2004) Fauna de anfíbios, répteis e mamíferos do estado do Rio de Janeiro, Sudeste do Brasil. Publicações Avulsas do Museu Nacional 104: 3-23.
- Sabaj Pérez MH (2014) Standard symbolic codes for institutional resource collections in herpetology and ichthyology: an Online Reference. <http://www.asih.org>. Accession date: 2016 1 19.
- Salles ROL, Silva-Soares T (2010) Répteis do município de Duque de Caxias, Baixada Fluminense, Rio de Janeiro, Sudeste do Brasil. Biotemas 23 (2): 135-144.
- Sentelhas PS, Marin FR, Ferreira AS, Santos de Sá EJ (2003) Banco de dados climáticos do Brasil. <http://www.bdclima.cnpm.embrapa.br/index.php>. Accession date: 2015 6 08.
- Silva-Soares T, Ferreira RB, Salles RDOL, Rocha CFD (2011) Continental, insular and coastal marine reptiles form the municipality of Vitória, state of Espírito Santo, Southeastern Brazil. Check List 7 (3): 290-298.
- SOS Mata Atlântica, INPE (2014) Atlas dos remanescentes florestais da Mata Atlântica - Período de 2012- 2013. Fundação SOS Mata Atlântica, 61 pp.
- Strussmann C, Sazima I (1993) The snake assemblage of the Pantanal at Poconé, western Brazil: faunal composition and ecological summary. Studies on Neotropical Fauna and Environment 28 (3): 157-168. DOI: [10.1080/01650529309360900](https://doi.org/10.1080/01650529309360900)
- Tabarelli M, Pinto LP, Silva JMC, Hirota MM, Bedê LC (2005) Desafios e oportunidades para a conservação da biodiversidade na Mata Atlântica brasileira. Megadiversidade 1: 132-138.
- Viana VM, Pinheiro LAFV (1998) Conservação da biodiversidade em fragmentos florestais. Série técnica IPEF 12: 25-42.
- Viviani VR, Rocha MY, Oskar H (2010) Fauna de besouros bioluminescentes (Coleoptera: Elateroidea: Lampyridae; Phengodidae, Elateridae) nos municípios de Campinas, Sorocaba-Votorantim e Rio Claro-Limeira (SP, Brasil): biodiversidade e influência da urbanização. Biota Neotropica 10 (2): 103-116. DOI: [10.1590/s1676-06032010000200013](https://doi.org/10.1590/s1676-06032010000200013)
- Vrcibradic D, Rocha CFD, Kiefer MC, Hatano FH, Fontes AF, Almeida-Gomes M, Van Sluys M (2011) Herpetofauna, Estação Ecológica Estadual do Paraíso, State of Rio de Janeiro, southeastern Brazil. Check List 7 (6): 745-749. DOI: [10.15560/11013](https://doi.org/10.15560/11013)
- Whittaker RJ, Araújo MB, Jepson P, Ladle RJ, Watson JME, Willis KJ (2005) Conservation biogeography: assessment and prospect. Diversity and Distributions 11: 3-23. DOI: [10.1111/j.1366-9516.2005.00143.x](https://doi.org/10.1111/j.1366-9516.2005.00143.x)
- Zaher H, Grazziotin FG, Cadle JE, Murphy RW, Moura-Leite JC, Bonatto SL (2009) Molecular phylogeny of advanced snakes (Serpentes, Caenophidia) with an emphasis on South American Xenodontines: a revised classification and description of new taxa. Papéis Avulsos de Zoologia 49: 115-153. DOI: [10.1590/s0031-10492009001100001](https://doi.org/10.1590/s0031-10492009001100001)

Supplementary materials

Suppl. material 1: Snake richness from urban forest fragments of Niterói and surroundings, state of Rio de Janeiro, Southeastern Brazil

Authors: Nathalie Citeli, Breno Hamdan, Thaís Barreto Guedes

Data type: Appendix 1

Brief description: Voucher numbers of specimens analyzed from the Municipality of Niterói housed in the following scientific collections: Coleção Científica Instituto Vital Brazil (IVB), Museu Nacional da Universidade Federal do Rio de Janeiro (MNRJ), and Museu de Zoologia da Universidade Estadual de Campinas (ZUEC).

Filename: Appendix I-Citeli et al..docx - [Download file](#) (98.19 kb)

Suppl. material 2: Raw data of occurrence of snakes in areas of the Atlantic Forest, Southeastern Brazil

Authors: Nathalie Citeli, Breno Hamdan, Thaís Barreto Guedes

Data type: Binary matrix in Excel

Brief description: Binary matrix that was the base to run the cluster analysis with Jaccard's coefficient and to draw the graphics provided in the results section (Fig. 31 and 32).

Filename: Citeli et al-Snakes.xlsx - [Download file](#) (13.25 kb)