

Table 3-1. Benthic foraminifera identified in the 2005 Tomales Bay samples.

*Ammonia beccarii* (Linné)  
*Bolivina pacifica* Cushman and McCulloch  
*Bolivina spissa* Cushman  
*Bolivina subexcavata* Cushman and Wickenden  
*Bolivina vaughani* Natland  
*Buccella frigida* (Cushman)  
*Bulimina denudata* Cushman and Parker  
*Buliminella elegantissima* (d'Orbigny)  
*Cibicides* sp.  
*Elphidiella hannai* (Cushman and Grant)  
*Elphidium crispum* (Linné)  
*Elphidium excavatum* (Terquem)  
    *Elphidium excavatum* var. *seleyensis* (Heron-Allen and Earland)  
    *Elphidium excavatum* var. *clavatum* Cushman  
*Fissurina cucurbitasema* Loeblich and Tappan  
*Fissurina lucida* (Williamson)  
*Fissurina* spp.  
*Florilus labradoricus* (Dawson)  
*Fursenkoina pontoni* (Cushman)  
*Glabratella* sp.  
*Hopkinsina pacifica* Cushman  
*Miliolinella oblonga* (Montagu)  
*Nonionella basispinata* (Cushman and Moyer)  
*Nonionella stella* Cushman and Moyer  
*Quinqueloculina* sp.  
*Reophax nana* Rhumbler  
*Rotorbinella* sp.  
*Spiroplectammia biformis* (Parker and Jones)  
*Suggrunda eckisi* Natland  
*Textularia earlandi* Phleger  
*Trichohyalus ornatissima* (Cushman)  
*Trochammina charlottensis* Cushman  
*Trochammina inflata* (Montagu)  
*Trochammina kelleetae* Thalmann  
*Trochammina hadai* Uchio  
*Trochammina vesicularis* Göes