

The systematics of the genus *Anapagurus* Henderson, 1886, and a new genus for *Anapagurus drachi* Forest, 1966 (Crustacea: Decapoda: Paguridae)

J. García-Gómez

García-Gómez, J. The systematics of the genus *Anapagurus* Henderson, 1886, and a new genus for *Anapagurus drachi* Forest, 1966 (Crustacea, Decapoda, Paguridae).

Zool. Verh. Leiden 295, 15.xi.1994: 1-131, figs. 1-44.—ISSN 0024-1652/ISBN 90-73239-36-2.

J. García-Gómez, InterAmerican Center, Wolfson Campus, Miami-Dade Community College, 627 S.W. 27th Avenue, Miami, Florida 33135-2966, U.S.A.

Key words: Hermit crabs; *Anapagurus*; systematic revision; new species; symbiotic associations; *Forestopagurus* new genus.

A worldwide systematic review of the genus *Anapagurus* is presented. In addition to the rediagnoses of the genus and all known species, five new species, *A. alboranensis*, *A. adriaticus*, *A. vossi*, *A. atlantidii*, and *A. congolensis*, are described. All 18 species are illustrated and a key to the species is given. Several new diagnostic characters found in the right chelipeds (1- the shape and relative size of the mesial and lateral lobes that flank the dorsoproximal margin of the palm; 2- the shape and relative size of the median sinus of the ventroproximal palmar margin; 3- the shape, armature and relative size of the dorsomesial distal carpal angle; and 4- the shape and extension of both ventrodistal and ventroproximal carpal margins, which determine the relative ventral length); the relative length and setation of the ultimate antennular segment; and the relative length between the right uropodal endopod and the distal protopodal lobe have been used. A number of systematic problems are clarified. *Anapagurus pusillus* Henderson, 1888 is reinstated. *Anapagurus brevicarpus* A. Milne Edwards & Bouvier, 1892 is found to be a junior synonym of *A. chiroacanthus* (Lilljeborg, 1856). *Anapagurus acutus* and *A. marginatus* described by A. Milne Edwards & Bouvier, 1893, are excluded from *Anapagurus* and considered species incertae sedis. *Anapagurus drachi* Forest, 1966, is removed from *Anapagurus*, and placed in a new genus, *Forestopagurus*.

Contents

Introduction	4
Materials and methods	5
Morphological considerations	7
<i>Anapagurus</i> Henderson, 1886	8
Key to the species of <i>Anapagurus</i>	11
<i>Anapagurus laevis</i> (Bell, 1846)	13
<i>Anapagurus hyndmanni</i> (Bell, 1846)	22
<i>Anapagurus chiroacanthus</i> (Lilljeborg, 1856)	29
<i>Anapagurus pusillus</i> Henderson, 1888, sensu stricto	40
<i>Anapagurus japonicus</i> Ortmann, 1892	44
<i>Anapagurus curvidactylus</i> Chevreux & Bouvier, 1892	48
<i>Anapagurus bicorniger</i> A. Milne Edwards & Bouvier, 1892	53
<i>Anapagurus longispina</i> A. Milne Edwards & Bouvier, 1900	58
<i>Anapagurus bonnieri</i> Nobili, 1905	63
<i>Anapagurus breviaculeatus</i> Fenizia, 1937	66
<i>Anapagurus hendersoni</i> Barnard, 1947	72

<i>Anapagurus wolffi</i> Forest, 1961	77
<i>Anapagurus petiti</i> Dechancé & Forest, 1962	81
<i>Anapagurus alboranensis</i> spec. nov.	85
<i>Anapagurus adriaticus</i> spec. nov.	91
<i>Anapagurus vossi</i> spec. nov.	97
<i>Anapagurus atlantidii</i> spec. nov.	104
<i>Anapagurus congolensis</i> spec. nov.	110
Biogeography	116
<i>Forestopagurus</i> gen. nov.	117
Acknowledgements	120
References	121
Index	129

Introduction

The genus *Anapagurus* was described by Henderson (1886a) as a subgenus of *Spiropagurus* Stimpson, 1858, based on differences in the sexual tubes, chelipeds and walking pereopods of three species collected in the Firth of Clyde, Scotland that he removed from *Pagurus*, *P. hyndmanni*, *P. laevis*, and *P. ferrugineus*. The same year, Norman (1886) considered the latter species a junior synonym of *P. chiroacanthus*. Two years later, Henderson (1888) elevated *Anapagurus* to the generic rank and added two new species, *A. pusillus* from off the Azores and Canary Islands, and *A. australiensis* [= *Micropagurus acantholepis* (Stimpson, 1858); see Gunn & McLaughlin, 1988] from Port Jackson, Australia.

In 1892 four new species were added to *Anapagurus*, *A. curvidactylus* from off Dakar, Senegal by Chevreux & Bouvier; *A. brevicarpus* from off northern Spain and Arguin Bank, Mauritania, and *A. bicorniger* from Golfo de Cádiz, Spain by A. Milne Edwards & Bouvier; and *A. pusillus* var. *japonica* (= *A. japonicus*) from Kadsiyama, Japan by Ortmann.

A year later A. Milne Edwards & Bouvier (1893) examined two females collected by the "Blake" in the West Indian region and found them to represent two new species, which they described and assigned to *Anapagurus*, *A. acutus* and *A. marginatus*. In 1900 the same authors described *A. laevis* var. *longispina* (= *A. longispina*), and a questionable taxon *Anapagurus?* *dubius* (= *Acanthopagurus dubius*; see De Saint Laurent, 1968, 1969).

In two consecutive years Nobili (1905, 1906a) described respectively, *Anapagurus bonnieri* from the Persian Gulf, and *A. polynesiensis* (= *Micropagurus polynesiensis*; see Haig & Ball, 1988) from Rikitea, French Polynesia.

In the study of the hermit crabs of the Gulf of Naples, Fenizia (1937) described *A. breviaculeatus*, and ten years later Barnard (1947) added a South African species, *A. hendersoni*. The final three new taxa, prior to the present study, assigned to *Anapagurus* were described in the 1960s, *A. wolffi* Forest, 1961; *A. petiti* Dechancé & Forest, 1962; and *A. drachi* Forest, 1966. Table 1 summarizes this brief review of the literature on *Anapagurus* and shows the origins of several major taxonomic problems that exist in the genus.

Ingle (1985) proposed to reappraise the adult taxonomy of the pagurid hermit

Table 1. Summary of the history of the genus *Anapagurus* Henderson, 1886.

* <i>Anapagurus hyndmanni</i> (Bell, 1846)				
* <i>Anapagurus laevis</i> (Bell, 1846)				
* <i>Anapagurus chiroacanthus</i> (Lilljeborg, 1856) ←	←			
<i>Anapagurus ferrugineus</i> (Norman, 1863) ————— (1)		(2)		
* <i>Anapagurus pusillus</i> Henderson, 1888, sensu stricto				
<i>Anapagurus australiensis</i> Henderson, 1888 --- (3) →				<i>Micropagurus acantholepis</i> (Stimpson, 1858)
* <i>Anapagurus curvidactylus</i> Chevreux & Bouvier, 1892				
<i>Anapagurus brevicarpus</i> A. Milne Edwards & Bouvier, 1892				
* <i>Anapagurus bicorniger</i> A. Milne Edwards & Bouvier, 1892				
<i>Anapagurus pusillus</i> var. <i>japonica</i> Ortmann, 1892 ----- → * <i>Anapagurus japonicus</i>				
<i>Anapagurus acutus</i> A. Milne Edwards & Bouvier, 1893 ----- (2) → Species incertae sedis.				
<i>Anapagurus marginatus</i> A. Milne Edwards & Bouvier, 1893 ----- (2) → Species incertae sedis.				
<i>Anapagurus laevis</i> var. <i>longispina</i> A. Milne Edwards & Bouvier, 1900 ----- → * <i>Anapagurus longispina</i>				
<i>Anapagurus? dubius</i> A. Milne Edwards & Bouvier, 1900 ----- (4) → <i>Acanthopagurus dubius</i>				
* <i>Anapagurus bonnieri</i> Nobili, 1905				
<i>Anapagurus polynesiensis</i> Nobili, 1905 ----- (5) → <i>Micropagurus polynesiensis</i>				
* <i>Anapagurus breviaculeatus</i> Fenizia, 1937				
* <i>Anapagurus hendersoni</i> Barnard, 1947				
* <i>Anapagurus wolffi</i> Forest, 1961				
* <i>Anapagurus petiti</i> Dechancé & Forest, 1962				
<i>Anapagurus drachi</i> Forest, 1966 ----- (2) → <i>Forestopagurus</i> gen. nov.				
* <i>Anapagurus alboranensis</i> spec. nov.				
* <i>Anapagurus adriaticus</i> spec. nov.				
* <i>Anapagurus vossi</i> spec. nov.				
* <i>Anapagurus atlantidii</i> spec. nov.				
* <i>Anapagurus congolensis</i> spec. nov.				

References: (1) Norman, 1886: 7; (2) present study; (3) Gunn & McLaughlin, 1988: 68; (4) De Saint Laurent, 1969: 731; (5) Haig & Ball, 1988: 184.

* Species presently assigned to *Anapagurus*

crabs occurring within the area of 30°N-80°N and 30°W-30°E in the northeastern Atlantic and the Mediterranean Sea. In this yet to be published work, Ingle intended to include those species of *Anapagurus* from this geographic area.

The objective of the present study is to do a worldwide revision of *Anapagurus* and to diagnose its species considering newly recognized characters, as well as the significance of intraspecific variation and interspecific overlap in morphological characters.

Materials and methods

Due to the geographical distribution of *Anapagurus*, this study has been largely dependent on specimens from foreign museums, although some U.S. collections were also used. A total of 2473 specimens were examined, including *Anapagurus laevis* (204), *A. hyndmanni* (530), *A. chiroacanthus* (1005), *A. pusillus* (21), *A. japonicus* (2), *A. curvidactylus* (33), *A. bicorniger* (146), *A. longispina* (53), *A. bonnieri* (8), *A. breviaculeatus* (220), *A. hendersoni* (44), *A. wolffi* (6), *A. petiti* (59), *A. alboranensis* spec. nov. (27), *A. adriaticus* spec. nov. (75), *A. vossi* spec. nov. (11), *A. atlantidii* spec. nov. (6), and *A. congolensis* spec.

nov. (23). The material was borrowed and subsequently returned to the following institutions or private collections: British Museum (Natural History), London, England (BMNH); private collection of Carlo Froglio, Ancona, Italy (CF); Instituto di Ricerche sulla Pesca Maritima, Ancona, Italy (IRPEM); Institut Royal des Sciences Naturelles de Belgique, Bruxelles, Belgium (IRSNB-IG); Museum of Comparative Zoology, Harvard University, Massachusetts, U.S.A. (MCZ); Zoological Museum of the Moscow State University, Moscow, U.S.S.R. (MMSU); Museum National d'Histoire Naturelle, Paris, France (MNHN); Musée Océanographique, Monaco (MOM); Museo Civico di Storia Naturale, Milano, Italy (MSNM); Göteborgs Naturhistoriska Museum, Göteborg, Sweden (NHMG); Swedish Museum of Natural History, Stockholm, Sweden (NHNM); Indonesian National Institute of Oceanology, Jakarta, Indonesia (NIOJ); Osaka Museum of Natural History, Osaka, Japan (OMNH); Oxford University Museum, Oxford, England (OUM); Nationaal Natuurhistorisch Museum (formerly Rijksmuseum van Natuurlijke Historie), Leiden, Netherlands (RMNH); South African Museum, Cape Town, South Africa (SAM); Forschungsinstitut Natur Museum, Senckenberg, Germany (SMF); Zoological Museum, Tel-Aviv University, Tel-Aviv, Israel (TAU); Universidad de Málaga, Departamento de Zoología, Málaga, Spain (UMDZ); Rosenstiel School of Marine and Atmospheric Science, University of Miami, U.S.A. (UMML); National Museum of Natural History, Smithsonian Institution, Washington D.C., U.S.A. (USNM); Kumamoto University, Faculty of Education, Kumamoto, Japan (ZLKU); Museum für Naturkunde der Humboldt Universität zu Berlin, Germany (ZMB); and Zoologisk Museum, København, Denmark (ZMUC). Specimens examined in this study are listed by institutional acronym following Leviton et al. (1985) and Leviton & Gibbs (1988), with the exceptions of a few institutions that were not listed.

The specimens were examined using a Wild M-5 binocular stereomicroscope and the illustrations were prepared using a Wild M-5 drawing tube. The shield length (SL) has been measured from the tip of the rostrum to the midpoint of the posterior margin of the shield; measurements to the nearest 0.1 mm were made using an ocular micrometer. In the "Material" sections, shield lengths in millimeters are given in square brackets following the number of specimens. The number of specimens examined of the various species of *Anapagurus* and the range of shield length is presented in Table 2.

Detailed examination of the present extensive collections has made it possible to recognize reliable diagnostic characters and to evaluate the range of morphological variations. A separate section on "Morphological Considerations" has been included to describe diagnostically important structures and to clarify the terminology used. The systematic account includes: 1) synonymy and diagnosis of *Anapagurus*, 2) key to the species of *Anapagurus*, 3) complete synonymies and diagnoses of the 13 previously described species, 4) complete descriptions of the five new species, 5) detailed illustrations of all species, and 6) diagnosis of *Forestopagurus* gen. nov.

For each lot of specimens the following information is given: number of specimens arranged by sex, range of shield length, station number, collecting locality, depth of capture in meters (m), bottom type, date of collection, name of collector, and place of deposition. The material examined is arranged by geographic area and listed from north to south, and west to east. A distributional map is provided for each species showing the type locality and the present and recorded stations. The horizontal

Table 2. Number of specimens of *Anapagurus* examined in the present study and range of shield length (mm, in parentheses).

Species	all specimens combined	male	female		intersex	juvenile
			non-ovigerous	ovigerous		
<i>A. laevis</i>	204(1.0-6.0)	112(1.6-6.0)	31(1.2-3.9)	60(1.8-4.9)	-	1(1.0)
<i>A. hyndmanni</i>	530(1.0-5.0)	339(1.1-5.0)	49(1.2-3.9)	142(1.2-3.6)	-	-
<i>A. chiroacanthus</i>	1005(0.6-7.5)	506(0.6-3.0)	166(0.8-5.6)	309(0.8-3.2)	24(1.0-7.5)	-
<i>A. pusillus</i>	21(0.7-3.0)	6(1.2-2.6)	4(0.7-2.3)	11(0.9-3.0)	-	-
<i>A. japonicus</i>	2(1.3,1.9)	1(1.9)	1(1.3)	-	-	-
<i>A. curvidactylus</i>	33(0.8-2.1)	20(1.0-2.1)	2(0.8,1.9)	11(1.0-1.6)	-	-
<i>A. bicorniger</i>	146(1.2-6.0)	77(1.2-6.0)	15(1.4-2.7)	54(1.3-2.9)	-	-
<i>A. longispina</i>	53(1.2-2.8)	36(1.2-2.8)	8(1.6-2.7)	9(1.3-2.5)	-	-
<i>A. bonnieri</i>	8(1.2-2.0)	5(1.3-2.0)	-	3(1.2-1.7)	-	2(0.6)
<i>A. breviaculeatus</i>	220(0.6-3.7)	133(0.7-3.7)	32(0.8-2.9)	53(1.0-3.0)	-	-
<i>A. hendersoni</i>	44(1.1-3.5)	27(1.9-4.4)	6(1.1-3.3)	11(2.0-3.7)	-	-
<i>A. wolffi</i>	6(0.8-1.0)	2(0.8,1.0)	1(0.9)	3(0.9-1.0)	-	-
<i>A. petiti</i>	59(1.3-3.7)	32(1.3-3.7)	12(1.4-2.8)	15(1.8-3.3)	-	-
<i>A. alboranensis</i> spec. nov.	27(1.1-3.4)	21(1.5-2.9)	2(1.1,1.7)	4(2.0-3.4)	-	-
<i>A. adriaticus</i> spec. nov.	75(1.1-4.4)	44(1.3-4.4)	11(1.1-3.8)	20(1.6-4.0)	-	-
<i>A. vossi</i> spec. nov.	11(0.9-1.2)	8(0.9-1.5)	-	3(1.0-1.2)	-	-
<i>A. atlantidii</i> spec. nov.	6(0.9-1.5)	3(0.9-1.0)	-	3(1.3-1.5)	-	-
<i>A. congolensis</i> spec. nov.	23(1.8-3.4)	17(1.9-3.4)	3(2.2-2.3)	3(1.8-2.6)	-	-

Grand total 2473

and vertical distributions of the species are discussed in the section of "Biogeography". Information on the carinoecia and symbiotic and other possible associations are given in separate sections of related organisms of each species account.

In the section "Related organisms" for each species of *Anapagurus* the number of specimens of the associated organism is provided besides the accession number or the acronym of the depository institution with the station number.

The symbols ♂ and ♀, and the abbreviation ovig., used in the sections on "Material" refer to male, non ovigerous, and ovigerous females, respectively. The abbreviation "AAK" stands for R/V "Academician A. Kovalevskii". The terminology applied to the lineae, sulci, and calcified areas of the cephalothorax follows in part that of Pilgrim (1973) and Morgan & Forest (1991).

Morphological considerations

In general, the terminology for morphological structures follows that of McLaughlin (1974, 1980). However, it has been necessary to clarify the terminology and to describe several new diagnostically important structures.

The systematic confusion that has persisted among the species of *Anapagurus* can be attributed to inadequate diagnostic characters. There has been a tendency in the past to assign significant diagnostic importance to characters of the right cheliped. However, recent studies (McLaughlin, 1974; Lemaitre et al., 1982) have shown that the right cheliped in several genera of hermit crabs is affected by sexual dimorphism, age, environmental conditions, and type of carinoecia used. The present studies

have shown that the right cheliped of species of *Anapagurus* has also been similarly influenced. Consequently interspecific overlaps in the ranges of related characters have occurred. Therefore, a combination of diagnostic characters must be used in separating the different species.

Species of three pagurid genera possess certain distinct projections between the ocular acicles, which are similar in their positions. In *Micropagurus acantholepis* (Stimpson, 1858) there are two lobular projections which have been called "bifurcated bracteole" by Stimpson (1907: 229) and "interocular lobes" (Gunn & McLaughlin (1988: 68, fig. 1a). In *Anapagurus bicorniger* and *A. petiti*, Dechancé & Forest (1962: 295, 297, figs. 1, 2) described two triangular lamellae which were called "sailles triangulaires interoculaires". While describing *Pagurus moluccensis*, Haig & Ball (1988: 187, fig. 14b) remarked on a "subrectangular interocular base with two elongate processes separated by deep, narrow median cleft". All these projections extend from the same structure that has been variously referred to as the "ophthalmic ring" (Stimpson, 1907), "anneau oculaire" (Bouvier, 1896, 1940), "ocular plate" (Snodgrass, 1951: 26, fig. 3E-e), and "ophthalmic somite" (Haig & Ball, 1988: 187); all of which imply that the structure is part of the eye. However, observing the molts, which give a more clear view of the head morphology, and following the opinion of McLaughlin (personal communication), I must conclude that the structure located between the ocular acicles does not constitute part of the eye. Dechancé & Forest (1962: 295) called the structure "la pièce interoculaire", which is more indicative of its location, and treated it as separate structure from the eye. Therefore the term "interocular plate" (fig. 1A) is adopted here and used throughout the text when referring to that structure.

Although slight variations were observed in the left chelipeds of species of *Anapagurus*, the considerable variations in the right chelipeds are taxonomically most important. The following new diagnostically important structures, which are apparently not affected by growth or sexual dimorphism, were found in the right chelipeds: (1) the shape and relative size of the mesial and lateral lobes that flank the dorsoproximal margin of palm (fig. 1B); (2) the shape and relative size of the median sinus of the ventroproximal palmar margin (fig. 1C); (3) the shape, armature and relative size of the dorsomesial distal carpal angle (fig. 1D); and (4) the shape and extension of both ventrodistal and ventroproximal carpal margins, which determine the relative ventral length (fig. 1C).

During this study other morphological characters, such as the relative length and setation of the ultimate antennular segment (fig. 1A), and the relative length between the right uropodal endopod and the distal protopodal lobe (fig. 1E), have been found to be of diagnostic significance.

Anapagurus Henderson, 1886

Spiropagurus (*Anapagurus*) Henderson, 1886a: 337; 1886b: 73 (by implication; see remarks). Type species, by subsequent selection by Holthuis (1962: 242); *Pagurus laevis* Bell, 1846: 184.
Anapagurus Henderson, 1888: 73 (in part); Milne Edwards & Bouvier, 1893: 119 (in part).

Diagnosis.—Carapace with only shield well calcified. Anterior margin of shield slightly produced, rounded. Lateral projections with short submarginal spine, occasionally unarmed. Corneae dilated. Ocular acicles subtriangular to subovate, termin-




Fig. 1. Diagrammatic *Anapagurus*, showing terms used in text. A, shield and cephalic appendages; B-D, right cheliped; B, dorsal view of chela; C, ventral view of carpus; D, mesial view of carpus; E, dorsal view of right uropod.

nating in short submarginal, occasionally marginal spine. Interocular plate unarmed or with 2 chitinous spines. Antennal peduncle with supernumerary segmentation; acicle well-developed, slightly arcuate. Maxillule with stiff bristle on moderately well-developed internal endopodal lobe; external lobe lacking or weakly developed. Third maxilliped with 1 accessory tooth on well developed crista dentata. Sternite of 3rd maxillipeds leveled and with shallow median cleft. Eleven pairs of phyllobanchiae. Chelipeds unequal, right the larger; dactyls opening in horizontal plane. Ambulatory legs long and slender. Anterior lobe of sternite of third pereopods subrectangular. Fourth pereopods subchelate; dactyl without preungual process; propodal rasp with single row of closely spaced corneous scales. Sternite of 5th pereopods forming 2 widely spaced lobes. Males with coxa of left 5th pereopod somewhat enlarged and with well-developed, curved sexual tube; coxa of right 5th pereopod sometimes with short sexual tube, or occasionally with gonopore partly covered by tufts of long setae; with 3 unpaired pleopods. Females with paired gonopores; 4 unpaired pleopods, 2nd to 4th pleopods biramous, 5th uniramous. Uropods asymmetrical. Telson with transverse suture; posterior lobes asymmetrical, separated by median cleft; terminal margins each with 2 to 4 spines; lateral margins unarmed, occasionally each with 1 to 4 spinules.

Genus size.— Eighteen species are recognized here as belonging to *Anapagurus* Henderson, 1886. They are marked with an asterisk (*) in Table 1.

Distribution.— With the exception of *Anapagurus bonnieri*, which has been collected in the Indian Ocean, and *A. japonicus* which lives in southern Japanese waters, the remaining 16 taxa of *Anapagurus* have an eastern Atlantic distribution, extending from the northwestern coast of Norway to the southern coast of South Africa. The depth range is 0-1262 m.

Remarks.— In the family Paguridae there are 22 genera with males having one or two sexual tubes (extruded vas deferens from the coxae of the 5th pereopods), highly advanced characteristics that easily separate them from other hermit crabs (Table 3).

Among the seven genera with well-developed, left sexual tubes, four were defined in part by the presence of an accessory tooth on the crista dentata of the 3rd maxilliped: *Spiropagurus* Stimpson, 1858; *Anapagurus* Henderson, 1886; *Micropagurus* McLaughlin, 1986; and *Pygmaeopagurus* McLaughlin, 1986; and three lack the accessory tooth, *Iridopagurus* De Saint Laurent-Dechancé, 1966; *Anapagrides* De Saint Laurent Dechancé, 1966; and *Enneobranchus* García-Gómez, 1988. A telson without a transverse suture, but with the terminal margin entire and multispinose ocular acicles immediately separates species of *Micropagurus* from species of *Spiropagurus*, *Anapagurus* and *Pygmaeopagurus*. The monotypic *Pygmaeopagurus*, represented by *P. hadrochirus* McLaughlin, is easily distinguished from species of *Spiropagurus* and *Anapagurus* by its massive right cheliped; ocular peduncles broad basally and tapering to bluntly pointed corneae; an obtusely triangular rostrum; and the absence of the male right gonopore. *Anapagurus* is distinguished from *Spiropagurus* by unequal chelipeds (the disparity in size is often very striking in adult males); more slender ambulatory legs with dactyls proportionately shorter and slightly setose; and, in males, a shorter curved left sexual tube.

Anapagurus was originally described by Henderson (1886a) as a subgenus of *Spiropagurus* when he listed its three species: *Spiropagurus* (*Anapagurus*) *hyndmanni*, *S. (A.) laevis*, and *S. (A.) ferrugineus*. In its appendix, Henderson noted that Norman

Table 3. Genera of the family Paguridae with males having sexual tubes.

<i>Acanthopagurus</i> De Saint Laurent, 1968
<i>Anapagrides</i> De Saint Laurent, 1966
<i>Anapagurus</i> Henderson, 1886
<i>Catapaguroides</i> A. Milne Edwards & Bouvier, 1892
<i>Catapagurus</i> A. Milne Edwards, 1880
<i>Cestopagurus</i> Bouvier, 1897
<i>Decaphyllus</i> De Saint Laurent, 1968
<i>Enneobranchus</i> García-Gómez, 1988
<i>Forestopagurus</i> gen. nov.
<i>Goreopagurus</i> McLaughlin, 1986
<i>Iridopagurus</i> De Saint Laurent-Dechancé, 1966
<i>Micropagurus</i> McLaughlin, 1986
<i>Nanopagurus</i> McLaughlin, 1986
<i>Nematopaguroides</i> Forest & De Saint Laurent, 1967
<i>Nematopagurus</i> A. Milne Edwards & Bouvier, 1892
<i>Ostraconotus</i> A. Milne Edwards, 1880
<i>Pagurodes</i> Henderson, 1888
<i>Parapagurodes</i> McLaughlin & Haig, 1973
<i>Pygmaeopagurus</i> McLaughlin, 1986
<i>Solenopagurus</i> De Saint Laurent-Dechancé, 1968
<i>Spiropagurus</i> Stimpson, 1858
<i>Trichopagurus</i> De Saint Laurent, 1968

(1886) had considered *Pagurus ferrugineus* a junior synonym of *Spiropagurus* (*Anapagurus*) *chiroacanthus*. Although in this paper and in a later publication (1886b) Henderson did not write explicitly *Spiropagurus* (*Anapagurus*), he made it clear that he considered *Anapagurus* a subgenus of *Spiropagurus*.

In 1893 A. Milne Edwards & Bouvier described *A. acutus* and *A. marginatus* based on two females collected in the tropical western Atlantic. Forest & De Saint Laurent (1967: 161) questioned the position of these species in *Anapagurus* and noted that their chelipeds resemble those of their newly erected *Nematopaguroides*. In the course of the present study, my examination of these two poorly preserved females and the descriptions and illustrations given by A. Milne Edwards & Bouvier (1893) have revealed important morphological differences in the armature of chelipeds and telson that exclude both species from *Anapagurus*. Both are herein considered species incertae sedis.

The study of the paratypes of *Anapagurus drachi* Forest, 1966 has revealed several important morphological differences (see under *Forestopagurus* gen. nov.: remarks) that set this species apart from *Anapagurus*, and a new genus is herein established for this taxon.

Key to the species of *Anapagurus*

1. Interocular plate with 2 chitinous spines 2
- Interocular plate unarmed 3
2. Antennal acicles unarmed. Carpus of right cheliped with 2 widely spaced spines on distal half of lateral face *A. petiti* Dechancé & Forest, 1962

- Antennal acicles with short distal spine. Carpus of right cheliped without spines on lateral face *A. bicorniger* A. Milne Edwards & Bouvier, 1892
- 3. Lateral margins of sternal lobes of 2nd pereopods parallel
..... *A. hendersoni* Barnard, 1947
- Lateral margins of sternal lobes of 2nd pereopods convergent 4
- 4. Ventrolateral distal angle of coxae of chelipeds with spines 5
- Coxae of chelipeds with ventrolateral distal angle unarmed 6
- 5. Right cheliped with spinulose ridge proximally near dorsomesial margin of palm; dorsomesial margin of carpus with row of 3 or 4 spines on distal half
..... *A. atlantidii* spec. nov.
- Right cheliped without spinulose ridge proximally near dorsomesial margin of palm; dorsomesial margin of carpus with irregular row of closely spaced spines 7
- 6. Ocular acicles with marginal spine
..... *A. longispina* A. Milne Edwards & Bouvier, 1900
- Ocular acicles with submarginal spine 8
- 7. Dactyls of ambulatory legs with longitudinal sulci on lateral and mesial faces
..... *A. hyndmanni* (Bell, 1846)
- Dactyls of ambulatory legs without longitudinal sulci
..... *A. curvidactylus* Chevreux & Bouvier, 1892
- 8. Ultimate antennal segment not reaching base of corneae. Lateral projections of shield unarmed. Laterodistal margin of 1st antennal segment unarmed
..... *A. wolffi* Forest, 1961
- Ultimate antennal segment reaching beyond base of corneae. Lateral projections of shield with short submarginal spine. Laterodistal margin of 1st antennal segment with short spine 9
- 9. Dactyl of right chela with dorsomesial and ventrolateral ridges 10
- Dactyl of right chela lacking ridges 11
- 10. Dorsolateral margin of left cheliped spinose. Terminal margins of telson, each with 3 or 4 spines *A. chiroacanthus* (Lilljeborg, 1856)
- Dorsolateral margin of left cheliped unarmed. Terminal margins of telson, each with 2 spines *A. japonicus* Ortmann, 1892
- 11. Corneae dilated 12
- Corneae not dilated 13
- 12. Ultimate antennular segment with laterodistal tuft of 5-7 long setae. Terminal margins of telson, each with 2 or 3 spines and long submarginal stiff setae
..... *A. vossi* spec. nov.
- Ultimate antennular segment with laterodistal tuft composed of 1 long and 1 or 2 short setae. Terminal margins of telson, each with 4 spines and very short submarginal stiff setae 14
- 13. Lobes of sternite of 2nd pereopods with lateral spinules. Dorsomesial distal angle of carpus of right cheliped produced into subcircular lobe. Right cheliped of males at most with vestigial ventromesial meral protuberance
..... *A. adriaticus* spec. nov.
- Lobes of sternite of 2nd pereopods unarmed. Dorsomesial distal angle of carpus of right cheliped produced into subtriangular or subrectangular lobe. Right cheliped of males with well-developed, ventromesial meral protuberance 15

14. Ocular acicles ovate, mesial margins strongly expanded. Carpus of right cheliped with dorsomesial irregular row of spines. Lobes of sternite of 2nd pereopods latero-distally expanded *A. laevis* (Bell, 1846)
- Ocular acicles roundly triangular, mesial margins not expanded. Carpus of right cheliped with dorsomesial row of strong spines on proximal $\frac{4}{5}$, and with 2 small spines distally. Lobes of sternite of 2nd pereopods not latero-distally expanded *A. congolensis* spec. nov.
15. Rostrum overreaching lateral projections. Dorsodistal margin of carpus of right cheliped lacking projection adjacent to dorsomesial distal angle *A. bonnieri* Nobili, 1905
- Rostrum exceeded by lateral projections. Dorsodistal margin of carpus of right cheliped with triangular or roundish projection adjacent to dorsomesial distal angle 16
16. Shield broader than long. Ultimate antennular segment exceeding length of penultimate segment by $\frac{1}{4}$ own length; ventral margin with row of tufts of long setae *A. pusillus* Henderson, 1888
- Shield longer than broad. Ultimate antennular segment exceeding length of penultimate segment by $\frac{1}{3}$ to $\frac{1}{2}$ own length; ventral margin with row of tufts of very short setae 17
17. Right cheliped with prominent, meral spinose protuberance in males
..... *A. breviaculeatus* Fenizia, 1937
- Right cheliped with weakly developed, meral spinulose protuberance in males *A. alboranensis* spec. nov.

Anapagurus laevis (Bell, 1846)
(figs. 2, 3, 44g)

Pagurus laevis Thompson, 1844: 267, nomen nudum (see remarks).

Pagurus laevis Bell, 1846: 184, 185, and unnumbered figure (see remarks); White, 1850: 26 (listed); Norman, 1861: 274; Carrington & Lovett, 1882: 391; Haddon & Shackleton, 1891: 637.

Eupagurus laevis; Stimpson, 1858: 236 (listed); 1907: 216 (listed).

Spiropagurus laevis; Norman, 1886: 7 (listed); Scott, 1888: 258 (listed); 1897: 172; Haddon & Shackleton, 1891: 637.

Spiropagurus (Anapagurus) laevis; Henderson 1886a: 338; 1886b: 74 (by implication; see remarks for *Anapagurus*).

Anapagurus laevis; Henderson, 1888: 73 (listed); Bouvier, 1891: 393, 394; 1896: 152; 1922: 31 (in part; see remarks); 1940: 145-147, fig. 100 bis, 107A; Chevreux & Bouvier, 1892b: 90, 91; A. Milne Edwards & Bouvier, 1892: 214 (in part); 1899: 66; 1900: 217, pl. XXVII: figs. 9, 10 (in part; see remarks); Tattersall, 1905: 55; Norman, 1905: 8 (listed); Hansen, 1908: 29; Lagerberg, 1908: 62 (listed); Norman & Brady, 1911: 263; Patience, 1911: 74; Selbie, 1921: 44-48, pl. VII: figs. 7-9, pl. VIII: figs. 1-3; Balss, 1924: 768, 769 (listed); Nobre, 1931: 240, fig. 135; 1936: 148, pl. 49: fig. 125; Monod, 1932: 487 (listed); De Miranda y Rivera, 1933: 30, 60 (listed); Dollfus & Carayon, 1942: 445, figs. 1-4; Calgren, 1945: 77; Thorson, 1946: 340; Zariquey Alvarez, 1946: 122, 123, fig. 157; 1955: 402; 1968: 256, 257, figs. 90r, 91j; Holthuis, 1950: 139-141, fig. 50; 1962: 242 (type species by subsequent selection, *Pagurus laevis* Bell, 1846); Pike, 1953: 226, 227, 230; 1961: 226, 227, 229, 230, 232, 233, 237-239, figs. 2B, 4, 7B; Forest, 1958: 100 (see remarks); 1965: 355, 356, figs. 12, 14, 15, 20; Gordan, 1956: 303 (listed); MacDonald, Pike & Williamson, 1957: 237-240, fig. 8; Fuller, 1958: 164; Pike & Williamson, 1958: 6-8, figs. 26-29; 1959: 562, 563, fig. 4; 1960: 525; Dechancé & Dufaure, 1959: 1566-1568; Peres & Picard, 1964: 104; Ross, 1967: 306 (listed); Bourdon, 1968: 195; Hazlett, 1968a: 138; 1968b: 248; Gilpin-Brown, 1969:

- 126; Gilat, 1969: 16, 54; Nielsen, 1970: 19, fig. 2A-D; Guille, 1971: 132; Naylor, 1972: 70, 78; Jensen & Bender, 1973: 190 (listed); Türkay, 1976: 36 (in part; see remarks); Neves, 1977: 179, 180; Manuel, 1981: 73; García Raso, 1982: 501, 502; Mataallanas, 1982: 248, 250, 251; Cranmer, Dyer & Fry, 1984: 342 (listed); Ingle, 1985: 764 (listed); Peres, 1985: 211, 225; Høeg & Lutzen, 1985: 36, 41; Lindley, 1986: 546, 548; Muirhead, Tyler & Thurston, 1986: 131; Markham, 1986: 154; Riedl, 1986: 498, pl. 178: unnumbered figure; Basford, Eleftheriou, & Raffaelli, 1989: 400.
- Anapagurus laevis*; A. Milne Edwards & Bouvier, 1894: 72, pl. XI: figs. 16-28 (in part; see remarks)(mis-spelling of *A. laevis*).
- ? *Anapagurus laevis*; Balss, 1926: 375, 377, 379; 1957: 1751; Mouchet, 1931: 14, 88-90, figs. 85, 86; Lewinsohn, 1976: 245, 246; Shiber & Fattah, 1978: 131 (listed); Kocatas, 1981: 162; Doumenc, Chintiroglou & Koukouras, 1985: 517 (see remarks).
- ? *Anapagurus levis*; Fenizia, 1937: text and figure legends for figs. 15, 16 (misspelling of *A. laevis*).
- not *Anapagurus laevis*; Forest, 1955: 131, 132, fig. 31, pl. VI: fig. 6 (= *A. congolensis* spec. nov.; fig. 32, pl. VI: figs. 7, 8 (= *A. hendersoni* Barnard, 1947) (see remarks).
- not *Anapagurus laevis*; Forest, 1961: 239 (= *A. wolffi* Forest, 1961; *A. vossi* spec. nov.); figs. 8, 12, 16 (= *A. atlantidii* spec. nov.)
- not ? *Anapagurus laevis*; Holthuis & Gottlieb, 1958: 72 (see remarks).
- ? not *Anapagurus laevis*; Fenizia, 1937: 22-24, figs. 15, 16; Sarojini & Nagabhushanam, 1969: 387, 388.

Material.— Syntypes: OUM; not examined. **Northeastern Atlantic:** Norway: 8 ♂♂ [2.8-4.5], 9 ovig. [2.6-3.6], Excursion RMNH station 5, 63°42'N, 9°51'E, 35-50 m, 20.viii.1963, RMNH D 20254; 3 ♂♂ [2.0-3.9], 3 ♀♀ [2.6-3.9], Excursion RMNH station 7, 63°42'N, 9°55'E, 50 m, 20.viii.1963, RMNH D 20255; 1 ovig. [2.1], Excursion RMNH station 17, 63°41'N, 9°21'E, 90 m, 23.viii.1963, RMNH D 20261; 1 ♀ [3.0], Excursion RMNH, 63°39'N, 9°30'E, 5-10 m, 31.viii.1961, RMNH D 17519; 1 ovig. [2.6], Excursion RMNH station 127, 63°41'N, 9°42'E, 25-50 m, RMNH D 24217; 1 ovig. [4.2], Excursion RMNH station 105, 63°43'N, 9°53'E, 150-200 m, 24.viii.1965, RMNH D 24218; 1 ♂ [3.9], Scandinavian Excursion station 7, Stjørnfjord, 20.viii.1963, RMNH D 24219; 1 ♀ [2.9], Molde Fjord, collector Schottländer, ZMB 20266; 2 ovig. [2.7, 3.4], off Bergen, viii.1907, collector Utrecht, RMNH D 4831; 1 ovig. [4.9], station 1225, off Storeggen, collector Uddström, NHRM 1179; 1 ♂ [5.9], station 1227, off Storeggen, collector Uddström, NHRM 1181; 1 ovig. [3.9], off Bergen, 182.9-292.6 m, 1877, collector O. Johanson, NHRM 2082; 1 ovig. [4.3], station L 369, off Bergen, 182.9-219.5 m, 1877, collector M. Olsson, NHRM 2084; 1 ovig. [4.7], off Bergen, 164.6-292.6 m, 1880, collector G. Nilsson, NHRM 2419; 2 ♂♂ [5.1, 6.0], 2 ovig. [2.9, 3.8], station L763, off Bergen, 274.3 m, 1880, collector T. Anderson, NHRM 2449; 1 ovig. [4.1], station L763, off Sorfjord, 36.6 m, vii.1880, collector Bovallius, NHRM 2472; 1 ♂ [3.4], 1 ovig. [3.1], off Storeggen, 109.7-128.0 m, NHRM 1444; 1 ♂ [3.5], station 1766, off Godö, 164.6 m, 1867, collector V. Yhlen, NHRM 1507; 1 ovig. [2.8], station 1767, off Rondö, 109.7 m, 1867, collector V. Yhlen, NHRM 1508; 1 ♂ [1.7], station 1771, off Jederen, 457.2 m, sandy bottom, 1867, collector V. Yhlen, NHRM 1512; 1 ovig. [2.9], station 1772, off Jederen, 457.2 m, sandy bottom, 1867, collector V. Yhlen, NHRM 1513; 1 ♀ [2.3], station 1775, off Jederen, 457.2 m, sandy bottom, 1867, collector V. Yhlen, NHRM 1516; 1 ovig. [2.4], station 1777, off Bodypet, 36.6 m, 1867, collector V. Yhlen, NHRM 1518; 1 ovig. [3.5], station 1776, off Sukkertoppen, 109.7 m, 1867, NHRM 1520; 1 ♀ [2.1], station 2284, off Storeggen, 128 m, rocky bottom, 1867, NHRM 2009; 6 ♂♂ [1.9-3.8], 3 ♀♀ [2.2-2.4], 1 ovig. [2.9], Excursion RMNH, 59°15.5'N, 5°11'E, 30-35 m, 9.ix.1961, RMNH D 17520; 1 ♂ [4.4], 3 ovig. [2.6-3.3], 1884, collector A.M.Norman, USNM 6809. **Sweden:** juvenile [1.0], station 1224, off Väderöarna, collector Goës, NHRM 1178; 1 ♂ [3.6], off Gåsö ränna, vi.1913, NHRM 4378; 1 ovig. [2.8], Kristinebergs Zoological Station, off Väderöarna, 20 m, bottom of rocks and gravel, 7.vii.1926, NHRM 4646; 1 ♂ [3.1], off Väderöarna, coraline bottom, 3.vii.1930, collector S. Bock, NHRM 5172; 1 ♂ [2.7], off Kummelbanken, 1900, collector T. Tullberg, NHRM 4173. **Denmark:** Kattegat: 1 ovig. [3.7], "Hauch" station 58, ca. 57°45'N, ca. 10°45'E, 132 m, fine clay with fine sand, 19.viii.1884, ZMUC; 1 ovig. [3.9], "Hauch" station 239, ca. 57°10'N, ca. 11°40'E, 55 m, fine clay with coarse sand, 21.viii.1885, ZMUC; 1 ♂ [4.0], "Hauch" station 243, ca. 57°05'N, ca. 11°40'E, 43 m, gravels with little clay, 24.viii.1885, ZMUC; 1. ovig. [4.0], "Hauch" station 474, ca. 57°05'N, ca. 11°40'E, 44 m, a little clay, 23.viii.1886, ZMUC. **British Isles:** 2 ♂♂ [2.5, 3.0], station 1226, off Nordsjön, Shetland Islands, collector Verngren, NHRM 1180; 1 ♂ [4.6], 1 ovig. [2.8], "N.L. Grönlands Expedition

1889", 61°16'N, 1°18'W, 150 m, bottom of sand, stones, and shells, 31.v.1889, NHRM 3930; 1 ♂ [4.2], 2 ovig. [2.8, 3.1], off Nordsjön, Shetland Islands, collector A.M. Norman, NHRM 6889; 2 ♂♂ [4.4, 4.7], 2 ovig. [2.7, 3.0], off Shetland Islands, collector A.M. Norman, USNM 17006; 4 ♂♂ [2.0-4.0], 2 ovig. [2.1], "Princesse Alice II" station 1043, 59°03'N, 1°47'45"W, 88 m, 13.ix.1898, MOM 380727; 2 ♀♀ [2.0, 2.2], station 101, 58°56'N, 1°21'W, 115 m, 27.vi.1908, RMNH D 29773; 1 ♂ [4.2], "Deutsche Tiefsee Expedition" station 3, off Aberdeen, 5.vii.1898, ZMB 16477; 6 ♂♂ [2.8-3.9], 1 ♀ [2.8], 1 ovig. [3.6], off Plymouth, MSNM 1870; 1 ♂ [3.4], 2 ♀♀ [3.2, 3.5], off Plymouth, MSNM 2074; 1 ♀ [3.3], off Plymouth, ZMUC; 25 ♂♂ [2.5-4.4], 1 ♀ [3.2], 10 ovig. [2.5-4.0], Firth of Clyde, collector J. Murray, USNM 16991. **Bay of Biscay:** 1 ♂ [1.6], 1 ♀ [1.2], "Hirondelle" station 42, 46°47'N, 6°12'30"W, 136 m, bottom of fine sand, 18.vii.1886, MOM 380021; 1 ♀ [1.4], "Hirondelle" station 44, 46°27'N, 6°30'W, 166 m, silty sand bottom, 20.vii.1886, MOM 380033; 8 ♂♂ [2.1-3.2], 2 ♀♀ [2.1, 2.2], "Hirondelle" station 45, 45°48'N, 5°58'W, 160 m, bottom of fine sand, 21.vii.1886, MOM 380042; 1 ♂ [2.4], 3 ♀♀ [2.6-3.9] "Hirondelle" station 46, 46°24'42"N, 5°55'30"W, 155 m, bottom of gray sand, 26.vii.1886, MOM 380054; 3 ♂♂ [1.6-2.7], 1 ♀ [2.2], "Hirondelle" station 85, 46°31'N, 6°52'W, 180 m, silty sand bottom, 28.v.1887, MOM 380118; 1 ♂ [2.7], "Princesse Alice" station 503, from 47°10'N, 8°08'W to 47°12'N, 8°12'W, 1262-748 m, bottom of clay and silty sand, 29.viii.1894, MOM 380271; 1 ovig. [3.0], "Princesse Alice II" station 1463, from 45°20'N, 3°17'W to 45°24'N, 3°07'W, 950-150 m, 27.vii.1902, MOM 380881; 1 ♀ [2.4], "Hirondelle" station 57, 43°44'30"N, 8°32'30"W, 240 m, bottom of rocks, pebbles, and sand, 5.viii.1886, MOM 382509; 4 ♂♂ [2.8-4.0], 1 ♀ [3.0], station 346, 43°51.0'N, 02°11.9'W, 460-520 m, 3.x.1970, MNHN 1995. **Spain:** 1 ♂ [1.8], Excursion RMNH station 1.157, off Isla Jidoiro, Ría de Arosa, 42 m, 30.vii.1962, RMNH D 19364; 3 ♂♂ [2.0-3.1], 1 ovig. [2.8], Excursion RMNH station 1.418, off Punta Figueiriño, Isla Salvora, Ría de Arosa, 50 m, 13.vii.1963, RMNH D; 1 ♀ [2.5], Excursion RMNH station 1.415, off Punta Figueiriño, Isla Salvora, Ría de Arosa, 50-54 m, 13.vii.1963, RMNH D 20260; 1 ♂ [3.8], 1 ♀ [2.1], Excursion RMNH station 1.417, off Punta Porto Chuncos, Península del Grove, Ría de Arosa, 27 m, 23.vii.1963, RMNH D; 1 ovig. [2.1], Excursion RMNH station 1.349, off Punta Barbafeitas, Isla de Arosa, Ría de Arosa, 45 m, 9.vii.1963, RMNH D 20256; 1 ♀ [2.1], "Josefa Cambeira" station 1.841, 42°26.5'N, 08°57.3'W, 35-65 m, 6.viii.1964, collector L.B. Holthuis, USNM 121796; 1 ♂ [3.4], "Josefa Cambeira" station 1.722, 42°33.0'N, 08°55.5'W, 70-50 m, collector L.B. Holthuis, USNM 121795; 1 ovig. [2.9], "Josefa Cambeira" station 121794, 42°30.4'N, 08°56.2'W, 45 m, collector L.B. Holthuis, USNM 121794; 1 ♀ [2.2], Excursion RMNH station 1.187, off Punta del Castillo, Ría de Arosa, 11 m, sandy bottom, 1.viii.1962, RMNH D 18599; 1 ♂ [3.4], "Josefa Cambeira" station 1.842, off Punta del Castillo, Isla Salvora, Ría de Arosa, 45-55 m, 2 ♂♂ [2.9, 3.6], 1 ♀ [2.7], "Josefa Cambeira" station 1.898, off Light Loba de Cordeiro, Ría de Arosa, 35-45 m, 11.viii.1964, RMNH D 24221. **Portugal:** 1 ♂ [1.9], 2 ovig. [1.8], 39°17'N, 9°30'W, 66 m, 2.ix.1951, collector K.W. Schuitema, RMNH D 7979; 1 ♂ [1.6], "Meteor" station 9c90, 37°14.7'N, 9°1.5'W, 114-117 m, 22.vi.1967, collector Thiel, SMF 4799; 3 ♂♂ [2.5-3.6], "Talisman" 39°53'N, 10°52'W, 99 m, bottom of silt and shells, 6.vi.1883, MCZ 6580; 1 ♂ [2.5], "Talisman", 39°53'N 10°52'W, 99 m, bottom of silt and shells, 6.vi.1883, USNM 22922. **Mediterranean Sea: Spain:** 2 ♂♂ [3.4, 3.8], off Málaga, 35-43 m, 1979, collector J.E. García Raso, UMDZ. **France:** 1 ♂ [3.0], station 40, off Beaulieu, 30 m, MOM 382511. **Monaco:** 1 ♂ [3.1], 1 ♀ [3.3], 1 ovig. [2.4], "Navire Eider", 50 m off Bas Moulins, 15 m, 17.ii.1909, collector Gangui, MOM 382512. **Off Mauritania:** 1 ♂ [2.3], "Tydemann" station 3.116, off Cap Blanc, 20°20'N, 17°42'W, 105 m, 27.x.1978, RMNH D. **Uncertain localities:** 1 ♂ [3.5], 1 ovig. [3.0], By Fjord, 29.ix.1907, collector C. Sopta, RMNH D 3712; 1 ♂ [3.6], "Thor" station 284, off NW Thagras Fyrshire, 90 m, 13.x.1904, ZMUC.

Diagnosis.—Shield broader than long. Rostrum slightly exceeded by lateral projections. Lateral projections with short submarginal spine.

Ocular peduncles approximately $\frac{3}{5}$ length of shield; with corneae strongly dilated. Ocular acicles ovate with mesiodistal margins strongly expanded, with short, submarginal spine. Interocular plate unarmed.

Antennular peduncles exceeding distal margin of corneae by approximately entire length of ultimate segment. Ultimate segment exceeding length of penultimate segment by approximately $\frac{1}{3}$ own length, with dorsolateral row of tufts of short

setae and 2 distal setae, 1 moderately long, 1 short; dorsomesial and ventral margins each with row of tufts of moderately long setae; basal segment with long, lateromedial spine and 1 short, ventromesial, distal spine.

Antennal peduncles overreaching corneae distally by approximately $\frac{2}{3}$ length of ultimate segment. Antennal acicles terminating in small spine, usually overreaching corneae distally.


Right cheliped with moderately narrow chela. Dactyl approximately $\frac{3}{4}$ length of palm, slightly curved ventrally; subcircular in cross section, with no ridges. Palm subequal to length of carpus, dorsomesial margin with spinulose ridge on proximal half; dorsal surface convex, with scattered granules or spinules and short setae, prominent, spinulose ridge near dorsomesial margin proximally; dorsolateral margin with spinules extending onto fixed finger and becoming obsolete near claw; dorso-proximal margin with well-developed lobes, laterally and mesially, larger mesial lobe with marginal spinules; ventroproximal margin with skewed, median, subquadrate sinus. Carpus exceeding length of merus by approximately $\frac{1}{4}$ own length; dorsomesial margin with row of spines of different sizes; dorsomesial, distal angle produced into concave, large subrectangular structure, with more raised, ventral, spinulose side; dorsodistal margin with strong, raised, triangular projection adjacent to dorsomesial, distal angle; dorsolateral margin with 1 short spine on proximal third; ventral surface inflated with tufts of short, stiff setae; ventromesial margin subcircular in mesial view; moderately long, ventral length between shallow, subtriangular, ventrodistal and shallow, subcircular, ventroproximal margins. Ventromesial margin of merus with spinules and 1 short, distal spine; ventrolateral margin with 3 to 5 long spines; dorsal margin with short, distal spine. Coxae of chelipeds with ventrolateral, distal angle unarmed.

Dactyl of left cheliped granulose or spinulose and tufts of moderately long setae. Dorsomesial margin of carpus with row of 7 to 11 strong, closely spaced spines increasing in size distally; dorsolateral margin with row of 3 to 6 short, closely spaced spines on proximal half and row of spinules distally.

Dactyls of ambulatory legs exceeding length of propodi by approximately $\frac{1}{4}$ own length, with dorsomesial and ventromesial rows of closely spaced, moderately long setae. Propodi exceeding length of carpi by approximately $\frac{1}{3}$ own length; ventromesial, distal angle with corneous spines, 1-3 short (2nd), 1 very short or unarmed (3rd). Carpi $\frac{2}{3}$ length of meri; dorsal margins with row of short spines (2nd) or with 1 proximal and 1 distal spine (3rd). Meri with 1 spinule on distal third, dorsally (2nd, 3rd) and ventrally (2nd). Lobes of sternite of 2nd pereopods unarmed, laterodistally expanded. Anterior lobe of sternite of 3rd pereopods subrectangular, with well-defined, distal angles.

Males with moderately long and broad, left sexual tube, smoothly recurved distally, and conical short, backwardly directed right sexual tube; pleopods 3 and 4

Fig. 2. *Anapagurus laevis* (Bell, 1846). ♂ [4.0 mm], "Thor" station 243, of Thagras Fyrshire, ZMUC: A, shield and cephalic appendages; B, left antennule (lateral view); C, right cheliped (dorsal view); D, carpus of same (ventral view); E, joint of palm and carpus of same (mesial view); F, left cheliped (dorsal view); G, left 2nd pereopod (mesial view); H, tip of dactyl of same; I, left 3rd pereopod (mesial view); J, tip of dactyl of same; K, sternum; L, extremity of abdomen, uropods and telson; M, left sexual tube (lateral view). Scales equal 1.0 mm: a: A, C-G, I, K, M; b: B, H, J, L.



unequally biramous, pleopod 5 uniramous. Right, distal protopodal lobe approximately $\frac{1}{2}$ length of uropodal endopod. Telson with posterior asymmetrical lobes, separated by narrow, moderately deep cleft; terminal margins each with 4 short spines; lateral margins with moderately long setae, unarmed, occasionally with 1 spine.

Colour.— “The general colour is yellowish testaceous, and there is a distinct red mark extending the whole length of the hand and bifurcating towards the fingers” (Bell, 1846).

Behaviour.— Hazlett studied the dislodging (1968a) and sexual (1968b) behaviour of *A. laevis*.

Related organisms.— The following gastropod and scaphopod shells occupied by *A. laevis* have been reported: *Natica*, *Murex*, *Dentalium* by Carrington & Lovett (1982); *Turritella* (6 specimens), *Nasse* (=*Hinia*), *Trochus*, *Murex*, and *Natice* (=*Natica*) by A. Milne Edwards & Bouvier (1900); *Natica nitida* by Dollfus & Carayon (1942); and *Turritella communis* Risso by Pike & Williamson (1959). In the present collections the following shells inhabited by *A. laevis* have been found: *Buccinum humphreysianum* Bennett, 1825 (1 specimen, RMNH D 20256); *Buccinum undatum* Linnaeus (3 specimens: USNM 16991, ZMUC station 243, and “Thor” station 284); *Colus jeffreysianus* (Fischer) (1 specimen, NHRM 3930); *Gibbula divaricata* Linnaeus (1 specimen, USNM 16991); *Hinia reticulata* (Linnaeus) (1 specimen, NHRM 2472); *Littorina littorea* Linnaeus (2 specimens, USNM 16991); *Lunatia guillemini* (Payraudeau) (1 specimen, USNM 16991); *Natica catena* da Costa (3 specimens: RMNH D 3712, NHRM 1518, and NHRM 4649); *Polinices grossularius* Marche Marchad, 1957 (1 specimen, RMNH D “Tydeman” 27.x.1978); Turridae (1 specimen, ZMB 20266); *Turritella biplicata* (Bronn) (2 specimens: RMNH D 3712 and RMNH D 7979); *Turritella communis* (11 specimens: MSNM 1870, MSNM 2074, RMNH D 4831, NHRM 3930, USNM 16991, and ZMUC-20.xi.1922); and unidentified shell (NHRM 1180).

The following organisms found attached outside shells inhabited by *A. laevis* have been reported: Cnidaria: *Epizoanthus incrustatus* Duben & Koren, 1847: [Haddon & Shackleton, 1891; Balss, 1924 (listed); Carlgren, 1945; Füller, 1958 (listed); Jensen & Bender 1973 (listed); Manuel, 1981; Cranmer, Dyer & Fry, 1984 (listed); and Muirhead, Tyler & Thurston, 1986].— *Paracalliaclitis lacazaei* Dechancé & Dufaure, 1959: new species collected off Banyuls-sur-Mer.— ?*Adamsia palliata* (Boahadsch, 1761): Doumenc, Chintiroglou & Koukouras, 1985 (Aegean Sea). Polychaete *Nereis fucata* (Savigny): Pike & Williamson, 1952; Gilpin-Brown, 1969. In the present collections a number of specimens of *A. laevis* (in parentheses) have been found in shells incrusted by the following organisms: Cnidarian *Epizoanthus incrustatus*: (1) RMNH D 29773 and (1) NHRM 2449. Poriferan *Suberites domuncula* (Oliv): (1) RMNH D 29773 (new record). Serpulid polychaetes: (1) NHRM 3930; (8) USNM 16991; and (1) ZMUC-20.xi.1922. Cirripedian *Balanus* sp: (1) USNM 16991.

Anapagurus laevis has been reported as a host for the following parasites: Isopoda: *Athelges paguri* Rathke: Tattersall, 1905; Patience, 1911; Pike, 1953 and 1961; Naylor, 1972; *Pseudionae hyndmanni* Bate & Westwood, 1868: Dollfus & Carayon, 1942; Pike, 1953 and 1961; Bourdon, 1968; Markham, 1986 (listed); *Liriopsis pygmaea* Rathke (hyperparasite on *Peltogaster paguri* infesting *A. laevis*): Pike, 1961; Ross, 1967 (listed); Naylor, 1972. Cirripedia: *Peltogaster paguri* Rathke, 1842: Nielsen, 1970; Høeg & Lützen, 1985; *Peltogasterella sulcata* (Lilljeborg, 1859): Nielsen, 1970 (as *Gemmosaccus sul-*

catus); Høeg & Lutzen, 1985. In the present collections the following parasites infesting *A. laevis* have been found: Isopoda: *Pseudione hyndmanni* in right branchial chamber of a male collected off N.W. Thagras Fyrshite, "Thor" station 284, ZMUC; no external effects of the parasitism were observed. Cirripedia: *Peltogaster paguri* (1 externa per host): 1 male lacking left sexual tube and pleopod 4 and another male with no external effects (RMNH D 20254); 1 female with reduced pleopods 2, 3, lacking pleopod 4 (RMNH D 20255); 1 male with short left sexual tube, lacking pleopod 3 (NHRM 1507); 1 male lacking left sexual tube and pleopod 4, reduced pleopod 3 (USNM 16991); *Peltogasterella sulcata* (9 externae per host): 1 male with short left sexual tube, feminization of pleopods (MSNM 1870).

The following organisms were found attached to different parts of specimens of *A. laevis*: Serpulid polychaetes: dorsally on carpus of right cheliped (NHRM 3930); ventrally on palm and carpus of right cheliped (USNM 17006). Foraminifera: 2 tests on shield, and 1 dorsally on propodus of right 2nd pereopod (NHRM 1516); 6 tests dorsally on right cheliped: 4 on palm and 2 on carpus (USNM 17006); 1 test dorsally on carpus of right cheliped (USNM 121794). Bryozoa: 1 colony middorsally on palm of right cheliped (USNM 17006). Matallanas (1982) found *A. laevis* the most abundant organism in the stomach of the fish *Macroramphosus scolopax* (Linnaeus) collected off the Catalan coast of Spain.

Life history.— Mouchet (1931) described and illustrated the spermatophores presumable of *A. laevis* from specimens collected off Salammbô, Tunisia. Thorson (1946) covered the breeding and reproduction aspects of this species. The zoeal stages and glaucothoe of *A. laevis* have been described by MacDonald, Pike & Williamson (1957), and Pike & Williamson (1958, 1960). Lindley (1986) reported the vertical larval distribution of *A. laevis* in the Great Sole Bank, Celtic Sea.

Distribution.— *Anapagurus laevis* has been collected in the eastern Atlantic from off Trondheimsfjord, 63°43'N, 9°53'E, Norway to off Mauritania, 17°02'N, 18°59'W, (Chevreux & Bouvier, 1892). The occurrence of *A. laevis* is herein confirmed in the western Mediterranean (i.e., Spain, France, Monaco) and questionably reported from the eastern Mediterranean. References of *A. laevis* off Naples (Balss, 1926; Fenizia, 1937), Adriatic Sea (Balss, 1957), Aegean Sea (Kocatas, 1981; Doumenec, Chintiroglou & Koukouras, 1985), and off Salammbô, Tunisia (Mouchet, 1931) are questionably assigned to Bell's taxon (see remarks). Depth range: 4.6 (Hansen, 1908)-1262 m (A. Milne Edwards & Bouvier, 1899); most frequently found in depths of 20 to 200 m.

Affinities.— Though *Anapagurus laevis* is most closely allied to *A. vossi* spec. nov. and *A. congolensis* spec. nov. also shares the overall appearance and armature of *A. adriaticus* spec. nov. (see section on affinities for *A. adriaticus* spec. nov.). The relationship between Bell's taxon and the two new taxa is inferred from the similarity observed between small adults of comparable size. These three taxa share among others, the following characters, e.g., the strongly dilated corneae and the much broader than long shield. The ultimate antennular segment with laterodistal tufts of fewer (2 or 3) setae, the dorsomesial distal spine of carpus of right cheliped with the same inclination (approximately 45°) as the other dorsomesial spines, and the distal armature of telson with short, submarginal, stiff setae easily distinguish *A. laevis* from *A. vossi* spec. nov. *Anapagurus laevis* can be separated from *A. congolensis* spec. nov., by the former's ovate ocular acicles with mesial margins strongly expanded, carpus of right cheliped with dorsomesial row of spines of different sizes, lobes of




Fig. 3. Distribution of *Anapagurus laevis* (Bell, 1846). Star: type locality. Solid circles: based on present records. Numbers: based on previous records (circled numbers: material examined; numbers?: questionable material): 1-Hansen, 1908; 2-Bouvier, 1922; 3-Basford, Eleftheriou & Raffaelli, 1989; 4-Thorson, 1946; 5-Norman & Brady, 1911; 6, 11-Bouvier, 1891; 7-Henderson, 1886; 8-Tattersall, 1905; 9-Selbie, 1921; 10-Carrington & Lovett, 1882; A. Milne Edwards & Bouvier, 12-1894; 12-1899; 12, 15, 16, 20, 22, 23-1900; 13, 14-Neves, 1977; 14-Türkay, 1976; 17-García Raso, 1982; Zariquey Alvarez, 18-1946; 19-1955; 21, 25?-Mouchet, 1931; 24?-Balss, 1926; 26?-Kocatas, 1981; Doumenc, Chintiroglou & Koukouras, 1985; 27?-Lewinsohn, 1976; 28-Chevreux & Bouvier, 1892.

sternite of 2nd pereopods laterodistally expanded, and in males the ventromesial, distal propodal angle of 2nd pereopods with 1-3 short spines.

Anapagurus laevis and *A. adriaticus* spec. nov. share the overall appearance and armature, but the former species is distinguished by its much broader shield; more dilated corneae; the expanded mesial margins of the ocular acicles; the unarmed, laterodistally expanded lobes of the sternites of the 2nd pereopods; and the much larger, right sexual tube.

Remarks.—Thompson (1844) listed the names of *Pagurus laevis* and other congeners (*P. cuanensis*, *P. ulidianus*, and *P. hyndmanni*) without descriptions, figures or references to other published works, and therefore are considered nomen nudum. Bell (1846) described and illustrated *P. laevis* and the other congeners but credited authorship of these new taxa to Thompson. And for more than a century thereafter the authorship of these species was attributed to Thompson. This nomenclatorial mistake was finally corrected by Forest in 1958. Stimpson (1858) listed Bell's taxon as *Eupagurus laevis*, and subsequently Henderson (1886a) assigned it to *Anapagurus*, a newly erected subgenus of *Spiropagurus*. In 1922, Bouvier reported several pagurids collected by the "Princesse Alice II" as *A. laevis*. My examination of one of these specimens, a male collected at station 1262 (MOM 380834) reveals that it is not conspecific with *A. laevis* but rather belongs to *A. breviaculeatus*. In a preliminary report (1892) on the collections made during the campaigns of the "Hirondelle", A. Milne Edwards & Bouvier (1894) reported on several specimens they believed assignable to *A. laevis*. My reexamination of these specimens has shown that three species were confounded under this name. One specimen from station 226 is referable to *A. longispina* (the male) and one to *Pagurus pubescens* (A. Milne Edwards & Bouvier, 1892) (the female; new catalogue number: MOM 383821). In further studies of pagurids collected during the expeditions of the "Travailleur" and "Talisman", these authors described in 1900 the new variety *A. laevis* var. *longispina* based on a single male specimen. During my reexamination, I found that three additional males from that same station were also referable to his taxon. From the description and illustrations of *A. laevis* from the Gulf of Naples given by Fenizia (1937), it seems that he was not referring to Bell's taxon, but rather to *A. breviaculeatus* and an undescribed *Anapagurus*. Because his specimens have not been available for examination, they are questionably assigned to not *A. laevis*. Based on characters enumerated by Fenizia in his 1937 paper to distinguish *A. laevis* from *A. breviaculeatus*, Holthuis & Gottlieb (1958) questionably reported *A. laevis* from off the coast of Israel, stating that, according to their material, these two taxa could not be considered distinct species. I have examined this Israeli material and found that instead of being conspecific with Bell's taxon all specimens belong to *A. breviaculeatus* with the exception of one ovigerous female (sample 191, RMNH D 13736) and one male (sample 201, RMNH D 13728) from Haifa Bay, and one ovigerous female (sample 695, RMNH D 13744) from Atlit, which are referable to *A. longispina*, and one male from Haifa Bay (sample 191, RMNH D 13742) that belongs to *A. adriaticus* spec. nov. References to *A. laevis* from the eastern Mediterranean are found not only in the systematic literature (Lewinsohn, 1976; Balss, 1926 and 1957) but also in studies of the comparative anatomy of spermato-phores (Mouchet, 1931), anthozoan systematics (Doumenc, Chintiroglou & Koukouras, 1985) and in listings (Shiber & Fattah, 1978; Kocatas, 1981). Since it is impossible to determine the identities of the specimens reported by these authors, the specimens

are questionably assigned to *A. laevis*.

Among the many specimens examined throughout the world's oceans, *A. laevis* has not been found outside the Atlantic. Therefore Bell's taxon is herein considered to be restricted to this ocean. Sarojini & Nagabhushanam (1969) reported the occurrence of *A. laevis* from the intertidal region of Lawson's Bay (India). The description and illustrations provided by these authors are insufficient to establish the identities of their specimens which were not available for examination.

In the study of the hermit crabs collected by the Belgian "Mbizi" Expedition off the African coast of Congo, Forest (1955) recorded as *A. laevis* certain specimens with more slender chelipeds and walking legs than those found in French specimens. I have had the opportunity to examine these specimens, and found that they are not conspecific with Bell's taxon, but rather belong to a new taxon, *A. congolensis* spec. nov. Furthermore, in his remarks about an ovigerous female (cited as a male in table of material examined, page 131), collected by the "Mercator" off the Island of Das-sen, South Africa, Forest expressed doubt as to his identifications to *A. laevis* by listing numerous differences that he found between his material and Bell's taxon. I have examined the latter specimen and have found that it should have been assigned to *A. hendersoni*.

In the study of the pagurids collected by the "Atlantide" off the tropical western coast of Africa, Forest (1961) identified several very small but adult, specimens as *A. laevis*. He found similarities between these small, adult, tropical African specimens and juvenile *A. laevis* from temperate, northeastern Atlantic waters. The present study of these African specimens leads me to the conclusion that they are not conspecific with Bell's taxon but instead represent three different taxa, *A. wolffi* and two undescribed species, *A. vossi* spec. nov. and *A. atlantidii* spec. nov.

And finally in a study of the Decapoda Reptantia collected by the "Meteor" along the coasts of Portugal and Morocco, Türkay (1976) reported two specimens as *A. laevis*. Examination of one of them, an ovigerous female, collected at station 9c 90 (SMF 4799) has shown that it actually belongs to *A. longispina*.

Anapagurus hyndmanni (Bell, 1846)
(figs. 4, 5)

Pagurus Hyndmanni Thompson, 1844: 182, nomen nudum (see remarks for *A. laevis*)

Pagurus Hyndmanni Bell, 1846: 182, 183, and unnumbered figure; White, 1850: 25 (listed); Carrington & Lovett, 1882: 390; Sinel, 1907: 216.

Eupagurus Hyndmanni; Stimpson, 1858: 236 (listed).

Spiropagurus (Anapagurus) Hyndmanni; Henderson, 1886a: 337; 1886b: 74 (by implication; see remarks for *Anapagurus*).

Spiropagurus Hyndmanni; Norman, 1886: 7 (listed); Scott, 1888: 258 (listed); 1897: 172.

Anapagurus hyndmanni; Henderson, 1888: 73 (listed); Alcock, 1905: 186 (listed); Mouchet, 1931: 80-88, 183-185, figs. 75-84, 148-150; Balss, 1944a: 464; 1944b: 599; 1956: 1439; Matthews, 1953: 265; Gordan, 1956: 302 (listed); MacDonald, Pike & Williamson, 1957: 241-243, fig. 9; Forest, 1958: 100; Pike & Williamson, 1958: 7, 8, figs. 30-33, 52; 1959: 564; Bourdon, 1960: 134, 145; Pike, 1961: 237; Pérès & Picard, 1964: 88, 115; Cardona Bendito, 1965: 144, 145, pl. II: fig. A; Zariquey Alvarez, 1968: 259 (listed); Neves, 1977: 181, 182; García Raso, 1982: 502, fig. 2; Ingle, 1985: 764 (listed); Dauvin & Joncourt, 1989: 593.

Anapagurus Hyndmanni; Bouvier, 1891: 393; 1896: 152, figs. 6, 36, 37 (pictorial key); 1940: 151, figs. 105,

107C; Norman, 1905: 8 (listed); Norman & Brady, 1911: 264; Balss, 1912: 110; Selbie, 1921: 49-53, pl. VIII: figs. 4-7; Pérez, 1930: 3, figs. 4, 5; Nobre, 1931: 223, 224, figs. 123, 124; 1936: 139, pl. 44: fig. 112, pl. 46: fig. 118.

Anapagurus curvidactylus; A. Milne Edwards & Bouvier, 1892: 214, 215; 1900: 225 (not *Anapagurus curvidactylus* Chevreux & Bouvier, 1892); Alcock, 1905: 187 (in part); Balss, 1921: 46 (in part); De Miranda y Rivera, 1933: 30, 60 (listed); Bouvier, 1940: 149 (in part); Zariquey Alvarez, 1968: 258 (in part).

Eupagurus hyndmanni; Stimpson, 1907: 216 (listed).

Anapagurus hyndmanni; Gurney, 1942: 258: figs. 106, 107; Anadón, 1975: 206, 210, 214 (misspelling of *A. hyndmanni*).

Material.—Syntypes: 2 specimens from 2 Irish localities: off Portaferry and Belfast Bay, unknown deposition; not examined. **Northeastern Atlantic: British Isles:** 3 ♂♂ [1.9-3.6], 4 ♀♀ [2.2-2.8], Birterbuy (=Bertraghboy) Bay, Ireland, 21.xi.1891, collector A.M. Norman, ZMUC; 4 ♂♂ [3.3-3.6], 1 ♀ [2.6], Birterbuy (=Bertraghboy) Bay, Ireland, 1874?, collector A.M. Norman, USNM 6808; 5 ♂♂ [2.7-3.2], collector A.M. Norman, USNM 17007. **France:** 1 ♂ [2.4], N off Boulogne, viii.1953, collector C. Swennen, RMNH D 12520; 5 ♂♂ [2.4-5.0], 3 ovig. [2.5-2.7], off Roscoff, 25-30 m, 26.vi.1953, collector J.H. Sock, RMNH D 9916; 1 ♂ [2.7], 1 ovig. [1.9], off Roscoff, 17.ix.1958, RMNH D 12331; 10 ♂♂ [1.7-3.1], 5 ♀♀ [1.0-2.4], 2 ovig. [2.1, 2.5], off Roscoff, 16-19.ix.1958, RMNH D 12530; 1 ♀ [1.5], 2 ovig. [1.6, 2.6], off Finistère, viii; ix.1928, collector W.L. Wrede, RMNH D 3589; 1 ♂ [3.2], 2 ovig. [2.0, 2.5], Baie de Morlaix, 20.v.1958, collector W. Vervoort, RMNH D 12267; 10 ♂♂ [1.1-2.6], 3 ♀♀ [1.6-2.6], 1 ovig. [2.3], off Roscoff, bottom covered with *Zostera marina*, 29.x.1977 -26.ii.1978, collector J. Stassen, RMNH D 35233; 1 ovig. [2.6], off Roscoff, 7.viii.1952, collector H. Nouvel, RMNH D 36308; 1 ♂ [3.8], "Hirondelle" station 41, 47°19'45"N, 5°25'W, 19 m, silty bottom, 17.vii.1886, MOM 382508; 2 ♂♂ [2.6, 3.1], 2 ovig. [2.1, 2.6], Baie de Concarneau, MNHN 1827; 2 ovig. [2.1, 2.4], Baie de Concarneau, 15.ix.1958, RMNH D 12527; 3 ♂♂ [1.9-2.0], 1 ovig. [2.1], off Pointe de Moussterlin, near Quimper, 13.ix.1958, RMNH D 12529; 188 ♂♂ [1.1-3.8], 20 ♀♀ [1.8-3.6], 106 ovig. [1.8-3.6], Baie de Forest, 5-15 m, 12.ix.1958, RMNH D 12447; 1 ♂ [2.2], off îles de Glénan, 5.x.1884, collector M.J. de Guerne, USNM 22923. **Spain:** 1 ♂ [3.0], "Travailleur", along Barquero Bay, 200 m, 1882, ZMUC; 2 ♂♂ [1.7, 3.6], 1 ♀ [1.3], 1 ovig. [3.6], station 1.472, Ría de Arosa, 23.vii.1963, USNM 121788; 1 ♂ [2.2], "Josefa Camberra" station 1.579, 42°30'4"N, 08°56'.2"W, Ría de Arosa, 45 m, 17.vii.1964, USNM 121789; 3 ♂♂ [1.6-1.9], 1 ♀ [1.2], 3 ovig. [1.2-1.6], "Flor da Ponte" station 1.634 (F12), 42°30'.5"N, 08°56'.0"W, Ría de Arosa, 18-22 m, 22.vii.1964, USNM 121790; 1 ♂ [2.9], "Flor da Ponte" station 1.681, 42°32'.5"N, 08°54'.0"W, Ría de Arosa, 28.vii.1964, USNM 121791; 4 ♂♂ [2.6-3.4], 1 ♀ [2.7], "Flor da Ponte" station 1.717, halfway between Lobeira de Cambados and Golfeina, Ría de Arosa, 16 m, 30.vii.1964, USNM 121792; 1 ♂ [3.1], 2 ♀♀ [1.4], Excursion RMNH station 6.1.118, Ría de Arosa, 11.8-13 m, 26.vii.1962, RMNH D 18597; 1 ♂ [1.7], Excursion RMNH station 170, Punta de la Come, Ría de Arosa, 9 m, 20.vii.1962, RMNH D 18598; 3 ♂♂ [1.5-1.7], 1 ♀ [1.5], Excursion RMNH station 1.187, near Punta del Castillo, Ría de Arosa, 11 m, 1.viii.1962, RMNH D 18599; 9 ♂♂ [2.0-3.6], 1 ♀ [2.6], 3 ovig. [2.0-3.2], Excursion RMNH station 1.155, off El Grove, Ría de Arosa, 21 m, 30.vii.1962, RMNH D 18600; 3 ♂♂ [1.6-3.4], 3 ♀♀ [1.3-1.9], Excursion RMNH station 1.189, near Centolliro, Ría de Arosa, 12 m, 1.viii.1962, RMNH D 18601; 1 ♂ [2.6], Excursion RMNH station 1.48, off Punta Cabio, Ría de Arosa, 6.5 m, 16.vii.1962, RMNH D 18602; 5 ♂♂ [2.6-3.3], Excursion RMNH station 1.429, N off Pombeiro, Península del Grove, Ría de Arosa, 15.vii.1963, RMNH D 20388; 1 ♂ [1.3], Excursion RMNH station 1.287, off Isla Palgesa, Ría de Arosa, 6.5 m, 4.vii.1963, RMNH D 20389; 4 ♂♂ [2.2-2.6], 1 ♀ [2.4], 2 ovig. [1.6, 1.8], Excursion RMNH station 1.247, off Isla Salvora, Ría de Arosa, 15 m, bottom of coarse sand, 2.vii.1963, RMNH D 20390; 1 ♂ [1.8], 1 ♀ [1.6], Excursion RMNH station 1.471, 0.5 km off Punta Porto Chunco, Península del Grove, Ría de Arosa, 27 m, 23.vii.1963, RMNH D 20391; 1 ♂ [2.7], Excursion station 1.264, 0.5 km ESE off Isla Ratas, S of Caramiñal, Ría de Arosa, 6 m, clayish bottom, 3.vii.1963, RMNH D 20392; 11 ♂♂ [1.8-3.5], 1 ♀ [1.1], 2 ovig. [2.5, 3.3], Excursion RMNH station 1.288, off San Martín, Península del Grove, Ría de Arosa, 11 m, 4.vii.1963, RMNH D 20393; 3 ♂♂ [1.1-2.1], Excursion RMNH station 1.433, off Salvores, Ría de Arosa, sandy bottom, 15.vii.1963, RMNH D 20394; 16 ♂♂ [1.9-3.8], Excursion RMNH station 1.472, 0.4 km SE off Lobeira de Cambudos, S of Isla de Arosa, Ría de Arosa, 20 m, 23.vii.1963, RMNH D 20395; 1 ♂ [3.3], Excursion RMNH station 1.366, 0.2 km E off Punta de Bois, Isla Salvora, Ría de

Arosa, 8.5 m, bottom of cascajo in coarse sand, 10.vii.1963, RMNH D 20396; 1 ♂ [2.3], Excursion RMNH station 1.259, NNE off Punta Cabio, S of Caramiñal, Ría de Arosa, 19 m, 3.vii.1963, RMNH D 20397; 2 ♂♂ [1.1, 1.4], Excursion RMNH station 1.371, 0.3 km NE off Isla Salvora, Ría de Arosa, 9 m, bottom of cascajo in coarse sand, 10.vii.1963, RMNH D 20398; 1 ♂ [1.5], 1 ovig. [1.8], Excursion RMNH station 1.289, 0.8 km SSE off Punta Banquira, Isla de Arosa, Ría de Arosa, 4.vii.1963, RMNH D 20399; 1 ♂ [3.1], Excursion RMNH station 1.430, off Playa San Vicente, Península del Grove, Ría de Arosa, 15.vii.1963, RMNH D 20400; 1 ♂ [2.2], Excursion RMNH station 1.530, 0.5 km NW off Nido de Corvo, Isla de Arosa, Ría de Arosa, 5.5 m, 30.vii.1963, RMNH D 20401; 1 ♂ [1.3], 1 ovig. [1.5], Excursion RMNH station 1.344, 0.1 km N off Punta Campelo, Isla de Arosa, Ría de Arosa, 27 m, 8.vii.1963, RMNH D 20402; 1 ♂ [2.9], 1 ovig. [1.8], "Flor da Ponte" station 1.656, Ría de Arosa, 10-20 m, 24.vii.1964, RMNH D 24263; 2 ♂♂ [3.1], "Flor da Ponte" station 1.317, Ría de Arosa, 10 m, 30.vii.1964, RMNH D 24264; 6 ♂♂ [2.7-3.9], 2 ♀♀ [1.7, 1.8], Excursion RMNH station 1.472, off Lobeira de Cambados, S of Isla de Arosa, Ría de Arosa, 20 m, 23.vii.1963, RMNH D. Portugal: 1 ♂ [1.9], 1 ovig. [2.9], off Berlengus, 39°17'N, 9°30'W, 66 m, 2.ix.1951, collector K.W. Schuitema, RMNH D 7980; 2 ♂♂ [2.2, 2.4], off Cabo da Roca, 39°17'N, 9°30'W, 37 m, 2.ix.1951, collector K.W. Schuitema, RMNH D 7981; 1 ♂ [3.4], Excursion RMNH station 3, mouth of Río Sado, Setúbal, 5.25 m, bottom covered with algae, 29.iv.1971, RMNH D 36309; 7 ♂♂ [2.2-2.6], 2 ovig. [2.0, 2.1], Excursion RMNH, off Cabo Arnel, Ría de Faro, Faro, 29.vii.1962, RMNH D 18596. Mediterranean Sea: Spain: 3 ♂♂ [2.3-3.8], 3 ovig. [2.5-3.5], off Fuengirola, Málaga, 19-28 m, collector J.E. García Raso, UMDZ. One ovig. [3.3] determined as *Anapagurus laevis* and collected with 2 ♂♂ *Pagurus bernhardus*; no data available; USNM 63400.

Diagnosis.— Shield approximately as long as broad. Rostrum slightly exceeded by lateral projections. Lateral projections with short to moderately long, submarginal spine.


Ocular peduncles approximately $\frac{3}{5}$ length of shield, with corneae slightly dilated. Ocular acicles narrowly triangular, with long, distal, submarginal spine. Interocular plate unarmed.

Antennular peduncles exceeding distal margin of corneae by $\frac{1}{2}$ or more than $\frac{1}{2}$ length of penultimate segment. Ultimate antennular segment about twice as long as penultimate segment, with dorsolateral and dorsomesial rows of short setae, and moderately long, dorsolateral, distal seta; ventral margins of ultimate and penultimate segments, each with row of tufts of very long setae that exceed length of ventral antennular flagellum; basal segment with lateromedial spine.

Antennal peduncles overreaching distal margin of corneae by approximately $\frac{1}{2}$ length of ultimate segment. Antennal acicles terminating in short spine, approximately reaching distal margin of corneae.

Chela of right cheliped broad (females) or moderately broad (males). Fingers curved ventrally, strongly so in mature males. Dactyl subequal to length of palm, triangular in cross section, with strong, dorsomedial and ventromesial, spinulose ridges; all surfaces with small spinules; ventral surface with tufts of long setae. Palm

Fig. 4. *Anapagurus hyndmanni* (Bell, 1846). A-L: ♂ (SL = 3.3 mm). Excursion RMNH station 1.366, off Isla Salvora, Ría de Arosa, Spain, RMNH D 20396: A, shield and cephalic appendages; B, left antennule (lateral view); C, right cheliped (dorsal view); D, carpus of same (ventral view) with a serpulid polychaete tube; E, joint of palm and carpus of same (mesial view); F, left cheliped (dorsal view); G, right 2nd pereopod (mesial view) with a serpulid polychaete tube on propodus; H, tip of dactyl of same; I, right 3rd pereopod (mesial view); J, sternum; K, extremity of abdomen, uropods and telson (folliculinid type 1 on exopod of left uropod); L, left sexual tube (lateral view). M, folliculinid type 2 on abdomen of ♂ (SL = 2.9 mm), "Flor da Ponte" station 1.681, 42°32.5'N, 08°54.0'W, Ría de Arosa, Spain, USNM 121791. Scales equal 1.0 mm: a: A, C-G, I, J; b: L; c: B, H, K; d: M.



approximately $\frac{2}{3}$ length of carpus, dorsomesial margin with spinulose ridge on proximal half; dorsal surface somewhat convex, with numerous spinules and scattered, short setae, without ridge near dorsomesial margin proximally; dorsolateral margin with row of spinules increasing in size distally but becoming obsolete near claw; dorsoproximal margin with well-developed lobes, laterally and mesially, larger mesial lobe with concavity to fit the dorsomesial, distal lobe of carpus when chela is flexed; ventroproximal margin with skewed, subquadrate sinus near ventromesial margin. Carpus exceeding merus by approximately $\frac{1}{3}$ own length, shorter in females; dorsomesial margin with irregular row of closely spaced, short spines; dorsomesial, distal angle produced into spinulose subcircular structure; dorsodistal margin with very weak, subcircular projections adjacent to dorsomesial and dorsolateral, distal angles; mesial face somewhat depressed; dorsolateral face rounded, slightly inflated, without delimiting, dorsolateral margin; dorsolateral, distal angle produced into small subcircular, spinulose lobe; ventromesial margin subcircular in mesial view; moderately long, ventral length between shallow, subcircular, ventrodistal and ventroproximal margins; mesial and dorsolateral faces each with spinules and scattered, short setae; ventral surface with spinules and scattered setae, short in females, very long on proximal half in males. Dorsal margin of merus with row of moderately long setae and moderately long, distal spine; ventromesial margin with short distal spine and slightly developed protuberance (males); ventrolateral margin with 4 or 5 strong spines; ventral surface with spinules and scattered setae, short in females, very long in males. Ischium with 1 spinule on ventrolateral margin partly covered by tufts of setae, short in females, long in males; ventromesial margins serrate. Coxae of chelipeds with 1 short, ventrolateral, distal spine.

Dactyl of left cheliped with median row of spinules on dorsal surface; all surfaces with numerous tufts of moderately long setae. Dorsomesial and dorsolateral margins of carpus, each with 2 or 3 strong, widely spaced spines on distal half.

Ambulatory legs with dactyls exceeding propodi by $\frac{1}{5}$ to $\frac{1}{4}$ own length, lateral and mesial faces each with row of closely spaced tufts of short, stiff setae near dorsal and ventral margins, and with longitudinal sulcus, broad (2nd) or narrow (3rd). Propodi exceeding carpi by nearly $\frac{1}{3}$ own length; dorsal surfaces each with row of closely spaced tufts of short, stiff setae, with short spinules (2nd) or unarmed (3rd); lateral and mesial faces each with row of widely spaced, stiff, short setae near dorsal and ventral margins; ventromesial distal angle unarmed but with tuft of stiff, short setae. Carpi approximately $\frac{2}{3}$ length of meri; dorsal surfaces with row of tufts of short, stiff setae and short spines (2nd) or 1 or 2 short spinules on proximal half and 2 or 3 short distal spines (3rd). Dorsal surfaces of meri serrate (2nd) or unarmed (3rd), each with row of tufts of moderately long setae; ventral margins with row of spinules (2nd) or unarmed (3rd), each with row of moderately long setae. Lobes of sternite of 2nd pereopods unarmed, distally rounded. Anterior lobe of sternite of 3rd pereopods subrectangular; anterior margin with tufts of moderately long setae. Males with moderately long, left sexual tube, recurved to level of coxa of 5th right pereopod; coxa of right 5th pereopod with short, curved sexual tube backwardly directed, partly covered by tufts of moderately long setae; three uniramous pleopods. Distal, protopodal lobe of right uropod equaling length of endopod, giving appearance of a chelate structure.

Telson with asymmetrical, posterior lobes separated by wide, shallow cleft; terminal margins each with 4 short spines; lateral margins unarmed but with tufts of moderately long setae.

Colour.— The colour of *Anapagurus hyndmanni* is pale, dull red and its ova are green. (Carrington & Lovett, 1882). The cornea is deeply pigmented (Selbie, 1921).

Most specimens of the present preserved (ethanol) collections have the chelipeds and ambulatory legs with blotches of orange, and the antennal flagella with bands (2 to 4 articles long) of deep orange, occasionally red.

Related organisms.— The following gastropod shells occupied by *A. hyndmanni* have been recorded: *Turritella terebra* by Bell (1846), *Turritella* sp. by Carrington & Lovett (1882), *Littorina* sp. by Sinel (1907), and *Nassarius* sp. (= *Hinia* sp.) by Neves (1977). In the present collections the following shells occupied by *A. hyndmanni* have been found: *Hinia reticulata* (Linnaeus, 1758) (8 specimens: RMNH D 7980, RMNH D 9916, RMNH D 25233, USNM 12170, and ZMUC- 21.xi.1891); *Gibbula pennanti* (Philippi, 1836) (5 specimens: RMNH D 24262, RMNH D 35233, and USNM 121791); *Gibbula albida* Gmelin (1 specimen, ZMUC-21.xi.1891); *Ocinebrina aciculata* (Lamarck, 1822) (2 specimens, RMNH D 35233); *Turritella biplicata* (1 specimen, USNM 121788); and unidentified shell (RMNH D 35233). In the present collections the following organisms were found on shells occupied by *A. hyndmanni*: Bryozoa: on *Turritella biplicata* (USNM 121788); on *Gibbula albida* and *Hinia reticulata* (ZMUC- 21.xi.1891). Protozoan folliculinid type 1 (see below) outside one shell of *Hinia reticulata* and inside two shells of *Ocinebrina aciculata* and one of *Gibbula pennanti* (RMNH D 35233). Annelida: Polychaeta: Serpulidae: outside and/or inside two shells of *Gibbula pennanti*, and outside two shells of *Ocinebrina aciculata* and one shell of *Hinia reticulata* (RMNH D 35233); and outside the shell of *Gibbula albida* (ZMUC- 21.xi.1891); and Terebellidae: RMNH D 35233 (near apex of unidentified shell (RMNH D 35233).

Anapagurus hyndmanni has been reported as a host for the following parasites: nematomorph *Nectonema agile* Verrill and turbellarian *Fecampia erythrocephala* by Mouchet (1931); isopod *Athelges tenuicaudis* by Bourdon (1960) and Pike (1961); and rhizocephalan *Peltogasterella sulcata* (as *Gemmosaccus sulcatus*) by Bourdon (1960). The ciliated protozoan *Folliculina* (= *Pebrilla*) *paguri* (Giard, 1888) reported by Balss (1956) on *A. hyndmanni* is the only recorded folliculinid in the genus *Anapagurus*. In the present collections, 57 specimens of *A. hyndmanni* were hosts of two species of folliculinids, herein referred to as type 1 (fig. 4K), with ovate test, and type 2 (fig. 4M) with a medial transverse constriction and a small, apical knob. The folliculinids usually ranged in number from 1 to 15 (occasionally 24 or 26) specimens per host; most tests were cemented to the posterior abdomen, telson, uropods, and occasionally to the shield. Hosts usually have either one of the two species of folliculinids, however both taxa were found in three specimens of *A. hyndmanni* (RMNH D 9916 and RMNH D 20395). Folliculinids type 1 were found in 14 specimens of *A. hyndmanni* (RMNH D 9916, 12447, 12530, 18597, 18598, 18599, 20390, 24263 and 35233). Folliculinids type 2 were associated with 40 specimens of *A. hyndmanni* (RMNH D 1827, 12267, 12331, 12447, 12530, 18597, 18600, 18601, 20388, 20393, 20395 and 24264; Excursion RMNH station 1.472; and USNM 121791).

One, occasionally two tubes of serpulid polychaetes were attached commonly to the carpus of the right cheliped ventrally or dorsally, occasionally to the propodus of




Fig. 5. Distribution of *Anapagurus hyndmanni* (Bell, 1846). Star: type locality. Solid circles: based on present records. Numbers: based on previous records: 1, 3, 6, 7, 8, 11-Carrington & Lovett, 1882; 2-Balss, 1926; 3-Norman & Brady, 1911; 4, 5, 12-Bouvier, 1891; 10-Henderson, 1886; 13-A. Milne Edwards & Bouvier, 1892, 1900 (material examined); 14-Cardona Bendito, 1965; 15, 16-Nobre, 1931, 1936.

the right 2nd pereopod mesially (RMNH D 12447, 18597, 20395 and 20396).

Filamentous green algae (Chlorophyta) were found attached to the carpus of the left cheliped dorsally, the merus of the ambulatory legs, and the antennal acicles and flagella (RMNH D 12447, 20388 and 20390).

Distribution.— Northeastern Atlantic: from the Shetland Islands (Carrington & Lovett, 1882) to Costa do Algarve (southern Portugal). Mediterranean Sea: off Fuentegirola, Málaga (southern Spain). Depth range: 0 to 200 m; most frequently found in depths of 5 to 30 m.

Affinities.— In the general similarity of chelipeds (e.g., coxae with ventrolateral distal spines; the right chelae of males with ventrally curved fingers and heavily setose carpus, merus, and ischium ventrally) and in the unarmed, ventromesial, distal angle of propodi of 2nd pereopods, *A. hyndmanni* appears close to *A. curvidactylus*. *Anapagurus hyndmanni* may be distinguished from the latter by its much longer, ultimate antennular segment bearing ventral setae longer than ventral antennular flagellum, longer antennal acicles, the presence of longitudinal sulci on the lateral and mesial faces of the ambulatory legs, and the much shorter left sexual tube.

Remarks.— A. Milne Edwards & Bouvier (1892) assigned a relatively large specimen (SL= 3.0 mm) collected by the "Travailleur" off Barquero in northern Spain at a depth of 200 m, to *A. curvidactylus*. In 1900, these authors remarked that the specimen had certain differences (i.e., larger right chela, dactyls of ambulatory legs with longitudinal sulci) from the typical Chevreux & Bouvier's taxon; they attributed these differences to the age of the specimen. I have been able to compare this specimen with the extensive collection at Leiden from Ría de Arosa (northwest Spain) assigned to *A. hyndmanni*, and believe the A. Milne Edwards & Bouvier's specimen is conspecific with Bell's taxon. The Alcock (1905), Balss (1921), Bouvier (1940), and Zariquey Alvarez's (1968) references are to *A. curvidactylus* and include the above specimen of *A. hyndmanni* from Barquero.

In the study of the decapod and stomatopod crustaceans of Portugal, Nobre illustrated *Anapagurus Hyndmanni* with a photograph (fig. 123, 1931 edition; pl. 44: fig. 112, 1936 edition) that was probably reversed in printing so that the actual right cheliped appeared as the left.

Anapagurus chiroacanthus (Lilljeborg, 1856)
(figs. 6, 7)

Pagurus chiroacanthus Lilljeborg, 1856: 118 [type locality: off Kullaberg, Scania (=Skåne), southwest Sweden]; Henderson, 1886b: 75.

Pagurus ferrugineus Norman, 1861: 273, 274, pl. XIII: figs. 1-3.

Pagurus chiroacantha; Sars, 1872: 256 (misspelling of *P. chiroacanthus*).

Spiropagurus (Anapagurus) ferrugineus; Henderson, 1886a: 338, 1886b: 75 (by implication; see remarks for *Anapagurus*).

Spiropagurus (Anapagurus) chiroacanthus; Henderson, 1886a: 353; Gurney, 1942: 153.

Anapagurus chiroacanthus; Henderson, 1888: 73 (listed); Bouvier, 1896: 153, figs. 40 and 41; Norman, 1905: 8 (listed); Lagerberg 1908: 61, pl. I: fig. 10; Norman & Brady, 1911: 264; Elmhirst, 1912: 23; Selbie, 1921: 48; Balss, 1926: 375, 377, 379; 1956: 1433, 1459, 1466; Nilsson-Cantell, 1926: 2-11; Fenizia, 1937: 29-31, fig. 20; Bouvier, 1940: 148, fig. 102; Gurney, 1942: 261; Reinhard, 1942: 412; Thorson, 1946: 339, 340; Zariquey Alvarez, 1946: 122, 123 (listed); 1955: 402; 1968: 257, 258; Buddenbrock, 1948: 1034, 1037; Pike, 1953: 29, 233; 1961: 226, 236, 237; Forest, 1955: 44, 127, 128, pl. 6, fig. 1; 1958:

100; 1961: 240, figs. 6, 10, 14, 17; 1965: 355; Gordan, 1956: 302 (listed); MacDonald, Pike & Williamson, 1957: 243-246, fig. 10; Füller, 1958: 173; Pike & Williamson, 1958: 7, 8, figs. 34-38, 53; 1959: 564; 1960: 525; Bourdon, 1968: 195; Hazlett, 1968a: 138; 1968b: 246, 247, 249; 1981: 8; Samuelsen, 1970: 44; Naylor, 1972: 70; Neves, 1977: 180, 181, fig. 9C; Stachowitzsch, 1980: 74, 93; Calafiore, 1981: 20, 25, 26; Kocatas, 1981: 162; Schembri, 1982: 872; Crothers & Crothers, 1983: 763, 783; Salmon, 1983: 148; Ingle, 1985: 764 (listed); 1990: 105-134; Høeg & Lützen, 1985: 32, 39; Markham, 1986: 154.

Pagurus chiracanthus; Aurivillius, 1891: 33, pl.V: fig. 5 (misspelling of *P. chiroacanthus*).

Anapagurus brevicarpus A. Milne Edwards & Bouvier, 1892: 215; 1898: 1247 (listed); 1899: 67; 1900: 223, pl.XXV: figs. 23-26, pl. XXVIII: fig. 11; Bouvier, 1896: 153, figs. 38, 39 (pictorial key); 1922: 31; 1940: 150, 151, figs. 104, 107 D (listed); Monod, 1932: 487 (listed); De Miranda y Rivera, 1933: 30, 60 (listed); Forest, 1958: 100 (listed); Zariquey Alvarez, 1968: 258, 259 (listed); Ingle, 1985: 764 (listed).

Eupagurus chiroacanthus; Norman, 1886: 7; Sars, 1899: 213; Stimpson, 1907: 215 (listed).

Spiropagurus chiroacanthus; Sars, 1890: 155, pl. iii; 1899: 267, pl. 89: fig. X; Bouvier, 1891: 68, 75; Gurney, 1939: 98 (listed).

Anapagurus chiracanthus; Balss, 1924: 783 (misspelling of *A. chiroacanthus*).

? *Anapagurus brevicarpus*; Mouchet, 1931: 9092, fig. 87; Forest, 1961: 243; Cardona Bendito, 1965: 146.

Anapagurus chiroacanthus var. *gracilis* Fenizia, 1937: 31, 32, fig.21; Pike & Williamson, 1959: 564.

Anapagurus chiroacanthus var. *cristatus* Fenizia, 1937: 32, 33, fig. 22.

Material.— Syntypes, BMNH; not examined. Northeastern Atlantic: Norway: 1 ♂ [1.9], 2 ♀♀ [2.0, 2.1], “Eugenie Expedition” station N: R21, off Farsund, NHRM 1437; 1 ♂ [2.9], “Ing. O. Gladans Expedition”, off Christiansand, 91 m, shelly bottom, 1871, collector Lindahl, NHRM 1504; 4 ♂♂ [1.6-2.4], 1 ♀ [2.2], Leirviksfjord, 128 m, shelly bottom, vii.1880, collector C. Bovallius, NHRM 2469; 2 ♂♂ [1.8, 1.9], off Haugesund, 43.9 m, sandy bottom, vii.1880, collector C. Bovallius, NHRM 2478; 2 ♂♂ [1.4, 2.1], 2 ovig. [1.9], off Haugesund, 73 m, shelly bottom, 5.vii.1880, collector C. Bovallius, NHRM 3486; 2 ♂♂ [1.5], 1 ovig. [2.1], off Norwegian coast, collector A. M. Norman, USNM 17008; 1 ♂ [2.8], Skagerrak, collector Malm, ZMB 18603; 1 ♂ [1.8], 60°33'N, 4°50'E, 12-20 m, bottom of shell, gravel and *Laminaria*, 1.ix.1981, RMNH D; 1 ♂ [2.3], 60°36'N, 4°55'E, 30-50 m, sandy bottom, 4.ix.1981, RMNH D; 3 ♂♂ [0.9-2.2], 1 ♀ [0.9], 3 ovig. [0.9], 1 intersex [1.2], 60°15'N, 5°13.5'E, 10-16 m, 11.ix.1961, RMNH D 19128; 9 ♂♂ [1.0-2.1], 2 ♀♀ [1.2, 5.6], 5 ovig. [1.2-2.4], 60°11'N, 5°2'E, 23-35 m, 6.ix.1961, RMNH D 19129; 1 ♂ [2.6], 1 ovig. [2.2], 63°39'N, 9°30'E, 5-10 m, 31.viii.1961, RMNH D 19130; 1 ovig. [1.6], 63°40'N, 9°14'E, 200 m, 29.viii.1961, RMNH D 19132; 2 ♂♂ [2.0], 2 ♀♀ [1.4, 1.5], 1 ovig. [0.8], off S Bergen, ca 1 m, 7.ix.1961, RMNH D 19133; 4 ♂♂ [2.0-2.9], 2 ♀♀ [1.0, 1.8], 3 ovig. [2.5- 2.7], 60°15.5'N, 5°11'E, 30-35 m, 9.ix.1961, RMNH D 19426; 3 ♂♂ [2.1-2.3], 63°37'N, 9°27'E, 30 m, 31.viii.1961, RMNH D 19427; 2 ♂♂ [1.5, 2.8], 59°15.5'N, 5°11.5'E, 18 m, 9.ix.1961, RMNH D 19429; 2 ♂♂ [1.4, 2.0], 63°43'N, 9°48'E, 7-50 m, 28.viii.1961, RMNH D 19432; 1 ♂ [2.3], 63°36'N, 9°27'E, 10-25 m, 22.viii.1963, RMNH D 20218; 2 ♂♂ [1.9, 2.3], 1 ♀ [2.0], 1 ovig. [1.9], 63°39'N, 9°49'E, 200 m, 27.viii.1965, RMNH D 21858; 2 ♂♂ [2.0, 2.3], 63°38'N, 9°18'E, 30 m, 26.viii.1965, RMNH D 24169; 1 ♂ [2.7], 63°42'N, 9°27'E, 20-70 m, sandy bottom, 26.viii.1965, RMNH D 24170; 3 ♂♂ [1.5-2.7], 1 ♀ [1.2], 63°41'N, 9°42'E, 25-50 m, sandy bottom, 31.viii.1965, RMNH D 24172; 1 ♀ [2.4], 63°41'N, 9°41'E, 15-20 m, bottom of *Laminaria*, 22.viii.1967, RMNH D 24173; Sweden: 7 ♂♂ [2.0-3.0], 6 ♀♀ [1.3-2.4], 4 ovig. [1.9-2.8], 3 intersex [2.2-7.5], off Sorriga, Gullmarn, Bohuslän, bottom of shells and sponges, collector S. Lovén, NHRM 1020; 2 ♂♂ [2.5, 2.7], 3 ♀♀ [1.9-2.5], 9 ovig. [1.5-3.2], off Sorriga, Gullmarn, Bohuslän, bottom of shells and sponges, collector S. Lovén, NHRM 1021; 6 ♂♂ [1.8-2.9], 12 ♀♀ [2.0-2.7], 28 ovig. [1.9-3.0], off Sorriga, Gullmarn, Bohuslän, bottom of shells and sponges, collector S. Lovén, NHRM 1022; 1 ♂ [2.7], off Sorriga, Bohuslän, NHRM 1041; 2 ♂♂ [2.8, 2.9], 2 ovig. [2.6], off Gullmarn, Bohuslän, NHRM 1171; 1 ovig. [1.8], off Stavanger, collector V. Friesen, NHRM 1182; 5 ♂♂ [1.1-1.5], off Sorriga, Bohuslän, NHRM 1407; 1 ♂ [1.8], off Gullmarn, Bohuslän, shallow depth, bottom of algae, NHRM 1411; 1 ♂ [1.8], off Bokevik, Bohuslän, 12.8 m, NHRM 1412; 10 ♂♂ [1.2-2.0], 5 ♀♀ [1.5-1.8], 7 ovig. [1.4-1.9], 1 mile off Koster, 36.6 m, rocky bottom, vii.1865, NHRM 1414; 1 ♂ [1.3], 3/4 mile off Nord Koster, Bohuslän, 36.6 m, sandy bottom, 1865, NHRM 1418; 2 ♂♂ [1.0, 1.8], off Kosterfjorden, Bohuslän, 27.4-36.6 m, sandy bottom, 1865, collector Ljungman, NHRM 1424; 16 ♂♂ [1.4-2.4], 8 ♀♀ [1.3-1.5], 8 ovig. [1.2-2.0], off Kosterfjorden, Bohuslän, 9.1-27.4 m, rocky bottom, 1865, collector Ljungman, NHRM

1425; 13 ♂♂ [1.4-2.5], 2 ♀♀ [1.6, 1.9], 6 ovig. [1.1-1.8], off Kosterfjorden, Bohuslän, 27.4-36.6 m, sandy bottom, 1865, collector Ljungman, NHRM 1427; 5 ♂♂ [1.2-1.9], 2 ♀♀ [1.3], 3 ovig. [1.5-1.8], 5 intersex [1.0-1.6], off Kosterfjorden, Bohuslän, 9.1-27.4 m, rocky bottom, 1865, collector Ljungman, NHRM 1428; 8 ♂♂ [1.8-2.6], 8 ♀♀ [1.5-2.2], 5 ovig. [1.7-2.0], 1 intersex [1.8], off Stora Väderön, Bohuslän, 45.7 m, 1869, collector Olsson, NHRM 1429; 5 ♂♂ [1.0-2.7], 5 ♀♀ [1.5-2.7], 3 ovig. [1.5-2.6] off Lindö, Bohuslän, 54.9 m, 1869, collector Olsson, NHRM 1433; 1 ovig. [2.0], off Styrsoranna, Bohuslän, 54.9 m, 1869, collector Olsson, NHRM 1435; 1 ♂ [2.0], 1 ovig. [1.9], 54.9 m, 1869, collector Olsson, NHRM 1436; 2 ♂♂ [1.6, 1.9], off St. Väderön, Bohuslän, 100.6 m, vi.1869, collector Olsson, NHRM 1505; 2 ♂♂ [1.9, 2.1], off Väderöarna, Bohuslän, 27.4-54.9 m, sandy bottom, collectors C. Bovallius & Théel, NHRM 1529; 7 ♂♂ [1.1-2.1], 2 ♀♀ [1.0, 1.2], 3 ovig. [1.4-1.8], off Väderöarna, Bohuslän, 54.9 m, sandy bottom, vii.1877, collectors C. Bovallius & Théel, NHRM 1530; 7 ♂♂ [0.9-2.0], 10 ♀♀ [0.9-1.5], 3 ovig. [1.2-1.4], 1 intersex [2.2], off Väderöarna, 54.9 m, sandy bottom, vii.1877, collectors C. Bovallius & Théel, NHRM 1531; 5 ♂♂ [1.8-2.4], 1 ♀ [2.1], 1 ovig. [1.6], off Väderöarna, Bohuslän, 54.9 m, sandy bottom, 12.vii.1877, collectors C. Bovallius & Théel, NHRM 1532; 1 ovig. [1.4], off Väderöarna, Bohuslän, 109.7 m, bottom of corals, vii.1877, collectors C. Bovallius & Théel, NHRM 1537; 1 ovig. [2.3], "Gunhild" station 75, Väderöarna, Bohuslän, 73.2-91.4 m, bottom of corals, 28.vii.1877, collectors C. Bovallius & Théel, NHRM 1538; 2 ♂♂ [1.6, 1.9], off Kilsund, Bohuslän, 91.4 m, shelly bottom, vii.1877, collectors C. Bovallius & Théel, NHRM 1542; 1 ♂ [2.7], off Lysekil, Bohuslän, 45.7 m, vii.1877, collectors C. Bovallius & Théel, NHRM 1545; 2 ♀♀ [1.7], 2 ovig. [1.9, 2.0], off Kosterfjorden, Bohuslän, 21.9-32.9 m, rocky bottom, viii.1865, collector Ljungman, NHRM 1872; 1 ♀ [2.9], "Gunhild" station unknown, off Utanför, Helsingborg, 32.9-40.2 m, bottom of sand and clay, 17.vii.1878, collectors Théel & Trybom, NHRM 2250; 1 ♀ [2.1], 1 ovig. [2.2], off Väderöarna, Bohuslän, 5.viii.1879, NHRM 2342; 2 ♂♂ [1.8, 2.1], 1 ♀ [2.1], off Väderöarna, Bohuslän, 1877, NHRM 2352; 9 ♂♂ [1.4-2.5], 1 ♀ [1.4], 1 ovig. [1.4], off Väderöarna, Bohuslän, viii.1881, NHRM 2539; 1 ♀ [2.9], 3 ovig. [2.4-2.9], 1 intersex [2.0], off Gullmarn, Bohuslän, viii.1881, collector Svederus, NHRM 2549; 4 ♂♂ [2.0-2.7], 1 ovig. [1.9], off Väderöarna, Bohuslän, viii.1882, collector Aurivillius, NHRM 2642; 1 ♀ [2.0], off Hallö, Bohuslän, 36.6 m, 1884, NHRM 2758; 1 ♂ [2.7], 1 ovig. [3.0], off Kristinebergs Zoological Station, Bohuslän, 1885, NHRM 2821; 1 ♂ [2.0], off Gullmarn, Strumpeskagen, Bohuslän, 17.vii.1876, collector C. Bovallius, NHRM 3152; 1 ♂ [1.5], off Gullmarn, Strumpeskagen, Bohuslän, 91.4 m, 5.vii.1876, collector C. Bovallius, NHRM 3221; 3 ovig. [2.1-2.6], off Flatholmen, Bohuslän, 54.9 m, 8.vii.1876, collector C. Bovallius, NHRM 3223; 4 ♂♂ [1.2-2.5], 3 ♀♀ [1.7-2.2], 3 ovig. [1.8-2.1], off Nord Koster, Bohuslän, 1869, collector C. Bovallius, NHRM 3224; 1 ♂ [2.5], 1 ovig. [2.5], off Stångehufvud, Lysekil, Bohuslän, 19.vii.1876, NHRM 3225; 1 ♂ [2.1], 1 ovig. [1.9], off Stångehufvud, Lysekil, Bohuslän, 19.vii.1876, NHRM 3226; 2 ♂♂ [2.1, 2.9], 1 ovig. [1.8], off Stångehufvud, Lysekil, Bohuslän, 19.vii.1876, NHRM 3227; 1 ♂ [2.8], 1 ♀ [2.9], 1 intersex [2.2], off Ullholmsrännan, Bohuslän, 31.v.1890, collector C. Aurivillius, NHRM 3484; 1 ovig. [2.7], off Harpöbrott, Bohuslän, v.1890, collector C. Aurivillius, NHRM 3485; 2 ♂♂ [2.7, 2.9], off Gullmarn, Flatholmsrännan, 17.vii.1890, collector C. Aurivillius, NHRM 3519; 2 ♂♂ [2.2, 2.6], 5 ovig. [2.0-2.7], off Kristineberg Zoological Station, Bohuslän, 1893, NHRM 3682; 4 ♂♂ [1.6-2.6], off Kosterfjorden, Kostersten, Bohuslän, 109.7-182.9 m, 3.vii.1895, collector C. Aurivillius, NHRM 3685; 1 ♂ [2.0], 1 ♀ [2.3], 1 ovig. [1.7], off Väderöarna, Bohuslän, bottom of clay, 29.vii.1896, collector I. Arvidsson, NHRM 3686; 2 ♀♀ [1.7, 2.4], 2 ovig. [1.9, 2.1], off Kummelbanken, 1900, collector T. Tullberg, NHRM 4173; 1 ♂ [2.4], off Kummelbanken, 1900, collector T. Tullberg, NHRM 4186; 4 ♂♂ [1.8-3.0], 1 ♀ [2.8], 4 ovig. [2.5-2.7], 1 intersex [1.9], north off Flatholmen, Gullmaren, 80 m, 1.vii.1906, collector T. Lagerberg, NHRM 4255; 2 ovig. [2.4, 2.5], off Gåsöränna, 40 m, 27.vi.1906, collector T. Lagerberg, NHRM 4256; 3 ♂♂ [2.2-2.5], 1 ♀ [1.5], 3 ovig. [1.8-3.1], 1 intersex [2.1], off Svartskär-Vinga, 26 m, 31.vii.1906, collector T. Lagerberg, NHRM 4261; 12 ♂♂ [1.8-3.0], 24 ovig. [1.8-2.7], off Gullmaren, vii.1923, collector C.A. Nilsson-Cantell, NHRM 4676; 64 ♂♂ [1.6-3.0], 5 ♀♀ [1.4-2.7], 28 ovig. [1.5-3.1], 1 intersex [1.9], off Gåsöränna, Gullmaren, vii.1923, collector C.A. Nilsson-Cantell, NHRM 4677; 26 ♂♂ [1.6-3.0], 2 ♀♀ [2.1, 2.9], 21 ovig. [1.7-2.0], 2 intersex [2.1, 2.6], off Smedjebron, Bohuslän, 16.vi.1923, collector C.A. Nilsson-Cantell, NHRM 4678; 1 ♀ [1.5], off Bovallstrand, Lygnösundet, Bohuslän, 12 m, sandy bottom, 23.vii.1935, collector S. Bock, NHRM 5174; 1 ovig. [2.8], off Kristineberg, 1905, collector Y. Schwartz, NHRM 5772; 1 ♂ [2.7], off west coast, MCZ 12688; 1 ♂ [2.0], 1 ovig. [1.8], 58°12'N, 11°20'E, 20-35 m, sandy bottom, 9.ix.1981, RMNH D; 3 ♂♂ [1.5-2.7], 58°16'N, 11°25'E, 35

m, bottom of gravel, 10.ix.1981, RMNH D; 14 ♂♂ [1.8-2.8], 20 ♀♀ [1.4-2.6], 2 ovig. [2.1, 2.2], 3 intersex [1.4-2.1], 58°16'N, 11°25'E, 30 m, 16.ix.1961, RMNH D 19131; 7 ♂♂ [1.4-2.3], 58°13.5'N, 11°25'E, 30-40 m, 16.ix.1961, RMNH D 19425; 3 ♂♂ [2.4-2.8], 58°14.5'N, 11°30.5'E, 7 m, 15.ix.1961, RMNH D 19428; 15 ♂♂ [1.5-2.9], 18 ♀♀ [1.6-2.7], 2 ovig. [2.5], 1 intersex [2.0], 58°15.5'N, 11°25.5'E, 25 m, 16.ix.1961, RMNH D 19431; 5 ♂♂ [2.1-2.8], 2 ♀♀ [2.0, 2.1], 1 intersex [2.0], 58°14'N, 11°25'E, 33-40 m, 16.ix.1961, RMNH D 19433; 2 ♂♂ [1.8], 58°12.5'N, 11°19'E, 35 m, 19.ix.1961, RMNH D 19434; 14 ♂♂ [1.6-2.7], 3 ♀♀ [1.9-2.1], 1 ovig. [2.0], 58°16'N, 11°25'E, 25 m, 29.viii.1961, RMNH D 24168; 1 ♂ [2.6], 1 ♀ [2.3], 1 intersex [1.9], 58°16'N, 11°29'E, 20-30 m, 15.ix.1965, RMNH D 24171; **Denmark:** 13 ♂♂ [2.2-2.8], 4 ovig. [1.9-2.8], off Lilla Middelgrund, 29.3-31.1 m, sandy bottom, 10.vii.1878, collector Théel, NHRM 2240; 1 ♂ [1.6] off Torrbo Skär-Skagen, 49.4 m, sandy bottom, 8.vii.1878, collector Théel, NHRM 2245; 2 ovig. [1.6, 1.9], off Skagen-Nidingarna, 34.7 m, bottom of gravel and stones, 9.vii.1878, collector Théel, NHRM 2246; 1 ♂ [1.7], 1 ♀ [2.6], off Aalborg-Bucht, 20-30 m, 29.vi-1.vii.1976, collector Pettke, SMF 7582; **France:** 2 ♀♀ [1.8, 2.0], "Princess Alice II" station 2974, off Belle Ile", 47°23'N, 3°25'W, 85 m, 27.vii.1910, MOM 380997; **Azores:** 2 ovig. [1.0, 1.6], "Princess Alice" station 594, off Point S. Antonio, Terceira, 54 m, rocky bottom, 22.vii.1895, MOM 380322; 3 ♂♂ [0.6-2.5], 1 ♀ [2.5], "Josephine Expedition" station d 45, off São Miguel, 9.1-18.3 m, rocky bottom, 16-17.vii.1869, NHRM 6976; 5 ♂♂ [1.4-1.7], 1 ovig. [1.4], "Josephine Expedition" station d 72 (c), off Ponta Delgada, 7.3-18.3 m, 1869, NHRM 6996; 2 ♂♂ [1.6, 1.8], "Josephine Expedition" station d 76 (b), off Ponta Delgada, 18.3 m, bottom of gravel, 1869, NHRM 6997. **Mediterranean Sea:** **Spain:** 1 ♂ [1.8], 1 ovig. [1.2], Bay of Cadaqués, 4-16.viii.1949, collector L.B. Holthuis, RMNH D 6774; 2 ♂♂ [1.1, 1.4], 2 ♀♀ [0.8, 0.9], 1 ovig. [1.2], Bay of Cadaqués, 20-50 m, 11-18.viii.1954, collector L.B. Holthuis, RMNH D 10284; 1 ♂ [1.3], Bay of Cadaqués, 30-40 m, 29.vii.1955, collector L.B. Holthuis, RMNH D 10913; 1 ♂ [1.7], Bay of Cadaqués, 26-31.vii.1955, collector L.B. Holthuis, RMNH D 10914; 2 ♂♂ [1.4], Bay of Cadaqués, 45 m, 1.viii.1957, collector L. B. Holthuis, RMNH D 12270; 1 ovig. [2.2], Bay of Cadaqués, up to 1 m, on seagrass bed, 31.vii.1957, collector L.B. Holthuis, RMNH D 12283; 1 ♂ [1.6], Bay of Cadaqués, 20-30 m, 14.viii.1959, collector L.B. Holthuis, RMNH D 14883; 5 ♂♂ [1.2-1.8], 1 ♀ [1.2], 1 ovig. [1.3], Bay of Cadaqués, 30-90 m, sandy bottom, 27.vii-2.viii.1961, collector L.B. Holthuis, RMNH D 16514; 5 ♂♂ [1.4-2.0], 1 ♀ [1.5], 3 ovig. [1.6-2.0], off Formentera, Balearic Islands, 64 m, 1871, collector F. Söderlund, NHRM 12058; 2 ♂♂ [1.3, 1.4], 1 ♀ [1.9], 91.4 m, 1871, collector F. Söderlund; 2 ♂♂ [2.3, 2.4], off San Antonio, Ibiza, Balearic Islands, 45.7 m, 1871, collector F. Söderlund, NHRM 12061; **France:** 4 ♂♂ [1.8-2.0], "L'Eider" station 3422, from off Beaulieu to Roquebrune-Cap-Martin, x-xi.1926, MNHN 1817; **Monaco:** 1 ♂ [2.0], off Cap d'Ail, Port-Marquet, 15 m, 17.iv.1953, collector H. Nouvel, RMNH D; **Italy:** 3 ♂♂ [2.1-2.6], off Sori, Golfo di Genova, 36.6-54.9 m, coralline bottom, 3.x.1887, collector Théel, NHRM 10575; 1 ♂ [2.4], off Naples, 7.xii.1937, MOM 382514; 1 ♂ [2.0], 1 ovig. [2.0], off Naples, 7.xii.1937, MOM 382515; 1 ♂ [1.3], Golfo di Taranto station 308, 39 m, sandy bottom, 26.viii.1966, collector A. Vatova, RMNH D 24174; **Yugoslavia:** 1 ovig. [1.7], off Rovinj, 20 m, 15.viii.1960, RMNH D 16159; 1 ♂ [2.2], off Rovinj, 32-38 m, 3.vi.1962, RMNH D 19430; 1 ovig. [1.8], off Rovinj, 32-38 m, 2.vi.1962, RMNH D 19435; **Strait of Sicily:** 12 ♂♂ [1.9-2.5], 1 ♀ [2.4], 3 ovig. [2.0-2.1], "AAK" station 1261, 37°26.8'N, 12°11.6'E, 57 m, bottom of sand and clay, 24.iv.1978, collector V.V. Murina, MMSU; 2 ♂♂ [1.9, 2.8], "AAK" station 1262/55, 37°40.0'N, 12°03.0'E, 21 m, bottom of clay and sand, 24.iv.1978, collector V.V. Murina, MMSU; 6 ♂♂ [1.6-2.2], 1 ♀ [2.0], 1 ovig. [1.5], "AAK" station 1265, 35°27.0'N, 11°39.3'E, 70 m, bottom of clay and sand, 26.iv.1978, collector V.V. Murina, MMSU; 1 ♂ [1.8], 1 ovig. [1.9], "AAK" station 1266/63, 35°22.0'N, 11°44.2'E, 65 m, sandy bottom, 26.iv.1978, collector V. V. Murina, MMSU; 2 ♂♂ [1.5, 1.8], 1 ♀ [1.9], 1 ovig. [1.5], "AAK" station 1267, 35°10.0'N, 12°14.4'E, 50 m, bottom of sand and algae, 27.iv.1978, collector V.V. Murina, MMSU; 2 ♂♂ [2.0, 2.2], "AAK" station 1268, 35°10.8'N, 12°34.0'E, 47-50 m, sandy bottom, 27.iv.1978, collector V.V. Murina, MMSU; 8 ♂♂ [1.9-2.6], 3 ovig. [1.7-2.0], "AAK" station 1269/70, 35°13.0'N, 12°38.5'E, 57 m, sandy bottom, 28.iv.1978, collector V.V. Murina, MMSU; 2 ♂♂ [2.1, 2.2], 1 ovig. [1.9], "AAK" station 1273/82, 35°56.3'N, 15°00.0'E, 91 m, bottom of sand and shells, 30.iv.1978, collector V.V. Murina, MMSU; 2 ♂♂ [1.9, 2.0], "AAK" station 1288, 35°38.8'N, 15°1.7'E, 97 m, bottom of clay and sand, 2.ix.1979, collector V.V. Murina, MMSU; 2 ♂♂ [1.8, 1.9], "AAK" station 1295, 35°22.7'N, 15°06.8'E, 80 m, bottom of clay and sand, 4.ix.1979, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 1 ♂ [2.0], 1 ♀ [1.9], "AAK" station 1296, 37°30.7'N, 12°06.5'E, 86 m, sandy bottom, 10.ix.1979, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 2 ♂♂ [2.2, 2.5],

"AAK" station 1297, 37°28.9'N, 12°12.0'E, 62 m, sandy bottom, 11.ix.1979, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 1 ovig. [2.7], "AAK" station 1299 (42/4), 37°28.9'N, 12°12.0'E, 62 m, sandy bottom, 11.ix.1979, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 1 ♂ [2.0], "AAK" station 1311/57, 35°09.2'N, 12°28.2'E, 48 m, sandy bottom, 17.ix.1979, collector V.V. Murina, MMSU; 17 ♂♂ [1.6-2.1], 5 ♀♀ [1.5-2.0], 4 ovig. [1.5-1.7], "AAK" station 1313, 35°17.2'N, 12°04.5'E, 53 m, bottom of sand and *Udotea*, 17.ix.1979, collector V.V. Murina, MMSU; 1 ♂ [1.9], 2 ovig. [1.5, 1.7], "AAK" station 1390, 35°18.3'N, 12°04.4'E, 55 m, sandy bottom, 2.ix.1980, collector V.V. Murina, MMSU; 1 ♂ [1.4], "AAK" station 1392, 35°30.7'N, 12°01.2'E, 72 m, bottom of sand and *Lithothamnium*, 3.ix.1980, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 1 ♂ [2.3], 1 ♀ [1.8], "AAK" station 1394, 60 m, sandy bottom, 6.ix.1980, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 3 ♂♂ [1.9-2.3], "AAK" station 1397, 37°19.3'N, 12°07.9'E, 80 m, sandy bottom, 7.ix.1980, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 1 ♂ [2.1], 2 ovig. [1.9, 2.6], "AAK" station 1398, 37°23.6'N, 12°16.5'E, 70 m, sandy bottom, 7.ix.1980, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 3 ♂♂ [1.4-2.8], "AAK" station 1400, 37°28.1'N, 12°13.5'E, 65 m, bottom of sand and dead and living corals, 8.ix.1980, collectors V.V. Murina & V.D. Tchukhtchin, MMSU. **Southeastern Atlantic: Cape Verde:** 1 ♂ [1.1], 1 ovig. [0.9], off Port Grande (= Mindelo), São Vicente, 16.5 m, 5.iii.1890, NHRM 7362; 1 ♂ [0.8], "Atlantide" station 40, Bay of São Pedro, São Vicente, 32 m, bottom of sand and foraminiferans, 11.xii.1945, ZMUC; 1 ovig. [1.2], "Atlantide" station 43, Bay of Praia, São Tiago, 8 m, bottom of sand and gray silt, 13.xii.1945, ZMUC; 1 ♂ [1.4], 1 ovig. [1.2], "Galathea" station 62, off Namibia, 22°19'N, 17°05'W, 62 m, bottom of shells and bryozoans, 2.xi.1950, ZMUC.

Diagnosis.— Shield slightly longer than broad. Rostrum slightly exceeding lateral projections. Lateral projections with short submarginal spine.

Ocular peduncles approximately $\frac{4}{5}$ length of shield; with corneae slightly dilated. Ocular acicles narrowly triangular, with strong submarginal spine. Interocular plate unarmed.

Antennular peduncles exceeding distal margin of corneae by approximately entire length of ultimate segment. Ultimate antennular segment exceeding length of penultimate segment by approximately $\frac{1}{2}$ own length, with dorsolateral row of short setae and 1 moderately long seta distally; ventral margin with row of tufts of moderately long setae; basal segment with laterodistal spine.

Antennal peduncles approximately reaching distal margin of corneae. Antennal acicles terminating in short spine reaching proximal margin of corneae.

All surfaces of right cheliped with numerous tufts of long or moderately long setae. Chela moderately narrow (males) or broad (females). Dactyl approximately $\frac{2}{3}$ length of palm, triangular in cross section, with elevated, dorsomedial, spinulose ridge and ventromesial, granular ridge. Palm approximately $\frac{4}{5}$ length of carpus, much narrower in males, with spinulose or granular ridge on proximal half of dorsomesial margin; dorsal surface convex, spinulose or granular, with prominent, spinulose ridge (more so in females) near dorsomesial margin proximally; dorsolateral margin with row of spinules extending onto fixed finger but becoming obsolete near claw; dorsoproximal margin with weakly developed lateral and mesial lobes; ventroproximal margin with median, subrectangular sinus. Carpus exceeding merus approximately by $\frac{1}{4}$ own length, much shorter in females; dorsomesial margin with irregular row of closely spaced spines; dorsomesial distal angle produced into moderately large, subcircular structure with a more raised ventral side; dorsodistal margin lacking triangular projection adjacent to dorsomesial distal angle; dorsal surface granular; dorsolateral margin spinulose or granular and with 1, occasionally 2, acute spines; dorsolateral, distal angle produced into small, subcircular lobe; mesial face

slightly compressed, granular or spinulose; ventromesial margin subcircular in mesial view; ventral surface granular or spinulose and with numerous tufts of very long, stiff setae; ventral length between shallow, subcircular, ventrodistal and ventroproximal margins moderately long in males but short in females; lateral face granular or spinulose. Dorsal margin of merus with small distal spine (small specimens) or unarmed (large specimens); ventromesial margin with small distal spine (both sexes) and well-developed setose protuberance (males); ventrolateral margin with 3 to 5 strong spines; ventral surface with spinules or granules and numerous tufts of very long, stiff setae. Ischium with 1 spinule on ventrolateral margin partly covered by tufts of long, stiff setae. Coxae of chelipeds with ventrolateral, distal angle unarmed.

Dactyl of left cheliped smooth but with tufts of long setae. Dorsomesial margin of carpus with row of 4 to 6 strong, widely spaced spines; dorsolateral margin with 3 or 4 strong, widely spaced spines.


Ambulatory legs with dactyls exceeding propodi by approximately $\frac{1}{4}$ own length, with dorsomesial row of widely spaced, moderately long, stiff setae interspersed with short setae; mesial faces each with row of widely spaced, moderately long, spine-like setae interspersed with moderately long setae near ventral margin; lateral faces each with row of widely spaced, moderately long setae near dorsal and ventral margins. Propodi exceeding carpi by approximately $\frac{1}{3}$ own length; dorsal surfaces each with row of tufts of closely spaced, moderately long, stiff setae; mesial and lateral faces each with row of widely spaced, moderately long setae near ventral margin; ventromesial distal angle with 1 moderately long, spine-like seta. Carpi approximately $\frac{2}{3}$ length of meri; dorsal surfaces each with row of tufts of moderately long, stiff setae and 1 proximal and 1 distal spine (2nd) or only 1 distal spine (3rd). Ventral margins of meri with row of tufts of closely spaced, long, stiff setae and 1 spinule on distal half (2nd) or unarmed (3rd). Lobes of sternite of 2nd pereopods unarmed, distally rounded. Anterior lobe of sternite of 3rd pereopods subrectangular (males) or subcircular (females); anterior margin with tufts of long setae. Males with well-developed, left sexual tube, recurved upward to level of coxa; coxa of right 5th pereopod with gonopore on distoventral surface, partly covered by tufts of moderately long setae; three uniramous pleopods. Distal protopodal lobe of right uropod less than $\frac{1}{2}$ length of endopod.

Telson with asymmetrical posterior lobes separated by narrow, shallow cleft; terminal margins each with 3 or 4 short spines; lateral margins unarmed but with tufts of moderately long setae.

Colour.—“Colour reddish brown, uniform all over” (Norman, 1861; as *Pagurus ferrugineus*).

“...; when freshly taken it (as *Anapagurus ferrugineus*) may however be recognized by the bright colour of the antennules, the stalks of which are red, while the flagella are a brilliant blue.” (Henderson, 1885).

Fig. 6. *Anapagurus chiroacanthus* (Lilljeborg, 1856). A-I: ♂ (SL = 2.5 mm), “AAK” station 1297, 37°28.9'N, 12°12.0'E, MMSU: A, shield and cephalic appendages; B, right antennule (lateral view); C, right cheliped (dorsal view); D, carpus of same (ventromesial view); E, left cheliped (dorsal view); F, left 2nd pereopod (mesial view); G, left 3rd pereopod (mesial view); H, sternum; I, extremity of abdomen, uropods, and telson. J: ♂ (SL = 2.0 mm), off Smedjebron, Bohuslän, NHRM 4678: left sexual tube (lateral view). Scales equal 1.0 mm: a: A, C-H, J; b: B, I.



"...; negli esemplari del golfo di Napoli (la cornea) ha colore fondamentale giallicio, con numerose macchie circolari, di colore rosso vermiglio" (Fenizia, 1937: 29-30). (= The corneae of the specimens from the Gulf of Naples are basically yellow with numerous bright red circular spots). "Il colore generale del corpo è di un rosso più chiaro" (Fenizia, 1937: 33; as *A. chiroacanthus* var. *cristatus*). (= The general colour of the body is light red).

Behaviour.— Several aspects of the behaviour of *A. chiroacanthus* have been studied, including dislodging (Hazlett, 1968a), sexual (Hazlett, 1968b; Salmon, 1983), field patterns of shell occupancy (Hazlett, 1981), and feeding (Schembri, 1982).

Related organisms.— The specimens of *A. chiroacanthus* studied by Fenizia (1937) were found inhabiting the gastropod shells of *Turritella*, *Ziziphinus*, and *Nassa* (= *Hinia*); the shell of *Nassa incrassata* (Strom, 1768) (= *Hinia incrassata*) was covered with poriferan *Suberites*. Pike & Williamson (1958) found *A. chiroacanthus* in *Nassarius* (= *Hinia*) and *Gibbula tumida* (Montagu, 1803). In the present collections the following shells were found occupied by *A. chiroacanthus*: Cerithidae (1 specimen, MMSU—"AAK" station 1311-57); *Epitonium clathrus* (Linnaeus) (1 specimen, NHRM 1020); *Epitonium* sp. (2 specimens NHRM 452 and 494; *Gibbula pennanti* (1 specimen, RMNH D 19130; 5 specimens, NHRM 442, 443, 582, 608, and 620); *Gibbula tumida* (1 specimen, USNM 17008); *Hinia reticulata*: (33 specimens, NHRM 1414, 1429, 1542, 1872, 2246, 2549, 3223, 3519, 3682, 4173, 6996 and 6997); *Littorina* sp. (6 specimens, NHRM 1425); Naticidae (2 specimens, NHRM 1414 and 4173); *Ocinebrina aciculata* (5 specimens, NHRM 6976, 6996 and 10575); *Raphitoma linearis* (Montagu, 1803) (1 specimen, NHRM 2469); *Turritella biplicata* (2 specimens, ZMUC; 2 specimens, MMSU—"AAK" station 1390; 12 specimens, NHRM 1390, 1425, 1429, 1531, 1532, 2240, 3223, 4173, 4255 and 4261); and *Turritella communis* (2 specimens, MMSU "AAK" station 1397 and 1398; 4 specimens, NHRM 1531, 2240 and 2549).

Gastropod shells covered by poriferans [e.g., *Suberites domuncula* and/or *S. ficus* (Esper)] and inhabited by *A. chiroacanthus* have been recorded by Aurivillius (1891), Sars (1924), Balss (1956), and Füller (1958). In the present collections 21 gastropod shells occupied by *A. chiroacanthus* were covered by *S. domuncula* (NHRM 1020, 1022, 1041, 2021, 2342, 2549, 3226, 3485, 3519, 3682 and 5772). Other organisms found on shells occupied by *A. chiroacanthus* are: anthozoan *Epizoanthus* sp. (RMNH D 19426); barnacles, bryozoans, foraminiferans, hydrozoans and serpulid polychates (MMSU "AAK" station 1398; NHRM 4255); and *Anomia epippium* (Linnaeus), barnacles and serpulid polychaetes (NHRM 4186).

Anapagurus chiroacanthus has been reported as being infested by two bopyrid isopods, *Pseudione hyndmanni* by Pike (1961), Bourdon (1968), and Markham (1986), and *Athelges tenuicaudis* Sars, 1899, by Elmhirst (1912), Thorson (1946), and Pike (1950, 1961). During the present study, only 9 specimens of the abdominal parasite, *Athelges tenuicaudis* have been found in *A. chiroacanthus* (NHRM 1020, 1021, 1429, 4261 and 4677). One of the females infested (NHRM 1429: SL= 2.0 mm) lacks the right gonopore, and one intersex specimen (NHRM 4677: SL= 1.9 mm) lacks the gonopores and the sexual tube. The other hosts show no external changes.

Two species of rhizocephalans, *Peltogaster paguri* and *Peltogasterella sulcata*, have been recorded as parasites of *A. chiroacanthus* by Balss (1926), Buddenbrook (1948), Høeg & Lützen (1985), Nilsson-Cantell (1926) and Reinhard (1942). The last two

authors described the effects of these parasites in *A. chiroacanthus*. In the present collections, one, and occasionally two, externae per host of *P. paguri* have been found in 31 specimens of *A. chiroacanthus* (MMSU "AAK" stations 1268, 1313 and 1390; RMNH D 19131, 19425, 19431, 19433 and 24171; NHRM 1020, 1022, 1414, 1425, 1427, 1428, 1429, 1531, 3227, 3484, 4255, 4261 and 4678). Eight *A. chiroacanthus* were found with 2-6 externae of *P. sulcata* per host (NHRM 1022, 1428, 1429, 1872, 2549, 2758 and 4678). Most parasitized specimens had varying degrees of changes in the presumed secondary sexual characteristics, such as the absence of pleopods 2 and 3 (under the externae), the feminization of the male pleopods, the reduction or absence of sexual tubes and lack of female gonopores.

Two ovigerous females of *A. chiroacanthus* collected off Tunisia (MMSU "AAK" station 1269-70) had 1 or 2 folliculinid protozoans type 1 (see section on related organisms for *A. hyndmanni*) attached to their abdomens dorsally. This is the first record of folliculinids infesting *A. chiroacanthus*.

On 54 specimens of *A. chiroacanthus* collected in the Kattegat waters (NHRM 1429, 1436, 1532, 1538, 2240, 2250, 2549, 2642, 3224, 3225, 3227, 3484, 3682, 3685, 3686, 4255, 4256, 4261, 4676, 4677 and 4678) and 1 off Ponta Delgada, Azores (NHRM 6996), there were foraminiferans. They usually ranged in number from 1 to 3 tests, occasionally 6 and 9; most tests are attached dorsally to the carpus and palm of the right cheliped and on the shield of the hosts. Occasionally they are found dorsally on the carpus of left cheliped and walking legs. This kind of association with foraminiferans is the first record for the genus *Anapagurus*.

Life history.— Larvae of *A. chiroacanthus* have been studied by Bouvier (1891), Gurney (1942), Thorson (1946), MacDonald, Pike & Williamson (1957), Pike & Williamson (1958, 1960), and Calafiore (1981). Ingle (1990) provided the description of its complete larval and post-larval development. Thorson (1946), and Pike & Williamson (1959) covered the breeding aspect of this species.

Distribution.— *A. chiroacanthus* has been recorded in the northeastern Atlantic from Raunefjorden, southwest of Bergen, Norway (Lagerberg, 1908; Samuelsen, 1970) to off Cape Verde Islands (Forest, 1961). Lilljeborg (1856) described this species as *Pagurus chiroacanthus* found off Kullaberg, southwest of Sweden. This species has also been recorded from the British Seas by Norman (1861; as *Pagurus ferrugineus*), Henderson (1886; as *P. ferrugineus*), Norman (1911), Elmhirst (1912), and Pike & Williamson (1959); off the Azores (A. Milne-Edwards & Bouvier, 1899; as *A. brevicarpus*); off Portugal (Neves, 1977); off Cape Bojador, Morocco (Forest, 1955) and off Rio de Oro (21°30'N, 17°05'W; Forest, 1961).

Additionally, the species has been recorded in the Mediterranean Sea: off the Balearic Islands (Forest, 1965), Bahía de Cadaqués and off Cabo de Creus (Zariquey Alvarez, 1955, 1968), Golfo di Napoli (Balss, 1926; Fenizia, 1937), and in the Aegean Sea, off Turkey (Kocatas, 1981).

The specimen collected off Trondheimsfjorden (63°42'N, 9°27'E); west Norway; RMNH D 24170) extends the northern range of the species. All localities off the coasts of Denmark, France, Monaco, Yugoslavia, Gulf di Taranto and Strait of Sicily are new records for *A. chiroacanthus*. Depth range: 0-200 m; most frequently found in depths of 10 to 60 m.

Affinities.— *Anapagurus chiroacanthus* is similar to *A. japonicus*, sharing the dorso-




Fig. 7. Distribution of *Anapagurus chiroacanthus* (Lilljeborg, 1856) Star: type locality. Solid circles: based on present records. Numbers: based on previous records (circled numbers: material examined): 1-Samuelson, 1970; 2, 4, 7-Norman, 1861; 3-Norman, 1911; 5, 6, 8-Pike & Williamson, 1959; 7-Henderson, 1886; Elmhirst, 1912; 9-Bouvier, 1922; 10-A. Milne Edwards & Bouvier, 1899; 11-Neves, 1977; 12-Forest, 1965; 13-Zariquey Alvarez, 1955; 1968; 14-Balss, 1926; Fenizia, 1937; 15-Kocatas, 1981; 16-Forest, 1955; 17, 18-Forest, 1961.

mesial and ventrolateral ridges of the dactyl of the right chela, and in males three uniramous pleopods. *Anapagurus chiroacanthus* can be separated from *A. japonicus* by the more setose chelipeds and walking legs, the spinose, dorsolateral margin of left cheliped, the much shorter but more numerous spines of the terminal margins of the telson, and in males the much longer, ventrocarpal length of the right cheliped.

Anapagurus chiroacanthus shares the setose appearance of *A. hendersoni* but the former species can be distinguished by the weaker armature of the chelipeds, walking legs, and telson, and the male uniramous pleopods 3 and 4.

Remarks.—Among specimens collected off Kullaberg, Skåne (southwestern Sweden), Lilljeborg (1856) had specimens that he described as *Pagurus chiroacanthus*. Norman (1861), studying the British crustaceans, described a new species, *Pagurus ferrugineus*, which he compared with the British congeners, *P. cuanensis*, *P. thomsoni* (=*P. pubescens*) and *P. laevis*. Previous to this latter work, Stimpson (1858) had erected *Spiropagurus*, the first pagurid genus with extruded vas deferens, so Norman either ignored the Stimpson's taxon, examined only females or simply overlooked the presence of sexual tube(s).

A. Milne Edwards & Bouvier (1892), briefly described *A. brevicarpus* based on two female specimens collected at different localities [north of Spain (170 m) and north of Arguin Bank (135–250 m)]. In 1900, these authors more completely described and illustrated this species. Forest (1961) reexamined the remains (right cheliped, left 3rd pereopod, and abdomen) of the syntype female collected north of Spain ("Travailleur" station 5, 43°59'N, 8°59'W, 170 m, silty bottom, 9.vii.1882, MNHN?) and, together with A. Milne Edwards and Bouvier's illustrations (1900: pl XXV: figs. 23–26, and pl. XXVIII: fig. 11), questionably considered *A. brevicarpus* a junior synonym of *A. chiroacanthus*.

The present extensive collections of *A. chiroacanthus*, which comprise nearly one thousand specimens, have revealed that among other characters, the right cheliped displays strong, morphological and morphometric sexual dimorphisms (e.g. in males the much narrower palm with less developed ridge near mesial margin proximally, and in females the much shorter carpus). Furthermore, I have had the opportunity to reexamine the following specimens (all females) of *A. brevicarpus* from reported collections: (1) the remains (right 2nd and 3rd pereopods) of the other syntype specimen collected north of Arguin Bank ("Talisman" station 90, 21°51'N, 19°48'W, 135–250 m, bottom of greenish sand and silt, 13.vii.1883, MCZ 6579); (2) the specimen reported by A. Milne Edwards & Bouvier (1899) collected off Ponta do São Antonio, Terceira, Azores ("Princess Alice" station 594, 54 m, rocky bottom, 22.vii.1895, MOM 380322); and (3) the specimen cited by Bouvier (1922) and collected off Paragés de Belle Ile ("Princess Alice II" station 2974, 47°23'N, 3°25'W, 85 m, plateau bottom, 27.vii.1910, MOM 380997). I have concluded that all of these specimens are conspecific with *A. chiroacanthus*.

The specimens that Mouchet (1931) used in her studies of the spermatophores of *A. brevicarpus*, and those reported by Cardona Bendito (1965) from Ría de Vigo, Spain, have not been available for examination, and therefore are questionably considered conspecific with *A. chiroacanthus*.

Anapagurus pusillus Henderson, 1888, sensu stricto
(figs. 8, 9)

Anapagurus pusillus Henderson, 1888: 73 (in part), pl. VII: fig. 7 (type locality, herein restricted by lectotype selection: "Challenger" station VII p, 28°35'00"N, 16°05'00"W, off Gomera Island, Canary Islands, 143 m, bottom of volcanic sand); Alcock, 1905: 187 (in part); Balss, 1924: 768; Monod, 1932: 487 (in part); Gordan, 1956: 303 (in part; see remarks).

not *Anapagurus pusillus*; Stebbing, 1910: 358 (= *A. hendersoni* Barnard, 1947); Balss, 1912: 110 (= *A. hendersoni* Barnard, 1947); Odhner, 1923: 13 (= *A. longispina* A. Milne-Edwards & Bouvier, 1900) (see remarks).

Anapagurus laevis; Forest, 1955: 131 (in part; see remarks).

Material.—Lectotype herein selected: ♂ [2.1], off Gomera Island, Canary Islands, "Challenger" station VII p: 28°35'00"N, 16°05'00"W, 143 m, bottom of volcanic sand, 10.ii.1873, BMNH 1888: 33; Paralectotypes: 2 ♂♂ [1.5, 1.7], 1 ♀ [1.7], 2 ovig. [1.7, 1.8], data as for lectotype. Off Azores: 1 ♀ [1.7], "Josephine Expedition" station e16, off Villa Franca, São Miguel, 54.9-91.4 m, bottom of gravel, 26.vii.1869, NHRM 6978; 1 ♂ [1.5], 1 ♀ [0.7], 2 ovig. [1.0, 1.6], "Josephine Expedition" station e54 (g), off Faial, up to 27.4 m, 1869, NHRM 6986; 2 ♂♂ [1.2, 2.6], 1 ♀ [2.3], 5 ovig. [1.7-2.0], "Josephine Expedition" station e62 (a), off Faial, 54.9-146.3 m, bottom of black gravel and shells, 3.viii.1869, NHRM 6989; 1 ovig. [0.9], "Josephine Expedition" station d 68 (c), 18.3m, bottom of gravel, 1869, NHRM 6995. Off S Portugal: 1 ovig. [3.0], Ria de Faro, near Cabo de Arnel, 29.vii.1962, RMNH D 19365.

Diagnosis.—Shield broader than long. Rostrum slightly overreached by lateral projections. Lateral projections with short, submarginal spine.


Ocular peduncles $\frac{2}{3}$ to $\frac{3}{4}$ length of shield; with corneae moderately dilated. Ocular acicles roundly triangular, with strong submarginal spine. Interocular plate unarmed.

Antennular peduncles exceeding distal margins of corneae by approximately $\frac{1}{2}$ length of ultimate segment. Ultimate segment exceeding length of penultimate segment by $\frac{1}{4}$ own length, with 1 long and 1 short seta on dorsolateral, distal angle; ventral margin with row of tufts of long setae; basal segment with lateromedial and laterodistal spines.

Antennal peduncles overreaching ocular peduncles by approximately $\frac{1}{3}$ length of ultimate segment. Antennal acicles usually not reaching corneae proximally.

Right cheliped with moderately broad chela. Dactyl $\frac{2}{3}$ to $\frac{3}{4}$ length of palm; dorsomesial margin with row of tubercles or spinules; all surfaces with tufts of short setae. Palm slightly longer than carpus; dorsomesial margin with slightly developed ridge on proximal half with row of closely spaced tubercles and tufts of short setae; dorsal surface convex, with scattered granules and tufts of short setae, and short, tuberculate ridge near dorsomesial margin proximally; dorsolateral margin with spinules increasing in size on fixed finger but not extending to claw; dorsoproximal

Fig. 8. *Anapagurus pusillus* Henderson, 1888. Lectotype ♂ (SL = 2.1 mm), BMNH-1888: 33: A, shield and cephalic appendages; B, left antennule (lateral view); C, right cheliped (dorsal view); D, fingers of same showing dentition of fixed finger; E, fingers of same showing dentition of dactyl; F, carpus of same (ventral view); G, carpus of same (mesial view); H, left cheliped (dorsal view); I, left 2nd pereopod (mesial view); J, left 3rd pereopod (mesial view); K, sternum; L, extremity of abdomen, uropods, and telson. M: ♂ (SL = 2.6 mm), "Josephine Expedition" station e 62(a), off Faial, Azores, NHRM 6989: left sexual tube (lateral view). Scales equal 1.0 mm: a: C, F-J, M; b: A; c: B, D, E, K, L.



margin with weakly developed, mesial and lateral lobes; ventroproximal margin with a slightly calcified, subcircular sinus on either side of median, skewed, subquadrate sinus. Carpus equaling or slightly exceeding length of merus; dorsomesial margin with row of strong acute spines; dorsomesial, distal angle produced into concave, subrectangular structure; dorsodistal margin with triangular projection adjacent to dorsomesial distal angle; dorsolateral margin with row of spinules; lateral face unarmed; ventromesial margin subcircular in mesial view; ventral surface with tufts of stiff setae, long in males, short in females; short ventral length between deep, subtriangular, ventrodistal margin and subcircular, ventroproximal margin. Ventromesial margin of merus with tuberculate protuberance (males) or tubercles (females); ventral surface with tubercles, setation denser and longer in males; dorsal margin with acute, distal spine. Coxae of chelipeds with ventrolateral, distal angle unarmed.

Dactyl of left cheliped smooth but with tufts of moderately long setae. Dorsomesial and dorsolateral margins of carpus, each with 3 or 4 strong, widely spaced spines.

Dactyls of ambulatory legs exceeding propodi by approximately $\frac{1}{4}$ their own length, with dorsomesial and ventromesial rows of widely spaced, short, stiff setae. Propodi exceeding carpi by $\frac{1}{3}$ own length; ventromesial distal angle with 1 or 2 spines (2nd) or unarmed (3rd). Carpi $\frac{1}{2}$ to $\frac{2}{3}$ length of meri; dorsal margins with 2-4 spines (2nd) or single, distal spine (3rd). Meri unarmed. Lobes of sternite of 2nd pereopods unarmed, distally rounded. Anterior lobe of sternite of 3rd pereopods broadly subrectangular, slightly skewed to left.

Males with moderately long and slender, left sexual tube, recurved upward to exterior, and vestigial right sexual tube; pleopods 3 and 4 unequally biramous, pleopod 5 uniramous. Distal protopodal lobe of right uropod approximately $\frac{1}{2}$ length of endopod.

Telson with posterior, asymmetrical lobes, separated by wide, shallow cleft; terminal margins each with 3 or 4 short spines.

Colour.— Unknown.

Related organisms.— "The majority (of the specimens were) firmly ensconced in shells of *Turritella bicingulata*, Lam., one individual in a shell of *Nassa limata*, Chem." (Henderson, 1888: 74). In the present collections, specimens of *A. pusillus* were found inhabiting the following gastropod shells: *Natica catena* (4 specimens, NHRM 6986 and 6989); *Calliostoma zizyphinum* (Linnaeus, 1758) (2 specimens, NHRM 6986); *Hinia reticulata* (3 specimens, two of the shells partly covered with bryozoans and serpulid polychaetes, NHRM 6989); and *Cerithiopsis tubercularis?* (Montagu) (1 specimen, NHRM 6995).

Distribution.— Besides the type locality, *Anapagurus pusillus* has been collected off the Azores and off the south coast of Portugal. Depth range: 0-146.3 m.

Affinities.— *Anapagurus pusillus* is most closely allied to *A. breviaculeatus* and *A. alboranensis* spec. nov.. The much broader shield and much shorter, ultimate antennular segment with heavily setose ventral margin easily distinguish *A. pusillus* from *A. breviaculeatus* and *A. alboranensis* spec. nov.

Remarks.— Henderson (1888) described *A. pusillus* using specimens from the Azores and Gomera (Canary Islands). He also referred a male specimen from Simon's Bay (False Bay, South Africa) with "some hesitation" to this species. Barnard (1947)




Fig. 9. Distribution of *Anapagurus pusillus* Henderson, 1888. Circled star: type locality, material examined. Solid circles: based on present records.

examined the Simon's Bay specimen, decided it was distinct from Henderson's taxon, and described it as a new species, *A. hendersoni*.

Stebbing (1910) quoted Henderson's (1888) description of *A. pusillus* when referring to the Simon's Bay specimen. As previously indicated, Barnard (1947) demonstrated that this specimen was distinct from *A. pusillus*. Balls (1912) reported *A. pusillus* from two specimens collected by the Valdivia off Cape Saint Francis, South Africa. My examinations of these specimens show that they belong to *A. hendersoni*. My examination of a female specimen from Port Alexander, Angola, identified by Odhner in 1923 as *A. pusillus*, shows that it belongs to *A. longispina*.

Forest (1955) identified specimens from the southwestern coast of Africa collected during expeditions of the "Mbizi" and "Mercator" as *A. laevis*. After examining Henderson's (1988) specimens of *A. pusillus* from the Azores and Canary Islands, Forest (1955) concluded that these too represented *A. laevis* and placed Henderson's (1888) taxon in synonymy. He did, however, concur with Barnard (1947) that Henderson's (1888) Simon's Bay specimen represented a distinct species. Having personally reexamined this specimen, I agree that it was appropriately designated a separate species. It agrees well with Barnard's (1950) supplemental material from the area, excepting one ovigerous female that represents an unidentified species of *Pagurus*. However, reexamination of Henderson's (1888) specimens from the Canary Islands has convinced me that they are not conspecific with *A. laevis*. Although I have not examined Henderson's material from the Azores, *A. pusillus* is reinstated, based upon the lectotype and paralectotypes herein designated.

The listings of Alcock (1905), Monod (1932), and Gordan (1956) included two species under the name *A. pusillus*.

Anapagurus japonicus Ortmann, 1892
(figs. 10, 11)

Anapagurus pusillus var. *japonica* Ortmann, 1892: 296, pl.12: fig. 11 [type locality: Kadsiyama (Katsuyama), Uraga Strait, Japan, 73 m].

Anapagurus pusillus var. *japonicus*; Alcock, 1905: 187 (listed); Gordan, 1956: 303 (listed).

Anapagurus japonicus; Miyake, 1961: 24; 1978: 139, fig. 55; Miyake, Sakai & Nishikawa, 1962: 126 (listed).

Anapagurus japonisus; Miyake, 1965: 648, fig. 1099 (misspelling of *A. japonicus*)

Material.— Holotype: ♂, Kadsiyama (Katsuyama, Uraga Strait), Honshu, Japan, 73 m, 1880, collector Döderlein, unknown deposition; not examined. Japan: 1 ♂ [1.9], off Uzé, Tomioka, Kumamoto Prefecture, Kyushu, 29.v.1967, collector T. Fujino, ZLKU 11976; 1 ♀ [1.3], Kushimotochō, Wakayama Prefecture, Honshu, iii.1977, collector H. Uchida, OMNH Ar 2127.

Diagnosis.— Shield approximately as long as broad. Rostrum approximately equaling lateral projections. Lateral projections with short submarginal spine.


Ocular peduncles approximately $\frac{3}{4}$ length of shield, with corneae slightly dilated. Ocular acicles broadly triangular, with strong submarginal spine. Interocular plate unarmed.

Antennular peduncles overreaching ocular peduncles by approximately entire length of ultimate segment. Ultimate segment exceeding length of penultimate segment of approximately $\frac{1}{3}$ own length, with dorsolateral row of short setae and 2 laterodistal setae, 1 long and 1 short, ventral margin with row of moderately long setae.

Antennal peduncles slightly exceeding length of ocular peduncles. Antennal acicles overreaching proximal margin of corneae.

Right cheliped massive, with broadly oval chela. Dactyl approximately $\frac{1}{2}$ length of palm, triangular in cross section, with elevated, dorsomedial, granular ridge and ventromesial, spinulose ridge, all surfaces granular and with tufts of short setae. Palm slightly longer than carpus, dorsomesial margin with well-developed spinulose ridge on proximal half; dorsal surface convex, with spinules (more numerous and larger near mesial margin), and very long and prominent, granular ridge near mesial margin proximally; dorsolateral margin with row of granules not extending to claw; dorsoproximal margin without mesial and lateral lobes; ventroproximal margin with skewed, median, subrectangular sinus; all surfaces with scattered tufts of short setae. Carpus exceeding merus by $\frac{1}{3}$ of its own length, inflated, semicircular ventrally; dorsomesial margin with irregular row of closely spaced spines and spinules and moderately long setae; dorsomesial, distal angle produced into very large, spinulose, concave, subrectangular structure; dorsodistal margin lacking triangular projection adjacent to dorsomesial, distal angle; dorsal surface spinulose; dorsolateral margin unarmed but with row of widely spaced, short, stiff setae; dorsolateral,

Fig. 10. *Anapagurus japonicus* Ortmann, 1892. ♂ (SL = 1.9 mm), off Uzé, Kyushu, Japan, ZLKU 11976: A, shield and cephalic appendages; B, left antennule (lateral view); C, right cheliped (dorsal view); D, carpus of same (ventral view); E, carpus of same (mesial view); F, left cheliped (dorsal view); G, right 3rd pereopod (mesial view); H, sternum; I, extremity of abdomen, uropods and telson; J, left sexual tube (lateral view). Scales equal 1.0 mm: a: J; b: A, C-I; c: B.



distal angle produced into small lobe; mesial face compressed, spinulose and with moderately long setae; ventromesial margin semicircular in mesial view; ventral surface spinulose and with tufts of moderately long setae; very short ventral length between deep, subtriangular, ventrodistal margin and deep, subcircular, ventroproximal margin; lateral face convex, spinulose. Dorsal margin of merus with row of tufts of moderately long setae and partially obscured distal spines; ventromesial margin spinulose (females) or with prominent, spinose protuberance (males); ventrolateral margin with 3 or 4 strong spines; ventral surface with spinules and scattered tufts of moderately long setae. Ischium unarmed but with scattered tufts of short setae. Coxae of chelipeds with ventrolateral, distal angle unarmed.

Dactyl of left cheliped smooth but with tufts of moderately long setae. Dorsomesial margin of carpus with 6 or 7 strong closely spaced spines; dorsolateral margin unarmed.

Right 3rd pereopod of one male (ZLNU 11976) (only ambulatory leg examined) with dactyl exceeding propodus by approximately $\frac{1}{6}$ own length, with mesial face with row of closely spaced, moderately long, stiff setae near dorsal and ventral margins; lateral face with row of widely spaced, short setae near dorsal and ventral margins. Propodus exceeding carpus $\frac{1}{2}$ own length; dorsal surface with row of tufts of short setae; mesial and lateral faces each with row of short setae near ventral margin; ventromesial, distal angle with short spine and tuft of short setae. Carpus approximately $\frac{2}{3}$ length of merus; dorsal surface with row of moderately long setae terminating in strong spine. Merus unarmed but dorsal and ventral surfaces each with tufts of moderately long setae. Lobes of sternite of 2nd pereopods unarmed, distally rounded. Anterior lobe of sternite of 3rd pereopods trapezoidal; anterior margin with tufts of long setae.

Male with left sexual recurved upward to level posterior carapace distally; coxa of 5th right pereopod with gonopore on distoventral surface, obscured by tufts of moderately long setae; three uniramous pleopods. Distal protopodal lobe of right uropod approximately $\frac{1}{2}$ length of endopod.

Telson with asymmetrical, posterior lobes separated by wide, shallow cleft; terminal margins each with 2 strong spines; lateral margins unarmed but with tufts of moderately long setae.

Colour.— "(in formalin). Body and legs uniformly light yellow-brown" (Miyake, 1978: 141).

Related organisms.— The male holotype was found inhabiting a trochid shell of *Ziziphinus unicus* Dunker (Ortmann, 1892: 297). Miyake (1978), in his study of the Anomura of Sagami Bay, recorded a male *A. japonicus* inhabiting the gastropod shell of *Lancea* (= *Scaliola*) *bella* A. Adams; the hermit crab was infested with a bopyrid isopod in the right branchial chamber. He also found two males in the gastropod shells of *Minolia punctata* A. Adams and *Dimidaculus stramineus* (Gray). In the present collection, an undetermined bopyrid isopod was found within the right branchial chamber of a female *A. japonicus* (OMNH Ar 2127). Except for the typical swelling of the branchiostegite, no other external changes in the host were observed.

Distribution.— *Anapagurus japonicus*, an apparently endemic Japanese species, has been recorded by Miyake (1961, 1965, 1978) off Amakusa, Western Kyushu (20-50 m), Chijiwa Bay, Nagasaki-ken (27-72 m), off Tsuyazaki, Sea of Genkai (28-39 m), and




Fig. 11. Distribution of *Anapagurus japonicus* Ortmann, 1892. Star: type locality. Solid circles: based on present records. Numbers: based on previous records: 1-4-Miyake, 1978; 4-Miyake, 1961.

Sagami Bay (50-80 m). Depth range: 20-80 m.

Affinities.—As earlier noted, *A. japonicus* appears most closely related to *A. chiroacanthus*. *Anapagurus japonicus* may be distinguished from the latter by its less setose chelipeds and ambulatory legs, the unarmed, dorsolateral margin of the left cheliped, and the much longer but fewer spines of the terminal margins of the telson.

Remarks.—Ortmann (1892) recorded a male from Kadsiyama, Japan as a new variant of *Anapagurus pusillus*, and this taxon was subsequently listed as such by Alcock (1905) and Gordan (1956). Later, Miyake (1961) elevated it to specific rank, an action with which I concur.

Since S. Miyake retired from Kyushu University 18 years ago its decapod collection has been neglected. Furthermore, the collections were moved to another building, resulting in many specimens being misplaced, thus there appears to be little chance to find the specimens of *A. japonicus* listed by Miyake under ZLKU numbers (K. Baba, personal communication). Only two specimens of *A. japonicus* were available for my study, one male (ZLKU 11976) and one female (OMNH Ar 2127). The present diagnosis of the right, third pereopod, the only leg present in the male was provided by that specimen.

Anapagurus curvidactylus Chevreux & Bouvier, 1892
(figs. 12, 13)

Anapagurus curvidactylus Chevreux & Bouvier, 1892a: 253; 1892b: 91-93, pl. II: figs. 2-8 (type locality, herein restricted by lectotype selection: "Melita" station 109, west off Gorée, Senegal, bottom of fragmented shells, 16.iii.1890); Bouvier, 1896: 152, figs. 33-35 (pictorial key); 1940: 149, 150, figs. 103, 107B (in part); Rathbun, 1900: 307 (listed); Alcock, 1905: 187 (in part); Balsas, 1921: 46; Monod, 1932: 487; Forest, 1955: 128, fig. 29, pl. VI: fig. 2; 1961: 241, figs. 9, 13, 18 (in part; see remarks); 1966: 162; Gordan, 1956: 302 (listed); Cardona Bendito, 1965: 146; Anadón, 1975: 206, 214; Zariquey Alvarez, 1968: 258; Ingle, 1985: 764 (listed).

Anapagurus curvidactylus var.? Forest, 1955: 129, fig. 30, pl. VI: figs. 3-5 (see remarks).


not *Anapagurus curvidactylus*; A. Milne Edwards & Bouvier, 1892: 214, 215; 1900: 225 (=*A. hyndmanni*); De Miranda y Rivera, 1933: 30, 60 (=*A. hyndmanni*) (listed).

Material.—Lectotype, herein selected: ♂ [2.0], "Melita" station 108, off Pointe Belair, Baie de Dakar, Senegal, 8 m, bottom of sand and mud, 15.iii.1890, MCZ 6578; Paralectotype: 1 ♂ [2.0], "Melita" station 109, W off Gorée, Senegal, 15 m, bottom of fragmented shells, 16.iii.1890, USNM 22921. **Eastern Atlantic: Europe:** 10 ♂♂ [1.8-2.1], 1 ♀ [1.9], 3 ovig. [1.2-1.6], "Flor da Ponte" station 1.634 (=F.12), 0.4 km off Los Mezos, Ría de Arosa, N.W. Spain, 18-22 m, 22.vii.1964, RMNH D 24262; 1 ♂ [2.5], "Flor da Ponte" station 1.656, 0.7 km off Punta Quilme, Isla de Arosa, Ría de Arosa, N.W. Spain, 3 m, 24.vii.1964, RMNH D 24265; **Africa:** 2 ovig. [1.0, 1.4], "Mercator", 7.4 km off Angra da Cintra, Rio de Oro, 27-36 m, 30.xii.1937, IRSNB-IG 11.553; 1 ♂ [1.2], "Mercator", along coast between Dakar and Hann, Baie de Dakar, Senegal, 13.xi.1935, IRSNB-IG 10.910; 2 ♂♂ [1.7, 1.8], 1 ovig. [1.3], Anse Bernard, Senegal, 8-10 m, 13.ii.1953, collector J. Forest, MNHN 488; 1 ovig. [1.5], "Atlantide" station 44, off French Guinea, 10°22'N, 16°22'W, 41 m, bottom of sand and shells, 17.xii.1945, ZMUC; 1 ovig. [1.4], "Atlantide" station 45, off French Guinea, 9°23'N, 15°07'W, 30-34 m, sandy bottom, 18.xii.1945, ZMUC; 1 ♂ [1.8], "Atlantide" station 147, off French Guinea, 9°28'N, 14°58'W, 45 m, 14.iv.1946, ZMUC; 1 ♀ [0.8], 3 ovig. [1.0-1.1], "Mercator", 3.7 km off l'Ile Annobon, 16-35 m, 29.i.1938, MNHN 492; 3 ♂♂ [1.0-1.5], "Mercator", Baie des Eléphants, Angola, 20 m, 4.ii.1937, IRSNB-IG 11.204.

Diagnosis.—Shield slightly broader than long. Rostrum somewhat exceeding lateral projections. Lateral projections with short submarginal spine.

Ocular peduncles approximately $\frac{3}{4}$ length of shield, with cornea slightly dilated. Ocular acicles narrowly triangular, with short, distomesial, submarginal spine. Interocular plate unarmed.

Fig. 12. *Anapagurus curvidactylus* Chevreux & Bouvier, 1892. Lectotype ♂ (SL = 2.0 mm), MCZ 6578: A, shield and cephalic appendages; B, left antennule (lateral view); C, right cheliped (dorsal view); D, carpus of same (ventral view); E, carpus of same (mesial view); F, left cheliped (dorsal view); G, right 2nd pereopod (mesial view); H, right 3rd pereopod (mesial view); I, sternum; J, extremity of abdomen, uropods and telson; K, left sexual tube (lateral view). Scales equal 1.0 mm: a: C-H, K; b: A, I; c: B, J.



Antennular peduncles exceeding distal margin of corneae approximately $\frac{3}{4}$ length of ultimate segment. Ultimate antennular segment exceeding length of penultimate segment by approximately $\frac{3}{5}$ own length, with 1 moderately long, dorsolateral, distal seta; ventral margin of ultimate and penultimate segments, each with row of tufts of setae, long, but not exceeding length of ventral antennular flagellum; basal segment with lateromedial spine.

Antennal peduncles overreaching distal margin of corneae approximately by $\frac{2}{5}$ length of ultimate segment. Antennal acicles terminating in short spine, approximately reaching proximal half of corneae.

Chela of right cheliped narrow (males) or moderately broad (females). Fingers curved ventrally, strongly so in mature males. Dactyl subequal to length of palm, triangular in cross section, with dorsomedial and ventromesial, granular or spinulose ridges; all surfaces with small spinules; ventral surface with tufts of long setae. Palm approximately $\frac{3}{5}$ length of carpus, with granular or spinulose ridge on proximal half of dorsomesial margin; dorsal surface somewhat convex, with numerous, small spinules and scattered, short setae, without ridge near dorsomesial margin proximally; dorsolateral margin with row of spinules extending onto fixed finger but becoming obsolete near claw; dorsoproximal margin with well-developed lobes, laterally and mesially, larger mesial lobe with concavity to fit the dorsomesial, distal lobe of carpus when chela is flexed; ventroproximal margin with skewed, subquadrate sinus near ventromesial margin. Carpus exceeding merus by approximately $\frac{2}{5}$ own length, shorter in females; dorsomesial margin with irregular row of closely spaced, short spines; dorsomesial, distal angle produced into concave, subcircular structure; dorso-distal margin with weak subcircular projection adjacent to dorsomesial, distal angle; mesial face somewhat compressed; dorsolateral face rounded, slightly inflated, without delimiting, dorsolateral margins; dorsolateral, distal angle produced into small, subcircular lobe; ventromesial margin subcircular in mesial view; moderately long, ventral length between shallow, subcircular, ventrodistal and ventroproximal margins; mesial and dorsolateral faces each with small spinules and scattered short setae; ventral surface with spinules and scattered, stiff, short (females) or long (males) setae. Dorsal margin of merus with row of moderately long setae and short, distal spine; ventromesial margin with short, distal spine and slightly developed protuberance (males); ventrolateral margin with 4 to 6 strong spines; ventral surface with spinules and scattered setae, short (females) or long (males). Ischium with 1 spinule on ventrolateral margin partly covered by tufts of setae, short (females) or long (males); ventromesial margins minutely serrate. Coxae of chelipeds with 1, occasionally 2, short, ventrolateral distal spines.

Dactyl of left cheliped smooth but with tufts of long, stiff setae. Dorsomesial margin of carpus with row of 8 to 10 short, closely spaced spines on distal half; dorsolateral margin with row of 8 to 10 short, closely spaced spines becoming obsolete distally.

Ambulatory legs with dactyls exceeding propodi by approximately $\frac{1}{4}$ own length, with lateral and mesial faces each with row of widely spaced, short setae near dorsal and ventral margins, dorsomesial setae stiffer. Propodi exceeding carpi by nearly $\frac{1}{3}$ own length; dorsal surfaces serrate, each with row of tufts of widely spaced short, stiff setae; lateral and mesial faces each with row of widely spaced, short setae near dorsal and ventral margins; ventromesial distal angle unarmed but with 1 short,

simple seta. Carpi approximately $\frac{2}{3}$ length of meri; dorsal surfaces with row of tufts of short, stiff setae and short spines increasing in size distally (2nd) or 1 to 3 distal spines (3rd). Dorsal surfaces of meri each with row of tufts of moderately long setae; ventral margins with row of spinules and tufts of long setae (2nd) or unarmed but with row of moderately long setae (3rd). Lobes of sternite of 2nd pereopods unarmed, distally rounded. Anterior lobe of sternite of 3rd pereopods subrectangular; anterior margin with tufts of moderately long setae.

Males with very long, left sexual tube, strongly recurved to level dorsodistal margin of branchiostegite; coxa of right 5th pereopod with gonopore on distoventral surface, partly covered by tufts of moderately long setae; three uniramous pleopods. Right distal protopodal lobe equaling length of uropodal endopod.

Telson with posterior, asymmetrical lobes separated by wide, shallow cleft; terminal margins each with 3 or 4 short spines; lateral margins unarmed but with tufts of moderately long setae.

Colour.— "L'animal est blanc avec une bande brun jaunatre à la base des doigts des pattes ambulatoires, ainsi qu'à la base et au sommet du propode des mêmes pattes". (Chevreux & Bouvier, 1892).

Related organisms.— In the present collections, specimens of *A. curvidactylus* have been found inhabiting the following gastropods shells: *Hinia reticulata* (5 specimens, RMNH D 24262); and *Turritella biplicata* (1 specimen, MNHN 492).

One ovigerous female of *A. curvidactylus* collected at station 44 by the "Atlantide" (ZMUC) was found in a gastropod shell so encrusted by a colonial bryozoan, that it extend past the shell aperture in the form of a helicospiral coil (Taylor & Cook, 1981). According to Taylor *et al.* (1989), this association between hermit crab and bryozoan is probably mutualistic.

On three males of *A. curvidactylus* collected in Ría de Arosa, Spain (RMNH D 24262), there were folliculinid protozoans type 1 (see section on related organisms for *A. hyndmanni*); tests were cemented dorsally to the abdomen, telson and protopod of the right uropodal exopod. Colonial hydrozoans were found dorsally on the carpus of the right cheliped on one of these specimens. No external changes caused were observed that could be attributed to these symbionts. The above are the first records of folliculinids and hydrozoans associated with *A. curvidactylus*.

Distribution.— Eastern Atlantic: from Ría de Arosa, northwest Spain to Baie des Eléphants, Angola. Depth range: 0-45 m.

Affinities.— As previously stated, *A. curvidactylus* is most closely related to *A. hyndmanni*. The former species can be distinguished by the much shorter ultimate antennular segment bearing ventral setae shorter than the ventral antennular flagellum, the shorter antennal acicles, the absence of longitudinal sulci on the lateral and mesial faces of the ambulatory dactyls, and the much longer, left sexual tube.

Remarks.— In a preliminary note on the pagurids collected by the "Melita" off the Canary Islands and Senegal, Chevreux & Bouvier (1892a) briefly described *A. curvidactylus*. Subsequently (1892b), they more completely described and illustrated this taxon. A. Milne Edwards & Bouvier (1892, 1900) identified a male specimen from the "Travailleur" collections off Barquero, Bay of Viscay, Spain, as *A. curvidactylus*; however my subsequent examination of this specimen has shown it to represent *A. hyndmanni*. De Miranda y Rivera (1933) just listed A. Milne Edwards & Bouvier's




Fig. 13. Distribution of *Anapagurus curvidactylus* Chevreux & Bouvier, 1892. Circled star: type locality, material examined. Solid circles: based on present records. Numbers: based on previous records (circled numbers: material examined): 1-Cardona Bendito, 1965; 2?-Zariquey Alvarez, 1968 (questionable material); 3, 5, 8, 10-Forest, 1955; 6-Chevreux & Bouvier, 1892a, 1892b; 7-Forest, 1961; 9-Forest, 1966.

(1892) reference of *A. curvidactylus* from the Cantabric region, which indeed refers to *A. hyndmanni*. References by other authors (Alcock, 1905; Bouvier, 1940) who cited A. Milne Edwards & Bouvier's work (1892, 1900) are listed as "in part" in the synonymy. I have also had the opportunity to examine the four females collected by the "Mercator" along the Island of Annobon that Forest (1955) doubtfully assigned to a variety of *A. curvidactylus*, and have confirmed that they indeed do belong to Chevreux & Bouvier's taxon. I have additionally examined the specimens collected by the "Atlantide" along the coasts of tropical west Africa that Forest (1961) doubtfully identified as *A. curvidactylus*, and have found that they are conspecific with *A. wolffi*.

Anapagurus bicorniger A. Milne Edwards & Bouvier, 1892
(figs. 14, 15)

Anapagurus bicorniger A. Milne Edwards & Bouvier, 1892: 215, 216 (type locality: Golfo de Cádiz, "Talisman" station 2, 100 m); 1898: 1247 (listed); 1899: 66; 1900: 220-223, pl. XXV: figs. 14-22; Bouvier, 1896: 152, fig. 30; 1940: 152, figs. 106, 107 E; Alcock, 1905: 187 (listed); Balss, 1926: 375-377, fig. 1; De Miranda y Rivera, 1933: 30, 60 (listed); Fenizia, 1937: 33, 34 (in part; see remarks), fig. 24, not pp. 35, 36, figs. 25, 26 (= *A. petitii*); Gordan, 1956: 302 (listed); Forest, 1958: 100; Holthuis & Gottlieb, 1958: 73; Pike & Williamson, 1960: 528-531, fig. 9 (larvae); Dechancé & Forest, 1962: 294-296, figs. 1, 3-5, 9, 11 (in part; see remarks); Zariquey Alvarez, 1968: 259 (in part; see remarks); Gilat, 1969: 16, 54; Neves, 1977: 182, 183; Calafiore, 1981: 20 (larvae); Kocatas, 1981: 162; García Raso, 1982: 503, 504, fig. 4; Ingle, 1985: 764 (listed).

Not *Anapagurus bicorniger*; Zariquey Alvarez, 1946: 122, 123 (=*A. petitii*); 1955: 402 (=*A. petitii*).

Material.—Syntypes: "Talisman" station 2, Golfo de Cádiz, South off Cabo de São Vicente, 36°53'N, 10°52'W, 99 m, bottom of silt and shells, 6.vi.1883: 2 ♂♂ [5.8, 6.0], 1 ovig. [2.0] (USNM 22920); 2 ♂♂ [1.2, 2.1], 2 ovig. [2.2] (MCZ 6577); 1 ♀ [1.8], 1 ovig. [2.1] (RMNH D 1658); 1 ♂ [2.4], 1 ♀ [2.4], 1 ovig. [2.0] (ZMUC). Mediterranean Sea: Off Balearic Islands: 1 ovig. [1.3], "AAK" station 1364, 38°43'N, 2°30.8'E, 125 m, bottom of *Lithothamnium* with sand and clay, 27.iv.1980, collector N.A. Valovaya, MMSU. Golfo de Valencia: 1 ♂ [1.5], 1 ♀ [1.4], 1 ovig. [1.9], "AAK" station 1247, 40°26.2'N, 1°09.2'E, 90 m, bottom of clay and sand, 9.iv.1978, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 1 ovig. [1.4], "AAK" station 1251, 40°17.0'N, 0° 46.9'E, 77 m, bottom of sand and clay, 15.iv.1978, collectors V. V. Murina & V.D. Tchukhtchin, MMSU; 1 ovig. [1.8], "AAK" station 1252, 40°11.6'N, 0°55.4'E, 90 m, bottom of sand and clay, 15.iv.1978, collector V.D. Tchukhtchin, MMSU; Golfo de Rosas: 1 ovig. [2.2], South of Cabo de Creus, 73-91 m, 3.ix.1945, collector R. Zariquey Alvarez, MNHN 469; Ligurian Sea: 1 ovig. [2.1], "Princess Alice" station 633, 43°44'10"N, 5°06'50"E, 69 m, 12.v.1896, MOM 38-0353; 1 ♂ [1.9], 2 ovig. [1.8, 2.2], off Roquebrune, 72-80 m, 30.vii.1963, RMNH D 19707; 1 ♂ [2.2], off Portofino, 1914, collector L. Nick, SMF 8764. Tyrrhenian Sea: 1 ♂ [2.1], 2 ♀♀ [1.7, 2.0], "Dana" station 4088, off Castellamare, 64 m, 26.v.1930, collector R. Spack, ZMUC. Adriatic Sea: 4 ♂♂ [2.1-2.5], 3 ♀♀ [1.9-2.2] off Rodi Garganico, 20 m, 8.viii.1978, collector C. Froglia, IRPEM-D 589; 1 ♂ [2.5], 1 ovig. [2.2], 42°25'N, 14°30'E, 60-65 m, 18.viii.1978, collector C. Froglia, IRPEM-D598; 2 ♂♂ [1.8, 2.0], 3 ovig. [2.1-2.4], 43°30'N, 13°48'E, 20 m, 18.x.1972, collector C. Froglia, IRPEM-D382; 2 ♂♂ [2.7], 1 ovig. [2.2], off Ancona, 13 m, v.1982, collector C. Froglia, MSNM 2384; 1 ♂ [2.6], 43°38.1'N, 13°26.4'E, 5 m, 12.iv.1973, IRPEM; 1 ♂ [2.7], 12 miles north of Ancona, viii.1987, collector C. Froglia, UMML; 10 ♂♂ [2.2-3.1], 9 ovig. [2.5-2.8], off Senigallia, 14 m, 2.ix.1986, collector C. Froglia, IRPEM; 1 ♂ [2.5], 1 ovig. [2.8], off Senigallia, 13 m, 30.ix.1975, collector C. Froglia, CF470; 5 ♂♂ [2.2-2.7], 1 ♀ [2.6], 2 ovig. [2.2, 2.7], between Esino and Misa, 13 m, 29.vii.1976, collector C. Froglia, IRPEM-D497; 2 ♂♂ [2.6], 2 ovig. [2.2, 2.3], 43°42'N, 13°30'E, 21 m, 2.iv.1973, collector C. Froglia, IRPEM-D367; 1 ♂ [2.7], 43°47'N, 13°21'E, 20 m, 24.x.1972, collector C. Froglia, IRPEM-D368; 1 ♂ [2.6], 2 ♀♀ [2.6, 2.7], 2 ovig. [2.4, 2.9], 43°47'N, 13°21'E, 20 m, 24.x.1972, collector C. Froglia, IRPEM-D383; 6 ♂♂ [2.3-3.0], 2 ♀♀ [2.6], 4 ovig. [2.4-2.8], 43°47'N, 13°23'E, 23 m, 6.xi.1973, collector C. Froglia, CF 162; 4 ♂♂ [2.8-3.1], 2 ovig. [2.5, 2.6], between

Fano and Senigallia, 22-25 m, 11.viii.1975, collector C. Froglia, CF 474; 3 ♂♂ [2.6-3.0], from 43°56'N, 13°15'E to 43°52.6'N, 13°17.7'E, 34-36 m, 11.viii.1975, collector C. Froglia, IRPEM-D394; 6 ♂♂ [2.6-2.7], 1 ovig. [2.5], 43°54'N, 13°26'E, 55 m, 7.v.1973, collector C. Froglia, IRPEM-D366; 1 ♂ [2.2], 1 ovig. [2.7], off Rovinj, 32 m, 10.ix.1985, collector Exk. Uni Frankfurt, SMF 13973; 1 ovig. [2.0], off Split, 30-35 m, bottom of sand, 5.iv.1932, NHRM 13953; 1 ♂ [1.6], off Split, 40 m, bottom of bryozoans, RMNH D 16162. Aegean Sea: 1 ovig. [1.5], off Izmir, RMNH D 20709. Off Israel: 1 ♂ [1.7], 4 ovig. [1.4-2.0], off Ashdod, 50 m, RMNH D; 2 ♂♂ [1.3, 1.4], 1 ovig. [1.7], off Palmahim, RMNH D; 13 ♂♂ [1.2-2.1], 2 ♀♀ [1.5, 1.8], 5 ovig. [1.6-1.9], 31°55'N, 34°33'E, 80 m, 24-26.i.1977, TAU-NS-17125 (AR).

Diagnosis.— Shield broader than long. Rostrum slightly exceeded by lateral projections. Lateral projections with short submarginal spine.


Ocular peduncles approximately $\frac{3}{4}$ length of shield, with corneae moderately dilated. Ocular acicles narrowly subtriangular, with short submarginal spine. Interocular plate with 2 chitinous spines usually not reaching proximal half of ocular acicles.

Antennular peduncles exceeding distal margin of corneae by approximately $\frac{2}{3}$ length of ultimate segment. Ultimate segment exceeding length of penultimate segment by approximately $\frac{1}{2}$ own length, with dorsolateral row of short setae and 2 moderately long setae on distal half, ventral margin with row of very short setae; basal segment with lateromedial, laterodistal and dorsodistal spines.

Antennal peduncles exceeding distal margin of corneae by approximately $\frac{1}{4}$ length of ultimate segment. Antennal acicles with distal short spine, not reaching corneae distally.

Right cheliped with moderately narrow chela; in mature males prominent hiatus between dactyl and fixed finger distally. Dactyl approximately $\frac{3}{4}$ length of palm, with granules more numerous on mesial surface and near dorsomesial margin and tufts of moderately long setae. Palm slightly longer than carpus; dorsomesial margin with numerous spinules and moderately long setae and with slightly developed ridge in proximal half; dorsal surface somewhat convex, with scattered spinules and tufts of short setae, and short, low, spinulose ridge near dorsomesial margin proximally; dorsolateral margin with row of spinules extending onto fixed finger and becoming obsolete near claw; dorsoproximal margin with weakly developed mesial and lateral lobes; ventroproximal margin with median, subquadrate sinus. Carpus approximately equal to length of merus; dorsomesial margin with irregular row of numerous short spines; dorsomesial distal angle produced into subrectangular structure with spinulose margins; dorsodistal margin with weak, triangular projection adjacent to dorsomesial distal angle; dorsolateral margin with row of granules or spinules; lateral face unarmed; ventromesial margin subcircular in mesial view; ventral surface granular and with tufts of long setae; very short ventral length between deep, subtriangular, ventrodistal margin and deep, subcircular, ventroproximal mar-

Fig. 14. *Anapagurus bicorniger* A. Milne Edwards & Bouvier, 1892. A-K ♂ (SL = 3.0 mm), off Senigallia, Adriatic Sea, IRPEM: A, shield and cephalic appendages; B, right antennule (lateral view); C, right cheliped (dorsal view); D, carpus of same (ventral view); E, carpus of same (mesial view); F, left cheliped (dorsal view); G, right 2nd pereopod (mesial view); H, ventromesial distal angle of propodus of same; I, right 3rd pereopod (mesial view); J, sternum; K, extremity of abdomen, uropods and telson; L, syntype ♂ (SL = 2.1 mm), MCZ 6577: left sexual tube (lateral view). Scales equal 1.0 mm: a: G, I; b: A, C-F, J, L; c: B; d: H, K.



gin. Ventromesial margin of merus with prominent spinose or spinulose protuberance in males; dorsal margin unarmed distally. Coxae of chelipeds with ventrolateral, distal angle unarmed.

Dactyl of left cheliped granulose or spinulose and with tufts of moderately long setae. Dorsomesial margin of carpus with row of closely spaced spines on distal half, increasing in size distally; dorsolateral margin with 1 strong spine on proximal half.

Dactyls of ambulatory legs exceeding propodi by approximately $\frac{1}{4}$ own length, dorsomesial and ventromesial margins each with closely spaced, moderately long setae. Propodi exceeding carpi $\frac{1}{2}$ to $\frac{2}{3}$ own length; ventromesial distal angle in adult males with 4–8 corneous spines of various sizes (2nd) or unarmed (3rd); in females, both (2nd and 3rd) unarmed. Carpi $\frac{1}{2}$ to $\frac{2}{3}$ length of meri; dorsal margins each with 1 proximal and 1 distal spine; meri with 2, or occasionally 3, widely spaced spinules on dorsal margin and 1 spinule on distal third ventrally. Lobes of sternite of 2nd pereopods unarmed, slightly expanded laterodistally. Anterior lobe of sternite of 3rd pereopods subrectangular.

Males with moderately long, left sexual tube and conical, short, backwardly directed, right sexual tube. Pleopod 3 unequally biramous, pleopods 4 and 5 uniramous. Distal protopodal lobe of right uropod less than $\frac{1}{2}$ length of endopod.

Telson with posterior asymmetrical lobes, separated by wide, median cleft; terminal margins each with 3 strong spines.

Colour.— A. Milne-Edwards & Bouvier (1892) and Bouvier (1940) described the colour as generally whitish.

Related organisms.— A. Milne-Edwards & Bouvier (1900: 222) reported turritellids as the gastropods shells inhabited by some of the syntypes of *A. bicorniger*. In the present collections, specimens of *A. bicorniger* have been found in the following shells: *Turritella biplicata* (1 specimen, MMSU-AAK-1247), and *Amyclina* sp. (6 specimens, RMNH D; and 6 specimens, TAU-NS-17125-AR).

One of the *Amyclina* sp. inhabited by a male *A. bicorniger* (RMNH) was completely covered by colonial hydroids.

One male of *A. bicorniger* (IRPEM-D394) was parasitized by one *Peltogaster paguri*. No external effects of the parasite were observed. This represents the first record of a rhizocephalan in *A. bicorniger*.

Folliculinid protozoans type 1 (see section on related organisms for *A. hyndmani*) were found on 26 specimens of *A. bicorniger* collected at different stations in the Adriatic Sea (MCSN 2384; IRPEM D368, D382, D383, D394, D497, and D589; CF 162, 470 and 474), and on one specimen from the Gulf of Rosas (MNHN-469). The folliculinids usually ranged in number from 1 to 12, occasionally 39 and 81, specimens per host; most tests are cemented dorsally to the posterior carapace, abdomen, and along the groove outlining the last abdominal segment. These are the first records of folliculinids associated with *A. bicorniger*.

Distribution.— Northeastern Atlantic: from off the Portuguese coast of Guia (Neves, 1977) to the Golfo de Cádiz. Mediterranean Sea: from the Golfo de Valencia, Spain, to off the coast of Palmahim, Israel. Larvae of *A. bicorniger* were studied in the Gulf of Naples (Pike & Williamson, 1960) and in the Strait of Messina (Calafiore, 1981). Depth range: 2–200 m.

Affinities.— The interocular-plate spines immediately distinguish *A. bicorniger*




Fig. 15. Distribution of *Anapagurus bicorniger* A. Milne Edwards & Bouvier, 1892. Circled star: type locality, material examined. Solid circles: based on present records. Numbers: based on previous records (circled numbers, material examined). 1-Neves, 1977; 2, 3-García Raso, 1982; 4-Dechancé & Forest, 1962; 5-A. Milne Edwards & Bouvier, 1899; 6-Fenizia, 1937; 7-Pike & Williamson, 1960; 8-Calafiore, 1981; 9-Kocatas, 1981; 10-Holthuis & Gottlieb, 1958.

and *A. petiti* from other congeners. *Anapagurus bicorniger* can be distinguished from *A. petiti* by the much narrower ocular acicles, the shorter interocular-plate spines, the much longer and armed antennal acicles, the more numerous but smaller carpal spines of the right cheliped, and the more numerous and longer ventral setae of the dactyls of the ambulatory legs.

Remarks.—A. Milne Edwards & Bouvier (1892) briefly described *A. bicorniger* from specimens collected in the Golfo de Cádiz. These authors simply listed *A. bicorniger* in their 1898 work and subsequently (1899) reported its occurrence in the Mediterranean for the first time. A. Milne Edwards & Bouvier (1900) provided illustrations and a more detailed specific description based on specimens collected off Monaco. Since its original description, *A. bicorniger* has been easily distinguished from the other congeners by the two interocular-plate, chitinous spines. It was not until 1962 that Dechancé & Forest realized that Balss (1926) and Fenizia (1937) had incorrectly identified specimens as *A. bicorniger* that actually represented their new species *A. petiti*. My reexamination leads me to believe that while Fenizia (1937) did fail to recognize the differences between some of his specimens and *A. bicorniger*, Balss (1926) was correct in his identification. There is an overlap in the number of the propodal ventromesial distal spines on the 2nd pereopods of adult males in *A. bicorniger* (4-8 spines) and *A. petiti* (6-10 spines). Furthermore in both taxa these propodal, ventromesial, distal spines are interspersed with spine-like setae making the counting difficult. The carpal armature of the 2nd pereopods is also variable in both taxa. The present collections have shown that the ventromesial, spinose, meral protuberance of the right cheliped in adult males, which has been used to separate *A. bicorni-*

ger and *A. petiti*, is also present in the latter taxon. Although Dechancé & Forest (1962) and Zariquey Alvarez (1968) correctly included in *A. petiti* those specimens cited by Zariquey Alvarez (1946 and 1955) as *A. bicorniger*, they overlooked the structural overlap in the 2nd pereopods in both species and perhaps were mistaken in citing Balss (1926) as "in part" (pro parte) in their synonymy of A. Milne Edwards & Bouvier's taxon.

Anapagurus longispina A. Milne Edwards & Bouvier, 1900
(figs. 16, 17)

Anapagurus laevis; A. Milne Edwards & Bouvier, 1892: 214 (in part; see remarks); 1894: 72 (in part; see remarks); 1900: 217 (in part; see remarks), not pl. XXVIII: figs. 9, 10; Holthuis & Gottlieb 1958: 72 (in part; see remarks); Türkay, 1976: 36 (in part; see remarks).

Anapagurus laevis var. *longispina* A. Milne Edwards & Bouvier, 1900: 219 (type locality: "Talisman" station 2, off Cabo de São Vicente, 39°53'N, 10°52'W, Golfo de Cádiz, SW Portugal, 99 m, bottom of silt and shells); Bouvier, 1905: 1 (listed); 1922: 31; Gordan, 1956: 303 (listed).

Anapagurus longispina; Forest, 1965: 359, figs. 19, 21, 22 (not citation of figures 12 and 13 in synonymy of A. Milne Edwards & Bouvier 1900's work)(see remarks); Zariquey Alvarez, 1968: 257 (not citation of figures 12 and 13 in synonymy of A. Milne Edwards & Bouvier 1900's work)(see remarks); Kocatas, 1981: 162 (not examined); Ingle, 1985: 764 (listed).

Material.— Holotype: ♂, "Talisman" station 2, off Cabo de São Vicente, 39°53'N, 10°52'W, Golfo de Cádiz, SW Portugal, 99 m, bottom of silt and shells, 6.vi.1883, unknown deposition; not examined. **Northeastern Atlantic:** 1 ovig. [1.9], "Josephine" Expedition station c72, off the Azores, 36°48.5'N, 14°12.2'W, 296.3 m, 1869, NHRM 6999; 1 ♂ [1.4], "Hirondelle" station 226, Strait of Pico-Fayal, Azores, 130 m, 14.viii.1888, MOM 380157; 1 ovig. [1.5], "Meteor" station 9c90, off Portuguese coast, 37°14.7'N, 9°1.5'W, 114-117 m, 22.vi.1967, collector H. Thiel, SMF 21926; 1 ♂ [2.0], "Meteor" station 19-210, Josephine Seamount, 36°41'N, 14°16.18'W, 223-237 m, 11.iii.1970, collector H. Thiel, SMF 5986; 1 ♂ [2.6], "Meteor" station 9c-123a, Josephine Seamount, 36°42.3'N, 14°15.2'W, 210-231 m, 2.vii.1967, collector H. Thiel, SMF 5987; 1 ♂ [2.3], "Meteor" station 9c-130, Josephine Seamount, 36°41.4'N, 14°14.8'W, 216-225 m, 3.vii.1967, collector H. Thiel, SMF 5988; 2 ♂♂ [2.0, 2.1] (USNM 22922) and 1 ♂ [2.0] (MCZ 6580), "Talisman" station 2, off Cabo de São Vicente, 39°53'N, 10°52'W, Golfo de Cádiz, 99 m, bottom of silt and shells, 6.vi.1883; 1 ♂ [1.5], "Princesse Alice" station 2034, Banc Seine, 33°47', 14°21'W, 185 m, 26.vii.1905, MOM 380921. **Mediterranean Sea:** 1 ♂ [1.7], "AAK" station 1192-129, 40°03.0'N, 0°56.1'E, 105 m, bottom of sand and fragmented shells, 2.vi.1974, collector V.V. Murina, MMSU; 1 ovig. [1.9], "AAK" station 1261, 37°26.8'N, 12°11.6'E, 57 m, bottom of sandy clay, collector V.V. Murina, MMSU; 1 ♂ [2.8], 2 ovig. [2.4, 2.5], "AAK" station 1296, 37°30.7'N, 12°06.5'E, 86 m, sandy bottom, 10.ix.1979, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 4 ♂♂ [1.9-2.5], 1 ♀ [2.7], "AAK" station 1305, 37°31.9'N, 11°49.9'E, 70 m, bottom of clayed sand, 14.ix.1979, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 1 ♂ [1.9], "AAK" station 1308, 37°08.9'N, 12°05.2'E, 65 m, bottom of sand and fragments of *Lithothamnium*, 15.ix.1979, collector V.V. Murina, MMSU; 1 ♂ [2.4], "AAK" station 1398, 37°23.6'N, 12°16.5'E, 70 m, sandy bottom, 7.ix.1980, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 1 ovig. [2.0], "AAK" station 1265, 35°27.0'N, 11°39.3'E, 70 m, bottom of clayed sand, 26.iv.78 collector V.V. Murina, MMSU; 6 ♂♂ [1.6-2.4], 3 ♀♀ [1.9-2.4], "AAK" station 1392, 35°7'N, 12°01.2'E, 72 m, bottom of sand and *Lithothamnium*, 3.ix.1980, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 2 ♂♂ [1.6-2.0], AAK 1273-82, 35°56.3'N, 15°00.0'E, 91 m, 30.iv.1978, sandy bottom with fragments of shells and *Lithothamnium*, collector V.V. Murina, MMSU; 4 ♂♂ [1.9-2.4], 1 ♀ [1.8], "AAK" station 1286, 35°49.0'N, 14°53.8'E, 91 m, sandy bottom with fragmented shells, 1.ix.1979, collector V.V. Murina; 5 ♂♂ [1.9-2.2], 3 ♀♀ [1.6-2.1], "AAK" station 1288, 35°38.8'N, 15°01.7'E, 97 m, bottom of clayed sand, 2.ix.1979, collector V.V. Murina, MMSU; 2 ♂♂ [1.6, 1.9], 31°32'N, 32°56'E, 73.2 m, bottom of sand and mud, 1.xi.1975, collector C. Lewinsohn, TAU 15905; 1 ♂ [1.2], sample 201, station 6, Haifa Bay, Israel, 70 m, 16.v.1955, collector E. Gottlieb, RMNH D 13728; 1 ovig. [1.4], sample 191, station 6→

5 (6), Haifa Bay, Israel, 79 m, 5.iv.1955, collector E. Gottlieb, RMNH D 13736; 1 ovig. [1.3], sample 695, off Atlit, Israel, 90 m, 9.ix.1948, collector A. Wirzubski, RMNH D 13744. **Southeastern Atlantic:** 1 ovig. [2.1], off Porto Alexandre, Angola, 109.7 m, 11.viii.1912, collector Skoog, NHMG 14062.

Diagnosis.— Shield slightly broader than long. Rostrum approximately equaling lateral projections. Lateral projections with short submarginal spine.

Ocular peduncles approximately $\frac{3}{4}$ length of shield; with corneae moderately to strongly dilated. Ocular acicles narrowly triangular, with long marginal spine. Interocular plate unarmed.

Antennular peduncles exceeding distal margin of corneae approximately $\frac{2}{3}$ length of ultimate segment. Ultimate segment exceeding length of penultimate segment by approximately $\frac{2}{5}$ own length, with dorsolateral, dorsomesial and ventral margins, each with row of moderately long setae; dorsolateral distal angle with tuft of setae composed on 1 long seta and 2 or 3 moderately long setae decreasing in length laterally; basal segment unarmed but with tuft of long setae.

Antennal peduncles exceeding distal margin of corneae by approximately $\frac{1}{2}$ length of ultimate segment. Antennal acicles terminating in moderately long spine, usually reaching corneae distally.

Right cheliped with narrow chela. Dactyl approximately $\frac{3}{5}$ length of palm, with tufts of moderately long setae and spinules more numerous on mesial surface and near dorsomesial margin. Palm slightly shorter than carpus, with somewhat developed spinulose ridge on dorsomesial margin proximally; dorsal surface moderately convex, spinulose, with scattered, short setae, with low, broad spinulose ridge near dorsomesial margin proximally; dorsolateral margin with row of closely spaced spinules extending and becoming more numerous on fixed finger but becoming obsolete near claw; dorsoproximal margin with weakly developed lateral lobe and almost vestigial mesial lobe; ventroproximal margin with median, skewed subrectangular sinus. Carpus exceeding merus approximately by $\frac{1}{3}$ own length; dorsomesial margin with row of 4 or 5 acute spines interspersed with spinules and moderately long setae; dorsomesial distal angle produced into moderately large, concave subquadrate structure; dorsodistal margin with weak, triangular projection adjacent to dorsomesial distal angle; dorsal surface depressed, spinulose or granular; dorsolateral margin with row of 3 or 4 moderately long spines; dorsolateral distal angle produced into small lobe; lateral face moderately convex, with numerous spinules and tufts of short setae; mesial face slightly compressed with scattered granules and moderately long setae; ventromesial margin subcircular in mesial view; ventral surface spinulose and with tufts of long, stiff setae; moderately short ventral length between deep, subtriangular ventrodistal margin and deep, subcircular ventroproximal margin. Dorsal margin of merus with acute distal spine; ventromesial margin with spinules and acute distal spine; ventrolateral margin with 3 or 4 acute spines; ventral surface spinulose, with tufts of long, stiff setae. Ischium with 1 spinule on ventrolateral margin; ventromesial margin minutely serrate. Coxae of chelipeds with ventrolateral distal angle unarmed.

Dactyl of left cheliped smooth but with tufts of moderately long setae. Dorsomesial margin of carpus with 3 or 4 strong, widely spaced spines; dorsolateral margin with row of 5 or 6 strong, closely spaced spines.

Ambulatory legs with dactyls exceeding propodi by approximately $\frac{1}{5}$ own length, with dorsomesial row of moderately long setae increasing in length distally,

and mediomesial and ventromesial rows of short setae; lateral faces each with row of widely spaced, short setae near dorsal and ventral margins. Propodi approximately twice length of carpi; dorsal surfaces each serrate and with row of short setae; mesial and lateral faces glabrous; ventromesial margin with widely spaced, very short setae; ventromesial distal angle with corneous spines decreasing in size distally, 4 to 6 long (2nd, males), 3 to 5 short (2nd, females), 1 very short (3rd, both sexes). Carpi $\frac{2}{3}$ to $\frac{2}{5}$ length of meri; dorsal surfaces serrate, each with 1 proximal and 1 distal spine and row of short setae. Dorsal margin of meri with 2 spinules, 1 on proximal half and 1 on distal third (2nd); ventral margins serrate, with 1 spinule on distal third (2nd) or unarmed (3rd); both margins with rows of short setae. Lobes of sternite of 2nd pereopods unarmed, distally rounded. Anterior lobe of sternite of 3rd pereopods subrectangular; anterior margin with tufts of long setae.

Males with left sexual tube directed diagonally toward exterior, and then upwardly recurved to level of dorsodistal margin of branchiostegite. Coxa of 5th right pereopod with short sexual tube backwardly directed, partly covered with tufts of moderately long setae. Pleopods 3 and 4 unequally biramous, pleopod 5 uniramous. Distal protopodal lobe of right uropod approximately $\frac{1}{2}$ length of endopod.

Telson with strongly asymmetrical posterior lobes separated by moderately broad, shallow cleft; terminal margins each with 3 or 4 short spines; lateral margins unarmed but with tufts of moderately long setae.


Colour.—Unknown.

Related organisms.—The specimens of *A. longispina* collected off Cabo de São Vicente (A. Milne Edwards & Bouvier, 1900) were found in the gastropod shells of *Turritella*, *Natica*, and *Nassa* (=*Hinia*). In the present collections, specimens of *A. longispina* were found inhabiting the gastropod shells of *Turritella biplicata* (2 specimens, MMSU- "AAK" station 1296), *Gibbula tumida* (1 specimen, MMSU- "AAK" station 1305), and *Cytheromorula* (=*Evokesia*?) sp. (1 specimen, NHRM 6999); and a pectinariid polychaete tube (MMSU- "AAK" 1296).

There is no external evidence of symbionts in any of the specimens examined.

Distribution.—*Anapagurus longispina* has been reported: in the northeastern Atlantic, off the Azores (A. Milne Edwards & Bouvier, 1894; as *A. laevis*), off southwest Portugal (A. Milne Edwards & Bouvier, 1900; as *A. laevis* and *A. laevis* var. *longispina*.—Türkay, 1976; as *A. laevis*), and in Seine Seamount, east off Madeira Island (Bouvier, 1922).—in the Mediterranean Sea, off Azores (Forest, 1965), off Toulon (A. Milne Edwards & Bouvier, 1900; as *A. laevis*), and in the Aegean Sea (Kocatas, 1981). The following localities are new records of *A. longispina*: in the Mediterranean Sea, Golfo of Valencia, Strait of Sicily, and off the coasts of Malta, Sinai and Israel; and in the southeastern Atlantic, off Port Alexander, Angola. Depth range: Forest (1965) recorded the bathymetric range of *A. longispina* as 99–445 m; the present record of 57 m shows that this species also occurs in shallower waters.

Fig. 16. *Anapagurus longispina* A. Milne Edwards & Bouvier, 1900. ♂ (SL = 2.0 mm), "Talisman", 39°53'N, 10°52'W, MCZ 6580: A, shield and cephalic appendages; B, right antennule (lateral view); C, right cheliped (dorsal view); D, carpus of same (ventral view); E, carpus of same (ventromesial view); F, left cheliped (dorsal view); G, left 2nd pereopod (mesial view); H, left 3rd pereopod (mesial view); I, sternum; J, extremity of abdomen, uropods and telson; K, left sexual tube (lateral view). Scales equal 1.0 mm: a: K; b: A, C–I; c: B; d: J.






Fig. 17. Distribution of *Anapagurus longispina* A. Milne Edwards & Bouvier, 1900. Star: type locality. Solid circles: based on present records. Numbers: based on previous records (circled numbers: material examined): 1-A. Milne Edwards & Bouvier, 1894 (as *A. laevis*); 2-Türkay, 1976 (as *A. laevis*); 3-5-A. Milne Edwards & Bouvier, 1900 (as *A. laevis*); 4-Forest, 1965; 6-Kocatas, 1981; 7-Holthuis & Gottlieb, 1958; 8-Bouvier, 1922.

Affinities.—As have other congeners, *A. longispina* has been confounded with *A. laevis*. The relationship between the two species is inferred from the similarity observed between the strongly dilated corneae. *A. longispina* can be readily separated from *A. laevis* by the depressed, dorsal carpal surface of the right cheliped of the former, the much longer and slender ultimate antennular segment, the sternal anterior lobes of the 2nd pereopods not extended distolaterally, and, in males, the much longer ventromesial distal spines of the propodi of the 2nd pereopods. Ocular acicles that terminate in long marginal spine easily distinguish *A. longispina* from other species of *Anapagurus*.

Remarks.—A. Milne Edwards & Bouvier (1900) observed the specimens of *A. laevis* collected by the "Travailleur" and the "Talisman" and noticed a male with long ocular acicles and named it *A. laevis* var. *longispina*. In one of the specimens collected by the "Travailleur" off Cap Sicié, near Toulon (station 9) Forest (1965) found several diagnostic characters that convinced him that *longispina* was a distinct species.

My examination of the collections of earlier authors has shown that certain specimens of *A. longispina* have been misidentified as *A. laevis*: A. Milne Edwards & Bouvier, 1892 and 1894 (one male collected at station 226 by the "Hirondelle", MOM 380157), 1900 (three males collected at station 2 by the "Talisman": (one) MCZ 6580 and (two) USNM 22922); Holthuis & Gottlieb, 1958 (three specimens collected in Haifa Bay and off Atlit, Israel, RMNH D 13728, 13736, and 13744); and Türkay, 1976 (one ovigerous female collected at station 9c90 by the "Meteor," SMF 4799).

Forest (1965) and Zariquey Alvarez (1968) incorrectly cited figures 12 and 13 in the synonymy of A. Milne Edwards & Bouvier 1900's work; the latter authors did not illustrate the new subspecies of *A. laevis*.

Anapagurus bonnieri Nobili, 1905
(figs. 18, 19)

Anapagurus Bonnieri Nobili, 1905: 239; 1906b: 89-91, pl V: figs. 19, 19a, not 19b (type locality: "J. Bonnier & Ch. Pérez Mission" station XLVII: from 25°10'N, 55°10'E to 24°55'N, 54°40'E, Persian Gulf, 12.8-27.4 m, on pearl oyster bank).

Anapagurus bonnieri; Gordan, 1956: 302 (listed); Lewinsohn, 1969: 87, fig. 15; Vine, 1986: 108.
Anapagurus sp; Haig & Ball, 1988: 179, fig. 9.

Material.— Types not examined. **Red Sea:** 4 ♂♂ [1.3-2.0], 1 ovig. [1.7], off Dahlak Archipelago, Ethiopia, 15°35'30"N, 40°40'30", 12.8-14.6 m, 23.x.1965, TAU 1931. **Indonesia:** 1 ♂ [1.4], 2 ovig. [1.2, 1.4], "Alpha Helix" Saparua station 1, off Pulau Saparua, 3°36"S, 128°39.5"E, 5-7 m, on white sand, 29.v.1975, collector E.E. Ball, NIOJ.

Diagnosis.— Shield approximately as long as broad. Rostrum slightly exceeding lateral projections. Lateral projections with short submarginal spine.

Ocular peduncles approximately $\frac{3}{4}$ length of shield, with corneae slightly dilated. Ocular acicles narrowly triangular, with strong submarginal spine. Interocular plate unarmed.

Antennular peduncles overreaching ocular peduncles by approximately $\frac{3}{4}$ length of ultimate segment. Ultimate segment exceeding length of penultimate segment by approximately $\frac{1}{3}$ own length, with 1 dorsolateral, short seta distally, ventral margin with row of tufts of setae, each tuft composed of 1 short seta between 2 long setae.

Antennal peduncles slightly exceeding or equaling length of ocular peduncles. Antennal acicles usually reaching to proximal margin of corneae.

Right cheliped with moderately broad chela. Dactyl $\frac{1}{3}$ to $\frac{3}{4}$ length of palm, with granules, larger near dorsomesial margin and tufts of long setae; dorsal surface slightly elevated in midline. Palm equaling or slightly exceeding length of carpus with dorsomesial margin with well-developed, granular ridge (spinulose ridge in larger specimens) and tufts of moderately long setae; dorsal surface somewhat convex, with granules (more numerous and larger near mesial margin) and tufts of short setae and long, prominent, granular ridge near dorsomesial margin proximally; dorsolateral margin well delimited, with row of granules increasing in size on fixed finger but not extending to claw; dorsoproximal margin with weakly developed, mesial and lateral lobes; all surfaces with scattered tufts of short setae; ventroproximal margin with skewed, median, subrectangular sinus and 2 small, slightly calcified sinuses. Carpus slightly exceeding length of merus, inflated; dorsomesial margin with row of closely spaced, acute spines and tufts of moderately long, stiff setae; dorsomesial distal angle produced into moderately large subtriangular structure, with a more raised ventral side; dorsodistal margin without triangular projection adjacent to dorsomesial distal angle; dorsal surface glabrous; dorsolateral margin with 1 acute spine on distal half, dorsolateral distal angle produced into small, subcircular lobe; mesial face slightly compressed, with tufts of moderately long setae; ventromesial margin subconical in mesial view; ventral surface granular and with tufts of long setae, short ventral length between deep, subtriangular, ventrodistal margin and deep, subcircular, ventroproximal margin; lateral face convex, unarmed. Dorsal margin of merus

with row of tufts of moderately long setae and partially obscured, strong distal spine; ventromesial margin with acute distal spine (small specimens) or tubercle (large specimens), and well-developed, tuberculate protuberance; ventrolateral margin with 3 or 4 strong spines (small specimens) or tubercles (large specimens); ventral surface with numerous tubercles and tufts of long, stiff setae. Ischium with 1 spinule on ventrolateral margin partly covered by short setae; ventromesial margin unarmed but with tufts of long, stiff setae. Coxae of chelipeds with ventrolateral distal angle unarmed.

Dactyl of left cheliped smooth but with tufts of moderately long setae. Dorsomesial and dorsolateral margins of carpus, each with 3 or 4 strong, widely spaced spines.

Ambulatory legs with dactyls exceeding propodi by approximately $\frac{1}{6}$ own length, with dorsomesial row of closely spaced, short, stiff setae; mesial faces each with row of closely spaced, stiff setae near ventral margin; lateral faces each with row of widely spaced setae near unarmed ventral margin. Propodi exceeding carpi $\frac{1}{3}$ to $\frac{1}{2}$ own length; dorsal surfaces each with row of short setae; mesial and lateral faces each with row of short setae near ventral margin; ventromesial distal angle with 1 short spine. Carpi approximately $\frac{2}{3}$ length of meri; dorsal surfaces with row of 5-7 strong spines (2nd) or 2 or 3 strong spines (3rd, small specimens), occasionally with 4 or 5 short spines and 2 distal strong spines (3rd, large specimens), and row of tufts of short stiff setae. Ventral margin of meri with 1 or 2 spinules on distal half (2nd) or unarmed (3rd) and row of tufts of short setae. Lobes of sternite of 2nd pereopods unarmed, distally rounded. Anterior lobe of sternite of 3rd pereopods subrectangular to subcircular; anterior margin with tufts of long setae.

Males with well-developed, left sexual tube, recurved upward to level of coxa; coxa of right 5th pereopod with gonopore on distoventral surface partly covered by tufts of long setae; three uniramous pleopods. Distal protopodal lobe of right uropod subequal to length of endopod.

Telson with asymmetrical posterior lobes separated by narrow deep cleft; terminal margins each with 2 to 4 strong spines; lateral margins unarmed but with tufts of long setae.


Colour.— "Basically transparent except for abundant white chromatophores" (Haig & Ball, 1988: 179; as *Anapagurus* sp.)

Behaviour.— "They [these hermit crabs] can take great leaps backwards (6-8 cm) and move very rapidly over the sand when frightened" (Haig & Ball, 1988: 179; as *Anapagurus* sp.).

Distribution.— In addition to the type locality, *A. bonnieri* has been collected off the Dahlak Archipelago (Ethiopia), Red Sea and off Pulau Saparua, Indonesia. Depth range: 5-15 m.

Affinities.— *Anapagurus bonnieri* shares a number of characters with *A. pusillus*,

Fig. 18. *Anapagurus bonnieri* Nobili, 1905. A-I: ♂ (SL = 2.0 mm), 15°35'30"N, 40°40'30"E, Red Sea, ISRSE 1931: A, shield and cephalic appendages; B, left antennule (lateral view); C, right cheliped (dorsal view); D, carpus of same (ventral view); E, carpus of same (mesial view); F, left cheliped (dorsal view); G, right 2nd pereopod (mesial view); H, right 3rd pereopod (mesial view); I, sternum. J, K: ♂ (SL = 1.4 mm), "ALPHA HELIX" Saparua station 1, 3°36'S, 148°39.5'E, NIOJ: J, uropods and telson; K, left sexual tube (lateral view). Scales equal 1.0 mm: a: K; b: A, C-I; c: B; d: J.






Fig. 19. Distribution of *Anapagurus bonnieri* Nobili, 1905. Star: type locality. Circled numbers: based on previous records, material examined: 1-Lewinsohn 1969; 2-Haig & Ball, 1988 (as *Anapagurus* sp.).

A. breviaculeatus and *A. alboranensis* spec. nov., however the former taxon is easily separated from the other three by its longer rostrum, which exceeds the lateral projections; the right cheliped with the dorsoproximal margin of the palm with weakly developed mesial and lateral lobes, the dorsodistal margin of the carpus lacking the projection adjacent to the dorsomesial distal angle, and the subconical ventromesial margin of the carpus in mesial view.

Remarks.— Lewinsohn (1969) mentioned that Nobili (1906b) erroneously illustrated a second pereopod (pl. V: fig. 19b) showing its dactyl with a long groove, a character that I too found absent in all of the specimens of *A. bonnieri* that I examined.

In their study of hermit crabs from northern Australian and eastern Indonesian waters, Haig & Ball (1988) listed a number of specimens from Pulau Saparua (Indonesia) as *Anapagurus* sp. I have been able to reexamine these and find that they are *A. bonnieri*.

Anapagurus breviaculeatus Fenizia, 1937
(figs. 20, 21)

Anapagurus laevis; Bouvier, 1922: 31 (in part; see remarks); Holthuis & Gottlieb, 1958: 72 (in part; see remarks).

Anapagurus breviaculeatus Fenizia, 1937: 25-28, figs. 17-19 (type locality: off La Pietra, from off Island of Nisida to the Gulf of Pozzuoli, Italy; 5-15 m); Bouvier, 1940: 147, 148, fig. 101 (see remarks); Zar-quietay Alvarez, 1955: 402; 1968: 257 (listed; see remarks); Gordan, 1956: 302 (listed); Balss, 1957: 1698 (listed); Forest, 1958: 100; 1965: 356-359, figs. 13, 16-18; 1967: 7; Pike & Williamson, 1958: 7, 8,

figs. 39-41; 1960: 525-528, fig. 8; Dechancé, 1962: 373, fig. 5; Peres & Picard, 1964: 63, 71; Hazlett, 1968b: 247, 248, fig. 5; Guille, 1971: 102, 111, 113, 119; Stevcic, 1972: 102, 103; Pastore, 1972: 107 (listed); Salmon, 1983: 148; García Raso, 1984: 259; 1985: 14 (listed); Ingle, 1985: 764 (listed); Ros, Romero, Ballesteros & Gili, 1985: 259.

Anapagurus breviaculatus; Forest, 1965: figure legends for figs. 13, 16-18; Kocatas, 1981: 162 (misspelling of *A. breviaculeatus*).

Material.—♂ (carapace length= 4.6 mm), off La Pietra, from off Island of Nisida to the Gulf of Rozzulli, Italy, 5-15 m, 1936?, Museo Zologico della R. Universita di Napoli; not examined. **Northeastern Atlantic:** 1 ovig. [2.2], "Flor da Ponte" station 1.655, off NW Spain, Ría de Arosa, 0.2 km south off Los Mezos Light, 42°30.5'N, 08°55.8'W, 17 m, 24.vii.1964, collector L.B. Holthuis, USNM 121793. **Mediterranean Sea: Balearic Sea:** 1 ♂ [3.6], 3 ovig. [2.2-2.8], Bahía de San Antonio, Ibiza, 45.7 m, 1871, collector A. Lindahl, NHRM 6965; 7 ♂♂ [1.7-3.7], 3 ♀♀ [1.7-2.7], 2 ovig. [2.8, 2.9], Bahía de San Antonio, Ibiza, 45.7 m, 1871, collector F. Söderlund, NHRM 12061; 4 ♂♂ [1.5-3.6], 2 ♀♀ [2.0], 1 ovig. [2.8], off Formentera, 45.7 m, 1871, collector F. Söderlund, NHRM 12056; 2 ♂♂ [2.0, 2.8], 3 ♀♀ [1.5-2.2], off Formentera, 54.9 m, 1871, collector F. Söderlund, NHRM 12057; 1 ♂ [1.5], 1 ♀ [2.8], 1 ovig. [2.9], off Formentera, 64 m, 1871, collector F. Söderlund, NHRM 12058; 2 ♂♂ [2.9, 3.6], 3 ♀♀ [2.3-2.8], off Formentera, 73.2 m, 1871, collector F. Söderlund, NHRM 12059; 2 ♂♂ [2.6, 3.2], off Formentera, 91.4 m, 1871, collector F. Söderlund, NHRM 12060. **Bahía de Cadaqués:** 4 ♂♂ [0.7-1.2], 1 ♀ [0.8], 80-100 m, 7-15.viii.1949, collector L. B. Holthuis, RMNH D 6775; 1 ♀ [1.6], 20-50 m, 11-18.viii.1954, collector L.B. Holthuis, RMNH D 10285; 1 ovig. [1.0], 10-25 m, 31.vii.1957, collector L.B. Holthuis, RMNH D 12280; 3 ♂♂ [0.8-1.2], off Isla Arenella, ~ 45 m, 1.viii.1957, collector L.B. Holthuis, RMNH D 12281; 1 ♂ [1.9], off Playa del Ros, 0-0.5 m, bottom of seagrass, 1.viii.1957, collector L.B. Holthuis, RMNH D 12282; 1 ♂ [1.7], off Playa d'En Pere, 0-1 m, bottom of seagrass, 31.vii.1957, collector L.B. Holthuis, RMNH D 12283; 1 ♂ [2.8], 1 ♀ [1.5], 1 ovig. [1.2], 20-30 m, 14.viii.1959, collector L.B. Holthuis, RMNH D 14877; 1 ♀ [1.9], Cala Jugadora, south of Cabo de Creus, bottom of seagrass, 17.viii.1959, collector L.B. Holthuis, RMNH D 14878; 1 ♂ [1.5], off Playa del Ros, 0-1 m, bottom of seagrass, 13.viii.1959, collector L.B. Holthuis, RMNH D 14879; 1 ♂ [1.0], off Farallons, 20-30 m, 12.viii.1959, collector L.B. Holthuis, RMNH D 14881; 1 ♂ [2.9], 1 ovig. [2.3], off Playa d'En Pere, 0-1 m, bottom of seagrass, 27.vii.1961, collector L.B. Holthuis, RMNH D 16587; 2 ♂♂ [2.2], 1 ♀ [1.8], 30 m, coarse sand, 2.viii.1961, collector L.B. Holthuis, RMNH D 16588; 3 ovig. [1.0-2.1], Port Lligat, ~30 m, sandy bottom, 1.viii.1961, collector L.B. Holthuis, RMNH D 16589; 5 ♂♂ [1.5-2.9], 1 ♀ [1.3], 2 ovig. [2.1, 2.5], 30-40 m, coarse sand, 27.vii-2.viii.1961, collector L.B. Holthuis, RMNH D 16590. **Ligurian Sea:** 1 ♂ [3.2], "Hirondelle" station 1262, off Monaco, 43°43'N, 7°25'20"E, 48 m, 15.v.1902, MOM 380834; 2 ♂♂ [2.6, 2.7], off Sori, Italy, 45.7 m, x.1887, collector H. Théel, NHRM 6951; 2 ♂♂ [2.3, 2.7], off Sori, Italy, 36.6-54.9 m, 3.x.1887, collector H. Théel, NHRM 13178; 1 ovig. [1.9], off Cape Caccia, NW Sardinia, 8 m, bottom of gross sand, 9.v.1982, collector B. Dittrich, RMNH D 350; 1 ♂ [1.0], 1 juvenile [0.6], off Cape Caccia, NW Sardinia, 11.5 m, sandy bottom, 19.viii.1982, collector B. Dittrich, RMNH D 371; 1 juvenile [0.6], off Cape Caccia, NW Sardinia, 25 m, sand bottom, 6.ix.1982, RMNH D 4216. **Tyrrhenian Sea:** 2 ♂♂ [1.9, 2.0], 1 ovig. [2.2], off Naples, MOM 382513; 2 ♂♂ [1.6, 2.5], 2 ovig. [1.3, 1.9], off Villa Rocca Romana, Naples, 15 m, 8.v.1959, collected by Stazione Zoologica di Napoli, RMNH D 14880; 1 ♂ [2.3], Bocca Piccola, Capri, 80-100 m, 17.v.1959, collected by Stazione Zoologica di Napoli, RMNH D 14882; 1 ♂ [2.1], off east coast of Sardinia, Italy, 39°08'N, 08°20.2'E, 10 m, 5.iv.1972, collector G. Orel, CF 387. **Gulf of Taranto:** 1 ♂ [1.9], 1 ovig. [2.0], Porto Cesareo, 40°15'N, 17°54'E, collector Larenzan, MNHN 3166; 1 ♂ [2.3], 1 ovig. [1.9], off Island of Eolie, 11.vi.1974, collectors C. Froglio, G. Gianetti & B. Antolini, IRPEM D392. **Strait of Sicily:** 9 ♂♂ [2.5-3.4], 1 ♀ [2.0], 1 ovig. [2.3], "AAK" station 1261, 37°26.8'N, 12°11.6'E, 57 m, sandy clay bottom, 24.iv.1978, collector V.V. Murina, MMSU; 3 ♂♂ [2.3-3.3], 1 ♀ [2.0], "AAK" station 1262 (55), 37°40.0'N, 12°03.0'E, 21 m, clayed sand bottom, 24.iv.1978, collector V. V. Murina, MMSU; 1 ovig. [2.7], "AAK" station 1296, 37°30.7'N, 12°06.5'E, 86 m, sandy bottom, 10.ix.1979, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 1 ♂ [3.5], 2 ♀♀ [2.7], 1 ovig. [2.6], "AAK" station 1297, 37°28.0'N, 12°12.0'E, 62 m, sandy bottom, 11.ix.1979, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 2 ♂♂ [2.9, 3.2], 1 ovig. [2.2], "AAK" station 1298, 37°26.3'N, 12°10.5'E, 55 m, sandy bottom, 11.ix.1979, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 1 ovig. [1.8], "AAK" station 1299, 37°28.9'N, 12°12.0'E, 62 m, sandy bottom, 11.ix.1979, collector

tors V.V. Murina & Sviridova, MMSU; 2 ♂♂ [2.2, 2.9], 1 ♀ [2.9], "AAK" station 1398, 37°23.6'N, 12°16.5'E, 70 m, sandy bottom, 7.ix.1979, collectors V.V. Murina & V.D. Tchukhtchin, MMSU. Off East of Tunisia: 2 ♂♂ [2.0, 2.4], "AAK" station 1208 (159), 35°23.2'N, 11°56.4'E, 62 m, sandy bottom, 9.vi.1974, collector V.V. Murina, MMSU; 2 ♂♂ [2.0, 2.6], "AAK" station 1265, 35°27.0'N, 11°39.3'E, 70 m, clayed sand bottom, 26.iv.1978, collector V.V. Murina, MMSU; 2 ♂♂ [1.7, 2.3], 1 ♀ [2.4], 1 ovig. [1.9], "AAK" station 1267, 35°10.0'N, 12°14.4'E, 50 m, sandy bottom covered with algae, 27.iv.1978, collector V. V. Murina, MMSU; 4 ♂♂ [2.1-2.6], 3 ♀♀ [1.6-2.1], "AAK" station 1268, 35°10.8'N, 12°34.0'E, 47-50 m, sandy bottom, 27.iv.1978, collector V.V. Murina, MMSU; 4 ♂♂ [1.9-2.4], 1 ♀ [2.1], 1 ovig. [2.4], "AAK" station 1269 (70), 35°13.0'N, 12°38.5'E, 57 m, sandy bottom, 28.iv.1978, collector V.V. Murina, MMSU; 1 ovig. [2.0], "AAK" station 1271, 35°46.2'N, 12°48.5'E, 70 m, bottom of organogenic sand, 29.iv.1978, collector V.V. Murina, MMSU; 4 ♂♂ [1.9-2.4], 4 ovig. [2.0-2.3], "AAK" station 1313, 35°17.2'N, 12°04.5'E, 53 m, sandy bottom covered with *Udotea*, 17.ix.1979, collector V.V. Murina, MMSU; 1 ♂ [2.6], "AAK" station 1314, 35°31.4'N, 12°02.6'E, 76 m, sandy bottom, 18.ix.1979, collector V.V. Murina, MMSU; 1 ♂ [2.2], 1 ovig. [2.6], 72 m, sandy bottom with fragments of *Lithothamnium*, 3.ix.1980, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 2 ♂♂ [1.9, 2.1], "AAK" station 1268, 35°10.8'N, 12°34.0'E, 47-50 m, sandy bottom, 27.iv.1978, collector V.V. Murina, MMSU. Off east of Malta: 2 ♂♂ [2.2, 2.9], 1 ovig. [1.6], "AAK" station 1273 (82), 35°56.3'N, 15°00.0'E, 91 m, sandy bottom with fragments of shells and *Lithothamnium*, 30.iv.1978, collector V.V. Murina, MMSU. Adriatic Sea: 3 ♂♂ [2.6-3.1], 2 ovig. [2.7, 3.0], Gulf of Manfredonia, Italy, 41°29.8'N, 15°59'E, 11 m, 7.vii.1974, collectors C. Froglia & G. Bombace, IRPEM D376; 2 ovig. [2.3, 2.6], 41°30.2'N, 15°58.7'E, 9 m, 7.vii.1974, collectors C. Froglia & P. Zucchiri, IRPEM D377; 1 ♂ [2.5], 1 ovig. [1.9], 41°26.6'N, 16°05.4'E, 12 m, 5.vii.1974, collectors C. Froglia & G. Bombace, IRPEM D381; 2 ♂♂ [1.8, 2.5], 1 ovig. [1.2], off Rovinj, Yugoslavia, 5.vi.1962, RMNH D 19366; 3 ♂♂ [1.0-1.8], 1 ovig. [1.2], "F.B. Burin", off Rovinj, Yugoslavia, 28 m, 5.ix.1985, SMF 13908; 1 ♀ [1.3], "F.B. Burin", off Rovinj, Yugoslavia, 32 m, 10.ix.1985, collector Exk. Uni Frankfurt, SMF 13972; 1 ovig. [1.9], "F.B. Burin", off Rovinj, Yugoslavia, 36 m, 15.ix.1987, SMF 21923; 1 ♂ [2.3], "F.B. Burin", off Rovinj, Yugoslavia, 28 m, 17.x.1987, SMF 21924; 1 ♂ [1.8], "Sixten Bock", off Split, Yugoslavia, 30-35 m, bottom of sand and shells, iii.1932, NHRM 13571; 1 ♂ [1.6], "Sixten Bock", off Split, Yugoslavia, 30-35 m, bottom of sand and shells, 6.iv.1932, NHRM 13572. Aegean Sea: 1 ovig. [2.2], 39°04'N, 24°07'E, 20-25 m, 9.viii.1978, collector M. Türkay, SMF 12608; 1 ♂ [2.4], 39°19'N, 24°3'E, 6-8 m, 6.vii.1978, collector M. Türkay, SMF 12609; 1 ovig. [2.0], 39°21.6'N, 24°09.6'E, 20 m, 6.viii.1978, collector M. Türkay, SMF 12610; 1 ♂ [1.7], off Izmir, Turkey, 27.vi.1969, collector A. Kocatas, RMNH D 20709. Off Israel: 1 ♂ [3.0], sample 218, station 14, Haifa Bay, 18 m, 17.v.1955, collector E. Gottlieb, RMNH D 13724; 2 ♂♂ [1.0, 1.2], 2 ♀♀ [0.9, 1.0], sample 250, station 7→6 (7), Haifa Bay, 43, 2.viii.1955, collector E. Gottlieb, RMNH D 13725; 1 ♂ [1.0], sample 137, station 9, Haifa Bay, 42-44 m, 8.vii.1954, collector E. Gottlieb, RMNH D 13726; 1 ♂ [1.9], sample 213, station 4→3 (4), Haifa Bay, 38 m, 17.v.1955, collector E. Gottlieb, RMNH D 13727; 2 ♂♂ [0.8, 0.9], 1 ♀ [1.3], 1 ovig. [1.2], sample 374, station 5→6 (5), Haifa Bay, 55-60 m, 24.iv.1956, collector E. Gottlieb, RMNH D 13729; 1 ♂ [0.9], sample 171, station 14, Haifa Bay, 18 m, 30.xi.1954, collector E. Gottlieb, RMNH D 13730; 1 ♂ [1.0], sample 286, station 6→10 (6), Haifa Bay, 73 m, 21.ix.1955, collector E. Gottlieb, RMNH D 13731; 1 ♂ [0.9], 1 ♀ [0.8], 1 ovig. [1.1], sample 233, station 4→5 (5), Haifa Bay, 49 m, 7.vi.1955, collector E. Gottlieb, RMNH D 13732; 1 ♂ [1.4], sample 219, station 14, Haifa Bay, 18 m, 17.v.1955, collector E. Gottlieb, RMNH D 13733; 1 ovig. [1.3], sample 102, station 3, Haifa Bay, 31 m, 2.vi.1954, collector E. Gottlieb, RMNH D 13734; 1 ♂ [1.2], sample 112, station 7, Haifa Bay, 37-38 m, 3.vi.1954, collector E. Gottlieb, RMNH D 13735; 1 ♂ [1.1], sample 191 station 6→5 (6), Haifa Bay, 79 m, 5.v.1955, collector E. Gottlieb, RMNH D 13736; 1 ♂ [1.2], 1 ovig. [1.2], sample 260, station 5→4 (4), Haifa Bay, 35 m, 18.viii.1955, collector E. Gottlieb, RMNH D 13737; 1 ♂ [1.8], 1 ovig. [1.5], sample 405, station 12→9 (12), Haifa Bay, 37-68 m, 13.v.1956, collector E. Gottlieb, RMNH D 13738; 1 ♂ [1.5], sample 257, station 9→8 (9), Haifa Bay, 42 m, 2.viii.1955, collector E. Gottlieb, RMNH D 13739; 1 ♂ [1.6], sample 123, station 10→6 (10), Haifa Bay, 73 m, 6.ix.1954, collector E. Gottlieb, RMNH D 13740; 1 ♂ [1.0], sample 52, station 2, Haifa Bay, 18 m, 23.viii.1953, collector E. Gottlieb, RMNH D 13741; 1 ♂ [1.0], sample 694, off Atlit, 54 m, 9.ix.1948, collector A. Wirszubski, RMNH D 13743; 1 ♂ [1.2], sample 682, off Caesarea, 72 m, 2.viii.1948, collector A. Wirszubski, RMNH D 13745; 2 ♂♂ [1.2], sample 691, off Tel Aviv, 36 m, 4.viii.1948, collector A. Wirszubski, RMNH D 13746.

Diagnosis.—Shield slightly longer than broad. Rostrum slightly exceeded by lateral projections. Lateral projections with short submarginal spine.

Ocular peduncles approximately $\frac{3}{4}$ length of shield; with corneae slightly dilated. Ocular acicles subtriangular, terminating in short submarginal spine. Interocular plate unarmed.

Antennular peduncles exceeding distal margin of corneae, approximately by $\frac{3}{5}$ length of ultimate segment. Ultimate segment exceeding length of penultimate segment by approximately $\frac{1}{3}$ own length, with dorsolateral row of very short setae and 1 distal, moderately long seta; ventral margin with row of very short setae; basal segment with lateromedial and laterodistal (occasionally bifid) spines.

Antennal peduncles slightly exceeding distal margin of corneae. Antennal acicles reaching corneae proximally, with small terminal spine.

Right cheliped with moderately narrow chela. Dactyl approximately $\frac{3}{4}$ length of palm, ventrally curved, strongly so in adult males; subcircular in cross section, with no ridges. Palm subequal to length of carpus, dorsomesial margin with granular ridge on proximal half; dorsal surface slightly convex, with scattered granules and short setae and with spinose ridge near dorsomesial margin proximally, prominent in females but almost obsolete in males; dorsolateral margin with granules (males) or spinules (females) becoming obsolete near claw; dorsoproximal margin with weakly developed mesial and lateral lobes; ventroproximal margin with skewed, median, subquadrate sinus. Carpus exceeding length of merus by approximately $\frac{1}{4}$ own length; dorsomesial margin with irregular row of numerous short spines (males) or fewer but stronger spines (females); dorsomesial distal angle produced into concave, subrectangular lobe; dorsodistal margin with a projection adjacent to dorsomesial distal angle, triangular (females and immature males) or roundish (adult males) dorsolateral margin with row of 4-10 widely spaced spinules (females and immature males) or unarmed (adult males); ventral surface inflated, with tufts of stiff setae, long in males or short in females; short ventral length between moderately deep, subcircular ventrodistal and ventroproximal margins; ventromesial margin subcircular in mesial view. Ventromesial margin of merus with prominent, spinose protuberance (males) or tubercles (females), and with 1 short distal spine; ventrolateral margin with 3 to 5 strong spines; dorsal margin with short distal spine (small specimens) or unarmed (large specimens). Ischium with 1 spinule on ventrolateral margin partly covered by tufts of long, stiff setae. Coxae of chelipeds with ventrolateral distal angle unarmed.

Dactyl of left cheliped smooth but with tufts of moderately long setae. Dorsomesial margin of carpus with row of 5 to 7 strong, widely spaced spines; dorsolateral margin with row of 3 or 4 strong, widely spaced spines.

Dactyls of ambulatory legs exceeding length of propodi by approximately $\frac{1}{4}$ own length, with dorsomesial row of closely spaced, moderately long setae and ventromesial row of closely spaced, short, stiff setae. Propodi exceeding length of carpi by approximately $\frac{1}{2}$ own length; ventromesial distal angle with corneous spines, 2-4 moderately long (2nd pereopods, males), 1 or 2 short (2nd pereopods, females), or unarmed but with stiff setae (3rd pereopods, both sexes). Carpi $\frac{2}{3}$ length of meri; dorsal margins with 4-9 spinules (2nd) or serrate and 1 distal spine (3rd). Meri unarmed or occasionally with 1 spinule on distal third, dorsally and/or ventrally (2nd). Lobes of sternite of 2nd pereopods unarmed, distally rounded. Anterior lobe

of sternite of 3rd pereopods subrectangular, narrower in females.

Males with moderately long and slender, left sexual tube, strongly recurved distally, and conical, short, backwardly directed, right sexual tube; pleopods 3 and 4 unequally biramous, pleopod 5 uniramous. Distal protopodal lobe of right uropod less than $\frac{1}{2}$ length of endopod.

Telson with posterior, asymmetrical lobes, separated by narrow cleft; terminal margins each with 4 short spines.

Colour.— "Il colore in generale e verde sul céfalotorace, specialmente nelle regioni branchiali; l'addome ha un bel colore verde scuro. Le zampe sono di colore ceruleo, percorse da fasce trasverse di colore verde pallido e da fasce longitudinali colour terra di Siena" (Fenizia, 1937: 27). (= Cephalothorax green especially in branchial regions; abdomen a beautiful dark green. Ambulatory legs sky blue with pale-green, transverse bands and ochre-brown, longitudinal stripes.)

Behaviour.— The sexual behaviour of *A. breviaculeatus* has been studied by Hazlett (1968) and Salmon (1983).


Related organisms.— Fenizia (1937: 27) cited *Nassa mutabilis*, *Turritella*, and *Nassa incrassata* as the gastropod shells occupied by the syntypes of *A. breviaculeatus*. In the present collections the following shells were found inhabited by *A. breviaculeatus*: *Turritella biplicata* (3 specimens, NHRM 12056, 12057 and 12061), naticid (1 specimen, NHRM 12061), *Lunatia guillemini* (1 specimen, NHRM 12057), *Raphitoma purpurea* (Montagu) (1 specimen, RMNH D 1370), *Natica catena* (1 specimen, RMNH D 14877), *Pusia tricolour* (Gmelin) (1 specimen, RMNH D 14879), and 2 unidentified shells (NHRM 12061).

Each of two females of *A. breviaculeatus* (RMNH D 10285 and MMSU "AAK" 1297), each had 6 externae of *Peltogasterella sulcata* (Lilljeborg, 1859) on the abdomen. While there were not apparent external effects of the parasites in one of the specimens, the parasitic infestation changed the normal feminine biramous 2nd pleopods to uniramous in the other. One *Peltogaster paguri* Rathke, 1842, was found attached below pleopod 3 of a male *A. breviaculeatus* (RMNH D 14877), but no external modifications were observed. These comprise the first records of rhizocephalan infestation of *A. breviaculeatus*.

One female of *A. breviaculeatus* (RMNH D 16588) was parasitized by a bopyrid isopod, *Athelges tenuicaudis* Sars, 1899, which did not change any of the presumed secondary sexual characters of the host. It represents the first record of a bopyrid isopod in *A. breviaculeatus*.

One or two folliculinid protozoans type 1 (see section on related organisms for *A. hyndmanni*) per host were attached dorsally to the abdomen (usually near 4th and 5th pleopods) of 9 specimens of *A. breviaculeatus* (MMSU "AAK" stations 1261, 1269-70, 1392 and 1398; RMNH D 14880 and 16588; IRPEM D381; and SMF 13908). These comprise the first records of folliculinids in *A. breviaculeatus*.

Fig. 20. *Anapagurus breviaculeatus* Fenizia, 1937. ♂ (SL = 2.9 mm), 41°29.8'N, 15°59'E, Gulf of Manfredonia, Adriatic Sea, IRPEM-D376: A, shield and cephalic appendages; B, right antennule (lateral view); C, right cheliped (dorsal view); D, carpus of same (ventromesial view); E, left cheliped (dorsal view); F, left 2nd pereopod (mesial view); G, left 3rd pereopod (mesial view); H, sternum; I, extremity of abdomen, uropods and telson; J, left sexual tube (lateral view). Scales equal 1.0 mm: a: C-G; b: A, H, J; c: B, I.






Fig. 21. Distribution of *Anapagurus breviaculeatus* Fenizia, 1937. Star: type locality. Solid circles: based on present records. Numbers: based on previous records (circled number: material examined): 1-García Raso, 1984, 1985; 2-Forest, 1965; 3-Zariquey Alvarez, 1955; 4-Dechancé, 1962; Guille, 1971; 5-Pike & Williamson, 1958, 1960; Hazlett, 1968; 6-Forest, 1967; 7-Stevicic, 1972; 8-Kocatas, 1981; 9-Holthuis & Gottlieb, 1958.

Distribution.—*Anapagurus breviaculeatus* has been collected in the Mediterranean Sea, from the Alboran Basin (García Raso, 1984) to off the coast of Israel. This latter locality is a new record for *A. breviaculeatus*. The ovigerous female in Ría de Arosa, 0.2 km south off Los Mezoz Light, Spain, 42°30.5'N, 08°55.8'W (USNM 121793) is the first record of this species outside of the Mediterranean Sea. Depth range: 0-100 m.

Affinities.—As previously indicated, *A. breviaculeatus* appears to be most closely related to *A. pusillus* and *A. alboranensis* spec. nov., but is easily separated by its much longer ocular peduncles and the much longer ultimate antennular segment.

Remarks.—In examining the specimens that Bouvier (1922) reported as *A. laevis*, I found that he confounded two distinct taxa. One male from "Princesse Alice II" station 1262 proved to be *A. breviaculeatus* rather than *A. laevis*. Bouvier (1940), in his comparison of *A. breviaculeatus* and *A. laevis*, incorrectly attributed shorter ocular peduncles to *A. breviaculeatus* and added various spines to his figure of this species, stated as having been redrawn from Fenizia's (1937: fig. 17) original illustration. Years later, Zariquey Alvarez (1968) quoted Bouvier's erroneous statement about the ocular peduncles of *A. breviaculeatus*. As mentioned under *A. laevis*, the specimens reported from the coast of Israel by Holthuis & Gottlieb (1958) as *A. laevis* actually were a composite of *A. breviaculeatus*, *A. longispina*, and *A. adriaticus* spec. nov.

Anapagurus hendersoni Barnard, 1947
(figs. 22, 23)

Anapagurus pusillus Henderson, 1888: 74 (in part; see remarks); Alcock, 1905: 187 (in part; listed); Stebbing, 1910: 358; Balss, 1912: 110, 113, 115; Monod, 1932: 487 (in part; listed); Gordan, 1956: 303 (in

part; listed) (not *Anapagurus pusillus* Henderson, 1888).

Anapagurus hendersoni Barnard, 1947: 377 (in part; see remarks) (Type locality herein assigned by present lectotype designation: Simon's Bay, False Bay, Cape Province, South Africa, "Challenger", 33 m); 1950: 465 fig. 85: f-g; Gordan, 1956: 302 (listed); Day, Field & Penrith, 1970: 56 (Saldanha Bay, Cape Province, South Africa, 9-82 m); Kensley, 1974: 64 (in part; see remarks); 1978: 252 (in part; see remarks); 1981: 31 (listed).

Anapagurus laevis; Forest, 1955: 131 (in part: only specimen collected by "Mercator" 1 mile NE of Island of Dassen), fig. 32, pl. VI: figs. 7, 8].

Material.— Lectotype, herein designated: ♂ [2.2], "Challenger": Simon's Bay, False Bay, Cape Province, South Africa, 33 m, xii. 1873, BMNH 1888: 33; Paralectotypes: 1 ♂ [2.4], "Pieter Faure": off Bird Island Passage, Cape Province, 18 m, 1897, SAM A 1500; 2 ♂♂ [2.0, 2.4], 1 ♀ [2.9], 1 ovig. [2.8], "Pieter Faure": off Cape Natal, W by N, 10 km, 88 m, 1897, SAM A 1501; 7 ♂♂ [1.9-3.5], 2 ovig. [2.3, 3.0], "Pieter Faure": off Umhloti River mouth, Natal, NW $\frac{1}{2}$ W, 24 km, 183 m, 1897, SAM A 1502; 1 ovig. [2.2], "Pieter Faure": off Cape Natal, W by N, 10 km, 99 m, 1897, SAM A 1503. 1 ♂ [2.2], 1 ovig. [2.8], "Valdivia" station 100: Francis Bay, 34°8'S, 24°59'E, 29.x.1898, ZMB 16476; 1 ♂ [3.3], 1 ovig. [2.5], "Th. Mortensen Java-South Africa Expedition" station 24, off Durban, 29°48'30"S, 31°18'E, 226 m, bottom sandy mud, 22.viii.1929, ZMUC; 3 ♂♂ [2.7-3.4], 1 ovig. [2.4], "Th. Mortensen Java-South Africa Expedition" station 31, off Durban, 29°54'25"S, 31°09'45"E, 128 m, bottom of sand and gravel, 29.viii.1929, ZMUC; 1 ♂ [2.4], 1 ovig. [2.1], "Th. Mortensen Java-South Africa Expedition" station 60, 2 $\frac{1}{4}$ miles off Cape Point, False Bay, 57 m, sandy bottom, 19.xii.1929, ZMUC; 2 ♀♀ [1.1, 1.7], "Th. Mortensen Java-South Africa Expedition" station 72, Chapmans Bay, 53 m, sandy bottom, 20.xii.1929, ZMUC; 1 ovig. [2.3], "Mercator" (11th cruise), 1 mile N.E. of Island of Dassen, 33-37 m, 15.i.1937, MNHN 511; 1 ♀ [3.1], "University of Cape Town Ecological Survey" station F8 653 B, 34°07.5'S, 18°31'E, 27.5 m, 22.ii.1947, collector H. Boschma, RMNH D 34771; 10 ♂♂ [2.4-4.4], 2 ♀♀ [2.7, 3.3], 2 ovig. [2.0, 3.7], UCT Ecol. Survey, False Bay, 34°08'S, 18°31.5'E, 27-28 m, 22.ii.1947, SAM 758A.

Diagnosis.— Shield usually as long as broad, with tufts of moderately long setae. Rostrum equaling in length lateral projections. Lateral projections with short submarginal spine.

Ocular peduncles approximately $\frac{1}{2}$ length of shield, with corneae moderately dilated. Ocular acicles roundly triangular, slightly expanded mesially, with strong acute submarginal spine. Interocular plate unarmed.

Antennular peduncles exceeding distal margin of corneae approximately $\frac{3}{4}$ length of ultimate segment. Ultimate segment exceeding length of penultimate segment by $\frac{1}{4}$ to $\frac{1}{3}$ own length, with dorsolateral row of short setae and laterodistal fringe of setae composed of 1 long and few short setae; ventral margin with row of tufts of long setae.

Antennal peduncles exceeding distal margin of corneae by approximately $\frac{1}{2}$ length of ultimate segment. Antennal acicles not reaching corneae distally.


Right cheliped with narrow chela. Dactyl $\frac{2}{3}$ to $\frac{3}{4}$ length of palm, with irregular rows of spinules on dorsomesial margin and tufts of long setae. Palm approximately $\frac{2}{3}$ length of carpus, dorsomesial margin with irregular rows of spinules; dorsal surface somewhat convex with scattered spinules and with short, high spinose ridge near dorsomesial margin proximally; dorsolateral margin with row of closely spaced, acute, erect spines increasing in size toward fixed finger, progressively decreasing on fixed finger and becoming obsolete near claw; dorsoproximal margin with well-developed, mesial, spinulose lobe and weakly developed lateral lobe; mesial face with well-developed ridge on proximal half; all surfaces with numerous tufts of moderately long, occasionally short setae; ventroproximal margin with

slightly developed median, subrectangular sinus flanked by 2 irregular, slightly calcified sinuses, left sinus larger than right. Carpus exceeding merus by $\frac{1}{3}$ own length; dorsomesial margin with closely spaced, acute, erect, occasionally bifid, spines of diverse sizes; dorsomesial distal angle produced into a subtriangular structure, with a concave surface and a more raised ventral side; dorsodistal margin with strong, triangular projection adjacent to dorsomesial distal angle, projection set at approximately 45° angle to longitudinal axis of carpus; dorsal surface slightly depressed, with numerous spinules and numerous tufts of short setae; dorsolateral margin with irregular rows of acute spinules; dorsolateral distal angle produced into small, subcircular lobe; mesial face compressed with spinules more numerous ventrally and numerous tufts of long setae; ventromesial margin subcircular in mesial view; ventral surface with numerous spinules and tufts of long setae; long ventral length between deep, subtriangular, ventrodistal margin and shallow subcircular ventroproximal margin; lateral face convex with spinules and numerous tufts of short setae. Merus with dorsal margin with row of moderately long setae terminating in partially obscured, strong spine; ventromesial margin with low, spinulose protuberance on proximal half and row of spinules terminating in 1 or 2 strong spines on distal; ventrolateral margin with row of 3 to 6 strong spines; ventral surface with numerous spinules and tufts of long setae. Ischium with 1 spinule on ventrolateral margin partly covered by short setae; ventromesial margin with row of spinules and tufts of long setae. Coxae of chelipeds with ventrolateral distal angle unarmed.

Dactyl of left cheliped smooth but with tufts of moderately long stiff setae. Dorsomesial and dorsolateral margins of carpus, each with row of 8 to 12 strong, closely spaced spines.

Ambulatory legs with dactyls exceeding propodi by approximately $\frac{1}{4}$ own length, with dorsomesial row of closely spaced, short, stiff setae; mesial faces each with 2 rows of tufts of long, stiff setae, one near dorsal margin and other close to ventral margin; lateral faces each with median row of short stiff setae, and row of tufts of long setae near ventral margin; ventral margins unarmed but each with 1 distal stiff seta, nearly equaling length of claw. Propodi exceeding carpi by $\frac{1}{3}$ to $\frac{1}{2}$ own length; dorsal surfaces each with numerous tufts of short, stiff setae; mesial faces each with median row of tufts of short setae and row of numerous tufts of long setae near ventral margin; ventromesial distal angle with 1 or 2 short, corneous spines partly obscured by moderately long, stiff setae. Carpi $\frac{1}{2}$ to $\frac{3}{4}$ length of meri; dorsal surfaces each with row of 2-7 short spines (2nd) or 1 short distal spine (3rd) and row of tufts of short, stiff setae. Ventral margins of merus with row of spinules (2nd) or unarmed (3rd), and tufts of long setae. Lobes of sternite of 2nd pereopods unarmed, expanded laterodistally. Anterior lobe of sternite of 3rd pereopods subrectangular to rectangular; anterior margin with numerous tufts of long setae.

Fig. 22. *Anapagurus hendersoni* Barnard, 1947. ♂ (SL = 3.4 mm), 34°08'S, 18°31.5'E, False Bay, South Africa, SAM-758A: A, shield and cephalic appendages; B, left antennule (lateral view); C, right cheliped (dorsal view); D, carpus of same (ventral view); E, carpus of same (ventromesial view); F, left cheliped (dorsal view); G, right 2nd pereopod (lateral view); H, left 3rd pereopod (mesial view); I, sternum; J, extremity of abdomen, uropods and telson; K, telson. L: ♂ (SL = 3.3 mm), Th. Mortensen Java-South Africa Expedition, station 24, 29°48'30"S, 31°18'E, off Durban, South Africa, ZMUC: left sexual tube (lateral view). Scales equal 1.0 mm: a: F-H; b: E; c: A, C, D, I, L; d: J; e: B; f: K.






Fig. 23. Distribution of *Anapagurus hendersoni* Barnard, 1947. Circled star: type locality, material examined. Solid circles: based on present records. Numbers: based on previous records (circled numbers: material examined): 1, 8, 9-Kensley, 1974; 2-Day *et al.*, 1970; 3-Forest, 1955; 4, 5, 7, 8, 9-Barnard, 1950; 5, 8-Kensley, 1978; 6-Balss, 1912.

Males with well-developed, left sexual tube, recurved upward to exterior and short, backwardly directed, right sexual tube; pleopods 3 and 4 unequally biramous, pleopod 5 uniramous. Distal protopodal lobe of right uropod subequaling length of endopod.

Telson with slightly asymmetrical posterior lobes, separated by wide, deep cleft; terminal margins each with 3 or 4 strong spines; lateral margins with 1-4 spinules.

Colour.— “Carapace anteriorly pale pinkish, 4th joints [meri] of chelipeds and legs with pale brownish transverse markings, 6th joints [propodi] of legs violaceous, eyes black, ova scarlet” (Barnard, 1950: 466).

Related organisms.— The lectotype was found inhabiting the shell of the gastropod *Trochus benzi* Krauss (Henderson, 1888: 74).

A female of *A. hendersoni* (RMNH D 34771) is the host for the holotype of the rhizocephalan, *Peltogaster contortus* Boschma; no external modification caused by this parasite was observed.

Distribution.—*A. hendersoni* apparently is an endemic South African species; it has been collected from Saldanha Bay, Cape Province on the west coast to off Cape Natal, Natal Province on the east coast. Depth range: 9–226 m.

Affinities.—The dense pilosity of *A. hendersoni* is very similar to that of *A. chiroacanthus*, but the former taxon can be easily distinguished from the latter and other congeners by the stronger armature of its chelipeds and telson, and by the almost parallel lateral margins of the sternal lobes of the 2nd pereopods.

Remarks.—Among the specimens used by Henderson (1888) in his description of *A. pusillus*, he included "with some hesitation" a male collected in Simon's Bay, South Africa. Furthermore, plate VII, figure 7 of the same paper is herein questionably referred to *A. laevis* (Bell, 1846). Alcock (1905) and Monod (1939) subsequently simply listed *A. pusillus* without reference to the questionable specimen. Stebbing (1910) in his catalogue of South African Crustacea clearly referred the Simon's Bay specimen to *A. pusillus*.

Barnard in 1947, briefly described *A. hendersoni* based on Henderson's Simon's Bay specimen and additional South African material collected from Table Bay to Natal. In 1950 the same author more fully described and illustrated the species. Examination of the syntypes of *A. hendersoni* has shown that Barnard confounded two distinct species as *A. hendersoni*; the lack of the sexual tube in one male specimen (SAM A 8225) originally included in the Barnard's type series shows that it is neither conspecific nor congeneric with *A. hendersoni*. In his list of type specimens of decapod crustaceans in the collections of the South African Museum, Kensley (1974) simply listed all of Barnard's (1947) material. Kensley (1978) in his records of *A. hendersoni* collected at station 31 by the Th. Mortensen Java-South Africa Expedition, included an ovigerous female that represents an unidentified species of *Pagurus*.

Having examined the specimen from the "Mercator" expedition collected 1 mile NE of Dassen Island, I found that its assignment to *A. laevis* by Forest (1955) was incorrect. This specimen represents *A. hendersoni*.

Anapagurus wolffi Forest, 1961
(figs. 24, 25)

Anapagurus wolffi Forest, 1961: 241–246, figs. 7, 11, 15, 19 (type locality: "Atlantide" station 60, 5°06'N, 9°34'W, 78 m, bottom of sand and silt).

Anapagurus laevis; Forest, 1961: 239 (in part; see remarks).

Anapagurus curvidactylus; Forest, 1961: 241 (in part; see remarks).

Material.—Holotype: ♂ [1.0], "Atlantide" station 60, 5°06'N, 9°34'W, off Liberia, 78 m, bottom of sand and silt, 9.i.1946, ZMUC; Paratypes: 1 ovig. [0.9], data as for holotype; 1 ♀ [0.9], "Atlantide" station 49, 7°29'N, 13°38'W, off Sierra Leone, 74–79 m, bottom of muddy sand, 30.xii.1945, ZMUC. 1 ovig. [0.9], "Atlantide" station 49, 7°29'N, 13°38'W, off Sierra Leone, 74–79 m, bottom of muddy sand, 30.xii.1945, ZMUC; 1 ♂ [0.8], "Atlantide" station 51, 7°14'N, 12°57'W, off Sierra Leone, 108 m, sandy bottom, 31.xii.1945, ZMUC; 1 ovig. [1.0], "Atlantide" station 58, 5°50'N, 10°31'W, 95 m, bottom of muddy sand, 8.i.1946, ZMUC.

Diagnosis.— Shield broader than long. Rostrum slightly exceeding unarmed lateral projections.

Ocular peduncles subequal to length of shield; with corneae moderately dilated. Ocular acicles equilaterally triangular, with short submarginal spine. Interocular plate unarmed.

Antennular peduncles exceeding distal margin of corneae approximately $\frac{3}{4}$ length of ultimate segment. Ultimate antennular segment exceeding length of penultimate segment by approximately $\frac{1}{2}$ own length; dorsolateral margin of ultimate segment with 1 short seta on distal half and 1 moderately long seta distally; basal segment with dorsomesial distal spine.


Antennal peduncles reaching proximal margin of corneae. Antennal acicles unarmed, reaching proximal half of corneae; laterodistal margin of first segment unarmed.

Right cheliped with moderately broad chela; distal half of fingers with longitudinal hiatus. Dactyl approximately $\frac{3}{4}$ length of palm, with tufts of moderately long setae; with granules on mesial surface and near dorsomesial margin. Palm equal to length of carpus, with row of spinules on dorsomesial margin; dorsal surface somewhat convex with scattered tufts of moderately long setae, with spinules near mesial and lateral margins, lacking ridge near dorsomesial margin proximally; dorsolateral margin with row of spinules extending onto fixed finger and becoming obsolete near claw; dorsoproximal margin with weakly developed mesial and lateral lobes; mesial face with vestigial ridge on proximal half; ventroproximal margin with almost obsolete median sinus flanked by 2 slightly calcified sinuses, right sinus larger than left. Carpus exceeding length of merus approximately $\frac{1}{4}$ own length; dorsomesial margin with 4 or 5 acute spines interspersed with spinules; dorsomesial distal angle produced into small lobe; dorsodistal margin with vestigial projection adjacent to dorsomesial distal angle; dorsolateral margin with 1 acute spine in distal half; lateral face unarmed but with tufts of moderately long setae; ventromesial margin obtusely triangular in mesial view; ventral surface granular and with tufts of long setae; very short ventral length between very deep, subcircular ventrodistal and ventroproximal margins. Dorsal margin of merus with small distal spine; ventromesial margin with acute distal spine; ventrolateral margin with row of spinules and 2 acute distal spines; ventral surface with spinules. Coxae of chelipeds with ventrolateral distal angle unarmed.

Dactyl of left cheliped smooth but with tufts of moderately long setae. Dorsomesial and dorsolateral margins of carpus, each with 3 or 4 strong, widely spaced spines.

Dactyls of ambulatory legs exceeding propodi by approximately $\frac{1}{4}$ own length; dorsomesial and ventromesial margins of dactyls, each with widely spaced, moderately long setae. Propodi exceeding carpi $\frac{1}{2}$ to $\frac{2}{3}$ own length, each with 1 short ven-

Fig. 24. *Anapagurus wolffi* Forest, 1961. Holotype ♂ (SL = 1.0 mm), "Atlantide" station 60, 5°06'N, 9°34'W, ZMUC: A, shield and cephalic appendages; B, right antennule (lateral view); C, right cheliped (dorsal view); D, carpus of same (ventral view); E, carpus of same (ventromesial view); F, left cheliped (dorsal view); G, right 2nd pereopod (mesial view); H, left 3rd pereopod (mesial view); I, sternum; J, extremity of abdomen, uropods and telson; K, left sexual tube (lateral view). Scales equal 1.0 mm: a: K; b: A-J.






Fig. 25. Distribution of *Anapagurus wolffi* Forest, 1961. Circled star: type locality, material examined. Circled numbers: based on previous records, material examined: Forest, 1961: 1 as *A. wolffi* and *A. lac-vis*; 2, 3 as *A. curvidactylus*.

tromesial distal seta. Carpi approximately $\frac{2}{3}$ length of meri; dorsal margins with 3 or 4 widely spaced spines (2nd) or 1 proximal and 1 distal spine (3rd). Meri unarmed but each with widely spaced tufts of moderately long setae. Lobes of sternite of 2nd pereopods unarmed, distally rounded. Anterior lobe of sternite of 3rd pereopods subrectangular, slightly skewed.

Males with moderately long, left sexual tube. Coxa of 5th right pereopod with ventral gonopore distally. Pleopods 3 to 5 unequally biramous. Distal protopodal lobe of right uropod less than $\frac{1}{2}$ length of endopod.

Telson with posterior asymmetrical lobes, separated by narrow cleft; terminal margins each with 2 strong spines and tufts of moderately long, submarginal, stiff setae; lateral margins unarmed but each with moderately long setae.

Colour.—Unknown.

Related organisms.—“Les deux individus de la (“Atlantide”) station 60 étaient logés dans des fragments cylindriques de tubes de *Dentalium*” (Forest, 1961).

Distribution.—*Anapagurus wolffi* has been collected in western African waters off the coasts of Sierra Leone and Liberia. Depth range: 74-108 m.

Affinities.—Forest (1961) observed that *A. wolffi* and *A. chiroacanthus* shared a number of characters, i.e., the aspect of the anterior part of the body, the outline and thickness of the right chela, and the appearance of the ambulatory legs. However *A.*

wolffi is easily distinguished, not only from *A. chiroacanthus* but also from the other congeners, in having: the shortest antennal peduncles (last segment barely reaching cornea proximally); the unarmed lateral projections of shield, laterodistal margin of 1st antennal segment, and ventromesial distal angle of propodal 2nd pereopods; the absence of a ridge near the dorsomesial margin of the palm proximally; and the shortest ventral carpal length of the right cheliped.

Remarks.—Forest (1961) described *A. wolffi* from specimens collected by the Danish Expedition to the coasts of tropical west Africa. During my examination of the "Atlantide" collection, I found that one ovigerous female, previously identified as *A. laevis* (station 49), and one male and one ovigerous female (stations 51 and 58) doubtfully identified as *A. curvidactylus*, were conspecific with *A. wolffi*.

Anapagurus petitii Dechancé & Forest, 1962
(figs. 26, 27)

Anapagurus bicorniger; Fenizia, 1937: 33, 34 (in part; see remarks), 35, 36, figs. 23, 25, 26; Zariquey Alvarez, 1946: 122, 123; 1955: 402.

Anapagurus petitii Dechancé & Forest, 1962: 297-299, figs. 2, 6-8, 10, 12 (in part; see remarks) (type locality: Banyuls-sur-Mer, Cap l'Abeille; 12-15 m); Zariquey Alvarez, 1968: 259, 260, fig. 90q (in part; see remarks); Guille, 1971: 102, 105, 119, 123; Stevcic, 1972: 102, 103; Kocatas, 1981: 162; García Raso: 1982: 504, fig. 3; Williamson, 1982: 92; Ingle, 1985: 764 (listed).

SPECIES N.1: Pike & Williamson, 1960: 531-533, fig. 10.

Material.—Holotype: ♂ (carapace length= 4.5 mm), Banyuls-sur-Mer, Cap l'Abeille, France, 12-15 m, v.1958, collector M. Dechancé (MNHN); not examined. **Mediterranean Sea: Off the Balearic Islands:** Formentera: 1 ovig. [3.0], 45.7 m, 1871, collector F. Söderlund, NHRM 12056; 6 ♂♂ [1.5-2.3], 2 ♀♀ [1.7, 2.1], 2 ovig. [2.2], 54.9 m, 1871, collector F. Söderlund, NHRM 12057; 1 ♂ [1.9], 1 ovig. [2.6], 64 m, 1871, collector F. Söderlund, NHRM 12058; 1 ovig. [3.0], 73.2 m, 1871, collector F. Söderlund, NHRM 12059; 1 ♂ [2.8], 2 ovig. [2.3, 2.5], 91.4 m, 1871, collector F. Söderlund, NHRM 12060; San Antonio, Ibiza: 1 ♂ [2.8], 45.7 m, 1871, collector A. Lindahl, NHRM 6965; 4 ♂♂ [1.8-2.8], 1 ♀ [2.0], 45.7 m, 1871, collector F. Söderlund, NHRM 12061. **Bahía de Cadaqués:** 2 ♂♂ [1.7, 1.9], 4 ♀♀ [1.6-2.1], viii.1954, collector R. Zariquey Alvarez, MNHN 1536; 1 ♂ [1.4], 45 m, viii.1957, collectors L.B. Holthuis & R. Zariquey Alvarez, RMNH D; 6 ♂♂ [1.3-2.6], 30-40 m, gross sand, 27.vii to 2.viii.1961, collector L.B. Holthuis, RMNH D 16559; 1 ♀ [1.7], 20 m, bottom of sand and silt covered with *Posidonia*, 28.vii.1961, collector L.B. Holthuis, RMNH D 16562; 1 ♂ [2.1], 1 ♀ [1.7], 30-40 m, 31.vii.1961, collector L.B. Holthuis, RMNH D 16561; 2 ♂♂ [2.2, 3.3], 2 ♀♀ [1.7, 1.8], 30 m, viii.1961, collector L.B. Holthuis, RMNH D 16560. **Off Monaco:** 1 ♂ [2.5], 1 ♀ [2.8], 1 ovig. [1.8], 15 m, 17.iii.1909, MOM 382512. **Off Roquebrune-Cap-Martin:** 1 ovig. [2.3], 30-50 m, bottom of silt covered with *Posidonia*, 5.iv.1951, collector H. Nouvel, RMNH D. **Off Sori:** 1 ♂ [3.7], 1 ovig. [1.8], 18.3 m, ix.1887, collector H. Théel, NHRM 6942; 1 ♂ [2.3], 36.6-54.9 m, coralline bottom, 3.x.1887, collector E. Théel, SMNH 13178. **Gulf of Naples:** 1 ♂ [3.0], sandy bottom, 25.vii.1959, collector R. Bott, SMF 2551; 3 ovig. [2.9-3.3], MSNM 1867. **Off SW of Lam-pedusa:** 1 ♂ [2.5], "AAK" station 1313, 35°17.2'N, 12°04', 5'E, 53 m, sandy bottom covered with *Udotea*, 17.ix.1979, collector V.V. Murina, MMSU. **Golfo di Taranto:** 1 ♂ [3.4], station 199, 27 m, 4.viii.1966, collector A. Vatova, RMNH D 23605. **Adriatic Sea:** 1 ♀ [1.4], off Rovinj, 28 m, 13.ix.1983, SMF 12100; 2 ovig. [2.4, 3.0], off Split, 30-35 m, sandy bottom, 20.iii.1932, collector S. Bock, NHRM 13564; 1 ♂ [2.6], off Split, 30-35 m, sandy bottom, 5.iv.1932, collector S. Bock, NHRM 13953.

Diagnosis.—Shield broader than long. Rostrum slightly exceeded by lateral projections. Lateral projections with short submarginal spine. Ocular peduncles approximately $\frac{2}{3}$ length of shield; with corneae moderately dilated.

Ocular acicles broadly subovate, with short submarginal spine. Interocular plate with 2 chitinous spines usually overreaching proximal half of ocular acicles.

Antennular peduncles exceeding distal margin of corneae approximately $\frac{2}{3}$ length of ultimate segment. Ultimate antennular segment exceeding length of penultimate segment by approximately $\frac{1}{2}$ own length, with dorsolateral row of short setae interspersed with moderately long setae; ventral margin with row of very short setae; basal segment with laterodistal and dorsodistal spines.

Antennal peduncles exceeding distal margin of corneae by approximately $\frac{1}{2}$ length of ultimate segment. Antennal acicles unarmed, usually reaching proximal half of ocular peduncles.


Right cheliped with moderately narrow chela. Dactyl $\frac{1}{2}$ to $\frac{2}{3}$ length of palm, with tufts of moderately long setae and scattered granules on mesial surface and dorsal surface near dorsomesial margin. Palm slightly longer than carpus; dorsomesial margin with well-developed ridge on proximal half with row of spinules and moderately long setae; dorsal surface somewhat convex, with scattered tufts of short setae, with very low, serrate ridge near dorsomesial margin proximally; dorsolateral margin with row of moderately long setae; dorsoproximal margin with weakly developed mesial and lateral lobes; ventroproximal margin with median subquadrate sinus. Carpus approximately $\frac{3}{4}$ length of merus; dorsomesial margin with 3 or 4 widely spaced, long spines interspersed with spinules; dorsomesial distal angle produced into concave, semicircular structure; dorsodistal margin with weak, triangular projection adjacent to dorsomesial angle; dorsolateral margin with row of 3 or 4 widely spaced, long spines; distal half of lateral face usually with 2 widely spaced, long spines; ventromesial margin subcircular in mesial view; ventral surface with tufts of long setae; very short ventral length between deep, subtriangular ventrodigital margin and deep subcircular ventroproximal margin. Ventromesial margin of merus with tubercles or spines in males; dorsal margin with 2 spinules distally. Coxae of chelipeds with ventrolateral distal angle unarmed.

Dactyl of left cheliped smooth but with tufts of moderately long, stiff setae. Dorsomesial and dorsolateral margins of carpus, each with row of 5 to 8 strong, closely spaced spines.

Dactyls of ambulatory legs exceeding propodi by approximately $\frac{1}{4}$ own length, with dorsomesial row of closely spaced, moderately long setae and ventromesial row of widely spaced, short setae. Propodi exceeding carpi $\frac{2}{3}$ own length; ventromesial distal angle with 5 to 10 spines (2nd, males), 1 or 2 spinules (2nd, females; 3rd, males) or unarmed (3rd, females); dorsal margin of carpi each with 1 proximal and 1 distal spines; meri with 2, occasionally 3 widely spaced spinules on dorsal margin, and 1 spinule (2nd) or unarmed (3rd) on distal third ventrally. Lobes of sternite of 2nd pereopods unarmed, distally rounded. Anterior lobe of sternite of 3rd pereopods semicircular.

Males with moderately long, left sexual tube and conical, short, backwardly

Fig. 26. *Anapagurus petiti* Dechancé & Forest, 1962. ♂ (SL = 3.0 mm), Gulf of Naples, Italy, FNMS 2551: A, shield and cephalic appendages, B, right antennule (lateral view); C, right cheliped (dorsal view); D, carpus of same (ventromesial view); E, left cheliped (dorsal view); F, left 2nd pereopod (mesial view); G, left 3rd pereopod (mesial view); H, sternum; I, extremity of abdomen, uropods and telson; J, left sexual tube (lateral view). Scales equal 1.0 mm: a: A, C-H, J; b: B, I.






Fig. 27. Distribution of *Anapagurus petiti* Dechancé & Forest, 1962. Star: type locality. Solid circles: based on present records. Numbers: based on previous records: 1-García Raso, 1982; 2, 3-Zariquey Alvarez, 1968; 3-Zariquey Alvarez, 1946, 1955 (as *A. bicorniger*); 4-Fenizia, 1937 (as *A. bicorniger*); 5-Kocatas, 1981.

directed, right sexual tube; pleopods 3 and 4 unequally biramous, pleopod 5 uniramous. Distal protopodal lobe of right uropod approximately $\frac{1}{2}$ length of endopod.

Telson with posterior, asymmetrical lobes, separated by wide, median cleft; terminal margins each with 3 or 4 short spines.

Colour.—Unknown.

Related organisms.—One female of *A. petiti* (NHRM 12057) was found in a shell of *Turritella biplicata* covered by hydroids, foraminiferans and serpulid polychaetes; another female (RMNH D 16560) was inhabiting the shell of a naticid completely covered by colonial hydroids.

Folliculinid protozoans type 1 (see section on related organisms for *A. hyndmani*) were found dorsally on the posterior carapace and abdomen of 5 specimens of *A. petiti* (MOM 382512; and NHRM 12057, 12060 and 13953). These protozoans ranged in number from 1 to 5 specimens per host. This is the first recorded folliculinid in association with *Anapagurus petiti*.

Distribution.—*Anapagurus petiti* has been found in the Mediterranean Sea, from off Málaga, Spain (García Raso, 1982) to off Turkey, in the Aegean Sea (Kocatas, 1981). Depth range: 10 to 92 m.

Affinities.—The 2 widely spaced, strong spines on distal half of lateral face of carpus of right cheliped immediately distinguish *A. petiti* from other members of *Anapagurus*. As earlier discussed *A. petiti* is most closely related to *A. bicorniger*, however, *A. petiti* is easily distinguished from the latter species by the much broader subcircular ocular acicles, the longer interocular plate spines, the much shorter and unarmed antennal acicles, the less numerous but stronger carpal spines of the right cheliped, and the less numerous and shorter ventral setae of the dactyls of the ambulatory legs.

Remarks.— As previously indicated (see remarks under *A. bicorniger*), Dechancé & Forest (1962) were correct in their assumption that Fenizia (1937) had confounded two taxa under the name *A. bicorniger*. Balss's (1926) description and illustration pertain only to *A. bicorniger*. Zarliquiey Alvarez's (1968) report of *A. bicorniger* pertained only to *A. petiti*.

In their description of *A. petiti*, Dechancé & Forest (1962) compared its adult and glaucothoe stages with those of *A. bicorniger*. These authors further stated that the larval and young stages assigned to SPECIES N.I by Pike & Williamson (1960) belong to their new taxon.

Anapagurus alboranensis spec. nov.
(figs. 28-30)

Material.— Holotype: ♂ [1.8], Alboran Sea, off Málaga, Spain, 35-43 m, 1979, collector J.E. García-Raso, UMDZ; Paratypes: Mediterranean Sea: Alboran Sea: 2 ♂♂ [2.4, 2.9], 2 ovig. [2.2, 3.4], data as for holotype. Balearic Sea: 1 ♂ [2.8], off San Antonio, Ibiza, Balearic Islands, 45.7 m, 1871, collector F. Söderlund, NHRM 4540; 1 ♂ [1.9], off Formentera, Balearic Islands, 64 m, 1871 collector F. Söderlund, NHRM 4539. Ligurian Sea: 1 ♂ [2.0], "Princesse Alice II" station 040, off Beaulieu, France, 30 m, 1907, MOM 382511; 1 ♂ [2.3], off Sori, Italy, 36.6-54.9 m, coraline bottom, 3.x.1887, collector E. Théel, NHRM 4541. Tyrrhenian Sea: 1 ♂ [2.4], off Naples, Italy, MSNM 1869; Strait of Sicily: 1 ♂ [2.6], "AAK" station 1400, 37°28.1'N, 12°13.5'E, 65 m, sandy bottom with living and dead corals, 8.ix.1980, collectors V.V. Murina & V.D. Tchukhtchin, MMSU. Adriatic Sea: 1 ♂ [1.8], 43°30'N, 13°48'E, 20 m, 18.x.1972, collector C. Froglio, IRPEM D382; 1 ♂ [1.6], 1 ♀ [1.1], off Rovinj, Yugoslavia, 30 m, 2.vi.1962, RMNH D 19367; 1 ♂ [2.4], off Rovinj, Yugoslavia, 3.viii.1981, collector Exk. Uni Frankfurt, SMF 11086; 1 ♂ [2.1], 2 ovig. [2.0], "F.B. Burin", off Rovinj, Yugoslavia, 31 m, 18.ix.1987, collector Baumkurre, SMF 21920; 2 ♂♂ [1.5, 1.8], "F.B. Burin", off Rovinj, Yugoslavia, 30 m, 18.ix.1987, collector Baumkurre, SMF 21922; 1 ♂ [2.3], "F.B. Burin", off Rovinj, Yugoslavia, 31 m, 18.ix.1987, collector Baumkurre, SMF 21921; 1 ♂ [1.7], off Istrien Peninsula, Yugoslavia, 9.ix.1974, collector M. Türkay, SMF 7007; 1 ♂ [1.9], 1 ♀ [1.7], "Sixten Bock", off Split, Yugoslavia, 30-35 m, bottom of sand and shells, 5.iv.1932, NHRM 4542; 1 ♂ [1.6], "Sixten Bock", off Split, Yugoslavia, 30-35 m, bottom of sand and shells, iii.1932, NHRM 4543. Northeastern Atlantic: 1 ♂ [1.7], off Sanlúcar de Barrameda, SW Spain, 1975, collector E. Comanescu, SMF 7870.

Description.— Shield slightly longer than broad. Rostrum slightly overreaching lateral projections. Lateral projections with short submarginal spine.

Ocular peduncles approximately $\frac{2}{3}$ length of shield; with corneae slightly dilated; dorsomesial face with 1-3 stiff setae distally. Ocular acicles roundly triangular, with submarginal spine, separated basally by approximately basal width of 1 acicle. Interocular plate unarmed.

Antennular peduncles overreaching ocular peduncles by $\frac{2}{3}$ to $\frac{3}{4}$ length of ultimate segment. Ultimate segment exceeding length of penultimate segment by approximately $\frac{1}{2}$ own length; dorsolateral margin glabrous or with row of tufts of very short setae and laterodistal pair of setae composed of 1 long and 1 short seta; ventral margin with row of tufts of short setae. Basal segment with lateromedial and laterodistal spines.


Antennal peduncles overreaching distal margin of corneae by approximately $\frac{1}{2}$ length of ultimate segment. Fifth and fourth segments with few, stiff setae. Third segment with small, ventrolateral distal spine obscured by long stiff setae. Second segment with dorsolateral distal angle produced, terminating in strong, bifid spine; dor-

somesial distal angle with strong spine. First segment with small spine at laterodistal margin; ventromesial angle produced, with 2 or 3 small spines laterally. Antennal acicles slightly arcuate, reaching to proximal half of corneae; dorsomesial margin with row of tufts of stiff setae; terminating in small spine. Antennal flagella long, reaching dactyl of right cheliped proximally; dorsoventrally flattened, with 1-4 short (less than 1 article in length) setae on both lateral and mesial distal angles of each article.

Maxillule with stiff bristle on moderately well-developed, internal endopodal lobe, external lobe not developed; epipod produced and with tufts of long setae. Maxilla with endopod overreaching scaphognathite. First maxilliped with endopod approximately $\frac{1}{2}$ length of exopod. Third maxilliped with long spine on dorsodistal meral margin. Sternite of 3rd maxillipeds with shallow median cleft.

Right cheliped with dactyl $\frac{2}{5}$ to $\frac{2}{3}$ length of palm; cutting edge with (proximal to distal) 2 or 3 short, calcareous teeth, 3 prominent, calcareous teeth in triangular arrangement, 2 or 3 short, calcareous teeth, 1 prominent, calcareous tooth, and 3 or 4 short, calcareous teeth interspersed with corneous teeth; terminating in small corneous claw; dorsal surface slightly elevated in midline; all surface granulose and with tufts of long setae; granules more numerous and larger on dorsomesial margin. Palm slightly longer than carpus, somewhat inflated dorsoventrally; dorsal surface slightly convex, with scattered granules and tufts of short setae, and short, low granulose ridge near dorsomesial margin proximally; dorsomesial margin with somewhat prominent ridge proximally and row of closely spaced granules and tufts of short setae; dorsoproximal margin with well-developed lobes, laterally and mesially (mesial lobe usually larger); dorsolateral margin with row of closely spaced granules extending onto fixed finger; ventroproximal margin with median, skewed, subquadrate sinus; all surfaces with scattered granules and tufts of short setae; cutting edge of fixed finger with 2 proximal rows of calcareous teeth, converging to form single row of 1 large and several small, calcareous teeth interspersed with corneous teeth distally; terminating in short, corneous claw; all surfaces of fixed finger with granules (more numerous on lateral face) and scattered tufts of setae (longer and more numerous on ventral face). Carpus exceeding merus by $\frac{3}{4}$ own length; dorsomesial margin with irregular row of closely spaced, occasionally widely spaced, acute spines and tufts of stiff setae; dorsomesial distal angle produced into large, subrectangular structure, with concave distal margin; dorsodistal margin with a roundish, triangular projection adjacent to dorsomesial distal angle; dorsolateral margin with 1 acute spine in distal half; dorsolateral distal angle produced into small lobe; lateral face unarmed; dorsomesial margin subcircular in mesial view; ventral surface with numerous closely spaced granules and very long, stiff setae; moderately long ventral length between shallow subcircular ventrodistal margin and deep, subcircular ventroproximal margin; all other surfaces granulose and with scattered, short setae.

Fig. 28. *Anapagurus alboranensis* spec. nov. ♂ (SL = 2.4 mm), off Naples, Italy, MSNM 1869: A, shield and cephalic appendages; B, left antennule (lateral view); C, right cheliped (dorsal view); D, carpus and merus of same (ventromesial view); E, carpus of same (mesial view); F, left cheliped (dorsal view); G, left 2nd pereopod (mesial view); H, right 3rd pereopod (mesial view); I, sternum; J, extremity of abdomen, uropods and telson; K, left sexual tube (lateral view). Scales equal 1.0 mm: a: C-H, K; b: A, I; c: B; d: J.






Fig. 29. *Anapagurus alboranensis* spec. nov. ♂ (SL = 2.4 mm), off Naples, Italy, MSNM 1869: left mouth-parts: A, maxillule (internal view); B, maxilla (internal view); C, 1st maxilliped (internal view); D, 2nd maxilliped (external view); E, 3rd maxilliped (internal view). Scale equals 1.0 mm.

Merus inflated dorsoventrally; dorsal margin with row of long stiff setae, terminating in partially obscured, acute spine; ventrolateral margin with 2-5 acute spines increasing in size distally; ventromesial margin usually with short spine (small specimens) or 1 short spine and 2 additional spinules (large specimens), and spinulose protuberance; ventral surface with numerous tubercles and tufts of very long, stiff setae completely covering posterior carpal arthrodial membrane. Ischium with 1 spinule on ventrolateral margin partly covered by tufts of long, stiff setae; ventromesial margin with row of spinules and stiff setae; dorsodistal angle with short protuberance partly obscured by long, stiff setae. Ventromesial margin of coxa with long setae and strong, acute spine at distal angle; ventrolateral distal angle unarmed.

Left cheliped, reaching distal half of palm of right. Dactyl approximately equaling length of palm; terminating in acute, corneous claw; cutting edge with row of corneous teeth; all surfaces with scattered granules and tufts of long, stiff setae. Palm slightly shorter than carpus, somewhat inflated ventrally and laterally; dorsolateral margin with closely spaced granules extending onto fixed finger; dorsal surface with very low, granulose ridge near dorsomesial margin proximally; all surfaces with scattered granules and long stiff setae; fixed finger with granulose dorsal and lateral surfaces, ventral surface smooth, with scattered tufts of long setae on all faces; cutting edge of fixed finger with row of corneous teeth interspersed with calcareous teeth, terminating in acute corneous claw. Carpus slightly shorter than merus, inflated ventrally; dorsomesial margin with 3 to 5 acute spines interspersed with spinules; dorsomesial distal angle produced into moderately large, subrectangular structure; dorsolateral margin with 2 to 5 acute spines interspersed with spinules; dorsolateral distal angle produced into moderately large lobe; dorsal surface with tufts of long, stiff setae. Dorsal margin of merus with tufts of long, stiff setae; ventrolateral margin with 2 to 5 long, acute spines; ventromesial margin with slightly developed spinulose protuberance and 1 short distal spine; ventral face granulose and with scattered tufts of long, stiff setae. Ischium with ventromesial row of spinules, occasionally unarmed, and long, stiff setae; dorsodistal angle with short protuberance partly obscured by long, stiff setae. Coxa with long setae on ventromesial margin and single, strong, acute spine at distal angle; ventrolateral distal angle unarmed.

Dactyls of ambulatory legs exceeding propodi by approximately $\frac{1}{4}$ own length; dorsal margins each with row of moderately long, widely spaced, stiff setae, more numerous and closely spaced on distal half (3rd); mesial faces each with row of closely spaced, stiff setae near ventral margin; lateral faces with row of widely spaced setae near ventral margin. Propodi exceeding carpi $\frac{1}{2}$ to $\frac{2}{3}$ own length; dorsal surfaces each with row of tufts of moderately short, stiff setae; mesial and lateral faces each with row of short setae near ventral margin; ventromesial distal angle with 2, occasionally 1, corneous spine (2nd) or unarmed (3rd). Carpi approximately $\frac{2}{3}$ length of meri; dorsal surfaces each with row of 2 to 7 spines (2nd) or 1 to 2, commonly 2, spines (3rd), and row of tufts of short, stiff setae; ventral surfaces with scattered, short setae. Ventrolateral margins of meri with one spinule on distal half (2nd) or unarmed (3rd), and row of tufts of short setae; dorsal surfaces usually unarmed or with 1 spinule on proximal third (left 2nd) and row of tufts of short, stiff setae. Ischia with tufts of long, stiff setae on dorsal and ventral margins. Lobes of sternite of 2nd pereopods unarmed, distally rounded. Anterior lobe of sternite of 3rd pereopods




Fig. 30. Distribution of *Anapagurus alboranensis* spec. nov. Circled star: type locality. Solid circles: based on paratypic material.

subrectangular; anterior margin with long setae.

Males with left sexual tube directed diagonally toward exterior, then recurved upward to level of coxa. Coxa of 5th right pereopod with short sexual tube downwardly directed, partly covered with tufts of long, stiff setae. Pleopods 3 and 4 unequally biramous; pleopod 5 uniramous. Distal protopodal lobe of right uropod approximately $\frac{1}{2}$ length of endopod.

Terminal margins of telson each with 4 closely spaced, short spines; lateral margins each with 1 spinule or bristle and long setae; dorsal surface with scattered, short setae.

Colour.—Unknown.

Etymology.—Named in reference to the type locality, the Alboran Sea.

Related organisms.—A male of *A. alboranensis* spec. nov. (MOM 382511) was found inhabiting the gastropod shell of *Buccinum humphreysianum*.

On 3 specimens of *A. alboranensis* spec. nov. collected in the Adriatic Sea, off Split, Yugoslavia (SMF), there were 1-4 folliculinid protozoans type 1 (see section on related organisms for *A. hyndmanni*) per host; tests were cemented to the abdomen dorso-distally and to the shield; no external modification caused by these symbionts was observed. Each of two males from the type locality had 3 folliculinid protozoans type 2 (see section on related organisms for *A. hyndmanni*) attached to their abdomens. This parasitic infestation apparently changed a secondary sex character of one of the hosts, i.e., the feminization of the pleopods.

Distribution.—*Anapagurus alboranensis* spec. nov. has been collected in the Mediterranean Sea, from the Alboran Sea to the Adriatic Sea. Outside the Mediterranean Sea, this species was found off Sanlúcar de Barrameda, southwest Spain. Depth range: 30-65 m.

Affinities.—As formerly noted (see section on affinities for *A. pusillus*), *A. alboranensis*

nensis spec. nov. resembles *A. pusillus* and *A. breviaculeatus* in the general appearance of the chelipeds and ambulatory legs. The new species can be distinguished by the presence of a spinule or bristle on the lateral margin of the posterior lobes of the telson, a shorter left sexual tube, and, at least from *A. breviaculeatus*, by the shorter ocular peduncles and the much shorter ultimate antennular segment.

Anapagurus adriaticus spec. nov.
(figs. 31-33)

Anapagurus laevis; Holthuis & Gottlieb, 1958: 72 (in part; see remarks) [not *Anapagurus laevis* (Bell, 1846)].

Material.—Holotype: ♂ [3.0], 42°27.7'N, 15°48'E, off Isole Tremiti, Adriatic Sea, 83 m, 9.vii.1972, collector C. Froglia, RMNH D 42360. Paratypes: Mediterranean Sea: Balearic Sea: 5 ♂♂ [1.3-2.8], 2 ♀♀ [1.4, 2.4], 8 ovig. [1.9-2.4], "AAK" station 1246, 40°14.5'N, 1°09.2'E, 110 m, bottom of clayed sand with fragmented shells, 9.iv.1978, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 5 ♂♂ [1.7-2.4], 1 ♀ [1.9], 4 ovig. [1.6-2.1], "AAK" station 1247, 40°26.2'N, 1°09.2'E, 90 m, bottom of clayed sand, 9.iv.1978, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 1 ♂ [3.3], off Formentera, Balearic Islands, 91.4 m, 1871, collector F. Söderlund, NHM 4538; 6 ♂♂ [2.2-3.1], 2 ♀♀ [1.9, 2.4], "AAK" station 1192 (129), 40°03.3'N, 0°56.1'E, 105 m, bottom of sand and fragmented shells, 2.vi.1974, collector V.V. Murina, MMSU. Ligurian Sea: 1 ♂ [4.4], "Hirondelle II" station 3680, SW off Pointe Carqueirane, near Toulon, France, 60-80 m, 20.iv.1915, MOM 382510; 1 ♂ [2.7], 1 ovig. [2.6], "E. Gilat" station 15, off Roquebrune-Cap Martin, SE France, 72-80 m, 30.vii.1903, RMNH D 20257. Off W Sardinia: 1 ♂ [2.2], from 38°50.8'N, 8°27.2'E to 38°54.7'N, 8°28.5'E, 67-80 m, 20.ix.1973, collectors C. Froglia & G. Orel. Tyrrhenian Sea: 1 ♀ [3.1], 1 ovig. [3.0], off Naples, MCSN 1868; 1 ♂ [3.1], off Naples, MSNM 1869; 1 ♀ [3.2], from 38°07.1'N, 12°50.8'E to 38°06.2'N, 12°55.6'E, Golfo di Castellammare, NW Sicily, 350-480 m, 16.vi.1974, collectors R.B. Manning & C. Froglia, IRPEM D393. Strait of Sicily: 1 ♂ [1.9], "AAK" station 1265, 35°27.0'N, 11°30.3'E, 70 m, clayed sand bottom, 26.iv.1978, collector V.V. Murina, MMSU; 1 ♂ [2.8], "AAK" station 1392, 35°30.7'N, 12°01.2'E, 72 m, sandy bottom with *Lithothamnium*, 3.ix.1980, collectors V.V. Murina & V.D. Tchukhtchin, MMSU; 2 ♂♂ [1.8, 2.0], "AAK" station 1393, 37°04.5'N, 12°06.0'E, 110 m, clayed sand bottom, 6.ix.1980, collectors V.V. Murina & V.D. Tchukhtchin, MMSU. Adriatic Sea: 4 ♂♂ [2.1-3.1], 1 ovig. [2.6], 42°27.7'N, 15°48.0'E, off Isole Tremiti, 83 m, 9.vii.1972, IRPEM D537; 1 ♂ [1.9], 1 ♀ [2.7], 42°53'N, 14°32'E, 145 m, 5.x.1972, collector C. Froglia, USNM; 1 ♂ [2.6], 42°34'N, 15°50'E, 130 m, 4.v.1972, collector C. Froglia, IRPEM D375; 1 ovig. [2.9], 44°02.9'N, 13°46.3'E, 67 m, 11.viii.1975, collector C. Froglia, USNM; 1 ♂ [3.4], 1 ovig. [2.9], 43°05'N, 14°20'E, 100 m, 8.vii.1985, collector C. Froglia IRPEM D613; 1 ovig. [4.0], off San Benedetto, 85 m, 5.v.1972, collector C. Froglia, UMML 32: 5819; 1 ♂ [3.4], 1 ♀ [3.8], 1 ovig. [3.6], 43°40'N, 14°13'E, 78 m, 27.iv.1972, IRPEM D156; 1 ♂ [2.4], 42°53'N, 14°34'E, 140 m, 10.x.1972, IRPEM D370; 1 ovig. [3.6], 43°59'N, 13°50'E, 73 m, 11. x.1973, CF 88. Off Israel: 1 ♂ [2.0], station 6-5 (b), Haifa Bay, 79 m, 5.iv.1955, collector Gottlieb, RMNH D 13742. Southeastern Atlantic: 1 ♂ [2.5], "John Elliott Pillsbury" station 45, 5°05'N, 4°04.5'W, Gulf of Guinea, 342-346 m, 30.v.1964, USNM 259325.

Description.—Shield approximately as long as broad. Rostrum equaling lateral projections. Lateral projections with strong submarginal spine.

Ocular peduncles approximately $\frac{2}{3}$ length of shield; with corneae slightly dilated; dorsomesial face with 1-3 stiff setae distally. Ocular acicles roundly triangular, terminating in acute submarginal spine; dorsal surface concave and with row of short setae and moderately long submarginal setae on both lateral and mesial borders; separated basally by approximately basal width of 1 acicle. Interocular plate unarmed.


Antennular peduncles overreaching ocular peduncles by approximately entire length of ultimate segment. Ultimate segment exceeding length of penultimate segment by more than $\frac{1}{2}$ own length, with dorsolateral row of very short setae and laterodistal pair of setae, 1 long and 1 short; ventral margin with row of tufts of very short setae. Basal segment with lateromedial and laterodistal spines.

Antennal peduncles overreaching corneae distally by approximately $\frac{1}{2}$ length of ultimate segment. Fifth and fourth segments with few, long, stiff setae. Third segment with moderately long, ventrolateral distal spine obscured by long, stiff setae. Second segment with dorsolateral distal angle produced, terminating in strong, bifid spine; dorsomesial distal angle with strong spine. First segment with small spine at laterodistal margin, ventromesial angle produced and with 2 or 3 small spines laterally. Antennal acicles slightly arcuate, terminating in small spine; usually reaching corneae distally; dorsomesial margin with row of tufts of stiff setae. Antennal flagella nearly reaching tip of right cheliped; dorsoventrally flattened, with 1-3 short (less than 1 article in length) setae on both lateral and mesial distal angles of each article, and 2 long (1 or 2 articles in length) setae every 2 to 4 articles on proximal half.

Maxillule with stiff bristle on moderately well-developed, internal endopodal lobe, external lobe slightly developed; epipod produced and with tufts of long setae. Maxilla with endopod overreaching scaphognathite by $\frac{1}{2}$ own length. First maxilliped with endopod approximately $\frac{1}{2}$ length of exopod; flagellum distinctly 3-segmented distally. Third maxilliped with long spine on dorsodistal meral margin; carpus with short spine on dorsodistal meral margin. Sternite of 3rd maxillipeds with shallow median cleft and long setae.

Right cheliped with dactyl approximately $\frac{2}{3}$ length of palm; cutting edge with 3 proximal, prominent, calcareous teeth in triangular arrangement and short, calcareous teeth interspersed with corneous teeth distally; terminating in small, corneous claw, slightly overlapped by fixed finger; dorsal surface slightly elevated in midline; all surfaces granular and with tufts of long setae, granules more numerous and larger on dorsomesial margin. Palm slightly longer than carpus, somewhat inflated dorsoventrally; dorsal surface slightly convex, and with long, low spinulose ridge near dorsomesial margin proximally; dorsomesial margin with prominent ridge on proximal half, with row of closely spaced granules and tufts of short setae; dorsoproximal margin with well-developed lobes, laterally and mesially (mesial lobe larger and with acute spine); dorsolateral margin with row of closely spaced granules extending onto fixed finger; ventroproximal margin with median, subrectangular sinus; all surfaces with scattered granules and tufts of short setae; cutting edge of fixed finger with 2 proximal rows of calcareous teeth, converging to single row of 1 prominent and several small, calcareous teeth interspersed with corneous teeth distally; terminating in short corneous claw; all surfaces of fixed finger with granules and tufts of moderately long setae. Carpus slightly exceeding length of merus; dorsomesial margin with irregular row of closely spaced, acute spines and tufts of stiff setae; dorso-

Fig. 31. *Anapagurus adriaticus* spec. nov. ♂ (SL = 3.4 mm), 43°05'N, 14°20'E, IRPEM-D 613: A, shield and cephalic appendages; B, right antennule (lateral view); C, right cheliped (dorsal view); D, carpus of same (ventral view); E, carpus of same (mesial view); F, left cheliped (dorsal view); G, right 2nd pereopod (mesial view); H, right 3rd pereopod (mesial view); I, sternum; J, uropods and telson; K, left sexual tube (lateral view). Scales equal 1.0 mm: a: C-H; b: A, I, K; c: J; d: B.




mesial distal angle produced into moderately large lobe with more raised, granular ventral side; dorsodistal margin with acute, triangular projection adjacent to dorso-mesial distal angle; dorsolateral margin with irregular row of short spines and 1 larger, acute spine on distal half; dorsolateral distal angle produced into small lobe; ventromesial margin truncate in mesial view; ventral surface inflated, with few granules and tufts of short setae; moderately long ventral length between shallow, subtriangular, ventrodistal margin and deep, subcircular, ventroproximal margin; mesial surface slightly compressed, smooth; all other surfaces with granules and scattered, short setae. Dorsal margin of merus with row of long, stiff setae and acute distal spine partially obscured by tufts of long, stiff setae; ventrolateral margin with 2 or 3 acute spines strongest distally; ventromesial margin usually with 1 or 2 short spines, males occasionally with vestigial, spinulose protuberance; ventral surface with numerous spinules and tufts of short, stiff setae. Ischium with 1 spinule on ventrolateral margin partly covered by long, stiff setae; ventromesial margin with row of spinules and stiff setae; dorsodistal angle with short protuberance and short stiff setae. Coxa with long setae on ventromesial margin and single, strong, acute spine at distal angle; ventrolateral distal angle unarmed.

Left cheliped almost reaching base of dactyl of right. Dactyl slightly exceeding length of palm; terminating in acute corneous claw overlapped by claw of fixed finger; cutting edge with row of corneous teeth; proximal half with scattered granules, or short spinules in large specimens dorsally and mesially; all surfaces with tufts of long, stiff setae. Palm approximately $\frac{2}{3}$ length of carpus, somewhat inflated ventrally; dorsolateral margin with closely spaced granules (spinules in large specimens) extending onto fixed finger; dorsal surface with very low, spinulose ridge near dorsomesial margin proximally; all surfaces with scattered granules, or spinules in large specimens, and long, stiff setae; fixed finger with scattered granules on dorsal and lateral surfaces, ventral surface smooth, with scattered tufts of long setae on all faces; cutting edge of fixed finger with row of corneous teeth interspersed with calcareous teeth; terminating in acute, corneous claw. Carpus approximately equaling length of merus, slightly inflated ventrally; dorsomesial margin with 4 to 9 acute spines, dorsomesial distal angle produced into acute subtriangular lobe; dorsolateral margin with 4 to 8 acute spines, dorsolateral distal angle produced into acute subtriangular lobe; dorsal surface with tufts of long, stiff setae. Dorsal margin of merus with tufts of long, stiff setae; ventrolateral margin with 2 or 3, occasionally 4 or 5, acute (or short) spines; ventromesial margin with 1 short spine; ventral face granular, or spinulose in large specimens, and with scattered tufts of long, stiff setae. Ischium with row of spinules and long, stiff setae on ventromesial margin; ventrolateral margin with 1 spinule partly covered by long, stiff setae; dorsodistal angle with short protuberance partly obscured by long, stiff setae. Coxa with long setae on ventromesial margin and single short spine at distal angle; ventrolateral distal angle unarmed.

Second and 3rd pereopods overreaching right cheliped. Dactyls exceeding propodi by approximately $\frac{1}{3}$ own length; dorsal margins each with row of moderately

Fig. 32. *Anapagurus adriaticus* spec. nov. ♂ (SL = 3.4 mm), 43°05'N, 14°20'E, IRPEM-D 613: right mouthparts: A, maxillule (external view); B, maxilla (internal view); C, 1st maxilliped (external view); D, 2nd maxilliped (external view); E, 3rd maxilliped (external view). Scale equals 1.0 mm.






Fig. 33. Distribution of *Anapagurus adriaticus* spec. nov. Circled star: type locality. Solid circles: based on paratypic material. Circled number 1: based on previous record, material examined: Holthuis & Gottlieb, 1958 (as *A. laevis*).

long, closely spaced, stiff setae; mesial faces each with row of short, closely spaced, stiff setae near ventral margin; lateral faces with row of few, widely spaced setae near dorsal and ventral margins. Propodi exceeding carpi approximately $\frac{1}{3}$ own length; dorsal surfaces serrate, each with row of tufts of moderately short, stiff setae; mesial and lateral faces each with row of few, widely spaced, short setae near ventral margin; ventromesial distal angle with 1 to 4 moderately long, corneous spines (2nd), 1 short, corneous spine or occasionally unarmed (3rd). Carpi approximately $\frac{2}{3}$ length of meri; dorsal surface with row of 4 to 9 spines (2nd) or with serrate margin and 1 proximal and 1 distal, corneous spine (3rd), and row of tufts of short, stiff setae; lateral and ventral surfaces with scattered short setae. Ventrolateral margins of meri serrate, with 1 spinule on distal half (2nd) or unarmed (3rd) and row of tufts of short setae; dorsal surfaces serrate and with spinule in proximal and distal thirds (2nd) or with 1 spinule in distal third (3rd), and row of tufts of short, stiff setae. Ischia with tufts of short, stiff setae on dorsal and ventral margins. Lobes of sternite of 2nd pereopods rounded distally, unarmed or with row of spinules on lateral margins in large specimens. Anterior lobe of sternite of 3rd pereopods subrectangular, anterior margin with long setae.

Males with well-developed, left sexual tube directed diagonally toward exterior and recurved upward to level of coxa. Coxa of 5th right pereopod with short sexual tube directed backward and partly covered with tufts of long, stiff setae. Pleopods 3 and 4 unequally biramous, pleopod 5 uniramous. Distal protopodal lobe of right uropod approximately $\frac{1}{2}$ length of endopod.

Terminal margins of telson each with 4 closely spaced spines on terminal margins, decreasing in size mesially; lateral margins unarmed but each with tufts of long setae; dorsal surface with scattered, short setae.

Colour.— Unknown.

Etymology.— The species name refers to the type locality, off Isole Tremiti, Adriatic Sea.

Related organisms.— In the present study specimens of *A. adriaticus* spec. nov. have been found in shells of *Turritella biplicata* (4 specimens, MMSU "AAK" stations 1247 and 1192-129); *Natica acyonix* Marche Marchad (2 specimens, UMML; and MMSU "AAK" station 1192-129); *Trophonopsis muricatus* (Montagu) (2 specimens, MMSU "AAK" station 1192-129; and NHRM 12060); *Taranais moerchi* (Malm) (1 specimen, MMSU "AAK" station 1192-129); and *Cantharidus striatus* (Linnaeus) (1 specimen, MMSU "AAK" station 1214-169).

There is no external evidence of symbionts in any of the specimens examined.

Distribution.— *Anapagurus adriaticus* spec. nov. has been collected in the Mediterranean Sea from the Balearic Sea to off the coast of Israel. One male collected in the Gulf of Guinea is the only record of this new species outside the Mediterranean Sea. Depth range: 30-480 m; most frequently found in depths of 65 to 110 m.

Affinities.— As earlier discussed, *A. adriaticus* spec. nov. and *A. laevis* share the overall appearance and armature but the new species is distinguished by its much narrower shield; less dilated corneae; the roundly triangular ocular acicles, with no expanded mesial margins; and the armed, distally rounded lobes of the sternite of the 2nd pereopods in large specimens.

Remarks.— As previously stated, Holthuis & Gottlieb (1958) questionably assigned to *A. laevis* specimens collected off the coast of Israel. Examination of these specimens has shown that, except for one male from Haifa Bay that actually is *A. adriaticus* spec. nov., the others belong to *A. breaviaculeatus*.

Anapagurus vossi spec. nov.

(figs. 34-36)

Anapagurus laevis; Forest, 1961 : 239 (in part; see remarks) [not *Anapagurus laevis* (Bell, 1846)], not figs. 8, 12, 16 (= *Anapagurus atlantidii* spec. nov.).

Material.— Holotype: ♂ [1.0], "John Elliott Pillsbury" station 24, off Ghana, Africa, from 4°56'N, 0°47.5'W to 4°56'N, 0°50'W, 35-37 m, 28.v.1964, collector UMML, MCZ Crustacea No. 13337. Paratypes: 1 ♂ [1.2], "Atlantide" station 154, off Portuguese Guinea, Africa, 11°54'N, 17°14'W, 55-80 m, bluish mud bottom, 17.iv.1946, ZMUC; 1 ovig. [1.0], "Atlantide" station 153, off French Guinea, Africa, 10°49'N, 16°39'W, 42 m, bottom of gray sand, 16.iv.1946, ZMUC; 1 ovig. [1.2], "Atlantide" station 49, off Sierra Leone, Africa, 7°29'N, 13°38'W, 74-79 m, muddy sand bottom, 30.xii.1945, ZMUC; 2 ♂♂ [0.9, 1.0] [found in the stomach of *Arnoglossus imperialis* (Rafinesque, 1810)], "Atlantide" station 70, off Gold Coast, Africa, 4°50'N, 2°49'W, 65 m, bottom of mud and bryozoans, 15.i.1946, ZMUC; 2 ♂♂ [1.4, 1.5], 1 ovig. [1.2], "John Elliott Pillsbury" station 68, off Liberia, Africa, from 4°23'N, 8°05.5'W to 4°24'N, 8°07.5'W, 70 m, 3.vi.1964, USNM 259326; 1.5 ♂ [1.5], "John Elliot Pillsbury" station 68, off Liberia, Africa, from 4°23'N, 8°05.5'W to 4°24'N, 8°07.5'W, 70 m, 3.vi.1964, UMML 32: 5820; 1 ♂ [1.4], "Atlantide" station 123, off Gabon, 2°03'S, 9°05'E, 49-50 m, bottom of mud, shells and corals, 5.iii.1946, ZMUC.

Description.— Shield broader than long. Rostrum equaling lateral projections. Lateral projections with short submarginal spine.

Ocular peduncles approximately $\frac{3}{4}$ length of shield; with corneae strongly dilat-

ed; dorsomesial face with 1-3 stiff setae. Ocular acicles with mesial margins more strongly expanded than lateral margins; terminating in acute submarginal spine; dorsal surface concave and with few, short setae, long submarginal setae on both lateral and mesial borders; separated basally by approximately basal width of 1 acicle. Interocular plate unarmed.


Antennular peduncles overreaching ocular peduncles by approximately entire length of ultimate segment. Ultimate segment exceeding length of penultimate segment by approximately $\frac{1}{2}$ own length; with dorsolateral row of short setae and laterodistal tuft of 5-7 long setae decreasing in size laterally; dorsomesial margin with row of short setae; ventral margin with row of tufts of very short setae. Basal segment unarmed but with tufts of moderately long setae.

Antennal peduncles overreaching cornea distally by approximately $\frac{1}{3}$ length of ultimate segment. Fifth and fourth segments with few long stiff setae. Third segment with short ventrolateral distal spine obscured by long stiff setae. Second segment with dorsolateral distal angle produced, terminating in single, strong spine; dorsomesial distal angle with moderately long spine. First segment with short spine at laterodistal margin; ventromesial angle produced, with 2 or 3 small spines laterally. Antennal acicles slightly arcuate, usually reaching cornea distally; dorsomesial margin with row of tufts of stiff setae; terminating in small spine. Antennal flagella nearly reaching dactyl of right cheliped proximally; dorsoventrally flattened, with 1-3 short (less than 1 article in length) setae on both lateral and mesial distal angles of each article and 2 moderately long (1 article in length) setae every 2 to 4 articles.

Maxillular endopod with stiff seta distally, no internal and external lobes. Maxilla with endopod overreaching scaphognathite by $\frac{1}{4}$ own length. First maxilliped with endopod approximately $\frac{1}{2}$ length of exopod. Dorsodistal margin of carpus and propodus of third maxilliped, each with very short spine; basis with 3 acute spines; mesial margin of exopod with 1 or 2 moderately long setae. Sternite of 3rd maxillipeds with shallow median cleft and long setae.

Right cheliped with dactyl approximately subequal to length of palm; cutting edge with 3 prominent, proximal, calcareous teeth in triangular arrangement and several shorter distal teeth distributed as follows: 3 or 4 short, calcareous teeth, 1 prominent, calcareous tooth and several very short calcareous teeth interspersed with corneous teeth; terminating in small, corneous claw, slightly overlapped by fixed finger; mesial face granular and with tufts of long setae; all other surfaces smooth but, also with tufts of long setae. Palm slightly shorter than carpus, somewhat inflated dorsoventrally; dorsal surface slightly convex, with long low spinulose ridge near dorsomesial margin proximally; dorsomesial margin with slightly developed ridge on proximal half and various rows of closely spaced granules; dorsoproximal margin with weakly developed lobes, laterally and mesially; dorsolateral margin well delimited, with row of closely spaced granules extending to fixed finger;


Fig. 34. *Anapagurus vossi* spec. nov. ♂ (SL = 1.5 mm), "John Elliot Pillsbury" station 68, from 4°23'N, 8°05.5'W to 4°24'N, 8°07.5'W, USNM: A, shield and cephalic appendages; B, right antennule (lateral view); C, right cheliped (dorsal view); D, carpus of same (ventral view); E, carpus of same (ventromesial view); F, left cheliped (dorsal view); G, right 2nd pereopod (mesial view); H, right 3rd pereopod (mesial view); I, sternum; J, extremity of abdomen, uropods and telson; K, left sexual tube (lateral view). Scales equal 1.0 mm: a: A-H; b: I, K; c: J.



ventroproximal margin with median, low subrectangular sinus flanked by 2 slightly calcified sinuses, right subcircular sinus larger and deeper than the left; lateral and mesial faces with scattered granules; ventral surface smooth; all surfaces with scattered tufts of short setae; cutting edge of fixed finger with 2 proximal rows of calcareous teeth converging to single row of 1 prominent and several small calcareous teeth interspersed with corneous teeth distally; terminating in short, corneous claw; all surfaces of fixed finger with tufts of long setae. Carpus approximately equaling length of merus; distal half of dorsomesial margin with 4 strong, acute spines interspersed with spinules, distal spine above dorsomesial distal lobe distinctly directed upward, dorsomesial distal angle produced into moderately large, concave structure, more extended forward ventrally than dorsally; dorsodistal margin with weak triangular projection adjacent to dorsomesial angle; dorsolateral margin with row of spinules and 1 short spine on proximal third; dorsolateral distal angle produced into a structure similar to, but approximately half the size of the dorsomesial distal structure; ventromesial margin subcircular in mesial view; ventral surface inflated with numerous granules and tufts of long setae; short ventral length between subtriangular ventrodistal margin and subcircular ventroproximal margin; mesial surface slightly compressed, granular, with numerous tufts of long, stiff setae; ventral surface granular, with numerous tufts of long stiff setae; dorsal and lateral surfaces granular, with scattered tufts of short setae. Merus with row of long, stiff setae on dorsal margin and partially obscured short, distal spine; ventrolateral margin usually with 2 widely spaced, strong, acute spines, occasionally with 5 closely spaced, acute spines increasing in size distally; ventromesial margin usually with 1 short spine, and a low granular protuberance; ventral surface with numerous granules and tufts of short, stiff setae. Ischium with 1 spinule on ventrolateral margin partly covered by short, stiff setae; ventromesial margin unarmed, but with row of moderately long, stiff setae. Coxa with ventromesial margin with long setae and strong acute spine at distal angle; ventrolateral distal angle unarmed.

Left cheliped usually overreaching base of dactyl of right. Dactyl slightly exceeding length of palm; terminating in acute corneous claw overlapped by claw of fixed finger; cutting edge with row of corneous teeth; all surfaces smooth, but with tufts of long, stiff setae. Palm approximately $\frac{2}{3}$ length of carpus, somewhat inflated ventrally; dorsolateral margin abruptly curving and narrowing at base of fixed finger, well-delimited by closely spaced granules extending into fixed finger; dorsal surface granular with very low, wide granular median ridge proximally and with scattered tufts of short setae; all other surfaces granular, with tufts of setae, short on lateral face, long on ventral and mesial faces; fixed finger with scattered granules on dorsal and lateral surfaces, ventral surface smooth, with scattered tufts of long setae on all faces; cutting edge of fixed finger with row of corneous teeth interspersed with calcareous teeth, terminating in acute, corneous claw. Carpus exceeding merus by approximately $\frac{1}{4}$ own length, slightly inflated ventrally; dorsomesial margin with 4 or 5 acute spines and tufts of long setae, dorsomesial distal angle produced into subcircular

Fig. 35. *Anapagurus vossi* spec. nov. ♂ (SL = 1.5 mm), "Pillsbury" station 68, from 4°23'N, 8°05.5'W to 4°24'N, 8°07.5'W, USNM: right mouthparts: A, maxillule (external view); B, maxilla (internal view); C, 1st maxilliped (external view); D, 2nd maxilliped (external view); E, 3rd maxilliped (external view). Scale equals 1.0 mm.



lobe; dorsolateral margin with numerous, closely spaced acute spines of diverse sizes and tufts of long setae, dorsolateral distal angle produced into subtriangular lobe; dorsal surface somewhat depressed, glabrous. Merus with tufts of long, stiff setae on dorsal margin; ventrolateral margin with 2 widely spaced, strong, acute spines; ventromesial margin with 1 short spine; ventral face granular and with scattered tufts of long, stiff setae. Ischium with 1 spinule on ventrolateral margin partly covered by short, stiff setae; ventromesial margin unarmed, but with row of moderately long stiff setae. Coxa with ventromesial margin with long setae and strong, acute spine at distal angle; ventrolateral distal angle unarmed.

Second and 3rd pereopods overreaching right cheliped. Dactyls exceeding propodi by approximately $\frac{1}{4}$ own length; dorsal margins each with row of long, closely spaced, stiff setae; mesial faces each with row of short, closely spaced, stiff setae near ventral margin; lateral faces each with row of few widely spaced setae near dorsal and ventral margins. Propodi exceeding carpi $\frac{1}{3}$ to $\frac{1}{2}$ own length; dorsal surfaces each with row of tufts of short, stiff setae; mesial and lateral faces each with row of few widely spaced, short setae near ventral margin; ventromesial distal angle with corneous spines, 3 or 4 long (2nd pereopods, males), 2 short (2nd pereopods, females), 1 or 2 very short (3rd pereopods, both sexes). Carpi approximately $\frac{1}{2}$ length of meri; dorsal margins each with 1 proximal and 1 distal spine and row of tufts of short, stiff setae; lateral and ventral surfaces each with scattered, short setae. Ventrolateral margins of meri each with row of tufts of short, stiff setae, 1 spinule on distal half (2nd) or unarmed (3rd); dorsal surfaces unarmed, but with row of tufts of short, stiff setae. Ischia with tufts of short, stiff setae on dorsal and ventral margins. Lobes of sternite of 2nd pereopods unarmed, distally rounded. Anterior lobe of sternite of 3rd pereopods subrectangular, anterior margin with long setae.

Males with relatively short and broad, left sexual tube recurved upward to level of coxa of 5th pereopod. Coxa of 5th right pereopod with well-developed, short sexual tube backwardly directed, partly covered with tufts of long, stiff setae. Pleopod 3 unequally biramous; pleopods 4 and 5 uniramous. Right distal protopodal lobe overreaching $\frac{1}{2}$ length of uropodal endopod.

Terminal margins of telson with 2 or 3 unequal, widely spaced spines and long, submarginal, stiff setae; lateral margins unarmed but each with tufts of long setae; distal half of dorsal surface heavily setose.

Colour.—Unknown.

Etymology.—This species is named for the late Dr. Gilbert L. Voss in recognition for his guidance, moral support and friendship throughout my graduate studies.

Related organisms.—Two males of *A. vossi* spec. nov., inhabiting the gastropod shells of *Nassa semistriata* Brocchi, were found in the stomach of a bothid fish, *Arnoglossus imperialis*. There is no external evidence of symbionts in any of the specimens examined.

Distribution.—Specimens of *A. vossi* spec. nov. were collected off the tropical western coast of Africa from Portuguese Guinea to Gabon. Depth range: 35-80 m.

Affinities.—As earlier discussed (see section on affinities for *A. laevis*), *A. vossi* spec. nov. shares a number of characters with *A. laevis* and *A. congolensis* spec. nov. *Anapagurus vossi* spec. nov. can be distinguished from the latter two taxa by the ultimate antennular segment with a laterodistal tuft of five to seven long setae, the right




Fig. 36. Distribution of *Anapagurus vossi* spec. nov. Circled star: type locality. Solid circle: based on present record. Circled numbers: based on previous records, material examined: Forest, 1961 (as *A. laevis*).

cheliped with a carpal, dorsomesial, distal spine upwardly directed, the distal armature of the telson with long, submarginal, stiff setae, and in males the relatively shorter and thicker, left sexual tube, and much longer and more conical, right sexual tube.

Remarks.—As previously stated, Forest (1961) reported several specimens collected by the "Atlantide" off the tropical western coast of Africa as *A. laevis*. From my examination of these specimens I have concluded that none represent this taxon, but rather represent three different taxa, *A. wolffi*, *A. vossi* spec. nov., and *A. atlantidii* spec. nov.

R.W. Ingle (personal communication, 1991) is going to describe a presumably new species of *Anapagurus* from Adventure Bank, Mediterranean that appears to have many features of *A. vossi* spec. nov. It has particularly noticeably broadened corneae, but differs in having one or more spines on the lower meral margin of the third pereopod, a dorso-distal carpal and meral spine on the third maxilliped endopod, and the propodal palm of the right cheliped being much broader and with small scattered granules (tubercles). In addition, Ingle's specimens have a different setal armature of the antennule first peduncular segment.

Anapagurus atlantidii spec. nov.
(figs. 37-39)

Anapagurus laevis; Forest, 1961: 239 (in part; see remarks), figs. 8, 12, 16 [not *Anapagurus laevis* (Bell, 1846)].

Material.— Holotype: ♂ [0.9], "Atlantide" station 146, off French Guinea, Africa, 9°27'N, 14°48'W, 50 m, bottom of shells and foraminiferans, 13.iv.1946, ZMUC; Paratypes: 1 ♂ [0.9], 3 ovig. [1.3-1.5], same data as holotype; 1 ♂ [1.0], "Atlantide" station 145, off French Guinea, Africa 9°20'N, 14°15'W, 32 m, 13.iv.1946, ZMUC.

Description.— Shield broader than long. Rostrum slightly overreaching lateral projections. Lateral projections with short submarginal spine.


Ocular peduncles approximately $\frac{3}{4}$ length of shield; with corneae moderately dilated; dorsomesial face with 1 or 2 short, stiff setae. Ocular acicles roundly triangular, terminating in acute submarginal spine; dorsal surface concave and with few short setae, long submarginal setae on both lateral and mesial borders; separated basally by approximately basal width of 1 acicle. Interocular plate unarmed.

Antennular peduncles overreaching ocular peduncles by approximately entire length of ultimate segment. Ultimate segment exceeding length of penultimate segment by approximately $\frac{1}{3}$ own length, with dorsolateral row of short setae and laterodistal set of 2 or 3 long setae; dorsomesial margin with row of short setae; ventral margin with row of tufts of moderately long setae. Basal segment with short lateromedial spine.

Antennal peduncles overreaching corneae distally by approximately $\frac{1}{4}$ length of ultimate segment. Fifth and fourth segments with few long, stiff setae. Third segment with short, ventrolateral, distal spine obscured by long, stiff setae. Second segment with dorsolateral distal angle produced, terminating in strong, bifid spine; dorsomesial distal angle with moderately strong spine. First segment with small spine at laterodistal margin; ventromesial angle produced and with 2 or 3 small spines laterally. Antennal acicles slightly arcuate, terminating in small spine; usually reaching proximal half of corneae, dorsomesial margin with row of tufts of stiff setae. Antennal flagella usually overreaching tip of right cheliped; dorsoventrally flattened, with 1-3 short (less than 1 article in length) setae on both lateral and mesial distal angles of each article and 2 moderately long (equal to or more than 1 article in length) setae every 2 to 4 articles.

Maxillular endopod with stiff seta distally, no internal and external lobes. Maxilla with endopod slightly overreaching scaphognathite. First maxilliped with endopod approximately $\frac{1}{2}$ length of exopod. Dorsodistal margin of merus of third maxilliped with short spine; basis with 3 acute spines; mesial margin of exopod with 1 or 2


Fig. 37. *Anapagurus atlantidii* spec. nov. A-J: ovig. (SL = 1.5 mm), "Atlantide" station 146, 9°27'N, 14°48'W, ZMUC: A, shield and cephalic appendages; B, right antennule (lateral view); C, right cheliped (dorsal view); D, carpus of same (ventral view); E, carpus of same (ventromesial view); F, left cheliped (dorsal view); G, left 2nd pereopod (mesial view); H, left 3rd pereopod (mesial view); I, sternum; J, extremity of abdomen, uropods and telson. K, L: holotype ♂ (SL = 0.9 mm), "Atlantide" station 146, 9°27'N, 14°48'W, ZMUC: K, sternum; L, left sexual tube (lateral view). Scales equal 1.0 mm: a: C-F; b: A, B, G-I; c: J-L.



moderately long setae. Sternite of 3rd maxillipeds with shallow, median cleft and long setae.

Right cheliped with dactyl subequal to length of palm; cutting edge with 3 prominent, proximal, calcareous teeth in triangular arrangement and several, shorter, distal teeth distributed as follows: 2 or 3 short, calcareous teeth, 1 prominent, calcareous tooth and several, very short, calcareous teeth interspersed with corneous teeth; terminating in small, corneous claw; slightly overlapped by fixed finger; mesial face granular and with tufts of long, stiff setae; all other surfaces smooth but with tufts of long setae. Palm subequal to length of carpus, somewhat inflated dorsoventrally; dorsal surface slightly convex, with very small granules and long, low spinulose ridge near dorsomesial margin proximally; dorsomesial margin with slightly developed ridge on proximal half and row of closely spaced granules and tufts of short setae; dorsoproximal margin with well-developed mesial lobe and weakly developed lateral lobe; dorsolateral margin well delimited by row of closely spaced spinules extending onto fixed finger; ventroproximal margin with median, low subrectangular sinus flanked by 2 slightly calcified sinuses, right subcircular sinus larger and deeper than left; lateral and mesial faces with closely spaced granules; ventral surface with very small granules; mesial surface with tufts of long setae, all other surfaces with scattered tufts of short setae; cutting edge of fixed finger with 2 proximal rows of calcareous teeth converging distally to single row of 1 prominent and several small, calcareous teeth interspersed with corneous teeth; terminating in short corneous claw; all surfaces of fixed finger with tufts of long setae. Carpus approximately equaling length of merus; dorsomesial margin with widely spaced long, stiff setae, and 3 or 4 long spines on distal half increasing in size distally, distalmost spine upwardly directed; dorsomesial distal angle produced into concave, subrectangular structure; dorsodistal margin with weak, triangular projection adjacent to dorsomesial distal angle; dorsolateral margin with row of spinules and widely spaced, moderately long, stiff setae, 1 short, acute spine on proximal third; dorsolateral distal angle produced into small lobe; mesial surface slightly compressed, granular, with numerous tufts of long, stiff setae; ventromesial margin conical in mesial view; ventral surface inflated, granular and with numerous tufts of long, stiff setae, except for a glabrous, slightly depressed median part; short ventral length between deep, subtriangular, ventrodistal margin and deep subcircular ventroproximal margin; dorsal surface slightly depressed, with scattered granules and tufts of short setae; lateral surface somewhat convex, granular and with scattered tufts of short setae. Merus with row of long stiff setae on dorsal margin and partially obscured, short, acute distal spine; ventrolateral margin with 3 or 4 acute spines; ventromesial margin usually with 1 acute spine, and low, spinulose protuberance; ventral surface with numerous granules and tufts of short, stiff setae. Ischium with 1 spinule on ventrolateral margin partly covered by short, stiff setae; ventromesial margin unarmed, but with row of moderately long, stiff setae. Coxa with 1 spinule on ventrolateral distal angle; ven-

Fig. 38. *Anapagurus atlantidii* spec. nov. A-E: ovig. (SL = 1.5 mm), "Atlantide" station 146, 9°27'N, 14°48'W, ZMUC: right mouthparts: A, maxillule (external view); B, maxilla (internal view); C, 1st maxilliped (external view); D, 2nd maxilliped (external view); E, 3rd maxilliped (external view). F. ovig. (SL=1.3 mm), "Atlantide" station 146, 9°27'N, 14°48'W, ZMUC: specimen in a gastropod shell completely encrusted by a colonial bryozoan. Scales equal 1.0 mm: a: F; b: A-E.



tromesial margin with tufts of moderately long, stiff setae, partly covering 1 short spine at distal angle; ventrolateral distal angle unarmed.

Left cheliped reaching distal fifth of palm of right. Dactyl exceeding palm by approximately $\frac{1}{4}$ own length; terminating in acute corneous claw overlapped by claw of fixed finger; cutting edge with row of corneous teeth; all surfaces with scattered spinules and tufts of long, stiff setae. Palm approximately $\frac{2}{3}$ length of carpus, somewhat inflated ventrally; dorsolateral margin well delimited by closely spaced spinules extending onto fixed finger; dorsal surface with spinules, more numerous on proximal half and with scattered tufts of short setae; very low, wide, spinulose median ridge proximally; dorsoproximal margin flanked by well-developed mesial lobe and poorly developed lateral lobe; dorsomesial margin with row of closely spaced spinules and tufts of moderately long, stiff setae; lateral surface spinulose and with scattered tufts of short setae; mesial and ventral surfaces smooth but with tufts of long, stiff setae; fixed finger with scattered spinules on dorsal and lateral surfaces, ventral surface smooth; scattered tufts of long setae on all faces; cutting edge of fixed finger with row of corneous teeth interspersed with calcareous teeth, terminating in acute, corneous claw. Carpus subequal to length of merus, slightly inflated ventrally; dorsomesial margin with 4 or 5 acute spines interspersed with spinules and tufts of long setae, dorsomesial distal angle produced into subcircular lobe; dorsolateral margin with 4 or 5 widely spaced, acute spines interspersed with spinules and tufts of long setae, dorsolateral distal angle produced into subtriangular lobe; dorsal surface somewhat depressed, glabrous. Merus with tufts of long, stiff setae on dorsal margin; ventrolateral margin with 2 closely spaced, strong, acute spines; ventromesial margin with 1 short spine; ventral face spinulose and with scattered tufts of long, stiff setae. Ischium with 1 spinule on ventrolateral margin partly covered by short, stiff setae; ventromesial margin unarmed, but with row of moderately long, stiff setae. Coxa with 1 spinule on ventrolateral distal angle; ventromesial margin with tufts of moderately long, stiff setae and 1 short, acute spine at distal angle; ventrolateral distal angle unarmed.

Second and 3rd pereopods overreaching right cheliped. Dactyls exceeding propodi by approximately $\frac{1}{3}$ own length; dorsal margins each with row of moderately long, widely spaced, stiff setae; mesial faces each with row of short, widely spaced, stiff setae near ventral margin; lateral faces each with row of few widely spaced setae near dorsal and ventral margins. Propodi exceeding carpi approximately $\frac{1}{3}$ own length; dorsal surfaces spinulose, each with row of widely spaced, short, stiff setae; mesial and lateral faces each with row of few, widely spaced, short setae near ventral margin; ventromesial distal angle with 1 corneous spine (2nd) or unarmed (3rd). Carpi approximately $\frac{2}{3}$ length of meri; dorsal margin each with 2 or 3 spines interspersed with spinules (2nd) or granular and with 1 dorsodistal spine (3rd), lateral and mesial faces glabrous; ventral surfaces each with scattered short setae. Ventrolateral margins of meri each with row of tufts of short setae, 2nd serrate and with 1 spinule on distal half, 3rd unarmed; dorsal surface with row of tufts of short, stiff setae and 1 short spine on distal half. Ischia with tufts of short, stiff setae on dorsal and ventral margins. Lobes of sternite of 2nd pereopods unarmed, distally rounded. Anterior lobe of sternite of 3rd pereopods subtrapezoidal to subrectangular, anterior margin with long setae.




Fig. 39. Distribution of *Anapagurus atlantidii* spec. nov. Circled star: type locality. Circled number 1: based on previous record, material examined: Forest, 1961 (as *A. laevis*).

Males with relatively short and broad, left sexual tube recurved upward to level of coxa of 5th pereopod. Coxa of 5th right pereopod with gonopore partly obscured by tufts of long setae. Three uniramous pleopods. Right distal protopodal lobe approximately $\frac{1}{2}$ length of uropodal endopod.

Terminal margins of telson with 4 subequal, closely spaced spines and tufts of moderately long, submarginal, stiff setae; lateral margins unarmed but each with moderately long setae; distal half of dorsal surface with scattered short setae.

Colour.— Unknown.

Etymology.— This species is named for the research vessel "Atlantide" in recognition of its marine investigations off the West African coast that provided the specimens for the species description.

Related organisms.— One of the ovigerous females of *A. atlantidii* spec. nov. collected at station 146 by the "Atlantide" (ZMUC) was found inhabiting a gastropod shell completely encrusted by an unidentified colony of bryozoan, similar to the association previously reported in this paper for *A. curvidactylus*.

Distribution.— Specimens of *A. atlantidii* spec. nov. have been collected only in tropical waters off the coast of French Guinea, Africa. Depth range: 32-50 m.

Affinities.— The presence of ventrolateral, distal, coxal spines on the chelipeds in *A. atlantidii* spec. nov. is shared with *A. hyndmanni* and *A. curvidactylus*, however, the new species can be easily distinguished from the other two taxa by its right cheliped with a palmar, spinulose ridge near its dorsomesial margin proximally, its carpal

dorsomesial row of spines with a distalmost spine directed upward, and its much shorter ventral carpal length.

Remarks.— During his study of the “Atlantide” collections, Forest (1961), as previously noted, confounded three taxa under the name *Anapagurus laevis*. The specimens collected at stations 145 and 146 by the “Atlantide” are herein assigned to the new taxon, *A. atlantidii* spec. nov.

Anapagurus congolensis spec. nov.
(figs. 40-42)

Anapagurus laevis; Forest, 1955: 131 (in part; see remarks) [Not *Anapagurus laevis* (Bell, 1846)].

Material.— Holotype: ♂ [2.3], “Mbizi” station 172, 30 miles SW off Pointe Noire, Congo, Africa, 5°15'S, 11°29'E, 225-240 m, bottom of sand and silt, 2.iv.1949, IRSNB-IG-16.808; Paratypes: 1 ♂ [2.9], “Mbizi” station 9, 26 miles WSW off Moita Seca, Congo, Africa, 6°21'S, 11°53'12"E, 100 m, bottom of sand, silt and brown coral, 4-5.viii.1948, IRSNB-IG-16.808; 4 ♂♂ [2.2-2.8], “Mbizi” station 10, 42 miles SW off Moita Seca, Congo, Africa, from 6°28'42"S, 11°36'30"E to 6°16'30"S, 11°45"E, 125 m, bottom of brown sandy silt, 5-6.viii.1948, IRSNB-IG-16.808; 8 ♂♂ [1.9-3.3], 3 ♀♀ [2.2-2.3], 2 ovig. [1.8, 2.3], “Mbizi” station 166, 35 miles W by S off Pointe Noire, Congo, Africa, 4°57'S, 11°16'E, 170 m, bottom of sand and silt, 28.iii.1949, IRSNB-IG-16.808; 2 ♂♂ [2.6, 3.4], 1 ovig. [2.6], “Mbizi” station 171, 32 miles W off Pointe Noire, Congo, Africa, 4°48"S, 11°30'E, 135 m, bottom of green sand and silt, 1-2.iv.1949, IRSNB-IG-16.808; 1 ♂ [2.6], “Mbizi” station 172, 30 miles SW off Pointe Noire, Congo, Africa, 5°15'S, 11°29'E, 225-240 m, bottom of sand and silt, 2.iv.1949, IRSNB-IG-16.808.


Description.— Shield broader than long. Rostrum slightly overreached by lateral projections. Lateral projections with short submarginal spine.

Ocular peduncles approximately $\frac{2}{3}$ length of shield; with corneae moderately dilated; dorsomesial face with 1 or 2 short, stiff setae. Ocular acicles roundly triangular, terminating in acute submarginal spine; dorsal surface concave and with few short setae; with long submarginal setae on both lateral and mesial borders; separated basally by slightly less than basal width of 1 acicle. Interocular plate unarmed.

Antennular peduncles overreaching ocular peduncles by approximately $\frac{3}{4}$ length of ultimate segment. Ultimate segment exceeding length of penultimate segment by approximately $\frac{1}{2}$ own length, with dorsolateral row of moderately long setae and laterodistal tuft of 2 or 3 long setae; dorsomesial margin with row of short setae; ventral margin with row of tufts of very short setae. Basal segment with short lateromedial and laterodistal spines.

Antennal peduncles overreaching distal margin of corneae by approximately $\frac{1}{4}$ length of ultimate segment. Fifth and fourth segment with few, long, stiff setae. Third segment with short, ventrolateral, distal spine obscured by long, stiff setae. Second segment with dorsolateral distal angle produced, terminating in strong, bifid spine; dorsomesial distal angle with moderately strong spine. First segment with small

Fig. 40. *Anapagurus congolensis* spec. nov. ♂ (SL = 2.6 mm), “Mbizi” station 172, 5°15'S, 11°29'E, IRSNB-IG-16.808: A, shield and cephalic appendages; B, right antennule (lateral view); C, right cheliped (dorsal view); D, carpus of same (ventromesial view); E, left cheliped (dorsal view); F, left 2nd pereopod (mesial view); G, left 3rd pereopod (mesial view); H, sternum; I, extremity of abdomen, uropods and telson; J, left sexual tube (lateral view). Scales equal 1.0 mm: a: A, C-H; b: B, I, J.




spine at laterodistal margin; ventromesial angle produced and with 2 or 3 small spines laterally. Antennal acicles slightly arcuate, terminating in small spine; usually not reaching corneae distally, dorsomesial margin with row of tufts of stiff setae. Antennal flagella usually overreaching tips of walking legs; dorsoventrally flattened, with 1-3 short (less than 1 article in length) setae on both lateral and mesial distal angles of each article, and 2 moderately long (1 article in length) setae every 2 to 4 articles.

Maxillular endopod with distal, stiff bristle, lacking internal and external lobes. Maxilla with endopod not reaching scaphognathite distally. First maxilliped with endopod approximately $\frac{1}{2}$ length of exopod. Dorsodistal margin of merus of third maxilliped with short spine; basis unarmed; mesial margin of exopod with numerous moderately long setae. Sternite of 3rd maxillipeds with shallow median cleft and long setae.

Right cheliped with dactyl subequal to length of palm; cutting edge with 3 prominent, proximal, calcareous teeth in triangular arrangement, and several shorter, distal teeth distributed as follows: 2 or 3 short, calcareous teeth, 1 prominent, calcareous tooth and several very short, calcareous teeth interspersed with corneous teeth; terminating in small, corneous claw; slightly overlapped by fixed finger; mesial face granular and with tufts of moderately long, stiff setae; all other surfaces smooth but with tufts of moderately long setae. Palm approximately $\frac{3}{4}$ length of carpus, somewhat inflated dorsoventrally; dorsomesial margin with row of closely spaced spinules and tufts of short setae; dorsal surface slightly convex with very small granules, with long, low spinulose ridge near dorsomesial margin proximally; dorsolateral margin well delimited by row of closely spaced spinules extending onto fixed finger; dorsoproximal margin with well-developed mesial lobe armed with 2 or 3 spinules and weakly developed lateral lobe; mesial face with slightly developed ridge on proximal half; ventroproximal margin with median, low subrectangular sinus flanked by 2 small, slightly calcified sinuses; lateral and mesial faces with closely spaced spinules; ventral surface with very small granules; mesial surface with tufts of moderately long setae, all other surfaces with scattered tufts of short setae; cutting edge of fixed finger with 2 proximal rows of calcareous teeth converging to single row of 1 prominent and several small, calcareous teeth interspersed with corneous teeth distally; terminating in short corneous claw; all surfaces of fixed finger with tufts of moderately long setae. Carpus approximately $\frac{1}{4}$ longer than merus; dorsomesial margin with 8 to 12 closely spaced spines increasing in size distally except for 2 most distal, smaller spines; dorsomesial distal angle produced into concave, subrectangular structure with more raised ventral side; dorsodistal margin with strong triangular projection adjacent to dorsomesial distal angle; dorsal surface slightly depressed, with scattered spinules and tufts of short setae; dorsolateral margin with row of spinules and 1 acute spine on proximal third and with moderately long, stiff setae; dorsolateral distal angle produced into small lobe; ventromesial margin subcircular in mesial view; ventral surface inflated, spinulose and with tufts of short, stiff

Fig. 41. *Anapagurus congolensis* spec. nov. ♂ (SL = 2.6 mm). "Mbizi" station 172, 5°15'S, 11°29'E, IRSNB-IG-16.808: mouthparts (right, external view): A, maxillule; B, maxilla; C, 1st maxilliped; D, 2nd maxilliped; E, 3rd maxilliped. Scale equals 1.0 mm.



setae; moderately long ventral length between deep, subtriangular ventrodistal margin and subcircular ventroproximal margin; mesial surface slightly compressed, spinulose, with moderately long, stiff setae; lateral surface somewhat convex, spinulose and with scattered tufts of short setae. Dorsal margin of merus with row of moderately long, stiff setae partly covering 1 strong distal spine; ventrolateral margin with row of spinules and 2 strong distal spines; ventromesial margin usually with 1 strong spine and low spinulose protuberance; ventral surface with numerous spinules and tufts of short, stiff setae. Ischium with 1 spinule on ventrolateral margin partly covered by short, stiff setae; ventromesial margin with row of spinules and moderately long, stiff setae. Ventromesial margin of coxa with tufts of moderately long, stiff setae, partly covering 1 strong spine at distal angle; ventrolateral distal angle unarmed.

Left cheliped reaching distal fifth of palm of right. Dactyl exceeding palm by approximately $\frac{1}{5}$ own length; terminating in acute corneous claw overlapped by claw of fixed finger; cutting edge with row of corneous teeth; all surfaces with scattered spinules and tufts of long, stiff setae. Palm $\frac{1}{2}$ to $\frac{2}{3}$ length of carpus, somewhat inflated ventrally; dorsolateral margin well delimited by closely spaced spinules extending onto fixed finger; dorsal surface with spinules (more numerous on proximal half) and scattered tufts of short setae; a very low, wide, spinulose median ridge proximately; dorsoproximal margin flanked by weakly developed mesial and lateral lobes; dorsomesial margin with row of closely spaced spinules and tufts of moderately long, stiff setae; lateral surface spinulose and with scattered tufts of short setae; mesial surface spinulose and with tufts of moderately long, stiff setae; ventral surface smooth but with tufts of moderately long, stiff setae; fixed finger with scattered spinules on dorsal and lateral surfaces, ventral surface smooth, with scattered tufts of long setae on all faces; cutting edge of fixed finger with row of corneous teeth interspersed with calcareous teeth, terminating in acute, corneous claw. Carpus subequal to length of merus, slightly inflated ventrally; dorsomesial margin with 4 to 6 acute spines interspersed with spinules and tufts of moderately long setae, dorsomesial distal angle produced into subcircular lobe; dorsolateral margin with 7 to 9 strong, acute spines and tufts of long setae, dorsolateral distal angle produced into subcircular lobe; dorsal surface somewhat depressed, glabrous. Dorsal margin of merus with tufts of moderately long, stiff setae; ventrolateral margin with row of spinules on proximal half and 2 widely spaced, strong spines on distal half; ventromesial margin with 1 strong spine; ventral face spinulose and with scattered tufts of moderately long, stiff setae. Ischium with ventromesial margin serrate and row of moderately long, stiff setae. Coxa with ventromesial margin with tufts of moderately long, stiff setae and 1 acute spine at distal angle; ventrolateral distal angle unarmed.

Second and 3rd pereopods overreaching right cheliped. Dactyls exceeding propodi by approximately $\frac{1}{4}$ own length; dorsal margins each with row of long, closely spaced, stiff setae near ventral margin; lateral faces with row of few widely spaced setae near dorsal and ventral margins. Propodi exceeding carpi approximately $\frac{1}{3}$ own length; dorsal surfaces smooth, each with row of closely spaced, short, stiff setae; mesial and lateral faces each with row of few, widely spaced, short setae near ventral margin; ventromesial distal angle with corneous spines, 3 or 4 moderately long (2nd, males), 2 short (2nd, females), 1 very short (3rd, both sexes), and tufts of




Fig. 42. Distribution of *Anapagurus congolensis* spec. nov. Circled star: type locality. Circled numbers: based on previous records, material examined: Forest, 1955 (as *A. laevis*).

short, stiff setae. Carpi approximately $\frac{2}{3}$ length of meri; dorsal surface with row of 5 to 8 spines (2nd) or serrate margin flanked by 1 proximal and 1 distal spines (3rd); lateral, mesial and ventral faces glabrous. Ventrolateral margin of meri serrate (2nd) or unarmed (3rd) and row of tufts of short setae; dorsal surface with 1 short spine on distal half (2nd) or unarmed (3rd) and row of tufts of short, stiff setae. Ischia with tufts of short, stiff setae on dorsal and ventral margins. Lobes of sternite of 2nd pereopods unarmed, distally rounded. Anterior lobe of sternite of 3rd pereopods subrectangular; anterior margin with long setae.

Males with relatively short and broad, left sexual tube recurved upward to level of coxa of 5th pereopod. Coxa of 5th right pereopod with gonopore partly obscured by tufts of short setae. Pleopods 3 and 4 unequally biramous; pleopod 5 uniramous. Distal protopodal lobe of right uropod subequal to length of endopod.

Terminal margins of telson with 4 subequal, closely spaced spines; lateral margins each with 1 spinule and moderately long setae; dorsal surface with scattered short setae.

Colour.— Unknown.

Etymology.— The species name refers to the type locality, off Pointe Noire, Congo, Africa.

Related organisms.— There is no external evidence of symbionts in any of the specimens examined.

Distribution.— All the specimens of *A. congolensis* spec. nov. were collected by the Belgian MBIZI expedition along the tropical western coast of Africa, from off Pointe Noire to Moita Seca. Depth range: 100-240 m.

Affinities.— As previously discussed (see section on affinities for *A. laevis*), *A. congolensis* spec. nov. is closely allied to *A. laevis* and *A. vossi* spec. nov. *Anapagurus congolensis* spec. nov. can be separated from *A. laevis* by: the former's roundly ocular acicles with unexpanded mesial margins; the right cheliped with carpal dorsomesial spines increasing in size distally except for the two most distal smaller spines; the lobes of the sternite of the 2nd pereopods being distally rounded; and in males, the ventromesial distal angle of the propodus of the 2nd pereopods with three or four moderately long spines (see section on affinities for *A. vossi* spec. nov.).

Remarks.— Among the hermit crabs collected during the expeditions of the "Mbizi" and "Mercator" along the southwestern coast of Africa, Forest (1955) illustrated and discussed specimens that he ascribed to *A. laevis*. From my reexamination of this material, I have determined that the specimens from the "Mbizi" expedition represent the new species herein described as *A. congolensis* spec. nov.. The single specimen from the "Mercator" expedition, I have assigned to *A. hendersoni*.

Biogeography

Horizontal distribution.— Most species of *Anapagurus* have an eastern Atlantic distribution and only two have been found elsewhere. Although Briggs (1974) in his book, Marine Zoogeography, placed considerable emphasis on the fishes, the general distributional patterns proposed by him appear to coincide well with those of other marine animals, and especially with those of *Anapagurus*. Following his classification of zoogeographical regions I propose five distributional groupings of species.

(1) Boreal-warm temperate-tropical species. One species, *A. chiroacanthus* (Fig. 7) is well adapted to three different zoogeographical regions of the eastern Atlantic, the boreal or cold temperate, the warm temperate and the tropical. It has the broadest horizontal distribution, occurring between ca 63°N, off the western coast of Norway, and ca 16°N off the Cape Verde Islands. The species has an east-west northern distribution that includes the Azores and extends into the eastern Mediterranean Sea.

(2) Boreal-warm temperate species. Two species of *Anapagurus* are distributed in two distinct regions of the eastern Atlantic, the boreal or cold temperate and the warm temperate: *A. laevis* (Fig. 3) and *A. hyndmanni* (Fig. 5). *Anapagurus laevis* has been collected between ca 63°N, off the western coast of Norway, and ca 20°N, off the coast of Mauritania. In the Mediterranean, *A. laevis* is apparently restricted to the western part. The distribution of *A. hyndmanni* from the British Isles and the Faeroe Islands to the south coast of Spain in the Mediterranean Sea, between ca 62°N and 36°N, appears to be more restricted than that of *A. laevis*.

(3) Warm temperate species. Nine species of *Anapagurus* are known to occur in three well-separated warm temperate regions: a) Mediterranean-Atlantic: *A. adriaticus* spec. nov. (Fig. 33) and *A. petiti* (Fig. 27) are restricted to the Mediterranean, occurring in both the eastern and western sectors; *A. pusillus* (Fig. 9) has been found only outside the Mediterranean Sea in the Azores, Canary Islands and off the south coast of Portugal; *A. bicorniger* (Fig. 15), *A. longispina* (Fig. 17), *A. breviaculeatus* (Fig. 21) and *A. alboranensis* spec. nov. (Fig. 30) are distributed in the Mediterranean, as

well as, in the eastern Atlantic; b) Southern African: *A. hendersoni* (Fig. 23), an apparently South African endemic species has been recorded from Saldanha Bay on the Atlantic west coast to off Cape Natal on the eastern coast in the Indian Ocean. Kensley (1981) summarized several studies of the Plio-Pleistocene molluscan fauna of the western coast of South Africa indicating that a more tropical, warm-water fauna dominated up to the last interglacial in a region which is today considered cold temperate. Apparently *A. hendersoni* is a relict of this tropical fauna that may have survived in well-protected warmer water basins such as Saldanha Bay and False Bay; and c) Japan: *A. japonicus* (Fig. 11), a possibly endemic Japanese species has been collected off the Islands of Honshu, Shikoku and Kyushu. The Uraga Strait, its type locality, is its northernmost record. Ekman (1967) called this region, the "Japanese Mediterranean" because of certain faunal similarities with the Mediterranean.

(4) Warm temperate-tropical species. One species, *A. curvidactylus* (Fig. 13) is broadly distributed in the eastern Atlantic from warm temperate waters off the northwestern coast of Spain to tropical waters of Angola, between ca 43°N and 10°S.

(5) Tropical species. Five species of *Anapagurus* are presumably restricted to two tropical regions widely separated by Africa: a) Eastern Atlantic: four species of *Anapagurus* have similar distributions in tropical African waters: *A. wolffi* (Fig. 25), *A. vossi* spec. nov. (Fig. 36), *A. atlantidii* spec. nov. (Fig. 39) and *A. congolensis* spec. nov. (Fig. 42); and b) Indo-West Pacific: *A. bonnieri* (Fig. 19) that has been collected in three localities: the Red Sea, the Persian Gulf, and off Pulau Saparua, Indonesia.

Vertical distribution (Fig. 43).— Although most species of *Anapagurus* are commonly found on the continental shelf between the intertidal area and about 200 m, some, including *A. longispina*, *A. hendersoni*, *A. adriaticus* spec. nov., *A. congolensis* spec. nov., and *A. laevis*, also occur on the continental slope extending beyond 200 m to depths as great as 1262 m. *Anapagurus laevis* has the broadest bathymetric distribution (4.6 to 1262 m) in the genus *Anapagurus*, and extends to the greatest depth.

Though *A. hyndmanni*, *A. chiroacanthus*, *A. pusillus*, and *A. curvidactylus* have been collected in the intertidal zone; *A. bonnieri* is known to have the shallowest depth range of all recognized species of *Anapagurus*, 5 to 15 m.

Forestopagurus gen. nov.

Anapagurus; Forest, 1966: 162 (in part).

Type species.— *Anapagurus drachi* Forest, 1966. Gender masculine.

Diagnosis.— Carapace with pterygostomial plate somewhat more calcified than remainder of branchiostegite. Posterior carapace also with noticeably calcified plates delimited by cervical groove, lineae anomurica and transversalis, and by cardiobranchialis and cardiac sulci. Anterior margin of shield slightly produced, rounded. Lateral projections with short submarginal spine. Corneae dilated. Ocular acicles ovate, terminating in small marginal spine. Interocular plate unarmed. Antennal peduncles with supernumerary segmentation, acicle well developed. Maxillule with stiff bristle on moderately well-developed internal endopodal lobe; external lobe absent. Third maxilliped with accessory tooth on well-developed crista dentata. Sternite of third maxillipeds convex and without median cleft. Eleven pairs of phyllobranchiae. Chelipeds markedly unequal; dactyls opening in horizontal plane. Anterior lobe of ster-




Fig. 43. Bathymetric distribution of *Anapagurus* species. The horizontal lines indicate the depth range for each species. Depth zones of the ocean floor are after Briggs (1974).




Fig. 44. *Forestopagurus drachi* (Forest, 1966). A-F: ovig. (SL = 1.6 mm), "Calypso" station 94, 1°38'25"N, 7°22'05"E, MNHN 1791. *Anapagurus laevis* (Bell, 1846). G: ♂ (SL = 3.8 mm), off Málaga, Spain, UMDZ. A-E: mouth parts (right, external view): A, maxillule; B, maxilla; C, 1st maxilliped; D, 2nd maxilliped; E, 3rd maxilliped. F, G: sternite of 3rd maxilliped. Scales equal 1.0 mm: a: G; b: F; c: A-E.

nite of third pereopods unarmed. Propodal rasp of fourth pereopods with single row of closely-spaced corneous scales. Sternite of fifth pereopods with two closely-spaced lobes. In males, coxa of right fifth pereopod with gonopore partly covered by tufts of long setae; coxa of left pereopod somewhat enlarged and with moderately long, curved sexual tube; without paired or unpaired pleopods. Females with paired gonopores; three unpaired biramous pleopods, pl₂-pl₄. Uropods asymmetrical. Telson with transverse suture; terminal margin entire, unarmed.

Etymology.— This genus is named for Dr Jacques Forest, a major contributor to our knowledge of hermit crabs, and who, as curator of the Musée National d'Histoire Naturelle (Paris), made the specimens available.

Distribution.— Off Principe Island, Africa; 31 m.

Remarks.— Forest (1966) described *A. drachi* from the Gulf of Guinea, and with "reservation and perhaps provisionally" included the species in the genus *Anapagurus*. From my examinations of the paratypes of *A. drachi* and the extensive collections of *Anapagurus* used in the present revision I have confirmed his doubts. All pleopods are lacking in the male paratype and the fifth left pleopod is wanting in the female paratypes. Other differences that set this species apart from *Anapagurus* include the unarmed telson which lacks a median cleft, the convex anterior margin of the sternite of the third maxillipeds without median cleft, the convex anterior margin of the sternite of the third maxillipeds without median cleft, the moderately long curved left sexual tube, and the noticeably calcified regions of the branchiostegites and the posterior carapace.

Pleopod loss in males is not restricted to *Forestopagurus*. Within the genus *Pagurus*, males of *P. prideaux* Leach, 1815, lack unpaired pleopods. A similar condition is reported for species of *Porcellanopagurus*, *Paguritta*, and *Ostraconotus*. As in *F. drachi*, females of these taxa have their pleopod number reduced to three. The evolutionary mechanism responsible for this loss is unknown.

Acknowledgments

I sincerely wish to express my gratitude to many individuals who have helped in the completion of this study. I am particularly indebted to the late Dr G.L. Voss for his teaching in the field of systematics, as well as his guidance, moral support and friendship throughout my graduate studies. It gives me much pleasure to thank Dr P.A. McLaughlin, a truly great carcinologist, for her generous help, advice, and encouragement throughout my professional career. She introduced me to crustacean systematics, initially suggested the topic of my dissertation, and unselfishly gave me her time, knowledge, and moral support. Many thanks are due to the other members of my dissertation committee, professors N.A. Voss, L.E. Brand, D.K. Odell, and D.R. Moore. Many thanks go to Dr Sandra Gilchrist, New College of the University of South Florida, Sarasota, Florida, for serving as the external examiner at my defense. Their suggestions and comments considerably improved the final result of this study.

I gratefully acknowledge the support of Miami-Dade Community College, Wolfson Campus, InterAmerican Center, Miami, Florida, as the first submitted draft of this manuscript was written during my professional development leave as Associate Professor of Natural Sciences.

Specimens for this study were provided through the efforts of the following individuals representing numerous institutions: Dr R.W. Ingle and Miss S.H. Halsey (BMNH); Dr M. Türkay (SMF); Dr L. Orrhage (NHMG); Drs G. Gianbruno and C. Pesarini (MCSN); Dr A.B. Johnston (MCZ); Drs J. Forest and Y. Coineau (MNHN); Dr C. Carpine (MOM); Ms J. Haig (NIOJ); Drs K. Muzik and K. Mizuki (OMNH); Drs L. B. Holthuis and C.H.J.M. Fransen (RMNH); Dr E. Louw and Miss M.G. van der Merwe (SAM); Dr Å. Andersson (NHRM); Dr J.E. García Raso (UMDZ); Drs R.B. Manning and M. Schotte (USNM); Dr K. Baba (ZLKU); Dr H.E. Gruner (ZMB); Dr T. Wolff (ZMUC); Dr Z. Duris (MMSU); and Dr B. Galil (TAU). Without their help this study would not have been possible.

I am especially thankful to Dr J. Leal who helped me with the identifications of some of the gastropod shells and to Dr J.C. Markham who identified the bopyrid isopods. I am also indebted to Ms O. Mallo, Mr G. Briz, Ms R. Buisman, Mr E. Infante, and Drs A. Mitsui and C.G.H. Rooth for translation services; Mr R. Araujo for his assistance in using the computerized mapping software; MSS L. Alberto, M. Y. Gómez and O.L. Machin for word processing services; Messrs D.J. Heuer and R. Suárez for printing services; and Mr A. Sosa for assisting in the preparation of the finalized draft.

I am most grateful to Mrs K.K. Hale, S. Jeffreys, and V. Hoffman, and Dr H. D. Albertson, of the Library of the Rosenstiel School of Marine and Atmospheric Science University of Miami for their invaluable cooperation in obtaining much of the required literature.

My personal gratitude goes to my mother for her love and moral support always.

Note (added in proof).—In a recent work by Ray Ingle (1993, Hermit Crabs of the Northeastern Atlantic Ocean and Mediterranean Sea: an illustrated key. Chapman & Hall Identification Guide 4: 1-vi, 1-495.—London) the description of *Anapagurus smythii* spec. nov., collected in Adventure Bank, Mediterranean Sea (37°20'N, 12°10'E, 168 m), increased the number of species of *Anapagurus* to 19.

References

- Alcock, A., 1905. Catalogue of the Indian decapod Crustacea in the collection of the Indian Museum. Part II. Anomura. Fasciculus I. Pagurides: i-xi, 1-197.—Indian Museum, Calcutta.
- Anadón, R., 1975. Aportación al conocimiento de la fauna bentónica de la ría de Vigo (NW de España) I. Pycnogónidos y crustáceos de Panjón.—Invest. Pesq. 39 (1): 199-218.
- Aurivillius, C.W.S., 1891. Über Symbiose als Grund accessorischer Bildungen bei marinem Gastropodengehäusen.—Konglia Svenska Vetenskaps-Akademiens Handlingar 24 (9): 1-37.
- Balss, H., 1912. Paguriden. In: Carl Chum (editor). Wissenschaftliche Ergebnisse der Deutschen Tiefsee-Expedition auf dem Dampfer "Valdivia" 1898-1899, 20 (2): 85-124.
- Balss, H., 1921. Crustacea VI: Decapoda Anomura (Paguridae) und Brachyura (Dromiacea bis Brachynatha I.) In: W. Michaelsen (editor), Beiträge zur Kenntnis der Meeresfauna Westafrikas 3: 37-67.—Friederichsen & Co., Hamburg.
- Balss, H., 1924. Über Anpassungen und Symbiose der Paguriden. Eine zusammenfassende Übersicht.—Zeitschrift fuer Morphologie und Ökologie de Tiere 1: 752-792.
- Balss, H., 1926. Ueber einige Decapoden Des Golfs von Neapel (Fam. Crangonidae, Paguridae).—Pubbl. Stn. Zool. Napoli 7 (3) 10: 375-381.
- Balss, H., 1944a. Decapoda. III. Morphologischer Teil. In: H.G. Bronns, Klassen und Ordnungen des Tierreichs.—Akademische Verlagsgesellschaft Becker & Erler Kom.-Ges., Leipzig (5) 1 (7) 3: 321-480.

- Balss, H., 1944b. Decapoda. III. Morphologischer Teil. In: H.G. Bronns, Klassen und Ordnungen des Tierreichs.— Akademische Verlagsgesellschaft Becker & Erler Kom.- Ges., Leipzig (5) 1 (7) 5: 577-669.
- Balss, H., 1956. Decapoda. VI. Ökologie. In: H.G. Bronns, Klassen und Ordnungen des Tierreichs.— Akademische Verlagsgesellschaft Geest & Portig K.- Ges., Leipzig (5) 1 (7) 11: 1369-1504.
- Balss, H., 1957. Decapoda. IX. Geographische Verbreitung. In: H.G. Bronns, Klassen und Ordnungen des Tierreichs.— Akademische Verlagsgesellschaft Geest & Portig K.- Ges., Leipzig (5) 1 (7) 13: 1673-1770.
- Barnard, K.H., 1947. Descriptions of new species of South African Decapod Crustacea, with notes on synonymy and new records.— Ann. Mag. Nat. Hist. (11th series) 13 (102) XLI: 361-392.
- Barnard, K.H., 1950. Descriptive Catalogue of South African Decapod Crustacea (Crabs and Shrimps).— Ann. S. Afr. Mus. 38: 1-837.
- Basford, D.J., A. Eleftheriou & D. Raffaelli, 1989. The epifauna of the northern North Sea (56°-61°N).— J. Mar. Biol. Assoc. U. K. 69 (2): 387-407.
- Bell, T., 1844-53. A History of the British stalk-eyed Crustacea: i-LXV, 1-386.— John Van Voorst, London (Issued separately in parts: pp. 144-260, 1846).
- Bourdon, R., 1960. Rhizocéphales et Isopodes parasites des Décapodes Marcheurs de la Baie de Quiberon.— Bull. Soc. Sci. Nancy 19 (2/3): 134-153.
- Bourdon, R., 1968. Les Bopyridae des mers Européennes.— Mém. Mus. Natl. Hist. Nat. Paris, nouvelle série A, Zoologie, 50 (2): 77-424.
- Bouvier, E.L., 1891. Étude de quelques Paguriens recueillis par M. Jules de Guerne sur les côtes de France et de Norvège.— Mém. Soc. Zool. Fr. pour l'année 1891, 4: 393-407.
- Bouvier, E.L., 1896. Les Pagurinés des mers d'Europe (Crustacés). Tableaux dichotomiques des genres et des espèces (Fin).— La Feuille des Jeunes Naturalistes, 3e série, 26e année, 308 et 309: 149-155.
- Bouvier, E.L., 1905. Sur les Crustacés Décapodes (abstraction faite des Carides) recueillis par le yacht "Princesse-Alice" au cours de la campagne de 1905.— Bull. Mus. Océanogr. Monaco 55: 1-4.
- Bouvier, E.L., 1922. Observations complémentaires sur les Crustacés Décapodes (Abstraction faite des Carides) provenant de campagnes de S.A.S le prince de Monaco.— Résult. Camp. Sci. Monaco 62: 1-106.
- Bouvier, E.L., 1940. Décapodes marcheurs.— Faune de France, 37: 1-404.
- Brandt, F., 1851. Krebse. In: A.T. von Middendorff, Reise in den Äussersten Norden und Osten Sibiriens während der Jahre 1843 und 1844, 2 (1) (Zoologie): 77-148.
- Briggs, J.C., 1974. Marine zoogeography: i-xiv, 1-475.— McGraw-Hill Book Company, New York.
- Buddenbrock, W. von, 1948. Decapoda. V. Physiologie. In: H. G. Bronns, Klassen und Ordnungen des Tierreichs.— Akademische Verlagsgesellschaft Geest & Portig K.-Ges., Leipzig (5) 1 (7) 8: 1007-1150.
- Calafiore, N., 1981. Le larvae dei Crostacei Decapodi (escluso Brachiuri) dello stretto di Messina e dei mari limitrofi.— Mem. Biol. Mar. Oceanogr. 11 (1): 17-36.
- Cardona Bendito, A., 1965. Decápodos marinos de la ría de Vigo.— Invest. Pesq. 28: 133-159.
- Carlgren, O., 1945. Polypdyr (Coelenterata). III. Koraldyr.— Danmarks Fauna 51: 1-168.
- Carrington, J.T. & E. Lovett, 1882. Notes and observations on British stalkeyed Crustacea.— The Zoolologist, serie 3, 6: 386-391.
- Chevreux, E., & E.L. Bouvier, 1892a. Voyage de la goëlette "Melita" aux Canaries et au Sénégal. Note préliminaire sur les Paguriens.— Bull. Soc. Zool. Fr. 16 (5): 252-256.
- Chevreux, E., & E.L. Bouvier, 1892b. Voyage de la goëlette "Melita" aux Canaries et au Sénégal, 1889-1890. Paguriens.— Mém. Soc. Zool. Fr. pour l'année 1892, 5: 83-144.
- Cranmer, G.J., M.F. Dyer & P.D. Fry, 1984. Further results from headline camera surveys in the North Sea.— J. Mar. Biol. Assoc. U. K. 64 (2): 335-342.
- Crothers, J. & M. Crothers, 1983. A key to the crabs and crab-like animals of British inshore waters.— Field Stud. 5 (5): 753-806.
- Dauvin, J.-C. & M. Joncourt, 1989. Energy values of marine benthic invertebrates from the western English Channel.— J. Mar. Biol. Assoc. U. K. 69 (3): 589-595.
- Day, J.H., J.G. Field & M.J. Penrith, 1970. The benthic fauna and fishes of False Bay, South Africa.— Trans. R. Soc. S. Afr. 39 (1): 1-108.
- Dechancé, M., 1962. Description des stades larvaires et de la Glaucothoé de *Spiropagurus elegans* Miers.— Bull. Mus. Natl. Hist. nat. Paris, 2e série, 34 (5): 371-386.

- Dechancé, M. & J.P. Dufaure, 1959. Une nouvelle association entre une Actinie et un Pagure.— C. R. Hebd. Séances Acad. Sci. Paris 249: 1566-1568.
- Dechancé, M. & J. Forest, 1962. Sur *Anapagurus bicorniger* A. Milne Edwards et E.L. Bouvier, et *A. petitii* sp. nov. (Crustacea Decapoda Paguridae).— Bull. Mus. Natl. Hist. nat. Paris, 2e série, 34: 293-307.
- De Miranda y Rivera, A., 1933. Ensayo de un catálogo de los crustáceos decápodos marinos de España y Marruecos Español.— Publ. Inst. Esp. Oceanogr., Notas y Resúmenes, serie 2, 67: 1-72.
- Dollfus, R.P. & J. Carayon, 1942. *Pseudione hyndmani* (Spence Bate et J.O. Westwood) chez *Anapagurus laevis* V. Thompson. Discussion de la spécificité parasitaire de ce *Pseudione* [Isop. Epicaridea].— Bull. Mus. Natl. Hist. nat. Paris (2) 14: 445-450.
- Doumenc, D., C. Chintiroglou & A. Koukouras, 1985. Actinies de la mer Égée: méthodes d'identification, zoogéographie.— Bull. Mus. Natl. Hist. nat. Paris, 4e série, 7, section A: Zoologie, 3: 497-529.
- Ekman, S., 1967. Zoogeography of the Sea: i-xiv, 1-417.— Sidgwick & Jackson Ltd., London.
- Elmhirst, R., 1912. Report of the Superintendent to the General Committee for 1912: List of recent additions to the Local Marine Fauna: Isopoda.— Annu. Rep. Mar. Biol. Assoc. West of Scotland: 23.
- Fenizia, G., 1937. I Paguridi del golfo di Napoli. 2^o Contributo. Specie e varietà dei generi *Catapaguroides* M. Edwards e Bouvier, *Anapagurus* Henderson e *Nematopagurus* M. Edwards e Bouvier.— Ann. Mus. Zool. R. Univ. Napoli (Nuova Serie) 7 (2): 1-40.
- Forest, J., 1955. Crustacés décapodes, Pagurides. Expédition océanographique belge dans les eaux côtières africaines de l'Atlantic sud (1948-1949).— Résult. sci. 3 (4): 21-147.
- Forest, J., 1958. Sur la nomenclature des Pagures des mers francaises.— Bull. Mus. Natl. Hist. nat. Paris, 2e série, 30 (1): 94-100.
- Forest, J., 1961. Pagurides de l'Afrique occidentale.— Atl. Rep. 6: 203-250.
- Forest, J., 1965. Campagnes du "Professeur Lacaze-Duthiers" aux Baléares: juin 1953 et aout 1954. Crustacés Décapodes.— Vie Milieu, Sér. B, 16 (1-B): 325-414.
- Forest, J., 1966. Résultats scientifiques des campagnes de la "Calypso", 7 (26)- Campagne dans le golfe de Guinée et aux les Principe, São Tomé et Annobon (1956) (suite). 17. Crustacés Décapodes; Pagurides.— Ann. Inst. Océanogr., nouvelle série, 44: 125-172.
- Forest, J., 1967. Sur une collection des Crustacés Décapodes de la région de Porto Cesareo. Description de *Portumnus pestai* sp. nov.— Thalassia Salentina 2: 1-29.
- Forest, J. & M. de Saint Laurent, 1967. Campagne de la "Calypso" au large des côtes Atlantiques de l'Amérique du Sud (1961-1962). 6. Crustacés Décapodes: Pagurides.— Ann. Inst. Oceanogr. Monaco 45 (2): 47-169.
- Füller, H., 1958. Symbiose im tierreich: 1-227.— A. Ziems Verlag, Wittenberg-Lutherstadt.
- García-Gómez, J., 1988. A new genus and three new species of hermit crabs (Crustacea: Decapoda: Paguridae) from the western Atlantic ocean.— Bull. Mar. Sci. 42 (1): 44-64.
- García Raso, J.E., 1982. Contribución al estudio de los Pagurídos (Crustacea, Decapoda, Anomura) en el litoral sudmediterráneo español.— Invest. Pesq. 46 (3): 493-508.
- García Raso, J.E., 1984. Primeras aportaciones al conocimiento de la fauna de Crustáceos Decápodos litorales de la Isla de Alborán (España).— Bol. Asoc. Esp. Entomol. 8: 253-263.
- García Raso, J.E., 1985. Nuevas aportaciones a la fauna de crustáceos decápodos de la isla de Alborán (España).— Bol. Soc. Portug. Entomol., Suplemento No 1: 11-18.
- Gilat, E., 1969. Study of an ecosystem in the coastal waters of the Ligurian Sea. III. Macrobenthic communities.— Bull. Inst. Océanogr. Monaco 69 (1396): 1-76.
- Gilpin-Brown, J.B., 1969. Host-adoption in the commensal polychaete *Nereis fucata*.— J. Mar. Biol. Assoc. U. K. 49: (1) 121-127.
- Gordan, J., 1958. A bibliography of pagurid crabs, exclusive of Alcock, 1905.— Bull. Am. Mus. Nat. Hist. 108 (3): 253-352.
- Guille, A., 1971. Bionomie benthique du plateau continental de la côte catalane française. IV. Densités, biomasses et variations saisonnière de la macrofaune.— Vie Milieu 22 (1B): 93-157.
- Gunn, S.W. & P.A. McLaughlin, 1988. The rediscovery of *Pagurus acantholepis* (Stimpson) (Decapoda: Anomura: Paguridae).— Mem. Mus. Victoria 49 (1): 67-71.
- Gurney, R., 1939. Bibliography of the larvae of decapod Crustacea: i-viii, 1-123.— Ray Society, London.
- Gurney, R., 1942. Larvae of decapod Crustacea: i-viii, 1-306.— Ray Society, London.
- Haddon, A.C. & A.M. Shackleton, 1891. A revision of the British Actinia. Part II: the Zoolantheae.— Sci. Trans. Royal Dublin Soc., serie 2, 4 (12): 609-672.

- Haig, J. & E.E. Ball, 1988. Hermit crabs from North Australian and Eastern Indonesian waters (Crustacea Decapoda: Anomura: Paguroidea) collected during the 1975 "Alpha Helix" Expedition.— Rec. Aust. Mus. 40: 151-196.
- Hansen, H.J., 1908. Crustacea Malacostraca. I.— The Danish Ingolf-Expedition 3 (2): 1-120.
- Hazlett, B.A., 1968a. Dislodging behavior in European pagurids.— Pubbl. Stn. Zool. Napoli 36 (2): 138-139.
- Hazlett, B.A., 1968b. The sexual behavior of some European hermit crabs (Anomura: Paguridae).— Pubbl. Stn. Zool. Napoli 36 (2): 238-252.
- Hazlett, B.A., 1981. The behavioral ecology of hermit crabs.— Ann. Rev. Ecol. Syst. 12: 1-22.
- Henderson, J.R., 1886a. Higher Crustacea of the Firth of Clyde. XXXIV. The Decapod and Schizopod Crustacea of the Firth of Clyde.— Proceedings and Transactions of the Natural History of Glasgow, new series 1: 315-353.
- Henderson, J.R., 1886b. A Synopsis of the British Paguridae.— Proc. R. Phys. Soc. Edinburgh 9 (4): 65-75.
- Henderson, J.R., 1888. Report on the Anomura collected by H.M.S. "Challenger" during the years 1873-1876.— Report on the scientific results of the voyage of H.M.S. "Challenger" during the years 1873-76. Zoology 27 (1): i-xi, 1-221.
- Høeg, J.T., & J. Lützen, 1985. Crustacea Rhizocephala.- Marine Invertebrates of Scandinavia 6: 1-92.— Norwegian University Press, Oslo, Norway.
- Holthuis, L.B., 1950. Decapoda (K IX) A. Natantia, Macrura Reptantia, Anomura en Stomatopoda (K X). In: H. Boschma, Fauna van Nederland 15: 1-166.
- Holthuis, L.B., 1962. Forty-seven genera of Decapoda (Crustacea); Proposed addition to the Official List. Z.N. (S.) 1499.— Bull. Zool. Nomenclature 19 (4): 232-253.
- Holthuis, L.B. & E. Gottlieb, 1958. An annotated list of the decapod Crustacea of the Mediterranean coast of Israel, with an appendix listing the Decapoda of the eastern Mediterranean.— Bull. Res. Counc. Isr. Sect. B: Zool. 7B (1-2): 1-126.
- Ingle, R.W., 1985. Northeastern Atlantic and Mediterranean hermit crabs (Crustacea: Anomura: Paguroidea: Paguridae). I. The genus *Pagurus* Fabricius, 1775.— J. Nat. Hist. 19 (4): 745-769.
- Ingle, R.W., 1990. Larval and post-larval development of *Anapagurus chiroacanthus* (Lilljeborg, 1856) Anomura: Paguroidea: Paguridae.— Bull. Br. Mus. (Nat. Hist.). Zool. 56 (2): 105-134.
- Jensen, K. & K. Bender, 1973. Invertebrates associated with snail shells inhabited by *Pagurus bernhardus* (L.) (Decapoda).— Ophelia 10 (2): 185-192.
- Kensley, B., 1974. Type specimens of Decapoda (Crustacea) in the collections of the South African Museum.— Ann. S. Afr. Mus. 66 (4): 55-80.
- Kensley, B., 1978. Decapod crustaceans collected in southern African waters by the Th. Mortensen Java-South Africa Expedition (Crustacea, Decapoda).— Steenstrupia 4 (21): 249-261.
- Kensley, B., 1981. On the zoogeography of Southern African decapod Crustacea, with a distributional checklist of the species.— Smithson. Contrib. Zool. 338: i-iv, 1-64.
- Kocatas, A., 1981. Liste préliminaire et répartition des crustacés décapodes des eaux turques.— Rapp. Comm. Int. Explor. Sci. Mer Méditerr. 27 (2): 161-162.
- Lagerberg, T., 1908. Sveriges Decapoder.— Goteborgs Kungl. Vetenskapsoch Vitterhets-samhallet Handlingar. Tjarde Foliden. Serien B, serie 4, 11 (2): i-x, 1-117.
- Lemaitre, R., P.A. McLaughlin, & J. García-Gómez, 1982. The Provenzanoi group of hermit crabs (Crustacea, Decapoda, Paguridae) in the western Atlantic.— Bull. Mar. Sci. 32 (3): 670-701.
- Leviton, A.E., R.H. Gibbs, Jr., E. Heal & C.E. Dawson, 1985. Standards in herpetology and ichthyology: Part I. Standard symbolic codes for institutional resource collections in herpetology and ichthyology.— Copeia 1985 (3): 802-832.
- Leviton, A.E. & R.H. Gibbs, Jr., 1988. Standards in herpetology and ichthyology. Standard symbolic codes for institutional resource collections in herpetology and ichthyology. Supplement No. 1: Additions and corrections.— Copeia 1988 (1): 280-282.
- Lewinsohn, C., 1969. Die Anomuren des Roten Meeres (Crustacea Decapoda: Paguridea, Galatheidea, Hippidea). The Second Israel South Red Sea Expedition, 1965. Report No. 6.— Zool. Verh. Leiden 104: 1-213.
- Lewinsohn, C., 1976. Crustacea Decapoda von der Insel Rhodos, Griechenland.— Zool. Meded. Leiden 49 (17): 237-254.

- Lilljeborg, W., 1856. Om Hafs-Crustaceer vid Kullabergi Skåne.— Ofversigt af Kongl. Vetenskapsakademiens Forhandlingar 12 (3): 117-138.
- Lindley, J.A., 1986. Vertical distributions of decapod crustacean larvae and pelagic post-larvae over Great Sole Bank (Celtic Sea) in June 1983.— Mar. Biol. 90 (4): 545-549.
- MacDonald, J.D., R.B. Pike, & D.I. Williamson, 1957. Larvae of the British species of *Diogenes*, *Pagurus*, *Anapagurus* and *Lithodes* (Crustacea, Decapoda).— Proc. Zool. Soc. Lond. 128 (2): 209-257.
- Manuel, R.L., 1981. British Anthozoa. In: D.M. Kermack & R.S.K. Barnes (editors), *Synopses of the British Fauna (New Series)* 18: 1-241.— Academic Press, London.
- Markham, J.C., 1986. Evolution and zoogeography of the Isopoda Bopyridae, parasites of Crustacea Decapoda. In: F.R. Schram (General Editor), *Crustacean Issues* 4, R.H. Gore & K.L Heck (Editors), *Crustacean Biogeography*, 2. *Dispersal*: 143-164.
- Matallanas, J., 1982. Aspectos generales del regimen alimentario de *Macroramphosus scolopax* (Linnaeus 1758) (Pisces, Macroramphosidae) en las costas catalanas. (Mediterráneo occidental).— Cah. Biol. Mar. 23 (3): 243-252.
- Matthews, D.C., 1953. The development of the pedunculate spermatophore of a hermit crab, *Dardanus asper* (De Haan).— Pac. Sci. 7 (3): 255-266.
- McLaughlin, P.A., 1974. The hermit crabs (Crustacea Decapoda, Paguridea) of northwestern North America.— Zool Verh. Leiden 130: 1-396.
- McLaughlin, P.A., 1980. Comparative morphology of recent Crustacea: i-xiv, 1-177.— W.H. Freeman & Company, San Francisco.
- McLaughlin, P.A., 1981a. Revision of *Pylopagurus* and *Tomopagurus* (Crustacea: Decapoda: Paguridae), with the descriptions of new genera and species. Part I. Ten new genera of the Paguridae and a redescription of *Tomopagurus* A. Milne Edwards & Bouvier.— Bull. Mar. Sci. 31 (1): 1-30.
- McLaughlin, P.A., 1981b. Revision of *Pylopagurus* and *Tomopagurus* (Crustacea: Decapoda: Paguridae), with the descriptions of new genera and species: Part II. *Rhodochirus* McLaughlin and *Phimochirus* McLaughlin.— Bull. Mar. Sci. 31 (2): 329-365.
- McLaughlin, P.A., 1986. Three new genera and species of hermit crabs (Crustacea: Anomura: Paguridae) from Hawaii.— J. Crust. Biol. 6 (4): 789-803.
- Milne Edwards, A. & E.L. Bouvier, 1892. Observations préliminaires, sur les Paguriens récueillis par les expéditions du "Travailleur" et du "Talisman".— Ann. Sci. Nat. Zool., ser. 7 (13): 185-226.
- Milne Edwards, A. & E.L. Bouvier, 1893. Reports on the results of dredging, under the supervision of Alexander Agassiz, in the Gulf of Mexico (1877-78), in the Caribbean Sea (1878-79), and along the Atlantic coast of the United States (1880), by the U.S. Coast Survey Steamer "Blake" Lieut.- Com. S.D. Sigsbee, U.S.N. and Commander J.R. Bartlett, U.S.N., Commanding 33.— Mem. Mus. Comp. Zool. Harvard College 14(3): 1-172.
- Milne Edwards, A. & E.L. Bouvier, 1894. Crustacés décapodes provenant des campagnes du yacht l'"Hirondelle" (1886, 1887, 1888). Première partie. Brachyures et Anomoures.— Résult. Camp. Sci. Monaco 7: 1-112.
- Milne Edwards, A. & E.L. Bouvier, 1898. A propos des Crustacés brachyures et anomoures provenant des six dernières campagnes scientifiques effectuées par S.A. le Prince de Monaco.— C. R. Hebd. Séances Acad. Sci., zool. cxxxvi: 1245-1247.
- Milne Edwards, A. & E.L. Bouvier, 1899. Crustacés décapodes provenant des campagnes de l'"Hirondelle" (Supplement) et de la "Princesse Alice" (1891-1897). Brachyures et Anomoures.— Résult. Camp. Sci. Monaco 13: 1-106.
- Milne Edwards, A. & E.L. Bouvier, 1900. Crustacés Décapodes, I. Brachyures et Anomoures. Expéditions scientifiques du "Travailleur" et du "Talisman" pendant les années 1880, 1881, 1882, 1883: 1-396.
- Miyake, S., 1961. Fauna and Flora of the Sea around the Amakusa Marine Biological Laboratory. Part II. Decapod Crustacea.— Amakusa Mar. Biol. Lab. Contrib. 139: i-iv, 1-30 (in Japanese).
- Miyake, S., 1965. Anomura. In: New illustrated encyclopedia of the fauna of Japan. Hokuryukan, Tokyo, 2: 630-652 (in Japanese).
- Miyake, S., 1978. The Crustacean Anomura of Sagami Bay: i-ix, 1-200.— Biological Laboratory Imperial Household, Japan (in English). Japanese text and index, 161 pp.

- Miyake, S., K. Sakai & S. Nishikawa, 1962. A fauna-list of the decapod Crustacea from the coasts washed by the Tsushima warm current.— Rec. Oceanogr. Works Jpn. (Special No. 6): 121-131.
- Monod, T., 1932. Sur quelques Crustacés de l'Afrique Occidentale (Liste des Décapodes mauritaniens et des Xanthidés ouest-africains).— Bull. Com. Etud. Hist. Sci. Afr. Occidentale Française, 25 (2, 3): 456-548.
- Morgan, G.J. & J. Forest, 1991. A new genus and species of hermit crab (Crustacea, Anomura, Diogenidae) from the Timor Sea, north Australia.— Bull. Mus. Natl. Hist. nat. Paris, 4e série, 13 (1, 2): 189-202.
- Mouchet, S., 1931. Spermatophores des Crustacés Décapodes Anomoures et Brachyoures et castration parasitaire chez quelques Pagures.— Ann. Stn. Océanogr. Salammbô 6: 1-203.
- Muirhead, A., P.A. Tyler, & M.H. Thurston, 1986. Reproductive biology and growth of the genus *Epi-zoanthus* (Zoanthidea) from the north-east Atlantic.— J. Mar. Biol. Assoc. U. K. 66 (1): 131-143.
- Naylor, E., 1972. British marine isopods. Keys and notes for the identification of the species. The Linnean Society of London, Academic Press, A new series. Synopses of the British Fauna, 3: 1-86.
- Neves, A.M., 1977. Crustáceos decápodes marinhos de Portugal continental existentes no Museu Bocage. III. Anomura.— Arq. Mus. Bocage, 2a série, 6 (10): 153-206.
- Nielsen, S.-O., 1970. The effects of the rhizocephalan parasites *Peltogaster paguri* Rathke and *Gemmosacus sulcatus* (Lilljeborg) on five species of paguridan hosts (Crustacea Decapoda).— Sarsia, 42: 17-32.
- Nilsson-Cantell, C.A., 1926. Über Veränderungen der sekundären Geschlechtsmerkmale bei Paguriden durch die Einwirkung von Rhizocephalen.— Ark. Zool. 18A (13): 1-21.
- Nobili, G., 1905. Quatre Décapodes nouveaux du Golfe Persique (récoltes de MM. J. Bonnier et Ch. Péres).— Bull. Mus. Natl. Hist. nat. Paris 11 (2): 238-239.
- Nobili, G., 1906a. Diagnoses préliminaires de Crustacés, Décapodes et Isopodes nouveaux recueillis par M. le Dr. G. Seurat aux îles Touamotou.— Bull. Mus. Natl. Hist. nat. Paris 12: 256-270.
- Nobili, G., 1906b. Crustacés Décapodes et Stomatopodes. Mission J. Bonnier et Ch. Péres (Golfe Persique, 1901).— Bull. Sci. Fr. Belg. 40: 13-159.
- Nobre, A., 1931. Crustáceos decápodes e stomatópodes Marinhos de Portugal: i-iv, 1-307.— Imprensa Portuguesa, Pôrto.
- Nobre, A., 1936. Crustáceos decápodes e stomatópodes marinhos de Portugal, 2a edção.— Fauna Mar. Port. 4: i-viii, 1-213.
- Norman, A.M., 1861. Contributions to British Carcinology. I. Characters of underdescribed Podophthalmia and Entomostraca.— Ann. Mag. Nat. 3rd series, 8 (46) 29: 273-281.
- Norman, A.M., 1886. Museum Normanianum, or A Catalogue of the Invertebrata of Europe, and the Arctic and North Atlantic Oceans, which are contained in the collection of the Rev. Canon A.M. Norman, M.A., D.C.L., F.L.S. III Crustacea: 1-26.— Morton, Printer, Houghton-Le-Spring.
- Norman, A.M., 1905. Museum Normanianum, or A Catalogue of the Invertebrata of the Arctic and North Atlantic Temperate Ocean and Palaearctic Region, which are contained in the collection of the Rev. Canon A.M. Norman, M.A., D.C.L., LL.D., F.R.S., F.L.S., & C. III. Crustacea. Second Edition: 1-47.— Thos. Cadcleugh & Son, Durham.
- Norman, A.M., & G.S. Brady, 1911. The Crustacea of Northumberland and Durham.— Transactions of the Natural History Society of Northumberland, Durham and Newcastle-upon-Tyne, new serie, 3: 252-417.
- Odhner, T., 1923. Marine Crustacea Podophthalmata aus Angola und Südafrika.— Göteborgs Kungl. Vetenskaps-och Vitterhets-Samhälles, Ser. 4, 27(5): 1-39 (Meddelanden från Göteborgs Musei zoologiska avdelning 31).
- Ortmann, A., 1892. Die Decapoden-Krebse des Strassburger Museums, 4. Die Abtheilungen Galatheidea und Paguridea.— Zool. Jahrb. 2: 241-326.
- Pastore, M.A., 1972. Decapoda Crustacea in the Gulf of Taranto and the Gulf of Catania with a discussion of a new species of Dromidae (Decapoda Brachyoura) in the Mediterranean Sea.— Thalassia Jugosl. 8 (1): 105-117.
- Patience, A., 1911. Bopyridae. In: L.A.L. King, Clyde Marine Fauna. Supplementary List.— Mar. Biol. Assoc. West Scotl. Annual Report, 1911: 74.
- Péres, J.M., 1985. History of the Mediterranean biota and the colonization of the depths. In: R. Margalef (editor), Key environments, Western Mediterranean, 7: 1232.— Pergamon Press, Oxford.

- Pérès, L.M. & J. Picard, 1964. Nouveau Manuel de Bionomie Benthique de la Mer Méditerranée.— Rec. Trav. Stn. Mar. Endoume 31 (47): 1-137.
- Pérez, C. 1930. Zoologie-Asymétrie viscérale et dimorphisme des spermatophores chez quelques Pagures.— C. R. des séances Acad. Sci. Zool. 190: 1-3.
- Pike, R.B., 1953. The Bopyrid parasites of the Anomura from British and Irish waters.— J. Linn. Soc. Lond. Zool. 42 (285): 219-237.
- Pike, R.B., 1961. Observations of Epicaridea obtained from hermit crabs in British waters, with notes on the longevity of the host-species.— Ann. Mag. Nat. Hist. Series 13, 4 (40): 225-240.
- Pike, R.B. & D.I. Williamson, 1958. Crustacea Decapoda: Larvae XI. Paguridea, Coenobitidea, Dromiidea, and Homolidea.— Cons. Int. Explor. Mer. Zooplankton sheet 81: 1-9.
- Pike, R.B. & D.I. Williamson, 1959. Observations on the distribution and breeding of British hermit crabs and the stone crab (Crustacea: Diogenidae, Paguridae and Lithodidae).— Proc. Zool. Soc. Lond. 132 (4): 551-567.
- Pike, R.B. & D.I. Williamson, 1960. Larvae of decapod Crustacea of the families Diogenidae and Paguridae from the Bay of Naples.— Pubbl. Stn. Zool. Napoli 31 (3): 493-552.
- Pilgrim, R.L.C., 1973. Axial skeleton and musculature in the thorax of the hermit crab, *Pagurus bernhardus* [Anomura: Paguridae].— J. Mar. Biol. Assoc. U. K. 53: 363-396.
- Rathbun, M.J., 1900. The Decapod crustaceans of West Africa.— U.S. Natl. Mus. Proc. XXII (1199): 271-316.
- Reinhard, E.G., 1942. Studies on the life history and host-parasite relationship of *Peltogaster paguri*.— Biol. Bull. (Woods Hole) 83 (3): 401-415.
- Riedl, R., 1986. Fauna y Flora del Mar Mediterráneo: 1-858.— Ediciones Omega, Barcelona.
- Ros, J.D., J. Romero, E. Ballesteros & J.M. Gili, 1985. Diving in Blue Water. The Benthos. In: R. Margalef (Editor), Key Environments, Western Mediterranean 8: 233-295.— Pergamon Press, Oxford.
- Ross, D.M., 1967. Behavioural and ecological relationships between sea anemones and other invertebrates.— Oceanogr. Mar. Biol. Annu. Rev. 5: 291-316.
- Saint Laurent, M. de, 1968. Révision des genres *Catapaguroides* et *Cestopagurus*, et description de quatre genres nouveaux. I. *Catapaguroides* A. Milne Edwards et Bouvier et *Decaphyllus* nov. gen. (Crustacés Décapodes Paguridae).— Bull. Mus. Natl. Hist. nat. Paris, 2e série, 39 (5): 923-954.
- Saint Laurent, M. de, 1969. Révision des genres *Catapaguroides* et *Cestopagurus* et description de quatre genres nouveaux. III. *Acanthopagurus* de Saint Laurent (Crustacés Décapodes Paguridae).— Bull. Mus. Natl. Hist. nat. Paris, 2e série, 41 (3): 731-742.
- Saint Laurent-Dechancé, M. de, 1966a. *Iridopagurus*, genre nouveau de Paguridae (Crustacés Décapodes) des mers tropicales américaines.— Bull. Mus. Natl. Hist. nat. Paris, 2e série, 38 (3): 151-173.
- Saint Laurent-Dechancé, M. de, 1966b. Remarques sur la classification de la famille des Paguridae et sur la position systématique d'*Iridopagurus* de Saint Laurent. Diagnose d'*Anapagrides* gen. nov.— Bull. Mus. Natl. Hist. nat. Paris, 2e série, 38 (3): 257-265.
- Salmon, M., 1983. Courtship, mating systems and sexual selection in decapods. In: S. Rebach & D.W. Dunham (editors), Studies in Adaptation. The Behavior of higher Crustacea 4 (6): 143-169.— John Wiley & Sons, New York.
- Samuelson, T.J., 1970. The biology of six species of Anomura (Crustacea, Decapoda) from Raunefjorden, western Norway.— Sarsia 45: 25-52.
- Sars, G.O., 1890. Bidrag til kundskaben om Decapodernes Forvandlinger. II. *Lithodes-Eupagurus-Spiropagurus-Galathodes-Galathea-Munida-Porcellana-(Nephrops)*.— Archiv for Mathematik og Naturvidenskab 13: 133-201.
- Sars, G.O., 1899. An account of the Crustacea of Norway with short descriptions and figures of all the species. II. Isopoda: i-x, 1-270.— The Bergen Museum, Norway.
- Sarojini, R. & R. Nagabhushanam, 1969. The occurrence of the hermit crab, *Anapagurus laevis* (Thompson) in India.— J. Mar. Biol. Assoc. of India 10 (1): 387, 388.
- Schembri, P.J., 1982. Feeding behaviour of fifteen species of hermit crabs (Crustacea: Decapoda: Anomura) from the Otago region, Southeastern New Zealand.— J. Nat. Hist. 16: 859-878.
- Scott, T., 1888. A revised list of the Crustacea of the Firth of Forth.— Ann. Rep. Fish. Board Scotl. (3) 6: 253-262.

- Scott, T., 1897. Report on a Collection of Marine Dredgings and other Natural History Materials made on the West Coast of Scotland by the late George Brook, F.L.S.— Proc. R. Phys. Soc. Edinburgh 13: 166-193, pl. V.
- Selbie, C.M., 1921. The Decapoda Reptantia of the coasts of Ireland. Part II. Paguridea.— Fisheries, Ireland, Sci. Invest. 1921 (1): 1-68.
- Shiber, J.G. & L. Fattah, 1978. Hermit crabs from the coast of Ras Beirut, Lebanon.— Cah. Biol. Mar. 19 (2): 127-134.
- Sinel, J., 1907. A contribution to our knowledge of the Crustacea of the Channel Islands.— Trans. Guernsey Soc. Nat. Sci. Local Res. 1906, 5: 212-225.
- Snodgrass, R.E., 1951. Comparative studies on the head of mandibulate arthropods: i-x, 1-118.— Comstock Publishing Company, Inc., New York.
- Stachowitsch, M., 1980. The epibiotic and endolithic species associated with the gastropod shells inhabited by the hermit crabs *Paguristes oculatus* and *Pagurus cuanensis*.— Pubbl. Stn. Zool. Napoli I, Mar. Ecol. 1 (1): 73-101.
- Stebbing, T.R.R., 1910. General Catalogue of South African Crustacea (Part V. of S.A. Crustacea, for the Marine Investigations in South Africa).— Ann. S. Afr. Mus. 6 (4) 6: 281-599.
- Stevicic, Z., 1972. Révision et complément de la liste inventaire des Crustacés Décapodes adriatiques.— Thalassia Jugosl. 8 (1): 101-104.
- Stimpson, W., 1858. Prodromus descriptionis animalium evertebratorum, quae in Expeditione ad Oceanum Pacificum Septentrionalem, a Republica Federata missa, Cadwaladaro Ringgold et Johanne Rodgers Ducibus, observavit et descripsit. Pars VII. Crustacea Anomoura. I. Teleosomi. Reprint (December 1858) from: Proc. Acad. Nat. Sci. Phila. 1858: 225-252.
- Stimpson, W., 1907. Report on the Crustacea (Brachyura and Anomura) collected by the North Pacific Exploring Expedition, 1853-1856.— Smithson. Misc. Collect. 49 (1717): 1-240.
- Tattersall, W.M., 1905. The Marine Fauna of the Coast of Ireland. Part V. Isopoda.— Fisheries, Ireland, Sci. Invest. 1904, 2: 1-90.
- Taylor, P.D. & P.L. Cook, 1981. *Hippoporidra edax* (Busk, 1859) and a revision of some fossil and living *Hippoporidra* (Bryozoa).— Bull. Br. Mus. (Nat. Hist.) Geol. Ser. 35: 243-251.
- Taylor, P.D., P.J. Schembri & P.L. Cook, 1989. Symbiotic associations between hermit crabs and bryozoans from the Otago region, southeastern New Zealand.— J. Nat. Hist. 23(5): 1059-1085.
- Thompson, W., 1844. Report on the fauna of Ireland.— Report of the Thirteenth Meeting of the British Association for the Advancement of Science in 1843: 267.
- Thorson, G. 1946. Reproduction and larval development of Danish marine bottom invertebrates, with special reference to the planktonic larvae in the sound (Øresund) (with a section on lamellibranch larvae by C. Barker Jørgensen): 1-523.— C.A. Reitzels Forlag, Kopenhagen.
- Türkay, M., 1976. Decapoda Reptantia von der portugiesischen und marokkanischen Küste Auswertung der Fahrten 8, 9c (1967), 19 (1970), 23 (1971) und 36 (1975) von F.S. "Meteor".— "Meteor" Forsch.-Ergebnisse, serie D, 23: 23-44.
- Vine, P., 1986. Red Sea Invertebrates: 1-224.— IMMEL Publishing, London.
- White, A., 1850. Catalogue of British Crustacea. List of the specimens of British animals in the collection of the British Museum.— Catalogues of the Zoological Collection in the British Museum. IV. British Zoology. 16. List of the British animals, 4: i-iv, 1-141.
- Williamson, D.I., 1982. Larval Morphology and Diversity. In: L.G. Abele (editor), The Biology of Crustacea, 2. Embryology, Morphology and Genetics, 2: 43-110.— Academic Press, New York.
- Zariquey Alvarez, R., 1946. Crustáceos decápodos mediterráneos. Manual para la clasificación de las especies que pueden capturarse en las costas mediterráneas españolas. Instituto Español de Estudios Mediterráneos.— Publicaciones sobre biología mediterránea, 2: 1-181.
- Zariquey Alvarez, R., 1955. Études carcinologiques méditerranéennes I: Crustáceos decápodos de la région de Cadaqués (España).— Vie Milieu 6 (3): 397-409.
- Zariquey Alvarez, R., 1968. Crustáceos Decápodos ibéricos.— Invest. Pesq. 32: i-xv, 1-510.

Received: 17.vi.1993

Accepted: 19.v.1994

Edited: C.H.J.M. Fransen

Index

- acantholepis, Micropagurus* 4, 8
Acanthopagurus 4
Acanthopagurus dubius 4
acus, Anapagurus 3, 4, 11
adriaticus, Anapagurus 3, 4, 5, 12, 19, 21, 72, 91, 97, 116, 117
alboranensis, Anapagurus 3, 4, 5, 13, 42, 66, 72, 85, 90, 116
Anapagrides 10
Anapagurus adriaticus 3, 4, 5, 12, 19, 21, 72, 91, 97, 116, 117
Anapagurus acutus 3, 4, 11
Anapagurus alboranensis 3, 4, 5, 13, 42, 66, 72, 85, 90, 116
Anapagurus atlantidii 3, 4, 5, 12, 14, 22, 97, 103, 104, 109, 110, 117
Anapagurus australiensis 4
Anapagurus bicorniger 3, 4, 5, 8, 12, 53, 56, 57, 58, 81, 84, 85, 116
Anapagurus bonnieri 3, 4, 5, 10, 13, 63, 64, 66, 117
Anapagurus Bonnieri 63
Anapagurus breviaculatus 67
Anapagurus breviaculeatus 3, 4, 5, 13, 21, 42, 66, 67, 70, 72, 91, 116
Anapagurus brevicarpus 3, 4, 30, 37, 39
Anapagurus chiracanthus 30
Anapagurus chiroacanthus 3, 5, 12, 29, 30, 36, 37, 39, 47, 77, 80, 81, 116, 117
Anapagurus chiroacanthus var. *cristatus* 30, 36
Anapagurus chiroacanthus var. *gracilis* 30
Anapagurus congolensis 3, 4, 5, 13, 14, 19, 22, 102, 110, 116, 117
Anapagurus curvidactylus 3, 4, 5, 12, 23, 29, 48, 51, 53, 77, 81, 109, 117
Anapagurus curvidactylus var.? 48
Anapagurus drachi 3, 4, 5, 117, 120
Anapagurus? dubius 4
Anapagurus ferrugineus 34
Anapagurus hendersoni 3, 4, 5, 12, 14, 22, 39, 40, 43, 72, 73, 77, 116, 117
Anapagurus hyndmani 23
Anapagurus hyndmanni 3, 4, 5, 12, 22, 23, 27, 29, 37, 48, 51, 53, 56, 70, 84, 90, 109, 116, 117
Anapagurus Hyndmanni 22, 29
Anapagurus japonicus 3, 4, 5, 10, 12, 37, 39, 44, 46, 47, 48, 117
Anapagurus japonicus 44
Anapagurus laevis ... 3, 5, 13, 14, 18, 19, 21, 22, 24, 40, 43, 58, 60, 62, 66, 72, 73, 77, 81, 91, 97, 102, 104, 116
Anapagurus laevis var. *longispina* 4, 58, 60, 62
Anapagurus levii 14
Anapagurus longispina 3, 4, 5, 12, 21, 22, 40, 43, 58, 60, 62, 72, 116, 117
Anapagurus marginatus 3, 4, 11
Anapagurus petiti 4, 5, 8, 11, 53, 57, 58, 81, 84, 85, 116
Anapagurus polynesiensis 4
Anapagurus pusillus 3, 4, 5, 13, 40, 42, 43, 47, 64, 72, 73, 77, 90, 91, 116, 117
Anapagurus pusillus var. *japonica* 4, 44
Anapagurus pusillus var. *japonicus* 44
Anapagurus sp. 64, 66
Anapagurus vossi 3, 4, 5, 12, 14, 19, 22, 97, 102, 103, 116, 117
Anapagurus wolffi 4, 5, 12, 14, 22, 53, 77, 80, 81, 103, 117
australiensis, Anapagurus 4
bicorniger, Anapagurus 3, 4, 5, 8, 12, 53, 56, 57, 58, 81, 84, 85, 116
bonnieri, Anapagurus 3, 4, 5, 10, 13, 63, 64, 66, 117
Bonnieri, Anapagurus 63
breviaculatus, Anapagurus 67

<i>breviaculeatus</i> , <i>Anapagurus</i>	3, 4, 5, 13, 21, 42, 66, 67, 70, 72, 91, 116
<i>brevicarpus</i> , <i>Anapagurus</i>	3, 4, 30, 37, 39
<i>chiracanthus</i> , <i>Anapagurus</i>	30
<i>chiroacanthus</i> , <i>Anapagurus</i>	3, 5, 12, 29, 30, 36, 37, 39, 47, 77, 80, 81, 116, 117
<i>chiroacanthus</i> var. <i>cristatus</i> , <i>Anapagurus</i>	30, 36
<i>chiroacanthus</i> var. <i>gracilis</i> , <i>Anapagurus</i>	30
<i>congolensis</i> , <i>Anapagurus</i>	3, 4, 5, 13, 14, 19, 22, 102, 110, 116, 117
<i>curvidactylus</i> , <i>Anapagurus</i>	3, 4, 5, 12, 23, 29, 48, 51, 53, 77, 81, 109, 117
<i>curvidactylus</i> var.?, <i>Anapagurus</i>	48
<i>drachi</i> , <i>Anapagurus</i>	3, 4, 5, 117, 120
<i>dubius</i> , <i>Acanthopagurus</i>	4
<i>dubius</i> , <i>Anapagurus</i> ?	4
<i>Enneobranchus</i>	10
<i>Eupagurus chiroacanthus</i>	30
<i>Eupagurus hyndmanni</i>	23
<i>Eupagurus Hyndmanni</i>	22
<i>Eupagurus laevis</i>	13, 21
<i>Forestopagurus</i>	3, 4, 6, 11, 117, 120
<i>ferrugineus</i> , <i>Anapagurus</i>	34
<i>hendersoni</i> , <i>Anapagurus</i>	3, 4, 5, 12, 14, 22, 39, 40, 43, 72, 73, 77, 116, 117
<i>hyndmani</i> , <i>Anapagurus</i>	23
<i>hyndmanni</i> , <i>Anapagurus</i>	3, 4, 5, 12, 22, 23, 27, 29, 37, 48, 51, 53, 56, 70, 84, 90, 109, 116, 117
<i>Hyndmanni</i> , <i>Anapagurus</i>	22, 29
<i>Iridopagurus</i>	10
<i>japonicus</i> , <i>Anapagurus</i>	3, 4, 5, 10, 12, 37, 39, 44, 46, 47, 48, 117
<i>japonicus</i> , <i>Anapagurus</i>	44
<i>laevis</i> , <i>Anapagurus</i> ...3, 5, 13, 14, 18, 19, 21, 22, 24, 40, 43, 58, 60, 62, 66, 72, 73, 77, 81, 91, 97, 102, 104, 116	
<i>laevis</i> var. <i>longispina</i> , <i>Anapagurus</i>	4, 58, 60, 62
<i>levis</i> , <i>Anapagurus</i>	14
<i>longispina</i> , <i>Anapagurus</i>	3, 4, 5, 12, 21, 22, 40, 43, 58, 60, 62, 72, 116, 117
<i>marginatus</i> , <i>Anapagurus</i>	3, 4, 11
<i>Micropagurus</i>	10
<i>Micropagurus acantholepis</i>	4, 8
<i>Micropagurus polynesiensis</i>	4
<i>Nematopaguroides</i>	11
<i>Ostraconotus</i>	120
<i>Pagurita</i>	120
<i>Pagurus</i>	4, 43, 77, 120
<i>Pagurus bernhardus</i>	24
<i>Pagurus chiracanthus</i>	30
<i>Pagurus chiroacanthus</i>	4, 29, 30, 37, 39
<i>Pagurus cuanensis</i>	21, 39
<i>Pagurus ferrugineus</i>	4, 11, 29, 34, 37, 39
<i>Pagurus hyndmanni</i>	21
<i>Pagurus Hyndmanni</i>	22
<i>Pagurus laevis</i>	4, 8, 13, 21, 39
<i>Pagurus moluccensis</i>	8
<i>Pagurus prideaux</i>	120
<i>Pagurus pubescens</i>	39
<i>Pagurus pubescentulus</i>	21
<i>Pagurus thomsoni</i>	39
<i>Pagurus ulidianus</i>	21
<i>polynesiensis</i> , <i>Anapagurus</i>	4
<i>Porcellanopagurus</i>	120

<i>pusillus</i> , <i>Anapagurus</i>	3, 4, 5, 13, 40, 42, 43, 47, 64, 72, 73, 77, 90, 91, 116, 117
<i>pusillus</i> var. <i>japonica</i> , <i>Anapagurus</i>	4, 44
<i>pusillus</i> var. <i>japonicus</i> , <i>Anapagurus</i>	44
<i>Pygmaeopagurus</i>	10
<i>Pygmaeopagurus hadrochirus</i>	10
<i>Spiropagurus</i>	4, 8, 10, 11, 21, 39
<i>Spiropagurus chiroacanthus</i>	30
<i>Spiropagurus (Anapagurus) chiroacanthus</i>	11, 29
<i>Spiropagurus Hyndmanni</i>	22
<i>Spiropagurus laevis</i>	13
sp., <i>Anapagurus</i>	64, 66
SPECIES N.1	81
<i>vossi</i> , <i>Anapagurus</i>	3, 4, 5, 12, 14, 19, 22, 97, 102, 103, 116, 117
<i>wolffi</i> , <i>Anapagurus</i>	4, 5, 12, 14, 22, 53, 77, 80, 81, 103, 117