


640
5666
REPT

CUBAN REPTILES:
ORIGINAL CITATIONS, HOLOTYPEs,
AND GEOGRAPHIC RANGE



Lourdes Rodríguez Schettino
Instituto de Ecología y Sistemática
Ministerio de Ciencia, Tecnología y Medio Ambiente



SMITHSONIAN
HERPETOLOGICAL INFORMATION
SERVICE
NO. 125

2000

SMITHSONIAN
HERPETOLOGICAL
INFORMATION
SERVICE .

The SHIS series publishes and distributes translations, bibliographies, indices, and similar items judged useful to individuals interested in the biology of amphibians and reptiles, but unlikely to be published in the normal technical journals. Single copies are distributed free to interested individuals. Libraries, herpetological associations, and research laboratories are invited to exchange their publications with the Division of Amphibians and Reptiles.

We wish to encourage individuals to share their bibliographies, translations, etc. with other herpetologists through the SHIS series. If you have such items please contact George Zug for instructions on preparation and submission. Contributors receive 50 free copies.

Please address all requests for copies and inquiries to George Zug, Division of Amphibians and Reptiles, National Museum of Natural History, Smithsonian Institution, Washington DC 20560 USA. Please include a self-addressed mailing label with requests.

INTRODUCTION

The Cuban terrestrial reptile fauna is composed of 120 known species, of which 97 species are endemic (80.8%). Cuban reptiles occur throughout the country, not only in all natural vegetation types but also in urban and agricultural places. They eat many invertebrates and are the prey for other vertebrates. For these reasons, reptiles play an important role in the food webs of Cuban ecosystems and must be well understood in order to ensure their correct management and conservation.

Since the publication of the most recent checklist of the West Indian herpetofauna (Powell *et al.*, 1996), several new Cuban reptiles have been described (Díaz *et al.*, 1996; Pérez-Beato, 1996-1997; Thomas and Hedges, 1998a, b; Thomas *et al.*, 1998) and some subspecies have been elevated to species (Thomas and Hedges, 1998a, b). As Powell *et al.* (1996) stated, no previous list had provided complete citations of the original descriptions of species. They provided a full list to the species level. Nevertheless, the subspecies level is often useful for understanding variation within species. Consequently, I give here a list with complete citations for all Cuban terrestrial reptiles described up to 1998, including the recognized species and subspecies.

For each taxon I report: A) the correct name in use with author and date; B) the complete citation of the original description; C) the page number for the name, between parentheses; D) the holotype or syntypes; and E) the geographic range within Cuba. The generic names *Chamaeleolis* and *Cadea* are maintained here despite the opinions of Hass *et al.* (1993) and Hedges (1996), respectively, because I believe that both have enough diagnostic characters for maintaining their taxonomic status and recognizing them as genera.

The abbreviations used are as follows: AMNH, American Museum of Natural History, New York; ANSP, Academy of Natural Sciences, Philadelphia; BMNH, British Museum (Natural History), London; BYU, Brigham Young University, Provo; CARE, Colección de Alberto R. Estrada; CAS-SU, Stanford University (in the California Academy of Sciences, San Francisco); CM, Carnegie Museum of Natural History, Pittsburgh; ChM, Charleston Museum, Charleston; CZACC, Colecciones Zoológicas, Instituto de Ecología y Sistemática, La Habana; HZM, Universität Hamburg, Zoologische Museum, Hamburg; IZ, Instituto de Zoología, La Habana; MCTH, Museo Carlos de la Torre y Huerta, Holguín, Cuba; MCZ, Museum of Comparative Zoology, Harvard University, Cambridge; MNHN, Muséum National d'Histoire Naturelle, Paris; MNHNCU, Museo Nacional de Historia Natural de Cuba, La Habana; UIMNH, University of Illinois, Museum of Natural History, Urbana; UMMZ, Museum of Zoology, University of Michigan, Ann Arbor; USNM, National Museum of Natural History, Washington; UZM, Universitets Zoologiske Museum, Kjøbenhavn; ZMB, Museum für Naturkunde, Humboldt-Universität, Berlin.

I thank Kevin de Queiroz for his help in acquiring some of the papers cited here. Also, Kevin de Queiroz, Vilma Rivalta, Ada Chamizo, Luis F. de Armas, and Luis V. Moreno for their useful comments on the manuscript.

CUBAN REPTILES: ORIGINAL CITATIONS & DISTRIBUTIONS

Order Testudines
Family Emydidae

Trachemys decussata decussata (Gray), 1831. Gray, J. E. 1831. *Synopsis Reptilium*. Treuttel, Wurz, Sowerby, and Woods, Londres, viii + 84 pp. (p. 28). Holotype: BMNH 1947.3.4.79. Central and eastern Cuba.

Trachemys decussata angusta (Barbour and Carr), 1940. Barbour, T. and Carr, A. 1940. Antillean terrapins. *Mem. Mus. Comp. Zool.* 54(5):381-415. (p. 402). Holotype: MCZ 34340. Western Cuba; Isla de la Juventud.

Order Crocodilia
Family Crocodylidae

Crocodylus rhombifer Cuvier, 1807. Cuvier, G. 1807. Sur les différentes espèces de crocodiles vivans et sur leurs caractères distinctifs. *Ann. Mus. Nat. Hist. Paris* 10:8-86. (p. 51). Holotype: Unlocated: probably in MNHN but not designated. Ciénaga de Zapata; Ciénaga de Lanier.

Crocodylus acutus Cuvier, 1807. Cuvier, G. 1807. Sur les différentes espèces de crocodiles vivans et sur leurs caractères distinctifs. *Ann. Mus. Nat. Hist. Paris* 10:8-86. (p. 55). Holotype: Unlocated. Cuba.

Caiman crocodilus fuscus (Cope), 1868. Cope, E. D. 1868. On the crocodilian genus *Perosuchus*. *Proc. Acad. Nat. Sci. Philadelphia* 20:203. (p. 203). Holotype: Unknown. Ciénaga de Lanier.

Order Squamata - Lizards
Family Amphisbaenidae

Amphisbaena cubana Gundlach and Peters in Peters, 1878. Peters, W. 1878 Über vier neue amerikanische Amphisbaena-Arten. *Monatsber. Akad. Wiss. Berlin* 1878:778-781. (p. 780). Holotype: ZMB 9383. From Cienfuegos to San Carlos, Guantánamo province; Isla de la Juventud.

Amphisbaena barbouri Gans and Alexander, 1962. Gans, C. and Alexander, A. A. 1962. Studies on Amphisbaenids (Amphisbaenia, Reptilia). 2. On the amphisbaenids of the Antilles. *Bull. Mus. Comp. Zool.* 128(3):65-158. (p. 97). Holotype: MCZ 12136. From La Habana province to Cienfuegos province.

Amphisbaena carlgansi Thomas and Hedges, 1998. Thomas, R. and Hedges, S. B. 1998. A new amphisbaenian from Cuba. *J. Herpetol.* 32(1):92-96. (p. 93). Holotype: MNNHCU-4421. Meseta de Cabo Cruz, Granma province.

Cadea blanoides Stejneger, 1916. Stejneger, L. 1916. Notes on amphisbaenian nomenclature. *Proc. Biol. Soc. Washington* 29:85. (p. 85). Holotype: BMNH 1946.8.2.20. From Sierra de los Órganos, Pinar del Río province to Matanzas province; Isla de la Juventud.

Cadea palirostrata Dickerson, 1916. Dickerson, M. C. 1916. Description of a new amphisbaenian collected by the late Dr. Charles S. Mead in 1911 on the Isle of Pines, Cuba. *Amer. Mus. Nat. Hist.* 35(34):659-662. (p. 659). Holotype: AMNH 2717. Isla de la Juventud.

Family Anguidae

Diploglossus delasagra (Cocteau), 1838 or 1839. Cocteau, J. T. 1838 or 1839. Reptiles. In *Historia Física, Política y Natural de la Isla de Cuba* (R. de la Sagra, ed.), (p. 110). Syntypes: MNHN 2856, 2858, 2859; RNH 3626. From Pinar del Río province to Gibara in Holguín province; Isla de la Juventud.

Diploglossus nigropunctatus Barbour and Shreve, 1937. Barbour, T. and Shreve, B. 1937. Novitates Cubanae. *Bull. Mus. Comp. Zool.* 80(9):377-387 + 4 láms. (p. 378). Holotype: MCZ 42504. Northeastern Guantánamo.

Diploglossus garridoi Thomas and Hedges, 1998. Thomas, R. and Hedges, S. B. 1998. New anguid lizard (*Diploglossus*) from Cuba. *Copeia* 1998(1):97-103. (p. 97). Holotype: MNHNCU 4420. El Manguito, Granma province.

Family Gekkonidae

Gonatodes albogularis fuscus Hallowell, 1855. Hallowell, E. 1855. Contributions to South American herpetology. *J. Acad. Nat. Sci. Philadelphia* 2(3):33-36. (p. 33). Holotype: Unlocated. Cuba.

Hemidactylus haitianus Meerwarth, 1901. Meerwarth, H. 1901. Die westindischen Reptilien und Batrachier des naturhistorischen Museums in Hamburg. *Mitt. Naturhist. Mus. Hamburg* 18:1-41. (p. 17). Holotype: HZM (destroyed). Cuba.

Hemidactylus mabouia Moreau de Jonnès, 1818. Moreau de Jonnès, A. 1818. Monographie du mabouia des murailles, ou *Gecko Mabouia* des Antilles. *Bull. Sci. Soc. Philomath. Paris* 1818:138-139. (p. 138). Holotype: MNHN 6573. Guantánamo city.

Hemidactylus turcicus turcicus (Linnaeus), 1758. Linnaeus, K. von 1758. *Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Tomus I, Editio Decima, reformata, 824 pp.* (p. 202). Holotype: Unlocated. Coastal places between La Habana city and Santiago de Cuba city.

Sphaerodactylus elegans elegans Mac Leay, 1834. Mac Leay, W. S. 1834. A few remarks No. XIV. *Proc. Zool. Soc. London* II(14):9-12. (p. 12). Holotype: Unlocated. Cuba; Isla de la Juventud; Archipiélago de los Canarreos, Archipiélago de Sabana-Camagüey; and Archipiélago de los Jardines de la Reina.

Sphaerodactylus argus argus Gosse, 1850. Gosse, P. 1850. Descriptions of a new genus and six new species of saurian reptiles. *Ann. Mag. Nat. Hist.* 2(6):344-348. (p. 347). Syntypes: BMNH 47.12.24.56, 47.12.24.59. From Juraguá, Cienfuegos province, to Santa Clara; from Los Negros to Cabo Cruz, Granma province; Santiago de Cuba city; Archipiélago de los Jardines de la Reina.

Sphaerodactylus torrei torrei Barbour, 1914. Barbour, T. 1914. A contribution to the zoogeography of the West Indies, with special references to amphibians and reptiles. *Mem. Mus. Comp. Zool.* 44(2):209-359. (p. 260). Holotype: MCZ 6916. From Santiago de Cuba city to Playa Juraguá.

Sphaerodactylus torrei spielmani Grant, 1958. Grant, C. 1958. A new gekkonid lizard (*Sphaerodactylus*) from Cuba. *Herpetologica* 14(4):225-227. (p. 225). Holotype: UIMNH 44105. West of Guantánamo Bay.

Sphaerodactylus scaber Barbour and Ramsden, 1919. Barbour, T. and Ramsden, C. 1919. The herpetology of Cuba. *Mem. Mus. Comp. Zool.* 47(2):71-213. (p. 126). Holotype: MCZ 12304. From Sagua la Grande, Villa Clara province, to Sierra de Najasa, Camagüey province; Cayo Conuco, Villa Clara province.

Sphaerodactylus intermedius Barbour and Ramsden, 1919. Barbour, T. and Ramsden, C. 1919. The herpetology of Cuba. *Mem. Mus. Comp. Zool.* 47(2):71-213. (p. 211). Holotype: MCZ 12305. From Rincón de Guanabo, La Habana province, to Hato Nuevo, Matanzas province.

Sphaerodactylus oliveri Grant, 1944. Grant, C. 1944. New *Sphaerodactylus* from Cuba and the Isla de Pinos. *Herpetologica* 2(6):118-125. (p. 118). Holotype: UMMZ 93310 or CAS-SU 14683. From eastern Cienfuegos to Trinidad and Topes de Collantes.

Sphaerodactylus storeyae Grant, 1944. Grant, C. 1944. New *Sphaerodactylus* from Cuba and the Isla de Pinos. *Herpetologica* 2(6):118-125. (p. 125). Holotype: CAS-SU 9296. Punta del Este, Isla de la Juventud.

Sphaerodactylus ramsdeni Ruibal, 1959. Ruibal, R. 1959. A new species of *Sphaerodactylus* from Oriente, Cuba. *Herpetologica* 15(2):89-93. (p. 89). Holotype: MCZ 8536. Sierras de la Gran Piedra, Santiago de Cuba province, and Sierra del Guaso, Guantánamo province.

Sphaerodactylus ruibali Grant, 1959. Grant, C. 1959. Another new *Sphaerodactylus* from Guantanomo, Cuba. *Herpetologica* 15(1):53. (p. 53). Holotype: UIMNH 44246. From western Guantánamo Bay to Macambo.

Sphaerodactylus nigropunctatus alayoi Grant, 1959. Grant, C. 1959. A new *Sphaerodactylus* from Guantanomo, Cuba. *Herpetologica* 15(1):49-53. (p. 49). Holotype: UIMNH 44215. Vicinity of Guantánamo city.

- Sphaerodactylus nigropunctatus granti* Thomas and Schwartz, 1966. Thomas, R. and Schwartz, A. 1966. The *Sphaerodactylus decoratus* complex in the West Indies. *Brigham Young Univ. Sci. Bull.* 7(4):1-26. (p. 10). Holotype: BYU 17233. Archipiélago de Sabana-Camagüey; from Nuevitas, Camagüey province, to Levisa, Holguín province, on North and to Guantánamo on South.
- Sphaerodactylus nigropunctatus lissodesmus* Thomas and Schwartz, 1966. Thomas, R. and Schwartz, A. 1966. The *Sphaerodactylus decoratus* complex in the West Indies. *Brigham Young Univ. Sci. Bull.* 7(4):1-26. (p. 12). Holotype: MCZ 57344. Sierra de Cubitas, Camagüey province.
- Sphaerodactylus nigropunctatus strategus* Thomas and Schwartz, 1966. Thomas, R. and Schwartz, A. 1966. The *Sphaerodactylus decoratus* complex in the West Indies. *Brigham Young Univ. Sci. Bull.* 7(4):1-26. (p. 13). Holotype: MCZ 81110. From Guantánamo Naval Base to Caimanera.
- Sphaerodactylus nigropunctatus ocujal* Thomas and Schwartz, 1966. Thomas, R. and Schwartz, A. 1966. The *Sphaerodactylus decoratus* complex in the West Indies. *Brigham Young Univ. Sci. Bull.* 7(4):1-26. (p. 16). Holotype: USNM 138015. From Niquero, Granma province, to Jutisí, east of Santiago de Cuba Bay.
- Sphaerodactylus notatus atactus* Schwartz, 1966. Schwartz, A. 1966. Geographic variation in *Sphaerodactylus notatus* Baird. *Rev. Biol. Trop.* 13(2):161-185. (p. 166). Holotype: AMNH 92820. Cuba; Isla de la Juventud; Archipiélago de los Canarreos; Cayos de San Felipe.
- Sphaerodactylus bromeliacum* Peters and Schwartz, 1972. Peters, G. and Schwartz, A. 1972. Ein neuer, Bromelien bewohnender Kugelfingergecko (Gekkonidae: *Sphaerodactylus*) aus Oriente/Cuba. *Mitt. Zool. Mus. Berlin* 48(2):393-399. (p. 395). Holotype: ZMB 42827. Yunque de Baracoa.
- Sphaerodactylus armasi* Schwartz and Garrido, 1974. Schwartz, A. and Garrido, O. H. 1974. A new Cuban species of *Sphaerodactylus* (Gekkonidae) of the *nigropunctatus* complex. *Proc. Biol. Soc. Washington* 8(30):337-344. (p. 339). Holotype: IZ 4089. From western Guantánamo Bay to Maisí, Guantánamo province.
- Sphaerodactylus celicara* Garrido and Schwartz, 1982. Garrido, O. H. and Schwartz, A. 1982. A new species of *Sphaerodactylus* (Reptilia: Sauria: Gekkonidae) from eastern Cuba. *Proc. Biol. Soc. Washington* 95(2):392-397. (p. 392). Holotype: IZ 5613. From Baracoa to Maisí, Guantánamo province.
- Sphaerodactylus docimus* Schwartz and Garrido, 1985. Schwartz, A. and Garrido, O. H. 1985. The Cuban lizards of the genus *Sphaerodactylus* (Sauria, Gekkonidae). *Milwaukee Public Mus. Contr. Biol. Geol.* 62:1-67. (p. 22). Holotype: MCZ 8510. Jutisí, Santiago de Cuba province.

Sphaerodactylus schwartzi Thomas, Hedges and Garrido, 1992. Thomas, R., Hedges, S. B. and Garrido, O. H. 1992. Two new species of *Sphaerodactylus* from eastern Cuba (Squamata: Gekkonidae). *Herpetologica* 48(3):358-367. (p. 359). Holotype: MNHNCU 3438. Southwestern Guantánamo province.

Sphaerodactylus cricoderus Thomas, Hedges, and Garrido, 1992. Thomas, R., Hedges, S. B. and Garrido, O. H. 1992. Two new species of *Sphaerodactylus* from eastern Cuba (Squamata: Gekkonidae). *Herpetologica* 48(3):358-367. (p. 362). Holotype: MNHNCU 238. Uvero, Santiago de Cuba province.

Sphaerodactylus pimienta Thomas, Hedges, and Garrido, 1998. Thomas, R., Hedges, S. B., and Garrido, O. H. 1998. A new gecko (*Sphaerodactylus*) from the Sierra Maestra of Cuba. *J. Herpetol.* 32(1):66-69. (p. 66). Holotype: MHNHC 4417. Northern slope of Sierra Maestra.

Tarentola americana americana Gray, 1831. Gray, J. E. 1831. *A synopsis of the species of the Class Reptilia*. Appendix to E. Griffith Cuvier's Animal Kingdom. Whittaker, Treucher, and Co., Londres, 110 pp., 55 pls. (p. 48). Holotype: AMNH 6700. Cuba.

Family Iguanidae

Cyclura nubila nubila (Gray), 1831. Gray, J. E. 1831. *A synopsis of the species of the class Reptilia*. Appendix to E. Griffith Cuvier's Animal Kingdom. Whittaker, Treacher, and Co., Londres, 110 pp., 55 pls. (p. 39). Holotype: BMNH 1946.8.29.88. Cuba.

Family Polychrotidae

Anolis equestris equestris Merrem, 1820. Merrem, B. 1820. *Tentamen Systematis Amphibiorum*. Marburg, xv + 191 pp. (p. 45). Holotype: Unlocated. From San Diego de los Baños, Pinar del Río province, to Sagua la Grande, Villa Clara province.

Anolis equestris thomasi Schwartz, 1958. Schwartz, A. 1958. A new subspecies of *Anolis equestris* from Eastern Cuba. *Herpetologica* 14(1):1-7. (p. 3). Holotype: AMNH 78148. From Camagüey province to Banes, Holguín province, and Bayamo, Granma province.

Anolis equestris buidei Schwartz and Garrido, 1972. Schwartz, A. and Garrido, O. H. 1972. The lizards of the *Anolis equestris* complex in Cuba. *Studies Fauna Curaçao other Carib. Isl.* 39(134):1-86. (p. 34). Holotype: IZ 1294 = CZACC 4.3193. Península de Hicacos.

Anolis equestris persparsus Schwartz and Garrido, 1972. Schwartz, A. and Garrido, O. H. 1972. The lizards of the *Anolis equestris* complex in Cuba. *Studies Fauna Curaçao other Carib. Isl.* 39(134):1-86. (p. 36). Holotype: AMNH 78116. Eastern Cienfuegos; Villa Clara and Sancti Spíritus provinces.

Anolis equestris juraguensis Schwartz and Garrido, 1972. Schwartz, A. and Garrido, O. H. 1972. The lizards of the *Anolis equestris* complex in Cuba. *Studies Fauna Curaçao other Carib. Isl.* 39(134):1-86. (p. 39). Holotype: IZ 1152 = CZACC 4.3210. Juraguá, west to Cienfuegos Bay.

- Anolis equestris verreonensis* Schwartz and Garrido, 1972. Schwartz, A. and Garrido, O. H. 1972. The lizards of the *Anolis equestris* complex in Cuba. *Studies Fauna Curaçao other Carib. Isl.* 39(134):1-86. (p. 44). Holotype: IZ 488 = CZACC 4.3221. From Vereón to Cabo Cruz, Granma province.
- Anolis equestris potior* Schwartz and Thomas, 1975. Schwartz, A. and Thomas, R. 1975. A checklist of West Indian amphibians and reptiles. *Carnegie Mus. Nat. Hist. Special Publ.* 1:1-216. (p. 81). Holotype: IZ 3098 = CZACC 4.3417. Cayo Santa María, Archipiélago de Sabana-Camagüey.
- Anolis equestris cincoleguas* Garrido, 1981. Garrido, O. H. 1981. Nueva subespecie de *Anolis equestris* (Sauria: Iguanidae) para Cuba, con comentarios sobre la distribución y afinidades de otras poblaciones del complejo. *Poeyana* 232:1-15. (p. 3). Holotype: IZ 5398. Cayos de las Cinco Leguas, Matanzas province.
- Anolis luteogularis luteogularis* Noble and Hassler, 1935. Noble, G. K. and Hassler, W. G. 1935. A new giant *Anolis* from Cuba. *Copeia* 3:113-115. (p. 113). Holotype: AMNH 46502. From eastern Península de Guanahacabibes to Güines, south of La Habana province.
- Anolis luteogularis hassleri* Barbour and Shreve, 1935. Barbour, T. and Shreve, B. 1935. Notes on Cuban anoles. *Occ. Papers Boston Soc. Nat. Hist.* 8:249-254. (p. 251). Holotype: MCZ 11178. Isla de la Juventud, northern region.
- Anolis luteogularis nivevultus* Schwartz and Garrido, 1972. Schwartz, A. and Garrido, O. H. 1972. The lizards of the *Anolis equestris* complex in Cuba. *Studies Fauna Curaçao other Carib. Isl.* 39(134):1-86. (p. 11). Holotype: IZ 339 = CZACC 4.3144. Península de Guanahacabibes.
- Anolis luteogularis delacruzi* Schwartz and Garrido, 1972. Schwartz, A. and Garrido, O. H. 1972. The lizards of the *Anolis equestris* complex in Cuba. *Studies Fauna Curaçao other Carib. Isl.* 39(134):1-86. (p. 16). Holotype: IZ 1277 = CZACC 4.3155. Santa Isabel, Isla de la Juventud.
- Anolis luteogularis sectilis* Schwartz and Garrido, 1972. Schwartz, A. and Garrido, O. H. 1972. The lizards of the *Anolis equestris* complex in Cuba. *Studies Fauna Curaçao other Carib. Isl.* 39(134):1-86. (p. 19). Holotype: IZ 388 = CZACC 4.3157. Isla de la Juventud, southern region.
- Anolis luteogularis coctilis* Schwartz and Garrido, 1972. Schwartz, A. and Garrido, O. H. 1972. The lizards of the *Anolis equestris* complex in Cuba. *Studies Fauna Curaçao other Carib. Isl.* 39(134):1-86. (p. 22). Holotype: IZ 402 = CZACC 4.3163. Cayo Cantiles, Archipiélago de los Canarreos.
- Anolis luteogularis calceus* Schwartz and Garrido, 1972. Schwartz, A. and Garrido, O. H. 1972. The lizards of the *Anolis equestris* complex in Cuba. *Studies Fauna Curaçao other Carib. Isl.* 39(134):1-86. (p. 25). Holotype: IZ 1295 = CZACC 4.3117. Santo Tomás, Península de Zapata.

Anolis luteogularis jaumei Schwartz and Garrido, 1972. Schwartz, A. and Garrido, O. H. 1972. The lizards of the *Anolis equestris* complex in Cuba. *Studies Fauna Curaçao other Carib. Isl.* 39(134):1-86. (p. 27). Holotype: IZ 369 = CZACC 4.3180. Playa Larga, Bahía de Cochinos.

Anolis luteogularis sanfelipensis Garrido, 1975. Garrido, O. H. 1975. Nuevos reptiles del Archipiélago Cubano. *Poeyana* 141:1-58. (p. 23). Holotype: IZ 2972 = CZACC 4.3358. Cayo Real, Cayos de San Felipe, off the southern coast of Pinar del Río province.

Anolis noblei noblei Barbour and Shreve, 1935. Barbour, T. and Shreve, B. 1935. Notes on Cuban anoles. *Occ. Papers Boston Soc. Nat. Hist.* 8:249-254. (p. 250). Holotype: MCZ 26653. Sierra de Nipe and Cupeyal, Holguín province.

Anolis noblei galeifer Schwartz, 1964. Schwartz, A. 1964. *Anolis equestris* in Oriente province, Cuba. *Bull. Mus. Comp. Zool.* 131(12):403-428. (p. 409). Holotype: MCZ 59326. Sierra Maestra; Sierra de la Gran Piedra.

Anolis baracoae Schwartz, 1964. Schwartz, A. 1964. *Anolis equestris* in Oriente province, Cuba. *Bull. Mus. Comp. Zool.* 131(12):403-428. (p. 419). Holotype: MCZ 57404. From Cayo Güín to Maisí, Guantánamo province.

Anolis smallwoodi smallwoodi Schwartz, 1964. Schwartz, A. 1964. *Anolis equestris* in Oriente province, Cuba. *Bull. Mus. Comp. Zool.* 131(12):403-428. (p. 412). Holotype: AMNH 89526. From eastern Sierra Maestra to west of Guantánamo Bay.

Anolis smallwoodi palardis Schwartz, 1964. Schwartz, A. 1964. *Anolis equestris* in Oriente province, Cuba. *Bull. Mus. Comp. Zool.* 131(12):403-428. (p. 416). Holotype: CM 33320. From eastern Guantánamo Bay to Monte Líbano and Baitiquirí.

Anolis smallwoodi saxuliceps Schwartz, 1964. Schwartz, A. 1964. *Anolis equestris* in Oriente province, Cuba. *Bull. Mus. Comp. Zool.* 131(12):403-428. (p. 422). Holotype: HZM 5376. From Moa, Holguín province, to Felicidad, Guantánamo province.

Anolis pigmaequestris Garrido, 1975. Garrido, O. H. 1975. Nuevos reptiles del Archipiélago Cubano. *Poeyana* 141:1-58. (p. 4). Holotype: IZ 2884 = CZACC 4.3344. Francés and Santa María Keys, Archipiélago de Sabana-Camagüey.

Anolis porcatus porcatus Gray, 1840. Gray, J. E. 1840. Catalogue of the species of reptiles collected in Cuba by W. S. MacLeay, Esq; with some notes on their habits extracted from his MS. *Ann. Mag. Nat. Hist.* 1(5):108-115. (p. 112). Syntypes BMNH 1946.8.12.7, 1946.8.12.66-70. Cuba; Archipiélago de los Colorados; Archipiélago de los Canarreos; Archipiélago de Sabana-Camagüey; and Archipiélago de los Jardines de la Reina.

Anolis porcatus aracelyae Pérez-Beato, 1996-1997. Pérez-Beato, O. 1996-1997. A new subspecies of *Anolis porcatus* (Sauria: Polychrotidae) from Western Cuba. *Rev. Biol. Trop.* 44(3)/45(1):295-299. (p. 296). Holotype: MNHNCU-1445. Pinar del Río province, excepting Península de Guanahacabibes.

Anolis isolepis isolepis Cope, 1861. Cope, E. D. 1861. Notes and descriptions of Anoles. *Proc. Acad. Nat. Sci. Philadelphia* 13:208-215. (p. 214). Syntypes: USNM lost. Sierra de Trinidad; Sierra de Jatibonico; Sierra Maestra; Macizo de Sagua Baracoa.

Anolis isolepis altitudinalis Garrido, 1985. Garrido, O. H. 1985. Nueva subespecie de *Anolis isolepis* (Lacertilia: Iguanidae) para Cuba. *Doñana, Acta Vertebrata* 12(1):41-49. (p. 42). Holotype: CZACC 4.7028. Pico Turquino.

Anolis allisoni Barbour, 1928. Barbour, T. 1928. Reptiles from the Bay Islands. *Proc. New England Zool. Club* 10:55-61. (p. 58). Holotype: MCZ 26725. From Playa Caimito, south of La Habana province, to Banes, Holguín province, and Cabo Cruz, Granma province.

Anolis angusticeps angusticeps Hallowell, 1856. Hallowell, E. 1856. Notes on the reptiles in the collection of the Museum of the Academy of Natural Sciences. *Proc. Acad. Nat. Sci. Philadelphia* 8:221-238. (p. 228). Holotype: ANSP 7789. Cuba; Archipiélago de los Colorados; Archipiélago de los Canarreos; Archipiélago de Sabana-Camagüey; and Archipiélago de los Jardines de la Reina.

Anolis paternus paternus Hardy, 1967. Hardy, J. D., Jr. 1967. Geographic variation in the West Indian lizard, *Anolis angusticeps*, with the description of a new form, *Anolis angusticeps paternus* from the Isle of Pines, Cuba (Reptilia: Iguanidae). *Carib. J. Sci.* 6(1-2):23-31. (p. 25). Holotype: USNM 142156. Isla de la Juventud, northern region.

Anolis paternus pinarensis Garrido, 1975. Garrido, O. H. 1975. Variación de *Anolis angusticeps* Hallowell (Lacertilia: Iguanidae) en el occidente de Cuba y en Isla de Pinos. *Poeyana* 144:1-18. (p. 8). Holotype: IZ 4073 = CZACC 4.3431. Southwestern Pinar del Río province.

Anolis guazuma Garrido, 1983. Garrido, O. H. 1983. Nueva especie de *Anolis* (Lacertilia: Iguanidae) de la Sierra del Turquino, Cuba. *Carib. J. Sci.* 19(3-4):71-76. (p. 71). Holotype: CZACC 4.6128. Pico Turquino.

Anolis alayoni Estrada and Hedges, 1995. Estrada, A. R. and Hedges, S. B. 1995. A new species of *Anolis* (Sauria: Iguanidae) from eastern Cuba. *Carib. J. Sci.* 31(1-2):65-72. (p. 65). Holotype: MHNHCU-2746. Mountains of Holguín and Guantánamo provinces.

Anolis garridoi Díaz, Estrada, and Moreno, 1996. Díaz, L. M., Estrada, A. R., and Moreno, L. V. 1996. A new species of *Anolis* (Sauria: Iguanidae) from the Sierra de Trinidad, Sancti Spíritus, Cuba. *Carib. J. Sci.* 32(1):54-58. (p. 54). Holotype: MHNHCU-4285. Topes de Collantes.

Anolis loysiana Duméril and Bibron, 1837. Duméril, A. M. C. and Bibron, G. 1837. *Erpétologie Générale, ou Histoire Naturelle Complète des Reptiles* 4. Librairie Encyclopédique de Roret, Paris, 572 pp. (p. 100). Holotype: MNHN 2465. Cuba.

Anolis argillaceus Cope, 1862. Cope, E. D. 1862. Contributions to neotropical saurology. *Proc. Acad. Nat. Sci. Philadelphia* 14:176-188. (p. 176). Holotype: USNM lost. From Nuevitas, Camagüey province, to Maisí, Guantánamo province.

- Anolis centralis centralis* Peters, 1970. Peters, G. 1970. Zur Taxonomie und Zoogeographie der Kabanischen Anolinen Eidechsen (Reptilia, Iguanidae). *Mitt. Zool. Mus. Berlin* 46(1):197-234. (p. 215). Holotype: ZMB 41616. From Sierra de Cubitas, Camagüey province, to Gibara, Holguín province, and Bayamo, Granma province.
- Anolis centralis litoralis* Garrido, 1975. Garrido, O. H. 1975. Distribución y variación de *Anolis argillaceus* Cope (Lacertilia: Iguanidae) en Cuba. *Poeyana* 142:1-28. (p. 12). Holotype: IZ 3472 = CZACC 4.3390. From Santiago de Cuba to Baitiquirí, Guantánamo province.
- Anolis pumilus* Garrido, 1988. Garrido, O. H. 1988. Nueva especie para la ciencia de *Anolis* (Lacertilia: Iguanidae) de Cuba perteneciente al complejo *argillaceus*. *Doñana, Acta Vertebrata* 15(1):45-57. (p. 46). Holotype: IZ 4470. From Península de Guanahacabibes to Jatibonico, Sancti Spíritus province; Isla de la Juventud; Archipiélago de Sabana.
- Anolis lucius* Duméril and Bibron, 1837. Duméril, A. M. C. and Bibron, G. 1837. *Erpétologie Générale, ou Histoire Naturelle Complète des Reptiles*, 4. Librairie Encyclopédique de Roret, Paris, 572 pp. (p. 105). Holotype: MNHN 2466. From Boca de Jaruco, La Habana province, to Baire, Granma province.
- Anolis argenteolus* Cope, 1861. Cope, E. D. 1861. Notes and descriptions of anoles. *Proc. Acad. Nat. Sci. Philadelphia* 13:208-215. (p. 213). Holotype: USNM lost. From Sierra de Najasa, Camagüey province, to Maisí, Guantánamo province.
- Anolis vermiculatus* Duméril and Bibron, 1837. Duméril, A. M. C. and Bibron, G. 1837. *Erpétologie Générale, ou Histoire Naturelle Complète des Reptiles*, 4. Librairie Encyclopédique de Roret, Paris, 572 pp. (p. 128). Syntypes MNHN 2407, 2349. Sierra de los Órganos and Sierra del Rosario.
- Anolis bartschi* (Cochran), 1928. Cochran, D. 1928. A second species of *Deiroptyx* from Cuba. *Proc. Biol. Soc. Washington* 41:169-170. (p. 169). Holotype: USNM 75805. Sierra de los Órganos.
- Anolis alutaceus* Cope, 1861. Cope, E. D. 1861. Notes and descriptions of anoles. *Proc. Acad. Nat. Sci. Philadelphia* 13:208-215. (p. 212). Syntypes MCZ 10932, ?USNM 27485-87. Cuba; Isla de la Juventud.
- Anolis clivicola* Barbour and Shreve, 1935. Barbour, T. and Shreve, B. 1935. Notes on Cuban anoles. *Occ. Papers Boston Soc. Nat. Hist.* 8:249-254. (p. 251). Holotype: MCZ 39664. Pico Turquino and Pico Bayamesa.
- Anolis anfiloquioi* Garrido, 1980. Garrido, O. H. 1980. Revisión del complejo *Anolis alutaceus* (Lacertilia: Iguanidae) y descripción de una nueva especie de Cuba. *Poeyana* 201:1-41. (p. 17). Holotype: IZ 4183. Levisa, Holguín province; Baracoa; Guantánamo.

Anolis inexpectata Garrido and Estrada, 1989. Garrido, O. H. and Estrada, A. R. 1989. Nueva especie del complejo *Anolis alutaceus* (Lacertilia: Iguanidae) para Cuba. *Rev. Biol.* 3(1):57-66. (p. 59). Holotype: MHNHCU-291. From Moa, Holguín province, to Yateras, Guantánamo province.

Anolis macilentus Garrido and Hedges, 1992. Garrido, O. H. and Hedges, S. B. 1992. Three new grass anoles from Cuba (Squamata: Iguanidae). *Carib. J. Sci.* 28:1-2. (p. 22). Holotype: MHNHCU-2721. Meseta del Guaso, Guantánamo province.

Anolis vescus Garrido and Hedges, 1992. Garrido, O. H. and Hedges, S. B. 1992. Three new grass anoles from Cuba (Squamata: Iguanidae). *Carib. J. Sci.* 28:1-2. (p. 25). Holotype: MHNHCU-2729. Palmarito, mountains to the north of Imías.

Anolis alfaroi Garrido and Hedges, 1992. Garrido, O. H. and Hedges, S. B. 1992. Three new grass anoles from Cuba (Squamata: Iguanidae). *Carib. J. Sci.* 28:1-2. (p. 27). Holotype: MHNHCU-2725. La Munición, Yateras, Guantánamo province.

Anolis cyanopleurus cyanopleurus Cope, 1861. Cope, E. D. 1861. Notes and descriptions of anoles. *Proc. Acad. Nat. Sci. Philadelphia* 13:208-215. (p. 211). Syntypes USNM 62068-70. From Yateras to Baracoa, Guantánamo province.

Anolys cyanopleurus orientalis Garrido, 1975. Garrido, O. H. 1975. Distribución y variación del complejo *Anolis cyanopleurus* (Lacertilia: Iguanidae) en Cuba. *Poeyana* 143:1-60. (p. 16). Holotype: IZ 1564 = CZACC 4.3397. Maisí, Guantánamo province.

Anolis cupeyalensis Peters, 1970. Peters, G. 1970. Zur Taxonomie und Zoogeographie der Kubanischen Anolinen Eidechsen (Reptilia, Iguanidae). *Mitt. Zool. Mus. Berlin* 46(1):197-234. (p. 225). Holotype: ZMB 41059. San Felipe, northeastern Jatibonico, Ciego de Ávila province; Sierra de Nipe, Holguín province; Yateras, Guantánamo province.

Anolis spectrum Peters, 1863. Peters, W. 1863. Über einige neue Arten des Saurier-Gattung *Anolis*. *Monatsb. Akad. Wiss. Berlin* 1863:135-149. (p. 136). Syntypes: ZMB 421 a-b. Pica Pica, Pinar del Río province; Fontanar, Ciudad de La Habana; North of Matanzas; Sierra de Trinidad; Jobo Rosado, Yaguajay, Sancti Spíritus province.

Anolis vanidicus vanidicus Garrido and Schwartz, 1972. Garrido, O. H. and Schwartz, A. 1972. The Cuban *Anolis spectrum* complex (Sauria: Iguanidae). *Proc. Biol. Soc. Washington* 85(45):509-522. (p. 515). Holotype: AMNH 78400. Macizo de Guamuhaya.

Anolis vanidicus rejectus Garrido and Schwartz, 1972. Garrido, O. H. and Schwartz, A. 1972. The Cuban *Anolis spectrum* complex (Sauria: Iguanidae). *Proc. Biol. Soc. Washington* 85(45):509-522. (p. 517). Holotype: ChM 55.1.63. Vicinity of Santiago de Cuba city.

Anolis fugitivus Garrido, 1975. Garrido, O. H. 1975. Distribución y variación del complejo *Anolis cyanopleurus* (Lacertilia: Iguanidae) en Cuba. *Poeyana* 143:1-60. (p. 28). Holotype: IZ 3854 = CZACC 4.3417. Nuevo Mundo, Moa, Holguín province.

- Anolis juangundlachi* Garrido, 1975. Garrido, O. H. 1975. Distribución y variación del complejo *Anolis cyanopleurus* (Lacertilia: Iguanidae) en Cuba. *Poeyana* 143:1-60. (p. 34). Holotype: IZ 3755. Carlos Rojas, Matanzas province.
- Anolis mimus* Schwartz and Thomas, 1975. Schwartz, A. and Thomas, R. 1975. A check-list of West Indian amphibians and reptiles. *Carnegie Mus. Nat. Hist. Special Publ.* 1:1-216. (p. 93). Holotype: IZ 3917 = CZACC 4.3411. Sierra de la Gran Piedra; Sierra de Santa María del Loreto.
- Anolis sagrei sagrei* Duméril and Bibron, 1837. Duméril, A. M. C. and Bibron, G. 1837. *Erpétologie Générale, ou Histoire Naturelle Complète des Reptiles*, 4. Librairie Encyclopédique de Roret, Paris, 572 pp. (p. 149). Syntypes MNHN 2430, 6797, ?MCZ 2171. Cuba; Archipiélago de los Colorados; Archipiélago de los Canarreos; Archipiélago de Sabana-Camagüey; Archipiélago de los Jardines de la Reina.
- Anolis sagrei greyi* Barbour, 1914. Barbour, T. 1914. A contribution to the zoogeography of the West Indies, with special reference to amphibians and reptiles. *Mem. Mus. Comp. Zool.* 44(2):209-359. (p. 287). Holotype: MCZ 7890. Vicinity of Camagüey city.
- Anolis bremeri bremeri* Barbour, 1914. Barbour, T. 1914. A contribution to the zoogeography of the West Indies, with special reference to amphibians and reptiles. *Mem. Mus. Comp. Zool.* 44(2):209-359. (p. 288). Holotype: MCZ 7889. From La Fe to Taco Taco, Pinar del Río province.
- Anolis bremeri insulaepinorum* Garrido, 1972. Garrido, O. H. 1972. *Anolis bremeri* Barbour (Lacertilia: Iguanidae) en el occidente de Cuba e Isla de Pinos. *Carib. J. Sci.* 12(1-2):59-77. (p. 63). Holotype: IZ 1626 = CZACC 4.3137. Isla de la Juventud, northern region.
- Anolis homolechis homolechis* (Cope), 1864. Cope, E. D. 1864. Contributions on the herpetology of tropical America. *Proc. Acad. Nat. Sci. Philadelphia* 16:166-181. (p. 169). Holotype: BMNH 1946.8.5.78. Cuba; Isla de la Juventud; Cayo Real; Cayo Cantiles.
- Anolis homolechis turquinensis* Garrido, 1973. Garrido, O. H. 1973. Distribución y variación de *Anolis homolechis* Cope (Lacertilia: Iguanidae) en Cuba. *Poeyana* 120:1-68. (p. 9). Holotype: IZ 2900 = CZACC 4.3271. Pico Turquino.
- Anolis quadriocellifer* Barbour and Ramsden, 1919. Barbour, T. and Ramsden, C. 1919. The herpetology of Cuba. *Mem. Mus. Comp. Zool.* 47(2):71-213. (p. 158). Holotype: MCZ 11867. Península de Guanahacabibes.
- Anolis jubar jubar* Schwartz, 1968. Schwartz, A. 1968. The Cuban lizards of the *Anolis homolechis* complex. *Tulane Studies Zool.* 14(4):140-184. (p. 157). Holotype: AMNH 96529. Northwesrtern Camagüey province.
- Anolis jubar cuneus* Schwartz, 1968. Schwartz, A. 1968. The Cuban lizards of the *Anolis homolechis* complex. *Tulane Studies Zool.* 14(4):140-184. (p. 158). Holotype: AMNH 96536. Playa Santa Lucía and Cayo Sabinal, Camagüey province.

Anolis jubar balaenarum Schwartz, 1968. Schwartz, A. 1968. The Cuban lizards of the *Anolis homolechis* complex. *Tulane Studies Zool.* 14(4):140-184. (p. 161). Holotype: AMNH 95975. Cayos Los Ballenatos, Bahía de Nuevitas, Camagüey province.

Anolis jubar oriens Schwartz, 1968. Schwartz, A. 1968. The Cuban lizards of the *Anolis homolechis* complex. *Tulane Studies Zool.* 14(4):140-184. (p. 162). Holotype: AMNH 95976. From Belic, Granma province, to west of Playa Juraguá, Santiago de Cuba province.

Anolis jubar yaguajayensis Garrido, 1973. Garrido, O. H. 1973. Distribución y variación de *Anolis homolechis* Cope (Lacertilia: Iguanidae) en Cuba. *Poeyana* 120:1-68. (p. 15). Holotype: IZ 2372 = CZACC 4.3288. From eastern Caibarién, Villa Clara province, to Punta Caguanes, Ciego de Ávila province.

Anolis jubar gibarensis Garrido, 1973. Garrido, O. H. 1973. Distribución y variación de *Anolis homolechis* Cope (Lacertilia: Iguanidae) en Cuba. *Poeyana* 120:1-68. (p. 23). Holotype: IZ 2387 = CZACC 4.3293. From Gibara to Nicaro, Holguín province.

Anolis jubar maisiensis Garrido, 1973. Garrido, O. H. 1973. Distribución y variación de *Anolis homolechis* Cope (Lacertilia: Iguanidae) en Cuba. *Poeyana* 120:1-68. (p. 28). Holotype: IZ 1524 = CZACC 4.3297. Maisí, Guantánamo province.

Anolis jubar albertschwartzii Garrido, 1973. Garrido, O. H. 1973. Distribución y variación de *Anolis homolechis* Cope (Lacertilia: Iguanidae) en Cuba. *Poeyana* 120:1-68. (p. 33). Holotype: IZ 2621 = CZACC 4.3312. From Guantánamo Bay to Loma de Macambo.

Anolis jubar santamariae Garrido, 1973. Garrido, O. H. 1973. Distribución y variación de *Anolis homolechis* Cope (Lacertilia: Iguanidae) en Cuba. *Poeyana* 120:1-68. (p. 43). Holotype: IZ 2643 = CZACC 4.3325. Cayo Santa María, Archipiélago de Sabana-Camagüey.

Anolis jubar cocoensis Estrada and Garrido, 1990. Estrada, A. R. and Garrido, O. H. 1990. Nueva subespecie de *Anolis jubar* (Lacertilia: Iguanidae) para Cayo Coco y la Loma de Cunagua, Ciego de Ávila, Cuba. *Rev. Biol.* 4(1):71-79. (p. 73). Holotype: MHNHCU-381. Cayo Coco; Loma de Cunagua, Camagüey province.

Anolis guafe Estrada and Garrido, 1991. Estrada, A. R. and Garrido, O. H. 1991. Dos nuevas especies de *Anolis* (Lacertilia: Iguanidae) de la región oriental de Cuba. *Carib. J. Sci.* 27(3-4):146-161. (p. 148). Holotype: CARE 60516 = CZACC 4.60516. Coastal terraces of Cabo Cruz, Granma province.

Anolis confusus Estrada and Garrido, 1991. Estrada, A. R. and Garrido, O. H. 1991. Dos nuevas especies de *Anolis* (Lacertilia: Iguanidae) de la región oriental de Cuba. *Carib. J. Sci.* 27(3-4):146-161. (p. 156). Holotype: CARE 60503 = CZACC 4.60503. Cabo Cruz, Granma province.

Anolis mestrei Barbour, 1916. Barbour, T. 1916. A new *Anolis* from Cuba. *Proc. Biol. Soc. Washington* 29:19-20. (p. 19). Holotype: MCZ 11285. From Isabel Rubio, Pinar del Río province, to Sierra de Anafe, La Habana province.

Anolis rubribarbus Barbour and Ramsden, 1919. Barbour, T. and Ramsden, C. 1919. The herpetology of Cuba. *Mem. Mus. Comp. Zool.* 47(2):71-213. (p. 156). Holotype: MCZ 11941. From Sagua de Tánamo to Moa, Holguín province.

Anolis allogus Barbour and Ramsden, 1919. Barbour, T. and Ramsden, C. 1919. The herpetology of Cuba. *Mem. Mus. Comp. Zool.* 47(2):71-213. (p. 159). Holotype: MCZ 8544. Cuba.

Anolis ahli Barbour, 1925. Barbour, T. 1925. A new Cuban *Anolis*. *Occ. Papers Boston Soc. Nat. Hist.* 5:167-168. (p. 168). Holotype: MCZ 19905. Sierra de Trinidad.

Anolis imias Ruibal and Williams, 1961. Ruibal, R. and Williams, E. E. 1961. The taxonomy of the *Anolis homolechis* complex of Cuba. *Bull. Mus. Comp. Zool.* 125(8):211-246. (p. 237). Holotype: MCZ 42556. South of Guantánamo province.

Anolis delafuentei Garrido, 1982. Garrido, O. H. 1982. Nueva especie de *Anolis* (Lacertilia, Iguanidae) para Cuba. *Doñana, Acta Vertebrata* 9:131-137. (p. 132). Holotype: IZ 5907 = CZACC 4.7173. Topes de Collantes.

Anolis birama Garrido, 1990. Garrido, O. H. 1990. Nueva especie de *Anolis* de la sección *Beta* (Lacertilia: Iguanidae) para Cuba. *Rev. Biol.* 4(2):157-162. (p. 158). Holotype: MCTH-4. Birama, Granma province.

Anolis ophiolepis Cope, 1861. Cope, E. D. 1861. Notes and descriptions of anoles. *Proc. Acad. Nat. Sci. Philadelphia* 13:208-215. (p. 211). Holotype: Unlocated. Cuba; Isla de la Juventud. *Chamaeleolis chamaeleonides* Duméril and Bibron, 1837. Duméril, A. M. C. and Bibron, G. 1837. *Erpétologie Générale, ou Histoire Naturelle Complète des Reptiles*. Librairie Encyclopédique de Roret, Paris, 4, 572 pp. (p. 168). Holotype: MNHN 1004. From Península de Guanahacabibes to Sierra de Cubitas, Camagüey province, and to Sierra Maestra; Isla de la Juventud.

Chamaeleolis porcus Cope, 1864. Cope, E. D. 1864. Contributions on the herpetology of tropical America. *Proc. Acad. Nat. Sci. Philadelphia* 16:166-181. (p. 168). Holotype: ANSP 8133. Eastern Cuba (from Holguín to Baracoa).

Chamaeleolis barbatus Garrido, 1982. Garrido, O. H. 1982. Descripción de una nueva especie cubana de *Chamaeleolis* (Lacertilia: Iguanidae), con notas sobre su comportamiento. *Poeyana* 236:1-25. (p. 3). Holotype: IZ 5368. Sierra del Rosario.

Chamaeleolis guamuhaya Garrido, Pérez-Beato, and Moreno, 1991. Garrido, O. H., Pérez-Beato, O., and Moreno, L. V. 1991. Nueva especie de *Chamaeleolis* (Lacertilia: Iguanidae) para Cuba. *Carib. J. Sci.* 27(3-4):162-168. (p. 163). Holotype: MHNHC-500. Macizo de Guamuhaya.

Family Teiidae

Ameiva auberi auberi Cocteau, 1838 or 1839. Cocteau, J. T. 1838 or 1839. *Reptiles. In Historia Física, Política y Natural de la Isla de Cuba* (R. de la Sagra, ed.), España, (p. 51). Syntypes: MNHN 1112, 2647, 1788, 4178. From La Fé, Pinar del Río province, to Canasí, La Habana province, on N and Melena del Sur, La Habana province, on S.

Ameiva auberi cacuminis Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 56). Holotype: AMNH 83028. Península de Guanahacabibes.

Ameiva auberi denticola Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 57). Holotype: AMNH 79202. From Carabelita to Manuel Lazo, Pinar del Río province.

Ameiva auberi procer Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 60). Holotype: AMNH 78390. Sierra de los Órganos; from Sierra del Rosario to Cabañas.

Ameiva auberi paulsoni Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 63). Holotype: AMNH 83012. La Coloma, Pinar del Río province.

Ameiva auberi pullata Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 67). Holotype: AMNH 82953. Matanzas; Cárdenas; Península de Hicacos; San Miguel de los Baños; all of them in Matanzas province.

Ameiva auberi abducta Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 70). Holotype: AMNH 96331. Península de Hicacos.

Ameiva auberi gemmea Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 71). Holotype: AMNH 82972. From Playa Gaguza to Baños de Elguea, Villa Clara province.

Ameiva auberi extraria Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 73). Holotype: AMNH 82982. Archipiélago de Sabana.

Ameiva auberi extorris Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 76). Holotype: IZ 52 = CZACC 4.3125. Cayuelo de la Vela (Archipiélago de Sabana).

Ameiva auberi garridoi Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 77). Holotype: IZ 96 = CZACC 4.3129. From Isabela de Sagua to Sagua la Grande, Villa Clara province.

Ameiva auberi atrothorax Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 79). Holotype: AMNH 78035. From eastern Cienfuegos to Casilda; Los Caneyes (Santa Clara).

Ameiva auberi peradusta Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 83). Holotype: IZ 85 = CZACC 4.3133. From Playa Girón to Juraguá; Calimete, Matanzas province.

Ameiva auberi llanensis Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 84). Holotype: MCZ 59321. From east of Sierra de Cubitas to Camagüey city.

. *Ameiva auberi sublesta* Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 87). Holotype: AMNH 96393. Cayo Sabinal.

Ameiva auberi citra Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 89). Holotype: AMNH 96375. Playa Santa Lucía, Camagüey province.

Ameiva auberi granti Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 91). Holotype: AMNH 83784. From Gibara, and Holguín, Holguín province, to Baracoa, Guantánamo province.

Ameiva auberi sabulicolor Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 94). Holotype: AMNH 83941. From Guantánamo city to Maisí, Guantánamo province.

Ameiva auberi ustulata Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 97). Holotype: AMNH 83778. From Santiago de Cuba Bay to

Guantánamo Bay; San Luis; El Cobre; and Palma Soriano, all of them in Santiago de Cuba province.

Ameiva auberi hardyi Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 100). Holotype: USNM 138468. From Cabo Cruz, Granma province, to Jutisí and Siboney, Santiago de Cuba province.

Ameiva auberi secta Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 102). Holotype: AMNH 82997. Isla de la Juventud, northern region.

Ameiva auberi marcida Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 195). Holotype: AMNH 82991. Isla de la Juventud, southern region.

Ameiva auberi zugi Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 107). Holotype: AMNH 83003. Archipiélago de los Canarreos; Bahía de Cochinos.

Ameiva auberi galbiceps Schwartz, 1970. Schwartz, A. 1970. A systematic review of *Ameiva auberi* Cocteau (Reptilia, Teiidae) in Cuba and in the Bahamas. I. The Cuban subspecies. *Ann. Carnegie Mus.* 4(4):45-117. (p. 111). Holotype: AMNH 78058. Archipiélago de los Jardines de la Reina.

Ameiva auberi orlandoi Schwartz and McCoy, 1975. Schwartz, A. and McCoy, C. J. 1975. A new name for a Cuban *Ameiva* (Reptilia: Teiidae). *Herpetologica* 31(2):240. (p. 240). Garrido, O. H. 1975. Nuevos reptiles del Archipiélago Cubano. *Poeyana* 141:1-58. (p. 37). Holotype: IZ 3427 = CZACC 4.3378. Archipiélago de Sabana-Camagüey.

Ameiva auberi sanfelipensis Garrido, 1975. Garrido, O. H. 1975. Nuevos reptiles del Archipiélago Cubano. *Poeyana* 141:1-58. (p. 45). Holotype: IZ 2987 = CZACC 4.3383. Cayo Real and Cayo Juan García, off the southern coast of Pinar del Río province.

Ameiva auberi nigriventris Gali and Garrido, 1987. Gali, F. and Garrido, O. H. 1987. Two new subspecies of *Ameiva auberi* (Reptilia: Teiidae) from Cuba. *Carib. J. Sci.* 22(3-4):165-173. (p. 169). Holotype: IZ 5778. From southeastern La Fe to northwestern Cayo Piedras, Isla de la Juventud.

Ameiva auberi schwartzi Gali and Garrido, 1987. Gali, F. and Garrido, O. H. 1987. (p. 166). Holotype: IZ 4389. From Mariel to Guajaibón River, La Habana province.

Family Tropiduridae

Leiocephalus carinatus carinatus Gray, 1827. Gray, J. E. 1827. A description of a new genus and some new species of saurian reptiles, with a revision of the species of chamaeleons. *Phil. Mag. Ann.* 2(2):207-209. (p. 208). Holotype: BMNH 1946.8.29,75. From northern coast of Pinar del Río province to Península de Hicacos.

Leiocephalus carinatus aquarius Schwartz and Ogren, 1956. Schwartz, A. and Ogren, L. 1956. A collection of reptiles and amphibians from Cuba, with the description of two new forms. *Herpetologica* 12(2):91-110. (p. 100). Holotype: ChM 55.1.62. From Cabo Cruz, Granma province, to Maisí, Guantánamo province.

Leiocephalus carinatus zayasi Schwartz, 1959. Schwartz, A. 1959. The Cuban lizards of the species *Leiocephalus carinatus* (Gray). *Reading Public Mus. Art Gallery* 10:1-47. (p. 9). Holotype: AMNH 77756. Península de Guanahacabibes.

Leiocephalus carinatus mogotensis Schwartz, 1959. Schwartz, A. 1959. The Cuban lizards of the species *Leiocephalus carinatus* (Gray). *Reading Public Mus. Art Gallery* 10:1-47. (p. 12). Holotype: AMNH 77755. Sierra de Viñales.

Leiocephalus carinatus labrossytus Schwartz, 1959. Schwartz, A. 1959. The Cuban lizards of the species *Leiocephalus carinatus* (Gray). *Reading Public Mus. Art Gallery* 10:1-47. (p. 33). Holotype: AMNH 77757. From Bahía de Cochinos to Casilda; Sierra de Trinidad; Cayo Conuco, Villa Clara province.

Leiocephalus carinatus cayensis Schwartz, 1959. Schwartz, A. 1959. The Cuban lizards of the species *Leiocephalus carinatus* (Gray). *Reading Public Mus. Art Gallery* 10:1-47. (p. 36). Holotype: AMNH 77758. Archipiélago de los Jardines de la Reina.

Leiocephalus carinatus microcyon Schwartz, 1959. Schwartz, A. 1959. The Cuban lizards of the species *Leiocephalus carinatus* (Gray). *Reading Public Mus. Art Gallery* 10:1-47. (p. 43). Holotype: AMNH 81271. Isla de la Juventud.

Leiocephalus cubensis cubensis Gray, 1840. Gray, J. E. 1840. Catalogue of the species of reptiles collected in Cuba by W. S. Mac Leay, Esq.; with some notes on their habits extracted from his *MS. Ann. Mag. Nat. Hist.* 1(5):108-115. (p. 110). Holotype: BMNH XXIII, 98a. From Artemisa, La Habana province, to Moa, Holguín province; Península de Zapata.

Leiocephalus cubensis paraphrus Schwartz, 1959. Schwartz, A. 1959. Variation in lizards of the *Leiocephalus cubensis* complex in Cuba and the Isla de Pinos. *Bull. Florida State Mus.* 4(4):97-143. (p. 111). Holotype: AMNH 78005. Archipiélago de los Jardines de la Reina.

Leiocephalus cubensis gigas Schwartz, 1959. Schwartz, A. 1959. Variation in lizards of the *Leiocephalus cubensis* complex in Cuba and the Isla de Pinos. *Bull. Florida State Mus.* 4(4):97-143. (p. 113). Holotype: AMNH 81056. Isla de la Juventud, southern region.

Leiocephalus cubensis pambasileus Schwartz, 1959. Schwartz, A. 1959. Variation in lizards of the *Leiocephalus cubensis* complex in Cuba and the Isla de Pinos. *Bull. Florida State Mus.* 4(4):97-143. (p. 118). Holotype: AMNH 81068. Campo and Hicacos Kays (Archipiélago de los Canarreos).

Leiocephalus cubensis minor Garrido, 1970. En Varona, L. S., and Garrido, O. H. 1970. Vertebrados de los Cayos de San Felipe, Cuba, incluyendo una nueva especie de jutía. *Poeyana* 75:1-26. (p. 18). Holotype: IZ 2754 = CZACC 4.3120. Cayo Juan García, Archipiélago de los Canarreos.

Leiocephalus macropus macropus Cope, 1862. Cope, E. D. 1862. Contributions to neotropical saurology. *Proc. Acad. Nat. Sci. Philadelphia* 14:176-188. (p. 184). Syntypes MCZ 10930, USNM 12254, 25819-23, 25825-26. From western Santiago de Cuba Bay to Cajobabo, Guantánamo province.

Leiocephalus macropus immaculatus Hardy, 1958. Hardy, J. D., Jr. 1958. A new lizard of the genus *Leiocephalus* from Cuba (Sauria: Iguanidae). *J. Washington Acad. Sci.* 48(9):294-300. (p. 294). Holotype: USNM 138412. From Magdalena River, Granma province, to western Santiago de Cuba Bay.

Leiocephalus macropus hoplites Zug, 1959. Zug, G. R. 1959. Three new subspecies of the lizard *Leiocephalus macropus* Cope from Cuba. *Proc. Biol. Soc. Washington* 72:139-150. (p. 140). Holotype: AMNH 78020. Loma de Cunagua, Camagüey province.

Leiocephalus macropus hyacinthurus Zug, 1959. Zug, G. R. 1959. Three new subspecies of the lizard *Leiocephalus macropus* Cope from Cuba. *Proc. Biol. Soc. Washington* 72:139-150. (p. 145). Holotype: AMNH 78015. Sierra de Trinidad.

Leiocephalus macropus koopmani Zug, 1959. Zug, G. R. 1959. Three new subspecies of the lizard *Leiocephalus macropus* Cope from Cuba. *Proc. Biol. Soc. Washington* 72:139-150. (p. 146). Holotype: MCZ 55541. Península de Guanahacabibes.

Leiocephalus macropus aegialus Schwartz and Garrido, 1967. Schwartz, A. and Garrido, O. H. 1967. A review of the Cuban iguanid lizard *Leiocephalus macropus* Cope. *Reading Public Mus. Art Gallery* 14:3-41. (p. 15). Holotype: AMNH 83255. Playa Santa Lucía, Camagüey province.

Leiocephalus macropus phylax Schwartz and Garrido, 1967. Schwartz, A. and Garrido, O. H. 1967. A review of the Cuban iguanid lizard *Leiocephalus macropus* Cope. *Reading Public Mus. Art Gallery* 14:3-41. (p. 17). Holotype: IZ 556 = CZACC 4.3064. From northern Cabo Cruz, Granma province, to Punta Hicacos, Granma province.

Leiocephalus macropus asbolomus Schwartz and Garrido, 1967. Schwartz, A. and Garrido, O. H. 1967. A review of the Cuban iguanid lizard *Leiocephalus macropus* Cope. *Reading Public Mus. Art Gallery* 14:3-41. (p. 30). Holotype: IZ 568 = CZACC 4.3095. From Banes, Holguín province, to Maisí, Guantánamo province.

Leiocephalus macropus lenticulatus Garrido, 1973. Garrido, O. H. 1973. Nuevas subespecies de reptiles para Cuba. *Torreia* 30:1-31. (p. 10). Holotype: IZ 2782 = CZACC 4.3247. Western Gibara, Holguín province.

Leiocephalus macropus felinoi Garrido, 1979. Garrido, O. H. 1979. Nuevas subespecies de *Leiocephalus macropus* Cope (Lacertilia: Iguanidae) para Cuba. *Poeyana* 188:1-16. (p. 2). Holotype: IZ 4751 = CZACC 4.3453. Bacunayagua, Matanzas province.

Leiocephalus macropus torrei Garrido, 1979. Garrido, O. H. 1979. Nuevas subespecies de *Leiocephalus macropus* Cope (Lacertilia: Iguanidae) para Cuba. *Poeyana* 188:1-16. (p. 7). Holotype: IZ 3725 = CZACC 4.3442. San Miguel de los Baños, Matanzas province.

Leiocephalus onaneyi Garrido, 1973. Garrido, O. H. 1973. Nueva especie de *Leiocephalus* (Lacertilia, Iguanidae) para Cuba. *Poeyana* 188:1-19 (p. 4). Holotype: IZ 2869 = CZACC 4.3267. Loma de Macambo, Guantánamo province.

Leiocephalus raviceps raviceps Cope, 1862. Cope, E. D. 1862. Contributions to neotropical saurology. *Proc. Acad. Nat. Sci. Philadelphia* 14:176-188. (p. 183). Syntypes ANSP 8601-03, MCZ 10928, USNM 4162. From western Guantánamo Bay to Cajobabo, Guantánamo province.

Leiocephalus raviceps uzelli Schwartz, 1960. Schwartz, A. 1960. Variation in the Cuban lizard *Leiocephalus raviceps* Cope. *Proc. Biol. Soc. Washington* 73:67-82. (p. 70). Holotype: AMNH 79321. From southeastern Santiago de Cuba Bay to Guantánamo Bay.

Leiocephalus raviceps klinikowskii Schwartz, 1960. Schwartz, A. 1960. Variation in the Cuban lizard *Leiocephalus raviceps* Cope. *Proc. Biol. Soc. Washington* 73:67-82. (p. 77). Holotype: AMNH 83326. Península de Hicacos, Matanzas province.

Leiocephalus raviceps jaumei Schwartz and Garrido, 1968. Schwartz, A. and Garrido, O. H. 1968. An undescribed subspecies of *Leiocephalus raviceps* Cope (Sauria: Iguanidae) from Western Cuba. *Proc. Biol. Soc. Washington* 81:23-30. (p. 24). Holotype: IZ 1349 = CZACC 4.3083. San Waldo, Cortés, Pinar del Río province.

Leiocephalus raviceps delavarai Garrido, 1973. Garrido, O. H. 1973. Nuevas subespecies de reptiles para Cuba. *Torreia* 30:1-31. (p. 4). Holotype: IZ 2774 = CZACC 4.3238. From Puerto Padre, Las Tunas province, to western Gibara, Holguín province.

Leiocephalus stictigaster stictigaster Schwartz, 1959. Schwartz, A. 1959. Variation in lizards of the *Leiocephalus cubensis* complex in Cuba and the Isla de Pinos. *Bull. Florida State Mus.* 4(4):97-143 (p. 121). Holotype: AMNH 77864. Península de Guanahacabibes.

Leiocephalus stictigaster sierrae Schwartz, 1959. Schwartz, A. 1959. Variation in lizards of the *Leiocephalus cubensis* complex in Cuba and the Isla de Pinos. *Bull. Florida State Mus.* 4(4):97-143. (p. 126). Holotype: AMNH 77813. From Manuel Lazo to Las Pozas; Sierra de los Órganos; Sierra del Rosario; all of them in Pinar del Río province.

Leiocephalus stictigaster exotheotus Schwartz, 1959. Schwartz, A. 1959. Variation in lizards of the *Leiocephalus cubensis* complex in Cuba and the Isla de Pinos. *Bull. Florida State Mus.* 4(4):97-143. (p. 130). Holotype: AMNH 81088. Isla de la Juventud, northern region.

Leiocephalus stictigaster astictus Schwartz, 1959. Schwartz, A. 1959. Variation in lizards of the *Leiocephalus cubensis* complex in Cuba and the Isla de Pinos. *Bull. Florida State Mus.* 4(4):97-143. (p. 134). Holotype: AMNH 81095. Isla de la Juventud southern region.

Leiocephalus stictigaster lucianus Schwartz, 1960. Schwartz, A. 1960. A new subspecies of *Leiocephalus stictigaster* Schwartz from Central Cuba. *Proc. Biol. Soc. Washington* 73:103-106. (p. 104). Holotype: AMNH 83583. From Nuevitas to Playa Santa Lucía, Camagüey province.

Leiocephalus stictigaster paraspheX Schwartz, 1964. Schwartz, A. 1964. New subspecies of *Leiocephalus* from Cuba. *Quart. J. Florida Acad. Sci.* 27(3):211-222. (p. 212). Holotype: AMNH 92153. Cayo Sabinal.

Leiocephalus stictigaster ophiplacodes Schwartz, 1964. Schwartz, A. 1964. New subspecies of *Leiocephalus* from Cuba. *Quart. J. Florida Acad. Sci.* 27(3):211-222. (p. 217). Holotype: AMNH 92771. Sierra de Cubitas, southern region, Camagüey province.

Leiocephalus stictigaster naranjoi Schwartz and Garrido, 1968. Schwartz, A. and Garrido, O. H. 1968. Four new subspecies of *Leiocephalus stictigaster* from Cuba. *Natl. Mus. Canada Nat. Hist. Papers* 37:1-23. (p. 3). Holotype: IZ 200 = CZACC 4.3445. Trinidad; Casilda.

Leiocephalus stictigaster lipomator Schwartz and Garrido, 1968. Schwartz, A. and Garrido, O. H. 1968. Four new subspecies of *Leiocephalus stictigaster* from Cuba. *Natl. Mus. Canada Nat. Hist. Papers* 37:1-23. (p. 11). Holotype: IZ 1230. Motel Los Caneyes, Santa Clara.

Leiocephalus stictigaster celestes Schwartz and Garrido, 1968. Schwartz, A. and Garrido, O. H. 1968. Four new subspecies of *Leiocephalus stictigaster* from Cuba. *Natl. Mus. Canada Nat. Hist. Papers* 37:1-23. (p. 14). Holotype: IZ 1182 = CZACC 4.3091. From Bueycitos to Jiguaní, Granma province.

Leiocephalus stictigaster gibarensis Schwartz and Garrido, 1968. Schwartz, A. and Garrido, O. H. 1968. Four new subspecies of *Leiocephalus stictigaster* from Cuba. *Natl. Mus. Canada Nat. Hist. Papers* 37:1-23. (p. 18). Holotype: IZ 1236 = CZACC 4.3095. From Malagueta Bay, Las Tunas province, to Punta de Mulas, Holguín province.

Leiocephalus stictigaster septentrionalis Garrido, 1975. Garrido, O. H. 1975. Nuevos reptiles del Archipiélago Cubano. *Poeyana* 141:1-58 (p. 28). Holotype: IZ 3425 = CZACC 4.3369. Archipiélago de Sabana-Camagüey.

Family Xantusiidae

Cricosaura typica Gundlach and Peters, in Peters, 1863. Peters, W. 1863. Hr. W. Peters berichtete über eine neue von Hrn. Dr. Gundlach auf Cuba entdeckte Sauriergattung, *Cricosaura typica* Gundlach et Peters. *Monatsb. Akad. Wiss. Berlin* 1863:362-368. (p. 362). Syntypes: ZMB 4832, 5071. Southern region of Granma and Santiago de Cuba provinces.

Order Squamata - Serpentes
Family Boidae

Epicrates angulifer Bibron, 1840. Bibron, G. 1840. Reptiles. In *Historia Física, Política y Natural de la Isla de Cuba* 4 (R. de la Sagra, ed.), España, 143 pp. (p. 129). Holotype: MNHN 3292. Cuba; Isla de la Juventud; Archipiélago de los Canarreos; Archipiélago de los Jardines de la Reina; Archipiélago de los Colorados; Archipiélago de Sabana-Camagüey.

Family Colubridae

Alsophis cantherigerus cantherigerus (Bibron), 1840. Bibron, G. 1840. Reptiles. In *Historia física, política y natural de la Isla de Cuba* 4 (R. de la Sagra, ed.), España, 143 pp. (p. 133). Syntypes: MNHN 3545-46, 3561-3563. From Pinar del Río to Villa Clara provinces; Isla de la Juventud; Archipiélago de Sabana-Camagüey; Archipiélago de los Canarreos; Cayos de San Felipe.

Alsophis cantherigerus adspersus (Gundlach and Peters) in Peters, 1865. Peters, W. 1865. Über einige neue Saugethiere (*Mormops*, *Macrotus*, *Vesperus*, *Molossus*, *Capromys*), Amphibien (*Platydactylus*, *Otocryptis*, *Euprepes*, *Ungalia*, *Dromicus*, *Tropidonotus*, *Xenodon*, *Hylodes*), und Fische (*Sillago*, *Sibastes*, *Channa*, *Myctophum*, *Carascus*, *Barbus*, *Capoeta*, *Poecilia*, *Saurenchelys*, *Leptocephalus*). *Monatsber. Akad. Wiss. Berlin* 1864:381-399. (p. 388). Syntypes ZMB 5064 a-b. From east of Guantánamo city to Maisí, Guantánamo province.

Alsophis cantherigerus pepei Schwartz and Thomas, 1960. Schwartz, A. and Thomas, R. 1960. Four new snakes (*Tropidophis*, *Dromicus*, *Alsophis*) from the Isla de Pinos and Cuba. *Herpetologica* 16(2):73-90 (p. 87). Holotype: AMNH 83639. From Mayarí to La Mata, Guantánamo province.

Alsophis cantherigerus schwartzi Lando and Williams, 1969. Lando, R. V. and Williams, E. E. 1969. Notes on the herpetology of the U. S. Naval Base at Guantanomo Bay, Cuba. *Studies Fauna Curaçao Carib. Isl.* 31(116):159-201. (p. 192). Holotype: AMNH 83638. From Trinidad, Sancti Spíritus province, to Felicidad, Guantánamo province.

Antillophis andrei andrei (Reinhardt and Lütken), 1862. Reinhardt, J. and Lütken, C. F. 1862. Bidrag til det vestindiske Öeriges og navnlingen til de dansk-vestindiske Öers Herpetologie. *Vidensk. Meddel. Naturhist. Foren Kjøbenhavn for 1862* (10-18):153-291. (p. 214). Holotype: UZM R60766-R60767. From Pinar del Río to Camagüey provinces.

Antillophis andrei nebulatus (Barbour), 1916. Barbour, T. 1916. The reptiles and amphibians of the Isle of Pines. *Ann. Carnegie Mus.* 10(2):297-308. (p. 305). Holotype: MCZ 11092. Isla de la Juventud.

Antillophis andrei orientalis (Barbour and Ramsden), 1919. Barbour, T. and Ramsden, C. 1919.

The herpetology of Cuba. *Mem. Mus. Comp. Zool.* 47(2):71-213. (p. 196). Holotype: MCZ 11726. Eastern Cuba.

Antillophis andrei peninsulae (Schwartz and Thomas), 1960. Schwartz, A. and Thomas, R. 1960.

Four new snakes (*Tropidophis*, *Dromicus*, *Alsophis*) from the Isla de Pinos and Cuba.

Herpetologica 16(2):73-90. (p. 81). Holotype: AMNH 83235. Península de Guanahacabibes.

Antillophis andrei melopyrrha (Thomas and Garrido), 1967. Thomas, R. and Garrido, O. H. 1967.

A new subspecies of *Dromicus andreae* (Serpentes: Colubridae). *Ann. Carnegie Mus.*

39(16):219-226. (p. 219). Holotype: IZ 1080 = CZACC 4.3071. Cayo Cantiles, Archipiélago de los Canarreos.

Antillophis andrei morenoi Garrido, 1973. Garrido, O. H. 1973. Nuevas subespecies de reptiles

para Cuba. *Torreia* 30:1-31. (p. 18). Holotype: IZ 2737. Cayo Santa María, Archipiélago de Sabana-Camagüey.

Arrhyton taeniatum Günther, 1858. Günther, A. 1858. *Catalogue of colubrine snakes in the*

collection of the British Museum. Taylor and Francis, Londres, xvi + 281 pp. (p. 244).

Holotype: BMNH 1946.1.21.48. Cuba; Isla de la Juventud.

Arrhyton vittatum (Gundlach and Peters), 1862. Gundlach, J. C. and Peters, W. 1862. Hrn. W.

Peters legta eine mittheilung vor über zwei neue von Hrn. Dr. Gundlach auf Cuba entdeckte schlangen. *Tropidonotus cubanus* and *Criptodacus vittatus*. *Monatsb. Akad. Wiss. Berlin* 1861:1001-1004. (p. 1003). Holotype: ZMB 4096. Cuba; Isla de la Juventud.

Arrhyton dolichura Werner, 1909. Werner, F. 1909. Über neue oder seltense Reptilien des

Naturhistorischen Museums in Hamburg. I. Schlangen. *Mitt. Naturhist. Mus. Hamburg* 26:205-247. (p. 224). Holotype: HZM destroyed. Western Ciudad de La Habana province.

Arrhyton landoi Schwartz, 1965. Schwartz, A. 1965. A review of the colubrid snake genus

Arrhyton with a description of a new subspecies from southern Oriente province, Cuba. *Proc. Biol. Soc. Washington* 78:99-114. (p. 109). Holotype: MCZ 42505. From Pilón, Granma province, to Imías, Guantánamo province.

Arrhyton ainictum Schwartz and Garrido, 1981. Schwartz, A. and Garrido, O. H. 1981. A review of

the Cuban members of the genus *Arrhyton* (Reptilia, Serpentes, Colubridae). *Ann. Carnegie Mus.* 50(7):207-230. (p. 216). Holotype: IZ 4256. Amancio Rodríguez, Las Tunas province.

Arrhyton tanyplectum Schwartz and Garrido, 1981. Schwartz, A. and Garrido, O. H. 1981. A review

of the Cuban members of the genus *Arrhyton* (Reptilia, Serpentes, Colubridae). *Ann. Carnegie Mus.* 50(7):207-230. (p. 221). Holotype: AMNH 77782. From San Vicente to Pan de Azúcar,

Pinar del Río, province.

Arrhyton supernum Hedges and Garrido, 1992. Hedges, S. B. and Garrido, O. H. 1992. Cuban snakes of the genus *Arrhyton*: two new species and a reconsideration of *A. redimitum* Cope. *Herpetologica* 48(2):168-177. (p. 171). Holotype: MHNHCU-2704. Yunque de Baracoa and Monte Líbano, Guantánamo province.

Arrhyton procerum Hedges and Garrido, 1992. Hedges, S. B. and Garrido, O. H. 1992. Cuban snakes of the genus *Arrhyton*: two new species and a reconsideration of *A. redimitum* Cope. *Herpetologica* 48(2):168-177. (p. 174). Holotype: MHNHCU-3285. Eastsoutheastern Playa Girón.

Tretanorhinus variabilis variabilis Duméril, Bibron, and Duméril, 1854. Duméril, A. M. C., Bibron, G., and Duméril, A. H. A. 1854. *Erpétologie Générale, ou Histoire Naturelle Complète des Reptiles*. Librairie Encyclopédique de Roret, Paris 7(1):xiii + 780 pp. (p. 349). Syntypes: MHN 7161, 7346. From Laguna de Ariguanabo, La Habana province, to Guantánamo province.

Tretanorhinus variabilis wagleri (Jan), 1863. Jan, 1863. *Elenco sistematico degli Ofidi descritti e disegnati per l'Iconografia Generale* A. Lombardi, Milán, vii + 143 pp. Enumerazione sistematica degli ofidi appartenenti al gruppo Coronellidae. *Arch. Zool. Anat. Fisiol.* 2(2):215-330. (p. 284). Holotype: Unlocated. From Vallecito de San Juan to San Diego de los Baños; Sierra de los Órganos; Sierra del Rosario; all of them in Pinar del Río province.

Tretanorhinus variabilis insulaepinorum Barbour, 1916. Barbour, T. 1916. The reptiles and amphibians of the Isle of Pines. *Ann. Carnegie Mus.* 10(2):297-308. (p. 306). Holotype: MCZ 11190 or CM 311. Isla de la Juventud.

Tretanorhinus variabilis binghami Schwartz and Ogren, 1956. Schwartz, A. and Ogren, L. 1956. A collection of reptiles and amphibians from Cuba, with the description of two new forms. *Herpetologica* 12(2):91-110. (p. 105). Holotype: ChM 55.1.61. From Manzanillo, Granma province, to Río Yao, Guantánamo province.

Family Natricidae

Nerodia clarkii compressicauda (Kenicott), 1860. Kenicott, 1860. *Proc. Acad. Nat. Sci. Philadelphia* 12. Holotype: USNM 1348. Northern coast, from Punta Brava, La Habana province, to Playa Santa Lucía, Camagüey province; Cayo Las Brujas, Villa Clara province.

Family Tropidophiidae

Tropidophis maculatus (Bibron), 1840. Bibron, G. 1840. Reptiles. En *Historia Física, Política y Natural de la Isla de Cuba* 4 (R. de la Sagra, ed.), España, 143 pp. (p. 127). Holotype: MNHN 7184. From Guane, Pinar del Río province, to Cárdenas, Matanzas province; Isla de la Juventud.

Tropidophis pardalis (Gundlach), 1840. Gundlach, J. C. 1840. Ueber zwei von mir gesammelte Boen (*Boa pardalis*). *Arch. Naturges.* 6(1):359-361. (p. 361). Holotype: Unlocated. Cuba; Isla de la Juventud; Cayo Paredón Grande, Archipiélago de Sabana-Camagüey.

Tropidophis haetianus haetianus (Cope), 1879. Cope, E. D. 1879. Eleventh contribution to the herpetology of tropical America. *Proc. Amer. Phil. Soc.* 18:261-277. (p. 273). Syntypes USNM 10164, 10169. Guardalavaca, Holguín province.

Tropidophis melanurus melanurus (Schlegel), 1837. Schlegel, H. 1837. *Essai sur la physionomie des serpens*. Vol. 2 xv + 606 pp. M. H. Schonekat, Amsterdam (p. 389) + *Atlas*, 21 pls. J. Kips, J. Hz. and W. P. Van Stockum, La Haya. Holotype: ANSP 10308. Cuba.

Tropidophis melanurus ericksoni Schwartz and Thomas, 1960. Schwartz, A. and Thomas, R. 1960. Four new snakes (*Tropidophis*, *Dromicus*, *Alsophis*) from the Isla de Pinos and Cuba. *Herpetologica* 16(2):73-90. (p. 74). Holotype: AMNH 82897. Isla de la Juventud.

Tropidophis melanurus dysodes Schwartz and Thomas, 1960. Schwartz, A. and Thomas, R. 1960. Four new snakes (*Tropidophis*, *Dromicus*, *Alsophis*) from the Isla de Pinos and Cuba. *Herpetologica* 16(2):73-90. (p. 79). Holotype: AMNH 82893. La Coloma, Pinar del Río province.

Tropidophis semicinctus (Gundlach and Peters), 1865, in Peters, 1865. Peters, W. 1865. Über einige neue Saugethiere (*Mormops*, *Macrotus*, *Vesperus*, *Molossus*, *Capromys*), Amphibien (*Platydactylus*, *Otocryptis*, *Euprepes*, *Ungalia*, *Dromicus*, *Tropidonotus*, *Xenodon*, *Hylodes*), und Fische (*Sillago*, *Sibastes*, *Channa*, *Myctophum*, *Carascus*, *Barbus*, *Capoeta*, *Poecilia*, *Saurenchelys*, *Leptocephalus*). *Monatsber. Akad. Wiss. Berlin* 1864:381-399. (p. 388). Syntype: ZMB 5076. From Rancho Mundito, Pinar del Río province, to northwestern Trinidad.

Tropidophis pilsbryi pilsbryi Bailey, 1937. Bailey, J. R. 1937. A review of some recent *Tropidophis* material. *Proc. New England Zool. Club* 16:41-52. (p. 42). Holotype: ANSP 20822. Miranda, Santiago de Cuba province; Guantánamo.

Tropidophis pilsbryi galacelidus Schwartz and Garrido, 1975. Schwartz, A. and Garrido, O. H. 1975. A reconsideration of some Cuban *Tropidophis* (Serpentes, Boidae). *Proc. Biol. Soc. Washington* 88(9):77-90. (p. 81). Holotype: IZ 4052. From Soledad, Cienfuegos province, to Trinidad.

Tropidophis nigriventris nigriventris Bailey, 1937. Bailey, J. R. 1937. A review of some recent *Tropidophis* material. *Proc. New England Zool. Club* 16:41-52. (p. 45). Holotype: UMMZ 70888. Eastern Camagüey province.

Tropidophis nigriventris hardyi Schwartz and Garrido, 1975. Schwartz, A. and Garrido, O. H. 1975. A reconsideration of some Cuban *Tropidophis* (Serpentes, Boidae). *Proc. Biol. Soc. Washington* 88(9):77-90. (p. 86). Holotype: USNM 138510. From Soledad, Cienfuegos province, to Trinidad.

Tropidophis wrighti Stull, 1938. Stull, O. G. 1938. A revision of the genus *Tropidophis*. *Occ. Papers Mus. Zool. Univ. Michigan* 195:1-49. (p. 38). Holotype: USNM 12420. From Céspedes, Camagüey province, to Santiago de Cuba province.

Tropidophis feicki Schwartz, 1957. Schwartz, A. 1957. A new species of boa (genus *Tropidophis*) from western Cuba. *Amer. Mus. Novitates* 1839:1-8. (p. 3). Holotype: AMNH 76224. From Guane, Pinar del Río province, to Pan de Matanzas.

Tropidophis fuscus Hedges and Garrido, 1992. Hedges, S. B. and Garrido, O. H. 1992. A new species of *Tropidophis* from Cuba (Serpentes: Tropidophidae). *Copeia* 3:820-825. (p. 820). Holotype: MNHNCU-2705. Yateras and Camarones, Guantánamo province.

Family Typhlopidae

Typhlops lumbricalis (Linnaeus), 1758. Linnaeus, K. von, 1758. *Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Tomus I, Editio Decima, reformata*, 824 pp. (p. 288). Holotype: Unlocated. Cuba; Isla de la Juventud.

Typhlops biminiensis biminiensis Richmond, 1955. Richmond, N. D. 1955. The blind snakes (*Typhlops*) of Bimini, Bahama Islands, British West Indies, with description of a new species. *Amer. Mus. Novitates* 1734:1-7 (p. 2). Holotype: CM 32604. Miramar, Ciudad de La Habana; Santa Cruz del Norte, La Habana province; Rancho Luna, Cienfuegos province; Holguín; Santiago de Cuba; Maisí, Guantánamo province.

LITERATURE CITED

- Díaz, L. M., A. R. Estrada, and L. V. Moreno (1996): A new species of *Anolis* (Sauria: Iguanidae) from the Sierra de Trinidad, Sancti Spíritus, Cuba. *Carib. J. Sci.* 32(1):54-58.
- Hass, C. A., S. B. Hedges, and L. R. Maxson (1993): Molecular insights into the relationships and biogeography of West Indian anoline lizards. *Biochem. Syst. Ecol.* 21(1):97-114.
- Hedges, S. B. (1996): The origin of West Indian amphibians and reptiles. In *Contributions to West Indian herpetology: a tribute to Albert Schwartz* (R. Powell and R. W. Henderson, eds.). Society for the Study of Amphibians and Reptiles, Ithaca (New York). *Contributions to Herpetology* vol. 12, pp. 95-128.
- Pérez-Beato, O. 1996-1997. A new subspecies of *Anolis porcatus* (Sauria: Polychrotidae) from Western Cuba. *Rev. Biol. Trop.* 44(3)/45(1):295-299.
- Powell, R., R. W. Henderson, K. Adler, and H. A. Dundee (1996): An annotated checklist of West Indian Amphibians and Reptiles. In *Contributions to West Indian Herpetology: A tribute to Albert Schwartz* (R. Powell and R. W. Henderson, eds.). Society for the Study of Amphibians and Reptiles, Ithaca (New York). *Contributions to Herpetology* vol. 12, pp. 51-93 + 8 color pls.
- Thomas, R., and S. B. Hedges (1998a): A new amphisbaenian from Cuba. *J. Herpetol.* 32(1):92-96.
- Thomas, R., and S. B. Hedges (1998b): New anguid lizard (*Diploglossus*) from Cuba. *Copeia* 1998(1):97-103.
- Thomas, R., S. B. Hedges, and O. H. Garrido (1998): A new gecko (*Sphaerodactylus*) from the Sierra Maestra of Cuba. *J. Herpetol.* 32(1):66-69.