

**Large-scale morphological changes and sediment budget of the Western Scheldt estuary  
1955–2020**

**the impact of large-scale sediment management**

Elias, Edwin P.L.; van der Spek, Ad J.F.; Wang, Zheng Bing ; Cleveringa, Jelmer; Jeuken, Claire J. L.; Taal, Marcel; van der Werf, Jebbe

**DOI**

[10.1017/njg.2023.11](https://doi.org/10.1017/njg.2023.11)

**Publication date**

2023

**Document Version**

Final published version

**Published in**

Netherlands Journal of Geosciences: Geologie en Mijnbouw

**Citation (APA)**

Elias, E. P. L., van der Spek, A. J. F., Wang, Z. B., Cleveringa, J., Jeuken, C. J. L., Taal, M., & van der Werf, J. (2023). Large-scale morphological changes and sediment budget of the Western Scheldt estuary 1955–2020: the impact of large-scale sediment management. *Netherlands Journal of Geosciences: Geologie en Mijnbouw*, 102(1-3), Article 11. <https://doi.org/10.1017/njg.2023.11>

**Important note**

To cite this publication, please use the final published version (if applicable).  
Please check the document version above.

**Copyright**

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

**Takedown policy**

Please contact us and provide details if you believe this document breaches copyrights.  
We will remove access to the work immediately and investigate your claim.

## Review

**Cite this article:** Elias EPL, Van der Spek AJF, Wang ZB, Cleveringa J, Jeuken CJL, Taal M, and Van der Werf JJ. Large-scale morphological changes and sediment budget of the Western Scheldt estuary 1955–2020: the impact of large-scale sediment management. *Netherlands Journal of Geosciences*, Volume 102, e12. <https://doi.org/10.1017/njg.2023.11>

Received: 5 May 2023

Revised: 23 August 2023

Accepted: 6 September 2023

### Keywords:






Western Scheldt; sediment budget; coastal morphodynamics; impact of large-scale engineering

### Corresponding author:

Edwin P. L. Elias;

Email: [edwin.elias@deltares.nl](mailto:edwin.elias@deltares.nl)

# Large-scale morphological changes and sediment budget of the Western Scheldt estuary 1955–2020: the impact of large-scale sediment management

Edwin P. L. Elias<sup>1</sup> , Ad J. F. Van der Spek<sup>1,2</sup> , Zheng Bing Wang<sup>1,3</sup> ,  
Jelmer Cleveringa<sup>2,4</sup>, Claire J. L. Jeuken<sup>1</sup>, Marcel Taal<sup>1</sup>  and Jebbe J. Van der Werf<sup>1,5</sup> 

<sup>1</sup>Unit of Marine and Coastal Systems, Deltares, Delft, the Netherlands; <sup>2</sup>Faculty of Geosciences, Utrecht University, Utrecht, the Netherlands; <sup>3</sup>Faculty of Civil Engineering and Geosciences, Delft University of Technology, Delft, the Netherlands; <sup>4</sup>Arcadis Nederland, Zwolle, the Netherlands and <sup>5</sup>Department of Water Engineering & Management, University of Twente, Enschede, the Netherlands

## Abstract

In the Western Scheldt estuary, like in many estuaries, safe navigation, flood protection, and ecological targets require a balanced and sustainable sediment management. A thorough understanding of the morphodynamic functioning of the estuary and its response to changes in hydrodynamics (natural sediment transport) and large-scale interventions is imperative. This paper presents a detailed overview of over 65 years of morphological changes and a comprehensive sediment budget of the Western Scheldt estuary that is based on analysis of a unique series of frequent bathymetric surveys and historical data on human–sediment interactions of dredging, dredge disposal and sand mining. Solving the sediment budget reveals an annual sediment import of 2.2 million m<sup>3</sup>. The highest sediment imports occurred between 1980–1994 and 2005–2020 (2.9 and 3.7 million m<sup>3</sup>/year). A 1.3 million m<sup>3</sup>/year net export prevailed between 1994 and 2005. Natural variations in the hydrodynamics (e.g., tidal asymmetry and amplification) and sediment transports cannot explain the derived temporal variations in sediment import rates. Anthropogenic activities play a dominant role. Capital dredging of the main navigation channel has led to channel deepening and significantly increased dredge and disposal volumes. Disposal on tidal flats and in the secondary channel leads to a long-term storage of sand and, consequently, a local increase in bed level and a sand deficit in the remainder of the system that results in increased sediment imports. Large-scale disposal in the western part of the estuary can (temporarily) reverse the sediment exchange between the estuary and its mouth area, as observed between 1994 and 2005. Apparently, large-scale anthropogenic reallocation of sediment by dredging and/or disposal as part of navigation channel improvement, sand mining or nourishment essentially influences the morphological development of the Western Scheldt estuary. This reveals responsibilities as well as opportunities of sediment management for the Western Scheldt and similar estuaries worldwide.

## Introduction and objective


The Western Scheldt (Fig. 1) is the only remaining open natural estuary of the Delta coast in the SW Netherlands, a series of (former) distributaries of the rivers Rhine, Meuse and Scheldt. The estuaries of the Eastern Scheldt, Grevelingen and Haringvliet were entirely or partially closed as part of the Delta Project, a flood protection scheme built between 1954 and 2010 following the storm surge disaster of 1953 (Watson & Finkl, 1990, 1992). The damming of the estuaries led to significant morphodynamic changes that are still noticeable today (Elias et al., 2016; Van der Spek & Elias, 2021).

Like in many estuaries, in the Western Scheldt, man and nature compete for the same natural resources (De Vriend et al., 2011; Barbier et al., 2011; Biguino et al., 2023). The governments of the Netherlands and the Belgian state of Flanders, by treaty, manage the Scheldt estuary jointly. The often conflicting economic and environmental interests make sustainable management of these systems a complex and non-trivial task, especially if it crosses country borders and multiple governments are involved (Graveland et al., 2002; Debergh et al., 2009).

The Western Scheldt remained an open estuary due to the presence of essential shipping lanes to the harbours of Vlissingen, Terneuzen and Antwerp (Fig. 1). In 2019 alone, almost 15,000 seagoing vessels sailed through the estuary to the port of Antwerp (Plancke et al., 2022). Shipping requires a nautically safe, morphologically stable navigation channel of sufficient depth and width. The ongoing maintenance of the shipping channel, however, may conflict with the need for sufficient estuarine habitat. Such habitat is essential to maintain the estuary's biodiversity and associated high ecological value and to fulfill Natura 2000 criteria (Bastmeijer,

© The Author(s), 2023. Published by Cambridge University Press on behalf of the Netherlands Journal of Geosciences Foundation. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.





**Figure 1.** Overview of (A) the (former) estuaries that form the Delta region of the SW Netherlands. The contiguous ebb-tidal deltas of the (former) distributaries of the rivers Rhine, Meuse and Scheldt are together named Voordelta (fore delta) and (B) the main channels and shoals in the Western Scheldt estuary. The underlying DEM of the estuary is based on 2021 survey data. Depths are in meters relative to the Dutch ordnance datum NAP (Normaal Amsterdams Peil), which is about present-day mean sea level. Note that in this figure the tidal channels and shoals are indicated by their Dutch names. The majority of these names end with words that are easily recognised and translated: 'vlakte', 'plaat' or 'platen' = shoal(s); 'geul' or 'pas' = channel; 'gat' or 'put' = deep channel / scour hole; 'nauw' = narrow often deep channel.



2016). To compensate for habitat losses as a result of channel deepening, already in 2004, plans were presented to restore intertidal habitat by depoldering the Hedwigepolder (Soresma, 2013), and in 2022, the removal of the dykes finally started.

From a morphodynamic viewpoint, the net sediment import or sediment export rates are a vital indicator of the long-term “health” of estuarine and tidal-inlet systems. Excessive sediment export can lead to the loss of shoals and intertidal habitat, whereas sediment import can lead to channel deposition, hamper navigability and increase the need for maintenance dredging. Additionally, sand import into the estuary is considered a loss from the coastal zone and under the present Dutch coastal policy needs to be compensated through nourishments (Mulder et al., 2011; Lodder & Slinger, 2022). Understanding the sediment import and export rates is therefore essential for an effective and sustainable management of the estuary and the adjoining coastal zone.

It is well known that anthropogenic sediment interventions, such as sand mining, dredging and dredge relocation (disposal), directly impacted the hydrodynamics and morphology of the Western Scheldt (Wang et al., 2002; Taal et al., 2013; Vandenbruwaene et al., 2019, 2020), causing subsequent changes in the sediment transport dynamics (Wang & Winterwerp, 2001; Jeuken, 2000; Jeuken & Wang, 2010; Schrijver, 2021). However, the sediment exchange between the estuary and the mouth area and North-Sea coast is still insufficiently understood, with some studies reporting sediment import and others sediment export (Bakker & De Looff, 1977; Nederbragt & Liek, 2004; Haecon, 2006; Cleveringa, 2013; Dam & Cleveringa, 2013; Schrijver, 2021). The specific research goals of this study are, therefore:

1. to quantify the sediment exchange between the Western Scheldt estuary and the mouth area, and
2. to better understand the sediment budget changes of the Western Scheldt estuary and its response to changes in hydrodynamics, natural and anthropogenic sediment transport (due to sand mining and dredging and disposal strategies).

To achieve these goals, we summarise the morphodynamic evolution of the Western Scheldt estuary, analyse changes in basin hydrodynamics, and construct and analyse a comprehensive sediment budget of the estuary over the period 1955–2020. The sediment budget analysis uses detailed bathymetric surveys executed by Rijkswaterstaat, the water management authority of The Netherlands. Long-term datasets comprising frequent observations of anthropogenically influenced morphodynamic changes in estuaries are generally scarce. Therefore, an additional goal of this paper is to share this dataset with the international audience. Moreover, this study aims to highlight and summarise the body of reports on the Western Scheldt written in Dutch.

## Study area

### General characteristics

The Scheldt estuary and river are 350 km long and extend through the Netherlands, Belgium and France (Fig. 1). Tidal influence extends roughly 160 km upstream to the weir at Gentbrugge. Upstream of this weir, the system is strictly fluvial. The fluvial channel branches out into two main tributaries and extends into France. The total catchment area equals 21,000 km<sup>2</sup>. The funnel-shaped estuary seaward of the weir displays three morphologically distinct systems:

1. *Single-channel system*: The upstream reach between Gentbrugge and the Flemish-Dutch border north of Antwerp is

called Sea Scheldt or ‘Zeeschelde’ in Dutch and consists of a large single main channel and mudbanks along its embankments. In 2010, the average bed level of the Sea Scheldt was around NAP –12 m (Verheyen & Van Holland, 2013; Van Dam, 2014). The highest mean tidal range reaches a maximum of nearly 5.5 m near Schelle just south of Antwerp, but decreases significantly upstream, it is entirely blocked by the weir near Gentbrugge, see Van Braeckel et al. (2006) and Van Holland (2013) for detailed descriptions of this portion of the estuary.


2. *Multi-channel system*: The 95 km long estuary downstream of the Belgian-Dutch border is called the Western Scheldt (Fig. 1B). The width of the estuary varies between 2 and 8 km and channel depths generally range between 10 and 50 m, but locally scour holes up to 65 m in depth exist. The Western Scheldt exhibits multiple (2–3) parallel aligned channels, separated by shoal complexes and connected by smaller scale connecting channels (e.g., Van Veen et al., 2005; Van den Berg et al., 1996; Jeuken, 2000; Toffolon & Crosato, 2007; Van Dijk et al., 2021); see Fig. 1B for the present-day configuration of channels and shoals. The meandering ebb-channel is the main channel, whereas the parallel subordinate channels are flood channels. The ebb and flood channels tend to join in subtidal shoal areas. The general channel and shoal patterns have been described by Winterwerp et al. (2001) as a series of macro- and mesocells (Fig. 2). These cells are based on morphological characteristics and associated tide-averaged sediment transport patterns (Jeuken, 2000).

3. *The mouth area of the Western Scheldt*, seaward of Vlissingen, stretches from the island of Walcheren in the north into Belgian territorial waters in the south (Van Enckevoort, 1996). *The mouth area* is often referred to as an ebb-tidal delta. However, this is strictly speaking incorrect since most of the area lacks the typical morphodynamics of an ebb-tidal delta, namely, sand supply by ebb currents followed by redistribution by waves. Therefore, we use the non-specific description ‘mouth area’ consisting of the main channel Wielingen, the extensive subtidal shoal area Vlakte van de Raan to its north, and a dynamic channel-and-shoal area offshore southwest Walcheren (Fig. 1A). The southern morphological boundary is not clearly defined as the coastal zone merges into the Flemish Banks, consisting of shallow, shore-parallel sand ridges separated by well-developed troughs (Van Cauwenberghe, 1971). The transect from Vlissingen to Breskens is often used to distinguish between the mouth area and the estuary. However, Vandenbruwaene et al., (2013) indicate that this transition occurs 2 km seaward of this cross-section based on estuary width.

## Processes shaping the estuary

### Hydrodynamics

In the mouth area, waves and tides are the dominant forcing processes for flow and sediment transport. Waves drive sediment transports directly via currents due to radiation stress gradients generated by wave breaking of obliquely incident waves and due to wave asymmetry. Indirectly, waves enhance bed-shear stresses and stir-up sediment, allowing more sediment into suspension to be transported by the tidal and wind-driven flow. The wave climate is mild and mainly consists of locally, in the shallow North Sea basin, generated waves. Offshore, the annual mean significant wave height is around 1.3 m from the west-southwest, with a corresponding mean wave period of 5 s (Wijnberg, 2002). However, during storms, wind-generated wave heights occasionally exceed 6 m, and associated water-level surges of more than 2 m may occur. For example, the record surge height was NAP + 4.55 m



**Figure 2.** Overview of the macrocells in the Western Scheldt as used in this study. Macrocells (Mc) 1, 3, 4, 5, 6, and 7 are subdivided in part a, the main channel, and part b, the secondary channels and shoals. In the original definition by Winterwerp et al. (2001), Macrocell 2 is referred to as Mesocell 2.

in Vlissingen during the storm surge flooding of 1 February 1953. Wave-driven sediment transport is more important in the mouth area compared to the estuary, as the large and shallow Vlake van de Raan shoal effectively shelters the estuary from the waves. At wave buoy Deurloo (see Fig. 1A for location), the year-averaged significant wave height varies between 0.7 and 0.9 m. In the estuary, at stations Hansweert and Bath, year-averaged significant wave heights range between 0.2 and 0.3 m (Barneveld et al., 2018; Nederhoff, 2016).

Tides, forced by the semi-diurnal (progressive) tide in the North Sea, dominate the water motion and sediment transport in the estuary. The North Sea tide travels south to north along the Dutch coast and west to east into the estuary. The tidal wave amplifies and distorts as it travels up the estuary; the mean tidal range increases from 3.8 m at Vlissingen to 5.2 m at Antwerp, 78 km upstream (Jeuken, 2000). The phase shift between the horizontal and vertical tide increases by roughly 2.5–3 hours from the mouth to Antwerp. The associated flood volumes reduce in a landward direction from an average of 1100 million m<sup>3</sup> per tide at Vlissingen to 70 million m<sup>3</sup> per tide at Antwerp (Gerritsen & De Jong, 1983; Claessens, 1988; Baeyens et al., 1997). Peak tidal velocities range between 1–1.5 m/s, with maximum flood velocities consistently exceeding the maximum ebb velocities. Based on the large tidal range and low wave energy, the inlet classifies as tide-dominated (Davis & Hayes, 1984).

Fresh water inflows from the river Scheldt, tributaries and local drainage are small (<1% of the tidal prism) and average around 110 m<sup>3</sup>/s. Inflow is lowest in summer between 50–60 m<sup>3</sup>/s, and increases to 160–180 m<sup>3</sup>/s in winter (Ysebaert et al., 1993; Van Maldegem et al., 1993). Vandenbruwaene et al. (2013) state that for a typical dry event during summer, the discharge can be as low as 34 m<sup>3</sup>/s, whereas, for typical flushing events, mostly during winter, it can reach 253 m<sup>3</sup>/s. As a result of the low freshwater inflow and large tidal range, the estuary is generally well-mixed, with the mean salinity decreasing from around 30 PSU at the mouth to 1 PSU just south of Antwerp (Baeyens et al., 1997; Vandenbruwaene et al., 2013). The small salinity gradients in the estuary are not likely to directly impact sand transports.

#### Bed characteristics

The multi-channel system and the mouth area of the estuary primarily consist of sand with a median grain size between 150 and 250  $\mu\text{m}$ . Detailed measurements in the Wielingen channel

(Santermans, 2013) show a mud content between 5–20% for the fraction <2  $\mu\text{m}$  and 5–30% for the fraction 2–63  $\mu\text{m}$  near Zeebrugge. However, mud accumulation can be significant in the harbour basins. An analysis of dredge deposits shows that in the coastal access channel to Zeebrugge, 60–70% of the dredged material consists of mud with median grain diameters below 63  $\mu\text{m}$ . Towards the estuary, the mud content decreases, and sand fractions, defined as sediment with a  $d_{50} > 63 \mu\text{m}$ , dominate (88–95%). An analysis of the McLaren dataset (McLaren, 1993, 1994) shows that in the top-layer sediment composition the sand fraction dominates the main channel and shoal areas (77–95%) with a median grain size ranging between 150 and 250  $\mu\text{m}$ . Mud (<20% sand) covers less than 6% of the area of the Western Scheldt, sand (>80% sand) covers around 71% of the estuary bed, muddy sand (50–80% sand) about 12% and sandy mud (20–50% sand) 10%. More extensive mud deposits can be found along the shorelines and in shoal-sheltered areas (Fig. 3 and Table 1).

Van der Spek (1997) and Van der Vegt et al. (2021) point to the presence of erosion-resistant deposits in the subsurface of the Vlake van de Raan and Wielingen that contribute to their stability. In the estuary, several geologic formations form erosion-resistant layers. The bed composition of these layers ranges from clay and peat layers to cemented sandstones (Gruijters et al., 2004). The erosion-resistant layers are in many parts of the channels exposed on the channel flanks and floor and have noticeably influenced the morphodynamic evolution of the estuary (Dam & Cleveringa, 2013; Van der Vegt et al., 2020), but do not directly impact the sediment budget presented in this study.

#### Dredging, disposal and sand mining

Human activities play an important role in the morphological evolution of the estuary (Coen, 1988; Jeuken, 2000; Jeuken & Wang, 2010; Vandenbruwaene et al., 2019). Shore-protection works, including the building of dikes and reclamation of supratidal areas, started in the Middle Ages and evolved with time into large-scale reclamations and construction of dikes against flooding (see Section 4 and Supplementary material for more details). In addition to shore-protection works, dredging and disposal volumes increased over the last century (Belmans, 1988; Claessens et al., 1991; Kornman et al., 2003; Santermans, 2013). In the Western Scheldt, dredging already started in 1923 with the removal of the sill near Bath, and dredging has continued ever since to maintain and improve access to the various ports along the estuary.

**Table 1.** Estimated sand content (%) in the bed based on McLaren (1994) and subdivided into macrocells (Mc) and morphological elements as defined in Fig. 2.

Morphological elements	Sand content [%]						
	Mc1	Mc2	Mc3	Mc4	Mc5	Mc6	Mc7
Main channel	95	–	83	96	92	90	77
Main shoal	81	82	92	88	89	–	–
Secondary channel	–	84	87	84	88	–	–
Intertidal flats South	–	71	80	79	90	91	56
Intertidal flats North	95	–	64	64	74	74	64


Main channel: area below NAP –2 m in Mc1a,b, Mc3a, Mc4a, Mc5a, Mc6 and Mc7.

Main shoal: shoal area between main and secondary channel and depths between NAP +3 and –2 m.

Secondary (side) channel: depth below NAP –2 m in Mc2, Mc3b, Mc4b, Mc5b.

Fringing intertidal flats south: area along southern embankment, depth between NAP +3m and –2 m.

Fringing intertidal flats north: area along northern embankment, depth between NAP +3m and –2 m.

**Figure 3.** Bed composition map indicating the sand fraction percentage in the bed (top layer) based on the 1994 McLaren measurements. Data available from Rijkswaterstaat.


The navigation channel was deepened in 1973–1976, 1997–1998 and 2010 by lowering the shallowest sills. In addition to channel deepening, the width of the main channel is maintained at 300 m east of Hansweert and mostly 500 m to the west (only at the sill of Borssele a width of 300 m is maintained).

The first and largest channel deepening was executed between 1973 and 1976, dredging approximately 57 million m<sup>3</sup> of sediment resulting in the deepening of the shallowest sills by 2.5 to 3 m, which increased the depth of the navigation channel to NAP –14.5 m. Disposal occurred primarily in the secondary channels of the same macrocells and secondarily along eroding channel banks of the main navigation channel. During the second channel deepening (1997–1998), over 17 million m<sup>3</sup> of sediment was dredged, primarily from shallow sills, to increase the navigation channel depth to NAP –16 m. The dredged sediments were relocated to the secondary channels in the western part of the estuary. During the most recent channel deepening in 2010, 7.7 million m<sup>3</sup> of sediment dredging increased the main channel's depth to NAP –17.2 m. Disposal locations included the secondary channels, the deepest parts of the main channel and the shallow subtidal, seaward-located tips of three intertidal shoal complexes.

In a natural channel, an equilibrium exists between the channel's cross-sectional area and the local tidal prism (O'Brien, 1931, 1969). Pieters et al. (1991) indicate that before dredging, from 1920 to 1930, the equilibrium depth of the sills in the navigation channel was 8–10 m. As the dredged channel is far past

its equilibrium depth, it becomes an effective sediment trap and sediment deposition increases. As a result, maintenance dredging enlarged from less than 0.5 million m<sup>3</sup>/year before 1950 to 7–10 million m<sup>3</sup>/year at present (Fig. 4). Increasing maintenance dredging is a problem faced by most if not all industrialised inlets worldwide such as the Yangtze (Port of Shanghai), Elbe (Port of Hamburg), Weser (Port of Bremen) and Humber Estuaries (Van Maren et al., 2015; Chen et al., 2016; Kerner, 2007; Antwerp Port Authority, 2012; also see [www.tide-toolbox.eu](http://www.tide-toolbox.eu)). The dredging activities do not necessarily alter the magnitude of the annual sediment budget for the entire estuary as most of the sediments are relocated/disposed elsewhere within the estuary (Kornman et al., 2003; Pieters, 1993). Disposal strategies have changed over time. Before 1997, most of the dredging took place in the eastern part of the estuary, and this material was deposited in the nearby secondary channels (strategy 'East'). This strategy was economically most efficient, and it minimises the transfer of potentially contaminated sediment from the eastern part of the estuary to the western part. As dredging volumes increased, it was anticipated that this strategy would not be sustainable on the long term. To ensure the continued 'natural' dynamics of the channels and shoals in the estuary and to reduce dredging volumes, between 1997 and 2010, most sediment was therefore relocated to the secondary channels in the western part of the estuary, strategy 'East - West' (Liek, 2001; Jeuken & Wang, 2010). Kornman et al. (2003) concluded that this strategy did not result in reduced maintenance





**Figure 4.** Overview of dredging, disposal and sand-extraction volumes in the Western-Scheldt estuary between 1955 and 2020.

dredging. In 2010, a new disposal strategy called *Flexibel Storten* (which translates to Flexible Disposal) was implemented. The Flexible Disposal strategy aims to preserve the physical characteristics, such as the multichannel system and intertidal flat area, and thereby better retain the ecological value of the estuary (Roose et al., 2008; Plancke et al., 2010; Depreiter et al., 2012). Disposal mainly occurs in the same or downstream macrocell, while the exact disposal location is adjusted based on extensive monitoring and adapting the disposal locations to minimise negative impacts. In addition to ecological benefits, disposal in the vicinity of the dredge locations also results in cost savings and less pollution due to shorter transit times of the dredge vessels.

In addition to dredging and disposal, 122 million m<sup>3</sup> of sand was mined from the estuary between 1956 and 2014. In the Sea Scheldt, sand mining, 1 million m<sup>3</sup>/year on average, continues until today. As part of this mining occurs close to the Dutch-Belgian border, indirectly this mining may also result in sediment losses from the Dutch part of the estuary.

This study does not include the harbour basins along the Western Scheldt. These harbour basins act as (temporary) sinks for sediments, and extensive (70–90%) mud accumulations are found here (Van Maldegem et al., 1993; Van Kessel et al., 2012; Van Kessel et al., 2023). Frequent dredging is therefore required. Between 1965 and 1987, on average, 1.9 million tons/year of mud was disposed back into the estuary, between 2002 and 2010 disposal ranges 1.4 and 1.8 million tons/year (Dam, 2017). The proximity of the disposal areas close to the harbour basins predominantly introduces local effects on sediment transport patterns. Aggregated over the macrocells, the influence of the disposal is likely limited.

### Available bathymetric data and data accuracy

#### Data

The morphodynamic evolution of the Western Scheldt between 1955 and 2020 is studied using a series of bathymetric maps compiled from the Rijkswaterstaat databases. These map sheets cover the estuary and its mouth up to the seaward NAP-20 m contour (Fig. 5A). The underlying bathymetric data are collected using ship measurements (single beam and multi-beam) for the


domain's wet, subtidal parts. Since 2001 the dry parts have been surveyed using LiDAR. Strict protocols are in place to combine data sets and transfer them to a pre-designed 20 × 20 m grid divided into map sheets (Van der Zijp, 2001). Six map sheets cover the Western Scheldt and are available from 1955 onwards. Annual surveys have been available for this part of the estuary since 1996. Before it was surveyed bi-yearly, two subsequent surveys are needed to cover the estuary completely. The mouth area is surveyed less frequently, and more significant data gaps in the individual measurements occur. Elias et al. (2016) composed complete maps of the mouth area from 1964 to 2014, and additional maps are created with recent measurements. These maps were generally composed of two subsequent surveys taken 1 year apart. The resulting total dataset contains bathymetric maps that allow an analysis of the complete system (Western Scheldt and its mouth area) over 1970–2021 for the area covered by the polygons shown in Fig. 5.

#### Dealing with uncertainty and errors

The bathymetric data used in this study are not without flaws. A good understanding of the data accuracy is a prerequisite when assessing the sediment budget. Changes in survey techniques and instruments, positioning systems, variations in correction and registration methods and changes in interpolation techniques over time make it difficult to estimate the exact accuracy of the measurements and the resulting maps.

The vertical accuracy of the *Vaklodigen* data was estimated by Wiegmann et al. (2005) and Perluka et al. (2006) to range between 0.11 and 0.40 m. Storm et al. (1993) indicate that errors depend on the morphologic unit surveyed. For intertidal areas, channels and flats and channel slopes, stochastic (random) errors are 0.23, 0.19 and 0.39 m, respectively, and systematic errors (bias) are −0.20, −0.10, and −0.25 m. Marijs & Parée (2004) provide a detailed overview of possible measurement errors focusing on the Western Scheldt datasets. See Supplementary material for a detailed elaboration of uncertainty and errors.

The importance of horizontal accuracy for the sediment budget depends on the method used to assess the volume changes and the nature of the changes and the slope of the bed (greater importance



**Figure 5.** Overview of (A) map sheets on which bathymetric data is documented and (B) survey years of bathymetric maps used in this study.

for steeper slopes because the same horizontal change causes a greater vertical change when the slope is steeper). The impact of the horizontal positioning accuracy on the sediment budgets depends on the choice of the sections and the characteristics of the changes in bathymetry. For example, when considering the Western Scheldt as a single section in the sediment budget, a slight shift in the position of its boundaries or shifts of the positions of the channels do not affect the water volume within the section. Determining the inverse of the water volume in the Western Scheldt over time is sufficient to determine its sediment budget.

When considering units with similar morphodynamic behaviour (areas of erosion and deposition) in the sediment budget, the spatial variations in depths and depth gradients determine whether a shift in a position significantly affects the sediment budget (Elias et al., 2016). For a large area with little spatial variations in depth, a shift in the horizontal position results in a small volume change. This is the overall situation in the Voordelta, including the mouth of the Western Scheldt. Conversely, a shift in the horizontal position results in a significant volume change for a small area with large spatial variations in water depth. In the Western Scheldt, this is the prevailing situation. Therefore, the sediment budget sections within the Western Scheldt are chosen sufficiently large (e.g., macrocell level) to reduce the impacts of horizontal positioning (in)accuracies.

Without having access to the actual ship data, correction for measurement error is not a trivial task and can only be performed based on expert judgement. The most obvious flaws in the bathymetric data can be corrected manually. Elias et al. (2016) performed such correction for the mouth area based on visual inspection of the sequences of sedimentation-erosion maps. Careful inspection of these sequences allows identifying outliers and systematic inconsistencies in the data. Sounding sheets showing improbably large changes compared with preceding and following surveys in a few years were excluded from the analysis.

## Morphological evolution of the Western Scheldt

### A brief history

Comprehensive overviews of the natural processes and human interferences that have influenced the morphological evolution of the Western Scheldt estuary over the past two centuries are given by, e.g., Coen (1988), Van den Berg et al. (1996) and Van der Spek (1997) and summarised in the Supplementary material.

The Western Scheldt came into being between the 14<sup>th</sup> and 16<sup>th</sup> centuries as the river Scheldt connected to the tidal basin of the Honte (see Vos, 2015, p. 88–89 for a detailed account). By the 17<sup>th</sup>



century, the Western Scheldt estuary had primarily formed. At this time, expansive channels, tidal flats and marshes along the northern side of the estuary were present that still connected the Western and Eastern Scheldt estuaries. These marsh areas and the connecting channels silted up naturally or were dammed in the following centuries (Van der Spek, 1997; Dam *et al.*, 2022). Only the Verdrongen Land van Saeftinghe (see Fig. 1 for location) has remained a central tidal marsh area. Land reclamations decreased the intertidal storage area from 295 km<sup>2</sup> in 1650 to 196 km<sup>2</sup> in 1800 and to 104 km<sup>2</sup> in the recent Western Scheldt (Van der Spek, 1997).

In the estuary, meandering ebb channels dominate the morphological developments with annual migration rates of tens of meters between 1800 and 1930 (see maps in Supplementary material). This process was increasingly constrained around 1930 as the channels reached the protected embankments of the estuary (Van den Berg *et al.*, 1996). Migration now stopped, and channel banks steepened. As a result, bank failures and slides have been observed at various locations (De Bruin & Wilderom, 1961; Mastbergen *et al.*, 2019), prompting the further reinforcement of vulnerable channel walls with protective layers of gravel or slag.

Since the 1930s, the large-scale morphology of the Western Scheldt has been relatively stable, displaying a regular, repetitive pattern of mutually evasive meandering ebb channels and straight flood channels separated by sub- and intertidal shoals (e.g., Van Veen *et al.*, 2005; Van den Berg *et al.*, 1996; Jeuken, 2000; Toffolon & Crosato, 2007). Winterwerp *et al.* (2001) refer to this repetitive pattern as a series of meso- and macrocells (Fig. 2). Despite stability on the macro level, large-scale dynamics of the smaller-scale channels and shoals can still occur (Fig. 6 and Supplementary material). A significant change is the shift of the main channel in the central part of the estuary. Until 1982, the ebb channel along the northern embankment of the estuary (Middelgat) was the main navigation channel. However, a channel bend cut-off that originated in 1951 resulted in significant natural sediment deposition in the main ebb channel and erosion of the secondary flood-dominated channel (see, e.g., Jeuken, 2000). The increased dimensions of the flood channel enabled a formal relocation of the navigation channel to this channel in 1982. Due to continued channel migration tendencies and channel deepening local maintenance dredging is required. At the same time, the entrance of the flood channel has been a sediment disposal location since the 1980s. Significant changes in the western part of the estuary resulted from the closure of the side channels Sloe and Braakman. Before the closure, these side channels were connected to the main channels in the estuary and mouth. As a result of the complete closures that occurred in the 1950s, these channels are gradually infilling (Dam, 2017). Sedimentation in the now closed-off side channels Sloe and Braakman, which were sediment sinks up to 1955, is counted as export from (the present configuration of) the estuary. Bakker & De Loof (1977) show a sediment deposition of 20.6 million m<sup>3</sup> in Sloe and 31.1 million m<sup>3</sup> in Braakman between 1878 and 1952 and an additional 41.9 million m<sup>3</sup> accreted on the marsh area of Saeftinghe.

### **The mouth area of the estuary**

The mouth area of the Western Scheldt estuary stretches from the island of Walcheren in the north into Belgian territorial waters in the south (Figs. 1 and 6). The present-day lay-out of the mouth area (Fig. 6B), with the most prominent channel Wielingen in the south and a complex of tidal channels and shoals along the SW coast of


Walcheren, came into existence about a century ago when the Deurlo, a third channel that dissected the large subtidal shoal Vlake van de Raan, silted up. This morphological pattern persisted over the past 45 years. The formation of the two-channel system significantly impacted the morphodynamic processes in the northern part of the mouth area, where the main shoals and the channel Oostgat all rotated northward or moved landward (Van Enkevoort, 1996; Van der Slikke, 1998). The increasing depth of the channel Wielingen contributed to the erosion of the Zeeuws-Vlaanderen coastline.

Based on the sediment budget, Elias *et al.* (2016) computed a sediment loss in the Dutch part of the mouth area of 0.8 million m<sup>3</sup>/year over the time period 1965–2010 and a loss of 1.2 million m<sup>3</sup>/year between 1980 and 2010. Structural erosion occurs in the central part of the mouth area, the Vlake van de Raan, and is estimated to be 0.7 million m<sup>3</sup>/year since 1984.

Elias & Van der Spek (2015) estimated the volume loss of the entire estuary mouth, including the Belgian part to range between 200 and 278 million m<sup>3</sup> in the period 1980–2010. Large morphodynamic changes were observed around the harbour of Zeebrugge. The extension of the harbour jetties to 4 km length (1972–1986) resulted in large-scale scour around the jetties, and deepening of the entrance channel resulted in a nearly 50 million m<sup>3</sup> sediment volume loss. Deepening of the Wielingen channel did not directly impact the sediment budget as the dredged deposits were disposed along the channel. Significant morphodynamic changes have occurred along the southwest Walcheren coastline (Fig. 6C), mainly caused by the migration of channels and shoals. The Oostgat channel is deepening slightly over its entire length; its seaward part offshore the western tip of Walcheren extends to the north. This channel, being close to the coastline, has resulted in coastal and channel bank erosion, necessitating frequent nourishments of the adjacent beaches. Extensive dune, beach and channel-bank nourishments (7.1 million m<sup>3</sup>) occurred in 2009 and 2010. Despite the large gross morphodynamic changes of nearly 170 million m<sup>3</sup> between 1967 and 2011, the net 17 million m<sup>3</sup> accretion is small and near-equal to the sand added along the coastline by nourishments. Nourishments have been frequently executed since 1952, although the majority (80%) of the 17 million m<sup>3</sup> nourishment volume was supplied more recently (since 1991). Initially, the nourishments were executed as beach and dune nourishments, but since 2005 the erosion induced by Oostgat has been successfully balanced with channel-bank nourishments (9.1 million m<sup>3</sup> in total). The channel-bank nourishments have stabilised the shoreface and shifted the channel offshore (Nederbragt & Koomans, 2006; Dekker, 2012).

Substantial sediment losses, nearly 25 million m<sup>3</sup>, occurred along the Wielingen channel and the Zeeuws-Vlaanderen and Belgium coastlines between 1967 and 2011. These losses include the 15.3 million m<sup>3</sup> of sand added to the Zeeuws-Vlaanderen coastline since 1971.

A noticeable feature in the volume time series of the mouth of the estuary (Fig. 7, raw data line) is the nearly 150 million m<sup>3</sup> increase in sediment volume between 1964 and 1985 and the decreasing volumes since. Inaccuracies in the bathymetries play a role in this volume development as a close examination of the bathymetric maps revealed unrealistic bathymetric changes (Elias *et al.*, 2016). However, correcting these maps for measurement errors is not a trivial task. An estimate of the uncertainty range is obtained by assuming that recent trends also represent long-term



**Figure 6.** Bathymetry of the Western Scheldt estuary for the representative years (A) 1970/71 and (B) 2020/21, based on Vaklodingen 2020 (mouth) and 2021 (estuary). The morphological changes over this interval are shown by the sedimentation-erosion patterns in panel (C).


changes. Such estimates indicate significantly enlarged volumes before 1980 (Fig. 7, trend lines).

### Hydrodynamic changes

Tidal amplification, deformation and the morphological development of a natural estuary are closely linked (Dronkers, 1986; Friedrichs & Aubrey, 1988). Tidal amplification is the change in

tidal range over the estuary relative to the tidal range at its mouth and can be expressed by the amplitude-ratio of the semi-diurnal tide between two stations. Tidal deformation refers to the distortion of the tidal wave and changing the tidal asymmetry as the tide propagates through the estuary. Tidal asymmetry in shallow estuaries can be represented by the non-linear growth of the compound constituents and harmonics of the principal tidal components (Friedrichs & Aubrey, 1988; Speer et al., 1991). The





**Figure 7.** Overview of the computed volumes for raw and corrected data between 1964 and 2017 of the mouth area of the Western Scheldt relative to 2014.

ratio between the amplitude of the semi-diurnal tide (D2) and its overtide (D4) determines the strength, while the phase difference describes the nature of the asymmetry (ebb- or flood dominant) and can directly be determined by comparing the durations of flood and ebb flow (Dronkers, 1986) or by comparing the durations of rising tide and the falling tide from water levels (Wang *et al.*, 2019).

Asymmetry of the horizontal tide is an essential factor for residual sediment transport. Shorter and stronger flood flow leads to landward sediment transport, referred to as flood-dominant asymmetry of the horizontal tide. In an estuary with low river flow like the Western Scheldt, the asymmetry of the horizontal tide at a particular cross-section is directly related to the asymmetry of the vertical tide in the part of the estuary landwards of the cross-section.

The tidal amplification and deformation are analysed using the water level observations from Vlissingen, Terneuzen, Hansweert and Bath (see Fig. 1 for locations). Continuous hourly water level records between 1971 and 1987 are available for these stations, while the data since 1987 have 10-minute intervals. The daily tidal characteristics were determined by analysing individual 25-hour (until 1987) or 24-hour and 50-minute time series. The daily mean water level ( $a_0$ ) and amplitude and phase of the diurnal ( $a_1$  and  $\phi_1$ ), semi-diurnal ( $a_2$  and  $\phi_2$ ) and overtide ( $a_4$  and  $\phi_4$ ) were determined through Fourier analysis. The falling ( $T_f$ ) and rising tides ( $T_r$ ) durations are determined directly from the entire time series. Tidal asymmetry is described through the relative phase-difference between the quarter-diurnal and the semi-diurnal tides (Fig. 8C) and by the ratio between the rising and falling period,  $T_r/T_f$  (Fig. 8D).

#### Tidal amplification and phase lag

The semi-diurnal tidal amplitude increased in the landward direction since 1971 as the amplification factor, i.e., the amplitude ratio relative to the station Vlissingen at the mouth, exceeds 1 at all


three stations in the estuary (Fig. 8A). In the western part of the estuary, the mean amplification factor remained constant between 1970 and 2020. At station Bath, the amplification factor increased by 6% from nearly 1.15 in 1971 to 1.23 in 2020. The phase lag of the semi-diurnal tide remained constant between Vlissingen and Hansweert, but decreased over time between Hansweert and Bath (Fig. 8B). This decrease in phase lag represents a faster tidal wave propagation in the eastern part of the estuary. Both effects, increase in amplification and faster propagation, are an expected result of the increasing depth of the navigational channel.

#### Tidal asymmetry

Between 1970 and 2020, the relative phase difference between D4 and D2 at the mouth of the estuary (Vlissingen) showed fluctuations between  $0^\circ$  and  $-10^\circ$ , which indicates flood dominance (Fig. 8C). Flood dominance also occurs at station Terneuzen. At this station, the increasing phase difference from  $0^\circ$  around 1970 to  $-10^\circ$  at present indicates that the strength of the flood dominance increased. More considerable but opposite changes occurred at stations Hansweert and Bath. A large, positive ( $20^\circ$ ) phase difference at Hansweert indicates ebb dominance in 1970. This phase lag decreased to around  $2^\circ$  in 1985 and retained a mostly positive value until 2008. At present, a negative phase lag of  $-8^\circ$  indicates a flood-dominant behaviour. The change in phase lag for station Bath is opposite to Hansweert between 1970 and 2010 as the phase lag reduced from  $-30^\circ$  (strongly flood dominant) to around  $0^\circ$  (no dominance). Since 2010 trends are similar to Hansweert, with an increasing flood dominance. At present, all stations in the estuary show flood dominance with a phase lag between  $-8$  and  $-14^\circ$  in 2020.

Defining the tidal asymmetry as the ratio between the rising and falling tide produces values between 0.86 and 1 for all stations (Fig. 8D). These values imply that the entire estuary has been flood dominant since 1970, and except for station Bath, flood dominance has increased. The most significant increase occurred at Hansweert between 1970 and 1985 (after the first channel deepening).





**Figure 8.** Development of the tidal amplification and deformation in the estuary based on year-averaged values for (A) Amplitude ratio and (B) phase lag of the semi-diurnal tide in the estuary relative to the station Vlissingen at the mouth; (C) the phase difference between the quarter-diurnal (D4) and the semi-diurnal (D2) tides and (D) the ratio between the periods of rising ( $T_r$ ) and falling tides ( $T_f$ ). Grey shading indicates the time periods of main channel deepening.

Relatively, both indicators for tidal asymmetry show the same changes in the strength of the tidal asymmetry at the various stations. However, in an absolute sense, the two parameters do not indicate the same nature of tidal asymmetry. Based on the phase difference, the estuary is mostly flood dominant, while based on the duration ratio the estuary has been flood dominant over the whole period. This difference shows that just considering the relative phase lag between the quarter-diurnal and semi-diurnal tide is insufficient to characterise the tidal asymmetry in the Western Scheldt estuary; it requires the consideration of the entire water level signal to determine the duration ratio of the ebbing and flooding tide.

### Large-scale sediment budget for the estuary; 1955–2020

#### Method

##### Sediment budget

Sediment budgets provide a powerful method to structure all contributions to the sediment volume change of the estuary (Reid & Dunne, 1996; Rosati, 2005; Frings et al., 2019). The sediment budget was resolved volumetrically and not strictly as a mass balance. This approach was justified given the dominance of the

sand fraction, and all sources and sinks to the sediment budget can either be volumetrically derived from the data or literature. The sediment budget can be computed from the bed level measurements ( $dV$ ), the changes caused by human interventions ( $V_{DREDGE}$ ,  $V_{DISPOSAL}$  and  $V_{SANDMINING}$ ) and the exchange fluxes between the mouth area ( $V_{WS}$ ), Sea Scheldt ( $V_{SCHELDT}$ ) and Land van Saeftinghe ( $V_{SAEF}$ ):

$$dV = V_{WS} - (V_{SCHELDT} + V_{SAEF}) + (V_{DISPOSAL} - V_{DREDGE} - V_{SANDMINING})$$

This sediment budget is resolved on several scale levels: (1) the scale of the estuary, (2) the eastern (Mc1, Mc2, Mc3 and Mc4) and western parts (Mc5, Mc6 and Mc7) of the estuary and (3) on the level of the macrocells (Fig. 9). The upper height limit for the bathymetric measurements in the estuary was set at NAP +3.0 m. The considered area includes all intra- and supratidal shoals and marshes and minimises potential errors due to differences in coverage along the diked shorelines. For each macrocell, an addition subdivision in 5 morphological units was made. These units were defined as:

1. main channel: area below NAP  $-2$  m in Mc1a,b, Mc3a, Mc4a, Mc5a, Mc6 and Mc7,
2. main shoal: the shoal area between the main channel and the secondary channel and depths above NAP  $-2$  m,
3. secondary (side) channel: area below NAP  $-2$  m in Mc2, Mc3b, Mc4b, Mc5b,
4. fringing intertidal flats south: the area along the southern embankment, bed level between NAP  $+3$  m and  $-2$  m and
5. fringing intertidal flats north: the area along the northern embankment, bed level between NAP  $+3$  m and  $-2$  m.

Except for Macrocell 2, where no main channel is present, for each macrocell the morphological units were grouped in two elements that represent the (a) main navigation channel and (b) all other channels and shoals (the secondary channel & shoals).

#### *Additional sediment sources, sinks and assumptions*

*Saeftinghe.* The large marsh area Verdrongen Land van Saeftinghe is not well represented in the data before 2001, while LiDAR data after 2001 may include erroneous heights (Cleveringa, 2013), and is therefore not directly computed from the bathymetric data. An estimate for the volume losses ( $V_{SAEF}$ ) of 0.3 million  $m^3$ /year based on the values used in previous studies (Houtekamer, 1995; Storm & Pieters, 1994; Uit den Bogaard, 1995, Nederbragt & Liek, 2004; Haecon, 2006) was added to the sediment budget. A literature review of mud in the Sea Scheldt by Dams *et al.* (2016) concludes that mud accumulation in Saeftinghe ranges between 0.06 and 0.12 million  $m^3$ /year. This study used an average value of 0.09 million  $m^3$ /year for the mud contribution, which leaves 0.21 million  $m^3$ /year for the sand deposition.

*Sea Scheldt.* Bathymetric measurements of the Sea Scheldt are less frequent compared to the estuary and only available in 10-year intervals. The morphodynamic changes are dominated by large-scale dredging and disposal. Haecon (2006) estimated a 1.0 million  $m^3$ /year sediment loss to the Sea Scheldt between 1955 and 2004. A literature review of mud in Sea Scheldt by Dams *et al.* (2016) concluded that the annual mud transport to Saeftinghe and Sea Scheldt combined averaged 0.17 million  $m^3$ /year before 1987 and increased to 0.26 million  $m^3$ /year since 1987. This increase is related to large-scale mud extraction in the Sea Scheldt that started around this time. Vandenbruwaene *et al.* (2017) computed a net downstream mud transport of 0.25 million  $m^3$ /year between 2001 and 2011, and a net upstream sand transport of 0.61 million  $m^3$ /year. The recent studies of Planke *et al.* (2021) and Vos *et al.* (2021) compute an upstream sand transport of 1.0 million  $m^3$ /year and a downstream mud transport of 0.16 million  $m^3$ /year in the period 2011–2016. Between 2016 and 2019, both the mud and the sand transports are upstream directed with values of 0.16 million  $m^3$ /year and 0.48 million  $m^3$ /year, respectively.

Given the uncertainty in annual sediment exchange between the estuary and the Sea Scheldt, the sediment budget is resolved for four scenarios of sediment exchange (Table 2). In the ‘minimum’ scenario, no sediment exchange is assumed to occur (Scenario 1); in the ‘maximum’ scenario (Scenario 2), the values, as proposed by Haecon (2006), are applied. A ‘best estimate’ scenario (Scenario 3) uses the estimates of Dams *et al.* (2016) before 2001 and the results of Vandenbruwaene *et al.*, (2017), Planke *et al.* (2021) and Vos *et al.* (2021) for the more recent (2001–2020) period. Scenario 4 uses the same estimates as Scenario 3, but distinguishes between sand and mud transport.

The macrocell sediment budget uses Scenario 3 as an assumption for  $V_{SCHELDT}$  and  $V_{SAEF}$  at Mc7. This assumption

allows to compute the sediment transport rates between the macrocells (blue arrows). The sediment transport directions within the macrocells cannot be directly determined from the sediment budget and require expert-judgement-based interpretation of the bathymetric changes (Fig. 6). In the eastern part of the estuary, the main channel forms a separation between the major shoal areas, it therefore seems valid to assume that sediment exchange between the macrocells only occurs through the main channel. In the western part, the main channel forms a major sediment pathway. In addition, sediment transport from the secondary areas in Mc3 to the secondary areas in Mc4 is probable, as these areas are connected directly through a series of shallow channels and sills (Fig. 6). Under these assumptions, estimates of the intra-estuary transports can be determined from the sediment budget.

*Dredging, disposal and sand mining.* Sediment volumes for dredging ( $V_{dredging}$ ), disposal ( $V_{disposal}$ ) and sand mining ( $V_{sandmining}$ ) are based on Cleveringa (2013) and thereby similar to Haecon (2006) until 2001 (Fig. 4). However, these authors indicate that the accuracy and completeness of the data are unknown. Recent data are based on values reported in IMDC (2020, 2021) and Schrijvers (2021).

The combined dredge-disposal timeseries show that sand extraction prevails in the eastern part of the estuary; part of the sediment is disposed to the West, resulting in a sediment surplus there. Sand mining has resulted in 122 million  $m^3$  of sand volume loss since 1955.

#### *Bed-level change*


Large yearly variations in bed level occur between 1955 and 2020 (see Fig. 6 for a representative depiction of these changes). The gross yearly sedimentation and erosion volumes vary between 40 and 100 million  $m^3$  resulting in over 7000 million  $m^3$  of volumetric change (Fig. 10B). The total net change of  $-58$  million  $m^3$ , an average net erosion rate of 0.9 million  $m^3$ /year, is small compared to the gross changes (Fig. 10A and Table 3). The observed net erosion of 58 million  $m^3$  is smaller than the sand extraction of 142 million  $m^3$  that occurred. Sand mining (122 million  $m^3$ ) is the dominant factor for the observed losses. The small difference between dredging and disposal volume accounts for 20 million  $m^3$ . The extracted volumes are two times larger than the net sediment loss but only represent 2% of the observed total change. Based on the measured volumetric changes, the estuary loses sediment, but after accounting for dredging, disposal and sand mining, the estuary would have gained 84 million  $m^3$  of sediment (Fig. 10A, dashed line = estuary total corrected).

Erosion dominates the main channel (Fig. 10C) as the channel (water) volume increased by 270 million  $m^3$ . Maintenance dredging of sills and structural channel deepening, 529 million  $m^3$  in total, plays an important role in the increasing channel volume. Part of the dredged material was disposed in the secondary channels, contributing to their 187 million  $m^3$  net accretion. The shoal areas, the central shoals between the main and secondary channel and the shoals along the embankments of the estuary (shoals south and north), show a small 9 million  $m^3$  sediment gain.

An estimate of the sand versus mud contribution to the sediment budget was obtained using the average sand and mud contents derived from the McLaren (1994) dataset (Table 1). The measurements by McLaren (1994) show that a significant mud fraction can be present in parts of the estuary (see Fig. 3 and Table 1); however, the sand fraction dominates the total volumetric change of the estuary (Fig. 10A). The cumulative mud volumes

**Table 2.** Sediment transport rates for exchange between the estuary and the Verdrongen Land van Saeftinghe and Sea Scheldt in millions m<sup>3</sup>/year (negative values = upstream transport, sediment export from the estuary, positive values= downstream transport).

Scenario	Interval	Type	Saeftinghe	Sea Scheldt
1. Minimum	1955–2020	Sediment	−0.30	0
2. Maximum	1955–2020	Sediment	−0.30	−1.00
3. Best estimate	1955–2020	Sediment	−0.30	−0.67
4. Sand and mud	1955–2020	Sand	−0.21	−0.63
		Mud	−0.09	−0.04
	1955–1987	Sand	−0.21	−0.61
		Mud	−0.09	−0.08
	1987–2001	Sand	−0.21	−0.61
		Mud	−0.09	−0.17
	2001–2011	Sand	−0.21	−0.61
		Mud	−0.09	+0.25
	2011–2016	Sand	−0.21	−1.00
		Mud	−0.09	+0.16
2016–2020	Sand	−0.21	−0.48	
	Mud	−0.09	−0.16	



**Figure 9.** Principal description of the sediment budget used in this study.

show minor variations between 1955 and 2005 and a clear trend of accumulation since 2005.

*The western part of the Western Scheldt estuary (Macrocells 1–4)*  
 The total 19 million m<sup>3</sup> net sediment volume gain of the western part of the estuary is small, considering the net 65 million m<sup>3</sup> of disposal in this part of the estuary (Fig. 10E). Corrected for dredging, disposal and sand mining, the total sediment erosion/export in the western part is approximately 46 million m<sup>3</sup>. This number results mainly from Mc3 as a net, continuous sediment export (0.82 million m<sup>3</sup>/year) from this macrocell occurs (Fig. 10D). Significant sediment accumulation occurs in Mc2 where a continuous increase in the volume of 0.72 million m<sup>3</sup>/year


over the entire period can be observed (Fig. 10D). The volumetric change of Mc1 is small (−0.28 million m<sup>3</sup>/year) and results mainly from the increase in depth before 1970. Mc4 shows an increase in sediment volume until 1990 and alternating periods of erosion and sedimentation and a limited total net change since.

*The eastern part of the Western Scheldt estuary (Macrocells 5–7)*  
 The dominant effect of sediment management is especially apparent in the eastern part of the estuary. The net sediment volume loss of 77 million m<sup>3</sup> (Fig. 10F) is smaller than the sum of sand extraction (−401 million m<sup>3</sup>) and disposal volumes (+209 million m<sup>3</sup>). Erosion is related to the increasing depth of the main



**Table 3.** Measured volume change in million m<sup>3</sup>/year for the total estuary (negative values = sediment loss, positive values = sediment gain).


	Human interventions [million m <sup>3</sup> ]				Volume change [million m <sup>3</sup> ]		
	Dredging	Disposal	Sand mining	Total intervention	Sand	Mud	Total
1955–1970	–63	+61	–25	–27	+2	+2	+3
1970–1980	–97	+84	–28	–42	–31	–2	–33
1980–1994	–123	+124	–35	–34	–9	0	–8
1994–2005	–101	+100	–23	–24	–49	+2	–47
2005–2020	–144	+141	–11	–15	+20	+6	+27
<b>1955–2020</b>	<b>–528</b>	<b>+510</b>	<b>–122</b>	<b>–142</b>	<b>–67</b>	<b>+9</b>	<b>–58</b>

**Figure 10.** (A): Timeseries of changes in sediment volume between 1955 and 2020 for (A) the cumulative net volume change of the total estuary (grey shading indicates the time-periods of main channel deepening), (B) gross and net yearly volume changes, (C) net volume change for the various morphodynamic elements, and (D) net volume change for individual macrocells. Panels (E) and (F) respectively show the cumulative net volume changes for the western and eastern parts of the estuary.

channel (–131 million m<sup>3</sup>). The shoal areas and secondary channels show a small net gain of 54 million m<sup>3</sup>.

The volume timeseries shows continuous erosion since 1970, although the erosion rates vary over time (Fig. 10F). The most considerable erosion rates occurred between 1970 and

1980 (–3.4 million m<sup>3</sup>/year) and between 1994 and 2005 (–3.7 million m<sup>3</sup>/year). They are related to the channel deepening that took place during these time frames (Taal *et al.*, 2013). The volume loss is limited between 1980 and 1994 and from 2005 until 2020. The volume timeseries corrected for human interventions shows



**Figure 11.** (A): Timeseries for cumulative sediment transports through the transect Vlissingen-Breskens based on four exchange scenarios with the Sea Scheldt. Red shading indicates the variation in import rates based on the four scenarios. Representation of the single-box sediment budget (1955–2020) for (B) Scenario 1, (C) Scenario 2, (D) Scenario 3 in (1) the single-box model, (2) distinguishing between the main channel and secondary channels and shoals, (3) and subdivided in the western and eastern part of the estuary. (E) single-box sediment budget for Scenario 4, subdivided into sand and mud.

no net transport in or out of the eastern part of the estuary until 1980. Since 1980, however, the corrected time series reveals a net gain in the eastern part of the estuary of 3.0 million  $\text{m}^3/\text{year}$ . The gain increases to nearly 4.0 million  $\text{m}^3/\text{year}$  if sediment exports to the Sea Scheldt and the Verdrongen Land van Saefthinge are included.

**Sediment exchange between the estuary and the North Sea**

An estimate of the sediment exchange between the Western Scheldt estuary and the North-Sea is obtained by resolving the

sediment budget over the 1955–2020 period (Fig. 11). The sediment budget reveals an average net import of 1.6–2.6 million  $\text{m}^3/\text{year}$ . In the best-estimate scenario (Table 2, Scenario 3), a net import of 2.2 million  $\text{m}^3/\text{year}$  is computed. However, the import rate varies over the period when the intervals of Table 3 are applied. The largest import rates occurred between 1980–1994 and 2005–2020, respectively, 2.9 and 3.7 million  $\text{m}^3/\text{year}$ . A net export of 1.3 million  $\text{m}^3/\text{year}$  is computed between 1994 and 2005. Distinguishing between sand and mud in the computation shows that the sand and

sediment show near identical sediment import curves (Fig. 11A). This similarity is due to the relatively minor contribution of mud in the volumetric changes. Mud has a significant presence in certain parts of the estuary, especially along the banks, but the volumetric changes in these areas are limited. As a result, the computed mud import is small (0.3 million m<sup>3</sup>/year).

Subdividing the sediment budget into (i) main channel and (ii) secondary channels and shoals (Fig. 11, D2, and D3) reveals the large difference in response between these two elements. The main channel has deepened at  $-4.2$  million m<sup>3</sup>/year, with near-equal contributions of the channel in the western and eastern parts of the estuary. However, note that the net volume dredged in the main channel of the eastern part is 4 million m<sup>3</sup>/year, which implies a significant sediment supply from west to east to compensate for this loss. An average 0.97 million m<sup>3</sup>/year loss of sediment to Saefthinge and the Sea Scheldt adds up to an average transport of sediment from west to east of 3.0 million m<sup>3</sup>/year. The volume gain in the secondary channels (of Mc2 and 4 in particular) and shoals is 3.3 million m<sup>3</sup>/year, most of it (2.5 million m<sup>3</sup>/year) in the western part of the estuary.

### Understanding intra-estuary sediment exchanges

Further understanding of the intra-estuary sediment exchange is obtained by filling in the detailed sediment budget (Figs. 12 and 13).

Deepening of the navigation channel (1973–1976, 1997–1998 and 2010) by dredging the shallowest sills, in combination with maintenance dredging, resulted in a deepening of the entire main channel (Mc1a–Mc7a). Bed lowering rates of the individual macrocells vary between  $-0.4$  and  $-1.3$  million m<sup>3</sup>/year. In the eastern part of the estuary, all main channel macrocells (Mc5a, 6a and 7a) show bed lowering but still act as significant sediment sinks. The rates of channel maintenance dredging exceed the observed lowering of the bed level significantly. Therefore, averaged over the entire period 1955–2020, a net eastward transport and sediment import at the mouth prevails. Using scenario 4 as an assumption for exchange with the Sea Scheldt and the Verdrongen Land van Saefthinge yields an average sediment import rate of 2.2 million m<sup>3</sup>/year. With a value of 3 million m<sup>3</sup>/year, transports between the western and eastern parts of the estuary are 35% larger (Fig. 13A) than the sediment import rates.

The most considerable volume losses (caused by bed lowering) occur in the central parts of the main channel (Mc4a and Mc5a). Despite this bed lowering, the sediment budget reveals that only Mc4a is a primary sediment source. This macrocell likely forms a crucial link connecting the eastern and western parts of the estuary as almost 2/3 of the sediment needed to compensate for the sediment demand of the main channels in the eastern part of the estuary is delivered by Mc4a (Fig. 13A). Macrocell 4a forms a continuous sediment source, but its magnitude significantly increased around 1985. This increase is related to significant bed lowering associated with the bend cut-off initiated in 1951 and increasing annual disposal.

Most of the secondary parts of the macrocells (Mc1b, Mc2, Mc4b, Mc5b, Mc6b and Mc7b) show an increase in bed level between 1955 and 2020 (Fig. 12A). Only in Mc3b, a relatively small bed lowering at a rate of  $-0.3$  million m<sup>3</sup>/year can be observed. This bed lowering is remarkable given the 1.2 million m<sup>3</sup>/year of net disposal that took place here. The extensive sediment dispersion capacity of Mc3b is reflected by its 1.3 million m<sup>3</sup>/year source capacity (Fig. 13A). Maximum source values exceeding 2.7 million m<sup>3</sup>/year occurred between 1994 and 2005 when disposal peaked.

Decreased disposal after 2005 reduced the source magnitude to 1.5 million m<sup>3</sup>/year. Mc4b shows an opposite behaviour compared to Mc3b. In this macrocell, the net disposal is limited (0.36 million m<sup>3</sup>/year), but a significant increase in the bed level occurred ( $+2$  million m<sup>3</sup>/year). As a result, this macrocell forms the most significant sediment sink in the estuary. The infilling of the Middelgat channel governs the increasing bed level (Fig. 6A). The most significant increase in bed level occurred between 1994 and 2005 (Fig. 12E), which coincides with the peak in disposal in Mc3b. Additionally, it is likely that Mc3b supplied sediment to Mc4b.


A breakdown of the sediment budget in the periods, 1955–1970, 1970–1980, 1980–1994, 1994–2005, 2005–2020, reveals distinct sediment transport patterns, changes and rates (Fig. 13B–F). The most noticeable changes are the temporary switch from sediment import to sediment export at the mouth of the estuary between 1994 and 2005 and the increased sediment exchange between the eastern and western parts of the estuary since 1980.

Between 1955 and 1970, the sediment transport from the western part to the eastern part of the estuary approximated 1.3 million m<sup>3</sup>/year. This transport is significantly smaller than the 3.4–5.1 million m<sup>3</sup>/year exchange observed since 1980. Before 1970, most of the maintenance dredging and disposal took place in the eastern part of the estuary. The secondary parts (Mc5b, Mc6b and Mc7b) acted as sediment sources, with source values near equal to the disposal values. As a result, these parts provided the majority of the sediment demand ( $\sim 50\%$ ) of the main channel, resulting in limited transport between the western and eastern part of the estuary.


Since 1970, the secondary elements Mc5b, Mc6b and Mc7b all show small source and sink values, indicating that little sediment exchange occurred between these elements and the main channel. Disposal of sediment in the secondary areas now leads to long-term storage of sediment but also to a sediment deficit in the remainder of the system and increased sediment import. Since exchange between the secondary areas and the main channels did occur before 1970, this behaviour change is likely caused by dredging and disposal associated with the deepening and maintenance of the navigation channel. This limited sediment exchange between the secondary area and the main channel has remained since. With increased dredging rates in the main channel, sediment transport from the western to the eastern part of the estuary has increased considerably since 1980.

It is probable that the temporary reversal of the net transport direction from import to export at the mouth of the estuary between 1994 and 2005 is linked to the sediment dispersal strategy. Between 1997 and 2010, most of the dredged material was deposited in the western part of the estuary. In Mc3b and Mc4a alone, a combined 6.5 million m<sup>3</sup>/year source value arises that significantly exceeds the sediment sink value in the eastern part of the estuary. As a result, a surplus of sediment exists in the estuary's western part, which can only be balanced through sediment export at the mouth. Such temporary reversal in sediment transport direction does not occur between the estuary's western and eastern parts. Since 1980 transport has increased significantly to 3 to 4 million m<sup>3</sup>/year and has been exceeding 5 million m<sup>3</sup>/year since 2005 (Fig. 13D, E, F). Based on the observed transports, one can conclude that the western part of the estuary can act as a sediment source for the eastern part if sufficient sediment is available. The main channel of Mc4 plays a key role as a sediment source area. Since 1980 this macrocell has acted as a 2.2–3.8 million m<sup>3</sup>/year sediment source due to approximately equal source contributions of channel deepening and disposal.





**Figure 12.** Summary of the average annual sediment volume changes per macrocell, based on bed-level changes and dredging, disposal and sand-mining volumes, using the detailed sediment box model for the time periods (A) 1955–2020, (B) 1955–1970, (C) 1970–1980, (D) 1980–1994, (E) 1994–2005 and (F) 2005–2020. Macrocell subdivision: (a) main channel, (b) main shoal + secondary channel + side shoals north & south. Export rates to Verdrongen Land van Saefthinge and Sea Scheldt are based on Scenario 4.



**Figure 13.** Detailed, macrocell-based, sediment budget for the Western-Scheldt estuary. Blue arrows indicate the computed transports between macrocells over the time periods (A) 1955–2020, (B) 1955–1970, (C) 1970–1980, (D) 1980–1994, (E) 1994–2005 and (F) 2005–2020.

The largest sediment import rates are observed between 2005 and 2020. These high rates are surprising as sand mining considerably decreased during this period and was stopped after 2014. The sediment budget reveals that an over 2 million m<sup>3</sup>/year decrease in the erosion rate of Mc4a plays an essential role in the augmented sediment import rates as less sediment is available to fulfill the sediment demand of the sink areas. In addition, the change in dredge-disposal strategy is also expected to contribute to the augmented sediment import rates. Implementing the Flexible Disposal strategy in 2010 resulted in large-scale disposal in Mc5b. The over 1 million m<sup>3</sup>/year disposal resulted in similar sedimentation, effectively reducing sediment availability by a similar amount. This sediment storage capacity of Mc5b could also be observed in the 1970–1980 period. During this period, extensive channel deepening occurred, which temporarily disrupted the sediment transports as import rates reduced to 1.9 million m<sup>3</sup>/year. Over 2 million m<sup>3</sup>/year of disposal and bed-level increase occurred in Mc5b.

In all periods considered, Mc4a is the largest sediment source in the main channel. This is due to the significant increase in depth and due to the disposal in the channel. Disposal continuously increased from an average 0.2 million m<sup>3</sup>/year between 1955 and 1970, to 2.9 million m<sup>3</sup>/year recently (2005–2020). The neighbouring macrocell Mc3a shows the opposite behaviour and acts as a small sediment sink between 1980 and 1994 (−0.05 million m<sup>3</sup>/year) as the main channel deepened at the same rate. Since 1994 channel dredging has doubled, but channel deepening was limited. As a result, the sink value increased; presently, this macrocell forms the second largest sink in the system (−1.9 million m<sup>3</sup>/year).

Mc1a, near the mouth of the estuary, shows alternating periods of increasing and decreasing bed levels and sink or source function. These changes are likely due to natural variability as net sand removal through mining or dredging was limited (0.1–0.2 million m<sup>3</sup>/year). Disposal does not occur in Mc2. Therefore, the bed-level increase in the latter macrocell is related to natural sediment transports. The 0.72 million m<sup>3</sup>/year deposition rate may be considered the upper limit of coastal sediment supply from the adjacent Zeeuws-Vlaanderen coastline.

## Discussion and synthesis

### *Uncertainty in the sediment budget*

#### *Sediment composition: sand versus mud*

The presented sediment budget does not resolve the complete sand and mud budget. A crude estimate of the contributions of the two fractions was based on a single bed-composition map for which the sand-mud ratio for the various morphodynamic elements was determined. These ratios were kept constant over all measurements. Based on this approximation, we can conclude that the sediment import is dominated by the sand fraction (1.9 million m<sup>3</sup>/year) with only a minor contribution of the mud fraction (0.3 million m<sup>3</sup>/year), as the sand and sediment budgets follow similar trends and are approximate of similar magnitude. Therefore, the sediment budget also provides a reasonable estimate of the sand budget. However, the mud budget results are incomplete and require careful consideration of, e.g., the harbour basins, sediment porosity and bed composition (Van Kessel et al., 2023), and should not be used for further analysis.

#### *The porosity of the sediment*

Changes in the porosity of the sediment may result in changes in the observed sediment volume without any sediment transport. Variations in the porosity can result from physical and biological

processes such as bubble sand formation (De Boer, 1979), compaction and bioturbation. Compaction is expected to be of importance for the marshes. These processes can result in significant variations on small time scales but are less likely to influence the year-averaged results presented here significantly.

### *Subsidence*

In theory, subsidence of the subsurface influences the sediment budget since compensation of subsidence by deposition involves an extra sediment volume that cannot be determined from observed bed level changes. Therefore, sediment deposition rates are possibly underestimated in subsidence areas, especially over longer time scales. Major parts of the Netherlands subsidence, including the Western Scheldt. The exact amount of subsidence of the subsurface is challenging to establish. Measurements on the reference datum stations in the Netherlands suggest subsidence amounts to 20 mm/century (following numbers in Brand et al., 2004; De Bruijne et al., 2005) in the Western Scheldt region. The contribution of subsidence of the subsurface in terms of volume is limited to circa 60,000 m<sup>3</sup>/year for the Western Scheldt and Saeftinghe. Local compaction of Holocene peat and mud deposits may add to the apparent sediment loss through subsidence, but this is not easy to estimate. The contribution of subsidence is unidirectional and becomes essential when considering sediment budgets over more extended periods (>50 years). Based on data analysis from 1987 to 2017, Hijma and Kooi (2018) conclude that subsidence in the Western-Scheldt estuary results in an estimated loss of 0.059 million m<sup>3</sup>/year (±0.03 million m<sup>3</sup>/year). In the Dutch part of the mouth of the estuary, subsidence introduces a loss of 0.101 million m<sup>3</sup>/year (±0.053 million m<sup>3</sup>/year). Subsidence values for the Belgian part of the mouth are unknown, but we assume these losses are smaller than the value observed in the Netherlands. These small values do not significantly impact the sediment budget presented in this study and are therefore not included in the analysis.

### *Sediment budget of the Western Scheldt estuary*

Analyses of bathymetric data revealed that the Western Scheldt estuary lost 58 million m<sup>3</sup> of sediment between 1955 and 2020, which is −0.9 million m<sup>3</sup> per year. The gross annual volume change is two orders of magnitude greater: 105 million m<sup>3</sup> per year. Dredging, disposal of the dredged material and sand mining caused a significant part of this volume change. After accounting for these activities, the estuary gained 84 million m<sup>3</sup> of sediment (+1.3 million m<sup>3</sup> per year).

The volumetric changes of the various morphodynamic units in the estuary illustrate that erosion dominates the main channel: −4.2 million m<sup>3</sup> per year, with near-equal contributions of the western and eastern parts of the estuary. This increase in channel volume is caused mainly by the maintenance dredging of sills and structural deepening of the navigation channel. The secondary channels and shoals accreted at a rate of 3.3 million m<sup>3</sup> per year, three-quarters of which accumulated in the western part of the estuary.

When comparing the net volume changes in the eastern and western parts of the estuary, opposing trends were found: the western part gained 19 million m<sup>3</sup> in sediment volume, whereas the eastern part lost 77 million m<sup>3</sup>. The net sediment volume gain of the western part of the estuary is small, considering the net 65 million m<sup>3</sup> of disposal in the system. Taking into account losses from dredging, disposal and sand mining, we calculate a total sediment loss in the western part of 46 million m<sup>3</sup>. The deepening of the main channel causes sediment volume changes in the eastern part of the estuary. The most considerable erosion rates occurred from 1970 to



1980 and 1994 to 2005, during and after periods of channel deepening. The volume development corrected for human interventions shows no net transport in or out of the eastern part of the estuary until 1980 and a net gain of 3.0 million m<sup>3</sup> per year since. The annually dredged net volume in the eastern part of the estuary of 4 million m<sup>3</sup> implies an intensive sediment exchange from west to east.

### *Sediment exchange between estuary and mouth area*

Sediment import at the mouth of the estuary is expected to be higher than the observed volume change corrected for interventions in the estuary, since sediment losses to the Sea Scheldt and Land van Saefthinge need to be accounted for. These losses are estimated based on values presented in literature. The total sediment exchange rates at the mouth of the estuary were derived from resolving the estuary-wide sediment budget for various estimates (scenarios) and summarised in Fig. 11. Sediment import rates are calculated to range between 1.6 and 2.6 million m<sup>3</sup>/year between 1955 and 2020. Our best estimate scenario, using the results of detailed sediment budgets of the Sea Scheldt and accounting for the contribution of mud and sand, equals 2.2 million m<sup>3</sup>/year. The import rates are dominated by the sand fraction as the volumetric changes in the mud-rich areas along the estuary's banks are limited (0.3 million m<sup>3</sup> per year) compared to the change in the sand-dominated channels and shoals (1.9 million m<sup>3</sup>/year).

Significant variations in sediment exchange rates occurred between 1955 and 2020. Lower import rates occur between 1970 and 1980 (1.85 million m<sup>3</sup>/year), higher rates are observed between 1955–1970 and 1980–1994 (2.9 million m<sup>3</sup>/year) and maximum rates occur between 2005 and 2020 (3.7 million m<sup>3</sup>/year). Temporarily, a net export of 1.3 million m<sup>3</sup>/year occurred between 1994 and 2005. The temporary reversal from import to export at the mouth of the estuary between 1994 and 2005 is likely linked to a changed disposal strategy. Between 1997 and 2010, most of the dredged material was deposited in the western part of the estuary. As a result, the available source volume in the central part of the estuary significantly exceeded the sediment demand in the eastern part. Consequently, the surplus of sediment in the western part of the estuary was balanced through sediment export at the mouth.

### *Sediment transport mechanisms; natural versus anthropogenic forcing*

Both natural and anthropogenic sediment transport induce morphological changes in the Western Scheldt estuary. Natural sediment transport refers to sediment movement that can be related directly to the hydrodynamics with tidal asymmetry and tidal amplification being important indicators (Fig. 8). The observed development of the tidal amplification in the estuary can be explained by the morphological changes. The deepening of the channels increases the flow conveyance capacity, and the accretion at the shoals decreases the tidal storage, favouring increased tidal amplification. This can be observed, comparing Figs. 8 and 11, which illustrates:

- the increasing trend of tidal amplification for the entire period (1970–2020) corresponds with the deepening (decreasing sediment volume) trend of the estuary. Most tidal amplification and deepening changes occur in the eastern part of the estuary.
- the most notable changes, increasing tidal amplification and decreasing sediment volume, occurred in 1970–1982 and 1995–2009,

- the recent decreasing trend of the tidal amplification since around 2010 corresponds to the sediment accretion in the same period,

The development of the sediment transport and that of the tidal asymmetry do not correspond sufficiently. The vertical tide at all stations shows persistent flood dominance since 1970 as the period of the rising tide is shorter than that of the falling tide (Fig. 8). The temporary reversal in net sediment transport at the mouth that is computed from the sediment budget is apparently not caused by a change in tidal asymmetry. In addition, a continuous increase of the import rate at the mouth would be expected over the period 1970–1995, as the tidal amplification and flood dominance in the estuary were increasing. It is unlikely that other types of tidal asymmetry, for example due to the difference between the durations of the HW and LW slacks, that were not considered in the water level analysis explain the mismatch between the development of sediment transport and tidal asymmetry. This latter asymmetry is relevant for fine sediment, but has limited influence on the sand fraction that dominates the volumetric changes since 1970.

Anthropogenic transport is the most probable explanation for the observed variations in sediment import and export rates. Pieters *et al.* (1991) indicate that before dredging, from 1920 to 1930, the equilibrium depth of the sills in the navigation channel was around 8 to 10 m. As, due to the channel deepening, the present-day channel depth of NAP -16 m is far past its equilibrium depth, it becomes an effective sediment trap as the channel tries to retain an equilibrium between its cross-sectional area relative to the tidal prism (O'Brien, 1931, 1969). As a result, not only maintenance dredging increased from less than 0.5 million m<sup>3</sup>/year before 1950 to 7–10 million m<sup>3</sup>/year at present, but as most of the dredged material is disposed back in the estuary, the disposal increased at a similar rate. It is probable that this disposal and the disposal strategy plays an important role in the observed variations in sediment exchange rates at the mouth. Observations of sediment movement in the navigation channel after a flow slide (Van Schaick, 2015) and studies of the development of disposal on the Zuidgors flat (De Vet *et al.*, 2020) indicated that disposed material is more easily transported than the autochthonous sediment.

So far, evaluation of the dredging and disposal strategies was focused on retaining the multichannel character of the estuary and its associated high ecological value (Van Dijk *et al.*, 2021). The macrocell-based sediment budget (Fig. 13) provides further evidence of the importance of the disposal strategy on the sediment exchange at the mouth of the estuary. Before 1970, most of the maintenance dredging and disposal occurred in the estuary's eastern part. At that time, the secondary channels and shoals acted as sediment sources, satisfying about 50% of the sediment demand of the main channel. Consequently, the sediment transport between the western and eastern parts of the estuary was limited. Since 1970 the secondary channels and shoals supplied limited volumes to the main channel, despite the disposal of dredged sediments, causing a sand deficit in the remainder of the system and increased sediment import. The channel deepening possibly caused this behaviour change and has remained since. With increased dredging rates in the main channel, sediment transport from the western to the eastern part of the estuary has increased considerably since 1980. This observation suggests that the western part of the estuary acts as a sediment source for the eastern part if sufficient sediment is available.

Excessive disposal in the western part of the estuary can lead to a reversal of the sediment transport direction at the mouth from import to export, as observed between 1994 and 2005. However, this does not imply that the Western Scheldt estuary exports sediment and temporarily supplies sediment to the coastal zone. The development of the western part of the estuary (Mc1–4) and the mouth area are more connected than previously assumed. Depending on sediment availability in the western part of the estuary (sink or source), the mouth area facing the estuary delivers or stores sediments temporarily. Therefore, the mouth area should form an integral part of the analysis.

### Concluding remarks

A globally unique dataset of long-term bathymetric maps (1955–2020) of high resolution (in time and space) digital data, hydrodynamic observations and well-kept records of dredging, disposal and sand mining allowed for detailed investigations of the morphodynamics, sediment budget and forcing processes of the Western Scheldt estuary.

Understanding the morphodynamic functioning of the Western Scheldt estuary and its response to changes in hydrodynamics (natural sediment transport) and anthropogenic influence (e.g., channel deepening, sand mining and channel maintenance dredging and sediment relocation strategies) is essential for a sustainable management of this estuary.

The sediment budget reveals an average net sediment import of 2.2 million m<sup>3</sup>/year between 1955 and 2020 to balance sediment losses to the Sea Scheldt and Saefthinge (−0.97 million m<sup>3</sup>/year), net sand extraction (−2.14 million m<sup>3</sup>/year), main channel deepening (−4.21 million m<sup>3</sup>/year) and secondary channel and shoal accretion (+3.30 million m<sup>3</sup>/year). However, the sediment import rates varied over this interval and from 1994 to 2005 the sediment import reversed to export.

Natural variations in the hydrodynamics (e.g., tidal asymmetry and amplification) and sediment transports cannot explain the derived temporal variations in sediment import rates. Instead, anthropogenic sediment transport plays a dominant role. Deepening of the main channel led to an increase in maintenance dredging and disposal from less than 0.5 million m<sup>3</sup>/year before 1950 to 7–10 million m<sup>3</sup>/year at present. Since the first channel deepening, the tidal flats and secondary channels in the eastern part of the estuary do no longer actively exchange sediment with the main channel. As a result, sediment disposal in these areas leads to long-term storage of sand and, consequently, an increase in bed level and a sand deficit in the remainder of the system that induces increased sediment imports. Large-scale dredge disposal in the western part of the estuary can (temporarily) reverse the sediment exchange between the estuary and its mouth area, as observed between 1994 and 2005.

Safe navigation, shoreline management and the ecological targets of the Western Scheldt estuary require balanced sediment management at the scale of the macrocells and with a timescale of decades. The realisation that the disposal strategy directly impacts the sediment import and export rates provides opportunities for the future management of the estuary. Adverse effects of sea-level rise, such as drowning of tidal flats, may be mitigated by selecting a disposal strategy that promotes sediment import. However, such strategy may come at a cost, as the mouth area is the most probable source to deliver these sediments.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/njg.2023.11>.

**Acknowledgements.** The paper summarises the results of research conducted as part of the projects: ‘KPP Beheer en Onderhoud Kust’ and ‘KPP Schelde-estuarium’. The latter is part of ‘Research and Monitoring for the Flemish Dutch Scheldt Committee (VNSC)’. Both projects aim to improve our understanding of the evolution of the large-scale morphological behaviour of the Scheldt Estuary and the Dutch coastal system. Marco Schrijver, Gert-Jan Liek and Marian Lazar (all at Rijkswaterstaat Sea and Delta) are thanked for pleasant collaboration, advice, review and assistance in obtaining the datasets presented here.

The comments of the reviewers greatly improved the manuscript and are much appreciated.

Open access to the Vaklodgingen is provided by Deltares at: <https://svn.oss.deltares.nl/repos/openearthrawdata/trunk/rijkswaterstaat/vaklodgingen/>

### References

- Antwerp Port Authority*, 2012. Morphological management of estuaries – Case study of the Scheldt estuary – Base report. Antwerp: 44 pp.
- Baeyens, W., van Eck, B., Lambert, C., Wollast, R. & Goeyens, L.*, 1997. General description of the Scheldt estuary. *Hydrobiologia* **366**(1/3): 1–14.
- Bakker, Th.J.N.P. & De Looff, D.*, 1977. Onderzoek naar de mogelijkheden tot en de gevolgen van zandwinning in de Westerschelde. Report WWKZ-77.4. Rijkswaterstaat (The Hague): 58 pp.
- Barbier, E.B., Hacker, S.D., Kennedy, C., Koch, E.W., Stier, A.C. & Silliman, B.R.*, 2011. The value of estuarine and coastal ecosystem services. *Ecological Monographs* **81**(2): 169–193.
- Barneveld, H.J., Nicolai, R.P., van Veen, M., van Haaster, S., Boudewijn, T.J., de Jong, J.W., van Didderen, K., van de Haterd, R.J.W., Middenveld, P.P., Michielsen, S., van de Moortel, I., Velez, C. & de Wild, E.*, 2018. Analyserapport, T2015-rapportage Schelde-estuarium. Report PR3152. HKV Lijn in Water (Lelystad): 886 pp (in Dutch).
- Bastmeijer, K.*, 2016. Natura 2000 and the protection of wilderness in Europe. In: Bastmeijer, K. (ed.): *Wilderness Protection in Europe. The Role of International, European and National Law*. Cambridge University Press: 177–198. DOI: [10.1017/CBO9781107415287.008](https://doi.org/10.1017/CBO9781107415287.008).
- Belmans, H.*, 1988. Verdiepings- en onderhouds baggerwerken in Wester- en Zeeschelde. *Water* **43**: 184–194 (in Dutch).
- Biguino, B., Haigh, I.D., Dias, J.M. & Brito, A.C.*, 2023. Climate change in estuarine systems: patterns and gaps using a meta-analysis approach. *Science of the Total Environment* **858**: 159742. DOI: [10.1016/j.scitotenv.2022.159742](https://doi.org/10.1016/j.scitotenv.2022.159742).
- Brand, G.B.M., van Brussel, G., ten Damme, J.H. & Gerritsen, J.*, 2004. Herberekening van het primaire net van het NAP, Verbetering precisie en betrouwbaarheid ten behoeve van de nieuwe NAP publicatie. Report AGI/GAP-04/004. Rijkswaterstaat AGI (Delft) (in Dutch).
- Chen, Y., Dong, J., Xiao, X., Zhang, M., Tian, B., Zhou, Y., et al.* 2016. Land claim and loss of tidal flats in the Yangtze Estuary. *Scientific Reports* **6**(1): 24018. DOI: [10.1038/srep24018](https://doi.org/10.1038/srep24018).
- Claessens, J.*, 1988. Het hydraulisch regime van de Schelde. *Water* **43**: 163–169 (in Dutch, with English summary).
- Claessens, J., van Hoof, J. & de Ruig, J.H.M.*, 1991. Interactie morfologie en baggerwerken. *Water* **60**: 182–189 (in Dutch).
- Cleveringa, J.*, 2013. Grootchalige sedimentbalans van de Westerschelde. Project LTV- Veiligheid en Toegankelijkheid. LTV V&T-Rapport K-17. Report Arcadis (Emmeloord): 82 pp (in Dutch).
- Coen, I.*, 1988. Ontstaan en ontwikkeling van de Westerschelde. *Water* **43**: 156–162 (in Dutch).
- Dam, G.*, 2017. Lange-termijn sedimentbalans van de Westerschelde. Report 1778/U16516/D/GD. Svasek Hydraulics: 64 pp (in Dutch).
- Dam, G. & Cleveringa, J.*, 2013. De rol van het slib in de sedimentbalans van de Westerschelde. Report G3; 1630/U12376/C/GD. Svasek Hydraulics: 44 pp (in Dutch).
- Dam, G., van der Wegen, M., Taal, M. & van der Spek, A.*, 2022. Contrasting behaviour of sand and mud in a long-term sediment budget

- of the Western Scheldt estuary. *Sedimentology* **69**(5): 2267–2283. DOI: [10.1111/sed.12992](https://doi.org/10.1111/sed.12992).
- Dams, J., Vanlede, J., Plancke, Y., Verwaest, T. & Mostaert, F.**, 2016. Slibbalans Zeeschelde: Deelrapport 3 – Literatuurstudie. Version 4.0. Report WL 00\_029. Flanders Hydraulics Research & Vrije Universiteit (Antwerp & Brussel) (in Dutch).
- Davis, R.A. & Hayes, M.O.**, 1984. What is a wave-dominated coast? *Marine Geology* **60**(1-4): 313–329.
- De Boer, P.L.**, 1979. Convolute lamination in modern sands of the estuary of the Oosterschelde, the Netherlands, formed as the result of entrapped air. *Sedimentology* **26**(2): 283–294.
- De Bruijne, A., van Buren, J., Kösters, A. & van der Marel, H.**, 2005. De geodetische referentiestelsels van Nederland. Definitie en vastlegging van ETRS89, RD en NAP en hun onderlinge relaties / Geodetic reference frames in the Netherlands. Definition and specification of ETRS89, RD and NAP, and their mutual relationships. Publication 43, Nederlandse Commissie voor Geodesie / Netherlands Geodetic Commission (Delft): 117 pp (in both Dutch and English).
- De Bruin, M.P. & Wilderom, M.H.**, 1961. Tussen Afsluitdammen en Deltadijken; Part 1; Noord-Beveland. Van Benthem & Jutting (Middelburg): 304 pp (in Dutch).
- De Vet, P.L.M., van Prooijen, B.C., Colosimo, I., Ysebaert, T., Herman, P.M.J. & Wang, Z.B.**, 2020. Sediment disposals in estuarine channels alter the ecomorphology of intertidal flats. *Journal of Geophysical Research: Earth Surface* **125**(2): e2019JF005432. DOI: [10.1029/2019jfo05432](https://doi.org/10.1029/2019jfo05432).
- De Vriend, H.J., Wang, Z.B., Ysebaert, T., Herman, P.M.J. & Ding, P.X.**, 2011. Eco-morphological problems in the Yangtze estuary and the Western Scheldt. *Wetlands* **31**(6): 1033–1042. DOI: [10.1007/s13157-011-0239-7](https://doi.org/10.1007/s13157-011-0239-7).
- Debergh, H., Lescauwae, A.K., Deneudt, K., Mees, J. & Hernandez, F.**, 2009. Indicatoren van duurzame ontwikkeling voor het Schelde-estuarium. IDO-Schelde. Literatuurstudie voor de onderbouwing van de opmaak van een set indicatoren van duurzame ontwikkeling voor het Schelde-estuarium in het kader van de langetermijnvisie 2030. Eindversie juni 2009. Report Vlaams Instituut voor de Zee VLIZ (Oostende): 44 pp (in Dutch).
- Dekker, L.**, 2012. Monitoring vooroeversuppleties Oostgat, Zeeuws-Vlaanderen t/m November 2011 en Zwakke Schakels t/m maart 2012 (eindrapport). Report Rijkswaterstaat Zeeland (Middelburg): 58 pp (in Dutch).
- Depreiter, D., Sas, M., Beirincx, K. & Liek, G.J.**, 2012. Flexible Disposal Strategy: monitoring as a key to understanding and steering environmental responses to dredging and disposal in the Scheldt Estuary. In: Dorst, L.L., *et al.* (eds.): Taking Care of the Sea, Conference Proceedings of Hydro12. Hydrographic Society Benelux (Rotterdam): 8 pp. <https://doi.org/10.3990/2.232>
- Dronkers, J.**, 1986. Tidal asymmetry and estuarine morphology. *Netherlands Journal of Sea Research* **20**(2-3): 117–131.
- Elias, E.P.L. & van der Spek, A.J.F.**, 2015. Uitwerking sedimentbudget van de Westerscheldemonding. Report 1210304-012. Deltares (Delft) (in Dutch).
- Elias, E.P.L., van der Spek, A.J.F. & Lazar, M.**, 2016. The 'Voordelta', the contiguous ebb-tidal deltas in the SW Netherlands: large-scale morphological changes and sediment budget 1965-2013; impacts of large-scale engineering. *Netherlands Journal of Geosciences* **96**(3): 233–259. DOI: [10.1017/njg.2016.37](https://doi.org/10.1017/njg.2016.37).
- Friedrichs, C.T. & Aubrey, D.G.**, 1988. Non-linear tidal distortion in shallow well-mixed estuaries: a synthesis. *Estuarine, Coastal and Shelf Science* **27**(5): 521–545.
- Frings, R.M., Hillebrand, G., Gehres, N., Banhold, K., Schriever, S. & Hoffmann, T.**, 2019. From source to mouth: basin-scale morphodynamics of the Rhine River. *Earth-Science Reviews* **196**: 102830. DOI: [10.1016/j.earscirev.2019.04.002](https://doi.org/10.1016/j.earscirev.2019.04.002).
- Gerritsen, F. & de Jong, H.**, 1983. Stabiliteit en doorstroombprofielen in de Westerschelde. Report WWKZ-83.V008. Rijkswaterstaat (Vlissingen): 38 pp (in Dutch).
- Graveland, J., Dauwe, B. & Kornman, B.**, 2002. Waardering voor de Westerschelde: voorstel voor beoordelingscriteria gebaseerd op inventarisaties van de ecologische toestand, gebruik, beleid en beoordelingsmethoden. Report RIKZ-2002.53. Rijkswaterstaat (Middelburg): 93 pp (in Dutch).
- Grujters, S.H.L.L., Schokker, J. & Veldkamp, J.G.**, 2004. Kartering moeilijk erodeerbare lagen in het Schelde estuarium. Report 03.213.B1208. Netherlands Institute of Applied Geosciences NITG (Utrecht): 17 pp (in Dutch).
- Haecon**, 2006. Actualisatie van de zandbalans van de Zee- en Westerschelde. Report 1249760008/lvp (in Dutch).
- Hijma, M. & Kooi, H.**, 2018. Bodemdaling in het kustfundament en de getijdenbekkens (deel 2). Een update, case IJmuiden en kwantificering onzekerheden door geologische processen en menselijke activiteiten. Report 11202190-001. Deltares (Delft): 59 pp (in Dutch).
- Houtekamer, N.L.**, 1995. Het Verdrinken Land van Saefinghe: schor onder druk? Report AX 94.018/RIKZ 94.033. Rijkswaterstaat (Middelburg) (in Dutch).
- IMDC**, 2020. Monitoringprogramma Flexibel Storten Westerschelde en Monitoringprogramma Beneden-Zeeschelde. Voortgangsrapportage 2018-2019: Data- en analyserapport. Report I/RA/11498/20.096/MMO. IMDC (Antwerpen): 316 pp (in Dutch).
- IMDC**, 2021. Vaarwegbeheer 2016-2021 Bestelopdracht 14: Flexibel Storten Periode 3 - 2020-2021. Maandrapport december 2020 - januari 2021. Report I/RA/11498/21.004/CPA/. IMDC (Antwerpen): 101 pp (in Dutch).
- Jeuken, M.C.J.L.** 2000. On the morphologic behavior of tidal channels in the Westerschelde estuary. Ph.D. Thesis, Utrecht University: 378 pp.
- Jeuken, M.C.J.L. & Wang, Z.B.**, 2010. Impact of dredging and dumping on the stability of ebb-flood channel systems. *Coastal Engineering* **57**(6): 553–566. DOI: [10.1016/j.coastaleng.2009.12.004](https://doi.org/10.1016/j.coastaleng.2009.12.004).
- Kerner, M.** 2007. Effects of deepening the Elbe Estuary on sediment regimen and water quality. *Estuarine, Coastal and Shelf Science* **75**(4): 492–500. DOI: [10.1016/j.ecss.2007.05.033](https://doi.org/10.1016/j.ecss.2007.05.033).
- Kornman, B., Liek, G.J. & Schippers, H.**, 2003. Baggeren en Storten in de Westerschelde – een nieuwe kijk op het onderhoudsbaggerwerk. Project Zeekennis/MOVE. Report RIKZ/AB/2002.840x. Rijkswaterstaat (Middelburg) (in Dutch).
- Liek, G.A.**, 2001. Monitoring van de effecten van de verruiming 48'-43'; beschrijving van de fysische toestand van de Westerschelde t/m 2000, rapport 5. Report RIKZ/2001.023. Rijkswaterstaat (Middelburg) (in Dutch).
- Lodder, Q. & Slinger, J.**, 2022. The 'Research for Policy' cycle in Dutch coastal flood risk management: the Coastal Genesis 2 research programme. *Ocean and Coastal Management* **219**: 106066. DOI: [10.1016/j.ocecoaman.2022.106066](https://doi.org/10.1016/j.ocecoaman.2022.106066).
- Marijs, K. & Paré, E.**, 2004. Nauwkeurigheid vaklodgingen Westerschelde en monding - "de praktijk". Report ZLMD-04.N.004. Rijkswaterstaat (Zeeland): 46 pp (in Dutch).
- Mastbergen, D.R., Beinssen, K. & Nédélec, Y.**, 2019. Watching the beach steadily disappearing: the evolution of understanding of retrogressive breach failures. *Journal of Marine Science and Engineering* **7**(10): 368. DOI: [10.3390/jmse7100368](https://doi.org/10.3390/jmse7100368).
- McLaren, P.**, 1993. Patterns of sediment transport in the western part of the Westerschelde. Technical Report. GeoSea Consulting (Cambridge, UK).
- McLaren, P.**, 1994. Sediment Transport in the Western Scheldt between Baarland and Rupelmonde. Report GeoSea Consulting (Cambridge, UK).
- Mulder, J.P.M., Hommes, S. & Horstman, E.M.**, 2011. Implementation of coastal erosion management in the Netherlands. *Ocean and Coastal Management* **54**(12): 888–897.
- Nederbragt, G. & Koomans, R.L.**, 2006. Nourishment of the slope of a tidal channel - From experiment to practice. In: Sanchez-Arcilla, A. (Ed.), *Coastal Dynamics 2005 - State of the Practice*, Proceedings Coastal Dynamics 2005: 1–10. [https://doi.org/10.1061/40855\(214\)55](https://doi.org/10.1061/40855(214)55).
- Nederbragt, G. & Liek, G.**, 2004. Beschrijving zandbalans Westerschelde en monding. Report RIKZ/2004.020. Rijkswaterstaat (The Hague): 70 pp (in Dutch).
- Nederhoff, C.M.**, 2016. Eerstelijnsrapportage Westerschelde 2015. Meetgegevens van 1996 t/m 2015. Report 1209394-000. Deltares (Delft): 249 pp (in Dutch).
- O'Brien, M.P.**, 1931. Estuary tidal prism related to entrance areas. *Civil Engineering* **1**(8): 738–739.
- O'Brien, M.P.**, 1969. Equilibrium flow areas of inlets on sandy coasts. *Journal of the Waterways and Harbors Division. Proceedings of the American Society of Civil Engineers* **95**, WW1: 43–53.
- Perluka, R., Wiegmann, E.B., Jordans, R.W.L. & Swart, L.M.T.**, 2006. Opnametechnieken Waddenzee. Report AGI-2006-GPMP-004. Rijkswaterstaat (Delft): 47 pp (in Dutch).



- Pieters, T.**, 1993. Het Schelde estuarium, beheren of beheersen? Samenvatting van actuele beleids- en beheersrelevante inzichten uit het project OOST-WEST, betreffende de fysieke structuur van het Schelde estuarium. RWS, rapport DGW-93.032.
- Pieters, T., Storm, C., Walhout, T. & Ysebaert, T. (eds.)**, 1991. Het Schelde-Estuarium, meer dan een Vaarweg. Report GWWS-91.081. Rijkswaterstaat (Middelburg): 135 pp (in Dutch).
- Plancke, Y., Maris, T., Verleye, T. & Sandra, M.**, 2022. Scheldt estuary. In: Dauwe, S., et al. (eds.): Knowledge Guide Coast and Sea 2022 - Compendium for Coast and Sea. Flanders Marine Institute VLIZ (Ostend): 225–236.
- Plancke, Y., Vos, G., Meire, D. & Mostaert, F.**, 2021. Sedimentstrategie Schelde-estuarium: Deelrapport 4 – Sedimentbalans Zeeschelde, Rupel en Durme voor de periode 2011-2016. Versie 0.1. Report WL 19\_076\_4. Flanders Hydraulics Research (Antwerp) (in Dutch).
- Plancke, Y.M.G., Ides, S.J., Vos, G.R., Roose, F. & Peters, J.J.**, 2010. A new disposal strategy for the Western Scheldt, conciliating port accessibility and nature preservation. In: Setting the Course - 32nd PIANC Congress, Liverpool, UK. PIANC MMX Papers: 1–11.
- Reid, L.M. & Dunne, T.**, 1996. Rapid Evaluation of Sediment Budgets. Catena Verlag GmbH (Reiskirchen): 147 pp.
- Roose, F., Plancke, Y. & Ides, S.**, 2008. A Synthesis on the Assessment of an Alternative Disposal Strategy to Serve Sustainability in the Scheldt Estuary. In: CEDA Dredging Days 2008: Dredging facing Sustainability. Antwerp (Belgium): 1–13.
- Rosati, J.**, 2005. Concepts in sediment budgets. Journal of Coastal Research 21(2): 301–322.
- Santermans, J.**, 2013. Achtergrondrapport Baggeren en Storten. Analyse van sediment- eigenschappen, baggerstatistiek, stortintensiteitskaarten en zandwinning in de Wester- en Beneden-Zeeschelde. Achtergronddocument, versie 2.0. Report LTV V&T A-31 (in Dutch).
- Schrijver, M.C.**, 2021. De balans opgemaakt. De ontwikkelingen in de Westerschelde tot 2019 bezien vanuit de sedimentbalans. Report Rijkswaterstaat Zee en Delta (Middelburg): 92 pp (in Dutch).
- Soresma**, 2013. Ontwikkeling van een intergetijdengebied in Hedwige- en Prosperpolder: Besluit-MER /plan-MER Nederland – tekstbundel. Definitief MER\_Nederland\_revisie\_07. Report Soresma /1294103000 - Oranjewoud/225938: 627 pp (in Dutch).
- Speer, P.E., Aubrey, D.G. & Friedrichs, C.T.**, 1991. Non-linear hydrodynamics of shallow tidal inlet/bay systems. In: Parker, B.B. (ed.): Tidal Hydrodynamics. Wiley (New York): 321–339.
- Storm, C., Bollebakker, P., de Jong, J. & Mol, G.**, 1993. Nauwkeurigheid zandbalans Westerschelde 1965-1990 en aanbevelingen ter optimalisatie. Report RIKZ-94.008. Rijkswaterstaat (Middelburg): 38 pp (in Dutch).
- Storm, K. & Pieters, T.**, 1994. Het verdrinken land verlandt; Het Verdrinken land van Saeftinge een uniek gebied nu en in de toekomst. Brochure Rijkswaterstaat (in Dutch).
- Taal, M., Wang, Z.B. & Cleveringa, J.**, 2013. LTV Veiligheid en Toegankelijkheid G-13: Synthese en conceptueel model Basisrapport grootschalige ontwikkeling. Report 1204420-000. Deltares (Delft): 80 pp (in Dutch).
- Toffolon, M. & Crosato, A.**, 2007. Developing macroscale indicators for estuarine morphology: the case of the Scheldt estuary. Journal of Coastal Research 23: 195–212.
- Uit den Boogaard, L.A.**, 1995. Resultaten zandbalans Westerschelde 1955-1993. IMAU Rapport R95-08. Utrecht University: 280 pp (in Dutch).
- Van Braeckel, A., Piesschaert, F. & van den Bergh, E.**, 2006. Historische analyse van de Zeeschelde en haar getijgebonden zijrivieren (19de eeuw tot heden). Report INBO.R.2006.29. Instituut voor Natuur- en Bosonderzoek (Brussels) (in Dutch).
- Van Cauwenberghe, C.** 1971. Hydrografische analyse van de Vlaamse banken langs de Belgisch-Fransekust. Het Ingenieursblad 40(19): 563–576.
- Van Dam, T.**, 2014. Morfologie en hydrodynamica in de Beneden-Zeeschelde. Ontwikkelingen van 1960 tot 2010. Report TVD/14311/1111. Svasek Hydraulics (Rotterdam): 113 pp (in Dutch).
- Van den Berg, J.H., Jeuken, M.C.J.L. & van der Spek, A.J.F.**, 1996. Hydraulic processes affecting the morphology and evolution of the Western-Scheldt estuary. In: Nordstorm, K.F., & Roman, C.T. (eds.): Estuarine Shores: Evolution, Environments and Human Alterations. John Wiley (London): 157–184.
- Van der Slikke, M.J.**, 1998. Grootschalige en interne zandbalans Westerscheldemonding (1969-1993). Report IMAU R98-05. Utrecht University (in Dutch).
- Van der Spek, A.J.F.**, 1997. Tidal asymmetry and long-term evolution of Holocene tidal basins in The Netherlands: simulation of paleo-tides in the Schelde estuary. Marine Geology 141(1-4): 71–90.
- Van der Spek, A.J.F. & Elias, E.P.L.**, 2021. Half a century of morphological change in the Haringvliet and Grevelingen ebb-tidal deltas (SW Netherlands) - Impacts of large-scale engineering 1964-2012. Marine Geology 432: 106404. DOI: [10.1016/j.margeo.2020.106404](https://doi.org/10.1016/j.margeo.2020.106404).
- Van der Vegt, H., Mastbergen, D. & van der Werf, J.J.**, 2020. Moeilijk-erodeerbare lagen in de Westerschelde - Onzekerheden en gevolgen voor morfodynamiek. Report 1210301-015-ZKS-0014. Deltares (Delft) (in Dutch).
- Van der Vegt, H., van Onselen, E. & van der Spek, A.**, 2021. Erosie-resistente lagen in het Westerschelde mondingsgebied en de invloed daarvan op de pilot-suppletie. Report 11203725-001-ZKS-0012. Deltares (Delft) (in Dutch).
- Van der Zijp, N.J.**, 2001. Controle en bewerking Jaarlijkse Kustmetingen en Vakkloddingen: Functioneel Ontwerp. Rapport MobiData (Rotterdam): 56 pp (in Dutch).
- Van Dijk, W.M., Cox, J.R., Leuven, J.R.F.W., Cleveringa, J., Taal, M., Hiatt, M.R., Sonke, W., Verbeek, K., Speckmann, B. & Kleinhans, M.**, 2021. The vulnerability of tidal flats and multi-channel estuaries to dredging and disposal. Anthropocene Coasts 4: 36–60.
- Van Enckevoort, I.**, 1996. Morfologisch onderzoek Westerscheldemonding; deel 2. Morfologische ontwikkeling van de Westerschelde monding sinds 1800. Report IMAU R 96-21. Utrecht University: 56 pp (in Dutch).
- Van Holland, G.**, 2013. Ontwikkeling mesoschaal Zeeschelde. Report I/RA/11387/13.112/GVH. Consortium Deltares-IMDC-Svasek-Arcadis: 131 pp (in Dutch).
- Van Kessel, T., van der Werf, J.J. & Giardino, A.** 2012. Analyse havenstortvakken Westerschelde. Report 1205999-000. Deltares (Delft): 108 pp (in Dutch).
- Van Kessel, T., Vroom, J. & Taal, M.**, 2023. A conceptual framework for fine sediment dynamics in the Scheldt estuary - Providing a framework for data and model analyses and impact assessments. Report 11208065-000. Deltares (Delft): 59 pp.
- Van Maldegem, D.C., Mulder, H.P.J. & Langerak, A.**, 1993. A cohesive sediment balance for the Scheldt estuary. Netherlands Journal of Aquatic Ecology 27(2-4): 247–256.
- van Maren, D.S., van Kessel, T., Cronin, K. & Sittoni, L.**, 2015. The impact of channel deepening and dredging on estuarine sediment concentration. Continental Shelf Research 95: 1–14.
- Van Schaick, S.J.**, 2015. Morphological development after the July 2014 flow slide on the tidal flat of Walsoorden in the Western Scheldt. MSc. Thesis, Delft University of Technology: 111 pp.
- Van Veen, J., van der Spek, A.J.F., Stive, M.J.F. & Zitman, T.**, 2005. Ebb and flood channel systems in the Netherlands tidal waters. Journal of Coastal Research 21: 1107–1120.
- Vandenbruwaene, W., Levy, Y., Plancke, Y., Vanlede, J., Verwaest, T. & Mostaert, F.**, 2017. Integraal plan Boven-Zeeschelde: Deelrapport 3 – Sedimentbalans Zeeschelde, Rupel en Durme. Versie 5.0. Report WL13\_131. Flanders Hydraulics Research (Antwerp) (in Dutch).
- Vandenbruwaene, W., Pauwaert, Z., Meire, D., Plancke, Y., Deschamps, M. & Mostaert, F.**, 2019. Agenda voor de Toekomst – Historische evolutie getij en morfologie Schelde estuarium: Deelrapport 1 Evolutie van het getij over de periode 1888-2017. Versie 5.0. Report WL 14\_147\_1. Flanders Hydraulics Research (Antwerp) (in Dutch).
- Vandenbruwaene, W., Plancke, Y., Verwaest, T. & Mostaert, F.**, 2013. Interestuarine comparison: Hydro-geomorphology. Hydro- and geomorphodynamics of the TIDE estuaries Scheldt, Elbe, Weser and Humber. Version 4. Report WL 770\_62b. Flanders Hydraulics Research (Antwerp). [https://www.tide-toolbox.eu/pdf/reports/WL2013R770\\_62b\\_rev4\\_20130220\\_TIDE\\_IC\\_HGM.pdf](https://www.tide-toolbox.eu/pdf/reports/WL2013R770_62b_rev4_20130220_TIDE_IC_HGM.pdf).
- Vandenbruwaene, W., Stark, J., Plancke, Y. & Mostaert, F.**, 2020. Agenda voor de Toekomst – Historische evolutie getij en morfologie Schelde estuarium: Deelrapport 5 – Synthese. Versie 4.0. WL Report WL 14\_147\_5. Flanders Hydraulics Research Antwerp (in Dutch).

- Verheyen, B. & van Holland, G.**, 2013. Zeeschelde - nota analyse bodemligging. Basisrapport kleinschalige ontwikkeling K-19, Report LTV - Veiligheid en Toegankelijkheid Zeeschelde. IMDC (Antwerp): 109 pp (in Dutch).
- Vos, G., Plancke, Y., Meire, D. & Mostaert, F.**, 2021. Sedimentstrategie Schelde-estuarium: Deelrapport 5 – Sedimentbalans Zeeschelde, Rupel en Durme voor de periode 2016-2019. Versie 0.1. Report WL, 19\_076\_6. Flanders Hydraulics Research (Antwerp) (in Dutch).
- Vos, P.C.**, 2015. Origin of the Dutch coastal landscape. Long-term landscape evolution of the Netherlands during the Holocene, described and visualized in national, regional and local palaeogeographical map series. PhD thesis, Utrecht University. Barkhuis (Groningen): 359 pp.
- Wang, Z.B., Jeuken, C.J.L., Gerritsen, H., de Vriend, H.J. & Kornman, B.A.**, 2002. Morphology and asymmetry of vertical tide in the Westerschelde estuary. *Continental Shelf Research* **22**(17): 2599–2609.
- Wang, Z.B., Vandenbruwaene, W., Taal, M. & Winterwerp, H.**, 2019. Amplification and deformation of tidal wave in the Upper Scheldt Estuary. *Ocean Dynamics* **69**(7): 829–839. DOI: [10.1007/s10236-019-01281-3](https://doi.org/10.1007/s10236-019-01281-3).
- Wang, Z.B. & Winterwerp, J.C.** Impact of dredging and dumping on the stability of ebb-flood channel systems. In: *Proceedings River, Coastal and Estuarine Morphodynamics Conference RCEM2001 (IAHR)*, Japan, 2001.
- Watson, I. & Finkl, C.W.**, 1990. State of the art in storm-surge protection: the Netherlands Delta project. *Journal of Coastal Research* **6**: 739–765.
- Watson, I. & Finkl, C.W.**, 1992. Simplified technical summary of the complete Delta Works, including the Eastern Scheldt. *Journal of Coastal Research* **10**: 56.
- Wiegmann, E.B., Perluka, R., Oude Elberink, S. & Vogelzang, J.**, 2005. Vaklodgingen: De inwintechieken en hun combinaties. Report AGI-2005-GSMH-012. Rijkswaterstaat (Delft): 47 pp (in Dutch).
- Wijnberg, K.M.**, 2002. Environmental controls on decadal morphologic behaviour of the Holland coast. *Marine Geology* **189**(3-4): 227–247.
- Winterwerp, J.C., Wang, Z.B., Stive, M.J.F., Arends, A., Jeuken, M.C.J.L., Kuijper, C. & Thoolen, P.M.C.**, 2001. A new morphological schematisation of the Western Scheldt estuary, The Netherlands. In: *Proceedings 2nd IAHR symposium on River, Coastal and Estuarine Morphodynamics*, Sept 2001, Obihiro, Japan: 525–534.
- Ysebaert, T., Meire, P., Maes, D. & Buijs, J.**, 1993. The benthic macrofauna along the estuarine gradient of the Scheldt estuary. *Netherlands Journal of Aquatic Ecology* **27**(2-4): 327–341.