Original Research Article

Ex-situ conservation of Palms with special reference to endemic and IUCN Red List species in Acharya Jagadish Chandra Bose Indian Botanic Garden, Howrah, India

ABSTRACT

The present paper deals with the holistic approach of palms diversity, endemism, conservation status, economic importance of different species of palms in Acharya Jagadish Chandra Bose Indian Botanic Garden (AJCBIBG) are discussed.

Keywords: AJCBIBG, Ex-situ conservation, Endemic and Threatened Palms

1. INTRODUCTION

Palms are the most fascinating group of plants that attracts attention of scientific community all over the world [1]. The family Arecaceae (Palmae nom. alt.) is represented by 2600 species belongs to 181 genera under 28 tribes of 5 subfamilies [2] and primarily distributed in tropical and subtropical regions of the world. Palms have a variety of growth forms such as shrubs, lianas and trees [3] and they are considered to be the economically important plants after to the grasses and legumes by providing food, shelter and other utility commodities for the mankind especially for the rural communities in the tropics [4,5,6,7]. In India, the family Arecaceae represented by *c*. 219 species belongs to 71 genera under 5 subfamilies [8]. Among the 219 taxa present in India, 46 taxa belonging to 9 genera, accounting for 21 percentare endemic to country [9,10] which are mainly distributed in Peninsular India, North Eastern India and Andaman and Nicobar Islands and other 125 species are well introduced and cultivated [1] for their socio-economic values.

Palms shows rich species diversity and its ability to grow in a wide range of habitats from rainforests to deserts enables the garden to develop a dedicated section of palms i.e., 'Palmetum' in the Division no.5 of Acharya Jagadish Chandra Bose Indian Botanical Garden (AJCBIBG). It is worth mentioning that, the popular Large Palm House (LPH), an age-old greenhouse repository for equatorial palms was established by Sir George King and designed by the Govt. Architect, Mr. E.J. Martin during the year 1881-82. The plantation work under the conservatory from different parts of the globe had been started from 1883. The Palm House serves its purpose for ex-situ conservation of germplasm for a large quantity of endemic and threatened species. The main attraction in the LPH is Double Coconut Palm i.e., Lodoiceamaldivica(J.F.Gmel.) Pers. which is famous for its largest seed, average living span 1200 years [11]Other unique representatives of this family present in this garden are Egyptian branching palm [(Hyphaene thebaica (L.) Mart.)], Indian branching Palm [(Hyphaene dichotoma (D.White bis ex Nimmo) Furtado)] and Tali palm [(CoryphatalieraRoxb.)], of these later species is already extinct from its natural habitat/wild [12].

Palms are an integral part of socio-cultural and family-based economy of many communities due to its applicability in construction, food and fodder for human and animals, boat making, handcrafts, rituals, medicinal and therapeutics. While working on the regular projects viz., Curatorial work in the garden and maintenance and introduction and *ex-situ* conservation of RET species in AJCBIBG, to know the exact species (Palms) in the garden for further introduction of other RET species, authors aimed to document palm diversity in the AJCBIBG.

2. STUDY AREA

AJC Bose Indian Botanic Garden also known as 'Company Bagan', or 'Royal Botanic Garden' is one of the oldest and largest botanic gardens of South East Asia. It is also considered as one of the best landscaped gardens of the World. The garden is spread over a

sprawling expanse of over 110 hectares (273 acres). It has 24 interconnected lakes, and the lakes are connected to the Ganges through sluices for the regular inlet and outlet of water. The gardens exhibit a wide variety of species including endemic, threatened, medicinal, economical, ornamental plants etc. in 25 divisions[13,14,15,16]. In 1787 Col. Robert Kyd established this garden with a view to introduce socio-economically important trees from all over the globe and now it is serving the purpose for *ex-situ* conservation of endemic, threatened, economically important plants etc. At present this garden is a living repository of about 1400 species [16] including palms.

3. REVIEW OF LITERATURE

A scrutiny of literature revealed that initial documentation on Indian Palm diversity with 9 species was carried out by Van Rheededuring1692-1693 [17] in his Hortus Malabaricus. Later Linneaus[18] included these species in his Species Plantarum. Roxburgh [19] described 41 species of palms in his Flora Indica. Martius [20] included 15 Indian species in his Historia Naturalis Palmarum. Griffith [21] described 96 palms in his Palms of British East India. Beccari and Hooker[22] described 71 species of palms in Hooker's Flora of British India(1892 -1894). Later Beccari [23,24,25,26,27] was also published series of papers on Malayan, Asiatic and Indian palms. Blatter [28] was given huge information palms in his work on The Palms of British India and Ceylon. Taxonomic accounts of palms occurring in the respective regions or provinces appeared in the regional Floras done by different workers viz., Bengal [29], Upper Gangetic Plains [30], Bombay [31], Bihar and Orissa [32], Madras [33] and Andaman and Nicobar Islands [34].

Botanical Survey of India, Howrah was studied considerable volume of work on the taxonomy and phytogeography of Indian palms [35]. Mahabale[36] gave a detailed account on morphology in his Palms of India. Renuka [37,38] gave an account of Rattans of Western Ghats. Lakshmana [39] worked on rattans of South India. Basu and

Chakraverthy[1]published a manual of cultivated palms in India. Rawat [40] brought out a hand book Palmsfor India. Renuka et al. [41] published Rattans of India. Bhat [4] published Palms of Karnataka, in this book he described 56 species belonging to 39 genera of palms. Renuka and Sreekumar [5] brought out a field guide to the palms of India in this book they reported 105 species belonging to 22 genera from India. Some important scientific works regarding the taxonomy, phytogeography on Indian palms from different regions of India were published by various authors [8,11,15,42,43,44,45,46,47,48,49,50].

4. MATERIAL AND METHODS

Regular field visits to the Palmetum and other sections in the garden were conducted during 2022-2023 to cover all the palm diversity. During the regular field visit, palm species were captured with Nikon Coolpix P900 Digital Camera when it is in flower and fruit and few important specimens were collected further identification. All the collected plant specimens and images were identified with the help of various palm literature [1,5,21,22,28] and other recent pictorial guides and relevant taxonomic papers were consulted wherever is required. The identified plant names checked in The International Plant Name Index [51] and The Plants of the World Online [52] for accepted names, basionyms and synonyms.

5. RESULTS AND DISCUSSIONS

The documentation of palms in AJCBIBG revealed that 110 taxa belonging to 46 genera and 4 subfamilies (Table 1), of which 31 species under 16 genera were indigenous, which were introduced from various regions of India and a total 68 taxa were introduced from more than 50 regions/countries namely Africa, Arabian peninsula, Australia, Borneo, Brazil, Central America, China, East Asia, Florida, South East Haiti, Indonesia, Jamaica, Lesser Sunda Islands, Madagascar, Malesia, Mauritius, Mexico, New Guinea, Philippines, Rodrigues, Seychelles, South America, South East Asia, Taiwan, Thailand, Tonga, Virgin Islands (Puerto Rico) etc. The garden plays an important role by creating a hub for more than 40%

native species diversity of Palms in India. Also, the *ex-situ* conservation of various rare, endangered and threatened germplasm will pave pathway for future taxonomic research and enrichment of floristic diversity.



Fig. 1. a. Arial view of Palmetum; b. Large Palm House; c. Palmetum; d. Corypha taliera Roxb.; e. Bentinckia nicobarica (Kurz) Becc.; F. Calamus andamanicus Kurz; g. Hyphaene thebaica (L.) Mart.; h. Hyophorbe lagenicaulis (L.H.Bailey) H.E.Moore

5.1. Conservation of Endemic Palms

Endemic/threatened plants are of high biological and conservation value. In fact, endemic floristic elements of a country or geographical region throw light on the biogeography of the area, centres of speciation, areas of extinction, vicariance and adaptive evolution of the flora occurring in the area [53]. Insofar 11 species under 8 genera are represented in AJC Bose Indian Botanic Garden which are endemic to the different regions of our country (Table 1). The endemic species in the garden are: Arenga wightii Griff. (W. Ghats: Goa, Karnataka, Kerala, Maharashtra, Tamil Nādu), Bentinckiacondapanna Berry ex Roxb. (W. Ghats: Kerala, Tamil Nādu), Bentinckianicobarica (Kurz) Becc. (Nicobar Islands), Calamus andamanicus Kurz. (Andaman and Nicobar Islands), Calamus baratangensis Renuka & Vijayak. (Andaman Islands), Calamus rheedei Griff. (W. Ghats: Karnataka, Kerala), Daemonoropsrarispinosa Renuka & Vijayak. (Andaman and Nicobar Islands), Phoenix rupicola T. Anderson (E. Himalaya: West Bengal), PinangamaniiBecc. (Andaman and Nicobar Islands), Rhopaloblasteaugusta (Kurz) H.E.Moore (Nicobar Island), and TrachycarpustakilBecc. (W. Himalaya: Uttarakhand). These palms were collected from various phytogeographical regions and successfully established in the garden by many botanists/staff of the garden. Maintenance of the 24% Indian endemic palms in the garden has highly conservation value.

5.2. Conservation of Threatened Palms

Of the 219 recorded species in the country, it is found that the about 95 species (43%) of palms are in the IUCN Red List. Of the conserved 108 species in the garden, it is found that 46 species (43%) are IUCN Red List species. These species are categorised under 7 different threat categories (Table 1). Tali Palm (*Coryphataliera*Roxb.) was discovered in 1919 by William Roxburgh and he considered it to be endemic to Bengal [19,54,55]. The last record of this palm growing in the wild was in Birbhum district of West Bengal (India) in a village near Shantiniketan [1,55,56]. A single individual is available in University Campus of

Dhaka, Bangladesh and assessed as Extinct in the Wild (EW) in the year 1998 by Johnson [12], further he declared the identity of the species in University Campus of Dhaka is uncertain. At present a single individual is confirmed in the palmetum section of AJCBIBG, Howrah.

Fourteen (14) threatened species are conserved in the garden, of these three (03) are Critically Endangered (CR) species namely Carrossier Palm(Attalea crassispatha(Mart.) Burret) which is native of South East Haiti, Bottle Palm (Hyophorbelagenicaulis (L.H.Bailey) H.E.Moore) from Mauritius and Spindle Palm Hyophorbeverschaffeltii (W.Bull ex J.Dix) H.Wendl. from Rodrigues were introduced and conserved in the garden. Four (04) Endangered species namelyBentickia Palm(Bentinckianicobarica (Kurz) Becc.) introduced from Nicobar Islands (India); Blue laten Palm(Latania loddigesii Mart.) from Mauritius; Red laten Palm (Latania lontaroides (Gaertn.) H.E.Moore) from Reunion Islands; Halifax fan Palm(Livistonadrudei F. Muell. ex Drude) were introduced from Australia. Seven (7)Vulnerable species introduced from were various countries. Manila Palm(Adonidiamerrillii (Becc.) Becc.) from Philippines, Wight's Sago Palm(Arenga wightii Griff.) from South East Asia, Hill areca nut (Bentinckiacondapanna Berry ex Roxb.) introduced from India (Western Ghats: Kerala, Tamil Nādu), Triangular Palm(Dypsisdecaryi (Jum.) Beentje &J.Dransf.) from South East Madagascar, Taraw Palm(Livistonasaribus(Lour.) Merr. ex A.Chev.) from South East Asia, Black Palm(Normanbyanormanbyi (W. Hill) L. H. Bailey) from Australia, Nicobar Majestic Palm(Rhopaloblasteaugusta(Kurz) H.E. Moore) from India (Nicobar Islands) were introduced.

Apart from 14 threatened species 33 other IUCN category species are also conserved in the garden. Of the 33 species, Nearly Threatened (NT) species (3) are *Arenga tremula* (Blanco) Becc., *Hyphaene dichotoma* (D. White bis ex Nimmo) Furtado and *Phoenix*

rupicola T. Anderson. Least Concerned (LC) category species (28)are Acoelorraphewrightii(Griseb. H. Wendl.) & H. Wendl. ex Becc., & Archontophoenixalexandrae(F. Muell.) H. Wendl. Drude, Archontophoenixcunninghamiana(H. Wendl.) H. Wendl. & Drude, Areca triandraRoxb. ex Buch.-Ham., Arenga obtusifolia Mart., Arenga westerhoutii Griff., Attalea cohune Mart., Attalea speciosa Mart., Bactris major Jacq., Bismarckia nobilis Hildebrandt & H. Wendl., Caryota mitis Lour., Caryotaurens L., Chrysalidocarpusmadagascariensis (D.T.Fish) Becc., Coccothrinaxargentata (Jacq.) L. H. Bailey, Coryphautan Lam., Elaeisguineensis Jacq., Heterospatheelata Scheff., Hyphaene coriacea Gaertn., Hyphaene thebaica (L.) Mart., Livistona decora (W. Bull) Dowe, Phoenix loureiroiKunth, Phoenix reclinata Jacq., Ptychosperma elegans (R. Br.) Blume, Roystonea regia (Kunth) O. F. Cook, Sabal mauritiiformis (H. Karst.) Griseb. & H. Wendl, Sabal palmetto (Walter) Lodd. ex Schult. &Schult.f., Syagrusromanzoffiana (Cham.) Glassman, Washingtoniafilifera(T. Moore & Mast.) H. Wendl. ex de Bary Data Deficient (DD) species are Areca catechu L. and Coryphaumbraculifera L.

Table 1. List of Palms (Arecaceae) with their distribution, IUCN and endemic status, present in AJCBIBG

S. No.	Scientific Name	Common name	Distribution	IUCN/Ende
				mic status
1.	Acoelorraphewrightii(Griseb. &	Silver Saw Palm	West Indies,	LC
	H. Wendl.) H. Wendl. ex Becc.		Florida	
2.	Adonidiamerrillii (Becc.) Becc.	Manila Palm	Philippines	VU
3.	Aiphaneshorrida (Jacq.) Burret	Ruffle Palm	Trinidad to S.	
			America	
4.	Archontophoenix alexandrae (F.	Alexander Palm	Eastern Australia	LC
	Muell.) H. Wendl. & Drude			
5.	Archontophoenixcunninghamiana	Banglow Palm	Eastern Australia	LC
	(H. Wendl.) H. Wendl. & Drude			
6.	Areca catechu L.	Betel Nut Palm	South East Asia	DD
7.	Areca triandraRoxb. ex Buch	Wild Areca Palm	East Asia	LC
	Ham.			
8.	Arenga australasica(H. Wendl. &	Australian Arenga	Australia	
	Drude) S. T. Blake ex H. E.	Palm		

	Moore			
9.	Arenga caudata(Lour.) H. E. Moore	Hooker's Fish Tail Palm	South East Asia	
10.	Arenga engleriBecc.	Taiwan Sugar Palm	Taiwan	
11.	Arenga obtusifolia Mart.	Sumatra Sugar Palm	South East Asia	LC
12.	Arenga pinnata (Wurmb) Merr.	Sugar Palm	South East Asia	
13.	Arenga tremula (Blanco) Becc.	Philippine Dwarf Sugar Palm	South East Asia	NT
14.	Arenga undulatifoliaBecc.	Jaka Palm	South East Asia	
15.	Arenga westerhoutii Griff.	Sugar Palm	Eastern Himalaya	LC
16.	Arenga wightii Griff.	Wight's Sago Palm	South East Asia	VU/ Westerr Ghats
17.	Attalea cohune Mart.	Cohune Palm	Mexico, Central America	LC
18.	Attalea crassispatha (Mart.) Burret	Carrossier Palm	South East Haiti	CR
19.	Attalea speciosa Mart.	Babassu Palm	South America	LC
20.	Bactris major Jacq.	Peach Palm	South America	LC
21.	Bentinckiacondapanna Berry ex Roxb.	Hill Areca Nut	South India	VU/ Western Ghats
22.	Bentinckianicobarica (Kurz) Becc.	Bentickia Palm	Nicobar Islands	EN/ Nicobar Islands
23.	Bismarckia nobilis Hildebrandt & H. Wendl.	Bismarckia Palm	Madagascar	LC
24.	Borassus flabellifer L.	Toddy Palm	India, South East Asia	
25.	Butia capitata (Mart.) Becc.	Butia Palm	Brazil	
26.	Calamus andamanicus Kurz.	Andaman Cane Palm	Andaman Islands	Andaman and Nicobar Islands
27.	Calamus arborescens Griff.	Sweet Flag	India, South East Asia	
28.	Calamus baratangensis Renuka &Vijayak.	Baratang Island Palm	Andaman Islands	Andaman Islands
29.	Calamus erectusRoxb.	Cane Fruit Palm	India	
30.	Calamus floribundus Griff.	Myanmar Cane	E. Himalaya to N. Myanmar	
31.	Calamus gurubaBuchHam. ex Mart.	Climbing Palm	South East Asia	
32.	Calamus latifolius Roxb.	Gouri Bet	E. Nepal to Indo- China and Peninsula Malaysia	
33.	Calamus leptospadix Griff.	Himalayan Rattan Palm	India	
34.	Calamus longisetus Griff.	Rattan Palm	India, Thailand	
35.	Calamus melanochaetes(Blume)	Giant Devil Palm	E. Himalaya to	

	Miq.		Taiwan	
36.	Calamus metzianusSchltdl.	Odiyan-Chooral	SW. India	
37.	Calamus rheedei Griff.	Vetra	W. Ghats, Karnataka, Kerala	Western Ghats
38.	Calamus rotang L.	Common Rattan	SE. India, Sri Lanka	
39.	Calamus viminalis Willd.	Bitter Rattan Palm	NE. India to China	
40.	Carpentaria acuminata (H. Wendl. & Drude) Becc.	Carpentaria Palm	Australia	
41.	Caryota mitis Lour.	Burmese Fishtail Palm	India, South East Asia	LC
42.	Caryotaurens L.	Fishtail Wine Palm	India, Myanmar	LC
43.	Chamaedorea elegans Mart.	Parlor Palm	Mexico to Honduras	
44.	ChrysalidocarpuslutescensH. Wendl.	Golden Cane Palm	Madagascar	
45.	Chrysalidocarpusmadagascariens is (D. T. Fish) Becc.	Butterfly Palm	Madagascar	LC
46.	Coccothrinaxargentata(Jacq.) L. H. Bailey	Florida Silver Palm	Florida, Mexico	LC
47.	Cocos nucifera L.	Coconut	Indian sub- continent, Oceania	
48.	CoryphatalieraRoxb.	Tali Palm	India, Bangladesh	EW
49.	Coryphaumbraculifera L.	Talipot Palm	Indian sub- continent	DD
50.	Coryphautan Lam.	Cabbage Palm	Indian sub- continent	LC
51.	Cyrtostachysrenda Blume	Sealing Wax Palm	Thailand to W. Malesia	
52.	Daemonoropsjenkinsiana (Griff.) Mart.	Major Jenkins Palm	Indian sub- continent	
53.	Daemonoropsrarispinosa Renuka & Vijayak.	Rattan Palm	Andaman and Nicobar Islands	Andaman and Nicoba Islands
54.	Dictyosperma album (Bory) Scheff.	Princess Palm	Coastal Forest of Mascarene	
55.	Dypsisdecaryi(Jum.) Beentje & J. Dransf.	Triangular Palm	SE. Madagascar	VU
56.	Elaeisguineensis Jacq.	Oil Palm	Africa	LC
57.	Heterospatheelata Scheff.	Sagisi Palm	Australia	LC
58.	Hyophorbelagenicaulis (L. H. Bailey) H. E. Moore	Bottle Palm	Mauritius	CR
59.	Hyophorbe verschaffeltii (W. Bull ex J. Dix) H. Wendl.	Spindle Palm	Rodrigues	CR
60.	Hyphaene coriacea Gaertn.	Ilala Palm	Ethiopia to S. Africa, Madagascar	LC
61.	Hyphaene dichotoma (D. White bis ex Nimmo) Furtado	Branching Palm	India	NT

				1
62.	Hyphaene thebaica (L.) Mart	Egyptian Doub Palm	Arabain peninsula	LC
63.	Kerriodoxa elegans J. Dransf.	White Backed Palm	Thailand	
64.	Latania loddigesiiMart.	Blue Laten Palm	Mauritius	EN
65.	Latania lontaroides(Gaertn.) H.E.Moore	Red Laten Palm	Reunion Islands	EN
66.	Licuala grandis (T. Moore) H. Wendl.	Palas Palm	Vanuatu Islands	
67.	Licuala peltataRoxb. ex Buch Ham.	Swamp Fan Palm	Himalaya to South East Asia	
68.	Licuala spicata Becc.	Sarawak Mangrov e Palm	Borneo	
69.	Licuala spinosa Wurmb	Mangrove Fan Palm	South East Asia	
70.	Livistona chinensis (Jacq.) R. Br. ex Mart.	Chinese Fan Palm	East Asia	
71.	Livistona decora (W. Bull) Dowe	Ribbon Fan Palm	Australia	LC
72.	<i>Livistonadrudei</i> F. Muell. ex Drude	Halifax Fan Palm	Australia	EN
73.	LivistonajenkinsianaGriff.	Fan Palm	Indian sub- continent	
74.	Livistonasaribus(Lour.) Merr. ex A. Chev.	Taraw Palm	South East Asia	VU
75.	Lodoiceamaldivica(J. F. Gmel.) Pers.	Double Coconut	Seychelles	
76.	Normanbyanormanbyi(W. Hill) L. H. Bailey	Black Palm	Australia	VU
77.	Phoenix acaulis Roxb.	Dwarf Date Palm	Himalaya to Bangladesh	
78.	Phoenix dactylifera L.	Date Palm	Arabian Peninsula to Pakistan	
79.	Phoenix loureiroiKunth	Mountain Date Palm	Indian Subcontinent	LC
80.	Phoenix pusillaGaertn.	Ceylon Date Palm	South East Asia	
81.	Phoenix reclinata Jacq.	Wild Date Palm	Africa	LC
82.	Phoenix roebelenii O'Brien	Pygmy Date Palm	South East Asia	
83.	Phoenix rupicola T. Anderson	Cliff Date Palm	India	NT/ Eastern Himalaya, West Benga
84.	Phoenix sylvestris (L.) Roxb.	Silver Date Palm	India	
85.	Pinangacoronata (Blume) Blume	Ivory Cane Palm	Lesser Sunda Islands	
86.	PinangamaniiBecc.	Pinanga Palm	Andaman and Nicobar Islands	Andaman and Nicobar Islands
87.	Pritchardiapacifica Seem. & H. Wendl.	Fiji Fan Palm	Tonga	

88.	Ptychosperma elegans (R.Br.) Blume	Elegant Palm	Australia	LC
89.	Ptychospermamacarthurii(H. Wendl. ex H. J. Veitch) H. Wendl. ex Hook.f.	Macarthur Palm	Australia	
90.	PtychospermasanderianumRidl.	Sanderianum Palm	New Guinea	
91.	Rhapisexcelsa (Thunb.) A. Henry	Lady Palm	China	
92.	Rhapis humilis Blume	Slender Lady Palm	China	
93.	Rhopaloblasteaugusta (Kurz) H. E. Moore	Nicobar Majestic Palm	Nicobar Islands	VU/ Nicobar Islands
94.	Roystoneaborinquena O. F. Cook	Rico Royal Palm	Hispaniola to Virgin Islands	
95.	Roystonea regia (Kunth) O. F. Cook	Royal Palm	Florida, Mexico	LC
96.	Sabal mauritiiformis (H.Karst.) Griseb. & H.Wendl.	Bayleaf Palm	Central America	LC
97.	Sabal palmetto (Walter) Lodd. ex Schult. &Schult.f.	Cabbage Palm	USA	LC
98.	Salaccazalacca (Gaertn.) voss	Snake Palm	Indonesia	
99.	Saribusrotundifolius(Lam.) Blume	Footstool Palm	South East Asia	
100.	Syagrusromanzoffiana(Cham.) Glassman	Queen Palm	South America	LC
101.	Syagrusschizophylla(Mart.) Glassman	Arikury Palm	Brazil	
102.	Thrinax parviflora Sw.	Broom Palm	Jamaica	
103.	<i>Trachycarpusfortunei</i> (Hook.) H. Wendl.	Chinese Windmill Palm	India (Sikkim), China	
104.	Trachycarpusmartianus(Wall. ex Mart.) H. Wendl.	Martius Fan Palm	North East India and S.E. Asia	
105.	TrachycarpustakilBecc.	Kumaon Palm	W. Himalaya, Uttarakhand	Western Himalaya, Uttarakhand
106.	VeitchiaarecinaBecc.	Montogomery Palm	VanautuIslanda	
107.	WallichiadistichaT. Anderson	Distichous Fishtail Palm	North East India	
108.	Wallichiatriandra(J. Joseph) S. K. Basu	Tibetian Arenga Palm	SE. Tibet to Arunachal Pradesh	
109.	Washingtoniafilifera(T. Moore & Mast.) H. Wendl. ex de Bary	California Fan Palm	USA	LC
110.	WodyetiabifurcataA. K. Irvine	Foxtail Palm	Australia	

6. CONCLUSION

The palm diversity in AJC Bose Indian Botanic Garden possesses many horticultural and economic importance. Starting from very common Coconut (*Cocos nucifera* L.), date (*Phoenix dactylifera* L.) to the species extinct in wild *Coryphataliera*Roxb. Palm remains very useful in socio-economic context. *Cocos nucifera*, *Phoenix dactylifera* L., *Borassus flabellifer*L. are taken as fruits; *Elaeisguineensis* Jacq. vegetable oil is extracted; *Attalea speciosa* Mart., *Bentinckianicobarica* (Kurz) Becc. and *Rhopaloblasteaugusta* (Kurz) H. E. Moore are useful in construction purposes indigenously, the leaves of *Borassus* L. can be useful in fan making, plate making; *Adonidiamerrillii* (Becc.) Becc., *Hyophorbelagenicaulis* (L. H. Bailey) H. E. Moore, *Hyphaene thebaica* (L.) Mart., Royal Palm, foxtail palm etc. are useful in landscaping.

It is worthy to mention that at present situation when everyone is looking for a sustainable, organic, environment friendly option in their daily lifestyle coconut coir andvegetable oil, plates and fan made from palm leaves will definitely be in huge demand in coming future. Government is also focusing on making nature friendly options over the plastic or readymade plates which not only boost small industry and Self-Help Group (SHGs) making these items also promote 'Make in India' campaign. Conservation of this highly important representative will create an awareness among common people as well create area of research.

REFERENCES

- Basu SK, Chakraverty RK. A Manual of Cultivated Palms in India. Botanical Survey of India, Calcutta; 1994.
- 2. Baker WJ, Dransfield J. Beyond Genera Palmarum: progress and prospects in palm systematics. Botanical Journal of the Linnean Society. 2016; 182(2): 207-233.

- 3. Muscarella R, Emilio T, Phillips OL, Lewis SL, Slik F, Baker WJ, Couvreur TL, Eiserhardt WL, Svenning JC, Affum- Baffoe K, Aiba SI. The global abundance of tree palms. Global Ecology and Biogeography. 2020; 29(9): 1495-1514.
- 4. Bhat KG. Palms of Karnataka. K. Gopalakrishna Bhat. 'Madhuca', Srinivasa Nagara, Chitpady, Udupi. 2011; 1-100.
- Renuka C, Sreekumar VB. A field guide to the palms of India. Kerala Forest Research Institute. 2012.
- 6. Qureshimatva Q, Gamit SB, Solanki HA. Arecaceae family in Gujrat, its morphological character-based identification keys. India Forester. 2020; 146 (2): 113–124.
- 7. Mehmud S, Roy H. Diversity and distribution of palms (Arecaceae) in Assam, India. Check List. 2021; 17(1): 69-93.
- 8. Hameed SS. In Mao AA, Dash S. (eds.). Flowering Plants of India: An Annotated Checklist. Volume 3. Botanical Survey of India, Kolkata. 2020.
- 9. Kulkarni AR, Mulani RM. Indigenous palms of India. Current science. 2004; 86(12): 1598-1603.
- Singh P, Karthigeyan K, Lakshminarasimhan P, Dash SS. Endemic Vascular Plants of India. Botanical Survey of India, Kolkata. 2015.
- 11. Hameed SS. A maiden fruit set in the 'Double Coconut' [*Lodoiceamaldivica* (J.F.Gmel.) Pers.- Arecaceae] in India by artificial pollination. Phytotaxonomy. 2015; 14: 69-75.
- Johnson D. *Coryphataliera*. The IUCN Red List of Threatened Species 1998:
 e.T38493A10118302.http://dx.doi.org/10.2305/IUCN.UK.1998.RLTS.T38493A1011830
 2.en. Accessed on 21 November 2023.
- 13. Chowdhery HJ, Pandey DS. 2007. Plants of Indian Botanic Garden. Botanical Survey of India.

- 14. Debnath H, Mahapatra HS, Hameed SS, Sreekumar PV. Census of plants in AJC Bose Indian Botanic Garden: A Report. Botanical Survey of India. Kolkata. 2014.
- 15. Hameed SS. Artificial pollination and fruit set in double coconut growing in India. Current Science. 2016; 110(6): 976-978.
- 16. Anonymous. Acharya Jagadish Chandra Bose Indian Botanic Garden. EIACP, Botanical Survey of India, Kolkata. 2023; 1-30.
- 17. Rheede DH. Hortus Indicus Malabaricus. Johannis van Some¬ren, et Joannis van Dyck, Amsterdam.1678.
- 18. Linnaeus C. Species Plantarum. Stockholm. 1753.
- Roxburgh W. Flora Indica: Descriptions of Indian Plants. Serampore: Printed for W. Thacker and co., Calcutta. 1832.
- 20. Martius, C E P van 1823-1850. Historia Naturalis Palmarum. Vol. 1-3. Munich.
- 21. Griffith W. Palms of British East India: posthumous papers bequeathed to the Honourable the East India Company. Periodical Experts Book Agency. 1850.
- 22. Beccari O. Palmae. In Hooker J.D. (ed.), The Flora of British India. L. Reeve and Co. London.1892-93; 6: 402-483.
- 23. Beccari O. Asiatic palms-Lepidocaryeae. Part I. The species of Calamus. Annals of the Royal Botanic Garden, Calcutta. 1908; 11: 1–518, Pl. i–ii, 1-238.
- 24. Beccari O. Asiatic palms-Lepidocaryeae. Part II. The species of Daemonorops. Annals of the Royal Botanic Garden, Calcutta. 1911; 12(1): 1 237, Pl. i–ii, 1-109.
- 25. Beccari O. Asiatic palms-Lepidocaryeae. The species of Calamus. Supplement to Part I. Annals of the Royal Botanic Garden Calcutta. 1913; 11 (Appendix): 1-142.
- 26. Beccari O. Asiatic palms-Lepidocaryeae. The species of Calamus. Supplement to Part I. Annals of the Royal Botanic Garden Calcutta. 1914; 11 (Appendix): Pl. 1-83.

- 27. Beccari O. Asiatic palms-Lepidocaryeae. Part III. The species of the genera Ceratolobus, Calospatha, Plectocomia, Plectocomiopsis, Myrialepis, Zalacca, Pigaffeta, Korthalsia, Metroxylon, Eugeissona. Annals of the Royal Botanic Garden Calcutta. 1918; 12(2): 1–231 Pl. i–iv. 1-120.
- 28. Blatter E. The palms of British and Ceylon. Oxford University Press London. 1926; 1-203.
- 29. Prain D. Bengal Plants. Vol. 2. Botanical Survey of India, Calcutta. 1903 (Reprint 1963).
- 30. Duthie JF. Flora of Upper Gangetic Plain. Vol. 2 Botanical Survey of India, Calcutta. 1903 (Reprint 1960).
- 31. Cooke T. The Flora of the Presidency of Bombay Vol. 3 Botanical Survey of India, Calcutta. 1907 (Reprint 1958).
- 32. Haines BH. The Botany of Bihar and Orissa. Vol. 3 Botanical Survey of India, Calcutta. 1924 (Reprint 1961).
- 33. Fischer CEC. In: The Flora of the Presidency of Madras. Vol. 3 Botanical Survey of India, Calcutta. 1928 (Reprint 1957).
- 34. Parkinson CE. The Forest Flora of the Andaman Islands: An Account of the Trees, Shrubs and Principal Climbers of the Islands. India, Superintendent, Government Central Press. 1923.
- 35. Basu, SK. Landscape with Palms. Horticultural Bull. 1975; 22:1-3.
- 36. Mahabale TS. Palms of India. Maharashtra Association for the Cultivation of Science, Research Institute, Pune. 1982.
- 37. Renuka C. Rattans of Western Ghats: A Taxonomical Manual. K. F. R. I., Peechi, Kerala. 1992; 1-62.
- 38. Renuka C. A manual of the rattans of Andaman and Nicobar Islands. Kerala Forest Research Institute, Peechi, Kerala. 1992; 1-172.

- 39. Lakshmana AC. Rattan of South India. Evergreen Publisher, Bangalore, India. 1993; 1-180.
- 40. Rawat D. Palms for India: Small Tips to Achieve Big Results in Growing, SahayogHortica (P) Ltd. 2008.
- 41. Renuka C. Rattans of India: taxonomy, biology and utilization. Kerala Forest Research Institute. 2010.
- 42. Basu SK. A census of palms cultivated in the Indian Botanic Garden, Howrah. Principes. 1978; 22: 127–135.
- 43. Basu SK. The present status of rattan palms in India-an overview, pp. 77-94. In: Wong,K. M. and Manokaron, N. (Eds.). Proceedings of the Rattan Seminar. Kualalumpur,Malayasia. 1985.
- 44. Basu SK. *Corypha* palms in India. Journal of Economic and Taxonomic Botany. 1986; 11(2): 477-486.
- 45. Basu SK. Rattans (Canes) in India: A monographic revision. Rattan Information Centre, Kepong, Kuala Lumpur, Malaysia. 1992; 1-141.
- 46. Renuka C. Palms of Kerala. Kerala Forest Research Institute, Peechi, Kerala. 1999; 1-72.
- 47. Renuka C. All India coordinated Project on the Taxonomy of Palms. Kerala Forest Research Institute. 2011; 1-209.
- 48. Mondal S, Chowdhury M. Rattans diversity in West Bengal, India. Advances in Plant Sciences. 2018; 31(2): 159-165.
- 49. Mondal S. Taxonomy, phenology and ethnobotany of palms in West Bengal. Ph.D. Thesis (Unpublished). University of North Bengal. 2019.
- 50. Hameed SS, SP Panda MU Sharief. Is *Hyphaene dichotoma* (White) Furtado, an indigenous 'Branching Palm' species under habitat threat? Ecology, Environment and Conservation. 2023; 29: 365-367.

- 51. IPNI (International Plant Names Index) 2023. Published on the Internet http://www.ipni.org, The Royal Botanic Gardens, Kew, Harvard University Herbaria & Libraries and Australian National Botanic Gardens. (Retrieved: November 2023).
- 52. POWO. Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew. 2023. Accessed on 1 November 2023. Available: http://www.plantsoftheworldonline.org.
- 53. Ahmedullah M, Nayar MP. Endemic Plants of the Indian Region. Vol. 1. Botanical Survey of India, Calcutta. 1987.
- 54. Benthall AP. The Trees of Calcutta and its neighborhood. Bishen Singh Mahendra Pal Singh, Dehra Dun. 1933 (Reprint 1984).
- 55. Basu SK. Threatened Palms of India some case studies. Journal of Economic and Taxonomic Botany. 1985; 7: 493–497.
- 56. Khan MS, Hassan MA Basu SK. Rescue of an "extinct" palm in Bangladesh. Species 36:9. Newsletter of the Species Survival Commission, IUCN-World Conservation Union.2001.