



## Establishment of a non-native xyleborine ambrosia beetle, *Xyleborus monographus* (Fabricius) (Coleoptera: Curculionidae: Scolytinae), new to North America in California

ROBERT J. RABAGLIA<sup>1\*</sup>, SHERI L. SMITH<sup>2</sup>, PAUL RUGMAN-JONES<sup>3</sup>, MARC F. DIGIROLOMO<sup>4</sup>, CURTIS EWING<sup>5</sup> & AKIF ESKALEN<sup>6</sup>

<sup>1</sup>USDA Forest Service, Forest Health Protection, Washington, DC, 20250, U.S.A.

✉ [bob.rabaglia@usda.gov](mailto:bob.rabaglia@usda.gov); <https://orcid.org/0000-0001-8591-5338>

<sup>2</sup>USDA Forest Service, Forest Health Protection, Vallejo, CA 94592, U.S.A.

✉ [sheri.smith2@usda.gov](mailto:sheri.smith2@usda.gov); <https://orcid.org/0000-0002-8581-1026>

<sup>3</sup>Department of Entomology, University of California, Riverside, California, 92521, U.S.A.

✉ [paulrj@ucr.edu](mailto:paulrj@ucr.edu); <https://orcid.org/0000-0002-2949-5022>

<sup>4</sup>USDA Forest Service, Forest Health Protection, Durham, New Hampshire, 03824, U.S.A.

✉ [marc.f.digirolomo@usda.gov](mailto:marc.f.digirolomo@usda.gov); <https://orcid.org/0000-0001-7596-3109>

<sup>5</sup>California Department of Forestry and Fire Prevention, Forest Pest Management, Davis, California, 95618, U.S.A.

✉ [curtis.ewing@fire.ca.gov](mailto:curtis.ewing@fire.ca.gov); <https://orcid.org/0000-0002-4567-1655>

<sup>6</sup>Department of Plant Pathology, University of California, Davis, California, 95616, U.S.A.

✉ [aeskalen@ucdavis.edu](mailto:aeskalen@ucdavis.edu); <https://orcid.org/0000-0002-8829-7413>

\*Corresponding author

### Abstract

Specimens of an ambrosia beetle, *Xyleborus monographus* (Fabricius), were found infesting oak trees in California. This is the first record of this species established in North America. Based on collection information, this species most likely has been established in the Napa County area for several years. A modified key to *Xyleborus* in North America, and diagnosis of the species is provided.

**Key words:** Invasive ambrosia beetle, adventive species, oaks, key to species, cytochrome c oxidase subunit I, ambrosia fungi

### Introduction

Bark and ambrosia beetles (Coleoptera: Curculionidae: Scolytinae) are one of the most commonly intercepted, introduced, and established taxa in many countries (Haack & Rabaglia 2013) and in North America several of these species have had significant impacts on natural and urban forests (Fraedrich *et al.* 2008; Eskalen *et al.* 2013; Smith & Hulcr 2015; Stouthamer *et al.* 2017). Due to their wide host range, cryptic nature, fungus farming, and inbreeding, ambrosia beetles in the tribe Xyleborini have been one of the most successful groups at colonizing new and non-native habitats (Atkinson *et al.* 1990; Rabaglia *et al.* 2006; Smith & Hulcr 2015; Gomez *et al.* 2018a). Nearly half of the more than 60 species of non-native scolytines established in the continental United States are xyleborine ambrosia beetles (Gomez *et al.* 2018a, 2018b). This paper reports on the first establishment of another xyleborine ambrosia beetle, *Xyleborus monographus* (Fabricius), in North America.

In September 2019, specimens of an ambrosia beetle in a valley oak, *Quercus lobata*, were collected in Calistoga, Napa County, California. These specimens were compared to beetles in the collection at the United States Museum of Natural History, Smithsonian Institution and determined to be *Xyleborus monographus*. Additional specimens were collected in Calistoga on October 16 and 17, 2019 (see below for collection details). These collections were thought to represent the first records of this species established in North America; however, previous specimens collected in September 2017 from the same area (see below for collection details) were recently identi-

fied as *X. monographus*. By early 2020, infested trees were found throughout a 15-mile-long area in Napa County, as well as neighboring Lake and Sonoma Counties, indicating a well-established population in the area.

In 2018, one specimen of *X. monographus* was detected in a multi-funnel trap baited with an ‘oak pinhole borer’ lure (consisting of ethanol, leaf alcohol, sulcatol and sulcatone [Synergy Semiochemicals, Delta, B.C., Canada]) in Fairview, Multnomah County, Oregon (W. Williams, Oregon Department of Forestry, unpublished data), however, trapping in 2019 did not detect additional beetles nor were infested trees found in the Portland area.

Molecular confirmation of the species identification was also sought by sequencing a section of the cytochrome c oxidase subunit I gene (COI) commonly used in DNA barcoding studies (Hebert *et al.* 2003). Two beetles from the initial collection were subject to non-destructive DNA extraction and sequencing following Stouthamer *et al.* (2017). Subsequent BLAST searches of the GenBank database (Benson *et al.* 2013) revealed 100% similarity to COI sequences derived from known European specimens of *Xyleborus monographus* (GenBank accessions: HQ953471, KM286137, and KM285995; also see BOLD Barcode Index Number - BOLD:AAO0226 [Ratnasingham & Hebert 2013]). In addition, a section of the D2 region of 28S ribosomal RNA was also sequenced according to Stouthamer *et al.* (2017). Our 28S sequences represent the first deposition in GenBank attributed to *X. monographus*, but the most similar existing matches were 28S sequences attributed to the congeneric species *X. affinis* [GU808581] and *X. bispinatus* [HM099741]; both ~98.6% similar. Sequences generated from the specimens were deposited in GenBank (accession numbers MN974135-36 [COI] and MN970511-12 [28S]) and the coinciding morphological vouchers were deposited in the Entomology Research Museum, University of California, Riverside, CA.

Specimens are deposited in the following collections:

NMNH—US National Museum of Natural History, Washington, DC

RJRC—Robert J. Rabaglia Collection, Annapolis, MD

UCRC—Entomology Research Museum, University of California, Riverside, CA

EMEC—Essig Museum of Entomology, University of California, Berkeley, CA

### *Xyleborus monographus* (Fabricius, 1792)

Figure 1

*Bostrichus monographus* Fabricius, 1792: 365.

*Xyleborus monographus* (Fabricius), Eichhoff 1864: 704.

*Bostrichus tuberculosus* Herbst, 1793: 113. Synonymy Eichhoff 1878: 397.

**Diagnosis.** Specimens of *Xyleborus monographus* can be distinguished from most species of *Xyleborus* in North America by the tubercles on declivital interstriae 1 that are distinctly larger than those on other interstriae. It is very similar to the eastern species *X. celsus* Eichhoff, but *X. monographus* is smaller (3.0–3.2 mm vs approximately 3.6–4.5 mm for *X. celsus*), and the width of interstriae 2 on the declivity is approximately half the width of interstriae 3 on *X. monographus*, vs twice the width of interstriae 3 on the declivity of *X. celsus*.

This is only the fourth species of *Xyleborus* reported from California; the other three species are native. It can be distinguished from *X. ferrugineus* (F.) which lacks tubercles on interstriae 1 and has tubercles on interstriae 3 larger than on others. *Xyleborus intrusus* Blandford and *X. xylographus* (Say) both have tubercles on interstriae 1 but they are not larger than those on interstriae 3. There are five other species of xyleborine ambrosia beetles in California, *Anisandrus dispar* (F.), *Cyclorhipidion bodoanum* (Reitter), *Euwallacea fornicatus* (Eichhoff), *E. kuroshio* Gomez & Hulcr, and *Xyleborinus saxesenii* (Ratzeburg), which can be distinguished from *Xyleborus* and each other by the keys and images in Gomez *et al.* (2018a) and updated in Smith *et al.* (2019).

The key to female *Xyleborus* species in America north of Mexico in Gomez *et al.* (2018a) is modified below to include *X. monographus*. Alterations are in bold.

3. Tubercles on declivital interstriae 1 distinctly larger than tubercles on other interstriae . . . . . 4
- Tubercles on declivital interstriae 1 either similar in size to tubercles on other interstriae or absent (except at base or apex) . . . . . 5
4. **Elytral disc and declivity setose; all declivital interstriae armed by strong tubercles at base; declivital interstriae 1 armed by two very large pointed tubercles, declivital interstriae 3 armed by several smaller tubercles . . . . . 4'**

- Elytral disc and declivity glabrous; all declivital interstriae armed by small granules, gradually decreasing in size toward apex; interstriae 1 near apex armed by one or two small tubercles ..... *glabratus* Eichhoff
- 4'. **Total body length 3.6–4.5 mm; declivital interstriae 2 1.5–2x as wide as interstriae 3, punctures deep and large (Fig. 2B); in *Carya* species.....** *celsus* Eichhoff
- **Total body length 3.1–3.2 mm; width of declivital interstriae 2 .5–1x as wide as interstriae 3, punctures shallow and small (Fig. 2A); mostly in *Quercus* species.....** *monographus* (Fabricius)

**Description:** Female- Length 3.1–3.2 mm, 3 times as long as wide; color reddish brown. Frons convex, surface reticulate, not smooth, punctures sparse, shallow; setae sparse, longer near epistomal margin. Pronotum about 1.5 times as long as wide, anterior rounded and convex, coarsely asperate on anterior half, basal half smooth with shallow, sparse punctures. Elytra approximately twice as long as wide, and slightly less than twice as long as pronotum, disc shining, striae shallowly impressed, punctures shallow, interstriae with fewer, shallow punctures; declivity steep, less than 25% of elytral length, flat, surface dull, striae punctures small and shallow, in rows curving away from suture at middle of declivity, then towards suture at apex, interstriae 1 wide, smooth with two large tubercles at middle of declivity, one smaller denticle at base of declivity, interstriae 2 smooth, about half as wide as interstriae 3, two small denticles on interstriae 3 at about the same level as those on interstriae 1. Elytral setae on striae minute, in rows, interstitial setae longer and fine.

Male. Not examined.

**Distribution.** The distribution records are based on Wood & Bright (1992) and supplements (Bright & Skidmore 1997, 2002; Knižek 2011; Bright 2014).

**Africa:** Algeria, Morocco; **Asia:** Azerbaijan, Iran, Iraq, South Korea, Turkey; **Europe:** Albania, Austria, Belgium, Bulgaria, Croatia, Czech Republic, Denmark, Estonia, France, Great Britain, Germany, Greece, Hungary, Italy, Latvia, Luxemburg, Macedonia, Montenegro, Netherlands, Norway, Poland, Portugal, Romania, Russia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Ukraine.

New records in **North America:** United States: California, Napa Co., Calistoga, 14 September, 2017, M. Garbelotto coll., ex. *Quercus lobata* (2, NMNH); as previous except., 10 September, 2019, L. Burkhardt coll., ex. *Quercus lobata* (1, USNM; 2, UCRC [UCRC\_ENT 00528716 and UCRC\_ENT 00528703]); as previous except: Silverado Trail, 16 October, 2019, Sheri Smith coll., ex. *Quercus lobata* (2, RJRC); as previous except: 17 October, 2019, Sheri Smith coll., ex. *Quercus lobata* (2, RJRC); as previous except: Bothe Napa Valley State Park, 38.551877, -122.522836, 29 January, 2020, Cutis Ewing coll., ex. *Quercus kelloggii* (10, EMEC1332925-34); as previous except: 38.552298, -122.522948, 29 January, 2020, Cutis Ewing coll., ex. *Quercus douglasii* (1, EMEC1332924); as previous except.: Silverado Tr. & Brannan St., 38.5856, -122.5724, 18 November, 2019, Curtis Ewing coll., ex. *Quercus lobata* (5, EMEC 1332915-19); as previous except: vineyard between Silverado Tr. & Rt. 128, 38.57737, -122.57563, 18 November, 2019, Curtis Ewing coll., ex. *Quercus lobata* (4, EMEC 1332920-23); as previous except: Middletown Hwy., Mayacmas Mts., 38.6670, -122.5974, 10 January, 2020, Curtis Ewing coll., ex. *Quercus lobata* (4, EMEC 1332935-38); California, Lake Co., Middletown, Graham Ln., 38.7525, -122.6217, 10 January, 2020, Curtis Ewing coll., ex. *Quercus lobata* (1, EMEC 1332939); as previous except: Calistoga, Silverado Tr. & Glass Mt. Rd., 38.53529, -122.59057, 29 January, 2020, Curtis Ewing coll., ex. *Quercus lobata*, (5, EMEC 1332940-44); as previous except: Middletown Hwy., Mayacmas Mts., 38.6670, -122.5974, 02 February, 2020, Curtis Ewing coll., ex. *Quercus lobata* (4, EMEC 1332945-48).

**Hosts and biology.** In Europe, the most commonly reported hosts of *X. monographus* are various species of oaks (*Quercus*) and other genera of Fagaceae (*Fagus* and *Castanea*). Wood & Bright (1992) report *Quercus* spp., and state it is uncommon in *Castanea vesca* and *Fagus orientalis*. Bright and Skidmore (1997) cites Koch (1992) and lists: *Betula pendula* (= *B. verrucosa*), *Carpinus betulus*, *Castanea sativa*, *Fagus sylvatica*, *Fraxinus excelsior*, *Juglans regia*, *Prunus avium*, *Quercus canariensis*, *Q. castaneifolia* var. *incana*, *Q. ceris*, *Q. coccifera*, *Q. ilex*, *Q. lusitanica*, *Q. petraea*, *Q. pubescens*, *Q. pyrenaica*, *Q. robur*, *Q. rubra*, *Q. suber*, *Ulmus laevis*. Schedl (1964) states it is most frequently found in *Quercus*, but lists several of the non-oak species above as hosts also.

In California, the original infested trees were mostly valley oaks (*Quercus lobata*), but some blue oaks (*Q. douglasii*) also were found infested in the area. A very limited infestation was found in a single limb of California black oak (*Quercus kelloggii*) with extensive heart rot.

Schedl (1964) stated that he found most attacks by this species in trunks of downed oaks felled in winter or early spring, and in branches larger than 20 cm in diameter. He also stated that most attacks occurred on the sides or undersides of logs, and only rarely on the upper, sun-exposed surfaces. In California, we found a similar attack pattern by this beetle on valley oaks. Most of the trunks were heavily colonized by the beetle, but we also have seen

attacks on mostly larger branches and in branches as small as 6.35 cm. diameter in the upper crowns of apparently healthy oaks.


Data from Schedl (1964) and his reference to Eichhoff (1881) and Escherich (1929), indicated two generations per year for this species in Germany, but he questioned if this was the case throughout Europe, and cited Palm (1959) who found only one generation per year in Sweden. More recent work in Greece (Markalas & Kalapandia 1997) and Slovakia (Galko *et al.* 2014) found one generation per year based on trapping data. These later two studies also found peak trap catch in late May and June. In Israel, adults were active for nearly the entire study period, March–September, and no activity peaks were detected, suggesting multiple overlapping generations (Buse *et al.* 2013).

There have been several studies in Europe that have tested the response of ambrosia beetles to ethanol-baited funnel traps (Markalas & Kalapandia 1997; Galko *et al.* 2014 and references therein), and they have shown positive response of *X. monographus* to ultra-high release (UHR) ethanol-baited traps.

As with all xyleborine ambrosia beetles, *X. monographus* exhibits sib-mating, with haploid and wing-less males (Kirkendall 1993). Schedl (1964) reports a sex ratio of 8.5:1 females to males. Other species of xyleborines have been reported to have ratios similar to this or more females to males (Smith & Hulcr 2015), and additional studies may show a more female biased sex ratio for this species as well.



**FIGURE 1.** Dorsal and lateral habitus of *Xyleborus monographus*. Scale bar represents 0.5 mm.



**FIGURE 2.** Elytral declivities of *Xyleborus monographus* (A) and *X. celsus* (B). I-2 and I-3: Interstriae 2 and 3 respectively. Scale bars represent 0.2 mm.

**Fungal symbionts.** Ambrosia beetles carry symbiotic fungi which they introduce into the xylem and are used as food for adults and larvae (Beaver 1989). Species in the Xyleborini, as do most other ambrosia beetles, have special structures, mycangia, in which the spores of their symbiotic fungi are carried. In *Xyleborus monographus*, as in all *Xyleborus* species, the mycangia are in the mandibles (Schedl 1964). In most cases the fungal associate is not pathogenic to the host tree, however, the fungal associate of the non-native *X. glabratus* Eichhoff (the red bay ambrosia beetle), *Raffaelea lauricola* is very pathogenic and has caused extensive mortality of several species of Lauraceae in the southeastern US (Fraedrich *et al.* 2008; Harrington *et al.* 2010). Gebhardt *et al.* (2004) found *Raffaelea montetyi* associated with *X. monographus* in Germany. They also found this fungus in *X. dryographus* (Ratzeburg) and *Platypus cylindrus* (F.). Inácio *et al.* (2012) tested the pathogenicity of *R. montetyi* strains from Portugal on cork oak (*Quercus suber*) saplings and had 100% mortality within 60 days.

Ambrosial fungal species from both beetle and plant tissues infested by *X. monographus* were collected with methods of Eskalen *et al.* (2013). Specimens were collected from infested valley oaks in Calistoga, CA. A total of 10 beetles and infested wood samples were collected from each of three infested trees. Fungal isolations from symptomatic tissues and female beetle mycangia were recovered following the methods of Lynch *et al.* (2016). Based on the morphological characterization and BLAST's query comparison of the ITS sequence data in the pres-

ent study and other isolates in GenBank, the fungal species, *Raffaelea montetyi* (UCD8134), *Paecilomyces formosus* (UCD8140), *Fusarium solani* (UCD8043), undescribed species of *Fusarium* (UCD8376) and *Leptographium* sp. (UCD8382), and a yeast species, *Saccharomyces microspore* (UCD8112) were recovered.

Further identification of the fungal species using multi loci gene sequence analyses is underway. Currently, pathogenicity tests of fungal species on young valley oak trees are being conducted.

## Conclusion

At this time, it is hard to predict what impacts this species may have on the oak resources of California or North America. Most of the more than 30 species of xyleborine ambrosia beetles established in North America have had little impact on the health of forests, but, as noted above, *Xyleborus glabratus* and species in the *Euwallacea fornicatus*-complex, along with their fungal associates, have had a significant impact on trees in the southeastern US and California, respectively. Sometimes the symbiotic fungi that ambrosia beetles vector from host to host can be plant pathogens and the mutualistic relationship becomes destructive when introduced to new habitats from a native habitat such as Southeast Asia (Hulcr & Stelinski 2017). Although *X. monographus* and fungal associates have not shown pathogenicity in their native range, how these species interact with new, naïve hosts in North America still needs to be investigated.

## Acknowledgements

We would like to thank Jacob Schneider and Joseph Schneider of Pacific Tree Care for their help in locating infested trees in Napa County. We also appreciate the assistance of Christina Freeman, California State Parks, the Frediani Family, Calistoga, Solage, Auberge Resorts, Calistoga, and Spotswoode Winery, St. Helena for allowing access to their properties to examine infested trees. Thanks to M. Lourdes Chamorro for allowing access to examine specimens at the National Museum of Natural History, Washington, DC. We appreciate the information provided by Wyatt Williams, Oregon Department of Forestry for information on the collection of *X. monographus* in 2018, and James LaBonte, Oregon Department of Agriculture, for confirming the identity of that specimen.

## References

- Atkinson, T.H., Rabaglia, R.J. & Bright, D.E. (1990) Newly detected exotic species of *Xyleborus* (Scolytidae) in eastern North America, with a revised key to species. *The Canadian Entomologist*, 122, 93–104.  
<https://doi.org/10.4039/Ent12293-1>
- Beaver, R.A. (1989) Insect-fungus relationships in bark and ambrosia beetles. *Symposium of the Royal Entomological Society*, 14, 121–143.  
<https://doi.org/10.1016/B978-0-12-751800-8.50011-2>
- Benson, D.A., Cavanaugh, M., Clark, K., Karsch-Mizrachi, I., Lipman, D.J., Ostell, J. & Sayers, E.W. (2013) GenBank. *Nucleic acids research*, 41, D36–D42.  
<https://doi.org/10.1093/nar/gks1195>
- Bright, D.E. & Skidmore, R.E. (1997) *A Catalog of Scolytidae and Platypodidae (Coleoptera), Supplement 1 (1990–1994)*. NRC Research Press, Ottawa, 368 pp.
- Bright, D.E. & Skidmore, R.E. (2002) *A Catalog of Scolytidae and Platypodidae (Coleoptera), Supplement 2 (1995–1999)*. NRC Research Press, Ottawa, 611 pp.
- Bright, D.E. (2014) A catalog of Scolytidae and Platypodidae (Coleoptera), Supplement 3 (2000–2010), with notes on subfamily and tribal reclassifications. *Insecta Mundi*, 0356, 1–336.
- Buse, J., Assmann, T., Friedman, A.-L.-L., Rittner, O. & Pavlicek, T. (2013) Wood-inhabiting beetles (Coleoptera) associated with oaks in a global biodiversity hotspot: a case study and checklist for Israel. *Insect Conservation and Diversity*, 6, 687–703.  
<https://doi.org/10.1111/icad.12023>
- Eichhoff, W.J. (1864) Über die Mundtheile und die Fühlerbildung der europäischen Xylophagi sens strict. *Berliner Entomologische Zeitschrift*, 8, 17–46.  
<https://doi.org/10.1002/mmnd.18640080103>
- Eichhoff, W.J. (1878) Ratio, descriptio, emendatio eorum Tomicinorum qui sunt in Dr medic. Chapuisii et autoris ipsius col-

- lectionibus et quos praeterea recognovit scriptor. *Mémoires de la Société Entomologique de Liège*, Série 2e, 8, 1–531, pls. I–V.
- Eichhoff, W. (1881) *Die europäischen Borkenkäfer*. Springer, Berlin, 315 pp.  
<https://doi.org/10.5962/bhl.title.8968>
- Escherich, K. (1923) *Forstinsekten Mitteleuropas*. Paul Parey, Berlin, 663 pp.  
<https://doi.org/10.5962/bhl.title.27087>
- Eskalen, A., Stouthamer, R., Lynch, S.C., Twizeyimana, M., Gonzalez, A. & Thibault, T. (2013) Host range of *Fusarium dieback* and its ambrosia beetle Coleoptera: Scolytinae vector in southern California. *Plant Disease*, 97, 938–951.  
<https://doi.org/10.1094/PDIS-11-12-1026-RE>
- Fabricius J.C. (1792) *Entomologia systematica emendata et aucta. Secundum classes, ordines, genera, species adjectis synonymis, locis, observationibus, descriptionibus. Tomus I. Pars II*. C.G. Proft, Hafniae, xx + 538 pp.  
<https://doi.org/10.5962/bhl.title.122153>
- Fraedrich, S.W., Harrington, T.C., Rabaglia, R.J., Ulyshen, M.D., Mayfield, A.E., Hanula, J.L., Eickwort, J.M. & Miller, D.M. (2008) A fungal symbiont of the redbay ambrosia beetle causes a lethal wilt in redbay and other Lauraceae in the southeastern United States. *Plant Disease*, 92 (2), 215–224.  
<https://doi.org/10.1094/PDIS-92-2-0215>
- Galko, J., Nikolov, C., Kimoto, T., Kunca, A., Gubka, A., Vakula, J., Zubrik, M. & Ostrihosn, M. (2014) Attraction of ambrosia beetles to ethanol baited traps in a Slovakian oak forest. *Biologia*, 69 (10), 1376–1383.  
<https://doi.org/10.2478/s11756-014-0443-z>
- Gomez, D.F., Rabaglia, R.J., Fairbanks, K.E.O. & Hulcr, J. (2018a) North American Xyleborini north of Mexico: a review and key to genera and species (Coleoptera, Curculionidae, Scolytinae). *ZooKeys*, 768, 19–68.  
<https://doi.org/10.3897/zookeys.768.24697>
- Gomez, D.F., Skelton, J., Steining, M.S., Stouthamer, R., Rugman-Jones, P., Sittichaya, W., Rabaglia, R.J. & Hulcr, J. (2018b) Species within the *Euwallacea fornicatus* (Coleoptera: Curculionidae) complex revealed by morphometric and phylogenetic analyses. *Insect Systematics and Diversity*, 2 (6), 2, 1–11.  
<https://doi.org/10.1093/isd/ixy018>
- Haack, R.A. & Rabaglia, R.J. (2013) Exotic bark and ambrosia beetles in the USA: Potential and current invaders. In: Peña, J.E. (Ed.), *Potential Invasive Pests of Agricultural Crops*. CAB International, Boston, pp. 48–74.  
<https://doi.org/10.1079/9781845938291.0048>
- Harrington, T. C., Agheyeveva, D.N. & Fraedrich, S.W. (2010) New combinations of *Raffaelea*, *Ambrosiella*, and *Hyalorhinocladia*, and four new species from the red bay ambrosia beetle, *Xyleborus glabratus*. *Mycotaxon*, 111, 337–361.  
<https://doi.org/10.5248/111.337>
- Hebert, P.D.N., Cywinska, A., Ball, S.L. & deWard, J.R. (2003) Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London, Series B, Biological Sciences*, 270, 313–321.  
<https://doi.org/10.1098/rspb.2002.2218>
- Herbst, J.F.W. (1793) Der Käfer, fünfter Theil. In: Jablonsky, C.G. (Ed.), *Natursystem aller bekannten in- und ausländischen Insekten*. J. Pauli, Berlin, pp. 1–392.
- Hulcr, J. & Stelinski, L.L. (2017) The ambrosia symbiosis: from evolutionary ecology to practical management. *Annual Review of Entomology*, 62, 285–303.  
<https://doi.org/10.1146/annurev-ento-031616-035105>
- Inácio, M.L., Henriques, J., Lima, A. & Sousa, E. (2012) Ophiostomatoid fungi associated with cork oak mortality in Portugal. *Integrated Protection in Oak Forests*, 76, 89–92.
- Kirkendall, L.R. (1993) Ecology and evolution of biased sex ratios in bark and ambrosia beetles. In: Wrensch, D.L. & Ebbert, M.A. (Eds.), *Evolution and diversity of sex ratio: insects and mites*. Chapman & Hall, New York, pp. 235–345.  
[https://doi.org/10.1007/978-1-4684-1402-8\\_8](https://doi.org/10.1007/978-1-4684-1402-8_8)
- Knížek, M. (2011) Scolytinae. In: Löbl, I. & Smetana, A. (Eds.), *Catalogue of Palearctic Coleoptera. Vol. 7. Curculionoidea I*. Apollo Books, Stenstrup, pp. 204–251.
- Lynch, S.C., Twizeyimana, M., Mayorquin, J., Wang, D., Na, F., Kayim, M., Kasson, M., Thu, P.Q., Bateman, C., Rugman-Jones, P., Hulcr, J., Stouthamer, R., & Eskalen, A. (2016) Identification, pathogenicity, and abundance of *Paracremonium pembeum* sp. nov. and *Graphium euwallaceae* sp. nov.- two newly discovered mycangial associates of the polyphagous shot hole borer (*Euwallacea* sp.) in California. *Mycologia*, 108(2), 313–329.  
<https://doi.org/10.3852/15-063>
- Markalas, S. & Kalapanida, M. (1997) Flight patterns of some Scolytidae attracted to flight barrier traps baited with ethanol in an oak forest in Greece. *Anzeiger Schdlingskde. Pflanzenschutz, Umweltschutz*, 70, 55–57.  
<https://doi.org/10.1007/BF01996922>
- Palm, T. (1959) Die Holz- und Rinden-Käfer der süd- und mittelschwedischen Laubbäume. *Opuscula Entomologica*, Supplementum 16, 1–374.
- Rabaglia, R.J., Dole, S.A. & Cognato, A.I. (2006) Review of American Xyleborina (Coleoptera: Curculionidae: Scolytinae) occurring North of Mexico, with an illustrated key. *Annals of the Entomological Society of America*, 99, 1034–1056.  
[https://doi.org/10.1603/0013-8746\(2006\)99\[1034:ROAXCC\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2006)99[1034:ROAXCC]2.0.CO;2)
- Ratnasingham, S. & Hebert, P.D.N. (2013) A DNA-Based Registry for All Animal Species: The Barcode Index Number (BIN)

- System. *PLoS ONE*, 8 (8), e66213.  
<https://doi.org/10.1371/journal.pone.0066213>
- Schedl, K. (1964) Biologie des gehöckerten Eichenholzbohrers, *Xyleborus monographus* Fab. (Scolytidae, Coleoptera). *Zeitschrift für Angewandte Entomologie*, 53, 411–428.  
<https://doi.org/10.1111/j.1439-0418.1963.tb02902.x>
- Smith, S.M. & Hulcr, J. (2015) *Scolytus* and other economically important bark and ambrosia beetles. In: Vega, F.E. & Hofstetter, R.W. (Eds.), *Bark Beetles: Biology and Ecology of Native and Invasive Species*. Academic Press, London, pp. 496–532.  
<https://doi.org/10.1016/B978-0-12-417156-5.00012-5>
- Smith, S.M., Gomez, D.F., Beaver, R.A., Hulcr, J. & Cognato, A.I. (2019) Reassessment of the species in the *Euwallacea fornicatus* (Coleoptera: Curculionidae: Scolytinae) complex after the rediscovery of the “lost” type specimen. *Insects*, 10, 261.  
<https://doi.org/10.3390/insects10090261>
- Stouthamer, R., Rugman-Jones, P., Thu, P.Q., Eskalen, A., Thibault, T., Hulcr, J., Wang, L.J., Jordal, B.H. Chen, C.Y., Cooperband, M., Lin, C.S., Kamata, N., Lu, S.S., Masuya, H., Mendel, Z., Rabaglia, R.J., Sanguansub, S., Shih, H.H., Sittichaya, W. & Zong, S. (2017) Tracing the origin of a cryptic invader: Phylogeography of the *Euwallacea fornicatus* (Coleoptera: Curculionidae: Scolytinae) species complex. *Agricultural and Forest Entomology*, 19, 366–375.  
<https://doi.org/10.1111/afe.12215>
- Wood, S.L. & Bright, D.E. (1992) A Catalog of Scolytidae and Platypodidae (Coleoptera), Part 2: Taxonomic Index. *Great Basin Naturalist Memoirs*, 13, 1–1553.