

Canadian Translation of Fisheries and Aquatic Sciences

No. 5152

Feeding of the ringed seal (Pusa hispida Schr.)

G.A. Fedoseev

Original title: Pitanie Kol'chatoi Nerpy

In: Izv. TINRO 59: 216-223, 1965


Original language: Russian

Available from:

Canada Institute for Scientific and Technical Information  
National Research Council  
Ottawa, Ontario, Canada K1A 0S2

1985

13 typescript pages





MULTILINGUAL SERVICES DIVISION - DIVISION DES SERVICES MULTILINGUES

TRANSLATION BUREAU

BUREAU DES TRADUCTIONS

LIBRARY IDENTIFICATION - FICHE SIGNALÉTIQUE

CTFAS 5152

Translated from - Traduction de  
RUSSIAN  
Into - En  
ENGLISH

Author - Auteur  
G.A. Fedoseev

Title in English or French - Titre anglais ou français

FEEDING OF THE RINGED SEAL  
(*Pusa hispida* Schr.)

Title in foreign language (Transliterate foreign characters)  
Titre en langue étrangère (Transcrire en caractères romains)

PITANIE KOL'CHATOI NERPY

Reference in foreign language (Name of book or publication) in full, transliterate foreign characters,  
Référence en langue étrangère (Nom du livre ou publication), au complet, transcrire en caractères romains.

Izvestiya TINRO

Reference in English or French - Référence en anglais ou français

TINRO, Proceedings

Publisher - Éditeur	DATE OF PUBLICATION DATE DE PUBLICATION			Page Numbers in original Numéros des pages dans l'original
	Year Année	Volume	Issue No. Numéro	216 - 223
Place of Publication Lieu de publication				Number of typed pages Nombre de pages dactylographiées
Magadan, USSR	1965	59		13

Requesting Department / Ministère-Client: DFO

Translation Bureau No. / Notre dossier n°: 1655879

Branch or Division / Direction ou Division: SIPB

Translation (Initials) / Traducteur (Initiales): GAD

Person requesting / Demandé par: T.G. Smith, Ste Anne de Bellevue

Your Number / Votre dossier n°: \_\_\_\_\_

Date of Request / Date de la demande: April 16th, 1985

MAY 3 1985

UNEDITED TRANSLATION  
For information only  
TRADUCTION NON REVISEE  
Information seulement



MULTILINGUAL SERVICES DIVISION – DIVISION DES SERVICES MULTILINGUES

TRANSLATION BUREAU

BUREAU DES TRADUCTIONS

Client's No.—N <sup>o</sup> du client	Department — Ministère DFO	Division/Branch — Division/Direction SIPB	City — Ville Ottawa
Bureau No.—N <sup>o</sup> du bureau 1655879	Language — Langue Russian	Translator (Initials) — Traducteur (Initiales) GAD	MAY 3 1985

Izvestiya TINRO, Vol. 59, Magadan, USSR, 1965, pp 216-223.

FEEDING OF THE RINGED SEAL  
(Pusa hispida Schr.)

By G.A. Fedoseev

In the literature, the feeding of the ringed seal is discussed /216\* only for the Canadian sector of the Arctic (McLaren, 1958), the Barents and Kara seas (K.K. Chapskii, 1940; G.F. Kurcheva, 1948; M.P. Vinogradov, 1949). For the rest - the Sea of Okhotsk, the Bering Sea, the East Siberian and Laptev seas, information on feeding, as well as other biological data, is either nonexistent or extremely inadequate.

This work discusses the feeding of the ringed seal of the Sea of (Pusa hispida ochotensis Pall.) and makes an attempt to compare it with the feeding of the other subspecies.

Comparison of the feeding of the ringed seal in different regions provides some explanation of the unique seasonal distribution of its subspecies and a more complete picture of the ecology of this seal as a species on the whole.

The work is based on original data collected in 1960-1963 in the Sea of Okhotsk and some published data.

In studying feeding, the contents of 159 stomachs of Okhotsk ringed seal (of 550 examined) were analysed. The bulk of the data was obtained from the regions of Tauisk Bay, and Babushkin Kekurnyi, Shel'ting and

\*The numbers in the right-hand margin indicate the corresponding pages in the original. - Transl.

and Ushka Gulfs. There was no marked difference in the species makeup of the contents of seal stomachs for different regions and years. For this reason, all the original material on seal feeding has been combined.

The stomach contents of ringed seal were weighed in a pan balance (5 kg). In some cases, individual components were weighed on dispensing balances. Components of feeding were determined both in the field and the office. Only the easily recognizable components were identified in the field. Identification was based on 1960 samples and was done by Ya. I. Zhitlo, a member of the Magadan branch of TINRO.

The literature on the feeding of the Okhotsk ringed seal in most cases gives only a list of the feeding components found without giving a quantitative analysis. P.G. Nikulin (1937) noted that the feeding of the ringed seal consists of small fish (navaga, smelt) and crustaceans. S.P. Naumov (1941), in discussing the feeding of the Okhotsk ringed seal, noted that in June 1929, he found mainly shrimp (Sclerocrangon) in seal stomachs. Gammarids (gammarus) and navaga remains were found in some stomachs. In July off the western shores of Sakhalin Island, the Okhotsk ringed seal, according to S.P. Naumov, pursues smelt and partly feeds on crustaceans, while in the second half of the summer and in the fall (until October), it consumes gobies, whitefish, navaga and crustaceans. S.Yu. Freiman (1936) also mentioned the feeding of ringed seal on fish, /217 noting that the approaches of this seal to the shores of Gizhiginsk Bay are connected with the presence of navaga. In June 1939, G.A. Pikharev (1946) found food in only 16 of 377 ringed seal stomachs examined. Pikharev explained the low number of stomachs with contents by the fact that digestion is rapid in seals. In describing food components, he noted crustaceans were found in 10 stomachs, among them: Thysanoessa raschii,

Themisto compresso f. bispinosa, Mesidothea, Gammarus schmidtii and Anonym nügax.

Only fish was found in 5 stomachs: Korean cod (Theragra chalcogramma), smelt (Hypomesus olidus Pall.) and herring (Clupea harengus Pall.). Food was mixed in one stomach.

Observations on the feeding of ringed seal (Table 2, Fig. 1) indicate a rather large diversity of feeding components and marked seasonal changes. Higher crustaceans make up the bulk of ringed seal feeding. Euphausiids of the genus Thysanoessa are found the most frequently, having been found in 137 of 159 food-containing stomachs (87%). Amphipoda and Decapoda are also found relatively frequently in relation to other food components, in 70 (44%) and 52 (32%) of stomachs. Other crustaceans are found considerably more rarely: Mysidacea have been found in 21 stomachs (14%), Isopoda in 7 (4%) and Copepoda in 3 (1%). Fishes were found in 44 stomachs (28%).

In the spring, the ringed seal feeds actively on Euphausiacea. In 128 individuals examined in March-April 1961 and 1963, food was found in 98 (77%) and consisted almost exclusively of euphausiids. The amount of these entomostracans in ringed seal stomachs often reached 700g or more (Table 2, Fig. 1).

During the moult (May-June), the feeding activity of the Okhotsk ringed seal is markedly diminished, but does not stop, as might appear at first glance from stomach analysis. The fact is that during the moult, the animals spend much time on ice floes, and since their food is rapidly digested, the stomach is empty in most animals taken. For example, in 397 animals which we examined in May-June, only 47 had food remains in the stomach and 39 in the intestine. In all, food remains were found in




Fig. 1. Weight ratio (in %) of food components of the Okhotsk ringed seal by seasons of the year:

1 - euphausiids, 2 - Mysidacea; 3 - Decapoda; 4 - fishes; 5 - Amphipoda; 6 - Copepoda; 7 - Isopoda.

a - February-April; b - May-June; c - November-December.

only 21% of animals examined. It is noteworthy that the number of food 218 components increases considerably during the moult (Fig. 1). This is due to the fact that in May and especially June, when mass moult occurs in the ringed seal, the animals spend a large part of the day on drift ice and must feed on any food which they encounter. The animals become very thin during the moult (May-June) (Fig. 2). With the end of moult (end of June - first ten days of July), the seal begins to feed actively. During the period of intensive feeding, the seal is distributed over the greatest part of its range. The seals are constantly afloat. With the end of the seal take in the summer and early fall, it becomes difficult to study its feeding, about which we have only indirect observations on distribution and behaviour.




Fig. 2. Condition of *Pusa hispida* Schr.  
 a - body weight, kg; b - age, years.  
 1 - March-April; 2 - May-June [average data].

Based on the reports of a number of investigators and our own observations, in July-September the ringed seal for the most part goes away from shores and accumulating mainly in regions of zooplankton development on the continental shelf, is annually found singly and in small groups (20-60 head) in almost the entire northern part of the Sea of Okhotsk. There are also reports in the literature about accumulations of ringed seal in the summer months. For example, S.Yu. Freiman (1935) reports that he found enormous accumulations of this seal afloat from Otlichitel'nyi Cape to Aldoma in August 1929 and that in his opinion the seal was pursuing a school of some sort of small fish, apparently capelin. P.G. Pikulin (1937) observed a large accumulation of Okhotsk ringed seal afloat in the region of Reineke and Men'shikov islands.

Thus the constant dwelling of the Okhotsk ringed seal in the summer in regions of mass development of zooplankton suggests that in this period,

just as in the spring, the ringed seal feeds on planktonic forms of crustaceans and fish, concentrating to forage in these regions. We are inclined to consider that in the summer in the northern part of the Sea of Okhotsk, the ringed seal feeds mainly on euphausiids, since, according to L.A. Ponomareva (1963), the accumulation of these entomostracans in surface layers reaches its maximum at this time. This hypothesis is confirmed by the following fact: at the end of the winter and in the spring, the accumulation of euphausiids is considerably smaller than in the summer, while the food of the ringed seal still consists almost /219 exclusively of these entomostracans, despite the accumulations of spawning navaga and herring at the same time. This fact indicates that euphausiids are evidently the preferred food of the ringed seal.

TABLE 1

Distribution of examined ringed seal stomachs in time and degree of fullness

<u>Time of year</u>	<u>Number of stomachs</u>			<u>Weight of stomach contents, in g</u>	
	<u>Total</u>	<u>including with food</u>	<u>%</u>	<u>Limits of variation</u>	<u>average</u>
February-April	128	98	77	60-950	370
May-June	397	47	12	40-560	210
November-December	25	14	56	150-600	350
<u>Total</u>	550	159	29	40-950	320

The period of intensive feeding is not limited to the summer forage, but includes the fall and part of the winter. In October and the beginning of November, the ringed seal concentrates en masse in bays, gulfs and coves. In our opinion, the fall approaches of the ringed seal to the coastal zone are a regular process of seasonal change in range, caused by the movement of seals into regions with different feeding conditions.



Judging by stomach dissections, in the fall-winter months, the main food components of the ringed seal are navaga, smelt, herring, sometimes gobies, sand lance and other fishes. Mainly nectobenthic forms of crustaceans are found: shrimps (Hipolitidae), (Pandalidae) and even mollusks (Gastropoda). The planktonic forms of crustaceans are found more rarely in stomachs at this time than at other periods. This is apparently due to a decrease in the abundance of these entomostracans due to their consumption by fishes, whales, birds and seals, and also to the migration of plankton from the surface layers to the deep as the result of the fall cooling of the surface waters.

The marked decrease in the condition of ringed seal in May-June (Fig. 2) is due to the moult. According to our measurements, the thickness of the fat layer hardly changes from November until the end of April and is 5-8 cm.

The food composition hardly changes with age. Fry and yearlings are an exception. Their food consists for the most part of small crustaceans - euphausiids and amphipods. Even in the fall, fishes and large crustaceans are found in the food of young ringed seals less often than in adult animals.

TABLE 2

List of food components of ringed seal.

<u>Food component</u>	<u>Canadian sector of Arctic</u>	<u>European sector of Arctic</u>	<u>Sea of Okhotsk</u>
<u>Copepoda</u>	+	-	+
Calanus hyperboreus Kroyer	+	-	-
Pareuchaeta norvegica Boeck	+	-	-
<u>Cirripedia</u>	+	-	-
Balanidae	+	-	-

TABLE 2 (Cont'd)

/220

<b>Разноногие раки — Amphipoda</b>	+	+	+
<i>Gammarus setosus</i> Demet	+	—	—
<i>Gammarus oceanicus</i> Segerst	+	—	—
<i>Gammarus wilkitzkii</i> Birula	+	—	—
<i>Gammarellus homari</i> Fabr	+	—	—
<i>Gammaracanthus loricatus</i> Sabine	+	+	—
<i>Anonyx nugax</i> Phipps	+	+	+
<i>Socarnes bidenticulatus</i> Bate	+	—	—
<i>Stegocephalus inflatus</i> Kröyer	+	—	—
<i>Ampelisca eschrichti</i> Kröyer	+	—	+
<i>Acanthostepheia</i> sp.	+	+	—
<i>Atylus carinatus</i> Fabr.	+	—	—
<i>Rhachotropis aculeata</i> Lepechin	+	—	—
<i>Pontogeneia inermis</i> Kröyer	+	—	—
<i>Amphithopsis longicaudata</i> Boeck	+	—	—
<i>Ischyrocerus anguipes</i> Kröyer	+	—	—
<i>Hyperia galba</i> Mantagu	+	—	—
<i>Themisto libellula</i> Mant.	+	+	+
<i>Pseudalibrotus biulaj</i> (gurjanova)	—	+	—
<i>Pseudalibrotus</i> sp.	—	+	—
<b>Десятиногие раки — Decapoda</b>	+	+	+
<i>Pasiphaea pacifica</i> Rathbun	—	—	+
<i>Pandalus</i> sp.	—	—	+
<i>Pandalus goniurus</i> Stimpson	+	—	—
<i>Pandalus montagu</i> Leach.	—	+	+
<i>Spirontocaris</i> sp.	—	+	+
<i>Spirontocaris murdochi</i> Rathbun	—	—	+
<i>Spirontocaris spinus</i> Sawyer	+	—	—
<i>Spirontocaris phippsi</i> Kröyer	+	—	—
<i>Eualus fabricii</i> Kröyer	+	—	—
<i>Eualus gaimardi</i> Milne-Edwards	+	—	—
<i>Lebbeus groenlandica</i> Fabr.	+	—	—
<i>Lebbeus polaris</i> Sabine	+	—	—
<b>Расщепленноногие раки — Mysidacea</b>	+	+	+
<i>Mysis oculata</i> Fabr	+	+	—
<i>Mysis mifta</i> Lilljeborg	+	—	—
<b>Черноглазки — Euphausiacea</b>	+	+	+
<i>Thysanessa inermis</i> Kröyer	+	—	—
<i>Thysanessa raschii</i> Sars.	+	+	+
<b>Равноногие раки — Isopoda</b>	—	+	+
<i>Mesidothea entomon</i>	—	+	+
<i>Idothea ochotensis</i> Brandt	—	—	+
<b>Брюхоногие моллюски — Gastropoda</b>	+	—	+
<i>Margarites helicina</i> Phipps	+	—	+
Turritellidae	+	—	+
<b>Двустворчатые моллюски — Bivalvia</b>	+	—	+
<i>Limacina helicina</i> Phipps	+	—	+
<i>Nucula tenuis</i> Montagu	+	—	+
<b>Головоногие моллюски — Cephalopoda</b>	+	+	—
<b>Многощетинковые черви — Polychaeta</b>	+	+	+
Maldanidae	+	—	+



Arctic cod is the main fish component of the feeding of ringed seal in most Arctic seas. It is one of the fishes which are most accessible to ringed seal. Another distinctive feature of the feeding of the ringed seal in Arctic seas is the fact that Mysis oculata makes up a large portion of the crustaceans in its diet. These entomostracans can form accumulations in the freshened waters of bays, gulfs and the pro-estuarine reaches of rivers. The mass accumulation of Mysis oculata in these regions is due to the presence of a peculiar "polar front" at the juncture of river and sea waters in the pro-estuarine reaches of large rivers.

In the northern part of the Sea of Okhotsk, there is no extensive freshening of sea water because of the absence of large rivers and Mysis oculata does not form mass accumulations. Euphausids make up the bulk of the crustaceans in ringed seal feeding. In the northern part of the Sea of Okhotsk, the ringed seal finds accumulations of euphausids in the surface layer of polynyas and leads. Euphausids are concentrated in March-April for mating, which is followed in four-six weeks by spawning, for which these entomostracans rise to the surface layers (L.A. Ponomareva, 1963).

The peculiarities of the distribution of food objects also determines the peculiar distribution of the ringed seal in a number of regions of the Arctic and the Far East. This peculiarity consists in the fact that the ringed seal is less confined to shores in the northern part of the Sea of Okhotsk than in the regions of the Canadian north, /222 the Barents and Kara Seas. This is especially evident on comparison of the distribution of seals during whelping and rearing of young stock.

In the Okhotsk seal, whelping sites lie for the most part far from shores beyond the selvage of shore ice, among mobile hummocky ice alternating with leads. With respect to feeding, these sites are the best for whelping since the first accumulations of euphausiids appear here in the spring. The existence of available foods is especially important for young stock in the transition to independent feeding.

The Arctic seas present a completely different picture. Both in Canada, off the shores of Baffin Island, and in the seas of the European north, the ringed seal whelps predominantly close to shores on motionless ice of bays and gulfs, making holes in the ice. Protection from wind and frost are undoubtedly important in the selection of whelping sites, but the food factor is no less important.

These findings on the feeding of the ringed seal show that small crustaceans sometimes not exceeding 10 mm in size, play a considerable role in its feeding. The adaptation of the ringed seal to the capture of such small food is of interest in this connection. McLaren (1958) considers that the ringed seal takes in small crustaceans with the water and filters them out. This seems quite plausible to us, as the structure of the dental system in the ringed seal is quite adapted for this. When the jaws close, the teeth adjoin with gaps through which the ringed seal apparently filters the water, leaving behind the entomostracans taken with it.

#### CONCLUSION

The ringed seal of the Sea of Okhotsk, like other subspecies of this seal, feeds on higher crustaceans and fish. Both play an important role in its feeding depending on the time of year. In the spring and summer, the ringed seal feeds mainly on crustaceans, and in the fall and partly in the winter, on fish.

In addition to similarities in feeding, different subspecies of ringed seal also show definite differences, perhaps due to specific conditions in biocoenoses of particular seas inhabited. In Arctic seas, Mysis oculata, Themisto <sup>bellina</sup> lebulala, and gammaracanthus loricatus are the main crustaceans in the diet, Thysanoessa raschii in the Sea of Okhotsk. The arctic cod is the main food fish in Arctic seas, navaga, smelt and herring in the Sea of Okhotsk.

The peculiarities in the distribution of food objects determines the peculiar distribution of the ringed seal in a number of regions, in that the ringed seal is less confined to shores in the Sea of Okhotsk than in a number of Arctic regions.

#### REFERENCES

1. Vinogradov, M.P. Marine mammals of the Arctic. Trudy Arkticheskogo in-ta, vol. 202, 1949.
2. Kurcheva, G.F. The feeding of the ringed seal in the southwestern part of the Kara Sea. Manuscript, 1948.
3. Naumov, S.P. Pinnipeds of the Sea of Okhotsk. Uchenye zapiski Moskovskogo gosud. pedagogich. in-ta, Vol. XXXIV, no. 2, 1941. /223
4. Nikulin, P.G. Observations on pinnipeds of the Sea of Okhotsk and Sea of Japan. Izv. TINRO, Vol. 10, Vladivostok, 1937.
5. Pikharev, G.A. Seals of the southwestern part of the Sea of Okhotsk. Izv. TINRO, Vol. 20, Vladivostok, 1941.
6. Pikharev, G.A. The feeding of the Okhotsk ringed seal. Izv. TINRO, Vol. XXII, Vladivostok, 1946.
7. Ponomareva, L.A. Euphausiids of the northern half of the Pacific Ocean, their distribution and ecology of mass species. Publishing House of the USSR Academy of Sciences, 1963.
8. Freiman. Materials on the commercial biology of seals of the Far East. Trudy VNIRO, vol. 111, Moscow, 1935.

9. Chapskii, K.K. The ringed seal of the western seas of the Soviet Arctic. Trudy nauchno-issledov. in-ta Glavnogo upravleniya Severnogo morskogo puti (Proceedings of the Scientific Research Institute of the Main Administration of the Northern Sea Route), Vol. 145, Leningrad, 1940.
10. McLaren, I.A. The biology of the ringed seal (*Phoca hispida*) in the eastern Canadian Arctic Bull. Fish. Res. Board of Canada, No. 118; 1958.

## ЛИТЕРАТУРА

1. / Виноградов М. П. Морские млекопитающие Арктики. Труды Арктического института, том 202, 1949.
2. Курчева Г.А.Ф. Питание кольчатой нерпы юго-западной части Карского моря. Рукопись, 1948.
3. Наумов С. П. Ластоногие Охотского моря. Ученые записки Московского госуд. педагогич. ин-та, т. XXXIV, вып. 2, 1941.
4. Никулин П. Г. Наблюдения над ластоногими Охотского моря и Японского. Изв. ТИНРО, т. 10, Владивосток, 1937.
5. Пихарев Г. А. Тюлени юго-западной части Охотского моря. Изв. ТИНРО, т. 20, Владивосток, 1941.
6. Пихарев Г. А. О питании акибы. Изв. ТИНРО, т. XXII, Владивосток, 1946.
7. Пономарева Л. А. Эвфаузииды северной половины Тихого океана, их распространение и экология массовых видов. Изд. АН СССР, 1963.
8. Фрейман. Материалы к промысловой биологии тюленей Дальнего Востока. Труды ВНИРО, т. 111, Москва, 1935.
9. Чапский К. К. Нерпа западных морей Советской Арктики. Труды научно-исследов. ин-та Главного управления Северного морского пути, т. 145, Ленинград, 1940.
10. McLaren I. A. The biology of the ringed seal (*Phoca hispida*) in the eastern Canadian Arctic. Bull. Fish. Res. B-d Canada. № 118; 1958.