

# Stellar Formation and Evolution

恆星形成與演化

Wen Ping Chen

<http://www.astro.ncu.edu.tw/~wchen/Courses/Stars/Default.htm>

## Table of Contents

- 1 ... Properties of Stars
- 2 ... Star Formation
- 3 ... Stellar Structure
- 4 ... Nuclear Reactions
- 5 ... Main-Sequence Evolution
- 6 ... Post-main Sequence
- 7 ... Compact Objects
- 8 ... Supernovae & Others

- ✓ What is a “star”?
- ✓ How hot is the surface of the Sun? How is this known? The Sun is gaseous, so how come it has a “surface”?
- ✓ How hot is the center of the Sun? How is this known?
- ✓ How long can the Sun remain as a shining body? How is this known?
- ✓ Describe the radial structure of the Sun. How is this known?

### Stellar Formation and Evolution --- Syllabus

**Instructor:** Professor Wen-Ping Chen

**Office:** 906

**Class Time:** Tuesday evening 5 to 8 scheduled (subject to change)

**Class venue:** Room 914

This course deals with the time variations of the structures of a star's interior and atmosphere. We will discuss the important physical processes governing the life of a star --- from its birth out of a dense, cold molecular cloud core, to shining with the star's own thermonuclear fuels, to rapid changes in structures when these fuels are no longer available, to the end of a star's life, with matter in extremely compact states.

What it may take for a star billions of years, will take us one semester to cover the following subjects:

- Observational Properties of Stars
- Molecular Clouds and the Interstellar Medium
- Cloud Collapse and Fragmentation
- Stars and Statistical Physics
- Protostars and Jets
- Circumstellar Disks and Planet Formation
- Evolution onto the Main Sequence
- Binaries and Star Clusters
- On the Main Sequence --- Nuclear Reactions
- Effects of Rotation
- Instabilities --- Thermally, Dynamically and Convectively
- Post-MS Evolution of Low-Mass Stars --- RG, AGB, HB, PNe
- Post-MS Evolution of Massive Stars --- SN and SNR
- Mass Loss, Stellar Pulsation and Cepheid Variables
- Compact Objects --- White Dwarfs, Neutron Stars, and Black holes

**Text:**

“*An Introduction to the Theory of Stellar Structure and Evolution*”, by Dina Prialnik, Cambridge, 2<sup>nd</sup> Ed. 2009

### References

All the references you have found useful for the course *Stellar Atmosphere and Structure* will be also of use in this course. The following are the ones I have been using or were published in recent years.

- ✓ *Physics of Stellar Evolution and Cosmology*, by H. Goldberg & Michael Scadron, 1982, Gordon and Breach
- ✓ *Stellar Structure and Evolution*, by R. Kippenhahn & W. Weigert, 1990, Springer-Verlag
- ✓ *Introduction to Stellar Astrophysics*, Vol 3 --- Stellar Structure and Evolution, by Erika Bohm-Vitense, 1992, Cambridge
- ✓ *Stellar Structure and Evolution*, by Huang, R.Q. 黄潤乾, Guoshin, 1990  
This book, originally in Chinese, has an English version, and has recently been revised. The Chinese version (恆星物理) has also been revised
- ✓ *The Physics of Stars*, by A.C. Phillips, 1994, John Wiley & Sons
- ✓ *Stellar Evolution*, by Amos Harpaz, A K Peters, 1994
- ✓ *The Stars --- Their Structure and Evolution*, R. J. Tayler, 1994, Cambridge
- ✓ *Theoretical Astrophysics, Vol II: Stars and Stellar Systems* by Padmanabhan, T., a hefty, mathematical 3 volume set; comprehensive coverage of basic astrophysical processes in vol. 1, stars in vol. 2, and galaxies and cosmology in vol. 3, 2001, Cambridge
- ✓ *Evolution of Stars and Stellar Populations*, by Maurizio Salaris and Santi, Cassisi, 2005, Wiley
- ✓ *The Formation of Stars*, by Steven W. Stahler & Francesco Palla, 2004, Wiley
- ✓ *From Dust to Stars*, by Norbert S. Schulz, 2005, Spinger
- ✓ *Stellar Physics, 2: Stellar Evolution and Stability*, by Bisnovaty-Kogan, 2<sup>nd</sup> Ed., 2010, Springer (translated from Russian)

For star formation, the book "*Molecular Clouds and Star Formation*", edited by Chi Yuan (袁旗) & Junhan You (尤峻漢) and published by World Scientific in 1993, should be a good reference. Unfortunately this book is currently out of print, but Prof Yuan kindly donated his editor copy.

In addition to written midterm (30% grade) and final (30%) exams, there will be homework assignments, plus in-class exercises or projects (35%).

For an extensive listing of books on "stars" ... <http://www.ericweisstein.com/encyclopedias/books/Stars.html>

## Course Goals

- To know the properties of various phases of the interstellar matter;
- To understand how stars form out of molecular clouds; under what conditions;
- To understand the physical properties of stars, and to know how these properties change with time as a star evolves;
- To understand the basic physics underlying complex stellar evolution models;
- To know how to interpret observational parameters of stars;
- To understand how stars of different masses evolve and what the end products of their evolution are.

**Stellar structure:** balance of forces

**Stellar evolution:** (con)sequence of thermonuclear reactions in different parts of a star

### Often used fundamental constants


#### Physical

$a$	radiation density constant	$7.55 \times 10^{-16} \text{ J m}^{-3} \text{ K}^{-4}$
$c$	velocity of light	$3.00 \times 10^8 \text{ m s}^{-1}$
$G$	gravitational constant	$6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$
$h$	Planck's constant	$6.62 \times 10^{-34} \text{ J s}$
$k$	Boltzmann's constant	$1.38 \times 10^{-23} \text{ J K}^{-1}$
$m_e$	mass of electron	$9.11 \times 10^{-31} \text{ kg}$
$m_H$	mass of hydrogen atom	$1.67 \times 10^{-27} \text{ kg}$
$N_A$	Avogadro's number	$6.02 \times 10^{23} \text{ mol}^{-1}$
$\sigma$	Stefan Boltzmann constant	$5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$ (= $ac/4$ )
$R$	gas constant ( $k/m_H$ )	$8.26 \times 10^3 \text{ J K}^{-1} \text{ kg}^{-1}$
$e$	charge of electron	$1.60 \times 10^{-19} \text{ C}$

Check out <http://pdg.lbl.gov/2006/reviews/astrorpp.pdf>

### Astronomical

$L_{\odot}$	Solar luminosity	$3.86 \times 10^{26} \text{ W}$
$M_{\odot}$	Solar mass	$1.99 \times 10^{30} \text{ kg}$
$T_{eff\odot}$	Solar effective temperature	5780 K
$T_{c\odot}$	Solar Central temperature	$1.6 \times 10^7 \text{ K}$ (theoretical)
$R_{\odot}$	Solar radius	$6.96 \times 10^8 \text{ m}$
$m_{\odot}$	apparent mag of Sun	-26.7 mag (V)
$M_{\odot}$	absolute mag of Sun	+4.8 mag (V)
$\theta$	apparent size of Sun	32'
$\langle \rho \rangle$	mean density of Sun	$1.4 \text{ g cm}^{-3}$
$(B-V)_{\odot}$	Color of the Sun	0.6 mag
<i>Parsec (unit of distance)</i>		$3.09 \times 10^{16} \text{ m}$



Galactic Ecology

## Properties of Stars

### Vocabulary

- **Luminosity** [ $\text{erg s}^{-1}$ ]  $L$  = bolometric luminosity = power
- **Spectral luminosity** [ $\text{erg s}^{-1} \mu\text{m}^{-1}$ ]  $L_\lambda$   $d\lambda = -(c/v^2) dv$
- **flux** [ $\text{erg s}^{-1} \text{cm}^{-2}$ ]  $f$
- **flux density** [ $\text{erg s}^{-1} \text{cm}^{-2} \mu\text{m}^{-1}$ ]  $f_\lambda$  or  $f_\nu$  1 Jansky (Jy) =  $10^{-23}$  [ $\text{erg s}^{-1} \text{cm}^{-2} \text{Hz}^{-1}$ ]  
 $f(\nu=0) = 3640$  Jy
- **Brightness/intensity** [ $\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$ ]  $B$   $S_\nu$  [ $\mu\text{Jy}$ ] =  $10^{(23.9-AB)/2.5}$
- **Specific intensity** [ $\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1} \text{Hz}^{-1}$ ]  $I_\nu$
- **Energy density** [ $\text{erg cm}^{-3}$ ]  $u = (4 \pi/c) J$   
 $J = \text{mean intensity} = (1/4\pi) \int I d\Omega$   $m_{AB} = -2.5 \log_{10} \left( \frac{f_\nu}{3631 \text{ Jy}} \right)$
- **Magnitude** ... apparent, absolute, bolometric, AB

## Our Sun ---- the best studied star



## Observable properties of stars



Basic parameters to compare between theories and observations

- ◆ Mass ( $M$ )
- ◆ Luminosity ( $L$ )
- ◆ Radius ( $R$ )
- ◆ Effective temperature ( $T_e$ )  $L = 4\pi R^2 \sigma T_e^4$
- ◆ Distance  $\rightarrow$  measured flux  $F = L / 4\pi d^2$

$M$ ,  $R$ ,  $L$  and  $T_e$  not independent

- $L$  and  $T_{\text{eff}}$  Hertzsprung-Russell (HR) diagram or color-magnitude diagram (CMD)
- $L$  and  $M$  mass-luminosity relation


<http://www.astrohandbook.com/links.html>



From wikipedia by Richard Powell based on *Hipparcos* data and Gliese catalog

[https://www.cosmos.esa.int/web/gaia/news\\_20150807](https://www.cosmos.esa.int/web/gaia/news_20150807)

For (nearby) star databases [http://www.projectrho.com/public\\_html/starmaps/catalogues.php](http://www.projectrho.com/public_html/starmaps/catalogues.php)



$$M_{\text{Jupiter}} \sim 0.001 M_{\odot}$$

**Stars:**

$$M > 0.08 M_{\odot}$$

**Brown Dwarfs:**

$$0.08 M_{\odot} > M > 13 M_{\text{J}}$$

**Planet-mass Objects:**


$$M < 13 M_{\text{J}}$$

## To measure the stellar distance

- ◆ Nearest stars  $d > 1\text{pc} \rightarrow p < 1''$
- ◆ For a star at  $d=100\text{ pc}$ ,  $p= 0.01''$
- ◆ Ground-based observations angular resolution  $\sim 1''$ ; *HST* has  $0.05''$
- ◆ *Hipparcos* measured the parallaxes of  $10^5$  bright stars with  $p \sim 0.001'' \rightarrow$  reliable distance determinations for stars up to  $d=100\text{ pc}$
- $\sim 100$  stars with good parallax distances

Preliminary Version of the Third Catalogue of  
Nearby Stars  
Gliese & Jahreiss (1991)

CDS catalog number: V/10A  
2964/3803 complete entries



*GAIA* will measure  $10^9$  stars!

## In most cases, the distance is estimated

- ◆ Stars with the same spectra are assumed to have identical set of physical parameters (spectroscopic parallax). For example, a G2V star should have the same absolute magnitude as the Sun.
- ◆ By comparison of the apparent brightness of an object with the known brightness of that particular kind of objects

$$m_\lambda - M_\lambda = 5 \log d - 5 + A_\lambda$$

$A_\lambda$  is usually unknown; it depends on the intervening dust grains that scatter and absorb the star light, and also depends on the distance to the object

- ◆ Main-sequence fitting; moving-cluster method; Cepheid variables
- ◆ Other methods for Galactic molecular clouds, galaxies, etc.






FIG. 1.—Normalized interstellar extinction curves from the far-IR through the UV. Several general features of the curves are noted. The solid and dotted curves are estimates for the case  $R \equiv A(V)/E(B-V) = 3.1$  derived in the Appendix of this paper and by Cardelli et al. (1989), respectively. The dashed curve shows the average Galactic UV extinction curve from Seaton (1979).

## To measure the stellar size

- ◆ Angular diameter of sun at 10 pc  
 $= 2R_{\odot}/10\text{pc} = 5 \times 10^{-9}$  radians  
 $= 10^{-3}$  arcsec
- ◆ Even the *HST* (0.05") barely capable of measuring directly the sizes of stars, except for the nearest supergiants
- ◆ Radii of  $\sim 600$  stars measured with techniques such as interferometry, (lunar) occultation or for eclipsing binaries









### To measure the stellar temperature

- ◆ What is  $T_{\text{eff}}$ ? What is the “surface” of a star?
- ◆ What is  $T$  anyway? Temperature is often defined by other physical quantities through an equation (“law”) (by radiation or by particles) blackbody, radiation, color, excitation, ionization, kinetic, electron, conductive ...
- ◆ Only in thermal equilibrium are all these temperatures the same.
- ◆ Photometry (spectral energy distribution) gives a **rough** estimate of  $T$ , e.g., fluxes/magnitudes measured at different wavelengths, such as the “standard” Johnson system *UBVRI*
- ◆ There are many photometric systems, using broad bands, intermediate bands, special bands, at optical or infrared wavelengths, etc.


Band	U	B	V	R	I
$\lambda/\text{nm}$	365	445	551	658	806
$\Delta\lambda/\text{nm}$	66	94	88	138	149





Running (slope) between  $B$  and  $V$  bands, i.e., the  $(B - V)$  **color (index)** → photospheric temperature


The larger the value of  $(B - V)$ , the redder (cooler) the star.



**Figure 1.8** Theoretical monochromatic flux emerging from an A type star with  $T_{\text{eff}} = 8000$  K. The first four Balmer absorption lines, as well as the Balmer jump, are identified in this figure. Thousands of other absorption atomic lines can also be seen. This theoretical flux was obtained with the Phoenix stellar atmosphere code (Hauschildt, P.H., Allard, F. and Baron, E., *The Astrophysical Journal*, 512, 377 (1999)) while using the elemental abundances found in the Sun. The flux at the surface of a blackbody with  $T = 8000$  K (dotted curve) is also shown.

LeBlanc



- ◆ Calibration for  $B - V = f(T_e)$
- ◆ The observed  $(B - V)$  must be corrected for interstellar extinction in order to derive the stellar intrinsic  $(B - V)_0$
- ◆ More accurate determination of T by spectra and stellar atmosphere models, e.g., the Kurucz's model




### Color Excess

$$E_{B-V} = (B - V)_{\text{obs}} - (B - V)_{\text{int}}$$

$$(B - V)_{\odot} = 0.656 \pm 0.005$$



Sloan Digital Sky Survey



Different temperature, elements (at different excitation and ionization levels) → different set of spectral lines





Figure 11-5  
 Discovering the Universe, Seventh Edition  
 © 2006 W.H. Freeman and Company


## Line ratios → Temperature




I --- neutral atoms; II --- ionized once; III --- ionized twice; ...

e.g., H I =  $H^0$  ... H II =  $H^+$  ... He III =  $He^{+2}$  ... Fe XXVI =  $Fe^{+25}$


**Hot stars** --- peaked at short wavelengths (UV); mainly He lines, some H lines



**Warm stars** --- peaked in the visible wavelengths; H lines prominent




**Cool stars** --- peaked at long wavelengths (IR); molecular lines/bands




## Brown dwarfs and Planetary Objects

L, T and Y types




Oppenheimer et al. 1998


## Brown dwarfs and Planetary Objects




<http://www.exoclimes.com/paper-outlines/exoplanets-and-brown-dwarfs-ii/>



One of the SDSS color-color diagrams



[http://spiff.rit.edu/classes/phys440/lectures/color/sdss\\_color\\_color\\_b.gif](http://spiff.rit.edu/classes/phys440/lectures/color/sdss_color_color_b.gif)



T dwarfs are blue in IR.

**Figure 16.15** Near-infrared color-color plot of M dwarfs (filled circles), L dwarfs (open circles), and T dwarfs (filled triangles). The objects are from a variety of regions. Note that the typical measurement errors for the L- and T-dwarfs are quite large, about 0.13 mag.

Stahler & Palla

## To measure the stellar luminosity

- ◆ **Absolute Magnitude**  $M$  defined as apparent magnitude of a star if it were placed at a distance of 10 pc

$$m_{\lambda} - M_{\lambda} = 5 \log(d_{\text{pc}}) - 5$$

But there is extinction ...

$$m_{\lambda} - M_{\lambda} = 5 \log(d_{\text{pc}}) - 5 + A_{\lambda}$$

- ◆ **Bolometric magnitude** – the absolute magnitude integrated over all wavelengths. We define the bolometric correction

- ◆ **Bolometric Correction**

$$BC = M_{\text{bol}} - M_v$$

$$M_{\text{bol}}^{\odot} = +4.74$$

is a function of the spectral type (*min at the F type, why?*) and luminosity of a star.

That is, we can apply BC (always negative, why?) to a star to estimate its luminosity (from the photosphere).

**Apparent Magnitude**  $m = -2.5 \log (\text{Flux}) + \text{ZeroPoint}$

- The Vega system: 0.0 mag (latest ~0.3 mag) at every Johnson band
- Gunn system: no Vega; use of F subdwarfs as standards (metal poor so smooth spectra), e.g., BD + 17 4708
- The AB system:  $AB_v = -2.5 \log_{10} f_v - 48.60$
- STMAG system: used for HST photometry  
 $STMAG_\lambda = -2.5 \log_{10} f_\lambda - 21.1$

Table 15.7. Calibration of MK spectral types.

<i>Sp</i>	<i>M(V)</i>	<i>B - V</i>	<i>U - B</i>	<i>V - R</i>	<i>R - I</i>	<i>T<sub>eff</sub></i>	<i>BC</i>
MAIN SEQUENCE, V							
O5	-5.7	-0.33	-1.19	-0.15	-0.32	42 000	-4.40
O9	-4.5	-0.31	-1.12	-0.15	-0.32	34 000	-3.33
B0	-4.0	-0.30	-1.08	-0.13	-0.29	30 000	-3.16
B2	-2.45	-0.24	-0.84	-0.10	-0.22	20 900	-2.35
B5	-1.2	-0.17	-0.58	-0.06	-0.16	15 200	-1.46
B8	-0.25	-0.11	-0.34	-0.02	-0.10	11 400	-0.80
A0	+0.65	-0.02	-0.02	0.02	-0.02	9 790	-0.30
A2	+1.3	+0.05	+0.05	0.08	0.01	9 000	-0.20
A5	+1.95	+0.15	+0.10	0.16	0.06	8 180	-0.15
F0	+2.7	+0.30	+0.03	0.30	0.17	7 300	-0.09
F2	+3.6	+0.35	0.00	0.35	0.20	7 000	-0.11
F5	+3.5	+0.44	-0.02	0.40	0.24	6 650	-0.14
F8	+4.0	+0.52	+0.02	0.47	0.29	6 250	-0.16
G0	+4.4	+0.58	+0.06	0.50	0.31	5 940	-0.18
G2	+4.7	+0.63	+0.12	0.53	0.33	5 790	-0.20
G5	+5.1	+0.68	+0.20	0.54	0.35	5 560	-0.21
G8	+5.5	+0.74	+0.30	0.58	0.38	5 310	-0.40
K0	+5.9	+0.81	+0.45	0.64	0.42	5 150	-0.31
K2	+6.4	+0.91	+0.64	0.74	0.48	4 830	-0.42
K5	+7.35	+1.15	+1.08	0.99	0.63	4 410	-0.72
M0	+8.8	+1.40	+1.22	1.28	0.91	3 840	-1.38
M2	+9.9	+1.49	+1.18	1.50	1.19	3 520	-1.89
M5	+12.3	+1.64	+1.24	1.80	1.67	3 170	-2.73
GIANTS, III							
G5	+0.9	+0.86	+0.56	0.69	0.48	5 050	-0.34
G8	+0.8	+0.94	+0.70	0.70	0.48	4 800	-0.42
K0	+0.7	+1.00	+0.84	0.77	0.53	4 660	-0.50
K2	+0.5	+1.16	+1.16	0.84	0.58	4 390	-0.61
K5	-0.2	+1.50	+1.81	1.20	0.90	4 050	-1.02
M0	-0.4	+1.56	+1.87	1.23	0.94	3 690	-1.25
M2	-0.6	+1.60	+1.89	1.34	1.10	3 540	-1.62
M5	-0.3	+1.63	+1.58	2.18	1.96	3 380	-2.48

Table 15.7. (Continued.)

<i>Sp</i>	<i>M(V)</i>	<i>B - V</i>	<i>U - B</i>	<i>V - R</i>	<i>R - I</i>	<i>T<sub>eff</sub></i>	<i>BC</i>
SUPERGIANTS, I							
O9	-6.5	-0.27	-1.13	-0.15	-0.32	32 000	-3.18
B2	-6.4	-0.17	-0.93	-0.05	-0.15	17 600	-1.58
B5	-6.2	-0.10	-0.72	0.02	-0.07	13 600	-0.95
B8	-6.2	-0.03	-0.55	0.02	0.00	11 100	-0.66
A0	-6.3	-0.01	-0.38	0.03	0.05	9 980	-0.41
A2	-6.5	+0.03	-0.25	0.07	0.07	9 380	-0.28
A5	-6.6	+0.09	-0.08	0.12	0.13	8 610	-0.13
F0	-6.6	+0.17	+0.15	0.21	0.20	7 460	-0.01
F2	-6.6	+0.23	+0.18	0.26	0.21	7 030	-0.00
F5	-6.6	+0.32	+0.27	0.35	0.23	6 370	-0.03
F8	-6.5	+0.56	+0.41	0.45	0.27	5 750	-0.09
G0	-6.4	+0.76	+0.52	0.51	0.33	5 370	-0.15
G2	-6.3	+0.87	+0.63	0.58	0.40	5 190	-0.21
G5	-6.2	+1.02	+0.83	0.67	0.44	4 930	-0.33
G8	-6.1	+1.14	+1.07	0.69	0.46	4 700	-0.42
K0	-6.0	+1.25	+1.17	0.76	0.48	4 550	-0.50
K2	-5.9	+1.36	+1.32	0.85	0.55	4 310	-0.61
K5	-5.8	+1.60	+1.80	1.20	0.90	3 990	-1.01
M0	-5.6	+1.67	+1.90	1.23	0.94	3 620	-1.29
M2	-5.6	+1.71	+1.95	1.34	1.10	3 370	-1.62
M5	-5.6	+1.80	+1.60	2.18	1.96	2 880	-3.47

Allen's Astrophysical  
Quantities (4<sup>th</sup> edition)

**Table 15.8. Calibration of MK spectral types.<sup>a</sup>**

$S_p$	$M/M_\odot$	$R/R_\odot$	$\log(g/g_\odot)$	$\log(\bar{\rho}/\bar{\rho}_\odot)$	$v_{\text{rot}}$ (km s <sup>-1</sup> )
MAIN SEQUENCE, V					
O3	120	15	-0.3	-1.5	
O5	60	12	-0.4	-1.5	
O6	37	10	-0.45	-1.45	
O8	23	8.5	-0.5	-1.4	200
B0	17.5	7.4	-0.5	-1.4	170
B3	7.6	4.8	-0.5	-1.15	190
B5	5.9	3.9	-0.4	-1.00	240
B8	3.8	3.0	-0.4	-0.85	220
A0	2.9	2.4	-0.3	-0.7	180
A5	2.0	1.7	-0.15	-0.4	170
F0	1.6	1.5	-0.1	-0.3	100
F5	1.4	1.3	-0.1	-0.2	30
G0	1.05	1.1	-0.05	-0.1	10
G5	0.92	0.92	+0.05	-0.1	< 10
K0	0.79	0.85	+0.05	+0.1	< 10
K5	0.67	0.72	+0.1	+0.25	< 10
M0	0.51	0.60	+0.15	+0.35	
M2	0.40	0.50	+0.2	+0.8	
M5	0.21	0.27	+0.5	+1.0	
M8	0.06	0.10	+0.5	+1.2	

**Table 15.8. (Continued.)**

$S_p$	$M/M_\odot$	$R/R_\odot$	$\log(g/g_\odot)$	$\log(\bar{\rho}/\bar{\rho}_\odot)$	$v_{\text{rot}}$ (km s <sup>-1</sup> )
GIANTS, III					
B0	20	15	-1.1	-2.2	120
B5	7	8	-0.95	-1.8	130
A0	4	5		-1.5	100
G0	1.0	6	-1.5	-2.4	30
G5	1.1	10	-1.9	-3.0	< 20
K0	1.1	15	-2.3	-3.5	< 20
K5	1.2	25	-2.7	-4.1	< 20
M0	1.2	40	-3.1	-4.7	
SUPERGIANTS, I					
O5	70	30:	-1.1	-2.6	
O6	40	25:	-1.2	-2.6	
O8	28	20	-1.2	-2.5	125
B0	25	30	-1.6	-3.0	102
B5	20	50	-2.0	-3.8	40
A0	16	60	-2.3	-4.1	40
A5	13	60	-2.4	-4.2	38
F0	12	80	-2.7	-4.6	30
F5	10	100	-3.0	-5.0	< 25
G0	10	120	-3.1	-5.2	< 25
G5	12	150	-3.3	-5.3	< 25
K0	13	200	-3.5	-5.8	< 25
K5	13	400	-4.1	-6.7	< 25
M0	13	500	-4.3	-7.0	
M2	19	800	-4.5	-7.4	

Note  
<sup>a</sup>A colon indicates an uncertain value.

*Allen's Astrophysical Quantities (4<sup>th</sup> edition)*

**Table 15.9. Zero-age main sequence.**

$(B - V)_0$	$(U - B)_0$	$M_v$	$(B - V)_0$	$(U - B)_0$	$M_v$
-0 <sup>m</sup> 33	-1 <sup>m</sup> 20	-5 <sup>m</sup> 2	+0.40	-0.01	+ 3.4
-0.305	-1.10	-3.6	+0.50	0.00	+ 4.1
-0.30	-1.08	-3.25	+0.60	+0.08	+ 4.7
-0.28	-1.00	-2.6	+0.70	+0.23	+ 5.2
-0.25	-0.90	-2.1	+0.80	+0.42	+ 5.8
-0.22	-0.80	-1.5	+0.90	+0.63	+ 6.3
-0.20	-0.69	-1.1	+1.00	+0.86	+ 6.7
-0.15	-0.50	-0.2	+1.10	+1.03	+ 7.1
-0.10	-0.30	+0.6	+1.20	+1.13	+ 7.5
-0.05	-0.10	+1.1	+1.30	+1.20	+ 8.0
0.00	+0.01	+1.5	+1.40	+1.22	+ 8.8
+0.05	+0.05	+1.7	+1.50	+1.17	+10.3
+0.10	+0.08	+1.9	+1.60	+1.20	+12.0
$(B - V)_0$	$(U - B)_0$	$M_v$	$(B - V)_0$	$(U - B)_0$	$M_v$
+0.15	+0.09	+2.1	+1.70	+1.32	+13.2
+0.20	+0.10	+2.4	+1.80	+1.43	+14.2
+0.25	+0.07	+2.55	+1.90	+1.53	+15.5
+0.30	+0.03	+2.8	+2.00	+1.64	+16.7
+0.35	0.00	+3.1			

*Allen's Astrophysical Quantities (4<sup>th</sup> edition)*

Main-Sequence Stars (Luminosity Class V)									
Sp. Type	$T_e$ (K)	$L/L_\odot$	$R/R_\odot$	$M/M_\odot$	$M_{\text{bol}}$	$BC$	$M_V$	$U - B$	$B - V$
O5	42000	499000	13.4	60	-9.51	-4.40	-5.1	-1.19	-0.33
O6	39500	324000	12.2	37	-9.04	-3.93	-5.1	-1.17	-0.33
O7	37500	216000	11.0	—	-8.60	-3.68	-4.9	-1.15	-0.32
O8	35800	147000	10.0	23	-8.18	-3.54	-4.6	-1.14	-0.32
B0	30000	32500	6.7	17.5	-6.54	-3.16	-3.4	-1.08	-0.30
B1	25400	9950	5.2	—	-5.26	-2.70	-2.6	-0.95	-0.26
B2	20900	2920	4.1	—	-3.92	-2.35	-1.6	-0.84	-0.24
B3	18800	1580	3.8	7.6	-3.26	-1.94	-1.3	-0.71	-0.20
B5	15200	480	3.2	5.9	-1.96	-1.46	-0.5	-0.58	-0.17
B6	13700	272	2.9	—	-1.35	-1.21	-0.1	-0.50	-0.15
B7	12500	160	2.7	—	-0.77	-1.02	+0.3	-0.43	-0.13
B8	11400	96.7	2.5	3.8	-0.22	-0.80	+0.6	-0.34	-0.11
B9	10500	60.7	2.3	—	+0.28	-0.51	+0.8	-0.20	-0.07
A0	9800	39.4	2.2	2.9	+0.75	-0.30	+1.1	-0.02	-0.02
A1	9400	30.3	2.1	—	+1.04	-0.23	+1.3	+0.02	+0.01
A2	9020	23.6	2.0	—	+1.31	-0.20	+1.5	+0.05	+0.05
A5	8190	12.3	1.8	2.0	+2.02	-0.15	+2.2	+0.10	+0.15
A8	7600	7.13	1.5	—	+2.61	-0.10	+2.7	+0.09	+0.25
F0	7300	5.21	1.4	1.6	+2.95	-0.09	+3.0	+0.03	+0.30
F2	7050	3.89	1.3	—	+3.27	-0.11	+3.4	+0.00	+0.35
F5	6650	2.56	1.2	1.4	+3.72	-0.14	+3.9	-0.02	+0.44
F8	6250	1.68	1.1	—	+4.18	-0.16	+4.3	+0.02	+0.52

Carroll &amp; Ostelie

Main-Sequence Stars (Luminosity Class V)									
Sp. Type	$T_e$ (K)	$L/L_\odot$	$R/R_\odot$	$M/M_\odot$	$M_{\text{bol}}$	$BC$	$M_V$	$U - B$	$B - V$
G0	5940	1.25	1.06	1.05	+4.50	-0.18	+4.7	+0.06	+0.58
G2	5790	1.07	1.03	—	+4.66	-0.20	+4.9	+0.12	+0.63
Sun <sup>a</sup>	5777	1.00	1.00	1.00	+4.74	-0.08	+4.82	+0.195	+0.650
G8	5310	0.656	0.96	—	+5.20	-0.40	+5.6	+0.30	+0.74
K0	5150	0.552	0.93	0.79	+5.39	-0.31	+5.7	+0.45	+0.81
K1	4990	0.461	0.91	—	+5.58	-0.37	+6.0	+0.54	+0.86
K3	4690	0.318	0.86	—	+5.98	-0.50	+6.5	+0.80	+0.96
K4	4540	0.263	0.83	—	+6.19	-0.55	+6.7	—	+1.05
K5	4410	0.216	0.80	0.67	+6.40	-0.72	+7.1	+0.98	+1.15
K7	4150	0.145	0.74	—	+6.84	-1.01	+7.8	+1.21	+1.33
M0	3840	0.077	0.63	0.51	+7.52	-1.38	+8.9	+1.22	+1.40
M1	3660	0.050	0.56	—	+7.99	-1.62	+9.6	+1.21	+1.46
M2	3520	0.032	0.48	0.40	+8.47	-1.89	+10.4	+1.18	+1.49
M3	3400	0.020	0.41	—	+8.97	-2.15	+11.1	+1.16	+1.51
M4	3290	0.013	0.35	—	+9.49	-2.38	+11.9	+1.15	+1.54
M5	3170	0.0076	0.29	0.21	+10.1	-2.73	+12.8	+1.24	+1.64
M6	3030	0.0044	0.24	—	+10.6	-3.21	+13.8	+1.32	+1.73
M7	2860	0.0025	0.20	—	+11.3	-3.46	+14.7	+1.40	+1.80

Carroll &amp; Ostelie

Giant Stars (Luminosity Class III)									
Sp. Type	$T_e$ (K)	$L/L_\odot$	$R/R_\odot$	$M/M_\odot$	$M_{\text{bol}}$	$BC$	$M_V$	$U - B$	$B - V$
O5	39400	741000	18.5	—	-9.94	-4.05	-5.9	-1.18	-0.32
O6	37800	519000	16.8	—	-9.55	-3.80	-5.7	-1.17	-0.32
O7	36500	375000	15.4	—	-9.20	-3.58	-5.6	-1.14	-0.32
O8	35000	277000	14.3	—	-8.87	-3.39	-5.5	-1.13	-0.31
B0	29200	84700	11.4	20	-7.58	-2.88	-4.7	-1.08	-0.29
B1	24500	32200	10.0	—	-6.53	-2.43	-4.1	-0.97	-0.26
B2	20200	11100	8.6	—	-5.38	-2.02	-3.4	-0.91	-0.24
B3	18300	6400	8.0	—	-4.78	-1.60	-3.2	-0.74	-0.20
B5	15100	2080	6.7	7	-3.56	-1.30	-2.3	-0.58	-0.17
B6	13800	1200	6.1	—	-2.96	-1.13	-1.8	-0.51	-0.15
B7	12700	710	5.5	—	-2.38	-0.97	-1.4	-0.44	-0.13
B8	11700	425	5.0	—	-1.83	-0.82	-1.0	-0.37	-0.11
B9	10900	263	4.5	—	-1.31	-0.71	-0.6	-0.20	-0.07
A0	10200	169	4.1	4	-0.83	-0.42	-0.4	-0.07	-0.03
A1	9820	129	3.9	—	-0.53	-0.29	-0.2	+0.07	+0.01
A2	9460	100	3.7	—	-0.26	-0.20	-0.1	+0.06	+0.05
A5	8550	52	3.3	—	+0.44	-0.14	+0.6	+0.11	+0.15
A8	7830	33	3.1	—	+0.95	-0.10	+1.0	+0.10	+0.25

Carroll &amp; Ostelie

F0	7400	27	3.2	—	+1.17	-0.11	+1.3	+0.08	+0.30
F2	7000	24	3.3	—	+1.31	-0.11	+1.4	+0.08	+0.35
F5	6410	22	3.8	—	+1.37	-0.14	+1.5	+0.09	+0.43
G0	5470	29	6.0	1.0	+1.10	-0.20	+1.3	+0.21	+0.65
G2	5300	31	6.7	—	+1.00	-0.27	+1.3	+0.39	+0.77
G8	4800	44	9.6	—	+0.63	-0.42	+1.0	+0.70	+0.94
K0	4660	50	10.9	1.1	+0.48	-0.50	+1.0	+0.84	+1.00
K1	4510	58	12.5	—	+0.32	-0.55	+0.9	+1.01	+1.07
K3	4260	79	16.4	—	-0.01	-0.76	+0.8	+1.39	+1.27
K4	4150	93	18.7	—	-0.18	-0.94	+0.8	—	+1.38
K5	4050	110	21.4	1.2	-0.36	-1.02	+0.7	+1.81	+1.50
K7	3870	154	27.6	—	-0.73	-1.17	+0.4	+1.83	+1.53
M0	3690	256	39.3	1.2	-1.28	-1.25	+0.0	+1.87	+1.56
M1	3600	355	48.6	—	-1.64	-1.44	-0.2	+1.88	+1.58
M2	3540	483	58.5	1.3	-1.97	-1.62	-0.4	+1.89	+1.60
M3	3480	643	69.7	—	-2.28	-1.87	-0.4	+1.88	+1.61
M4	3440	841	82.0	—	-2.57	-2.22	-0.4	+1.73	+1.62
M5	3380	1100	96.7	—	-2.86	-2.48	-0.4	+1.58	+1.63
M6	3330	1470	116	—	-3.18	-2.73	-0.4	+1.16	+1.52

Carroll &amp; Ostelie

Supergiant Stars (Luminosity Class Approximately Ia <sub>b</sub> )									
Sp. Type	$T_e$ (K)	$L/L_{\odot}$	$R/R_{\odot}$	$M/M_{\odot}$	$M_{\text{bol}}$	$BC$	$M_V$	$U - B$	$B - V$
O5	40900	1140000	21.2	70	-10.40	-3.87	-6.5	-1.17	-0.31
O6	38500	998000	22.4	40	-10.26	-3.74	-6.5	-1.16	-0.31
O7	36200	877000	23.8	—	-10.12	-3.48	-6.6	-1.14	-0.31
O8	34000	769000	25.3	28	-9.98	-3.35	-6.6	-1.13	-0.29
B0	26200	429000	31.7	25	-9.34	-2.49	-6.9	-1.06	-0.23
B1	21400	261000	37.3	—	-8.80	-1.87	-6.9	-1.00	-0.19
B2	17600	157000	42.8	—	-8.25	-1.58	-6.7	-0.94	-0.17
B3	16000	123000	45.8	—	-7.99	-1.26	-6.7	-0.83	-0.13
B5	13600	79100	51.1	20	-7.51	-0.95	-6.6	-0.72	-0.10
B6	12600	65200	53.8	—	-7.30	-0.88	-6.4	-0.69	-0.08
B7	11800	54800	56.4	—	-7.11	-0.78	-6.3	-0.64	-0.05
B8	11100	47200	58.9	—	-6.95	-0.66	-6.3	-0.56	-0.03
B9	10500	41600	61.8	—	-6.81	-0.52	-6.3	-0.50	-0.02
A0	9980	37500	64.9	16	-6.70	-0.41	-6.3	-0.38	-0.01
A1	9660	35400	67.3	—	-6.63	-0.32	-6.3	-0.29	+0.02
A2	9380	33700	69.7	—	-6.58	-0.28	-6.3	-0.25	+0.03
A5	8610	30500	78.6	13	-6.47	-0.13	-6.3	-0.07	+0.09
A8	7910	29100	91.1	—	-6.42	-0.03	-6.4	+0.11	+0.14

Carroll &amp; Ostelie

F0	7460	28800	102	12	-6.41	-0.01	-6.4	+0.15	+0.17
F2	7030	28700	114	—	-6.41	0.00	-6.4	+0.18	+0.23
F5	6370	29100	140	10	-6.42	-0.03	-6.4	+0.27	+0.32
F8	5750	29700	174	—	-6.44	-0.09	-6.4	+0.41	+0.56
G0	5370	30300	202	10	-6.47	-0.15	-6.3	+0.52	+0.76
G2	5190	30800	218	—	-6.48	-0.21	-6.3	+0.63	+0.87
G8	4700	32400	272	—	-6.54	-0.42	-6.1	+1.07	+1.15
K0	4550	33100	293	13	-6.56	-0.50	-6.1	+1.17	+1.24
K1	4430	34000	314	—	-6.59	-0.56	-6.0	+1.28	+1.30
K3	4190	36100	362	—	-6.66	-0.75	-5.9	+1.60	+1.46
K4	4090	37500	386	—	-6.70	-0.90	-5.8	—	+1.53
K5	3990	39200	415	13	-6.74	-1.01	-5.7	+1.80	+1.60
K7	3830	43200	473	—	-6.85	-1.20	-5.6	+1.84	+1.63
M0	3620	51900	579	13	-7.05	-1.29	-5.8	+1.90	+1.67
M1	3490	60300	672	—	-7.21	-1.38	-5.8	+1.90	+1.69
M2	3370	72100	791	19	-7.41	-1.62	-5.8	+1.95	+1.71
M3	3210	89500	967	—	-7.64	-2.13	-5.5	+1.95	+1.69
M4	3060	117000	1220	—	-7.93	-2.75	-5.2	+2.00	+1.76
M5	2880	165000	1640	24	-8.31	-3.47	-4.8	+1.60	+1.80
M6	2710	264000	2340	—	-8.82	-3.90	-4.9	—	—

Carroll &amp; Ostelie

TABLE II

Adopted calibration of MK spectral types in absolute magnitudes  $M_V$

Sp	ZAMS	V	IV	III	II	Ib	Iab	Ia
O5	-4.6	-5.6	-5.8	-6.0	-6.3	-6.6	-6.9	-7.2
O6	-4.0	-5.4	-5.7	-5.9	-6.3	-6.6	-6.9	-7.2
O7	-3.9	-5.2	-5.5	-5.8	-6.2	-6.5	-6.8	-7.2
O8	-3.7	-4.9	-5.2	-5.6	-6.1	-6.4	-6.7	-7.2
O9	-3.5	-4.5	-4.9	-5.3	-5.9	-6.3	-6.6	-7.2
B0	-3.1	-4.0	-4.4	-4.9	-5.6	-6.1	-6.5	-7.2
B1	-2.3	-3.3	-3.9	-4.5	-5.2	-5.9	-6.4	-7.2
B2	-1.6	-2.5	-3.1	-3.7	-4.5	-5.3	-6.1	-7.2
B3	-1.0	-1.7	-2.3	-3.0	-3.8	-4.6	-5.4	-7.2
B5	-0.1	-0.8	-1.2	-1.7	-2.4	-3.1	-3.8	-7.2
B6	0.3	-0.5	-0.9	-1.3	-1.9	-2.6	-3.3	-7.2
B7	0.6	-0.2	-0.6	-1.0	-1.6	-2.3	-3.0	-7.2
B8	1.0	0.1	-0.3	-0.7	-1.3	-2.0	-2.7	-7.2
B9	1.4	0.5	0.1	-0.4	-1.0	-1.7	-2.4	-7.2
A0	1.6	0.8	0.4	-0.1	-0.7	-1.4	-2.1	-7.2
A1	1.7	1.1	0.7	0.2	-0.2	-0.9	-1.6	-7.2
A2	1.8	1.3	0.9	0.4	-0.1	-0.8	-1.5	-7.3
A3	1.9	1.5	1.0	0.5	0.0	-0.7	-1.4	-7.3
A5	2.3	1.9	1.4	0.8	-0.2	-0.9	-1.6	-7.5
A7	2.6	2.3	1.7	1.1	-0.2	-0.9	-1.6	-7.7
F0	3.0	2.8	2.2	1.5	-0.7	-1.4	-2.1	-7.9
F2	3.2	3.1	2.4	1.8	-1.0	-1.7	-2.4	-8.0
F5	3.7	3.6	2.6	2.0	-1.6	-2.3	-3.0	-8.0
F8	4.2	4.1	2.8	2.2	-2.2	-2.9	-3.6	-8.1
G0	4.5	4.4	2.9	2.3	-2.8	-3.5	-4.2	-8.2
G2	4.7	3.0	1.1	-0.4	-1.1	-1.8	-2.5	-8.2
G5	5.1	3.1	1.0	-0.4	-1.1	-1.8	-2.5	-8.2
G8	5.6	3.2	0.9	-0.5	-1.2	-1.9	-2.6	-8.1
K0	6.0	3.2	0.8	-0.5	-1.2	-1.9	-2.6	-7.9
K1	6.2	3.2	0.8	-0.5	-1.2	-1.9	-2.6	-7.7
K2	6.5	0.7	-0.5	-1.2	-1.9	-2.6	-3.3	-7.6
K3	6.7	0.6	-0.5	-1.2	-1.9	-2.6	-3.3	-7.5
K4	7.0	0.5	-0.6	-1.3	-2.0	-2.7	-3.4	-7.4
K5	7.3	0.3	-0.6	-1.3	-2.0	-2.7	-3.4	-7.2
K7	8.1	0.0	-0.7	-1.4	-2.1	-2.8	-3.5	-7.0
M0	8.9	-0.6	-0.8	-1.2	-1.6	-2.0	-2.4	-6.9
M1	9.4	-0.8	-0.9	-1.3	-1.7	-2.1	-2.5	-6.8
M2	10.0	-0.9	-1.0	-1.4	-1.8	-2.2	-2.6	-6.7
M3	10.5	-1.0	-1.1	-1.5	-1.9	-2.3	-2.7	-6.7
M4	11.5	-0.6	-1.1	-1.4	-1.8	-2.2	-2.6	-6.7
M5	13.5	-0.1	-1.1	-1.4	-1.8	-2.2	-2.6	-6.7

TABLE III

Adopted temperatures and bolometric corrections for MK spectral types

Sp	log $T_{eff}$		Bol. Correction			
	V	III	I-II	V	III	I-II
O5	4.626		4.618		-4.15	-3.80
O6	4.593		4.585		-3.90	-3.55
O7	4.568		4.556		-3.65	-3.30
O8	4.550		4.535		-3.40	-3.15
O9	4.525		4.512		-3.15	-2.95
B0	4.498		4.431		-2.95	-2.50
B1	4.423		4.371		-2.60	-2.15
B2	4.362		4.307		-2.20	-1.75
B3	4.286		4.243		-1.85	-1.40
B5	4.188		4.137		-1.30	-0.90
B6	4.152		4.100		-1.05	-0.75
B7	4.107		4.068		-0.80	-0.60
B8	4.061		4.041		-0.55	-0.45
B9	4.017		4.013		-0.35	-0.35
A0	3.982		3.991		-0.25	-0.25
A1	3.973		3.978		-0.16	-0.16
A2	3.961		3.964		-0.10	-0.10
A3	3.949		3.949		-0.03	-0.03
A5	3.924		3.919		0.02	0.05
A7	3.903		3.897		0.02	0.09
F0	3.863		3.869		0.02	0.13
F2	3.845		3.851		0.01	0.11
F5	3.813		3.813		-0.02	0.08
F8	3.789	3.782	3.778		-0.03	0.03
G0	3.774	3.763	3.756		-0.05	0.00
G2	3.763	3.740	3.732		-0.07	-0.05
G5	3.740	3.712	3.699	-0.09		-0.22
G8	3.720	3.695	3.663	-0.13		-0.28
K0	3.703	3.681	3.643	-0.19		-0.37
K1	3.695	3.663	3.633			-0.43
K2	3.686	3.648	3.623	-0.30		-0.49
K3	3.672	3.628	3.613			-0.57
K4	3.663	3.613	3.613			-0.86
K5	3.643	3.602	3.585	-0.62		-1.15
K7	3.602			-0.89		-1.25
M0	3.591	3.591	3.568	-1.17		-1.40
M1	3.574	3.580	3.556	-1.45		-1.60
M2	3.550	3.574	3.544	-1.71		-1.95
M3	3.531	3.562	3.518	-1.92		-2.24
M4	3.512	3.550	3.491	-2.24		-2.6
M5	3.491	3.531	3.470	-2.55		-3.1
M6		3.512		-4.4		-4.0

Straižys & Kuriliene (1981)

TABLE IV

Bolometric absolute magnitudes  $M_{bol}$  for MK spectral types

Sp	ZAMS	V	IV	III	II	Ib	Iab	Ia
O5	-8.7	-9.8	-10.0	-10.2	-10.3	-10.4	-10.7	-11.0
O6	-8.0	-9.3	-9.6	-9.8	-9.9	-10.2	-10.4	-10.8
O7	-7.5	-8.8	-9.1	-9.3	-9.5	-9.8	-10.1	-10.5
O8	-7.2	-8.3	-8.6	-8.9	-9.2	-9.6	-9.8	-10.4
O9	-6.7	-7.6	-8.1	-8.4	-8.9	-9.3	-9.6	-10.2
B0	-6.2	-7.0	-7.4	-7.9	-8.1	-8.6	-9.0	-9.7
B1	-4.9	-5.8	-6.3	-6.8	-7.4	-8.0	-8.6	-9.4
B2	-4.0	-4.7	-5.3	-5.9	-6.8	-7.6	-8.6	-9.0
B3	-2.8	-3.6	-4.1	-4.7	-5.6	-6.6	-7.8	-8.6
B5	-1.4	-2.1	-2.5	-3.0	-3.8	-4.8	-6.1	-7.1
B6	-0.9	-1.6	-2.0	-2.4	-3.2	-4.2	-5.6	-6.6
B7	-0.2	-1.0	-1.4	-1.8	-2.6	-3.6	-5.0	-6.0
B8	0.4	-0.4	-0.8	-1.2	-2.0	-3.0	-4.4	-5.4
B9	1.0	0.1	-0.2	-0.8	-1.6	-2.6	-3.8	-4.8
A0	1.4	0.7	0.2	-0.3	-1.1	-2.1	-3.3	-4.3
A1	1.6	0.9	0.5	-0.1	-0.9	-1.9	-2.9	-3.9
A2	1.7	1.2	0.7	0.1	-0.7	-1.7	-2.7	-3.7
A3	1.9	1.5	1.0	0.4	-0.4	-1.4	-2.4	-3.4
A5	2.3	1.9	1.4	0.8	-0.2	-1.2	-2.2	-3.2
A7	2.6	2.3	1.8	1.1	-0.7	-1.7	-2.7	-3.7
F0	3.0	2.9	2.2	1.6	-1.4	-2.4	-3.4	-4.4
F2	3.2	3.1	2.4	1.8	-1.8	-2.8	-3.8	-4.8
F5	3.7	3.6	2.6	2.0	-2.5	-3.5	-4.5	-5.5
F8	4.2	4.1	2.8	2.2	-3.2	-4.2	-5.2	-6.2
G0	4.4	4.4	2.9	2.3	-3.8	-4.8	-5.8	-6.8
G2	4.6	4.6	2.9	1.0	-2.4	-3.4	-4.4	-5.4
G5	5.1	3.0	0.8	-0.5	-1.5	-2.5	-3.5	-4.5
G8	5.5	3.1	0.6	-0.7	-1.7	-2.7	-3.7	-4.7
K0	5.8	3.0	0.5	-0.8	-1.8	-2.8	-3.8	-4.8
K1	5.9	3.0	0.4	-0.9	-1.9	-2.9	-3.9	-4.9
K2	6.0	0.2	-0.3	-1.0	-2.0	-3.0	-4.0	-5.0
K3	6.2	-0.1	-0.4	-1.1	-2.1	-3.1	-4.1	-5.1
K4	6.4	-0.4	-0.7	-1.4	-2.4	-3.4	-4.4	-5.4
K5	6.7	-0.9	-1.2	-1.9	-2.9	-3.9	-4.9	-5.9
K7	7.3	-1.4	-1.7	-2.4	-3.4	-4.4	-5.4	-6.4
M0	7.5	-1.8	-2.1	-2.8	-3.8	-4.8	-5.8	-6.8
M1	7.9	-2.4	-2.7	-3.4	-4.4	-5.4	-6.4	-7.4
M2	8.3	-2.6	-2.9	-3.6	-4.6	-5.6	-6.6	-7.6
M3	8.8	-2.9	-3.2	-3.9	-4.9	-5.9	-6.9	-7.9
M4	9.3	-3.1	-3.4	-4.1	-5.1	-6.1	-7.1	-8.1
M5	11.0	-3.2	-3.5	-4.2	-5.2	-6.2	-7.2	-8.2
M6		-3.6	-3.9	-4.6	-5.6	-6.6	-7.6	-8.6

TABLE VI

Stellar masses  $\log M/M_{\odot}$  for different MK spectral types derived from the evolutionary tracks

Sp	ZAMS	V	IV	III	II	Ib	Iab	Ia
O5	1.60	1.81	1.85	1.89	1.90	1.92	1.99	
O6	1.48	1.70	1.76	1.80	1.80	1.87	1.91	2.00
O7	1.40	1.59	1.65	1.68	1.71	1.76	1.83	1.92
O8	1.34	1.48	1.54	1.60	1.65	1.72	1.76	1.90
O9	1.28	1.38	1.45	1.49	1.58	1.66	1.72	1.83
B0	1.20	1.30	1.34	1.40	1.40	1.48	1.56	1.70
B1	1.04	1.11	1.18	1.23	1.28	1.38	1.46	1.64
B2	0.92	0.99	1.04	1.08	1.18	1.30	1.38	1.54
B3	0.78	0.84	0.88	0.94	1.11	1.23	1.32	1.45
B5	0.62	0.68	0.72	0.75	1.00	1.18	1.26	1.40
B6	0.56	0.61	0.64	0.68	0.94	1.15	1.26	1.38
B7	0.49	0.53	0.57	0.60	0.91	1.11	1.23	1.36
B8	0.43	0.48	0.49	0.52	0.88	1.08	1.20	1.34
B9	0.36	0.41	0.45	0.49	0.85	1.04	1.20	1.32
A0	0.32	0.35	0.39	0.43	0.81	1.04	1.18	1.30
A1	0.31	0.34	0.36	0.41	0.78	1.00	1.18	1.30
A2	0.29	0.32	0.34	0.39	0.75	0.98	1.15	1.30
A3	0.27	0.30	0.32	0.36	0.75	0.97	1.11	1.30
A5	0.23	0.26	0.29	0.33	0.74	0.95	1.11	1.30
A7	0.20	0.22	0.26	0.30	0.73	0.94	1.15	1.32
F0	0.16	0.16	0.20	0.23	0.72	0.93	1.20	1.38
F2	0.13	0.13	0.16	0.20	0.72	0.93	1.20	1.40
F5	0.08	0.08	0.13	0.18	0.72	0.93	1.26	1.40
F8	0.04	0.04	0.11		0.72	0.93	1.28	1.41
G0	0.02	0.02	0.10		0.72	0.93	1.30	1.43
G2	0.00	0.00	0.10	0.33	0.72	0.93	1.30	1.45
G5	-0.02	0.08	0.39	0.73	0.94	1.32	1.46	
G8	-0.04	0.08	0.42	0.76	0.94	1.32	1.46	
K0	-0.07	0.11	0.46	0.78	0.96	1.30	1.45	
K1	-0.10	0.13	0.46	0.78	0.96	1.30	1.45	
K2	-0.10		0.45	0.79	0.98	1.28	1.43	
K3	-0.12		0.38	0.80	1.00	1.30	1.43	
K4	-0.15		0.36					
K5	-0.19		0.37	0.83	1.08			

TABLE VII

Calibration of MK spectral types in surface gravities (log g)

Sp	ZAMS	V	IV	III	II	Ib	Iab	Ia
O5	4.13	3.90	3.86	3.82	3.76	3.74	3.69	
O6	4.16	3.86	3.80	3.76	3.69	3.64	3.60	3.53
O7	4.18	3.85	3.80	3.74	3.64	3.57	3.52	3.45
O8	4.17	3.87	3.81	3.75	3.62	3.53	3.49	3.39
O9	4.21	3.95	3.82	3.74	3.58	3.50	3.44	3.31
B0	4.22	4.00	3.88	3.74	3.39	3.27	3.19	3.05
B1	4.28	4.00	3.86	3.71	3.31	3.17	3.01	2.87
B2	4.28	4.06	3.88	3.68	3.19	3.00	2.84	2.68
B3	4.31	4.06	3.89	3.71	3.12	2.79	2.68	2.49
B5	4.32	4.10	3.98	3.81	2.90	2.52	2.40	2.22
B6	4.32	4.09	3.96	3.84	2.77	2.42	2.29	2.13
B7	4.35	4.07	3.95	3.82	2.77	2.33	2.21	2.02
B8	4.34	4.07	3.92	3.79	2.79	2.27	2.11	1.97
B9	4.34	4.03	3.94	3.75	2.81	2.20	2.04	1.88
A0	4.32	4.07	3.91	3.75	2.85	2.23	2.01	1.81
A1	4.35	4.10	3.96	3.78	2.88	2.22	1.96	1.76
A2	4.32	4.16	3.98	3.78	2.87	2.23	1.92	1.71
A3	4.34	4.20	4.03	3.83	2.85	2.20	1.86	1.65
A5	4.36	4.22	4.06	3.86	2.81	2.14	1.74	1.53
A7	4.36	4.26	4.10	3.86	2.75	2.08	1.65	1.38
F0	4.32	4.28	4.05	3.83	2.67	2.00	1.51	1.25
F2	4.30	4.26	4.01	3.81	2.63	1.92	1.39	1.15
F5	4.32	4.28	3.93	3.74	2.48	1.81	1.22	1.00
F8	4.39	4.35	3.89	3.28	1.71	1.06	0.83	
G0	4.39	4.39	3.84	2.29	1.62	0.95	0.72	
G2	4.40	4.40	3.77	3.20	2.20	1.53	0.86	0.61
G5	4.49	4.49	3.71	3.07	2.04	1.45	0.71	0.45
G8	4.55	4.64	2.95	1.84	1.30	0.60	0.30	
K0	4.57	3.57	2.89	1.74	1.20	0.54	0.25	
K1	4.55	3.55	2.78	1.66	1.16	0.54	0.25	
K2	4.55	2.63	1.59	1.10	0.48	0.23		
K3	4.56	2.36	1.52	1.00	0.46	0.19		
K4	4.57	2.16						
K5	4.57	1.93	1.20	0.77	0.35	0.10		
K7	4.62							
M0	4.61	1.63	1.01	0.61	0.30	0.00		
M1	4.67	1.41	0.84	0.51	0.19	-0.07		
M2	4.69	1.31	0.70	0.39	0.09	-0.13		
M3	4.71	1.12	0.38	0.10	-0.16	-0.34		
M4	4.77	0.98						
M5	5.06	(0.76)						
M6		(0.52)						

TABLE VIII

Stellar radii log R/R<sub>⊙</sub> for different MK spectral types

Sp	ZAMS	V	IV	III	II	Ib	Iab	Ia
O5	0.95	1.17	1.21	1.25	1.28	1.30	1.36	
O6	0.87	1.13	1.19	1.23	1.27	1.33	1.37	1.43
O7	0.82	1.08	1.14	1.18	1.25	1.31	1.37	1.45
O8	0.80	1.02	1.08	1.14	1.23	1.31	1.35	1.47
O9	0.75	0.93	1.03	1.09	1.22	1.30	1.36	1.48
B0	0.70	0.86	0.94	1.04	1.20	1.32	1.40	1.54
B1	0.59	0.77	0.87	0.97	1.20	1.32	1.44	1.60
B2	0.54	0.68	0.80	0.92	1.21	1.37	1.49	1.65
B3	0.45	0.61	0.71	0.83	1.21	1.43	1.53	1.69
B5	0.36	0.50	0.58	0.68	1.27	1.55	1.65	1.81
B6	0.34	0.48	0.56	0.64	1.30	1.58	1.70	1.84
B7	0.29	0.45	0.53	0.61	1.28	1.60	1.72	1.88
B8	0.26	0.42	0.50	0.58	1.26	1.62	1.76	1.90
B9	0.23	0.41	0.47	0.59	1.23	1.63	1.79	1.93
A0	0.22	0.36	0.46	0.56	1.20	1.62	1.80	1.96
A1	0.19	0.33	0.41	0.53	1.16	1.60	1.82	1.98
A2	0.20	0.30	0.40	0.52	1.15	1.59	1.83	2.01
A3	0.18	0.26	0.36	0.48	1.16	1.60	1.84	2.04
A5	0.15	0.23	0.33	0.45	1.18	1.62	1.90	2.10
A7	0.13	0.19	0.29	0.43	1.21	1.65	1.97	2.19
F0	0.13	0.15	0.29	0.41	1.24	1.68	2.06	2.28
F2	0.13	0.15	0.29	0.41	1.26	1.72	2.12	2.34
F5	0.09	0.11	0.31	0.43	1.30	1.77	2.23	2.41
F8	0.04	0.06	0.33		1.38	1.82	2.32	2.50
G0	0.03	0.03	0.34		1.43	1.87	2.39	2.57
G2	0.01	0.01	0.38	0.78	1.48	1.92	2.44	2.64
G5		-0.04	0.41	0.88	1.56	1.96	2.52	2.72
G8		-0.08	0.43	0.95	1.67	2.03	2.57	2.79
K0		-0.11	0.48	1.00	1.73	2.09	2.59	2.81
K1		-0.11	0.50	1.05	1.77	2.11	2.61	2.81
K2		-0.11		1.12	1.81	2.15	2.61	2.81
K3		-0.12		1.22	1.85	2.21	2.63	2.83
K4		-0.15		1.31				
K5		-0.17		1.44	2.03	2.37	2.69	2.89
K7		-0.20						
M0		-0.22		1.64	2.12	2.48	2.72	2.92
M1		-0.27		1.78	2.21	2.55	2.78	2.99
M2		-0.30		1.83	2.27	2.61	2.85	3.03
M3		-0.36		1.92	2.44	2.76	2.98	3.16
M4		-0.42		1.98				
M5		-0.72		(2.04)				
M6				(2.16)				

Straižys & Kuriliene (1981)

Table 7.5. Filter wavelengths, bandwidths, and flux densities for Vega.<sup>a</sup>

Filter name	$\lambda_{\text{iso}}^b$ ( $\mu\text{m}$ )	$\Delta\lambda^c$ ( $\mu\text{m}$ )	$F_\lambda$ ( $\text{W m}^{-2} \mu\text{m}^{-1}$ )	$F_\nu$ (Jy)	$N_\phi$ ( $\text{photons s}^{-1} \text{m}^{-2} \mu\text{m}^{-1}$ )
V	0.5556 <sup>d</sup>	...	$3.44 \times 10^{-8}$	3 540	$9.60 \times 10^{10}$
J	1.215	0.26	$3.31 \times 10^{-9}$	1 630	$2.02 \times 10^{10}$
H	1.654	0.29	$1.15 \times 10^{-9}$	1 050	$9.56 \times 10^9$
K <sub>s</sub>	2.157	0.32	$4.30 \times 10^{-10}$	667	$4.66 \times 10^9$
K	2.179	0.41	$4.14 \times 10^{-10}$	655	$4.53 \times 10^9$
L	3.547	0.57	$6.59 \times 10^{-11}$	276	$1.17 \times 10^9$
L'	3.761	0.65	$5.26 \times 10^{-11}$	248	$9.94 \times 10^8$
M	4.769	0.45	$2.11 \times 10^{-11}$	160	$5.06 \times 10^8$
8.7	8.756	1.2	$1.96 \times 10^{-12}$	50.0	$8.62 \times 10^7$
N	10.472	5.19	$9.63 \times 10^{-13}$	35.2	$5.07 \times 10^7$
11.7	11.653	1.2	$6.31 \times 10^{-13}$	28.6	$3.69 \times 10^7$
Q	20.130	7.8	$7.18 \times 10^{-14}$	9.70	$7.26 \times 10^6$

1 Jansky =  $10^{-23} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1}$   
 $= 1.51 \times 10^7 \text{ photons s}^{-1} \text{ m}^{-2} (\Delta\lambda/\lambda)^{-1}$

Allen's Astrophysical Quantities (4<sup>th</sup> edition)

Band	$\lambda_0$	$d\lambda/\lambda$	$f_v (m=0)$	Reference
	$\mu\text{m}$		Jy	
U	0.36	0.15	1810	Bessel (1979)
B	0.44	0.22	4260	Bessel (1979)
V	0.55	0.16	3640	Bessel (1979)
R	0.64	0.23	3080	Bessel (1979)
I	0.79	0.19	2550	Bessel (1979)
J	1.26	0.16	1600	Campins, Reike, & Lebovsky (1985)
H	1.60	0.23	1080	Campins, Reike, & Lebovsky (1985)
K	2.22	0.23	670	Campins, Reike, & Lebovsky (1985)
g	0.52	0.14	3730	Schneider, Gunn, & Hoessel (1983)
r	0.67	0.14	4490	Schneider, Gunn, & Hoessel (1983)
i	0.79	0.16	4760	Schneider, Gunn, & Hoessel (1983)
z	0.91	0.13	4810	Schneider, Gunn, & Hoessel (1983)

<https://www.astro.umd.edu/~ssm/ASTR620/mags.html>

#### Notes

<sup>a</sup>Cohen et al. [1] recommend the use of Sirius rather than Vega as the photometric standard for  $\lambda > 20 \mu\text{m}$  because of the infrared excess of Vega at these wavelengths. The magnitude of Vega depends on the photometric system used, and it is either assumed to be 0.0 mag or assumed to be 0.02 or 0.03 mag for consistency with the visual magnitude.

<sup>b</sup>The infrared isophotal wavelengths and flux densities (except for  $K_s$ ) are taken from Table 1 of [1], and they are based on the UKIRT filter set and the atmospheric absorption at Mauna Kea. See Table 2 of [1] for the case of the atmospheric absorption at Kitt Peak. The isophotal wavelength is defined by  $F(\lambda_{\text{iso}}) = \int F(\lambda)S(\lambda) d\lambda / \int S(\lambda) d\lambda$ , where  $F(\lambda)$  is the flux density of Vega and  $S(\lambda)$  is the (detector quantum efficiency)  $\times$  (filter transmission)  $\times$  (optical efficiency)  $\times$  (atmospheric transmission) [2].  $\lambda_{\text{iso}}$  depends on the spectral shape of the source and a correction must be applied for broadband photometry of sources that deviate from the spectral shape of the standard star [3]. The flux density and  $\lambda_{\text{iso}}$  for  $K_s$  were calculated here. For another filter,  $K'$ , at  $2.11 \mu\text{m}$ , see [4].

<sup>c</sup>The filter full width at half maximum.

<sup>d</sup>The wavelength at V is a monochromatic wavelength; see [5].

#### References

1. Cohen, M. et al. 1992, *AJ*, **104**, 1650
2. Golay, M. 1974, *Introduction to Astronomical Photometry* (Reidel, Dordrecht), p. 40
3. Hanner, M.S., et al. 1984, *AJ*, **89**, 162
4. Wainscoat, R.J., & Cowie, L.L. 1992, *AJ*, **103**, 332
5. Hayes, D.S. 1985, in *Calibration of Fundamental Stellar Quantities*, edited by D.S. Hayes, et al., Proc. IAU Symp. No. 111 (Reidel, Dordrecht), p. 225

*Allen's Astrophysical Quantities* (4<sup>th</sup> edition)

## Exercise

Sirius, the brightest star in the night sky, has been measured  $m_B = -1.47$ ,  $m_V = -1.47$ . The star has an annual parallax of  $0.379''/\text{yr}$ .

1. What is its distance in parsec?
2. What is its absolute V-band magnitude?
3. From the absolute magnitude, what spectral type can be inferred for Sirius?
4. From the observed (B-V) color, what spectral type can be inferred?
5. What kinds of uncertainties/assumptions are associated with the above estimations?

## SIMBAD Astronomical Database

2015/4/21 sirius

---

sirius

other query Identifier Coordinate Criteria Reference Basic Script Output Help  
 modes : query query query query query submission options

Query : sirius

Available data : [Basic data](#) • [Identifiers](#) • [Plot & images](#) • [Bibliography](#) • [Measurements](#) • [External archives](#) • [Notes](#) • [Ann](#)

---

**Basic data :**

\* **alf CMa** -- Double or multiple star

Other object types: \* (\* ,BD,GC,HD,HIC,HIP,HR,SAO,UBV) , IR (AKARI ,IRAS,IRC,2MASS,RAFGL) , \*\* (\*\* ,MDS) , PM\* (LHS) , V\* (NSV) , UV (TD1)

ICRS coord. (ep=*J2000*) : **06 45 08.91728 -16 42 58.0171 ( Optical ) [ 11.70 10.90 90 ] A 2907ABa...474..653V**

FK5 coord. (ep=*J2000* eq=*2000*) : **06 45 08.917 -16 42 58.02 [ 11.70 10.90 90 ]**

FK4 coord. (ep=*B1950* eq=*1950*) : **06 42 56.72 -16 38 45.4 [ 67.39 63.09 0 ]**

Gal coord. (ep=*J2000*) : **227.2303 -08.8903 [ 11.70 10.90 90 ]**

Proper motions *mas/yr* : **-546.01 -1223.07 [ 1.33 1.24 0 ] A 2907ABa...474..653V**

Radial velocity / Redshift / cz : **V(km/s) -5.50 [0.4] / z(-) -0.000018 [0.000001] / cz -5.50 [0.40] (-) A 2907ABa...474..653V**

Parallax *mas* : **379.21 [1.58] A 2907ABa...474..653V**


Spectral type : **A1V+DA C 2813yCat...1.28235**

Fluxes (8) : **U -1.51 [-] C 2802yCat.2237...80  
 B -1.46 [-] C 2802yCat.2237...80  
 V -1.46 [-] C 2802yCat.2237...80  
 R -1.46 [-] C 2802yCat.2237...80  
 I -1.43 [-] C 2802yCat.2237...80  
 J -1.36 [-] C 2802yCat.2237...80  
 H -1.33 [-] C 2802yCat.2237...80  
 K -1.35 [-] C 2802yCat.2237...80**

<http://simbad.u-strasbg.fr/simbad/>

## To measure the stellar mass

- ◆ Stellar mass difficult to measure, direct measurements, except the Sun, only by binary systems  
*(but uncertain even for these, why?)*
- ◆ Then one gets the *mass-luminosity relation*  $L \propto M^\alpha$  where the slope  $\alpha=3$  to 5, depending on the mass range
- ◆ The main-sequence (MS) is a sequence of stellar mass under hydrostatic equilibrium
- ◆ Why are lower mass stars cooler on the surface and fainter in luminosity?




$$M_{\max} \sim 120 M_{\odot}$$

$$M_{\min} \sim 0.008 M_{\odot}$$

$$L_{\max} \sim 10^{+6} L_{\odot}$$

$$L_{\min} \sim 10^{-4} L_{\odot}$$





Luminosity versus mass for a selection of stars in binaries

Iben (2013)

## Luminosity class and surface gravity

$$\log g = \log GM/R^2$$

- Betelgeuse ... (M2 I)  $\log g \approx -0.6$  [cgs]
- Jupiter ...  $\log g = 3.4$
- Sun (G2 V) ...  $\log g = 4.44$
- G $\ell$ 229B ... (T6.5)  $\log g \approx 5$
- Sirius B... (WD)  $\log g \approx 8$



Composite Hertzsprung-Russell Diagram. Stars of different absolute luminosity,  $L$  - right axis, or bolometric absolute magnitude,  $M_{bol}$  - left axis, are plotted as a function of surface temperature,  $T_s$  - bottom axis, or spectral type - top axis. (Adapted from L. Goldberg and E.R. Dyer, Science in Space, eds. L.V. Berkner and H. Odishaw (1961).)

Lang "Data"


## Exercise

1. What is the spectral type of Alpha Scorpii?
2. What is its apparent magnitude? Expected absolute magnitude? Bolometric luminosity?
3. What is its distance estimated from its apparent magnitude? Measured directly by parallax? Why do these differ?
4. What is the expected diameter of the star in km, in  $R_{\odot}$  and in AU? What is then the expected angular diameter seen from Earth? Can it be resolved by the *HST*?

(Always show your work clearly, and cite the references.)

## To measure the stellar abundance

- ◆ By spectroscopy
- ◆ Stellar composition  $X, Y, Z$  = mass fraction of H, He and all other elements ("metals")  $Z$ : *metallicity*  $X+Y+Z=1$
- ◆ Solar abundance:  $X_{\odot} = 0.747$ ;  $Y_{\odot} = 0.236$ ;  $Z_{\odot} = 0.017$
- ◆ One often compares the iron abundance of a star to that of the sun. Iron is not the most abundant (only 0.001), but easy to measure in spectra. *Why?*



Data from: Katharina Lodders (2003) *Apl*, 591, 1220


$$[\text{Fe}/\text{H}] = \log_{10} \left( \frac{N_{\text{Fe}}}{N_{\text{H}}} \right)_{\text{star}} - \log_{10} \left( \frac{N_{\text{Fe}}}{N_{\text{H}}} \right)_{\odot}$$

$$\log \left( \frac{N_{\text{Fe}}}{N_{\text{H}}} \right)_{\odot} = -4.33$$



i.e., 1 iron atom for 20,000 H atoms

$$[M/H] \approx \log(Z/Z_{\odot})$$

Younger stars tend to be more metal-rich. Stars older than 10 Gyr almost all have  $[Fe/H] \lesssim -0.5$ ; stars younger than 5 Gyr have  $[Fe/H] \gtrsim -0.5$ .



Edvardsson et al. (1993)



H																	He				
Li	Be															B	C	N	O	F	Ne
Na	Mg															Al	Si	P	S	Cl	Ar
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr				
Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe				
Cs	Ba	Hf		Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn				
Fr	Ra	La		Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu				
		Ac		Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr				

B Big Bang   
 L Large stars   
 S Super-novae  
C Cosmic rays   
 S Small stars   
 M Man-made

Cosmic element factories --- the Big Bang, stellar nucleosynthesis, supernova explosions, and compact mergers


[http://en.wikipedia.org/wiki/Abundance\\_of\\_the\\_chemical\\_elements](http://en.wikipedia.org/wiki/Abundance_of_the_chemical_elements)

## To measure the stellar age

- ◆ Very tricky. Often one relies on measurements of  $M_V$ ,  $T_{\text{eff}}$ ,  $[\text{Fe}/\text{H}]$ , and then uses some kind of theoretically computed isochrones to interpolate the age (and mass)
- ◆ Crude diagnostics include
  - ✓ Lithium absorption line, e.g., 6707Å
  - ✓ Chromospheric activities, e.g., X-ray or Ca II emission
  - ✓ Evolving off the main sequence
- ◆ ... hence subject to large uncertainties

### References:

Edvardsson et al., 1993, A&A, 275, 101  
 Nordström et al., 2004, A&A, 418, 989





An MS star of the same spectral type


A PMS (young) star

**Figure 16.9** Lithium absorption in a pre-main-sequence star. Shown is a portion of the optical spectrum of BP Tau, a T Tauri star of spectral type K7, corresponding to an effective temperature of 4000 K. Also shown, for comparison, is a main-sequence star of the same spectral type, 61 Cyg B. Only in the first star do we see the Li I absorption line at 6708 Å. Both objects also have a strong line due to neutral calcium.

Stahler



## Pre-main sequence evolutionary models (tracks)




<http://stev.oapd.inaf.it/>

Bertelli et al, 2008, A&A, 484, 815


## Stellar populations

- ◆ Population I ..... Stars in the Galactic disk; like the Sun;  
*metal rich*
- ◆ Population II ..... Stars like those in the globular clusters;  
*metal poor*
- ◆ Population III .... Stars formed in the early universe; perhaps very hot and luminous; *metal free*



Distribution of Star Populations in Milky Way

<http://en.wikipedia.org/wiki/Metallicity>



Typical properties of Stellar Populations in the Milky Way

	Population I		Population II	
	very young	young	old	very old
Scale height [kpc]	60	100	500	2000
$\Sigma_w$ [ $\text{km s}^{-1}$ ]	8	10	25	75
$Z$	$> 0.02$	0.01	0.005	$< 0.002$
Age (rel. to the Universe)	$< 0.05$	0.25	0.75	1
Distribution	generally in aggregates		spherical	




Figure 2 (a) The radial abundance gradient in the galactic disk. Mean metallicities from DDO and *UBV* photometry from Janes (1979) (*triangles*) are plotted versus galactocentric distance relative to the Sun. Also shown are results from Washington photometry of classical Cepheids by Harris (1981) (*solid circles*) and high-dispersion abundance analysis of G to M supergiants by Luck & Bond (1980) (*open circles*); (b) The relation between age and metallicity for the open cluster samples of Janes (1979, Table 8). Ages are taken from McClure & Twarog (1978), Jennens & Helfer (1975), Cannon (1970), and sources quoted by Janes. (Reliable ages were not found for six clusters.) Open circles distinguish clusters with galactocentric radius larger than the solar value by more than 1 kpc. No correction has been made for any vertical abundance gradient.

Mould 1982 ARA&A, 20, 91

**Star clusters** are good laboratories to study stellar evolution, because member stars in a star cluster

- ◆ are (almost) of the same age;
- ◆ are (almost) at the same distance;
- ◆ evolve in the same Galactic environments;
- ◆ have the same chemical composition;
- ◆ are dynamical bound.

Two distinct classes:

- ✓ **globular clusters** (100+ in the MW)
- ✓ **open clusters** (a few  $10^3$  known in the MW)

How do these two classes differ in terms of shape, size, spatial distribution, number of member stars, and stellar population?



### Open Clusters

$10^2$  to  $10^3$  member stars;  $\sim 10$  pc across; loosely bound; open shape;  
 young population I;  
 located mainly in spiral arms;  
 $>1000$  open clusters known in the MW

### Globular Clusters

$10^5$  to  $10^6$  member stars; up to 100 pc across; tightly bound;  
 centrally concentrated;  
 spherical shape; old population II;  
 located in the Galactic halo;  
 200 globular clusters known in the MW



Stars in M80 are mostly old, metal poor members of Population II.








Figure 4.7. Hertzsprung-Russell color-magnitude diagrams for two very different samples of stars. (a) Diagram of 16 631 stars of various ages and various distances that are within about 330 LY of the sun, from the  $V$ -band and parallax distance measures by the Hipparcos satellite and ground-based color measurements. Horizontal-branch (HB) stars overlie the giant branch. (b) Color-magnitude diagram from ground-based studies of  $\sim 12\,200$  stars in globular cluster M3. The stars in the cluster are at a common distance of  $\sim 32\,000$  LY, and thus apparent magnitudes suffice for the ordinate of this H-R diagram. The stars are mostly of the common age of  $\sim 12$  Gyr. The data come in two distinct samples. A deep photographic sample yields most of the stars in the diagram both above and below the diagonal line. Short exposures with a charge-coupled device (CCD) camera provide fluxes for the rarer brighter stars plotted only above the dashed line representing  $B = 18.6$ . The scatter just above the diagonal line is mostly due to increased uncertainty at the fainter end of the CCD sample. The distance modulus of M3 is 14.93. Thus,  $M_V = 0$  in (a) corresponds to  $V = 14.93$  in (b). Note the absence of bright main-sequence stars in M3 and also the well-defined horizontal branch. [(a) ESA SP-1200, Hipparcos catalog (1997) in J. Kovalevsky, *ARAA* **36**, 121 (1998); (b) F. Ferraro *et al.*, *A&A* **320**, 757 (1997); photographic data from R. Buonomo *et al.*, *A&A* **290**, 69 (1994).]

Bradt "Astrophysics Processes"

OB associations  
in the solar  
neighborhood



De Zeeuw +1999



Range of possible GC ages (Chaboyer & Krauss 2003)

<http://www.astro.caltech.edu/~george/ay127/readings/KraussChaboyer2003.pdf>

