


THE BRITISH JOURNAL OF PSYCHIATRY



Published by authority of
The Royal College of Psychiatrists

NOTICE TO CONTRIBUTORS

The *British Journal of Psychiatry* is published monthly under the auspices of the Royal College of Psychiatrists. The *Journal* publishes original work in all fields of psychiatry. Communications on general editorial matters, including manuscripts for publication, printed matter and books for review, should be sent to *The Editor, The Royal College of Psychiatrists, 17 Belgrave Square, London SW1X 8PG.*

Contributions are accepted for publication on condition that their substance has not been published or submitted for publication elsewhere. The *Journal* does not hold itself responsible for statements made by contributors. Unless so stated, material in the *Journal* does not necessarily reflect the views of the Editor or of the Royal College of Psychiatrists.

Articles published become the property of the *Journal* and can be published elsewhere in full or part only with the Editor's written permission.

Manuscripts Three high quality copies (one of which should be the original typescript) should be submitted. Articles must be typed on one side of the paper only with double spacing and wide margins, and the pages must be numbered.

The **title** should be brief and to the point. A sub-title may be used to amplify the main title. The names of the authors should appear on the title page; their names, degrees, and affiliations should be given at the end of the paper.

A **summary** should be given at the *beginning* of the article. It should be brief and factual and not normally more than 120 words.

References should be listed alphabetically at the end of the paper, *the titles of journals being given in full.* For the reference list, authors should follow the style of the *Journal* and study the illustrations set out below. Titles of books and of journals will be printed in italics and should therefore be *underlined* in the typescript.

KENDELL, R. E. (1974) The stability of psychiatric diagnoses. *British Journal of Psychiatry*, **124**, 352–8.

RUTTER, M., TIZARD, J. & WHITMORE, K. (1970) *Education, Health and Behaviour*, p 14. London: Longman.

SCOTT, P. D. (1964) Definition, classification, prognosis and treatment. In *Pathology and Treatment of Sexual Deviation* (ed. I. Rosen). Oxford University Press.

DURKHEIM, E. (1897) *Le Suicide*. Paris. Translated 1952 as *Suicide: A Study in Sociology*, by J. A. Spaulding and C. Simpson, pp 191–206. London: Routledge and Kegan Paul.

In the text, references should be made by giving in brackets the name of the author and the year of publication, e.g. (Smith, 1971); or 'Smith (1971) showed that . . .'


Symbols and Abbreviations Follow 'Units, Symbols and Abbreviations, a Guide for Biological and Medical Editors and Authors' (1971, The Royal Society of Medicine, 1 Wimpole Street, London W1M 8AE). Terms or abbreviations which might not be understood by the average reader should be explained.

Tables and Figures Each table and figure must be on a separate sheet and its desired position in the text should be indicated (Table II here). Figures should be original drawings or glossy photos (not photocopies); the author's name and the title of the paper should be written in pencil on the back. Tables and figures should be self-explanatory, with adequate headings and footnotes. Units of measure must always be clearly indicated.

Editing Manuscripts accepted for publication are subject to copy-editing and to editorial changes required for conformity with *Journal* style.

Proofs A proof will be sent to the senior author of an article. Corrections other than printer's errors may be disallowed or charged to the authors. Reprints prepared at the same time as the *Journal* should be ordered for *all* authors from the printers when the proof is returned to the Editor.

General advice to authors *In the assessment of papers submitted to the Journal, great importance is attached to conciseness and clarity.* Authors should study 'General Notes on the Preparation of Scientific Papers', published by the Royal Society (6 Carlton House Terrace, London SW1Y 5AG), 1974 edition. They should check the accuracy of all references in their manuscript and ensure that dates and spellings correspond in the text and reference list.



in acute schizophrenia

Largactil* i.m. takes quiet control

Phenothiazine therapy is still among the most effective for acute, agitated schizophrenia. And the standard phenothiazine, especially for catatonic and paranoid forms, is still the original and oldest-established — 'Largactil'.

**rapidly suppresses psychomotor
excitement, agitation, hyperactivity**

'Largactil' chlorpromazine
tablets 10 mg PL 0012/5108
 25 mg PL 0012/5109
 50 mg PL 0012/5110
 100 mg PL 0012/5111

injection
solution 1% PL 0012/5082
 2.5% PL 0012/5308
 syrup PL 0012/5083
Forte suspension PL 0012/5001
 suppositories PL 0012/5084

*trade mark




M&B May & Baker



A member of
the Rhône-Poulenc
Group of Companies

further information is available from May & Baker Ltd Dagenham Essex RM10 7XS

MA6433



Ativan^{*}


(lorazepam, Wyeth)

stands out in the treatment of anxiety

Unlike most other benzodiazepines, ATIVAN has a direct one-step metabolism and is converted almost entirely to one inactive metabolite. ATIVAN thus ensures a more rapid and predictable anxiolytic response from your patients with fewer unwanted effects.

Ativan – rapid and predictable anxiolytic response

AT122

 ATIVAN is presented as yellow tablets containing 2.5 mg lorazepam. (A blue oblong tablet and an injectable form are also available.) Further information is available on request to the Company.
Wyeth Laboratories John Wyeth & Brother Ltd., Taplow, Maidenhead, Berks. *trade marks

ST. ANDREW'S HOSPITAL, NORTHAMPTON

PRIVATE PSYCHIATRIC CARE—A COMPREHENSIVE FACILITY

St. Andrew's aims to provide the best in psychiatric care in a comfortable and relaxed environment. Staffed by eight full-time Consultant Psychiatrists, three Clinical Psychologists and with a full complement of skilled nursing and rehabilitation staff, the hospital offers a broad range of psychiatric treatments. Care is provided on a short, medium and long term basis and the following specific units are available:

SHORT TERM ACUTE TREATMENT
 ALCOHOL TREATMENT
 BEHAVIOUR MODIFICATION
 DAY HOSPITAL

PSYCHOGERIATRIC
 BRAIN DAMAGE AND EPILEPSY INVESTIGATION
 PSYCHOTHERAPY
 MENTAL HANDICAP

Out-patient facilities are available at the hospital, at 148 and 152 Harley Street, London W.1., Oxford and Bedford.

St. Andrew's is an Independent Hospital and subscribers to the main Private Contributory Schemes may claim benefits within the terms of these schemes.

*Further details and brochures may be obtained from the Medical Director,
 St. Andrew's Hospital, Northampton.
 Telephone 0604 21311.*




**Years ahead
with clinical experience
and controlled research studies**

ModecateTM
(fluphenazine decanoate)

internationally accepted as the standard depot treatment for schizophrenia

Full prescribing information is available from:
Technical Department, E. R. Squibb & Sons Ltd., Regal House, Twickenham, TW1 3QT.





**Some antidepressants
enable your patients
to take life
more easily**

In depression the risk of suicide is increased.
Many antidepressant drugs increase this risk considerably.
Fluanxol is effective, non-sedating and quick-acting in the
treatment of depression. In addition clinical reports
show that in overdosage it lacks the acute toxicity of
other antidepressants – up to 3 months'

Fluanxol supply of tablets have
been taken in unsuccessful
suicide attempts.

controls the depression and reduces the risk of fatality.

Lundbeck Limited, Lundbeck House, Hastings Street, Luton LU1 5BE. Telephone Luton 411482 Telex 825325

Lundbeck

The Retreat, York

for Psychiatric Illnesses

Founded in 1792 by the Quaker, William Tuke, who established the tradition of concern for patients as individuals. The Retreat is a 250 bedded private registered nursing home surrounded by extensive grounds on the outskirts of the historic City of York. It is easily reached by rail and motorways.

Care and treatment is offered for most types of psychiatric illness on the short or long term in a sympathetic and friendly atmosphere. Patients suffering from neuroses, psychoses, alcoholism and dementia are treated in surroundings suitable for their individual needs.

The Nursing Home is a registered charity and is able to offer inclusive care in shared accommodation from £12.65 per day or in single rooms at slightly increased charge. The Nursing Home is recognised by the main private patient schemes.

For further details apply to The Medical Director, The Retreat, York YO1 5BN (Telephone 0904-54551).



Disipal has made her a little more responsive to her phenothiazine therapy.

The addition of Disipal to phenothiazine therapy enables optimum therapeutic response to be achieved without unacceptable side effects. Disipal also elevates the patient's mood, thus relieving the depression so often associated with major tranquillizer therapy.

Drug of choice

Following a three month double blind crossover trial, the authors concluded that, "orphenadrine is the drug of choice in the treatment of drug-induced extra-pyramidal reactions and depression."¹

Increased response

Furthermore, the authors postulate that "the introduction of orphenadrine in the treatment of a patient whose response to phenothiazines is not maintained, might well result in further benefit."¹

For patients on major tranquillizer therapy

Disipal

- * controls extra-pyramidal reactions
- * elevates patient mood.

1. Capstick N., J. Int. Med. Res., 1976, 4 (6), 435. Disipal, orphenadrine hydrochloride B.P., is a registered trade mark.

Full prescribing information on request from:—

Brocades (Great Britain) Ltd


Brocades House, Pyrford Road, West Byfleet, Weybridge, Surrey KT14 6RA
Telephone: Byfleet 45536/42291 Telex: 917301

Sparine allows
senile patients to grow old
without growing troublesome



Many patients become confused, agitated and difficult to manage as they grow older. They often become unco-operative and aggressive towards those who wish to care for them and place a considerable extra burden on nursing staff. SPARINE (promazine hydrochloride B.P.) reduces agitation without causing over-sedation and helps to make them more co-operative, alert and active.

**Sparine calms and controls
elderly, agitated patients**

Full prescribing information is available on request
Wyeth Laboratories, Taplow, Maidenhead, Berks. 
*trade marks Sp. 37.

Ticehurst House Wadhurst, Sussex TN5 7JA

First-class medical and nursing care are available at Ticehurst House for the treatment of both short stay and longer term psychiatric patients.

ALCOHOLISM

Comprehensive resources also exist to help those with alcohol-related disorders in the **Newington Unit** at Ticehurst under a new Director.

Further information is
available from the Matron.
Telephone: 0580 200391

**A Nestor
Nursing Home**



BOWDEN HOUSE CLINIC

Harrow-on-the-Hill, Middlesex

Tel: 01-864 0221

Only 20-25 Mins. from London
via Westway Extension

Founded in 1911 by Dr. H. Crichton-Miller

A non-profit making Charity outside the National
Health Service

A private clinic (all patients having single rooms) for the treatment of patients suffering from neuroses, early psychoses, psychosomatic disorders, drug addiction and alcoholism.


Treatment is supervised by experienced psychiatrists whose services are inclusive in the patients' fees.

A full physical examination and pathological investigations are made in the first week.

Facilities are also available to Consultants wishing to treat their own patients independently.

Apply for details: Administrative Secretary

Applications for admission to the Matron



★ Even when the classification is not crystal clear

Whether the patient is agitated or lethargic, depression is nearly always dramatic. The present can't be coped with – and the future is something to be feared.

For patients like this, Evadyne combines a profound anti-depressant effect with a rapid anxiolytic action.

Within three to four days, anxiety is relieved and the outlook starts to look more rosy. The patient discovers that she can communicate and co-operate once more. As early as the second week, the symptoms of depression are relieved and the way ahead seems clearer.

Evadyne (butriptyline hydrochloride) is available as Tablets of 25 mg. and 50 mg. in packs of 100.

Full prescribing information is available on request.

Evadyne

Provides a whole new outlook on the management of depression

Ayerst Laboratories Limited, Invincible Road, Farnborough, Hants GU14 7QH. Telephone: Farnborough 511981

Evadyne is a registered trade mark.

University of Bristol
Departments of Mental Health and
Extra-Mural Studies

PSYCHOTHERAPY WORKSHOP

14th-19th May, 1978

This workshop is intended for psycho-
logists, psychiatrists, social workers, and
general practitioners who have a few
years' experience of psychotherapy and
possibly (but not necessarily) some
training.

£85.00 resident membership.

Further particulars and application forms
from: The Assistant Director, Department
of Extra-Mural Studies, University of
Bristol, 32 Tyndall's Park Road, Bristol
BS8 1HR.

(Tel. Bristol 24161 ext. 649).



MISSOURI
INSTITUTE
of
PSYCHIATRY

Post Doctoral Fellowships

The Missouri Institute of Psychiatry in St. Louis announces a major expansion of its Psychiatric Post-Doctoral Fellowship Program. A university based, multidisciplinary, psychiatric, educational and research center, the Missouri Institute of Psychiatry is also affiliated with several State Department of Mental Health and community hospitals. Qualified faculty and all fellows receive academic appointments in the Department of Psychiatry, University of Missouri-Columbia, School of Medicine. The Post-Doctoral Program offers 2-3 years of post residency clinical experience and research oriented training in the areas of Psychobiology and Mental Health Information Systems. A fellowship in Administrative Psychiatry is also available. Clinical and basic science research training is under the guidance of internationally recognized experts in psychopharmacology, biochemistry, neurophysiology and computer applications, backed by outstanding library facilities. Salaries - \$27,000. For further information, interested physicians who have completed a Residency or equivalent in Psychiatry or Neurology should write:

John Barton, M.D.
Department of Psychiatry
University of Missouri-Columbia
School of Medicine
Missouri Institute of Psychiatry
5400 Arsenal Street
St. Louis, Missouri 63139

The University of Missouri is an equal opportunity employment institution.



WATERFORD HOSPITAL St. John's Newfoundland Canada

Applications are invited from suitably qualified persons for the post of Staff Psychiatrist (full time) at the above Psychiatric Hospital.

This hospital provides a service for the Province of Newfoundland and Labrador, and is the only one wholly concerned with Psychiatry. There are, however, a number of psychiatric units in general hospitals throughout the province. The Waterford is affiliated with Memorial University, St. John's and is involved in the residency training program of the Department of Psychiatry of the University.

There are 400 beds for acute and chronic cases, and a newly built section has recently become partly operational, which when fully occupied will increase the total in patient occupancy to 500. The hospital operates an extensive Community Care program and houses the Province's Forensic Unit.

Candidates must have the MRC Psych. Special experience in alcoholism, adolescent psychiatry or subnormality would be an advantage, but not essential.

Starting salary is from \$28,500 to \$34,500 depending on experience and whether fully or provisionally registrable in Newfoundland. There is a pension scheme and a number of fringe benefits. Short stay contracts will be considered of 1 to 2 years should the applicant so wish. Assistance in relocation is available.

Applications should include a curriculum vitae and the names of three referees and be addressed to:

THE MEDICAL DIRECTOR,
Waterford Hospital,
Waterford Bridge Road
St. John's, Newfoundland,
Canada
A1C 5T9.

PSYCHIATRIST

Physician with Psychiatric Speciality qualifications required for new 12-bed community mental health facility serving a population of 35,000 people with in-patient and out-patient programs in modern 116-bed district hospital with 24 active medical staff members.

6 sessions per week at \$140 per session at mental health facility, balance in active private practice downtown under B.C. Medical Plan.

Nelson is a pleasantly located community on Kootenay Lake in south-eastern British Columbia, Canada, well served with sport, recreational and community facilities.

Direct enquiries to:

R. H. Procter, Administrator,
Kootenay Lake District Hospital,
3 View Street,
Nelson, B. C., Canada
V1L 2V1

Epidemiological Approaches in Child Psychiatry

edited by P. J. Graham

December 1977, x+402pp., £10.00/\$19.50 0.12.294550.6

The papers contained in this volume—originally presented at a meeting held at the Ciba Foundation attended by workers from many different countries—provide an up-to-date account of work currently being undertaken in the epidemiology of child psychiatry. Behaviour and emotional problems are so prevalent that they form a major contribution to the total number of handicapping disorders in children. An understanding of how they occur, how common they are and what the best methods are for delivering services is an essential part of any comprehensive health care system. This book provides information not only for those working in the research field but also for those psychiatrists, psychologists, social workers and health planners who wish to inform themselves of the latest developments. The work includes an account of recent and otherwise unpublished findings in relation to urban/rural differences in rates of disorder, strategies for prevention, as well as ethological studies looking at the ways in which the mothers with different earlier experiences handle their children.

Academic Press

London New York San Francisco

A Subsidiary of Harcourt Brace Jovanovich, Publishers

24-28 Oval Road, London NW1, England

111 Fifth Avenue, New York, NY 10003, USA



4th EDITION OF THE READING LIST IN PSYCHIATRY

The 4th edition of the READING LIST is now available on application to the printers. Please complete the order form below.

To: HEADLEY BROTHERS LTD, ASHFORD, KENT TN24 8HH

Please post copies of the 4th Edition of the Reading List in Psychiatry (40p per copy including postage) for which I enclose a cheque/postal order payable to Headley Brothers Ltd.

Name

Address

.....

.....

Put depression safely behind them with **Bolvidon**



**effective,
safer
tetracyclic
antidepressant**


BOLVIDON


A product of original
Organon research

Bolvidon: Mianserin hydrochloride 10mg

Unlike tricyclic antidepressants

- * Free from anticholinergic effects
- * Remarkable lack of side effects
- * Wide safety margin
- * Lacks cardiotoxic effects

Full prescribing information available on request:



Organon Laboratories Ltd,
Crown House, Morden, Surrey SM4 5DZ
Telephone: 01-542 6611

THE BRITISH JOURNAL OF PSYCHIATRY

(THE JOURNAL OF MENTAL SCIENCE)

Published by Authority of the Royal College of Psychiatrists

© 1978

Vol 132

MARCH 1978

EDITOR

J. L. Crammer

ASSOCIATE EDITORS

Alexander Walk Martin Roth

ASSISTANT EDITORS

Brian Barraclough Sidney Crown Hugh Freeman Alan Kerr
Julian Leff Raymond Levy Philip Snaith

BOARD OF ASSESSORS

C. R. Bagley
J. H. J. Bancroft
Klaus Bergmann
Robert Bluglass
P. K. Bridges
George W. Brown
Edward Chesser
John Cooper
John Copeland
J. A. N. Corsellis
T. J. Crow
D. Eccleston
T. J. Fahy
G. W. Fenton
Fay Fransella
C. D. Frith
Dennis Gath
David Goldberg
P. J. Graham
K. L. Granville-Grossman
John Gunn

Max Harper
Dorothy H. Heard
W. Alan Heaton-Ward
Bernard Heine
Oscar W. Hill
Steven Hirsch
B. D. Hore
H. Hunter
J. G. Ingham
L. Iversen
Ivan Janota
A. L. Johnson
H. Gwynne Jones
R. E. Kendall
Israel Kolvin
R. Kumar
D. W. Liddell
C. J. Lucas
Pamela C. B. MacKinnon
C. McCance
B. M. Mandelbrote

R. H. S. Mindham
P. A. P. Moran
Alan Norton
D. J. Pallis
E. S. Paykel
F. Post
R. T. C. Pratt
Philip H. Rack
Peter H. Schurr
C. P. Seager
David M. Shaw
B. F. Sheffield
J. Shields
Alwyn Smith
E. B. O. Smith
John Steiner
Peter H. Venables
A. Wakeling
Dermot Walsh
Lorna Wing
Sula Wolff

Publishers

HEADLEY BROTHERS LTD, Ashford, Kent TN24 8HH

Prothiaden

What's the significance of the new 75mg tablet?

Presentation Prothiaden is dothiepin hydrochloride, an antidepressant of the tricyclic group.

Prothiaden is available as sugar coated tablets, each containing 75mg of dothiepin hydrochloride. The tablets are red in colour and bear the overprint 'P 75' in white. It is also available in hard gelatin capsules, each containing 25mg of dothiepin hydrochloride. The capsules are red/brown in colour with the overprint 'P25' in white.

Uses Prothiaden is indicated in the treatment of depression and the anxiety frequently associated with depressive illness.

Dosage and Administration Prothiaden should be given in a dosage of 75 to 150mg daily. The following dosage schemes are suggested:

Mild to moderate depression, 25mg three times daily or 75mg at night.
Moderate to severe depression, 150mg daily in divided doses, or as a single dose at night. (If the latter regimen is adopted, it is preferable to use a smaller dose for the first few days).

In certain circumstances, i.e. in hospital use, Prothiaden has been given at dosages up to 225mg daily. No dietary restrictions are necessary during treatment with Prothiaden.

Contra-indications Warnings etc. Prothiaden has anticholinergic properties; therefore, it may precipitate urinary retention in susceptible individuals and its use should be avoided in patients with existing or potential urinary retention. Patients with closed-angle glaucoma should not be given Prothiaden and the occurrence of a painful red eye in a patient receiving the drug may indicate acute closed-angle glaucoma; this requires urgent treatment. In patients with chronic simple glaucoma, the risk of Prothiaden causing a rise in intraocular pressure is relatively small provided adequate anti-glaucoma therapy is being used. Caution is advised when treating epileptic patients and those with cardiovascular disorders. From studies in animals, it was concluded that Prothiaden had no teratogenic effects in the species tested. Nevertheless, as with any relatively new drug, the use of Prothiaden during pregnancy should be avoided if possible.

Use with other drugs. Prothiaden should not be given concurrently with MAO inhibitors, nor should it be given within 14 days of ceasing treatment with an MAO inhibitor. Prothiaden may alter the pharmacological effects of some concurrently administered drugs: CNS depressants, including alcohol and narcotic analgesics, will be potentiated, as will the effects of adrenaline and noradrenaline (it should be borne in mind that some local anaesthetic preparations contain these sympathomimetics). The hypotensive effect of certain antihypertensive agents (e.g. bethanidine, debrisoquine, guanethidine) may be reduced.

Side effects. Generally, the side-effects associated with Prothiaden have been mild and controlled by reduced dosage. The following have been reported—dryness of mouth, constipation, disturbed accommodation, lassitude, dizziness, orthostatic hypotension, palpitations, somnolence, tremor, headache.

Overdosage. The symptoms of overdosage with Prothiaden may include sedation, dry mouth, blurring of vision, tachycardia, tremor, sweating, nausea, vomiting, confusion. The main dangers from overdosage arise from unconsciousness, convulsions, abnormal cardiac rhythms, hypotension and depression of respiration. The smallest dose of Prothiaden alone which resulted in the death of an adult was reported to be 0.75–1.0g (30 to 40 x 25mg capsules). The largest dose from which recovery took place was reported to be 5.0g (200 x 25mg capsules). In view of the many factors which influence the outcome of an overdose, these figures should not be considered in isolation.

Treatment of overdosage. Gastric lavage, in the unconscious patient or where the cough reflex is depressed, after the lungs should be protected by a cuffed endotracheal tube. Repeated gastric/intestinal aspiration may remove drug and metabolites excreted into the gut via the bile. General support of the circulation with continuous ECG monitoring is advised. Abnormalities of cardiac rhythm and epileptic convulsions may occur and should be treated accordingly. Forced diuresis and haemodialysis are not recommended. Bed-rest is advisable, even after clinical recovery.

Pharmaceutical Precautions—Recommended storage conditions 5°C to 20°C.

Legal Category POM.

Package Quantities—Prothiaden Tablets (75mg) 100, 500. Prothiaden Capsules (25mg) 100, 600.

Basic NHS Prices: 100 x 25mg £157 600 x 25mg £903
100 x 75mg £452 500 x 75mg £2170

Product Licence Number Prothiaden capsules PL 0096/5007. Prothiaden Tablets PL 0096/0046.

The Crookes Laboratories Ltd., Basingstoke, Hants. 

References: 1. Pearce, J. B. & Linford Rees, W. J., *Int. Med. Res.* 1974, **2**, 12.
2. Rees, J. A. & Cryer, P. C. *Curr. Med. Res. Opin.* 1976, **4**, 416.

... 'dothiepin in a single daily dose is better tolerated and produces a greater overall response than a divided dose regimen of amitriptyline.'

2

... 'with dothiepin both the incidence and severity of side-effects were much less than with amitriptyline.'

2.

... 'A single daily dose of dothiepin (Prothiaden) has been shown to produce a better clinical response and to be better tolerated than a divided daily dose regimen.' 1

Prothiaden 75mg

more effective, better tolerated
treatment for depression and anxiety.