

22 May 2012
INTERNATIONAL DAY
FOR BIOLOGICAL DIVERSITY
Marine Biodiversity



MARINE BIODIVERSITY IN INDIA


Venkataraman K, Raghunathan C, Raghuraman R, Sreeraj CR

Zoological Survey of India



MARINE BIODIVERSITY IN INDIA

Venkataraman K, Raghunathan C, Raghuraman R, Sreeraj CR



Zoological Survey of India

CITATION

Venkataraman K, Raghunathan C, Raghuraman R, Sreeraj CR; 2012. *Marine Biodiversity* : 1-164 (Published by the Director, Zool. Surv. India, Kolkata)

Published : May, 2012

ISBN 978-81-8171-307-0

© Govt. of India, 2012

Printing of Publication Supported by NBA

Published at the Publication Division by the Director, Zoological Survey of India, M-Block, New Alipore, Kolkata-700 053
Printed at Calcutta Repro Graphics, Kolkata-700 006.



Dr. Balakrishna Pisupati
Chairman



सत्यमेव जयते

भारत सरकार
Govt. of India
NATIONAL BIODIVERSITY AUTHORITY



FOREWORD

The marine ecosystem is home to the richest and most diverse faunal and floral communities. India has a coastline of 8,118 km, with an exclusive economic zone (EEZ) of 2.02 million sq km and a continental shelf area of 468,000 sq km, spread across 10 coastal States and seven Union Territories, including the islands of Andaman and Nicobar and Lakshadweep. Indian coastal waters are extremely diverse attributing to the geomorphologic and climatic variations along the coast. The coastal and marine habitat includes near shore, gulf waters, creeks, tidal flats, mud flats, coastal dunes, mangroves, marshes, wetlands, seaweed and seagrass beds, deltaic plains, estuaries, lagoons and coral reefs. There are four major coral reef areas in India-along the coasts of the Andaman and Nicobar group of islands, the Lakshadweep group of islands, the Gulf of Mannar and the Gulf of Kachchh. The Andaman and Nicobar group is the richest in terms of diversity.

Mangrove ecosystems are found along both the east and west coasts of India, covering an estimated area of 4,120 sq km. Important mangrove areas are in the Sundarbans, Bhitarkanika, Krishna and Godavari delta of Andhra Pradesh, Andaman and Nicobar Islands, Gulf of Kachchh, and the Pichavaram-Vedaranyam area of Tamil Nadu coast. Seagrass beds are found along the coasts of Tamil Nadu, Lakshadweep islands, Andaman and Nicobar Islands, and the Sundarbans. Eight hundred and forty four species of seaweeds are found in shallow waters all along the Indian coast, particularly in Tamil Nadu, Gujarat, Goa, Maharashtra and Lakshadweep.


The biodiversity of marine habitats is under threat around the world oceans. Although, seas cover majority of our planet's surface, far less is known about the marine biodiversity than the terrestrial systems. Conservation of marine biodiversity has become an international issue since the extinction of the Steller's sea cow in 1868. Until we have complete information about the diversity of marine habitats and the impact of human activities etc. on them, the conservation of marine biodiversity remains a distant dream.

Considering the significance of marine biodiversity conservation, the Zoological Survey of India has brought out this book on "Marine Biodiversity in India" as an informal and non-technical prose depicting illustration of marine biotic communities in Indian seas, towards the understanding and benefit of the general public. This book brings together the representative of most of the faunal groups including few floral communities of marine realm. Real life pictures of underwater creatures taken by the authors, presented in this book, would definitely form a source of invaluable source of awareness.

More and more knowledge about the seas of India would evolve with time as new generations of people interact with the seas around. Humans are continuously and rapidly reducing the number of marine species resulting in depletion of their habitats and communities, leading to the destruction of the marine ecosystem.


In this milieu, I appreciate Drs K Venkataraman, Director and C. Raghunathan, Scientist C, Zoological Survey of India, in bringing out this valuable source of information on Marine Biodiversity to public domain. I sincerely hope that this effort succeeds in creating an awareness to conserve the marine ecosystem for the future generation of India.

April 30, 2012


(Balakrishna Pisupati)
Chairman,
National Biodiversity Authority



Dr. K. VENKATARAMAN
Director



सत्यमेव जयते

Government of India
Ministry of Environment and Forests
ZOOLOGICAL SURVEY OF INDIA

M-Block, New Alipore, Kolkata – 700 053

PREFACE

Homo sapiens have a very biased view of planet Earth; its proper name should be *Oceanus* or *Water* (J.Hope and M. Zacharias, 2011). The oceans are the dominant feature of our planet covering nearly 71 per cent of its surface. The oceans are responsible for the regulatory control of conditions on land; the oceans modulate and moderate the terrestrial climate. It is no exaggeration that life in the oceans are self sustainable and not dependent on land of our dear planet. The oceanic realm is an inexhaustible cornucopia for the humankind. Oceans provide great wealth and diversity of food, medicines and raw materials. Oceanic resources like fish, seaweed and other organisms serve as an important means of livelihood to millions of people.

Marine ecosystem encompasses coastal areas, sea and thousands of diverse living organisms, which are interdependent. In the three dimensional classification of water and seabed, the water mass is technically called the pelagic zone consisting of the neritic and the oceanic provinces. The marine environment, as a whole, comprises of living and non-living resources, which are presently being exploited by maritime nations. Out of the total 32 animal phyla, 15 are represented by the taxa covering 12856 species in marine ecosystem of India. They may constitute either migratory or resident species. The migratory organisms include pelagic crustaceans, coelenterates, cephalopods, fishes, reptiles, birds and mammals. The benthic macro fauna comprises resident species of polychaetes, bivalves, gastropods, sipunculates and mud burrowing fishes. Free swimmers or nekton are important components of marine biodiversity. Out of the total 22,000 described finfish species, about 4000 species occur in the Indian Ocean, of which 1800 species are reported in Indian seas.

There are potentially many ways to collate and discuss threats to marine biodiversity, but threats can be broadly categorized as a result of over harvesting, pollution, habitat loss, introduced species and global climate change. Many marine areas have a range of biota rivaling or exceeding that of tropical forests. However, the diversity of life in our oceans is now being drastically altered by increasing irreversible anthropogenic activities. Humanity's response to our deteriorating marine environment has been predictably slow, reactive and piecemeal. Delays in responding to this environmental crisis are exacerbated due to the fact that most marine environments are still viewed as a global common resource, where there is little incentive to any one nation to address these issues, and those problems must be solved at an international level.

Amidst the burgeoning threats, it is mandatory to understand our own marine bio-resources at layman level. Hence it is aimed at bringing out this book on Marine Biodiversity with the pictures of undersea life for the benefit of commons. I hope this book will be useful for students and general public to understand our marine wealth.

May, 2012

(K. Venkataraman)

CONTENTS



Introdcution
1-3



Sea fans
19-25



Hard corals
38-51



Ribbon worms
64-65



Zooplankton
4



Sea pens
26-27



Black corals
52-54



Segmented worms
66-71



Sponges
5-11



Sea anemones
28-32



Jelly fish
55-56



Peanut worms
72



Hydroids
12-13



Tube Anemones
33




Siphonophores
57



Chitons
73-74



Hydrocorals
14



Zooanthids
34-35



Comb jellies
58



Snails
75-84



Soft corals
15-18



Corallimorpharians
36-37



Flat worms
59-63



Bivalves
85-89

CONTENTS



Sea slugs
90-99



Hermit crabs
113



Sea urchins
126-129



Phytoplankton
162




Cephalopods
100-102



Lobsters
114



Sea cucumbers
130-134



Sea grasses
163



Barnacles
103-104



Feather stars
115-120



Sea squirts
135-137



Sea weeds
164-165



Mantis shrimps
105



Sea stars
121-123



Fishes
138-156



Mangroves
166-168



Shrimps
106-108



Brittle stars
124



Marine Reptiles
157-159



Crabs
109-112



Sand dollars
125



Marine Mammals
160-161



Introduction



From bacteria to baleen whales, our planet is home to tens of millions of different life forms at least; biologists can only guess at the true number of species. Everywhere you look in the ocean there are living things. How many and what kind of organisms there are depend on where you go—that is, on the specific nature of the habitat. Every habitat has distinct characteristics that determine which organisms live there and which do not. The amount of light, for example, determines whether algae and plants can grow. The type of bottom, the temperature and salinity of the water, waves, tides, currents, and many other aspects of the environment profoundly affect marine life.



There are so many different living things in the sea that it almost boggles the mind. From microscopic bacteria to gigantic whales, marine organisms come in all shapes, sizes, and colors. The ways in which they live are just as varied. Making sense of all this diversity might seem a hopeless task. Fortunately, there is a unifying concept that helps make the bewildering diversity of life comprehensible. This concept is the theory of evolution. Remember that scientists do not use the term “theory” lightly. Evolution, the gradual alteration of a species' genetic makeup, is supported by a vast body of evidence. It is as well established as the theory of gravity. The way in which evolution occurs, on the other hand, never ceases to fascinate biologists. In most simple form marine biodiversity is 'Life on Earth', including plants, animals and microorganisms, the genes that they contain and the ecosystems that they form.

Accelerated loss of coastal and marine biodiversity components over the last few decades has been of great concern. Environmental changes, over exploitation and habitat loss are among the major causes of species loss that, according to certain estimates, is of the order of a species a day. Equally important as knowledge of what lives in the seas, is a prediction of what would live there in the future. This is especially true of regions where rapid loss of habitats and decline in water quality could be drastically altering the species diversity.

Marine biologists are often frustrated by how hard it is to actually see what is going on in the ocean. They can collect samples with nets and dredges, make measurements with automated instruments, and do experiments in the laboratory, and in these ways have learned a great deal about marine life.



We humans are visual creatures, though, and no amount of sampling, measurement, or experimentation can completely substitute for actually watching organisms in their natural habitats. One solution is to enter the ocean and observe with our own eyes. Scuba and research submersibles have tremendously aided the study of the ocean, allowing us not only to see the organisms we study, but also to conduct experiments in the natural environment. These methods have

their limitations, though. Scuba divers can penetrate only the shallowest parts of the ocean, and then only for short periods on the order of a few hours. Even though SCUBA diving has dramatically expanded the possibilities for the study of the sub tidal environment by marine biologists. The present booklet elucidate the few examples of marine biodiversity of India attempted by the researchers of Zoological Survey of India.

Zooplankton

Plankton (Singular plankter) literally mean animals which cannot swim against water currents. They keep drifting in the aquatic environment in the direction of the wind or water current. Zooplankton are the heterotropic component of the plankton community, which range from microscopic to few feet in size. Even though there are many permanent members, much of its composition is made by the juveniles of some animal groups. They form an important link in the aquatic food chain as 'secondary producers' accumulating the carbon for transferring to the other level of food chain, the consumers. Interestingly many large animals like some whales are exclusively dependent on this group of animals for their sustenance.



Microsetella norvegica



Acartia sp.

Sponges



Clathria mimia / Fire sponge

▲ One of the common encrusting sponges seen throughout the Indo-Pacific region. The species is easily recognized by the white veined pattern branching from the oscula (opening).

Phylum : Porifera


The animal consists of jelly-like matrix sandwiched between two thin layers of cells. Sponges do not have nervous, digestive or circulatory systems. All are sessile aquatic animals. More commonly, the mesohyl is stiffened by mineral spicules, by spongin fibers or both. Demosponges use spongin, and in many species, silica spicules and in some species, calcium carbonate exoskeletons.



Monanchora arbuscula / Red encrusting sponge

Seen in wave-exposed or shaded, vertical and overhanging locations, the species is sheet-like, although a bushy form occurs in protected or deeper environments.

There is one large osculum (opening) and many smaller ostia (pores). The ring of branched papillations around the orifice gives the appearance of octopus suckers.



Crella cyathophora / Honey comb sponge



Haliclona fascigera / Pink tube sponge

This is almost exclusively finding on corals, and is great competitor for the corals. This species kills the coral colony in due course of time. ▶



Chalinula nematifera / Coral encrusting sponge

Colonies of this species form yellow puffy balls with large craters on the surface.



Paratetilla bacca | Golf ball sponge



Acanthella klethra | Orange ball sponge



Stylissa massa | Mango sponge

A tough leathery sponge, it grows in a plate like, laminar shape or crescent shaped form.

The exterior surface is smooth and fleshy resulting from the lack of large spicules.



Plakortis simplex | Chicken liver sponge



Carteriospongia foliascens | Fan sponge



◀ Colonies are solitary black ball with a large central cavity. Outer surface often heavily encrusted with epizoic organisms.

Ircinia strobilina | Black ball sponge



This is one of the largest and easily recognized sponges. Attains the height of more than a meter. ▶

Xestospongia testudinaria | Barrel sponge



Eudendrium sp. | Funnel hydroid

Phylum : Cnidaria
Class : Hydrozoa

Hydroids are mostly benthic feather like colonies, in which the polypoid generation predominates. Many species have a jelly fish like sexual stage known as hydromedusa.

◀ The polyps are trumpet shaped or funnel shaped each with one whorl of narrow tentacles. The aeolid nudibranch *Flabellina rubrolineata* feeds on this hydroid.

Hydroids



Macrorhynchia philippina | Fire or stinging hydroid

Hydrocorals

Phylum: Cnidaria


Hydrocorals are common constituents of coral reefs. The fire corals are often associated with reef flats while the lace corals are mostly restricted to caves and vertical walls.

Frequently encountered in caves and shaded vertical surfaces.



Distichopora sp. | Lace coral

A copiously branched stylasterid with slender ultimate branches. Commonly encountered in shaded locations.



Stylaster sp. | Lace coral

Due to toxins injected by the stinging cells, this species is capable of inflicting very irritating stings to scantily dressed divers.



Millepora sp. | Branching fire coral

Soft corals




Sarcophyton sp. / Mushroom leather coral


Phylum : Cnidaria
Class : Anthozoa
Subclass : Octocorallia
Order : Alcyonacea

The soft corals are an order of cnidarians which do not produce calcium carbonate skeletons and so are neither reef-building corals nor do they lay new foundations for future corals. Instead they contain minute, spiny skeletal elements called sclerites. Aside from their scientific utility in species identification, sclerites give these corals some degree of support and give their flesh a spiky, grainy texture that deters predators.

The polyp bearing capitulum arising from a smooth basal stalk. When the larger polyps are retracted, the colony takes on a very smooth leathery appearance.



Dendronephthya sp. | Divaricate tree coral



◀ In this species each polyp is incapable of totally withdrawing into the branches of the coral. The colonies have a prickly appearance due to sharp supporting bundles of sclerites on each polyp.




Lobophytum sp. | Lobed leather coral

◀ The upper surface is mostly lobed, the lobes being rounded or finger type. The polyps can retract completely into the body of the coral. Very common in the reef areas.



Cladiella australis | Knobby leather coral

▶ The upper surface is bumpy or has many round fingers like lobes.



Verrucella sp.



Order : Gorgonacea

A sea fan, also known as gorgonian, is an order of sessile colonial cnidarian found throughout the oceans of the world, especially tropics and subtropics. Gorgonians are similar to the sea pen, another soft coral. Individual tiny polyps form colonies that are normally erect, flattened, branching, and reminiscent of a fan. They may be brightly coloured, often purple, red or yellow.

Sea fans



Mopsilla sp.



▲ Large colorful and picturesque sea fan up to one meter or more in height. This is perhaps the most well know species of sea fan.



Melithaea ochracea | Neon coral sea fan



Nicella flabellata



Echinogorgia flora



Verucella sp.



Isis hippuris | Golden sea fan

▲ This is a species of sea fan found in the Indo-West Pacific region. It produces hippuristanol, a molecule with apparent anti-cancer functions.

Sea pens



Cavernularia pusilla | Flower sea pen

Order: Pennatulacea

Sea pens are highly specialized octocorals that are well adapted for life in soft sediments such as sand or mud. Many sea pens are nocturnal.



Veretillum sp.



Virgularia gustaviana



Virgularia sp.

They have earned the name because many species, but certainly not all, look like an antique quill pen.

Sea anemones

Subclass: Hexacorallia


Order: Actiniaria

Sea anemones are a group of water-dwelling, predatory animals of the order Actiniaria; they are named after the anemone, a terrestrial flower. Sea anemones and hard corals are anatomically similar, differing mainly in the presence of a calcium carbonate skeleton in the latter. Majority of them in the reef have symbiotic associations with anemone fishes.



▲ Large anemones sometimes over 1.5 meters in diameter.

Heteractis crispa | Leathery sea anemone



▲ A beautifully colored anemone. Perhaps the most commonly photographed of all reef anemones. It prefers exposed locations with sufficient current and surge.

Heteractis magnifica | Magnificent sea anemone



Actinostephanus sp.



Triactis sp.



Cryptodendrum adhaesivum | Adhesive sea anemone

▲ These are usually differently colored with blue, yellow, pink, green or brown.



Bolocerooides sp.



Heteractis sp.

The tentacles have ring like swellings at intervals along the length giving them a bead like appearance.




Heteractis aurora | Beaded sea anemone

Individual anemones are attached to the shells of hermit crabs. They are symbiotic with hermit crabs of the genera *Dardanus* and *Eupagurus*.



Calliactis miriam / Hermit crab anemone

Tube anemones



Class: Anthozoa

Subclass: Ceriantipatharia

Order: Ceriantharia

Tube-dwelling anemones or cerianthids look very similar to sea anemones, but belong to an entirely different subclass of anthozoans. They are solitary, living buried in soft sediments. Tube anemones live and can withdraw into tubes, which are made of a fibrous material.



Palythoa sp.

Subclass: Hexacorallia

Order: Zoantharia

Small anemone like animals. Most are colonial, being connected by a basal stolon or a common membranous mat.

Zoanthids



Protopalycha sp.

Corallimorpharians



Subclass: Zoantharia

Order: Corallimorpharia

Corallimorpharians are cnidarians closely related to the true sea anemones. They are mostly tropical, with a narrow column topped with a wide oral disc. The tentacles are usually short or very short, arranged in rows radiating from the mouth. Many species occur together in large groups.

◀ Fully expanded animals form plate like discs.
Contracted individuals are rounded and bowl like.

Amplexidiscus fenestrafer | Giant mushroom anemone

This corallimorpharian has numerous large mammiform to cylindrical tubercles on the face of the disc. Infrequently encountered in cracks or depressions.



Discosoma sp. | Disc anemone



Hard corals

Phylum: Cnidaria

Class: Anthozoa

There is something special about coral reefs. The warm, clear water, spectacular colors, and multitude of living things captivate almost everyone who sees a reef. Coral reefs rival that other great tropical community, the rain forest, in their beauty, richness, and complexity. Tropical rain forests and coral reefs are also similar in that the basic physical structure of both communities is produced by organisms. Both reef-building corals and the giant trees of a rain forest create a three-dimensional framework that is home to an incredible assortment of organisms.

Coral reefs are made of vast amounts of calcium carbonate (CaCO_3), limestone that is deposited by living things. Of the thousands of species in coral reef communities, only a fraction produces the limestone that builds the reef. The most important of these reef building organisms, are corals.


Although corals can catch small fish and animals, such as plankton, using stinging cells on their tentacles, most corals obtain the majority of their energy and nutrients from photosynthetic unicellular algae called zooxanthellae. Such corals require sunlight and grow in clear, shallow water, typically at depths shallower than 60 meters.



Brush coral | *Acropora hyacinthus*



Briar coral | *Anacropora reticulata*



Thorn coral | *Stylocoeniella armata*



Thorn coral | *Stylocoeniella guentheri*



Birds Nest Coral | *Seraitopora hystrix*



Cauliflower coral | *Stylophora pistillata*



Pearl Bubble coral | *Physogyra lichtensteini*



Torch coral | *Euphyllia glabrescens*



Fluorescence grass coral | *Galaxea fascicularis*



Galaxy coral | *Galaxea astreata*



Porcelain coral | *Leptoseris explanata*



Stony pillar coral | *Psammacora digitata*



Lettuce coral | *Pavona cactus*




Elephant skin coral | *Pachyseris ruguosa*



Mushroom coral | *Fungia scutaria*



Stone-leaf coral | *Lithophyllon undulatum*



Spiny Cup Coral | *Pectinia pectinata*



Elephant nose coral | *Mycedium elephantotus*



Ruffled coral | *Merulina ampliata*



Velvet horn coral | *Hydnophora microconos*



Tree coral | *Dendrophyllia robusta*



Pagoda coral | *Turbinaria mesenterina*





Sinuuous cup coral | *Symphyllia radians*



Brain coral | *Lobophyllia hemprichii*



Knob coral | *Favia lacuna*



Honey comb coral | *Favites abidita*



Massive stony coral | *Porites arnaudi*



Flower pot coral | *Goniopora tenuidens*

Black Corals

Class: Anthozoa

Subclass: Ceriantipatharia

Order: Antipatharia

They normally occur in the tropics. Though black coral's living tissue is brilliantly colored, it takes its name from the distinctive black or dark brown color of its skeleton. Also unique to black coral are the tiny spines that cover the surface of the skeleton, the origin of the nickname little thorn coral. Black coral is listed in Appendix II of the Convention on International Trade in Endangered Species (CITES).



Feather black coral | *Plumapathes pennacea*

▲ Can be sparsely or densely branched with pinnulated colonies and is less than half a meter in height.

These black corals are long, slender and whip like. Frequently encountered in deeper reefs.



Whip coral/Wire coral | *Cirripathes* sp.



▲ Whip coral | *Cirripathes anguina*
Mainly encountered in the deeper areas of reef flats.



▲ Wire coral | *Stichopathes* sp.
Unbranched and sinuous colonies. Frequently encountered on reef margins.

Jelly fishes

Phylum: Cnidaria

Subphylum: Medusozoa

Jellyfish are found in every ocean, from the surface to the deep sea. Some hydrozoan jellyfish, or hydromedusae, are also found in fresh water; freshwater species are less than an inch in diameter, are colorless and do not sting. Large, often colorful, jellyfish are common in coastal zones worldwide.



Aurelia sp.

◀ The tentacles are capable of inflicting an intense sting, which can last for many hours.



▲ Upside down jelly fish *Cassiopeia Andromeda*

It lies upside down in shallow waters usually in sandy areas, absorbing sunlight for photosynthesis by the symbiotic algae in its tissues.



▲ Moon jelly fish *Aurelia aurita*

Aurelia aurita is one of a group of more than ten morphologically nearly identical jellyfish species in the genus *Aurelia*. In general, it is nearly impossible to identify *Aurelia* medusae without genetic sampling.

Siphonophores

Despite its outward appearance, the Man o' War is not a true jellyfish but a siphonophore, which differ from jellyfish in that they are not actually a single creature, but a colonial organism made up of many minute individuals called zooids. The Portuguese Man of War is responsible for up to 10,000 human stings in Australia each summer, particularly on the east coast. Stings usually cause severe pain to humans, leaving whip-like, red welts on the skin that normally last 2 or 3 days after the initial sting, though the pain should subside after about an hour.



Portuguese Man of War | *Physalia physalis*

Class : Hydrozoa

Order : Siphonophora

They are colonial, but the colonies can superficially resemble jellyfish; although they appear to be a single organism, each specimen is actually a colony of Siphonophora. The best known species is the dangerous Portuguese Man of War.




Benthic comb jelly | *Ceoplana meteoris*

▲ The threads like tentacles are long and feather like. Usually found on soft sediment.

Comb jellies

Phylum: Ctenophora

Their most distinctive feature is the "combs", groups of cilia that they use for swimming, and they are the largest animals that swim by means of cilia. Ctenophores may be abundant during the summer months in some coastal locations, but in other places they are uncommon and difficult to find. In bays where they occur in very high numbers, predation by ctenophores may control the populations of small zooplanktons such as copepods, which might otherwise wipe out the phytoplankton.



Ceoplana sp.

Flat worms

Phylum: Platyhelminthes

Class: Turbellaria

Order: Polycladida

Polyclad flatworms include some of the most flamboyant colorful animals of the sea. As thin as a leaf and usually oval in shape, these animals may first catch your attention by their spectacular color patterns.



Cycloporus venetus

▲ Often found feeding on colonial ascidians from the reef slope.



Commonly encountered species, but as yet unnamed and undescribed.


Acanthozoon sp.



Found out and about on reef crest and reef slope, sometimes swimming in the water column.

Pseudobiceros bedfordi

Velvety black colored species. Commonly found feeding on tunicates.



Thysanozoon nigropapillosum



Pseudobiceros stellae

Found under rubble on the reef crest; animal active at night.



Pseudoceros gamblei

Found on the reef crest and intertidal areas under rocks and rubble.

Found under rubble on the reef crest.



Pseudoceros concinnus

Due to the color pattern
this species is very
difficult to be get
noticed in the reef.



Pseudoceros goslineri

Ribbon worms



Usually the adult is more than a meter in length. It is nocturnal inhabiting shallow, sandy habitats.

Phylum : Nemertea

Although most are less than 20 centimetres long, one specimen has been estimated at 54 metres, which would make it the longest animal ever found. A few live in the open ocean while the rest find or make hiding places on the bottom. Most are carnivores, feeding on annelids, clams and crustaceans, and may kill annelids of about their own size. They sometimes take fish, both living and dead.

Five lined ribbon worm | *Baseodiscus quinquelineatus*



▲ This slender nemertean exceeds one meter in length when fully extended.

Barred ribbon worm *Baseodiscus hemprichii*

Segmented worms

Phylum: Annelida

Class: Polychaeta

Polychaetes are a class of annelid worms, generally marine. Each body segment has a pair of fleshy protrusions called parapodia that bear many bristles, called chaetae, which are made of chitin. Polychaetes are extremely variable in both form and lifestyle and include a few taxa that swim among the plankton. They are the indicator group of the marine ecosystem health.

The tentacular crown is horseshoe shaped. The tubes are generally buried in sandy substrata.



Fan worm | *Sabellastarte indica*



◀ The tube of this species is largely embedded in sand at the edge of a shallow patch reef.

Social feather duster | *Bispira brunnea*





Christmas tree worm | *Spirobranchus giganteus*

The tentacles are extremely variable in color: yellow, blue, purple, red, orange or brown. Lives in calcareous tubes which penetrate living coral heads (Mainly *Porites* sp.). An indicator species about the eutrophication in the reef.





Polynoid worm *Asterphilia* sp.

▲ A commensal species found on the asteroids and holothuroids.



▲ Scale worm | *Gastrolepida clavigera*

This species is an obligate commensal on sea cucumbers, especially species of *Bohadschia* and *Thelenota*.



Sand worm | *Nereis* sp.

▲ *Nereis* possesses setae and parapodia for locomotion. Locomotor chaetae are for crawling, and are the bristles that are visible on the exterior of the polychaeta.



Thread polychaete

This species inhabits dead and living reefs and forms burrows in blocks of coral.



Sipunculus sp.

Phylum: Sipuncula

Sipunculids are relatively common, and live in shallow waters, either in burrows or in discarded shells like hermit crabs do. Some bore into solid rocks to make a shelter for themselves. Although typically less than 10 cm long, some sipunculids may reach several times that length. Sipunculid worm jelly is a delicacy in the town of Xiamen in Fujian province of China.

Peanut worms

This species is found on the underside of coral rubble that is partially buried in coarse coral gravel on shallow patch reefs.



Themiste sp.



Acanthopleura spinosa

Chitons

Phylum: Mollusca

Class: Polyplacophora

Chitons are primitive, flattened mollusks with eight shell valves. These plates overlap somewhat at the front and back edges, and yet the plates articulate well with one another. Because of this the plates provide good protection for impacts from above.



◀ Owing to the reduction of the shell plates and increased girdle, this species looks more like a worm than a mollusk.

Worm chiton | *Cryptoplax larvaeformis*

Snails

Phylum: Mollusca

Class: Gastropoda

The class gastropods contain a vast total of named species, second only to the insects in overall number. The fossil history of this class goes all the way back to the Late Cambrian. Although the name "snail" can be, and often is, applied to all the members of this class, commonly this word means only those species with an external shell large enough that the soft parts can withdraw completely into it. Those gastropods without a shell, and those with only a very reduced or internal shell, are usually known as slugs.

Egg cowry | *Cymbovula deflexa*

The animal is found exclusively on its food gorgonian *Rumphella* sp.





Funnel hydroid | *Mitra stictica*



Kettle mitre | *Mitra cucumerina*



Nassarius snail | *Nassarius* sp.



Isabelle's Mitre | *Cancilla Isabella*



Gnawed or eroded cowry | *Cypraea erosa*



Lynx Cowrie | *Cypraea lynx*



Money cowry | *Cypraea moneta*



Tiger cowry | *Cypraea tigris*



Yellow Helmet | *Cassis cornuta*



Emperor/Queen Helmet | *Cassis madagascariensis*



Chank shell | *Turbinella pyrum*



Brown turban | *Turbo bruneus*



Top shell | *Trochus niloticus*



Velvet snail | *Coriocella hibyae*



Oblong ovulid cowry | *Phenacovolva tokioi*



Phylum: Mollusca

Class: Bivalvia

Bivalves have a shell consisting of two asymmetrically rounded halves called valves that are mirror images of each other, joined at one edge by a flexible ligament called the hinge. The shell is typically bilaterally symmetrical, with the hinge lying in the sagittal plane. Bivalves are unique among the molluscs, having lost their odontophore and radula in their transition to filter feeding.

It is found cemented to walls and open reef faces in shallow to deep areas.

Variable thorny oyster | *Spondylus varians*

Bivalves



Pen shell | *Atrina pectinate*

▲ The animals are largely embedded in coarse sand and rubble with only top of the shell and mantle visible. Generally found in shallow waters.



◀ Lives in crevices of living corals. Only the shell opening and mantle visible.

Coral bivalve | *Pedum spondyloideum*



Commonly found under rocks in sandy and rubble habitats. Swims actively with the siphoning of shells ▶ when disturbed.

File shell | *Limaria orientalis*



Fluted giant clam | *Tridacna squamosa*



Crocus giant clam | *Tridacna crocea*

Giant giant clam | *Tridacna gigas*

GIANT CLAM

The giant clam is the largest living bivalve mollusc. This is one of the most endangered clam species. One of a number of large clam species native to the shallow coral reefs of the South Pacific and Indian oceans, they can weigh more than 200 kilograms measure as much as 120 cm across, and have an average lifespan in the wild of 100 years or more.



Sea slugs

Phylum : Mollusca

Class : Gastropoda

Informal group : Opisthobranchia

Sea slug is a common name used for several different groups of saltwater snails that either lack a shell or have only an internal shell, in other words this name is used for various lineages of marine gastropod mollusks that are either not shell-bearing or appear not to be. The phrase "sea slug" is often applied to nudibranchs, members of the order nudibranchia within the opisthobranch gastropods. Many nudibranchs are colorful and are a noticeable part of the underwater fauna. However the phrase "sea slug" is also applied to various other groups within the Opisthobranchia including the sea hares, the sacoglossans, various families of bubble snails (Cephalaspidea) and others, some of which have reduced shells, and many of which are shell-less.



Phyllidiella zeylanica



Chelidonura pallida



Cerberilla annulata



Aldisa erwinkoehleri



Chromodoris striatella




Costasiella paweli



Chromodoris colemani



Chromodoris fidelis





Discodoris boholiensis



Flabellina rubrolineata



Elysia grandifolia



Glossodoris cincta



Fryeria marindica



Gastropecten bicornutum



Halgerda bacalusia



Gumnodoris striata



Hypselodoris nigrostriata



Spanish Dancer | *Hexabranthus sanguineus*

This species of very large, strong-swimming nudibranch is one of the largest of all nudibranchs: specimens of well over 40 cm in length have been reported. The species name, *sanguineus*, refers to its bright red coloration, but a yellow variant also exists.



◀ Egg mass
The egg mass of Spanish dancer is commonly known as Sea rose. To protect its egg cluster from predators, the Spanish dancer deposits with its eggs some of the toxins that it produces for its own defense.




Jorunna rubescens's Egg Mass



Jorunna rubescens



Phyllidia ocellata




Phyllidiella zeylanica



Hypsodoris bullockii



Thuridilla caerulea



Phyllaplysia engeli



Philinopsis pilsbryi



Thorunna africana

Cephalopods

Phylum: Mollusca

Class: Cephalopoda

These exclusively marine animals are characterized by bilateral body symmetry, a prominent head, and a set of arms or tentacles modified from the primitive molluscan foot. Fishermen sometimes call them ink fish, referring to their common ability to squirt ink.

This species can be recognized by a black spot surrounded by another thin black ring on the base of the arm web. This animal is active during day and is found in shallow reefs.



Common octopus | *Octopus cyanea*

Cuttlefish | *Sepia* sp.

▲ This small species is characterized by large branched papillae over its surface. This species is found in shallow sandy areas. The internal shell of cuttlefish is a calcium supplementary feed for pet birds.



Broad club cuttlefish | *Sepia latimanus*

▲ This is the only large cuttlefish in most of the tropical Indo-Pacific. Mostly singles or pairs are often observed active during the day time.



Subphylum: Crustacea

Class: Maxillopoda

Subclass: Thecostraca

Infraclass: Cirripedia

Barnacles are exclusively marine, and tend to live in shallow and tidal waters, typically in erosive settings. They are sessile and suspension feeders. Barnacles are encrusters, attaching themselves permanently to a hard substrate. Barnacles are of economic consequence as they often attach themselves to man-made structures, sometimes to the structure's detriment. Particularly in the case of ships, they are classified as fouling organisms.

▲ Acorn barnacle | *Balanus* sp.

Shaped like a volcano. When submerged, the thoracic feeding appendages or cirri may be observed reaching out of the orifice, in a grasping motion.

Barnacles



◀ Though it seems to be a bivalve, the species is made up of five white plates. Found attached to floating objects such as buoys, boat hulls and driftwood.

Pelagic goose-neck barnacle | *Lepas* sp.

Mantis shrimps

Observed in coral rubble bottoms.



Lysiosquilla sp.

Subphylum: Crustacea

Class: Malacostraca

Order: Stomatopoda

Mantis shrimps are marine crustaceans. They are neither shrimp nor mantids, but receive their name purely from the physical resemblance to both the terrestrial praying mantis and the shrimp. Mantis shrimp appear in a variety of colors, from shades of browns to bright neon colors.



Pseudosquilla sp.

Burrows in sand and gravel on coral reefs and sand flats.

Shrimps

Order : Decapoda
(Shrimps, Crabs and Lobsters)

Decapods, like other crustaceans have hard, jointed exoskeletons. Their growth involves molting. The decapods (ten-footed) are an order of crustaceans, including many familiar groups, such as crabs, lobsters, prawns and shrimp. Most decapods are scavengers.



Banded Coral cleaner Shrimp | *Stenopus hispidus*

▲ This common cleaner shrimp waits in crevices or out in the open at its cleaning station, waving its antennae to attract fishes. As fish approach to be cleaned, the shrimp touches them with the antennae until the fish becomes quite, allowing the shrimp to move about its body in search of parasites.



Sexy anemone shrimp | *Thor amboinensis*

▲
It lives on corals, sea anemones and other reef communities. They perform a dance like movement which they are famous for.

Found in association with sea anemones, especially *Cryptodendrum adhaesivum*.



Hinge beak prawn | *Rhynchocinetes durbanensis*

Lives deep in crevices and holes. Usually occurring in large numbers together.



Anemone shrimp | *Periclimenes brevicarpalis*

Crabs



Ghost crab | *Ocypode ceratophthalma*

▲ This species can be distinguished from other related crabs by the eyestalks extending beyond the eyes into long points. They can run very fast in the intertidal sandy areas.



Coral Crab | *Tetralia nigrolineata*



▲ This crab is often seen associated with stag horn corals (*Acropora* sp).



▲ This species is associated with soft corals environment.

Stone crab | *Liomera rugata*



Red spotted coral crab | *Trapezia rufopunctata*

▲ This species is obligate symbionts of pocilloporid corals.



Elephant squat lobster | *Allogalathea elegans*

▲ Found in association with feather stars. The color of the individual mimics the host, so color depends on the host feather star's color.



Spotted porcelain crab | *Neopetrolisthes maculatus*

▲ Lives in the shelter of sea anemones. Individuals can enter and leave the mouth of the anemone freely.



Fiddler crab | *Uca tetragonon*



Ghost crab | *Ocypode cordimana*

Hermit crabs



Dardanus sp. | Hermit crab


Order: Decapoda

Infraorder: Anomura

Superfamily: Paguroidea

Hermit crabs are decapod crustaceans. Most of the species possess an asymmetrical abdomen which is concealed in an empty gastropod shell that is carried around by the hermit crab. The vulnerable abdomen is protected from predators by a salvaged empty seashell carried by the hermit crab, into which its whole body can retract. Most frequently hermit crabs use the shells of sea snails (although the shells of bivalves and scaphopods and even hollow pieces of wood and stone are used by some species).

Lobsters



Painted crayfish | *Panulirus versicolor*

Order: Decapoda

Infraorder: Astacidea

Family: Nephropidae

Lobsters are economically important as seafood, forming the basis of a global industry that nets more than US \$ 1 billion annually. They are also revered for their flavor and texture.

▲ This species is nocturnal and hiding in crevices during the daylight.

Feather stars

Phylum : Echinodermata

Class : Crinoidea

Crinoids are characterized by a mouth on the top surface that is surrounded by feeding arms. They have a U-shaped gut, and their anus is located next to the mouth. Although the basic echinoderm pattern of fivefold symmetry can be recognized, most crinoids have many more than five arms.



Robust Feather Star *Himerometra robustipinna*



Green Feather Star *Himerometra* sp.




Noble Feather Star | *Comanthina nobilis*



Variable Bushy Feather Star | *Comanthina schlegeli*



Many-armed Feather Star | *Comaster multibrachiatus*



Comaster multifidus



Common Feather Star | *Comanthus* sp.



Bennett's Feather Star | *Oxycomanthus bennetti*




Crevice crinoid | *Comanthus parvicirrus*

Sea stars

Phylum: Echinodermata

Class: Asteroidea

Starfish occur across a broad depth range from the intertidal to abyssal depths (>6000 m). Starfish are among the most familiar of marine animals and possess a number of widely known traits, such as regeneration and feeding on mussels. Starfish possess a wide diversity of body forms and feeding methods. Sea stars and other echinoderms pump water directly into their bodies, via the water vascular system, as they find it. This makes them vulnerable to all forms of water pollution, as they have little ability to filter the water of toxins and contaminants. Oil spills and similar events often take a toll on echinoderm populations that carry consequences for the ecosystem.



Brown mesh sea star | *Nardoa galathae*





Warty mesh sea star | *Nardoa tuberculata*



Cushion star | *Choriaster granulatus*



Indian Sea Star | *Fromia indica*



Crown-of-Thorns | *Acanthaster planci*

▲ Potential predator of corals and prey upon the coral polyps. Out breaks of crown-of-thorn star fish often leads considerable damages to the corals of Indo-Pacific.

Brittle stars

Phylum: Echinodermata

Class: Ophiuroidea

They crawl across the seafloor using their flexible arms for locomotion. Brittle stars use their arms for locomotion. They do not, like sea stars, depend on tube feet, which are mere sensory tentacles without suction. Brittle stars move fairly rapidly by wriggling their arms which are highly flexible and enable the animals to make either snake-like or rowing movements. Brittle stars live in areas from the low-tide level downwards.



Banded Brittle Star | *Ophiacantha indica*



Green Brittle starfish | *Ophiarachna incrassate*



Toothed Brittle Star | *Ophiocoma dentata*

Sand dollars

Phylum: Echinodermata

Class: Echinoidea

Order: Clypeasteroidea

The term sand dollar refers to the species of extremely flattened, burrowing echinoids. Some species within the order, not quite as flat, are known as sea biscuits. Coordinated movements of the spines enable sand dollars to move across the seabed. The velvety spines of live sand dollars appear in a variety of colors—green, blue, violet, purple—depending on the species. The tests of dead individuals are often found on beaches, the textured skin missing and the skeleton bleached white by sunlight.



Laganum Sand Dollar | *Laganum depressum*



Sand Dollar | *Peronella lessueri*

Sea urchins

Subphylum: Echinozoa

Class: Echinoidea

Sea urchins are small, spiny, globular animals which inhabit all oceans. Common colors include black and dull shades of green, olive, brown, purple, and red. They move slowly, feeding mostly on algae. Sea otters, wolf eels, triggerfish, and other predators feed on them. Population density varies by habitat with more dense populations being found in barren areas as compared to kelp stands. Even in these barren areas, greatest densities are also found in shallow water. Populations are also generally found in deeper water if wave action is present.



Diadem Urchin | *Echinothrix diadema*




Pencil Urchin | *Phyllacanthus imperialis*



Crown Urchin | *Prinocidaris verticillata*



Burrowing Urchin | *Echinometra mathaei*



Hatpin Urchin | *Echinothrix calamaris*



Lance urchin | *Phyllacanthus imperialis*



Pincushion Tuxedo Urchin | *Mespilia globulus*

Sea cucumbers

Phylum: Echinodermata

Subphylum: Echinozoa

Class: Holothuroidea

They are marine animals with a leathery skin and an elongated body containing a single, branched gonad. Sea cucumbers are found on the sea floor worldwide. There are many commercially important species of sea cucumber that are harvested and dried for export for use in Chinese cuisine as Hoi sam. Some varieties of sea cucumber (known as gamat in Malaysia or teripang in Indonesia) are said to have excellent healing properties.



▲ This species mimics the nudibranch *Phyllidia ocellata*; which helps them from predators. This species when juvenile has this colorful phase, but as it grows it becomes a typical boring brownish sea cucumber.

Blotched Sea Cucumber | *Bohadschia graffei* (Juvenile)




Edible sea cucumber | *Holothuria edulis*

Most common sea cucumber of Indo-Pacific region. Animals may form dense aggregations in shallow waters.

As the name suggests this is one of the edible species of sea cucumbers served as “trepang” or “beche de mer”.



Black lolly fish | *Holothuria atra*



It is called Curry fish as it is among the sea cucumbers that are edible and harvested for the restaurant trade. Found in shallow water rubble and sandy habitat.

Pineapple sea cucumber | *Thelenota ananas*

This is a large species, reaching half a meter in length. It inhabits the interface between reefs and sand.





▲ This is the longest sea cucumber which may exceed even 2 meter in length. Found in shallow sandy habitats and grass beds.

Spotted Worm Cucumber | *Synapta maculata*

Sea squirts

Phylum: Chordata

Subphylum: Tunicata

Tunicates, also known as urochordates, are members of the subphylum Tunicata, a group of underwater sac like filter feeders with incurrent and excurrent siphons that is classified within the phylum Chordata. Tunicates begin life in a mobile larval stage that resembles a tadpole, later developing into a barrel-like and usually sedentary adult form.



Clavelina moluccensis



Atrium robustum



Botrioloides leachi



Perophora modificata



Didemnum molle



Didemnum sp.



Phallusia Arabica



Clavelina sp.

Fishes

Kingdom : Animalia

Phylum : Chordata

Clade (Unranked) : Craniata

A fish is any gill-bearing aquatic vertebrate (or craniate) animal that lacks limbs with digits. Most fish are "cold-blooded", or ectothermic, allowing their body temperatures to vary as ambient temperatures change. Fish are abundant in most bodies of water. Fish, especially as food, are an important resource worldwide. Commercial and subsistence fishers hunt fish in wild fisheries or farm them in ponds or in cages in the ocean. They are also caught by recreational fishers, kept as pets, raised by fish keepers, and exhibited in public aquaria.


Forms large feeding groups in the inshore and outer reefs.



Powder blue surgeonfish | *Acanthurus leucosternon*

Yellow back fusilier | *Caesio xanthonota*

Form large aggregations often mix with other fusiliers.



Scrawled file fish | *Aluterus scriptus*

◀ Solitary fish, seen in the coastal, lagoon and outerreefs.

Group of parrot fishes feeding on the corals (*Chlorurus* sp.)



Titan triggerfish | *Balistoides viridescens*

Anemone fishes

Twenty-eight species are recognized, one in the genus *Premnas*, while the remaining are in the genus *Amphiprion*. In the wild they all form symbiotic mutualisms with sea anemones. The clownfish feeds on small invertebrates which otherwise potentially could harm the sea anemone, and the fecal matter from the clownfish provides nutrients to the sea anemone



Spine cheek anemone fish | *Premnas biaculeatus*




Clown anemone fish | *Amphiprion percula*

A. akallopisos, like *A. percula*, is nearly always associated with *Heteractis magnifica* and *Stichodactyla gigantea*. It resides in shallow inshore reefs as deep as 15 m with a moderate to strong current. The skunk clownfish can also be kept in captivity by aquarists.



Skunk anemone fish | *Amphiprion akallopisos*

Painted frogfish | *Antennarius pictus*

Frog fishes

Frogfish have a stocky appearance, a typical of fish. They are often brightly colored, white, yellow, red, green, or black or spotted in several colors in order to blend in with their coral surroundings. Coloration can also vary within one species, making it difficult to differentiate between them.

Giant frogfish | *Antennarius commersoni*



Dragon sea moth | *Eurypegasis draconis*



Twin spot lionfish | *Dendrochirus biocellatus*




Common lion fish | *Pterois volitans*



Porcupine fish | *Diodon hystrix*

Puffer or Porcupine fishes

Puffer fish are generally believed to be the second-most poisonous vertebrate in the world, after the Golden Poison Frog. Certain internal organs, such as liver, and sometimes their skin are highly toxic to most animals when eaten, but nevertheless the meat of some species is considered a delicacy in Japan, Korea and China.



Star puffer fish | *Arothron stellatus*

Butterfly fishes

Butterfly fish look like smaller versions of angelfish but unlike these lack pre-opercle spines at the gill covers. The conspicuous coloration of butterfly fish may be intended for interspecies communication. Butterfly fish have uninterrupted dorsal fins with tail fins that may be rounded or truncated, but are never forked.



White collar butterfly fish | *Chaetodon collare*



Black coral goby | *Bryaninops tigris*



Red finned triple fin | *Ucla xenogrammus*



Three lined blenny | *Ecsenius trilineatus*



Striped triple fin | *Helcogramma striatum*



Black fin sand perch | *Parapercis snyderi*



Reef lizard fish | *Synodus variegatus*



▲ Forms large drifting aggregations in the coastal reefs and outer slopes.

Big eye snapper | *Lutjanus lutjanus*



Black fin barracuda | *Sphyræna genie*



Robust ghost pipefish | *Solenostomus cyanopterus*



Ornate ghost pipefish | *Solenostomus paradoxus*

Ghost pipe fishes

Ghost pipe fishes must rank among the most exotic creatures inhabiting the earth's shallow seas.



Dwarf hawk fish | *Cirrhichthys falco*



Black belt hog fish | *Bodianus mesothorax*



Tomato grouper | *Cephalopholis sonnerati*



Saddle grouper | *Cephalopholis sexmaculata*



Black spotted moray | *Gymnothorax favagineus*



Emperor angelfish | *Pomacanthus imperator*



Long fin spade fish | *Platax teira*

Marine reptiles

Phylum: Chordata

Clade: Amniota


Class: Reptilia

Marine reptiles are reptiles which have become secondarily adapted for an aquatic or semi-aquatic life in a marine environment. Some marine reptiles, such as ichthyosaurs and mosasaurs, rarely ventured onto land and gave birth in the water. Others, such as sea turtles and saltwater crocodiles, return to shore to lay their eggs. Some marine reptiles also occasionally rest and bask on land.

Sea snakes are venomous elapid snakes that inhabit marine environments for most or all of their lives. All have paddle-like tails and many have laterally compressed bodies that give them an eel-like appearance. Unlike fish, they do not have gills and must surface regularly to breathe.



Yellow lipped sea krait | *Laticauda colubrina*



Member species of the family Crocodylidae are large aquatic reptiles that live throughout the tropics in Africa, Asia, the Americas and Australia. Crocodiles tend to congregate in freshwater habitats like rivers, lakes, wetlands and sometimes in brackish water.

This is the largest of all living reptiles. It is found in suitable habitats in Northern Australia, the eastern coast of India and parts of Southeast Asia.



Saltwater crocodile | *Crocodylus porosus*

Sea turtles are marine reptiles that inhabit all of the world's oceans except the Arctic. Sea turtles are almost always submerged in water, and, therefore, have developed an anaerobic system of respiration. Although all sea turtles breathe air, under dire circumstances they may divert to anaerobic respiration for long periods of time.

It is the only species in the genus *Chelonia*. Their common name derives from the usually green fat found beneath their carapace. Like other sea turtles, they migrate long distances between feeding grounds and hatching beaches. Many islands worldwide are known as Turtle Island due to green sea turtles nesting on their beaches.



Green sea turtle | *Chelonia mydas*

Marine mammals

Phylum: Chordata

Class: Mammalia

Marine mammals are a diverse group of 120 species of mammal that are primarily ocean-dwelling or depend on the ocean for food. Marine mammals evolved from land dwelling ancestors and share several adaptive features for life at sea such as generally large size, hydrodynamic body shapes, modified appendages and various thermoregulatory adaptations. Whales are the largest mammals ever. Different species are, however, adapted to marine life to varying degrees.



Bottlenose dolphin | *Tursiops truncatus*

It is the only living representative of the once-diverse family Dugongidae; its closest modern relative, Steller's sea cow (*Hydrodamalis gigas*), was hunted to extinction in the 18th century. It is also the only sirenian in its range, which spans the waters of at least 37 countries throughout the Indo-Pacific, though the majority of dugongs live in the northern waters of Australia between Shark Bay and Moreton Bay. The dugong is the only strictly-marine herbivorous mammal, as all species of manatee utilize fresh water to some degree.

The IUCN lists the dugong as a species vulnerable to extinction, while the Convention on International Trade in Endangered Species limits or bans the trade of derived products. Despite being legally protected in many countries, the main causes of population decline remain anthropogenic and include hunting, habitat degradation, and fishing-related fatalities.




Sea cow | *Dugong dugon*



Phytoplankton

Phytoplankton are the autotrophic component of the plankton community. Most phytoplankton are too small to be individually seen with the unaided eye. However, when present in high enough numbers, they may appear as a green discoloration of the water due to the presence of chlorophyll within their cells. Phytoplankton are photosynthesizing microscopic organisms that inhabit the upper sunlit layer of almost all oceans and bodies of fresh water. They are agents for "primary production," the incorporation of carbon from living organisms, a process that sustains the aquatic food web.



Grammatophora undulata



Oscillatoria margarilifera

Seagrasses

Kingdom : Plantae
Clade : Angiosperms
Clade : Monocots
Order : Alismatales

Sea grasses are flowering plants which grow in marine, fully-saline environments. These unusual marine flowering plants are called sea grasses because the leaves are long and narrow and are very often green, and because the plants often grow in large "meadows" which look like grassland: in other words many of the species of sea grasses superficially resemble terrestrial grasses.



Turtle grass | *Thalassia* sp.



Paddle grass | *Halophila* sp.

Seaweeds

Kingdom: Protista

Seaweed is a loose, colloquial term encompassing macroscopic, multicellular, benthic marine algae. The term includes some members of the red, brown and green algae. Seaweeds can also be classified by use (as food, medicine, fertilizer, industrial, etc.). In addition, some tuft-forming bluegreen algae (Cyanobacteria) are sometimes considered as seaweeds — "seaweed" is a colloquial term and lacks a formal definition.



Cactus algae | *Halimeda* sp.



Mermaid's wine cup | *Acetabularia* sp.



Peacock seaweed | *Padina* sp.



Spiny-leaf seaweed | *Turbinaria* sp.



Cactus algae | *Halimeda* sp.



Button seaweed | *Dictyosphaeria* sp.

Mangroves

Mangroves are various kinds of trees up to medium height and shrubs that grow in saline coastal sediment habitats in the tropics and subtropics. The mangrove biome, or mangal, is a distinct saline woodland or shrubland habitat characterized by depositional coastal environments, where fine sediments (often with high organic content) collect in areas protected from high-energy wave action. Mangroves dominate three quarters of tropical coastlines. About 110 species are considered mangroves. However, a given mangrove typically features only a small number of tree species.

Small to medium sized trees grow upto 15 m tall. Occur in less saline parts of mangrove forests on deeply muddy soil, along tidal creeks with slow moving water. Found along rivers, occurring upstream to where the tidal influence is still felt.



Mangrove apple | *Sonneratia caseolaris*

Large tree grows to 20 m tall. Found solitary at the front of the mangrove belt and usually on sandy soil and sometimes mixes with seaward *Avicennia marina* individuals. Occasionally prefers rock and gravel and can tolerate the full strength of sea water.



Mangrove apple | *Sonneratia alba*

Long fruited stilted red mangrove | *Rhizophora mucronata*

Most important and widespread mangrove species; may grow upto 15 m tall and more tolerant of sandy and firmer ground than any other species of *Rhizophora*. Grows well in soft mud too and is believed to be among the few that can survive complete daily inundation.



Straight stemmed tall tree may grow upto 15 - 25 m. Thrives under a wide range of inter tidal conditions, including salinity levels from near freshwater to full strength seawater and tolerates wide range of soil types. Typically, the most common in middle and upper inter tidal zones.

Large leafed orange mangrove | *Bruguiera gymnorrhiza*

There are so many different living things in the sea that it almost boggles the mind. From microscopic bacteria to gigantic whales, marine organisms come in all shapes, sizes, and colours. It is estimated that more than 2 lakhs species may be available in the seas of the world, of which 80,000 has been identified and in India so far 15000 has been identified. This is a comprehensive book to the fascinating marine life of India. Forty six different categories of marine life are illustrated with more than 300 underwater photographs. This book can serve as a perfect reference material for marine enthusiasts, divers, students, underwater photographers' etc. The book provides the identifications of species, based on the collective experience of the authors, who have been working for many years on the marine animals. All the groups are illustrated with few photographs of those animals which are common in Indian Seas. The illustrations of the book are a perfect source to learn about the life of the ocean. This book can help an aficionado to become underwater photographer. Almost all photographs are taken by the authors from Andaman and Nicobar Islands with few exceptions which were contributed by expert underwater photographers.