Blue Bird

Tiamulin Type B Medicated Swine Feed

For manufacture of Type C medicated swine feed only

ACTIVE INGREDIENT:

INGREDIENTS:

rice hulls, mineral oil

Indications	Amount of Type B Medicated Feed Per Ton of Feed	Tiamulin hydrogen fumarate in Complete Type C Medicated Feed Per Ton	Withdrawal Period (days)
For treatment of swine dysentery associated with <i>Brachyspira</i> (formerly <i>Serpulina or Treponema</i>) <i>hyodysenteriae</i> susceptible to tiamulin	20 lb	200 g	7
For control of porcine proliferative enteropathies (ileitis) associated with <i>Lawsonia intracellularis</i>	3.5 lb	35 g	2
For control of swine dysentery associated with <i>Brachyspira</i> (formerly <i>Serpulina or Treponema</i>) <i>hyodysenteriae</i> susceptible to tiamulin	3.5 lb	35 g	2

WARNINGS:

Observe withdrawal time shown in table above. Keep out of reach of children. Avoid contact with skin. Direct contact with skin or mucous membranes may cause irritation.

CAUTION:

- 1. Do not feed undiluted.
- 2. Do not use in feeds for animals other than swine. The effects of tiamulin on swine reproductive performance, pregnancy and lactation have not been determined.
- 3. Swine being treated with tiamulin should not have access to feeds containing polyether ionophores (e.g., lasalocid, monensin, narasin, salinomycin and semduramicin) as adverse reactions may occur.
- 4. If signs of toxicity occur, discontinue use of medicated feed.

Manufactured By: Blue Bird Feed Company Robin, IN 99999 NAH/DEN10-MP/BBB-BGL/5 Net Weight: Lot Number:

09/12