

# Virile crayfish (*Orconectes virilis*)

## Ecological Risk Screening Summary

US Fish and Wildlife Service, February 2011  
Revised, June 2015



Photo: © Jan Bosselaers, in Loughman and Simon (2011). Licensed under a Creative Commons Attribution License.

## 1 Native Range, and Status in the United States

---

### Native Range

From Benson (2015):

“Missouri, upper Mississippi, lower Ohio, and Great Lakes drainages”

### Status in the United States

From Benson (2015):

“Collected in Butte, Colusa, San Joaquin, and Yolo counties (Riegel 1959), and established in San Francisco Bay, California (Ruiz 2000). Unspecified locations in Connecticut, Maine, Massachusetts, New Hampshire, Pennsylvania, Rhode Island, Vermont, and Virginia; many creeks in Maryland, New York, and West Virginia.”

## Means of Introductions in the United States

From CABI (2015):

“The use of crayfish as live bait by anglers is attributed as a major factor in the spread of non-native crayfish species within the USA (Lodge et al., 2000; DiStefano et al., 2009).”

## Remarks

From Benson (2015):

“Found in streams with moderate flow and turbidity, abundant cover, stable water levels. Breeding occurs in July and eggs are laid in the spring. Potential as a human food resource.”

## 2 Biology and Ecology

---

### Taxonomic Hierarchy and Taxonomic Standing

From ITIS (2015):

“Kingdom Animalia  
Subkingdom Bilateria  
Infrakingdom Protostomia  
Superphylum Ecdysozoa  
Phylum Arthropoda  
Subphylum Crustacea  
Class Malacostraca  
Subclass Eumalacostraca  
Superorder Eucarida  
Order Decapoda  
Suborder Pleocyemata  
Infraorder Astacidea  
Superfamily Astacoidea  
Family Cambaridae  
Subfamily Cambarinae  
Genus Orconectes  
Subgenus Orconectes (*Gremicambarus*)  
Species *Orconectes virilis* (Hagen, 1870)”

“Taxonomic Status: valid”

### Size, Weight, and Age Range

From ANSIS (2007):

“Adults reach a size of 45–125 mm [Momot 1967; Pflieger 1996]”

“Males usually grow larger than females [Momot 1967; Bovbjerg 1970; Pflieger 1996]”

## **Environment**

From ANSIS (2007):

“Primarily found in streams and usually in areas with rocky bottoms [Bovbjerg 1970; Pflieger 1996]

Prefer fertile, warm, moderately turbid streams with lots of cover [Pflieger 1996]

Prefer cobble substrate and rocky crevices in streams [Bovbjerg 1970; Hill and Lodge 1994]

Often uses rocks, logs, and other organic debris as cover [Bovbjerg 1970; Pflieger 1996]

Occasionally dig pits in river banks and under rocks especially when water levels are low [Bovbjerg 1970]”

## **Climate/Range**

From ANSIS (2007):

“Can tolerate temperatures 0° C and 32° C [Momot 1967]

Temperatures between 24° C and 25° C are preferred [Peck 1985]

Locomotion is slowed or stopped at temperatures below 10° C [Momot 1967]

Quiescence occurs in low temperatures [Bovbjerg 1970; Pflieger 1996]”

## **Distribution Outside the United States**

Native

From ANSIS (2007):

“Native range extends from New Hampshire and Maine, across the Midwest including much of the Mississippi River and its tributaries, as far west as Colorado, south to Texas and north to Ontario, Canada [Aiken 1965]”

Introduced

From CABI (2015):

“It was deliberately introduced into France in 1897 and Sweden in 1960 but both attempts were unsuccessful. Currently in Europe, *O. virilis* is only known to be present in the Netherlands and the UK. It was introduced to both countries in 2004, and is likely to be as a result of the aquarium trade. In the Netherlands the species has only recently become widespread (Ahern et al., 2008).”

## **Means of Introduction Outside the United States**

From CABI (2015):

“The two introduced *O. virilis* populations within Europe are thought to be linked to released aquarium specimens.”

## Short description

From ANSIS (2007):

“Overall reddish brown or olive brown color [Pflieger 1996]  
Numerous yellow bumps on the medial (side closest to the head) sides of the pinchers [Pflieger 1996]  
Pinchers often have orange or orange-red tips [Pflieger 1996]  
Dark specks are often found on pinchers [Pflieger 1996]  
Paired black blotches along the abdomen, especially prominent in young and individuals that have recently molted [Pflieger 1996]”

## Biology

From GISD (2010):

### “Nutrition

*Orconectes virilis* is omnivorous and consumes a variety of live and dead animal and plant material (ANSIS, 2007). It is known to consume macroinvertebrates such as snails and insects, as well as, small fish, fish eggs, tadpoles, and macrophytes. *O. virilis* is believed to obtain most of its nutrition by scavenging dead animals (Collicut, 1998).

### Reproduction

Breeding of *Orconectes virilis* begins in the fall to early spring and extends until the adults retreat to deep water and become inactive. Breeding sometimes occurs for a brief period in the spring when the water begins warming (Collicut, 1998; ANSIS, 2007). Females store sperm for up to months and fertilize their eggs in the spring. Eggs are attached to swimmerets in a large ball resembling a raspberry. The eggs hatch one to two months after they are laid. Young hatchlings look like miniature adults and can probably grow to about 2-3cm long by the fall. (Collicut, 1998).

### Lifecycle stages

The maximum lifespan of *Orconectes virilis* is 3 years. They undergo several molts in their first few months. *O. virilis* typically reaches a length of 23-56 mm in first year and a length of 58-84 mm by the second. They reach maturity during their second summer usually around 24mm.(ANSIS, 2007). *O. virilis* undergoes form alternation between the sexually mature Form I during the mating season and the sexually immature Form II outside of the mating season ([Mathews] et al, 2008).”

## Human uses

From GISD (2010):

“*Orconectes virilis* is a popularly consumed food in the United States and increasingly more so in Europe (Collicut, undated). *O. virilis* a commonly used fish bait and is sold as such in bait shops. It is also sold commercially in the pet trade as pets or prey for predaceous aquarium fishes (Hunner, 1997). Beginning in approximately 1950 *O. virilis* was stocked as a biological control in the Colorado River watershed of western New Mexico and northeastern Arizona, primarily for vegetation control (Johnson, 1986).”

## Diseases

From Davidson et al. (2010):

“We explored bacteria, nematodes, and a virus as potential biological control agents for *O. virilis* while avoiding harm to native species. White Spot Syndrome Virus (WSSV) from shrimp was found to be highly pathogenic and readily passed by cannibalistic behavior but not by water transmission.”

From CABI (2015):

“The population of *O. virilis* from the River Lee, England was tested for crayfish plague and was found to have one of the highest infestation rates of any population of crayfish found in the UK.”

**Crayfish plague and white spot disease are OIE-reportable diseases.**

## Threat to humans

From CABI (2015):

“While there have been direct economic impacts of other introduced crayfish species such as damage to river banks, and flood defences; there are no specific examples for *O. virilis*.”

## 3 Impacts of Introductions

---

From ANSIS (2007):

“Males cause a decrease in the length and growth of some macrophytes [Chambers et al. 1990]  
Drastically modify the aquatic macrophyte and macroinvertebrate communities, which in turn may lead to a decline and reconfiguration of fish communities [Lodge and Lorman 1987; Chambers et al. 1990; Hanson et al. 1990]  
Greatly reduce the number of snails in the community [Hanson et al. 1990; Lodge et al. 1994]  
Cause a reduction in the number of small invertebrates present [Hanson et al. 1990]  
May consume eggs of sunfish, bluegill, and trout leading to reduced population sizes [Horns and Magnuson 1981; Dorn and Mittelbach 2004]”

From GISD (2010):

“*O. virilis* is known to alter and reduce macrophyte biomass and diversity (Chambers et al, 1990; Davidson et al, 2010; Olden et al, 2009). It may displace native crayfish as in the case of the endangered *Pacifastacus nigrescens* in California (Light et al, 1995); *Orconectes obscurus* and *Cambarus bartonii* in North Carolina (Cooper et al, 1998); and *Orconectes limosus* and *Cambarus bartonii bartonii* in Maryland (USA) (Schwartz et al, 1963). Field and laboratory observations indicate that it may compete with and/or cause the decline of native fish species such as juvenile desert suckers (*Catostomus clarkii*), the Sonora sucker (*Catostomus insignis*), and the federally threatened Little Colorado spinedace (*Lepidomeda vittata*) in Arizona (Davidson et al, 2010; Bryan et al, 2002); the flannelmouth sucker (*Catostomus latipinnis*) in Colorado (Carpenter, 2005); and the White Sands pupfish (*Cyprinodon tularosa*) in New Mexico

(Rogowski & Stockwell, 2006). It has been reported to contribute to the decline of the Chiricahua leopard frog (*Rana chiricahuensis*) in Arizona (Davidson et al, 2010). *O. virilis* is also known to cause the decline of native snails (ANSIS, 2007), as in the case of three forks spring snail (*Pyrgulopsis trivialis*) in Arizona (Davidson et al, 2010) and Canadian native snails *Stagnicola elodes* and *Physa gyrina* in laboratory studies (Hanson et al, 1990).”

“Additional impacts associated with *O. virilis* include the reduction of insects and other macroinvertebrates (Davidson et al, 2010; Hanson et al, 1990), the alteration of the structure and composition of littoral zones (Chambers et al, 1990), the increase in turbidity of waters (Davidson et al, 2010), and impacts to irrigation networks and levees as a result of their burrowing near head gates and weir boxes (Godfrey, 2002).”

From Martinez (2012):

“No crayfish species are native to the Colorado River Basin (CRB), including the portion of the state of Colorado west of the Continental Divide. Virile crayfish [*Orconectes virilis* (Hagen, 1870)], a recent invader in the middle Yampa River in northwestern Colorado, displayed an abrupt increase in abundance in the early 2000s, which coincided with a drought, a severe decline in the abundance of small-bodied and juvenile native fishes, and a dramatic increase in the abundance of nonnative smallmouth bass [*Micropterus dolomieu* (Lacepède, 1802)]. The annual density of virile crayfish was 6.4/m<sup>2</sup> in 2005 and 9.3/m<sup>2</sup> in 2006. The annual biomass density of virile crayfish was 9.0g/m<sup>2</sup> in 2005 and 15.8 g/m<sup>2</sup> in 2006, representing a riverwide biomass of 122 kg/ha, which equaled that of other macroinvertebrates and fish combined (120.7 kg/ha). Efforts to recover and preserve native fishes in the Upper Colorado River Basin (UCRB), particularly in the Yampa River, have been hampered by nonnative predatory fishes, but the implications of crayfish may have been overlooked and underestimated. Stream conditions during the drought apparently facilitated proliferation by virile crayfish in the middle Yampa River, likely contributing to hyperpredation on native fishes by invasive smallmouth bass.”

From Carpenter (2000):

“The virile crayfish (*Orconectes virilis*), an aggressive polytrophic species, has been introduced into many Arizona streams. ... In Sabino Creek, I manipulated crayfish densities in isolated pools to examine effects of crayfish on growth, mortality, and recruitment of Gila chub (*Gila intermedia*). Regardless of crayfish density, Gila chub declined slightly in weight and condition. Mortality and recruitment did not differ between densities of crayfish. I examined crayfish effects on benthic macroinvertebrates, a submerged aquatic macrophyte and associated invertebrates, and three fish species in a small stream in the White Mountains by fencing eight stream sections to prevent movement. The three fishes were speckled dace (*Rhinichthys osculus*), Sonora sucker (*Catostomus insignis*), and desert sucker (*C. clarki*). Molluscs > 10 mm and macrophytes were less abundant at sites with a high density of crayfish than at sites with low crayfish densities. Insect diversity was lower in high- vs. low-density sites. No treatment effect was observed on growth or condition of individually marked fish. Short-term laboratory experiments demonstrated predatory interactions and competition for shelter between crayfish and Gila chub, desert sucker, and speckled dace. Crayfish used shelter more than fish, displaced fish from shelter, and frequently attacked fish. Fish never attacked crayfish, and only once

displaced crayfish from shelter. In predation experiments, crayfish preyed upon all species, but preyed most heavily upon desert suckers. Fish never altered use of the water column in the presence of crayfish. Density manipulation experiments in a laboratory measured food competition between crayfish and two native fishes. Growth of Gila chub was less affected by crayfish than by increased density of Gila chub. Thus crayfish are not strong competitors with Gila chub for food. However, growth of flannelmouth sucker (*Catostomus latipinnis*) was negatively impacted by presence of crayfish. These laboratory experiments provide evidence that introduced crayfish can reduce fish growth by competition for food, and that native fishes are vulnerable to crayfish predation.”

## 4 Global Distribution

---



**Figure 1.** Global distribution of *O. virilis*. Map from GBIF (2015).

## 5 Distribution within the United States




Figure 2. Distribution of *O. virilis* in the US. Map from Benson (2015).


## 6 Climate Matching

### Summary of Climate Matching Analysis


The climate match (16 climate variables; Euclidean Distance) was high throughout the US, except for parts of the Deep South, isolated patches in the Interior West, and a narrow band along the Pacific Coast. Climate 6 match indicated that the continental US has a high climate match. The range for a high climate match is 0.103 and greater; climate match of *O. virilis* is 0.921.

Crayfishes have been observed to establish populations in climates different from that found within their native range (M. Hoff, U.S. Fish and Wildlife Service, personal communication). The climate match shown here may be an underestimate of climate suitability for the establishment of *O. virilis*.





**Figure 3.** RAMP (Sanders et al. 2014) source map showing weather stations selected as source locations (red) and non-source locations (gray) for *O. virilis* climate matching. Source locations from GBIF (2015). Only established locations were used.



**Figure 4.** Map of RAMP (Sanders et al. 2014) climate matches for *O. virilis* in the continental United States based on source locations reported by GBIF (2015). 0= Lowest match, 10=Highest match.

## 7 Certainty of Assessment

The biology and ecology of *O. virilis* are well-known. Negative impacts from introductions and spread of this species are adequately documented in the scientific literature. No further information is needed to evaluate the negative impacts the species is having where introduced. Certainty of this assessment is high.

## 8 Risk Assessment

### Summary of Risk to the Continental United States

*Orconectes virilis* is a crayfish native to the northern US that has been introduced to other parts of the contiguous US, Mexico, and Europe. Its introductions have resulted in declines of native macrophytes, macroinvertebrates, and fish. The species is known to carry and/or be susceptible to two OIE-reportable diseases. High climate match and history of invasiveness argues for a high risk of spread and impact of this species. Overall risk is high.

## **Assessment Elements**

- **History of Invasiveness: High**
- **Climate Match: High**
- **Certainty of Assessment: High**
- **Overall Risk Assessment Category: High**

## 9 References

---

**Note: The following references were accessed for this ERSS. References cited within quoted text but not accessed are included below in Section 10.**

- Aquatic Nuisance Species Information System (ANSIS). 2007. *Orconectes virilis*. US Army Corps of Engineers, Vicksburg, Mississippi. Available: [http://el.erdc.usace.army.mil/ansrp/ANSIS/html/orconectes\\_virilis\\_northern\\_crayfish.htm](http://el.erdc.usace.army.mil/ansrp/ANSIS/html/orconectes_virilis_northern_crayfish.htm). (June 2015).
- Benson, A. 2015. *Orconectes virilis*. USGS Nonindigenous Aquatic Species Database, Gainesville, Florida. Available: <http://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=215>. (June 2015).
- CABI. 2015. *Orconectes virilis* (virile crayfish) [original text by A. Ellis]. Invasive Species Compendium. CAB International, Wallingford, UK. Available: <http://www.cabi.org/isc/datasheet/72034>. (June 2015).
- Carpenter, J. 2000. Effects of introduced crayfish on selected native fishes of Arizona. Doctoral dissertation. University of Arizona, Tucson, Arizona.
- Davidson, E. W., J. Snyder, D. Lightner, G. Ruthig, J. Lucas, and J. Gilley. 2010. Exploration of potential microbial control agents for the invasive crayfish, *Orconectes virilis*. *Biocontrol Science and Technology* 20(3):297-310.
- Global Biodiversity Information Facility (GBIF). 2015. GBIF backbone taxonomy: *Orconectes virilis* (Hagen, 1870). Global Biodiversity Information Facility, Copenhagen. Available: <http://www.gbif.org/species/2227064>. (June 2015).
- Global Invasive Species Database (GISD). 2010. *Orconectes virilis* (crustacean). IUCN Invasive Species Specialist Group, Gland, Switzerland. Available from: <http://www.issg.org/database/species/ecology.asp?si=218> (June 2015).
- Integrated Taxonomic Information System (ITIS). 2015. *Orconectes virilis* (Hagen, 1870). Integrated Taxonomic Information System, Reston, Virginia. Available: [http://www.itis.gov/servlet/SingleRpt/SingleRpt?search\\_topic=TSN&search\\_value=97425](http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=97425). (June 2015).
- Loughman, Z. J., and T. P. Simon. 2011. Zoogeography, taxonomy, and conservation of West Virginia's Ohio River floodplain crayfishes (Decapoda, Cambaridae). *ZooKeys* 74:1-78.
- Martinez, P. J. 2012. Invasive crayfish in a high desert river: implications of concurrent invaders and climate change. *Aquatic Invasions* 7(2):219-234.
- Sanders, S., C. Castiglione, and M. Hoff. 2014. Risk Assessment Mapping Program: RAMP. US Fish and Wildlife Service.
-

## 10 References Quoted But Not Accessed

---

**Note: The following references are cited within quoted text within this ERSS, but were not accessed for its preparation. They are included here to provide the reader with more information.**

- Ahern, D., J. England, and A. Ellis. 2008. The virile crayfish, *Orconectes virilis* (Hagen, 1870) (Crustacea: Decapoda: Cambaridae), identified in the UK. *Aquatic Invasions* 3(1):102-104.
- Aiken, D. E. 1965. Distribution and ecology of three species of crayfish from New Hampshire. *American Midland Naturalist* 73:240-244.
- Bovbjerg, R. V. 1970. Ecological isolation and competition exclusion in two crayfish (*Orconectes virilis* and *Orconectes immunis*). *Ecology* 51:225-236.
- Bryan, S. D., A. T. Robinson, and M. G. Sweetser. 2002. Behavioral responses of a small native fish to multiple introduced predators. *Environmental Biology of Fishes* 63(1):49-56.
- Carpenter, J. 2005. Competition for food between an introduced crayfish and two fishes endemic to the Colorado River basin. *Environmental Biology of Fishes* 72(3):335-342.
- Chambers, P. A., J. M. Hanson, J. M. Burke, and E. E. Prepas. 1990. The impact of the crayfish *Orconectes virilis* on aquatic macrophytes. *Freshwater Biology* 24(1):81-91.
- Collicut, D. 1998. Biology of northern crayfish. *Nature North*. Available: <http://www.naturenorth.com/fall/crayfish/Fcray.html>. (July 2003).
- Collicut, undated [cited by GISD 2010, source did not provide full citation]
- Cooper, J. E., A. L. Braswell, and C. McGrath. 1998. Noteworthy distributional records for crayfishes (Decapoda: Cambaridae) in North Carolina. *Journal of the Elisha Mitchell Scientific Society* 114(1):1-22.
- DiStefano, R. J., M. E. Litvan, and P. T. Horner. 2009. The bait industry as a potential vector for alien crayfish introductions: problem recognition by fisheries agencies and a Missouri evaluation. *Fisheries* 34(12):586-597.
- Dorn, N. J., and G. G. Mittelbach. 2004. Effects of a native crayfish (*Orconectes virilis*) on the reproductive success and nesting behavior of sunfish (*Lepomis* spp.). *Canadian Journal of Fisheries and Aquatic Sciences* 61:2135-2143.
- Godfrey, L. 2002. UC pest management guidelines. University of California, Davis. Available: <http://www.ipm.ucdavis.edu/PMG/r682500211.html>. (July 2003).
- Hanson, J. M., P. A. Chambers, and E. E. Prepas. 1990. Selective foraging by the crayfish *Orconectes virilis* and its impact on macroinvertebrates. *Freshwater Biology* 24:69-80.

- Hill, A. M., and D. M. Lodge. 1994. Diel changes in resource demand: competition and predation in species replacement among crayfishes. *Ecology* 75:2118-2126.
- Horns, W. H., and J. J. Magnuson. 1981. Crayfish predation on lake trout eggs in Trout Lake, Wisconsin. *Rapports et procès-verbaux des réunions / Conseil permanent international pour l'exploration de la mer* 178:299-303.
- Hunner 1997 [cited by GISD 2010, source did not provide full citation]
- Johnson, J. E. 1986. Inventory of Utah USA crayfish with notes on current distribution. *Great Basin Naturalist* 46(4):625-631.
- Light, T., D. C. Erman, C. Myrick, and J. Clarke. 1995. Decline of the Shasta crayfish (*Pacifastacus fortis* Faxon) of northeastern California. *Conservation Biology* 9(6):1567-1577.
- Lodge, D. M., M. W. Kershner, J. E. Aloï, and A. P. Covich. 1994. Effects of an omnivorous crayfish (*Orconectes rusticus*) on a freshwater littoral food web. *Ecology* 75:1265-1281.
- Lodge, D. M., and J. G. Lorman. 1987. Reductions in submersed macrophyte biomass and species richness by the crayfish *Orconectes rusticus*. *Canadian Journal of Fisheries and Aquatic Sciences* 44:591-597.
- Lodge, D. M., C. A. Taylor, D. M. Holdich, and J. Skurdal. 2000. Nonindigenous crayfish threaten North American freshwater biodiversity. *Fisheries* 25:7-19.
- Mathews, L. M., L. Adams, E. Anderson, M. Basile, E. Gottardi, M. A. Buckholt. 2008. Genetic and morphological evidence for substantial hidden biodiversity in a freshwater crayfish species complex. *Molecular Phylogenetics & Evolution* 48(1):126-135.
- Momot, W. T. 1967. Population dynamics and productivity of the crayfish, *Orconectes virilis*, in a marl lake. *American Midland Naturalist* 78:55-81.
- Olden, J. D., E. R. Larson, and M. C. Mims. 2009. Home-field advantage: native signal crayfish (*Pacifastacus leniusculus*) out consume newly introduced crayfishes for invasive Chinese mystery snail (*Bellamya chinensis*). *Aquatic Ecology* 43(4):1073-1084.
- Peck, S. K. 1985. Effects of aggressive interaction on temperature selection by the crayfish *Orconectes virilis*. *American Midland Naturalist* 114:159-167.
- Pflieger, W. L. 1996. The crayfishes of Missouri. Missouri Department of Conservation, Jefferson City, Missouri.
- Riegel, J. A. 1959. The systematics and distribution of crayfishes in California. *California Fish and Game* 45(1):29-50.

- Rogowski, D. L., and C. A. Stockwell. 2006. Assessment of potential impacts of exotic species on populations of a threatened species, White Sands pupfish, *Cyprinodon tularosa*. *Biological Invasions* 8(1):79-87.
- Ruiz, G. M., P. W. Fofonoff, J. T. Carlton, M. J. Wonham, and A. H. Hines. 2000. Invasion of coastal marine communities in North America: apparent patterns, processes, and biases. *Annual Review of Ecology and Systematics* 31:481-531.
- Schwartz, F. J., R. Rubelmann, and J. Allison. 1963. Ecological population expansion of the introduced crayfish, *Orconectes virilis*. *The Ohio Journal of Science* 63(6):266-273.