


Biological Sciences Report
USGS BSR 2001-0008
OCS Study MMS 2002-034

Northeastern Gulf of Mexico Coastal and Marine Ecosystem Program

Community Structure and Trophic Ecology of Fishes on the Pinnacles Reef Tract



U.S. Department of the Interior
Minerals Management Service
Gulf of Mexico OCS Region



U.S. Department of the Interior
U.S. Geological Survey

Cover illustration credits:

High resolution multi-beam swath bathymetry map produced by Dr. James V. Gardner, USGS, Menlo Park, CA.

Fish photographs by D. C. Weaver and W. F. Smith-Vaniz.

Biological Sciences Report
USGS BSR 2001-0008
OCS Study MMS 2002-034

Northeastern Gulf of Mexico Coastal and Marine Ecosystem Program

Community Structure and Trophic Ecology of Fishes on the Pinnacles Reef Tract

Douglas C. Weaver, George D. Dennis, and Kenneth J. Sulak

June 2002

Florida Caribbean Science Center
U.S. Geological Survey
7920 NW 71st Street
Gainesville, Florida 32653-3071

MMS U.S. Department of the Interior
Minerals Management Service
Gulf of Mexico OCS Region



U.S. Department of the Interior
U.S. Geological Survey

PROJECT COOPERATION

This study was undertaken to meet information needs identified by the U.S. Geological Survey (USGS), Outer Continental Shelf Ecosystem Program in concert with the Minerals Management Service (MMS).

DISCLAIMER

This report was prepared by the Florida Caribbean Science Center of the USGS. This report has been technically reviewed by USGS and MMS, and has been approved for publication. Approval does not signify that the contents necessarily reflect the views and policies of the USGS, nor does mention of trade names or commercial products constitute endorsement or recommendation for future use. It is, however, exempt from review and compliance with the MMS editorial standards.

REPORT AVAILABILITY

Extra copies of this report may be obtained from the MMS Public Information Unit at the following address:

U.S. Department of the Interior
Minerals Management Service
Public Information Office (MS 5034)
1201 Elmwood Park Boulevard
New Orleans, LA 70123-2394

Telephone: (504) 736-2519 or
1-800-200-GULF

The following datasets are available as read-only files on the Florida Caribbean Science Center Website at <http://www.fcsc.usgs.gov/index.html>:

Located in: Pinnacles Tech Report/
Pinnacles Master Station Log.xls
Pinnacles Master Specimen Log.xls
USGS Technical Report 2001-0008.doc
USGS Technical Report 2001-0008.pdf

SUGGESTED CITATION

Weaver, D. C., G. D. Dennis, and K. J. Sulak. 2001. Northeastern Gulf of Mexico Coastal and Marine Ecosystem Program: Community Structure and Trophic Ecology of Demersal Fishes on the Pinnacles Reef Tract; Final Synthesis Report. U.S. Department of the Interior, Geological Survey, USGS BSR-2001-0008 and Minerals Management Service Gulf of Mexico OCS Region, New Orleans, LA, OCS Study MMS 2002-034, 92 pp. + apps.

PRINCIPAL INVESTIGATORS AND COLLABORATORS

Dr. Kenneth J. Sulak (Principal Investigator (PI), Project Lead Scientist) ¹
ken_sulak@usgs.gov

Douglas C. Weaver (Co-PI; Project Research Biologist) ^{1,2}
doug_weaver@usgs.gov

Dr. George D. Dennis (Co-PI; Project Research Biologist) ¹
george_dennis@usgs.gov

Collaborators:

Dr. Steve W. Ross ³

Dr. William F. Smith-Vaniz ¹

¹Florida Caribbean Science Center, U.S. Geological Survey, 7920 NW 71st Street, Gainesville, FL 32653-3071.

²Department of Fisheries and Aquatic Sciences, University of Florida, 7922 NW 71st Street, Gainesville, FL 32653-3071.

³North Carolina National Estuarine Research Reserve (NCNERR), Center for Marine Science Research, 1 Marvin Moss Lane, Wilmington, NC 28403.

CONTENTS

	Page
Figures	ix
Tables	xiii
Executive Summary	xv
Chapter 1: Community Structure of Pinnacles Reef Tract Fishes	
Background and Introduction	1
Objectives	6
Methods	22
Results	34
Discussion	62
Chapter 2: Trophic Ecology of Pinnacles Reef Tract Fishes	
Introduction	69
Objectives	70
Methods	70
Results and Discussion	71
Acknowledgments	83
Literature Cited	85
Appendix A. Summary of ROV surveys for 1997-2001.	A-1
Appendix B. A list of reef-fish species from natural reefs in the northern Gulf of Mexico with a comparison among the northwestern Gulf, Pinnacles, and West Florida shelf.	B-1
Appendix C. Food habits of reef fishes of the Pinnacles Reef Tract by trophic guild.	C-1
Appendix D. Characteristic ichthyofauna of Pinnacles Reef Tract, displaying the biodiversity of fishes by trophic guild.	D-1

LIST OF FIGURES

Figure	Page
1.1. Oblique view of the Alabama Alps study site and 36 Fathom Ridge area along the Pinnacles Reef Tract, showing the variability in outer shelf reef morphology (map provided by J. Gardner, USGS, Menlo Park, CA).	2
1.2. Northeastern Gulf of Mexico outer continental shelf with polygon marking the combined MMS MAMES and MASPTHMS study areas, enclosing the Pinnacles Reef Tract, including NEGOM-CMEP and USGS study sites (adapted from CSA and TAMU 2001).	4
1.3. Locations of NEGOM-CMEP megasites and monitoring sites (adapted from CSA and TAMU 2001).	5
1.4. USGS Pinnacles Reef Tract study sites in relation to NEGOM-CMEP megasites (adapted from CSA and TAMU 2001).	8
1.5. A. High-resolution multi-beam swath bathymetry for 40 Fathom Fishing Ground area with study sites. B. Acoustic backscatter for same area differentiating substrate type (provided by J. Gardner, USGS, Menlo Park, CA).	9
1.6. High-resolution multi-beam swath bathymetry for Roughtongue Reef study site (provided by J. Gardner, USGS, Menlo Park, CA).	10
1.7. High-resolution multi-beam swath bathymetry for Cat's Paw Reef study site (provided by J. Gardner, USGS, Menlo Park, CA).	12
1.8. High-resolution multi-beam swath bathymetry for Yellowtail Reef study site (provided by J. Gardner, USGS, Menlo Park, CA).	13
1.9. High-resolution multi-beam swath bathymetry for Near Shoreline Ridge area with study sites (provided by J. Gardner, USGS, Menlo Park, CA).	14
1.10. High-resolution multi-beam swath bathymetry for Double Top Reef study site (provided by J. Gardner, USGS, Menlo Park, CA).	15
1.11. High-resolution multi-beam swath bathymetry for 36 Fathom Ridge area with study sites (provided by J. Gardner, USGS, Menlo Park, CA).	17
1.12. High-resolution multi-beam swath bathymetry for Alabama Alps study site (provided by J. Gardner, USGS, Menlo Park, CA).	18

LIST OF FIGURES

(continued)

Figure	Page
1.13. High-Resolution multi-beam swath bathymetry for Ludwick and Walton Pinnacles area with study sites (provided by J. Gardner, USGS, Menlo Park, CA).	19
1.14. Distribution of sampling effort at Roughtongue Reef (NEGOM-CMEP Site 1) for cruises 97-01 through 2000-01.	27
1.15. Sabiki jigs used to collect small reef fishes during this study, including <i>Pronotoqrammus martinicensis</i> and <i>Hemanthias vivanus</i> .	28
1.16. A. Suction sampler device schematic. B. NURC-UNCW Phantom DS4 ROV with suction sampler. C. A typical collection of fishes made by the suction sampler.	30
1.17. Still frames taken from video surveys at the Pinnacles Reef Tract. A. <i>Pronotoqrammus martinicensis</i> school along crevice in reef top at Yellowtail Reef, 63 m. B. Close-up of mixed group of <i>P. martinicensis</i> and <i>Hemanthias vivanus</i> along reef face. C. Mixed school of anthiines along reef face at CMEP-Site 5, 75 m. D. Single <i>P. martinicensis</i> along reef face at Scamp Reef, 105 m.	40
1.18. Still frames taken from video surveys at the Pinnacles Reef Tract. A. Mixed school of <i>Hemanthias vivanus</i> and <i>Chromis enchrysur</i> with a single <i>Pristigenys alta</i> on reef top biotope at Yellowtail Reef, 62 m. B. <i>Holacanthus bermudensis</i> and <i>Chaetodon aya</i> along northern edge of reef top biotope at Yellowtail Reef, 63 m. C-D. Reef top biotope with well-developed sessile invertebrate community of gorgonians and crinoids, Yellowtail Reef, 62 m.	41
1.19. Still frames taken from video surveys at the Pinnacles Reef Tract. A. <i>Seriola dumerili</i> schooling above reef flat top, Yellowtail Reef, 62m. B. <i>Gonioplectrus hispanus</i> at reef base, CMEP-Site 5, 78 m. C. Female <i>Mycteroperca phenax</i> on reef face at Scamp Reef, 105 m. D. Dominant male <i>M. phenax</i> exhibiting grey-head color phase along reef face, Scamp Reef, 105 m.	42
1.20. Typical ichthyofauna contributing to community structure on the Pinnacles Reef Tract, northeastern Gulf of Mexico, based on ROV data.	49

LIST OF FIGURES
(continued)

Figure	Page
1.21. Pinnacles Shallow Reef Tract fish faunal composition for reef crest biotope.	50
1.22. Pinnacles Shallow Reef Tract fish faunal composition for reef base biotope.	51
1.23. Pinnacles Shallow Reef Tract fish faunal composition for talus zone biotope.	52
1.24. Cluster of reef sites using percent similarity and unweighted pair-group mean average combinatorial strategy.	54
1.25. Cluster analysis of taxa using percent similarity and unweighted pair-group mean average combinatorial strategy.	56
1.26. Constancy and fidelity for site and taxa groups defined by cluster analysis.	57
2.1. Qualitative food web model for reef fishes of the Pinnacles Reef Tract constructed from stomach content analysis.	77
2.2. Quantitative food web model for reef fishes of the Pinnacles Reef Tract constructed from stomach content analysis.	78

LIST OF TABLES

Table	Page
1.1. Summary of USGS Pinnacles Reef Tract study locations.	7
1.2. Summary of research activities conducted from 1997-2000.	24
1.3. Previous investigations of the ichthyofauna of the Pinnacles Reef Tract, comparing number of reef fishes observed by ROV in each study.	34
1.4. List of fishes documented in the Pinnacles Reef Tract from 1997 - 2000.	35
1.5. Abundance of top 30 taxa observed in ROV surveys partitioned by survey period (day/night) and reef tract (shallow/deep) for reef habitat.	44
1.6. Results of chi-square goodness of fit test for reef fish abundance by survey period (day/night) and reef tract (shallow/deep).	45
1.7. Day-night comparison of selected community parameters and top six taxa (abundance) paired by reef based on ROV data.	46
1.8. Biotope association of Pinnacles Reef Tract fishes based on ROV surveys during Cruise 97-01, combined for all study sites.	48
1.9. Summary statistics for ROV data by site group based on contributing to cluster analysis.	55
1.10. Abundance of top 15 predatory fishes caught by baited hook and line angling from 1997-2000.	59
1.11. Abundance of top 15 fish taxa caught in traps from 1997-2000.	60
1.12. Abundance of top 15 fish taxa caught in trawls in 1997.	61
2.1. Feeding guilds defined by characteristic prey type.	72
2.2. Dietary analysis of characteristic reef fishes of the Pinnacles Reef Tract by trophic guild.	73
2.3. Feeding guilds of deep reef fishes along the Pinnacles Reef Tract.	75

EXECUTIVE SUMMARY

Topographic prominences form reef-like features along the outer continental shelf (OCS) edge and continental slope of the southeastern United States and Gulf of Mexico. An extensive set of reef features, known as the Pinnacles Reef Tract, occurs off Mississippi and Alabama at depths between 60 and 110 m. In this area recent research has focused largely on distribution of reef structures, their geology and physical characteristics, and processes relating to their origin. Community ecology studies have been limited in scope, confined mainly to distribution of sessile megafaunal and incidental observations of fishes. This study's goal was to determine patterns of fish community structure and trophic relationships on Pinnacles reef and reef-associated biotopes. We examined variation in hard-bottom reef fish assemblages with reef topography and geographic location, to identify faunal differentiation associated with reef profile, biotopes, area, and depth. In addition, trophodynamic investigations were undertaken using food habits and food web structure of Pinnacles Reef Tract fishes. Analysis of food web structure was used to identify the relative importance of resident reef fishes (autochthonous) versus off-reef trophic resources (allochthonous). This study was a component of the overall Minerals Management Service (MMS) Northeastern Gulf of Mexico Coastal and Marine Ecosystem Program (NEGOM-CMEP), designed to compliment other synoptic programs. It focused on hard-bottom features that serve as essential fish habitat for reef fishes on the OCS in areas of oil and gas development.

Ten sites within the combined Mississippi-Alabama Marine Ecosystems Study and Mississippi-Alabama Shelf Pinnacle Trend Habitat Mapping Study (MAMES/MASPHTHMS) area were visited on the initial cruise. An additional eight study sites were visited on subsequent cruises. High Resolution Multi-Beam Swath bathymetric maps were completed in the Pinnacles Reef Tract to provide detailed maps of the geomorphology of study reefs. These high-resolution images have been used to guide subsequent sampling. The combined sampling effort for all cruises comprised 326 stations, apportioned into 112 angling, 63 trap, 22 bottom trawl, 58 remotely operated vehicle (ROV), 15 dredge/core/grab, and 37 plankton stations, with a small number of stations representing bottom surveys, CTD casts, light meter deployments, and exploratory sampling gear. All sampling combined returned over 6,000 specimens for food habits analyses, taxonomic verification and documentation, and subsequent life history analyses. Shipboard photographs documenting 113 fish species were obtained. ROV surveys resulted in about 85 hours of reef and reef-associated biotope observations and were used for analysis of resident fish fauna.

The results of this study have expanded the number of fishes documented from the Pinnacles Reef Tract, and is the first extensive collection of deep (greater than 60 m) reef fishes made in this region, and possibly worldwide. Of the 159 species documented from collected specimens or via videotape, 88 were identified from ROV censuses, and 70 can be classified as obligate reef-fishes belonging to the Caribbean reef-fish fauna. Such obligate reef fishes have life histories tied to three-dimensional hard-bottom habitats, and would not otherwise be found on the OCS open shelf. An additional 32 species are facultative reef associates that range broadly over OCS habitats, but occur on reef habitat,

when available. ROV surveys revealed the dominance of planktivorous reef fishes on high profile reef features. Anthiine fishes (small planktivorous seabasses) numerically dominate the deep reef fish community, and may serve as a keystone species by foraging in the water column and transferring energy to the deep reef ecosystem. The diurnal nature of the fauna is evident in that the total number of individuals was significantly lower at night especially on shallow reefs. While small diurnal planktivores are considered to be an important link between open water plankton and the reef, nocturnal planktivores may play a relatively important role in the overall energetics of the reef community. Based on ROV analysis of fish faunal composition, species richness and relative abundance of reef fishes differs among biotopes (reef top, reef crest, reef face, reef base, talus zone, and sand flats). Cluster analysis grouped reefs by depth trend and relief.

At least 250 reef-associated fishes have been documented from offshore banks, reefs, and other hard-bottom habitats in the northern Gulf of Mexico based on previous literature. The suite of primary reef fishes inhabiting the Pinnacles Reef Tract represents a depauperate, winnowed fraction of the Caribbean reef-fish fauna. It can be hypothesized that the known Pinnacles reef fish diversity is diminished relative to the Caribbean due to: 1) distance from the source fauna, 2) depth limitation, 3) temperature limitation, 4) habitat limitation due to turbidity, 5) trophic limitations, 6) fishing pressure, or 7) incomplete sampling. Environmental conditions per se do not appear to control reef fish distribution in the area. Based on the analysis presented here the lack of a full range of reef habitats, such as coral and coralline algae dominated communities, is the greatest influence on taxonomic structure. In addition, heavy fishing pressure may have substantially altered the reef fish community. Trophic structure can assist in evaluation of overfishing of large predators and the associated shift in the reef fish community. The numerical dominance by small and medium planktivorous reef fish taxa in shelf-edge reef communities in the Pinnacles Reef Tract has likely been magnified by removal of piscivores through recent increases in commercial fishing activity in this region.

Trophodynamic investigations undertaken in the present study include feeding habits and food web structure of Pinnacles Reef Tract fishes. Defining trophic pathways is the key to developing an understanding of the ecological framework in the reef fish community. Analysis of food web structure also identifies relative importance of resident reef fishes versus off-reef trophic resources such as adjacent soft-substrate and pelagic prey in the diet of large predatory fishes, predator-prey relationships among fishes and invertebrates, and may also identify potential pathways of energy transfer from fishes to the benthic invertebrate assemblage through the release of fecal material.

A food web model was developed for the Pinnacles fish community, based on numerical contribution of prey to the diet of each predator. Over 1000 individuals were examined for food habits, with approximately 50% containing prey items. Fish were assigned to feeding guilds based on stomach contents to identify similarity in prey resources among species, sources of prey, and overall patterns in trophic structure. The numerically dominant species (greater than 80% of the individuals observed by ROV) on high profile reefs, rough-tongue bass and red barbier, have diets dominated by calanoid copepods,

gastropod larvae, and a variety of other pelagic mesoplankton. The generalized carnivore guild was represented by numerous taxa including tattler and a variety of serranids, scorpaenids, muraenids, and holocentrids. This guild contains the most taxa of deep reef fishes, although individual species are not as numerically abundant as planktivores and benthic carnivores in reef communities.

Anthiine fishes may serve as a keystone predator by foraging in the water column and transferring energy to the deep reef ecosystem. Anthiines in turn become prey for a variety of reef predators. Dense populations of Pinnacles fishes appear to exceed levels that could be supported solely by locally produced prey resources. Fishes on deep reefs must therefore be “subsidized” by passive transport of prey from surrounding benthic or pelagic habitats on water currents, or via active feeding migrations of predators to adjacent areas of the water column or the underwater landscape. Large predators, including greater amberjack and red snapper, utilize prey from pelagic, reef, and soft-bottom habitats and provide pathways of energy transfer.

The numerical dominance of planktivorous reef fish taxa in shelf-edge reef communities and vermilion snapper in our hook and line samples in the Pinnacles Reef Tract have likely been magnified by removal of piscivores through recent increases in commercial fishing activity in this region and historical fishing practices throughout the northwestern Gulf of Mexico since the mid 1800’s. While our observations of trophic structure in the deep reef community closely resemble those observed in the Indo-Pacific, the reef fish community structure may have been greatly altered by human activity through fishing during the past century.

Fishes and invertebrates have been characterized at the Pinnacles Reef Tract by three major studies and all major reef features have been precisely mapped. This area may be the most well-known deep hard bottom area in the world. Thus we postulate that the majority of the reef fish species that occur at depths between 60 and 110 m are now documented. Yet due to the large area and diversity of habitat further scientific inquiry will continue to add to our knowledge of the fish fauna. Our present knowledge of the Pinnacles ichthyofauna can be used to characterize the fundamental taxonomic composition and trophic structure of deep reef-fish communities throughout southeastern United States and Gulf of Mexico OCS areas and therefore be useful in assessing future anthropogenic impacts on faunal structure.

CHAPTER 1

COMMUNITY STRUCTURE OF PINNACLES REEF TRACT FISHES

BACKGROUND AND INTRODUCTION

Topographic prominences form reef-like features along the continental shelf edge and continental slope of the southeastern United States and Gulf of Mexico. Such reefs have been known in the northern Gulf for over 70 years (Trowbridge 1930, Shepard 1937, Ludwick and Walton 1957, Moore and Bullis 1960). Shepard (1937) discussed the distribution of salt domes throughout the northwestern Gulf of Mexico based on fathometer surveys conducted by the U. S. Coast and Geodetic Survey in 1936, and Goedicke (1955) applied the term "pinnacles" to describe outer shelf features in the northern Gulf, including East and West Flower Garden Bank. Subsequently, Ludwick and Walton (1957) followed suit, using the term pinnacles to refer to steep-sided reef-like structures on the outer shelf in the northeastern Gulf. Historically, the term pinnacles included the entire class of reef structures rising from the open shelf, whether the top of the structure is spire-like, flat, or ragged, and regardless of the areal size or height of the overall structure (**Fig. 1.1**).

Shelf-edge prominences, including northeastern Gulf pinnacles, include structures of very heterogeneous origins (Gittings et al. 1992a, Gardner et al. 2001). West of the Mississippi, many shelf-edge banks are formed by the extrusion of salt domes (diapirs) from the surrounding seafloor, and are either capped with contemporary communities of reef-building organisms such as scleractinian corals, or are characterized by exposed carbonate rocks that are colonized by a diverse assemblage of sessile invertebrates (Rezak et al. 1985, Gittings et al. 1992a, Hardin et al. 2001). In the western Gulf off south Texas, shelf-edge prominences are non-diapiric drowned fossil coral-algal reefs (Rezak et al. 1985). East of the Mississippi, relatively few diapiric structures occur and are primarily located along the western portion of the Mississippi-Alabama shelf (Gittings et al. 1992a).

Most shelf-edge (60 to 110 m) features in this region appear to be non-diapiric, drowned, or fossil reefs initially formed during low sea level stands during the Pleistocene, and colonized by invertebrate communities [Continental Shelf Associates (CSA) 1992, Sager et al. 1992, Gittings et al. 1992a, Hancock 1997, Continental Shelf Associates and Texas A&M University (CSA & TAMU) 2001, Gardner et al. 2001]. Hard-substrate features supporting reef-like communities also occur on slide scar features that are exposed by slumping of the upper continental slope (Gardner et al. 2001) or on banks built by ahermatypic corals such as *Lophelia pertusa* (Linnaeus 1758) (Moore and Bullis 1960, Rogers 1999).

With few exceptions (e.g., Flower Garden Banks) outer continental shelf (OCS) reefs throughout the northern Gulf of Mexico lie in waters that are either too deep for sufficient light penetration, or seasonally too cold or too turbid to support hermatypic corals and




Figure 1.1. Oblique view of the Alabama Alps study site and 36 Fathom Ridge area along the Pinnacles Reef Tract, showing the variability in outer shelf reef morphology (map provided by J. Gardner, USGS, Menlo Park, CA). Vertical exaggeration = 10X. Isobaths in meters.

most forms of algae (Rezak et al. 1985). In the northeastern Gulf, OCS deep reefs (>60 m) are predominantly fossil structures that do not harbor hermatypic corals, extensive coralline or leafy algae communities (CSA 1992, Gittings et al. 1992a, Sager et al. 1992, Hancock 1997, CSA and TAMU 2001). There appears to be considerable production of biogenic debris by benthic organisms that colonize these reefs, as evidenced by coarse composition of sediment and high acoustic backscatter samples near features (Sager et al. 1992, Sager and Schroeder 2001, Gardner et al. 2002, Gardner et al. 2001). Moreover, many topographic features display evidence of extensive historical subaerial erosion (i.e., flattening of reef tops at lower sea level stages) and submarine sculpturing by scouring and bioerosion (resulting in the formation of basal undercuts, hollows and caves). Although the major coral reef builders are absent in pinnacle megafaunal communities, some biological deposition continues due to growth of ahermatypic corals, bryozoans, calcareous worms, and other calcium depositing and cementing organisms on the surface of these features (Sager et al. 1992, Hancock 1997, CSA and TAMU 2001).

Reef communities of the OCS have been comprehensively studied in the northwestern Gulf (Bright et al. 1984, Rezak et al. 1990, Gittings et al. 1992b, Gittings and Hickerson 1998). However, assessment of fish community structure has been primarily taxonomic and qualitative (Bright and Cashman 1974, Rezak et al. 1985), or semi-quantitative (Dennis and Bright 1988a). Long-term monitoring on the Flower Garden Banks has concentrated on corals, with limited attention to quantitative assessment of fish populations (Gittings et al. 1992b).


Early investigations on the distribution of OCS reefs in the northeastern Gulf began in the middle of the 20th century. Relative to the technology of the time, the pioneering research of Ludwick and Walton (1957) was remarkable in its scope and breadth, laying the foundation

for most subsequent geological and biological research on OCS deep reefs in the Gulf of Mexico. Using single-beam acoustic transects, Ludwick and Walton mapped the location of OCS pinnacles all along the northern Gulf from the Mississippi Delta to Port St. Joe, Florida. They produced the first bathymetric chart of the zone of pinnacles and reconstructed basal topography underlying the pinnacles. They measured physical and chemical seawater properties, mapped sediments and foraminiferal populations, and identified the major biological contributors to fossil reef formation from dredged reef rock. Ludwick and Walton also identified living sessile macrofauna and provided the first underwater photographs of pinnacles epifauna and bottom morphology.

Subsequent to Ludwick and Walton's work, major study programs of the Mississippi-Alabama Pinnacles Reef Tract (referred to hereafter as Pinnacles Reef Tract) have proceeded under the auspices of the Minerals Management Service (MMS) including the Mississippi-Alabama Marine Ecosystems Study (MAMES) (Laswell et al. 1990, Brooks 1991), Mississippi-Alabama Shelf Pinnacle Trend Habitat Mapping Study (MASPTHMS) (CSA 1992), and Northeastern Gulf of Mexico Coastal and Marine Ecosystem Program (NEGOM-CMEP) (CSA and TAMU 2001). These studies have conducted bathymetric and habitat reconnaissance, side-scan mapping, oceanographic, physical, and biological characterization and monitoring of pinnacle features and their hard-bottom communities. These investigations have provided the site locations, bathymetric charts, physical background, and macrofaunal framework for this study of pinnacle reef fish communities (**Figs. 1.2, 1.3**). Our community structure study complements work accomplished simultaneously under the NEGOM-CMEP program.

In the northeastern Gulf, recent research on OCS reefs has focused largely on distribution of reef structures, their geology and physical characteristics, and processes relating to their origin (Brooks 1991; CSA 1992, Sager et al. 1992, Sager and Schroeder 2001). Community ecology studies have been limited in scope, confined mainly to distribution of sessile megafaunal invertebrates (Gittings et al. 1992a, CSA and TAMU 2001), and incidental observations of fish communities (Shipp and Hopkins 1978; Gittings et al. 1991, Thompson et al. 1999). Snyder (2001) investigated some aspects of reef fish community structure (taxonomic diversity, habitat association) during the NEGOM-CMEP program.

The present study affords a more comprehensive evaluation of community structure, biotope affinities, and trophic ecology of the Pinnacles Reef Tract fish fauna. Here we examine variation in hard-bottom reef fish assemblages with reef topography and geographic location, to identify faunal differentiation associated with reef profile, biotopes, area, and depth. We also identify food habits and trophic structure of Pinnacles fishes (Chapter 2). The present study led to the initiation of important high-resolution multi-beam swath bathymetry of Pinnacles habitat/topography as is depicted in the study sites description.



4

Figure 1.2. Northeastern Gulf of Mexico outer continental shelf with polygon marking the combined MMS MAMES and MASPTHMS study areas, enclosing the Pinnacles Reef Tract, including NEGOM-CMEP and USGS study sites (adapted from CSA and TAMU 2001).




Figure 1.3. Locations of NEGOM-CMEP megasites (large numerals) and monitoring sites (small numerals) (adapted from CSA and TAMU 2001).

OBJECTIVES

A component of the overall NEGOM-CMEP Program, this study was designed to investigate aspects of fish community structure and compliment other synoptic programs. The study focused on hard-bottom features that serve as essential fish habitat for reef fishes on the OCS in areas of oil and gas development. The overall goal was to determine patterns of fish composition, community structure, and trophic relationships on Pinnacles reef and reef-associated biotopes. Project objectives were as follows:

- Using a broad range of sampling and video methods, qualitatively define taxonomic composition of fishes associated with steep-sided, reef-like, hard-bottom structures comprising the Pinnacles Reef Tract of the northeastern Gulf of Mexico.
- Quantitatively define species composition, abundance, dominance rank order, and species richness of fishes according to reef type, relief, and biotope. In addition, the fish fauna of the near reef zone and surrounding soft bottom area were also characterized.
- Define food habits and trophic structure of Pinnacles reef fishes (Chapter 2).

Study Sites

Bathymetric charts produced during the MAMES (Laswell et al. 1990, Brooks 1991), MASPTHMS (CSA 1992), and NEGOM-CMEP programs (CSA and TAMU 2001) were used to locate areas of interest during the initial USGS reconnaissance cruise in 1997. Ten potential sites within the combined MAMES/MASPTHMS polygon were visited on the initial cruise (**Table 1.1, Fig. 1.4**). An additional eight study sites were visited on subsequent cruises. During May/June 2000, High Resolution Multi-Beam Swath (HRMBS) bathymetric maps were completed in the Pinnacles Reef Tract to provide detailed maps of the geomorphology of study reefs (Gardner et al. 2000, Gardner et al. 2001, Gardner et al. 2002). High-resolution images resulting from that survey are included in this report, and were used to guide subsequent sampling.

Many of the sites for this study were selected within NEGOM-CMEP megasites to make use of data collected there by other investigators (**Figs. 1.3, 1.4**). Study reefs were selected from both the shallow reef trend (65-80 m) and deep reef trend (85-110 m). Eight main study reefs (five shallow, three deep) selected for fish community structure and trophodynamics study are described below:

1) **Roughtongue Reef**¹ - is a roughly elliptical (400 m major base diameter), high profile, flat-top structure with steep vertical sides (**Figs. 1.5A, 1.6**). The reef lies within NEGOM-CMEP Megasite 1 in the eastern part of the study area, in Destin Dome Lease Blocks 532 and 533, and corresponds to NEGOM-CMEP Site 1 (CSA and TAMU 2001) (**Fig. 1.3**). The general area containing this and the next two target reefs has historically

¹ names were designated by USGS for convenient reference and are not official geographic names

Table 1.1. Summary of USGS Pinnacles Reef Tract study locations. Geographic coordinates indicate general reef location.

Reef Name	Latitude (N) DD	Longitude (W) DD	Base Depth (m)	Crest Depth (m)	Max Relief (m)	Relief Category	Reef Area (ha)	Corresponding NEGOM-CMEP Location
Alabama Alps	29.2518	88.3373	88	72	16	High	20.0	Megasite 5, Site 7
Cat's Paw Reef	29.4396	87.5870	78	64	14	High	23.9	Megasite 1
Corkscrew Reef	29.4426	87.5445	78	66	12	High	13.8	none
Double Top Reef	29.3920	87.9830	80	68	12	High	3.0	Megasite 3
Far Tortuga	29.5571	87.4616	70	66	4	Intermediate	74.8	none
Ludwick and Walton Pinnacle 1	29.3268	87.7715	110	100	10	High	9.0	Megasite 4
L&W Pinnacle 2	29.3273	87.7680	110	100	10	High	7.0	Megasite 2, Site 4
L&W Pinnacle A	29.3428	87.7406	106	100	6	High	5.2	Megasite 2
L&W Pinnacle B	29.3386	87.7546	104	98	6	High	11.0	Megasite 2
L&W Pinnacle C	29.3400	87.7466	110	102	8	High	6.7	Megasite 2
L&W Pinnacle D	29.3350	87.7591	104	100	4	Intermediate	2.5	Megasite 2
L&W Pinnacle H	29.3265	87.7625	110	104	6	High	5.3	Megasite 2
NEGOM-CMEP Site 5	29.3938	87.9810	80	68	12	High	0.7	Megasite 3, Site 5
Patch Reef Field	29.4415	87.6915	71-77	69-75	3	Low	1000.0	none
Porgy Reef	29.4371	87.6180	78	68	10	High	19.4	Megasite 1
Roughtongue Reef	29.4415	87.5785	78	64	14	High	15.4	Megasite 1, Site 1
Scamp Reef	29.3250	87.7765	110	100	10	High	4.6	Megasite 2
Shark Reef	29.3965	88.0193	77	74	3	Low	1.2	Megasite 3
Solitary Mound 1	29.4558	87.6641	71	66	5	Intermediate	0.8	none
Solitary Mound 2	29.4643	87.6320	72	64	8	High	12.6	none
Triple Top Reef	29.3978	87.9915	76	68	8	High	3.6	Megasite 3
Yellowtail Reef	29.4403	87.5751	68	60	8	High	13.9	Megasite 1




Figure 1.4. USGS Pinnacles Reef Tract study sites in relation to NEGOM-CMEP megasites (open boxes) (adapted from CSA and TAMU 2001).




Figure 1.5. **A.** High-resolution multi-beam swath bathymetry for 40 Fathom Fishing Ground area with study sites. **B.** Acoustic backscatter for same area differentiating sediment type. Horizontal lines are due to sampling artifacts. Red indicates high backscatter and blue low backscatter (provided by J. Gardner, USGS, Menlo Park, CA). Isobaths in meters.




Figure 1.6. High-resolution multi-beam swath bathymetry for Roughtongue Reef study site (provided by J. Gardner, USGS, Menlo Park, CA). Vertical exaggeration = 9X. Transect line drawn from point A to B depicts reef in cross-section. Isobaths in meters.

been called the 40 Fathom Fishing Ground by fishermen, and in the MAMES program this area is referred to as the Eastern Delta Mounds (CSA and TAMU 2001). Roughtongue Reef belongs to the shallow pinnacle trend, with a base depth of 80 m (**Fig. 1.6**). Extensive areas of accumulated sediments occur on the reef flat top and a diverse, dense invertebrate assemblage occurs throughout the interior (Hardin et al. 2001). Vertical rock faces, boulders, and rock outcrops occurring along the sides and surrounding the feature display eroded surfaces, and typically have lower invertebrate densities, dominated by the solitary coral *Rhizopsammia manuelensis* Chevalier 1966 and other small ahermatypic species (Hardin et al. 2001). Attached to the main reef is a smaller mound immediately to the south. The reef is surrounded by an extensive area of high acoustic backscatter resulting from coarse reef-derived sediment or hard substrates (**Fig. 1.5B**, red area). The USGS designated name refers to the common name for the small planktivorous serranid, *Pronotogrammus martinicensis*, the roughtongue bass, which is extremely abundant on this reef.

2) **Cat's Paw Reef** is a group of six small, medium to high profile, flat-topped mounds, arranged in the pattern of a "cat's paw" print, with 5-10 m relief (**Figs. 1.5, 1.7**). This cluster of mounds lies about 1000 m west of Roughtongue Reef in the 40 Fathom Fishing Ground, NEGOM-CMEP Megasite 1 area (**Figs. 1.4, 1.5, 1.7**) and is within Destin Dome Lease Block 532. Individual reef formations within the feature have flat top communities present with limited sediment cover, highly eroded and sculpted rock surfaces with vertical faces along edges of features. Small soft corals are abundant on horizontal surfaces, solitary coral colonies (including *R. manuelensis*), with spiral sea whips, antipatharians, and crinoids also common. Coarse sand substratum was present around reef features, and silt was not evident on sediment surfaces.

3) **Yellowtail Reef**, a single, elliptical (200 m base diameter), high-profile, flat-top structure, that reaches the shallowest crest depth (60 m) of all study sites (**Table 1.1, Fig. 1.8**). This structure also belongs to the 40 Fathom Fishing Ground group, and lies within NEGOM-CMEP Megasite 1 (**Fig. 1.5A**) within Destin Dome Lease Block 532. It forms the northwestern end of a reef arc, with Cat's Paw Reef at the center, and Roughtongue Reef lying at the southeastern end. Like other reef features in the group, an extensive flat top area is present, characterized by accumulated sediments and a dense invertebrate assemblage dominated by octocorals, antipatharians, sponges, and coralline algae. Rock outcrops characterize the northern extent of the feature, and these areas are heavily colonized by sessile invertebrates and coralline algae. The USGS designated name refers to the yellowtail reeffish, *Chromis enchrysurus*, which is particularly abundant on this reef.

4) **Double Top Reef** is a horseshoe shaped (100 m base diameter), high profile structure that consists of multiple flat-top mounds with steep vertical sides (**Figs. 1.9, 1.10**). This structure lies within NEGOM-CMEP Megasite 3 (**Figs. 1.3, 1.4**) in the central part of the study area, in Main Pass Lease Block 223. A secondary mound attached to the northeast corresponds to NEGOM-CMEP Site 5 (**Fig. 1.9**). In the MAMES program this general area is referred to as the Near Shoreline Ridge (Brooks 1991). Double Top Reef belongs to the shallow pinnacle trend in the northeastern Gulf, and also includes a similarly-shaped series of mounds in the study area referred to as Triple Top Reef, and an adjacent, low profile




Figure 1.7. High-resolution multi-beam swath bathymetry for Cat's Paw Reef study site (provided by J. Gardner, USGS, Menlo Park, CA). Vertical exaggeration = 9X. Transect line drawn from point A to B depicts reef in cross-section. Isobaths in meters.




Figure 1.8. High-resolution multi-beam swath bathymetry for Yellowtail Reef study site (provided by J. Gardner, USGS, Menlo Park, CA). Vertical exaggeration = 9X. Transect line drawn from point A to B depicts reef in cross-section. Isobaths in meters.




Figure 1.9. High-resolution multi-beam swath bathymetry for Near Shoreline Ridge area with study sites (provided by J. Gardner, USGS, Menlo Park, CA).




Figure 1.10. High-resolution multi-beam swath bathymetry for Double Top Reef study site (provided by J. Gardner, USGS, Menlo Park, CA). Vertical exaggeration = 9X. Transect line drawn from point A to B depicts reef in cross-section. Isobaths in meters.

feature referred to as Pancake Reef that corresponds to NEGOM-CMEP Site 6 (**Fig. 1.9**). These features also have flat top communities characterized by high sediment cover and dense invertebrate assemblages dominated by octocorals and antipatharians, with few solitary corals. Vertical rock walls and overhangs are dominated by *R. manuelensis* and other solitary corals, and are highly sculpted and eroded. The USGS designated names refer to the topography of these reef structures. Double Top Reef and Triple Top Reef are not NEGOM-CMEP study sites.

5) **Alabama Alps** is a long, narrow, north-south aligned, high profile mound approximately 1000 m in length (**Figs. 1.11, 1.12**). This structure lies within NEGOM-CMEP Megasite 5 in the western part of the study area, in Main Pass Lease Block 286, and corresponds in part to NEGOM-CMEP Site 7. In the MASPTHMS program this same area is referred to as Lagniappe Delta Shallow, and has historically been called the 36 Fathom Ridge by fisherman. Alabama Alps forms the northwestern terminus of a long northwest to southeast-aligned ridge and pinnacle arc paralleling the shelf edge (**Fig. 1.1**), and belongs to the shallow pinnacle trend of the northeastern Gulf. The top of this feature has sections of relatively flat terrain with scattered sections of sediment cover, particularly in the southern portion of the feature (**Figs. 1.11, 1.12**). Invertebrate assemblages on the flat sections are dominated by octocorals, antipatharians, and sponges (Hardin et al. 2001). The sides of the feature range from vertical walls to large attached monoliths where the solitary coral *R. manuelensis* is the dominant sessile invertebrate with crinoids, antipatharians, coralline algae, sponges, and other solitary corals present (Hardin et al. 2001). The USGS designated name refers to the precipitous terrain, particularly the near-vertical west-face scarp of the structure, and its position off the state of Alabama.

6) **Ludwick and Walton Pinnacle 1** is the central member of a group of five medium to high-profile, spire-top, shelf-edge structures with 10 m maximum relief and a base depth of 110 m. This group belongs to the deep shelf-edge pinnacle trend in the northeastern Gulf (**Fig. 1.13**). These pinnacles form a short east-west aligned arc on the shelf-slope break, bordering the northern edge of a massive shelf-edge slump feature. Pinnacle 1 is one of those profiled and contoured by Ludwick and Walton (1957). It lies within the MAMES Western Delta Mounds area, NEGOM-CMEP Megasite 2 and within Destin Dome Lease Block 661. A fairly uniform coverage of coarse sand, rock fragments, shell fragments, and other calcareous debris surrounds the base of feature with occasional small rocky reef outcrops and patch reefs encrusted with *R. manuelensis*, octocorals, antipatharians, and crinoids. Emergent rocky features with vertical walls, rock ridges, and rock arches are distributed across the reef. Vertical rock faces have highly eroded surfaces, and are densely covered with *R. manuelensis*, with low coverage of other solitary corals, octocorals, sponges, and antipatharians.

7) **Ludwick and Walton Pinnacle 2** is another of the deep shelf-edge pinnacle group. This structure, lying immediately to the east of Pinnacle 1, also was profiled and contoured by Ludwick and Walton (1957). It also lies within the MAMES Western Delta Mounds area, NEGOM-CMEP Megasite 2, and Destin Dome Lease Block 661. The




Figure 1.11. High-resolution multi-beam swath bathymetry for 36 Fathom Ridge area with study sites (provided by J. Gardner, USGS, Menlo Park, CA). Isobaths in meters.




Figure 1.12. High-resolution multi-beam swath bathymetry for Alabama Alps study site (provided by J. Gardner, USGS, Menlo Park, CA). Vertical exaggeration = 9X. Transect line drawn from point A to B depicts reef in cross-section. Isobaths in meters.




Figure 1.13. High-resolution multi-beam swath bathymetry for Ludwick and Walton Pinnacles area with study sites (provided by J. Gardner, USGS, Menlo Park, CA). Vertical exaggeration = 10X. Distance from east to west = 6 km. Isobaths in meters.

feature is a broad mound that gradually slopes from the center, and is characterized by low relief hardbottom interspersed with a sediment veneer (Hardin et al. 2001). Scattered across the surface are individual rocky features reaching heights of 3 m. The elevated rocky features were colonized by dense populations of *R. manuelensis*, other solitary corals, octocorals, crinoids, and basket stars, while low relief hard bottom regions were characterized primarily by octocorals, antipatharians, and crinoids (Hardin et al. 2001). This pinnacle corresponds to NEGOM-CMEP Site 4.

8) Scamp Reef is a member of the Ludwick and Walton Pinnacles deep shelf-edge group with a precipitous southern reef face (**Fig. 1.13**). This structure, lying immediately to the west of Pinnacle 1, also was profiled and contoured by Ludwick and Walton (1957). It lies within the MAMES Western Delta Mounds area, NEGOM-CMEP Megasite 2 (**Fig. 1.4**), and within Viosca Knoll Lease Block 654. This feature has extensive vertical rock outcrops with profiles in excess of 5 m. Spectacular arches, overhangs and rugged topography occur along the southern face of the reef, with exposed rock colonized by *R. manuelensis*, antipatharians, crinoids, octocorals, and ahermatypic coral colonies. The USGS name Scamp Reef refers to the abundance of the scamp grouper, *Mycterperca phenax*, that reside at this site.

9) Other Study Sites. Additional study sites were visited during single occasions on reconnaissance cruise 97-01 or subsequent research cruises, but were not intensively sampled during this project (**Fig. 1.4**). The majority of these sites are located in the eastern part of the study area, and are within or adjacent to CMEP Megasite 1.

Porgy Reef is a broad, gradually sloping mound, approximately 500 m in diameter, located west of Roughtongue Reef, and within the boundary of Megasite 1. Silty sand substrates surround the reef, and the interior of the feature is composed of sculpted rock outcrops. No visible sediments were accumulated on the reef interior, and a flat top invertebrate community was absent. Invertebrate assemblages were dominated by *R. manuelensis*, corkscrew sea whips, antipatharians, octocorals, and crinoids on exposed rock surfaces.

Shark Reef is an elongate, east to west oriented, low profile reef feature approximately 250 m in length, located within CMEP Megasite 3, west of Double Top Reef and Triple Top Reef study sites. A well-developed invertebrate community was present on vertical walls and overhangs with small brown solitary corals and *R. manuelensis* abundant. Horizontal surfaces were heavily silted, and many sessile invertebrates appeared to be dead, with basal parts remaining but few living colonies visible. Pencil urchins were common on silted reef surfaces.

Solitary Mound 1 and **Solitary Mound 2** are located to the northwest of Megasite 1. Solitary Mound 1 is a gently sloping mound of intermediate height less than 100 m in diameter. A coarse sand substratum surrounds the reef features, and rock surfaces are extremely eroded/sculpted. Large rocky caverns and depressions occur on the sides of the reef with *R. manuelensis*, stony corals, with numerous corkscrew sea whips and crinoids on the reef crest, and crinoids and soft corals common on vertical rock surfaces. Rocky outcrops around the base of the feature are colonized by *R. manuelensis*, octocorals, and

long-spined urchins. Solitary Mound 2 is part of an intermittent ridge, and is an elongate feature approximately 800 m in length. This feature has a well-developed reef flat top with extensive sand and silt cover, where large red soft corals and other octocorals were abundant. Small rocky outcrops were present along the interior of the feature, along with numerous pits in the accumulated sediments.

Patch Reef Field is located to the east of Megasite 1 and is described in detail by Gittings et al. (1991). It is an area with numerous, low profile hard bottom outcroppings. Reef features in this area are only 2 to 3 m in maximum profile. Sessile invertebrates are relatively common on rock surfaces. At higher elevations, the reef community resembles larger features, with abundant solitary corals, octocorals, antipatharians, and sponges. Eroded, sculpted reef faces are encrusted with *R. manuelensis* and other solitary corals on overhangs, and corkscrew sea whips and antipatharians are common along the top and sides of rock outcroppings. As noted by Gittings et al. (1991), a thin layer of silt was present on reef surfaces, and many areas of the reef base, reef face, and small patch reefs were heavily silted. Flocculent silt was also present in the near-bottom water column. Most of the footage on this dive was over open sand bottom, with many small individual patch reefs encountered during the dive, as also observed by Gittings et al. (1991).

Far Tortuga Reef is the site farthest east in this study and is a broad, gradually sloping reef of intermediate profile, approximately 800 m in diameter. Scattered rock outcrops on the feature have typical invertebrate assemblages, with small octocorals and antipatharians abundant on rock surfaces, and *R. manuelensis*, crinoids and sponges also common. The reef feature was composed of relatively small, low profile outcrops interspersed with sandy patches, with a silt veneer over coarse sand around the reef outcrops. High profile formations or a flat-top community were not encountered.

Corkscrew Reef is a series of three high profile, gradually sloping mounds, each approximately 300 m in diameter that are located directly to the east of Megasite 1 (**Table 1.1**). This site is characterized by coarse sand with low silt surrounding the reef, and coarse sand on the reef. Reef surfaces were highly eroded, sculpted rock faces with *R. manuelensis* and large solitary corals on surfaces. Patchy rock outcrops with low invertebrate cover and abundant corkscrew sea whips and antipatharians on the top of reef features. No well-developed flat top community was visible at this site, with the rocky crests of reef features harbored dense assemblages of corkscrew sea whips and other antipatharians.

Ludwick and Walton Pinnacles A-D, H includes a number of additional sites distributed throughout the Ludwick and Walton deep shelf-edge pinnacles tract, ranging from 200 to 1000 m in diameter (**Fig. 1.13, Table 1.1**). Reefs along the Ludwick and Walton Pinnacles tract are surrounded by sand and silt substrate at the base of the feature, with elevated highly sculpted and eroded rock surfaces and extensive caves and depressions. Dominant invertebrates on the deep pinnacles are *R. manuelensis*, other solitary corals, octocorals, antipatharians, and crinoids. Invertebrate density is highest on vertical surfaces and rocky crests. Flat top reef communities with accumulated sediments and dense invertebrate assemblages characteristic of high profile shallow reefs were not observed at any of these sites.

METHODS

Cruise Activities

The initial reconnaissance and sampling cruise USGS 97-01 was conducted on the R/V SUNCOASTER 4-13 August 1997 (92 stations). The remaining six USGS-funded cruises (234 stations) were completed on the R/V TOMMY MUNRO (**Table 1.2**). The combined sampling effort for all cruises comprised 326 stations, apportioned into 112 angling, 63 trap, 22 bottom trawl, 58 remotely operated vehicle (ROV), 15 dredge/core/grab, and 37 plankton stations, with a small number of stations representing bottom surveys, CTD casts, light meter deployments, and exploratory sampling gear (**Table 1.2**). All sampling combined returned over 6,000 specimens for food habits analyses, taxonomic verification and documentation, and subsequent life history analyses. Shipboard photographs documenting 113 species were obtained. ROV tapes resulted in about 85 hours of observation of reef and reef-associated biotopes and the resident fish fauna for analysis.

Sampling was conducted during all seven research cruises spanning the time interval of 1997-2000 (**Table 1.2**). Positions for all sampling sites were determined by Differential Global Positioning System (DGPS) and charts were provided by CSA-TAMU collaborators. Sampling conducted during the 2000-01 cruise was further georeferenced using digital maps of HRMBS topography and vessel position tracked in ArcView software to display station location (**Fig. 1.14**). Operations were conducted continuously 24 hours per day.

Hook and Line Angling

During all cruises, demersal and pelagic fishes from hard-bottom reefs were obtained using hook and line angling with bait and large hooks (for macro-carnivores) and angling with unbaited, multi-hook “Sabiki” jigs (for planktivores and micro-carnivores) (**Fig. 1.15**). Sabiki jigs are constructed of a variety of small hook sizes (5 to 20 mm in length) with added strips of iridescent film to imitate shrimps or other small planktonic organisms, and each pre-packaged unit contains from 4 to 7 hooks. Sabiki jigs were deployed using 142 to 340 g lead sinkers, depending upon water depth and currents, and were successfully used to collect fishes to 110 m depth. Fishes were primarily collected during daylight hours, as the majority of individuals captured using this technique were visual planktivores.

Traps

A variety of sizes and mesh traps including large (1.8 m X 1.5 m X 0.6 m) chevron traps with 25 mm X 12 mm mesh to box traps (1.2 m X 1.2 m X 0.6 m) of 25 mm hexagonal wire were used to obtain fish specimens to confirm visual identification and sample cryptic taxa not easily observed by the ROV. Some traps were equipped with 3 or 6 mm Vexar mesh liner to retain small specimens. Traps were deployed in trap lines made up of two to four traps. Also attached to these lines were smaller collection devices such as minnow traps, PVC bundles, and bottles in an attempt to attract and capture small shelter seeking fishes.

Trawls

A 4.9 m semi-balloon trawl with a 3.8 cm mesh body and a 0.6 cm mesh liner was used to collect bottom fishes near reefs at 13 stations and a 7.6 m semi-balloon trawl with a 3.8 cm mesh body and a 0.6 cm mesh liner at five stations. Both trawls were equipped with a “Texas” roller rig made up of 7.6 cm rubber disks on a chain in front of the head rope to reduce hangs on reef structures.

Specimen Disposition

In-situ collections provided voucher specimens to verify and document species identifications, and material for food habits studies. On board the research vessel, individual voucher fish specimens were carefully preserved, pinned out to display coloration and diagnostic features, and photographed using digital and 35 mm cameras on a water bath light table (Randall 1961). Photographs were taken by D. Weaver, W. Smith-Vaniz, and J. Caruso. Specimens were preserved in 10% buffered formalin or frozen in the field, and later transferred to 70% ethanol in the laboratory. Selected voucher specimens were catalogued in the National Museum of Natural History (NMNH) and Florida Museum of Natural History (FMNH) ichthyology collections. The remaining voucher material, metadata, and voucher photographs are maintained at the Florida Caribbean Science Center (FCSC).

Other gears

Additional shipboard activities included plankton sampling to document pelagic prey availability to the deep reef fish community, and collection of physicochemical parameters of the water column. These data will be reported elsewhere.

ROV Methodology

Collection of specimens was complemented by use of ROVs to conduct habitat reconnaissance and document faunal composition via color video camera. All dives were recorded to videotape for subsequent laboratory analyses. A Phantom DS4 ROV, equipped with a color video camera, was the primary system used for in-situ observations. Three different Phantom ROVs were employed during the seven Pinnacles research cruises. United States Navy (97-01) and NMFS (99-03) ROVs were operated by NMFS Mississippi Laboratories during the first three years of the program, and the NOAA/NURP Phantom DS4 was operated by NURC-UNCW on the remaining cruises. Due to technical difficulties and image quality, only ROV data from cruises 97-01, 99-03, and 2000-01 were used for quantitative purposes (Appendix A). ROV observations were used to define and compare fish faunal composition at each main target study reef by depth, reef height (high, medium and low profile), and biotope.

Table 1.2. Summary of research activities conducted from 1997-2000. Number of stations listed by study site and sampling gear.

Cruise No. Date	Sampling Gear	Main Study Sites					Other Study Sites												
		Alabama Alps	Cat's Paw Reef	Double Top Reef	Rough Tongue Reef	Triple Top Reef	Yellowtail Reef	Patch Reef Field	Corkscrew Reef	CMEP Site 5	Far Tortuga Reef	L & W Pinnacle 1	L & W Pinnacle 2	L & W Pinnacles A-H	Scamp Reef	Porgy Reef	Shark Reef	Solitary Mound 1	Solitary Mound 2
97-01 4-13 Aug 1997	ROV Day		1	2	4	1	5		1	1							2	1	
	ROV Night				1		1								1				1
	Angling AM				2		4												
	Angling PM			2	3		1		1		1				2			1	
	Angling Night				2	1	1								1	1			1
	Plankton																		
	Traps				2	1	1		1								1		
	Trawls			2		1	1		1										2
	CTD			1	2	1	1				1								
98-02 8-10 Oct 1998	ROV Day																		
	ROV Night																		
	Angling AM			1	4				1										
	Angling PM			1	1		2												
	Angling Night				1		1												
	Plankton																		
	Traps				3	3													
	Trawl																		
CTD																			
99-01 18-23 Feb 1999	ROV Day																		
	ROV Night																		
	Angling AM	3	2		2														
	Angling PM	1	1		1		1												
	Angling Night				1														
	Plankton	5	1		1														
	Traps	4	1		1														
Trawl																			
CTD																			

Table 1.2. (continued).

Cruise No.	Date	Sampling Gear	Main Study Sites						Other Study Sites											
			Alabama Alps	Cat's Paw Reef	Double Top Reef	Roughtongue Reef	Triple Top Reef	Yellowtail Reef	Patch Reef Field	Corkscrew Reef	CMEP- Site 5	Far Tortuga Reef	L & W Pinnacle 1	L & W Pinnacle 2	L & W Pinnacles A-H	Seamp Reef	Porgy Reef	Shark Reef	Solitary Mound 1	Solitary Mound 2
99-02 30 May- 4 June 1999	ROV Day				1		1													
	ROV Night																			
	Angling AM	2	2	2			2									2				
	Angling PM	1	3	2	1		1									1		1		
	Angling Night		1		1						1									
	Plankton	2	4	2	2		4									1				
	Traps	4	4		6	4						1								
	Trawl																			
CTD																				
99-03 20-22 Aug 1999	ROV Day	2			1		2						2							
	ROV Night	1									1									
	Angling AM	1					1													
	Angling PM	3																		
	Angling Night	1																		
	Plankton	2					1													
	Traps		1																	
	Trawl				1															
CTD	1			1		1					1									

Table 1.2. (continued).

Cruise No. Date	Sampling Gear	Main Study Sites							Other Study Sites										
		Alabama Alps	Cat's Paw Reef	Double Top Reef	Roughtongue Reef	Triple Top Reef	Yellowtail Reef	Patch Reef Field	Corkscrew Reef	CMEP- Site 5	Far Tortuga Reef	L & W Pinnacle 1	L & W Pinnacle 2	L & W Pinnacles A-H	Seamp Reef	Porgy Reef	Shark Reef	Solitary Mound 1	Solitary Mound 2
2000-01 5-10 Mar 2000	ROV Day	2					3			1				1	1				
	ROV Night	1												1	1				
	Angling AM	1			3		3		1					1					
	Angling PM	2			1		3							1	1				
	Angling Night				1		1												
	Plankton	2			1		1												
	Traps	6			2		7						1	5					
	Trawl	4																	
CTD																			
2000-02 5-6 Sept 2000	ROV Day				1		2												
	ROV Night																		
	Angling AM				1		1												
	Angling PM						1												
	Angling Night																		
	Plankton				1		1												
	Traps																		
	Trawl																		
CTD																			




Figure 1.14. Distribution of sampling effort at Roughtongue Reef (NEGOM-CMEP Site 1) for cruises 97-01 through 2000-01. Each symbol represents the position of the research vessel during sampling gear deployment. Gray circles-angling, black triangles-traps, open squares-ROV dives.



Figure 1.15. Sabiki jigs used to collect small reef fishes during this study, including *Pronotoqrammus martinicensis* (top and middle) and *Hemanthias vivanus* (bottom).

Transecting methodology were drawn from those previously employed in ROV/submersible/camera sled studies that have shown that videotape analysis of large mobile demersal fishes requires a moving image and a low oblique perspective to capture diagnostic features of morphology and coloration (Grassle et al. 1975, Cohen and Pawson 1977, Uzmann et al. 1977, Rice et al. 1979, Parker and Ross 1986, Hecker 1987, Butler et al. 1991, Sulak and Ross 1993, 1996, Adams et al. 1995, Felley and Vecchione 1995). ROV transects were conducted at a constant speed of 0.10-0.15 m/sec across the bottom, with altitude of the ROV held to a maximum of 1.0 m (distance between sea floor and video camera lens, varying somewhat according to terrain). Horizontal swath width was resolved as the lower border of the video frame with the video camera zoom set to full wide-angle, and pan and tilt angles set to about 25° and 20°, respectively. Video was recorded on board the ship on S-VHS or Hi-8 videotape with audio annotation.

In situ collections of small reef fishes were accomplished using a suction sampler fitted to the ROV (**Fig. 1.16**) that dispensed a suspended rotenone solution (co-designed by D. Weaver and L. Horn, UNC-Wilmington). Powdered rotenone (Prentiss, Inc.) was suspended in 3 L of concentrated Ivory Liquid dishwashing detergent, and mixed with 8 L of seawater. Approximately 3 kg of standard table salt and 10 kg of granular sugar was added to increase the specific gravity of the solution to help maintain contact with the bottom. The suspended rotenone was passed through a standard kitchen strainer to remove large clumps, and loaded into the dual reservoir chambers of the sampler. A standard marine bilge pump, encased in an aluminum housing and powered from the surface, was used to dispense the rotenone solution and collect fishes at select reef sites during ROV dives. Fishes were retained in a 5 mm mesh liner fitted inside the first PVC chamber (**Fig. 1.16**).

Videotape Analysis

Videotapes were transferred to S-VHS format and analyzed on a commercial quality S-VHS VCR with variable speed editor/jogger control (slow-motion, frame by frame advance) to facilitate species identification at slow speed advance and enumerate individuals. Videotape analysis was used to quantify fish abundance by taxa and biotope. Incidental observations of pelagic fishes also were recorded and individuals identified. In addition to moving videotape transects, the zoom function of the NURC-UNCW ROV video camera was periodically used to capture close-up images of individuals to facilitate identification during cruises 2000-01 and 2000-02. These still video segments were used to verify species identifications made during transect surveys. All fish species were identified to the lowest possible taxonomic unit, with confirmation from close-up video images and by reference to voucher specimens collected by angling, trapping, or ROV suction sampler.

Habitat Parameters

In addition to defining fish community structure, videotape screening was used to make qualitative notes on the physical habitat and megafaunal invertebrates associated with particular biotopes and fish assemblages. Where appropriate (e.g., high profile, flat

A





Figure 1.16. A. Suction sampler device schematic. B. NURC-UNCW Phantom DS4 ROV with suction sampler. C. A typical collection of fishes made by the suction sampler.

topped features) fish observed were assigned to one of six biotopes: reef top, reef crest, reef base, reef talus (circum-reef sediment apron), and soft bottom.

Reef Top is referred to as flat reef tops or reef flats by Gittings et al. (1992a) and describes the biotope that develops on large, flat-topped mounds such as Roughtongue Reef, Yellowtail Reef, and CMEP-Site 5. This biotope is identified by extensive pockets of accumulated sediments and rocky debris, interspersed with low profile rock outcrops, ridges, and crevices. Invertebrate assemblages on the reef flat top community exhibit high density and species richness, and are dominated by erect sponges, octocorals (particularly sea fans, including *Nicella* sp.), comatulid crinoids, antipatharians, gorgonocephalid basket stars, bryozoans, and coralline algae (Gittings et al. 1992a, Hardin et al. 2001). Our reef top category corresponds to top interior locations of Hardin et al. (2001) and continuous hard bottom category of Snyder (2001).

Reef Crest biotopes were defined as the interface between the near vertical, rocky reef face and the sediment covered flat top characterized by dense assemblages of invertebrates. Reef crest biotopes typically were characterized by extensive rocky outcrops with small areas of sediment cover and low invertebrate densities. We distinguished the reef crest ecotone from the adjacent flat reef top and vertical reef face biotopes to identify the possible influence of position in the leading edge of water currents on the reef fish community. Our reef crest category corresponds to the top edge location listed by Hardin et al. (2001).

Reef Face as referred to by Gittings et al. (1992a), are rugged, often near-vertical rocky surfaces that characterize the steep-sided edges of large flat-topped features and the steep walls of narrow, spire shaped deep water pinnacles. Reef face communities are characterized by a lower overall density of epifauna than reef tops, with an abundance of ahermatypic corals, including *R. manuelensis*, *Madrepora* sp., and *Madracis/Oculina* sp., comatulid crinoids, a variety of octocoral fans, antipatharians including the spiral whip *Stichopathes lutkeni* Brooks, 1889, coralline algae (to about 75 m), and sea urchins (Gittings et al. 1992a, Hardin et al. 2001). This biotope is described as reef side by Hardin et al. (2001) and Snyder (2001).

Reef Base was defined as the ecotone between the steep reef face and the talus zone, where the rugged rocky face of the reef was often undercut and formed small caves and overhangs. This region contained both rocky vertical faces with abundant solitary corals and the coarse sediments and rocky debris/talus resulting from bioerosion occurring on the reef above. This biotope is also identified by Hardin et al. (2001) and Snyder (2001).

Reef Talus (circum-reef sediment apron) is listed as debris fields or rubble flats by Gittings et al. (1991, 1992a). These are flat areas of reef debris and coarse carbonate sediments occurring on the flanks of large, high-relief mounds (Gittings et al. 1992a, Sager and Schroeder 2001). Coarse sediments and debris are produced by shell material resulting from bioerosion, and rock fragments from the main reef. Small rocky outcrops in this biotope are often encrusted with solitary corals, small octocoral fans, and antipatharians (Gittings et al. 1992a, Hardin et al. 2001). The reef talus zone is

identifiable on HRMBS maps as the area of highest acoustic backscatter (red) to the south and west of hard bottom features (Gardner et al. 2000) (**Fig. 1.5B**).

Soft Bottom/Sand Plain refers to the relatively flat substrate surrounding reef features, characterized by fine to coarse quartz sand without visible portions of shell hash or rock rubble. These areas are often flat and featureless, but occasionally are contoured by ripples, gentle waves, or numerous excavated burrows, pits, and mounds that add to the topography. Sessile invertebrates in this habitat are limited to occasional small octocorals or antipatharians that attach to buried rock surfaces. Non-reef associated soft bottom is distinguishable in acoustic backscatter maps as areas of lowest backscatter (**Fig. 1.5B**, blue).

Data Analysis

Survey Period and Reef Tract Comparisons

ROV dives were divided into day and night periods, and data from crepuscular periods were excluded from this analysis. A two-way chi-square goodness of fit test was used to compare abundance between survey period and reef tract. A sequential Bonferroni procedure was used to adjust significance values for multiple tests (Sokal and Rohlf, 1995). At five reefs, Alabama Alps, Ludwig & Walton Pinnacles, Scamp, Yellowtail, and Roughtongue Reefs, both day and night data were obtained by ROV. Although sample sizes were small, a paired t-test was used to test day-night differences in total fish abundance, number of taxa, and top six taxa abundance (Sokal and Rohlf, 1995). All data were standardized by effort measured as video survey time in minutes.

Cluster Analysis

We used cluster analysis to help explore and define the structure of fish assemblages among reefs and biotopes. Data were summarized by reef and survey period. Data from Ludwick and Walton Pinnacles 1, 2, A, B and C were combined into a single reef category. Raw counts were standardized by effort expressed as number of fish per minute of video survey time for comparisons among reefs. Survey time was not recorded by biotope so data were double standardized for analysis. Taxa were restricted to those that occurred at two or more sites to reduce the potential effect of bias from rare species (Wolda, 1981). Two measures of similarity, percent similarity and Morisita, were used for comparison as they emphasize different attributes (Bloom, 1981; Wolda, 1981). An agglomerative hierarchical clustering strategy was used with an unweighted pair-group mean average sorting algorithm to obtain grouping of reefs and taxa (Boesch, 1977). The COMPAH computer program was used to perform the analysis (Gallagher, 1998). Clusters were subjectively divided into groups to facilitate comparisons. We compared clusters from the two similarity measures. If a reef or taxa was found in different groups between clusters it was considered poorly resolved and a weak member of the group; thus not indicative of the group structure.

Percent Similarity

$$\% \text{ SIM}_{ij} = \sum_{k=1}^S \min \left(\frac{x_{ik}}{\sum_{k=1}^S x_{ik}}, \frac{x_{jk}}{\sum_{k=1}^S x_{jk}} \right) .$$

where, x_{ik} = Abund. of species k in sample i .
 S = Number of species.

Morisita-Horn Similarity

$$MHS_{ij} = 2 \sum_{k=1}^S \frac{x_{ik} x_{jk}}{(d_i + d_j) N_i N_j} .$$

where, S = Number of species.
 x_{ik} = Abundance of species k in sample i .
 N_i = Total individuals in sample i .

$$d_i = \frac{\sum_{k=1}^S x_{ik}^2}{N_i^2} .$$

Both sites (reefs) and taxa were clustered and nodal analysis was performed to enhance the interpretation of the data (Boesch, 1977). The relationship between reefs and taxa groups was measured as constancy, the faithfulness of a taxa group to a particular reef group and fidelity, the degree to which taxa are limited to reef groups. Constancy and fidelity were arbitrarily categorized from high to low to better elucidate relationships between reefs and taxa.

Constancy

$$C_{ij} = \frac{a_{ij}}{(N_i N_j)}$$

Fidelity

$$F_{ij} = \frac{(a_{ij} ? N_j)}{(N_j ? a_{ij})}$$

where a_{ij} number of occurrences of taxa
 from taxa group i in site group j .
 N_i total number taxa in taxa group i .
 N_j total number taxa in taxa group j .

RESULTS

Faunal Composition

The results of this study have expanded the number of fishes documented from the Pinnacles Reef Tract, and is the first extensive collection of deep reef fishes made in this region, and possibly worldwide (**Table 1.3**). Of the 159 species documented (**Table 1.4**) from collected specimens or via videotape, 88 were identified from ROV censuses, and 65 can be classified as obligate reef-fishes belonging to the Caribbean reef-fish fauna. Such obligate reef fishes have life histories tied to three-dimensional, hard-bottom habitats, and would not otherwise be found on the OCS open shelf (**Figs. 1.17-1.19**). An additional 32 species are facultative reef associates that range broadly over OCS habitats, but occur on reef habitat, when available. The remaining species include primarily Carolinian soft-substrate species, including both coastal and shelf-dwelling fishes, along with a small number of pelagic and deep midwater species that were found over reef features with varying frequency and regularity.

Table 1.3. Previous investigations of the ichthyofauna of the Pinnacles Reef Tract, comparing number of reef fishes observed by ROV in each study.

Study	Reef Fish Taxa	Total Fish Taxa	Voucher Specimens?
Darnell, 1991b	39	70	No
Snyder, 2001	49	76	No
This study	65	88	Yes

Table 1.4. List of fishes documented in the Pinnacles Reef Tract from 1997-2000.

Scientific and common names follow Robins et al. (1991) except where noted with asterisk. Columns denoted as **PRF** - primary reef fishes, **A** - hook and line angling, **As** - Sabiki angling, **Ro** – ROV observed, **Rc** – ROV collected, **Tp** – Trap, **Tl** – Trawl, and **P** – voucher specimen photograph.

<u>Scientific Name</u>	<u>Common Name</u>	<u>PRF</u>	<u>A</u>	<u>As</u>	<u>Ro</u>	<u>Rc</u>	<u>Tp</u>	<u>Tl</u>	<u>P</u>
Lamnidae - mackerel sharks									
<i>Isurus oxyrinchus</i>	shortfin mako		X						X
Scyliorhinidae - cat sharks									
<i>Scyliorhinus retifer</i>	chain dogfish						X	X	X
Triakidae – houndsharks									
<i>Mustelus canis</i>	smooth dogfish		X		X				X
<i>Mustelus sinsumexicanus</i>	gulf smoothhound						X		
Carcharhinidae - requiem sharks									
<i>Carcharhinus falciformis</i>	silky shark		X						
<i>Rhizoprionodon terraenovae</i>	Atlantic sharpnose shark		X						
Sphyrnidae – hammerheads									
<i>Sphyrna lewini</i>	scalloped hammerhead		X						
Squatinaidae - angel sharks									
<i>Squatina dumeril</i>	Atlantic angel shark							X	X
Rajidae - skates									
<i>Raja eglanteria</i>	clearnose skate							X	
<i>Raja texana</i>	roundel skate							X	X
Muraenidae – morays									
<i>Gymnothorax hubbsi</i> ¹	lichen moray	X				X			X
<i>Gymnothorax kolpos</i>	blacktail moray	X	X		X		X		X
<i>Gymnothorax moringa</i>	spotted moray	X					X		X
<i>Gymnothorax nigromarginatus</i>	blackedge moray						X	X	X
<i>Gymnothorax saxicola</i>	honeycomb moray			X				X	X
<i>Muraena retifera</i>	reticulated moray	X	X				X		X
Ophichthidae - snake eels									
<i>Ophichthus gomesii</i>	shrimp eel				X			X	
<i>Ophichthus puncticeps</i>	palespotted eel						X	X	X
Congridae - conger eels									
<i>Conger oceanicus</i>	conger eel						X		
<i>Rhynchoconger gracilior</i> *	whiptail conger							X	
Nettostomatidae - duckbill eels									
<i>Hoplunnis diomedianus</i>	blacktail pike-conger				X			X	
Clupeidae - herrings									
<i>Etrumeus teres</i>	round herring							X	
Argentinidae – argentines									
<i>Glossanodon pygmaeus</i>	pygmy argentine							X	
Aulopidae – flagfins									
<i>Aulopus filamentosus</i>	yellowfin flagfin			X					X
Synodontidae - lizardfishes									
<i>Saurida normani</i>	shortjaw lizardfish					X		X	X
<i>Synodus foetens</i>	inshore lizardfish							X	
<i>Synodus intermedius</i>	sand diver	X			X			X	X
<i>Trachinocephalus myops</i>	snakefish							X	X
Myctophidae - lanternfishes					X				
Carapidae - pearlfishes									
<i>Echiodon dawsoni</i>	chain pearlfish		X ²						

¹new Gulf of Mexico record

²stomach content

*revised nomenclature

Table 1.4. (continued).

Scientific Name	Common Name	PRF	A	As	Ro	Rc	Tp	Tl	P
Ophidiidae - cusk-eels									
<i>Brotula barbata</i>	bearded brotula	X	X		X		X		X
<i>Lepophidium brevibarbe</i>	blackedge cusk-eel				X			X	
<i>Lepophidium jeannae</i>	mottled cusk-eel							X	X
<i>Ophidion beani</i>	longnose cusk-eel							X	
<i>Otophidium omostigmum</i>	polka-dot cusk-eel							X	
Moridae - codlings									
<i>Physiculus fulvus</i>	metallic codling							X	
Bregmacerotidae - codlets									
<i>Bregmaceros cantori</i>	pale codlet				X			X	X
Phycidae* – phycid hakes									
<i>Urophycis floridana</i>	southern hake						X		X
<i>Urophycis regia</i>	spotted hake						X		X
Batrachoididae – toadfishes									
<i>Opsanus pardus</i>	leopard toadfish		X		X		X		X
<i>Porichthys plectrodon</i>	Atlantic midshipman							X	X
Antennariidae – frogfishes									
<i>Antennarius ocellatus</i>	ocellated frogfish				X				
<i>Antennarius radiosus</i>	singlespot frogfish							X	X
Ogcocephalidae – batfishes									
<i>Halieutichthys aculeatus</i>	pancake batfish							X	X
<i>Ogcocephalus corniger</i>	longnose batfish				X			X	X
<i>Ogcocephalus declivirostris</i>	slantbrow batfish							X	X
Holocentridae – squirrelfishes									
<i>Corniger spinosus</i>	spinycheek soldierfish	X		X	X				X
<i>Holocentrus adscensionis</i>	squirrelfish	X		X	X				X
<i>Ostichthys trachypoma</i>	bigeye soldierfish	X			X				
<i>Sargocentron bullisi</i> *	deepwater squirrelfish	X		X	X				X
Caproidae - boarfishes									
<i>Antigonia capros</i>	deepbody boarfish	X			X				
Syngnathidae - pipefishes									
<i>Hippocampus</i> sp.	seahorse						X ²		
Aulostomidae – trumpetfishes									
<i>Aulostomus maculatus</i>	trumpetfish	X				X			X
Fistulariidae – cornetfishes									
<i>Fistularia petimba</i>	red cornetfish	X		X	X				X
Scorpaenidae - scorpionfishes									
<i>Pontinus rathbuni</i>	highfin scorpionfish	X		X				X	X
<i>Scorpaena agassizii</i>	longfin scorpionfish							X	X
<i>Scorpaena dispar</i>	hunchback scorpionfish	X	X		X				X
Triglidae – searobins									
<i>Bellator militaris</i>	horned searobin							X	X
<i>Prionotus paralatus</i>	Mexican searobin							X	X
<i>Prionotus roseus</i>	bluespotted searobin							X	X
<i>Prionotus rubio</i>	blackwing searobin							X	X
<i>Prionotus stearnsi</i>	shortwing searobin				X			X	X

PRF – primary reef fish **A** – Angling **As** – Sabiki angling **Ro** – ROV observed **Rc** –ROV collected

Tp – Trap **Tl** – Trawl **P** – Photograph

²stomach content

*revised nomenclature

Table 1.4. (continued).

<u>Scientific Name</u>	<u>Common Name</u>	<u>PRF</u>	<u>A</u>	<u>As</u>	<u>Ro</u>	<u>Rc</u>	<u>Tp</u>	<u>Tl</u>	<u>P</u>
Serranidae - sea basses									
<i>Anthias tenuis</i>	threadnose bass	X			X	X			X
<i>Centropristis ocyurus</i>	bank sea bass		X		X		X		X
<i>Centropristis philadelphica</i>	rock sea bass		X		X				X
<i>Epinephelus drummondhayi</i>	speckled hind	X	X				X		X
<i>Epinephelus flavolimbatus</i>	yellowedge grouper		X		X		X		X
<i>Epinephelus niveatus</i>	snowy grouper	X	X		X		X		X
<i>Epinephelus nigritus</i>	warsaw grouper	X					X		X
<i>Gonioplectrus hispanus</i>	Spanish flag	X	X		X				X
<i>Hemanthias leptus</i>	longtail bass	X		X	X				X
<i>Hemanthias vivanus</i>	red barbier	X		X	X	X			X
<i>Liopropoma eukrines</i>	wrasse bass	X		X	X	X			X
<i>Mycteroperca microlepis</i>	gag	X	X		X		X		X
<i>Mycteroperca phenax</i>	scamp	X	X		X		X		X
<i>Paranthias furcifer</i>	creole-fish	X		X	X		X		X
<i>Pronotogrammus martinicensis</i> *	rougtongue bass	X		X	X	X			X
<i>Rypticus maculatus</i>	whitespotted soapfish	X	X		X				X
<i>Serranus atrobranchus</i>	blackear bass				X			X	X
<i>Serranus notospilus</i>	saddle bass				X			X	X
<i>Serranus phoebe</i>	tattler	X		X	X				X
Grammatidae – basslets									
<i>Lipogramma regia</i> ¹	royal basslet	X			X				
Opistognathidae – jawfishes									
<i>Opistognathus lonchurus</i>	moustache jawfish	X		X	X				X
Priacanthidae - bigeyes									
<i>Priacanthus arenatus</i>	bigeye	X	X		X				X
<i>Pristigenys alta</i>	short bigeye	X	X		X				X
Apogonidae - cardinalfishes									
<i>Apogon pseudomaculatus</i>	twospot cardinalfish	X		X	X				X
Malacanthidae - tilefishes									
<i>Caulolatilus chrysops</i>	goldface tilefish	X	X				X		X
<i>Caulolatilus intermedius</i>	anchor tilefish		X						X
<i>Malacanthus plumieri</i>	sand tilefish	X	X		X		X		X
Pomatomidae - bluefishes									
<i>Pomatomus saltatrix</i>	bluefish		X		X				X
Echeneidae - remoras									
<i>Echeneis naucrates</i>	sharksucker		X						
Carangidae - jacks									
<i>Caranx crysos</i>	blue runner		X						X
<i>Caranx hippos</i>	crevalle jack		X						
<i>Selar crumenophthalmus</i>	bigeye scad		X					X	
<i>Seriola dumerili</i>	greater amberjack	X	X		X		X		X
<i>Seriola fasciata</i>	lesser amberjack		X		X		X		X
<i>Seriola rivoliana</i>	almaco jack	X	X				X		X
<i>Seriola zonata</i>	banded rudderfish		X						X
<i>Trachurus lathami</i>	rough scad			X	X				X

PRF – primary reef fish **A** – Angling **As** – Sabiki angling **Ro** – ROV observed **Rc** –ROV collected

Tp – Trap **Tl** – Trawl **P** – Photograph

¹new Gulf of Mexico record

*revised nomenclature

Table 1.4. (continued).

Scientific Name	Common Name	PRF	A	As	Ro	Rc	Tp	Tl	P
Lutjanidae - snappers									
<i>Lutjanus campechanus</i>	red snapper	X	X		X		X		X
<i>Pristipomoides aquilonaris</i>	wenchman		X		X			X	X
<i>Rhomboplites aurorubens</i>	vermilion snapper	X	X	X	X		X		X
Sparidae – porgies									
<i>Calamus leucosteus</i>	whitebone porgy		X				X		X
<i>Calamus nodosus</i>	knobbed porgy	X	X						X
<i>Lagodon rhomboides</i>	pinfish				X		X	X	X
<i>Pagrus pagrus</i>	red porgy	X	X		X		X		X
<i>Stenotomus caprinus</i>	longspine porgy				X			X	X
Sciaenidae - drums									
<i>Cynoscion arenarius</i>	sand seatrout		X						X
<i>Leiostomus xanthurus</i>	spot				X			X	X
<i>Micropogonias undulatus</i>	Atlantic croaker						X	X	X
<i>Pareques iwamotoi</i>	blackbar drum	X			X		X		X
<i>Pareques umbrosus</i>	cubbyu	X			X		X		X
Mullidae - goatfishes									
<i>Mullus auratus</i>	red goatfish							X	X
<i>Upeneus parvus</i>	dwarf goatfish				X			X	X
Chaetodontidae - butterflyfishes									
<i>Chaetodon aya</i>	bank butterflyfish	X			X	X	X		X
<i>Chaetodon ocellatus</i>	spotfin butterflyfish	X			X		X		X
<i>Chaetodon sedentarius</i>	reef butterflyfish	X		X	X		X		X
Pomacanthidae - angelfishes									
<i>Holacanthus bermudensis</i>	blue angelfish	X			X		X		X
<i>Holacanthus ciliaris</i>	queen angelfish	X			X				
<i>Holacanthus tricolor</i>	rock beauty	X			X				
Pomacentridae - damselfishes									
<i>Chromis enchrysurus</i>	yellowtail reeffish	X		X	X	X			X
<i>Chromis insolata</i>	sunshinefish	X			X				
<i>Chromis scotti</i>	purple reeffish	X			X				
Labridae - wrasses									
<i>Bodianus pulchellus</i>	spotfin hogfish	X		X	X				X
<i>Decodon puellaris</i>	red hogfish	X		X	X			X	X
<i>Halichoeres bathyphilus</i>	greenband wrasse	X		X	X				X
Uranoscopidae - stargazers									
<i>Kathetostoma albigutta</i>	lancer stargazer							X	X
Blenniidae - combtooth blennies									
<i>Parablennius marmoratus</i>	seaweed blenny	X	X ²		X				
Gobiidae - gobies									
<i>Coryphopterus punctipictophorus</i>	spotted goby	X			X				
<i>Risor ruber</i>	tusked goby	X							X ³
Sphyraenidae – barracudas									
<i>Sphyraena borealis</i>	northern sennet		X						X
Gempylidae* - snake mackerels									
<i>Neopinnula americana</i>	American sackfish				X				
Trichiuridae - cutlassfishes									
<i>Trichiurus lepturus</i>	Atlantic cutlassfish				X			X	X

PRF – primary reef fish **A** – Angling **As** – Sabiki angling **Ro** – ROV observed **Rc** –ROV collected

Tp – Trap **Tl** – Trawl **P** – Photograph

²stomach content

³box core

*revised nomenclature

Table 1.4. (continued).

Scientific Name	Common Name	PRF	A	As	Ro	Rc	Tp	Tl	P
Scombridae - mackerels									
<i>Acanthocybium solandri</i>	wahoo		X						
<i>Auxis thazard</i>	frigate mackerel			X					
<i>Euthynnus alletteratus</i>	little tunny		X						
<i>Sarda sarda</i>	Atlantic bonita		X						
<i>Scomberomorus cavalla</i>	king mackerel		X		X				
Stromateidae - butterfishes									
<i>Peprilus burti</i>	gulf butterfish					X			
Bothidae - lefteye flounders									
<i>Engyophrys senta</i>	spiny flounder							X	X
<i>Trichopsetta ventralis</i>	sash flounder							X	X
Paralichthyidae* - sand flounders									
<i>Ancylosetta dilecta</i>	three-eye flounder					X		X	X
<i>Citharichthys cornutus</i>	horned whiff					X		X	X
<i>Cyclopsetta fimbriata</i>	spotfin flounder					X		X	X
<i>Paralichthys squamilentus</i>	broad flounder							X	X
<i>Syacium papillosum</i>	dusky flounder					X		X	X
Achiridae - American soles									
<i>Gymnachirus texae</i>	fringed sole							X	X
Soleidae* - soles									
<i>Symphurus diomedianus</i>	spottedfin tonguefish					X		X	X
<i>Symphurus parvus</i>	pygmy tonguefish							X	
Balistidae - triggerfishes									
<i>Balistes capriscus</i>	gray triggerfish	X	X						X
Monacanthidae* - filefishes									
<i>Monacanthus tuckeri</i>	slender filefish	X		X					X
Ostraciidae - boxfishes									
<i>Acanthostracion quadricornis*</i>	scrawled cowfish					X			
Tetraodontidae - puffers									
<i>Canthigaster rostrata</i>	sharpnose puffer	X		X	X	X			X
<i>Canthigaster jamestyleri</i> ¹	goldface toby	X		X	X				X
<i>Lagocephalus laevigatus</i>	smooth puffer		X						X
<i>Sphoeroides dorsalis</i>	marbled puffer							X	X
<i>Sphoeroides spengleri</i>	bandtail puffer		X		X				X
Diodontidae - porcupinefishes									
<i>Chilomycterus antillarum</i>	web burrfish	X				X			
<i>Chilomycterus schoepfii</i>	striped burrfish					X			
<i>Diodon holocanthus</i>	balloonfish	X				X			
Molidae – ocean sunfishes									
<i>Mola mola</i>	ocean sunfish					X			

PRF – primary reef fish **A** – Angling **As** – Sabiki angling **Ro** – ROV observed **Rc** –ROV collected

Tp – Trap **Tl** – Trawl **P** – Photograph

¹new Gulf of Mexico record

*revised nomenclature




Figure 1.17. Still frames taken from video surveys at the Pinnacles Reef Tract. **A.** *Pronotogrammus martinicensis* school along crevice in reef top at Yellowtail Reef, 63 m. **B.** Close-up of mixed group of *P. martinicensis* and *Hemanthias vivanus* along reef face. **C.** Mixed school of anthiines along reef face at CMEP-Site 5, 75 m. **D.** Single *P. martinicensis* along reef face at Scamp Reef, 105 m. **C** and **D** illustrate invertebrates that dominate the drowned reef community [ahermatypic coral colonies, *Rhizopsammia manuelensis* (black clusters), and crinoids].




Figure 1.18. Still frames taken from video surveys at the Pinnacles Reef Tract. **A.** Mixed school of *Hemanthias vivanus* and *Chromis enchrysurus* with a single *Pristigenys alta* on reef top biotope at Yellowtail Reef, 62 m. **B.** *Holacanthus bermudensis* and *Chaetodon aya* along northern edge of reef top biotope at Yellowtail Reef, 63 m. **C-D.** Reef top biotope with well-developed sessile invertebrate community of gorgonians and crinoids, Yellowtail Reef, 62 m.




Figure 1.19. Still frames taken from video surveys at the Pinnacles Reef Tract. **A.** *Seriola dumerili* schooling above reef flat top, Yellowtail Reef, 62m. **B.** *Gonioplectrus hispanus* at reef base, CMEP-Site 5, 78 m. **C.** Female *Mycteroperca phenax* on reef face, Scamp Reef, 105 m. **D.** Dominant male *M. phenax* exhibiting grey-head color phase along reef face, Scamp Reef, 105 m.

ROV Survey

Videotape data from only three cruises (USGS 97-01, 99-03 and 2000-01) were of sufficient quality for quantitative analysis of reef fish abundance. Fishes observed during other ROV surveys where visibility was poor, or while collecting fishes using the suction sampler (USGS 2000-01 and 2000-02), were added to the list of species occurring on the Pinnacles Reef Tract but not included in the analysis. Eighty-eight species were identified during ROV surveys, with 17 taxa observed only by this method (**Table 1.4**).

ROV surveys revealed the dominance of planktivorous reef fishes on high profile reef features (**Table 1.5**, Appendix A). Anthiines were the principal taxa observed by day, and communities were dominated by either *Hemanthias vivanus* or *Pronotoqrammus martinicensis*, depending on reef location and biotope. Day ROV surveys yielded higher (9719 individuals) overall abundance of fishes than night surveys (2624 individuals) (**Table 1.5**, Appendix A1), indicating most reef species are diurnally active and hide in crevices or burrows at night. About equal numbers of taxa were observed during the day (68) and night (66) though almost twice as much effort was applied during the day. Day surveys had 20 unique taxa, of which none were frequent. Night surveys had 18 unique taxa. Three taxa, *Bregmaceros cantori*, myctophids, and *Trichiurus lepturus*, were common, but not observed on all night dives. These typically non-reef species may make a more substantial contribution to the reefs than previously expected. We compared reef fish abundance by survey period and reef tract simultaneously as these factors might be confounded. Only data from reef habitat were used in this analysis. The chi-square goodness of fit test showed that most taxa had a significant difference in abundance between survey period and reef tract (**Table 1.6**). The diurnal nature of the fauna is evident in that the total number of individuals was significantly lower at night especially on shallow reefs. Seven taxa were significantly more abundant on shallow reefs during the day (**Table 1.6**). Only *Synodus* spp. was significantly lower during the day due to its absence from shallow water reefs. Two taxa, *P. martinicensis* and *Anthias tenuis* were significantly more abundant on deep reefs during the day. Five taxa were most abundant on shallow and six on deep reefs (**Table 1.6**). *Pareques umbrosus* was significantly lower on shallow reefs at night while *Chaetodon aya*, *Decodon puellaris*, and *Serranus phoebe* were lower on deep reefs.

When day/night comparisons were made by reef there was a highly significant difference in total fish abundance between day and night with higher numbers during the day (**Table 1.7**). However, there was no significant difference in number of taxa observed between day and night. Of the top six taxa, four were more abundant during the day but none were significantly so due to the small sample size (**Table 1.7**). *Pronotoqrammus martinicensis* tends to lie on the reef surface at night being inactive but visible. In contrast *H. vivanus* hides in crevices at night, thus hidden from the ROV survey. Myctophids, a nocturnally migrating taxon, were significantly more abundant at night overall, as was *T. lepturus*, but due to patchy distributions there was no significant difference among reefs.

Table 1.5. Abundance of top 30 taxa observed in ROV surveys partitioned by survey period (day/night) and reef tract (shallow/deep) for reef habitat.

Rank	Taxa	Overall No.	Percent of Total Reef Habitat			
			Day		Night	
			Shallow	Deep	Shallow	Deep
	<i>Pronotoqrammus</i>					
1	<i>martinicensis</i>	5027	25.1	77.3	46.7	27.4
2	<i>Hemanthias vivanus</i>	3881	54.4	3.6	15.9	0.3
3	Gobiidae	499	3.9	<0.1	0.4	
4	Myctophidae	335			0.1	23.1
5	<i>Anthias tenuis</i>	272	<0.1	8.3		0.8
6	<i>Chromis enchrysur</i>	240	3.0	<0.1	3.1	0.1
7	<i>Trichiurus lepturus</i>	239				16.5
8	<i>Pristigenys alta</i>	231	1.1	1.8	4.6	3.2
9	<i>Bregmaceros cantori</i>	179			2.6	9.9
10	<i>Chaetodon aya</i>	161	1.2	1.3	2.8	0.7
11	<i>Halichoeres bathyphilus</i>	130	1.7	0.3	0.3	0.1
12	<i>Trachurus lathami</i>	123	0.3		2.0	5.5
12	<i>Decodon puellaris</i>	123	1.0	0.3	0.4	0.1
14	<i>Pareques umbrosus</i>	115	1.0	0.7	0.3	1.7
15	<i>Serranus notospilus</i>	98				
16	<i>Serranus phoebe</i>	84	0.8	0.2	0.9	
17	<i>Canthigaster rostrata</i>	83	0.6	0.1	3.1	0.1
18	<i>Corniger spinosus</i>	80	0.5	0.5	1.8	0.5
19	<i>Synodus</i> spp.	77			0.2	1.2
20	<i>Liopropoma eukrines</i>	74	0.8	0.5	0.3	0.2
21	<i>Scorpaena dispar</i>	66	0.2	0.3	2.6	0.3
22	<i>Saurida</i> spp.	64				
23	<i>Seriola dumerili</i>	63	0.1	0.4	0.9	2.1
24	<i>Chaetodon sedentarius</i>	45	0.3	0.2	1.7	
25	<i>Paranthias furcifer</i>	39	0.6		0.1	
26	<i>Rhomboplites aurorubens</i>	36	0.3		1.4	0.1
27	<i>Apogon pseudomaculatus</i>	35	0.2	0.0	1.3	0.4
27	<i>Mycteroperca phenax</i>	35	0.3	0.3		0.3
29	<i>Chromis scotti</i>	29	0.4		0.1	
30	Scorpaenidae	28	0.0	0.3	0.2	1.0
	Total No. Individuals	12343	6574	3145	1175	1449
	No. Taxa	66	60	42	48	43
	No. Unique Taxa		11	5	3	7
	Video Survey Time (min)		689	329	296	313

Table 1.6. Results of chi-square goodness of fit test for reef fish abundance by survey period (day/night) and reef tract (shallow/deep). Significantly higher or lower abundance is indicated by factor.

Rank	Taxa	Day		Night		Overall
		Shallow	Deep	Shallow	Deep	<i>P</i>
1	<i>Pronotogrammus martinicensis</i>		Higher			<0.0001
2	<i>Hemanthias vivanus</i>	Higher				<0.0001
3	Gobiidae	Higher				<0.0001
4	Myctophidae				Higher	<0.0001
5	<i>Anthias tenuis</i>		Higher			<0.0001
6	<i>Chromis enchrysur</i>	Higher				<0.0001
7	<i>Trichiurus lepturus</i>				Higher	<0.0001
8	<i>Pristigenys alta</i>					ns
9	<i>Bregmaceros cantori</i>				Higher	<0.0001
10	<i>Chaetodon aya</i>				Lower	0.0005
11	<i>Halichoeres bathyphilus</i>	Higher				<0.0001
12	<i>Trachurus lathami</i>				Higher	<0.0001
12	<i>Decodon puellaris</i>				Lower	<0.0001
14	<i>Pareques umbrosus</i>			Lower		0.0003
15	<i>Serranus notospilus</i>					ND
16	<i>Serranus phoebe</i>				Lower	<0.0001
17	<i>Canthigaster rostrata</i>			Higher		<0.0001
18	<i>Corniger spinosus</i>					ns
19	<i>Synodus</i> spp.	Lower				<0.0001
20	<i>Liopropoma eukrines</i>	Higher				<0.0001
21	<i>Scorpaena dispar</i>			Higher		<0.0001
22	<i>Saurida</i> spp.					ND
23	<i>Seriola dumerili</i>				Higher	<0.0001
24	<i>Chaetodon sedentarius</i>			Higher		<0.0001
25	<i>Paranthias furcifer</i>	Higher				<0.0001
26	<i>Rhomboplites aurorubens</i>			Higher		<0.0001
27	<i>Apogon pseudomaculatus</i>			Higher		0.0006
27	<i>Mycteroperca phenax</i>					ns
29	<i>Chromis scotti</i>	Higher				<0.0001
30	Scorpaenidae				Higher	<0.0001
	No. Individuals	Higher	Higher			<0.0001
	No. Taxa			Higher		<0.0001

ns – not significant ND – no data for comparison

Table 1.7. Day-night comparison of selected community parameters and top six taxa (abundance) paired by reef based on ROV data. Sample size equals five. Significant differences are indicated in bold.

Mean per Minute Video				
	Day	Night	t	P
No. Individuals	16.48	4.00	3.28	0.01
No. Taxa	0.36	0.32	0.29	0.39
<i>P. martinicensis</i>	6.35	1.38	1.48	0.11
<i>H. vivanus</i>	5.44	0.24	2.10	0.05
Myctophids	0	0.68	-1.55	0.10
<i>A. tenuis</i>	0.85	0.02	1.00	0.19
Gobiids	0.17	0.01	1.37	0.12
<i>T. lepturus</i>	0	0.33	-1.01	0.18

Taxonomic groups that were not effectively sampled or video documented during the course of the study included juvenile anthiines and gobiids, due to their extremely small size. Gobies were the third most abundant species observed by ROV censuses (**Table 1.5**), and may be represented by a number of species, but only the spotted goby, *Coryphopterus punctipectophorus*, was documented by close-up video during this study (**Table 1.4**).

Biotopes

From ROV analyses of fish faunal composition, we recognize that species richness and relative abundance of reef fishes differs among biotopes [reef zones (including reef top, reef crest, reef face, and reef base), talus zone, and sand flat habitats, **Table 1.8, Figs. 1.20-1.23**], and these differences occur across the various study reefs. All reef biotopes are dominated by anthiine serranids, but showed partitioning. *Hemanthias vivanus* was most abundant in reef top and reef crest biotopes while *P. martinicensis* was increasingly abundant on the reef face and reef base (**Table 1.8**). Additional reef fish taxa that numerically dominated reef top and reef crest biotopes included gobiids, the pomacentrid *C. enchrysur*, the priacanthid *Pristigenys alta* and the labrids *Halichoeres bathyphilus* and *D. puellaris*. The reef face biotope was dominated by anthiine serranids. The schooling serranid *Paranthias furcifer*, *D. puellaris*, and the holocentrid *Corniger spinosus* also characterize this biotope. Fishes common on the reef base biotope included species that typically are associated with crevices and holes, including the holocentrid *Ostichthys trachypoma*, the sciaenid *Pareques umbrosus* and the serranid *Liopropoma eukrines*.

Fishes observed on the reef top were represented by more taxa (28) than reef crest, face, and base biotopes (19, 22 and 22, respectively). High species richness is likely due to the diversity of microhabitats available, including rock outcrops, accumulated coarse sand, and dense assemblages of sessile invertebrates such as sponges, gorgonians and antipatharians that form spatially complex structure. The diversity of habitat present on the reef top also presents a wide variety of prey resources that are associated with unconsolidated sediments and a diverse invertebrate assemblage.

A wide range of fishes also frequents the reef top biotope during night hours. Mobile demersal visual piscivores (groupers, snappers) are found on the reef top during evening hours, and schools of amberjack commonly occur on flat-topped features at night. Night ROV surveys also documented the occurrence of soft bottom sciaenids (e.g., *Leiostomus xanthurus*) and carangids (*Trachurus lathami*) utilizing the reef top. Midwater planktivores from the adjacent slope, including *B. cantori*, *T. lepturus*, and lanternfishes (Myctophidae), have been occasionally observed over reef features at night, and were observed in great abundances on deeper sites such as the Ludwick and Walton Pinnacles (Appendix A4).

While reef biotopes were similar in overall species composition, they differed greatly from the talus zone and sand flat (**Table 1.8, Figs. 1.22-1.23**). Species assemblages were very similar among reef biotopes (percent similarity > 0.40) and dissimilar from the talus zone and sand flat (percent similarity = 0.14) (**Fig. 1.20**). Overall number of taxa was lower in the talus zone (15) and sand flat (16) than reef biotopes. Gobiids were particularly abundant in the circum-reef talus zone (70.1%), with *D. puellaris*, and the serranids *Serranus phoebe* and *S. notospilus* were also dominant (**Table 1.8, Appendix A**). Lizardfishes (*Synodus* spp.) were also common in this biotope.

Table 1.8. Biotope association of Pinnacles Reef Tract fishes based on ROV surveys during Cruise 97-01, combined for all study sites. Numbers of individuals observed at each biotope during videotape analysis, and percent of total, are given. Top five taxa by biotope are indicated in bold.

Taxa	REEF TOP		REEF CREST		REEF SLOPE		REEF BASE		TALUS ZONE		SAND FLAT	
	No.	Percent	No.	Percent	No.	Percent	No.	Percent	No.	Percent	No.	Percent
<i>Hemanthias vivanus</i>	1698	58.8	314	59.7	442	39.6	39	8.7			-	-
<i>Pronotogrammus martinicensis</i>	618	21.4	135	25.7	536	48.1	308	68.8	1	0.5	4	1.2
Gobiidae	161	5.6	27	5.1	10	0.9	5	1.1	129	70.1	71	20.5
<i>Chomis enchrysur</i>	88	3.0	7	1.3	5	0.4	2	0.4	2	1.1	-	-
<i>Halichoeres bathyphilus</i>	57	2.0	9	1.7	-	-	3	0.7	7	3.8	-	-
<i>Pristigenys alta</i>	43	1.5	1	0.2	8	0.7	2	0.4	1	0.5	-	-
<i>Decodon puellaris</i>	37	1.3	1	0.2	13	1.2	8	1.8	18	9.8	19	5.5
<i>Chaetodon aya</i>	26	0.9	6	1.1	9	0.8	5	1.1	-	-	-	-
<i>Canthigaster rostrata</i>	24	0.8	-	-	2	0.2	-	-	-	-	-	-
<i>Trachurus lathami</i>	19	0.7	-	-	-	-	-	-	-	-	3	0.9
<i>Serranus phoebe</i>	15	0.5	3	0.6	3	0.3	6	1.3	8	4.3	5	1.4
<i>Liopropoma eukrines</i>	13	0.5	-	-	-	-	9	2.0	1	0.5	-	-
<i>Scorpaena dispar</i>	12	0.4	3	0.6	4	0.4	3	0.7	2	1.0	2	0.6
<i>Rhomboplites aurorubens</i>	6	0.2	-	-	9	0.8	-	-	-	-	-	-
<i>Parablennius marmoratus</i>	6	0.2	1	0.2	-	-	-	-	-	-	-	-
<i>Sargocentron bullisi</i>	6	0.2	-	-	-	-	-	-	-	-	-	-
<i>Priacanthus arenatus</i>	5	0.2	-	-	-	-	3	0.7	1	0.5	-	-
<i>Pareques umbrosus</i>	5	0.2	-	-	-	-	26	5.8	-	-	-	-
<i>Mycteroperca microlepis</i>	5	0.2	1	0.2	5	0.4	-	-	-	-	-	-
<i>Chaetodon sedentarius</i>	5	0.2	5	1.0	3	0.3	2	0.4	-	-	-	-
<i>Seriola dumerili</i>	4	0.1	3	0.6	-	-	-	-	-	-	-	-
<i>Holacanthus bermudensis</i>	4	0.1	1	0.2	2	0.2	-	-	-	-	-	-
<i>Corniger spinosus</i>	4	0.1	5	1.0	12	1.1	6	1.3	-	-	-	-
<i>Serranus notospilus</i>	3	0.1	-	-	-	-	4	0.9	3	1.6	84	24.2
<i>Centropristis ocyurus</i>	3	0.1	-	-	-	-	1	0.2	-	-	2	0.6
<i>Synodus</i> sp.	2	0.1	-	-	-	-	-	-	6	3.3	40	11.5
<i>Lutjanus campechanus</i>	2	0.1	-	-	2	0.2	-	-	-	-	-	-
<i>Mycteroperca phenax</i>	2	0.1	-	-	11	1.0	1	0.2	-	-	-	-
<i>Pontinus rathbuni</i>	-	-	2	0.4	-	-	-	-	1	0.5	-	-
<i>Pagrus pagrus</i>	-	-	1	0.2	-	-	-	-	-	-	-	-
<i>Paranthias furcifer</i>	-	-	1	0.2	15	1.3	-	-	-	-	-	-
<i>Pareques iwamotoi</i>	-	-	-	-	5	0.4	2	0.4	-	-	-	-
<i>Rypticus maculatus</i>	-	-	-	-	2	0.2	-	-	-	-	-	-
<i>Opsanus pardus</i>	-	-	-	-	2	0.2	-	-	-	-	-	-
<i>Gonioplectrus hispanus</i>	-	-	-	-	2	0.2	-	-	-	-	-	-
<i>Ostichthys trachypoma</i>	-	-	-	-	-	-	11	2.5	-	-	-	-
<i>Hemanthias leptus</i>	-	-	-	-	-	-	1	0.2	-	-	-	-
Unidentified flatfish	-	-	-	-	-	-	1	0.2	-	-	7	2.0
<i>Saurida</i> sp.	-	-	-	-	-	-	-	-	-	-	62	17.9
<i>Pristipomoides aquilonaris</i>	-	-	-	-	-	-	-	-	-	-	7	2.0
<i>Hoplunnis</i> sp.	-	-	-	-	-	-	-	-	-	-	7	2.0
<i>Syacium papillosum</i>	-	-	-	-	-	-	-	-	-	-	6	1.7
<i>Upeneus parvus</i>	-	-	-	-	-	-	-	-	-	-	3	0.9
<i>Bregmaceros cantori</i>	-	-	-	-	-	-	-	-	1	0.5	2	0.6
<i>Ancylosetta dilecta</i>	-	-	-	-	-	-	-	-	1	0.5	-	-
Other taxa	13	0.5	-	-	13	1.2	-	-	2	1.0	22	6.6
Totals	2886	100.0	526	100.0	1115	100.0	448	100.0	184	100.0	346	100.0




Figure 1.20. Typical ichthyofauna contributing to community structure on the Pinnacles Reef Tract, northeastern Gulf of Mexico, based on ROV data. Cluster diagram shows relationship among biotopes.




Figure 1.21. Pinnacles Shallow Reef Tract fish faunal composition for reef crest biotope.




Figure 1.22. Pinnacles Shallow Reef Tract fish faunal composition for reef base biotope.

REEF BIOTOPE ASSEMBLAGES




Figure 1.23. Pinnacles Shallow Reef Tract fish faunal composition for talus zone biotope.

Sand flats were also dominated by gobies, with the small serranid *S. notospilus*, lizardfishes (*Synodus* spp. and *Saurida* spp.), *D. puellaris* and flatfishes also common (**Table 1.8**). Anthiine serranids were either absent (*H. vivanus*) or in low abundance (*P. martinicensis*) at these biotopes.

Reef Groups and Species Associations

Fish assemblages from ROV data were compared among reefs using cluster analysis. Five groups of reefs were resolved at a 0.30 similarity level (**Fig. 1.24**). **Table 1.9** compares reef type and community parameters among these groups. Taxa clustering resulted in eight groups at a 0.12 similarity level (**Fig. 1.25**).

Site groups 3 to 5 separate reefs with distinctive faunal compositions. Deeper reef night samples formed the most unique cluster (Site Group 5) with high constancy (faithfulness to a reef group) of Taxa Group G such as myctophids, *T. lepturus*, *B. cantori*, and *T. lathami* (**Fig. 1.25**). The Yellowtail Reef night survey was exceptional in the high constancy with Taxa Group G and very low abundance of *P. martinicensis*. This species is found on the bottom at night and was regularly seen during the night at other sites. Its low abundance at Yellowtail Reef may be due to the low relief of the area surveyed at night. Far Tortuga and Shark Reef (Site Group 3), both large mounds with gentle, silted slopes and no high vertical faces, shared high abundance of gobies and *D. puellaris*.

Site Group 2 contained most of the high relief shallow reef trend sites with the exception of Solitary Mounds. This site group had high constancy with Taxa Groups B and F the largest suite of taxa and strong fidelity (the degree to which taxa are limited to a reef group) with Taxa Group F that included the common taxa, *H. vivanus* and *P. martinicensis* (**Fig. 1.26**). Solitary Mounds fit with this group due to its high abundance of *C. enchrysur*. Site Group 2 reefs on average had high abundances and low species richness (**Table 1.9**).

Site Group 1 sites included deep reefs, night surveys of high diversity shallow water sites, and shallow low relief sites. This indicates that the deep reef fauna is most closely related to the nocturnal fauna of the shallow reef tract. No taxa group had high constancy for this site group, but there was strong fidelity with Group F taxa (**Fig. 1.26**). The low abundance of Taxa Group B, especially *C. enchrysur* and *D. puellaris*, in Site Group 1 differentiates it from Site Group 2. Alabama Alps day and night surveys were very similar (percent similarity = 0.83) sharing 32% (20) of the taxa. In particular *A. tenuis*, *P. umbrosus*, and *S. dumerili* contributed to this similarity. Cat's Paw and Corkscrew Reefs were not well resolved moving among clusters depending on similarity measure employed.




Figure 1.24. Cluster of reef sites using percent similarity and unweighted pair-group mean average combinatorial strategy. [N] designates night dives.

Table 1.9. Summary statistics for ROV data by site group based on cluster analysis. [N] denotes night dives at these sites.

Site Group	Reef	Reef Trend	Relief	Survey Time (min)	Total No. Fish	Total No. Taxa	Mean No. Fish*	Mean No. Taxa*
1								
	Alabama Alps [N]	Deep	High	97.0	482	28	4.97	0.06
	Alabama Alps	Deep	High	61.2	1697	25	27.75	0.01
	Ludwick & Walton Pinnacles ¹	Deep	High	244.8	1088	32	4.44	0.03
	Scamp Reef	Deep	High	23.3	360	16	15.44	0.04
	Cat's Paw Reef	Shallow	High	27.4	81	7	2.96	0.09
	Patch Reef Field	Shallow	Low	58.6	85	12	1.45	0.14
	Roughtongue Reef [N]	Shallow	High	200.2	966	41	4.83	0.04
	Corkscrew Reef	Shallow	High	40.9	363	24	8.88	0.07
	Porgy Reef [N]	Shallow	High	58.5	130	14	2.22	0.11
2								
	CMEP5	Shallow	High	11.3	580	21	51.42	0.04
	Yellowtail Reef	Shallow	High	171.6	2535	40	14.77	0.02
	Double Top Reef	Shallow	High	106.2	354	16	3.33	0.05
	Roughtongue Reef	Shallow	High	90.9	1516	29	16.69	0.02
	Solitary Mounds ²	Shallow	Intermediate	79.6	584	25	7.34	0.04
3								
	Far Tortuga	Shallow	Intermediate	23.5	72	11	3.06	0.15
	Shark Reef	Shallow	Low	78.6	103	17	1.31	0.17
4								
	Yellowtail Reef [N]	Shallow	High	37.2	79	24	2.12	0.30
5								
	Ludwick & Walton Pinnacles ³ [N]	Deep	High	146.2	766	25	5.24	0.03
	Scamp reef [N]	Deep	High	69.9	200	20	2.86	0.10

¹includes Pinnacles 1, A, B, and C. ²includes Mounds 1 and 2. ³includes Pinnacles 1 and A.

*per minute video




Figure 1.25. Cluster analysis of taxa using percent similarity and unweighted pair-group mean average combinatorial strategy. Clusters were truncated at the group level. Taxa are represented by code, first three letters of genera plus four letters of species name, or first seven letters for genus and higher taxa.





Figure 1.26. Constancy and fidelity for site and taxa groups defined by cluster analysis. Size of cells is related to number of taxa or sites in each group.

Angling

Various remote sampling gears are effective for different groups of taxa. Whereas underwater video documents the fauna much more holistically, some taxa are attracted to or repelled by the ROV light. In addition, cryptic and burrowing species are underrepresented. We used several additional gear types to compliment the ROV survey and enhance our knowledge of the system. Sixty-nine species were taken by hook and line (**Table 1.4**). The gear contributed substantially to our knowledge of community composition, as 21 species were only taken by this gear (particularly sharks, jacks, and mackerels). The numerically dominant taxa collected by baited hook and line are similar to previous surveys (CSA 1992), as *R. aurorubens* and *P. pagrus* were the most frequently caught large predators in both studies (**Table 1.10**). The numerical dominance of *R. aurorubens* and *P. pagrus* also was reported by Grimes et al. (1982) for shelf-edge reefs of the Carolinas. Hook and line sampling was important in estimating large predator abundance, since many of the larger predators are either attracted to the lights of the ROV (carangids, large serranids) or repelled (*R. aurorubens*, *P. pagrus*) during visual surveys biasing abundance estimates. While *R. aurorubens* and *P. pagrus* dominated hook and line samples, they were rarely observed during fish surveys, and typically swim away from the lights when encountering the ROV. We know from comparison of video survey and angling collections that the latter provide an incomplete and highly biased picture of faunal composition, particularly when using standard hook and line angling gear with large, baited hooks.

Traps

Traps collected 38 fish taxa with six species- *Conger oceanicus*, *Epinephelus nigritus*, *Gymnothorax moringa*, *Hippocampus* sp. (stomach contents), *Urophycis floridana*, and *U. regia* - unique to the gear (**Table 1.4**). Of the common taxa, the gear added substantially to the collections of *Pareques* spp., *Brotula barbata*, and eels such as *Muraena retifera* and *Gymnothorax kolpos* (**Table 1.11**). Traps are most effective samplers of cryptic and burrowing species and fish that locate prey by chemosensory means (e.g., eels).

Trawls

Trawls could only be deployed on low relief bottom and thus sampled near-reef sand flat biotope. Fifty-five taxa were collected (**Table 1.4**) with 27 unique to the gear. The faunal separation of the reef and flat bottom communities can be seen in the increased importance of ophidiids (cusk eels), triglids (sea robins), and flatfish (Bothidae and Paralichthyidae) in the trawl catch (**Table 1.12**). *Saurida* spp. and *Serranus notospilus* were dominant in the trawl catch and in ROV surveys on surrounding bottom. Few gobies were taken by the trawl due to mesh size and their propensity to rapidly flee to nearby burrows. Trawls were the only method effective in sampling juvenile *D. puellaris* and *S. notospilus*.

Table 1.10. Abundance of top 15 predatory fishes caught by baited hook and line angling from 1997-2000.

Rank	Species	No.	Percent
1	<i>Rhomboplites aurorubens</i>	450	60.4
2	<i>Pagrus pagrus</i>	114	15.3
3	<i>Lutjanus campechanus</i>	94	12.6
4	<i>Seriola dumerili</i>	22	3.0
5	<i>Centropristis ocyurus</i>	9	1.2
6	<i>Caulolatilus chrysops</i>	9	1.2
7	<i>Malacanthus plumieri</i>	7	0.9
8	<i>Opsanus pardus</i>	6	0.8
9	<i>Paranthias furcifer</i>	4	0.5
10	<i>Mycteroperca phenax</i>	4	0.5
11	<i>Seriola rivoliana</i>	3	0.4
12	<i>Mycteroperca microlepis</i>	3	0.4
13	<i>Muraena retifera</i>	3	0.4
14	<i>Epinephelus niveatus</i>	3	0.4
15	<i>Epinephelus flavolimbatus</i>	3	0.4
Total		745	100

Table 1.11. Abundance of top 15 fish taxa caught in traps from 1997-2000.

Rank	Species	No.	Percent
1	<i>Rhomboplites aurorubens</i>	233	34.9
2	<i>Pagrus pagrus</i>	126	18.9
3	<i>Pareques umbrosus</i>	71	10.6
4	<i>Centropristis ocyurus</i>	42	6.3
5	<i>Lutjanus campechanus</i>	33	4.9
6	<i>Brotula barbata</i>	31	4.6
7	<i>Seriola dumerili</i>	15	2.2
8	<i>Muraena retifera</i>	15	2.2
9	<i>Gymnothorax saxicola</i>	14	2.1
10	<i>Mycteroperca phenax</i>	13	1.9
11	<i>Micropogonias undulatus</i>	10	1.5
12	<i>Mycteroperca microlepis</i>	7	1.0
13	<i>Gymnothorax kolpos</i>	7	1.0
14	<i>Pareques iwamotoi</i>	5	0.7
15	<i>Holacanthus bermudensis</i>	5	0.7
Total		668	100

Table 1.12. Abundance of top 15 fish taxa caught by trawl from 1997.

Rank	Species	No.	Percent
1	<i>Saurida</i> spp.	140	14.0
2	<i>Serranus notospilus</i>	122	12.2
3	<i>Prionotus paralatus</i>	79	7.9
4	<i>Pristipomoides aqualonaris</i>	76	7.6
5	<i>Stenotomus caprinus</i>	72	7.2
6	<i>Syacium papillosum</i>	70	7.0
7	<i>Prionotus stearnsi</i>	47	4.7
8	<i>Serranus atrobranchus</i>	43	4.3
9	<i>Centropristis philadelphica</i>	37	3.7
10	<i>Synodus</i> spp.	36	3.6
11	<i>Citharichthys cornutus</i>	34	3.4
12	<i>Ophidion holbrooki</i>	27	2.7
13	<i>Trichopsetta ventralis</i>	20	2.0
14	<i>Lepophidium brevibarbe</i>	20	2.0
15	<i>Hali euthichthys aculeatus</i>	18	1.8
Total		998	100

DISCUSSION

Within the depth range of the Pinnacles reef tract (60-110 m), previous studies in the northeastern Gulf have indicated that the fish fauna includes components of the Pan-Caribbean reef fish fauna, the Carolinian inshore shelf fauna, and the soft-bottom deep-water outer shelf/upper slope fauna (Shipp and Hopkins 1977, Darnell 1991b, Gittings et al. 1991, McEachran 1991, Snyder 2001). Faunal composition and habitat affinities were comparatively evaluated with previous studies on Gulf of Mexico and western Atlantic reef and offshore continental shelf fish species (e.g., Springer and Bullis 1956, Bullis and Thompson 1965, Bright and Cashman 1974, Smith et al. 1975, Walls 1975, Hastings et al. 1976, Smith 1976, Sonnier et al. 1976, Gilmore 1977, Shipp and Hopkins 1978, Williams and Shipp 1980, Boland et al. 1983, Clarke 1986, Dennis and Bright 1988a, b, Hoese and Moore 1998, Thompson et al. 1999, Snyder 2001). At least 250 reef-associated fishes have been documented from offshore banks, reefs, and other hard-bottom habitats in the northern Gulf of Mexico based on previous literature (Appendix B). The suite of primary reef fishes inhabiting the Pinnacles Reef Tract (**Table 1.4**) represents a depauperate, winnowed fraction of the total reef-fish fauna (Appendix B). The selectively winnowed reef-fish fauna of the Pinnacles Reef Tract appears to represent a characteristic assemblage of deep-reef fishes broadly distributed in the Pan-Caribbean area (Colin, 1974, Colin, 1976). It is important to establish the reasons for this winnowing, particularly if the potential effects of anthropogenic activities on the OCS are to be rationally assessed and monitored. It can be hypothesized that the known Pinnacles reef fish diversity is diminished relative to the Caribbean due to: 1) distance from the source fauna, 2) depth limitation, 3) temperature limitation, 4) habitat limitation due to turbidity, 5) trophic limitations, 6) fishing pressure, or 7) incomplete sampling. However, the first alternative can largely be eliminated as mid-shelf and shelf edge banks off Texas and Louisiana are as far or farther from the Caribbean source region, but support a large number of reef species (Appendix B). Over 150 reef fish species reported to occur in the northwestern Gulf and the Florida Middle Ground have not been documented in the Pinnacles Reef Tract (Appendix B). Thus, a large number of Caribbean reef-fish species do arrive on reefs in the northern Gulf either by larval transport on ocean currents (e.g., Loop Current intrusions), or by active migration. Therefore, habitat or other limitations, and not geographic location alone, appear to be responsible for the absence (or rarity) of many reef fishes on the Pinnacles.

Many Caribbean reef-fish species do not occur at increasing depths within a given geographic area, even in regions of high water clarity and warm temperatures that extend beyond 100 m (Colin 1974, 1976, Dennis and Bright 1988a). Depth per se does not have a direct effect on fish assemblages but covaries with light, temperature, and other physical parameters. The effects of temperature versus depth are difficult to distinguish, and highly variable according to geographic location. The northern Gulf of Mexico is exposed to winter cold fronts that can drastically lower the temperature of inshore waters (Kelly and Bender 2001). The position of the Pinnacles Reef Tract on the OCS provides temperature buffering, with deep Gulf waters moderating temperature fluctuations. Winter bottom temperature averaged 18.1°C with a low of 15.4°C during the study period (Kelly and Bender, 2001). These temperatures are just

below levels required for reef development and are similar to conditions at mid-shelf banks in the northwestern Gulf of Mexico that lack active reef growth (Bright et al. 1984, Rezak et al. 1985). While reef fish recruits can colonize outside their normal geographic range they often do not persist due to limiting environmental conditions. However, water temperatures are not low enough at the Pinnacles Reef Tract to be a limiting factor to colonization for reef fishes. *Chaetodon ocellatus* (spotfin butterflyfish), a tropical reef species, does not survive winter temperatures below 10°C though numerous recruits colonize New Jersey estuaries annually (McBride and Able 1998). Similarly, reef fishes in the northern Adriatic can tolerate temperatures from 7-10°C during winter months (Kotrschal and Reynolds 1982). These temperatures are well below winter minima recorded at the Pinnacles Reef Tract and the temperature range becomes less variable with depth contributing to stable environmental conditions (Rezak et al. 1985, Kelly and Bender 2001).

A persistent nepheloid layer characterized by high turbidity was identified as a controlling factor for hard bottom communities in the northwestern Gulf (Rezak et al. 1985). The nepheloid layer increases light attenuation resulting in decreased epibiota and reef fish species richness and abundance below 80 m (Dennis and Bright, 1988a, Rezak et al. 1990). Previous studies have suggested that the Mississippi River plume influences the distribution and abundance of sessile invertebrates within 70 km of the river delta, and may produce a gradient of sedimentation and water column turbidity throughout the Pinnacles Reef Tract (Gittings et al. 1992a, Hardin et al. 2001, Kelly and Bender 2001). In addition, turbidity may affect the feeding efficiency of planktivores (Coker, 1993). In the northeastern Gulf, nepheloid layers are infrequent, though in conjunction with episodic Mississippi freshwater plumes and upwelling result in increased light attenuation (Kelly and Bender, 2001).

The combination of low winter temperatures and low light levels resulting from increased water depth and turbidity on the Pinnacles Reef Tract results in a reduction of algal and hard coral growth, and subsequent habitat availability for reef fishes. While planktivorous reef fishes dominate Pinnacles reefs, the absence of *Chromis multilineata* (brown chromis), *Chromis cyanea* (blue chromis), and *Clepticus parrae* (creole wrasse), planktivores that form dense schools on reefs in the northwestern Gulf, suggests that it is a result of factors associated with depth and not food limitation that accounts for the absence of such species on the Pinnacles Reef Tract. Of the 150 species listed in Appendix B, only 25 are known from below 60 m depths, equivalent to the shallowest Pinnacles reefs. Conversely, a number of Pinnacles species, notably the dominant anthiine serranids, are absent from the shallow coral cap regions of the Flower Garden Banks (18-45 m) and also are not present at the Florida Middle Ground, where depths range from 26-48 m (Smith et al. 1975, Clarke 1986, Brooks and Doyle 1991). Increased water depth and associated light attenuation may induce food limitations that account for the absence or rarity of other taxa. For example, herbivorous scarids and acanthurids, which form large, free-roaming schools that graze algae from the reef substratum on shallow coral reefs throughout the Caribbean, and occur at both the Flower Gardens and Florida Middle Ground, are absent from the Pinnacles Reef Tract. Similarly, many species of pomacentrids (*Stegastes* spp.) that

propagate algal gardens on dead coral surfaces are absent from the Pinnacles Reef Tract fauna, but are present on shallower banks in the northwestern Gulf where light levels are sufficient for growth of filamentous and leafy algae (Dennis and Bright, 1988a). Reduced light levels also results in the absence of a well-developed algal-sponge zone, a dominant biotope occurring at the Flower Gardens at similar depth range (50-80 m) to the shallow Pinnacles reefs (Rezak et al. 1985). Reef fishes that are abundant on the algal-sponge zone at the Flower Gardens, including *Centropyge argi* (cherubfish), *Sparisoma atomarium* (greenblotch parrotfish), and *Serranus annularis* (orangeback bass) are absent on the Pinnacles Reef Tract.

Substrate complexity or rugosity contributes substantially to patterns of reef fish distribution and abundance (Syms and Jones 2000). Friedlander and Parrish (1998) found that epifauna played little role in determining fish assemblages; substrate complexity was a much more important factor. In low relief soft bottom habitats, pits and burrows are particularly important in determining fish distribution (Felley et al. 1989). On deep reefs, the dominant planktivores are visual predators active only during daylight hours. They retreat into crevices and shelter holes at night, and also during the day upon approach of potential predators. The relative abundance of small planktivores and benthivores may depend not only on food availability, but also on relative abundance of shelter holes (= substrate rugosity). While many studies have found correlations between hole size/number and structural complexity with reef fish abundance or habitat selection (Friedlander and Parrish 1998, Syms and Jones 2000, Snyder 2001), when empirically tested no limitation on shelter holes was found for a shallow water reef fish (Robertson and Sheldon 1979).

Most Pinnacle Reef Tract features are high relief drowned reefs giving rise to a range of biotopes (Gardner et al. 2001). Geological features associated with diapirism, including gas and brine seeps that greatly influence the biotic communities in the northwestern Gulf are rare in the Pinnacles Reef Tract (Gittings et al. 1992a, Gardner et al. 2001). Gittings et al. (1992a) and Hardin et al. (2001) have observed different assemblages of sessile macro-invertebrates in the Pinnacles Reef Tract related to structural form (e.g., horizontal summits versus vertical or rugged features). Study sites that have extensive flat-top areas including Roughtongue Reef, Yellowtail Reef, and Alabama Alps, develop a high diversity invertebrate assemblage that is dominated by octocorals, antipatharians, sponges, bryozoans, coralline algae (on shallower reef sites) and crinoids, and is typically characterized by relatively high percent cover of invertebrates (Hardin et al. 2001) (**Fig. 1.18**). Vertical reef faces, overhangs, and rock outcrops and boulders occurring at the reef base are dominated by the solitary coral *R. manuelensis* (**Fig. 1.17**). Results of our ROV analyses indicate strong fish faunal differentiation based on physically defined biotope type (**Table 1.5; Figs. 1.21-1.23**, Appendix A). Fine-scale factors such as habitat complexity (Friedlander and Parrish 1998) may control absolute fish abundance within a particular biotope type, but relative abundance appears more a function of basic biotope type.

Fishes observed on the reef top included the highest number of taxa across all biotopes, and this species richness is likely due to the diversity of microhabitats available,

including rock outcrops, sand pockets, and dense assemblages of invertebrates such as sponges, octocorals (including large sea fans), and antipatharians that form spatially complex structures. Also the reef top represents the greatest reef area; thus the species-area relationship may be a factor (MacArthur and Wilson, 1963). The diversity of habitat present on the reef top also presents a wide variety of prey resources that are associated with unconsolidated sediments and a diverse invertebrate assemblage. This biotope not only supports planktivores (including *C. enchrysur*), but also benthic carnivores (*H. bathyphilus* and *D. puellaris*), epibenthic browsers (*C. rostrata*, *C. sedentarius* and *H. bermudensis*), and generalized carnivores (*S. phoebe* and *L. eukrines*) (see Chapter 2 for more details on trophic categories).

The reef crest and face are important foraging habitat not only for planktivores, but also for epibenthic carnivores, and mobile piscivores that feed on both planktivores and benthivores. The rugose, highly sculptured substrate provides shelter for small fishes. Stony corals largely replace soft corals, crinoids, antipatharians, and sponges on vertical reef faces, limiting the structural complexity of the reef surface. The reef base, composed of caves and undercuts, is often occupied by species such as *G. hispanus*, *P. umbrosus*, *C. spinosus*, *O. trachypoma*, and *L. eukrines*. Extending out from the reef base is the circum-reef talus apron. This biotope is characterized by small to medium-size benthic carnivores (e.g., labrids, serranids, *P. umbrosus*) that forage for benthic and epibenthic prey, and not plankton. Once on the open sand flat habitat of the shelf proper, fish density is low and fish are primarily active at night and aggregated around any structure (Felley et al. 1989). The distinct fish assemblages within each biotope and the different areal extent of biotopes among reefs needs further study to provide a more detailed picture of factors controlling deep reef community structure. We are extending our future research in this area.

Reef profile and topographic complexity remain somewhat enigmatic in their respective roles in controlling community structure of reef fishes. Beyond a critical threshold, the overall height of a pinnacle structure may be fairly unimportant. What is critical is that the structure be of sufficient height to reduce exposure to turbidity and sedimentation, and perturb flow on the reef face and crest, and over the reef top (Gittings et al. 1992a, MacDonald and Peccini 2001). Messing et al. (1990) found that lithoherm banks in the northeastern Straits of Florida hydrodynamically accelerated ambient 2-7 cm/s bottom currents to greater than 100 cm/s over the upcurrent reef crest. These authors schematically illustrate both the compression of current streamlines over a lithoherm resulting in current acceleration (Messing et al. 1990, Fig. 3), and the resultant upcurrent to downcurrent biozonation of suspension-feeding sessile macroinvertebrates (Messing et al. 1990, Fig. 4). Fairly continuous currents from the southwest sweep the steep-sided Pinnacle reefs as reflected in the orientation of soft corals (MacDonald and Peccini 2001). Flow models indicate that acceleration of currents over shelf-edge pinnacles may increase availability of pelagic prey in the vicinity of high profile features (Kelly and Bender 2001). The "wall of mouths" concept developed on the Great Barrier Reef (Hamner et al. 1988) suggests an upcurrent to downcurrent pattern of faunal differentiation due to plankton removal from the impinging current and the importance of plankton availability in controlling fish faunal assemblages among biotopes. The

characterization of biotopes and their associated fish communities on a finer scale that couples water currents and biological/geological characteristics of the benthic habitat may ultimately be necessary to explain the occurrence and composition of the reef fish assemblage. We are presently following this line of inquiry.

Although hermatypic corals with zooxanthellae and leafy algae are a major carbon source for fishes on shallow reefs (Ogden 1982), recent research has revealed that plankton also may be of great importance in certain regions (Hamner et al. 1988). A characteristic set of deep-water planktivores, together with a suite of geographically ubiquitous, deep-water browsers and benthic carnivores, dominate the Pan-Caribbean deep-water fish fauna (Colin, 1974, Colin, 1976; Dennis and Bright, 1988a, this study). Indeed, a model of trophic partitioning of the reef-fish fauna in deep water, based on the Pinnacles Reef Tract example, may have broad application throughout deep reef communities of the world's oceans (see Chapter 2 for details). Trophic partitioning of the Pinnacles fauna closely parallels the earlier findings of Thresher and Colin (1986) for the deep (>75-90 m) reef community of Enewetak Atoll in the southwestern Pacific. Below 90 m, the Enewetak fish fauna was similarly dominated by planktivorous anthiine serranids (>90% of all individuals), together with planktivorous labrids and pomacentrids (*Chromis* spp.). All other trophic guilds were very poorly represented in deep water, except for piscivores (1-10% by number) that feed on the abundant planktivores. Patterns are similar in the Pinnacles Reef Tract except for the low abundance of piscivores.

Both the dominant sessile macro-invertebrates (e.g., gorgonians and antipatharians) and reef fishes on Pinnacles Reef Tract reefs are planktivores, and spatial and temporal access to plankton appear to be major factors in the control of community structure and organism distribution (Hardin et al. 2001, MacDonald and Peccini 2001). Consistency of supply and volume of the planktonic food source may be a strong controlling factor for these fish assemblages. Intercepting the incoming current, the upcurrent reef face and crest should be the most favorable biotopes for swarms of diurnal planktivores (predominantly anthiine serranids). We hypothesize that this is true based on ROV observations, but a quantitative assessment is needed.

Direct anthropogenic effects, such as fishing, also may have altered the reef fish community in the Pinnacles Reef Tract. Commercial fisheries for *Lutjanus campechanus* began in the 1870's, first based in Pensacola, Florida, and expanding to numerous fleets based throughout the Gulf by the early 1930's (Jarvis 1935), targeting snappers and groupers on offshore banks. Darnell (1991b) reports declines in biomass of bottom fishes on the shelf and decreased harvest rates of *L. campechanus*, and attributes these patterns to overfishing and habitat loss. Snyder (2001) points out declining catch rates of large groupers (*Epinephelus* spp. and *Mycteroperca* spp.), *L. campechanus*, and *S. dumerili* throughout the northeastern Gulf since the early 1970's, and a rise in catch rates of *R. aurorubens* during this same time period. These patterns have been linked to the widespread use of bottom longlines in outer shelf waters of the Gulf (Richards and McGowan 1989) that target large demersal fishes, and may explain the low abundance of these species in our collections and ROV surveys. The apparent decrease in the abundance of large piscivores in the Pinnacles Reef Tract due to fishing

activities may have had the further consequence of increasing the absolute and relative abundance of their primary prey, the anthiine serranid planktivores or other species of lower trophic levels, such as *R. aurorubens*, and lead to alteration of ecosystem structure and function (Parsons 1992, 1996, Pauly et al. 1998). Thus, caution should be exercised in generalizing patterns in community structure for ecosystems that have been subject to anthropogenic alteration.

Fishes and invertebrates have been characterized at the Pinnacles Reef Tract by three major studies and all major reef features have been precisely mapped. This area may be the most well-known deep hard bottom area in the world. Thus we postulate that the majority of the reef fish species that occur at depths between 60 and 110 m are now documented. Yet due to the large area and diversity of habitat further scientific inquiry will continue to add to our knowledge of the fish fauna. Our present knowledge of the Pinnacles ichthyofauna can be used to characterize the fundamental taxonomic composition and trophic structure of deep reef-fish communities throughout southeastern United States and Gulf of Mexico OCS areas and therefore be useful in assessing future anthropogenic impacts on faunal structure.

CHAPTER 2

TROPHIC ECOLOGY OF PINNACLES TRACT REEF FISHES

INTRODUCTION

Defining trophic pathways is the key to developing an understanding of the ecological framework in the reef fish community. Analysis of food web structure also identifies relative importance of resident reef fishes (autochthonous) versus off-reef trophic resources (allochthonous) such as adjacent soft-substrate and pelagic prey in the diet of large predatory fishes, predator-prey relationships among fishes and invertebrates, and may also identify potential pathways of energy transfer from fishes to the benthic invertebrate assemblage through the release of fecal material. Trophic guild categories are general terms to describe the location of their main source of prey and identify pelagic versus benthic trophic pathways. Within each feeding guild most members will utilize a wide spectrum of prey based on availability/encounter rates, prey size, and prey mobility. Detailed studies of the food habits of deep reef fishes have not been reported in the literature, and few reports of shallow reef fish diets in the Gulf of Mexico are available (e.g., Bullock and Smith 1991, Nelson and Bortone 1996, Weaver 1996). Investigations on food habits of reef fishes in the South Atlantic Bight primarily have focused on commercially important species (Manooch 1977, Grimes 1979, Ross 1982, Matheson et al. 1986, Dodrill et al. 1993, Lindquist and Clavijo 1993, Lindquist et al. 1994, Sedberry and Cuellar 1993). The reef fish, invertebrate, and plant assemblages that characterize live bottom communities in the Gulf of Mexico vary dramatically with depth (Rezak et al. 1985, 1990), limiting comparisons among shallow and deep reef communities from published literature. Therefore, identifying food habits of deep reef fishes and developing food web models from the literature would yield an inaccurate, simplistic view. A full understanding requires sampling of fishes from deepwater reef communities to document dietary patterns. Diets of marine fishes are known to vary dramatically with body size, local microhabitat, and season (Crabtree et al. 1991, Mullaney and Gale 1995, Sheridan and Trimm 1983, Starck 1971), and therefore collection of fishes provides the only method to develop a detailed description of trophic pathways in the deep reef environment. Trophodynamic investigations undertaken in the present study include feeding habits and food web structure of Pinnacles Reef Tract fishes.

OBJECTIVES

- Delineate food habits of numerically dominant fish species.
- Construct a trophic model based on quantitatively-defined food webs that model the Pinnacles fish community structure and function, and convey predictions about primary sources of forage base and energy for reef and reef-associated biotopes.

METHODS

There is an extensive body of literature regarding methods for analysis of fish stomach contents and food habits, and conventional quantitative parameters of broad application have been reviewed (Hyslop 1980). Specimens sampled for food habits from USGS 97-01 through 2000-01 cruises were kept on ice, frozen, or preserved in formalin until processed in the laboratory at the Florida Caribbean Science Center, Gainesville, FL (FCSC). Only specimens from hook and line sampling were analyzed and specimens with empty or everted stomachs were excluded. Trap specimens were not included in the analysis. Fishes were dissected in the laboratory and the whole digestive tracts (stomachs plus intestines) were removed. Prey items were identified to lowest possible taxonomic level. For simplicity, dietary analyses in this report are given as percent number, calculated by dividing the number of prey within each taxon by the total number of prey items for each of the characteristic reef fishes collected excluding empty stomachs. Stomach contents were used to identify the ecological role of each taxon in the reef fish community and used to identify feeding guilds following previous authors (Hobson 1982, Bohnsack et al. 1999).

Food Web Model

A food web model was developed for the Pinnacles fish community. A quantitative model was based on numerical contribution of prey to the diet of each predator. Primary goals of food web models are: 1) to compare relative contribution of allochthonous materials (i.e., plankton and adjacent soft-bottom off-reef organisms) to the energy budget of the dominant members of the reef fish community; and 2) to determine the relative importance of reef fishes in the diet of large reef-associated carnivores. Diets of the characteristic reef fishes were incorporated into a working food web model following the methods of Winemiller (1990) to identify the dominant trophic pathways among the reef fish community. Trophic position of each predator was calculated using the formula:

$$T_i = 1.0 + \sum_{j=1}^n T_j (p_{ij})$$

where T_i is the trophic level of an individual predator species i , summing relative importance of prey taxon j where p_{ij} is the proportion of the total diet (based on numerical abundance) for species i consisting of prey taxon j . Primary producers are assigned a value of $T=0.0$, herbivores $T=1.0$, and a consumer eating half plant and half herbivore tissue would be $T=1.5$, and so forth, as T_i 's are computed sequentially from bottom to top. Calculated T_i values were used as the y axis for placement of consumer nodes of the food web. The x coordinate of each consumer was weighted according to the proportion of prey from different habitat compartments. Proportions of pelagic prey (plankton) were multiplied by 0.0 (far left), reef-based prey by 0.5 (middle), and soft bottom prey by 1.0 (far right), to determine relative dependence on each habitat for prey (Winemiller 1990). This method gives estimates of relative dependence on allochthonous versus autochthonous materials (i.e., a position close to the left portion of the web is dependent on pelagic prey, the middle portion resident reef prey, and the right side off-reef soft bottom prey). Links connecting each compartment were then calculated with respect to movement of energy from prey to predator (uplinks) (Winemiller 1990). Links among the different nodes (predator-prey relation) can be expressed with uniform width and relative strength (qualitative models) or line width varied to reflect proportional contribution of prey in the diet (quantitative models). Terminology of plankton size classes and constituent members follow Christensen (1995).

RESULTS AND DISCUSSION

Over 1000 individuals were examined for food habits with approximately 50% containing prey items. Many specimens everted their stomachs due to the change in pressure when brought to the surface and the majority of the large predators (jacks, snappers, and groupers) had empty stomachs. In addition, many resident reef fishes were poorly sampled by hook and line angling, reflected by low sample sizes. Feeding guilds were defined by characteristic prey (**Table 2.1**). Dietary patterns of the characteristic reef fish taxa are presented in this report (**Table 2.2**, Appendix C). Fish were assigned to feeding guilds based on stomach contents to identify similarity in prey resources among species, sources of prey, and overall patterns in trophic structure (**Table 2.3**). Use of the conventional terms, such as macroinverteviores and microinverteviores, by other authors to describe feeding guilds is avoided due to the wide size range of prey selected by most species (e.g., *Serranus phoebe* feeds on adult crabs and fishes as well as decapod and fish larvae).

The numerically dominant species (greater than 80% of the individuals observed by ROV) on high profile reefs, *Pronotogrammus martinicensis* and *Hemanthias vivanus*, have diets dominated by calanoid copepods, gastropod larvae, and a variety of other pelagic mesoplankton organisms (**Table 2.2**, Appendix C). Macroplanktivores that prey on larger size classes of plankton include *Rhomboplites aurorubens*, *Pristigenys alta*, *Apogon pseudomaculatus*, and *Chaetodon aya*. Benthic feeders included *Pagrus pagrus*, *Pareques umbrosus*, *Decodon puellaris*, and *Halichoeres bathyphilus*.

Table 2.1. Feeding guilds defined by characteristic prey type.

Feeding Guild

Mesoplanktivore- Mixed zooplankton in the microzooplankton to mesoplankton (20-2000 μ) size classes. Includes copepods, ostracods, larvaceans, gastropod veligers, pteropods, crustacean larvae, invertebrate eggs, and fish eggs.

Macroplanktivore- Mixed zooplankton in the macroplankton (>2000 μ) size class. Includes hyperiid amphipods, shrimp zoea, stomatopod zoea, crab megalopa, fish larvae, salps, and squids.

Benthic Carnivores (~microinvertivores)-primarily infauna/epifaunal invertebrates including crabs, bivalves, gastropods, brittle stars, sea urchins, and polychaetes.

Epibenthic Browsers- encrusting, sessile invertebrates including hydrozoans, bryozoans, soft corals, crinoids, and sponges.

Generalized Carnivores (~macroinvertivores)- a variety of mobile, epibenthic prey that may include fishes, crabs, shrimps, and gastropods.

Piscivores- includes fishes and squids.

Table 2.2. Dietary analysis of characteristic reef fishes of the Pinnacles Reef Tract by trophic guild. Values given are based on numerical contribution to the diet (%N).

		Mesoplanktivores					Macroplanktivores					Benthic Carnivores						
		<i>Bregmaceros cantori</i>	<i>Chromis enchrysurus</i>	<i>Hemanthias vivanus</i>	<i>Paramthias furcifer</i>	<i>Promologrammus martinicensis</i>	<i>Apogon pseudonaculatus</i>	<i>Chaetodon aya</i>	<i>Priacanthus arenatus</i>	<i>Pristigerys alta</i>	<i>Rhomboplites aurorubens</i>	<i>Bodianus pulchellus</i>	<i>Cantolattus chrysops</i>	<i>Decodon puellaris</i>	<i>Halichoeres bathyphilus</i>	<i>Pagrus pagrus</i>	<i>Pareques iwamotoi</i>	<i>Pareques umbrosus</i>
Sample Size		2	20	47	1	120	5	13	15	9	72	4	2	2	10	28	1	4
Phylum	Prey Item	Percent Number																
	Invertebrate eggs	-	19.1	3.0	-	11.0	-	-	-	-	-	-	-	-	-	-	-	-
Protoctista	Foraminiferans	-	-	-	-	-	-	-	-	-	-	-	-	-	6.9	-	-	-
	Radiolarians	-	-	-	-	1.5	-	-	-	-	-	-	-	-	-	-	-	-
Porifera	Sponges	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.1	-	-
Cnidaria	Hydrozoans	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Soft Corals	-	-	-	-	-	-	4.8	-	-	-	-	-	-	-	-	-	-
Mollusca	Heteropods	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Gastropods	-	-	-	-	-	-	-	-	-	-	14.3	-	31.3	-	16.0	33.3	-
	Pelecypods	-	-	-	-	-	-	-	-	-	-	38.1	20.0	37.5	48.3	3.2	-	-
	Pteropods	-	-	-	-	-	-	-	1.3	-	6.9	-	-	-	-	-	-	-
	Chitons	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Squids	-	-	-	-	-	-	-	32.9	16.0	0.2	-	-	-	3.4	-	-	-
Annelida	Polychaetes	-	13.3	17.7	35.7	10.3	-	-	-	14.3	-	45.0	6.3	3.4	4.3	-	9.1	
Arthropoda	Copepods	60.0	32.4	78.8	28.6	69.7	-	-	-	15.2	-	-	-	-	-	-	-	
	Amphipods	20.0	-	-	-	0.2	-	5.1	-	-	15.2	-	-	-	-	-	-	
	Barnacle larvae	-	-	-	-	0.1	-	-	-	1.4	-	-	-	-	-	-	-	
	Crab larvae	-	1.1	-	14.3	0.4	12.5	90.1	14.5	16.0	3.7	-	-	12.5	13.8	2.1	-	
	Crabs	-	-	-	-	-	25.0	-	1.3	2.0	-	19.0	5.0	6.3	6.9	34.0	33.3	
	Isopods	-	-	-	-	-	37.5	-	-	-	-	-	-	-	-	-	-	
	Ostracods	-	-	0.1	-	1.7	-	-	-	2.1	-	-	-	-	-	-	-	
	Shrimp larvae	20.0	2.1	-	21.4	-	25.0	-	1.3	16.0	10.1	-	-	-	6.9	-	-	
	Shrimps	-	-	-	-	-	-	-	3.9	40.0	-	-	-	-	-	2.1	-	
	Stomatopod larvae	-	-	-	-	-	-	-	27.6	6.0	8.0	-	-	6.3	-	6.4	-	
	Stomatopods	-	-	-	-	-	-	-	-	-	-	-	25	-	4.3	-	-	
Bryozoa	Bryozoans	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Echindermata	Echinoids	-	-	-	-	-	-	-	-	-	-	-	-	-	-	20.2	-	
	Ophiuroids	-	-	-	-	-	-	-	-	-	-	9.5	5.0	-	6.9	-	33.3	
	Holothuroids	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.1	-	
	Crinoids	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Chordata	Larvaceans	-	32.0	0.1	-	1.9	-	-	-	-	1.1	-	-	-	-	-	-	
	Salps	-	-	-	-	0.5	-	-	-	-	9.9	-	-	-	-	-	-	
	Fish eggs	-	-	-	-	2.4	-	-	-	-	1.4	-	-	-	-	-	-	
	Fish larvae	-	-	-	-	-	-	-	14.5	-	10.6	-	-	-	-	-	-	
	Fishes	-	-	-	-	-	-	-	2.6	2.0	-	19.0	-	-	3.4	5.3	-	

Table 2.2. (continued).

		<u>Epibenthic browsers</u>			<u>Generalized Carnivores</u>									<u>Piscivores</u>			
	Sample Size	<i>Holacanthus bermudensis</i>	<i>Chaetodon sedentarius</i>	<i>Canthigaster rostratus</i>	<i>Centropomus ocyurus</i>	<i>Epinephelus flavolimbatus</i>	<i>Holocentrus adscensionis</i>	<i>Liopropoma eukrines</i>	<i>Lutjanus campechanus</i>	<i>Malacanthus plumieri</i>	<i>Opsanus pardus</i>	<i>Pontinus rathbuni</i>	<i>Sargocentron ballisi</i>	<i>Serranus phoebe</i>	<i>Aulostomus maculatus</i>	<i>Gymnothorax saxicola</i>	<i>Seriola dumerili</i>
Phylum	Prey Item	4	8	2	6	2	3	1	26	7	5	1	2	44	1	1	6
		Percent Number															
	Invertebrate eggs	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Protoctista	Foraminiferans	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Radiolarians	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Porifera	Sponges	8.3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Cnidaria	Hydrozoans	-	71.4	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Soft Corals	41.7	14.4	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Mollusca	Heteropods	-	-	-	-	-	-	-	5.1	-	-	-	-	-	-	-	-
	Gastropods	-	-	3.2	11.8	-	-	-	25.0	33.3	-	-	-	-	-	-	-
	Pelecypods	-	-	7.8	-	-	-	-	18.8	-	-	-	-	-	-	-	-
	Pteropods	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Chitons	-	-	2.1	-	-	-	-	-	-	-	16.7	-	-	-	-	-
	Squids	-	-	-	-	-	-	2.6	-	-	-	-	1.4	-	-	-	-
Annelida	Polychaetes	-	-	-	-	-	-	-	-	-	-	16.7	1.4	-	-	-	-
Arthropoda	Copepods	-	5.2	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Amphipods	-	-	-	-	-	-	-	-	-	-	-	1.4	-	-	-	-
	Barnacle larvae	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Crab larvae	-	-	-	-	42.9	-	-	-	-	-	33.3	5.6	-	-	-	-
	Crabs	-	-	-	76.5	-	-	5.1	18.8	25.0	-	41.7	-	-	-	-	-
	Isopods	-	-	-	-	-	-	10.3	-	-	-	-	-	-	-	-	-
	Ostracods	-	2.2	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Shrimp larvae	-	2.1	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Shrimps	-	-	-	5.9	33.3	14.3	7.7	-	-	60.0	-	5.6	-	-	-	-
	Stomatopod larvae	-	-	-	-	-	-	-	-	-	-	33.3	-	-	-	-	-
	Stomatopods	-	-	-	-	-	-	2.6	-	-	-	1.4	-	-	-	-	-
Bryozoa	Bryozoans	41.7	-	-	-	-	-	-	6.3	-	-	-	-	-	-	-	-
	Echinoids	-	1.5	-	-	-	-	-	-	8.3	-	-	-	-	-	-	-
	Ophiuroids	-	-	-	5.9	-	-	-	12.5	-	-	1.4	-	-	-	-	-
	Holothuroids	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Crinoids	8.3	-	78.6	-	-	-	-	-	8.3	-	-	-	-	-	-	-
Chordata	Larvaceans	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Salps	-	-	-	-	-	-	10.3	-	-	-	-	-	-	-	-	-
	Fish eggs	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Fish larvae	-	-	-	-	-	-	-	-	-	-	2.8	-	-	-	-	-
	Fishes	-	-	-	-	66.7	42.9	100	56.4	18.8	25.0	40.0	-	37.5	100	100	100

Table 2.3. Feeding guilds of fishes along the Pinnacles Reef Tract. Determination of feeding guild is modified from Hobson (1982) and Bohnsack et al. (1997, 1999) and based on stomach contents identified in this study, or from food habits by Randall (1967) for closely related taxa.

Feeding Guild

Mesoplanktivores- *Anthias tenuis* (threadnose bass), *Chromis enchrysur* (yellowtail reef fish), *Hemanthias vivanus* (red barbier), *Paranthias furcifer* (creole-fish), *Pronotogrammus martinicensis* (rougtongue bass), *Trachurus lathami* (rough scad).

Macroplanktivores- *Apogon pseudomaculatus* (twospot cardinalfish), *Chaetodon aya* (bank butterflyfish), *Priacanthus arenatus* (bigeye), *Pristigenys alta* (short bigeye), *Rhomboplites aurorubens* (vermilion snapper).

Benthic Carnivores- *Balistes capriscus* (gray triggerfish), *Calamus leucosteus* (whitebone porgy), *Calamus nodosus* (knobbed porgy), *Caulolatilus chrysops* (goldface tilefish), *Decodon puellaris* (red hogfish), *Halichoeres bathyphilus* (greenband wrasse), *Pagrus pagrus* (red porgy), *Pareques iwamotoi* (blackbar drum), and *P. umbrosus* (cubbyu).

Epibenthic Browsers – *Canthigaster rostrata* (sharppnose puffer), *Chaetodon sedentarius* (reef butterflyfish), *Chaetodon ocellatus* (spotfin butterflyfish), *Holacanthus bermudensis* (blue angelfish), *Parablennius marmoratus* (seaweed blenny).

Generalized Carnivores- *Bodianus pulchellus* (spotfin hogfish), *Centropristis ocyurus* (bank sea bass), *Corniger spinosus* (spinycheek soldierfish), *Epinephelus flavolimbatus* (yellowedge grouper), *Epinephelus nigritus* (Warsaw grouper), *Epinephelus niveatus* (snowy grouper), *Holocentrus adscensionis* (squirrelfish), *Liopropoma eukrines* (wrasse bass), *Lutjanus campechanus* (red snapper), Muraenidae, *Opsanus pardus* (leopard toadfish), *Malacanthus plumieri* (sand tilefish), *Sargocentron bullisi* (deepwater squirrelfish), *Pontinus rathbuni* (highfin scorpionfish), *Rypticus maculatus* (whitespotted soapfish), *Scorpaena dispar* (hunchback scorpionfish), *Serranus phoebe* (tattler).

Piscivores- *Aulostomus maculatus* (trumpetfish), *Fistularia petimba* (red cornetfish), Muraenidae, *Mycteroperca microlepis* (gag), *M. phenax* (scamp), *Seriola dumerili* (greater amberjack), *Seriola fasciata* (lesser amberjack), *Seriola rivoliana* (almaco jack).

Epibenthic browsers included *Chaetodon sedentarius* and *Holacanthus bermudensis* whose diets were composed of sessile invertebrates. Piscivores were represented by *Mycteroperca* spp., *Seriola* spp., *Lutjanus campechanus*, muraenids, *Aulostomus maculatus*, and *Fistularia petimba*. The generalized carnivore guild was represented by numerous taxa including *S. phoebe*, and a variety of serranids, scorpaenids, muraenids, and holocentrids. As discussed by Snyder (2001), this guild contains the most taxa of deep reef fishes, although individual species are not as numerically abundant as planktivores and benthic carnivores in reef communities.

Planktivorous anthiine fishes may serve as a keystone predator by foraging in the water column and transferring energy to the deep reef ecosystem. Anthiines in turn become prey for a variety of reef predators. Stomach contents indicate that newly settled juveniles and young-of-the-year anthiines are an important seasonal component of the diet for a variety of reef fishes, including *L. campechanus*, carangids, and serranids. Trophic pathways among the deep reef community indicate primary links to the pelagic community through direct consumption of planktonic organisms and may be supplemented by surrounding benthic communities through consumption of eggs and larvae of soft-bottom fishes, a variety of invertebrates, and potentially by spawning populations of migratory species from nearshore reef communities (e.g., *Mycteroperca microlepis*). In contrast, Darnell (1991a) concluded that zooplankton contributed only 2% of the overall foods for the soft-bottom fish community of the Mississippi-Alabama shelf, and that small epibenthic organisms, infauna, and larger demersal organisms constituted the majority of the food resources for the soft bottom assemblage.

Initial analysis of food web structure (**Figs. 2.1, 2.2**) for the deep reef community further identifies potential pathways of energy flow and trophic links between hard- and soft-bottom habitats that may be useful in identifying changes in trophic structure associated with anthropogenic activities (e.g., the oil/gas exploration and fishing) in areas that are of special concern to resource managers. The current reef fish community exhibits trophic pathways that are primarily supported by planktonic prey, as planktivorous reef fishes dominated all reef biotopes observed during ROV surveys and *R. aurorubens*, a macroplanktivore, was the numerically dominant predator taken by angling and traps.

While generalized carnivores are the most speciose feeding guild on the Pinnacles Reef Tract, they were relatively rare in ROV surveys and collections made during this study. Planktivores (primarily anthiines and the pomacentrid *C. enchrysurus*) numerically dominated all reef biotopes during ROV surveys with benthic carnivores (Gobiidae, *H. bathyphilus*, *D. puellaris*, and *P. umbrosus*) second in importance. Generalized carnivores were represented by *S. phoebe* and *L. eukrines*, but individual taxa made up 2% or less by number on reef biotopes. Food web models based on relative abundance of fishes indicate that the primary pathway of energy to the deep reef community is by direct consumption of copepods, amphipods, fish larvae, decapod larvae, and gelatinous zooplankton that prey on smaller planktonic prey groups in the water column (**Figs. 2.1, 2.2**). Secondary pathways that support benthic carnivores are presumably based on detrital pathways that support infaunal and epifaunal invertebrates. The numerically dominant predators, *R. aurorubens* and *P. pagrus*, feed directly on planktonic and soft-




Figure 2.1. Qualitative food web model for reef fishes of the Pinnacles Reef Tract constructed from stomach content analysis. Position on the food web model is based on relative proportion of pelagic prey (x axis) and trophic position (y axis).




Figure 2.2. Quantitative food web model for reef fishes of the Pinnacles Reef Tract constructed from stomach content analysis. Position on the food web model is based on relative proportion of pelagic prey (x axis) and trophic position (y axis). Relative sizes of nodes (circles) depict abundance of reef fishes based on ROV data analysis and hook and line collections and relative widths of links depicted are based on numerical contribution of prey in the diet of each predator.

bottom organisms, respectively, and do not have diets characterized by reef-associated fishes or invertebrates. *Seriola* spp., *L. campechanus*, and *S. phoebe* prey upon anthiine serranids, and therefore also appear to be supported primarily by pelagic trophic pathways.

While few groupers examined during this study retained identifiable stomach contents, anthiine serranids presumably form a significant part of their diet, as *A. tenuis*, *H. vivanus*, and *P. martinicensis* have been reported from the stomach contents of deepwater groupers (Bullock and Smith 1991). Predation on anthiines would provide a direct link to pelagic food webs.

Trophic pathways within reef systems have been the subject of study since the 1950's (Odum and Odum 1955, Hiatt and Strasburg 1960), yet trophic interactions and mechanisms of energy transfer within the reef fish community are still poorly known (Bohnsack and Sutherland 1985, Nelson and Bortone 1996). In their original work on the trophic structure of coral reefs, Odum and Odum (1955) concluded that the coral reef community at Enewetak Atoll was primarily supported by benthic and zooxanthellate algae, and suggested that incoming plankton was not a significant component of organic matter to the community. Tropical oceans that bathe most coral reefs are oligotrophic, therefore most shallow reef systems might be self-supporting, although more recent studies have identified the important trophic links to zooplankton in tropical coral reef communities (Hamner et al. 1988). Benthic algae typically supports a large population of grazing reef fishes. These herbivores are dominant members of many shallow coral reef systems (Hiatt and Strasburg 1960, Ogden 1982, Ebeling and Hixon 1991), yet are absent from the Pinnacles Reef Tract. Porter and Porter (1977) suggested that coral reef communities support a self-generated food web, and tightly recycle nutrients. Pinnacles deep reef communities are characterized by an abundance of planktivorous fishes and suspension-feeding invertebrates (Hardin et al. 2001), utilizing allochthonous prey resources in the water column to sustain populations, and are therefore very different in trophic structure when compared to many shallow coral reef communities.

The relative importance of plankton in food web structure of reef communities has received limited attention, particularly for more eutrophic warm-temperate reef ecosystems in the Atlantic and Gulf of Mexico. Advection of plankton from open ocean water masses to the continental shelf has been noted (Hopkins et al. 1981). Bottom currents also bring plankton from adjacent shelf waters to the shelf break (Kelly and Bender 2001). Copepods and gelatinous zooplankton are the common grazers of phytoplankton and nanoplankton (<5 μ) in the open water column and may be important in transfer of open water primary production to other components of marine ecosystems such as the benthic fish assemblage (Fourtier et al. 1994, Kleppel et al. 1996). Along a shallow windward reef face in Australia, transport of plankton from the open ocean was considered the major input of organic material to the fish and invertebrate community (Hamner et al. 1988). On temperate reefs in California, over 67% of fishes are obligate or facultative planktivores (Ebeling and Hixon 1991). Planktivorous fishes also dominate fish communities on New Zealand rocky reefs (Kingsford and MacDiarmid 1988, Kingsford 1989). Feeding on plankton is an important trophic pathway in many reef fish

communities, particularly those exposed to strong ocean currents. Planktivory is considered an important route of energy flow to the reef community by two pathways: 1) direct consumption of zooplankton by planktivorous fishes and sessile macroinvertebrates, and 2) consumption of planktivores by larger piscivores or deposition of fecal material to the reef (Hobson 1991). On Indo-Pacific coral reefs, fecal material provides an alternative source of nutrients to reef inhabitants as many fishes ingest fecal material of *Chromis atropectoralis*, a planktivore that releases feces of high energy content (Bailey and Robertson 1982). Hamner et al. (1988) also noted that fecal pellets of planktivores were observed on the reef surface more frequently than zooplankton. *Chromis punctipinnis*, a common member of the reef fish community in California, has been shown to deposit substantial quantities of fecal material to rocky reefs each night (Bray et al. 1981). Many invertebrates (e.g., crabs, shrimps, and brittle stars) rapidly ingest deposited fecal material deposited by planktivorous fishes (Rothans and Miller 1991) and in turn these mobile invertebrates are preyed upon by fishes. Sedberry and Cuellar (1993) discussed the potential role in energy transfer by *R. aurorubens* through release of feces over the reef surface. This pathway may be important in trophic connections between planktivorous reef fishes and benthic invertebrate assemblages in deep reef communities throughout the western Atlantic. Reef-dwelling planktivores that dominate the Pinnacles reef community may supplement the diet of many motile reef invertebrates that are the dominant prey of other resident fishes.

Hard-bottom communities are generally characterized by much higher fish biomass than surrounding, unconsolidated sediments (Sedberry and Van Dolah 1984, Parker and Ross 1986). However, studies of productivity on shallow (<38 m) hard-bottom reef systems along the southeastern U. S. reveal that these reef communities are heterotrophic and not nearly as productive as their tropical coral reef counterparts. Typical productivity/respiration ratios for the southeastern U.S. live bottom reefs average 0.68 versus 1.45 on Indo-Pacific coral reefs (Longhurst and Pauly 1987, Hopkinson et al. 1991). Shallow hard bottom communities require secondary production via pelagic zooplankton or benthic primary production in adjacent sandy habitats as a food source to support reef organisms (Hopkinson et al. 1991). Deep reefs in the Gulf of Mexico should exhibit even lower primary productivity/respiration ratios than shallow water communities. Therefore, dense populations of Pinnacles fishes appear to exceed levels that could be supported solely by locally produced prey resources. Fishes on deep reefs must therefore be “subsidized” by passive transport of prey from surrounding benthic or pelagic habitats on water currents, or via active feeding migrations of predators to adjacent areas of the water column or the underwater landscape (Polis et al. 1997). On shallow coral reef ecosystems, grunts of the genus *Haemulon* migrate away from the reef to forage on benthic invertebrates in adjacent sea grass beds and sand flat areas at night, and form inactive schools over coral reefs by day (Meyer et al. 1983, Parrish 1989). Temperate reef fishes also forage in off-reef habitats and consume invertebrates in neighboring soft-bottom habitats (Posey and Ambrose 1994, Weaver 1996). In both tropical and temperate reef systems, planktivorous reef fishes aggregate at discrete areas and rely on water currents to transport prey resources from the surrounding water column (Bray 1981, Hamner et al. 1988). Passive transport of planktonic prey to resident fishes, or

active migration away from reefs to forage on benthic or pelagic prey resources represents allochthonous trophic subsidies. Spatial subsidies, where primary and secondary productivity of adjacent habitats provides prey resources that are transported to the reef community enhance populations of resident predators, result in increased predator density at levels well above those sustainable from local resources (Polis et al. 1997). Local enhancement of fish populations at the Pinnacles occurs through two major pathways: 1) consumption of plankton by numerically dominant anthiines and *R. aurorubens* as they actively forage into the water column or 2) foraging in adjacent soft-bottom sediments by benthic carnivores such as *P. pagrus*, *S. phoebe*, and labrids.

While small diurnal planktivores, such as *P. martinicensis* and *H. vivanus*, are considered to be an important trophic link between open water plankton and the reef community, resident nocturnal planktivores may play a relatively minor role in the overall energetics of the reef community. Although nocturnal planktivores are widely distributed on reefs, most resident species occur in low abundance (i.e., *P. alta*), and are not considered to be important prey for piscivorous fishes due to their deep bodies and well-developed fin spines (Hobson 1991). Hobson (1991) also noted that reef-dwelling nocturnal planktivores were likely to be less important in transferring energy between open-water communities and the reef ecosystem, since they forage close to the reef on emergent prey at night. Many of the emergent plankters such as amphipods and crab zoea collected at night were common prey in the diet of nocturnal planktivores (Alldredge and King 1977, Porter and Porter 1977). Amphipods, isopods, and a variety of other small crustaceans make up substantial populations of cryptofauna that live within the reef substrata during the day (Klumpp et al. 1988).

However, on the deep reef tract of the Pinnacles, other nocturnal planktivores appear to play an important role in reef trophodynamics. Small midwater species such as *B. cantori*, *T. lathami*, and myctophids are periodically abundant on high profile reefs, and would provide an additional prey resource for nocturnal carnivores. While rarely observed in ROV surveys, *R. aurorubens* appears to be the numerically dominant predator and forages by day and night on plankton (Grimes 1979). Other species that feed on the reef surface at night include *Leiostomus xanthurus*, a species normally associated with soft bottom habitats. Foraging activities of these species may transport considerable energy between hard bottom, soft bottom and pelagic habitats during feeding migrations.

Large predators, including *S. dumerili* and *L. campechanus*, also utilize prey from pelagic, reef, and soft-bottom habitats and provide pathways of energy transfer. The presence of *B. cantori*, *T. lathami* and urochordates in the diet of *L. campechanus* indicates direct links to the pelagic food web, while the presence of soft bottom fishes, including lizardfishes and sea robins in the diet of *S. dumerili* indicates foraging activities in soft bottom habitats. Large predators, including groupers, will often migrate between reef structures and opportunistically consume soft-bottom and pelagic prey (Weaver 1996) and provide conduits of energy flow between reef and off-reef communities.

Potential impacts of the petroleum industry in outer shelf regions may differentially affect pelagic versus benthic feeders if drilling wastes and other discharges influence near-bottom turbidity, sedimentation, and prey availability. Identification of trophic structure may also assist in evaluation of overfishing of large predators and the associated shift in the reef fish community. Shelf-edge reef sites are the primary spawning locations for many commercially important groupers and snappers, and have been heavily fished in recent years (Richards and McGowan 1989, Gilmore and Jones 1992, Coleman et al. 1996, Koenig et al. 2000). Koenig et al. (2000) reported a decrease in large predator abundance (primarily *Mycteroperca phenax* and *S. dumerili*), and attributed this change to increased fishing activity in this region. Based on ROV surveys, *P. martinicensis* and *H. vivanus* comprised 6% (total planktivores made up 6.4%) of the reef fish community in 1980, and 71% (total planktivores ~84%) of the reef fish community in 1997 surveys on the east coast of Florida (Koenig et al. 2000), a pattern that parallels the current reef fish community at the Pinnacles. This increase in lower trophic levels in the deep reef fish community also parallels a reported decline in the mean trophic level of commercial fisheries worldwide (Pauly et al. 1998). The numerical dominance by small planktivorous reef fish taxa in shelf-edge reef communities and *R. aurorubens* in our hook and line samples in the Pinnacles Reef Tract have likely been magnified by removal of piscivores through recent increases in commercial fishing activity in this region and historical fishing practices throughout the northwestern Gulf of Mexico since the mid 1800's. Snyder (2001) discusses the decline in catch rates of the large predatory species that characterize offshore reef ecosystems since the 1970's, and an increase in catch rates of *R. aurorubens* during this same time period. While our observations of trophic structure in the deep reef community closely resemble those observed in the Indo-Pacific (Thresher and Colin 1986), the influence on fishing activity will be difficult to assess in determining a natural community, and the current patterns of reef fish community structure may have been greatly altered by human activity through fishing during the past century.

ACKNOWLEDGMENTS

We thank the following individuals for invaluable collaboration and assistance in the field during research cruises: G. Yeargin, D. DeVries, T. Glancy, P. Schofield, S. Meister, B. Eleby, S. Ross, M. Randall, P. Stevens, S. Szedlmayer, R. McBride, J. Caruso, C. Harper, L. Parker, J. Barichivich, G. Hill, G. Fitzhugh, C. Gurshin, J. Lancaster, C. Furman and M. Zatzoff. We also thank Captain Paul Beaugez and the Crew of the R/V TOMMY MUNRO, and Captain Jerry Love and the Crew of the R/V SUNCOASTER for their logistical support in research cruises. ROV support was provided by L. Horn and T. Potts, National Undersea Research Center (NURC) University of North Carolina-Wilmington (UNCW), and C. Harper, National Oceanographic and Atmospheric Administration/National Marine Fisheries Service (NOAA/NMFS)-Stennis Space Center and NMFS Mississippi laboratories. Additional logistical support was provided by the NURC-UNCW Research Center, NOAA-NMFS, the Marine Resources Research Institute of the South Carolina Department of Natural Resources, the Florida Marine Research Institute of the Florida Fish and Wildlife Conservation Commission, the University of Florida (UF), the University of South Florida, the University of North Carolina at Wilmington, Auburn University, and the Louisiana Universities Marine Consortium (LUMCON). Maps, data, and general information on the Pinnacles region was provided by the Continental Shelf Associates (CSA), I. MacDonald, K. Spring, D. Snyder, and W. Schroeder. Detailed bathymetry maps for this report were provided by J. Gardner and P. Dartnell, USGS-Menlo Park. Research on the Pinnacles Reef Tract was funded by the Outer Continental Shelf Ecosystem Program (K. Sulak, PI) of the U.S. Geological Survey, in Cooperation with the Minerals Management Service (MMS), Gulf of Mexico Region. High-resolution bathymetric mapping was supported by USGS Interactive BRD/GD Science Program (K. Sulak, J. Gardner, and C. Sherwood, Co-PI's) with supplementary support from USGS Geologic Division, NMFS, and MMS. Additional funding and logistical support for doctoral research were provided to D. Weaver by the Walt Disney Conservation Fund, the National Fish and Wildlife Foundation, and the Department of Fisheries and Aquatic Sciences-UF during cruises 2000-01 and 2000-2. G. Brewer provided advice during all phases of this project, and reviewed the earlier drafts of this report. T. Ahlfeld, S. Childs, and G. Boland of the Minerals Management Service provided review and comments of the draft report.

LITERATURE CITED

- Adams, P.B., J.L. Butler, C.H. Baxter, T.E. Laidig, K.A. Dahlin, and W.W. Wakefield. 1995. Populations estimates for Pacific coast groundfishes from video transects and swept area trawls. U.S. Fish. Bull. 93:446-455.
- Allredge, A. and J.M. King. 1977. Distribution, abundance and substrate preferences of demersal reef zooplankton at Lizard Island Lagoon, Great Barrier Reef. Mar. Biol. 41:317-333.
- Bailey, T.G. and D.R. Robertson. 1982. Organic and caloric levels of fish feces relative to its consumption by coprophagous reef fishes. Mar. Biol. 69:45-50.
- Bloom, S.A. 1981 Similarity indices in community studies: potential pitfalls. Mar. Ecol. Prog. Ser. 5:125-128.
- Boesch, D.F. 1977. Application of numerical classification in ecological investigations of water pollution. EPA Ecological Research Series. EPA-600/3-77-033, 114 pp.
- Bohnsack, J. A. and D. L. Sutherland. 1985. Artificial reef research: a review with recommendations for future priorities. Bull. Mar. Sci. 37(1):11-39.
- Bohnsack, J.A., D.E. Harper, D.B. McClellan, D.L. Sutherland, and M.W. White. 1997. Resource surveys of fishes within Looe Key National Marine Sanctuary. NOAA Tech. Memo. NOS-MEMD-5. 108 pp.
- Bohnsack, J.A., D.B. McClellan, D.E. Harper, G.S. Davenport, G.J. Konoval, A.M. Eklund, J.P. Contillo, S.K. Bolden, P.C. Fischel, G.S. Sandorf, J.C. Javech, M.W. White, M.H. Pickett, M.W. Hulsbeck, J.L. Tobias, J.S. Ault, G.A. Meester, S.G. Smith, and J. Luo. 1999. Baseline data for evaluating reef fish populations in the Florida Keys, 1979-1998. NOAA Tech. Memo. NMFS-SEFSC-427. 61 pp.
- Boland, G.S., B.J. Gallaway, J.S. Baker, and G.S. Lewbel. 1983. Ecological effects of energy development on reef fish of the Flower Garden banks. Final Report to National Marine Fisheries Service, Galveston Laboratory. LGL Ecological Research Associates, 466 pp.
- Bray, R.N. 1981. Influence of water currents and zooplankton densities on daily foraging movements of blacksmith, *Chromis punctipinnis*, a planktivorous reef fish. U.S. Fish. Bull. 78:829-841.
- Bray, R.N. A.C. Miller, and G.G. Geesey 1981. The fish connection: a trophic link between planktonic and rocky reef communities? Science 214:204-205.
- Bright, T.J. and C.W. Cashman. 1974. Fishes, pp. 341-409. In: T. J. Bright and L. H. Pequegnat (eds.), Biota of the West Flower Garden Bank. Gulf Publishing Company, Houston, TX. 435 pp.
- Bright, T.J., J.W. Tunnell, L.H. Pequegnat, T.E. Burke, C.W. Cashman, D.A. Cropper, J.P. Ray, R.C. Tresslar, J. Teerling, and J.B. Willis. 1974. Biotic zonation on the West Flower Garden Bank. Pp. 4-52. In: T.J. Bright and L.H. Pequegnat (eds.), Biota of the West Flower Garden Bank. Gulf Publishing Company, Houston, TX. 435 pp.
- Bright, T. J., G. P. Kraemer, G. A. Minnery, and S. T. Viada. 1984. Hermatypes of the Flower Garden Banks, northwestern Gulf of Mexico: a comparison to other western Atlantic reefs. Bull. Mar. Sci. 34:461-76.
- Brooks, G. R., and L. J. Doyle. 1991. Geologic development and depositional history of the Florida Middle Ground: a mid-shelf, temperate-zone reef system in the

- northeastern Gulf of Mexico. From shoreline to abyss: contributions in marine geology in honor of Francis Parker Shepard. Ed. R. H. Osborne, 189-203.
- Brooks, J.M., ed. 1991. Mississippi-Alabama Marine Ecosystem Study: Data Summary and Synthesis. Volume II: Technical Narrative. U.S. Department of the Interior, Minerals Management Service, Gulf of Mexico OCS Regional Office, New Orleans, LA. OCS Study MMS-91-063. 862 pp.
- Bullis, H.R., Jr. and J.R. Thompson. 1965. Collections made by the exploratory fishing vessels Oregon, Silver Bay, Combat and Pelican during 1956-1960 in the southwestern Atlantic. U.S. Fish & Wildlife Serv. Sci. Rept. No. 510, 310 pp.
- Bullock, L.H. and M.F. Godcharles. 1982. Range extensions for four sea basses (Pisces: Serranidae) from the eastern Gulf of Mexico, with color notes on *Hemanthias lepturus* (Ginsburg). NE Gulf Sci. 5(2):53-56.
- Bullock, L.H. and G.B. Smith. 1991. Seabasses (Pisces: Serranidae). Memoirs of the Hourglass Cruises, Fla. Mar. Res. Inst. No. 8, 243 pp.
- Butler, J.L., W.W. Wakefield, P.B. Adams, B.H. Robison, and C.H. Baxter. 1991. Application of line transect methods to surveying demersal communities with ROVs and manned submersibles. Proc. OCEANS '91 Conf., Honolulu, HI. 1:689-696.
- Cashman, C.W. 1973. Contributions to the ichthyofaunas of the West Flower Garden reef and other reef sites in the Gulf of Mexico and western Caribbean. Ph.D. dissertation. Texas A&M Univ. College Station, TX. 247 pp.
- Christensen, V. 1995. A model for trophic interactions in the North Sea in 1981, the Year of the Stomach. Dana Rept. 11(1):1-28.
- Clarke, D. 1981. Ichthyofauna. pp. 117-124. In: T.S. Hopkins, W. Schroeder, T. Hilde, L. Doyle, and J. Steinmetz (eds.), Northern Gulf of Mexico Topographic Features Study. Final Report. Vol. 5. Texas A&M University Tech. Rept. No. 81-2-T, BLM Contract No. AA551-CT8-35, 150 pp.
- Clarke, D.G. 1986. Visual census of fish populations at the Florida Middle Ground. NE Gulf Sci. 8(1):65-81.
- Cohen, D.M. and D.L. Pawson. 1977. Observations from the DSRV ALVIN on populations of benthic fishes and selected larger invertebrates in and near DWD-106. U.S. Department of Commerce, NOAA, Dumpsite Evaluation Rept. 77-1, 2:423-450.
- Coleman, F.C., C.C. Koenig, and L.A. Collins. 1996. Reproductive styles of shallow-water grouper (Pisces: Serranidae) in the eastern Gulf of Mexico and the consequences of fishing spawning aggregations. Environ. Biol. Fishes 47:129-141.
- Colin, P. 1974. Observation and collection of deep-reef fishes off the coasts of Jamaica and British Honduras (Belize). Mar. Biol. 24:29-38.
- Colin, P. 1976. Observations of deep-reef fishes in the Tongue-of-the-Ocean, Bahamas. Bull. Mar. Sci. 26(4):603-604.
- Continental Shelf Associates, Inc. (CSA). 1992. Mississippi-Alabama shelf pinnacle trend habitat mapping study OCS Study/MMS 92-0026. U.S. Department of the Interior, Minerals Management Service, New Orleans, LA, 75 pp.
- Continental Shelf Associates, Inc. and Texas A&M University, Geochemical and Environmental Research Group (CSA and TAMU). 2001. Mississippi/Alabama

- Pinnacle Trend Ecosystem Monitoring, Final Synthesis Report. U.S. Department of the Interior, Geological Survey, USGS/CR-2000-007 and Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA, OCS Study MMS 2001-080. 415 pp + apps.
- Crabtree, R.E., H.J. Carter, and J.A. Musick. 1991. The comparative feeding ecology of temperate and tropical deep-sea fishes from the western north Atlantic. *Deep-Sea Res.* 38:1277-1298.
- Cuker, B.E. 1993. Suspended clay alter trophic interactions in the plankton. *Ecology* 74(3):944-953.
- Darcy, G.H. and E.J. Gutherz. 1984. Abundance and density of demersal fishes on the West Florida shelf, January 1978. *Bull. Mar. Sci.* 34(1): 81-105.
- Darnell, R.M. 1991a. Demersal Fish Food Analysis. Chapter 9, pp. 9-1 to 9-76. In: J.M. Brooks (ed.), *Mississippi-Alabama Marine Ecosystem Study: Data Summary and Synthesis. Volume II: Technical Narrative.* OCS Study MMS-91-0063. U.S. Department of the Interior, Minerals Management Service, Gulf of Mexico OCS Regional Office, New Orleans, LA. 862 pp.
- Darnell, R.M. 1991b. Summary and Synthesis. Chapter 15, pp. 15-1 to 15-145. In: J.M. Brooks (ed.), *Mississippi-Alabama Marine Ecosystem Study: Data Summary and Synthesis. Volume II: Technical Narrative.* OCS Study MMS-91-0063. U.S. Department of the Interior, Minerals Management Service, Gulf of Mexico OCS Regional Office, New Orleans, LA. 862 pp.
- Darnell, R.M. and J.A. Kleypas. 1987. *Eastern Gulf Bio-atlas.* U.S. Dept. Interior, Minerals Management Service. OCS Study MMS-86-0041, 548 pp.
- Dennis, G.D. and T.J. Bright. 1988a. Reef fish assemblages on hard banks in the northwestern Gulf of Mexico. *Bull. Mar. Sci.* 43(2):280-307
- Dennis, G.D. and T.J. Bright. 1988b. New records of fishes in the northwestern Gulf of Mexico, with notes on some rare species. *NE Gulf Sci.* 10(1):1-18.
- Dodrill, J., C.S. Manooch, and A.B. Manooch. 1993. Food and feeding behavior of adult snowy grouper, *Epinephelus niveatus* (Valenciennes) (Pisces: Serranidae), collected off the central North Carolina coast with ecological notes on major food groups. *Brimleyana* 19:101-135.
- Ebeling, A.W. and M.A. Hixon. 1991. Tropical and temperate reef fishes: comparison of community structures, pp. 509-563. In: P.F. Sale (ed.), *The Ecology of Fishes on Coral Reefs.* Academic Press, San Diego, CA. 754 pp.
- Felley, J.D. and M. Vecchione. 1995. Assessing habitat use by nekton on the continental slope using archived videotapes from submersibles. *U.S. Fish. Bull.* 93:262-273.
- Felley, J.D., M. Vecchione, G.R. Gaston, and S.M. Felley. 1989. Habitat selection by demersal nekton: analysis of videotape data. *NE Gulf Sci.* 10(2):69-84.
- Fourtier, L., J. Le Fevre, and L. Legendre. 1994. Export of biogenic carbon to fish and to the deep ocean: the role of large planktonic microphages. *J. Plank. Res.* 16(7):809-839.
- Friedlander, A.M. and J.D. Parrish. 1998. Habitat characteristics affecting fish assemblages on a Hawaiian coral reef. *J. Exp. Mar. Biol. Ecol.* 224:1-30.
- Gallagher, E.D. 1998. COMPAAH program and documentation. 54 pp. www.es.umb.edu/edgwebp.htm.

- Gardner, J.V., K.J. Sulak, P. Dartnell, L. Hellequin, B. Calder, and L.A. Mayer. 2000. Cruise Report RV Ocean Surveyor Cruise O-1-00-GM. The Bathymetry and Acoustic Backscatter of the Pinnacles Area, Northern Gulf of Mexico May 23, through June 10, 2000 Venice, LA to Venice, LA. U.S. Geological Survey Open-File Report 00-350, 36 pp. <http://geopubs.wr.usgs.gov/open-file/of00-350>.
- Gardner, J.V., P. Dartnell, K.J. Sulak, B. Calder, and L. Hellequin. 2001. Physiography and Late Quaternary-Holocene processes of the northeastern Gulf of Mexico outer continental shelf off Mississippi and Alabama. *Gulf of Mexico Sci.* 19(2): 132-157.
- Gardner, J.V., P. Dartnell, and K.J. Sulak. 2002. Multibeam mapping of the Pinnacles Region, Gulf of Mexico. U.S. Geological Survey Open-File Report 02-006, <http://geopubs.wr.usgs.gov/open-file/of02-006/>.
- Gilmore, R.G. 1977. Fishes of the Indian River Lagoon and adjacent waters, Florida. *Bull. Fla. State Mus., Biol. Sci.* 22(3):101-148.
- Gilmore, R.G. and R.J. Jones. 1992. Color variation and associated behavior in the epinepheline groupers *Mycteroperca microlepis* (Good and Bean) and *M. phenax* Jordan and Swain. *Bull. Mar. Sci.* 51:83-103.
- Gittings, S.R. and E. Hickerson, eds. 1998. Dedicated Issue: Flower Garden Banks National Marine Sanctuary. *Gulf of Mexico Sci.* 16(2):128-237.
- Gittings, S.R., T.J. Bright, and W.W. Schroeder. 1991. Topographic features characterization-biological. Chapter 13, pp. 13-1 to 13-117. In: J.M. Brooks (eds.), *Mississippi-Alabama Marine Ecosystem Study: Data Summary and Synthesis. Volume II: Technical Narrative.* OCS Study MMS-91-0063. U.S. Department of the Interior, Minerals Management Service, Gulf of Mexico OCS Regional Office, New Orleans, LA. 862 pp.
- Gittings, S.R., T.J. Bright, W.W. Schroeder, W.W. Sager, J.S. Laswell and R. Rezak. 1992a. Invertebrate assemblages and ecological controls on topographic features in the northeast Gulf of Mexico. *Bull. Mar. Sci.* 50(3):435-455.
- Gittings, S.R., G.S. Boland, K.J.P. Deslarzes, D.K. Hagman and B.S. Holland. 1992b. Long-term monitoring at the East and West Flower Garden Banks. OCS Study MMS 92-0006. U.S. Department of the Interior, Minerals Management Service, New Orleans, LA, 206 pp.
- Goedicke, T.R. 1955. Origin of the pinnacles on the continental shelf and slope of the Gulf of Mexico. *Tex. J. Sci.* 6:149-159.
- Grassle, J.F., H.L. Sanders, R.R. Hessler, G.T. Rowe, and T. McLellen. 1975. Pattern and zonation: a study of the bathyal megafauna using the research submersible Alvin. *Deep-Sea Res.* 22:457-481.
- Grimes, C.B. 1979. Diet and feeding ecology of the vermilion snapper, *Rhomboplites aurorubens* (Cuvier) from North Carolina and South Carolina waters. *Bull. Mar. Sci.* 29:53-61.
- Grimes, C.B., C.S. Manooch, and G.R. Huntsman. 1982. Reef and rock outcropping fishes of the outer continental shelf of North Carolina and South Carolina, and ecological notes on the red porgy and vermilion snapper. *Bull. Mar. Sci.* 32(10):277-289.

- Hamner, W.M., M.S. Jones, J.H. Carleton, I.R. Hauri and D. McB. Williams. 1988. Zooplankton, planktivorous fish, and water currents on a windward reef face: Great Barrier Reef, Australia. *Bull. Mar. Sci.* 42(3):459-479.
- Hancock, K.M. 1997. Paleocology of late Pleistocene carbonate buildups on the Mississippi-Alabama continental shelf. Unpublished Master's thesis. Univ. Alabama, Tuscaloosa, AL. 93 pp.
- Hardin, D.D., K.D. Spring, S.T. Viada, A.D. Hart, B.D. Graham, and M.B. Peccini. 2001. Hard Bottom Communities. Chapter 7, pp. 209-268. In: Continental Shelf Associates, Inc. and Texas A&M University, Geochemical and Environmental Research Group. (eds.), Mississippi/Alabama Pinnacle Trend Ecosystem Monitoring, Final Synthesis Report. U.S. Department of the Interior, Geological Survey, USGS/CR-2000-007 and Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA, OCS Study MMS 2001-080. 415 pp + apps.
- Hastings, R.W. 1979. The origin and seasonality of the fish fauna on a new jetty in the northeastern Gulf of Mexico. *Bull. Fla. State Mus., Biol. Sci. No. 24*, 117 pp.
- Hastings, R.W., L.H. Ogren, and M.T. Mabry. 1976. Observations on the fish fauna associated with offshore platforms in the northeastern Gulf of Mexico. *U.S. Fish. Bull.* 74(2):387-402.
- Hecker, B. 1987. Epifauna on the U.S. South Atlantic slope and rise, pp. 208-287. In: J.A. Boehme et al. (eds.), Study of biological processes on the U.S. South Atlantic slope and rise. Phase 2 Final Report, U.S. Department of the Interior, Minerals Management Service, MMS86-0096.
- Hiatt, R.W. and D.W. Strasburg. 1960. Ecological relationships of the fish fauna on coral reefs of the Marshall Islands. *Ecol. Monogr.* 30(1):65-126.
- Hobson, E.S. 1982. The structure of fish communities on warm-temperate and tropical reefs. pp. 160-166 In: G.R. Huntsman, W.R. Nicholson, and W.W. Fox (eds.), The biological basis for reef fishery management. NOAA Tech. Memo. NMFS-SEFC-80.
- Hobson, E.S. 1991. Trophic relations of fishes specialized to feed on zooplankters above coral reefs, pp. 69-95. In: P.F. Sale (ed.), The ecology of fishes on coral reefs. Academic Press, San Diego, CA. 754 pp.
- Hoese, H.D. and R.H. Moore. 1998. Fishes of the Gulf of Mexico. 2nd Edition. Texas A&M Univ. Press, College Station, TX, 422 pp.
- Hopkins, T.L., D.M. Milliken, L.M. Bell, E.J. McMichael, J.J. Heffernan, and R.V. Cano. 1981. The landward distribution of oceanic plankton and micronekton over the west Florida continental shelf as related to their vertical distribution. *J. Plank. Res.* 3:645-658.
- Hopkinson, C.S., R.D. Fallon, B. Jansson, and J.P. Schubauer. 1991. Community metabolism and nutrient cycling at Gray's reef, a hard bottom habitat in the Georgia Bight. *Mar. Ecol. Prog. Ser.* 73:105-120
- Hyslop, E.J. 1980. Stomach contents analysis - a review of methods and their application. *J. Fish Biol.* 17:411-429.
- Jarvis, N.D. 1935. Fishery for red snappers and groupers in the Gulf of Mexico. Bureau Fisher. Invest. Rept. No. 26, 29 pp.
- Kelly, F.J. and L.C. Bender. 2001. Physical Oceanography/Hydrography. Chapter 6, pp 127-208, In: Continental Shelf Associates, Inc. and Texas A&M University,

- Geochemical and Environmental Research Group (eds.), Mississippi/Alabama Pinnacle Trend Ecosystem Monitoring, Final Synthesis Report. U.S. Department of the Interior, Geological Survey, USGS/CR-2000-007 and Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA, OCS Study MMS 2001-080. 415 pp + apps.
- Kingsford, M.J. 1989. Distribution patterns of planktivorous reef fish along the coast of northeastern New Zealand. *Mar. Ecol. Prog. Ser.* 54:13-24.
- Kingsford, M.J. and A.B. MacDiarmid. 1988. Interactions between planktivorous reef fish and zooplankton in temperate waters. *Mar. Ecol. Prog. Ser.* 48:103-117.
- Kleppel, G.S., C.A. Burkhardt, K. Carter, and C. Tomas. 1996. Diets of calanoid copepods on the west Florida continental shelf: relationships between food concentration, food composition and feeding activity. *Mar. Biol.* 127:209-217.
- Klumpp, D.W., A.D. McKinnon, and C.N. Mundy. 1988. Motile cryptofauna of a coral reef: abundance, distribution and trophic potential. *Mar. Ecol. Prog. Ser.* 45:95-108.
- Koenig, C.C., F.C. Coleman, C.B. Grimes, G.R. Fitzhugh, K.M. Scanlon, C.T. Gledhill, and M. Grace. 2000. Protection of fish spawning habitats for the conservation of warm-temperate reef-fish fisheries of shelf-edge reefs of Florida. *Bull. Mar. Sci.* 66(3):593-616.
- Kotrschal, K. and W. W. Reynolds. 1982. Behavioral ecology of northern Adriatic reef fishes in relation to seasonal temperature regimes. *Contrib. Mar. Sci.* 25:99-106.
- Laswell, J.S., W. Sager, W.W. Schroeder, R. Rezak, K.S. Davis, and E.G. Garrison. 1990. Atlas of high-resolution geophysical data. Mississippi-Alabama Marine Ecosystems Study. U.S. Department of Interior, Minerals Management Service, New Orleans, LA, OCS Study/MMS 90-0000. 42 pp.
- Lindquist, D.G. and I.E. Clavijo. 1993. Quantifying deep reef fishes from a submersible and notes on a live collection and diet of red barbier, *Hemanthias vivanus*. *J. Elisha Mitchell Sci. Soc.* 109(3):135-140.
- Lindquist, D.G., L.B. Calhoun, I.E. Clavijo, M.H. Posey, S.K. Bolden, L.A. Pike, S.W. Burk, and P.A. Cardullo. 1994. Reef fish stomach contents and prey abundance on reef and sand substrata associated with adjacent artificial and natural reefs in Onslow Bay, North Carolina. *Bull. Mar. Sci.* 55:308-318.
- Longhurst, A.R. and D. Pauly. 1987. *Ecology of Tropical Oceans*. Academic Press, San Diego, CA. 407 pp.
- Ludwick, J.C. and W.R. Walton. 1957. Shelf edge, calcareous prominences in the northeastern Gulf of Mexico. *Bull. Amer. Assoc. Petrol. Geol.* 41(9):2054-2101.
- MacArthur, R.H. and E.O. Wilson. 1967. *The theory of island biogeography*. Princeton Univ. Press., Princeton, NJ. 203 pp.
- MacDonald, I.R. and M.B. Peccini. 2001. GIS and Microhabitat Studies. Chapter 9, pp 299-326. In: Continental Shelf Associates, Inc. and Texas A&M University, Geochemical and Environmental Research Group (eds.), Mississippi/Alabama Pinnacle Trend Ecosystem Monitoring, Final Synthesis Report. U.S. Department of the Interior, Geological Survey, USGS/CR-2000-007 and Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA, OCS Study MMS 2001-080. 415 pp + apps.

- Manooch, C.S. 1977. Food habits of the red porgy, *Pagrus pagrus* Linnaeus (Pisces: Sparidae) from North Carolina and South Carolina. *Bull. Mar. Sci.* 27:776-787.
- Matheson, R.H., G.R. Huntsman, and C.S. Manooch. 1986. Age, growth, mortality, food and reproduction of the scamp, *Mycteroperca phenax*, collected off North Carolina and South Carolina. *Bull. Mar. Sci.* 38(2):300-312.
- McBride, R.S. and R.W. Able. 1998. Ecology and fate of butterflyfishes, *Chaetodon* spp. in the temperate western North Atlantic. *Bull. Mar. Sci.* 63(2):401-416.
- McEachran, J.D. 1991. Distribution of fishes. Chapter 8, pp. 8-1 to 8-81. In: J.M. Brooks (ed.), *Mississippi-Alabama Marine Ecosystem Study: Data Summary and Synthesis. Volume II: Technical Narrative*. U.S. Department of the Interior, Minerals Management Service, Gulf of Mexico OCS Regional Office, New Orleans, LA. OCS Study MMS-91-0063. 862 pp.
- Messing, C.G., A.C. Neumann, and J.C. Lang. 1990. Biozonation of deep-water lithoherms and associated hardgrounds in the northeastern Straits of Florida. *Palaios* 5:15-33.
- Meyer, J.L., E.T. Schultz, and G.S. Helfman. 1983. Fish schools: an asset to corals. *Science* 220:1047-1049.
- Moore, D.R. and H.R. Bullis. 1960. A deep-water coral reef in the Gulf of Mexico. *Bull. Mar. Sci.* 10(1):125-128.
- Mullaney, M.D. Jr. and L.D. Gale. 1995. Ecomorphological relationships in ontogeny: anatomy and diet in gag, *Mycteroperca microlepis* (Pisces: Serranidae). *Copeia* 1996(1):167-180.
- Nelson, B.D. and S.A. Bortone. 1996. Feeding guilds among artificial reef fishes in the northern Gulf of Mexico. *Gulf of Mexico Sci.* 14(2):66-80.
- Odum, H.T. and E.P. Odum. 1955. Trophic structure and productivity of a windward coral reef community on Eniwetok Atoll. *Ecol. Monogr.* 25(3):291-320.
- Ogden, J.C. 1982. Fisheries management and the structure of coral reef fish communities, pp. 147-159. In: G.R. Huntsman, W.R. Nicholson, and W.W. Fox (eds.), *The biological basis for reef fishery management*. NOAA Tech. Memo. NMFS-SEFC-80.
- Parker, R.O. and S.W. Ross. 1986. Observing reef fishes from submersibles off North Carolina. *NE Gulf Sci.* 8:31-49.
- Parrish, J.D. 1989. Fish communities of interacting shallow-water habitats in tropical oceanic regions. *Mar. Ecol. Prog. Ser.* 58:143-160.
- Parsons, T.R. 1992. The removal of marine predators by fisheries and the impact of trophic structure. *Mar. Poll. Bull.* 26:51-53.
- Parsons, T.R. 1996. The impact of industrial fisheries on the trophic structure of ecosystems, pp. 352-357. In: G.A. Polis and K.O. Winemiller (eds.), *Food Webs: Integration of Patterns and Dynamics*. Kluwer Academic Publishers, Boston, MA.
- Pattengill, C.V., B.X. Semmens, and S.R. Gittings. 1997. Reef fish trophic structure at the Flower Gardens and Stetson Bank, NW Gulf of Mexico. *Proc. 8th Inter. Coral Reef Symp.* Panama, 1:1023-1028.
- Pauly, D., V. Christensen, J. Dalsgaard, R. Froese, and F. Torres. 1998. Fishing down marine food webs. *Science* 279:860-863.

- Polis, G.A., W.B. Anderson, and R.D. Holt. 1997. Toward an integration of landscape and food web ecology: The dynamics of spatially subsidized food webs. *Ann. Rev. Ecol. Syst.* 28:289-316.
- Porter, J.W. and K.G. Porter. 1977. Quantitative sampling of demersal plankton migrating from different coral reef substrates. *Limnol. Oceanogr.* 22(3):553-556.
- Posey, M.H. and W.G. Ambrose. 1994. Effects of proximity to an offshore hard-bottom reef on infaunal abundances. *Mar. Biol.* 118:745-753.
- Powell, D., L. Dwinell, and S. Dwinell. 1972. An annotated listing of the fish reference collection at the Florida Department of Natural Resources Marine Research Laboratory. Fla. Dept. Nat. Resour. Mar. Res. Lab. Spec. Sci. Rept., No. 36, 179 pp.
- Putt, R.E., D.A. Gettleson, and N.W. Phillips. 1986. Fish assemblages and benthic biota associated with natural hard-bottom areas in the northwestern Gulf of Mexico. *NE Gulf Sci.* 8(1):51-63.
- Randall, J.E. 1961. A technique for fish photography. *Copeia* 1961(1):241-242.
- Randall, J.E. 1967. Food habits of reef fishes of the West Indies. *Stud. Trop. Oceanogr.* 5:665-847.
- Rezak, R., T.J. Bright, and D.W. McGrail. 1985. Reefs and banks of the northwestern Gulf of Mexico: their geological, biological, and physical dynamics. Wiley and Sons, New York, NY. 259 pp.
- Rezak, R., S.R. Gittings and T.J. Bright. 1990. Biotic assemblages and ecological controls on reefs and banks of the northwest Gulf of Mexico. *Amer. Zool.* 30:23-35.
- Rice, A.L., R.G. Aldred, D.S.M. Billet, and M.H. Thurston. 1979. The combined use of an epibenthic sledge and a deep-sea camera to give quantitative relevance to macro-benthos samples. *Ambio Spec. Rept.* 6:59-72.
- Richards, W.J. and M.F. McGowan. 1989. Biological productivity in the Gulf of Mexico: identifying the causes of variability in fisheries, pp. 287-325. In: K. Sherman and L.M. Alexander (eds.), *Biomass Yields and Geography of Large Marine Ecosystems*. Westview Press, Boulder, CO.
- Robertson, D.R. and J.M. Sheldon. 1979. Competitive interactions and the availability of sleeping sites for a diurnal coral reef fish. *J. Exp. Mar. Biol. Ecol.* 40:285-298.
- Robins, C.R. and G.C. Ray. 1986. A field guide to Atlantic coast fishes of North America. Houghton Mifflin Co., New York, NY. 354 pp.
- Robins, C.R., R.M. Bailey, C.E. Bond, J.R. Brooker, E.A. Lachner, R.N. Lea, and W.B. Scott. 1991. Common and scientific names of fishes from the United States and Canada. *Amer. Fish. Soc. Spec. Publ. No. 20*, 183 pp.
- Rogers, A. D. 1999. The biology of *Lophelia pertusa* (Linnaeus 1758) and other deep-water reef-forming corals and impacts from human activities. *Int. Rev. Hydrobiol.* 84:315-406.
- Ross, J.L. 1982. Feeding habits of the gray tilefish, *Caulolatilus microps* (Goode and Bean, 1878) from North and South Carolina waters. *Bull. Mar. Sci.* 32:448-454.
- Rothans, T.C. and A.C. Miller. 1991. A link between biologically imported particulate organic nutrients and the detritus food web in reef communities. *Mar. Biol.* 110:145-150.

- Sager, W.W., W.W. Schroeder, J.S. Laswell, K.S. Davis, R. Rezak, and S.R. Gittings. 1992. Mississippi-Alabama outer continental shelf topographic features formed during the late Pleistocene-Holocene transgression. *Geo-Marine Lett.* 12:41-48.
- Sager, W.W. and W.W. Schroeder. 2001. Geological Characterization. Chapter 3, pp 23-94. In: Continental Shelf Associates, Inc. and Texas A&M University, Geochemical and Environmental Research Group (eds.), Mississippi/Alabama Pinnacle Trend Ecosystem Monitoring, Final Synthesis Report. U.S. Department of the Interior, Geological Survey, USGS/CR-2000-007 and Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA, OCS Study MMS 2001-080. 415 pp + apps.
- Sedberry, G.R. and N. Cuellar. 1993. Planktonic and benthic feeding by the reef-associated vermilion snapper, *Rhomboplites aurorubens* (Teleostei, Lutjanidae). *U.S. Fish. Bull.* 91:699-709.
- Sedberry, G.R. and R.F. Van Dolah. 1984. Demersal fish assemblages associated with hard bottom habitat in the South Atlantic Bight of the U.S.A. *Environ. Biol. Fishes* 11(4):241-258.
- Shepard, F.P. 1937. "Salt" domes related to Mississippi submarine troughs. *Bull. Amer. Assoc. Petrol. Geol.* 48:1349-1362.
- Sheridan, P.F. and D.L. Trimm. 1983. Summer foods of Texas coastal fishes relative to age and habitat. *U.S. Fish. Bull.* 81(3):643-647.
- Shipp, R.L. and T.S. Hopkins. 1978. Physical and biological observations of the northern rim of the Desoto Canyon made from a research submersible. *NE Gulf Sci.* 2(2):113-121.
- Smith, G.B. 1976. Ecology and distribution of eastern Gulf of Mexico fishes. Fla. Dept. Nat. Resour. Mar. Res. Publ. No. 19, 78 pp.
- Smith, G.B., H.M. Austin, S.A. Bortone, R.W. Hastings, and L.A. Ogren. 1975. Fishes of the Florida Middle Ground with comments on ecology and zoogeography. Fla. Dept. Nat. Resour. Mar. Res. Publ. No. 9, 14 pp.
- Smith-Vaniz, W.F., B.B. Collette, and B.E. Luckhurst. 1999. Fishes of Bermuda. *Amer. Soc. Ichthyol. & Herpetol. Spec. Publ. No. 4*, 424 pp.
- Snyder, D.B. 2001. Fish Communities. Chapter 8, pp 269-298. In: Continental Shelf Associates and in: Continental Shelf Associates, Inc. and Texas A&M University, Geochemical and Environmental Research Group (eds.), Mississippi/Alabama Pinnacle Trend Ecosystem Monitoring, Final Synthesis Report. U.S. Department of the Interior, Geological Survey, USGS/CR-2000-007 and Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA, OCS Study MMS 2001-080. 415 pp + apps.
- Sokal, R.R. and F.J. Rohlf. 1995. *Biometry*. 3rd edition. Freeman & Co., New York, NY, 887 p.
- Sonnier, F., H.D. Hoese, and J. Teerling. 1976. Observations on the offshore reef and platform fish fauna of Louisiana. *Copeia* 1976(1):105-111.
- Springer, S. and H.R. Bullis, Jr. 1956. Collections made by the "Oregon" in the Gulf of Mexico. Lists of crustaceans, mollusks, and fishes identified from collections made by the exploratory fishing vessel "Oregon" in the Gulf of Mexico and adjacent seas 1950 through 1955. *U.S. Fish & Wildlife Serv. Sci. Rept. No. 196*, 134 pp.

- Springer, V.G. and K.D. Woodburn. 1960. An ecological study of the fishes of the Tampa Bay area. Fla. State Board Conserv. Mar. Lab. Prof. Pap. Ser. No. 1. 104 pp.
- Starck, W.A. II. 1971. Biology of the gray snapper, *Lutjanus griseus* (Linnaeus), in the Florida Keys. Stud. Trop. Oceanogr. 10:11-150.
- Sulak, K.J. and S.W. Ross. 1993. Analysis of submersible videotapes for demersal fish faunas from the continental slope in "The Point" region, In: R.J. Diaz and J.A. Blake (eds.), Benthic processes in an unusual area of the U.S. Atlantic Continental Slope off Cape Hatteras. U.S. Department of the Interior, Minerals Management Service, MMS Report 30672. Appendix.
- Sulak, K.J. and S.W. Ross. 1996. Lilliputian bottom fish fauna of the Hatteras middle continental slope. J. Fish Biol. 49(Supplement A):91-113.
- Syms, C. and G.P. Jones. 2000. Disturbance, habitat structure, and the dynamics of a coral-reef fish community. Ecology 8(1):2714-2729.
- Thompson, M.J., W.W. Schroeder, and N.W. Phillips. 1999. Ecology of live bottom habitats of the northeastern Gulf of Mexico: a community profile. U.S. Department of the Interior, U.S. Geological Survey, Biological Resources Division, USGS/BRD/CR-1999-001 and Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA, OCS Study MMS 99-004. 74 pp.
- Thresher, R. E. and P. L. Colin. 1986. Trophic structure, diversity and abundance of fishes of the deep reef (30-300m) at Enewetak, Marshall Islands. Bull. Mar. Sci. 38:253-272.
- Trowbridge, A.C. 1930. Building of Mississippi Delta. Bull. Amer. Assoc. Petrol. Geol. 14:867-901.
- Uzmann, J.R., R.A. Cooper, R.B. Theroux, and R.L. Wigley. 1977. Synoptic comparison of three sampling techniques for estimating abundance and distribution of selected megafauna: submersible versus camera sled versus otter trawl. Mar. Fish. Rev. 39:11-19.
- Walls, J.G. 1975. Fishes of the northern Gulf of Mexico. TFH Publ., Neptune City, NJ. 432 pp.
- Weaver, D. 1996. Feeding ecology and ecomorphology in three sea basses (Pisces: Serranidae) in the northeastern Gulf of Mexico. M.S. Thesis, University of Florida, Gainesville, FL. 94 pp.
- Williams, J.T. and R.L. Shipp. 1980. Observations on fishes previously unrecorded or rarely encountered in the northeastern Gulf of Mexico. NE Gulf Sci. 4(1):17-27.
- Winemiller, K.O. 1990. Spatial and temporal variation in tropical fish trophic networks. Ecol. Monogr. 60(3):331-367.
- Wolda, H. 1981. Similarity indices, sample size, and diversity. Oecologia 50:296-302.

APPENDIX A

Summary of ROV surveys from 1997-2001

Legend:

A=Alabama Alps
C=Cat's Paw Reef
D=Double Top Reef
CS=Corkscrew Reef
C5=CMEP Monitoring Site 5
F=Far Tortuga Reef
L=Ludwick&Walton Pinnacle
P=Porgy Reef
PF=Patch Reef Field
R=Roughtongue Reef
S=Shark Reef
SC=Scamp Reef
SM=Solitary Mound
T=Triple Top Reef
Y=Yellowtail Reef

Appendix A1. Summary of reef-associated fishes observed during 1997 (97-01) ROV surveys by habitat. Data are expressed as number of fish observed per minute of video for each site. ND-No Data and (n)-night survey.

Station No.	04	08	09	11	12	20	22	27	33	35	36	44	45	50
Base Depth (m)	68	68	78	78	78	78	106	104	110	110	110	77	71	72
Study Site	Y	Y	R	R	R	C	L-A	L-B	L-C	L-1	L-1	PF	SM-1	SM-2
Taxa														
<i>Antigonia capros</i>	-	-	-	-	-	-	-	-	-	0.05	-	-	-	-
Apogonidae	-	0.09	-	-	-	-	-	-	-	-	-	-	-	-
<i>Apogon pseudomaculatus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Aulostomus maculatus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Bodianus pulchellus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Bregmaceros cantori</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Canthigaster rostrata</i>	0.14	-	0.08	0.10	-	-	-	-	-	-	-	-	0.04	-
Carangidae	-	-	-	-	-	-	-	-	-	0.05	-	-	-	-
<i>Centropristis ocyurus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Chaetodon aya</i>	-	0.09	-	0.36	0.09	-	0.03	-	0.06	0.05	0.18	0.03	-	-
<i>Chaetodon ocellatus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	0.03
<i>Chaetodon sedentarius</i>	-	0.18	-	0.05	0.09	0.04	-	-	-	-	-	-	-	0.06
<i>Chilomycterus schoepfii</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Chromis enchrysur</i>	4.58	-	0.08	0.57	0.19	-	-	-	-	-	-	-	0.09	0.51
<i>Corniger spinosus</i>	-	0.18	-	-	-	0.04	0.03	-	0.01	0.19	-	0.03	0.02	-
<i>Cyclopsetta fimbriata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Decodon puellaris</i>	0.14	-	0.31	-	0.09	-	0.11	-	0.01	0.09	-	0.03	0.35	-
<i>Diodon holocanthus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Epinephelus flavolimbatus</i>	-	-	-	-	-	-	-	-	-	0.05	-	-	-	-
<i>Epinephelus niveatus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Fistularia petimba</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Gobiidae	0.56	0.18	2.33	1.20	-	0.04	-	-	0.01	-	-	0.09	1.56	-
<i>Gonioplectrus hispanus</i>	-	0.09	-	-	-	-	-	-	0.02	-	0.18	-	-	-
<i>Gymnothorax kolpos</i>	-	-	-	0.05	-	-	-	-	-	-	-	-	-	-
<i>Halichoeres bathyphilus</i>	0.28	0.35	0.93	1.30	0.09	-	-	-	-	-	0.04	0.02	-	-
<i>Hemanthias leptus</i>	-	-	-	-	-	-	0.03	-	0.04	-	0.04	-	-	-
<i>Hemanthias vivanus</i>	28.75	-	10.11	25.08	31.50	-	-	-	-	1.30	0.97	0.41	7.78	-
<i>Holocanthus bermudensis</i>	-	-	-	-	-	-	-	-	-	-	-	-	0.02	-
<i>Holocanthus tricolor</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Holocentrus adsensionus</i>	-	-	-	0.05	-	-	-	-	-	-	-	-	-	-
<i>Lactophrys quadricornis</i>	-	-	-	-	-	-	-	-	-	-	-	-	0.02	-
<i>Leiostomus xanthurus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Liopropoma eukrines</i>	-	0.09	-	0.26	0.19	-	0.06	-	-	-	0.04	0.02	0.06	-
<i>Lutjanus campechanus</i>	-	-	-	-	0.19	-	-	-	0.04	-	-	-	0.09	0.03
<i>Monacanthus spp.</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Muraenidae	-	-	-	-	-	-	-	-	-	-	0.04	-	-	-
<i>Mycteroperca microlepis</i>	-	-	-	-	-	-	-	-	0.12	-	-	-	0.02	-
<i>Mycteroperca phenax</i>	-	-	-	-	0.37	-	-	-	-	0.05	0.04	-	0.06	-
Myctophidae	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Ozococephalus spp.</i>	-	-	0.08	-	-	-	0.03	-	-	-	-	-	-	0.03
<i>Opsanus pardus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	0.03
<i>Ostichthys trachypoma</i>	-	-	-	-	-	-	0.31	-	-	-	0.04	-	-	-
<i>Pagrus pagrus</i>	-	-	-	-	-	-	-	-	0.02	-	-	-	-	-
<i>Parablennius marmoratus</i>	-	-	0.08	0.05	-	-	-	-	-	-	-	-	-	-
<i>Paranthias furcifer</i>	-	-	-	-	1.12	-	-	-	-	-	-	-	-	-
<i>Pareques iwamotoi</i>	-	-	-	-	-	-	0.06	-	0.07	-	0.04	-	-	-
<i>Pareques umbrosus</i>	-	0.62	-	-	-	-	-	-	0.02	-	-	0.02	-	0.03
<i>Peprilus burti</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Pontinus spp.</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Priacanthus arenatus</i>	-	-	-	-	-	0.07	0.08	-	-	-	-	-	-	-
<i>Pristigenys alta</i>	0.28	0.27	0.08	0.10	0.09	0.04	-	-	0.05	0.51	0.09	0.02	0.11	0.45
<i>Pronotogrammus martinicensis</i>	0.42	16.93	0.86	3.02	2.52	2.66	7.18	-	3.33	1.72	6.52	0.73	0.84	0.03
<i>Rhomboplites aurorubens</i>	-	-	-	-	-	-	-	-	-	-	-	-	0.02	0.12
<i>Rypticus maculatus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Sargocentron bullisi</i>	-	-	-	0.10	0.09	-	-	-	-	-	-	-	-	-
<i>Scorpaena dispar</i>	-	-	-	0.05	-	0.07	0.14	-	0.05	-	-	0.02	-	0.06
<i>Seriola dumerili</i>	0.14	0.09	-	-	-	-	-	-	0.04	-	-	-	0.04	-
<i>Seriola fasciata</i>	-	-	-	-	0.09	-	-	-	-	-	-	-	-	-
<i>Seriola rivoliana</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Serranus notospilus</i>	-	-	-	-	-	-	0.11	-	-	-	-	-	-	-
<i>Serranus phoebe</i>	0.14	0.09	-	-	0.19	-	0.03	-	-	-	-	0.03	0.04	0.03
<i>Sphoeroides spengleri</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Smodus sp.</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Trachurus lathami</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	0.57
Trichiuridae	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Unidentified flatfish	-	-	-	-	-	-	0.03	-	-	-	-	-	-	-
<i>Upeneus naryus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	0.03
Total Fish per Minute	35.42	19.24	14.93	32.36	36.90	2.95	8.21	-	3.92	4.10	8.24	1.45	11.17	2.01
No. Minutes	7.2	11.3	12.9	19.2	10.7	27.3	35.9	ND	84.4	21.5	22.7	58.6	46.3	33.3

Appendix A1. (continued).

53	54	56	63	64	65	69	72	76	81	83	86	87	90	Station No.
76	78	78	77	77	78	78	78	78	70	68	68	68	68	Base Depth (m)
T	D	D	S	S	R	R(n)	CS	P(n)	F	Y(m)	Y	Y	Y	Study Site
-	-	-	-	-	-	-	-	-	-	-	-	-	-	Taxa
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Antigonia capros</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	Apogonidae
-	-	-	-	-	-	0.04	0.02	0.07	-	0.08	-	-	-	<i>Apogon pseudomaculatus</i>
-	-	-	-	-	-	0.01	-	-	-	-	-	-	-	<i>Aulostomus maculatus</i>
-	-	-	-	-	0.02	-	-	-	-	-	0.03	0.02	-	<i>Bodianus pulchellus</i>
-	-	-	-	-	-	0.09	-	0.32	-	-	-	-	-	<i>Bregmaceros cantori</i>
-	-	-	-	-	0.19	0.06	0.02	0.03	-	0.05	-	0.21	0.07	<i>Canthigaster rostrata</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	Carangidae
-	0.09	-	-	-	-	-	-	-	0.09	-	-	-	-	<i>Centropristis ocyurus</i>
-	0.06	0.02	0.02	-	0.15	0.06	0.29	0.03	0.17	0.11	0.17	0.19	0.28	<i>Chaetodon aya</i>
-	-	-	-	-	-	0.01	-	-	-	-	-	-	-	<i>Chaetodon ocellatus</i>
-	-	-	0.02	0.10	0.02	0.06	0.02	0.03	-	0.05	0.14	0.02	-	<i>Chaetodon sedentarius</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	0.07	<i>Chilomycterus schoepfii</i>
-	0.03	-	0.03	-	0.35	0.08	0.05	-	-	0.21	0.28	0.11	0.21	<i>Chromis enchrysur</i>
-	-	-	0.02	-	0.06	0.14	0.15	-	-	0.05	0.10	0.13	0.07	<i>Corniger spinosus</i>
-	-	-	-	-	-	-	-	-	-	0.03	-	-	-	<i>Cyclosetta fimbriata</i>
-	0.14	0.02	0.17	0.10	0.10	-	0.10	0.02	0.04	0.03	0.17	0.06	0.28	<i>Decodon puellaris</i>
-	-	-	-	-	-	-	-	-	-	0.03	-	-	-	<i>Diodon holocanthus</i>
-	-	-	-	-	-	-	0.02	-	-	-	-	-	-	<i>Epinephelus flavolimbatus</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Epinephelus niveatus</i>
-	-	-	-	-	-	0.01	-	-	-	-	-	-	-	<i>Fistularia petimba</i>
-	0.43	0.02	0.17	0.97	-	-	0.10	0.05	0.89	0.05	0.38	0.49	0.07	Gobiidae
-	-	-	-	-	-	-	0.02	-	-	-	-	-	-	<i>Gonioplectrus hispanus</i>
-	-	-	-	-	-	-	-	0.05	-	-	-	-	-	<i>Gymnothorax kolpos</i>
-	0.02	-	-	-	0.37	-	0.27	-	0.26	0.03	0.10	0.19	0.28	<i>Halichoeres bathyphilus</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Hemanthias leptus</i>
-	1.89	1.43	0.19	-	8.24	0.09	1.34	-	-	0.08	1.59	4.79	8.78	<i>Hemanthias vivanus</i>
-	-	-	-	-	0.08	-	0.07	-	-	-	-	-	-	<i>Holacanthus bermudensis</i>
-	-	-	-	-	0.02	0.01	-	-	-	-	-	-	-	<i>Holacanthus tricolor</i>
-	-	-	-	-	-	0.02	-	-	-	-	-	-	-	<i>Holocentrus adsensionus</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Lactophrys quadricornis</i>
-	-	-	-	-	-	0.10	-	-	-	-	-	-	-	<i>Leiostomus xanthurus</i>
-	-	0.10	-	-	0.02	-	0.15	-	0.13	-	0.03	0.15	0.50	<i>Liopropoma eukrines</i>
-	-	-	-	-	-	-	-	-	-	0.05	-	-	-	<i>Lutjanus campechanus</i>
-	-	-	-	-	-	-	-	-	-	0.03	-	-	-	<i>Monacanthus spp.</i>
-	-	-	-	-	-	-	0.02	-	-	-	-	-	-	Muraenidae
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Mycteroperca microlepis</i>
-	-	-	-	-	-	-	0.12	-	-	-	-	0.15	0.07	<i>Mycteroperca phenax</i>
-	-	-	-	-	-	0.01	-	-	-	-	-	-	-	Myctophidae
-	-	-	-	-	-	-	-	-	-	0.08	-	-	-	<i>Oecocephalus spp.</i>
-	-	-	-	-	-	0.01	-	0.02	0.09	0.05	-	0.02	0.07	<i>Opsanus pardus</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Ostichthys trachypoma</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Pagrus pagrus</i>
-	-	-	-	-	0.04	-	-	-	-	-	-	0.04	-	<i>Parablennius marmoratus</i>
-	-	-	-	-	-	0.01	-	-	-	-	-	0.08	0.14	<i>Paranthias furcifer</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Pareques iwamotoi</i>
-	0.14	-	0.03	-	0.02	0.02	0.90	-	-	0.03	-	-	-	<i>Pareques umbrosus</i>
-	-	-	-	-	-	0.01	-	-	-	-	-	-	-	<i>Peprilus burti</i>
-	-	0.02	-	0.05	-	0.01	-	0.03	-	-	0.03	-	-	<i>Pontinus spp.</i>
-	-	-	-	-	0.06	0.01	0.02	-	-	-	-	-	-	<i>Priacanthus arenatus</i>
-	0.08	0.02	0.03	-	0.08	0.28	0.20	-	0.17	0.21	0.24	0.04	-	<i>Pristigenys alta</i>
-	0.68	0.82	0.32	0.20	2.56	1.29	4.74	1.20	0.72	0.05	3.42	3.84	6.79	<i>Promotogrammus martinicensis</i>
-	0.03	0.17	0.02	0.15	-	0.05	-	0.17	-	-	-	-	-	<i>Rhomboplites aurorubens</i>
-	-	-	0.03	-	-	-	-	-	-	-	-	-	-	<i>Rypicus maculatus</i>
-	-	-	-	-	0.06	-	-	-	-	0.13	-	0.02	-	<i>Sargocentron bullisi</i>
-	-	0.10	0.05	-	0.02	0.14	0.02	-	-	0.21	-	-	-	<i>Scorpaena dispar</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Seriola dumerili</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Seriola fasciata</i>
-	-	-	-	-	-	-	0.05	-	-	-	-	-	-	<i>Seriola rivoliana</i>
-	-	-	0.05	-	-	-	0.07	-	-	-	-	-	-	<i>Serranus notospilus</i>
-	0.03	0.10	-	0.15	0.10	0.04	0.10	-	0.47	0.08	-	0.04	0.14	<i>Serranus phoebe</i>
-	-	-	0.02	-	-	-	-	-	-	-	-	-	0.07	<i>Sphaeroides spengleri</i>
-	0.03	-	-	-	0.02	0.02	-	-	0.04	-	-	-	-	<i>Synodus sp.</i>
-	-	-	-	-	-	0.10	-	0.12	-	0.08	-	-	-	<i>Trachurus lathami</i>
-	-	-	-	-	-	-	-	0.07	-	0.29	-	-	-	Trichiuridae
-	-	-	-	-	-	-	-	-	-	-	-	-	-	Unidentified flatfish
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Upenus parvus</i>
-	3.68	2.86	1.17	1.79	12.61	2.82	8.89	2.22	3.06	2.12	6.71	10.61	17.91	Total Fish per Minute
ND	65.0	41.3	59.1	19.6	48.1	123.2	41.0	58.5	23.5	37.2	28.9	97.2	14.1	No. Minutes

Appendix A2. Summary of fishes associated with the talus zone observed during 1997 (97-01) ROV surveys. Data are as number of fish observed per minute of video. ND-No Data and (n)-night survey.

Station No.	04	08	09	11	12	20	22	27	33	35	36	44	45	50
Base Depth (m)	68	68	78	78	78	78	106	104	110	110	110	77	71	72
Study Site	Y	Y	R	R	R	C	L-A	L-B	L-C	L-1	L-1	PF	SM-1	SM-2
Taxa														
<i>Ancylopsetta dilecta</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Chromis enchrysur</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Decodon puellaris</i>	-	2.20	-	-	-	-	-	-	-	-	-	-	-	-
Gobiidae	-	8.81	-	-	-	-	-	-	0.68	-	-	-	-	-
<i>Halichoeres bathyphilus</i>	-	0.31	-	-	-	-	-	-	-	-	-	-	-	-
<i>Liopropoma eukrines</i>	-	0.16	-	-	-	-	-	-	-	-	-	-	-	-
Opistognathidae	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Pontinus</i> sp.	-	-	-	-	-	-	-	-	0.14	-	-	-	-	-
<i>Priacanthus arenatus</i>	-	0.16	-	-	-	-	-	-	-	-	-	-	-	-
<i>Pristigenys alta</i>	-	0.16	-	-	-	-	-	-	-	-	-	-	-	-
<i>Pronotogrammus martinicensis</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Scorpaena dispar</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Serranus notospilus</i>	-	0.16	-	-	-	-	-	-	0.27	-	-	-	-	-
<i>Serranus phoebe</i>	-	0.16	-	-	-	-	-	-	-	-	-	-	-	-
<i>Synodus</i> sp.	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Total Fish per Minute	-	12.74	-	-	-	-	-	-	1.35	-	-	-	-	-
No. Minutes	ND	6.4	ND	ND	ND	ND	ND	ND	7.4	ND	ND	ND	ND	ND

Appendix A3. Summary of fishes associated with the sand flats observed during 1997 (97-01) ROV surveys. Data are as number of fish observed per minute of video. ND-No Data and (n)-night survey.

Station No.	04	08	09	11	12	20	22	27	33	35	36	44	45	50
Base Depth (m)	68	68	78	78	78	78	106	104	110	110	110	77	71	72
Study Site	Y	Y	R	R	R	C	L-A	L-B	L-C	L-1	L-1	PF	SM-1	SM-2
Taxa														
<i>Bregmaceros cantori</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Centropristis ocyurus</i>	-	0.21	-	-	-	-	-	-	-	-	-	-	-	-
<i>Citharichthys cornutus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Decodon puellaris</i>	-	-	-	-	-	-	-	-	0.03	-	-	0.29	1.00	-
Gobiidae	-	0.63	-	-	-	-	0.08	0.04	0.31	0.56	-	0.18	0.25	-
<i>Halichoeres bathyphilus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Halieutichthys aculeatus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Hemanthias vivanus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Hoplunnis</i> sp.	-	-	-	-	-	-	-	0.08	-	-	-	-	-	-
<i>Lagodon rhomboides</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Leiostomus xanthurus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Lepophidium brevibarbe</i>	-	-	-	-	-	-	-	0.01	-	-	-	-	-	-
<i>Ogcocephalus</i> spp.	-	-	-	-	-	-	-	0.01	0.03	-	-	-	-	-
Ophichthidae	-	-	-	-	-	-	-	0.01	-	-	-	-	-	-
Opistognathidae	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Pagrus pagrus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Paralichthys squamilentus</i>	-	-	-	-	-	-	-	-	0.03	-	-	-	-	-
<i>Parequetus umbrosus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Pontinus</i> sp.	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Prionotus stearnsi</i>	-	-	-	-	-	-	-	0.01	-	-	-	-	-	-
<i>Pristipomoides aquilonaris</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Raja</i> sp.	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Rhynchoconger</i> sp.	-	-	-	-	-	-	-	0.01	-	-	-	-	-	-
<i>Saurida</i> sp.	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Scorpaena dispar</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Serranus atrobranchus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Serranus notospilus</i>	-	1.26	-	-	-	-	0.17	0.33	0.23	0.07	-	0.11	1.24	-
<i>Serranus phoebe</i>	-	-	-	-	-	-	0.17	0.02	-	-	-	0.04	-	-
<i>Sphoeroides</i> sp.	-	0.21	-	-	-	-	-	-	-	-	-	-	-	-
<i>Syacium papillosum</i>	-	-	-	-	-	-	-	0.01	0.05	-	-	-	-	-
<i>Symphurus</i> sp.	-	-	-	-	-	-	-	0.01	-	-	-	-	-	-
<i>Synodus</i> sp.	-	-	-	-	-	-	-	0.02	0.03	0.07	-	-	-	-
<i>Trachurus lathami</i>	-	-	-	-	-	-	-	0.04	-	-	-	-	-	-
Trichiuridae	-	-	-	-	-	-	-	0.01	-	-	-	-	-	-
Unidentified flatfish	-	-	-	-	-	-	-	0.01	0.03	-	-	-	-	-
<i>Upeneus parvus</i>	-	-	-	-	-	-	-	0.04	-	-	-	-	-	-
Total Fish per Minute	-	2.09	-	-	-	-	0.42	0.73	0.75	0.70	-	0.73	2.49	-
No. Minutes	ND	4.8	ND	ND	ND	ND	12.0	84.7	38.5	14.3	ND	27.5	4.0	ND

Appendix A2. (continued).

53	54	56	63	64	65	69	72	76	81	83	86	87	90	Station No.
76	78	78	77	77	78	78	78	78	70	68	68	68	68	Base Depth (m)
T	D	D	S	S	R	R(n)	CS	P(n)	F	Y(n)	Y	Y	Y	Study Site
-	-	-	-	-	-	-	-	-	0.05	-	-	-	-	Taxa
-	-	-	-	-	-	-	-	-	-	-	0.25	-	-	<i>Ancylosetta dilecta</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Chromis enchrysur</i>
-	-	-	-	-	-	-	0.10	-	0.05	-	0.25	0.21	0.86	<i>Decodon puellaris</i>
-	-	-	-	-	-	-	3.12	-	1.03	-	4.43	0.21	-	Gobiidae
-	-	-	-	-	-	-	-	-	-	-	-	0.64	-	<i>Halichoeres bathyphilus</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Liopropoma eukrines</i>
-	-	-	-	-	-	-	0.10	-	0.05	-	-	-	-	Opistognathidae
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Pontinus</i> sp.
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Priacanthus arenatus</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Pristigenys alta</i>
-	-	-	-	-	-	-	-	-	-	-	0.12	-	-	<i>Pronotogrammus martinicensis</i>
-	-	-	-	-	-	-	-	-	0.05	-	0.12	-	-	<i>Scorpaena dispar</i>
-	-	-	-	-	-	-	0.31	-	-	-	-	-	-	<i>Serranus notospilus</i>
-	-	-	-	-	-	-	-	-	0.05	-	0.74	-	-	<i>Serranus phoebe</i>
-	-	-	-	-	-	-	0.21	-	0.09	-	0.37	-	-	<i>Synodus</i> sp.
-	-	-	-	-	-	-	3.84	-	1.30	-	6.28	1.06	0.86	Total Fish per Minute
ND	ND	ND	ND	ND	ND	ND	9.6	ND	21.6	ND	8.1	4.7	1.2	No. Minutes

Appendix A3. (continued).

53	54	56	63	64	65	69	72	76	81	83	86	87	90	Station No.
76	78	78	77	77	78	78	78	78	70	68	68	68	68	Base Depth (m)
T	D	D	S	S	R	R(n)	CS	P(n)	F	Y(n)	Y	Y	Y	Study Site
-	-	-	-	-	-	-	-	-	-	-	-	-	-	Taxa
0.07	-	-	-	-	-	-	-	0.33	-	-	-	-	-	<i>Bregmaceros cantori</i>
-	-	-	0.08	-	-	-	-	-	-	-	-	-	-	<i>Centropristis ocyurus</i>
-	-	-	-	-	-	-	-	-	-	-	0.18	-	-	<i>Citharichthys cornutus</i>
-	-	-	-	-	-	-	0.19	-	-	-	-	-	-	<i>Decodon puellaris</i>
0.16	-	0.18	1.48	-	-	-	0.55	0.83	-	-	1.06	-	-	Gobiidae
-	-	-	-	-	-	-	0.08	-	-	-	0.18	-	-	<i>Halichoeres bathyphilus</i>
0.03	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Halieutichthys aculeatus</i>
-	-	-	-	-	-	-	0.04	-	-	-	-	-	-	<i>Hemanthias vivanus</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Hoplunnis</i> sp.
-	-	-	0.08	-	-	-	-	-	-	-	-	-	-	<i>Lagodon rhomboides</i>
-	-	-	0.08	-	-	-	-	0.08	-	-	-	-	-	<i>Leiostomus xanthurus</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Lepophidium brevibarbe</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Ogcocephalus</i> spp.
-	-	-	-	-	-	-	-	-	-	-	-	-	-	Ophichthidae
-	-	-	-	-	-	-	0.04	-	-	-	-	-	-	Opistognathidae
-	-	-	-	-	-	-	-	0.08	-	-	-	-	-	<i>Pagrus pagrus</i>
-	-	-	0.08	-	-	-	-	-	-	-	-	-	-	<i>Paralichthys squamilentus</i>
-	-	-	-	-	-	-	-	0.08	-	-	-	-	-	<i>Parequetus umbrosus</i>
-	-	-	-	-	-	-	-	0.17	-	-	-	-	-	<i>Pontinus</i> sp.
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Prionotus stearnsi</i>
0.07	-	-	-	-	-	-	0.16	-	-	-	-	-	-	<i>Pristipomoides aquilonaris</i>
-	-	-	-	-	-	-	0.04	-	-	-	-	-	-	<i>Raja</i> sp.
-	-	-	-	-	-	-	-	0.08	-	-	1.94	-	-	<i>Rhynchoconger</i> sp.
2.03	-	-	-	-	-	-	-	-	-	-	0.35	-	-	<i>Saurida</i> sp.
-	-	-	0.08	-	-	-	0.08	-	-	-	0.18	-	-	<i>Scorpaena dispar</i>
0.03	-	-	-	-	-	-	-	-	-	-	0.18	-	-	<i>Serranus atrobranchus</i>
0.03	1.43	0.18	-	-	-	-	0.51	0.42	-	-	1.06	-	-	<i>Serranus notospilus</i>
-	-	-	-	-	-	-	0.04	-	-	-	-	-	-	<i>Serranus phoebe</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Sphoeroides</i> sp.
0.13	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Syacium papillosum</i>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Symphurus</i> sp.
0.07	1.43	0.73	-	-	-	-	0.04	0.50	-	-	5.11	-	-	<i>Synodus</i> sp.
-	-	0.18	-	-	-	-	-	-	-	-	-	-	-	<i>Trachurus lathami</i>
-	-	-	-	-	-	-	-	0.08	-	-	0.18	-	-	Trichiuridae
-	-	0.18	-	-	-	-	-	-	-	-	0.35	-	-	Unidentified flatfish
-	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Upeneus parvus</i>
2.63	2.86	1.45	1.88	-	-	-	1.75	2.67	-	-	10.74	-	-	Total Fish per Minute
30.5	0.7	5.5	12.8	ND	ND	ND	25.7	11.8	ND	ND	5.7	ND	ND	No. Minutes

Appendix A4. Summary of Cruise USGS 99-03 and 2000-01(0001) ROV surveys. Values given are number of individuals observed per minute of tape time. Night surveys are indicated by (n).

Station No.	0001-028	0001-029	0001-036	0001-055	0001-057	0001-060	9903-2132	0001-016	0001-018	0001-019	0001-033	0001-052	0001-059
Base Depth (m)	69	69	88	110	105	79	109	80	110	110	88	110	105
Study Site	Y	Y	A	L-H	SC	C5	L-1(n)	R (n)	L-2 (n)	L-2 (n)	A (n)	L-A (n)	SC (n)
Taxa													
<i>Anthias tenuis</i>	-	0.02	-	-	-	-	-	-	-	-	0.11	-	-
<i>Apogon pseudomaculatus</i>	-	0.22	0.05	-	-	-	-	0.04	-	-	0.05	-	0.01
<i>Aulostomus maculatus</i>	-	0.04	0.05	-	-	-	-	-	-	-	-	-	-
<i>Bodianus pulchellus</i>	-	0.11	-	-	-	-	-	-	-	-	-	-	-
<i>Bregmaceros cantori</i>	-	-	-	-	-	-	-	-	-	2.22	0.03	-	-
<i>Brotula barbata</i>	-	-	-	-	0.04	0.09	-	-	-	-	0.01	-	-
<i>Canthigaster jamestyleri</i>	0.11	-	-	-	-	-	-	0.01	-	-	-	-	-
<i>Canthigaster rostrata</i>	0.32	0.22	0.15	-	-	-	-	0.32	-	-	0.01	-	-
Carapidae	-	-	-	-	-	-	-	-	-	0.02	-	-	-
<i>Centropristis ocyurus</i>	-	-	-	-	-	0.44	-	-	-	-	-	-	-
<i>Chaetodon aya</i>	0.43	0.20	0.70	0.04	0.13	0.44	-	0.26	-	0.03	0.05	0.18	0.03
<i>Chaetodon sedentarius</i>	0.11	0.04	0.10	-	-	-	-	0.10	-	-	-	-	-
<i>Chilomycterus antillarum</i>	-	-	-	-	-	-	-	-	-	0.02	-	-	-
<i>Chromis enchrysur</i>	1.51	1.40	0.05	-	-	0.27	-	0.23	-	-	0.02	-	-
<i>Chromis insolata</i>	-	0.04	-	-	-	-	-	-	-	-	-	-	-
<i>Chromis scotti</i>	-	0.47	-	-	-	0.27	-	0.01	-	-	-	-	-
<i>Corniger spinosus</i>	-	0.11	0.15	-	0.04	0.27	-	0.03	-	-	0.07	-	-
<i>Coryphopterus punctipectophorus</i>	0.22	0.15	-	-	-	-	-	0.03	-	-	-	-	-
<i>Decodon puellaris</i>	-	0.02	-	0.05	-	-	-	0.04	-	-	-	-	0.01
<i>Epinephelus niveatus</i>	-	-	-	-	-	0.09	-	-	-	-	-	-	-
<i>Fistularia petimba</i>	-	-	-	-	-	-	-	-	-	-	-	-	0.01
<i>Gonioplectrus hispanus</i>	-	-	-	-	0.04	0.09	-	-	-	-	-	-	-
<i>Gymnothorax hubbsi</i>	0.11	-	-	-	-	-	-	-	-	-	-	-	-
<i>Gymnothorax kolpos</i>	-	-	-	-	-	-	-	-	-	-	-	0.18	-
<i>Halichoeres bathyphilus</i>	0.11	0.20	0.05	0.10	-	-	-	0.04	-	-	0.01	-	-
<i>Hemanthias leptus</i>	-	-	-	-	0.26	-	-	-	-	-	-	-	-
<i>Hemanthias vivanus</i>	36.97	6.74	0.10	-	2.57	25.71	-	2.25	-	-	0.04	0.18	-
<i>Holacanthus bermudensis</i>	0.11	0.04	-	0.01	0.09	0.09	-	-	-	-	0.01	-	-
<i>Holocentrus adscensionis</i>	0.11	-	-	-	-	-	-	-	-	-	-	-	-
<i>Lactophrys quadricornis</i>	-	-	-	-	-	0.09	-	-	-	-	-	-	-
<i>Liopropoma eukrines</i>	-	0.13	0.05	0.09	0.09	0.18	-	0.05	-	-	0.02	0.18	-
<i>Lutjanus campechanus</i>	-	-	0.05	-	-	-	-	-	-	-	0.01	-	-
<i>Malacanthus plumieri</i>	-	-	-	-	-	-	-	0.01	-	-	-	-	-
Muraenidae	-	0.02	-	-	-	-	-	-	-	-	0.01	-	-

Appendix A4. (continued).

Station No.	0001-028	0001-029	0001-036	0001-055	0001-057	0001-060	9903-2132	0001-016	0001-018	0001-019	0001-033	0001-052	0001-059
Base Depth (m)	69	69	88	110	105	79	109	80	110	110	88	110	105
Study Site	Y	Y	A	L-H	SC	C5	L-1(n)	R (n)	L-2 (n)	L-2 (n)	A (n)	L-A (n)	SC (n)
Taxa													
<i>Mustelus</i> sp.	-	-	-	-	-	-	0.04	-	-	0.02	0.01	-	0.01
<i>Mycteroperca microlepis</i>	-	-	-	-	0.04	-	-	-	-	-	-	-	-
<i>Mycteroperca phenax</i>	-	0.02	-	0.01	0.13	-	-	-	-	0.02	0.01	0.18	0.03
Myctophidae	-	-	-	-	-	-	0.24	-	0.65	0.10	-	27.37	2.15
<i>Neopinnula americana</i>	-	-	-	-	-	-	-	-	-	0.03	-	-	-
<i>Ogcocephalus</i> sp.	-	-	-	-	-	0.09	-	0.01	-	-	0.01	-	0.01
<i>Opistognathus lonchurus</i>	-	0.02	-	-	-	-	-	-	-	-	-	-	-
<i>Opsanus pardus</i>	-	-	-	-	-	-	-	0.01	-	-	-	-	-
<i>Ostichthys trachypoma</i>	-	-	-	-	0.09	-	-	-	0.13	0.03	-	1.46	-
<i>Pagrus pagrus</i>	-	-	-	0.11	0.09	-	0.04	0.01	-	-	-	-	0.01
<i>Paranthias furcifer</i>	-	0.37	-	-	-	-	-	-	-	-	-	-	-
<i>Pareques iwamotoi</i>	-	-	-	-	-	0.09	-	-	-	-	0.02	0.36	0.10
<i>Pareques umbrosus</i>	-	0.02	0.15	0.04	0.21	0.44	-	-	-	-	0.20	0.55	0.04
<i>Pomatomus saltatrix</i>	-	-	-	-	-	-	-	-	-	-	-	-	0.01
<i>Priacanthus arenatus</i>	-	-	-	-	-	0.27	-	0.01	-	-	-	-	-
<i>Pristigenys alta</i>	0.32	0.02	0.20	0.32	0.09	0.53	0.05	0.14	0.13	0.32	0.10	0.91	0.07
<i>Pronotogrammus martinicensis</i>	0.97	2.79	14.01	1.48	11.41	20.83	0.02	4.13	-	0.22	3.74	3.10	0.03
<i>Rhomboplites aurorubens</i>	-	-	-	-	-	-	-	-	0.04	0.02	-	-	-
<i>Rypticus maculatus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Sargocentron bullisi</i>	0.11	0.07	-	0.01	-	-	-	0.08	-	-	-	-	-
<i>Scomberomorus cavalla</i>	-	-	-	-	-	-	-	-	0.04	-	-	-	-
<i>Scorpaena dispar</i>	-	0.02	-	0.01	-	-	-	0.06	-	-	0.04	-	-
Scorpaenidae	-	-	-	0.06	0.13	0.18	0.02	0.03	0.04	0.11	0.01	-	0.07
<i>Seriola dumerili</i>	-	-	0.25	-	-	0.35	-	0.14	-	0.10	0.22	0.55	0.01
<i>Seriola rivoliana</i>	0.22	0.02	-	-	-	-	-	-	-	-	0.02	-	-
<i>Serranus phoebe</i>	0.22	0.04	0.05	0.04	-	0.62	-	0.04	-	-	-	-	-
<i>Sphoeroides spengleri</i>	-	-	-	-	-	-	-	0.01	-	-	-	-	-
<i>Stenotomus caprinus</i>	-	-	-	-	-	-	-	-	-	-	-	-	0.01
<i>Synodus</i> sp.	-	-	-	-	-	-	-	-	-	0.11	0.02	-	0.13
<i>Trachurus lathami</i>	-	-	-	-	-	-	-	0.03	0.52	0.87	0.09	0.73	-
<i>Trichiurus lepturus</i>	-	-	-	-	-	-	4.31	-	-	-	-	0.18	0.01
<i>Upeneus parvus</i>	-	-	-	-	-	-	-	-	-	0.05	-	0.91	0.07
Total Fish per Minute	1.84	3.39	14.66	2.09	12.18	23.40	4.71	4.71	1.57	1.98	4.49	36.31	2.79
No. Minutes	9.3	53.7	20.1	80.3	23.3	11.3	55.0	77.0	22.6	63.1	97.0	5.5	69.9

A-7

APPENDIX B

A list of reef-fish species from natural reefs in the northern Gulf of Mexico

Appendix B. A list of reef-fish species from natural reefs in the northern Gulf of Mexico with a comparison among the northwestern Gulf, Pinnacles, and West Florida shelf. Data sources: NW Gulf: Ca-Cashman 1973, Br-Bright et al. 1974, So-Sonnier et al. 1976, Bo-Boland et al. 1983, Pu-Putt et al. 1986, DB-Dennis and Bright 1988a, b, Pa-Pattengill et al. 1997, and HB-Hoese and Moore 1998. West Florida: SW-Springer and Woodburn, 1960, Po-Powell et al. 1972, Sm-Smith et al. 1975, Sm-Smith 1976, SH-Shipp and Hopkins 1978, Ha-Hastings 1979, WS-Williams and Shipp 1980, BG-Bullock and Godcharles 1982, DG-Darcy and Gutherz 1984, Cl-Clarke 1981, DK-Darnell and Kleypas 1987, and BS-Bullock and Smith 1991.

Scientific Name	Common Name	NW Gulf		Pinnacles 60-110 m	West Florida Shelf	
		<60 m	>60 m		<30 m	>30 m
Orectolobidae – carpet sharks						
<i>Ginglymostoma cirratum</i>	nurse shark	So				Sm76
Muraenidae – morays						
<i>Enchelycore carychroa</i>	chestnut moray	Ca				
<i>Enchelycore nigricans</i>	viper moray	Ca				
<i>Gymnothorax funebris</i>	green moray					Cl
<i>Gymnothorax hubbsi</i>	lichen moray			X		
<i>Gymnothorax kolpos</i>	blacktail moray			X		
<i>Gymnothorax miliaris</i>	goldentail moray	Pa				
<i>Gymnothorax moringa</i>	spotted moray	Ca	DB	X		Sm75
<i>Gymnothorax vicinus</i>	purplemouth moray	Ca				
<i>Muraena retifera</i>	reticulated moray			X		Sm75 SH
Synodontidae – lizardfishes						
<i>Synodus intermedius</i>	sand diver	DB	DB	X		Sm75
<i>Synodus saurus</i>	bluestripe lizardfish	Pa				Sm75
<i>Synodus synodus</i>	red lizardfish	Ca	DB			
Bythitidae – viviparous brotulas						
<i>Ogilbia cayorum</i>	key brotula					SW DK
Ophidiidae - cusk-eels						
<i>Brotula barbata</i>	bearded brotula		HM	X		
Batrachoididae						
<i>Opsanus pardus</i>	leopard toadfish	HM		X		SW/Sm75 SH

Appendix B. (continued).

Scientific Name	Common Name	NW Gulf		Pinnacles	West Florida Shelf	
		<60 m	>60 m	60-110 m	<30 m	>30 m
Holocentridae - squirrelfishes						
<i>Corniger spinosus</i>	spinycheek soldierfish			X		
<i>Holocentrus adscensionis</i>	squirrelfish	Ca	DB	X	Sm75	
<i>Holocentrus rufus</i>	longspine squirrelfish	Ca	DB			
<i>Myripristis jacobus</i>	blackbar soldierfish	Ca			Sm75	DG
<i>Neoniphon marianus</i>	longjaw squirrelfish	DB	DB			DK
<i>Ostichthys trachypoma</i>	bigeye soldierfish			X		SH/DG
<i>Plectrypops retrospinis</i>	cardinal soldierfish	Ca				
<i>Sargocentron bullisi</i>	deepwater squirrelfish			X	Sm75	SH/DK
<i>Sargocentron coruscum</i>	reef squirrelfish				Po	
<i>Sargocentron poco</i>	saddle squirrelfish	Ca				
<i>Sargocentron vexillarium</i>	dusky squirrelfish	Pa			Ha	DG
Caproidae - boarfishes						
<i>Antigonia capros</i>	deepbody boarfish			X		
Aulostomidae – trumpetfishes						
<i>Aulostomus maculatus</i>	trumpetfish	Ca		X	Sm75	
Fistulariidae – cornetfishes						
<i>Fistularia petimba</i>	red cornetfish			X		
<i>Fistularia tabacaria</i>	bluespotted cornetfish	Bo				
Scorpaenidae – scorpionfishes						
<i>Neomerinthe hemingwayi</i>	spinycheek scorpionfish			DK		DK
<i>Pontinus rathbuni</i>	highfin scorpionfish			X		
<i>Scorpaena bergi</i>	goosehead scorpionfish				DG	
<i>Scorpaena dispar</i>	hunchback scorpionfish		DB	DK	Po	DK
<i>Scorpaena inermis</i>	mushroom scorpionfish					DK
<i>Scorpaena plumieri</i>	spotted scorpionfish	So			Sm75	
<i>Scorpaenodes caribbaeus</i>	reef scorpionfish	Ca				

Appendix B. (continued).

Scientific Name	Common Name	NW Gulf		Pinnacles 60-110 m	West Florida Shelf	
		<60 m	>60 m		<30 m	>30 m
Serranidae - sea basses						
<i>Anthias nicholsi</i>	yellowfin bass					SH/WS ¹
<i>Anthias tenuis</i>	threadnose bass		?	X		
<i>Bathyanthias mexicana</i>	yellowtail bass		HM			DG/BS
<i>Cephalopholis cruentata</i>	graysby	Ca			Sm75	Cl
<i>Cephalopholis fulva</i>	coney	DB			Sm75	
<i>Dermatolepis inermis</i>	marbled grouper	Ca	DB			BG
<i>Epinephelus adscensionis</i>	rock hind	Ca	DB		Sm75	BS
<i>Epinephelus drummondhayi</i>	speckled hind			X	Sm75	
<i>Epinephelus guttatus</i>	red hind	Ca	DB		Sm75	BS
<i>Epinephelus itajara</i>	jewfish				SW/Sm75	
<i>Epinephelus morio</i>	red grouper	Bo/Pa			Sm75/DK	BS
<i>Epinephelus mystacinus</i>	misty grouper					BS
<i>Epinephelus niveatus</i>	snowy grouper		DB	X		
<i>Epinephelus nigritus</i>	warsaw grouper		DB	X		
<i>Epinephelus striatus</i>	Nassau grouper	DB				
<i>Gonioplectrus hispanus</i>	Spanish flag		Br	X		
<i>Hemanthias aureorubens</i>	streamer bass		HM	? ²		BS
<i>Hemanthias leptus</i>	longtail bass		DB	X		
<i>Hemanthias vivanus</i>	red barbier		DB	X		DG
<i>Hypoplectrus puella</i>	barred hamlet	?			Sm75	Cl/DK
<i>Liopropoma eukrines</i>	wrasse bass	Ca	DB	X	Po	SH/Cl
<i>Liopropoma rubre</i>	peppermint bass	Ca				
<i>Mycteroperca acutirostris</i>	comb grouper	Ca/So				
<i>Mycteroperca bonaci</i>	black grouper	DB			Sm75	DG
<i>Mycteroperca interstitialis</i>	yellowmouth grouper	Ca	DB		Sm75	

[?]presence uncertain.

¹photo from DeSoto Canyon indicates *P. martinicensis* whereas specimen from west Florida Shelf is *A. nicholsi*.

²reported by Gittings et al. 1991 but no specimens known to confirm its occurrence.

Appendix B. (continued).

Scientific Name	Common Name	NW Gulf		Pinnacles	West Florida Shelf	
		<60 m	>60 m	60-110 m	<30 m	>30 m
Serranidae (continued)						
<i>Mycteroperca microlepis</i>	gag		HM	X	Sm75	DG
<i>Mycteroperca phenax</i>	scamp		DB	X	Sm75	
<i>Mycteroperca tigris</i>	tiger grouper	DB			BG	
<i>Mycteroperca microlepis</i>	gag		HM	X	Sm75	DG
<i>Mycteroperca phenax</i>	scamp		DB	X	Sm75	
<i>Mycteroperca tigris</i>	tiger grouper	DB			BG	
<i>Mycteroperca venenosa</i>	yellowfin grouper	So			SW/Sm75	
<i>Paranthias furcifer</i>	creole-fish	Ca		X	Sm75	SH
<i>Plectranthias garrupellus</i>	apricot bass					BS
<i>Pronotogrammus martinicensis</i>	rougtongue bass		DB	X	Po	SH ¹
<i>Rypticus bistrispinus</i>	freckled soapfish			DK	WS	SH/WS
<i>Rypticus maculatus</i>	whitespotted soapfish	DB		X	Sm75	SH
<i>Rypticus subbifrenatus</i>	spotted soapfish	Ca				
<i>Serraniculus pumilio</i>	pygmy sea bass				SW	
<i>Schultzea beta</i>	school bass	HM		DK		DG
<i>Serranus annularis</i>	orangeback bass	DB	DB			BS
<i>Serranus notospilus</i>	saddle bass			X		DK/DG
<i>Serranus phoebe</i>	tattler	DB	DB	X		CI/SH
<i>Serranus subligarius</i>	belted sandfish	Pu			Sm75/DK	CI
<i>Serranus tabacarius</i>	tobaccofish	Pa			Sm75	BS
<i>Serranus tigrinus</i>	harlequin bass				Sm75	
<i>Serranus tortugarum</i>	chalk bass					DG
Grammatidae – basslets						
<i>Lipogramma regia</i>	royal basslet			X		
Opistognathidae – jawfishes						
<i>Opistognathus aurifrons</i>	yellowhead jawfish	DB			Sm75	CI
<i>Opistognathus lonchurus</i>	moustache jawfish		DB	X	Po	

Appendix B. (continued).

Scientific Name	Common Name	NW Gulf		Pinnacles 60-110 m	West Florida Shelf	
		<60 m	>60 m		<30 m	>30 m
Priacanthidae – bigeyes						
<i>Heteropriacanthus cruentatus</i>	glasseye snapper	Ca				
<i>Priacanthus arenatus</i>	bigeye	Ca	DB	X	Sm76	SH
<i>Pristigenys alta</i>	short bigeye		DB	X	Sm75	SH
Apogonidae – cardinalfishes						
<i>Apogon affinis</i>	bigtooth cardinalfish	Pa			Po	WS
<i>Apogon aurolineatus</i>	bridle cardinalfish	HM	DB		SW/Sm76	WS/DK
<i>Apogon binotatus</i>	barred cardinalfish				Sm75	
<i>Apogon maculatus</i>	flamefish	Ca	DB		Sm75	CI/DK
<i>Apogon pseudomaculatus</i>	twospot cardinalfish	So	DB	X	Sm75	SH
<i>Apogon quadrisquamatus</i>	sawcheek cardinalfish	Pa			Po/DK	WS/DK
<i>Apogon townsendi</i>	belted cardinalfish	Ca				
<i>Astrapogon alutus</i>	bronze cardinalfish				SW/Sm76	
<i>Astrapogon stellatus</i>	conchfish					DK
<i>Phaeoptyx conklini</i>	freckled cardinalfish	Ca				
<i>Phaeoptyx pigmentaria</i>	dusky cardinalfish	Ca				
<i>Phaeoptyx xenus</i>	sponge cardinalfish				SW/Sm75	
Malacanthidae - tilefishes						
<i>Caulolatilus chrysops</i>	goldface tilefish		HM	X		
<i>Caulolatilus cyanops</i>	blackline tilefish		DB			Po/DG
<i>Malacanthus plumieri</i>	sand tilefish	Ca	Br	X	Sm75	Po
Carangidae – jacks						
<i>Caranx bartholomaei</i>	yellow jack	So				
<i>Caranx latus</i>	horse-eye jack	Ca				
<i>Caranx lugubris</i>	black jack	So				
<i>Caranx ruber</i>	bar jack	Ca			Sm75	
<i>Seriola dumerili</i>	greater amberjack	Ca	Br	X	Sm75	
<i>Seriola rivoliana</i>	almaco jack	Ca		X	Sm75	

Appendix B. (continued).

Scientific Name	Common Name	NW Gulf		Pinnacles 60-110 m	West Florida Shelf	
		<60 m	>60 m		<30 m	>30 m
Lutjanidae – snappers						
<i>Apsilus dentatus</i>	black snapper		HM			
<i>Etelis oculatus</i>	queen snapper					DG
<i>Lutjanus apodus</i>	schoolmaster				SW	
<i>Lutjanus campechanus</i>	red snapper	Ca	Br	X	Sm75	SH
<i>Lutjanus cyanopterus</i>	cubera snapper	Pa			Sm75	
<i>Lutjanus griseus</i>	gray snapper	Ca			SW/Sm75	Cl
<i>Lutjanus jocu</i>	dog snapper	So				
<i>Lutjanus synagris</i>	lane snapper	HM	DB		Sm75	
<i>Ocyurus chrysurus</i>	yellowtail snapper	Ca			SW/Sm75	
<i>Rhomboplites aurorubens</i>	vermilion snapper	Pu	HM	X	Sm75	SW/DG
Haemulidae – grunts						
<i>Anisotremus virginicus</i>	porkfish	HM			SW/DK	
<i>Anisotremus surinamensis</i>	black margate	Pa				
<i>Haemulon aurolineatum</i>	tomtate	Ca			SW/Sm75	
<i>Haemulon melanurum</i>	cottonwick	Ca	Br			
<i>Haemulon macrostomum</i>	Spanish grunt	Pa			SW/DK	
<i>Haemulon plumieri</i>	white grunt	HM			SW/Sm75	
<i>Haemulon sciurus</i>	bluestriped grunt				Po	
<i>Haemulon striatum</i>	striped grunt		DB			Po/DG
Inermiidae – bonnetmouths						
<i>Emmelichthyops atlanticus</i>	bonnetmouth	Pa				
<i>Inermia vittata</i>	boga	Pa			DG	
Sparidae – porgies						
<i>Archosargus probatocephalus</i>	sheepshead	Pu			Sm76	
<i>Calamus bajonado</i>	jolthead porgy				Sm75	DK
<i>Calamus calamus</i>	saucereye porgy	Pa			DG	
<i>Calamus nodosus</i>	knobbed porgy	Ca		X	Sm75	DK

Appendix B. (continued).

Scientific Name	Common Name	NW Gulf		Pinnacles 60-110 m	West Florida Shelf	
		<60 m	>60 m		<30 m	>30 m
Sparidae (continued)						
<i>Calamus penna</i>	sheepshead porgy				DG	
<i>Pagrus pagrus</i>	red porgy		DB	X	Sm75	DK
Sciaenidae – drums						
<i>Equetus lanceolatus</i>	jackknife fish	Ca	Br		Sm75	Cl
<i>Equetus punctatus</i>	spotted drum	So	DB			
<i>Odontoscion dentex</i>	reef croaker	HM			Sm76	
<i>Pareques iwamotoi</i>	blackbar drum		HM	X		SH
<i>Pareques umbrosus</i>	cubbyu	Pu	DB	X	Sm75	SH/DK
Mullidae – goatfishes						
<i>Mulloidichthys martinicus</i>	yellow goatfish	Ca				
<i>Pseudupeneus maculatus</i>	spotted goatfish	Ca	Br		Sm76	
Chaetodontidae –butterflyfishes						
<i>Chaetodon aculeatus</i>	longsnout butterflyfish	Ca	DB			
<i>Chaetodon aya</i>	bank butterflyfish	Pa	Ca	X	Po	SH/DK
<i>Chaetodon capistratus</i>	foureye butterflyfish	So			Sm75	DK
<i>Chaetodon ocellatus</i>	spotfin butterflyfish	Ca		X	SW/Sm75	SH/DK
<i>Chaetodon sedentarius</i>	reef butterflyfish	Ca	Br	X	Sm75	SH/DK
<i>Chaetodon striatus</i>	banded butterflyfish	So			Sm76/DK	
Pomacanthidae – angelfishes						
<i>Centropyge argi</i>	cherubfish	Ca	DB		Sm75	
<i>Holacanthus bermudensis</i>	blue angelfish	Ca	Br	X	SW/Sm5	SH/DK
<i>Holacanthus ciliaris</i>	queen angelfish	Ca		X	SW/Sm75	SH
<i>Holacanthus tricolor</i>	rock beauty	Ca	Br	X	WS	
<i>Pomacanthus arcuatus</i>	gray angelfish	DB			Sm75/DK	
<i>Pomacanthus paru</i>	French angelfish	Ca	DB		SW/Sm75	

Appendix B. (continued).

Scientific Name	Common Name	NW Gulf		Pinnacles 60-110 m	West Florida Shelf	
		<60 m	>60 m		<30 m	>30 m
Kyphosidae - sea chubs						
<i>Kyphosus incisor</i>	Yellow chub	So				
<i>Kyphosus sectatrix</i>	Bermuda chub	Ca			Po/Cl	
Pomacentridae – damselfishes						
<i>Abudefduf saxatilis</i>	sergeant major	Pu			SW/Sm76	
<i>Chromis cyanea</i>	blue chromis	Ca			Sm75	
<i>Chromis enchrysur</i>	yellowtail reeffish	Ca	Br	X	Sm75	SH/DK
<i>Chromis insolata</i>	sunshinefish	DB		X	Po	
<i>Chromis multilineata</i>	brown chromis	Ca				
<i>Chromis scotti</i>	purple reeffish	Ca		X	Sm75	SH/DK
<i>Microspathodon chrysurus</i>	yellowtail damselfish	Ca				DK?
<i>Stegastes adustus</i>	dusky damselfish	Bo				
<i>Stegastes diencaeus</i>	longfin damselfish	Pa				
<i>Stegastes leucostictus</i>	beaugregory	Pu			DK	
<i>Stegastes partitus</i>	bicolor damselfish	Ca			Sm75	DK
<i>Stegastes planifrons</i>	threespot damselfish	Ca				WS
<i>Stegastes variabilis</i>	cocoa damselfish	Ca			SW/Sm75	Cl
Cirrhitidae – hawkfishes						
<i>Amblycirrhitus pinos</i>	redspotted hawkfish	Ca	DB			
Labridae – wrasses						
<i>Bodianus pulchellus</i>	spotfin hogfish	Ca	Br	X	Cl	SH/DG
<i>Bodianus rufus</i>	Spanish hogfish	Ca			Sm75	
<i>Clepticus parrae</i>	creole wrasse	Ca				
<i>Decodon puellaris</i>	red hogfish		HM	X		DK
<i>Doratonotus megalepis</i>	dwarf wrasse				Ha	
<i>Halichoeres bathyphilus</i>	greenband wrasse			X	Po	
<i>Halichoeres bivittatus</i>	slippery dick	Ca	DB		SW/Sm75	Cl
<i>Halichoeres caudalis</i>	painted wrasse	So			SW/Sm75	Cl/DK

Appendix B. (continued).

Scientific Name	Common Name	NW Gulf		Pinnacles	West Florida Shelf	
		<60 m	>60 m	60-110 m	<30 m	>30 m
Labridae (continued)						
<i>Halichoeres garnoti</i>	yellowhead wrasse	Ca				
<i>Halichoeres maculipinna</i>	clown wrasse	Ca				
<i>Halichoeres pictus</i>	rainbow wrasse				Po	CI/DK
<i>Halichoeres radiatus</i>	puddingwife	Ca	DB			
<i>Lachnolaimus maximus</i>	hogfish	So			SW/Sm75	CI
<i>Thalassoma bifasciatum</i>	bluehead	Ca			Sm75	
<i>Xyrichtys martinicensis</i>	rosy razorfish				Po	
<i>Xyrichtys splendens</i>	green razorfish	Pa				Po
Scaridae – parrotfishes						
<i>Scarus coelestinus</i>	midnight parrotfish	Pa			Ha	
<i>Scarus iserti</i>	striped parrotfish	So/Pa			Sm75	
<i>Scarus taeniopterus</i>	princess parrotfish	Ca				
<i>Scarus vetula</i>	queen parrotfish	Ca				
<i>Sparisoma atomarium</i>	greenband parrotfish	DB	DB		WS	
<i>Sparisoma aurofrenatum</i>	redband parrotfish	Ca			Sm75	
<i>Sparisoma chrysopteron</i>	redtail parrotfish				Ha	DG
<i>Sparisoma rubripinne</i>	redfin parrotfish				Ha	
<i>Sparisoma radians</i>	bucktooth parrotfish	HM			Sm75	DK
<i>Sparisoma viride</i>	stoplight parrotfish	Ca			Ha	
Labrisomidae - labrisomids						
<i>Labrisomus haitiensis</i>	longfin blenny				Sm75	
<i>Labrisomus nuchipinnis</i>	hairy blenny	HM				
<i>Starksia ocellata</i>	checkered blenny	Ca			SW/CI	DK
<i>Nemaclinus atelestos</i>	threadfin blenny	Ca				
<i>Paraclinus fasciatus</i>	banded blenny				Po	

B-10

Appendix B. (continued).

Scientific Name	Common Name	NW Gulf		Pinnacles 60-110 m	West Florida Shelf	
		<60 m	>60 m		<30 m	>30 m
Chaenopsidae – pikeblennies						
<i>Chaenopsis limbaughi</i>	yellowface pikeblenny				Po	Po
<i>Chaenopsis ocellata</i>	bluethroat pikeblenny				SW	SW/Po
<i>Emblemaria atlantica</i>	banner blenny		DB		Cl	Cl
<i>Emblemaria pandonis</i>	sailfin blenny	Ca			Sm76	
<i>Emblemaria piratula</i>	pirate blenny				WS	WS
Blenniidae- combtooth blennies						
<i>Hypleurochilus bermudensis</i>	barred blenny	Ca			Po	
<i>Hypleurochilus multifilis</i>	plumed blenny	HM				
<i>Hypleurochilus caudovittatus</i>	zebratail blenny				SW/DK	
<i>Hypsoblennius invemar</i>	tessellated blenny	HM				
<i>Ophioblennius atlanticus</i>	redlip blenny	Ca				
<i>Parablennius marmoreus</i>	seaweed blenny	HM/DB		X	Sm75	DK
<i>Scartella cristata</i>	molly miller	HM				
Gobiidae – gobies						
<i>Coryphopterus eidolon</i>	pallid goby	Pa				
<i>Coryphopterus glaucofraenum</i>	bridled goby				Sm75	
<i>Coryphopterus punctipectophorus</i>	spotted goby	HM/Pa		X	SW/Cl	
<i>Coryphopterus thrix</i>	sand goby	Ca				
<i>Evermannichthys spongicola</i>	sponge goby					Po
<i>Gnatholepis thompsoni</i>	goldspot goby	Ca				
<i>Gobiosoma grosvenori</i>	rockcut goby					DK
<i>Gobiosoma horsti</i>	yellowline goby				Sm75	DK
<i>Gobiosoma macrodon</i>	tiger goby				SW/DK	
<i>Gobiosoma oceanops</i>	neon goby	Ca			Sm75	Cl
<i>Gobiosoma xanthiprora</i>	yellowprow goby	X ³	X ³		Cl	Cl
<i>Gobulus myersi</i>	paleback goby				Cl	

B-11

³new record in NW Gulf

Appendix B. (continued).

Scientific Name	Common Name	NW Gulf		Pinnacles	West Florida Shelf	
		<60 m	>60 m	60-110 m	<30 m	>30 m
Gobiidae (continued)						
<i>Lythrypnus elasson</i>	dwarf goby				CI/WS	WS
<i>Lythrypnus nesiotes</i>	island goby	Ca			Po	WS/DK
<i>Lythrypnus phorellus</i>	convict goby	Ca				
<i>Lythrypnus spilus</i>	bluegold goby	Ca				
<i>Priolepis hipoliti</i>	rusty goby	Ca				
<i>Psilotris celsus</i>	highspine goby				CI	
<i>Risor ruber</i>	tusked goby	Ca		X	Po	WS/DK
Microdesmidae - dartfishes						
<i>Ptereleotris calliurus</i>	blue dartfish	DB	DB		SW/Sm75	CI
Ephippidae – spadefish						
<i>Chaetodipterus faber</i>	Atlantic spadefish	Pu		DK	Sm75	
Bothidae – lefteye flounder						
<i>Bothus lunatus</i>	peacock flounder	Pa				
<i>Bothus ocellatus</i>	eyed flounder	Pa			DK	DK
Acanthuridae – surgeonfishes						
<i>Acanthurus bahianus</i>	ocean surgeonfish	Ca				
<i>Acanthurus chirurgus</i>	doctorfish	So			Sm75	
<i>Acanthurus coeruleus</i>	blue tang	Ca			SW/Sm76	
Sphyraenidae – barracudas						
<i>Sphyraena barracuda</i>	great barracuda	Ca	DB		Sm76	
Balistidae – leatherjackets						
<i>Balistes capriscus</i>	gray triggerfish	Ca		X	Sw/Sm75	DG
<i>Balistes vetula</i>	queen triggerfish	Ca	DB		Sm76	
<i>Canthidermis maculata</i>	rough triggerfish	Ca				
<i>Canthidermis sufflamen</i>	ocean triggerfish	Ca	DB		Po	
<i>Melichthys niger</i>	black durgon	Ca				
<i>Xanthichthys ringens</i>	sargassum triggerfish	Ca	DB		DK	DG

B-12

Appendix B. (continued).

Scientific Name	Common Name	NW Gulf		Pinnacles 60-110 m	West Florida Shelf	
		<60 m	>60 m		<30 m	>30 m
Monacanthidae – filefishes						
<i>Aluterus heudelotii</i>	dotteral filefish				DK	DK
<i>Aluterus monoceros</i>	unicorn filefish	Bo/Pa				
<i>Aluterus scriptus</i>	scrawled filefish	Ca		DK	Cl	DK
<i>Cantherhines macrocerus</i>	whitespotted filefish	So				DK
<i>Cantherhines pullus</i>	orangespotted filefish	Ca				
<i>Monacanthus tuckeri</i>	slender filefish	Ca		X		
<i>Stephanolepis setifer</i>	pygmy filefish	Ca			Sm75	DK
Ostraciidae – boxfishes						
<i>Acanthostracion polygonia</i>	honeycomb cowfish	Pa			DG	
<i>Lactophrys bicaudalis</i>	spotted trunkfish	Pa				
<i>Lactophrys trigonus</i>	trunkfish				SW	
<i>Lactophrys triqueter</i>	smooth trunkfish	Ca			Po	
Tetraodontidae - puffers						
<i>Canthigaster rostrata</i>	sharpnose puffer	Ca	DB	X	Sm75	DK
<i>Canthigaster jamestyeri</i>	goldfaced puffer			X		
<i>Sphoeroides spengleri</i>	bandtail puffer	Pa	?	X	SW/Sm75	SH
Diodontidae – porcupinefishes						
<i>Chilomycterus antennatus</i>	bridled burrfish	Pa				
<i>Chilomycterus antillarum</i>	web burrfish			X		
<i>Diodon holocanthus</i>	balloonfish	So	DB	X	Po	
<i>Diodon hystrix</i>	porcupinefish	Pa				
Triacanthodidae						
<i>Parahollardia lineata</i>	jambeau			DK		DK/DG

B-13

APPENDIX C

Food habits of reef fishes of the Pinnacles Reef Tract

Appendix C. Food habits of reef fishes of the Pinnacles Reef Tract by trophic guild. Number of specimens examined that contained prey are given, along with range of standard length (SL). Values given are based on relative abundance (% number) of prey items in the diet as determined by stomach content analysis.

Trophic Guild: Mesoplanktivores

<i>Bregmaceros cantori</i> 2 specimens: 51 mm SL	codlet	<i>Paranthias furcifer</i> 1 specimen: 245 mm SL	creole-fish
FOOD		FOOD	
Copepods	60.0	Polychaetes	35.7
Amphipods	20.0	Copepods	28.6
Shrimp Larvae	20.0	Shrimp larvae	21.4
		Crab larvae	14.3
<i>Chromis enchrysur</i> 20 specimens: 66-90 mm SL	yellowtail reeffish	<i>Pronotogrammus martinicensis</i> 120 specimens: 70-152 mm SL	rougtongue bass
FOOD		FOOD	
Copepods	32.4	Copepods	69.7
Larvaceans	32.0	Invertebrate eggs	11.0
Invertebrate eggs	19.1	Polychaetes	10.3
Polychaetes	13.3	Fish eggs	2.4
Shrimp and shrimp larvae sergestids	2.1	Larvaceans	1.9
Crab larvae	1.1	Ostracods	1.7
		Radiolarians	1.5
<i>Hemanthias vivanus</i> 47 specimens: 67-110 mm SL	red barbier	Urochordates salps	0.5
FOOD		Crab larvae	0.4
Copepods	78.8	Amphipods hyperiid	0.2
Polychaetes	17.7	Barnacle larvae	0.1
Invertebrate eggs	3.0	Fish larvae	0.05
Larvaceans	0.14	squid	0.05
Ostracods	0.14		
Urochordates salps	0.07		
Amphipods hyperiid	0.04		
Radiolarians	0.03		
Pteropods	0.03		
Chaetognaths	0.02		
Crab larvae	0.01		
Fishes	0.01		
<i>Hemanthias vivanus</i>			
Fish larvae	0.01		
Fish eggs	0.01		

Appendix C. (continued).

Trophic Guild: Macroplanktivores

<i>Apogon pseudomaculatus</i> 5 specimens: 62-71 mm SL	twospot cardinalfish	<i>Priacanthus arenatus</i> 15 specimens: 196-295 mm SL	bigeye
FOOD		FOOD	
Isopods	37.5	Squids	32.9
Crabs	25.0	Stomatopod larvae	27.6
Shrimps and shrimp larvae	25.0	Fish larvae	14.5
Crab larvae	12.5	<i>Scomberomorus</i> sp.	
		<i>Bothus</i> sp.	
		<i>Synodus</i> sp.	
<i>Chaetodon aya</i> 13 specimen: 67-94 mm SL	bank butterflyfish	Crab larvae	14.5
FOOD		Shrimp	3.9
Crab and shrimp larvae	90.1	Fishes	2.6
Amphipods	5.1	nettostomatid	
Gorgonians	4.8	Shrimp larvae	1.3
		Pteropods	1.3
		Crabs	1.3
<i>Pristigenys alta</i> 9 specimens: 134-257mm SL	short bigeye	<i>Rhomboplites aurorubens</i> 72 specimens: 120-372 mm SL	vermillion snapper
FOOD		FOOD	
Shrimp	40.0	Amphipods	15.2
<i>Lucifer faxoni</i>		hyperiids	
Squids	16.0	Copepods	15.2
Shrimp larvae	16.0	Polychaetes	14.3
Crab larvae	16.0	Fish and fish larvae	10.6
Stomatopod larvae	6.0	<i>Bregmaceros cantori</i>	
Fishes	2.0	myctophid	
Crabs	2.0	<i>Prionotus</i> sp.(larvae)	
		<i>Synodus</i> sp. (larvae)	
		Shrimps and shrimp larvae	10.1
		<i>Lucifer faxoni</i>	
		Urochordates	9.9
		salps	
		Stomatopod larvae	8.0
		Pteropods	6.9
		Crab larvae	3.7
		Ostracods	2.1
		Fish eggs	1.4
		Barnacle larvae	1.4
		Larvaceans	1.1
		Squids	0.2

Appendix C. (continued).

Trophic Guild: Benthic Carnivores

<i>Bodianus pulchellus</i> 4 specimens: 155-161 mm SL	spotfin hogfish	<i>Decodon puellaris</i> 2 specimens: 125-132 mm SL	red hogfish
FOOD		FOOD	
Pelecypods	38.1	Pelecypods	37.5
Crabs	19.0	Gastropods	31.3
Fishes	19.0	Crab larvae	12.5
<i>H. vivanus</i>		Polychaetes	6.3
<i>P. martinicensis</i>		Crabs	6.3
Gastropods	14.3	pagurid	
Ophiuroids	9.5	Stomatopod larvae	6.3
<i>Caulolatilus chrysops</i> 2 specimens: 331-443 mm SL	goldface tilefish	<i>Halichoeres bathyphilus</i> 10 specimens: 97-107 mm SL	greenband wrasse
FOOD		FOOD	
Polychaetes	45.0	Pelecypods	48.3
Stomatopods	25.0	Crab larvae	13.8
Pelecypods	20.0	Ophiuroids	6.9
Ophiuroids	5.0	Crabs	6.9
Crabs	5.0	Shrimps and shrimp larvae	6.9
		Foraminifera	6.9
		Fishes	3.4
		Polychaetes	3.4
		Squids	3.4
<i>Pareques iwamotoi</i> 1 specimen: 234 mm SL	blackbar drum	<i>Pagrus pagrus</i> 28 specimens: 147-455 mm SL	red porgy
FOOD		FOOD	
Ophiuroids	33.3	Crabs	34.0
Gastropods	33.3	pagurids	
Crabs	33.3	<i>Munida</i> sp.	
		<i>Calappa</i> sp.	
		<i>Petrolisthes</i> sp.	
		Echinoids	20.2
		spatangoids	
		Gastropods	16.0
		Stomatopod larvae	6.4
		Fishes	5.3
		<i>Pristopomoides aquilonaris</i>	
		<i>Kathetostoma albigutta</i>	
		<i>Hemanthias vivanus</i>	
		Stomatopods	4.3
		Polychaetes	4.3
		Pelecypods	3.2
		Crab Larvae	2.1
		Shrimps	2.1
		alpheid	
		Holothuroids	1.1
		Sponges	1.1
<i>Pareques umbrosus</i> 4 specimens: 110-177 mm SL	cubbyu		
FOOD			
Crabs	36.4		
<i>Munida</i> sp.			
<i>Calappa</i> sp.			
xanthid			
Shrimps	36.4		
Shrimp larvae	9.1		
Ophiuroids	9.1		
Polychaetes	9.1		

Appendix C. (continued).

Trophic Guild: Generalized Carnivores

<i>Centropristis ocyurus</i> 6 specimens: 142-212 mm SL	bank sea bass	<i>Malacanthus plumieri</i> 7 specimens: 345-376 mm SL	sand tilefish
FOOD		FOOD	
Crabs	72.2	Gastropods	25.0
pagurids		Fishes	18.8
portunids		<i>Hemanthias vivanus</i>	
calappid		<i>Pronotogrammus martinicensis</i>	
<i>Munida</i> sp.		Crabs	18.8
Gastropods	11.1	Pelecypods	18.8
Ophiuroids	5.6	Ophiuroids	12.5
Shrimps and shrimp larvae	5.6	Bryozoans	6.3
Coral fragment	5.6		
<i>Epinephelus flavolimbatus</i> 2 specimens: 156-694 mm SL	yellowedge grouper	<i>Opsanus pardus</i> 5 specimens: 194-215 mm SL	leopard toadfish
FOOD		FOOD	
Fishes	66.7	Gastropods	33.3
myctophid		Crabs	25.0
Shrimps	33.3	Fishes	25.0
		<i>Pronogrammus martinicensis</i>	
		Crinoids	8.3
		Echinoids	8.3
<i>Holocentrus adscensionis</i> 3 specimens: 168-228 mm SL	squirrelfish	<i>Sargocentron bullisi</i> 2 specimens: 129-134 mm SL	deepwater squirrelfish
FOOD		FOOD	
Fishes	42.9	Crabs and crab larvae	33.3
<i>Hemanthias vivanus</i>		Stomatopod larvae	33.3
Crab larvae	42.9	Polychaetes	16.7
Shrimps	14.3	Chitons	16.7
alpheid			
<i>Liopropoma eukrines</i> 1 specimen: 70 mm SL	wrasse bass	<i>Serranus phoebe</i> 44 specimens: 107-168 mm SL	tattler
FOOD		FOOD	
Fishes	100.0	Crabs	41.7
<i>Coryphopterus</i> sp.		<i>Munida</i> sp.	
		Xanthids	
 		Fishes	37.5
<i>Lutjanus campechanus</i> 26 specimens: 192-465 mm SL	red snapper	<i>Hemanthias vivanus</i>	
FOOD		<i>Coryphopterus</i> sp.	
Fishes	56.4	Crab larvae	5.6
<i>Hemanthias vivanus</i>		Shrimps	5.6
<i>Serranus</i> sp.		alpheids	
<i>Bregmaceros cantori</i>		Fish larvae	2.8
<i>Hoplunnus</i> sp.		Ophiuroids	1.4
<i>Trachurus lathami</i>		Squids	1.4
Isopods	10.3	Stomatopods	1.4
Urochordates	10.3	Amphipods	1.4
Shrimps	7.7	Polychaetes	1.4
<i>Sicyona</i> sp.			
Heteropods	5.1		
Crabs	5.1		
Stomatopods	2.6		
Squids	2.6		

Appendix C. (continued).

Trophic Guild: Piscivores

<i>Aulostomus maculatus</i> 1 Specimen: 160 mm SL	trumpetfish	<i>Seriola dumerili</i> 6 specimens: 561-744 mm SL	greater amberjack
FOOD Fishes <i>Coryphopterus</i> sp.	100.0	FOOD Fishes <i>Synodus</i> sp. <i>Prionotus</i> sp. <i>Hemanthias vivanus</i> <i>Pronotogrammus martinicensis</i> <i>Trachurus lathami</i>	100.0
<i>Gymnothorax saxicola</i> 1 specimen: 453 mm SL	honeycomb moray		
FOOD Fishes <i>Hippocampus</i> sp.	100.0		

Trophic Guild: Epibenthic Browsers

<i>Chaetodon sedentarius</i> 8 specimens: 92-107mm SL	reef butterflyfish	<i>Holacanthus bermudensis</i> 4 specimens: 210-291mm SL	blue angelfish
FOOD Hydrozoans Gorgonians Copepods Urochordates tunicates Ostracods Shrimp larvae Echinoid spines	71.4 14.4 5.2 3.2 2.2 2.1 1.5	FOOD Bryozoans Gorgonians Crinoids Sponges	41.7 41.7 8.3 8.3
<i>Canthigaster rostrata</i> 2 specimens: 37-40 mm SL	sharpnose puffer		
FOOD Crinoids Bivalves Gastropods Chitons	78.6 7.8 3.2 2.1		

APPENDIX D

Characteristic ichthyofauna of Pinnacles Reef Tract,
displaying the biodiversity of fishes by trophic guild.

Appendix D. Characteristic ichthyofauna of Pinnacles Reef Tract, displaying the biodiversity of fishes by trophic guild. Maximum body size given follows Smith-Vaniz et al. (1999) or Robins and Ray (1986). Depth range is based on videotape analysis or site of collection recorded for each species.

Species	Common Name	Maximum Body Size (mm)	Activity Pattern	Study Sites	Depth Range(m)
Mesoplanktivores					
1. <i>Anthias tenuis</i>	threadnose bass	90 SL	D	A, R, Y	61- 88
2. <i>Bregmaceros cantori</i>	antenna codlet	60 SL	N	R, S	65-110
3. <i>Chromis enchrysur</i>	yellowtail reeffish	105 SL	D	D, R, T, Y	61- 80
4. <i>Hemanthias leptus</i>	longtail bass (adult)	450 SL	D	L, S	100-110
5. <i>Hemanthias leptus</i>	longtail bass (juvenile)				
6. <i>Hemanthias vivanus</i>	red barbier	120 SL	D	All Sites	61-110

FL=fork length, SL=standard length, and TL=total length.

D=Diurnal, N=Nocturnal, C=Crepuscular.

A=Alabama Alps, D=Double Top Reef, L=Ludwick and Walton Pinnacles, R=Roughtongue Reef, S=Scamp Reef, T=Triple Top Reef, Y=Yellowtail Reef.



Appendix D. (continued).

Species	Common Name	Maximum Body Size (mm)	Activity Pattern	Study Sites	Depth Range(m)
Mesoplanktivores (continued)					
7. <i>Paranthias furcifer</i>	creole-fish	350 SL	D	R, Y	61- 80
8. <i>Pronotogrammus martinicensis</i>	rougtongue bass	160 SL	D	All Sites	61-110
Macroplanktivores					
9 <i>Apogon pseudomaculatus</i>	twospot cardinalfish	105 TL	N	All Sites	61-110
10. <i>Chaetodon aya</i>	bank butterflyfish	150 TL	D	All Sites	61-110
11. <i>Opistognathus lonchurus</i>	moustache jawfish	100 TL	D	Y, R	61- 80
12. <i>Priacanthus arenatus</i>	bigeye	360 SL	N	A, L, R, Y	61-110

FL=fork length, SL=standard length, and TL=total length.

D=Diurnal, N=Nocturnal, C=Crepuscular.

A=Alabama Alps, D=Double Top Reef, L=Ludwick and Walton Pinnacles, R=Roughtongue Reef,

S=Scamp Reef, T=Triple Top Reef, Y=Yellowtail Reef.

D-5



Appendix D. (continued).

<u>Species</u>	<u>Common Name</u>	<u>Maximum Body Size (mm)</u>	<u>Activity Pattern</u>	<u>Study Sites</u>	<u>Depth Range(m)</u>
Macroplanktivores (continued)					
13. <i>Pristigenys alta</i>	short bigeye	330 FL	N	All Sites	61-110
14. <i>Rhomboplites aurorubens</i>	vermilion snapper	600 TL	D/N	All Sites	61-110
Generalized Carnivores					
15. <i>Bodianus pulchellus</i>	spotfin hogfish	330 FL	N	R, Y	61- 80
16. <i>Brotula barbata</i>	bearded brotula	330 FL	N	L, S	100-110
17. <i>Centropristis ocyurus</i>	bank sea bass	300 TL	D/N	D, Y	61- 80
18. <i>Corniger spinosus</i>	spinycheek soldierfish	200 TL	N	All Sites	61-110

FL=fork length, SL=standard length, and TL=total length.

D=Diurnal, N=Nocturnal, C=Crepuscular.

A=Alabama Alps, D=Double Top Reef, L=Ludwick and Walton Pinnacles, R=Roughtongue Reef, S=Scamp Reef, T=Triple Top Reef, Y=Yellowtail Reef.



Appendix D. (continued).



Species	Common Name	Maximum Body Size (mm)	Activity Pattern	Study Sites	Depth Range(m)
Generalized Carnivores (continued)					
19. <i>Epinephelus drummondhayi</i>	speckled hind (adult)	1100 TL	D/N	Y	61- 80
20. <i>Epinephelus drummondhayi</i>	speckled hind (juvenile)				
21. <i>Epinephelus flavolimbatus</i>	yellowedge grouper	1150 TL	D/N	L	100-110
22. <i>Epinephelus niveatus</i>	snowy grouper	1200 TL	D/N	D, L	70-110
23. <i>Epinephelus nigritus</i>	warsaw grouper	2300 TL	D/N	A	70- 88
24. <i>Gonioplectrus hispanus</i>	Spanish flag (adult)	230 TL	D/N	All Sites	61-110

FL=fork length, SL=standard length, and TL=total length.

D=Diurnal, N=Nocturnal, C=Crepuscular.

A=Alabama Alps, D=Double Top Reef, L=Ludwick and Walton Pinnacles, R=Roughtongue Reef,

S=Scamp Reef, T=Triple Top Reef, Y=Yellowtail Reef.



Appendix D. (continued).

Species	Common Name	Maximum Body Size (mm)	Activity Pattern	Study Sites	Depth Range(m)
Generalized Carnivores (continued)					
25. <i>Gonioplectrus hispanus</i>	Spanish flag (juvenile)	230 TL	D/N	All Sites	61-110
26. <i>Gymnothorax hubbsi</i>	lichen moray	300 TL	N	A, Y	61- 88
27. <i>Holocentrus adscensionis</i>	squirrelfish	345 SL	N	R, Y	61- 80
28. <i>Liopropoma eukrines</i>	wrasse bass	130 TL	D/N	All Sites	61-110
29. <i>Lutjanus campechanus</i>	red snapper	1000 TL	D/N	All Sites	61-110
30. <i>Malacanthus plumieri</i>	sand tilefish	600 TL	D	R, Y	61- 80

D-10

FL=fork length, SL=standard length, and TL=total length.

D=Diurnal, N=Nocturnal, C=Crepuscular.

A=Alabama Alps, D=Double Top Reef, L=Ludwick and Walton Pinnacles, R=Roughtongue Reef,

S=Scamp Reef, T=Triple Top Reef, Y=Yellowtail Reef.



Appendix D. (continued).


Species	Common Name	Maximum Body Size (mm)	Activity Pattern	Study Sites	Depth Range(m)
Generalized Carnivores (continued)					
31. <i>Opsanus pardus</i>	leopard toadfish	380 TL	D/N	R, Y	61- 80
32. <i>Pontinus rathbuni</i>	highfin scorpionfish	250 TL	D/N	R, Y	61- 80
33. <i>Rypticus maculatus</i>	whitespotted soapfish	200 TL	D/N	R, Y	61- 80
34. <i>Sargocentron bullisi</i>	deepwater squirrelfish	130 SL	N	L, R, Y	61-110
35. <i>Scorpaena dispar</i>	hunchback scorpionfish	300 TL	N	All Sites	61-110
36. <i>Serranus phoebe</i>	tattler	215 SL	D	All Sites	61-110

FL=fork length, SL=standard length, and TL=total length.

D=Diurnal, N=Nocturnal, C=Crepuscular.

A=Alabama Alps, D=Double Top Reef, L=Ludwick and Walton Pinnacles, R=Roughtongue Reef,

S=Scamp Reef, T=Triple Top Reef, Y=Yellowtail Reef.



Appendix D. (continued)

Species	Common Name	Maximum Body Size (mm)	Activity Pattern	Study Sites	Depth Range(m)
Piscivores					
37. <i>Aulostomus maculatus</i>	trumpetfish	750 TL	D	A, R, Y	61- 88
38. <i>Fistularia petimba</i>	red cornetfish	1800 TL	D/N	R, S	65-110
39. <i>Gymnothorax kolpos</i>	blacktail moray	900 TL	D/N	All Sites	61-110
40. <i>Muraena retifera</i>	reticulated moray	900 TL	D/N	All Sites	61-110
41. <i>Mycteroperca microlepis</i>	gag	1200 TL	D/C	All Sites	61-110
42. <i>Mycteroperca phenax</i>	scamp	900 TL	D/C	All Sites	61-110

D-14

FL=fork length, SL=standard length, and TL=total length.

D=Diurnal, N=Nocturnal, C=Crepuscular.

A=Alabama Alps, D=Double Top Reef, L=Ludwick and Walton Pinnacles, R=Roughtongue Reef,

S=Scamp Reef, T=Triple Top Reef, Y=Yellowtail Reef.



Appendix D. (continued).





Species	Common Name	Maximum Body Size (mm)	Activity Pattern	Study Sites	Depth Range(m)
Piscivores (continued)					
43. <i>Seriola fasciata</i>	lesser amberjack	680 FL	D/N	All Sites	61-110
44. <i>Synodus intermedius</i>	sand diver	380 SL	D/N	Y, L	61-110
Benthic Carnivores					
45. <i>Caulolatilus chrysops</i>	goldface tilefish	600 TL	D/N	L	100-110
46. <i>Caulolatilus intermedius</i>	anchor tilefish	600 TL	D/N	L	100-110
47. <i>Calamus leucosteus</i>	whitebone porgy	460 TL	D/N	Y, R	61- 80
48. <i>Calamus nodosus</i>	knobbed porgy	460 TL	D/N	Y, R	61- 80

FL=fork length, SL=standard length, and TL=total length.

D=Diurnal, N=Nocturnal, C=Crepuscular.

A=Alabama Alps, D=Double Top Reef, L=Ludwick and Walton Pinnacles, R=Roughtongue Reef,

S=Scamp Reef, T=Triple Top Reef, Y=Yellowtail Reef.



D-17

Appendix D. (continued)

Species	Common Name	Maximum Body Size (mm)	Activity Pattern	Study Sites	Depth Range(m)
Benthic Carnivores (continued)					
49. <i>Decodon puellaris</i>	red hogfish	300 TL	D	All Sites	61-110
50. <i>Halichoeres bathyphilus</i>	greenband wrasse (f)	230 TL	D	All Sites	61-110
51. <i>Halichoeres bathyphilus</i>	greenband wrasse (m)				
52. <i>Ogcocephalus declivirostris</i>	slantbrow batfish	165 TL	D/N	All Sites	61-110
53. <i>Pagrus pagrus</i>	red porgy	910 TL	D/N	All Sites	61-110
54. <i>Pareques iwamotoi</i>	blackbar drum	300 TL	D/N	A, L, S	70-110




D-18

FL=fork length, SL=standard length, and TL=total length.

D=Diurnal, N=Nocturnal, C=Crepuscular.

A=Alabama Alps, D=Double Top Reef, L=Ludwick and Walton Pinnacles, R=Roughtongue Reef,

S=Scamp Reef, T=Triple Top Reef, Y=Yellowtail Reef.



Appendix D. (continued)

Species	Common Name	Maximum Body Size (mm)	Activity Pattern	Study Sites	Depth Range(m)
Benthic Carnivores (continued)					
55. <i>Pareques umbrosus</i>	cubbyu	250 TL	D/N	All Sites	61-110
Epibenthic Browsers					
56. <i>Chaetodon sedentarius</i>	reef butterflyfish	150 TL	D	A, D, R, Y	61- 88
57. <i>Chaetodon ocellatus</i>	spotfin butterflyfish	200 TL	D	R, Y	61- 80
58. <i>Canthigaster jamestyeri</i>	goldface sharpnose puffer	110 TL	D	R, Y	61- 80
59. <i>Canthigaster rostrata</i>	sharpnose puffer	110 TL	D	A, R, Y	61- 88
60. <i>Holacanthus bermudensis</i>	blue angelfish	440 TL	D	All Sites	61-110

FL=fork length, SL=standard length, and TL=total length.

D=Diurnal, N=Nocturnal, C=Crepuscular.

A=Alabama Alps, D=Double Top Reef, L=Ludwick and Walton Pinnacles, R=Roughtongue Reef,

S=Scamp Reef, T=Triple Top Reef, Y=Yellowtail Reef.



55



58



56



59



57



60

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE June 2002	3. REPORT TYPE AND DATES COVERED Final Report 04/15/1997-09/30/2001	
4. TITLE AND SUBTITLE Northeastern Gulf of Mexico Coastal and Marine Ecosystem Program Community Structure and Trophic Ecology of Fishes on the Pinnacles Reef Tract			5. FUNDING NUMBERS	
6. AUTHOR(S) Douglas C. Weaver, George D. Dennis, and Kenneth J. Sulak				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Geological Survey Florida Caribbean Science Center 7920 NW 71st Str. Gainesville, Florida 32653			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Geological Survey Eastern Regional Office 1700 Leetown Road Kearneysville, WV 25430			10. SPONSORING / MONITORING AGENCY REPORT NUMBER USGS BSR 2001-0008 OCS Study MMS 2002-0034	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 Words) Topographic prominences form reef-like features along the outer continental shelf (OCS) edge and continental slope of the southeastern United States and Gulf of Mexico. An extensive set of reef features, known as the Pinnacles Reef Tract, occurs off Mississippi and Alabama at depths between 60 and 110 m. Eighteen sites within the combined Mississippi-Alabama Marine Ecosystems Study and Mississippi-Alabama Shelf Pinnacle Trend Habitat Mapping Study area were surveyed. All sampling combined returned over 6,000 specimens for food habits analyses, taxonomic verification and documentation, and subsequent life history analyses. The results of this study have expanded the number of fishes documented from the Pinnacles Reef Tract, and is the first extensive collection of deep (greater than 60 m) reef fishes made in this region, and possibly worldwide. Our present knowledge of the Pinnacles ichthyofauna can be used to characterize the fundamental taxonomic composition and trophic structure of deep reef-fish communities throughout southeastern United States and Gulf of Mexico OCS areas and therefore be useful in assessing future anthropogenic impacts on faunal structure.				
14. SUBJECT TERMS Pinnacles, geology, deep reef, hard bottom communities, fish, trophic ecology, food web, community ecology			15. NUMBER OF PAGES 94 pp. + apps	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT SAR	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-18
298-102



The Department of the Interior Mission

As the Nation's principal conservation agency, the Department of the Interior has responsibility for most of our nationally owned public lands and natural resources. This includes fostering sound use of our land and water resources; protecting our fish, wildlife, and biological diversity; preserving the environmental and cultural values of our national parks and historical places; and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and works to ensure that their development is in the best interests of all our people by encouraging stewardship and citizen participation in their care. The Department also has a major responsibility for American Indian reservation communities and for people who live in island territories under U.S. administration.



The Minerals Management Service Mission

As a bureau of the Department of the Interior, the Minerals Management Service's (MMS) primary responsibilities are to manage the mineral resources located on the Nation's Outer Continental Shelf (OCS), collect revenue from the Federal OCS and onshore Federal and Indian lands, and distribute those revenues.

Moreover, in working to meet its responsibilities, the **Offshore Minerals Management Program** administers the OCS competitive leasing program and oversees the safe and environmentally sound exploration and production of our Nation's offshore natural gas, oil and other mineral resources. The **MMS Royalty Management Program** meets its responsibilities by ensuring the efficient, timely and accurate collection and disbursement of revenue from mineral leasing and production due to Indian tribes and allottees, States and the U.S. Treasury.

The MMS strives to fulfill its responsibilities through the general guiding principles of: (1) being responsive to the public's concerns and interests by maintaining a dialogue with all potentially affected parties and (2) carrying out its programs with an emphasis on working to enhance the quality of life for all Americans by lending MMS assistance and expertise to economic development and environmental protection.



The U.S. Geological Survey Mission

The U.S. Geological Survey (USGS) is a world leader in the natural sciences through our scientific excellence and responsiveness to society's needs. The USGS serves the Nation by providing reliable scientific information to 1) describe and understand the Earth; 2) minimize loss of life and property from natural disasters; 3) manage water, biological, energy, and mineral resources; and 4) enhance and protect our quality of life.