(c)2001-2005 Mineral Data Publishing, version 1

Crystal Data: Tetragonal. Point Group: n.d. As small tabular crystals, and in foliated masses.

**Physical Properties:** Cleavage:  $\{001\}$ , distinct. Hardness = 2.5-3 D(meas.) = 7.14 D(calc.) = [7.30]

**Optical Properties:** Translucent. *Color*: Greenish yellow to yellow. *Luster*: Vitreous on cleavages, greasy on fractures.

Optical Class: Uniaxial (-); biaxial in part, probably due to intergrown heliophyllite.  $\omega = 2.32(2)$  (Li)  $\epsilon = 2.25(2)$ 

Cell Data: Space Group: n.d. a = 10.82 c = 25.60 Z = 8

X-ray Powder Pattern: Långban, Sweden.

2.85 (100), 3.66 (80), 2.72 (80), 2.07 (70), 1.591 (70), 1.92 (60), 1.652 (50)

Chemistry:

	(1)	(2)
$\mathrm{As_2O_3}$	10.60	12.01
PbO	83.45	81.32
Cl	8.00	8.61
$-O = Cl_2$	1.81	1.94
Total	100.24	100.00

(1) Långban, Sweden. (2) Pb<sub>6</sub>As<sub>2</sub>O<sub>7</sub>Cl<sub>4</sub>.

Occurrence: A rare mineral in metamorphosed Fe–Mn orebodies (Sweden).

**Association:** Heliophyllite, finnemanite, freedite, lead, copper, manganoan calcite, andradite, magnetite (Långban, Sweden).

**Distribution:** In Sweden, from Långban, in the Harstigen mine, near Persberg, and at Jakobsberg, Värmland. From Laurium, Greece, in slag.

**Name:** From the Greek for *unusual*, in allusion to the composition.

**References:** (1) Palache, C., H. Berman, and C. Frondel (1951) Dana's system of mineralogy, (7th edition), v. II, 1036–1037. (2) Welin, E. (1968) X-ray powder data for minerals from Långban and the related mineral deposits of Central Sweden. Arkiv Mineral. Geol., 4(30), 499–541.