



Flaccisagitta enflata

(Grassi, 1881)



Phylum
Order
Family

Chaetognatha
Aphragmophora
Sagittidae



Alvarino (1967), SERTC (2011), Pierrot-Bults & Chidgey (1988)

Synonyms
Sagitta enflata Grassi, 1881

Size

- Maximum adult body length 25 mm, tail 14-17% of length

Genus notes

- Limp, flaccid body
- Lateral fins show distinctive rayless zones
- No intestinal diverticula, hard to see
- Anterior fins separate from posterior fins (no connecting bridge of tissue) and start some distance behind ventral ganglion

Species notes

- Body flaccid, transparent, widest around mid length
- Hooks not serrated
- Anterior fins short, partially rayed, round, not close to ventral ganglion
- Posterior fins short, partially rayed, round
- Collarette absent
- Small eyes, with star-shaped pigment spot
- Seminal vesicles round and touching tail fin, well separated from posterior fins
- Ovaries short, reaching to middle of posterior fins; ova large


Distribution

Ecology

Source

Conway (2003)
Gibbons pers. comm. (2013)
Marine species identification portal (2011)

(Full reference available at <http://www.imas.utas.edu.au/zooplankton/references>)



A.Slotwinski, CSIRO © 2011



Compiled: C. H. Davies & A. S. Slotwinski 2012
Verified: M. J. Gibbons 2013 (preliminary)



Flaccisagitta enflata

(Grassi, 1881)

Phylum
Order
Family

Chaetognatha
Aphragmophora
Sagittidae

