

OPHTHALMIC FORMULATIONS CATALOG

Order from our FDA-Registered 503B Outsourcing Facility
and PCAB Accredited 503A Patient Specific Pharmacy



At Harrow, we are committed to earning your trust every day by providing quality products and exceptional service! We back this commitment up with a **100% money-back guarantee** on every order placed.

- John Saharek
Chief Commercial Officer



EXCEPTIONAL CUSTOMER SERVICE

MONDAY THROUGH FRIDAY **5:30AM - 8PM (PST)**
SATURDAY AND SUNDAY **8:30AM - 5PM (PST)**



Contact us via live Chat **7 days a week**
at [imprimisrx.com](https://www.imprimisrx.com)



Call us at **(844) 446-6979** or text us at
(858) 264-2082 for fast and easy order processing




Orders ship within **24 business hours**
(complete information required)



Auto refill available for all patient
specific prescriptions



Ask us about our **PharmaPack®** to
bundle and save



OUR COMMITMENT TO QUALITY

Our 503B-FDA Registered Outsourcing facility is **inspected by the FDA.**

We use bulk active and inactive substances obtained from **FDA-registered manufacturers**, accompanied by a certificate of analysis.

Release and stability testing performed at **FDA-registered labs.**

Sterility, endotoxin and potency testing performed on each batch.

Continuous environmental monitoring to maintain product quality standards.

WHAT SETS US APART?

	Standard 503A Pharmacy Requirements	503A ImprimisRx Pharmacy Requirements	503B Outsourcing Facility Requirements
API TESTING / RAW MATERIALS	No requirement	Verify manufacturer certificate of analysis	Identity test for each incoming lot
POTENCY TESTING	Random lot testing for potency	All finished product lots	All finished product lots
STERILITY TESTING	Sterile lots per USP <71>	Provide results for sterile lots per USP <71> for every batch	Provide results for sterile lots per USP <71> or equivalent for every batch
ENDOTOXIN TESTING	Sterile injectable lots per USP <85>	Sterile injectable lots per USP <85>	Sterile injectable lots per USP <85>
PRE-SHIPMENT QUARANTINE	Not required, but recommended	14 days for solution 21 days for suspension	14 days for solution 21 days for suspension
ENVIRONMENTAL TESTING	Every 6 months	Operator/ISO 5 with every batch. Facilities twice a month.	Continuous (24 hours/day; 7 days/week)
TEST RESULTS INCLUDED WITH ORDER	No requirement	Sterility results Endotoxin results	Sterility results Endotoxin results
BEYOND USE DATING (EXPIRATION)	Literature and pharmacist experience based, but not required. Stability study over time recommended.	Real time stability using validated stability indicating methods.	Stability study data from three lots using validated stability indicating methods.

OUR APPROACH

Committed to providing access to high-quality medications for you and your patients at affordable prices.

Providing High-Quality Formulations



ImprimisRx® is committed to delivering high-quality formulations from our patient specific 503A pharmacy and our FDA-Registered Outsourcing Facility.

Making Medications Accessible and Affordable




We work with leading ophthalmologists and optometrists to ensure patients have access to critical sight-saving medications at affordable prices.

Evolving Beyond Compounding



We are continuing to take our platform to the next level. Our offering includes FDA-approved brands in addition to compounded formulations.



Commonly Requested Formulations*



Tri-Moxi+®

INJECTABLE SUSPENSION

PRESERVATIVE-FREE

Steroid

Antibiotic

Triamcinolone Acetonide 15mg/mL
Moxifloxacin 1mg/mL



Unit Size	Single-use vial
503A	\$27 per vial
503B	\$540 (20 vials)
NDC #	71384-0746-06

Dex-Moxi®

INJECTABLE SOLUTION

PRESERVATIVE-FREE

Steroid

Antibiotic

Dexamethasone Phosphate 1mg/mL
Moxifloxacin 5mg/mL



Unit Size	Single-use vial
503A	\$31 per vial
503B	\$620 (20 vials)
NDC #	71384-0512-01

Dex-Moxi-Ketor®

INJECTABLE SOLUTION

PRESERVATIVE-FREE

Steroid

Antibiotic

NSAID

Dexamethasone Phosphate 1mg/mL
Moxifloxacin 0.5mg/mL
Ketorolac Tromethamine 0.4mg/mL



Unit Size	Single-use vial
503A	\$34 per vial
503B	\$680 (20 vials)
NDC #	71384-0513-01

These products carry serious risks. Risks associated with transzonular injection via cannula of “Dropless” products include: ciliary body hemorrhage, vitreous floaters, and blurred vision. Please see pages 23-27 for a summary of risks.

Free Shipping Included

*For professional use only. ImprimisRx specializes in customizing medications to meet unique patient and practitioner needs. No compounded medication is reviewed by the FDA for safety and efficacy. ImprimisRx does not compound essential copies of commercially available products. These ingredients carry risks. Visit imprimisrx.com/productinformation for full product details. References available upon request.

OTHER INJECTABLES

Moxifloxacin

INJECTABLE SOLUTION

PRESERVATIVE-FREE

Antibiotic

Moxifloxacin 5mg/mL



Unit Size	Single-use vial
503A	\$18.50 per vial
503B	\$370 (20 vials)
NDC #	71384-0511-08

Moxifloxacin

INJECTABLE SOLUTION

PRESERVATIVE-FREE

Antibiotic

Moxifloxacin 1mg/mL



Unit Size	Single-use vial
503A	\$18.50 per vial
503B	\$370 (20 vials)
NDC #	71384-0509-08

Phenyl-Lido

INJECTABLE SOLUTION

PRESERVATIVE-FREE / SULFITE-FREE

Mydriatic

Phenylephrine HCl 1.5%
Lidocaine HCl 1%



Unit Size	Single-use vial
503A	\$18.50 per vial
503B	\$370 (20 vials)
NDC #	71384-0650-01

Epi-Lido

INJECTABLE SOLUTION

PRESERVATIVE-FREE / SULFITE-FREE

Mydriatic

Epinephrine 0.025%
Lidocaine HCl 0.75% in BSS



Unit Size	Single-use vial
503A	\$18.50 per vial
503B	\$370 (20 vials)
NDC #	71384-0640-01
Storage	Store refrigerated

These products carry serious risks. Please see pages 23-27 for a summary of risks.

Free Shipping Included

*For professional use only. ImprimisRx specializes in customizing medications to meet unique patient and practitioner needs. No compounded medication is reviewed by the FDA for safety and efficacy. ImprimisRx does not compound essential copies of commercially available products. These ingredients carry risks. Visit imprimisrx.com/productinformation for full product details. References available upon request.

Commonly Requested Formulations*

Mydriatic 2

TOPICAL SOLUTION

Mydriatic

Tropicamide 1%
Phenylephrine HCl 2.5%



Unit Size	Single-use	10mL (multi-dose)
503A	\$11 per bottle	-
503B	\$220 (20 bottles)	\$300 (10 bottles)
NDC #	71384-0666-05	71384-0666-10



Mydriatic 3

TOPICAL SOLUTION

Mydriatic

Tropicamide 1%
Cyclopentolate 1%
Phenylephrine HCl 2.5%

Unit Size	Single-use
503A	\$11 per bottle
NDC #	71384-0631-01



Mydriatic 4

TOPICAL SOLUTION

Mydriatic

Tropicamide 1%
Proparacaine HCl 0.5%
Phenylephrine HCl 2.5%
Ketorolac tromethamine 0.5%

Unit Size	Single-use	5mL (multi-dose)
503A	\$11 per bottle	-
503B	\$220 (20 bottles)	\$148 (2 bottles)
NDC #	71384-0632-05	71384-0632-05

These products carry serious risks. Please see pages 23-27 for a summary of risks.

Free Shipping Included

*For professional use only. ImprimisRx specializes in customizing medications to meet unique patient and practitioner needs. No compounded medication is reviewed by the FDA for safety and efficacy. ImprimisRx does not compound essential copies of commercially available products. These ingredients carry risks. Visit imprimisrx.com/productinformation for full product details. References available upon request.

Commonly Requested Formulations*



NON-INVASIVE, NON-OPIOID

PATENTED SUBLINGUAL COMPOUNDED FORMULATION

MKO Melt®

SUBLINGUAL TROCHE

Anxiolytic

Analgesic

Antiemetic

Midazolam 3mg
Ketamine HCl 25mg
Ondansetron 2mg



Unit Size 1 Melt

503B \$388 (21 Melts)

NDC # 71384-0630-21

These products carry serious risks. Please see pages 23-27 for a summary of risks.

Free Shipping Included

*For professional use only. ImprimisRx specializes in customizing medications to meet unique patient and practitioner needs. No compounded medication is reviewed by the FDA for safety and efficacy. ImprimisRx does not compound essential copies of commercially available products. These ingredients carry risks. Visit imprimisrx.com/productinformation for full product details. References available upon request.

Commonly Requested Formulations*



Pred-Moxi-Brom®
TOPICAL SUSPENSION

Steroid Antibiotic NSAID

Prednisolone Acetate 1%
Moxifloxacin 0.5%
Bromfenac 0.075%

Unit Size	5mL	8mL
503A: Patient	\$57 per bottle	\$83.50 per bottle
503A: Office	\$45 per bottle	\$80 per bottle
503B	\$900 (20 bottles)	\$1,600 (20 bottles)
NDC #	71384-0310-05	71384-0310-08



Pred-Moxi-Brom
TOPICAL SOLUTION

Steroid Antibiotic NSAID

Prednisolone Phosphate 1%
Moxifloxacin 0.5%
Bromfenac 0.075%

Unit Size	8mL
503A: Patient	\$83.50 per bottle
503A: Office	\$80 per bottle
503B	\$1,600 (20 bottles)
NDC #	71384-0340-08



Pred-Gati-Brom®
TOPICAL SUSPENSION

Steroid Antibiotic NSAID

Prednisolone Acetate 1%
Gatifloxacin 0.5%
Bromfenac 0.075%

Unit Size	5mL	8mL
503A: Patient	\$57 per bottle	\$83.50 per bottle
503A: Office	\$45 per bottle	\$80 per bottle
NDC #	71384-0504-05	71384-0504-08



Pred-Gati-Brom
TOPICAL SOLUTION

Steroid Antibiotic NSAID

Prednisolone Phosphate 1%
Gatifloxacin 0.5%
Bromfenac 0.075%

Unit Size	5mL	8mL
503A: Patient	\$57 per bottle	\$83.50 per bottle
503A: Office	\$45 per bottle	\$80 per bottle
NDC #	71384-0550-05	71384-0550-08

These products carry serious risks. Please see pages 23-27 for a summary of risks.

Free Shipping Included

*For professional use only. ImprimisRx specializes in customizing medications to meet unique patient and practitioner needs. No compounded medication is reviewed by the FDA for safety and efficacy. ImprimisRx does not compound essential copies of commercially available products. These ingredients carry risks. Visit imprimisrx.com/productinformation for full product details. References available upon request.

Commonly Requested Formulations*



Pred-Moxi-Nepaf®
TOPICAL SUSPENSION

Steroid Antibiotic NSAID

Prednisolone Acetate 1%
Moxifloxacin 0.5%
Nepafenac 0.1%

Unit Size	5mL	8mL
503A: Patient	\$57 per bottle	\$83.50 per bottle
503A: Office	\$45 per bottle	\$80 per bottle
503B	\$900 (20 bottles)	\$1,600 (20 bottles)
NDC #	71384-0320-05	71384-0320-08



Pred-Nepaf®
TOPICAL SUSPENSION

Steroid NSAID

Prednisolone Acetate 1%
Nepafenac 0.1%

Unit Size	5mL
503A: Patient	\$55 per bottle
503A: Office	\$39 per bottle
503B	\$780 (20 bottles)
NDC #	71384-0400-05



Pred-Moxi®
TOPICAL SUSPENSION

Steroid Antibiotic

Prednisolone Acetate 1%
Moxifloxacin 0.5%

Unit Size	5mL
503A: Patient	\$55 per bottle
503A: Office	\$39 per bottle
503B	\$780 (20 bottles)
NDC #	71384-0300-05



Moxi-Brom®
TOPICAL SOLUTION

Antibiotic NSAID

Moxifloxacin 0.5%
Bromfenac 0.075%

Unit Size	5mL
503B	\$780 (20 bottles)
NDC #	71384-0311-05



Pred-Brom®
TOPICAL SUSPENSION

Steroid NSAID

Prednisolone Acetate 1%
Bromfenac 0.075%

Unit Size	5mL
503A: Patient	\$55 per bottle
503A: Office	\$39 per bottle
503B	\$780 (20 bottles)
NDC #	71384-0505-05



Pred-Acetate
TOPICAL SUSPENSION

PRESERVATIVE-FREE

Steroid

Prednisolone Acetate 1%

Unit Size	5mL
503A: Patient	\$49.50 per bottle
503A: Office	\$41 per bottle
503B	\$410 (10 bottles)
NDC #	71384-0501-05

These products carry serious risks. Please see pages 23-27 for a summary of risks.

Free Shipping Included

*For professional use only. ImprimisRx specializes in customizing medications to meet unique patient and practitioner needs. No compounded medication is reviewed by the FDA for safety and efficacy. ImprimisRx does not compound essential copies of commercially available products. These ingredients carry risks. Visit imprimisrx.com/productinformation for full product details. References available upon request.

Commonly Requested Formulations*



Any Single Drug
\$19/month[†]

Any Combo Drug
\$39/month[†]

Bim

Single

TOPICAL SOLUTION
PRESERVATIVE-FREE

Prostaglandin Analogue

Bimatoprost 0.01%



Unit Size 5mL

503A: Patient \$38.00 per bottle

Months Supply[†] 2 months

NDC # 71384-0541-05

Dor

Single

TOPICAL SOLUTION
PRESERVATIVE-FREE

Carbonic Anhydrase Inhibitor

Dorzolamide 2%



Unit Size 10mL

503A: Patient \$38.00 per bottle

Months Supply[†] 2 months

NDC # 71384-0516-10

Brim-Dor[®]

Combo

TOPICAL SOLUTION
PRESERVATIVE-FREE

Alpha Adrenergic Agonist

Carbonic Anhydrase Inhibitor

Brimonidine 0.15%
Dorzolamide 2%



Unit Size 10mL

503A: Patient \$78.00 per bottle

Months Supply[†] 2 months

NDC # 71384-0518-10

Tim-Bim

Combo

TOPICAL SOLUTION
PRESERVATIVE-FREE

Beta-blocker

Prostaglandin Analogue

Timolol 0.5%
Bimatoprost 0.01%



Unit Size 5mL

503A: Patient \$78.00 per bottle

Months Supply[†] 2 months

NDC # 71384-0542-05

These products carry serious risks. Please see pages 23-27 for a summary of risks.

Free Shipping Included

Important: Patients may need to take more than one eye drop product pursuant to multiple dosing regimens, as directed by his or her prescriber, in order for the active ingredients to remain effective throughout the day.

[†]Months supply can vary based on the dosing regimen prescribed by the doctor.

*For professional use only. ImprimisRx specializes in customizing medications to meet unique patient and practitioner needs. No compounded medication is reviewed by the FDA for safety and efficacy. ImprimisRx does not compound essential copies of commercially available products. These ingredients carry risks. Visit imprimisrx.com/productinformation for full product details. References available upon request.

Commonly Requested Formulations*



Triple/Quad Kit
Tim-Brim-Dor® 5mL
Tim-Brim-Dor-Bim 5mL

\$156.00 per kit
(2 months supply[†])

Tim-Dor-Bim
TOPICAL SOLUTION
PRESERVATIVE-FREE

Combo

Beta-blocker

Prostaglandin Analogue

Carbonic Anhydrase Inhibitor

Timolol 0.5%
Dorzolamide 2%
Bimatoprost 0.01%

Unit Size 5mL

503A: Patient \$78.00 per bottle

Months Supply[†] 2 months

NDC # 71384-0543-05



Tim-Brim-Dor®
TOPICAL SOLUTION
PRESERVATIVE-FREE

Combo

Beta-blocker

Alpha Adrenergic Agonist

Carbonic Anhydrase Inhibitor

Timolol 0.5%
Brimonidine 0.15%
Dorzolamide 2%

Unit Size 10mL (2X 5mL bottles)

503A: Patient \$78.00 per 10mL

Months Supply[†] 2 months

NDC # 71384-0520-05



Tim-Brim-Dor-Bim Combo

TOPICAL SOLUTION
PRESERVATIVE-FREE

Beta-blocker

Alpha Adrenergic Agonist

Carbonic Anhydrase Inhibitor

Prostaglandin Analogue

Timolol 0.5% Brimonidine 0.15%
Dorzolamide 2% Bimatoprost 0.01%

Unit Size 5mL

503A: Patient \$78.00 per bottle

Months Supply[†] 2 months

NDC # 71384-0544-05



These products carry serious risks. Please see pages 23-27 for a summary of risks.

Free Shipping Included

Important: Patients may need to instill more than one eye drop product pursuant to multiple dosing regimens, as directed by his or her prescriber, in order for the active ingredients to remain effective throughout the day.

[†]Months supply can vary based on the dosing regimen prescribed by the doctor.

*For professional use only. ImprimisRx specializes in customizing medications to meet unique patient and practitioner needs. No compounded medication is reviewed by the FDA for safety and efficacy. ImprimisRx does not compound essential copies of commercially available products. These ingredients carry risks. Visit imprimisrx.com/productinformation for full product details. References available upon request.

Commonly Requested Formulations*



Klarity-C Drops®

TOPICAL EMULSION

PRESERVATIVE-FREE

Immune Modulator



Cyclosporine 0.1%

Unit Size 5.5mL

503A: Patient \$59 per bottle or 3 for \$165

503B \$480 (10 bottles)

NDC # 71384-0514-05

No Prior Authorizations

No Coupons

No Pharmacy Hassles

Klarity Drops®

TOPICAL SOLUTION

PRESERVATIVE-FREE

Demulcent

Glycerin 1%



Unit Size 10mL

503A: Patient \$18

503B \$360 (20 bottles)

NDC # 71384-0500-10

These products carry serious risks. Please see pages 23-27 for a summary of risks.

Free Shipping Included

*For professional use only. ImprimisRx specializes in customizing medications to meet unique patient and practitioner needs. No compounded medication is reviewed by the FDA for safety and efficacy. ImprimisRx does not compound essential copies of commercially available products. These ingredients carry risks. Visit imprimisrx.com/productinformation for full product details. References available upon request.

Commonly Requested Formulations*



Klarity-A® Drops

TOPICAL SOLUTION

PRESERVATIVE-FREE

Antibiotic

Azithromycin 1%



Unit Size **3.5mL**

503A: Patient **\$54**

NDC # **71384-0220-03**

Klarity-L Drops®

TOPICAL SUSPENSION

PRESERVATIVE-FREE

Steroid

Loteprednol Etabonate 0.5%



Unit Size **5mL**

503A: Patient **\$54**

NDC # **71384-0210-05**

Klarity-CL® Drops

TOPICAL SUSPENSION

PRESERVATIVE-FREE

Immune Modulator

Steroid

**Cyclosporine 0.1%
Loteprednol Etabonate 0.2%**



Unit Size **5mL**

503A: Patient **\$69**

NDC # **71384-0321-05**

These products carry serious risks. Please see pages 23-27 for a summary of risks.

Free Shipping Included

*For professional use only. ImprimisRx specializes in customizing medications to meet unique patient and practitioner needs. No compounded medication is reviewed by the FDA for safety and efficacy. ImprimisRx does not compound essential copies of commercially available products. These ingredients carry risks. Visit imprimisrx.com/productinformation for full product details. References available upon request.

Commonly Requested Formulations*

OTHER DROPS

Atropine Sulfate 0.01%

TOPICAL SOLUTION

PRESERVATIVE-FREE

Anticholinergic



Unit Size 5mL
 503A \$62 for 1 bottle
 \$104 for 2 bottles
 \$117 for 3 bottles
 503B \$390 (10 bottles)
 NDC # 71384-0566-05

Atropine Sulfate 0.025%

TOPICAL SOLUTION

PRESERVATIVE-FREE

Anticholinergic



Unit Size 5mL
 503A \$62 for 1 bottle
 \$104 for 2 bottles
 \$117 for 3 bottles
 503B \$390 (10 bottles)
 NDC # 71384-0567-05

Atropine Sulfate 0.05%

TOPICAL SOLUTION

PRESERVATIVE-FREE

Anticholinergic



Unit Size 5mL
 503A \$62 for 1 bottle
 \$104 for 2 bottles
 \$117 for 3 bottles
 503B \$390 (10 bottles)
 NDC # 71384-0568-05

Fortisite®

Tobramycin 1.5% Vancomycin 5%

TOPICAL SOLUTION

Antibiotic



Unit Size 7mL
 503A: Patient \$299 per bottle
 \$249 for 2 or more bottles
 503B \$299 per bottle
 \$249 for 2 or more bottles
 NDC # 71384-0721-07
 Storage Store Refrigerated

These products carry serious risks. Please see pages 23-27 for a summary of risks.

Free Shipping Included

*For professional use only. ImprimisRx specializes in customizing medications to meet unique patient and practitioner needs. No compounded medication is reviewed by the FDA for safety and efficacy. ImprimisRx does not compound essential copies of commercially available products. These ingredients carry risks. Visit imprimisrx.com/productinformation for full product details. References available upon request.

List of Formulation NDCs (Sorted Alphabetically)

Formulations*	Concentration	Unit Size	NDC
Atropine sulfate PF	(0.01%)	5mL	71384-0566-05
Atropine sulfate PF	(0.025%)	5mL	71384-0567-05
Atropine sulfate PF	(0.05%)	5mL	71384-0568-05
Azithromycin in Klarity	(1%)	3.5mL	71384-0220-03
Bimatoprost	(0.01%)	5mL	71384-0541-05
Brim-Dor PF (Brimonidine and dorzolamide PF)	(0.15/2)%	10mL	71384-0518-10
Cyclosporine in Klarity (Cyclosporine ophthalmic emulsion PF)	(0.1%)	5.5mL	71384-0514-05
Cyclosporine and Loteprednol in Klarity (Cyclosporine/loteprednol etabonate ophthalmic suspension PF)	(0.1%/0.2%)	5mL	71384-0321-05
Dex-Moxi® (Dexamethasone sodium phosphate and moxifloxacin hydrochloride)	(1mg/0.5mg/mL)	Single-use vial	71384-0512-01
Dex-Moxi-Ketorolac (Dexamethasone sodium phosphate, moxifloxacin, and ketorolac tromethamine)	(1mg/0.5mg/ 0.4mg/mL)	Single-use vial	71384-0513-01
Dorzolamide PF	(2%)	10mL	71384-0516-10
Epi-Lido (Epinephrine/Lidocaine hydrochloride PF/SF in BSS)	(0.75/0.025)%	Single-use vial	71384-0640-01
Klarity Drops® (Glycerin 1% ophthalmic solution PF)	N/A	10mL	71384-0500-10
Loteprednol in Klarity (Loteprednol etabonate 0.5% ophthalmic suspension PF)	(0.5%)	5mL	71384-0210-05
MKO Melt® (Midazolam, ketamine HCl, and ondansetron)	(3/25/2)mg	1 Melt	71384-0630-21
Moxifloxacin	1mg/0.8mL	Single-use vial	71384-0509-08
Moxifloxacin	5mg/0.8mL	Single-use vial	71384-0511-08
Moxi-Brom® (Moxifloxacin and bromfenac)	(0.5/0.075)%	5mL	71384-0311-05
Mydriatric-2 single-use (Tropicamide and phenylephrine hydrochloride)	(1/2.5)%	Single-use	71384-0666-05
Mydriatric-2 multi-dose (Tropicamide and phenylephrine hydrochloride)	(1/2.5)%	10mL	71384-0666-10
Mydriatric-3 single-use (Tropicamide, cyclopentolate, and phenylephrine hydrochloride PF)	(1/1/2.5)%	Single-use	71384-0631-01

PF = Preservative-Free
SF= Sulfite Free

*For professional use only. ImprimisRx specializes in customizing medications to meet unique patient and practitioner needs. No compounded medication is reviewed by the FDA for safety and efficacy. ImprimisRx does not compound essential copies of commercially available products. These ingredients carry risks. Visit imprimisrx.com/productinformation for full product details. References available upon request.

List of Formulation NDCs (Sorted Alphabetically)

Formulations*	Concentration	Unit Size	NDC
Mydriatic-4 single-use (Tropicamide, proparacaine hydrochloride, phenylephrine hydrochloride, and ketorolac tromethamine)	(1/0.5/2.5/0.5)%	Single-use	71384-0632-05
Mydriatic-4 multi-dose (Tropicamide, proparacaine hydrochloride, phenylephrine hydrochloride, and ketorolac tromethamine)	(1/0.5/2.5/0.5)%	5mL	71384-0632-05
Phenyl-Lido (Phenylephrine hydrochloride/lidocaine hydrochloride PF/SF)	(1.5/1)%	Single-use vial	71384-0650-01
Pred Acetate suspension (Prednisolone acetate PF)	(1%)	5mL	71384-0501-05
Pred-Brom® suspension (Prednisolone acetate and bromfenac)	(1/0.075)%	5mL	71384-0505-05
Pred-Gati-Brom® solution (Prednisolone sodium phosphate, gatifloxacin, and bromfenac)	(1/0.5/0.075)%	5mL	71384-0550-05
Pred-Gati-Brom solution (Prednisolone sodium phosphate, gatifloxacin, and bromfenac)	(1/0.5/0.075)%	8mL	71384-0550-08
Pred-Gati-Brom® suspension (Prednisolone acetate, gatifloxacin, and bromfenac)	(1/0.5/0.075)%	5mL	71384-0504-05
Pred-Gati-Brom suspension (Prednisolone acetate, gatifloxacin, and bromfenac)	(1/0.5/0.075)%	8mL	71384-0504-08
Pred-Moxi® suspension (Prednisolone acetate and moxifloxacin)	(1/0.5)%	5mL	71384-0300-05
Pred-Moxi-Brom® suspension (Prednisolone acetate, moxifloxacin, and bromfenac)	(1/0.5/0.075)%	5mL	71384-0310-05
Pred-Moxi-Brom suspension (Prednisolone acetate, moxifloxacin, and bromfenac)	(1/0.5/0.075)%	8mL	71384-0310-08
Pred-Moxi-Brom® solution (Prednisolone phosphate, moxifloxacin, and bromfenac)	(1/0.5/0.075)%	8mL	71384-0340-08
Pred-Moxi-Nepaf® suspension (Prednisolone acetate, moxifloxacin, and nepafenac)	(1/0.5/0.1)%	5mL	71384-0320-05
Pred-Moxi-Nepaf suspension (Prednisolone acetate, moxifloxacin, and nepafenac)	(1/0.5/0.1)%	8mL	71384-0320-08
Pred-Nepaf® (Prednisolone acetate, nepafenac)	(1/0.1%)	5mL	71384-0502-05
Tim-Bim (Timolol and bimatoprost) PF	(0.5/0.1)%	5mL	71384-0542-05
Tim-Brim-Dor® (Timolol, brimonidine, and dorzolamide) PF	(0.5/0.15/2)%	10mL	71384-0520-05
Tim-Brim-Dor-Bim (Timolol, brimonidine, dorzolamide, and bimatoprost) PF	(0.5/0.15/2/0.01)%	5mL	71384-0544-05
Tim-Dor-Bim (Timolol, dorzolamide, and bimatoprost) PF	(0.5/0.15/0.1)%	5mL	71384-0543-05
Tobra-Vanc (Tobramycin and vancomycin)	(1.5/5)%	7mL	71384-0721-07
Tri-Moxi+® (Triamcinolone acetonide and moxifloxacin)	(9mg/0.6mg/0.6mL)	Single-use vial	71384-0746-06


PF = Preservative-Free

SF = Sulfite Free

*For professional use only. ImprimisRx specializes in customizing medications to meet unique patient and practitioner needs. No compounded medication is reviewed by the FDA for safety and efficacy. ImprimisRx does not compound essential copies of commercially available products. These ingredients carry risks. Visit imprimisrx.com/productinformation for full product details. References available upon request.

ORDERING ONLINE

From our
FDA-Registered
503B Outsourcing Facility




3 EASY STEPS

Visit imprimisrx.com/s/order to get started.


REGISTER

Click on the “Register” button to get started. Fill out your information and send it to us.




GET APPROVED

Within 24 hours, you should get an email from our customer support team confirming your prescriber account.



ORDER

Log in and start ordering. Select your favorites, create recurring orders, and get online discounts.



CUSTOMER SUPPORT

Don't hesitate to contact our customer support team. We're here for you!



Call
(844) 446-6979



Chat online
with an agent

PRESCRIBING ONLINE

Ship to Your Office or Patients




3 EASY STEPS

Visit maxrx.com to get started.


REGISTER

Click on the “Don’t have an account” button to get started.




GET APPROVED

Within 24 hours, you should get an email from our customer support team confirming your account.



ORDER

Log in and start ordering. Create patient profiles and track order status.



WHY USE MAXRX PRESCRIBER PORTAL™?

Simple order submission



Ability to track your prescriptions



24-hour processing



1. CREATE A PATIENT PROFILE OR SELECT AN EXISTING ONE

Start your order by selecting one of your patients or create another patient profile.

Account Name	Phone	Shipping Street	Shipping City	Shipping State/Province
1 Barb Akew	(111) 222-3456	642 Mustard Ln.	Oceanside	
2 Terrance Smith	(222) 333-4567	723 Olde Bryd Pt.	Vista	CA
3 Jane Lafferty	(333) 444-5678	8344 Fawn St.	Oceanside	CA
4 Cameron Lydell	(444) 555-6789	946 Takeanap Blvd.	San Diego	CA
5 Arthur Dennington	(555) 666-7890	5065 En Fuego Dr.	Escondido	CA
6 John Mullins	(666) 777-7890	67907 Percy Ave.	Escondido	CA

2. CHOOSE THE MEDICATIONS YOU WANT TO PRESCRIBE

Select the quantity, number of refills and prescribing instructions.

* Medication

Search: cycl

- Show All Results for "cycl"
- Klarity-C Drops® (Cyclosporine 0.1% ophthalmic emulsion PF)* - 0.10%
71384-0514-05
- Mydriatric 3 (Tropi-Cyclo-Phenyl)* - 1mL - 1/1/2.5%
71384-0631-01

ImprimisRx dispenses only to individually identified patients with valid prescriptions. No compounded medication is reviewed by the FDA for safety or efficacy. ImprimisRx does not compound essentially copies of commercially available products. References available upon request.

3. REVIEW AND SUBMIT YOUR PRESCRIPTION

Review your prescription to make sure everything is right and submit your order.




Subtotal Amount \$5,170.00	Order Status Order Submitted
Patient Cindi White	503A Bill To
	503A Ship To

Order Items (2)			
OrderItemAutoNumber	Product	SIG Instructions	Medical Necessity
000101	Pred-Acetate 1% PF	Apply to left and right eye twice ...	Allergy to Preservatives
000102	Klarity-C Drops® Cyclosporine ...	Use twice daily	Cannot use commercial

[View All](#)

4. TRACK THE STATUS OF YOUR PRESCRIPTION

Track the status of the prescription every step of the way.

All Orders ▾ 						
Refresh this list to view the latest data						<input type="text" value="Search this list..."/>  
	Order Number ↑ ▾	Prescriber Name ▾	Patient Name ▾	Order Date ▾	Order Status ▾	
1	O-0000000103			7/22/2020	Order Submitted	▾
2	O-0000000104			7/22/2020	Open	▾
3	O-0000000105			7/22/2020	Open	▾
4	O-0000000106			7/22/2020	Open	▾
5	O-0000000107			7/22/2020	Order Submitted	▾
6	O-0000000133			7/27/2020	Order Submitted	▾

You can trust **ImprimisRx[®]** to care for your patients



William F. Wiley, MD is a paid consultant of ImprimisRx

Compounded medications have an important role to play in contemporary ophthalmic surgery. They offer more choices for physicians and good value for patients. Registration as a 503B facility is an indication of high-quality manufacturing standards and federal oversight of the facility.

EMR PRESCRIBING

e-Scribe using your existing EMR system and send your prescription directly to our pharmacy.



1. LOCATE OUR PHARMACY

ImprimisRx NJ, LLC
1705 Route 46, Suite 4
Ledgewood, NJ 07852
866-792-7328

Quick Tip: Set the state to New Jersey to find “ImprimisRx NJ” in your list of pharmacies.

2. COMPLETE THE PATIENT INFORMATION

Make sure your patient profile has name, phone number, address and date of birth in your EMR so demographics can transmit to our system.

3. SEARCH AND SELECT MEDICATION

Search for ONE of the active ingredients in the desired ImprimisRx compound. Select the correct compound from your EMR’s AUTO-GENERATED list.

Example: Search “cyclosporine” for Cyclosporine in Klarity/Klarity-C

4. ENTER THE CORRECT QUANTITY

Formulations may have more than one size; be sure to select the correct fill size. Example: Klarity-C is sold in 3 bottles (5.5mL bottles), totaling 16.5mL. Please enter 16.5mL as the default for Klarity-C.

5. PROVIDE INSTRUCTIONS FOR USE

Please indicate dosing instructions—how the formulation will be administered, volume to be used, and how often the drops are to be applied.

<p>STEROID-ANTIBIOTIC-ANTI-INFLAMMATORY AGENT 5ML or 8ml (ie: Prednisolone-Gatifloxacin-Bromfenac - 5ml)</p> <p>Instill ___X___ drop(s) into affected eye ___X___ times a day for ___X___ days.</p>

6. INCLUDE YOUR NOTES TO THE PHARMACY

- Dispense ImprimisRx compound (ie: “Prednisolone Acetate-Gatifloxacin-Bromfenac” or “Pred-Gati-Brom”- 5mL)
**Needed only if formulation cannot be selected in EMR*
- Medical necessity (ie: patient cannot tolerate commercial formulation, patient needs double strength, etc.)
*Please note: financial/economic reason is not valid. *Required*
- Patient allergies
**Required*
- Ship to patient or doctor, bill to patient or doctor
**Required*
- Medication start date, surgery date
**If applicable*

NEED HELP?



Call
(844) 446-6979



Chat online
with an agent

Compound Formulation Active Pharmaceutical Ingredients and The Associated Risks and Contraindications

The risks and contraindications outlined below are those known to be associated with the active ingredients, as reflected in the prescribing information for FDA-approved products containing the same active ingredients, and the technique by which the products are to be administered.

Atropine Sulfate		Formulation
Possible Adverse Effects	Blurred vision, Visual disturbance, Light sensitivity, Eye stinging, Irritation, Eye pain, Palpitation/Arrhythmias, Flushing, Restlessness, Feeling irritable, Dry mouth, Dry skin	Atropine Sulfate PF
Potential Contraindications / Precautions	Do not give to children less than 3 months of age. Patients 65 years of age or older, use this preparation with care. Pregnant or breast-feeding, talk to your physician prior to use.	
Azithromycin		Formulation
Possible Adverse Effects	Eye irritation, Blurred vision, Contact dermatitis, Corneal erosion, Decreased visual acuity, Dysgeusia, Eye pain, Facial edema, Eye reactions (burning sensation, eye discharge, stinging), Nasal congestion, Sinusitis, Swelling of the eye, Skin rash	Klarity-A® Drops
Potential Contraindications / Precautions	Hypersensitivity to Azithromycin or any component of the formulation. Pregnant or breast-feeding, talk to your physician prior to use.	
Bimatoprost		Formulations
Possible Adverse Effects	Iris pigmentation changes, Eyelid skin darkening, Eyelash changes (increased length, thickness, pigmentation and number of lashes), Iritis/uveitis, Macular edema, including cystoid macular edema, Foreign body sensation, Punctate keratitis, Stinging, Conjunctival hyperemia, Blurred vision, Itching, Burning, Increased pigmentation of the iris, Excessive tearing, Eyelid discomfort/pain, Dry eye, Eye pain, Eyelid margin crusting, Erythema of the eye lid, Photophobia	Bim Tim-Bim Tim-Dor-Bim Tim-Brim-Dor-Bim
Potential Contraindications / Precautions	Patients less than 16 years, Lactation, Pregnancy, Child-bearing aged females	
Brimonidine Tartrate		Formulations
Possible Adverse Effects	Oral dryness, Ocular hyperemia, Burning and stinging on instillation, Headache, Blurred vision, Foreign body sensation, Fatigue/drowsiness, Conjunctival follicles, Ocular allergic reactions, Ocular pruritis, Corneal staining/erosion, Photophobia, Eyelid erythema, Ocular ache/pain, Upper respiratory symptoms, Dizziness, Blepharitis, Ocular irritation, Lid crusting, Conjunctival hemorrhaging, Abnormal taste, Insomnia, Depression, Hypertension, Anxiety, Palpitations/arrhythmias, Nasal dryness and syncope	Brim-Dor® Tim-Brim-Dor® Tim-Brim-Dor-Bim
Potential Contraindications / Precautions	Infants, Lactation, Pregnancy, Concomitant MAO inhibitor therapy	
Bromfenac		Formulations
Possible Adverse Effects	Abnormal sensation in eyes, Anaphylaxis, Anterior chamber eye Hemorrhage, Burning sensation, Conjunctival hyperemia, Corneal erosion, Corneal perforation, Corneal thinning, Corneal ulcer, Epithelial keratopathy, Eye irritation, Eye pain, Eye pruritus, Eye redness, Headache, Hypersensitivity reaction, Iritis, Keratitis, Prolonged bleeding, Stinging sensation	Moxi-Brom Pred-Brom® Pred-Gati-Brom® Pred-Moxi-Brom® Suspension Pred-Moxi-Brom Solution
Potential Contraindications / Precautions	Lactation, Pediatric patients, Pregnancy, Corneal disease, Thrombocytopenia, Rheumatoid arthritis	

Cyclopentolate		Formulation
Possible Adverse Effects	Blurred vision, Burning sensation, Coma, Death, Disorientation, Dry secretions, Eye irritation, Hallucination, Hyperactive behavior, Hyperpyrexia, Hypertension, Hyperhidrosis, Increased intraocular pressure, Photophobia, Psychotic reaction, Restlessness, Seizure, Superficial punctate keratitis, Tachycardia, Urinary retention, Vasodilation, Xerostomia	Mydratic 3
Potential Contraindications / Precautions	Infants	
Cyclosporine/Loteprednol Etabonate Ophthalmic Suspension PF		Formulation
Possible Adverse Effects	Blurred vision, Conjunctival hyperemia, Epiphora, Eye discharge, Eye pain, Headache, Foreign body sensation, Hypersensitivity, Pruritus, Stinging of the eyes, Swelling of the eyes, Burning sensation, Dry eyes, Runny nose, Urticaria, Visual disturbance, Sore throat, Light sensitivity, Cataracts or glaucoma with chronic use	Klarity-CL® Drops
Potential Contraindications / Precautions	Infants and children under the age of 16. Pregnant or breast-feeding, talk to your physician prior to use.	
Cyclosporine Ophthalmic Emulsion		Formulation
Possible Adverse Effects	Blurred vision, Conjunctival hyperemia, Epiphora, Eye discharge, Eye pain, Foreign body sensation, Hypersensitivity, Pruritus, Stinging of the eyes, Swelling of the eyes, Burning sensation, Urticaria visual disturbance	Klarity-C Drops®
Potential Contraindications / Precautions	Infants and patients less than 16 years of age, Lactation	
Dexamethasone Sodium Phosphate		Formulations
Possible Adverse Effects	Conjunctival irritation, Burning and stinging sensation, Fever, Eye irritation and infection, Airway obstruction, Anaphylaxis, Angioedema, Circulatory shock, Blindness, Post-injection flare, Wound healing suppression, Corneal perforation, Headache, Subcapsular posterior cataract, Ocular hypertension, Optic nerve damage, Visual field defect	Dex-Moxi® Dex-Moxi-Ketor®
Potential Contraindications / Precautions	Infants and patients less than 12 years of age, Lactation, Pregnancy	
Dextran		Formulation
Possible Adverse Effects	Mild eye burning or irritation, Itching or redness of your eye, Watery eyes, Blurred vision	Klarity Drops®
Potential Contraindications / Precautions	Self-medication: When used for self-medication, stop use and contact healthcare provider if changes in vision, eye pain, continued redness or irritation occur, or if the condition worsens or persists for more than 72 hours. To avoid contamination, do not touch tip of container to any surface. Replace cap after using. Do not use if solution is cloudy or changes color.	
Dorzolamide		Formulations
Possible Adverse Effects	Ocular burning, stinging, or discomfort, immediately following ocular administration, Bitter taste, Superficial punctate keratitis, Ocular allergic reaction, Conjunctivitis and lid reactions, Blurred vision, Eye redness, Tearing, Dryness, Photophobia, Headache, Nausea, Asthenia/fatigue, Skin rashes, Urolithiasis, Iridocyclitis	Brim-Dor® Dor Tim-Dor-Bim Tim-Brim-Dor® Tim-Brim-Dor-Bim
Potential Contraindications / Precautions	Neonates and Patients Less than 12 Years, Lactation, Pregnancy	
Epinephrine HCl		Formulation
Possible Adverse Effects	Increased intraocular pressure, Hypertension	Epi-Lido
Potential Contraindications / Precautions	Lactation, Pregnancy	

Gatifloxacin		Formulation
Possible Adverse Effects	Conjunctival irritation, Decreased visual acuity, Dysgeusia, Eye discharge, Eye irritation, Eye pain, Eye redness, Eyelid edema, Headache, Increased lacrimation, Keratitis, Papillary conjunctivitis	Pred-Gati-Brom®
Potential Contraindications / Precautions	Infants, Lactation, Pregnancy	
Glycerin		Formulation
Possible Adverse Effects	Blurred vision, Crusting of eyelid, Stinging of eyes (mild)	Klarity Drops®
Potential Contraindications / Precautions	Self-medication: When used for self-medication, stop use and contact healthcare provider if changes in vision, eye pain, continued redness or irritation occur, or if the condition worsens or persists for more than 72 hours. To avoid contamination, do not touch tip of container to any surface. Replace cap after using. Do not use if solution is cloudy or changes color.	
Ketamine HCl		Formulation
Possible Adverse Effects	Airway obstruction, Anaphylaxis, Anxiety, Apnea, Bradycardia, Cardiac arrhythmia, Confusion, Disorientation, Dysphoria, Hallucination, Hypotension, Insomnia, Nausea, Nystagmus, Psychotic reaction, Respiratory depression, Skin rash, Tonic clonic, Epilepsy, Vomiting	MKO Melt®
Potential Contraindications / Precautions	Lactation, Pregnancy	
Ketorolac Tromethamine		Formulations
Possible Adverse Effects	Burning sensation of eyes, Stinging of eyes, Conjunctival hyperemia, Corneal edema, Corneal erosion, Corneal infiltrates, Corneal perforation, Corneal thinning, Eye pain and irritation, Headache, Hypercreativity reaction, Iritis, Ocular edema, Ophthalmic inflammation, Superficial eye infection, Superficial keratitis, Tissue damage	Dex-Moxi-Ketor® Mydriatic 4
Potential Contraindications / Precautions	Patients less than 3 years of age, Lactation, Pregnancy, Child bearing aged females	
Lidocaine HCl		Formulations
Possible Adverse Effects	Conjunctival Hyperemia, Corneal Epithelial Changes, Headache	Epi-Lido Phenyl-Lido
Potential Contraindications / Precautions	Infants, Lactation, Pregnancy	
Loteprednol Etabonate		Formulation
Possible Adverse Effects	Headache, Eye burning, Irritation, itching, Dry eyes, Blurred eyesight, Feeling that something is in the eye, Runny nose, Sore throat, Light sensitivity, Cataracts or glaucoma with chronic use	Klarity-L® Drops
Potential Contraindications / Precautions	Infants and children under the age of 18. Pregnant or breast-feeding, talk to your physician prior to use.	
Midazolam		Formulation
Possible Adverse Effects	Hypoxemia, Nausea and vomiting, Agitation, Airway obstruction, Bradycardia, Hypoxia, Laryngospasm, Respiratory depression, Rhonchi, Sedation, Skin rash, Cardiac arrest, Respiratory arrest	MKO Melt®
Potential Contraindications / Precautions	Infants less than 6 months of age, Lactation, Pregnancy	

Moxifloxacin HCl		Formulations
Possible Adverse Effects	Conjunctival irritation, Fever, Eye Irritation, Airway obstruction, Anaphylaxis, Angioedema, Circulatory, Shock, Dyspnea, Facial edema, Fungal, Superinfection, Laryngeal edema, Loss of consciousness, Pharyngeal edema, Pruritus, Urticaria, Pigment dispersion syndrome	Dex-Moxi® Dex-Moxi-Ketor® Moxifloxacin PF Moxi-Brom Pred-Moxi® Pred-Moxi-Brom® Suspension Pred-Moxi-Brom Solution Pred-Moxi-Nepaf® Suspension Tri-Moxi+®
Potential Contraindications / Precautions	Infants, Lactation, Pregnancy	
Nepafenac		Formulations
Possible Adverse Effects	Abnormal sensation in eyes, Increased intraocular pressure, Anterior chamber eye hemorrhage, Burning sensation, Conjunctival hyperemia, Corneal erosion, Corneal perforation, Corneal thinning, Corneal ulcer, Epithelial keratopathy, Eye irritation, Eye pain, Eye pruritus, Eye redness, Headache, Hypersensitivity reaction, Iritis, Keratitis, Prolonged bleeding, Stinging sensation	Pred-Moxi-Nepaf® Suspension Pred-Nepaf® Suspension
Potential Contraindications / Precautions	Lactation, Pediatric patients, Pregnancy, Corneal disease, Thrombocytopenia (bleeding disorders)	
Ondansetron		Formulation
Possible Adverse Effects	Headache and malaise, Sedation, Wound healing impairment, Anxiety, Bradycardia, Constipation, Diarrhea, Dizziness, Fever, Gynecologic disease, Hypotension, Hypoxia, Pruritus, Shivering, Urinary retention, Skin rash, Anaphylaxis, Chest pain, Dyspnea, Laryngeal edema	MKO Melt®
Potential Contraindications / Precautions	Infants less than 6 months of age, Lactation, Pregnancy	
Phenylephrine HCl		Formulations
Possible Adverse Effects	Acute keratitis, Acute myocardial infarction, Allergic conjunctivitis, Blurred vision, Dermatitis, Diaphoresis, Dizziness, Eye irritation, Headache, Hypertension, Lacrimation, Pallor, Photophobia, Pigment granules liberation from the iris, Subarachnoid hemorrhage, Syncope, Tachycardia, Tremor	Mydriatic 2 Mydriatic 3 Mydriatic 4 Phenyl-Lido
Potential Contraindications / Precautions	Neonates and patients less than 12 years of age, Lactation, Pregnancy	
Prednisolone Acetate		Formulations
Possible Adverse Effects	Accommodation disturbance, Anterior uveitis, Blepharitis, Conjunctival hyperemia, Conjunctivitis, Corneal perforation or Ulcer, Decreased visual acuity, Eye infection, Glaucoma, Increased intraocular pressure, Keratitis, Mydriasis, Optic nerve damage, Visual field defect, Wound healing impairment	Prednisolone Acetate Pred-Brom® Pred-Gati-Brom® Pred-Moxi® Pred-Moxi-Brom® Suspension Pred-Moxi-Nepaf® Suspension Pred-Nepaf® Suspension
Potential Contraindications / Precautions	Infants, Lactation, Pregnancy, Cataract, Eye infection, Increased intraocular pressure	
Prednisolone Phosphate		Formulations
Possible Adverse Effects	Burning or stinging sensation, Accommodation disturbance, Conjunctival hyperemia, Decreased visual Acuity, Eye infection, Glaucoma, Increased intraocular pressure, Keratitis, Mydriasis, Optic nerve damage, Subcapsular posterior cataract, Visual field defect, Wound healing impairment, Corneal ulcer, Viral infection	Pred-Brom® Pred-Gati-Brom® Pred-Moxi-Brom Solution
Potential Contraindications / Precautions	Infants, Lactation, Pregnancy, Cataract, Eye infection, Increased intraocular pressure	

Proparacaine HCl		Formulation
Possible Adverse Effects	Blurred vision, Stinging of the eyes, Hypersensitivity reaction	Mydriatic 4
Potential Contraindications / Precautions	Infants, Lactation, Pregnancy	
Timolol Maleate		Formulations
Possible Adverse Effects	Ocular burning/stinging on instillation, Dry eyes, Itching, Foreign body sensation, Discomfort in the eye, Eyelid erythema, Conjunctival injection, Headache, Eye pain, Epiphora, Photophobia, Blurred or abnormal vision, Corneal fluorescein staining, Keratitis, Blepharitis, Cataract, Cardiac arrhythmia, Palpitation, Bradycardia, Syncope, Heart block, Cerebral vascular accident, Cerebral ischemia, Cardiac failure and Cardiac arrest	Tim-Bim Tim-Dor-Bim Tim-Brim-Dor® Tim-Brim-Dor-Bim
Potential Contraindications / Precautions	Patients Less than 12 Years, Lactation, Pregnancy, Bronchial Asthma or History of Sinus Bradycardia, Second or third-degree atrioventricular block, Overt heart failure, Cardiogenic shock.	
Tobramycin		Formulation
Possible Adverse Effects	Ocular burning/stinging on instillation, Lid itching and swelling, Conjunctival erythema, Anaphylactic reaction, Stevens-Johnson syndrome, Erythema multiforme. The following adverse reactions reported with systemic aminoglycosides: Neurotoxicity, Ototoxicity, Nephrotoxicity, Aminoglycosides may aggravate muscle weakness in patients with known or suspected neuromuscular disorders, such as myasthenia gravis or Parkinson's disease.	Tobra-Vanc
Potential Contraindications / Precautions	Patients less than 12 years of age, Nursing mothers, Pregnancy	
Triamcinolone Acetonide		Formulation
Possible Adverse Effects	Anaphylactoid shock, Blindness, Cataract, Corneal perforation, Cortical cataract, Conjunctival hemorrhage, Eye pain, Infection, Irritation pruritus, Abnormal sensation in eyes, Abnormal sensory symptoms, Edema, Bleeding at injection site, Glaucoma, Hypertension, Hypopyon, Increased intraocular pressure, Increased Lacrimation, Vitreous Detachment, Vitreous Opacity, Wound Healing Impairment	Tri-Moxi+®
Potential Contraindications / Precautions	Infants, Lactation, Pregnancy, Cataract, Eye infection, Increased intraocular pressure	
Tropicamide		Formulations
Possible Adverse Effects	Blurred vision, Cardiorespiratory arrest, Headache, Hypersensitivity reaction, Increased intraocular pressure, Muscle rigidity, Nausea, Pallor, Photophobia, Psychotic reaction, Superficial punctate, Keratitis, Tachycardia, Transient burning or stinging in the Eyes, Vomiting, Xerostomia	Mydriatic 2 Mydriatic 3 Mydriatic 4
Potential Contraindications / Precautions	Patients less than 12 years of age, Lactation, Pregnancy	
Vancomycin		Formulation
Possible Adverse Effects	Ocular burning/stinging on instillation, Lid itching and swelling, Conjunctival erythema, Anaphylactic reaction, Stevens-Johnson syndrome. The following adverse reactions reported with systemic vancomycin: Nephrotoxicity, Onset of pseudomembranous colitis symptoms, Ototoxicity, Neutropenia. Hemorrhagic occlusive retinal vasculitis (HORV) may be associated with the use of intraocular vancomycin (including intravitreal, intracameral bolus, and irrigating solution containing vancomycin). Although the precise etiology of HORV remains unproven, each surgeon should weigh the potential risk of vancomycin against the risk of endophthalmitis.	Tobra-Vanc
Potential Contraindications / Precautions	Patients less than 12 years of age, Nursing mothers, Pregnancy	



ImprimisRx® gives back.

ImprimisRx contributes to missions and indigent programs that provide cataract surgery to over 14,000 patients in need every year. For every Dropless and LessDrops prescription shipped, we donate a percentage to sight-saving missions and indigent patient support programs.

Ask about our indigent patient support program.

Contact us via chat at [imprimisrx.com](https://www.imprimisrx.com)

Photos courtesy of Dr. Mary Cathleen Schanzer and Dr. Lee Coleman.

A note from the Chief Commercial Officer at Harrow



John Saharek
Chief Commercial Officer

At Harrow, we are committed to earning your trust every day by providing quality products and exceptional service! We back this commitment up with a 100% money back guarantee on every order placed with our 503B FDA-Registered Outsourcing Facility and 503A Patient Specific Pharmacy.

Please let us know how we can continue to improve via the chat feature on imprimisrx.com.

Thank you for allowing us to serve you and your patients.



ImprimisRx®
1000 Aviara Pkwy, Suite 220
Carlsbad, CA 92011