

# Pathogens in *Crassostrea ariakensis* and other Asian oyster species: implications for non-native oyster introduction to Chesapeake Bay

J. A. Moss<sup>1</sup>, E. M. Burreson<sup>1</sup>, J. F. Cordes<sup>1</sup>, C. F. Dungan<sup>2</sup>, G. D. Brown<sup>1</sup>, A. Wang<sup>3</sup>,  
X. Wu<sup>4</sup>, K. S. Reece<sup>1,\*</sup>

<sup>1</sup>Virginia Institute of Marine Science, The College of William and Mary, Gloucester Point, Virginia 23062, USA

<sup>2</sup>Maryland Department of Natural Resources, Cooperative Oxford Laboratory, Oxford, Maryland 21654, USA

<sup>3</sup>School of Marine Science, Hainan University, 58 Renmin Street, Haikou, Hainan 570228, PR China

<sup>4</sup>College of Animal Sciences, Zhejiang University, 38 Zheda Road, Hangzhou, Zhejiang Province 310027, PR China

**ABSTRACT:** With the drastic decline of eastern oyster *Crassostrea virginica* populations in the Chesapeake Bay due to over-fishing, diseases and habitat destruction, there is interest in Maryland and Virginia in utilizing the non-native oyster species *Crassostrea ariakensis* for aquaculture, fishery resource enhancement, and ecological restoration. The International Council for the Exploration of the Sea (ICES) recommends that non-native species be examined for ecological, genetic and disease relationships in the native range prior to a deliberate introduction to a new region. Therefore, a pathogen survey of *C. ariakensis* and other sympatric oyster species was conducted on samples collected in the PR China, Japan and Korea using molecular diagnostics and histopathology. Molecular assays focused on 2 types of pathogens: protistan parasites in the genus *Perkinsus* and herpesviruses, both with known impacts on commercially important molluscan species around the world, including Asia. PCR amplification and DNA sequence data from the internal transcribed spacer region of the rRNA gene complex revealed the presence of 2 *Perkinsus* species not currently found in USA waters: *P. olseni* and an undescribed species. In addition, 3 genetic strains of molluscan herpesviruses were detected in oysters from several potential *C. ariakensis* broodstock acquisition sites in Asia. Viral gametocytic hypertrophy, *Chlamydia*-like organisms, a *Steinhausia*-like microsporidian, *Perkinsus* sp., *Nematopsis* sp., ciliates, and cestodes were also detected by histopathology.

**KEY WORDS:** Asia · *Crassostrea ariakensis* · *Crassostrea hongkongensis* · *Crassostrea gigas* · *Perkinsus olseni* · Molluscan herpesviruses · *Steinhausia* sp. · Parasites · ISH · PCR

Resale or republication not permitted without written consent of the publisher

## INTRODUCTION

The eastern oyster *Crassostrea virginica* is important both economically and ecologically in Chesapeake Bay, USA. Oyster populations have been in a severe state of decline in recent decades, due to combined effects of over-harvesting, habitat loss and disease pressures from marine pathogens (Mann et al. 1991). In the Commonwealth of Virginia, oyster landings exceeding 6 million bushels (1 bushel = 35.24 l) in the 1930s have declined to less than 20 000 bushels since

the 1990s (NRC 2003, Allen 2005). In 1995, a Virginia General Assembly resolution requested that the Virginia Institute of Marine Science (VIMS) determine the appropriate legal process for, and examine the feasibility of, introducing a non-native oyster species to enhance ecological benefits and revitalize the oyster industry in the Chesapeake Bay region. Initial results of research using the Pacific oyster *C. gigas*, which has been successfully introduced at several locations around the world, suggested that this oyster species would not perform well in the Chesapeake Bay (Calvo

\*Corresponding author. Email: kreece@vims.edu

et al. 1999). The search for another non-native oyster species that might be more suitable for the regional conditions was therefore initiated.

Considerable interest has recently focused on the Suminoe oyster *Crassostrea ariakensis*, which resembles the native oyster *C. virginica* in taste (Grabowski et al. 2003), and is tolerant of temperate to subtropical water temperatures and variable salinities. Field trials conducted in Virginia waters have documented lower mortality and faster growth in the Suminoe oyster, as compared with the native oyster. Disease surveys of triploid Suminoe oysters deployed in these parallel trials also suggest that in comparison to the native eastern oyster, *C. ariakensis* is relatively resistant to *Haplosporidium nelsoni* and *Perkinsus marinus* infections (Calvo et al. 2001, Orner 2005), the 2 major parasites that have decimated *C. virginica* oyster populations in Chesapeake Bay since the 1950s (Sindermann 1990).

Based on significant impacts that non-native introductions of various aquatic species have had around the world, the International Council for the Exploration of the Seas (ICES) developed a 'Code of Practice on Introductions and Transfers of Marine Organisms' (ICES 2005). The code describes a series of protocols to be followed prior to introductions of exotic animals. A thorough review of the ecological, genetic and disease impacts on native bivalves of the proposed introduced species in its natural range and in source locations is recommended. These and similar recommendations from other organizations were the impetus behind a survey that we conducted on the parasites of *Crassostrea ariakensis* and other sympatric oyster species in the native range of *C. ariakensis* in China, Japan and Korea.

Previous research has documented the harmful impact of both *Perkinsus* spp. and herpes-like viruses on molluscan species in Asia (Choi & Park 1997, Park & Choi 2001, Chang et al. 2005). This fact, along with the availability of molecular assays for these organisms (Renault et al. 2000a, Casas et al. 2002), prompted us to specifically screen for these pathogens in Asian oyster populations. Since the discovery of *Perkinsus marinus* in *Crassostrea virginica* along the Gulf of Mexico and Atlantic coasts of the USA in the late 1940s and early 1950s (Mackin et al. 1950, Ray 1952), *Perkinsus* spp. have been found worldwide, and many are reported to cause disease in commercially important mollusc species. *P. marinus* is notorious for its devastating effects on Atlantic and Gulf of Mexico, USA oyster populations (Andrews & Hewatt 1957, Burreson & Ragone Calvo 1996). In the Chesapeake Bay region, *P. chesapeakei* (= *P. andrewsi*) (Burreson et al. 2005) has been associated with disease outbreaks in *Mya arenaria* and *Tagelus plebeius* (Dungan et al. 2002). Clam and oys-

ter mortalities have also occurred in conjunction with *Perkinsus* spp. infections on the Atlantic and Mediterranean coasts of Europe (Da Ros & Cazonier 1985, Azevedo 1989, Figueras et al. 1992, Santmartí et al. 1995, Montes et al. 2001, Villalba et al. 2005), in Australia (Lester & Davis 1981, Goggin & Lester 1995) and in Korea (Choi & Park 1997, Park & Choi 2001).

In addition to *Perkinsus* spp. parasites, viruses and particularly molluscan herpesviruses have been found to cause massive mortality, mainly in hatchery-reared larvae and juvenile animals, leading to severe economic losses (Hine et al. 1992, Le Deuff et al. 1994, Arzul et al. 2001, Friedman et al. 2005). Numerous cases of herpes-like viruses affecting commercial marine molluscs have been reported around the world, the earliest in 1972 in *Crassostrea virginica* in Maine, USA (Farley et al. 1972). Herpes-like viruses have been reported in the Pacific oyster *C. gigas* (Hine et al. 1992, Nicolas et al. 1992, Renault et al. 1994, Friedman et al. 2005), the European flat oyster *Ostrea edulis* (Comps & Cochennec 1993, Renault et al. 2000a), the Australian flat oyster *O. angasi* (Hine & Thorne 1997), larvae of the Chilean oyster *Tiostrea chilensis* in New Zealand (Hine et al. 1998), the European carpet shell clam *Ruditapes decussatus* (Renault et al. 2001), the Manila clam *R. philippinarum*, (Renault 1998) and the scallop *Pecten maximus* in France (Arzul et al. 2001). A similar herpes-like virus may be responsible for mortality events in abalone *Haliotis diversicolor supertexta* in Taiwan (Chang et al. 2005).

Unlike the larvae and some juvenile oysters, adult *Crassostrea gigas* appear capable of surviving asymptotically with *Ostreid herpes virus-1* (OsHV-1) infections (Arzul et al. 2002), with no gross physiological signs detectable prior to death in the infected individuals. Histopathological signs indicative of infection include enlarged and abnormally shaped cell nuclei, and abnormal chromatin patterns throughout the connective tissue (Renault et al. 1994). The experience in French hatcheries has shown that this pathogen is likely to be vertically transmitted from broodstock to progeny, and can be very difficult to eradicate from facilities (Arzul et al. 2001). Consequently, if introduced along with *C. ariakensis*, molluscan herpesviruses could have a devastating impact on the remaining *C. virginica* populations in Chesapeake Bay, and on the growing aquaculture industry in the USA mid-Atlantic region.

In accordance with the ICES (2005) recommendation that non-native species be examined for diseases in their native range prior to a deliberate introduction into a new region, we conducted a pathogen survey of *Crassostrea ariakensis* and other sympatric oyster species in samples collected in China, Japan and Korea. *Perkinsus* spp. and other metazoan parasites were

observed by histology. Molecular diagnostics developed to target *Perkinsus* spp. and OsHV-1 identified 2 *Perkinsus* species not currently found in USA waters (*P. olseni* and an undescribed *Perkinsus* species), as well as 3 genetic variants of molluscan herpesviruses that are highly similar in DNA sequence to OsHV-1 in the genomic region analyzed. It should be noted that recent studies using these same molecular diagnostic assays have detected the 2 endemic *Perkinsus* species, *P. marinus* and *P. chesapeaki* in waters of eastern North America (Burreson et al. 2005, Audemard et al. 2006). No evidence of molluscan herpesviruses, however, has been detected in bivalve hosts along the Atlantic and Gulf of Mexico coasts of the USA (Friedman et al. 2005).

## MATERIALS AND METHODS

**Sample collection and preparation.** Samples of *Crassostrea ariakensis* and, inadvertently, several other oyster species (see below), were collected from coastal sites in China, Japan and Korea between 1999 and 2005 (Fig. 1 and Table 1). Additional oyster hemolymph samples were taken from hatchery-reared *C. ariakensis* including F1 northern China *C. ariakensis*

(NCA) spawned from broodstock collected from the Yellow River in China in 1999, F1 southern China *C. ariakensis* (SCA) spawned from broodstock collected from the Dafen River in China in 1999, and west coast USA *C. ariakensis* (WCA) spawned from broodstock imported to VIMS from Washington, USA in 1999. Adductor muscle, mantle, gill tissue and/or hemolymph from each individual were preserved in either DMSO (25 mM EDTA, 20% DMSO and saturated NaCl) or 95% ethanol for DNA extraction and PCR analysis. When samples were additionally preserved and processed for histological analysis (Table 1), a sterile blade was used to excise a transverse tissue section through the visceral mass, and histological samples were fixed in Davidson's solution (Shaw & Battle 1957). Because of the large size of the oysters in the 2002 Chinese samples, 2 tissue sections were preserved for each animal, one that included digestive gland, gill and mantle, and one that included adductor muscle, heart and kidney. Paraffin wax-infiltrated histological tissues were embedded, sectioned at 5 to 6  $\mu\text{m}$  thickness, and sections were stained with Mayer's hematoxylin and eosin for microscopic analyses.

**Nucleic acid extraction.** Genomic DNA was extracted individually for each oyster from excised mantle and gill snips using the DNeasy<sup>®</sup> Tissue Kit (Qiagen) following manufacturer's protocols. Hemolymph samples were centrifuged at  $16\,000 \times g$  for 5 min and precipitated floccules were then subjected to DNA extraction with the DNeasy<sup>®</sup> Tissue Kit. DNA was eluted in 50 to 200  $\mu\text{l}$  of elution buffer.

**Test for amplifiable DNA.** In order to ensure that amplifiable DNA was present in all extracted samples, genomic DNAs were tested using universal small subunit ribosomal RNA (SSU-rRNA) gene primers 16S-A (5'-CCGAATTCGTCGACAACCTGGTTGATCCTGCCAGT-3') and 16S-B (5'-GGATCCAAGCTTGATCCTTCTGCAGTTCCCTAC-3') (modified from Medlin et al. 1988) with an expected amplification product of approximately 1800 bp. Each PCR reaction contained the following: 20 mM Tris-HCl (pH 8.4), 50 mM KCl, 0.75 mM  $\text{MgCl}_2$ , 0.1 mM of each dNTP, 0.5  $\mu\text{M}$  of each primer, 0.0125 U  $\mu\text{l}^{-1}$  *Taq* polymerase, 0.2 mg  $\text{ml}^{-1}$  bovine serum albumin (BSA), and 0.5  $\mu\text{l}$  genomic DNA (10 to 50 ng total). Amplifications were performed with an initial denaturation of 94°C for 4 min, followed by 35 cycles at 94°C for 30 s, 45°C for 30 s, 65°C for 2 min, with a final elongation at 65°C for 2 min. Following amplification, 3  $\mu\text{l}$  of PCR product were analyzed by agarose gel electrophoresis (2%), stained with ethidium bromide and visualized under UV light. Images were recorded with an Alpha Innotech FluorChem<sup>®</sup> imaging system.

**Genus-specific *Perkinsus* spp. PCR assay.** Screening for *Perkinsus* spp. DNA was performed using *Perkin-*




Fig. 1. Map of sampling locations. See Table 1 for site abbreviations

Table 1. Sampling sites, abbreviations used, sample sizes, bivalve species present, analyses performed and general results (+ or –) of assays. Sampling sites in China are ordered from northern to southern locations, and are listed chronologically for multiple collections from a single site. Positive (+) assay results in the molecular assays indicate that either molluscan herpesvirus or *Perkinsus* sp. DNA was detected (see Table 2) and, for the histological analysis, that a potential pathogen was detected (see Table 3). C. = *Crassostrea*, SCA = F1 southern China *C. ariakensis*, NCA = F1 northern China *C. ariakensis*, WCA = west coast USA *C. ariakensis*

Year	Sampling location (sample abbreviation)	Natural (N) Cultivated (C)	Sample size (N)	Oyster species	Assays performed	Assay results (+ or –)
<b>Japan</b>						
1999	Itoki River, Kyushu (IR)	N	50	<i>C. ariakensis</i>	Molecular	+
2003	Ariake Sea (AK)	N	24	<i>C. gigas</i> , <i>C. sikamea</i>	Molecular	–
2004	Amakusa (AM)	N	60	<i>C. gigas</i>	Molecular	–
2004	Midori (MI)	N	68	<i>C. gigas</i>	Molecular	–
2005	Mie Prefecture (JPAB05)	C	246	<i>C. ariakensis</i> , <i>C. sikamea</i> , <i>C. gigas</i>	Molecular	+
<b>South Korea</b>						
2004	Seogwipo (SE)	Unknown	161	<i>Saccostrea</i> sp., <i>C. gigas</i> , unknown sp.	Molecular	–
2004	Kahwa River (KR)	Unknown	35	<i>C. ariakensis</i> , <i>C. gigas</i>	Molecular	+
2004	Sumjin River (SR)	Unknown	20	<i>C. ariakensis</i>	Molecular	+
2004	Kanghwa Island, Inchon (IN)	Unknown	20	<i>C. ariakensis</i>	Molecular	+
<b>PR China (PRC)</b>						
1999	Yellow River, Bohai Sea, Shandong (YR)	N	43	<i>C. ariakensis</i>	Molecular	+
2002	Dajiawa, Shandong (DJ)	N	26	<i>C. ariakensis</i> , <i>C. hongkongensis</i>	Molecular Histological	– –
2002	Longhai, Fujian (LH)	C	9	<i>C. hongkongensis</i>	Molecular Histological	– +
2002	Tong'an, Fujian (TA)	N	28	<i>C. ariakensis</i> , <i>C. hongkongensis</i>	Molecular Histological	– +
1999	Yamen River, Zhuhai, Guangdong (Z)	N	50	<i>C. hongkongensis</i>	Molecular	+
1999	Shouchang River, Yangjiang, Guangdong (YJ)	N	50	<i>C. hongkongensis</i>	Molecular	+
2002	Chengcun, Yangxi, Guangdong (CH)	C	37	<i>C. ariakensis</i> , <i>C. hongkongensis</i>	Molecular Histological	+ +
2002	Shenzhen, Guangdong (SZ)	N	13	<i>C. ariakensis</i> , <i>C. hongkongensis</i>	Molecular Histological	+ +
2002	Guandu, Zhanjiang, Guangdong (GD_C)	C	35	<i>C. hongkongensis</i>	Molecular	+
2002	Guandu, Zhanjiang, Guangdong (GD_N)	N	25	<i>C. hongkongensis</i>	Molecular Histological	+ +
2003	Zhanjiang, Guangdong (ZJ)	C	71	<i>C. ariakensis</i> , <i>C. hongkongensis</i>	Molecular	+
1999	Dafen River, Beihai, Guangxi Zhuang (DR)	N	50	<i>C. ariakensis</i> , <i>C. hongkongensis</i>	Molecular	+
2002	Dafen River, Beihai, Guangxi Zhuang (DR_02)	N	17	<i>C. ariakensis</i> , <i>C. hongkongensis</i>	Molecular Histological	– +
2002	Podi, Beihai, Guangxi Zhuang (PO)	C	39	<i>C. ariakensis</i> , <i>C. hongkongensis</i>	Molecular Histological	+ +
2003	Beihai, Guangxi Zhuang (BC)	N	64	<i>C. ariakensis</i> , <i>C. hongkongensis</i>	Molecular	+ +
2005	Beihai, Guangxi Zhuang (BC_05)	N	113	<i>C. ariakensis</i> , <i>C. hongkongensis</i>	Molecular Histological	+ –
2003	Lingshui, Hainan, Guangxi Zhuang (HN)	C	19	<i>Pinctada margaritifera</i> , <i>P. martensii</i> , unknown sp.	Molecular	+
<b>USA hatchery stocks</b>						
2000	SCA hatchery (SCA) (F1 of DR, PRC)	C	52	<i>C. ariakensis</i> , <i>C. hongkongensis</i>	Molecular	–
2002	NCA hatchery (NCA) (F1 of YR, PRC)	C	50	<i>C. ariakensis</i>	Molecular	+
2003	WCA hatchery (WCA)	C	49	<i>C. ariakensis</i>	Molecular	+

*sus* genus-specific primers that target the internal transcribed spacer (ITS) region of the rRNA gene complex (modified slightly from Casas et al. 2002), specifically PerkITS-85 (5'-CCGCTTTGTTTGGATCC-C-3') and PerkITS-750 (5'-ACATCAGGCCTTCTAAT-GATG-3'). Each PCR reaction contained the following: 20 mM Tris-HCl (pH 8.4), 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 0.2 mM of each dNTP, each primer at 1.0 μM, 0.025 U μl<sup>-1</sup> *Taq* polymerase, 0.05 mg ml<sup>-1</sup> BSA, and 0.5 μl genomic DNA (10 to 50 ng total). Amplifications were performed with an initial denaturation of 95°C for 4 min followed by 40 cycles of 95°C for 1 min, 53°C for 1 min, 65°C for 3 min, with final elongation at 65°C for 5 min. Following amplification, 4 μl of PCR product were analyzed as described above.

**Molluscan herpesvirus PCR assay.** The 'A' region of the molluscan herpesvirus genome encoding a gene of unknown function (Batista et al. 2007) was amplified using nested 'A' region primer pairs (Renault et al. 2000b). First a product of approximately 1000 bp was amplified, followed by an approximate 900 bp product in the nested reaction. For the A3 (5'-GCCAAC-CGTTGGAACCATAACAAGCG-3')/A4 (5'-GGGAAT-GAGGTGAACGAACTATAGACC-3') primer pair (external primers), the PCR reaction contained the following: 20 mM Tris-HCl (pH 8.4), 50 mM KCl, 1.0 mM MgCl<sub>2</sub>, 0.4 mg ml<sup>-1</sup> BSA, 0.8 mM of each dNTP, 0.24 μM of each primer, *Taq* polymerase at 0.24 U μl<sup>-1</sup>, and template DNA at 20 ng μl<sup>-1</sup>. Amplifications were performed with an initial denaturation at 94°C for 4 min, followed by 35 cycles of 94°C for 4 min, 50°C for 30 s, and 72°C for 30 s, with final elongation at 72°C for 5 min. Reaction conditions and reagent concentrations were the same for the internal amplification reaction using the A5 (5'-CGCCCCAACCACGATTTTTCACT-GACCC-3')/A6 (5'-CCCGTCAGATATAGGATGAGATTTG-3') primer pair, however 0.5 μl of the initial PCR reaction after the A3/A4 amplification was used in the A5/A6 reaction in place of genomic DNA. Following amplification using the A5/A6 primers, 5 μl of PCR product were analyzed by agarose gel electrophoresis as described above.

**In situ DNA probe hybridization (ISH) assays.** Paraffin wax-infiltrated tissues from 2 different animals, one collected from Podi, Beihai, China, in 1999, and another collected from Beihai, China, in 2005, were cut into 5 μm sections for *in situ* hybridization assays. A genus-specific, 5' digoxigenin-labeled genus-*Perkinsus* probe (Elston et al. 2004) was used to target unique SSU-rRNA sequences of *Perkinsus* spp. Digoxigenin-labeled oligonucleotides were obtained from Operon Biotechnologies. The ISH protocol of Stokes & Burreson (1995) was followed, with the modifications of Elston et al. (2004). Pronase at a final concentration of 1.25 mg ml<sup>-1</sup> was used for permeabiliza-

tion during a 30 min incubation. A probe concentration of 7 ng μl<sup>-1</sup> was used for hybridization. An anti-digoxigenin antibody linked to alkaline phosphatase was used in conjunction with NBT/BCIP for colorimetric detection of bound probe. Negative controls included duplicate histological sections of all tested samples, which received hybridization buffer without probe during hybridization incubations.

**PCR-RFLP identification of oyster host species.** Species identification of host oyster samples were carried out using a molecular diagnostic key based on the PCR amplification and restriction enzyme digestion of the ITS-1 gene region (Cordes & Reece 2005). PCR amplifications were carried out using the primers of Hedgecock et al. (1999). The PCR reaction contained the following: 20 mM Tris-HCl (pH 8.4), 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 0.2 mM of each dNTP, 0.1 μM of each primer, *Taq* polymerase at 0.25 U μl<sup>-1</sup> and template DNA at 0.2 ng μl<sup>-1</sup>. Amplifications were performed with an initial denaturation at 95°C for 3 min, followed by 30 cycles of 95°C for 1 min, 52°C for 2 min, and 72°C for 2 min, with a final elongation at 72°C for 5 min. Amplification products were digested with the diagnostic restriction enzymes *Hae*III or *Dde*I following manufacturer's protocols (New England Biolabs). Following amplification and digestion, 4 μl of the initial PCR product and 10 μl of the digested PCR product were electrophoresed on a 3% (1.5% agarose and 1.5% low-melt agarose) agarose gel, stained with ethidium bromide, and visualized under UV light. Banding patterns were compared to those of reference oyster species for species identifications.

**Cloning and sequencing.** PCR products of the *Perkinsus* spp. ITS region, and those amplified by primers designed to target OsHV-1 sequences, were cloned into the plasmid pCR®4-TOPO® and transformed into *Escherichia coli* using a TOPO TA Cloning® Kit (Invitrogen) following the manufacturer's protocols. Transformed bacterial colonies were screened for inserts using a boil-preparation method, followed by a PCR-based screening reaction using the M13 forward and reverse primer pairs supplied in the cloning kit, or by *Eco*RI digest following plasmid DNA isolation.

When using the *Eco*RI digest method, 4 ml of 2YT media were inoculated with transformed bacterial colonies and incubated for 12 to 15 h in a water bath at 37°C while shaking at 200 rpm. Plasmid DNA was purified from bacterial cultures using a Qiaprep Spin Miniprep Kit (Qiagen) and electrophoresed on a 2% agarose gel. *Eco*RI restriction digestions were performed in 15 μl reactions containing 3.0 μl of purified plasmid DNA, 10.2 μl of sterile distilled water, 1.5 μl of 10× reaction buffer and 0.3 μl of *Eco*RI restriction endonuclease. Plasmid DNA was digested at 37°C for 3 h, and 10 μl of the digested plasmid DNA was elec-

trophoresed on a 3% (1.5% agarose and 1.5% low melt agarose) agarose gel, stained with ethidium bromide and visualized under UV light.

For the PCR-based screening method, bacterial colonies were picked from agar plates using a sterile wooden toothpick and inoculated into 10 µl of sterile water in 200 µl plastic strip tubes. Inoculated water samples were boiled for 4 min at 94°C, and 0.5 µl of the boiled preparation were used in a PCR reaction using the M13 forward and reverse primers as described by Moss et al. (2006). Following amplification with the M13 primer pair, 3 µl of PCR product were electrophoresed on a 2% agarose gel, stained with ethidium bromide and visualized under UV light as described above.

Prior to sequencing, PCR products from clones containing the correct insert size were treated with shrimp alkaline phosphatase (SAP) and exonuclease I (*ExoI*) (Amersham Biosciences) in order to degrade nucleotides and single-stranded DNA (primers) remaining after PCR. We combined 5 µl of the M13 PCR product with 0.5 U of SAP and 5.0 U of *ExoI* and incubated at 37°C for 30 min, 80°C for 15 min and 15°C for 5 s.

Plasmid inserts or PCR products of plasmid inserts were sequenced bi-directionally, using the ThermoSequenase labeled primer cycle sequencing kit (Amersham Pharmacia) according to methods described by Reece & Stokes (2003), or using the Big Dye Terminator kit (Applied Biosystems) with M13 sequencing primers as described by Moss et al. (2006).

**Analysis of *Perkinsus* spp. and molluscan herpesvirus sequences.** *Perkinsus* spp. and molluscan herpesvirus sequences were compared to those deposited in GenBank and those compiled previously by researchers at VIMS, using BLAST (basic local alignment search tool) searches (Altschul et al. 1990) of the National Center for Biotechnology Information (NCBI) database.

Available ITS region sequences from *Perkinsus* spp. and 'A' fragment sequences of molluscan herpesviruses were downloaded from GenBank and included in phylogenetic analyses of the sequences obtained in this study. Representative *Perkinsus* spp. ITS region sequences included the following: *P. marinus* AY295177–AY295186; *P. chesapeaki* AF091541, AF440466, AF440468, AY876302, AY876304, AY876306, AY876308, AY876312, AY876314; *P. olseni* AF441207–AF441211, AF441213–AF441217; *P. mediterraneus* AY487834–AY487843; *P. honshuensis* DQ516696–DQ516702 and *P. qugwadi* AF15128. Representative molluscan herpesvirus sequences included the genome sequence of OsHV-1, AY509253, and sequences AY459364 and AY459362.

*Perkinsus* spp. ITS region and molluscan herpesvirus sequences were aligned separately using the CLUSTAL-W algorithm (Thompson et al. 1994) in

MacVector 8.0.1, with open and extend gap penalties of 7 and 3, respectively. Neighbor-joining and parsimony analyses of *Perkinsus* spp. ITS region sequences were conducted using PAUP\*4b10.0 (Swofford 2002). Bootstrap analyses were done with 10 random additions of 100 bootstrap replicates, with gaps treated as missing data. For jackknife analyses, 30% deletion was done with 10 random additions and 100 replicates, with gaps treated as missing data.

## RESULTS

### Host identifications

The species identification of each individual was determined using the molecular genetic PCR-RFLP key developed by Cordes & Reece (2005) (Table 1). Although only the host species *Crassostrea ariakensis* was targeted for this study, and the animals were identified by fishermen and scientists in Asia as *C. ariakensis*, many other *Crassostrea* species, and even a *Saccostrea* species, were found among the samples (Zhang et al. 2005, Cordes & Reece 2005).

### PCR-based screening results

PCR-based screening results from locations in Japan, China and Korea (Fig. 1) revealed that molluscan herpesviruses and *Perkinsus* spp. parasites are widespread in Asian populations of *Crassostrea ariakensis*, *C. hongkongensis* and other oyster species found at the same sites (Table 2). The Itoki River (Japan), Shouchang River and Chengcun (China), and Kahwa and Sumjin River (South Korea) sites had the highest prevalences of molluscan herpesviruses among locations sampled (Table 2). There was DNA evidence of *Perkinsus* spp. infections in *C. ariakensis* and other bivalve species at several Japanese and Chinese sampling sites. DNA from an undescribed *Perkinsus* species was detected in all samples collected between 1999 and 2005 from sites in southern China, indicating that the parasite is endemic to that region (Table 2).

*Perkinsus marinus* DNA was detected in VIMS hatchery stocks of *Crassostrea ariakensis*, however, none of the oysters screened from the VIMS hatchery was PCR-positive for molluscan herpesviruses (Table 2).

### Histological screening results

The viral, bacterial, protistan and metazoan parasites detected in oysters by histopathology at 8 sites

Table 2. *Perkinsus* spp. and molluscan herpesvirus (HV) pathogens. Prevalence data based on molecular assays. Only samples where either one or both of these pathogens were detected are listed. The bivalve species infected with either pathogen at a sampling site are indicated. See Fig. 1 and Table 1 for locations in SE Asia. VIMS = Virginia Institute of Marine Sciences, WCA = west coast USA *C. ariakensis*, NCA = F1 northern China *C. ariakensis*, *C.* = *Crassostrea*

Year	Sample	N	No. HV positive	% HV prevalence	No. <i>Perkinsus</i> sp. positive	% <i>Perkinsus</i> sp. prevalence	<i>Perkinsus</i> species	Bivalve species
<b>Japan</b>								
1999	Itoki River, Kyushu (IR)	50	13	26.0	9	18.0	<i>P. olseni</i>	<i>C. ariakensis</i>
2005	Mie Prefecture (JPAB05)	172	5	2.9	0	–		<i>C. ariakensis</i>
		29	2	6.9	0	–		<i>C. gigas</i>
		45	2	4.4	0	–		<i>C. sikamea</i>
<b>South Korea</b>								
2004	Kahwa River (KR)	35	10	28.6	0	–		<i>C. ariakensis</i>
2004	Sumjin (SR)	20	8	40.0	0	–		<i>C. ariakensis</i>
2004	Kanghwa Island, Incheon (IN)	20	1	5.0	0	–		<i>C. ariakensis</i>
<b>China</b>								
1999	Yellow River, Bohai Sea, Shandong (YR)	43	0	–	2	4.7	<i>P. olseni</i>	<i>C. ariakensis</i>
2002	Tong'an, Fujian (TA)	28	0	–	1	3.6	Undescribed	<i>C. ariakensis</i>
1999	Yamen River, Zhuhai, Guangdong (Z)	50	5	10.0	5	10.0	<i>P. olseni</i>	<i>C. hongkongensis</i>
1999	Shouchang River, Yangjiang, Guangdong (YJ)	50	10	20.0	1	2.0	<i>P. olseni</i>	<i>C. hongkongensis</i>
2002	Chengcun, Yangxi, Guangdong (CH)	37	4	10.8	6	16.2	Undescribed	<i>C. hongkongensis</i>
2002	Shenzhen, Guangdong (SZ)	13	0	–	1	7.6	Undescribed	<i>C. hongkongensis</i>
2002	Guandu, Zhanjiang, Guangdong, (GD_C)	35	0	–	4	11.4	Undescribed	<i>C. hongkongensis</i>
2002	Guandu, Zhanjiang, Guangdong (GD_N)	25	0	–	3	12.0	Undescribed	<i>C. hongkongensis</i>
2003	Zhanjiang, Guangdong, (ZJ)	1	0	–	1	100.0	Undescribed	<i>C. ariakensis</i>
		70	0	–	32	45.7	Undescribed	<i>C. hongkongensis</i>
1999	Dafen River, Beihai, Guangxi Zhuang (DR)	26	0	–	9	34.6	Undescribed	<i>C. ariakensis</i>
2002	Podi, Beihai, Guangxi Zhuang (PO) <sup>a</sup>	24	1	4.1	2	8.3	Undescribed	<i>C. hongkongensis</i>
		14	0	–	2	14.3	Undescribed	<i>C. ariakensis</i>
2003	Beihai, Guangxi Zhuang (BC)	24	0	–	2	8.3	Undescribed	<i>C. hongkongensis</i>
		59	0	–	11	18.6	Undescribed	<i>C. hongkongensis</i>
2005	Beihai, Guangxi Zhuang (BC_05)	12	0	–	4	33.3	Undescribed	<i>C. ariakensis</i>
		101	0	–	36	35.6	Undescribed	<i>C. hongkongensis</i>
2003	Lingshui, Hainan, Guangxi Zhuang (HN)	19	0	–	12	63.2	Undescribed	<i>Pinctada margaritifera</i> , <i>P. martensii</i> , unknown sp. —not distinguished
2002	VIMS NCA	49	0	–	10	20.4	<i>P. marinus</i>	<i>C. ariakensis</i>
2003	VIMS WCA	50	0	–	1	2.0	<i>P. marinus</i>	<i>C. ariakensis</i>

<sup>a</sup>No DNA remained for species identification of one oyster with *Perkinsus* sp. DNA in this sample

sampled in 2002 are listed in Table 3 and illustrated in Figs. 2 & 3. Most parasites were uncommon. Although molluscan herpesvirus DNA was detected by PCR in 4 of 37 animals screened from one of the samples that was also examined by histology, there was no histological evidence of herpes viral infection. In addition to the parasites listed in Table 3, a sample of 33 *Crassostrea hongkongensis* from southern China collected in 2003 had 18 of 26 female oysters (69%) infected with

a *Steinhausia*-like microsporidian in the ova (Fig. 2D). Unfortunately, the exact source location for this sample is unknown.

Among the 17 oysters sampled from Beihai (China) in 2005 that tested positive for *Perkinsus* spp. infections by PCR, only one showed lesions typical of *Perkinsus* sp. when examined histologically. In oyster BC05Ca-20, there were numerous mature *Perkinsus* sp. signet ring trophozoites of 5 to 10 µm diameter,




Fig. 2. Parasites observed by histopathology in *Crassostrea* sp. in China. (A) Viral gametocytic hypertrophy (arrow) in gonad of wild *C. hongkongensis* from Guandu, Zhanjiang, China. (B) *Chlamydia*-like inclusions (arrows) in digestive tubules of wild *Crassostrea hongkongensis* from Guandu, Zhanjiang, China. (C) Ciliates (arrows) attached to gill epithelium in cultured *C. hongkongensis* from Chengcun, Yangxi, Guangdong, China. (D) *Steinhausia*-like microsporidian (arrow) in ovum of wild *C. hongkongensis* from southern China. (E) Spores of *Nematopsis*-like gregarines (arrows) in cultured *C. hongkongensis* from Guandu, Zhanjiang, China. (F) Encapsulated metacestode (arrow) in gill tissue of wild *C. hongkongensis* or *C. ariakensis* from Dafen River, Beihai, Guangxi Zhuang, China



Table 3. Histological analysis of oysters collected at 7 locations in China in 2002. See Fig. 1 and Table 1 for locations

Sampling location	N	Natural (N), Cultured (C)	Viral gametocytic hypertrophy	<i>Chlamydia</i> -like organisms	<i>Perkinsus</i> spp.	Ciliates	<i>Nematopsis</i> sp.	Cestodes
Dajiawa, Shandong	29	N	0	0	0	4	0	0
Longhai, Fujian	28	C	0	3	0	2	0	0
Tong'an, Fujian	29	N	0	1	0	2	11	1
Chengcun, Yangxi, Guangdong	31	C	0	1	1	2	0	1
Shenzhen, Guangdong	26	N	0	1	0	1	0	0
Guandu, Zhanjiang, Guangdong	30	N	1	1	0	0	9	0
Guandu, Zhanjiang, Guangdong	60	C	1	6	3	0	14	0
Dafen River, Beihai, Guangxi Zhuang	30	N	0	1	0	2	0	6
Podi, Beihai, Guangxi Zhuang	30	C	0	2	2	2	0	5




Fig. 3. (A) Mayer's hematoxylin and eosin-stained histological section of a *Crassostrea hongkongensis* oyster showing *Perkinsus* sp. cells (arrows) in the stomach epithelium. (B) *In situ* hybridization with the *Perkinsus* sp. genus-specific probe to *Perkinsus* sp. cells (arrows) in the stomach epithelium

subdividing 5 to 10 μm diameter schizonts, and clusters of multiple immature sibling trophozoites of 3 to 4 μm diameter (Fig. 3). *Perkinsus* sp. parasite cells (552) enumerated in one section occurred (1) in multifocal stomach epithelium lesions (43%), (2) rare lesions in digestive gland epithelia, (3) among con-

nective tissues of the mantle (27%), the visceral mass (15%) and gills (10%) and (4) systemically circulating both free and phagocytosed within host hemocytes in the oyster vasculature (3%).

### ISH results

*In situ* hybridization probes designed to target *Perkinsus* species hybridized only to *Perkinsus* sp. cells in oyster samples that were PCR positive for *Perkinsus* sp. DNA. Fig. 3 shows 2 consecutive sections from *Crassostrea hongkongensis* from Beihai (China) BC05Ca-20 (described above) infected with the undescribed *Perkinsus* species. Fig. 3A is a Mayer's hematoxylin and eosin-stained section showing *Perkinsus* sp. cells in the stomach epithelium of the oyster. Fig. 3B shows positive *in situ* hybridization with the genus-specific probe to *Perkinsus* sp. cells. The *Perkinsus* genus-specific probe labeled cells in connective tissues of the gills and mantle, as well as in the epithelia of intestine and stomach. *In situ* hybridization reactions conducted without probe produced no signal in tissues of the same host oyster.

### *Perkinsus* spp. ITS region sequences

ITS region PCR amplification products were sequenced from a selected number of positive oyster samples. All sequences were deposited in GenBank (Table 4). Sequencing of the ITS region amplicons from the genus *Perkinsus* assay indicated that the Chesapeake Bay native *P. marinus* was found in VIMS hatchery stocks of *Crassostrea ariakensis*. This is not surprising, since hatchery-reared oysters are exposed to water coming from the adjacent York River, which is within the native range of *P. marinus*. ITS region sequences from oysters from the NCA and WCA

hatchery stocks formed a monophyletic clade with known *P. marinus* sequences, with 100% bootstrap support in both neighbor joining distance analysis and in maximum parsimony analysis (Figs. 4 & 5). *Perkinsus* sp. ITS region sequences amplified from DNAs extracted from *C. ariakensis* and *C. hongkongensis* oysters collected from several Asian sites, including the Yamen and Yellow rivers (China) and the Itoki River, Kyushu (Japan) formed a monophyletic clade with known *P. olseni* ITS region sequences in both neighbor joining distance analysis and maximum parsimony analysis (100% bootstrap support) (Figs. 4 & 5).

*Crassostrea ariakensis* and *C. hongkongensis* collected from locations in China along the southern coast from Tong'an, Fujian to the Dafen River, Beihai, appeared to be infected with an undescribed *Perkinsus* species. The genus-specific *Perkinsus* spp. primers amplified unique nucleotide sequence fragments of approximately 689 bp from numerous oysters. We sequenced 42 clones from 16 individual oysters selected from 6 different samples, and they were 99.3% similar (uncorrected-*p*) to each other. Overall, 28 of the 42 clones (67%) shared a common ITS rRNA sequence,

while the other 14 clones all were all unique. BLAST analyses of GenBank suggested that these were *Perkinsus* sp. sequences. Pairwise distances and molecular phylogenetic analyses confirmed that these sequences were closely related to those of known *Perkinsus* species. In both neighbor joining and maximum parsimony analyses, these new parasite ITS region sequences grouped with those of known *Perkinsus* spp. (Figs. 4 & 5). Mean pairwise distances (uncorrected-*p*) of ITS region nucleotide sequences within known *Perkinsus* spp. ranged from 0.2% in the undescribed *Perkinsus* sp. to 1.6% variation in *P. chesapeakei*. Only one ITS region nucleotide sequence was available for *P. qugwadi*, therefore mean intraspecific ITS variation could not be calculated. The ITS region of this new *Perkinsus* sp. was most distantly related to *P. qugwadi* (63.6% to 63.7% similarity), and most closely related to the recently described species, *P. honshuensis* (89.0% to 89.8% similarity) (Table 5). Nucleotide sequences from this apparently new parasite taxon form a well-supported sister group (100% bootstrap support within this species) to the larger *Perkinsus* spp. clade that includes *P. marinus*, *P. mediterraneus* and *P. olseni* (Figs. 4 & 5).

Table 4. *Perkinsus* spp. GenBank accession numbers associated with ITS rRNA and molluscan herpesvirus (HV) sequences generated in this study. See Fig. 1 and Table 1 for locations in SE Asia. VIMS = Virginia Institute of Marine Sciences, WCA = west coast USA *C. ariakensis*, NCA = F1 northern China *C. ariakensis*

Pathogen	Source	GenBank accession numbers
<i>Perkinsus olseni</i>	Itoki River, Kyushu, Japan (IR)	EF204075, EF204076
<i>Perkinsus olseni</i>	Yellow River, Bohai Sea, Shandong, China (YR)	EF204073, EF204074
<i>Perkinsus olseni</i>	Yamen River, Zhuhai, Guangdong, China (Z)	EF204070–EF204072
Undescribed <i>Perkinsus</i> sp.	Tong'an Fujian, China (TA)	EF204046–EF204050
Undescribed <i>Perkinsus</i> sp.	Chengcun, Yangxi, Guangdong, China (CH)	EF204034, EF204035, EF204043
Undescribed <i>Perkinsus</i> sp.	Shenzhen, Guangdong, China (SZ)	EF204015–EF204018, EF204029–EF204031, EF204036, EF204038, EF204039, EF204041
Undescribed <i>Perkinsus</i> sp.	Guandu, Zhanjiang, Guangdong, China (GD)	EF204022, EF204051–EF204056
Undescribed <i>Perkinsus</i> sp.	Dafen River, Beihai, Guangxi Zhuang, China (DR)	EF204021, EF204024–EF204028, EF204040, EF204044, EF204045
Undescribed <i>Perkinsus</i> sp.	Podi, Beihai, Guangxi Zhuang, China (PO)	EF204019, EF204020, EF204023, EF204032, EF204033, EF204042
<i>Perkinsus marinus</i>	VIMS hatchery (WCA)	EF204008–EF204011
<i>Perkinsus marinus</i>	VIMS hatchery (NCA)	EF204012–EF204014
HV	Ariake Sea, Mie, Japan (JPAB05)	EF221836–EF221839
HV	Kahwa River, South Korea (KR)	EF221840
HV	Sumjin River, South Korea (SR)	EF221841
HV	Yamen River, Zhuhai, Guangdong, China (Z)	EF221843
HV	Chengcun, Yangxi, Guangdong, China (CH)	EF221842
HV	Shouchang River, Yangjiang, Guangdong, China (YJ)	EF221844

### Molluscan herpesvirus sequences

Molluscan herpesvirus DNA was found in *Crassostrea ariakensis* populations at a site in the Itoki River, Ariake Bay (Japan), sampled in 1999, and was also found in the 2005 samples taken from Mie Prefecture (Japan) near the first sampling site. A portion of oysters collected at the Kahwa River and Sumjin River

sampling sites (South Korea) as well as in the Yamen River, Shouchang River, Dafen River, and Chengcun sites (China) were also positive for molluscan herpesvirus DNA (Table 3).

Subsets of oysters positive for molluscan herpesvirus DNA from the Yamen River, Shouchang River, and Chengcun (China) and Mie (Japan) were chosen as representatives from those populations, and their herpesvirus DNA was cloned and sequenced. The molluscan herpesvirus DNA amplified from *Crassostrea ariakensis* collected from the Chinese sites and from the 2 South Korean sites had variations among DNA sequences, and we found polymorphic sequences within and between oysters from different locations. The level of polymorphism between the viral sequences was low (3.7%), with a combined total of 34 individual randomly distributed single nucleotide differences over the entire 917 bp sequence from all of the clones analyzed. We also found 3 polymorphic nucleotide sites that generally had consistent polymorphisms across all individuals within a collection site, except for individuals sampled in 2005 from the Ariake Sea, Japan (Fig. 6). The molluscan herpesvirus DNA amplified from *C. ariakensis* specimens collected from the Yamen River and Shouchang River (China) were similar to viral sequences found in French *C. gigas* (LeDeuff & Renault 1999, Renault et al. 2000b), sharing a thymine at a polymorphic site 604 bp from the 5' end of the fragment (primers removed). The molluscan herpesvirus sequences amplified from Chengcun (China) samples of *C. ariakensis* and those from both the Kahwa and Sumjin rivers (South Korea) are similar to that found in *C. gigas* from Tomales Bay, California (USA) (Friedman et al. 2005) and in the Itoki River (Japan) *C. ariakensis*, sharing a cytosine at that particular polymorphic site. The viral DNA amplified from the Itoki River samples and from some of the 2005 Ariake Sea (Japan) samples appears to have a unique polymorphism at a site 115 bp from the 5' end of the fragment, sharing an adenine residue where all other viral sequences share a guanine. From the 2005 Ariake Sea (Japan) samples, molluscan herpesvirus PCR fragments were cloned and sequenced from 3 molluscan herpesvirus-positive *C. ariakensis* oysters and 1 positive *C. sikamea*. The DNA sequences from these oysters suggest that 2 of the animals (1 specimen of *C. ariakensis* and the *C. sikamea* specimen) were infected with the same strain of molluscan herpesvirus found in




Fig. 4. *Perkinsus* spp. ITS rRNA gene sequences. Neighbor joining tree determined by analysis of the ITS gene sequences of known *Perkinsus* spp. sequences and the ITS gene sequences of those *Perkinsus* spp. found in oysters collected in Japan and China. GenBank accession numbers associated with each sample from this study (**bold**) are listed in Table 4




Fig. 5. *Perkinsus* spp. ITS rRNA gene sequences. Maximum parsimony tree determined by analysis with 100 replicates of 10 random additions of the ITS gene sequences of known *Perkinsus* spp. sequences and the ITS gene sequences of those *Perkinsus* spp. found in oysters collected in Japan and China. Maximum parsimony bootstrap support values for each clade are given above the lines and jackknife support values are given below the lines. **Bold** support values indicate species clades. GenBank accession numbers associated with each sample from this study (**bold**) are listed in Table 4

the Japanese *C. ariakensis* collected in 1999 from the Itoki River. The other 2 *C. ariakensis* specimens were infected with the strain commonly seen in France and parts of China. Viral DNAs amplified from Chengcun (China) and Korean *C. ariakensis* shared an additional polymorphic site 761 bp from the 5' end of the A5/A6 fragment. These viral sequences share a guanine at this site, where all other viral sequences share a cytosine.

## DISCUSSION

Several potential disease agents, including 2 different *Perkinsus* species, molluscan herpesviruses and a *Steinhausia*-like microsporidian, were detected in oysters collected from sites in Asia where potential *Crassostrea ariakensis* broodstocks might originate. Standard ICES protocols should minimize the risk of introducing horizontally transmitted pathogens with

Table 5. *Perkinsus* spp. Range of sequence similarities and pairwise distances (uncorrected 'p' values) observed among rDNA ITS region sequences of currently accepted species from GenBank, and those of the undescribed species obtained in this study. The range of observed within-species sequence distance is given across the diagonal. Raw distance value ranges between species are given above the diagonal, and ranges for percent distance values are given below

Species	<i>Perkinsus</i>					Undescribed <i>Perkinsus</i> sp.	<i>Perkinsus qugwadi</i>
	<i>marinus</i>	<i>chesapeakei</i>	<i>mediterraneus</i>	<i>olseni</i>	<i>honshuensis</i>		
<i>P. marinus</i>	<b>0.000–0.004</b> <b>0.4%</b>	0.119–0.139	0.053–0.069	0.049–0.063	0.052–0.062	0.112–0.126	0.323–0.330
<i>P. chesapeakei</i>	11.9–13.9%	<b>0.000–0.016</b> <b>1.6%</b>	0.134–0.148	0.126–0.138	0.117–0.126	0.156–0.175	0.342–0.350
<i>P. mediterraneus</i>	5.3–6.9%	13.4–14.8%	<b>0.000–0.007</b> <b>0.7%</b>	0.050–0.065	0.029–0.040	0.119–0.129	0.331–0.335
<i>P. olseni</i>	4.9–6.3%	12.6–13.8%	5.0–6.5%	<b>0.000–0.003</b> <b>0.3%</b>	0.049–0.060	0.107–0.119	0.333–0.336
<i>P. honshuensis</i>	5.2–6.2%	11.7–12.6%	2.9–4.0%	4.9–6.0%	<b>0.000–0.006</b> <b>0.6%</b>	0.102–0.110	0.327–0.331
Undescribed <i>Perkinsus</i> sp.	11.2–12.6%	15.6–17.5%	11.9–12.9%	10.7–11.9%	10.2–11.0%	<b>0.000–0.002</b> <b>0.2%</b>	0.363–0.364
<i>P. qugwadi</i>	32.3–33.0%	34.2–35.0%	33.1–33.5%	33.3–33.6%	32.7–33.1%	36.3–36.4%	<b>0.0<sup>a</sup></b> <b>0.0%<sup>a</sup></b>

<sup>a</sup>Only one sequence for *P. qugwadi* was available




Fig. 6. Alignment of molluscan herpesvirus 'A' region sequences isolated from oysters showing the polymorphic sites unique to viral sequences at each location. Viral sequences found in France and in Tomales Bay, California, (USA) have been deposited previously in GenBank, and accession numbers are listed accordingly. For other sample codes, see Table 1

the host, since broodstock would be held under strict quarantine conditions, and only progeny of oysters brought to the USA would be introduced into the environment. However, neither the ICES protocols, nor a mechanical procedure such as cleansing gametes to remove superficial parasites, would prevent transmission of pathogens if they infect the gametes themselves. If vertical transmission of any pathogen identified in this survey occurred, they could be introduced to the Chesapeake Bay by vertical transmission from infected broodstock to F1 or F2 *C. ariakensis* progeny oysters, with the potential for serious negative impacts to already depleted native oyster populations. In addition, *C. ariakensis* could act as a reservoir host for exotic pathogens that may be introduced by other means. For example, ballast water may have been the

source of a possible exotic *Bonamia* sp. that caused a severe mortality event during 2003 in triploid *C. ariakensis* deployed for experimental purposes in North Carolina, USA waters (Burreson et al. 2004), but has not been documented to infect local native bivalves. The possibility of vertical transmission of viruses among aquatic organisms is recognized (Bootland et al. 1991, Lo et al. 1997, Tsai et al. 1999), and recent transmission studies in France with *C. gigas* detected OsHV-1 in 3 successive generations of oysters (Barbosa-Solomieu et al. 2005). In our study, 3 genetic variants of molluscan herpesviruses were detected by PCR.

Two of the pathogens observed histologically pose a threat of introduction to Chesapeake Bay via infected broodstock. Viral gametocytic hypertrophy and the

*Steinhausia*-like microsporidian were both observed in oyster ova. Viral gametocytic hypertrophy has been reported in *Crassostrea virginica* from Chesapeake Bay (Farley 1978), and a *Steinhausia*-like microsporidian has been reported in the clam *Macoma balthica* in Chesapeake Bay (Farley 1977). The parasites observed in Asian oysters, however, may be strains or species different from those in Chesapeake Bay, and could pose problems if introduced. There is substantial evidence for vertical transmission of some microsporidian parasites of invertebrate hosts (Kelly et al. 2003, Galbreath et al. 2004, Vizoso & Ebert 2004), therefore indicating a real possibility for vertical transmission of these Asian pathogens.

Although there is no current evidence to suggest that *Perkinsus* sp. parasites may be vertically transmitted, and therefore might not be introduced to a new area through importation of small numbers of infected broodstock that are held in quarantine, Karlsson (1991) found protozoan cells that were described as '*Perkinsus*-like', but were not a true *Perkinsus* sp., in male and female scallop gonads. Subsequent infection of the larvae post-spawning was observed, suggesting the possibility of vertical transmission (Karlsson 1991, Whyte et al. 1993). ICES protocols are being followed; however the potential for vertical transmission of the pathogens observed in this study and the associated risks of introducing the non-native host must be recognized. Consistent and careful disease testing of deployed oysters may be required if an introduction occurs. It should be noted that there was no evidence of molluscan herpesviruses in the current VIMS hatchery stocks of *C. ariakensis* that we screened, and although *Perkinsus* sp. DNA was detected in these samples, it was *P. marinus* DNA, a native species. There is currently no evidence of *P. olseni* or the new undescribed *Perkinsus* species in the VIMS stocks.

Most parasites observed in the histological analyses conducted here (Table 3) were generally uncommon, and are unlikely to be introduced to a new area via infected broodstocks. Histological observations indicated tissue tropisms by *Perkinsus* sp. pathogens among both connective tissues and digestive system epithelia of oysters from Beihai, China. These observations are consistent with the detection of *Perkinsus* sp. DNAs in this sample using molecular tools (see below).

Molecular detection assays identified *Perkinsus* spp. and molluscan herpesvirus DNA in many of the samples surveyed. *P. olseni* DNA was detected in several samples of Asian *Crassostrea ariakensis* and *C. hongkongensis*, and is known to be widely distributed among molluscs in the Pacific and eastern Atlantic oceans (Lester & Davis 1981). Following synonymization with *P. atlanticus* (Murrell et al. 2002), *P. olseni* has been reported to infect a variety of hosts from around

the world (Villalba et al. 2005), including the black-lipped and the green-lipped abalone in Australia (Lester & Davis 1981), the carpet shell clam in Portugal (Azevedo 1989), and the Manila clam in Spain, Portugal, northern China, Korea, and Japan (Dungan & Reece 2006). The discovery of *P. olseni* in *C. ariakensis* and *C. hongkongensis* in Japan and northern China expands the currently known geographic and host distribution of that parasite.

In addition to *Perkinsus olseni*, a previously undescribed *Perkinsus* species was found in populations of *Crassostrea ariakensis* and *C. hongkongensis* in southern China, as well as in some pearl oysters and unidentified bivalves. As discussed in several recent publications (Burreson et al. 2005, Reece & Dungan 2006), molecular methods are the only reliable way to distinguish *Perkinsus* spp., because host and environmental elements may influence parasite morphological characteristics. ITS region *Perkinsus* sp. sequences amplified from southern China host sample DNAs grouped with those sequences from other *Perkinsus* spp. However, as with each of the other species, they form a unique monophyletic clade within the genus, indicating that this is a unique species. The similarity among the sequences within this clade, and the genetic distance between these sequences and those of other *Perkinsus* spp. are consistent with the distances observed in previous studies that have used the ITS region sequences to discriminate species or strains (Brown et al. 2004, Burreson et al. 2005, Dungan & Reece 2006). Sequencing of multiple gene regions is recommended to confirm *Perkinsus* spp. phylogenies based on the ITS rRNA region. To date, phylogenies that were based on other regions, including the large subunit ribosomal RNA and actin gene sequences, have confirmed results obtained using ITS rRNA (Dungan & Reece 2006, Dungan et al. 2007).

More than 30 yr ago Farley et al. (1972) reported, based on transmission electron microscopy, a herpes-like virus infecting *Crassostrea virginica* oysters from Maine. A recent survey of oysters from the Atlantic, Gulf of Mexico and Pacific coasts of the USA, which used molecular diagnostic tools designed originally to detect OsHV-1, indicated that a herpes-like virus is currently found only on the Pacific coast of the United States in Tomales Bay, California and not along the Atlantic or Gulf of Mexico coasts (Friedman et al. 2005). Molluscan herpes-like viruses also occur in France (Nicolas et al. 1992, Comps & Cochenne 1993, Renault et al. 1994, 2000a, 2001, Arzul et al. 2002), Australia (Hine & Thorn 1997), New Zealand (Hine & Thorn 1997, Hine et al. 1998), and Taiwan (Chang et al. 2005). Previously, at least 2 genetic strains of molluscan herpesviruses were recognized; with DNA sequence polymorphisms in the 'A' region distinguish-

ing between the original strain (OsHV-1) found in France, another French variant was described later (Arzul et al. 2001), and a recently described variant strain was found in California (Friedman et al. 2005). Based on our observation of 3 total polymorphic sites within the 'A' region, we suggest that there are at least 2 different genetic strains of molluscan herpesvirus in Japan, one strain in Korea and 2 strains in China.

Initial sequencing of molluscan herpesvirus DNA detected in *Crassostrea ariakensis* from sites in Korea and in *C. hongkongensis* in Chengcun (China) suggested, based on the site at 604 bp, that these oysters could be infected with the same genetic strain that was detected in *C. gigas* from Tomales Bay, California (Friedman et al. 2005). However, additional DNA sequencing revealed that the viral DNAs amplified from oysters in Chengcun (China) and from Korean *C. ariakensis* share an additional polymorphic site, making them different from the California molluscan herpesvirus strain. Interestingly, molluscan herpesvirus sequences found in oysters from Shouchang River and Yamen River (China) had the same 'A' region sequence as OsHV-1 that was first detected in *C. gigas* from France (Le Deuff & Renault 1999, Renault et al. 2000b). However, molluscan herpesvirus sequences from Japanese oysters have a unique pattern of differences at the 3 polymorphic sites, suggesting that the 2 Japanese molluscan herpesvirus strains could be unique to that geographic region. Sequencing of additional gene regions of the molluscan herpesvirus found in Asia may further discriminate strains suggested by the polymorphisms observed here in the 'A' fragment. Overall, this study expands the current known host geographic range of molluscan herpesviruses to include Japan, China and Korea, where it infects *C. ariakensis*, *C. hongkongensis*, *C. gigas* and *C. sikamea*.

**Acknowledgements.** This research was supported by Virginia Sea Grant College Program award NA96RG0025 and NOAA/NMFS awards NA17FU2891 and NA04NMF4570431. Collection assistance was provided by X. Du of Zhanjiang Ocean University, PR China; J. Higano of the Environmental Management Division, National Research Institute of Aquaculture, Fisheries Research Agency, Mie, Japan; X. Guo of Rutgers University; H. An, National Fisheries Research and Development Institute, Pusan, Korea; and R. Carnegie and K. Johnson of VIMS. Oyster species identification assistance was kindly provided by J. Xiao of VIMS. Histological services and diagnoses were provided in part by R. Crockett and S. Denny of VIMS. VIMS contribution number 2876.

#### LITERATURE CITED

- Allen SK Jr (2005) Stalemate over the new oyster. *Virginia Mar Res Bull* 37(2):2–16
- Altschul SF, Gish W, Miller W, Meyers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215:403–410
- Andrews JD, Hewatt WG (1957) Oyster mortality studies in Virginia. II. The fungus disease caused by *Dermocystidium marinum* in oysters of Chesapeake Bay. *Ecol Monogr* 27:1–26
- Arzul I, Nicolas JL, Davidson AJ, Renault T (2001) French scallops: a new host for *Ostreid herpesvirus-1*. *Virology* 290:342–349
- Arzul I, Renault T, Thebault A, Gerard A (2002) Detection of oyster Herpes Virus DNA and proteins in asymptomatic *Crassostrea gigas* adults. *Virus Res* 84:151–160
- Audemard C, Ragone Calvo LM, Paynter KT, Reece KS, Burrenson EM (2006) Real-time PCR investigation of parasite ecology: *in situ* determination of oyster parasite *Perkinsus marinus* transmission dynamics in lower Chesapeake Bay. *Parasitol* 132:827–842
- Azevedo C (1989) Fine structure of *Perkinsus atlanticus* n. sp. (Apicomplexa, Perkinsea) parasite of the clam *Ruditapes decussatus* from Portugal. *J Parasitol* 75:627–635
- Batista FM, Arzul I, Pepin JF, Ruano F, Friedman CS, Boudry P, Renault T (2007) Detection of ostreid herpesvirus 1 DNA by PCR in bivalve mollusks: a critical review. *J Virol Meth* 139:1–11
- Barbosa-Solomieu V, Miossec L, Vazquez-Juarez R, Ascencio-Valle F, Boudry P, Renault T (2005) *Ostreid herpesvirus 1* detection among three successive generations of Pacific oysters (*Crassostrea gigas*). *Virus Res* 107:47–56
- Bootland LM, Dobos P, Stevenson RMW (1991) The IPNV carrier state and demonstration of vertical transmission in experimentally infected brook trout. *Dis Aquat Org* 10:13–21
- Brown G, Hudson KL, Reece KS (2004) Multiple polymorphic sites at the ITS and ATAN loci among and within cultured isolates of *Perkinsus marinus*. *J Eukaryot Microbiol* 51:312–320
- Burrenson EM, Ragone Calvo LM (1996) Epizootiology of *Perkinsus marinus* disease of oysters in Chesapeake Bay, with emphasis on data since 1985. *J Shellfish Res* 15:17–34
- Burrenson EM, Stokes NA, Carnegie RB, Bishop MJ (2004) *Bonamia* sp. (Haplosporidia) found in non-native oysters, *Crassostrea ariakensis*, in Bogue Sound, North Carolina. *J Aquat Anim Health* 16:1–9
- Burrenson EM, Reece KS, Dungan CF (2005) Molecular, morphological, and experimental evidence support the synonymy of *Perkinsus chesapeakei* and *Perkinsus andrewsi*. *J Eukaryot Microbiol* 52:258–270
- Calvo GW, Luckenbach MW, Allen SK Jr, Burrenson EM (1999) A comparative field study of *Crassostrea gigas* and *Crassostrea virginica* in relation to salinity in Virginia. *J Shellfish Res* 18:465–473
- Calvo GW, Luckenbach MW, Allen, SK Jr., Burrenson EM (2001) A comparative field study of *Crassostrea ariakensis* (Fujita 1913) and *Crassostrea virginica* (Gmelin) in relation to salinity in Virginia. *J Shellfish Res* 20:221–229
- Casas SM, Villalba A, Reece KS (2002) Study of perkinsosis in the carpet shell clam *Tapes decussates* in Galicia (NW Spain). I. Identification of the aetiological agent and *in vitro* modulation of zoosporulation by temperature and salinity. *Dis Aquat Org* 50:51–65
- Chang PH, Kuo ST, Lai SH, Yang HS, Ting YY, Hsu CL, Chen HC (2005) Herpes-like virus infection causing mortality of cultured abalone *Haliotis diversicolor supertexta* in Taiwan. *Dis Aquat Org* 65:23–27
- Choi KS, Park KI (1997) Report on the occurrence of *Perkinsus* sp. in the Manila clams, *Ruditapes philippinarum* in Korea. *J Aquaculture* 10:227–237

- Comps M, Cochenne N (1993) A herpes-like virus from the European oyster *Ostrea edulis* L. *J Invertebr Pathol* 62: 201–203
- Cordes JF, Reece KS (2005) Genetic identification of oyster species based on restriction fragment-length polymorphism (RFLP) analysis of two molecular markers amplified using the polymerase chain reaction (PCR). *J Shellfish Res* 24:319
- Da Ros L, Cazonier WJ (1985) *Perkinsus*, a protistan threat to bivalve culture in the Mediterranean basin. *Bull Eur Assoc Fish Pathol* 5:23–27
- Dungan CF, Reece KS (2006) *In vitro* propagation of two *Perkinsus* spp. parasites from Japanese manila clams *Venerupis philippinarum* and description of *Perkinsus honshuensis* n. sp. *J Eukaryot Microbiol* 53:316–326
- Dungan CF, Hamilton RM, Hudson KL, McCollough CB, Reece KS (2002) Two epizootic diseases in Chesapeake Bay commercial clams, *Mya arenaria* and *Tagelus plebius*. *Dis Aquat Org* 50:67–78
- Dungan CF, Reece KS, Moss JA, Hamilton RM, Diggles BK (2007) *Perkinsus olseni* *in vitro* isolates from the New Zealand clam *Austrovenus stutchburyi*. *J Eukaryot Microbiol* 54:263–270
- Elston RA, Dungan CF, Meyers TR, Reece KS (2004) *Perkinsus* sp. infection risk for Manila clams, *Venerupis philippinarum* (A. Adams and Reeve, 1850) on the Pacific coast of North and Central America. *J Shellfish Res* 23: 101–105
- Farley CA (1977) Neoplasms in estuarine mollusks and approaches to ascertain causes. In: Kraybill HF, Dawe CJ, Harshbarger JC, Tardiff RG (eds) *Aquatic pollutants and biologic effects with emphasis on neoplasia*. *Ann NY Acad Sci* 298:225–232
- Farley CA (1978) Viruses and virus like lesions in marine mollusks. *Mar Fish Rev* 40(10):18–20
- Farley CA, Banfield WG, Kasnic JRG, Foster WS (1972) Oyster herpes-type virus. *Science* 178:759–760
- Figueras A, Robledo JAF, Novoa B (1992) Occurrence of haplosporidian and *Perkinsus*-like infections in carpet-shell clams *Ruditapes decussates* (Linnaeus, 1758) of the Ría de Vigo (Galacia, NW Spain). *J Shellfish Res* 11: 377–382
- Friedman CS, Estes RM, Stokes NA, Burge CA and 5 others (2005) Herpes virus in juvenile Pacific oysters *Crassostrea gigas* from Tomales Bay, California, coincides with summer mortality episodes. *Dis Aquat Org* 63:33–41
- Galbreath JGMS, Smith JE, Terry RS, Becnel JJ, Dunn AM (2004) Invasion success of *Fibrillanosema cragonycis*, n.sp., n.g.: a novel vertically transmitted microsporidian parasite from the invasive host *Crangonyx pseudogracilis*. *Int J Parasitol* 34:235–244
- Goggin CL, Lester RJG (1995) *Perkinsus*, a protistan parasite of abalone in Australia: a review. *Mar Freshw Res* 46: 639–646
- Grabowski JH, Powers SP, Peterson CH, Powers MJ, Green DP (2003) Consumer ratings of non-native (*Crassostrea gigas* and *Crassostrea ariakensis*) vs. native (*Crassostrea virginica*) oysters. *J Shellfish Res* 22:21–30
- Hedgecock D, Li G, Banks MA, Kain Z (1999) Occurrence of the Kumamoto oyster *Crassostrea sikamea* in the Ariake Sea, Japan. *Mar Biol* 133:65–68
- Hine PM, Thorn T (1997) Replication of herpes-like viruses in haemocytes of adult flat oysters *Ostrea angasi*: an ultra-structural study. *Dis Aquat Org* 29:189–196
- Hine PM, Wesney B, Hay BE (1992) Herpesviruses associated with mortalities among hatchery-reared larval Pacific oysters *Crassostrea gigas*. *Dis Aquat Org* 12:135–142
- Hine PM, Wesney B, Besant P (1998) Replication of a herpes-like virus in larvae of the flat oyster *Tiostrea chilensis* at ambient temperatures. *Dis Aquat Org* 32:161–171
- ICES (International Council for the Exploration of the Sea) (2005) Code of practice on the introductions and transfers of marine organisms. ICES, Copenhagen
- Karlsson JD (1991) Parasites of the bay scallop, *Argopecten irradians* Lamark (1819). In: Schumay SE, Sandifer PA (eds) *An international compendium of scallop biology and culture*. World Aquaculture Workshop no. 1. World Aquaculture Society, Baton Rouge, LA, p 180–190
- Kelly A, Hatcher MJ, Dunn AM (2003) The impact of a vertically transmitted microsporidian, *Nosema granulosis* on the fitness of its *Gammarus deubeni* host under stressful environmental conditions. *Parasitol* 126:119–124
- Le Deuff RM, Renault T (1999) Purification and partial genome characterization of a herpes-like virus infecting the Japanese oyster, *Crassostrea gigas*. *J Gen Virol* 80:1317–1322
- Le Deuff RM, Nicola JL, Renault T, Cochenne N (1994) Experimental transmission of herpes-like virus to axenic larvae of the Pacific oyster, *Crassostrea gigas*. *Bull Eur Fish Pathol* 14:69
- Lester RJG, Davis GHG (1981) A new *Perkinsus* species (Apicomplexa, Perkinsea) from the abalone *Haliotis ruber*. *J Invertebr Pathol* 37:181–187
- Lo CF, Ho CH, Chen CH, Liu KF and 9 others (1997) Detection and tissue tropism of white spot syndrome baculovirus (WSBV) in captured brooders of *Panesus monodon* with a special emphasis on reproductive organs. *Dis Aquat Org* 30:53–72
- Mackin JG, Owen HM, Collier A (1950) Preliminary note on the occurrence of a new protistan parasite, *Dermocystidium marinum* n. sp. in *Crassostrea virginica* (Gmelin). *Science* 111:328–329
- Mann R, Burreson EM, Baker PK (1991) The decline of the Virginia oyster fishery in Chesapeake Bay: considerations for introduction of non-endemic species, *Crassostrea gigas* (Thunberg, 1773). *J Shellfish Res* 10:379–388
- Medlin L, Elwood HJ, Stickel S, Sogin ML (1988) The characterization of enzymatically amplified eukaryotic 16S-like rRNA-coding proteins. *Gene* 71:491–499
- Montes J, Durfort M, García-Valero J (2001) Parasitism by the protozoan *Perkinsus atlanticus* favours the development of opportunistic infections. *Dis Aquat Org* 46:57–66
- Moss JA, Burreson EM, Reece KS (2006) Advanced *Perkinsus marinus* infections in *Crassostrea ariakensis* maintained under laboratory conditions. *J Shellfish Res* 25:65–72
- Murrell A, Kleeman SN, Barker SC, Lester RJG (2002) Synonymy of *Perkinsus olseni* Lester & Davis, 1981 and *Perkinsus atlanticus* Azevedo, 1989 and an update on the phylogenetic position of the genus *Perkinsus*. *Bull Eur Assoc Fish Pathol* 22:258–265
- NRC (National Research Council of the National Academy of Sciences) (2003) *Non-native oysters in Chesapeake Bay*. National Academies Press, Washington, DC
- Nicolas JL, Comps M, Cochenne N (1992) Herpes-like virus infecting Pacific oyster larvae, *Crassostrea gigas*. *Bull Eur Assoc Fish Pathol* 12:11–13
- Orner DM (2005) Chesapeake Bay office integrated research program symposium report, fiscal year 2004. NOAA Chesapeake Bay Office, Annapolis, MD
- Park KI, Choi KS (2001) Spatial distribution of the protozoan parasite *Perkinsus* sp. found in the Manila clams, *Ruditapes philippinarum*, in Korea. *Aquaculture* 203:9–22
- Ray SM (1952) A culture method for the diagnosis of infections with *Dermocystidium marinum* Mackin, Owen and Collier in oysters. *Science* 116:360–361



- Reece KS, Dungan CF (2006) *Perkinsus* sp. infections of marine molluscs. In: Fish health section. Blue book. Suggested procedures for detection and identification of certain finfish and shellfish pathogens, 2005 edn. Chapter 5.2:1–17. Fish Health Section, American Fisheries Society, Bethesda, MD
- Reece KS, Stokes NA (2003) Molecular analysis of a haplosporidian parasite from cultured New Zealand abalone *Haliotis iris*. *Dis Aquat Org* 53:61–66
- Renault T (1998) Infections herpétique chez les invertébrés: détection de virus de type herpes chez les mollusques bivalves marins. *Virologie* 2:401–403
- Renault T, Cochenne N, Le Deuff RM, Chollet B (1994) Herpes-virus infecting Japanese oyster (*Crassostrea gigas*) spat. *Bull Eur Assoc Fish Pathol* 14:64
- Renault T, Le Deuff RM, Chollet B, Cochenne N, Gerard A (2000a) Concomitant herpes-like virus infections in hatchery reared larvae and nursery-cultured spat *Crassostrea gigas* and *Ostrea edulis*. *Dis Aquat Org* 42:173–183
- Renault T, Le Deuff RM, Lipart C, Delsert D (2000b) Development of a PCR procedure for the detection of a herpes-like virus infecting oysters in France. *J Virol Meth* 88:41–50
- Renault T, Lipart C, Arzul I (2001) A herpes-like virus infects a non-ostreid bivalve species: virus replication in *Ruditapes philippinarum* larvae. *Dis Aquat Org* 45:1–7
- Santmartí MM, García-Valero J, Montes JF, Pech A, Durfort M (1995) Seguimiento del protozoo *Perkinsus* sp., en las poblaciones de *Tapes decussatus* y *Tapes semidecussatus* del Delta del Ebro. *Actas V Congr Nac Acuicult* 5: 260–265
- Shaw BL, Battle HI (1957) The gross microscopic anatomy of the digestive tract of the oyster *Crassostrea virginica* (Gmelin). *Can J Zool* 35:325–347
- Sindermann CJ (1990) Principal diseases of marine fish and shellfish, Vol 2. Academic Press, San Diego, CA
- Stokes NA, Burreson EM (1995) A sensitive and specific DNA probe for the oyster pathogen *Haplosporidium nelsoni*. *J Eukaryot Microbiol* 42:350–357
- Swofford DL (2002) PAUP Phylogenetic analysis using parsimony (and other methods), Ver. 4.0b10. Sinauer Associates, Sunderland, MA
- Thompson JD, Higgins DG, Gibson TJ (1994) Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice. *Nucleic Acids Res* 22:4673–4680
- Tsai MF, Kou GH, Liu HC, Liu KF and 5 others (1999) Long-term presence of white spot syndrome virus (WSSV) in cultivated shrimp population without disease outbreaks. *Dis Aquat Org* 38:107–114
- Villalba A, Casas SM, López C, Carballal MJ (2005) Study of perkinsosis in the carpet shell clam *Tapes decussatus* in Galicia (NW Spain). II. Temporal pattern of disease dynamics and association with clam mortality. *Dis Aquat Org* 65:257–267
- Vizoso DB, Ebert D (2004) Within-host dynamics of a microsporidium with horizontal and vertical transmission: *Octosporea bayeri* in *Daphnia magna*. *Parasitol* 28:31–38
- Whyte SK, Cawthorn RJ, McGladdery SE, MacMillan RJ, Montgomery DM (1993) Cross-transmission studies of *Perkinsus karlssoni* (Apicomplexa) from bay scallops *Argopecten irradians* to native Atlantic Canadian shellfish species. *Dis Aquat Org* 17:33–39
- Zhang Q, Allen SK Jr, Reece KS (2005) Genetic variation in wild and hatchery stocks of the Suminoe oyster (*Crassostrea ariakensis*) assessed by PCR-RFLP and microsatellite markers. *Mar Biotechnol* 7:588–599

Editorial responsibility: Catherine Collins,  
Aberdeen, UK

Submitted: February 23, 2007; Accepted: May 23, 2007  
Proofs received from author(s): September 10, 2007