



Alleged Lessepsian foraminifera prove native and suggest Pleistocene range expansions into the Mediterranean Sea

Paolo G. Albano^{1,2,*}, Anna Sabbatini³, Jonathan Lattanzio³, Jan-Filip Päßler², Jan Steger², Quan Hua⁴, Darrell S. Kaufman⁵, Sönke Szidat⁶, Martin Zuschin², Alessandra Negri³

¹Department of Animal Conservation and Public Engagement, Stazione Zoologica Anton Dohrn, 80121 Naples, Italy

²Department of Palaeontology, University of Vienna, 1090 Vienna, Austria

³Department of Life and Environmental Science, Università Politecnica delle Marche, 60122 Ancona, Italy

⁴Australian Nuclear Science and Technology Organisation, Kirrawee DC, New South Wales 2232, Australia

⁵School of Earth and Sustainability, Northern Arizona University, Flagstaff, Arizona 86011, USA

⁶Department of Chemistry, Biochemistry and Pharmaceutical Sciences and Oeschger Centre for Climate Change Research, University of Bern, 3012 Bern, Switzerland

ABSTRACT: Biogeographical patterns are increasingly modified by the human-driven translocation of species, a process that accelerated several centuries ago. Observational datasets, however, rarely range back more than a few decades, implying that a large part of invasion histories went unobserved. Small-sized organisms, like benthic foraminifera, are more likely to have been reported only recently due to their lower detectability compared to larger-sized organisms. Recently detected native species of tropical affinity may have thus been mistaken for non-indigenous species due to the lack of evidence of their occurrence in pre-invasion records. To uncover the unobserved past of the Lessepsian invasion — the entrance of tropical species into the Mediterranean through the Suez Canal — we collected sediment cores on the southern Israeli shelf. We deployed state-of-the-art radiocarbon techniques to date 7 individual foraminiferal tests belonging to 5 alleged non-indigenous species and show that they are centuries to millennia old, thus native. Two additional species previously considered non-indigenous occurred in centennial to millennial-old sediments, suggesting their native status. The evidence of multiple tropical foraminiferal species supposed to be non-indigenous but proved native in the eastern Mediterranean suggests either survival in refugia during the Messinian Salinity Crisis (5.96–5.33 million years) or, more likely, dispersal from the tropical Atlantic and Indo-Pacific during the Pleistocene. In the interglacials of this epoch, higher sea levels may have allowed biological connectivity between the Mediterranean and the Red Sea for shallow-water species, showing that the Isthmus of Suez was possibly a more biologically porous barrier than previously considered.

KEY WORDS: Isthmus of Suez · Lessepsian invasion · Historical biogeography · Connectivity · Foraminifera · Mediterranean Sea · Radiocarbon dating

1. INTRODUCTION

Biogeographical patterns are the outcome of the processes of evolution, dispersal and extinction that acted over geological time scales. Such patterns are

increasingly modified by the human-driven translocation of species around the globe over the past decades to few millennia (Ojaveer et al. 2018). However, observational datasets of most biota are limited to yearly or decadal scales, implying that a large

*Corresponding author: pgalbano@gmail.com

part of the history of biological invasions went unrecorded. Therefore, early invasions may have gone undocumented and the non-indigenous species involved erroneously assumed native. Conversely, poorly detectable native species may have been recorded only in recent times and falsely considered as non-indigenous if biogeographically related to other regions. Both scenarios may obscure natural patterns and hamper the full understanding of the magnitude and rate of invasions, as well as their ecological consequences.

The paleontological record is increasingly becoming a fundamental source of information on unobserved past ecosystem states, enabling the reconstruction of patterns of diversity, dispersal and extinction, and thus uniquely contributing to the process of disentangling natural from human-modified biogeographical patterns (Kidwell & Tomašových 2013, Dietl et al. 2015, Yasuhara et al. 2020). The study of fossil pollen allowed the recognition of early translocations of plants by humans, e.g. by Norse settlers in southern Greenland in the 10th century (Fredskild 1978) or by European settlers in North America in the 16th and 17th centuries (McAndrews 1988) (see Jackson 1997 for a review). The absence from the local fossil record of synanthropic land snail shells found exclusively in urban areas on San Salvador Island (Bahamas) suggested their non-indigenous origin (Yanes 2012). Moreover, fossil pollen data have indicated that alleged non-indigenous plants are instead native in North America (Betancourt et al. 1984), the Galapagos Islands (van Leeuwen et al. 2008, Coffey et al. 2011) and on Easter Island (Flenley et al. 1991). Also in the marine environment, the paleontological record has proved useful in reconstructing the unobserved history of biological invasions (Yasuhara et al. 2012). For example, it has provided evidence of possible early translocations of dinoflagellates (McMinn et al. 1997, Irwin et al. 2003, Amorim & Dale 2006, Marret et al. 2009) and of the timing of the arrival of non-indigenous benthic foraminifera (McGann et al. 2012, Guastella et al. 2021), and has been helpful in quantifying time lags in the detection and identification of the drivers of invasion success of an introduced bivalve (Albano et al. 2018).

The Mediterranean Sea is a biodiversity hotspot (Coll et al. 2010) with an endemism rate of 45% (Costello et al. 2017) shaped by its peculiar geological and paleoclimatic history (Taviani 2002, Sabelli & Taviani 2014). A west-to-east gradient of declining diversity and abundance is apparently related to decreasing nutrient availability and increasing tem-

perature and salinity (Tortonese 1951, Coll et al. 2010). The magnitude of this decline remains uncertain due to the lower sampling effort in the eastern and southern sectors (Morri et al. 2009, Idan et al. 2018, Achilleos et al. 2020, Albano et al. 2020). This gradient is currently being disrupted by biological invasions, particularly through the opening of the Suez Canal in 1869, which broke a long-standing biogeographic barrier between the temperate Northern Atlantic and the Indo-West Pacific realms. This so-called 'Lessepsian invasion' (Por 1978, Galil 2009) has an opposite east-to-west gradient related to the location of the Suez Canal (Galil 2012, Nunes et al. 2014). Additionally, climate-driven native diversity collapses in the eastern Mediterranean Sea are further disrupting natural patterns (Rilov 2016, Albano et al. 2021, Steger et al. 2021).

The low sampling effort in the eastern Mediterranean implies that we know little about what its biodiversity looked like before and during the first century of the Lessepsian invasion, between the mid-19th and mid-20th centuries. Reliable and quantitative data have only become available in the last few decades (e.g. Galil & Lewinsohn 1981, Tom & Galil 1991, Edelist et al. 2011, Guarnieri et al. 2017). This knowledge gap has led to the assumption that each newly recorded species of tropical affinity in the eastern Mediterranean is a non-indigenous one.

Benthic foraminifera are single-celled marine eukaryotes, often with calcareous tests, that are commonly preserved in marine sediments. They are good indicators of environmental health and global change of marine ecosystems (Sabbatini et al. 2014). Hundreds of non-indigenous foraminifera have been reported from the Mediterranean Sea (Zenetos et al. 2008), but their status has rarely been assessed based on a historical baseline. In this study, we report the occurrence of tests of benthic foraminifera previously considered non-indigenous in deep sections of sediment cores collected off the Mediterranean coast of Israel and we thus question their non-indigenous status. Similarly, Stulpinaite et al. (2020) and Meriç et al. (2016, 2018) corrected the non-indigenous status of some foraminifera species based on occurrences in pre-Lessepsian (pre-AD 1869) sediments. In order to assess whether our tests are post-Lessepsian, we deployed state-of-the-art radiocarbon techniques to date them individually and ascertain their exact age. We show that all tests are of Holocene age—thus representing native species—and discuss the consequences of these results for the biogeography of the Mediterranean Sea.

2. MATERIALS AND METHODS

2.1. Core and sample collecting and analysis

Two ~1.2 to 1.5 m long sediment cores were collected at each of 2 sites at ~30 m (SC30, replicates SC30_1 and SC30_3) and ~40 m (SC40, replicates SC40_3 and SC40_4) depth off Ashqelon, on the Mediterranean shelf of southern Israel, with a gravity corer aboard the RV 'Mediterranean Explorer' in 2016 (Table S1, Fig. S1 in Supplement 1 at www.int-res.com/articles/suppl/m700p065_supp1.pdf). From a visual inspection, differences in grain size distribution along the cores were obvious but no sedimentary structures could be recognized. Cores were cut into 1 cm thick slices, which were air dried for at least 96 h.

One core each was used for foraminiferal analysis and sieved with a 150 μm mesh, which is a typical sieve size for the study of benthic foraminifera (e.g. Fontanier et al. 2002, Duros et al. 2011) and balances the need for intercepting small-sized tests with the time to process the samples. Sediments were then dried at 40°C for 48 h. One slice every 5 cm was used for foraminiferal analyses. This resolution is considered sufficient for marine sediments in areas with relatively low sedimentation rates as studied here, which are effectively mixed by physical and biological processes at decimeter scales (e.g. Tomašových et al. 2017, Gallmetzer et al. 2019). This strong degree of time-averaging is also nicely indicated by the presence of shells of *Corbula gibba* of very different ages in the same layers in our cores (see Fig. 1). Samples were split into aliquots containing at least 300 tests, which were subsequently picked, identified and counted. The foraminiferal data were standardized and the occurrence of species reported as non-indigenous in the literature recorded (Table 1).

The replicate core was used to determine the sediment grain size, again each fifth 1 cm thick slice. The fraction finer than 63 μm was analyzed with a Micromeritics SediGraph III 5120, after mixing with 5 ml of 2% sodium pyrophosphate ($\text{Na}_4\text{P}_2\text{O}_7$). Coarser fractions were sieved and weighed. Sediment classification follows Folk (1954).

2.2. Core age models

From each target increment—the top 5 cm of the cores, layers with major changes in sediment grain size and the core bottom (see Table 2)—3 to 5 valves of the native bivalve *C. gibba* with a minimum mass of 1 mg were selected for radiocarbon dating, yield-

ing a total of 29 and 28 valves for cores SC30_1 and SC40_4, respectively. *C. gibba* was the only macrofossil common enough throughout both cores and of sufficient mass to be dated with standard radiocarbon methods. To avoid dating individual bivalves twice, only right valves were used whenever possible. If a sufficient number was available within an increment, right valves to be dated were selected by random drawing; otherwise, all right valves within the increment were used, and randomly drawn, non-matching, left valves were added until a sufficient sample size was achieved. Selected valves were pre-cleaned by manually removing sediment and encrustations visible under a stereomicroscope.

Valves were dated by accelerator mass spectrometry (AMS) using powdered carbonate targets (Bush et al. 2013, Bright et al. 2021), with a typical analytical precision better than 0.6% (1 σ). Mollusk shells were subsampled by gently breaking and then selecting a small fragment. All samples were cleaned by sonicating and rinsing 3 times in deionized distilled water (DDI; 16.3 M Ω -cm). Samples were leached with 2 M HCl, with the extent of leaching dependent on sample mass: samples larger than 1 mg were leached to remove about 30% by mass and samples between ca. 0.5 and 1 mg were leached to remove about 15%. Samples were ultimately rinsed 3 times with DDI water then dried in a 50°C oven overnight. They were then ground to a fine powder using a small clean agate mortar and pestle. Between 0.15 and 0.50 mg of the carbonate powder of each sample was transferred to sterilized (3 h at 500°C) borosilicate glass culture tubes (6 mm outer diameter [OD] \times 50 mm). The sample carbonate was combined with 6 to 7 mg of niobium (Nb Puratronic, -325 mesh, 99.99%) powder using a spatula. The tubes were flushed with N₂ gas and capped with Supelco plastic column caps (1/4" [ca. 6.35 mm] OD) to reduce atmospheric exposure until the powder was pressed into targets. The metal and carbonate mixture was pressed into pre-drilled (0.160" [ca. 4.1 mm] depth) aluminum targets at 400 psi, rotated 90° and pressed again at 400 psi. The targets were sent to the Keck Carbon Cycle AMS Laboratory at the University of California Irvine for ¹⁴C analysis.

Radiocarbon ages were converted to calibrated ages or calendar years (cal. yr BP, i.e. before AD 1950) using the OxCal program v. 4.4.2 (Bronk Ramsey 2009) and the Marine20 data (Heaton et al. 2020) with a constant regional marine reservoir correction (ΔR) of -142 ± 66 yr, which is the weighted mean of 8 published pre-bomb ΔR values from Israel and Lebanon (Table S3).

Table 1. Native range and non-indigenous status in the Mediterranean Sea of the target foraminiferal species

Family	Species	Available data SC30	Native range according to the literature	Non-indigenous status in the Mediterranean Sea
Hauerinidae	<i>Articulina alticostata</i> (Cushman, 1944)	Occurrence and age	Indo-Pacific (Cushman 1944, Loeblich & Tappan 1994, Langer & Lipps 2006, Debenay 2012, Mamo 2016)	Reported from Turkey (Meriç et al. 2004, 2008b) and on this basis considered non-indigenous (Zenetos et al. 2010) but identification questioned by Stulpinaite et al. (2020)
Hauerinidae	<i>Cribromilolinella milletti</i> (Cushman, 1954)	Occurrence and age	Indo-Pacific (Cushman et al. 1954, Yim & He 1988)	A record as non-indigenous species is in the EASIN (European Alien Species Information Network) database without details. Considered not present in the Mediterranean Sea by Stulpinaite et al. (2020)
Hauerinidae	<i>Pseudotriloculina subgranulata</i> (Cushman, 1918)	Occurrence	Indo-Pacific including the Red Sea (Cushman 1918, Hottinger et al. 1993, Debenay & Cabioch 2007), possibly also Caribbean (Amergian 2019)	Reported from Israel (Yanko et al. 1998, Hyams- Kaphzan et al. 2008) and on this basis considered non-indigenous (Zenetos et al. 2012); unpublished records in sediment cores off Israel suggesting native status (Stulpinaite et al. 2020)
Hauerinidae	<i>Quinqueloculina erinacea</i> (Mikhalevich, 1976)	Occurrence	Indo-Pacific (Debenay 2012), Caribbean (Mikhalevich 1976)	New record for the Mediterranean Sea
Hauerinidae	<i>Quinqueloculina mosharrafai</i> (Said 1949)	Occurrence	Red Sea (Said 1949) and the Persian (Arabian) Gulf (Al-Enezi et al. 2020)	Reported from Israel (Hyams-Kaphzan 2014). Considered non-indigenous by Stulpinaite et al. (2020)
Hauerinidae	<i>Milolinella fichteliana</i> (d'Orbigny, 1839)	Occurrence and age	Indian Ocean and the Red Sea (Hottinger et al. 1993, Thissen & Langer 2017)	Reported from Turkey (Meriç et al. 2008a,b). Considered non-indigenous by Stulpinaite et al. (2020)
Spiroloculimidae	<i>Spiroloculina antillarum</i> (d'Origny, 1839)	Occurrence and age	Stulpinaite et al. (2020) recognized 2 morphospecies: the typical one from Cuba (d'Orbigny 1839) to which our specimens belong, and another occurring in the Red Sea (Hottinger et al. 1993) and the Indian Ocean (Thissen & Langer 2017)	Stulpinaite et al. (2020) considered the Caribbean and the Red Sea/Indian Ocean morphospecies non- indigenous and native, respectively, in the Mediterranean Sea. They reported the former from Croatia (Wiesner 1923) and Libya (Martinotti 1920), the latter from Turkey (Meriç et al. 2008a,b) and Israel (Avnaim-Katav 2010)
Spiroloculimidae	<i>Spiroloculina nummiformis</i> (Said 1949)	Occurrence	Indo-Pacific including the Red Sea (Said 1949, Hottinger et al. 1993, Parker & Gischler 2011, Debenay 2012)	Reported from Israel (Hyams-Kaphzan 2000, Hyams- Kaphzan et al. 2008). Considered non-indigenous by Stulpinaite et al. (2020)
Nubeculariidae	<i>Nodophthalmidium antillarum</i> (Cushman, 1922)	Occurrence and age	Indo-Pacific including the Red Sea (Hottinger et al. 1993, Makled & Langer 2011, Langer et al. 2013), Caribbean (Cushman 1922) and north- western Atlantic (Culver & Buzas 1980)	Reported from Turkey (Alavi 1980, Meriç et al. 2012, 2014), Lebanon (Moncharmont Zei 1968) and Israel (Hyams-Kaphzan 2000). Considered non-indigenous by Stulpinaite et al. (2020)
Ophthalmidiidae	<i>Edentostomina cultrata</i> (Brady, 1881)	Occurrence	Indo-Pacific (Brady 1881, Langer & Lipps 2006), New Zealand (Hayward et al. 1999), north-western Atlantic to Caribbean (Culver & Buzas 1980, 1982, Sen Gupta et al. 2009), north-eastern Atlantic (Dorst & Schönfeld 2013)	Recorded from Turkey (Alavi 1980, Oflaz 2006) and on this basis considered non-indigenous (Zenetos et al. 2010); unpublished records in sediment cores off Israel suggesting native status (Stulpinaite et al. 2020)

We quantified the central tendency and spread of the shell age-frequency distributions with the median and the inter-quartile range (IQR), respectively. Finally, the core age model was obtained with Bayesian statistics using the R package rbacon 2.4.2 (Blaauw & Christen 2011) with the parameters as specified in Table S2.

2.3. Radiocarbon dating of foraminifera tests

Seven foraminiferal tests belonging to the allegedly non-indigenous miliolids *Articulina alticostata* (1 test), *Cribromiliolinella milletti* (1 test), *Miliolinella fichteliana* (1 test), *Spiroloculina antillarum* (1 test) and *Nodophthalmidium antillarum* (3 tests) (Figs. S2 & S3, Table 1) were dated individually. Tests were taken from the deepest layers in the cores where these species occurred (suggesting older ages and thus potential native status) and those in good preservation state were selected for dating.

Laboratory procedures followed Gottschalk et al. (2018). Foraminiferal tests were loaded into septum-sealed glass vials and the air was replaced with helium. The samples were weakly pre-leached using 0.01 M hydrochloric acid for 3 min at room temperature and the resultant CO₂ was again replaced with helium. The carbonate material was then dissolved in 85% orthophosphoric acid to form CO₂, which was measured directly in the AMS MIni CARbon DAting System (MICADAS) using the gas ion source at the University of Bern, Switzerland (Szidat et al. 2014, Gottschalk et al. 2018). The samples were measured together with 6 standards each of both IAEA-C1 and IAEA-C2 for background correction, ¹⁴C normalization and δ¹³C fractionation correction using the software Bats (Wacker et al. 2010). Afterwards, a constant contamination of F¹⁴C = 0.30 ± 0.04 with a carbon mass of 0.68 ± 0.10 μg C was applied to correct for the procedural blank, as previously determined for dating foraminifera in this laboratory (Gottschalk et al. 2018). Uncertainties of ¹⁴C ages (see Table 3) include a full propagation of all contributions from AMS measurement, background correction, ¹⁴C normalization and δ¹³C fractionation correction, as well as the correction of the constant contamination, whereof the last contribution was the largest because of the small sizes of the samples (individual foraminiferal tests). Radiocarbon ages and their calibrated ages in cal. yr BP (see the age calibration procedure in Section 2.2) are reported in Table S4 in Supplement 2 at www.int-res.com/articles/suppl/m700p065_supp2.xlsx for the mollusks and in Table 3 for the foraminifera.

3. RESULTS

3.1. Core age and sediment profiles

Cores SC30 and SC40 span millennia and centuries, respectively, as a consequence of markedly different sedimentation rates (0.2 and 2.4 mm yr⁻¹, respectively). The age model of core SC30 suggests a median age of ca. 5500 cal. yr BP at its maximum depth (Fig. 1). Time averaging, which quantifies the temporal mixing of the assemblages and here expressed as inter-quartile range, is several hundred years at most sediment depths and reaches 709 yr at the core bottom (Table 2). The age model of core SC40 suggests a median age at its maximum depth of ca. 600 cal. yr BP. Time averaging is generally limited to a few centuries in the upper 30 cm and to decades only in deeper sections. At 14 cm sediment depth, we detected a layer that contains the oldest shells (1427 cal. yr BP) with the highest time averaging (337 yr) of the core. Although we could not observe sedimentary structures, this strong age inversion and high degree of time averaging suggests transport of older shells from elsewhere (e.g. from shallower sites; at the SC30 station, the surficial death assemblage has a median age of 505 cal. yr BP and the age range is between 127 and 3043 cal. yr BP; Table 2).

Core SC30 comprises 3 zones with different sediment types (Fig. 1): a top layer down to ca. 40 cm sediment depth where sand to muddy sand prevails, a second interval down to ca. 110 cm sediment depth where clay prevails, and the bottom of the core where the sediment is coarser again. This latter sediment change, dated around 2600 cal. yr BP, may be related to a decrease in the Nile River discharge due to the onset of drier conditions in central Africa and consequent reduction in pelitic input relative to coarser sediments from coastal erosion (Hassan et al. 2012). Most of core SC40 is homogeneously muddy throughout; the top 5 cm contain slightly coarser sediment (Fig. 1).

3.2. Foraminiferal test ages

The dated foraminiferal tests span between 731 and 8261 cal. yr BP in calibrated median age (Table 3). With the single exception of a test of *Nodophthalmidium antillarum* (specimen F008) with a 95% confidence interval (CI) calibrated age range marginally overlapping with the opening of the Suez Canal (but a median age of 1111 cal. yr BP), all other 95% CI calibrated ages are much older.




Fig. 1. Core age model (sloped lines, based on radiocarbon dating of *Corbula gibba* shells), age of foraminiferal tests (see Table 3), proportion of mud content (grains < 63 μm) and occurrence of foraminiferal species reported in the literature as non-indigenous in the Mediterranean Sea for 2 cores collected off Ashqelon, southern Mediterranean Israeli shelf, at (A) 30 (core SC30) and (B) 40 m (core SC40) depth. The probability distribution of the calibrated age of *C. gibba* shells is shown in blue violin plots. Calibrated ages (median and 95% CI) of the foraminiferal tests are indicated by red dots and error bars, respectively. cal. yr: calibrated years

Table 2. Summary statistics for the age–frequency distributions of the bivalve *Corbula gibba* downcore. cal. yr: calibrated years

Core	Core slices (cm)	Number of dates	Median age (cal. yr BP)	Minimum median age (cal. yr BP)	Maximum median age (cal. yr BP)	Inter-quartile range (yr)
SC30_1	5	4	505	127	3043	863
SC30_1	25	5	1881	1768	1944	97
SC30_1	45	5	2557	1789	3531	228
SC30_1	83–84	5	3274	3079	5045	116
SC30_1	103–104	5	4451	3281	5054	721
SC30_1	120–121	5	4910	3533	7284	709
SC40_4	1	5	104	34	393	201
SC40_4	14	4	381	170	1427	337
SC40_4	24–27	5	290	122	358	131
SC40_4	36	3	176	123	249	63
SC40_4	49	3	139	96	168	36
SC40_4	86–87	5	147	80	204	70
SC40_4	142–145	3	181	123	359	118

Table 3. Sample details, measured ^{14}C ages with 68% uncertainties and calibrated ^{14}C ages with 95% confidence intervals of foraminiferal tests from sediment cores off Ashqelon, southern Israel

Species	Foraminiferal test code (University of Bern code)	Core	Sediment depth (cm)	Carbon mass ($\mu\text{g C}$)	^{14}C age (yr BP)	Calibrated median age (cal. yr BP)	Calibrated age at 95% CI (cal. yr BP)
<i>Articulina alticostata</i>	F006 (BE-12479.1.1)	SC40_4	35	19	1183 \pm 130	731	464–1047
<i>Cribromiliolinella milletti</i>	F001 (BE-11692.1.1)	SC40_4	20	2	3335 \pm 538	3243	1926–4688
<i>Miliolinella fichteliana</i>	F004 (BE-11695.1.1)	SC30_1	110	30	3823 \pm 88	3784	3460–4103
<i>Spiroloculina antillarum</i>	F010 (BE-12483.1.1)	SC30_1	110	10	7798 \pm 271	8261	7667–8967
<i>Nodophthalmidium antillarum</i>	F007 (BE-12480.1.1)	SC40_4	35	3	1851 \pm 492	1486	504–2685
<i>Nodophthalmidium antillarum</i>	F008 (BE-12481.1.1)	SC40_4	35	2	1462 \pm 567	1111	0–2300
<i>Nodophthalmidium antillarum</i>	F009 (BE-12482.1.1)	SC40_4	35	5	1704 \pm 308	1271	629–1998

Additionally, *Miliolinella fichteliana* and *Spiroloculina antillarum* occur in SC30 at sediment depths greater than 1 m, where the core age model suggests ages between 4000 and 6000 cal. yr BP (Fig. 1). In SC40, *Edentostomina cultrata* occurs at 115 cm sediment depth, where the core age model suggests ages around 400 cal. yr BP (Fig. 1), and *Pseudotriloculina subgranulata* occurs in SC40 down to the core bottom at 145 cm sediment depth (ca. 600 cal. yr BP). Occurrences so deep below the seafloor and in layers dating to well before the opening of the Suez Canal suggest the pre-Lessepsian presence of these species in the Mediterranean Sea.

The non-indigenous *Quinqueloculina erinacea* and *Q. mosharrafaei* occur at 20 and 10 cm sediment depth, respectively, and *S. nummiformis* occurs at 5 cm sediment depth in SC40 (Fig. 1). These depths, however, are too shallow to exclude reworking of young material from the surface and thus we have no evidence that *Q. erinacea*, *Q. mosharrafaei* and *Spiroloculina nummiformis* occurred on the Mediterranean Israeli shelf before the opening of the Suez Canal.

4. DISCUSSION

4.1. Alleged non-indigenous foraminifera are native

The 7 foraminiferal tests we dated were all pre-Lessepsian (i.e. pre-AD 1869), suggesting that *Articulina alticostata*, *Cribromiliolinella milletti*, *Miliolinella fichteliana*, *Spiroloculina antillarum* (Caribbean morphotype, see Stulpinaite et al. 2020) and *Nodophthalmidium antillarum* are native Mediterranean species. Consistently, *M. fichteliana* has been reported from sediment cores collected between Rhodes and Crete, Greece, and dated between 4000 and ca. 25 000 cal. yr BP (Abu-Zied et al. 2008), and *S. antillarum* has been reported from Pleistocene sediments related to the last interglacial in Palma de Mallorca, Balears (Spain) (Mateu 1972), from Holocene sediments in cores collected at the archeological site of Elaiussa Sebaste, Mersin (Turkey) (Melis et al. 2015) and in other Pleistocene to Holocene sediments in Turkey (Meriç et al. 2018) (but see discussion of its taxonomy in Section 4.2).

In our material, almost all foraminiferal tests are older than bivalve shells in the same layers. Such age offsets between different taxa in sediment cores are a widespread pattern in both shallow and deep waters. For example, a multi-phyla dating experiment on bivalves, foraminifera, echinoids, brachyurans and otoliths from Holocene sediment cores collected in the Adriatic Sea showed age offsets of 4000 yr between the benthic foraminifera *Adelosina intricata* and *Elphidium crispum*, and offsets of 1000 to 3000 yr between these foraminifera and the bivalves *Corbula gibba* and *Gouldia minima* (Nawrot et al. 2022). Relative differences in the likelihood of burial may be the primary cause of the age differences between bivalves and foraminifera. Foraminiferal tests can be rapidly incorporated into the sediment by burrowers such as callianassid shrimps and polychaetes that can actively mix the sediment down to more than 1 m (Dworschak & Rodrigues 1997, Pervesler & Hohenegger 2006). When buried, tests are protected from most destructive forces. Re-exhumation by bioturbators thus brings to surface tests that are significantly older than other skeletal remains such as bivalve shells that, due to their larger size, have a lower probability of taking part in these burial–exhumation cycles (Martin et al. 1996). The burial–exhumation dynamic also blurs the timing of production peaks and leads to skeletons of comparable durability being of different age despite their occurrence in the same horizon (Tomašových et al. 2019a,b).

Even if not individually dated, the occurrence of the foraminifera *Pseudotriloculina subgranulata* and *Edentostomina cultrata*, previously considered non-indigenous, at 115 and 145 cm sediment depths in our SC40 core suggests their native status. This result confirms the proposition of Stulpinaite et al. (2020) based on undated sediment cores collected off Israel. Further alleged non-indigenous foraminifera were found in sediments dating back from the middle Pleistocene to the Holocene (Meriç et al. 2016, 2018). The assignment of so many allegedly non-indigenous foraminifera species to native status requires a broader reassessment of our knowledge of this group in the Mediterranean Sea.

4.2. Consequences for marine ecology

Small-sized organisms like foraminifera are particularly prone to under- and misreporting. There is evidence for a variety of taxa that small body size delays their discovery (Blackburn & Gaston 1998), as early naturalists were biased towards collecting and describ-

ing large, conspicuous species (Blackburn & Gaston 1994). Indeed, 70% of the species analyzed here were first reported from the Mediterranean Sea in the last 20 yr, 2 were reported in the second half of the 20th century, and only *S. antillarum* was first reported in 1920. This last species is also an example of the challenges faced when treating very small-bodied taxa whose taxonomy is not settled. Stulpinaite et al. (2020) recognized 2 morphospecies: the *S. antillarum* s.s. and an undescribed species occurring in the eastern Mediterranean and the Red Sea, but their records in the literature are under the same name and not always illustrated, obscuring their geographic distribution and origin.

Our results also highlight how well-dated sedimentary sequences can support ecology. Direct observation of ecosystems is mostly limited to short time scales (years to decades), much shorter than the life span of many species and most habitats (years to decades and centuries to millennia, respectively) (Yasuhara et al. 2012, Kidwell & Tomašových 2013). Multi-decadal time series are limited to very few places (e.g. the North Sea, Schroeder 2005, and the Arctic fjords, Kortsch et al. 2012). Even when sampling and reporting took place early, data may be qualitative and based on serendipitous collecting events rather than on well-designed sampling (Dauvin 2010), and these early efforts may have been strongly biased in favor of large-sized organisms, as reported above. In these conditions, discriminating a natural from an anthropogenic pattern may not be possible. The occurrence of identifiable and individually dated (sub-)fossils in stratigraphic context complements available ecological datasets (e.g. Yasuhara et al. 2012, 2020, Cramer et al. 2017, Tomašových et al. 2019a, Gallmetzer et al. 2019, O’Dea et al. 2020). In this study, we showed that the Holocene record helps in discriminating the native vs. non-indigenous status of species, and may prove valuable in determining the origin of the so-called cryptogenic species, that is, those that are currently not demonstrably native or non-indigenous (Carlton 1996). Such cryptogenic species could have been introduced before any monitoring occurred and thus were not detected as newcomers.

4.3. Consequences for the biogeography of the Mediterranean Sea

We demonstrate that 3 foraminifera species of Indo-Pacific affinity (*A. alticostata*, *C. milletti* and *M. fichteliana*) and 3 species with a broad distribution in the tropical seas of the Caribbean and the Indo-

Pacific province (*P. subgranulata*, *N. antillarum* and *E. cultrata*) that were previously considered non-indigenous in the Mediterranean are instead native. Their Mediterranean distribution is limited to the eastern basin. The type locality of *S. antillarum* is in the Caribbean, but here it is shown to also be native to the Mediterranean, where it has a broad distribution. Additionally, the benthic foraminifer *Amphistegina lobifera* Larsen, 1976, which is currently considered an invasive species in the Mediterranean Sea, occurs both in the Pleistocene (Meriç et al. 2016) and at sediment depths slightly deeper than ours within undated sediment cores from Haifa Bay (Stulpinaite et al. 2020), suggesting a Holocene age and that the alleged non-indigenous status of this species merits further investigation. We consider it unlikely that these species could have occurred in the basin during the Holocene, become extinct and then been re-introduced through the Suez Canal, as the few extinctions of marine species during the Holocene are directly or indirectly ascribed to human action, and foraminifera are neither associated with direct human pressures nor with cascading effects strong enough to drive them to regional extinction (Dulvy et al. 2009).

The occurrence in the Mediterranean Sea of tropical foraminifera with broad global distributions may have 2 alternative explanations. In a first scenario, such distributions are the outcome of vicariance processes caused by the closure of the Tethyan Seaways during the Cenozoic (Harzhauser et al. 2007, Bialik et al. 2019, Straume et al. 2020). Populations of Tethyan species were geographically separated, leading to disjunct occurrences in both the present-day Caribbean and the Indo-Pacific. Some Tethyan species may have survived in the Mediterranean despite its quasi-complete desiccation during the Messinian Salinity Crisis (Hsü et al. 1977, Roveri et al. 2014), later evolving into closely allied species (Basso et al. 1996, Athanasiadis 1999).

An alternative scenario is that species which disappeared from the Mediterranean during the Messinian Salinity Crisis, or evolved later, (re)entered the basin upon normalization of marine conditions. The timing of their range expansion into the Mediterranean may be as early as the Pliocene for pan-tropical foraminifera, but strictly Indo-Pacific taxa may have entered only later, considering the lack of evidence of connections between the Mediterranean and the Red Sea during the Pliocene (Taviani 2002). Indeed, there is geological and evolutionary evidence of such connections in the late Pleistocene. During the Last Interglacial high stand (Marine Isotope Stage [MIS] 5e, ca. 125 000 kyr), the global sea level was 5 to 9 m higher than at present

(Kopp et al. 2009, Dutton & Lambeck 2012, Roveri et al. 2016) and a shallow marine connection was probably in place across the Suez Isthmus enabling faunal exchanges between the Mediterranean and Red Sea (Plaziat et al. 1995). Such connections may have occurred in earlier times too, because populations of the benthic fish *Serranus cabrilla* that occur in both seas diverged approximately 194 000 yr ago at the end of the MIS 7 interglacial (Bos et al. 2020). More occurrences of sea levels higher than today in the Red Sea have been reconstructed for the last 500 kyr (Siddall et al. 2003), suggesting significant opportunities for the dispersal of shallow water species. Such high sea levels suggest that the Isthmus of Suez was probably a more porous biogeographic barrier than previously thought. Still, it would have acted as a strong filter in favor of shallow water euryhaline species (e.g. the intertidal Mediterranean gastropod *Pirenella conica* that now occurs in the northern Red Sea, Plaziat et al. 1995; or the intertidal Indo-Pacific bivalve *Brachidontes pharaonis* that has been recently hypothesized to have occurred in the Mediterranean before the opening of the Suez Canal, Belmaker et al. 2021), or species with particular ability to cross barriers. Fishes (e.g. *Serranus cabrilla*) can be easily perceived as effective dispersers, but foraminifera also have multiple dispersal pathways. Zygotes and embryonic juveniles are easily transported over large distances, with the latter capable of extending numerous filose pseudopodia that enhance flotation and thus the length of pelagic life (Alve 1999, Alve & Goldstein 2002, 2003). Additionally, foraminifera can be resuspended by storms or fish activity and then transported over long distances (Murray 2006). Last, the recently reported ichthyocory — the live passage of organisms through fish digestive tracts — for foraminifera (Guy-Haim et al. 2017) further increases their dispersal potential. An additional challenge for species entering the Mediterranean during interglacials is their survival during glacial times. However, although in the easternmost Mediterranean Sea summer sea surface temperatures during the last glacial maximum were 2°C lower than today, winter temperature anomalies were close to zero (Hayes et al. 2005), suggesting the occurrence of a refugium for thermophilic species. Indeed, survival in winter is one of the main conditions for warm water species to persist in cooler conditions (Amarasekare & Simon 2020).

The permeability of the Suez Isthmus barrier and these favorable climatic conditions may be the most likely scenario to explain the occurrence of tropical species in the Mediterranean and show that the biogeography of the Mediterranean Sea is still far from well understood.

Acknowledgements. This research is part of the project 'Historical ecology of Lessepsian migration' funded by the Austrian Science Fund (FWF) (P28983-B29, PI: P.G.A.). We thank Bella S. Galil for her support during the project, Baki Yokeş and Daniela Basso for providing papers, Maarten Blaauw for assistance with the rbacon package, and Jordon Bright and John Southon for radiocarbon dating the mollusks. Itay Katzman and the crew of the RV 'Mediterranean Explorer' helped throughout the fieldwork. We thank Moriaki Yasuhara, Vlasta Čosović and 1 anonymous reviewer. for their useful comments on the submitted version of the manuscript.

LITERATURE CITED

- Abu-Zied RH, Rohling EJ, Jorissen FJ, Fontanier C, Casford JSL, Cooke S (2008) Benthic foraminiferal response to changes in bottom-water oxygenation and organic carbon flux in the eastern Mediterranean during LGM to Recent times. *Mar Micropaleontol* 67:46–68
- Achilleos K, Jimenez C, Berning B, Petrou A (2020) Bryozoan diversity of Cyprus (eastern Mediterranean Sea): first results from census surveys (2011–2018). *Mediterr Mar Sci* 21:228–237
- Al-Enezi E, Khader S, Balassi E, Frontalini F (2020) Modern benthic foraminiferal diversity: an initial insight into the total foraminiferal diversity along the Kuwait coastal water. *Diversity (Basel)* 12:142
- Alavi SN (1980) Micropalaeontological studies of recent sediments from the Cilicia Basin (N.E. Mediterranean). PhD dissertation, University of London
- Albano PG, Gallmetzer I, Haselmair A, Tomašových A, Stachowitsch M, Zuschin M (2018) Historical ecology of a biological invasion: the interplay of eutrophication and pollution determines time lags in establishment and detection. *Biol Invasions* 20:1417–1430
- Albano PG, Azzarone M, Amati B, Bogi C, Sabelli B, Rilov G (2020) Low diversity or poorly explored? Mesophotic molluscs highlight undersampling in the Eastern Mediterranean. *Biodivers Conserv* 29:4059–4072
- Albano PG, Steger J, Bošnjak M, Dunne B and others (2021) Native biodiversity collapse in the Eastern Mediterranean. *Proc R Soc B* 288:20202469
- Alve E (1999) Colonization of new habitats by benthic foraminifera: a review. *Earth Sci Rev* 46:167–185
- Alve E, Goldstein ST (2002) Resting stage in benthic foraminiferal propagules: a key feature for dispersal? Evidence from two shallow-water species. *J Micropalaeontol* 21:95–96
- Alve E, Goldstein ST (2003) Propagule transport as a key method of dispersal in benthic foraminifera (Protista). *Limnol Oceanogr* 48:2163–2170
- Amarasekare P, Simon MW (2020) Latitudinal directionality in ectotherm invasion success. *Proc R Soc B* 287:20191411
- Amergian KE (2019) Can Florida's springs coast provide a potential refuge for calcifying organisms? Evidence from benthic foraminifera. MSc thesis, University of South Florida, Tampa, FL
- Amorim A, Dale B (2006) Historical cyst record as evidence for the recent introduction of the dinoflagellate *Gymnodinium catenatum* in the north-eastern Atlantic. *Afr J Mar Sci* 28:193–197
- Athanasiadis A (1999) *Mesophyllum macedonis*, nov. sp. (Rhodophyta, Corallinales), a putative Tethyan relic in the North Aegean Sea. *Eur J Phycol* 34:239–252
- Avnaim-Katav S (2010) Chronostratigraphy and Paleo-environment: reconstruction of Quaternary subsurface successions, Southern Haifa Bay. PhD thesis, University of Haifa, Haifa
- Basso D, Fravega P, Vannucci G (1996) Fossil and living corallineans related to the Mediterranean endemic species *Lithophyllum racemus* (Lamarck) Foslie. *Facies* 35:275–292
- Belmaker J, Abelson A, Haddas-Sasson M, Yamaguchi N, Shefer S, Geffen E (2021) Potential pitfalls in the definition of Lessepsian migrants: the case of *Brachidontes*. In: Jawad LA (ed) *The Arabian seas: biodiversity, environmental challenges and conservation measures*. Springer International Publishing, Cham, p 1293–1307
- Betancourt JL, Long A, Donahue DJ, Jull AJT, Zabel TH (1984) Pre-Columbian age for North American *Corispermum* L. (Chenopodiaceae) confirmed by accelerator radiocarbon dating. *Nature* 311:653–655
- Bialik OM, Frank M, Betzler C, Zammit R, Waldmann ND (2019) Two-step closure of the Miocene Indian Ocean Gateway to the Mediterranean. *Sci Rep* 9:8842
- Blaauw M, Christen JA (2011) Flexible paleoclimate age-depth models using an autoregressive gamma process. *Bayesian Anal* 6:457–474
- Blackburn TM, Gaston KJ (1994) Animal body size distributions change as more species are described. *Proc R Soc B* 257:293–297
- Blackburn TM, Gaston KJ (1998) Some methodological issues in macroecology. *Am Nat* 151:68–83
- Bos AR, Ogwang J, Bariche M, Horoiwa M, Megahed M, Ouf A, Yasuda N (2020) Anti-Lessepsian migration rectified: the Comber *Serranus cabrilla* (L. 1758) existed in the Red Sea prior to the Suez Canal opening. *Mar Biol* 167:126
- Brady HB (1881) Notes on some of the Reticularian Rhizopoda of the 'Challenger' Expedition. Part III. *Q J Microsc Sci* 21:31–71
- Bright J, Ebert C, Kosnik MA, Southon JR and others (2021) Comparing direct carbonate and standard graphite ¹⁴C determinations of biogenic carbonates. *Radiocarbon* 63:387–403
- Bronk Ramsey C (2009) Bayesian analysis of radiocarbon dates. *Radiocarbon* 51:337–360
- Bush SL, Santos GM, Xu X, Southon JR, Thiagarajan N, Hines SK, Adkins JF (2013) Simple, rapid, and cost effective: a screening method for ¹⁴C analysis of small carbonate samples. *Radiocarbon* 55:631–640
- Carlton JT (1996) Biological invasions and cryptogenic species. *Ecology* 77:1653–1655
- Coffey EED, Froyd CA, Willis KJ (2011) When is an invasive not an invasive? Macrofossil evidence of doubtful native plant species in the Galápagos Islands. *Ecology* 92:805–812
- Coll M, Piroddi C, Steenbeek J, Kaschner K and others (2010) The biodiversity of the Mediterranean sea: estimates, patterns, and threats. *PLOS ONE* 5:e11842
- Costello MJ, Tsai P, Wong PS, Cheung AKL, Basher Z, Chaudhary C (2017) Marine biogeographic realms and species endemism. *Nat Commun* 8:1057
- Cramer KL, O'Dea A, Carpenter C, Norris RD (2017) A 3000 year record of Caribbean reef urchin communities reveals causes and consequences of long-term decline in *Diadema antillarum*. *Ecography* 41:164–173
- Culver SJ, Buzas MA (1980) Distribution of recent benthic foraminifera off the North American Atlantic coast. *Smithson Contrib Mar Sci* 6

- Culver SJ, Buzas MA (1982) Distribution of recent benthic foraminifera in the Caribbean region. *Smithson Contrib Mar Sci* 14
- Cushman JA (1918) Foraminifera from Murray Island, Australia. In: Vaughan TW (ed) Some shoal-water bottom samples from Murray Island, Australia, and comparisons of them with samples from Florida and the Bahamas. *Papers from the Department of Marine Biology* 9. Carnegie Institution of Washington, Washington, DC, p 289–290
- Cushman JA (1922) Shallow-water Foraminifera of the Tortugas region. Publication No. 311, Carnegie Institution of Washington, Washington, DC
- Cushman JA (1944) The genus *Articulina* and its species. *Cushman Laboratory for Foraminiferal Research Spec Publ* 10:1–21
- Cushman JA, Todd R, Post RJ (1954) Recent Foraminifera of the Marshall Islands, Bikini and nearby atolls, Part II. *Oceanography (Biologic)*. Geological Survey Professional Paper 260-H:319–384
- d'Orbigny AD (1839) Foraminifères. In: de la Sagra R (ed) *Histoire physique, politique et naturelle de l'île de Cuba*. A. Bertrand, Paris, p 1–224
- ✦ Dauvin JC (2010) Benthic time-series observations in North-eastern European Marine Station: is a European label needed? *Mar Pollut Bull* 60:483–488
- Debenay JP (2012) A guide to 1,000 Foraminifera from southwestern Pacific: New Caledonia. IRD Éditions Institut de recherche pour le développement, Marseille
- Debenay JP, Cabioch G (2007) Recent and Quaternary foraminifera collected around New Caledonia. In: Payri CE, Richer de Forges B (eds) *Compendium of marine species from New Caledonia*. Institut de recherche pour le développement, Nouméa, New Caledonia, p 69–94
- ✦ Dietl GP, Kidwell SM, Brenner M, Burney DA, Flessa KW, Jackson ST, Koch PL (2015) Conservation paleobiology: leveraging knowledge of the past to inform conservation and restoration. *Annu Rev Earth Planet Sci* 43:79–103
- ✦ Dorst S, Schönfeld J (2013) Diversity of benthic Foraminifera on the shelf and slope of the NE Atlantic: analysis of datasets. *J Foraminiferal Res* 43:238–254
- Dulvy NK, Pinnegar JK, Reynolds JD (2009) Holocene extinctions in the sea. In: Turvey ST (ed) *Holocene extinctions*. Oxford University Press, Oxford, p 129–150
- ✦ Duros P, Fontanier C, Metzger E, Pusceddu A and others (2011) Live (stained) benthic foraminifera in the Whittard Canyon, Celtic margin (NE Atlantic). *Deep Sea Res I* 58: 128–146
- ✦ Dutton A, Lambeck K (2012) Ice volume and sea level during the last interglacial. *Science* 337:216–219
- ✦ Dworschak PC, Rodrigues SDA (1997) A modern analogue for the trace fossil *Gyrolithes*: burrows of the thalassinidean shrimp *Axianassa australis*. *Lethaia* 30:41–52
- ✦ Edelist D, Sonin O, Golani D, Rilov G, Spanier E (2011) Spatiotemporal patterns of catch and discards of the Israeli Mediterranean trawl fishery in the early 1990s: ecological and conservation perspectives. *Sci Mar* 75:641–652
- ✦ Flenley JR, King ASM, Jackson J, Chew C, Teller JT, Prentice ME (1991) The Late Quaternary vegetational and climatic history of Easter Island. *J Quaternary Sci* 6:85–115
- Folk RL (1954) The distinction between grain size and mineral composition in sedimentary rock nomenclature. *Geology* 62:344–359
- ✦ Fontanier C, Jorissen FJ, Licari L, Alexandre A, Anschutz P, Carbonel P (2002) Live benthic foraminiferal faunas from the Bay of Biscay: faunal density, composition, and microhabitats. *Deep Sea Res I* 49:751–785
- Fredskild B (1978) Palaeobotanical investigations of some peat deposits of Norse age at Quagssiarssuk, South Greenland. *Medd Gronl* 204:1–41
- ✦ Galil BS (2009) Taking stock: inventory of alien species in the Mediterranean sea. *Biol Invasions* 11:359–372
- ✦ Galil BS (2012) Truth and consequences: the bioinvasion of the Mediterranean Sea. *Integr Zool* 7:299–311
- ✦ Galil B, Lewinsohn C (1981) Macrobenthic communities of the Eastern Mediterranean continental shelf. *PSZNI: Mar Ecol* 2:343–352
- ✦ Gallmetzer I, Haselmair A, Tomašových A, Mautner AK and others (2019) Tracing origin and collapse of Holocene benthic baseline communities in the northern Adriatic Sea. *Palaeos* 34:121–145
- ✦ Gottschalk J, Szidat S, Michel E, Mazaud A and others (2018) Radiocarbon measurements of small-size foraminiferal samples with the mini carbon dating system (MICADAS) at the University of Bern: implications for paleoclimate reconstructions. *Radiocarbon* 60:469–491
- ✦ Guarnieri G, Frascchetti S, Bogi C, Galil BS (2017) A hazardous place to live: spatial and temporal patterns of species introduction in a hot spot of biological invasions. *Biol Invasions* 19:2277–2290
- ✦ Guastella R, Marchini A, Caruso A, Evans J and others (2021) Reconstructing bioinvasion dynamics through micropaleontologic analysis highlights the role of temperature change as a driver of alien foraminifera invasion. *Front Mar Sci* 8:675807
- ✦ Guy-Haim T, Hyams-Kaphzan O, Yeruham E, Almogi-Labin A, Carlton JT (2017) A novel marine bioinvasion vector: ichthyochory, live passage through fish. *Limnol Oceanogr Lett* 2:81–90
- ✦ Harzhauser M, Kroh A, Mandic O, Piller WE, Göhlich U, Reuter M, Berning B (2007) Biogeographic responses to geodynamics: a key study all around the Oligo–Miocene Tethyan Seaway. *Zool Anz* 246:241–256
- ✦ Hassan FA, Hamdan MA, Flower RJ, Keatings K (2012) The oxygen and carbon isotopic records in Holocene freshwater mollusc shells from the Faiyum paleolakes, Egypt: their paleoenvironmental and paleoclimatic implications. *Quat Int* 266:175–187
- ✦ Hayes A, Kucera M, Kallel N, Saffi L, Rohling EJ (2005) Glacial Mediterranean sea surface temperatures based on planktonic foraminiferal assemblages. *Quat Sci Rev* 24:999–1016
- Hayward BW, Grenfell HR, Reid CM, Hayward KA (1999) Recent New Zealand shallow-water benthic Foraminifera: taxonomy, ecological distribution, biogeography, and use in paleoenvironmental assessment. *Institute of Geological & Nuclear Sciences, Lower Hutt*
- ✦ Heaton TJ, Köhler P, Butzin M, Bard E and others (2020) Marine20—The marine radiocarbon age calibration curve (0–55,000 cal BP). *Radiocarbon* 62:779–820
- Hottinger L, Halicz E, Reiss Z (1993) Recent Foraminifera from the Gulf of Aquaba, Red Sea. *Dela, Slovenska akademija znanosti in umetnosti, Razred za naravoslovne vede - Opera, Academia Scientiarum et Artium Slovenica, Classis IV, Historia Naturalis* 33:1–179
- ✦ Hsü KJ, Montadert L, Bernoulli D, Cita MB and others (1977) History of the Mediterranean salinity crisis. *Nature* 267: 399–403
- Hyams-Kaphzan O (2000) Benthic foraminifera from the Mediterranean inner shelf (to 40 m) of Israel. PhD dis-

- sertation, Ben-Gurion University of the Negev, Be'er Sheva
- Hyams-Kaphzan O (2014) Characteristics of benthic foraminifera inhabiting rocky reefs in northern Israeli Mediterranean shelf. Geological Survey of Israel Report, GSI/36/2014, Jerusalem
- Hyams-Kaphzan O, Almogi-Labin A, Sivan D, Benjamini C (2008) Benthic foraminifera assemblage change along the southeastern Mediterranean inner shelf due to fall-off of Nile-derived siliciclastics. *Neues Jahrb Geol Paläontol Abh* 248:315–344
- Idan T, Shefer S, Feldstein T, Yahel R, Huchon D, Ilan M (2018) Shedding light on an East-Mediterranean mesophotic sponge ground community and the regional sponge fauna. *Mediterr Mar Sci* 19:84–106
- Irwin A, Hallegraeff GM, McMinn A, Harrison J, Heijnis H (2003) Cyst and radionuclide evidence demonstrate historic *Gymnodinium catenatum* dinoflagellate populations in Manukau and Hokiangia Harbours, New Zealand. *Harmful Algae* 2:61–74
- Jackson ST (1997) Documenting natural and human-caused plant invasions using paleoecological methods. In: Luken JO, Thieret JW (eds) *Assessment and management of plant invasions*. Springer, New York, NY, p 37–55
- Kidwell SM, Tomašových A (2013) Implications of time-averaged death assemblages for ecology and conservation biology. *Annu Rev Ecol Syst* 44:539–563
- Kopp RE, Simons FJ, Mitrovica JX, Maloof AC, Oppenheimer M (2009) Probabilistic assessment of sea level during the last interglacial stage. *Nature* 462:863–867
- Kortsch S, Primicerio R, Beuchel F, Renaud PE, Rodrigues J, Lønne OJ, Gulliksen B (2012) Climate-driven regime shifts in Arctic marine benthos. *Proc Natl Acad Sci USA* 109:14052–14057
- Langer MR, Lipps JH (2006) Assembly and persistence of foraminifera in introduced mangroves on Moorea, French Polynesia. *Micropaleontology* 52:343–355
- Langer MR, Thissen JM, Makled WA, Weinmann AE (2013) The foraminifera from the Bazaruto Archipelago (Mozambique). *Neues Jahrb Geol Paläontol Abh* 267:155–170
- Loeblich AR, Tappan H (1994) Foraminifera of the Sahul shelf and Timor Sea. *Cushman Laboratory for Foraminiferal Research Special Publication* 31:1–661
- Makled WA, Langer MR (2011) Benthic foraminifera from the Chuuk Lagoon Atoll system (Caroline Islands, Pacific Ocean). *Neues Jahrb Geol Paläontol Abh* 259:231–249
- Mamo BL (2016) Benthic Foraminifera from the Capricorn Group, Great Barrier Reef, Australia. *Zootaxa* 4215(1): 1–123
- Marret F, Mudie P, Aksu A, Hiscott RN (2009) A Holocene dinocyst record of a two-step transformation of the Neoeuxinian brackish water lake into the Black Sea. *Quat Int* 197:72–86
- Martin RE, Wehmiller JF, Harris MS, Liddell WD (1996) Comparative taphonomy of bivalves and foraminifera from Holocene tidal flat sediments, Bahia la Choya, Sonora, Mexico (northern Gulf of California): taphonomic grades and temporal resolution. *Paleobiology* 22:80–90
- Martinotti A (1920) Foraminiferi della spiaggia di Tripoli. *Atti Soc Ital Sci Nat Mus Civ Stor Nat Milano* 59: 249–334
- Mateu G (1972) Les foraminifères du Tyrrhénien de la baie de Palma de Majorque et les conditions bioécologiques de l'ancienne mer des Baléares. *Rapp P-V Réun CIESM Méditerran* 20:539–540
- McAndrews JH (1988) Human disturbance of North American forests and grasslands: the fossil pollen record. In: Huntley B, Webb TI (eds) *Vegetation history*. Kluwer, Dordrecht, p 673–697
- McGann M, Grossman EE, Takesue RK, Penttila D, Walsh JP, Corbett R (2012) Arrival and expansion of the invasive foraminifera *Trochammina hadai* Uchio in Padilla Bay, Washington. *Northwest Sci* 86:9–26
- McMinn A, Hallegraeff GM, Thomson P, Jenkinson AV, Heijnis H (1997) Cyst and radionuclide evidence for the recent introduction of the toxic dinoflagellate *Gymnodinium catenatum* into Tasmanian waters. *Mar Ecol Prog Ser* 161:165–172
- Melis R, Bernasconi MP, Colizza E, Rita FD and others (2015) Late Holocene palaeoenvironmental evolution of the northern harbour at the Elaiussa Sebaste archaeological site (south-eastern Turkey): evidence from core ELA6. *Turk J Earth Sci* 24:566–584
- Meriç E, Avşar N, Bergin F (2004) Benthic foraminifera of Eastern Aegean Sea (Turkey): systematics and autoecology. *Turkish Marine Research Foundation, Istanbul*
- Meriç E, Avşar N, Nazik A, Yokeç MB, Dinger F (2008a) A review of benthic foraminifera and ostracodes of the Antalya Coast. *Micropaleontology* 54:199–240
- Meriç E, Avşar N, Yokeç MB (2008b) Some alien foraminifera along the Aegean and southwestern coasts of Turkey. *Micropaleontology* 54:307–349
- Meriç E, Avşar N, Nazik A, Yokeç MB and others (2012) Ilica Bay (Çeşme-Izmir) benthic foraminifer-ostracod assemblages and Pacific Ocean – Red Sea originated foraminifera and abnormal individuals. *Bull Miner Res Explor* 145:62–78
- Meriç E, Avşar N, Yokeç MB, Dinçer F (2014) Atlas of recent benthic foraminifera from Turkey. *Micropaleontology* 60: 211–398
- Meriç E, Yokeç MB, Avşar N, Kiyak NG and others (2016) Did *Amphistegina lobifera* Larsen reach the Mediterranean via the Suez Canal? *Quat Int* 401:91–98
- Meriç E, Yokeç MB, Avşar N, Dinçer F (2018) Did *Spiroculina antillarum*, *Articulina carinata*, *Coscinospira hemphrichii*, *Peneroplis pertusus*, *P. planatus*, *Sorites orbiculus*, *Astacolus insolitus*, *Siphonina tubulosa*, *Amphistegina lessonii* and *A. lobifera* reach the Mediterranean via the Suez Canal? *Int J Environ Geoinform* 5: 378–385
- Mikhalevich VI (1976) New species of shelf foraminifera in the northeastern part of South America. *Zool Zh* 55: 445–449
- Moncharmont Zei M (1968) I foraminiferi di alcuni campioni di fondo prelevati lungo la costa di Beirut (Libano). *Boll Soc Nat Napoli* 77:3–34
- Morri C, Puce S, Bianchi CN, Bitar G, Zibrowius H, Bavestrello G (2009) Hydroids (Cnidaria: Hydrozoa) from the Levant Sea (mainly Lebanon), with emphasis on alien species. *J Mar Biol Assoc UK* 89:49–62
- Murray JW (2006) *Ecology and applications of benthic foraminifera*. Cambridge University Press, Cambridge
- Nawrot R, Berensmeier M, Gallmetzer I, Haselmair A, Tomašových A, Zuschin M (2022) Multiple phyla, one time resolution? Similar time averaging in benthic foraminifera, mollusk, echinoid, crustacean, and otolith fossil assemblages. *Geology* 50:902–906
- Nunes AL, Katsanevakis S, Zenetos A, Cardoso AC (2014) Gateways to alien invasions in the European seas. *Aquat Invasions* 9:133–144

- ✦ O'Dea A, Lepore M, Altieri AH, Chan M and others (2020) Defining variation in pre-human ecosystems can guide conservation: an example from a Caribbean coral reef. *Sci Rep* 10:2922
- Oflaz SA (2006) Taxonomy and distribution of the benthic Foraminifera in the Gulf of Iskenderun, Eastern Mediterranean. MSc thesis, Middle East Technical University, Ankara
- ✦ Ojaveer H, Galil BS, Carlton JT, Alleway H and others (2018) Historical baselines in marine bioinvasions: implications for policy and management. *PLOS ONE* 13: e0202383
- ✦ Parker JH, Gischler E (2011) Modern foraminiferal distribution and diversity in two atolls from the Maldives, Indian Ocean. *Mar Micropaleontol* 78:30–49
- ✦ Pervesler P, Hohenegger J (2006) Orientation of crustacean burrows in the Bay of Panzano (Gulf of Trieste, Northern Adriatic Sea). *Lethaia* 39:173–186
- ✦ Plaziat JC, Baltzer F, Choukri A, Conchon O and others (1995) Quaternary changes in the Egyptian shoreline of the Northwestern Red Sea and Gulf of Suez. *Quat Int* 29–30:11–22
- Por FD (1978) Lessepsian migration: the influx of Red Sea biota into the Mediterranean by way of the Suez Canal. Springer-Verlag, Berlin
- ✦ Rilov G (2016) Multi-species collapses at the warm edge of a warming sea. *Sci Rep* 6:36897
- ✦ Rovere A, Raymo ME, Vacchi M, Lorscheid T and others (2016) The analysis of Last Interglacial (MIS 5e) relative sea-level indicators: reconstructing sea-level in a warmer world. *Earth Sci Rev* 159:404–427
- ✦ Roveri M, Flecker R, Krijgsman W, Lofi J and others (2014) The Messinian Salinity Crisis: past and future of a great challenge for marine sciences. *Mar Geol* 352:25–58
- Sabbatini A, Morigi C, Nardelli MP, Negri A (2014) Foraminifera. In: Goffredo S, Dubinsky Z (eds) *The Mediterranean Sea: its history and present challenges*. Springer Netherlands, Dordrecht, p 237–256
- Sabelli B, Taviani M (2014) The making of the Mediterranean molluscan biodiversity. In: Goffredo S, Dubinsky Z (eds) *The Mediterranean Sea: its history and present challenges*. Springer Netherlands, Dordrecht, p 285–306
- Said R (1949) Foraminifera of the northern Red Sea. *Cushman Laboratory for Foraminiferal Research Spec Publ* 26:1–44
- Schroeder A (2005) Community dynamics and development of soft bottom macrozoobenthos in the German Bight (North Sea) 1969–2000. *Ber Polarforsch Meeresforsch* 494
- Sen Gupta BK, Smith LE, Machain-Castillo ML (2009) Foraminifera of the Gulf of Mexico. In: Felder DL, Camp DK (eds) *Gulf of Mexico—origins, waters, and biota*. Biodiversity. Texas A&M Press, College Station, TX, p 87–129
- ✦ Siddall M, Rohling EJ, Almogi-Labin A, Hemleben C, Meisner D, Schmelzer I, Smeed DA (2003) Sea-level fluctuations during the last glacial cycle. *Nature* 423:853–858
- Steger J, Bošnjak M, Belmaker J, Galil BS, Zuschin M, Albano PG (2021) Non-indigenous molluscs in the Eastern Mediterranean have distinct traits and cannot replace historic ecosystem functioning. *Glob Ecol Biogeogr* 31:89–102
- ✦ Straume EO, Gaina C, Medvedev S, Nisancioglu KH (2020) Northern Hemisphere oceanic gateways. *Gondwana Res* 86:126–143
- Stulpinaite R, Hyams-Kaphzan O, Langer MR (2020) Alien and cryptogenic Foraminifera in the Mediterranean Sea: a revision of taxa as part of the EU 2020 Marine Strategy Framework Directive. *Mediterr Mar Sci* 21:719–758
- ✦ Szidat S, Salazar GA, Vogel E, Battaglia M, Wacker L, Synal HA, Türler A (2014) ¹⁴C analysis and sample preparation at the new Bern laboratory for the analysis of radiocarbon with AMS (LARA). *Radiocarbon* 56:561–566
- Taviani M (2002) The Mediterranean benthos from late Miocene up to present: ten million years of dramatic climatic and geologic vicissitudes. *Biol Mar Mediterr* 9: 445–463
- ✦ Thissen JM, Langer MR (2017) Spatial patterns and structural composition of foraminiferal assemblages from the Zanzibar Archipelago (Tanzania). *Palaeontographica, Abt A: Palaeozoology – Stratigraphy* 308:1–67
- ✦ Tom M, Galil B (1991) The macrobenthic associations of Haifa Bay, Mediterranean coast of Israel. *PSZNI: Mar Ecol* 12:75–86
- ✦ Tomašových A, Gallmetzer I, Haselmair A, Kaufman DS, Vidović J, Zuschin M (2017) Stratigraphic unmixing reveals repeated hypoxia events over the past 500 yr in the northern Adriatic Sea. *Geology* 45:363–366
- ✦ Tomašových A, Gallmetzer I, Haselmair A, Kaufman DS, Mavrič B, Zuschin M (2019a) A decline in molluscan carbonate production driven by the loss of vegetated habitats encoded in the Holocene sedimentary record of the Gulf of Trieste. *Sedimentology* 66:781–807
- ✦ Tomašových A, Kidwell SM, Alexander CR, Kaufman DS (2019b) Millennial-scale age offsets within fossil assemblages: result of bioturbation below the taphonomic active zone and out-of-phase production. *Paleoceanogr Paleoclimatol* 34:954–977
- Tortonese E (1951) I caratteri biologici del Mediterraneo orientale e i problemi relativi. *Attual Zool* 7:207–251
- ✦ van Leeuwen JFN, Froyd CA, van der Knaap WO, Coffey EE, Tye A, Willis KJ (2008) Fossil pollen as a guide to conservation in the Galápagos. *Science* 322:1206
- ✦ Wacker L, Christl M, Synal HA (2010) Bats: a new tool for AMS data reduction. *Nucl Instrum Methods Phys Res B* 268:976–979
- Wiesner H (1923) *Die Miliolideen der östlichen Adria*. Prag-Bubeneč
- ✦ Yanes Y (2012) Anthropogenic effect recorded in the live-dead compositional fidelity of land snail assemblages from San Salvador Island, Bahamas. *Biodivers Conserv* 21:3445–3466
- Yanko V, Ahmad M, Kaminski M (1998) Morphological deformities of benthic foraminiferal tests in response to pollution by heavy metals: implications for pollution monitoring. *J Foraminiferal Res* 28:177–200
- ✦ Yasuhara M, Hunt G, Breitburg D, Tsujimoto A, Katsuki K (2012) Human-induced marine ecological degradation: micropaleontological perspectives. *Ecol Evol* 2:3242–3268
- ✦ Yasuhara M, Huang HH, Hull P, Rillo M and others (2020) Time machine biology: cross-timescale integration of ecology, evolution, and oceanography. *Oceanography* 33:16–28
- Yim WWS, He XX (1988) Holocene foraminifera in Hong Kong and their palaeoenvironmental significance. In: Whyte P, Aigner JS, Jablonski NG, Taylor G, Walker D, Pinxian W, So CL (eds) *The palaeoenvironment of East Asia from the Mid-Tertiary*. Proc 2nd Int Conf Paleoenvi-

ronment of East Asia, Vol. II: Oceanography, Palaeozoology and Palaeoanthropology. Centre of Asian Studies, University of Hong Kong, p 787–809

- ✦ Zenetos A, Meric E, Verlaque M, Galli P and others (2008) Additions to the annotated list of marine alien biota in the Mediterranean with special emphasis on foraminifera and parasites. *Mediterr Mar Sci* 9:119–166
- ✦ Zenetos A, Gofas S, Verlaque M, Cinar ME and others (2010) Alien species in the Mediterranean Sea by 2010.

A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. *Mediterr Mar Sci* 11:381

- ✦ Zenetos A, Gofas S, Morri C, Rosso A and others (2012) Alien species in the Mediterranean Sea by 2012. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part 2. Introduction trends and pathways. *Mediterr Mar Sci* 13:328–352

*Editorial responsibility: Toshi Nagata,
Kashiwanoha, Japan*
*Reviewed by: M. Yasuhara, V. Čosović and
1 anonymous referee*

Submitted: November 27, 2021
Accepted: September 14, 2022
Proofs received from author(s): October 27, 2022