

The rare subgroup C1 of *Morphysa* (Polychaeta, Eunicidae): re-description of species and first records in the Mediterranean Sea

NIKOLAOS KATSIARAS^{1,2,3}, NOMIKI SIMBOURA¹ & DROSOS KOUTSOUBAS²

¹Institute of Oceanography, Hellenic Centre for Marine Research, 46.7km Athinon-Souniou Ave, Anavissos 19013 Greece

²Department of Marine Sciences, University of the Aegean, University Hill, Mytilene 81100 Greece.

³Corresponding author. E-mail: nkatsiaras@hcmr.gr

Abstract

Morphysa is one of the largest genera in the family Eunicidae, containing 76 valid species. However, among them there are only five species that share the diagnostic characters of having only composite falcigers in inferior setae and branchiae confined in the anterior body. These species are relatively rare and less described while in their original descriptions their differentiation is mainly based on features believed to be age-specific today. In this study, the type-materials of *M. adenensis*, *M. purcellana* and *M. gemmata* have been re-examined and re-described with illustrations. For *M. adenensis*, new material from the Mediterranean Sea was also examined, where this group has not been recorded up to date. The above species were compared to each other and new differential characters are being highlighted. A revised key for the *Morphysa* species found in the Mediterranean is also provided.

Key words: *Morphysa adenensis*, *Morphysa purcellana*, *Morphysa gemmata*

Introduction

The family of Eunicidae Berthold, 1827 is one of the largest polychaete families (Fauchald 1970). To date, the genus *Morphysa* Quatrefages, 1865 alone contains 76 valid species (Fauchald & Bellan 2013) and there are several being recently described. The diagnostic features of the genus that were traditionally considered most important can be summarized as having five prostomial appendages, but no peristomial cirri. According to a wide review of the genus by Fauchald (1970), all species can be grouped based on the type of composite setae present and then subdivided by the distribution of the branchiae. Therefore, four major groups were distinguished: Group A that includes species with no composite setae present, Group B with only composite spinigers present, Group C with only composite falcigers present and finally, Group D with both composite falcigers and composite spinigers present. Each of these groups can be further divided into species having branchiae on a limited anterior region (Subdivision 1, e.g. Group C1), or over a long region of the body (Subdivision 2).

However, recent studies have proved that the presence of the different types of composite setae along the body can vary to some extent in the same species with age. For example, Lu & Fauchald (1998) observed that in *Morphysa bellii oculata* Treadwell, 1921 (which belongs to group D), the presence of composite falcigers range from very abundant all along the body in young stages, to limited at only posterior setigers in old stages. Nevertheless, morphology of setae varies along the eunicid body in a structured way (Zanol *et al.* 2014; Fauchald 1992) and therefore, apart from presence/absence of specific types, distribution patterns in respect of age stages should be considered in diagnosis.

The species of *Morphysa* recorded in the Mediterranean Sea to date are *Morphysa bellii* (Audouin & Milne Edwards, 1833), *Morphysa fallax* Marion & Bobretzky, 1875 and *Morphysa cinari* Sahin, 2014 that belong to Group D; *Morphysa kinbergi* McIntosh, 1910 and *Morphysa sanguinea* Montagu, 1815 that belong to Group B. Past records of *Morphysa disjuncta* Hartman, 1961 in the Mediterranean Sea are considered doubtful (Sahin 2014).

Recent sampling surveys (2009–2013) included in this work, at the Eastern Mediterranean seagrass infauna,

(1995) describes pectinate setae as having marginal teeth longer than inner in *M. conferta* and Gathof (1984) has illustrated it in this way for *M. sp. A*, but no information is given about the distributional patterns, so it is not absolutely clear if also the rest of C1 group species should remain under *Marphysa*.

The detailed re-examination and re-description of setal distribution for other *Marphysa* species is being recently encouraged (e.g. Hutchings & Karageorgopoulos, 2003; Hutchings *et al.* 2012). Both studies contributed, for example, to redefine and doubt the worldwide distribution of *M. sanguinea*. The Mediterranean records, in this study, stretch the geographical distribution of *M. adenensis* from Andaman Sea to Eastern Mediterranean (Fig. 10), but re-examination of other records may also lead to distribution redefinition. Moreover, if additional data (e.g. molecular) were available, distinction between Mediterranean material and specimens from the other ocean basins could be possible, as this was the case in studies that erected sibling species among morphologically identical populations and reduced the distribution of species formerly known as cosmopolitan (e.g. Knowlton 1993; Westheide & Schmidt 2003). When studying on species of Group C1 that are found so rarely, the basis of the conclusions are inevitably limited to phenotypic distinctions among few type-materials.

The relatively rare records of *M. adenensis* in a global scale, combined with the fact that the Mediterranean habitats were *P. oceanica* shoots (a micro-habitat which is much less investigated in the Eastern Mediterranean) make it probable to be a native, but overlooked species. In addition, the *P. oceanica* habitats that it has been found the past 4 years are very distant to each other. On the other hand, the Indo-pacific specimens were collected from a range of habitats that includes intertidal muddy substrata, which has been studied thoroughly in Eastern Mediterranean. Therefore, a recent introduction from the Indo-Pacific could also be speculated. Since clear evidence is absent, *M. adenensis* is reported as a cryptogenic species in the Mediterranean Sea at this stage.

The presence of *M. adenensis* in the Mediterranean Sea raises the number of *Marphysa* species of the area up to six. An updated artificial (**not phylogenetic**) key for the Mediterranean *Marphysa* spp. is provided:

1. Acicular setae bidentate. Composite falcigers only present or composite falcigers with composite spinigers present in anterior segments 2
- Acicular setae unidentate. Composite spinigers only present or composite spinigers with rare composite falcigers in far posterior segments 4
2. Composite spinigers and composite falcigers present 3
- Composite falcigers only present. Blade-length gradation up to more than twice as long in anterior segments. 1–2 pectinate setae of 4–8 teeth, present throughout. Acicular setae with vertical angle between teeth *M. adenensis*
3. Branchiae pectinate *M. bellii*
- Branchiae with 1–2 filaments *M. fallax*
4. Branchiae confined only in the anterior part of the body 5
- Branchiae distributed up to the posterior segments *M. sanguinea*
5. Composite spinigers only present. Acicular setae without terminal hood *M. kinbergi*
- Composite spinigers and rare composite falcigers in far posterior segments present. Acicular setae with terminal hood *M. cinari*

Acknowledgments

We would like to thank Emma Sherlock (Natural History Museum, London) for loaning the holotype of *M. gemmata* and the type of *M. purcellana*, also Dr. Terik Meziane (Museum National d' Histoire Naturelle, Paris) for loaning the holotype of *M. adenensis* and the anonymous referees for their constructive comments on the manuscript. We are also grateful to our colleagues at University of the Aegean, Andreas Alifragkis and Zoi Kokkini for their valuable help in collecting the material. Finally, we are greatful of Haralampos Sampalis for his aid in photo editing. Some samples were collected during the Water Framework Directive National Monitoring Project of Greece, financed by the European Union Funds through the National Strategic Reference Framework (NSRF).

References

- Audouin, J.V. & Milne Edwards, H. (1833) Classification des Annélides et description de celles qui habitent les côtes de la France. *Annales des sciences naturelles, Paris*, Series 1, 29, 195–269.
 Aungtonya, C., Thaipal, S. & Bussarawit, S. (2002) A list of Polychaetes (Annelida) in the reference collection database of the

- phuket marine biological center, Thailand. *Phuket Marine Biological Center Special Publication*, 24, 21–32.
- Bigot, L., Gremare, A., Amoroux, J.M., Frouin, P., Maire, O. & Gaertner, J.C. (2008) Assessment of the ecological quality status of soft-bottoms in Reunion Island (tropical southwest Indian Ocean) using AZTI marine biotic indices. *Marine Pollution Bulletin*, 56, 704–722.
<http://dx.doi.org/10.1016/j.marpolbul.2007.12.020>
- Carrera-Parra, L.F. & Salazar-Vallejo, S.I. (1998) A new genus and 12 new species of Eunicidae (Polychaeta) from the Caribbean Sea. *Journal of the Marine Biology Association of the UK*, 78, 145–182.
<http://dx.doi.org/10.1017/s0025315400040005>
- Day, J.H. (1953) The polychaetes fauna of South Africa. Part 2. Errant Species from Cope Shores and Estuaries. *Annals of the Natal Museum*, XII (3), 397–441.
- Day, J.H. (1960) The polychaete fauna of South Africa. Part 5: Errant species dredged off Cape coasts. *Annals of the South African Museum*, 45, 261–373.
- Day, J.H. (1962) Polychaeta from several localities in the Western Indian Ocean *Proceeding of the Zoological Society of London*, 139 (4), 627–656.
<http://dx.doi.org/10.1111/j.1469-7998.1962.tb01597.x>
- Day, J.H. (1963a) The polychaete fauna of South Africa. Part 8: New species and records from grab samples and dredgings. *Bulletin of the British Museum of Natural History Zoology*, 10 (7), 383–445.
- Day, J.H. (1967) *A monograph on the polychaeta of southern Africa. Part 1. Errantia*. British Museum (Natural History), London, 458 pp.
- Fauchald, K. (1970) Polychaetous annelids of the families Eunicidae, Lumbrineridae, Iphitimidae, Arabellidae, Lysaretidae and Dorvillieidae from Western Mexico. *Allan Hancock Monographs in Marine Biology*, 5, 1–335.
- Fauchald, K. (1992) A review of the genus Eunice (Polychaeta: Eunicidae) based upon type material. *Smithsonian Contributions to Zoology*, 523, 1–422.
<http://dx.doi.org/10.5479/si.00810282.523>
- Fauchald, K. & Bellan, G. (2013) *Marphysa Quatrafages*, 1865. In: Read, Q. & Fauchald, K. (Eds.), *World Polychaeta Database*. Available from: <http://www.marinespecies.org/aphia.php?p=taxdetails&id=129281> (accessed 2 February 2014)
- Gathof, J.M. (1984) Chapter 40 Eunicidae. In: Uebelacker, J.M. & Johnson, P.G. (Eds.), *Taxonomic guide to the polychaetes of the northern Gulf of Mexico. Vol. 6. Final Report to the Minerals Management Service, contract 14-12-001-29091*. Barry A. Vittor & Assoc. Inc. Mobile, Alabama, pp. 1–31. [pp. 40.1–40.30]
- Gravier, C. (1900) Contribution à l'étude des Annelides Polychaetes de la Mer Rouge. Première partie. *Nouvelles Archives du Museum d'Historie Naturelle Paris*, Series 4, 2 (2), 137–282.
- Hartman, O. (1961) Polychaetous annelids from California. *Allan Hancock Pacific Expeditions*, 25, 1–226.
- Hilbig, B. (1995) Family Eunicidae Savigny, 1818. In: Blake, J.A., Hilbig, B., Scott, P.H. (Eds.), *Taxonomic Atlas of the Benthic Fauna of the Santa Maria Basin and Western Santa Barbara Channel. 5 - The Annelida Part 2. Polychaeta: Phyllodocida (Syllidae and scale-bearing families), Amphinomida, and Eunicida*. Santa Barbara Museum of Natural History, Santa Barbara, pp. 263–278.
- Hutchings, P.A. & Karageorgopoulos, P. (2003) Designation of a neotype of *Marphysa sanguinea* (Montagu, 1813) and a description of a new species of Marphysa from Eastern Australia. *Hydrobiologia*, 1–3, 87–94.
<http://dx.doi.org/10.1023/a:1026124310552>
- Hutchings, P.A., Glasby, C.J. & Wijnhoven, S. (2012) Note on additional diagnostic characters of *Marphysa sanguinea* (Montagu, 1813) (Annelida: Eunicida: Eunicidae), a recently introduced species in Netherlands. *Aquatic Invasions*, 7 (2), 277–282.
<http://dx.doi.org/10.3391/ai.2012.7.2.014>
- Knowlton, N. (1993) Sibling species in the sea. *Annual Review of Ecology and Systematics*, 24, 189–216.
<http://dx.doi.org/10.1146/annurev.ecolsys.24.1.189>
- Lu, H. & Fauchald, K. (1998) *Mapphysa beilli* (Polychaeta: Eunicidae) and two related species, *M. oculata* and *M. totospina*, a new species, with notes on size-dependant features. *Proceeding of the biological society of Washington*, 111 (4), 829–842.
- Mak, P.M.S. (1980) The coral associated polychaetes of Hong Kong, with special references to the serpulids. In: Morton, B. & Tseng, C.K. (Eds.), *Proceedings of the First International Marine Biological Workshop: The Marine Flora and Fauna of Hong Kong and Southern China*. Hong Kong. Hong Kong University Press, Hong Kong, pp. 595–617.
- Marion, A.F. & Bobretzky, N. (1875) Étude des Annélides du golfe de Marseille. *Annales des sciences naturelles, Paris*, Series 6, 2, 1–106.
- McIntosh, W.C. (1910) A monograph of the British annelids. Polychaeta. Syllidae to Ariciidae. *Ray Society of London*, 2 (2), 233–524.
- Mohammad, M.B.M. (1973) New species and records of polychaete annelids from Kuwait, Arabian Gulf. *Zoological Journal of the Linnean Society*, 52, 23–44.
<http://dx.doi.org/10.1111/j.1096-3642.1973.tb01876.x>
- Montagu, G. (1815) Descriptions of several new or rare animals, principally marine, discovered on the South coast of Devonshire. *Transactions of the Linnean Society of London*, 11, 1–26.
<http://dx.doi.org/10.1111/j.1096-3642.1813.tb00035.x>
- Moore, J.P. (1911) The polychaetous annelids dredged by the U.S.S. 'Albatross' off the coast of southern California in 1904. III.

- Euphosynidae to Goniadidae. *Proceedings of the Academy of Natural Sciences, Philadelphia*, 63, 234–318.
<http://dx.doi.org/10.5962/bhl.title.12425>
- Sahin, K.G. (2014) *Morphysa cinari*, a new species of Eunicidae from the coasts of Turkey (eastern Mediterranean) and re-descriptions of *Morphysa kinbergi* McIntosh, 1910 and *Morphysa disjuncta* 1961. *Journal of Natural History*, 48 (33–34), 1989–2006.
<http://dx.doi.org/10.1080/00222933.2014.905125>
- Steiner, T.M. & Amaral, A.C.Z. (2000) Two new species of Morphysa Quatrefages 1865 (Eunicidae, Polychaeta) from the intertidal sandy beaches of the São Sebastião Channel, State of São Paulo (Brazil). *Bulletin of Marine Science*, 67 (1), 479–489.
- Treadwell, A.L. (1921) Leodicidae of the West Indian region. *Carnegie Institute of Washington Publication*, 15 (293), 1–131.
- Westheide, W. & Schmidt, H. (2003) Cosmopolitan versus cryptic meiofaunal polychaete species: an approach to a molecular taxonomy. *Helgoland Marine Research*, 57, 1–6.
<http://dx.doi.org/10.1007/s10152-002-0114-2>
- Willey, A. (1904) Littoral Polychaeta from the Cape of Good Hope. *Transactions of the Linnean Society of London, Series 2 (Zoology)*, 9 (6), 255–268
<http://dx.doi.org/10.1111/j.1096-3642.1904.tb00450.x>
- Zanol, J., Halanych, K.M. & Fauchald, K. (2014) Reconciling taxonomy and phylogeny in the bristleworm family Eunicidae (polychaeta, Annelida). *Zoologica Scripta*, 43, 79–100.
<http://dx.doi.org/10.1111.zsc.12034>