

## Aceric acid hydrolase 127A, *Bacteroides thetaiotaomicron* *BtAch127A* (GH127)

### Catalogue number:


CZ08031, 1 mg  
CZ08032, 3 × 1 mg

### Description

Aceric acid hydrolase 127A (*BtAch127A*), assigned the E.C. number 3.2.1.-, is a derivative of *Bacteroides thetaiotaomicron*. It is a 3-C-carboxy-5-deoxy-L-xylose (aceric acid) hydrolase. The recombinant *BtAch127A*, purified from *Escherichia coli*, is a single-domain Glycoside Hydrolase family 127 (GH127) enzyme (see more details at [www.cazy.org](http://www.cazy.org)). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl<sub>2</sub>, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

### Electrophoretic Purity

The molecular integrity and purity of *BtAch127A* (GH127) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).



**Figure 1.** SDS-PAGE analysis of *BtAch127A* (GH127) was conducted in Lane 1, employing a 14% polyacrylamide gel. The enzyme exhibits a singular band corresponding to a molecular weight of approximately 79.80 kDa. Lane M contains a Protein Marker for reference.

### Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

### Substrate specificity

*BtAch127A* (GH127) hydrolyses the L-Acef- $\alpha$ 1,3-L-Rhap linkage in rhamnogalacturonan II (RGII).

### Temperature and pH optima

The enzyme exhibits optimal activity within a pH range of 6.5-7.5 and at a temperature of 37°C. Maximal enzymatic activity is achieved at pH 7 and a consistent temperature of 37°C.

### Enzyme activity

The substrate specificity and kinetic properties of *BtAch127A* (GH127) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

### Reference

Ndeh *et al.* (2017) *Nature*. 544(7648):65-70.

Mahowald *et al.* (2009) *Proc Natl Acad Sci U S A*. 106(14):5859-5864.

### Customer Support

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at [info@nzytech.com](mailto:info@nzytech.com).

### Quality control assay

Protein purity is determined to be  $\geq 90\%$ , as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

V2301

For research use only