

Research Article

Biological introductions or native ranges: two curious cases of new distributional records in the Panama Canal

Sebastian Kvist^{1,2,*}, Danielle de Carle^{1,2}, Aydeé Cornejo³ and Alejandro Oceguera-Figueroa⁴

¹Department of Natural History, Royal Ontario Museum, 100 Queen's Park, Toronto, Ontario M5S 2C6, Canada

²Department of Ecology and Evolutionary Biology, University of Toronto, 25 Willcocks Street, Toronto, Ontario, M5S 2B4, Canada

³Colección Zoológica Dr. Eustorgio Méndez, Instituto Conmemorativo Gorgas de Estudios de la Salud, Calle 36 Este, Panamá, Panamá

⁴Laboratorio de Helmintología, Instituto de Biología, Universidad Nacional Autónoma de México. Ciudad Universitaria, Coyoacán, A.P. 70-153. Ciudad de México. C.P. 14510, Mexico

*Corresponding author

E-mail: sebastian.kvist@utoronto.ca

Received: 14 December 2017 / Accepted: 10 June 2018 / Published online: 3 July 2018

Handling editor: Andrew David

Abstract

We report two new, curious records of freshwater worms collected in the Panama Canal, Panama: the ribbon worm, *Apatronemertes albimaculosa* Wilfert and Gibson, 1974, and the leech, *Placobdella parasitica* (Say, 1824). Identification of these specimens relied on morphological examinations and, more heavily, on molecular barcoding using the mitochondrial COI locus. Both phylogenetic and genetic distance analyses were employed to robustly establish the identities of the specimens. Whereas the native range of *A. albimaculosa* is currently unknown, *P. parasitica* was first described from Minnesota, USA and is possibly native to southern Canada and most of the USA. Neither species has previously been recorded from Central America. We briefly discuss possible scenarios for the dispersal of both species, and conclude that the introduction of *P. parasitica* into Panamanian waters was likely human-mediated in association with the introduction of its turtle host. Insofar as ours is the first record of wild-caught *A. albimaculosa* worldwide, we do not exclude the possibility of Panama being within the native range of the species. However, it is also possible that the species has been transported with ballast water into the Panama Canal; a relatively common phenomenon.

Key words: introduced species, Nemertea, ribbon worms, Hirudinea, leech, phylogeny, DNA barcoding

Introduction

Linking the Atlantic Ocean to the Pacific Ocean, the Panama Canal is a major center for shipping activity, and one of the world's most heavily trafficked aquatic throughways (Mason and Rowlands 1938; Carse 2012). Every year, over 14,000 vessels traverse the narrow channel which stretches from Rio Chagres on the Caribbean slope in the north to Rio Grande on the Pacific slope in the south, effectively crossing the Continental Divide (Ruiz et al. 2006). This tremendous movement across the canal, coupled with the evacuation and loading of ballast water, has led to several cases of biological introductions to the Panama Canal, as well as the dispersal of organisms between the Atlantic and Pacific Oceans (Hanna 1966; Aron and Smith 1971; Jones 1972; Wasson et

al. 2001; Wonham et al. 2001; Smith et al. 2004; Cohen 2006; Ruiz et al. 2009; Schlöder et al. 2013). For example, the Harris mud crab *Rhithropanopeus harrisii* (Gould, 1841) native to the eastern coastal ranges of North America, currently seems to be well-established throughout several parts of the Panama Canal (Roche and Torchin 2007). Parasitic taxa, such as the Asian fish tapeworm *Schyzocotyle acheilognathi* (Yamaguti, 1934), have also been inadvertently introduced along with their hosts (Choudhury et al. 2013).

Nemertea (ribbon worms) is a predominantly marine taxon with only 22 out of the 1,280 species inhabiting freshwater (Gibson and Moore 1976; Kajihara et al. 2008; Sundberg and Gibson 2008; Strand and Sundberg 2015). The freshwater species *Apatronemertes albimaculosa* Wilfert and Gibson, 1974

(Nemertea: Heteronemertea: Lineidae) was originally described from aquarium plants (*Vallisneria spiralis* Linnaeus, 1758) in Düsseldorf, Germany (Wilfert and Gibson 1974), and was, by all accounts, artificially introduced from an unknown, but likely tropical location, together with plant material (Kajihara et al. 2016). Since then, the species has been reported from Austria, Spain, USA, and Japan (Senz 1993; Kajihara et al. 2016). In all cases, specimens were found in freshwater aquaria, suggesting that this species hitchhikes with plant material sold in the aquarium trade. Indeed, *A. albimaculosa* has been recorded in association with a wide variety of plants; a detailed list of potential carriers can be found in Kajihara et al. (2016). As a corollary, a different species of freshwater nemertean, *Prostoma graecense* (Böhming, 1892), has also been recorded from disjointed locations in Europe, Oceania and North America. Its odd distribution is a result of recent, accidental introduction by human activities (Lagunas-Calvo et al. 2018).

Equally prone to hitchhiking, the freshwater leech *Placobdella parasitica* (Say, 1824) (Annelida: Rhynchobdellida: Glossiphoniidae) is ectoparasitic on a wide range of freshwater turtles (Davy et al. 2009). The species feeds exclusively on blood from its hosts, and its known geographic range stretches from southern Canada in the north, to Texas, USA in the south (Klemm 1985; Sawyer 1986; Dodd Jr. 1988; Davy et al. 2009). According to historical and contemporary accounts (Sawyer 1986; Dodd Jr. 1988; Brooks et al. 1990; Ryan and Lambert 2005; Davy et al. 2009), *P. parasitica* is chiefly encountered attached to its hosts, suggesting that the host may limit its dispersal capabilities. *Placobdella parasitica* has been documented from a rather wide host range, including Blanding's turtle *Emydoidea blandingii* (Holbrook, 1838); northern map turtle *Graptemys geographica* (Lesueur, 1817); snapping turtle *Chelydra serpentina* (Linnaeus, 1758); wood turtle *Glyptemis insculpta* (Le Conte, 1830); painted turtle *Chrysemys picta* (Schneider, 1783); spotted turtle *Clemmys guttata* (Schneider, 1792); stinkpot *Sternotherus odoratus* (Latreille in Sonnini and Latreille, 1801); bog turtle *Glyptemys muhlenbergii* (Schoepff, 1801); eastern mud turtle *Kinosternon subrubrum* (Lacépède, 1788); pond slider *Trachemys scripta* (Thunberg in Schoepff, 1792); and spiny softshell turtle *Apalone spinifera* (Lesueur, 1827) (Davy et al. 2009 and references therein). Furthermore, despite robust collection efforts, *Placobdella parasitica* has never been recorded from Mexico or Central America (Cornejo et al. 2015; Oceguera-Figueroa and Pacheco-Chaves 2012; Tessler et al. 2018).

In the present study, we report the first records of the nemertean *Apatronemertes albimaculosa* and the leech *Placobdella parasitica* from the vicinity of the Panama Canal, and discuss potential circumstances leading to their disjointed distributions.

Material and methods

Specimen collection and DNA sequencing

Eleven leech specimens were collected in December 2015 from the exposed skin of the pond slider (*Trachemys scripta*) in Rio Chagres, Gamboa (09°06.96017'N; 79°41.60917'W). Four specimens of ribbon worms were collected in June 2017 from submerged logs and rocks in a small, unnamed pond directly adjacent to Lago Miraflores (Panama Canal) (09°00.680'N; 79°35.730'W). The fauna associated with the nemerteans included various crustaceans (mainly crabs and isopods), freshwater polychaetes, insect larvae, and smaller fish. Nemertean specimens were photographed prior to preservation and all specimens were preserved in 95% EtOH. A small portion of the posterior end (= caudal sucker of *P. parasitica* to avoid contamination by the blood meal) was cut and used for DNA sequencing. The present study targeted a 658 bp region of the mitochondrial locus cytochrome *c* oxidase subunit I (COI), commonly used for zoological DNA barcoding (see Hebert et al. 2003) and that has been successfully applied to the identifications of both leeches and nemerteans in previous studies (Lagunas-Calvo et al. 2018; Kvist et al. 2010). Extraction of DNA was carried out using the DNeasy Blood and Tissue Kit (Qiagen, Valencia, CA, USA) following the manufacturer's protocol. Sequences for COI were amplified for each sample using 25 µl PCR reactions, consisting of 16.39 µl ddH₂O; 2.5 µl buffer; 2.5 µl MgCl₂; 1 µl of each primer [HCO2198 (5'-TAAACTTCAGGGT GACCAAAAAATCA-3') and LCO1490 (5'-GGTC AACAAATCATAAAGATATTGG-3'); Folmer et al. 1994] at 10 µM concentration; 0.56 µl dNTPs at 10 mM concentration; 0.05 µl Platinum *Taq* polymerase (Invitrogen, Carlsbad, CA); and 1 µl total genomic DNA. PCR products were subsequently purified with ExoSAP-IT (Affymetrix, Santa Clara, CA) according to the manufacturer's protocol. Cycle sequencing reactions (10 µl total volume) were carried out using 0.5 µl each of ABI Big Dye Terminator V 3.1 and Big Dye 5x Sequencing Buffer (Applied Biosystems, Carlsbad, CA); 2 µl primer at 10 µM concentration; and 2.5 µl purified PCR product. Reaction mixtures were heated to 96 °C for 1 min. followed by 30 cycles of 96 °C (10 s), 50 °C (5s) and 60 °C (4 min). The samples were sequenced on




Figure 1. Select photos of collected specimens. (a) live habitus of a specimen of *Apatronemertes albimaculosa* from a pond adjacent to Lago Miraflores (Panama Canal), (b) preserved specimen of *Apatronemertes albimaculosa* from a pond adjacent to Lago Miraflores (Panama Canal), (c) dorsal view of preserved specimen of *Placobdella parasitica* from Rio Chagres, Gamboa, (d) ventral view of preserved specimen of *Placobdella parasitica* from Rio Chagres, Gamboa. Photographs by Alejandro Oceguera-Figueroa.

an ABI PRISM 3730XL DNA Analyzer (Applied Biosystems, Carlsbad, CA). All resulting sequences were deposited in GenBank (accession numbers MH128987–MH128991). Leech specimens were deposited in the Colección Nacional de Helmintos (CNHE: 10678) at the Universidad Nacional Autónoma de México, Mexico and nemerteans were deposited in the Colección Nacional de Invertebrados (CINV: 107) at the Universidad Nacional Autónoma de México, Mexico. Specimens were collected under permit SE/A-61-17 issued by MiAmbiente, Dirección de Áreas Protegidas y Vida Silvestre, Panama.

Morphological and molecular identifications

Morphological examinations were carried out using a Zeiss Discovery V8 (Toronto, Canada) dissecting microscope, and provisional identifications were affixed to the specimens. Nemerteans were identified to order-level, whereas the leeches were identified to species-level; see Figure 1 for select photographs of the specimens; for detailed morphological accounts of *A. albimaculosa*, see Wilfert and Gibson (1974) and Kajihara et al. (2016); and for accounts of *P. parasitica*, see Moser et al. (2013). Provisional

identifications were later corroborated and refined with the aid of DNA sequences, employing a standard BLASTn search approach against the non-redundant (nr) database on GenBank.

Additionally, phylogenetic trees were constructed for each taxon in order to provide robust identifications of the newly collected specimens under a tree-based framework. One analysis for each target taxon was conducted using a subset of the data from de Carle et al. 2017 (for the leech) and Kvist et al. 2015 (for the nemertean). The datasets were separately aligned using the online version of MAFFT ver. 7 (Katoh and Standley 2013) employing the “auto” option and default settings. PartitionFinder ver. 1.1.1 (Lanfear et al. 2012) was used to estimate the optimal partitioning scheme and model of nucleotide evolution under the Akaike Information Criterion, considering each codon position. Maximum likelihood analyses were then performed with RAxML ver. 8 (Stamatakis 2014) on the CIPRES platform (Miller et al. 2010). For the nemertean dataset, analyses used a GTR+Γ model of nucleotide evolution for three partitions (two of these also included a proportion of invariable sites parameter). For the leech dataset, two partitions (both following a GTR+Γ model of nucleotide evolution) were suggested by PartitionFinder and one of these included a proportion of invariable sites. The RAxML runs consisted of 1,000 iterations with 25 initial GAMMA rate categories that were allowed to fluctuate independently for each partition, and final optimization with four GAMMA shape categories. Support values for the clades were estimated via 100 iterations of the rapid bootstrap algorithm using the same settings as above.

Uncorrected *p*-distances were calculated in PAUP* ver. 4.0b10 (Swofford 2002) applying the following settings: function of minimal evolution, ignoring gaps for affected pairwise comparisons (although no gaps were present in the resulting alignments) and equal rates for variable sites.

Results

Morphological identifications

Based on external morphology only (no internal morphological analysis was carried out, as these are most often not needed for accurate specimen identifications [see Sundberg et al. 2016]), our nemertean specimens agree with the description of *Apatronemertes albimaculosa*. Importantly, live specimens showed a bright red coloration, with small flecks of white; a bluntly rounded head that is not demarcated from the remainder of the body; cephalic grooves extending from tip of the head to close to the mouth;

and a lack of eye-spots, frontal organ and caudal cirrus (Figure 1a, b). See Sundberg et al. (2016) for an in-depth discussion on the utility of external morphology for the identification of nemertean species.

The newly collected specimens of *Placobdella parasitica* are compatible with the original description (Say 1824), as well as the redescription (Moser et al. 2013) of the species. In particular, our specimens display a smooth (unpapillated) dorsum with a yellow medial line, and 8–12 darker, sometimes blue, anteroposterior lines on its venter (Figure 1 c, d).

Molecular identifications

DNA from three ribbon worm specimens and two leech specimens was successfully sequenced. The final alignments each consisted of 658 aligned sites for 54 (Nemertea) and 37 (Hirudinea) terminals. Initial BLASTn searches suggested that the ribbon worms belong to the species *Apatronemertes albimaculosa* (*e*-values = 0; identity scores 98%–99%), and that the leeches belong to *Placobdella parasitica* (*e*-values = 0; identity scores 99%). The resulting phylogenetic hypotheses corroborate these identifications (Figures 2 and 3), as both taxa nest within highly supported clusters of conspecific individuals with short branch lengths. Average uncorrected *p*-distances between our specimens of *Apatronemertes albimaculosa* and that of the only other nominal specimen deposited in GenBank, which was collected in Japan (accession number LC095809; Kajihara et al. [2016]), is $0.06\% \pm 0.009$. The same distance measure for the specimens of *Placobdella parasitica*, when compared to specimens from a variety of North American localities is $1.62\% \pm 0.27$; when compared only to specimens collected at the type locality of the species (see Moser et al. 2013), the same value is $1.53\% \pm 0.08$. It is worth noting that relatively high levels of genetic diversity seem to be common in *P. parasitica*: sequences for this species deposited in GenBank show a maximum pairwise COI distance of 2.04%. Notwithstanding this genetic disparity, specimens of *Placobdella parasitica* are easily identified using unequivocal morphological characters.

Discussion

One nemertean species, *Apatronemertes albimaculosa*, and one hirudinean species, *Placobdella parasitica*, are recorded for the first time from the Panama Canal, Panama. The nemertean species has hitherto only been recorded in conjunction with plant material related to the aquarium trade in Europe, Japan, and USA. Our new records represent the first




Figure 2. Maximum likelihood tree ($\ln L = -11845.231952$) resulting from the analysis of the nemertean COI dataset. Maximum likelihood bootstrap values above 50% are shown above each node, newly sequenced taxa are shown in bold font and the target taxa are highlighted by a grey background. GenBank accession numbers are noted before each taxon name.

case of wild-caught specimens, possibly suggesting that the species is native to the area surrounding the Panama Canal. The freshwater biota of the Panamanian Isthmus is known to be heavily influenced by marine ecosystems (Hildebrand 1939; Menzies 1968; San Martín et al. 1998; Ruiz et al. 2006; Galil et al. 2008), such that several species from predominantly marine taxonomic groups seem to be able to persist in freshwater habitats of Panama. The presence of *A. albimaculosa* in this freshwater canal may represent another example of this phenomenon. To further support this hypothesis, Wilfert and Gibson (1974) and Kajihara et al. (2016) suggest a tropical origin

for the plant material with which *A. albimaculosa* has historically been associated. The plant genus *Vallisneria*, for instance, is hypothesized to be very widespread (Good 1964; Les et al. 2003) and, whereas its native range has yet to be established (Les et al. 2008), it does occur in the neotropics of Central America (e.g. Les et al. 2008; Balcuñas and Center 1981). Alternatively, because of the sheer number of vessels that move through the Panama Canal – and, by extension, the high volume of ballast water and elevated potential for hull fouling, it is possible that the species was introduced to Panamanian waters from a different locality. There are already numerous




Figure 3. Maximum likelihood tree ($\ln L = -5950.074003$) resulting from the analysis of the hirudinean COI dataset. Maximum likelihood bootstrap values above 50% are shown above each node, newly sequenced taxa are shown in bold font and the target taxa are highlighted by a grey background. GenBank accession numbers are noted before each taxon name (note that GenBank numbers beginning with KF0588 represent specimens collected at the type locality [see Moser et al. 2013]).

accounts of non-endemics in the Panama Canal (e.g. Cohen 2006), and this species could potentially be added to the ever-growing list. The reproductive cycle of *Apatronemertes albimaculosa* certainly facilitates dispersal: the species lays eggs in a mucous sheet that is then attached to plant material (Wilfert and Gibson 1974), which may be sold in the aquarium trade or be taken up as ballast. It is unclear how resilient the eggs are to changes in external conditions (pollution, temperature, pH) that would inevitably arise from the invasion of new areas, but the adult worms seem to be rather susceptible to changes in temperature: exposure to temperatures below 10 °C is fatal (Kajihara et al. 2016). Based on our current knowledge of the species, it is impossible to establish the direction of introduction, yet its presence in disparate areas is clearly a byproduct of human activity.

The presence of *Placobdella parasitica* in Panama is less ambiguous. Given the species' abundance in North American ponds and lakes, and its absence south of the USA, our new record likely represents an introductory event (see Cornejo et al. 2015 for an account of current knowledge of Panamanian leeches). The turtle host, *Trachemys scripta*, was introduced to Panama, likely through the pet trade (Moll and Legler 1971). Both species occur throughout large parts of North America: *P. parasitica* occurs from the southernmost areas of Canada – between New Brunswick and Saskatchewan – as far south as Texas, USA, and is abundant throughout its range (Dodd Jr. 1988; Donaldson and Echternacht 2005; Davy et al. 2009); while *T. scripta* is distributed between southern Virginia and Florida, resulting in an expansive geographic overlap between host and parasite. In

both adult and juvenile stages, *Placobdella parasitica* is a more-or-less stationary ectoparasite, presumably only detaching from its hosts to reproduce. After mating, Glossiphoniid leeches show a high degree of parental care, brooding their eggs on their venters until the juveniles hatch, and are delivered to their first blood meal (Siddall et al. 2005). It is therefore difficult to envision a scenario in which the eggs of *P. parasitica* would be inadvertently taken up as ballast, and it seems even less likely that a parasitized turtle would persist in a ballast tank for extended periods, much less survive transportation through the filtering systems that mitigate ballast water intake. Equally improbable is the notion that the leeches would be transported attached to the hull of a vessel, as this species cannot tolerate long periods of exposure to salt water. We therefore hypothesize that *Placobdella parasitica* was inadvertently introduced via the pet trade along with its turtle host. It seems clear that the geographic distribution of this particular leech is underlain by intricacies and serendipity.

Acknowledgements

We thank Carlos Nieto from the Colección Zoológica Dr. Eustorgio Méndez for allowing us to access deposited material. We also thank Sergio Bermudez from ICGES for collecting the specimens of *Placobdella parasitica*, Edgar Pérez and Jaime Aguilar for providing fieldwork support and El Ministerio de Ambiente de Panamá for facilitating and providing collection permits (SE/A-61-17) during the field trip. A.C. thanks PAPIIT (grant # IA204114) for funding and S.K. was funded by a NSERC discovery grant, a Peer Review grant from the Royal Ontario Museum and a stipend from Olle Engkvist Byggmästare Foundation. Andrew David and two anonymous reviewers provided helpful feedback that benefitted the manuscript.

References

- Aron WI, Smith SH (1971) Ship canals and aquatic ecosystems. *Science* 174: 13–20, <https://doi.org/10.1126/science.174.4004.13>
- Balciunas JK, Center TD (1981) Preliminary host specificity tests of a Panamanian *Parapoxyn* 1 as a potential biological control agent for *Hydrilla* 2. *Environmental Entomology* 10: 462–467, <https://doi.org/10.1093/ee/10.4.462>
- Brooks RJ, Galbraith DA, Layfield JA (1990) Occurrence of *Placobdella parasitica* (Hirudinea) on snapping turtles, *Chelydra serpentina*, in southeastern Ontario. *Journal of Parasitology* 76: 190–195, <https://doi.org/10.2307/3283014>
- Carse A (2012) Nature as infrastructure: making and managing the Panama Canal watershed. *Social Studies of Science* 42: 539–563, <https://doi.org/10.1177/0306312712440166>
- Choudhury A, Zheng S, Pérez-Ponce de León G, Martínez-Aquino A, Brosseau C, Gale E (2013) The invasive Asian fish tapeworm, *Bothriocephalus acheilognathus* Yamaguti, 1934, in the Chagres River/Panama Canal drainage, Panama. *BioInvasions Record* 2: 99–104, <https://doi.org/10.3391/bir.2013.2.2.02>
- Cohen AN (2006) Chapter III - Species Introductions and the Panama Canal. In: Gollasch S, Galil BS, Cohen AN (eds), *Bridging Divides - Maritime Canals as Invasion Corridors*. Monographiae Biologicae, vol. 83. Springer, Berlin Heidelberg, New York, USA, pp 127–206
- Cornejo A, Oceguera-Figueroa A, Vega JAB (2015) Sanguijuelas (Annelida: Clitellata) de agua dulce de Panamá: comparación con la riqueza de especies de Centro América. *Puente Biológico* 7: 1–13
- Davy CM, Shim KC, Coombes SM (2009) Leech (Annelida: Hirudinea) infestations on Canadian turtles, including the first Canadian record of *Helobdella modesta* from freshwater turtles. *The Canadian Field-Naturalist* 123: 44–47, <https://doi.org/10.22621/cfn.v123i1.673>
- de Carle D, Oceguera-Figueroa A, Tessler M, Siddall ME, Kvist S (2017) Phylogenetic analysis of *Placobdella* (Hirudinea: Rhynchobdellida: Glossiphoniidae) with consideration of COI variation. *Molecular Phylogenetics and Evolution* 114: 234–248, <https://doi.org/10.1016/j.ympev.2017.06.017>
- Dodd Jr. CK (1988) Patterns of distribution and seasonal use of the turtle *Sternotherus depressus* by the leech *Placobdella parasitica*. *Journal of Herpetology* 22: 74–81, <https://doi.org/10.2307/1564358>
- Donaldson BM, Echternacht AC (2005) Aquatic habitat use relative to home range and seasonal movement of eastern box turtles (*Terrapene carolina carolina*: Emydidae) in eastern Tennessee. *Journal of Herpetology* 39: 278–284, [https://doi.org/10.1670/0022-1511\(2005\)039\[0278:AHURTH\]2.0.CO;2](https://doi.org/10.1670/0022-1511(2005)039[0278:AHURTH]2.0.CO;2)
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek RC (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294–299
- Galil BS, Nehring S, Panov V (2008) Waterways as invasion highways – Impact of climate change and globalization. In: Nentwig W (ed), *Biological Invasions*. Springer Berlin, Heidelberg, Germany, pp 59–74
- Gibson R, Moore J (1976) Freshwater nemerteans. *Zoological Journal of the Linnean Society* 58: 177–218, <https://doi.org/10.1111/j.1096-3642.1976.tb00828.x>
- Good R (1964) The geography of the flowering plants. 3d edn. Wiley, New York, USA, 518 pp
- Hanna GD (1966) Introduced mollusks of the western North America. *Occasional Papers of the California Academy of Sciences* 48: 1–108
- Hebert PD, Cywinski A, Ball SL, deWaard JR (2003) Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London B: Biological Sciences* 270: 313–321, <https://doi.org/10.1098/rspb.2002.2218>
- Hildebrand SF (1939) The Panama Canal as a passageway for fishes, with lists and remarks on the fishes and invertebrates. *Zoologica* 24: 15–45
- Jones ML (1972) The Panamic biota: some observations prior to a sea-level canal. *Bulletin of the Biological Society of Washington* 2: 1–270
- Kajihara H, Chernyshev AV, Sun SC, Sundberg P, Crandall FB (2008) Checklist of nemertean genera and species published between 1995 and 2007. *Species Diversity* 13: 245–274, <https://doi.org/10.12782/specdiv.13.245>
- Kajihara H, Takibata M, Grygier MJ (2016) Occurrence and molecular barcode of the freshwater heteronemertean *Apatronemertes albimaculosa* (Nemertea: Pilidiophora) from Japan. *Species Diversity* 21: 105–110, <https://doi.org/10.12782/sd.21.2.105>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* 30: 772–780, <https://doi.org/10.1093/molbev/mst010>
- Klemm DJ (1985) A guide to the freshwater Annelida (Polychaeta, Naidid and Tubificid Oligochaeta, and Hirudinea) of North America. Kendall/Hunt Publishing Co, Dubuque, Iowa, pp 1–198
- Kvist S, Chernyshev AV, Giribet G (2015) Phylogeny of Nemertea with special interest in the placement of diversity from Far East Russia and northeast Asia. *Hydrobiologia* 760: 105–119, <https://doi.org/10.1007/s10750-015-2310-5>
- Kvist S, Oceguera-Figueroa A, Siddall ME, Erséus C (2010) Barcoding, types and the *Hirudo* files: using information content to critically

- evaluate the identity of DNA barcodes. *Mitochondrial DNA* 21: 198–205, <https://doi.org/10.3109/19401736.2010.529905>
- Lagunas-Calvo O, Hernández-Mena DI, Rivas G, Contreras-Mirón S, Mendoza-Garfias B, Echeverría-Martínez OM, Oceguera-Figueroa A (2018) Rediscovery of the nemertean *Prostoma graecense* from Xochimilco, 75 years after its first and only record. *Revista Mexicana de Biodiversidad* 89: 39–45, <https://doi.org/10.22201/ib.20078706e.2018.1.2094>
- Lanfear R, Calcott B, Ho SY, Guindon S (2012) PartitionFinder: combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Molecular Biology and Evolution* 29: 1695–1701, <https://doi.org/10.1093/molbev/mss020>
- Les DH, Crawford DJ, Kimball RT, Moody ML, Landolt E (2003) Biogeography of discontinuously distributed hydrophytes: a molecular appraisal of intercontinental disjunctions. *International Journal of Plant Sciences* 164: 917–932, <https://doi.org/10.1086/378650>
- Les DH, Jacobs SW, Tipperry NP, Chen L, Moody ML, Wilstermann-Hildebrand M (2008) Systematics of *Vallisneria* (Hydrocharitaceae). *Systematic Botany* 33: 49–65, <https://doi.org/10.1600/03636440873887483>
- Mason CY, Rowlands A (1938) Panama Canal Traffic. *Economic Geography* 14: 325–337, <https://doi.org/10.2307/141526>
- Menzies RJ (1968) Transport of marine life between oceans through the Panama Canal. *Nature* 220: 802–803, <https://doi.org/10.1038/220802a0>
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: *Gateway Computing Environments Workshop (GCE)*. IEEE, pp 1–8, <https://doi.org/10.1109/GCE.2010.5676129>
- Moll EO, Legler JM (1971) The life history of a Neotropical slider turtle, *Pseudemys scripta* (Schoepff), in Panama. Bulletin 11. Los Angeles County Museum of Natural History, Los Angeles, pp 1–102
- Moser WE, Richardson DJ, Hammond CI, Lazo-Wasem EA (2013) Redescription of *Placobdella parasitica* (Say, 1824) Moore, 1901 (Hirudinida: Glossiphoniidae). *Bulletin of the Peabody Museum of Natural History* 54: 255–262, <https://doi.org/10.3374/014.054.0203>
- Oceguera-Figueroa A, Pacheco-Chaves B (2012) Registros de sanguíjuelas de Costa Rica y clave para la identificación de las especies con redescrición de *Cylacobdella costaricensis*. *Revista Mexicana de Biodiversidad* 83: 946–957, <https://doi.org/10.7550/rmb.31623>
- Roche DG, Torchin ME (2007) Established population of the North American Harris mud crab, *Rhithropanopeus harrisii* (Gould 1841) (Crustacea: Brachyura: Xanthidae) in the Panama Canal. *Aquatic Invasions* 2: 155–161, <https://doi.org/10.3391/ai.2007.2.3.1>
- Ruiz GM, Lorda J, Arnwine A, Lion K (2006) Chapter II Shipping patterns associated with the Panama Canal: effects on biotic exchange? In: Gollasch S, Galil BS, Cohen AN (eds), *Bridging Divides - Maritime Canals as Invasion Corridors. Monographiae Biologicae*, vol. 83. Springer, Berlin Heidelberg New York, pp 113–126, https://doi.org/10.1007/978-1-4419-5047-3_4
- Ruiz GM, Torchin ME, Grant K (2009) Using the Panama Canal to test predictions about tropical marine invasions. *Smithsonian Contributions to the Marine Sciences* 38: 291–300
- Ryan TJ, Lambert A (2005) Prevalence and colonization of *Placobdella* on two species of freshwater turtles (*Graptemys geographica* and *Sternotherus odoratus*). *Journal of Herpetology* 39: 284–287, <https://doi.org/10.1670/180-04N>
- San Martín G, López E, Camacho AI (1998) First record of a freshwater Pisionidae (Polychaeta): description of a new species from Panama with a key to the species of *Pistone*. *Journal of Natural History* 32: 1115–1127, <https://doi.org/10.1080/00222939800770561>
- Sawyer RT (1986) Leech biology and behaviour. New York: Oxford University Press, pp 419–793
- Say T (1824) D. Class Vermes. Order Cryptobranchia. Hirudo. In: W.H. Keating, Ed. *Narrative of an expedition to the source of St. Peter's River, Lake Winnepeek, Lake of the Woods, performed in the year 1823, by order of the Hon. J. C. Calhoun, Secretary of War, under the command of Stephen H. Long, Major U.S.T.E.: Compiled from the notes of Major Long, Messrs. Say, Keating and Colhoun. Volume 2*. Philadelphia: H.C. Carey and I. Lea, pp 266–268
- Schlöder C, Canning-Clode J, Saltonstall K, Strong EE, Ruiz GM, Torchin ME (2013) The Pacific bivalve *Anomia peruviana* in the Atlantic: a recent invasion across the Panama Canal? *Aquatic Invasions* 8: 443–448, <https://doi.org/10.3391/ai.2013.8.4.08>
- Senz W (1993) Nemertinen europäischer Küstenbereiche (nebst ergänzenden Angaben zur Anatomie von *Apatornemertes albimaculosa* Wilfert & Gibson, 1974). *Annalen des Naturhistorischen Museums in Wien, Serie B für Botanik und Zoologie* 94/95: 47–145
- Siddall ME, Budinoff RB, Borda E (2005) Phylogenetic evaluation of systematics and biogeography of the leech family Glossiphoniidae. *Invertebrate Systematics* 19: 105–112, <https://doi.org/10.1071/IS04034>
- Smith SA, Bell G, Bermingham E (2004) Cross-Cordillera exchange mediated by the Panama Canal increased the species richness of local freshwater fish assemblages. *Proceedings of the Royal Society of London B: Biological Sciences* 271: 1889–1896, <https://doi.org/10.1098/rspb.2004.2796>
- Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1312–1313, <https://doi.org/10.1093/bioinformatics/btu033>
- Strand M, Sundberg P (2015) Phylum Nemertea. In: Thorp JH, Rogers DC (eds), *Freshwater Invertebrates: Ecology and General Biology*, Fourth Edition. Academic Press, London, U.K., pp 205–209, <https://doi.org/10.1016/B978-0-12-385026-3.00011-5>
- Sundberg P, Gibson R (2008) Global diversity of nemerteans (Nemertea) in freshwater. *Hydrobiologia* 595: 61–66, <https://doi.org/10.1007/s10750-007-9004-6>
- Sundberg P, Andrade S, Bartolomaeus T, Beckers P, von Döhren J, Krämer D, Gibson R, Giribet G, Herrera-Bachiller A, Junoy J, Kajihara H, Kvist S, Kånnby T, Sun S-C, Thiel M, Turbeville JM, Strand M (2016) The future of nemertean taxonomy (phylum Nemertea) — a proposal. *Zoologica Scripta* 45: 579–582, <https://doi.org/10.1111/zsc.12182>
- Swofford DL (2002) PAUP* version 4.0 b10. Sinauer, Sunderland, MA
- Tessler M, Siddall M, Oceguera-Figueroa A (2018) Leeches from Chiapas, Mexico, with a new species of *Erpobdella* (Hirudinida, Erpobdellidae). *American Museum Novitates* 3895: 1–15, <https://doi.org/10.1206/3895.1>
- Wasson K, Zabin C, Bedinger L, Diaz MC, Pearse JS (2001) Biological invasions of estuaries without international shipping: the importance of intraregional transport. *Biological Conservation* 102: 143–153, [https://doi.org/10.1016/S0006-3207\(01\)00098-2](https://doi.org/10.1016/S0006-3207(01)00098-2)
- Wilfert M, Gibson R (1974) A new genus of hermaphroditic freshwater heteronemertean (Nemertinea). *Zoomorphology* 79: 87–112, <https://doi.org/10.1007/BF00298777>
- Wonham MJ, Walton WC, Ruiz GM, Frese AM, Galil B (2001) Going to the source: role of the invasion pathway in determining potential invaders. *Marine Ecology Progress Series* 215: 1–12, <https://doi.org/10.3354/meps215001>