



Zenith™ Red & Gold

syngenta®flowers

Marigold Triploid Zenith™

Culture Guide

Botanical name: *Tagetes triploid*

Product form: Seed

Containers: Packs, Pints, Quarts, Gallons

Habit: Mounded

Garden Specifications

Garden Height: 14" (35 cm) tall

Garden Width: 14" (35 cm) wide

Exposure: Full sun

USDA zone: 10

AHS zone: 12–1

Product use: Packs, Containers, Landscapes, Patio Pots, Combos

Germination

Stages 1 & 2

Germination time: 3–5 days

Media temp: 68–72 °F (20–22 °C)

Chamber: Optional

Light: Not required for germination

Seed cover: Yes

Moisture level: 5 - SATURATED: Soil is dark brown and shiny, free water is present at the surface of the soil, water drips freely from the bottom of the tray, and trays are heavy with a visible bend in the middle.

Rec. tray size: 288-cell tray

Seeds per cell: 1

Young Plant Production

Stages 3 & 4

Temperature

Day: 66–68 °F (19–20 °C)

Night: 66–68 °F (19–20 °C)

Average daily temperature: 67 °F (19 °C)

Lighting

Day extension lighting: Not necessary

Light intensity: 2,000–2,500 foot candles (400–500 micro mols)

Day length response: Day neutral

Daily light integral: 10–12 mols/day

Media pH: 5.8–6.2

Media EC: SME 0.5 to 0.75 mS/cm

Fertilizer: 75–125 ppm N

Moisture level: Media should be allowed to dry between irrigations. Alternate between moisture level 2 and 4.

2 - MEDIUM: Soil is light brown in color, no water can be extracted from soil, and soil will crumble apart.

4 - WET: Soil is dark brown but not shiny, no free water is seen at the surface of the soil, when pressed or squeezed water drips easily, and trays are heavy with a visible bend in the middle.

Pinching: No

Plant growth regulators (PGRs): Usually not needed. If necessary, spray with B-Nine® WSG at 1,500–2,500 ppm to control stem elongation.

Plug grow time: 3–4 weeks in a 288-cell tray

Comments: Very easy to germinate on the bench. If seedlings get stretched, they can be planted deep at transplanting, and new roots will form along the stems. Do not hold too long in the plug tray or stress-induced flowering may occur, and plants will flower before bulking up to a satisfactory size.

Finishing

Temperature

Day: 66–68 °F (19–20 °C)

Night: 62–64 °F (17–18 °C)

Average daily temperature: 65 °F (18 °C)

Lighting

Day extension lighting: Not necessary

Light intensity: 4,000–6,000 foot candles

Day length response: Day neutral

Daily light integral: 12–14 mols/day

Transplanting: Transplant directly into the finished container. Place the rooting media slightly below the level of media in the container. Make sure the root ball is covered and that the plug is situated in the center of the pot.

Media pH: 6.1–6.5

Media EC: SME EC: 1.5–2.1 mS/cm, PourThru EC: 2.3–3.2 mS/cm

Fertilizer: 125–175 ppm N

Pinching: No. Pinching is not recommended or needed.

Plant growth regulators (PGRs): Growth can be controlled naturally through proper moisture management. If needed to control height, spray with B-Nine® WSG at 2,500–5,000 ppm, Bonzi® at 15–30 ppm, or Sumagic® at 5–10 ppm. To help control growth in the retail environment drench with Bonzi® at 2–5 ppm or Sumagic® at 1–2 ppm when plants are up to size and flowering.

Tech tip: If plugs are stretched, plant them deep (up to the cotyledons is OK) to prevent floppiness. Maintain soil pH above 6.1 to avoid iron and/or manganese toxicity.



Try Chrysal Alesco®, a postharvest foliar spray, to protect ethylene sensitive crops during shipping and retail

Moisture level: Media should be allowed to dry between irrigations. Alternate between moisture level 2 and 4.

2 - MEDIUM: Soil is light brown in color, no water can be extracted from soil, and soil will crumble apart.

4 - WET: Soil is dark brown but not shiny, no free water is seen at the surface of the soil, when pressed or squeezed water drips easily, and trays are heavy with a visible bend in the middle.

Common pests: Aphids, Whiteflies, Thrips, Spider mites, Leafminers

Common diseases: Pythium, Botrytis, Bacterial leaf spot, Alternaria leaf spot

Scheduling

Size	Crop Time	Plants Per Pot
Large Packs	4–5 weeks	1 ppp
1.0 pint (4 inch)	5–6 weeks	1 ppp
1.0 quart (4.5 to 5 inch)	5–6 weeks	1 ppp
1.25 to 2.5 quart (5.5 to 6.5 inch, trade gallon)	6–7 weeks	2–3 ppp

Estimated finish crop time is from transplant of a 288-cell tray and finished at an average daily temperature (ADT) of 65 °F (18 °C).

Example crop schedule for a 1.0 pint

Weeks From Transplant	Description
2 weeks	Check media pH and adjust if under 6.1
3 weeks	If needed to control height, spray with B-Nine® WSG at 2,500–5,000 ppm, Bonzi® at 15–30 ppm, or Sumagic® at 5–10 ppm
5 weeks	To help control growth in the retail environment, drench with Bonzi® at 2–5 ppm, or Sumagic® at 1–2 ppm
6 weeks	Ready to ship



Zenith™ Lemon Yellow

syngenta® flowers

All photos are either the property of Syngenta or are used with permission.

© 2020 Syngenta. **Important: Always read and follow label instructions. Some products may not be registered for sale or use in all states or counties. Please check with your local Extension Service to ensure registration status. Some or all of the varieties may be protected under one or more of the following: Plant Variety Protection, United States Plant Patents and/or Utility Patents and may not be propagated or reproduced without authorization.** The trademarks or service marks displayed or otherwise used herein are the property of a Syngenta Group Company. All other trademarks are the property of their respective owners.