

A record of the mysid *Hemimysis lamornae mediterranea* (Crustacea: Mysida) from the western Mediterranean, with a complete morphological description

CARLOS SAN VICENTE¹, LIDIA DELGADO², ESTEBAN HERNANDEZ³ AND GUILLERMO GUERAO³

¹c/Nou núm 8, 43839-Creixell, Tarragona, Spain, ²Departament de Biologia Animal, Facultat de Biologia, Universitat de Barcelona, Avenida Diagonal 645, Barcelona, Spain, ³IRTA, Unitat Operativa de Cultius Aquàtics, Ctra. Poble Nou, Km 5.5, 43540-Sant Carles de la Ràpita, Tarragona, Spain

The mysid Hemimysis lamornae mediterranea Bacescu, 1936 is described from specimens sampled in the Ebro Delta, Spain, north-western Mediterranean. To date, this subspecies was only known from the Gulf of Naples and Marseille; this is the first record of the H. lamornae mediterranea on the Iberian Peninsula coast. Hemimysis lamornae mediterranea is distinguishable from its closest congeners, H. lamornae typica (Couch, 1856) and H. lamornae pontica (Czerniavsky, 1882) by the number of setae, the shape of the maxilla, and the relatively smaller number of spines on the uropod endopod and on the lateral margin of the telson.

Keywords: taxonomy, Mysidae, *Hemimysis*, Ebro Delta, Spain, western Mediterranean Sea

INTRODUCTION

Although the Mediterranean Sea is one of the best known geographical areas of the world, there are many regions and habitats that remain insufficiently studied, and several taxonomic groups are still poorly known from this region (Coll *et al.*, 2010). Improved species descriptions as well as data on geographical distributions are still a high priority. The Ebro Delta, Spain constitutes a good example of how a coastal area of great ecological significance in the western Mediterranean can still remain insufficiently known for some crustacean peracarid taxa such as mysids.

Hemimysis is a group of small-sized and usually red coloured mysids widely distributed throughout the north-eastern Atlantic Ocean, the Mediterranean Sea, the Black Sea and the Caspian Sea (Tattersall & Tattersall, 1951; Bacescu, 1954; Ledoyer, 1989). Usually, *Hemimysis* individuals form dense swarms in open waters (>6000 ind. m^{-3} ; Ketelaars *et al.*, 1999) or in dark submarine caves (more than 10^5 individuals m^{-3} ; Carola *et al.*, 1993). Some species have been extensively studied, mainly *H. anomala* involved in recent invasive events in the northern hemisphere (see Wittmann & Ariani, 2009; Minchin & Boelens, 2010 for references). The genetic structure of Mediterranean cave-dwelling *Hemimysis* populations has been studied for *H. margalefi* and *H. speluncola* (Lejeusne & Chevaldonné, 2006). This work confirms the strong influence of habitat disjunction (natural fragmentation) on population structure.

Hemimysis lamornae has a vast distribution from sub-Arctic Icelandic waters to the North Sea and western Baltic waters and to the Mediterranean and the Black Sea. The species breeds all the year (at least in captivity) and its biology is reasonably well studied (Cannon & Manton, 1927; Manton, 1928; Foxon, 1940; Tattersall & Tattersall, 1951). Bacescu (1936) distinguished 3 subspecies, according to size-range of individuals, number of thoracopod articles, armature of uropod and telson: *H. lamornae typica* (Couch, 1856) from the Atlantic European coast; *H. lamornae mediterranea* Bacescu, 1936 from the Mediterranean Sea; and *H. lamornae pontica* (Czerniavsky, 1882) from the Black Sea. As demonstrated by its extended distributional area, *H. lamornae* shows remarkable adaptative capacities.

The subspecies *H. lamornae mediterranea* is documented from the Ebro Delta (Sant Carles de la Ràpita, Spain). However, *H. lamornae mediterranea* was only known from the Gulf of Naples (Bacescu, 1936; Wittmann, 2001) and a submarine cave in Marseille (Ledoyer, 1989). The present paper deals with the recent discovery of a new population in the Ebro Delta, Sant Carles de la Ràpita, Spain. Previous descriptions only describe the main diagnostic characters of *H. lamornae mediterranea*. This present study provides a full description of both the male and female, as well as information on ontogenetic variability of diagnostic characters of this Catalanian population.

MATERIALS AND METHODS

Mysids were collected from a breakwater of the Sant Carles de la Ràpita harbour (Alfacs Bay, Ebro Delta, $40^{\circ}37'12''N$ $0^{\circ}35'34''E$) at a depth of 0.3–0.5 m. Samplings were carried out in June and November 2010, always during night-time

Corresponding author:

G. Guerao

Email: guillermo.guerao@irta.cat

(21:00–22:00 h), with a hand-held net (mesh-size: 500 μm ; mouth aperture diameter: 35 cm). Samples were preserved in 70% ethanol for laboratory examination.

Individuals of *H. lamornae mediterranea* were identified and separated from other species and classified into 6 demographic categories according to the following sexual characteristics: (1) juveniles (secondary sexual characteristics absent); (2) immature males (terminal segment of the antennular peduncle with non-hirsute appendix masculina on the ventral surface; with developing pleopod 4); (3) mature males (antennule with hirsute appendix masculina; pleopod 4 developed); (4) immature females (with developing oostegites at the bases of the two posterior thoracopods); (5) mature females (with a well-developed marsupium); and (6) brooding females (mature females with eggs or larvae in the marsupium).

The total length (TL) was measured from the tip of the rostrum to the posterior end of the telson, excluding spines. Dissection and measurements were taken with a Nikon SMZ800 stereo microscope equipped with an image analysing system (AnalySIS, SIS, Münster, Germany). An Olympus BH-2 microscope was used for the description of appendages morphology.

SYSTEMATICS

Order MYSIDA Haworth 1825

Family MYSIDAE Haworth, 1825

Subfamily MYSINAE Haworth, 1825

Genus *Hemimysis* Sars, 1879

Hemimysis lamornae mediterranea Bacescu, 1936

(Figures 1–6)

Hemimysis lamornae, Norman (1892), pp. 247–249, partim.

Hemimysis sp. Colosi (1929), pp. 428–430, figure 21.

Hemimysis lamornae mediterranea, Bacescu (1936), pp. 77–82, figure 3.

Hemimysis lamornae, Tattersall (1951), p. 163.

Hemimysis lamornae, Tattersall & Tattersall (1951), p. 335.

Hemimysis lamornae mediterranea, Ledoyer (1989), pp. 45–46, figure 3.

MATERIAL EXAMINED

Sant Carles de la Ràpita harbour, Ebro Delta, Spain: 9 brooding females (TL range: 5.5–6.2 mm); 4 immature females (TL range 4.6–5.1 mm); 4 mature males (TL range 4.8–5.8 mm); 10 immature males (TL range 3.3–4.4 mm); 3 juveniles (TL range 2–3.2 mm) and 1 intramarsupial postnauploid specimen; 30 April 2010. A total of 10 males and 10 females have been deposited in the Biological Collections of Reference of the Institut de Ciències del Mar (CSIC) in Barcelona, under accession codes ICM_20110207_01.

DESCRIPTION

General form robust. Carapace with a short subtriangular rostrum with rounded apex; antero-lateral angles rounded; posterior margin emarginated, leaving the last two thoracic somites exposed in dorsal view; cervical groove present. Eyes large, globular, slightly broader than the eyestalk, extending laterally beyond the limits of the carapace; cornea pigment black. Pleon with first five somites subequal in length; sixth somite twice as long as the fifth (Figure 1A–C). Antennular peduncle as long as antennal scale; first article longer than

broad, second article shortest, third article slightly longer than broad supporting an hirsute appendix masculina on the male (Figure 1E). Antennal peduncle 3-segmented, not extending beyond antennal scale; second article longest; inner distal margin of second ant third articles with 1–2 and one setae, respectively. Protopod of antenna with outer distal angle rounded. Antennal scale lanceolate, with apical suture, 3.5 times as long as maximum width, proximal half of outer margin naked and straight (Figure 1D, F). Labrum almost oval, longer than wide, posterior margin with two distinct areas consisting of a cluster of short setae and an area covered with small scale-like protrusions (Figure 2A, B). Mandibles well developed; three-segmented palp, 2nd article about twice as long as 3rd with 3–7 setae on dorsal margin and 21–22 setae on ventral margin (Figure 2C, F). Distal segment of mandible palp with 18 ventral spinose setae and one distal large conspicuous seta, dorsal margin with two setae (Figure 2D). Left mandible setal row consisting of four hirsute spines and right mandible setal row consisting of one hirsute spine and four entire spines (Figure 2E, G). Maxillule basis (outer lobe) apex armed with ten setae and five setae on ventral surface; coxal endite (inner lobe) with three apical setae and a row of small setae (Figure 2H). Maxilla with distal article of endopod longer than wide, setose on its distal two-third distal margin, apex armed with three cuspidate setae; exopod extending to 1/3 of distal article of endopod, with 22 setae on lateral margin; lateral margin of coxa armed with 11 setae (Figure 2I, J). First and second thoracic appendages formed as maxillipeds, similar to those of other species of the genus. First thoracopod with prominent lobe on the basis armed with 13 spinose setae and two rows of small setae on the proximal and distal margins; endopod with ischium and merus subequal in length, armed with 3 and 4 spinose setae on their medial margins, respectively; carpus with 7 spinose setae on its medial margin and 1 distal spinose seta on its distal margin; propodus with 3 spinose setae on its proximal medial margin, 3 spinose setae on its distal margin and 6 spinose setae on its lateral margin; dactylus shorter than propodus, armed with 8 spinose setae and 1 distal strong nail (Figure 3A, B). Second thoracopod longer than first; endopod with the merus armed with 3 setae on its medial margin; carpus and propodus subequal in length armed with a single medial seta and 2 medial distal setae, respectively; dactylus half length of propodus, armed distally with 3 strong and spinose nails (Figure 3C, D). Third to eighth thoracopods with the endopod longer than exopod; endopods with the inner margin of the basis rounded, pre-ischium short and more or less triangular in shape, ischium and merus subequal in length, carpo-propodus three-segmented, dactylus shorter than the distal segment of the carpo-propodus and armed with a slender nail; exopods nine segmented (Figure 3E–L). Male genital apophyses bearing hirsute apical setae (Figure 4A). Female pleopods rudimentary, unsegmented plates (Figure 4I–M). Male pleopods 1 and 2 stronger and larger than female pleopods; pleopod 3 with large sympod (Figure 4B–C); endopod short, unsegmented and armed with a few setae (Figure 4D); pleopod 4 with small unsegmented endopod; exopod five-segmented, distal articles bearing a long seta which is armed terminally with a shorter seta with small setae along one side (Figure 4E, G, H); pleopod 5 biramous, with two segmented sympod, and four articles both




Fig. 1. *Hemimysis lamornae mediterranea* from the Ebro Delta. (A) Female, dorsal view; (B) female, lateral view; (C) male, lateral view; (D) antenna of female, ventral view; (E) antennula of male, dorsal view; (F) antenna of male, ventral view. (A–B) Brooding female, TL = 6.8 mm; (C) mature male, TL = 6.2 mm; (D) brooding female, TL = 6.1 mm; (E–F) mature male, TL = 5.8 mm.

on endopod and exopod (Figure 4F). Exopod of uropod slender, about one and half times as long as the telson. Endopod of uropod shorter, extending beyond apex of telson for 1/4 of its length, armed on the inner margin near statocyst with two or three (juveniles) to five (mature female) setae (Figure 5A–D). Telson short, about sub-equal in length to the last abdominal somite, two times as long as broad (Figure 5E); lateral margins converging distally, armed on the distal half with 3–4 (postnauploide and juveniles) to seven (mature male) setae; telson cleft to about one fourth of its length, cleft armed with 6–8 (juveniles) to 14 (mature male and female) spines; apical lobes with a long strong seta at distal end (Figure 5F–J).

SIZE

Maximum size (total length) was 5.8 mm in adult males and 6.2 mm in females.

COLOUR

The Ebro Delta individuals have the body transparent with red-orange coloured specks evenly distributed on the carapace, eyestalks, medial dorsal and ventral pleon, thoracopods, pleopods, telson and uropods; with large black eyes (Figure 6).

DISTRIBUTION

Mediterranean Sea, known from Naples, Italy; Marseille, France and Ebro Delta, Spain.

HABITAT

Hemimysis lamornae mediterranea is a slightly euryhaline species that lives in shallow waters between and on rocks, algae (*Cystoseira*) or among *Zostera* (Bacescu, 1936, 1941). It emerges from outside on the sea bottom during the night and also in the nocturnal surface plankton (Wittmann, 2001). During the day, it is an abundant species in some sub-marine caves where it had a tendency to occur in abundance




Fig. 2. *Hemimysis lamornae mediterranea* from the Ebro Delta. (A) Labrum; (B) posterior margin of labrum; (C) right mandible; (D) mandibular palp, articles 2 and 3; (E) incisor to molar processes of right mandible; (F) left mandible; (G) incisor to molar processes of left mandible; (H) maxillule; (I) maxilla; (J) maxillar palp. (A–B, F–G & I–J) Brooding female, TL = 6.1 mm; (C–E) mature male, TL = 5.8 mm; (H) immature male, TL = 4.4 mm.

in the moderately lit areas being replaced in the darkest areas by *H. margalefi* and *H. spelunca* (Ledoyer, 1989).

REMARKS

Hemimysis lamornae mediterranea Bacescu, 1936 closely resembles the other two subspecies, *H. lamornae typica* (Couch, 1856) and *H. lamornae pontica* (Czerniavsky, 1882). The three subspecies can be distinguished by their different size as well as by some morphological details. However, Ledoyer (1989) expressed doubts about their validity and proposed to raise *H. lamornae mediterranea* as a valid separate species.

In the northern species, *H. lamornae typica*, the range of adults length was 8–10 mm, the cleft of the telson was one-quarter of its length, the lateral margins of the telson were armed with seven to ten setae, the endopods of the uropods were armed with 10–12 setae and the spinules arming the cleft of the telson were 36 (Sars, 1879; Tattersall & Tattersall, 1951).

Bacescu (1936) examined large numbers of specimens of *H. lamornae* from Romanian waters and the Gulf of Naples. By comparing these with the descriptions of specimens from northern waters of Europe, he found that certain differences between them remained constant. Adult




Fig. 3. *Hemimysis lamornae mediterranea* from the Ebro Delta. (A) Endopod of first maxilliped; (B) distal articles of endopod of first maxilliped; (C) endopod of second maxilliped; (D) distal article of endopod of second maxilliped; (E) third thoracopod; (F) distal articles of third thoracopod endopod; (G–K) fourth to eighth thoracopods; (L) distal articles of eighth thoracopod endopod. (A–B) Mature male, TL = 5.8 mm; (C–D & I–K) brooding female, TL = 6.1 mm; (E–H) brooding female, TL = 5.1 mm.

specimens from Romanian waters, including many breeding, and therefore adult, females agreed closely with Czerniavsky's description of *H. pontica*. The length of adults was 6–7 mm, the cleft of the telson was only one-seventh to one-eighth of the telson length, the lateral margins of the telson were armed with five to eight setae, the uropod endopods were armed with a single seta and the spinules arming the cleft of the telson were sixteen to seventeen. Bacescu (1936) found that these characters were quite constant and he did not find any intermediate forms between the Romanian and Napoli specimens. He considered that the northern type, as described by Sars (1879), Norman (1860) and Zimmer (1909), was the original species and called it *Hemimysis*

lamornae var. *typica*; that the Naples type, which he called *H. lamornae* var. *mediterranea* has been derived from it and that the Romanian form, *H. lamornae* var. *reducta* (synonymous with *H. pontica*, Czerniavsky, 1882) was derived from the Mediterranean form. Such a zoogeographical scheme was supported and amplified to new species by Ledoyer (1989).

The Mediterranean specimens of *H. lamornae mediterranea* from Naples were 6–8 mm in length, the cleft of the telson was one-fifth to one-sixth of the length of the telson, the cleft was armed with 22–30 spinules, the lateral margins bore five to seven setae, the inner margin of the endopod of the uropod was armed with three to five setae and the tarsus




Fig. 4. *Hemimysis lamornae mediterranea* from the Ebro Delta. (A) Eighth male thoracopod; (B–F) first to fifth male pleopods; (G, H) distal armed seta of penultimate and ultimate articles of 4th male pleopod exopod, respectively; (I–M) first to fifth female pleopod. (A, & D–H) Mature male, TL = 5.8 mm; (B–C) mature male, TL = 5 mm; (I–M) brooding female, TL = 6.0 mm.

of the endopods of the thoracic limbs was divided into only three, rarely four, subsegments (Bacescu, 1936).

The Ebro Delta specimens agree with the Bacescu's description of the Naples population of *H. lamornae mediterranea* but there is some difference with the Marseille specimens studied by Ledoyer (1989). The specimens from Marseille differ slightly from those of the Naples and Ebro Delta, especially in the number of setae on the inner margin of the uropod endopod (armed with 2–3 setae) and the thoracopod carpo-propodus (divided into four subsegments, three-segmented in the Naples and Ebro Delta populations). Nevertheless, the number of setae and the shape of the maxilla, the relatively small number of spines on the uropod endopod and on the lateral margin of the telson perfectly

distinguish this subspecies. Moreover, in the Ebro Delta population, *H. lamornae mediterranea* was not found coexisting with *H. margalefi*, a situation that may be common in underwater cave habitats (Ledoyer, 1989). This latter species is distinguishable from *H. lamornae mediterranea* by the shape and armature of the antennal scale, maxilla and telson.

Such morphological differences between populations may result from phenotypic plasticity in response to varying environmental conditions (e.g. temperature, salinity, food availability, flow regime, predator/prey interactions, etc.) within different geographical areas (Scheiner & Callahan, 1993). Intraspecific geographical variations have been observed in other mysids species, such as variation in the number of spines on the lateral margins of the telson




Fig. 5. *Hemimysis lamornae mediterranea* from the Ebro Delta. (A–B) Ventral view of uropod of two juveniles of different size; (C–D) ventral view of uropod of male and female, respectively; (E) dorsal view of telson and uropods of mature male; (F) distal end of telson of postnauploid; (G–H) distal end of telson of two juveniles of different size; (I–J) distal end of telson of immature and mature male, respectively. (A, G) Juvenile, TL = 2.0 mm; (B, H) juvenile, TL = 3.2 mm; (C, E, I) mature male, TL = 5.8 mm; (D) brooding female, TL = 6.1 mm; I, immature male, TL = 4.4 mm; (F) postnauploid premoult intramarsupial stage.

between populations of both *Praunus flexuosus* and *P. neglectus* (Mauchline, 1971), geographical differences in the proportions of the antennal scale of *Neomysis americana* (Williams *et al.*, 1974) or in the eye and telson morphology of *N. integer* and *Mesopodopsis slabberi* (Remerie *et al.*, 2005).

The ontogenetic morphological differences observed in *H. lamornae mediterranea* means to take care in the assignment of individuals to this subspecies, especially if adults are not available for examination. Following Wittmann (1992), without detailed species descriptions based on a diversity of characters, the morphological and typological species concepts may be erroneous.

To date, 10 mysid species (including *H. lamornae mediterranea*) are known from the estuarine and coastal zones of the

Ebro Delta: *Siriella clausi*, *Gastrosaccus mediterraneus*, *G. roscoffensis*, *G. sanctus*, *Leptomysis lingvura*, *L. mediterranea*, *Diamysis lagunaris*, *Mesopodopsis slabberi* and *Schistomysis assimilis* (Suau & Vives, 1957; Chinchilla & Comín, 1977; San Vicente & Munilla, 2000), a value probably underestimated in this area characterized by highly structuring environmental gradients and diversity of habitats.

Although it is difficult to draw a border between coastal and deep species, in well-known Mediterranean areas such as the Gulfs of Marseille and Naples, the reported number of species is much higher (Macquart-Moulin, 1965; Wittmann, 2001). Therefore, much remains to be studied to better understand the mysid biodiversity of one of the largest delta of the Mediterranean and European coasts.




Fig. 6. *Hemimysis lamornae mediterranea* from the Ebro Delta. (A) Brooding female, dorsal view; (B) brooding female, lateral view; (C) male, lateral view. Scale bar: 2 mm.

ACKNOWLEDGEMENT

This work was supported by the Ministry of Science and Research to G.G. (post-doctoral fellowship; INIA).

REFERENCES

- Bacescu M.** (1936) *Hemimysis lamornae*, subsp., *reducta* nov. subsp. et *Hemimysis anomala*, dans les eaux Roumaines de la Mer Noire (avec une étude comparative des formes de *Hem. lamornae* des autres mers: *H.l. typica* et *H.l. mediterranea* nov. subsp.). *Annales Scientifiques de l'Université de Jassy* 19, 331–338.
- Bacescu M.** (1941) Les mysidacés des eaux Méditerranéennes de la France (spécialement de Banyuls) et des eaux de Monaco. *Bulletin de l'Institut Océanographique* 795, 1–46.
- Bacescu M.** (1954) *Mysidacea*. In *Fauna Republicii Populare Romine, Crustacea*, IV (3). Academia Republicii Populare Romine, Bucuresti, pp. 1–126.
- Cannon H.G. and Manton S.M.** (1927) On the feeding mechanism of a mysid crustacean: *Hemimysis lamornae*. *Transactions of the Royal Society of Edinburgh* 55, 219–253.
- Carola M., Coma R., Riera T. and Zabala M.** (1993) Fecal pellets collection as a method for assessing egesta of the marine cave-dwelling mysid *Hemimysis speluncola*. *Scientia Marina* 57, 51–63.
- Chinchilla M. and Comín F.A.** (1977) Contribució al coneixement dels crustacis del delta de l'Ebre. *Treballs de l'Institut Català d'Història Natural* 8, 119–144.

- Coll M., Piroddi C., Steenbeek J., Kaschner K. and Ben Rais Lasram F.** (2010) The biodiversity of the Mediterranean Sea: estimates, patterns, and threats. *PLoS ONE* 5, e11842. doi:10.1371/journal.pone.0011842.
- Colosi G.** (1929) I Misidacei del Golfo di Napoli. *Pubblicazioni della Stazione Zoologica di Napoli* 9, 405–441.
- Couch R.Q.** (1856) On Crustacea new to the British fauna. *Zoologist: a Monthly Journal of Natural History, Series 1* 14, 281–5288.
- Czerniavsky V.** (1882) *Monographia Mysidarum inprimis Imperii Rossici*. Fasc. 1,2. Trudy Sankt-Peterburgskago Obshchestwo Estestwoispytatelei, 12, 1–170, 13, 1–85.
- Foxon G.E.H.** (1940) The reactions of certain mysids to stimulation by light and gravity. *Journal of the Marine Biological Association of the United Kingdom* 24, 89–97.
- Haworth A.H.** (1825) XVII. A new binary arrangement of the brachyurous Crustacea. *Philosophical Magazine Series 1* 65, 105–106.
- Ketelaars H.A.M., Lambregts-van de Clundert F.E., Carpentier C.J., Wagenvoort A.J. and Hoogenboezem W.** (1999) Ecological effects of the mass occurrence of the Ponto-Caspian invader, *Hemimysis anomala* G.O. Sars, 1907 (Crustacea: Mysidacea), in a freshwater storage reservoir in the Netherlands, with notes on its autecology and new records. *Hydrobiologia* 394, 233–248.
- Ledoyer M.** (1989) Les Mysidacés (Crustacea) des grottes sous-marines obscures de Méditerranée nord-occidentale et du proche Atlantique (Portugal et Madère). *Marine Nature* 2, 39–62.
- Lejeusne C. and Chevaldonné P.** (2006). Brooding crustaceans in a highly fragmented habitat: the genetic structure of Mediterranean marine cave-dwelling mysid populations. *Molecular Ecology* 15, 4123–4140.
- Macquart-Moulin C.** (1965) Les Mysidacés benthoplanctoniques du Golfe de Marseille. *Recueil des Travaux de la Station Marine d'Endoume. Faculté des Sciences de Marseille* 38, 129–253.
- Manton S.M.** (1928) On the embryology of a mysid crustacean, *Hemimysis lamornae*. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 216, 363–463.
- Mauchline J.** (1971) The biology of *Praunus flexuosus* and *P. neglectus* (Crustacea, Mysidacea). *Journal of the Marine Biological Association of the United Kingdom* 51, 641–652.
- Minchin D. and Boelens R.** (2010) *Hemimysis anomala* is established in the Shannon River Basin District in Ireland. *Aquatic Invasions* 5, Supplement1, S71–S78.
- Norman A.M.** (1860) On an undescribed crustacean of the genus *Mysis*. *Annals and Magazine of Natural History, Series 3* 6, 429–431.
- Norman A.M.** (1892) On British Mysidae, a family of Crustacea Schizopoda. *Annals and Magazine of Natural History, Series 6* 10, 242–263.
- Remerie T., Bourgois T. and Vanreusel A.** (2005) Morphological differentiation between geographically separated populations of *Neomysis integer* and *Mesopodopsis slabberi* (Crustacea, Mysida). *Hydrobiologia* 549, 239–250.
- San Vicente C. and Munilla T.** (2000) Misidáceos suprabentónicos de las playas catalanas (Mediterráneo nordoccidental). *Orsis* 15, 45–55.
- Sars G.O.** (1879) Carcinologiske Bidrag til Norges fauna. I. *Monographi over de ved Norges Kyster forekommende Mysider. Pt. 3. Christiania, A.W. Brøgger*, 1–131.
- Scheiner S.M. and Callahan H.S.** (1993) Measuring natural selection on phenotypic plasticity. *Evolution* 53, 1704–1713.
- Suau P. and Vives F.** (1957) Sobre la presencia de *Gastrosaccus sanctus* (Van Beneden) en la desembocadura del rio Ebro. *Investigación Pesquera VIII*, 121–126.

- Tattersall W.M.** (1951) A review of the Mysidacea of the United States National Museum. *Bulletin of the United States National Museum* 201, 1–292.
- Tattersall W.M. and Tattersall O.S.** (1951) *The British Mysidacea*. London: The Ray Society, 460 pp.
- Williams A.B., Bowman T.E. and Damkaeri D.M.** (1974). Distribution, variation, and supplemental description of the opossum shrimp, *Neomysis americana* (Crustacea: Mysidacea). *Fishery Bulletin* 72, 835–842.
- Wittmann K.J.** (1992) Cyclomorphosis in the temperate zone Mysidacea: evidence and possible adaptive taxonomical significance. In Köhn J., Jones M.B. and Moffat A. (eds) *Taxonomy, biology and ecology of (Baltic) mysids (Mysidacea: Crustacea)*. *International Expert Conference, September 1991, Hiddensee, Germany*. Rostock University, pp. 25–32.
- Wittmann K.J.** (2001) Centennial changes in the near-shore mysid fauna of the Gulf of Naples (Mediterranean Sea), with description of *Heteromysis riedli* sp. nov. (Crustacea, Mysidacea). *Marine Ecology* 22, 85–110.
- Wittmann K.J. and Ariani A.P.** (2009) Reappraisal and range extension of non-indigenous Mysidae (Crustacea, Mysida) in continental and coastal waters of eastern France. *Biological Invasions* 11, 401–407.
- and
- Zimmer C.** (1909) Die nordischen Schizopoden. Pages. In Brandt K. and Apstein C. (eds) *Nordisches Plankton, Zoologischer Teil, Dritter Band: Crustacea*. Kiel and Leipzig: Lipsius und Tischer, pp. 1–178.
- Correspondence should be addressed to:**
G. Guerao
IRTA, Unitat Operativa de Cultius Experimentals
43540-Sant Carles de la Ràpita, Tarragona, Spain
email: guillermo.guerao@irta.cat