



Contents lists available at ScienceDirect

Deep-Sea Research II

journal homepage: www.elsevier.com/locate/dsr2

DNA barcoding reveals new insights into the diversity of Antarctic species of *Orchomene sensu lato* (Crustacea: Amphipoda: Lysianassoidea)

C. Havermans^{a,b,*}, Z.T. Nagy^a, G. Sonet^a, C. De Broyer^a, P. Martin^a

^a Royal Belgian Institute of Natural Sciences, Rue Vautier 29, B-1000 Brussels, Belgium

^b Laboratory of Marine Biology, Place Croix du Sud 3, Catholic University of Louvain, B-1348 Louvain-la-Neuve, Belgium

ARTICLE INFO

Keywords:

DNA barcoding
Genetic diversity
Species
Amphipoda
Lysianassoidea
Southern Ocean

ABSTRACT

Recent molecular analyses revealed that several so-called “circum-Antarctic” benthic crustacean species appeared to be complexes of cryptic species with restricted distributions. In this study we used a DNA barcoding approach based on mitochondrial cytochrome oxidase I gene sequences in order to detect possible cryptic diversity and to test the circumpolarity of some lysianassoid species. The orchomenid genus complex consists of the genera *Abyssorchomene*, *Falklandia*, *Orchomenella*, *Orchomenyx* and *Pseudorchomene*. Species of this genus complex are found throughout the Southern Ocean and show a high species richness and level of endemism. In the majority of the studied species, a genetic homogeneity was found even among specimens from remote sampling sites, which indicates a possible circum-Antarctic and eurybathic distribution. In four investigated species (*Orchomenella* (*Orchomenopsis*) *acanthurus*, *Orchomenella* (*Orchomenopsis*) *cavimanus*, *Orchomenella* (*Orchomenella*) *franklini* and *Orchomenella* (*Orchomenella*) *pinguides*), genetically divergent lineages and possible cryptic taxa were revealed. After a detailed morphological analysis, *O. (O.) pinguides* appeared to be composed of two distinct species, formerly synonymized under *O. (O.) pinguides*. The different genetic patterns observed in these orchomenid species might be explained by the evolutionary histories undergone by these species and by their different dispersal and gene flow capacities.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

According to most estimations on global biodiversity, the majority of species living on this planet are currently undescribed (Novotny et al., 2002; Blaxter, 2003, 2004; Bouchet, 2006). Aiming to have a “complete” account of all living organisms would require more work than the present manpower and technology can handle. Moreover, in the context of the current biodiversity crisis and the declining number of taxonomists, several authors suggest the use of DNA barcoding to accelerate and simplify species identification (Hebert et al., 2003a,b; Blaxter, 2004; Janzen et al., 2005; Schander and Willassen, 2005; Schindel and Miller, 2005). DNA barcoding uses a short DNA sequence as the standard genetic marker for species identification (a ca. 648 bp segment near the 5' end of the mitochondrial cytochrome oxidase I gene,

COI, for animals). The barcode sequence from each unknown specimen is compared with a reference library of sequences derived from specimens of known identity (Hajibabaei et al., 2007). This sequence library is currently being established. This approach speeds up species identification and also facilitates the discovery of undescribed species (Witt et al., 2003). The efficiency of a barcoding marker in species delimitation depends on the separation between intra- and interspecific divergences (Hebert et al., 2003a,b; Meyer and Paulay, 2005; Waugh, 2007). In accordance with the biological species definition, intraspecific genetic distances have to be generally smaller (mostly by an order of magnitude) than interspecific genetic distances. This provides the basis for species delimitation (Waugh, 2007; Meier et al., 2008). In several animal taxa, the effectiveness of this approach has been confirmed, such as in birds (Hebert et al., 2004b), fish (Ward et al., 2005), molluscs (Meyer and Paulay, 2005), spiders (Barrett and Hebert, 2005) and several groups of butterflies (Hebert et al., 2004a; Janzen et al., 2005; Hajibabaei et al., 2006). In poorly studied groups, DNA barcoding can be performed prior to “conventional”, morphology-based taxonomic studies in order to quickly sort specimens into genetically divergent groups (Hajibabaei et al., 2007). However, the DNA barcoding approach is not without controversy when it is considered as a tool for classification and identification (e.g., Lipscomb et al., 2003; Moritz and Cicero, 2004; Will and Rubinoff, 2004). It has raised some debates about traditional

* Corresponding author at: Department of Freshwater Biology, Royal Belgian Institute of Natural Sciences, Rue Vautier 29, B-1000 Brussels, Belgium.
Tel.: +32 2 627 42 78; fax: +32 2 627 42 77.

E-mail addresses: chavermans@naturalsciences.be (C. Havermans), zoltan-tamas.nagy@naturalsciences.be (Z.T. Nagy), gontran.sonet@naturalsciences.be (G. Sonet), claude.debroyer@naturalsciences.be (C. De Broyer), patrick.martin@naturalsciences.be (P. Martin).

taxonomy becoming extinct and being replaced by DNA sequencing. However, DNA barcoding should not be considered as a substitute for conventional taxonomy; its principal utility is as a searchable label, by linking barcodes to fully described voucher specimens (Waugh, 2007). The coupling of a detailed morphological and ecological investigation with the barcode results is critical for species descriptions. Nevertheless, DNA barcoding has its limitations: its accuracy seems to depend on the taxonomic knowledge and the sample coverage of the group (e.g., Meyer and Paulay, 2005). Additionally, the phenomena of incomplete lineage sorting, genetic introgression, pseudogenes (e.g., Buhay, 2009) or bacterial infections (*Wolbachia*, e.g., Whitworth et al., 2007) can make species identification inadequate with this tool.

The Southern Ocean is considered as a hotspot of biodiversity and endemism for several orders of peracarid crustaceans (Malacostraca), which have undergone spectacular adaptive radiations (Watling and Thurston, 1989; Brandt, 1999, 2005; Lörz and Brandt, 2004; Lörz and Held, 2004). Peracarids comprise about 1500 strictly Antarctic species and, among them, amphipods represent the most speciose group with more than 815 gammaridean and corophiidean species recorded in the Southern Ocean *sensu lato* (De Broyer et al., 1999, 2003, 2007). The superfamily Lysianassoidea is one of the most dominant gammaridean amphipod groups in Antarctic waters, both in number of species and in abundance (Arnaud et al., 1986; De Broyer et al., 2001).

Unlike Antarctic benthic communities living in shallow water, little is known about the biodiversity of the Antarctic deep-sea region where many collected invertebrate species are new to science (Brandt et al., 2007). Moreover, species counts for the fauna of the Southern Ocean are suspected to be underestimated. Indeed, many Antarctic marine benthic invertebrates are currently considered to have a circum-Antarctic and/or eurybathic distribution (Arntz et al., 1994). The circum-Antarctic distribution can be explained by similar environmental conditions in the sea around the continent, as well as by the circumpolar current systems (Arntz et al., 2005). The high degree of eurybathy is considered as an evolutionary adaptation to the oscillation of the ice cap extension during the Antarctic glacial and interglacial cycles. Ice extensions and retreats could have been followed by a migration of taxa up and down the Antarctic continental shelf and slope (Brey et al., 1996). However, recent molecular analyses revealed that several of these species represent in fact complexes of morphologically similar (cryptic) species showing restricted distribution ranges. This is the case for several Antarctic organisms: isopods (Held, 2003; Held and Wägele, 2005; Raupach and Wägele, 2006; Raupach et al., 2007; Brökeland and Raupach, 2008), molluscs (Beaumont and Wei, 1991; Page and Linse, 2002; Allcock et al., 2004; Strugnell et al., 2008), crinoids (Wilson et al., 2007), pycnogonids (Mahon et al., 2008) and fish (Bernardi and Goswami, 1997; Smith et al., 2008).

The lysianassoid genus *Orchomene sensu lato* represents a good model for biodiversity studies due to its (relative) species richness, high degree of endemism, its abundance and important role in the Southern Ocean, and the presence at both shallow and abyssal depths. Following the most recent systematic classification (De Broyer et al., 2007), this orchomenid genus complex includes the genera *Abyssorchomene* De Broyer, 1984, *Orchomenella* G.O. Sars, 1895 (including the subgenera *Orchomenella* and *Orchomenopsis*), *Orchomenyx* De Broyer, 1984 and *Pseudorchomene* Schellenberg, 1926. A recent molecular phylogenetic study also suggested the inclusion of the monotypic genus *Falklandia* De Broyer, 1985 within this genus complex (Havermans et al., 2010). The genera *Falklandia*, *Orchomenyx* and *Pseudorchomene* are endemic to the Southern Ocean. Although two genera, *Orchomenella* and *Abyssorchomene*, may be considered as

cosmopolitan (Barnard and Karaman, 1991), they also comprise some species endemic to the Southern Ocean.

The phylogeny of the group was recently investigated (Havermans et al., 2010) and it was shown that the molecular phylogeny does not correspond to the morphological classification at the genus level. Several (sub)genera (*Abyssorchomene*, *Orchomenella*, *Orchomenopsis*) appeared to be non-monophyletic and some diagnostic characters used in this complex of genera are likely a result of convergent evolution. The scope of the current paper does not focus on this issue but rather focuses on the issue of species delimitation within this group. Our aim is to test whether the COI gene is an appropriate barcoding marker for these taxa. Our previous study showed that previously proposed taxonomic subdivisions should be revised and these taxa remain difficult to identify for the non-expert. These taxa, with a confuse taxonomy, represent an interesting case to test the validity of the barcoding approach. Finally, the circumpolarity and species boundaries will be investigated using genetic and biogeographic data in several orchomenid species such as *Orchomenella* (*Orchomenopsis*) *cavimanus* (Stebbing, 1888) and *Abyssorchomene* *scotianensis* (Andres, 1983), which were characterized so far by a circum-Antarctic and eurybathic distribution (De Broyer et al., 2007).

2. Material and methods

During recent expeditions with the R/V “Polarstern”, amphipod material was collected from the Magellanic region, the Scotia Sea, the eastern shelf of the Antarctic Peninsula, the Weddell Abyssal Plain, the Eastern Weddell Sea and Bouvet Island (ANTARKTIS XV-3, De Broyer et al., 1999; ANTARKTIS XIX-5, De Broyer et al., 2003; ANTARKTIS XXI-2, ANDEEP I, II, III, De Broyer et al., 2003, 2006; ANTARKTIS XXIII-8, d’Udekem d’Acoz and Robert, 2008). Additional samples from the Ross Sea (BIOROSS Cruise) and from King George Island (South Shetland Islands) were provided by the National Institute of Water and Atmospheric Research (NIWA, New Zealand) and the Polish Antarctic IPY Expedition 2007, respectively. Agassiz and bottom trawls, dredges, epibenthic sleds, grabs, multi-box corers and baited traps were used to collect amphipods. Samples were fixed in 96% or absolute ethanol.

The molecular analysis included 121 specimens belonging to ca. 19 species, identified by a preliminary morphological analysis. Specimens of the lysianassoid genus *Ambasiopsis* were used as outgroup. Genomic DNA was isolated from the sixth pereopod using the QIAamp DNA Mini Kit (Qiagen). Amplification of the COI marker was carried out by polymerase chain reaction using the universal primers LCO1490 and HCO2198 (Folmer et al., 1994). Purified PCR products were sequenced bidirectionally using an ABI 3130xl capillary DNA sequencer (Applied Biosystems). Detailed information on specimens used in this study is given in Table 1 and sequences were deposited in GenBank.

Alignments were made manually (alignments are available from the first author upon request). A neighbour-joining tree (Saitou and Nei, 1987) was estimated using MEGA 4 (Tamura et al., 2007) and sequence divergences were calculated using the Kimura 2-parameter (K2P) distance model (Kimura, 1980), the best metric system when distances are low (Nei and Kumar, 2000) (see supplementary material available at doi:10.1016/j.dsr2.2010.09.028). Branch support was evaluated using non-parametric bootstrapping (number of replicates was 2000). Frequency distribution histograms of pairwise inter- and intraspecific distances were calculated with R (version 2.7.0) using the APE package (Paradis et al., 2004) and plotted using geneplotter, graphics related functions for Bioconductor (Gentleman et al., 2004). For further estimations on divergence, TaxonDNA v.1.5a12 (Meier et al., 2006) was used.

Table 1
Data on specimens used for this study. Abbreviations: AP–Antarctic Peninsula, BB–Burdwood Bank, BI–Bouvet Island, JI–Joinville Island, KGI–King George Island, LB–Larsen B area, PAE–Polish Antarctic IPY Expedition 2007, SS–Scotia Sea, RS–Ross Sea, WS–Weddell Sea, n.d.–no data.

Species	Individual codes	Expedition	Station number	Locality	Longitude/latitude	Depth (m)	Accession no.
Uristidae							
<i>Abyssorchomene charcoti</i> (Chevreux, 1912)	AC-SS205	ANDEEP I&II	128	SS	62°43'S 55°30' W	205	GU109230
<i>Abyssorchomene charcoti</i> (Chevreux, 1912)	AC-0510075	ANDEEP I&II	128	SS	62°43'S 55°30' W	205	HM053979
<i>Abyssorchomene charcoti</i> (Chevreux, 1912)	AC-1110071	ANDEEP I&II	058	JI	60°59'S 55°43' W	113	HM053980
<i>Abyssorchomene charcoti</i> (Chevreux, 1912)	AC-1110072	ANDEEP I&II	127	JI	62°42'S 55°22' W	295	HM053981
<i>Abyssorchomene charcoti</i> (Chevreux, 1912)	AC-1403073	ANT XXI-2	103	WS	70°49'S 10°39'W	387	HM053982
<i>Abyssorchomene chevreuxi</i> (Stebbing, 1906)	AC-WS4700	ANDEEP III	110	WS	65°00'S 43°02'W	4700	GU109248
<i>Abyssorchomene chevreuxi</i> (Stebbing, 1906)	AC-P3076	ANDEEP I&II	131-1	AP	65°17'S 51°35'W	3076	GU109229
<i>Abyssorchomene chevreuxi</i> (Stebbing, 1906)	AC-2609074	ANDEEP I&II	131-1	AP	65°17'S 51°35'W	3076	HM053983
<i>Abyssorchomene chevreuxi</i> (Stebbing, 1906)	AC-0810074	ANDEEP III	81	WS	70°31'S 14°34'W	4409	HM053984
<i>Abyssorchomene chevreuxi</i> (Stebbing, 1906)	AC-26090710	ANDEEP I&II	131-1	AP	65°17'S 51°35'W	3076	HM053985
<i>Abyssorchomene nodimanus</i> (Walker, 1903)	AN-WS393	ANT XXI-2	167	WS	70°48'S 10°39'W	393	GU109241
<i>Abyssorchomene nodimanus</i> (Walker, 1903)	AN-WS387	ANT XXI-2	103	WS	70°49'S 10°39'W	387	GU109260
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-0304076	ANDEEP III	150	SS	61°48'S 47°27'W	1943	HM053986
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-SS1943	ANDEEP III	150	SS	61°48'S 47°27'W	1943	GU109255
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-SS270	ANT XIX-5	191	SS	57°41'S 26°24'W	270	GU109258
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-08100719	ANT XIX-5	191	SS	57°41'S 26°24'W	270	HM053987
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-08100722	ANT XIX-5	191	SS	57°41'S 26°24'W	270	HM053988
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-0506081	ANT XIX-5	191	SS	57°41'S 26°24'W	270	HM053989
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-2409073	ANT XXI-2	14	BI	54°37'S 03°06'E	515	HM053990
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-2109078	ANT XXI-2	14	BI	54°37'S 03°06'E	515	HM053991
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-0510071	ANT XXI-2	14	BI	54°37'S 03°06'E	515	HM053992
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-LB383	ANT XXIII-8	698-1	LB	65°59'S 60°24'W	383	GU109233
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-J12	ANT XXIII-8	684-1	SS	62°57'S 57°57'W	822	HM053993
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-G10	ANT XXIII-8	620	JI	60°56'S 55°49'W	334	HM053994
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-1110076	ANDEEP I&II	114	SS	61°44'S 60°45'W	2889	HM053995
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-08100721	ANT XIX-5	194	SS	57°40'S 26°25'W	308	HM053996
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-2609072	ANDEEP I&II	083	SS	61°07'S 56°08'W	349	HM053997
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-0810072	ANDEEP I&II	083	SS	61°07'S 56°08'W	349	HM053998
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-A1	ANT XXIII-8	n.d.	n.d.	n.d.	n.d.	HM053999
<i>Abyssorchomene plebs</i> (Hurley, 1965)	AP-31100710	ANT XV-3	T13	WS	70°29'S 07°57'W	550	HM054000
<i>Abyssorchomene rossi</i> (Walker, 1903)	AR-1010076	ANT XXI-2	288	WS	72°47'S 19°29'W	847	HM054001
<i>Abyssorchomene rossi</i> (Walker, 1903)	AR-3110078	ANT XV-3	T13	WS	70°29'S 07°57'W	550	HM054002
<i>Abyssorchomene rossi</i> (Walker, 1903)	AR-I19	ANT XXIII-8	698-1	LB	65°59'S 60°24'W	383	HM054003
<i>Abyssorchomene scotianensis</i> (Andres, 1983)	AS-SS3408	ANDEEP III	142	SS	62°11'S 49°29'W	3408	GU109242
<i>Abyssorchomene scotianensis</i> (Andres, 1983)	AS-2210075	ANDEEP I&II	131-1	AP	65°17'S 51°35'W	3076	HM054004
<i>Abyssorchomene scotianensis</i> (Andres, 1983)	AS-2609079	ANDEEP I&II	131-1	AP	65°17'S 51°35'W	3076	HM054005
<i>Abyssorchomene scotianensis</i> (Andres, 1983)	AS-2210072	ANDEEP I&II	131-1	AP	65°17'S 51°35'W	3076	HM054006
<i>Abyssorchomene scotianensis</i> (Andres, 1983)	AS-P3076	ANDEEP I&II	131-1	AP	65°17'S 51°35'W	3076	GU109240
<i>Abyssorchomene scotianensis</i> (Andres, 1983)	AS-05100710	ANDEEP I&II	131-1	AP	65°17'S 51°35'W	3076	HM054007
<i>Abyssorchomene scotianensis</i> (Andres, 1983)	AS-2210071	ANDEEP I&II	131-1	AP	65°17'S 51°35'W	3076	HM054008
<i>Abyssorchomene scotianensis</i> (Andres, 1983)	AS-2210074	ANDEEP I&II	131-1	AP	65°17'S 51°35'W	3076	HM054009
<i>Abyssorchomene scotianensis</i> (Andres, 1983)	AS-22100919	ANDEEP III	78	WS	71°09'S 14°00'W	2166	HM054010
<i>Abyssorchomene scotianensis</i> (Andres, 1983)	AS-0810073	ANDEEP III	80	WS	70°39'S 14°43'W	3088	HM054011
<i>Abyssorchomene scotianensis</i> (Andres, 1983)	AS-1110077	ANDEEP I&II	114	SS	61°44'S 60°45'W	2889	HM054012
<i>Abyssorchomene</i> sp. 1	An-SS3406	ANDEEP III	142	SS	62°11'S 49°29'W	3406	GU109239
<i>Abyssorchomene</i> sp. 1	An-0810076	ANDEEP III	80	WS	70°39'S 14°43'W	3088	HM054013
<i>Abyssorchomene</i> sp. 1	An-WS3088	ANDEEP III	80	WS	70°39'S 14°43'W	3088	GU109236
<i>Abyssorchomene</i> sp. 1	An-0810078	ANDEEP III	81	WS	70°31'S 14°34'W	4409	HM054014
<i>Abyssorchomene</i> sp. 1	An-08100710	ANDEEP III	78	WS	71°09'S 14°00'W	2166	HM054015

Table 1 (continued)

Species	Individual codes	Expedition	Station number	Locality	Longitude/latitude	Depth (m)	Accession no.
<i>Abyssorchomene</i> sp. 1	An-0707088	ANT XXIII-8	706-7	LB	65°26'S 61°26'W	828	HM054016
<i>Abyssorchomene</i> sp. 1	An-07070810	ANT XXIII-8	705-1	LB	65°33'S 61°37'W	310	HM054017
<i>Abyssorchomene</i> sp. 1	20814	Tangaroa (NIWA)	TAN0402-257	RS	66°12'S 162°26'E	1395	HM054018
<i>Abyssorchomene</i> sp. 2	An2-1010074	ANT XXI-2	14	BI	54°37'S 03°06'E	515	HM054019
<i>Abyssorchomene</i> sp. 2	An2-2202072	ANT XXI-2	14	BI	54°37'S 03°06'E	515	HM054020
Lysianassidae							
<i>Falklandia reducta</i> (Schellenberg, 1931)	FR-SS285	ANT XIX-5	252	SS	61°23'S 55°26' W	285	GU109256
<i>Orchomenella (Orchomenella) franklini</i> (Walker, 1903)	OF-SS259-1	ANT XXIII-8	614-15	JI	60°52'S 55°27'W	259	GU109226
<i>Orchomenella (Orchomenella) franklini</i> (Walker, 1903)	OF-0707082	ANT XXIII-8	614-15	JI	60°52'S 55°27'W	259	HM054021
<i>Orchomenella (Orchomenella) franklini</i> (Walker, 1903)	OF-SS259-3	ANT XXIII-8	614-15	JI	60°52'S 55°27'W	259	GU109235
<i>Orchomenella (Orchomenella) franklini</i> (Walker, 1903)	OF-0707084	ANT XXIII-8	614-15	JI	60°52'S 55°27'W	259	HM054022
<i>Orchomenella (Orchomenella) franklini</i> (Walker, 1903)	OF-0707087	ANT XXIII-8	n.d.	n.d.	n.d.	n.d.	HM054023
<i>Orchomenella (Orchomenella) franklini</i> (Walker, 1903)	OF-D1	ANT XXIII-8	614-3	JI	60°52'S 55°27'W	259	HM054024
<i>Orchomenella (Orchomenella) franklini</i> (Walker, 1903)	OF-I16	ANT XXIII-8	654-3	JI	61°22'S 56°03'W	363	HM054025
<i>Orchomenella (Orchomenella) franklini</i> (Walker, 1903)	OF-2610071	ANDEEP I&II	133-3	AP	65°19'S 54°14'W	1120	HM054026
<i>Orchomenella (Orchomenella) franklini</i> (Walker, 1903)	OF-24100720	ANDEEP I&II	133-3	AP	65°19'S 54°14'W	1120	HM054027
<i>Orchomenella (Orchomenella) franklini</i> (Walker, 1903)	OF-1010075	ANT XXI-2	245	WS	70°56'S 10°32'W	337	HM054028
<i>Orchomenella (Orchomenella) pinguides</i> (Walker, 1903)	OP-WS387	ANT XXI-2	103	WS	70°49'S 10°39'W	387	GU109247
<i>Orchomenella (Orchomenella) pinguides</i> (Walker, 1903)	OP-WS395	ANT XXI-2	108	WS	70°48'S 10°39'W	395	GU109259
<i>Orchomenella (Orchomenella) pinguides</i> (Walker, 1903)	OP-WS175	ANT XXI-2	39	WS	71°06'S 11°32'W	175	GU109237
<i>Orchomenella (Orchomenella) pinguides</i> (Walker, 1903)	OP-24100714	ANDEEP I&II	133-3	AP	65°19'S 54°14'W	1120	HM054029
<i>Orchomenella (Orchomenella) pinguides</i> (Walker, 1903)	20810	Tangaroa (NIWA)	TAN0402-206	RS	71°09'S 171°02'E	975	HM054030
<i>Orchomenella (Orchomenella) pinguides</i> (Walker, 1903)	20807	Tangaroa (NIWA)	TAN0402-103	RS	71°14'S 170°42'E	555	HM054031
<i>Orchomenella (Orchomenopsis) acanthurus</i> (Schellenberg, 1931)	OA-RS252	Tangaroa (NIWA)	TAN0402-133	RS	71°38'S 170°13'E	252	GU109263
<i>Orchomenella (Orchomenopsis) acanthurus</i> (Schellenberg, 1931)	20845	Tangaroa (NIWA)	TAN0402-134	RS	71°38'S 170°09'E	65	HM054032
<i>Orchomenella (Orchomenopsis) acanthurus</i> (Schellenberg, 1931)	OA-P137	ANT XXIII-8	605-3	AP	61°20'S 55°31'W	137	GU109266
<i>Orchomenella (Orchomenopsis) acanthurus</i> (Schellenberg, 1931)	OA-WS284	ANT XXI-2	132	WS	70°56'S 10°31'W	284	GU109225
<i>Orchomenella (Orchomenopsis) acanthurus</i> (Schellenberg, 1931)	OA-23040818	ANT XXI-2	146	WS	70°56'S 10°47'W	404	HM054033
<i>Orchomenella (Orchomenopsis) acanthurus</i> (Schellenberg, 1931)	OA-0707085	ANT XXIII-8	605	AP	61°20'S 55°29'W	151	HM054034
<i>Orchomenella (Orchomenopsis) acanthurus</i> (Schellenberg, 1931)	OA-0707086	ANT XXIII-8	605	AP	61°20'S 55°29'W	151	HM054035
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-3110073	ANDEEP I&II	042-2	SS	59°40'S 57°35'W	3683	HM054036
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-SS349	ANDEEP I&II	083	SS	61°07'S 56°08'W	349	GU109250
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-SS349-1	ANDEEP I&II	083	SS	61°07'S 56°08'W	349	GU109251
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-SS349-4	ANDEEP I&II	083	SS	61°07'S 56°08'W	349	GU109252
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-1909073	ANT XIX-5	162	SS	53°25'S 42°40'W	293	HM054037
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-2109075	ANT XIX-5	162	SS	53°25'S 42°40'W	293	HM054038
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-SS293	ANT XIX-5	162	SS	53°25'S 42°40'W	293	GU109264
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-WS515	ANT XIX-5	289	WS	72°49'S 19°30'W	515	GU109257
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-SS413	ANT XIX-5	157	BB	54°32'S 55°55'W	413	GU109262
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-SS1943	ANDEEP III	150	SS	61°48'S 47°27'W	1943	GU109254
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-0603075	ANT XV-3	T13	WS	70°29'S 07°57'W	550	HM054039
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-WS387	ANT XXI-2	103	WS	70°49'S 10°39'W	387	GU109244
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-WS847	ANT XXI-2	288	WS	72°47'S 19°29'W	847	GU109261
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-WS1017-5	ANDEEP III	74	WS	71°18'S 13°56'W	1017	GU109243
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-WS1017-4	ANDEEP III	74	WS	71°18'S 13°56'W	1017	GU109249
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-I2	ANT XXIII-8	683-1	AP	62°57'S 57°57'W	839	HM054040
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-0304075	ANDEEP III	80	WS	70°39'S 14°43'W	3088	HM054041
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	20829	Tangaroa (NIWA)	TAN0402-14	RS	71°43'S 171°45'E	451	HM054042
<i>Orchomenella (Orchomenopsis) cavimanus</i> (Stebbing, 1888)	OC-0707089	ANT XXIII-8	635	SS	60°56'S 55°55'W	231	HM054043
<i>Orchomenyx macronyx</i> (Chevreux, 1905)	OM-JI151	ANT XXIII-8	605-1	JI	61°20'S 55°29'W	151	GU109231
<i>Orchomenyx macronyx</i> (Chevreux, 1905)	OM-JI161	ANT XXIII-8	685-1	JI	62°33'S 55°41'W	161	GU109228
<i>Orchomenyx schellenbergi</i> (Thurston, 1972)	OS-23040820	PAE	2-07	KGI	58°27'S 62°09'W	210	HM054044
<i>Orchomenyx schellenbergi</i> (Thurston, 1972)	OS-KGI210	PAE	2-07	KGI	58°27'S 62°09'W	210	GU109265
<i>Orchomenyx schellenbergi</i> (Thurston, 1972)	OS-2304083	PAE	2-07	KGI	58°27'S 62°09'W	210	HM054045
<i>Orchomenyx tabarini</i> (Thurston, 1972)	OT-P211	ANT XXIII-8	689-5	AP	62°27'S 55°25'W	211	GU109227

<i>Pseudorchomene coatsi</i> (Chilton, 1912)	PC-1809076	ANDEEP I&II	083	SS	31°07'S 56°08'W	349	HM054046
<i>Pseudorchomene coatsi</i> (Chilton, 1912)	PC-05100712	ANDEEP I&II	083	SS	31°07'S 56°08'W	349	HM054047
<i>Pseudorchomene coatsi</i> (Chilton, 1912)	PC-SS349	ANDEEP I&II	083	SS	31°07'S 56°08'W	349	GU109245
<i>Pseudorchomene coatsi</i> (Chilton, 1912)	PC-2609071	ANDEEP I&II	083	SS	31°07'S 56°08'W	349	HM054048
<i>Pseudorchomene coatsi</i> (Chilton, 1912)	PC-SS2889-4	ANDEEP I&II	114	SS	61°44'S 60°45'W	2889	GU109232
<i>Pseudorchomene coatsi</i> (Chilton, 1912)	PC-SS2889-5	ANDEEP I&II	114	SS	61°44'S 60°45'W	2889	GU109234
<i>Pseudorchomene coatsi</i> (Chilton, 1912)	PC-2210079	ANT XV-3	T13	WS	70°29'S 07°57'W	550	HM054049
<i>Pseudorchomene coatsi</i> (Chilton, 1912)	PC-22100710	ANT XV-3	T13	WS	70°29'S 07°57'W	550	HM054050
<i>Pseudorchomene coatsi</i> (Chilton, 1912)	PC-2609076	ANT XXI-2	240	WS	70°48'S 10°39'W	406	HM054051
<i>Pseudorchomene coatsi</i> (Chilton, 1912)	PC-08100715	ANT XXI-2	103	WS	70°49'S 10°39'W	387	HM054052
<i>Pseudorchomene coatsi</i> (Chilton, 1912)	PC-1909075	ANT XIX-5	261	SS	72°16'S 58°15'W	723	HM054053
<i>Pseudorchomene coatsi</i> (Chilton, 1912)	Pn-WS847	ANT XXI-2	288	SS	72°47'S 19°29'W	847	GU109238
<i>Pseudorchomene</i> sp.	Pn-0304072	ANDEEP III	150	SS	61°48'S 47°27'W	1943	HM054054
<i>Pseudorchomene</i> sp.	Pn-0510077	ANDEEP III	150	SS	61°48'S 47°27'W	1943	HM054055
<i>Pseudorchomene</i> sp.	Pn-SS1943	ANDEEP III	150	SS	61°48'S 47°27'W	1943	GU109253
Adeliellid group							
<i>Ambasiopsis</i> sp.	As-08100711	ANT XXI-2	19	BI	54°31'S 03°14'E	260	HM054056
<i>Ambasiopsis</i> sp.	As-08100712	ANT XXI-2	19	BI	54°31'S 03°14'E	260	HM054057
<i>Ambasiopsis</i> sp.	As-BI260	ANT XXI-2	19	BI	54°31'S 03°14'E	260	GU109246

3. Results

The alignment of COI sequences included 658 positions, comprising 272 variable sites with the outgroup included, 247 variable sites without considering the outgroup. The amino acid translation. The mean base frequencies were A, 0.24; C, 0.13; G, 0.21; T, 0.42. The transition/transversion ratio was 1.566.

3.1. Intraspecific divergence

The mean K2P divergence in the intraspecific pairwise comparisons is 1.86% for all orchomenid species. Distinct intraspecific divergence patterns could be observed within the different species. Most of the species showed intraspecific pairwise distances lower than 2.4%, except for four species: *Orchomenella* (*Orchomenella*) *pinguides*, *Orchomenella* (*Orchomenella*) *franklini*, *O. (O.) cavimanus* and *Orchomenella* (*Orchomenopsis*) *acanthurus*. These higher divergence values may be due to unrecognized cryptic species and may thus not represent intraspecific divergences. In *O. (O.) cavimanus*, a gradient of intraspecific divergences could be observed, varying from 0% to 10.6%. When these four putative species complexes are not included, the mean K2P intraspecific divergence becomes 0.4%. In several species (e.g. *Abyssororchomene plebs*, *Abyssororchomene* sp. 1, *Pseudorchomene coatsi*), very low genetic divergences could be observed. In *Abyssororchomene* sp. 1, a mean intraspecific variation of 0.7% exists between specimens from the Scotia Sea (3406 m depth), the Antarctic Peninsula (Larsen B, 828 and 310 m), the eastern Weddell Sea (4409 m) and the Ross Sea (1395 m) (Fig. 1). *A. plebs* showed a mean K2P distance of 0.2% between specimens of the Antarctic Peninsula, the Scotia Sea, the eastern Weddell Sea, the Atlantic sector with Bouvet Island, as well as between specimens from shelf (270 m) and abyssal depths (2889 m) in the Scotia Sea (Fig. 1). Specimens of *P. coatsi* from the continental shelf (350 m) and from abyssal depths (2889 m) also show low genetic distances with a mean K2P distance of 0.2%.

3.2. Interspecific divergence

The mean interspecific K2P divergence between species (except the four potential species complexes) is 14.5%, ranging from 6.3% (between *P. coatsi* and *Pseudorchomene* sp.) to 20.1% (between *Abyssororchomene chevreuxi* and *O. (O.) acanthurus*). The frequency distribution of pairwise K2P distances within and between well-defined orchomenid species (without the putative species complexes) is shown in Fig. 2. Interspecific divergence exceeds intraspecific divergence to such an extent that a ‘‘gap’’ can be observed. This gap range is the interval between the highest intraspecific and the lowest interspecific distances (Astrin et al., 2006; Meier et al., 2008). In our case, the gap size is about 3.9%.

3.3. Species delimitation based on the neighbour-joining tree

The neighbour-joining analysis (Fig. 1) shows that conspecifics based on morphological identification always group together and thus confirms the monophyly of all species investigated by multiple specimens. This analysis also revealed clusters corresponding to undescribed species. These appeared to be distinct from known species by a detailed morphological analysis. The species *Abyssororchomene* sp. 1 and *Abyssororchomene* sp. 2 are separated from their related species by genetic distances in the range of the formerly defined interspecific distances. The mean divergence between *A.* sp. 1 and *A. chevreuxi* is of 10.2% and between *A.* sp. 2 and *A. chevreuxi* of 14.0%. Another undescribed species, *Pseudorchomene* sp., can be distinguished as a sister species of *P. coatsi* from both genetic and morphological points of view.




Fig. 1. Neighbour-joining tree of COI sequences based on K2P distances. Bootstrap supports (2000 replications) are shown on the branches. Clusters with low genetic divergences are collapsed (number of studied representatives are indicated in parentheses). In cases where species complexes of the genus *Orchomenella* were found, the locality is indicated for each specimen. Within *Orchomenella*, the subgeneric assignment was used in the figure. *Orchomenella* (*Orchomenella*) *pinguides* is divided into two clusters, A and B, which appeared to be distinct based on a morphological analysis. In addition, locality data of two *Abyssorchomene* species are also represented on maps, with the number of specimens for each location in parentheses.

Although distances between these two species (between 6.3% and 7.2%) are in the lower range of the interspecific distances, they are significantly higher than the highest intraspecific distance (2.4%).

3.4. Cryptic species

Within *O. (O.) pinguides*, *Orchomenella* (*O.*) *franklini*, *O. (O.) acanthurus* as well as *O. (O.) cavimanus*, we observed several clusters supported by high bootstrap values (Fig. 1). The frequency distributions of pairwise K2P distances in these species complexes are presented in Fig. 3. In *O. (O.) acanthurus*, distances range from 0% to 15.7%, which can be separated in three blocks

ranging from 0% to 0.6%, from 5.8% to 8.4% and from 12.3% to 15.5%. In *O. (O.) pinguides*, distances vary from 0.2% to 7.9%. In *O. (O.) franklini* distances range from 0.2% to 3.5% and from 8.4% to 10.1%. These distances clearly indicate that some specimens are separated from each other by genetic distances in the range of interspecific distances. In the fourth case, *O. (O.) cavimanus*, genetic K2P distances vary from 0% to 10.6%. Without the most divergent cluster of *O. (O.) cavimanus* (i.e. the uppermost one in Fig. 1), K2P distances decrease to 0.3–5.6%.

In addition, representatives of these *Orchomenella* species occur in (partial) sympatry. For example, in *O. (O.) franklini*, specimens coming from the same sample locations at Joinville Island pop up in clusters separated by high genetic distances




Fig. 2. Frequency distribution of pairwise K2P distances of "well-defined" Antarctic orchomenid species (fourteen ingroup species are included and putative species complexes are excluded).

(8.8–10.5%) (Fig. 1 and Table 1). In *O. (O.) acanthurus*, specimens coming from the same site at the Antarctic Peninsula appear in two clusters separated by distances of more than 15%. In *O. (O.) cavimanus*, a gradient of genetic divergences could be observed between one specimen of the Magellanic region (Burdwood Bank), specimens of the Scotia Sea, the Antarctic Peninsula, the Eastern Weddell Sea and a specimen of the Ross Sea. Specimens of the same sample locations were found scattered within the species cluster and no geographically related clusters could be observed (Fig. 1 and Table 1).

O. (O.) pinguides can be divided into at least two clusters (A and B, see Fig. 1), each comprising three specimens and separated by distances higher than 7%. Cluster A comprises specimens from the Ross Sea and the Antarctic Peninsula while the second includes specimens from the Eastern Weddell Sea. Within cluster B, one specimen is separated by a distance of more than 5%, while occurring in sympatry with the other two specimens. A detailed morphological investigation was conducted on the specimens belonging to the different clusters detected in *O. (O.) pinguides*. This required the revision of the type material of the species, as well as the type material of *Allogaussia lobata*, synonymized with *O. (O.) pinguides* by Hurley (1975; see complete taxonomic references and geographic records in De Broyer et al., 2007). Supported by the barcoding results, this revision permitted the detection of minor but consistent morphological differences between *O. (O.) pinguides* and *A. lobata*




Fig. 3. Frequency distribution of pairwise K2P distances of (A) *Orchomenella (Orchomenopsis) acanthurus*, (B) *Orchomenella (Orchomenella) pinguides*, (C) *Orchomenella (Orchomenella) franklini* and (D) *Orchomenella (Orchomenopsis) cavimanus*.

Table 2
Morphological differences between specimens of the two clusters observed within *Orchomenella* (*Orchomenella*) *pinguides*. Cluster A corresponds to *O. (O.) pinguides sensu stricto* and cluster B corresponds to *Orchomenella (Orchomenella) lobata*, a species previously synonymized with *O. (O.) pinguides*.

	<i>O. (O.) pinguides s.s. (Cluster A)</i>	<i>O. (O.) lobata (Cluster B)</i>
Epistome: front margin	Straight	Regularly convex
Epistome: proximal angle	Distinct, rounded	Absent
Gnathopod 1: length carpus vs propodus	± 75% (75–80%)	± 90% (81–100%)
Pereiopod 7: basis postero-distal margin	Regularly convex (or very weakly truncate)	Distinctly truncate
Pleosomite 3: dorso-distal angle	Moderately developed, regularly convex; weakly overvaulting urosomite 1	Moderately to strongly developed, subrectangular with angle rounded; strongly overvaulting urosomite 1
Urosomite 1: dorsal hump	Moderately to well developed, without weak mid-dorsal carina; strongly overvaulting urosomite 2	Well developed, with weak mid-dorsal carina; strongly overvaulting urosomite 2
Epimeral plate 3: proximal angle on hind margin	Distinct	Indistinct (margin nearly regularly rounded)
Epimeral plate 3: postero-distal angle	Indistinct, rounded	Distinct, well marked
Telson cleft	< 50% (38–50%)	> 50% (50–68%)

and thus re-establish the latter species as valid (as *Orchomenella (Orchomenella) lobata*). This morphological analysis will be presented in details elsewhere but is summarized in Table 2. The three specimens of cluster A initially identified as *O. (O.) pinguides* clearly belong to *O. (O.) lobata*. The specimens of cluster B were identified as *O. (O.) pinguides sensu stricto*. Based on the observation of morphological differences, which can be interpreted as interspecific variation and the high divergences separating these two clusters, this species complex seems to consist of two distinct species. Within cluster B, one specimen is separated from the other two by distances higher than 5% but it was not possible to separate them on a morphological basis. Both species, *O. (O.) lobata* and *O. (O.) pinguides*, have been recorded several times in sympatry (De Broyer, pers. comm.) and are characterized by a circumpolar distribution.

4. Discussion

4.1. “Barcoding gaps” and species delimitation

The clear barcoding gap observed in our COI dataset means that the assignment of a specimen to a particular species based on a “threshold” value of sequence divergence would mostly work for this group and would be also efficient to detect new and/or cryptic species (Hebert et al., 2003a, 2004b; Barrett and Hebert, 2005). Hebert et al. (2004a,b) proposed a standard sequence threshold of ten times the mean intraspecific divergence (K2P distance) to delimit animal species, which was also applied in studies on amphipods (e.g. Witt et al., 2006). In our case, this threshold would be 4.0%. However, the use of thresholds as an (exclusive) evidence ignores variation that may exist in different taxonomic groups. Meyer and Paulay (2005) assume that insufficient sampling on both intraspecific and interspecific levels can lead to false barcoding gaps. On the other hand, the main reason for an overlap between intra- and interspecific distances could be the poor taxonomic knowledge of a group, e.g. the presence of cryptic species that has been overlooked (Wiemers and Fiedler, 2007). This might be the case in the present study as well, where a barcoding gap apparently exists between well-defined species, but an overlap appears when considering the four putative species complexes. However, the species complex observed in *O. (O.) pinguides* appeared to be composed of two overlooked, distinct species. It is also possible that with a more extensive geographical sampling, which is the case in *O. (O.) cavimanus*, the intraspecific variation could increase further as individuals from more populations are sampled. By this, the barcoding gap range might decrease or become inexistent, which makes it impossible to designate a threshold value. Therefore, in such case, additional data from a morphological analysis or from

nuclear markers are essential to verify the species status. In our previous study (Havermans et al., 2010), phylogenetic analyses were conducted on several of the specimens used for this barcoding study, based on COI and the nuclear gene 28S rRNA. The monophyletic clusters identified with the neighbour-joining tree corresponded to the clades revealed by the phylogenetic study, even in the case of the species complexes. In these complexes, specimens were also separated by higher divergences than within-species variations. Considering this, the phylogenetic species concept could also be applied, which defines a species as the smallest resolvable separately evolving lineage or the smallest diagnosable cluster (Vogler and Monaghan, 2007). The clusters within the species complexes identified in this study would then be recognized as different species. It also remains not less important to critically examine the morphology of the specimens belonging to the species complexes. A first examination of *O. (O.) pinguides* has been accomplished and revealed the presence of two distinct species. At first view, no morphological differences could be observed within the other species complexes, but this requires a detailed examination of all specimens and their types and this is clearly out of the scope of this paper.

4.2. Genetic structures of orchomenid species

Even within this group of closely related species, completely different genetic structures could be observed. The mitochondrial data revealed distinct, monophyletic clusters in *O. (O.) pinguides*, *O. (O.) franklini*, *O. (O.) cavimanus* and *O. (O.) acanthurus*. After a detailed morphological analysis some differences between specimens of *O. (O.) pinguides* suggested the presence of two morphologically similar species, which were formerly synonymized under *O. (O.) pinguides*. However, specimens of the three other species complexes seemed difficult to separate on a morphological basis. However, the genetic divergences between the clusters within *O. (O.) cavimanus*, *O. (O.) acanthurus* and *O. (O.) franklini* are congruent with species-level divergences in the orchomenid genus complex. Held (2003) developed a set of criteria to provide evidence for cryptic speciation of serolid isopods of the Antarctic waters: (1) a bimodal distribution of pairwise distance measures with no intermediate values, (2) a differentiation at a level known for this gene from other undisputed species pairs closely related to the studied species, and (3) the persistence of an expressed genetic divergence in sympatry. In our case, (1) a clear gap is observed in the distribution of intra- and interspecific distances, (2) the genetic distances between the different clusters of *O. (O.) cavimanus*, *O. (O.) acanthurus* and *O. (O.) franklini* are in the same range as interspecific distances of closely related orchomenid species, and

is less than the lowest interspecific variation (6.3%) but higher than the threshold value. No well-supported clusters could be observed, since there was an overlap of genetic divergences. This might suggest that *O. (O.) cavimanus* is in the process of speciation and had not yet the time to diverge genetically to the point where species distinction is possible. However, the specimens separated by intermediate genetic distances occur in sympatry and no clear geographical populations can be distinguished. Furthermore, a specimen from the Magellanic region (Burdwood Bank) clustered within species from the Scotia Sea and the Antarctic Peninsula, suggesting a possible dispersal across the Antarctic Polar Front, which normally presents a physical dispersal barrier for marine biota (Crame, 1999).

Nevertheless, we cannot assume that present distributions necessarily reflect ancient ones, knowing that the several cooling and warming episodes of the Antarctic geological history would have led to changes in the ranges of Antarctic marine taxa (Clarke and Crame, 1997; Page and Linse, 2002). This should be investigated with more fine-scaled molecular methods at the population level and an in-depth morphological examination.

5. Conclusion

A species identification by DNA barcoding was carried out for this group of Antarctic amphipods and revealed species new to science as well as the discovery of three likely species complexes and two genetically and morphologically distinct species formerly synonymized under one. Our barcoding study has been shown to be efficient for these amphipod taxa and will facilitate future taxonomic studies. The new and cryptic species will be submitted to a more accurate morphological analysis, since taxonomy should imply a holistic approach where morphological, ecological and genetic evidence is used together to delimitate species.

The application of DNA barcoding could be used in the future for species diversity studies in this group and other lysianassooid groups, as a way for non-specialists to discriminate taxa that are otherwise difficult to identify. It will thus make species identifications faster and more accessible at a lower cost at the same time.

In poorly known amphipod groups, high intraspecific genetic divergences could indicate overlooked species or species complexes. This barcode application can provide a preliminary signal of species richness. Moreover, the discovery of cryptic diversity could have profound implications for evolutionary theories and biogeography and may be a potentially important factor influencing future conservation decisions (Witt et al., 2006; Bickford et al., 2007). Furthermore, the Census of Antarctic Marine Life (CAML) states that there is an urgent need for more genetic barcode studies on Antarctic organisms, in view of the rate of climate-driven habitat changes which might lead to extinctions (Grant and Linse, 2009).

This study indicated that the species richness of Antarctic amphipods is underestimated, not only for the poorly known deep-sea but also for the better studied shelf fauna. Given the fact that our sampling mainly focused on the Atlantic sector and to a lesser extent on the Ross Sea, we expect that the entire diversity is even much higher. Therefore, additional samples from other areas in Antarctica are needed to assess the real diversity, and evaluate whether these identified clusters have a true circumpolar distribution.

Acknowledgments

The first author is supported by an "Action II" grant (contract number WI/36/H04) from the Belgian Science Policy Office and the study was performed in the framework of the Scientific

Research Programme on the Antarctic (BIANZO II project) supported by the Belgian Science Policy Office. For genetic analyses, funding was provided by the Joint Experimental Molecular Unit (project BARCOLYS) that is supported by the Belgian Science Policy Office. We thank the Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, for the invitation to participate in the expeditions ANT XV-3, ANT XIX-5, ANDEEP I-II-III and ANT XXIII-8, and the crew of the R/V "Polarstern" for their professional help during the cruises. We are indebted to A. Jazdzewska (Laboratory of Polar Biology and Hydrobiology, University of Łódź, Poland) for providing samples from the Polish Antarctic IPY Expedition in 2007 (51/N-IPY/2007/0). We thank Dr. K. Schnabel (CM, Marine Invertebrate Collection, NIWA, Wellington, New Zealand) for providing specimens collected during the biodiversity survey of the western Ross Sea and Balleny Islands in 2004 (BIOROSS Cruise) undertaken by the National Institute of Water and Atmospheric Research and financed by the New Zealand Ministry of Fisheries. This is contribution no. 32 to the Census of Antarctic Marine Life.

Appendix A. Supplementary Material

Supplementary data associated with this article can be found in the online version at doi:10.1016/j.dsr.2010.09.028.

References

- Allcock, A.L., Breirley, A.S., Thorpe, J.P., Rodhouse, P.G., 2004. Restricted gene flow and evolutionary divergence between geographically separated populations of the Antarctic octopus *Pareledone turqueti*. *Marine Biology* 129, 97–102.
- Arnaud, P.M., Jazdzewski, K., Presler, P., Siciński, J., 1986. Preliminary survey of benthic invertebrates collected by Polish Antarctic Expeditions in the Admiralty Bay (King George Island, South Shetlands, Antarctica). *Polish Polar Research* 7, 7–24.
- Arntz, W.E., Brey, T., Gallardo, V.A., 1994. Antarctic zoobenthos. *Oceanography and Marine Biology: An Annual Review* 32, 241–304.
- Arntz, W.E., Thatje, S., Gerdes, D., Gili, J.M., Gutt, J., Jacob, U., Montiel, A., Orejas, C., Teixidó, N., 2005. The Antarctic–Magellan connection: macrobenthos ecology on the shelf and upper slope, a progress report. *Scientia Marina* 69, 237–269.
- Astrin, J.J., Huber, B.A., Misof, B., Klütch, C.F.C., 2006. Molecular taxonomy in pholcid spiders (Pholcidae, Araneae): evaluation of species identification methods using COI and 16S rRNA. *Zoologica Scripta* 35, 441–457.
- Barnard, J.L., Karaman, G., 1991. The families and genera of marine Gammaridean Amphipoda (except marine Gammaroids). Part 1. Records of the Australian Museum 13, 1–866.
- Barrett, R.D.H., Hebert, P.D.N., 2005. Identifying spiders through DNA barcodes. *Canadian Journal of Zoology* 83, 481–491.
- Beaumont, A.R., Wei, J.H.C., 1991. Morphological and genetic variation in the Antarctic limpet *Nacella concinna* (Strebel, 1908). *Journal of Molluscan Studies* 57, 443–450.
- Bernardi, G., Goswami, U., 1997. Molecular evidence for cryptic species among the Antarctic fish *Trematomus bernachii* and *Trematomus hansonii*. *Antarctic Science* 9, 381–385.
- Bickford, D., Lohman, D.J., Sohdi, N.S., Ng, P.K.L., Meier, R., Winker, K., Ingram, K.K., Das, I., 2007. Cryptic species as a window on diversity and conservation. *Trends in Ecology and Evolution* 22, 148–155.
- Blaxter, M.L., 2003. Molecular systematics: counting angels with DNA. *Nature* 421, 122–124.
- Blaxter, M.L., 2004. The promise of DNA taxonomy. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences* 359, 669–679.
- Bouchet, P., 2006. The magnitude of marine biodiversity. In: Duarte, C.M. (Ed.), *The Exploration of Marine Biodiversity. Scientific and Technological challenges*, Fundación BBVA, pp. 33–64.
- Brandt, A., 1999. On the origin and evolution of Antarctic Peracarida (Crustacea, Malacostraca). *Scientia Marina* 63, 261–274.
- Brandt, A., 2005. Evolution of Antarctic biodiversity in the context of the past: the importance of the Southern Ocean deep sea. *Antarctic Science* 17, 509–521.
- Brandt, A., Gooday, A.J., Brandão, S.N., Brix, S., Brökeland, W., Cedhagen, T., Choudhury, M., Cornelius, N., Danis, B., De Mesel, I., Diaz, R.J., Gillan, D.C., Ebbe, B., Howe, J.A., Janussen, D., Kaiser, S., Linse, K., Malyutina, M., Pawłowski, J., Raupach, M., Vanreusel, A., 2007. First insights into the biodiversity and biogeography of the Southern Ocean deep sea. *Nature* 447, 307–311.
- Brey, T., Dahm, C., Gorny, M., Klages, M., Stiller, M., Arntz, W.E., 1996. Do Antarctic invertebrates show an extended level of eurybathy? *Antarctic Science* 8 3–6.

- Vogler, A.P., Monaghan, M.T., 2007. Recent advances in DNA taxonomy. *Journal of Zoological Systematics and Evolutionary Research* 45, 1–10.
- Ward, R.D., Zemlak, T.S., Innes, B.H., Last, P.R., Hebert, P.D.N., 2005. DNA barcoding Australia's fish species. *Philosophical Transactions of the Royal Society B: Biological Sciences* 360, 1847–1857.
- Watling, L.E., Thurston, M.H., 1989. Antarctica as an evolutionary incubator: evidence from the cladistic biogeography of the amphipod family Iphimediidae. In: Crame, J.A. (Ed.), *Origins and Evolution of the Antarctic Biota*. The Geological Society, London, pp. 297–313.
- Waugh, J., 2007. DNA barcoding in animal species: progress, potential and pitfalls. *Phylogenetics and systematics*. *Bioessays* 29, 188–197.
- Whitworth, T.L., Dawson, R.D., Magalon, H., Baudry, E., 2007. DNA barcoding cannot reliably identify species of the blowfly genus *Protocalliphora* (Diptera: Calliphoridae). *Proceedings of the Royal Society, B, Biological Sciences* 274, 1731–1739.
- Wiemers, M., Fiedler, K., 2007. Does the DNA barcoding gap exist?—A case study in blue butterflies (Lepidoptera: Lycaenidae). *Frontiers in Zoology* 4, 8.
- Will, K.W., Rubinoff, D., 2004. Myth of the molecule: DNA barcodes for species cannot replace morphology for identification and classification. *Cladistics* 20, 47–55.
- Wilson, N.G., Hunter, R.L., Lockhart, S.J., Halanych, K.M., 2007. Multiple lineages and absence of panmixia in the “circumpolar” crinoid *Promachocrinus kerguelensis* from the Atlantic sector of Antarctica. *Marine Biology* 152, 895–904.
- Witt, J.D.S., Blinn, D.W., Hebert, P.D.N., 2003. The recent evolutionary origin of the phenotypically novel amphipod *Hyaella montezuma* offers an ecological explanation for morphological stasis in a closely allied species complex. *Molecular Ecology* 12, 405–413.
- Witt, J.D.S., Threlloff, D.L., Hebert, P.D.N., 2006. DNA barcoding reveals extraordinary cryptic diversity in an amphipod genus: implications for desert spring conservation. *Molecular Ecology* 15, 3073–3082.