

A REDESCRIPTION OF *ANTHOMASTUS AGILIS* TIXIER-DURIVAUT WITH A GENERIC PLACEMENT IN *SINULARIA* (COELENTERATA: OCTOCORALLIA), AND SOME REMARKS ON *SINULARIA RAMOSA* TIXIER-DURIVAUT.

PHIL ALDERSLADE

Northern Territory Museum of Arts and Sciences,  
GPO Box 4646, Darwin, NT, 0801, Australia.

ABSTRACT

The holotype of *Anthomastus agilis* (family Alcyoniidae) does not have siphonozooids nor the characteristic spiculation of *Anthomastus*, as originally reported. The specimen is redescribed and transferred to the genus *Sinularia*. Some additional information and sclerite drawings are given for *Sinularia ramosa*.

KEYWORDS: Coelenterata, Anthozoa, Alcyonacea, Octocorallia, Alcyoniidae, *Anthomastus*, *Sinularia*, soft corals, new combination, taxonomy, New Caledonia.

INTRODUCTION

During the course of researching the identity of an unusual dimorphic soft coral from Darwin, northern Australia, I had cause to examine the holotype of *Anthomastus agilis* which had been described by Tixier-Durivault (1970a:186-187). Apart from lacking the characteristically large autozooids of *Anthomastus*, it also became obvious that the spiculation was quite atypical of that genus. A far too cursory look at the proposed siphonozooids that were reported by Tixier-Durivault led to my erroneous "pers. comm." to Dr Gary Williams (1992:1, 23) that the material represented a new genus. A subsequent examination, regrettably too late to prevent publication, revealed that the siphonozooids recorded by Tixier-Durivault were actually inflated caecae of the subsurface canal system. Ironically, Tixier-Durivault (1951:11) had previously pointed out that other authors had made similar mistakes.

The holotype has the characteristic spiculation of a *Sinularia* and is herein transferred to that genus as an undescribed species. The somewhat unusual growth form is perhaps attributable to its apparent juvenility when preserved, and the inflated nature of the specimen, and

possibly also its most recent habitat, which was the Noumea Aquarium run by its founder M. Rend Catala.

It appears that Utinomi and Imahara (1976: 211) failed to take into account the difference in the respective sizes of the internal sclerites when they proposed that *A. agilis* was a probable synonym of *A. phalloides* Benham, 1928. The sclerites of *A. agilis* were described as up to 2.2 mm in length and those of *A. phalloides* only up to 0.25 mm. Williams (1992:19) recently transferred *A. phalloides* to the genus *Minabea*.



Fig. 1. *Sinularia agilis*, holotype. 1.5 times natural size.

Mme d'Hondt's 1992 paper (published in 1993) was only received subsequent to submitting the manuscript for this paper. Several electron micrographs of surface sclerites

of *Anthomastus agilis* are illustrated in Figure 1 of that publication and should be viewed in conjunction with the present description.




Fig. 2. *Simulium agilis*, holotype, sclerites from the surface of the lobes.




Fig. 3. *Simularia agilis*, holotype, sclerites from the surface of the lobes.




Fig. 4. *Sinularia agilis*, holotype, sclerites from the surface of the base.

SYSTEMATICS

Family Alcyoniidae Lamouroux  
Genus *Sinularia* May, 1898

*Sinularia agilis* (Tixier-Durivault, 1970a)  
new comb.  
(Figs 1-5)

*Anthomastus agilis* Tixier-Durivault,  
1970a:186-187.

Type Material. HOLOTYPE - Noumea, R. Catala,  
1966, Museum National d'Histoire Naturelle, Paris.

Description. The holotype (Fig. 1) is a single lobe,  
about 40 mm in length and 11 mm at its greatest upper  
diameter, with a small flattened lobe, 15 x 6 mm, that  
protrudes from one side. The

colony seems to have been preserved in a relatively  
inflated state and is soft and compressible.

The polyp apertures are about 0.5 mm across and are  
present over both the large and small lobes down to  
within approximately 15 mm of the base of the  
specimen. While some of the polyps have their  
contracted, but unfolded, tentacles protruding, most are  
retracted until they are flush with the lobe surface. The  
inflated caecae of the subsurface canal system appear as  
clear patches between the white heads of the  
club-shaped surface sclerites, and are present right  
down to the base of the specimen. Tixier-Durivault's  
illustration (1970a: fig. 10E) is misleading in its uniform  
representation of the shape of the patches which  
actually vary from small and more-or-less circular, to  
larger irregular and often elongate structures.  
Microscopical examination of small tissue fragments in  
both




Fig. 5. *Sinularia agilis*, holotype: A, sclerites from the interior of the lobes; B, sclerites from the interior of the base; C, detail of sclerite warring.

plan and sectional views revealed no apertures or mesenterial structures in the caecae, only aggregates of cells which are probably zooxanthellae.

The surface of the large lobe contains club shaped sclerites (Figs 2, 3) with irregularly and complexly warted heads. They vary from 0.070.30 mm in length with most <0.28 mm long. There are also a few curved spindles, about 0.25 mm long, with simple processes, but they are quite rare.

The interior of the large lobe contains long, pointed, narrow spindles up to 2.2 mm long (Fig. SA). Some of the smaller forms are branched. The complex warts are high and spiny (Fig. SC).

The surface of the base contains clubs and a few irregular forms (Fig. 4). They are 0.09 - 0.22 mm long, with most <0.17 mm. The handles of the clubs are thicker than those in the lobe surface.

The interior of the base contains large, robust spindles, up to 2.5 mm long, many of which are terminally bifurcated (Fig. SB). The complex warts are the same as those on the sclerites in the lobe interior, but they are far more densely placed.

**Remarks.** While comparing *Simularia agilis* to other species of *Simularia*, I was able to examine sclerite samples of the holotype of *S. ramosa* Tixier-Durivault, 1945. There is some similarity in the general shape of the sclerites




Fig. 6. *Simularia ramosa*, holotype: A, polyp sclerites; B, sclerites from the surface of the lobes.

between this species and *S. agilis*, and it is not inconceivable that *S. agilis* could have a similar mature colony form. The style of the sclerite waning is different, but perhaps it is within the bounds of intraspecific variation. Unfortunately, the latter is a subject on which little is known for the majority of octocorals, and nothing is known with respect to *S. ramosa*. Tixier-Durivault did report other occurrences of the species, from New Caledonia (1970a:293) and Vietnam (1970b:205), but regrettably none of the 12 specimens were described. The author, characteristically, just repeated her earlier original diagnostic text word for word, including the sclerite dimensions.

The holotype of *S. ramosa* was redescribed by Verseveldt (1980:112-113), but two important facts were overlooked. First, the polyps are armed with at least four to five pairs of rod-like and club-like sclerites (Fig. 6A) arranged en chevron. They are 0.09-0.17 mm long. Second, the surface of the lobes contains numerous, narrow, often curved, pointed spindles (Fig. 6B) that are ornamented with simple prominences. It is also worth noting that the majority of surface clubs are only 0.09-0.12 mm long. This is not obvious from the text, but it is why this species is included in Verseveldt's key under group III. This size is considerably smaller than for *S. agilis* which is here considered a distinct species.

#### ACKNOWLEDGEMENTS

My sincere thanks to Mme Marie-Jose d' Hondt for her continued support of my research, for allowing me to borrow type material, and for always being willing to share her knowledge. I am also grateful to Lorna Gravener for typing the manuscript.

#### REFERENCES

- Benham, W.B. 1928. On some alcyonarians from New Zealand waters. *Transactions and Proceedings of the New Zealand Institute* 59: 67-84.
- d'Hondt, M.-J. 1993 (1992). Description d' *Anthomastus globosus* n.sp. (Octocorallia, Alcyonacea) de Nouvelle-Calédonie. Remarques sur espèces du genre. *Bulletin du Museum National D'Histoire Naturelle, Paris*, 4<sup>o</sup> ser., 14, section A (311): 623-638.
- May, W. 1898. Die von Dr Stuhlmann im Jahre 1889 gesammelten ostafrikanischen Alcyonaceen des Hamburger Museums. *Jahrbuch des Hamburgischen Wissenschaftlichen Anstalten* 15(2): 1-38.
- Tixier-Durivault, A. 1945. Les Alcyonaires du Museum. 1. Famille des Alcyoniidae. 2. Genre *Sinularia*. *Bulletin du Museum National D'Histoire Naturelle* (2) 17(1): 243-250.
- Tixier-Durivault, A. 1951. Revision de la famille des Alcyoniidae. Le genre *Sinularia* May, 1898. *Memoires de l'Institut Royal des Sciences Naturelles de Belgique* (2) 40: 1-146.
- Tixier-Durivault, A. 1970a. Les Octocoralliaires de Nouvelle-Calédonie. L'Expedition française sur les récifs coralliens de la Nouvelle-Calédonie organisée sous l'égide de la fondation SingerPolignac 1960-1963 4: 171-350.
- Tixier-Durivault, A. 1970b. Les Octocoralliaires de Nha-Trang (Viet-Nam). *Cahiers du Pacifique* 14:115-236.
- Utinomi, H., and Imahara, Y. 1976. A new second species of dimorphic alcyonacean octocoral *Minabea* from the Bays of Sagami and Suruga, with the emendation of generic diagnosis. *Publications of the Selo Marine Biological Laboratory* 23 (3-5): 205-212.
- Verseveldt, J.1980. A revision of the genus *Sinularia* May (Octocorallia, Alcyonacea). *Zoologische Verhandelingen* 179: 1-91.
- Williams, G.C.1992. Revision of the soft coral genus *Minabea* (Octocorallia: Alcyoniidae) with new taxa from the Indo-West Pacific. *Proceedings of the California Academy of Sciences* 48(1): 1-26.

Accepted 8 March 1994