

Reconstruction of Family-Level Phylogenetic Relationships within Demospongiae (Porifera) Using Nuclear Encoded Housekeeping Genes

Malcolm S. Hill¹, April L. Hill¹, Jose Lopez², Kevin J. Peterson³, Shirley Pomponi⁴, Maria C. Diaz⁵, Robert W. Thacker⁶, Maja Adamska⁷, Nicole Boury-Esnault⁸, Paco Cárdenas⁹, Andia Chaves-Fonnegra², Elizabeth Danka¹, Bre-Onna De Laine¹, Dawn Formica², Eduardo Hajdu¹⁰, Gisele Lobo-Hajdu¹¹, Sarah Klontz¹², Christine C. Morrow¹³, Jignasa Patel², Bernard Picton¹⁴, Davide Pisani¹⁵, Deborah Pohlmann¹, Niamh E. Redmond¹², John Reed⁴, Stacy Richey¹, Ana Riesgo¹⁶, Ewelina Rubin², Zach Russell¹, Klaus Rützler¹², Erik A. Sperling¹⁷, Michael di Stefano¹, James E. Tarver¹⁸, Allen G. Collins^{12,19*}

1 Gottwald Science Center, University of Richmond, Richmond, Virginia, United States of America, **2** Nova Southeastern University Oceanographic Center, Dania Beach, Florida, United States of America, **3** Department of Biological Sciences, Dartmouth College, Hanover, New Hampshire, United States of America, **4** Harbor Branch Oceanographic Institute, Florida Atlantic University, Fort Pierce, Florida, United States of America, **5** Museo Marino de Margarita, Boulevard de Boca Del Rio, Boca del Rio, Nueva Esparta, Venezuela, **6** Department of Biology, University of Alabama at Birmingham, Birmingham, Alabama, United States of America, **7** Sars International Centre for Marine Molecular Biology, Thormøhlensgt, Bergen, Norway, **8** IMBE-UMR7263 CNRS, Université d'Aix-Marseille, Station marine d'Endoume, Marseille, France, **9** Department of Systematic Biology, Evolutionary Biology Centre, Uppsala University, Uppsala, Sweden, **10** Departamento de Invertebrados, Museu Nacional/Universidade Federal do Rio de Janeiro, Rio de Janeiro, Rio de Janeiro, Brazil, **11** Departamento de Genética, IBRAG, Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Rio de Janeiro, Brazil, **12** Department of Invertebrate Zoology, National Museum of Natural History, Smithsonian Institution, Washington, D.C., United States of America, **13** School of Biological Sciences, MBC, Queen's University, Belfast, United Kingdom, **14** National Museums Northern Ireland, Holywood, Northern Ireland, United Kingdom, **15** School of Earth Sciences and School of Biological Sciences, The University of Bristol, Bristol, United Kingdom, **16** Museum of Comparative Zoology, Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, Massachusetts, United States of America, **17** Harvard University, Department of Earth and Planetary Science, Cambridge, Massachusetts, United States of America, **18** School of Earth Sciences, University of Bristol, Bristol, United Kingdom, **19** National Systematics Laboratory of NOAA's Fisheries Service, National Museum of Natural History, Smithsonian Institution, Washington, D.C., United States of America

Abstract

Background: Demosponges are challenging for phylogenetic systematics because of their plastic and relatively simple morphologies and many deep divergences between major clades. To improve understanding of the phylogenetic relationships within Demospongiae, we sequenced and analyzed seven nuclear housekeeping genes involved in a variety of cellular functions from a diverse group of sponges.

Methodology/Principal Findings: We generated data from each of the four sponge classes (i.e., Calcarea, Demospongiae, Hexactinellida, and Homoscleromorpha), but focused on family-level relationships within demosponges. With data for 21 newly sampled families, our Maximum Likelihood and Bayesian-based approaches recovered previously phylogenetically defined taxa: *Keratosa*^P, *Myxospongiae*^P, *Spongillida*^P, *Haploscleromorpha*^P (the marine haplosclerids) and *Democlavia*^P. We found conflicting results concerning the relationships of *Keratosa*^P and *Myxospongiae*^P to the remaining demosponges, but our results strongly supported a clade of *Haploscleromorpha*^P+*Spongillida*^P+*Democlavia*^P. In contrast to hypotheses based on mitochondrial genome and ribosomal data, nuclear housekeeping gene data suggested that freshwater sponges (*Spongillida*^P) are sister to *Haploscleromorpha*^P rather than part of *Democlavia*^P. Within *Keratosa*^P, we found equivocal results as to the monophyly of Dictyoceratida. Within *Myxospongiae*^P, Chondrosida and Verongida were monophyletic. A well-supported clade within *Democlavia*^P, *Tetractinellida*^P, composed of all sampled members of Astrophorina and Spirophorina (including the only lithistid in our analysis), was consistently revealed as the sister group to all other members of *Democlavia*^P. Within *Tetractinellida*^P, we did not recover monophyletic Astrophorina or Spirophorina. Our results also reaffirmed the monophyly of order Poecilosclerida (excluding Desmacellidae and Raspailiidae), and polyphyly of Hadromerida and Halichondrida.

Conclusions/Significance: These results, using an independent nuclear gene set, confirmed many hypotheses based on ribosomal and/or mitochondrial genes, and they also identified clades with low statistical support or clades that conflicted with traditional morphological classification. Our results will serve as a basis for future exploration of these outstanding questions using more taxon- and gene-rich datasets.

Citation: Hill MS, Hill AL, Lopez J, Peterson KJ, Pomponi S, et al. (2013) Reconstruction of Family-Level Phylogenetic Relationships within Demospongiae (Porifera) Using Nuclear Encoded Housekeeping Genes. *PLoS ONE* 8(1): e50437. doi:10.1371/journal.pone.0050437

Editor: Senjie Lin, University of Connecticut, United States of America

Received: May 1, 2012; **Accepted:** October 22, 2012; **Published:** January 23, 2013

This is an open-access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the Creative Commons CC0 public domain dedication.

Funding: This work was supported by the National Science Foundation's Assembling the Tree of Life Program (http://www.nsf.gov/funding/pgm_summ.jsp?pims_id=5129; DEB awards 0829763, 0829783, 0829791, and 0829986). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: collinsa@si.edu

Introduction

Sponges belong to an ancient metazoan lineage with a fossil record that stretches back to the late Cryogenian >635 Myr ago [1–3]. Some estimates place their appearance at nearly 800 Myr ago [4,5]. As a sister group (or groups) to all the other animals in the metazoan tree of life, sponges represent a fulcrum point in the history of animal life lying at the junction between single-celled ancestors and the rest of Metazoa. Sponges have also been important ecosystem engineers throughout much of their history, e.g., as major reef-builders during the Upper Devonian, Upper Permian, and through a major portion of the Jurassic [6,7]. In modern oceans, poriferans continue to perform important ecological functions as water filterers, bioeroders, structural habitat providers, microbial symbiont incubators, dissolved organic carbon sinks, natural product biosynthesizers, chemical accumulators, and potential marine pathogen reservoirs [8–15]. As one of the most diverse taxa of extant sessile invertebrates [16], a detailed exploration of poriferan evolutionary relationships will yield important insights into many phases of metazoan history.

Due to their simple bodies with a paucity of easily accessible morphological traits, sponges are notoriously resistant to attempts at taxonomic classification [16]. Indeed, taxonomic controversy extends from the highest levels of classification (e.g., whether the phylum Porifera is monophyletic [17–20]) to whether particular genera belong to one or another family (e.g., [21]), or even whether different nominal species are truly distinct (e.g., [22,23]). In the mid-1980s, van Soest [24] presented a call to include explicitly phylogenetic perspectives in sponge systematics through cladistic analysis. Since that time, phylogenetic classification has permeated the field of sponge taxonomy (e.g., [25–38]). As currently envisioned, four classes comprise the phylum Porifera: Calcarea = (Calcispongiae plus the fossil group Heteractinida), Demospongiae, Homoscleromorpha, and Hexactinellida [39]. Ample evidence exists to conclude that each of these classes is monophyletic, and so each has been provided with an explicit phylogenetic definition [40]. Indeed, substantial evidence is accumulating for the existence of various sponge clades at different levels [40,41], and throughout this paper, we will differentiate between Linnean taxa and those clades that have been provided with explicitly phylogenetic definitions by italicizing phylogenetically defined taxa and following them with a superscript *p*, as in *Demospongiae^p* (i.e., *PhyloCode* designations).

A major challenge to scientists working in this field has been the identification of appropriate markers for addressing the daunting task of dealing with ancient divergences among the diverse assortment of poriferan taxa. Evolutionary relationships across the most diverse class of Porifera, Demospongiae, have mainly been addressed with three sets of phylogenetic markers: ribosomal DNA sequences [17,42], complete mitochondrial genome sequences [43], and amino acid sequences that code for seven nuclear

housekeeping genes [18,44]. A broad correspondence in inferences about demosponge phylogeny exists between these three sets of data (see discussion below), but both of the latter two sets of data have been sampled from a far more limited number of taxa. The Porifera Tree of Life project (www.portol.org) employs a variety of tools to integrate morphological and molecular data and to expand the diversity of sponge taxa used to elucidate all levels of sponge phylogeny. In this study, we report findings based on a significant expansion (38 new samples from 38 species representing 30 families, including 21 families newly sampled) of the nuclear housekeeping gene dataset first developed for metazoan-wide phylogenetic and molecular dating analyses [45,46] and later applied by Sperling et al. [18,44] to sponges, with a thorough taxonomic vetting process and a slightly modified phylogenetic analysis focused on relationships within *Demospongiae^p*.

Results

Extraction of high quality RNA for subsequent cDNA synthesis and cloning was a significant hurdle, curtailing use of some samples (e.g., lithistids), even though a large number of archived specimens were available for potential study [47]. Several hundred cDNAs were cloned and sequenced, but only 159 usable sequences were generated due to the amplification of non-sponge contaminants (Tables 1–2). We evaluated single gene phylogenies (ALD, ATPB, etc.) including all the members of each gene family that could be identified in GenBank (via reciprocal blasting) to identify and remove potential paralogs. Our dataset for phylogenetic analysis contains 2,033 amino acid characters and a total of 68 sponge species representing 48 of 137 accepted and recently proposed families of Porifera [38,40,48], including 51 species from 37 of 91 families recognized for Demospongiae (Table 1). The most appropriate models of amino acid evolution, as determined by ProtTest [49] for the various datasets (i.e., all genes, each individual gene, etc.), nearly always involved some variant of the LG matrix [50] (Table 3). Maximum likelihood mapping, performed for each gene under the best fitting model, among those implemented in TreePuzzle [51], showed that each of the seven considered genes convey enough phylogenetic signal to be considered potentially useful phylogenetic markers to resolve the relationships within Demospongiae (Figs. S1, S2, S3, S4, S5, S6, S7). Bayesian cross-validation [52] analyses showed that the CAT based models (CAT and CAT-GTR) fit our dataset significantly better than any empirical site-homogeneous time reversible model tested (WAG+G, and LG+G). Cross-validation also showed that the CAT-based models fit the data better than the more complex site-homogeneous time reversible model: the mechanistic amino acid-GTR (Table 4) model. Accordingly, hypothesized relationships obtained with homogeneous time-reversible models (e.g. LG or GTR), where differing from those obtained in our CAT and particularly CAT-GTR analysis, could be considered inferior.

Table 1. Annotated list of samples and sequences used for analysis. New sequences and samples are indicated in bold.

Higher Clades/Classification and Identification	Voucher #	ALD	ATPB	CAT	EF1a	MAT	PFK	TPI	PorToL ID
Keratosa^a, Dendroceratida									
Dictyodendrillidae	USNM_1148204	GQ332402	GQ330912	GQ336998	GQ330927	GQ330916	GQ330918	GQ330922	NA
Dictyodendrillidae	USNM_1133861	JQ606746	JQ606789	JQ606796	JQ606696	JQ680966	JQ680966	JQ680967	P153
Keratosa^a, Dictyoceratida									
Dysideidae	USNM_1148214	GQ332403	GQ330913	GQ336999	GQ330928	GQ330919	GQ330919	GQ330919	NA
Irciniidae	USNM_1153592			JQ680968	JQ606699	JQ606661	JQ606661	JQ606661	TOL24
Irciniidae	USNM_1148130	GQ331021	GQ330993	GQ331006	GQ330979	GQ330952	GQ330952	GQ330939	NA
Spongiidae	USNM_1154092		JQ606797	JQ606729	JQ606706				RWT1816
Thorectidae	USNM_1133719	JQ606755	JQ606799	JQ606731	JQ606708	JQ606668	JQ606668	JQ606775	P14
Myxospongiae^a, Chondrosida									
Chondrillidae	USNM_1148122	GQ332401	HM859880	GQ336997	GQ330926	GQ330915	GQ330915	HM859889	NA
Haliarctidae	USNM_1148131	GQ331020	GQ330992	HM859888	HM859888	GQ330965	GQ330965	GQ330938	NA
Myxospongiae^a, Verongida									
Aplysiniidae	USNM_1133710	JQ606737		JQ606713	JQ606687				P4
Aplysiniidae	USNM_1153593	JQ606736	JQ606781	JQ606712	JQ606685	JQ606671	JQ606671	JQ606671	TOL 25
Aplysiniidae	USNM_1148123	GQ331013	GQ330987	GQ331000	GQ330973	GQ330958	GQ330958	GQ330932	NA
Aplysiniidae	NA	GQ331026	HM859882	GQ331012		GQ330971	GQ330971	GQ330946	NA
Spongillida^a									
Spongillidae	NA	DQ087496	DQ087498	DQ087497	DQ087499	DQ087500	DQ087500	DQ087500	NA
Spongillidae	NA	AY580188	AY580189	AY580190	AY580191	AY580192	AY580193	AB000891	NA
Haploscleromorpha^a									
Callyspongiidae	USNM_1154088		JQ606785	JQ606716	JQ606690	JQ606672	JQ606656	JQ606760	RWT1812
Chalinidae	USNM_1133711	JQ606741		JQ606717	JQ606691	JQ606655	JQ606655	JQ606761	P5
Chalinidae	NA	GQ331014	GQ330988	GQ331001	GQ330974	GQ330959	GQ330949	GQ330933	NA
Chalinidae	NA	GQ331019	GQ330991	GQ331005	GQ330978	GQ330964	GQ330964	GQ330937	NA
Niphatidae	USNM_1153590	JQ606749	JQ606793	JQ680969	JQ606701	JQ606679	JQ606679	JQ606768	TOL20
Niphatidae	NA	(16)	(16)	(16)	(16)	(16)	(16)	(16)	NA
Petrosiidae	MCZ_DNA105722		KA659909	KA659907	KA659906	KA659904	KA659905	KA659901	NA
Petrosiidae	USNM_1154090	JQ606750		JQ606725	JQ606703	JQ606677	JQ606663	JQ606771	RWT1813
Phloeodictyidae	USNM_1133740	JQ606751	JQ606795	JQ606726					P34
Democlavia^a, Tetractinellida^a, Astrophorina									
Ancorinidae	USNM_1175047		JQ606794	JQ606725	JQ606703	JQ606677	JQ606677	JQ606770	JR190
Geodiidae	USNM_1133730	JQ606735	JQ606779	JQ606725	JQ606703	JQ606677	JQ606652	JQ606757	P24
Geodiidae	NA	GQ330990	GQ331004	GQ330977	GQ330963	GQ330963	GQ330936	GQ330936	NA
<i>incertae sedis</i>	USNM_1175067		JQ606702	JQ606702				JQ606769	JR15

Table 1. Cont.

Higher Clades/Classification and Identification	Voucher #	ALD	ATPB	CAT	EF1a	MAT	PFK	TPI	PorToL ID
Democlavia^{sp}, Tetractinellidae^{sp}, Spirophorina									
Sclerodermidae	<i>Microscloderma</i> sp. nov. (8)	USNM_1133739	JQ606784		JQ606689				P33
Tetillidae	<i>Cinachyrella apion</i> (9)	USNM_1153585	HM859881	HM859884	HM859886	GQ330960		GQ330934	NA
Democlavia^{sp}, Agelasida									
Agelasidae	<i>Agelas conifera</i>	USNM_1154089	JQ606734	JQ606778	JQ606711	JQ606684	JQ606651	JQ606756	RWT1814
Hymenhabdillidae	<i>Cymbaxinella^{sp} corrugata</i> (10)	USNM_1153725	JQ606739	JQ606782	JQ606714		JQ606653	JQ606758	TOL29
Democlavia^{sp}, Axinellida									
Raspailiidae	<i>Ectyoplasia ferox</i> (11)	USNM_1133718	JQ606753			JQ606680	JQ606666		P13
Democlavia^{sp}, Hadromerida									
Clonaidae	<i>Clona varians</i>	USNM_1154091	JQ606742	JQ606786	JQ606718	JQ606692	JQ606674	JQ606762	RWT1815
Placospongiidae	<i>Placospongia intermedia</i>	USNM_1133726	JQ606752	JQ606727	JQ606704	JQ606679	JQ606664	JQ606772	P20
Polymastiidae	<i>Polymastia tenax</i>	USNM_1133747	JQ606796	JQ606728	JQ606705		JQ606665	JQ606773	P40
Spirastrellidae	<i>Spirastrella</i> sp. (12)	USNM_1148132	GQ331017	GQ330989	GQ331003	GQ330976	GQ330962	GQ330950	NA
Suberitidae	<i>Suberites</i> sp.	USNM_1148202	GQ331024	GQ330997	GQ330984	GQ330969		GQ330944	NA
Tethyidae	<i>Tethya californiana</i> (13)	USNM_1148128	GQ331025	GQ330998	GQ331011	GQ330985	GQ330970	GQ330945	NA
Democlavia^{sp}, incertae sedis									
Desmacellidae	<i>Bienna caribea</i>	USNM_1175046	JQ606745	JQ606721	JQ606695		JQ606660		TOL27
Desmacellidae	<i>Desmacella pumilio</i>	USNM_1175045	JQ606738	JQ606780	JQ606686	JQ606673		JQ606763	JR19
Dictyonellidae	<i>Dictyonellidae</i> sp. nov.	USNM_1133716	JQ606740	JQ606783	JQ606715	JQ606688	JQ606654	JQ606759	P11
Halichondriidae	<i>Halichondria melanadocia</i>	USNM_1133755	JQ606747	JQ606790	JQ606722	JQ606697		JQ606765	P48
Halichondriidae	<i>Halichondria</i> sp.		GQ332404	GQ330914	GQ337000	GQ330929	GQ330920	GQ330924	NA
Democlavia^{sp}, Poecilosclerida									
Coelosphaeriidae	<i>Lissodendoryx colombiensis</i>	USNM_1133712	JQ606743	JQ606787	JQ606719	JQ606693	JQ606658		P6
Crambeidae	<i>Monanchora arbuscula</i> (14)	USNM_1148203	GQ331023	GQ330996	GQ331010	GQ330983	GQ330955	GQ330943	NA
Crambeidae	<i>Monanchora arbuscula</i>	USNM_1153736	JQ606744	JQ606788	JQ606720	JQ606694	JQ606675	JQ606764	TOL23
Hymedesmiidae	<i>Phorbas</i> sp. nov.	USNM_1133787	JQ606791	JQ606791	JQ606698	JQ606676		JQ606766	P80
Microcionidae	<i>Clathria (Clathria) prolifera</i> (15)	USNM_1148129	DQ087472	DQ087473	DQ087474	DQ087476	DQ087477	DQ087478	NA
Mycalidae	<i>Mycale laevis</i>	USNM_1133707	JQ606748	JQ606792	JQ606723	JQ606700	JQ606678	JQ606767	P1
Tedanidae	<i>Tedania ignis</i>	USNM_1153591	JQ606754	JQ606798	JQ606730	JQ606707	JQ606681	JQ606774	TOL21
Calciopongia^{sp}, Calcaronea^{sp}									
Amphoriscidae	<i>Leucilla nuttingi</i>	NA	GQ330994		GQ330980	GQ330966	GQ330953	GQ330940	NA
Leucosoleniidae	<i>Leucosolenia</i> sp.	USNM_1126268	DQ087465	DQ087466	DQ087467	DQ087468	DQ087469	DQ087470	NA
Leucosoleniidae	<i>Leucosolenia complicata</i>	pending	pending	pending	pending	pending	pending	pending	NA
Sycettidae	<i>Sycon lingua</i>	USNM_1148127	DQ087458	DQ087459	DQ087460	DQ087461	DQ087462		NA
Sycettidae	<i>Sycon coactum</i>	MCZ_DNA105723	KA659914	KA659917	KA659915	KA659916	KA659918	EKA659911	NA

Table 1. Cont.

Higher Clades/Classification and Identification		Voucher #	ALD	ATPB	CAT	EF1a	MAT	PFK	TPI	PortoL ID
Sycettidae	<i>Sycon ciliatum</i>	ZMBN_87981-2	pending	pending	pending	pending	pending	pending	pending	NA
Calcspongiae^a, Calcinea^a										
Clathrinidae	<i>Clathrina cerebrum</i>	NA	GQ331016			GQ330975	GQ330961			NA
Leucettidae	<i>Leucetta chagosensis</i>	NA	(17)	(17)		(17)	(17)			NA
Homoscleromorpha^a										
Oscarellidae	<i>Oscarella carmela</i>	NA	GQ332405		GQ337001	ACL97976	GQ330917	GQ330921	GQ330925	NA
Plakinidae	<i>Corticium candelebrum</i>	MCZ_DNA105720			KA659897		KA659898	KA659899	KA659900	NA
Plakinidae	<i>Plakartis angulospiculatus</i>	USNM_1148206	GQ331022		GQ331008	GQ330981	GQ330967		GQ330941	NA
Hexactinellida^a, Hexasterophora^a										
Aphrocallistidae	<i>Aphrocallistes vastus</i>	NA	GQ330986	GQ330999	GQ330972	GQ330957	GQ330947	GQ330931		NA
Aphrocallistidae	<i>Heterochone calyx</i>	NA	(18)	(18)	(18)	(18)				
Euplectellidae	<i>Hertwigia falcifera</i>	USNM_1175049		JQ606800	JQ606733		JQ606682	JQ606669	JQ606776	JR14
Rossellidae	<i>Acanthascus dawsoni</i>	NA	GQ330995	GQ331009				GQ330954	GQ330942	NA
Rossellidae	<i>Nodastrella asconemaoida</i>	USNM_1175065		JQ606801		JQ606709	JQ606683		JQ606777	JR11
Rossellidae	<i>Bathydorus</i> sp.	USNM_1175050		JQ606802	JQ606732	JQ606710		JQ606670		JR09
Non-Sponge Metazoans										
Cnidaria	<i>Nematostella vectensis</i>	NA	(16)	(16)	(16)	(16)	(16)	(16)	(16)	NA
Cnidaria	<i>Metridium senile</i>	NA	AAT06124	AAT06144	AAT06185	AAT06205	AAT06226	AAT06226	AAT06245	NA
Cnidaria	<i>Acropora millepora</i>	NA	(18)	(18)	(18)	(18)	(18)	(18)	(18)	NA
Placozoa	<i>Trichoplax adhaerens</i>	NA	(18)	(18)	(18)	(18)	(18)	(18)	(18)	NA

⁽¹⁾Formerly identified as *Darwinella muelleri* (Darwinellidae) in Sperling et al. (2007); specimen from the Gulf of Mexico.
⁽²⁾Formerly identified as *Dysidea camera* in Sperling et al. (2007), and as *Dysidea* sp. in GenBank.
⁽³⁾Formerly identified as *Chondrilla* sp. in Sperling et al. (2007) and as *Chondrilla nucula* in GenBank.
⁽⁴⁾Formerly labeled as *Clypeatula cooperensis* in Sperling et al. (2004) and *Ephydatia cooperensis* in GenBank, but synonymized with *Ephydatia fluviatilis* in WPD.
⁽⁵⁾Presently in WPD as *Strelleia fibrosa* as part of family Ancorinidae, but see Cárdenas et al. (2011) for updated classification.
⁽⁶⁾Formerly identified as *Geodia gibberosa* in Sperling et al. (2009) and in GenBank; *G. tumulosa* was resurrected by Cárdenas et al. (2011).
⁽⁷⁾*Characella* presently classified in the WPD within Pachastrellidae, but is incertae sedis according to Cárdenas et al. (2011).
⁽⁸⁾*Microscleroderma* and its family Scleritodermae presently classified in the WPD within Lithistida, well-known as a polyphyletic group, but is transferred to Spirophorida by Cárdenas et al. (2012).
⁽⁹⁾Formerly identified as *Cinachyrella alloclada* in Sperling et al. (2009) and in GenBank.
⁽¹⁰⁾Presently in WPD as *Axinella corrugata* as part of family Axinellidae within Halichondrida, but see Gazave et al. (2010) and Morrow et al. (2012), who updated its classification.
⁽¹¹⁾*Ectyoplasia* and *Raspailiidae* presently classified in the WPD within Poecilosclerida, but was transferred to Axinellida by Morrow et al. (2012).
⁽¹²⁾Formerly identified as *Damiira* sp. in Sperling et al. (2009) and in GenBank.
⁽¹³⁾Formerly identified as *Tethya aurantia* in Sperling et al. (2009), and as *Tethya actinia* in GenBank.
⁽¹⁴⁾Formerly identified as *Spirastrella coccinea* in Sperling et al. (2009) and in GenBank.
⁽¹⁵⁾Formerly labeled as *Microciona prolifera* in Peterson & Butterfield (2005) and in GenBank, and as *Clathria* (*Microciona*) *prolifera* in Sperling et al. (2009).
⁽¹⁶⁾Derived from genomic traces, as reported in Sperling et al. (2007).
⁽¹⁷⁾Derived from genomic traces, as reported in Sperling et al. (2010).
⁽¹⁸⁾Derived from genomic traces, as reported in Sperling et al. (2009).
doi:10.1371/journal.pone.0050437.t001

Table 2. Summary of genes and taxa for analysis* by poriferan clade.

	ALD	ATPB	CAT	EF1A	MAT	PFK	TPI	NHK7	NHK6	NHK5	NHK4
<i>Keratosa</i> ^P	5	6	6	7	2	5	4	6	6	7	7
<i>Myxospongiae</i> ^P	6	5	5	5	4	0	4	6	6	6	6
<i>Haploscleromorpha</i> ^P	7	7	8	7	5	6	7	9	9	9	9
<i>Spongillida</i> ^P	2	1	2	2	2	1	2	2	2	2	2
<i>Tetractinellida</i> ^P	2	5	3	5	3	1	5	5	5	6	6
Other <i>Democlavida</i> ^P	19	18	17	19	12	15	18	21	21	21	21
<i>Demospongiae</i>^P	41	42	41	45	30	28	40	49	49	51	51
<i>Calcispongiae</i> ^P	7	6	5	7	8	5	5	8	8	8	8
<i>Hexactinellida</i> ^P	2	6	5	4	4	4	4	6	6	6	6
<i>Homoscleromorpha</i> ^P	2	2	2	2	3	2	3	3	3	3	3
TOTAL	54	56	53	58	43	39	52	66	66	68	68

*NHK7 refers to the complete dataset, while NHK6-4 refer to datasets where the markers CAT, EF1A, and ATPB are successively removed.
doi:10.1371/journal.pone.0050437.t002

That said, just five of the resolved nodes in the Bayesian analysis contradict those in the ML-based topology and none of these have pp values > 0.90.

The partitioned ML analysis of the combined data had the same topology as that found when assuming a single model of amino acid evolution (LG+F+G). Additionally, no major differences were found when comparing a Bayesian analysis performed under LG+G, the ML analysis performed using LG+F+G, and the ML analysis performed using multiple partitions. We used this topology as the reference point for comparing the different analyses (Fig. 1). The Bayesian topology (Fig. 2) is highly consistent with the ML-based topology (Table 5). Each of the single-gene ML topologies (Figs. S8, S9, S10, S11, S12, S13, S14) differs from that derived from the combined dataset. An ordered ranking of how well the single-gene topologies match our overall hypothesis, based on nodal difference is: PFK, TPI, ALD, MAT, ATPB, CAT and EF1A (Table 5). This performance is also reflected in a tabulation of whether notable clades were recovered in the single-gene topologies (Table 6), where ATPB, CAT and EF1A recovers less

than half of a set of reference clades in the topology based on the combined data. ML analyses serially excluding CAT, EF1A, and ATPB resulted in topologies (Figs. S15, S16, S17) that are highly consistent with the tree based on the analysis of combined data (Table 5–6). A supertree analysis was performed to evaluate the extent to which the principal signal [53] in the single-gene partitions differed from the signal in the gene concatenation and the results showed a substantial level of agreement (Fig. S18).

Nodal support for the ML-based phylogenetic hypothesis (Fig. 1) varies widely; 46 of 70 nodes have bootstrap support (bs) exceeding 70%. Similarly, although generally higher in magnitude, posterior probability (pp) values in the Bayesian topology are not universally high, with 44 of 70 nodes having values exceeding 0.90 (Fig. 2).

To test whether some of our results could be attributed to tree reconstruction artifacts we performed a variety of analyses. We first built trees using differently fitting models (WAG, LG, GTR, CAT, and CAT-GTR) and compared their results. This analysis indicated an important area of disagreement with reference to the relationships between *Keratosa*^P and *Myxospongiae*^P (see discussion). We performed a posterior predictive analysis to identify compositionally heterogeneous taxa. This analysis indicated that many taxa in the dataset are, indeed, compositionally heterogeneous (Table S1). The 6-categories Dayhoff recoding strategy is commonly used to ease compositional heterogeneity. We recoded our dataset using the 6-categories Dayhoff strategy and performed

Table 3. Amino acid model selection, used for maximum likelihood searches on different datasets*.

Dataset	Most Appropriate Model	Criterion	Model Assumed
NHK7	LG+G+I+F	all AIC	LG+G+F
ALD	LG+G	AICc-1,2	LG+G
ATPB	WAG+G+I	all AIC	WAG+G
CAT	LG+G+I	AIC, AICc-1,3	LG+G
EF1A	LG+G+I+F	AIC, AICc-1,3	LG+G+F
MAT	LG+G+I	AICc-1,2	LG+G
PFK	LG+G	all AIC	LG+G
TPI	LG+G+I	all AIC	LG+G
NHK6	LG+G+I	all AIC	LG+G
NHK5	LG+G+I	all AIC	LG+G
NHK4	LG+G+I	AICc-1,2	LG+G

*NHK7 refers to the complete dataset, while NHK6-4 refer to datasets where the markers CAT, EF1A, and ATPB are successively removed.
doi:10.1371/journal.pone.0050437.t003

Table 4. Model cross validation performed using CAT-GTR as the reference model.

Models Compared	Mean Score	Standard Deviation
CAT+gamma	CAT-GTR+gamma	-66.0556* 27.2128
GTR+gamma	CAT-GTR+gamma	-203.2* 26.4986
LG+gamma	CAT-GTR+gamma	-201.862* 26.7209
WAG+gamma	CAT-GTR+gamma	-226.778* 32.4408

*A negative cross validation score indicates that the reference model (CAT-GTR) fits the data better than the tested model. This table indicates that CAT-GTR provides the best fit to the data (as the standard deviations around the means are not sufficient to define a confidence intervals including positive values).
doi:10.1371/journal.pone.0050437.t004




Figure 1. Hypothesis of demosponge relationships based on maximum likelihood analysis of seven nuclear housekeeping genes. Topology rooted on three cnidarians and the placozoan *Trichoplax*. Bootstrap indices (400 replicates) are shown at each node, with those exceeding 70 in bold. New taxa added as part of the PorToL project are indicated in bold; new taxa added from EST/genomics projects are indicated with a single asterisk; and taxa with new identifications after examination of the voucher specimen are marked with two asterisks. Clade names in italics followed by a superscript p have been phylogenetically defined in other studies (see text). doi:10.1371/journal.pone.0050437.g001

a posterior predictive analysis and found that the Dayhoff recoding eliminated almost all heterogeneity from the data (Table S2). CAT-GTR analyses of the Dayhoff recoded dataset found a tree (Fig. S19) that is highly comparable with the CAT-GTR tree of

Fig. 2 (non-recoded data). However, results of the Bayesian analysis using Dayhoff recoded data and assuming GTR (Fig. S20) contains a key difference. In the Dayhoff recoded GTR analysis *Myxospongiae*^p is not the sister group of *Keratosa*^p but the sister group




Figure 2. Hypothesis of demosponge relationships based on Bayesian analysis of seven nuclear housekeeping genes. Topology rooted on three cnidarians and the placozoan *Trichoplax*. Posterior probabilities are shown at each node, with those exceeding 0.90 in bold. New taxa added as part of the PorToL project are indicated in bold; new taxa added from EST/genomics projects are indicated with a single asterisk; and taxa with new identifications after examination of the voucher specimen are marked with two asterisks. Clade names in italics followed by a superscript p have been phylogenetically defined in other studies (see text). doi:10.1371/journal.pone.0050437.g002

of all the other Demospongiae (albeit with a low PP). Analyses performed after excluding compositionally heterogeneous species, fast-evolving sites, or the outgroups consistently reiterate the

results of our Bayesian analysis (compare Fig. 2 with Figs. S21, S22, S23).

Table 5. Nodal differences between reference topology (ML assuming LG+G+F) and topologies derived from different datasets* and analyses.

Dataset/Analysis	Percentage of Taxa in Common	Nodal Difference	Random Difference	Standard Deviation
ALD	76.4%	2.50	4.49	0.36
ATPB	83.3%	3.42	4.65	0.34
CAT	77.8%	3.48	4.55	0.34
EF1A	86.1%	3.73	4.60	0.37
MAT	65.3%	2.67	4.38	0.34
PFK	58.3%	1.91	4.15	0.34
TPI	77.8%	2.13	4.42	0.29
NHK6	100.0%	1.45	4.80	0.35
NHK5	97.2%	1.42	4.73	0.31
NHK4	97.2%	1.42	4.75	0.38
NHK7/Bayesian	100.0%	1.26	4.77	0.35

*NHK7 refers to the complete dataset, while NHK6-4 refer to datasets where the markers CAT, EF1A, and ATPB are successively removed.
doi:10.1371/journal.pone.0050437.t005

Discussion

Sponge Classes

Analyses of the seven nuclear housekeeping gene dataset provide strong support for each of the four major clades of sponges assigned the rank of class (Calcarea, Demospongiae, Hexactinellida, and Homoscleromorpha). Because we did not include non-metazoan outgroups our results cannot be used to assess sponge monophyly. Concerning the relationships among the four sponge classes, support is generally poor. Our tree does not recover *Silicea*^p (*Demospongiae*^p+*Hexactinellida*^p), which has been supported in a great deal of other works based on disparate datasets [4,18,19,28,54], but instead places *Calcispongiae*^p with *Hexactinellida*^p (Figs. 1–2), most likely erroneously with low support

(bs = 74%; pp = 0.68). Relationships within *Calcispongiae*^p and *Hexactinellida*^p are consistent with previous analyses [54–56]. As designed, our analyses do not provide any basis for inferring relationships among the sponge classes (as they do not include non-metazoan outgroups), but rather elucidate phylogenetic relationships within *Demospongiae*^p (Figs. 1–2).

Major Demosponge Clades

Hypotheses derived from our analyses of nuclear housekeeping gene data (Figs. 1–2) are fairly consistent with the so-called “G clades” originally derived from analysis of ribosomal DNA data [17], and largely recovered by mitochondrial genome [43] and nuclear housekeeping gene data [18]. G1 and G2 correspond to *Keratos*^p and *Myxospongiae*^p, respectively, following the names of

Table 6. Comparison of clades found in NHK7* ML topology with those revealed in single-gene and other analyses.*

Clades of Interest											Bayes		
	ML ALD	ML ATPB	ML CAT	ML EF1A	ML MAT	ML PFK	ML TPI	ML NHK6	M NHK5	ML NHK4	NHK7		
Cnidaria	yes	no	yes	no	yes	yes	no	yes	yes	yes	yes	yes	yes
<i>Calcispongiae</i> ^p	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes
<i>Homoscleromorpha</i> ^p	yes	no	No	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes
<i>Hexactinellida</i> ^p	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes
<i>Demospongiae</i> ^p	yes	no	no	no	no	no	no	yes	yes	yes	yes	yes	yes
<i>Keratos</i> ^p (G1)	yes	yes	yes	no	yes	yes	yes	yes	yes	yes	yes	yes	yes
<i>Myxospongiae</i> ^p (G2)	yes	no	yes	no	yes	–	no	yes	yes	yes	yes	yes	yes
G1+G2	yes	no	no	no	no	–	no	yes	yes	yes	yes	no	no
<i>Spongillida</i> ^p	yes	–	yes	yes	yes	–	yes	yes	yes	yes	yes	yes	yes
<i>Haploscleromorpha</i> ^p (G3)	yes	yes	no	yes	yes	no	yes	yes	yes	yes	yes	yes	yes
<i>Spongillida</i> ^p +G3	no	no	no	no	no	yes	no	yes	yes	yes	yes	yes	yes
<i>Democlavia</i> ^p (G4)	no	no	no	no	no	no	yes	yes	yes	yes	yes	yes	yes
<i>Tetractinellida</i> ^p	yes	yes	no	no	yes	–	yes	yes	yes	yes	yes	yes	yes
G3+G4+ <i>Spongillida</i> ^p	yes	no	no	no	yes	yes	no	yes	yes	yes	yes	yes	yes
Clades	12/14	5/13	5/14	5/14	10/14	7/10	8/14	14/14	14/14	14/14	14/14	13/14	
Percent	86%	38%	36%	36%	71%	70%	57%	100%	100%	100%	100%	93%	

*NHK7 refers to the complete dataset, while NHK6-4 refer to datasets where the markers CAT, EF1A, and ATPB are successively removed.
doi:10.1371/journal.pone.0050437.t006

Borchiellini et al. [17]. One key difference between the results of these studies concerns the placement of the clade containing all freshwater sponges, *Spongillida*^P, phylogenetically defined in Cárdenas et al. [40]. Traditionally, these sponges were classified as the suborder Spongillina within the order Haplosclerida. However, ribosomal DNA and mitochondrial genome data suggested that *Spongillida*^P falls as the earliest diverging lineage of the “G4” clade. Sperling et al. [18] found a similar clade, for which they provided a phylogenetic definition and the name *Democlavida*^P (=subclass Heteroscleromorpha of Cárdenas et al. [40]), with the exception that *Spongillida*^P was found as the sister group of the marine haplosclerids. The marine haplosclerid taxa have consistently been shown to be a well-supported clade that has recently been phylogenetically defined and named *Haploscleromorpha*^P [40].

This study finds strong support at nearly all deep nodes within *Demospongiae*^P (Figs. 1–2), even with our more diverse taxon sampling. The clear distinction of these clades indicates that the divergence among these groups is likely ancient [4]. Thus, future genomic exploration within *Demospongiae*^P will be guided by these emerging phylogenetic results so as to make best use of the comparative method. To be especially useful for rank-based taxonomy and nomenclature, type species within genera and type genera within families (e.g., our sampling of *Spongia officinalis*, *Halisarca dujardini*, and *Desmacella pumilio*) should be targeted whenever possible. Also, to the extent possible, type species should be collected from their respective type localities for maximum taxonomic and nomenclatural utility. For phylogenetic nomenclature, ‘specifiers’ (i.e., species, specimens or apomorphies used in *PhyloCode* definitions) should be targeted. Of course, when species are used as specifiers (which has so far usually been the case for poriferan names), their name-bearing type specimens are *de facto* specifiers (*PhyloCode*, Note 13.2.2.).

Nuclear housekeeping gene data strongly support an as yet unnamed clade containing the groups of demosponges with silica-mineralized skeletons: *Democlavida*^P, *Haploscleromorpha*^P, and *Spongillida*^P (Figs. 1–2), in accordance with other analyses of ribosomal genes [17], complete mitochondrial genomes [43], and a smaller dataset of nuclear housekeeping genes [18]. Our ML and Bayesian analyses provide equivocal results concerning the phylogenetic relationships of *Keratosa*^P and *Myxospongiae*^P. A sister group relationship between *Keratosa*^P and *Myxospongiae*^P has been suggested, with only modest support, based on analyses of 18S rRNA genes [17] and complete mitochondrial genomes [43] but has also been contradicted by earlier Bayesian analyses of nuclear housekeeping genes [4,18,44]. Our ML topology (Fig. 1) shows *Keratosa*^P and *Myxospongiae*^P [which both lack mineralized skeletons (with the exception of siliceous microscleres in *Chondrilla* within *Myxospongiae*^P: Chondrosida)] as a clade that is sister to the mineralized sponges. In contrast, the Bayesian analysis (Fig. 2) identifies *Myxospongiae*^P as the earliest diverging clade of *Demospongiae*^P, and shows *Keratosa*^P as the sister group to the mineralized groups. It is important to note, however, that all site-homogeneous models (LG and GTR) display the *Keratosa*^P+*Myxospongiae*^P clade, while the site-heterogeneous CAT and CAT-GTR models (which fit the data better) support *Myxospongiae*^P as the sister group of all the other demosponges. Thus, model selection is responsible for this disagreement. Because the best fitting models suggest *Myxospongiae*^P is sister to the remaining demosponges, the contradicting results obtained using LG, GTR and WAG (*Keratosa*^P+*Myxospongiae*^P) are likely artifactual.

Keratosa^P

This clade is composed of members of the demosponge orders Dictyoceratida and Dendroceratida. Our sampling includes members of five of the six families: Dysideidae, Irciniidae, Spongiidae and Thorectidae in the former, Dictyodendrillidae in the latter. Ribosomal data [17] indicate that Dendroceratida is monophyletic, but our results rely on a single genus (*Igemella*) so we cannot support or refute that result. The nuclear housekeeping gene data also fail to provide support for the monophyly of Dictyoceratida, a result that has also been obtained through the analysis of ribosomal data [35,57]. We have conflicting results concerning Dictyoceratida, with our ML-topology (Fig. 1) suggesting that dendroceratids are derived from within a paraphyletic Dictyoceratida and the Bayesian tree having a poorly supported monophyletic Dictyoceratida. The key taxon, from the perspective of this analysis, is the representative of Dysideidae. All the other dictyoceratids in our study, representing Irciniidae, Spongiidae, and Thorectidae, always form a well-supported clade. It is interesting to note that when the worst performing markers (CAT, EF1A, and ATPB) are sequentially removed from analysis, Dictyoceratida, including our representative of Dysideidae, forms a monophyletic group with strong support (Figs. S15, S16, S17).

Myxospongiae^P

Members of the orders Chondrosida and Verongida make up *Myxospongiae*^P. Our sampling includes both families of Chondrosida (Chondrillidae and Halisarcidae), the latter of which was previously placed in its own order Halisarcida (e.g., [58]). Within Verongida, just one of the four families of Verongida (i.e., Aplysinidae) is sampled. With the present taxon sampling, our analyses support monophyly of Chondrosida, a result not obtained by some analyses of ribosomal data [17,59], but found in others [35,60]. However, our analysis lacks a representative of *Chondrosia*, which has proven to be a difficult taxon in relation to the question of Chondrosida monophyly [17,59]. Similarly lacking a representative of the problematic *Chondrosia*, an analysis of complete mitochondrial genome data also supports a monophyletic *Chondrosida*^P [43], which has nevertheless recently been given a phylogenetic definition [40]. Within Verongida, nuclear housekeeping genes support monophyly of Aplysinidae, for which we were able to sample each of its component genera (Figs. 1–2). Relationships among the three aplysinid genera (*Verongula*, *Aplysina*, and *Aiolochoiroa*), however, are not well supported. Based on ribosomal data, Erwin and Thacker [61] found that Aplysinidae is not monophyletic because *Verongula* grouped with members of Pseudoceratinidae and members of *Aiolochoiroa* grouped with Ianthellidae and Aplysinellidae. The absence of pseudoceratinids, ianthellids and aplysinellids from our samples prevents our analyses from testing these hypotheses, but if Erwin and Thacker’s [61] findings are true, they would suggest that our sampling represents a more disparate group of Verongida (*Aplysina* in Aplysinidae and *Verongula* in Pseudoceratinidae) than is suggested by current taxonomy (*Aplysina* and *Verongula* in Aplysinidae). Indeed, this phylogenetic result (i.e., that *Aplysina* and *Verongula* belong to distinct families) was recently verified with mitochondrial and nuclear markers by Erpenbeck et al. [59].

Haploscleromorpha^P & *Spongillida*^P

From a broad perspective, one of the most important outstanding questions in demosponge phylogenetics is the phylogenetic placement of the freshwater sponges, *Spongillida*^P, which is phylogenetically defined in Cárdenas et al. [40]. Traditional taxonomy based on morphology [62] and earlier analyses of nuclear housekeeping genes [18] suggest a close relationship

between *Spongillida*^p and the marine haplosclerids, *Haploscleromorpha*^d. In contrast, both mitochondrial genome and ribosomal data suggest that *Spongillida*^p is sister to the rest of the *Democlavia*^d [17,35,43,63]. The results here, for the most part, agree with the former hypothesis and specifically indicate that *Spongillida*^p is the earliest diverging lineage of the traditional order Haplosclerida (with high support, Figs. 1–2). An exception to this result is one of the single gene analyses (ALD, Fig. S8), which found *Spongillida*^p branching among democlaviid taxa, albeit with no support. Limited taxon sampling, and in particular, the fact that our analyses do not include any representatives of the democlaviid family Scopalinidae (which was recently suggested by Morrow et al. [38] to have a close relationship to the freshwater sponges), could explain these contradicting results. In any event, it is fairly clear that *Spongillida*^p is a distinct lineage from the marine haplosclerids.

Our sampling within *Haploscleromorpha*^d represents five of the six accepted families. Monophyletic haplosclerid suborders Petrosina and Haplosclerina were not recovered (although support values are somewhat low at some of the deeper branches of the clade), corroborating the results of McCormack et al. [64] and Redmond et al. [35,37]. Not surprisingly, given that studies with denser taxon sampling have shown widespread polyphyly of subtaxa within this group [35,37,65], we find both Petrosiidae and Niphatidae to be polyphyletic. Even at the generic level, *Amphimedon* (Niphatidae) is revealed to be polyphyletic. *Amphimedon queenslandica*, whose genome has been sequenced [66], clusters with *Callyspongia vaginalis* (Callyspongiidae) with high support, suggesting that the taxonomy of this important model organism remains confused, corroborating evidence from ribosomal data [35,37].

Democlavia^p

Democlavia^d is the most species-rich (roughly 75% of demosponge species; [38]) and diverse of the major demosponge clades, and includes the traditional orders Agelasida, Astrophorida, Hadromerida, Halichondrida, Poecilosclerida, and Spirophorida [48], several of which are already thought to not be monophyletic (as discussed below). As such, the systematics of *Democlavia*^d presents many challenges, but important breakthroughs are being made in understanding the phylogeny of this clade based on increasingly taxon-rich analyses of ribosomal RNA and mitochondrial CO1 data [38]. Our nuclear housekeeping gene dataset and analyses provide an opportunity to test hypotheses arising from these alternative sets of data and suggest new hypotheses where previous results have provided no resolution.

Our analyses reveal a well-supported clade containing members of Astrophorina and Spirophorina (suborder designations for these taxa, following [40]), including our only sampled lithistid (*Microscleroderma* sp. nov.). Other analyses of ribosomal and mitochondrial data have revealed the same clade [17,35,42,67–69], the phylogenetically defined *Tetractinellida*^d [17,40]. Although modest in support, our analyses always suggest that *Tetractinellida*^d is sister to the remaining members of *Democlavia*^d. Our sampling of sub-order Astrophorina includes two of the six families, Ancorinidae (*Dercitus*, recently transferred from Pachastrellidae by Cárdenas et al. [70]) and Geodiidae (*Geodia tumulosa* and *Geostelletta fibrosa*), as well as an *incertae sedis* taxon, *Characella* aff. *connectens*, which was also formerly assigned to family Pachastrellidae. The latter three species form a well-supported clade, but no specific position for our representative of Ancorinidae within *Tetractinellida*^d is supported (Figs. 1–2). The family Pachastrellidae *sensu* Maldonado [71] is based on a plesiomorphic character (streptasters; [70]) so it is no surprise that our results confirm that *Characella* and *Dercitus* do not have an especially close relationship.

Our analyses include two representatives of Spirophorina – *Cinachyrella* sp., representing the family Tetillidae, and the lithistid *Microscleroderma* sp. nov., representing the family Scleritodermidae – but there is no support for the group being monophyletic. The lithistids are a taxonomically rich group sharing a common growth form (skeleton of interlocked desmas), with 13 recognized families. Lithistids have always presented taxonomic challenges from morphological perspectives (see [72]) and the redistribution of its members to different sponge clades has been proposed for quite some time [72,73] and continues [40]. In this vein, the lithistid family Desmanthidae appears to be closely related to Dictyonellidae [38]. The presence of sigmaspires in Scleritodermidae [72] is consistent with this group being reallocated to Spirophorina within *Tetractinellida*^d [40].

Another well-supported alliance of taxa includes most members of order Poecilosclerida that we have sampled, specifically representatives of Coelosphaeridae, Crambeidae, Hymedesmiidae, Microcionidae, Mycalidae, and Tedaniidae (Figs. 1–2). Monophyly of Poecilosclerida has been found in several analyses of ribosomal data [17,35,42,74], but more recent studies with greater taxon sampling have shown the group to be polyphyletic [38,75], as found here. Morrow et al. [38] demonstrated that the families Desmacellidae and Raspailiidae should be removed from Poecilosclerida. Our results support this action, as our representatives of these families branch deeper within *Democlavia*^d (Figs. 1–2). Unfortunately, these data do not provide strong support for relationships within this poecilosclerid group, which remains the most species-rich order and therefore one of the more challenging clades within *Demospongiae*^d.

The sister group to Poecilosclerida (*sensu* [38]) consists of most of our sampled hadromerids as well as the family Halichondriidae from the order Halichondrida. A similar relationship was derived in Morrow et al. [38]. Within this clade, three hadromerids, *Cliona* (Clionidae), *Placospongia* (Placospongiidae), and *Spirastrella* (Spirastrellidae) form a well-supported clade. In turn, this clade is revealed to have a relatively well-supported relationship with the families Halichondriidae and Suberitidae. The latter two families, currently classified within Halichondrida and Hadromerida, respectively, have long been known to have a close relationship [27]. Interestingly, the hadromerid *Tethya* (Tethyidae) consistently branches with this alliance of Suberitidae, Halichondriidae, and the hadromerids (representing Clionidae, Placospongiidae and Spirastrellidae) albeit with limited support. One other hadromerid in our analysis, *Polymastia tenax*, falls outside this clade, a peculiar result given that Polymastiidae is considered among the “core” components of Hadromerida [76]. In the 28S-based analysis of Morrow et al. [38], Polymastiidae emerged as a distinct clade, sister to Suberitidae plus Halichondriidae but with low support, whereas their analysis of CO1 data recovered a clade with Polymastiidae sister to the hadromerid families Tethyidae, Hemiasterellidae, and Clionidae, but again with only low support.

The monophyly of *Agelasida*^d is well supported. This result is obtained only after taking into account recent findings made by Gazave et al. [36], who provided a phylogenetic definition of the clade, and corroborated by Morrow et al. [38]. In light of polyphyly of *Axinella* (order Axinellida), Gazave et al. [36] erected the taxon *Cymbaxinella*^d for those species, including *Axinella corrugata* sampled here, with a close relationship to *Agelas* (family Agelasidae). With broader taxon sampling, Morrow et al. [38] established the new family Hymerhabdiidae for this same clade within *Agelasida*^d. In contrast with this study [38], however, nuclear housekeeping gene data do not provide further support for a sister group relationship between *Agelasida*^d and the clade containing the

core poecilosclerids, hadromerids and Halichondriidae. The only representative of order Axinellida in our analysis is *Ectyoplasia*; the species belongs to the family Raspailiidae, which was moved from Poecilosclerida to Axinellida by Morrow et al. [38]. That study [38] also found that representatives of Desmacellidae fell in two groups, a finding we also recovered given that *Desmacella* and *Biemna* did not exhibit a particularly close relationship. It is important to note that our analysis includes the type species of *Desmacella*. Nuclear housekeeping gene data provide modest support for a relationship between *Desmacella* and the family Dictyonellidae (Figs. 1–2).

Conclusions

As with any phylogenetic analysis, the hypotheses presented here do not represent the final statement on demosponge phylogeny. In particular, the aforementioned gaps in taxonomic sampling limit the extent to which these analyses are able to assess interesting and relevant hypotheses of demosponge relationships. Nonetheless, this analysis makes several important strides forward. First, our results bolster previous claims of the efficacy of the nuclear housekeeping gene marker set [44], albeit at a high cost in effort. Analyses of these data with enhanced taxon sampling confirm numerous phylogenetic hypotheses derived from ribosomal DNA and mitochondrial markers. Most importantly, this boosts overall confidence in the emerging picture of demosponge systematics and phylogenetics that has largely been based on ribosomal and mitochondrial markers, which are more readily obtained from sponge samples. Nevertheless, there are still key points of difference, for example the position of the freshwater *Spongillida*^d clade, that remain to be tested by new datasets, and numerous open questions not yet satisfactorily answered by any phylogenetic analyses, such as the position of *Tetractinellida*^d within *Democlavia*^d, and the relationships among *Keratosd*^d, *Myxospongiae*^d, and the clade consisting of *Democlavia*^d, *Haploscleromorpha*^d, and *Spongillida*^d. A final important advance of this study is that incorporates a diverse set of sponge systematists engaged in transforming the taxonomy (both *PhyloCode*-based and more traditional approaches) used to describe demosponge diversity. As a new understanding of demosponge relationships emerges, the names – and possibly the rules by which we erect and use them – must change [38–41].

Materials and Methods

Ethics Statement

In accordance with policy and legal requirements associated with specimens vouchered in the collections of the Smithsonian US National Museum of Natural History (NMNH), Harbor Branch Oceanographic Institute (HBOI), Harvard Museum of Comparative Zoology (MCZ), and Zoological Museum Bergen Norway (ZMBN), all collections involved in this study were obtained with all appropriate and relevant permits. Specifically, samples from Panama were collected under a Marine Collecting Permit provided by The Republic of Panama; samples from the State of Florida were collected under a Florida recreational resident saltwater fishing license issued from Florida Fish and Wildlife Conservation Commission; and one sample from Honduras was collected with the permission of Rosa del Carmen Garcia, Directora General de Pesca y Acuicultura. No permits were required to collect sponge specimens in US territorial waters outside state boundaries, the Catalan coast of Spain, Vancouver Island, Canada, or Norway.

Sample and sequence collection

Samples were collected from a variety of locations and stored as described below or obtained from frozen collections at the Harbor Branch Oceanographic Institute-Florida Atlantic University (Table 1; <http://PorToL.org/NHK7data>). To obtain RNA of sufficient quality and quantity, when possible, fresh material was collected and preserved via one of several methods. One involved placing fresh material in cold 75% ethanol with liquid changes occurring after 15 min, 1 hour and 4 hours. When available, material was also placed in RNAlater (Invitrogen), directly in TRIzol[®] (Invitrogen) reagent, following the manufacturer's instructions, or in liquid nitrogen. In most cases, the tissue placed directly in TRIzol[®] or frozen in liquid nitrogen yielded the highest quality and/or quantity of RNA. However, the most practical storage method in the field was 75% ethanol preservation and in most cases this was suitable for RNA extraction and subsequent polymerase chain reaction (PCR) amplifications from cDNA.

Following Sperling et al. [18,44] total RNA was isolated using a one-step TRIzol[®] method (Invitrogen), and cDNA was synthesized from 1–2 µg RNA using RETROSCRIPT[®] (Ambion) reverse transcriptase using both random decamers and oligo dT primers, which were then pooled. PCR was used to amplify 7 nuclear-encoded genes: aldolase (ALD), ATP synthase beta chain (ATPB), catalase (CAT), elongation factor 1-alpha (EF1alpha), methionine adenosyltransferase (MAT), phosphofructokinase (PFK), and triose-phosphate isomerase (TPI). All primer sequences for initial PCR of housekeeping genes can be found in Sperling et al. [44]. In many cases, however, it was necessary to use nested PCR primers if amplification and re-amplification of housekeeping gene products was not possible. Table S3 provides primer sequences for nested amplifications of individual housekeeping genes. Primary or nested amplification products were cloned into PCR cloning vectors (pGEM[®]-T, Promega or TOPO TA[®], Invitrogen) and individual clones were prepared for DNA sequencing using standard protocols.

After editing and trimming vector sequences with GENEIOUS [77], DNA sequences were assessed for gene and sponge identity via BLASTX or BLASTP queries [78], followed by preliminary single-gene phylogenetic analyses under the likelihood framework described below. The identification of likely paralogs followed standard procedures based on the generation of trees including all the members of each gene family that could be identified in GenBank (via reciprocal blasting). Within the context of these trees, paralogy groups were identified and only the sequences nesting within the selected orthology group were used. New sequences generated in this study have been submitted to GenBank (Table 1). Sequences are also available via the Porifera Tree of Life database (PorToL.org). In addition, voucher specimens for many of the sequences presented in Sperling et al. [18,44] were examined, resulting in several instances of updated taxonomic identification and classification (Table 1).

Nucleotide sequences were translated and aligned using MUSCLE [79] and visualized in SEAVIEW (v. 4.3) [80]. In addition to the new sequences, the initial alignment included data for sponges that had already been published (Table 1). Also, five species for which transcriptome data exist were also added to the dataset. Both mRNA and cDNA from *Corticium candelabrum*, *Petrosia ficiformis* and *Sycon coactum* were obtained using protocols available in Riesgo et al. [81]. *Sycon ciliatum* and *Leucosolenia complicata* sequences are derived from current genome and transcriptome sequencing projects for these species [82] and Adamska, unpublished). *De novo* assemblies of the reads obtained with Illumina GA (Illumina, Solexa, USA) were built with CLC Genomics Workbench 4 (CLCbio, MA, USA). Local blasts against the contig lists

generated were used to search for the housekeeping genes. Initially, 50 outgroup taxa representing Bilateria, Ctenophora, Cnidaria, Placozoa and non-metazoan Opisthokonta were included in the analyses. However, preliminary phylogenetic analyses, conducted as described below, indicated that inferred demosponge relationships were robust to outgroup choice and therefore outgroups in the final dataset were reduced to the cnidarian taxa (*Acropora*, *Metridium* and *Nematostella*) and the placozoan *Trichoplax*. Approximately 40 positions in the alignment were manually excluded from analyses because they represented insertions present in one or a small number (<5) of taxa.

Phylogenetic Analyses

For all gene trees we investigated the presence of significant clustering information using Maximum Likelihood Mapping [83] as implemented in TreePuzzle V. 5.2 [51]. The dataset was analyzed in both Bayesian and Maximum Likelihood (ML) frameworks. For the ML analyses, appropriate models of amino acid evolution were assessed using the Akaike Information Criterion (AIC), as implemented in ProtTest (v.2.4) [49]. The computing cluster of the Smithsonian's Laboratories of Analytical Biology was used to run the parallelized version of RaxML [84] to search for maximum likelihood (ML) topologies. We assumed the model that best fit our data according to the second-order AIC (AICc-1) with the exception that a proportion of invariant sites was not estimated (according to a recommendation in the RaxML manual). We also used RaxML to conduct bootstrap analyses (400 replicate searches) to assess nodal support. We searched for ML topologies using each gene separately as well as all genes combined. We analyzed the combined data a) assuming a single model for all the data and b) by assigning most appropriate models to each gene partition (mixed models).

Bayesian analyses were performed using the site-heterogeneous CAT-GTR+gamma in Phylobayes 3.3b [85]. This model was selected because Bayesian model selection, performed using 10-fold cross-validation [86], showed that CAT-GTR best fitted our dataset, outperforming CAT, GTR and LG. The considered models were: WAG, LG, GTR, CAT, and CAT-GTR (all models used a gamma correction to account for rate heterogeneity among sites). The CAT based models (in this case CAT and CAT-GTR [86]) are mixture models developed to better take into account site-specific features of protein evolution. These models are thus expected to fit the data better than homogeneous time reversible models like LG and GTR [86]. Indeed, CAT based models have previously been shown to fit amino acid datasets better than other models and they have been shown to be highly effective at reducing systematic biases, like long branch attraction, which are well known to be very pervasive in deep time phylogenetics. In Phylobayes two independent analyses were run for 30,000 cycles sampling every 100 points. The analyses were considered converged when the largest discrepancy observed across all bipartitions (i.e. the maxdiff statistics) dropped below 0.15, despite the Phylobayes manual's suggestion that a chain has reached convergence when maxdiff <0.3. Support values for the nodes recovered in the CAT-GTR analysis are expressed as posterior probabilities.

Comparisons were made between the different single-gene topologies and the Bayesian topology to the ML tree derived from the overall data. In addition, nodal differences were calculated, as measured by the root-mean-squared distance, in Topd (v.3.3) [87]. Taxa that were missing data for some genes were pruned from the combined tree prior to calculating nodal differences. Topd was also used to conduct randomization analyses to test whether similarities between the various topologies and the combined ML

topology were not greater than expected by chance. Finally, further ML searches were conducted by sequentially excluding the three genes that subtend the trees that are most distant from the tree derived from the concatenated dataset, as measured by subtracting the random nodal difference from the actual nodal difference. To further investigate the extent to which the principal signal [53] in the single-genes corroborated the results of concatenated Bayesian and ML analyses, we performed a supertree analysis. The supertree was built using the Matrix Representation with Parsimony method [53]. Input trees used for this analysis were, for each gene, the 400 bootstrap trees derived (see above) under ML. This set of 2800 input trees was bootstrapped to generate 100 replicate datasets, each of which scored 2800 trees using the software CLANN [88]. For each bootstrapped dataset a bootstrap supertree was recovered and a majority rule consensus of the recovered bootstrap supertrees was built to estimate nodal support.

Finally, analyses were performed to test for tree reconstruction artifacts. More precisely we investigated the potential effect of long-branch attraction and compositional attraction on our results. We first investigated the effect of using alternative model of evolution on our results. We thus built trees (within a Bayesian framework) using models (WAG, LG, GTR, CAT, and CAT-GTR, each with a gamma correction) providing different levels of fit to the data and compared the trees we obtained. We tested whether the taxa in our dataset were compositionally heterogeneous performing a posterior predictive analysis (see for example [18]) of compositional heterogeneity using Phylobayes under the CAT-GTR model. The posterior predictive analysis indicated that several taxa displayed a biased composition of their sites. This, if not addressed, can cause compositional artifacts. To test whether our results were affected by compositional biases we performed two analyses. First we analysed (under CAT-GTR) a dataset from which all compositionally heterogeneous taxa were excluded. This experiment has the downside of excluding potentially important taxa. Accordingly, a second experiment was performed in which our dataset was recoded using the Dayhoff scheme. Dayhoff recoding can alleviate compositional artifact, and a posterior predictive analysis of our Dayhoff-recoded dataset was performed (under CAT-GTR) to evaluate whether further compositionally biased taxa remained after the application of Dayhoff recoding. Finally, our Dayhoff recoded dataset was analysed using both a site-homogeneous (GTR) and a site heterogeneous (CAT-GTR) model.

To test for the potential effect of long-branch attraction artifacts we identified fast evolving sites in our dataset using the program Tiger [89]. After that, sites that Tiger deemed as being fast evolving (bins 7 to 10) were excluded and the slowly evolving sites analysed in isolation. In addition to the site-stripping analysis, we also performed an analysis where all the outgroups to Demospongiae (including Hexactinellida) were removed.

Supporting Information

Figure S1 Maximum Likelihood Mapping shows ALD has signal to resolve unambiguously over 90% of the quartets that make up the ALD-derived tree. ALD cannot resolve 4.4% of the quartets. (PDF)

Figure S2 Maximum Likelihood Mapping shows ATPB has signal to resolve unambiguously over 82% of the quartets that make up the ATPB-derived tree. ATPB cannot resolve 8% of the quartets. (PDF)

Figure S3 Maximum Likelihood Mapping shows CAT has signal to resolve unambiguously over 82% of the quartets that make up the CAT-derived tree. CAT cannot resolve 9% of the quartets.
(PDF)

Figure S4 Maximum Likelihood Mapping shows EF1a has signal to resolve unambiguously over 76% of the quartets that make up the EF1a-derived tree. EF1a cannot resolve 12.3% of the quartets.
(PDF)

Figure S5 Maximum Likelihood Mapping shows MAT has signal to resolve unambiguously nearly 83% of the quartets that make up the MAT-derived tree. MAT cannot resolve 10.2% of the quartets.
(PDF)

Figure S6 Maximum Likelihood Mapping shows PFK has signal to resolve unambiguously over 71% of the quartets that make up the PFKtree. PFK cannot resolve 20.6% of the quartets.
(PDF)

Figure S7 Maximum Likelihood Mapping shows TPI has signal to resolve unambiguously over 76% of the quartets that make up the TPI-derived tree. TPI cannot resolve 15.8% of the quartets.
(PDF)

Figure S8 Maximum Likelihood topology based on ALD, with assumed model of LG+gamma.
(PDF)

Figure S9 Maximum Likelihood topology based on ATPB, with assumed model of WAG+gamma.
(PDF)

Figure S10 Maximum Likelihood topology based on CAT, with assumed model of LG+gamma.
(PDF)

Figure S11 Maximum Likelihood topology based on EF1A, with assumed model of LG+F+gamma.
(PDF)

Figure S12 Maximum Likelihood topology based on MAT, with assumed model of LG+gamma.
(PDF)

Figure S13 Maximum Likelihood topology based on PFK, with assumed model of LG+gamma.
(PDF)

Figure S14 Maximum Likelihood topology based on TPI, with assumed model of LG+gamma.
(PDF)

Figure S15 Maximum Likelihood topology based on NHK6, with assumed model of LG+gamma.
(PDF)

Figure S16 Maximum Likelihood topology based on NHK5, with assumed model of LG+gamma.
(PDF)

Figure S17 Maximum Likelihood topology based on NHK4, with assumed model of LG+gamma.
(PDF)

Figure S18 Consensus supertree derived from the input trees that represents the signal in the collection of the individual trees.
(PDF)

Figure S19 Bayesian analysis of Dayhoff recoded data using CAT-GTR.
(PDF)

Figure S20 Bayesian analysis of Dayhoff recoded data using GTR.
(PDF)

Figure S21 Bayesian analysis using CAT-GTR, with all compositionally heterogeneous taxa excluded.
(PDF)

Figure S22 Bayesian analysis using CAT-GTR, excluding fast-evolving sites with Tiger software (“SlowFast Tree”).
(PDF)

Figure S23 Bayesian analysis using CAT-GTR, with no outgroups.
(PDF)

Table S1 Results of the Posterior Predictive Analysis of the combined data set (all 7 genes) under the CAT GTR model. Taxa with a star are heterogeneous in composition.
(PDF)

Table S2 An analysis of the Dayhoff recoded dataset (still under CAT-GTR). As expected, nearly all the heterogeneity is gone (compared to Table S1).
(PDF)

Table S3 Nested primers used to facilitate amplifications of 5 of the 7 genes analyzed in this work.
(PDF)

Acknowledgments

We are indebted to the many sponge biologists who contributed samples, expertise, and sequences to this project. Special thanks go to Sebastian Joffre and Crystal Romero for their assistance in sponge RNA preparations, Sebastian Basile, Tandace Burkhart and Renee Potens with sequence data curation and bioinformatics analyses, and Chris Freeman for assistance in collecting specimens. We thank Martin Dohrmann for identifying hexactinellid samples and providing helpful comments on an earlier version of this MS. We also thank Gonzalo Giribet and Sally Leys for guidance provided to obtain all the transcriptomic sequencing for the species *Petrosia ficiformis*, *Sycon coactum*, and *Corticium candelabrum*.

Author Contributions

Conceived and designed the experiments: MSH ALH JL KJP SP MCD RWT EAS AGC. Performed the experiments: MSH ALH JL MA AC ED BD DF JP D. Pohlmann NER SR AR ER ZR EAS Mds JET. Analyzed the data: MSH D. Pisani AGC. Contributed reagents/materials/analysis tools: SP JR. Wrote the paper: MSH ALH JL KJP MCD RWT MA NBE PC EH GLH SK CCM D. Pisani NER AR KR EAS AGC BP.

References

- Gehling JG, Rigby JK (1996) Long expected sponges from the neoproterozoic Ediacara fauna of South Australia. *J Paleol* 2: 185–195.
- Love GD, Grosjean E, Stalvics C, Fike DA, Grotzinger JP, et al. (2009) Fossil steroids record the appearance of Demospongiae during the Cryogenian period. *Nature* 457: 718–721.
- Brain CKB, Prave AR, Hoffmann K-H, Fallick AE, Botha A, et al. (2012) The first animals: ca. 760-million-year-old sponge-like fossils from Namibia. *S Afr J Sci* 108:658.
- Sperling EA, Robinson JM, Pisani D, Peterson KJ (2010) Where's the glass? Biomarkers, molecular clocks, and microRNAs suggest a 200-Myr missing Precambrian fossil record of siliceous sponge spicules. *Geobiology* 8: 24–36.

5. Erwin DH, Laflamme M, Tweedt SM, Sperling EA, Pisani D, et al. (2011) The Cambrian conundrum: early divergence and later ecological success in the early history of animals. *Science* 334: 1091–1097.
6. Krautter M, Conway KW, Barric JV, Neuweiler M (2001) Discovery of a “living dinosaur”: Globally unique modern hexactinellid sponge reefs off British Columbia, Canada. *Facies* 44: 265–282.
7. Milsom C, Rigby S (2010) *Fossils at a glance*, 2nd Edition. Wiley-Blackwell.
8. Hill MS (1998) Spongivory on Caribbean reefs releases corals from competition with sponges. *Oecologia*, 117: 143–150.
9. Rützler K (2004) Sponges On Coral Reefs: A Community Shaped By Competitive Cooperation. *Boll Mus Ist Biol Univ Genova* 68: 85–148.
10. De Goeij JM, van den Berg H, van Oostveen MM, Epping EHG, van Duyl FC (2008) Major bulk dissolved organic carbon (DOC) removal by encrusting coral reef cavity sponges. *Mar Ecol Prog Ser* 357: 139–151.
11. Erwin PM, Thacker RW (2008) Cryptic diversity of the symbiotic cyanobacterium *Synechococcus spongianum* among sponge hosts. *Mol Ecol* 17: 2937–47.
12. Venkateswara Rao J, Srikanth K, Pallela R, Gnaneshwar Rao T (2009) The use of marine sponge *Haliclona tenuiramosa* as bioindicator to monitor heavy metal pollution in the coasts of Gulf of Mannar, India. *Environ Monit Assess* 156: 451–459.
13. Negandhi K, Blackwelder PL, Ereskovsky AV, Lopez JV (2010) Florida Reef Sponges Harbor Coral Disease-Associated Bacteria. *Symbiosis* 51: 117–129.
14. Pawlik JR (2011) The chemical ecology of sponges on Caribbean reefs: Natural products shape natural systems. *BioScience* 61: 888–898.
15. Webster NS, Taylor MW (2012) Marine sponges and their microbial symbionts: love and other relationships. *Environ Microbiol* 14: 335–346. doi:10.1111/j.1462-2920.2011.02460.x
16. Hooper JNA, van Soest RWM (2002) *Systema Porifera. A guide to the classification of sponges*. New York: Kluwer Academic/Plenum Publishers. 1810 p.
17. Borchellini C, Chombard C, Manuel M, Alivon E, Vacelet J, et al. (2004) Molecular phylogeny of Demospongiae: implications for classification and scenarios of character evolution. *Mol Phylogenet Evol* 32: 823–837.
18. Sperling EA, Peterson KJ, Pisani D (2009) Phylogenetic-signal dissection of nuclear housekeeping genes supports the paraphyly of sponges and the monophyly of Eumetazoa. *Mol Biol Evol* 26: 2261–2274.
19. Philippe H, Derelle R, Lopez P, Pick K, Borchellini C, et al. (2009) Phylogenomics revives traditional views on deep animal relationships. *Curr Biol* 19: 706–712.
20. Philippe H, Brinkmann H, Lavrov DV, Littlewood DTJ, Manuel M, et al. (2011) Resolving difficult phylogenetic questions: why more sequences are not enough. *PLoS Biology* 9(3): e1000602. doi:10.1371/journal.pbio.1000602
21. Rützler K, Hooper JNA (2000) Two new genera of hadromerid sponges (Porifera, Demospongiae). *Zoosystema* 22: 337–344.
22. Ferrario F, Calcinaï B, Erpenbeck D, Galli P, Wörheide G (2010) Two *Pione* species (Hadromerida, Clionaidae) from the Red Sea: a taxonomical challenge. *Org Divers Evol* 10: 275–285.
23. de Paula, Zilberberg C, Hajdu E, Lóbo-Hajdu G (2012) Morphology and molecules on opposite sides of the diversity gradient: Four cryptic species of the *Cliona celata* (Porifera, Demospongiae) complex in South America revealed by mitochondrial and nuclear markers. *Mol Phylogenet Evol* 62: 529–541.
24. van Soest RWM (1990) Towards a phylogenetic classification of sponges. In: Rützler K, editor. *New Perspectives in Sponge Biology. Proceedings of the 3rd International Conference on the Biology of Sponges*, Woods Hole, 1985. Washington: Smithsonian Institution Press. pp. 344–350.
25. Lafay B, Boury-Esnault N, Vacelet J, Christen R (1992) An analysis of partial 28S ribosomal RNA sequences suggests early radiations of sponges. *BioSys* 28: 139–151.
26. West L, Powers D (1993) Molecular phylogenetic position of the hexactinellid sponges in relation to the Protista and Demospongiae. *Mol Mar Biol Biotech* 2: 71–75.
27. Chombard C (1998) *Les Demospongiae à asters: essai de phylogénie moléculaire. Homologie du caractère ‘aster’*. Ph.D. Thesis, Museum national d’Histoire naturelle de Paris.
28. Collins AG (1998) Evaluating multiple alternative hypotheses for the origin of Bilateria: An analysis of 18S rRNA molecular evidence. *Proc Natl Acad Sci* 95: 15458–15463.
29. Hajdu E (1999) Toward a phylogenetic classification of the Mycalids with anisochelae (Demospongiae: Poecilosclerida), and comments on the status of *Naviculina* Gray, 1867. *Mem Queensland Mus* 44: 225–238
30. Alvarez B, Crisp MD, Driver F, Hooper JNA, van Soest RWM (2000) Phylogenetic relationships of the family Axinellidae (Porifera: Demospongiae) using morphological and molecular data. *Zool Scr* 29: 169–198.
31. Borchellini C, Chombard C, Lafay B, Boury-Esnault N (2000) Molecular Systematics of sponges (Porifera). *Hydrobiologia* 420: 15–27.
32. Hajdu E, van Soest RWM, Hooper JNA (1994) Proposal for a phylogenetic subordinal classification of poecilosclerid sponges. In: van Soest RWM, van Kempen TMG, Brackman J-C, editors. *Sponges in Time and Space - Proceedings of the 4th International Porifera Congress*. Rotterdam: Balkema. pp. 123–139.
33. Lavrov DV, Lang BF (2005) Poriferan mtDNA and animal phylogeny based on mitochondrial gene arrangements. *Syst Biol* 54: 651–659.
34. Wörheide G, Solé-Cava AM, Hooper JNA (2005) Biodiversity, molecular ecology and phylogeography of marine sponges: patterns, implications and outlooks. *Integr Comp Biol* 43: 377–385.
35. Redmond NE, van Soest RWM, Kelly M, Raleigh J, Travers SAA, et al. (2007) Reassessment of the classification of the Order Haplosclerida (Class Demospongiae, Phylum Porifera) using 18S rRNA gene sequence data. *Mol Phylogenet Evol* 43: 344–352.
36. Gazave E, Carteron S, Chenuil A, Richelle-Maurer E, Boury-Esnault N, et al. (2010) Polyphyly of the genus *Axinella* and of the family Axinellidae (Porifera: Demospongiae^s). *Mol Phylogenet Evol* 57: 35–47.
37. Redmond NE, Raleigh J, van Soest RWM, Kelly M, Travers SAA, et al. (2011) Phylogenetic Relationships of the Marine Haplosclerida (Phylum Porifera) Employing Ribosomal (28S rRNA) and Mitochondrial (*cox1*, *nad1*) Gene Sequence Data. *PLoS ONE* 6(9): e24344.
38. Morrow CC, Pictou BE, Erpenbeck D, Boury-Esnault N, Maggs CA, et al. (2012) Congruence between nuclear and mitochondrial genes in Demospongiae: A new hypothesis for relationships within the G4 clade (Porifera: Demospongiae). *Mol Phylogenet Evol* 62: 174–190.
39. Hooper JNA, van Soest RWM, Pisera A (2011) *Phylum Porifera Grant, 1826*. In: Zhang, Z-Q, editor. *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa* 3148: 13–18.
40. Cárdenas P, Pérez T, Boury-Esnault N (2012) Sponge systematics facing new challenges. *Adv Mar Biol* 61: 79–209.
41. Wörheide G, Dohrmann M, Erpenbeck D, Larroux C, Maldonado M, et al. (2012) Deep phylogeny and evolution of sponges (Phylum Porifera). *Adv Mar Biol* 61: 1–78.
42. Nichols SA (2005) An evaluation of support for order-level monophyly and interrelationships within the class Demospongiae using partial data from the large subunit rDNA and cytochrome oxidase subunit I. *Mol Phylogenet Evol* 34: 81–96.
43. Lavrov DV, Wang X, Kelly M (2008) Reconstructing ordinal relationships in the Demospongiae using mitochondrial genomic data. *Mol Phylogenet Evol* 49: 111–124.
44. Sperling EA, Pisani D, Peterson KJ (2007) Poriferan paraphyly and its implications for Precambrian paleobiology. In Vickers-Rich P, Komarow P, editors. *The Rise and Fall of the Ediacaran Biota*. Geol Soc, Special Publications, London, pp. 355–368.
45. Peterson KJ, Lyons JB, Nowak KS, Takacs CM, Wargo MT, et al. (2004) Estimating metazoan divergence times with a molecular clock. *Proc Natl Acad Sci USA* 101: 6536–6541.
46. Peterson KJ, Butterfield NJ (2005) Origin of the Eumetazoa: testing ecological predictions of molecular clocks against the Proterozoic fossil record. *Proc Natl Acad Sci USA* 102: 9547–9552.
47. Reed JK, Pomponi SA (1997) Biodiversity and distribution of deep and shallow water sponges in the Bahamas. *Proc 8th Int Coral Reef Symp* 2:1387–1392.
48. van Soest, RWM, Boury-Esnault N, Hooper JNA, Rützler K, de Voogd NJ, et al. (2012) World Porifera database. Available: <http://www.marinespecies.org/porifera>. Accessed 2012 Aug 1.
49. Abascal F, Zardoya R, Posada D (2005) ProtTest: Selection of best-fit models of protein evolution. *Bioinformatics*: 21: 2104–2105. doi:10.1093/bioinformatics/bti263
50. Le SQ, Gascuel O (2008) LG: An Improved, General Amino-Acid Replacement Matrix. *Mol Biol Evol* 25: 1307–1320.
51. Schmidt HA, Strimmer K, Vingron M, von Haeseler A (2002) TREE-PUZZLE: maximum likelihood phylogenetic analysis using quartets and parallel computing. *Bioinformatics* 18: 502–504.
52. Stone M (1974) Cross-validators choice and assessment of statistical prediction. *J R Stat Soc Ser B* 36:111–147.
53. Pisani D, Wilkinson M (2002) Matrix representation with parsimony, taxonomic congruence, and total evidence. *Syst Biol* 51: 151–155.
54. Dohrmann M, Janussen D, Reitner J, Collins AG, Wörheide G (2008) Phylogeny and evolution of glass sponges (Porifera, Hexactinellida). *Syst Biol* 57: 388–405.
55. Manuel M, Borchellini C, Alivon E, Le Parco Y, Vacelet J, et al. (2003) Phylogeny and evolution of calcareous sponges: Monophyly of Calcinea and Calcarenea, high level of morphological homoplasy, and the primitive nature of axial symmetry. *Syst Biol* 52: 311–333.
56. Dohrmann M, Voigt O, Erpenbeck D, Wörheide G (2006) Non-monophyly of most supraspecific taxa of calcareous sponges (Porifera, Calcarea) revealed by increased taxon sampling and partitioned Bayesian analysis of ribosomal DNA. *Mol Phylogenet Evol* 40: 830–843.
57. Ma J, Yang Q (2007) Phylogeny and Morphogenesis of Dictyoceratida (Porifera: Demospongiae). *Acta Palaeontologica Sinica* 46: 480–484.
58. Ereskovsky A, Lavrov DV, Boury-Esnault N, Vacelet J (2011) Molecular and morphological description of a new species of *Halisarca* (Demospongiae: Halisarcida) from Mediterranean Sea and a redescription of the type species *Halisarca diyardini*. *Zootaxa* 2768: 5–31.
59. Erpenbeck D, Sutcliffe P, Cook SdC, Dietzel A, Maldonado M, et al. (2012) Horny sponges and their affairs: On the phylogenetic relationships of keratose sponges. *Mol Phylogenet Evol* 63: 809–816.
60. Vacelet J, Borchellini C, Pérez T, Bultel-Poncé V, Brouard J-P, et al. (2000) Morphological, chemical, and biochemical characterization of a new species of sponges without skeleton (Porifera, Demospongiae) from the Mediterranean Sea. *Zoosystema* 22: 313–326.
61. Erwin PM, Thacker RW (2007) Phylogenetic analyses of marine sponges within the order Verongida: a comparison of morphological and molecular data. *Invertebr Biol* 126: 220–234.

62. van Soest RWM, Hooper JNA (2002) Order Haplosclerida Topsent, 1928. In: Hooper JNA, van Soest RWM, editors. *Systema Porifera. A guide to the classification of sponges*. New York: Kluwer Academic/Plenum Publishers. pp. 831–832.
63. Itskovich VB, Belikov SI, Efremova SM, Masuda Y, Perez T, et al. (2007) Phylogenetic relationships between freshwater and marine Haplosclerida (Porifera, Demospongiae) based on full-length 18S rRNA and partial COXI gene sequences. In: Custódio MR, Lôbo-Hadju G, Hajdu Muricy G, editors. *Porifera Research: Biodiversity, Innovation and Sustainability. Série Livros. Museu Nacional, Rio de Janeiro*. pp. 383–391.
64. McCormack GP, Erpenbeck D, van Soest R (2002) Major discrepancy between phylogenetic hypotheses based on molecular and morphological criteria within the Order Haplosclerida (Phylum Porifera: Class Demospongiae). *J Zool System Evol Res* 40: 237–240.
65. Raleigh J, Redmond NE, Delehan E, Torpey S, van Soest RWM, et al. (2007) Mitochondrial Cytochrome oxidase 1 phylogeny supports alternative taxonomic scheme for the marine Haplosclerida. *J Mar Biol Ass UK* 87: 1577–1584.
66. Srivastava M, Simakov O, Chapman J, Fahey B, Gauthier MEA, et al. (2010) The *Amphimedon queenslandica* genome and the evolution of animal complexity. *Nature* doi:10.1038/nature09201
67. Addis JS, Peterson KJ (2005) Phylogenetic relationships of freshwater sponges (Porifera, Spongillina) inferred from analyses of 18S rDNA, COI mtDNA, and ITS2 rDNA sequences. *Zool Scr* 34: 549–557.
68. Holmes B, Blanch H (2007) Genus-specific associations of marine sponges with group I crenarchaeotes. *Mar Biol* 150: 759–772.
69. Erpenbeck D, Duran S, Rützler K, Paul V, Hooper JNA, et al. (2007) Towards a DNA taxonomy of Caribbean demosponges: a gene tree reconstructed from partial mitochondrial CO1 gene sequences supports previous rDNA phylogenies and provides a new perspective on the systematics of Demospongiae. *J Mar Biol Assoc UK* 87: 1563–1570.
70. Cárdenas P, Xavier JR, Reveillaud J, Schander C, Rapp HT (2011) Molecular phylogeny of the Astrophorida (Porifera, Demospongiae(p)) reveals an unexpected high level of spicule homoplasy. *PLoS ONE* 6(4), e18318. doi:10.1371/journal.pone.0018318
71. Maldonado M (2002) Family Pachastrellidae Carter, 1875. In: Hooper JNA, van Soest RWM, editors. *Systema Porifera. A guide to the classification of sponges*. New York: Kluwer Academic/Plenum Publishers. pp. 141–162.
72. Pisera A, Lévi C (2002) 'Lithistid' Demospongiae. In: Hooper JNA, van Soest RWM, editors. *Systema Porifera. A guide to the classification of sponges*. New York: Kluwer Academic/Plenum Publishers. pp. 299–301.
73. Kelly-Borges M, Pomponi S (1994) Phylogeny and classification of lithistid sponges (Porifera: Demospongiae): a preliminary assessment using ribosomal DNA sequence comparisons. *Mol Mar Biol Biotechnol* 3: 87–103.
74. Kober KM, Nichols SA (2007) On the phylogenetic relationships of Hadromerida and Poecilosclerid sponges. *J Mar Biol Assoc UK* 87: 1585–1598.
75. Erpenbeck D, List-Armitage SE, Alvarez B, Degnan BM, Hooper JNA, et al. (2007) The systematics of Raspailiidae (Demospongiae, Poecilosclerida, Microcionina) reanalysed with a ribosomal marker. *J Mar Biol Soc UK* 87: 1571–1576.
76. Hooper JNA, van Soest RWM (2002) Order Hadromerida Topsent, 1894. In: Hooper JNA, van Soest RWM, editors. *Systema Porifera. A guide to the classification of sponges*. New York: Kluwer Academic/Plenum Publishers. pp. 169–172.
77. Drummond AJ, Kearse M, Heled J, Moir R, Thierer T, et al. (2006) Geneious v1.2.1, Available: <http://www.geneious.com/>.
78. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, et al. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25: 3389–3402.
79. Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32:1792–1797.
80. Gouy M, Guindon S, Gascuel O (2010) SeaView version 4: a multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol* 27: 221–224.
81. Riesgo A, Pérez-Porro AR, Carmona S, Leys SP, Giribet G (2012) Optimization of preservation and storage time of sponge tissue samples to obtain quality mRNA for Next-Generation Sequencing with Illumina platforms. *Mol Ecol Resour* 12: 312–322.
82. Fortunato S, Adamski M, Bergum B, Guder C, Jordal S, et al. (2012) Genome-wide analysis of the sox family in the calcareous sponge *Sycon ciliatum*: multiple genes with unique expression patterns. *EvoDevo* 3:14.
83. Strimmer K, von Haeseler A (1997) Likelihood-mapping: a simple method to visualize phylogenetic content of a sequence alignment. *Proc Natl Acad Sci USA* 94: 6815–6819.
84. Stamatakis A (2006) RAXML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22: 2688–2690. doi:10.1093/bioinformatics/btl446
85. Lartillot N, Lepage T, Blanquart S (2009) PhyloBayes 3: a Bayesian software package for phylogenetic reconstruction and molecular dating. *Bioinformatics* 25: 2286–2288. doi:10.1093/bioinformatics/btp368
86. Lartillot N, Philippe H (2004) A Bayesian Mixture Model for Across-Site Heterogeneities in the Amino-Acid Replacement Process. *Mol Biol Evol* 21: 1095–1109.
87. Puigbò P, Garcia-Vallve S, McInerney JO (2007) TOPD/FMTS: a new software to compare phylogenetic trees. *Bioinformatics* 23: 1556–1558. doi:10.1093/bioinformatics/btm135
88. Creevey CJ, McInerney JO (2005) Clann: Investigating phylogenetic information through supertree analyses. *Bioinformatics* 21: 390–392.
89. Cummins CA, McInerney JO (2011) A method for inferring the rate of evolution of homologous characters that can potentially improve phylogenetic inference, resolve deep divergence and correct systematic biases. *Syst Biol* 60: 833–844.