


Deep-Sea Coral Taxa in the U.S. Southeast Region: Depth and Geographic Distribution (v. 2020)

by Thomas F. Hourigan¹, Stephen D. Cairns², John K. Reed³, and Steve W. Ross⁴

1. NOAA Deep Sea Coral Research and Technology Program, Office of Habitat Conservation, Silver Spring, MD
2. National Museum of Natural History, Smithsonian Institution, Washington, DC
3. Cooperative Institute of Ocean Exploration, Research, and Technology, Harbor Branch Oceanographic Institute, Florida Atlantic University, Fort Pierce, FL
4. Center for Marine Science, University of North Carolina, Wilmington

This annex to the U.S. Southeast chapter in “The State of Deep-Sea Coral and Sponge Ecosystems in the United States” provides a list of deep-sea coral taxa in the Phylum Cnidaria, Classes Anthozoa and Hydrozoa, known



to occur in U.S. waters from Cape Hatteras to the Florida Keys (Figure 1). Deep-sea corals are defined as azooxanthellate, heterotrophic coral species occurring in waters 50 meters deep or more. Details are provided on the vertical and geographic extent of each species (Table 1). This list is an update of the peer-reviewed 2017 list (Hourigan et al. 2017) and includes taxa recognized through 2019, including one newly described species. Taxonomic names are generally those currently accepted in the World Register of Marine Species (WoRMS), and are arranged by order, and alphabetically within order by family, genus, and species. Data sources (references) listed are those principally used to establish geographic and depth distribution.

Figure 1. U.S. Southeast region delimiting the geographic boundaries considered in this work. The region extends from Cape Hatteras to the Florida Keys and includes the Jacksonville Lithoherms (JL), Blake Plateau (BP), *Oculina*

Coral Mounds (OC), Miami Terrace (MT), Pourtales Terrace (PT), Florida Straits (FS), and Agassiz/Tortugas Valleys (AT). (Note: A number of additional species not included in this list occur in the Florida Straits in the waters of Cuba and the Bahamas, but have not been confirmed from adjacent U.S. waters).

Recommended citation: Hourigan TF, Cairns SD, Reed JK, Ross SW (2020) Deep-Sea Coral Taxa in the U.S. Southeast Region: Depth and Geographical Distribution (v. 2020). <https://deepseacoraldata.noaa.gov/library/2020-regional-deep-sea-coral-species-list>.

Table 1. List of known deep-sea coral species in Phylum Cnidaria, Class Anthozoa and Class Hydrozoa, and their reported distributions in the Southeast U.S. region. Blue shaded fields indicate newly described species since 2017. Bold text indicates changes to the list found in Hourigan et al. (2017), including additions or range extensions, denoted with an asterisk (*), and changes in taxonomy since 2017 denoted with a cross (†) (e.g., species that were listed in 2017, but have since been given a new name or alternative spelling). “NR” indicates a lack of reported distribution or depth information. References are numbered to correspond with citations following the table.

Distribution: NC = off North Carolina; SC = off South Carolina; GA = off Georgia; BP = Blake Plateau; FL = off Florida (NEFL = Northeast Florida, SEFL = Southeast Florida, FS = Florida Straits, JL = Jacksonville Lithoherms, OC = *Oculina* Mounds, MT = Miami Terrace, PT = Pourtalès Terrace, AT = Agassiz/Tortugas Valleys, DT = Dry Tortugas, and FK = Florida Keys). Note that locations in parentheses indicate specific localities within Florida where the species has been recorded. When a Museum collection record is the only reference for a species, the distribution listed may not be representative of the full distribution of the species.

Higher Taxon	Species	Distribution	Depth Range (m)	References
Class Anthozoa				
Subclass Hexacorallia				
Order Antipatharia				
Family Antipathidae	<i>Antipathes atlantica</i> Gray, 1857	NC-SC	20-91	1,2,3
	<i>Antipathes gracilis</i> Gray, 1860	NC-NEFL	31-88	1,2
	<i>Antipathes lenta</i> Pourtalès, 1871	SC-NEFL	20-100	1,3
	<i>Stichopathes luetkeni</i> Brook, 1889 (= <i>Stichopathes lutkeni</i> , alternative spelling)	SEFL(OC)	50-90	4,5
	<i>Stichopathes occidentalis</i> (Gray, 1857)	NC	67	1
	*<i>Stichopathes pourtalesi</i> Brook, 1889	FL(PT)	82-229	1,6
	<i>Stichopathes</i> sp.	NC-FS(PT)	30-314	1,5
Family Aphanipathidae	cf. <i>Phanopathes rigida</i> (Pourtalès, 1880) [= cf. <i>Antipathes rigida</i> (Pourtalès, 1880)]	FS(PT)	250-320	7
Family Cladopathidae	<i>Chrysopathes micracantha</i> Opresko & de Laia Loiola, 2008	BP(GA, NEFL)	658-871	1,8
	<i>Heteropathes americana</i> (Opresko 2003) (= <i>Heliopathes americana</i> Opresko 2003)	GA	1682-2200	1,9
Family Leiopathidae	<i>Leiopathes glaberrima</i> (Esper, 1788)	GA, FL	37-475	2,10,11
Family Myriopathidae	<i>Tanacetipathes barbadensis</i> (Brook, 1889)	FL	13-285	1
	<i>Tanacetipathes hirta</i> (Gray, 1857)	NC-FL	25-400	3,4
Family Schizopathidae	† <i>Bathypathes</i> sp. (= <i>Bathypathes</i> cf. <i>alternata</i> Brook, 1889)	BP, SEFL	364-716	1,7,12
	<i>Bathypathes patula</i> Brook, 1889	BP	762	1
	<i>Parantipathes tetrasticha</i> (Pourtalès, 1868)	FS(PT)	176-225	1,7
	*<i>Stauropathes</i> sp. cf. <i>S. punctata</i> (Roule, 1905)	BP	1217-1321	1
Family Stylopathidae	<i>Stylopathes litocrada</i> Opresko, 2006	FL(OC,PT)	91-380	1,13

Higher Taxon	Species	Distribution	Depth Range (m)	References
Order Scleractinia				
Family Anthemiphylliidae	<i>Anthemiphyllia patera patera</i> Pourtalès, 1878	SEFL	500-700	14
Family Caryophylliidae	<i>Anomocora fecunda</i> (Pourtalès, 1871)	FS	37-640	15,16
	<i>Anomocora marchadi</i> (Chevalier, 1966) (= <i>Asterosmilia marchadi</i> Chevalier, 1966)	SEFL	35-229	17
	<i>Anomocora prolifera</i> (Pourtalès, 1871) (= <i>Asterosmilia prolifera</i> Pourtalès, 1871)	SEFL,FS	30-329	14,17
	<i>Caryophyllia</i> (<i>Caryophyllia</i>) <i>ambrosia caribbeana</i> Cairns, 1979	FS(PT)	183-2360	1,14,17
	<i>Caryophyllia</i> (<i>C.</i>) <i>berteriana</i> Duchassaing, 1850	SEFL	99-1033	1,15
	<i>Caryophyllia</i> (<i>C.</i>) <i>paucipalata</i> Moseley, 1881	SEFL	293-752	1,15
	<i>Caryophyllia</i> (<i>C.</i>) <i>polygona</i> Pourtalès, 1878	GA, SEFL	310-1817	15
	<i>Cladocora debilis</i> Milne Edwards & Haime, 1849 ^a	FS	11-400	1,16
	<i>Coenosmilia arbuscula</i> Pourtalès, 1874	FS	74-622	1
	<i>Concentrotheca laevigata</i> (Pourtalès, 1871)	SC-FS(PT)	183-576	1,15
	<i>Crispatotrochus squiresi</i> (Cairns, 1979)	SC-SEFL	46-807	1,15
	<i>Dasmosmilia lymani</i> (Pourtalès, 1871)	NC-FS(PT)	37-366	1,15
	<i>Dasmosmilia variegata</i> (Pourtalès, 1871)	FS	110-421	1,15
	<i>Desmophyllum dianthus</i> (Esper, 1794)	NC-FS	183-2250	1,14
	<i>Labyrinthocyathus facetus</i> Cairns, 1979	SC-SEFL	385-768	1,15
	<i>Labyrinthocyathus langae</i> Cairns, 1979	GA-FS	506-810	1,15
	<i>Lophelia pertusa</i> (Linnaeus, 1758) †[= <i>Desmophyllum pertusum</i> (Linnaeus, 1758)] ^b	NC-FS(PT)	146-895	4,7,14,15,17
	<i>Oxysmilia rotundifolia</i> (Milne Edwards & Haime, 1849)	SC-FS	46-640	1,15
	<i>Paracyathus pulchellus</i> (Philippi, 1842)	NC-FS(PT)	17-250	1,15
	<i>Phacelocyathus flos</i> (Pourtalès, 1878)	FS	20-560	1
	<i>Phyllangia americana americana</i> Milne Edwards & Haime, 1849	NC-FL	0-183	4,5,16
	<i>Phyllangia pequegnatae</i> Cairns, 2000	GA	48-112	1,17
	<i>Polycyathus senegalensis</i> Chevalier, 1966	NC-SEFL	12-143	1,17
	<i>Pourtalosmilia conferta</i> Cairns, 1978	NC,SEFL	55-191	1,17
	<i>Premocyathus cornuiformis</i> (Pourtalès, 1868)	GA-FS	137-931	1
	<i>Rhizosmilia maculata</i> (Pourtalès, 1874) (= <i>Coenocyathus bartschi</i> Wells, 1947)	SEFL-DT	1-1317	1,17
	<i>Solenosmilia variabilis</i> Duncan, 1873	GA-FS	220-1383	12,14,15,17,18
	<i>Stephanocyathus</i> (<i>Odontocyathus</i>) <i>coronatus</i> (Pourtalès, 1867)	BP,FS	543-1250	1
	<i>Stephanocyathus</i> (<i>Stephanocyathus</i>) <i>diadema</i> (Moseley, 1876)	BP,FS	700-2683	1,15
	<i>Stephanocyathus</i> (<i>S.</i>) <i>laevifundus</i> Cairns, 1977	SEFL	300-1158	1,15
<i>Stephanocyathus</i> (<i>S.</i>) <i>paliferus</i> Cairns, 1977	FL-FS	220-715	15	
<i>Tethocyathus cylindraceus</i> (Pourtalès, 1868)	FS(PT)	183-649	1	
<i>Tethocyathus recurvatus</i> (Pourtalès, 1878)	FS(PT)	320-488	15,16	

Higher Taxon	Species	Distribution	Depth Range (m)	References
Family Caryophylliidae, cont.	<i>Tethocyathus variabilis</i> Cairns, 1979	SEFL	250-576	15,16
	<i>Thalamophyllia gombergi</i> Cairns, 1979	FS(PT)	155-220	1
	<i>Thalamophyllia riisei</i> (Duchassaing & Michelotti, 1860)	FS	4-914	1
	<i>Trochocyathus (Trochocyathus) rawsonii</i> Pourtalès, 1874	SC-FS(PT)	55-700	1,15
Family Deltocyathidae	<i>Deltocyathus calcar</i> Pourtalès, 1874	NC-FS(PT)	81-675	15
	<i>Deltocyathus eccentricus</i> Cairns, 1979	SC-FS(PT,AT)	183-910	15
	<i>Deltocyathus italicus</i> (Michelotti, 1838)	NEFL-FS	403-2634	14,16
	<i>Deltocyathus moseleyi</i> Cairns, 1979	SEFL	201-777	1,15
	<i>Deltocyathus pourtalesi</i> Cairns, 1979	SC-FS(PT)	311-567	1,15
Family Dendrophylliidae	<i>Balanophyllia (Balanophyllia) floridana</i> Pourtalès, 1868	NC-FS	13-220	1,17
	<i>Bathypsammia fallosocialis</i> Squires, 1959	SC-SEFL	213-805	1,15
	<i>Bathypsammia tintinnabulum</i> (Portalès, 1868)	SC-FS	210-1115	1,15
	<i>Cladopsammia manuelensis</i> (Chevalier, 1966)	FS(PT)	70-366	1,14
	<i>Dendrophyllia alternata</i> Pourtalès, 1880	SEFL(BP)	276-900	1
	<i>Eguchipsammia cornucopia</i> (Portalès, 1871)	FS(PT)	91-300	1,14
	<i>Eguchipsammia gaditana</i> (Duncan, 1873)	NC,GA	97-505	1,17
	<i>Eguchipsammia strigosa</i> Cairns, 2000	NC	25-77	1,17
	<i>Enallopsammia profunda</i> (Portalès, 1867)	NC(NBP)-FS(PT,AT)	305-1748	5,7,14,19
	<i>Enallopsammia rostrata</i> (Portalès, 1878)	SC	300-1646	14,15
	<i>Thecopsammia socialis</i> Pourtalès, 1868	NC(NBP)-FS(PT)	214-878	1,15
Family Flabellidae	<i>Flabellum (Flabellum) atlanticum</i> Cairns, 1979	SEFL	357-618	1
	<i>Flabellum (F.) floridanum</i> Cairns, 1991 (= <i>Flabellum fragile</i> Cairns, 1977)	FS(FK)	86-366	1
	*<i>Flabellum (Ulocyathus) angulare</i> Moseley, 1876	NC	2266-3186	1
	<i>Flabellum (U.) moseleyi</i> Pourtalès, 1880	FS	216-1097	1,15
	<i>Javana cailleti</i> (Duchassaing & Michelotti, 1864)	SEFL, FS	30-1809	14
	<i>Polymyces fragilis</i> (Portalès, 1868)	FS(PT)	75-822	14,15
Family Fungiacyathidae	<i>Fungiacyathus (Bathyactis) crispus</i> (Portalès, 1871)	NC-FL	183-1115	14
	<i>Fungiacyathus (B.) symmetricus</i> (Portalès, 1871)	SEFL,FS	183-1664	15
	<i>Fungiacyathus (Fungiacyathus) pusillus</i> (Portalès, 1868)	FS(PT)	285-685	15
Family Gardineriidae	<i>Gardineria paradoxa</i> (Portalès, 1868)	FL(FS)	91-700	15
Family Oculinidae	<i>Madrepora carolina</i> (Portalès, 1871)	NC-FL	53-1003; commonly 200-300	5,14,15,17,18
	<i>Madrepora oculata</i> Linnaeus, 1758	NC-FL	80-1500	4,7,14,15,17,18
	<i>Oculina tenella</i> Pourtalès, 1871	SEFL,FS	25-159	17
	<i>Oculina varicosa</i> Lesueur, 1821	NC-FL	3-150	1,4,14,15,17,18, 20

Higher Taxon	Species	Distribution	Depth Range (m)	References
Family Pocilloporidae	<i>Madracis asperula</i> Milne Edwards & Haime, 1849	FS	24-311	1,17
	<i>Madracis myriaster</i> (Milne Edwards & Haime, 1849)	NC-FS(PT)	20-1220 Commonly (150-300)	1,4,5,12,15,17
	† <i>Madracis pharensis</i> (Heller, 1868) ^f [= <i>Madracis pharensis f. pharensis</i> (Heller, 1868) in part]	NC	6-333	17
Family Rhizangiidae	<i>Astrangia poculata</i> (Ellis & Solander, 1786)	NC to FL	0-263	14
Family Schizocyathidae	<i>Pourtalocyathus hispidus</i> (Pourtalès, 1878)	FL-FS	349-1006	1,15
	<i>Schizocyathus fissilis</i> Pourtalès, 1874	SEFL	88-1300	16,17
Family Stenocyathidae	<i>Stenocyathus vermiformis</i> (Pourtalès, 1868)	GA(BP), FS(PT)	165-835	14
Family Turbinoliidae	<i>Cryptotrochus carolinensis</i> Cairns, 1988	NC (NBP)	320-338	1,21
	<i>Deltocyathoides stimpsonii</i> (Pourtalès, 1871)	SC(BP)-FS	110-553	1,15
	<i>Peponocyathus folliculus</i> (Pourtalès, 1868)	NC-FS	284-457	1,14

Higher Taxon	Species	Distribution	Depth Range (m)	References
Class Anthozoa				
Subclass Octocorallia				
Order Alcyonacea				
Family Acanthogorgiidae	* <i>Acanthogorgia armata</i> Verrill, 1878	NC, GA	27-538	1
	<i>Acanthogorgia aspera</i> Pourtalès, 1867	SBP-FS(PT)	56-729	1,22
Family Alcyoniidae	<i>Anthomastus agassizii</i> Verrill, 1922	GA,FL	320-3186	1,7,23,24
	<i>Anthomastus grandiflorus</i> Verrill, 1878	NC	137-457 (750-2919)	1,22,23
	<i>Bellonella rubistella</i> (Deichmann, 1936)	SC-SEFL	24-366	1,25
	* <i>Pseudoanthomastus</i> sp.	FL(SBP)	712	1
Family Anthothelidae	<i>Anthothela grandiflora</i> (Sars, 1856) ^c	FL(JL,FS)	744-868	1
	<i>Anthothela tropicalis</i> Bayer, 1961	BL-FS	173-805	1,26
	<i>Iciligorgia schrammi</i> Duchassaing, 1870	SEFL	11-366	4,5,27
Family Chrysogorgiidae	<i>Chrysogorgia herdendorfi</i> Cairns, 2001	BP	2178-2200	22,28
	<i>Chrysogorgia multiflora</i> Deichmann, 1936	GA-FS	1021-1200	22,28
	<i>Chrysogorgia spiculosa</i> (Verrill, 1883)	FS	914-2265	1,22,28
	<i>Chrysogorgia squamata</i> (Verrill, 1883)	BP,JL	431-1046	1,7,22
	<i>Distichogorgia sconsa</i> Bayer, 1979	BP	814	22,29
	<i>Flagelligorgia gracilis</i> Cairns & Cordeiro, 2017	NC-FS	196-567	1,30
	<i>Iridogorgia pourtalesii</i> Verrill, 1883	FS(PT)	(286) 558-1633	31,32
	* <i>Metallogorgia</i> sp. ^d	BP	1248	33,34
<i>Radicipes gracilis</i> (Verrill, 1884)	NC-SEFL	196-567	1,22	

Higher Taxon	Species	Distribution	Depth Range (m)	References
Family Clavulariidae	<i>Carijoa riisei</i> (Duchassaing & Michelotti, 1860)	SC-FK	0-104	4,5,25
	<i>Clavularia modesta</i> (Verrill, 1874)	SC,GA	29-861	1
	<i>Scleranthelia rugosa</i> (Pourtalès, 1867)	SC-SEFL	70-586	1,25
	<i>Telesto fruticulosa</i> Dana 1846	NC-NEFL	7-100	1,22,25
	<i>Telesto nelleae</i> Bayer, 1961	NC	27-227 (1153)	1,25,27
	<i>Telesto sanguinea</i> Deichmann, 1936	SC-FK	18-134	1,25
	<i>Telestula tubaria</i> (Wright & Studer, 1889)	FK	1281-1464	1
	† <i>Trachythela</i> sp. cf. <i>T. rudis</i> Verrill, 1922 [= <i>Clavularia</i> cf. <i>rudis</i> (Verrill, 1922)]	NEFL(BP)	805	1
Family Coralliidae	<i>Corallium medea</i> Bayer, 1964	SEFL	544-1426	1,22
	* <i>Hemicorallium bayeri</i> (Simpson & Watling, 2010)	SEFL	695-776	1,35
	<i>Hemicorallium niobe</i> (Bayer, 1964) (= <i>Corallium niobe</i> Bayer, 1964)	SEFL	668-1609	22
Family Ellisellidae	<i>Ellisella elongata</i> (Pallas, 1766) (= <i>Ellisella barbadensis</i> ; <i>Ctenocella (Viminella) barbadensis</i> ; <i>Viminella barbadensis</i>)	SC-FS(PT)	12-453	12,25,36
	<i>Ellisella grandis</i> (Verrill, 1901) (= <i>Ctenocella (Ellisella) grandis</i>)	FS	42-305	36
Family Gorgoniidae	<i>Eunicella albatrossi</i> Stiasny	BP	530-950	1
	<i>Eunicella modesta</i> Verrill, 1883	BP-SEFL	475-880	1,7,22
	<i>Leptogorgia cardinalis</i> (Bayer, 1961)	NEFL	19-77 (309)	25,36
	<i>Leptogorgia euryale</i> (Bayer, 1952)	SC	15-91	25
	<i>Leptogorgia hebes</i> Verrill, 1869	NC-NEFL	8-116	4,25
	<i>Leptogorgia punicea</i> (Milne Edwards & Haime, 1857)	NC-SEFL	20-105	25
	<i>Leptogorgia stheno</i> (Bayer, 1952)	FS	26-183	1,36
	<i>Leptogorgia virgulata</i> (Lamarck, 1815)	NC-SEFL	2-82 (220)	5,25,36
Family Isididae	† <i>Acanella</i> sp. cf. <i>A. arbuscula</i> (Johnson, 1862) ^e [= <i>Acanella</i> cf. <i>eburnea</i> (Pourtalès, 1868)]	SC	309-2100	1,16
	* <i>Cladaris</i> sp.	BP	717	1
	<i>Isidella</i> sp. 1	FL(PT,JL)	744-762	1,7
	<i>Keratoisis flexibilis</i> (Pourtalès, 1868)	BP-FS(PT)	170-816	1,7,12
	<i>Keratoisis grayi</i> (Wright, 1869) (= <i>Keratoisis ornata</i> Verrill, 1878)	BP	274-3235	1,22
	<i>Lepidisis caryophyllia</i> Verrill, 1883	FS	1003-1064	1,36
	<i>Lepidisis longiflora</i> Verrill, 1883	SEFL	752-1235	1
Family Nephtheidae	* <i>Duva florida</i> (Rathke, 1806)	BP	712-732	1
	<i>Gersemia fruticosa</i> (Sars, 1860)	FL	770 (600-3100)	1,23
	<i>Pseudodrifia nigra</i> (Pourtalès, 1868) (= <i>Capnella nigra</i> ; <i>Eunephthya nigra</i> ; <i>Nephthya nigra</i>)	SC-FS(PT)	60-1153	1,7,24,25
Family Nidaliidae	<i>Nidalia dissidens</i> Verseveldt & Bayer, 1988	SEFL	274-539	36
	<i>Nidalia occidentalis</i> Gray, 1835	NC-SEFL	37-440	1,4,5,25

Higher Taxon	Species	Distribution	Depth Range (m)	References
Family Paragorgiidae	<i>Paragorgia arborea</i> (Linnaeus, 1758)	NC	200-800	16
	<i>Paragorgia johnsoni</i> Gray, 1862 (= <i>Paragorgia boschmai</i> Bayer, 1964)	SEFL	522-608	1,22
Family Plexauridae ^f	<i>Bebryce cinerea</i> Deichmann, 1936	SC	76-549	25
	<i>Bebryce parastellata</i> Deichmann, 1936	NC	40-586	25,37
	<i>Muricea laxa</i> Verrill, 1864 ^g	SEFL	18-128	1,36
	<i>Muricea pendula</i> Verrill, 1868 ^g	NC-FK	13-125	22,25
	<i>Muriceides hirta</i> (Pourtalès, 1868) (= <i>Trachymuricea hirta</i> , <i>Acanthogorgia hirta</i>)	FS(PT)	681-716	7
	<i>Paramuricea</i> sp.	SC	78	25
	<i>Paramuricea multispina</i> Deichmann, 1936	SEFL,FS(PT)	106-1289	1,7,12,38
	<i>Paramuricea placomus</i> (Linnaeus, 1758)	SC-FS(PT)	462-470	1,12,22,38
	<i>Placogorgia mirabilis</i> Deichmann, 1936	FS(PT)	53-261	12
	<i>Scleracis guadalupensis</i> (Duchassaing & Michelotti, 1860)	NC,FL	18-715	25
	<i>Spinimuricea atlantica</i> (Johnson, 1862) (= <i>Echinomuricea cf. atlantica</i>)	FS(MT)	284-323	7
	<i>Swiftia casta</i> (Verrill, 1883)	GA,FS	(53) 102-937	1,12,31,39
	<i>Swiftia exserta</i> (Ellis & Solander, 1786)	JL,MT	21-177 (342, 494)	1,4,36
	<i>Swiftia koreni</i> (Studer, 1889)	JL,FS(PT)	(97) 221-985	1,36
	<i>Swiftia pourtalesii</i> Deichmann, 1936	FS	174-221	22,31,37
	*<i>Thesea citrina</i> Deichmann, 1936	NC	76-159	1
	*<i>Thesea gracilis</i> (Gray, 1868)	NC	73-110	1
	*<i>Thesea granulosa</i> Deichmann, 1936	FS(PT)	174-210	1
	<i>Thesea parviflora</i> Deichmann, 1936	NC (Cape Hatteras), FS(PT)	81-183	1,12
	<i>Thesea solitaria</i> (Pourtalès, 1868)	FS	185-318	36,40
<i>Villogorgia</i> sp. cf. <i>V. nigrescens</i> Duchassaing & Michelotti, 1860	FS(PT)	215	12	
Family Primnoidae	<i>Acanthoprímnoa goesi</i> (Aurivillius, 1931)	FS(PT)	137-595 (711)	22,41
	<i>Callogorgia americana</i> Cairns & Bayer 2002 (= <i>Callogorgia americana americana</i>)	FS(PT)	183-732	22,42,43
	<i>Callogorgia gracilis</i> (Milne Edwards & Haime, 1857)	SEFL	82-514	16,22,42,43
	<i>Calyptrophora gerdæ</i> Bayer, 2001	FS	229-556	16,22
	<i>Calyptrophora trilepis</i> (Pourtalès, 1868)	BP	593-911	22,44
	<i>Candidella imbricata</i> (Johnson, 1862)	BP,SEFL	514-2063	7,22,41
	<i>Narella bellissima</i> (Kükenthal, 1915)	SEFL	439-841	22
	<i>Narella pauciflora</i> Deichmann, 1936	SEFL	738-1542	22
	<i>Narella regularis</i> (Duchassaing & Michelotti, 1860)	SEFL	494-868	1,22

Higher Taxon	Species	Distribution	Depth Range (m)	References
Family Primnoidae, cont.	<i>Narella versluysi</i> (Hickson, 1909)	SEFL	668-796 (2118)	22,45
	<i>Paracalyptrophora duplex</i> Cairns & Bayer, 2004	FL	374-555	22,41
	<i>Plumarella aurea</i> (Deichmann, 1936)	SC-FL	310-878	16,22,41
	<i>Plumarella dichotoma</i> Cairns & Bayer, 2004	SC-FL	494-1065	22,41
	<i>Plumarella laxiramosa</i> Cairns & Bayer, 2004	NC,SC	348-572	22,41
	<i>Plumarella pellucida</i> Cairns & Bayer, 2004	NC-FS	549-1160	1,22,41
	<i>Plumarella pourtalesii</i> (Verrill, 1883)	NC-FS	183-893	1,7,12,22,41
	*<i>Thouarella (Euthouarella) grasshoffi</i> Cairns, 2006	BP	720-1760	1,46
	<i>Thouarella (Thouarella) bipinnata</i> Cairns, 2006	BP-SEFL	507-1000	1,22,46
Family Spongiodermidae	<i>Callipodium rubens</i> (Verrill, 1872) (= <i>Anthopodium rubens</i> Verrill, 1872)	NC,OC	9-92	5,27
	<i>Diodogorgia nodulifera</i> (Hargitt & Rogers, 1901)	GA-FL	20-183	1,4,5,25,27
	<i>Titanideum frauenfeldii</i> (Kölliker, 1865)	NC-FL	13-293	1,4,5,27,47
Order Pennatulacea				
Family Anthoptilidae	<i>Anthoptilum grandiflorum</i> (Verrill, 1879) (= <i>Benthoptilum sertum</i> Verrill, 1885)	NC,SC,FL(BP)	274-3651	1,48
	<i>Anthoptilum murrayi</i> Kölliker, 1880	SC	430-2491	1
Family Funiculinidae	<i>Funiculina quadrangularis</i> (Pallas, 1766)	BP	55-2866	39
Family Kophobelemnidae	<i>Kophobelemnon stelliferum</i> (Müller, 1776)	NC,SC	393-2199	1
	<i>Sclerobelemnon theseus</i> Bayer, 1959	SC	79	25
Family Pennatulidae	*<i>Gyrophyllum hironellei</i> Studer, 1891	BP	940-1086	1
	*<i>Pennatula aculeata</i> Danielssen, 1860	NC	2914	1,33
Family Protoptilidae	*<i>Protoptilum carpenteri</i> Kölliker, 1872	NC	1910	1
Family Renillidae	<i>Renilla reniformis</i> (Pallas, 1766)	NC-FL	1-73 off NC	1,25
Family Umbellulidae	<i>Umbellula lindahli</i> Kölliker, 1875 (= <i>Umbellula lindahlia</i> – alternative spelling)	BP	549-3338	1
Family Virgulariidae	<i>Acanthoptilum</i> sp.	NC-FL	49-279	1,25,49
	<i>Acanthoptilum pourtalesii</i> Kölliker, 1870 ^h	FS	22-80	37
	<i>Stylatula elegans</i> (Danielssen, 1860)	SC-FL	20-812	1,25,49
	<i>Virgularia presbytes</i> Bayer, 1955	NC-FL	9-110	4,25,27

Higher Taxon	Species	Distribution	Depth Range (m)	References
Class Hydrozoa				
Subclass Hydroidolina				
Order Anthoathecata				
Family Stylasteridae	<i>Crypthelia floridana</i> Cairns, 1986	SBP,FS(PT)	593-823	50
	<i>Crypthelia glossopoma</i> Cairns, 1986	FS(PT)	198-864	50
	<i>Distichopora foliacea</i> Pourtalès, 1868	GA,FS(PT)	174-527	12,50,51
	<i>Errina cochleata</i> Pourtalès, 1867	FS	194-534	52
	<i>Lepidopora biserialis</i> Cairns, 1986	FS	196-370	52
	<i>Lepidopora glabra</i> (Pourtalès, 1867)	FS	159-2021	51,52
	<i>Pliobothrus echinatus</i> Cairns, 1986	FS(PT)	164-750	1,12,50,51
	<i>Pliobothrus symmetricus</i> Pourtalès, 1868	SC-FS(PT)	139-1100; commonly 150-400	50,51
	<i>Stylaster complanatus</i> Pourtalès, 1867	GA	183-707	50
	<i>Stylaster duchassaingi</i> Pourtalès, 1867	FS(PT)	40-915	1,51,52
	<i>Stylaster erubescens</i> Pourtalès, 1868	SC-FS	146-965; commonly 650-850	12,50,51
	<i>Stylaster filogranus</i> Pourtalès, 1871	PT	175-549	12,51
	<i>Stylaster laevigatus</i> Cairns, 1986	SC	20-1170; commonly 300-400	50,51
	<i>Stylaster miniatus</i> (Pourtalès, 1868)	SC-FS(PT)	146-530	12,50,52
<i>Stylaster roseus</i> (Pallas, 1766)	SC(BP)	2-494; commonly <100	51,52	

Notes

- WoRMS lists the family for genus *Cladocora* as uncertain (*Scleractinia incertae sedis* – temporary name).
- Transfer of *Lophelia pertusa* to the genus *Desmophyllum* has been proposed recently based on genetic similarity of mitochondrial genomes and microsatellites (Addamo et al. 2016), and this change has been accepted by WoRMS. However, because of the significant morphological difference between these two genera and a difference of opinion even among molecular scientists, we suggest delaying this transfer until additional molecular studies are done on more genes.
- Moore et al. (2017) re-described *Anthothela grandiflora* (Sars, 1856) and placed many specimens previously identified as this species in a different genus: *Lateothela grandiflora* (Tixier-Durivault & D'Hondt, 1974). They did not examine the two specimens from the Southeast identified as *A. grandiflora*, however they occur within the range identified for this species. *A. grandiflora* and *L. grandiflora* co-occur in many localities, so either or both may be represented in this region.
- Records of *Metallogorgia* sp. are only from video collected from recent surveys in the region (NOAA Ship *Okeanos Explorer*, cruise EX1806), however the morphology of this genus is very distinctive, and specimens have been collected from nearby waters in the Bahamas (USNM 55918) and Cuba (USNM 100892).
- Saucier et al. (2017) have proposed that *Acanella eburnea* be synonymized with *A. arbuscula*. Reference: Saucier EH, Sajjadi A, France SC (2017) A taxonomic review of the genus *Acanella* (Cnidaria: Octocorallia: Isididae) in the North Atlantic Ocean, with descriptions of two new species. *Zootaxa* 4323:359-390.

- f. One plexaurid, *Pseudoplexaura porosa* (Houttuyn, 1772), which was included in our 2017 list, has been removed. Sanchez & Wirshing (2005) and Schubert et al. (2016) reported that *P. porosa* is a zooxanthellate species. The latter also indicate that this species occurs primarily shallower than 50m, so the previously reported deepest record (283m) may be incorrect.
- g. Schubert et al. (2016) have identified *Muricea laxa* and *M. pendula* as zooxanthellate octocorals, however, the depth range of these species in the Southeast U.S. is significantly deeper than most other zooxanthellate octocorals. Sánchez et al. (2019) identifies the genus *Muricea* as aposymbiotic.
- h. Deichmann (1936) identified that two specimens of *Acanthoptilum pourtalesii* Kölliker, 1870 were collected in the Florida Straits by Pourtales, but noted that she did not examine specimens of this species.

Acknowledgements

This list represents an update of Hourigan et al. (2017), which in turn built upon an initial list by Ross and Nizinski (2007). We would like to thank Drs. Sandra Brooke, Martha Nizinski, Andrea Quattrini, Marcelo Kitahara, Alberto Lindner, Dennis Opresko, and Daniel Wagner, who provided thoughtful reviews. Dr. Heather Coleman assisted greatly with the references and design.

Literature Cited

- Addamo AM, Vertino A, Stolarski J, Garcia-Jimenez R, Taviani M, Machordom A (2016) Merging scleractinian genera: the overwhelming genetic similarity between solitary *Desmophyllum* and colonial *Lophelia*. *BMC Evol Biol* 16:10
- Hourigan TF, Cairns SD, Reed JK, Ross SW (2017) Deep-Sea Coral Taxa in the U.S. Southeast Region: Depth and Geographical Distribution. Online resource: <https://deepseacoraldata.noaa.gov/library/2017-state-of-deep-sea-corals-report>. National Oceanic and Atmospheric Administration
- Ross SW, Nizinski MS (2007) State of Deep Coral Ecosystems of the U.S. Southeast Region: Cape Hatteras to Southeastern Florida. In: Lumsden SE, Hourigan TF, Bruckner AW, Dorr G (eds) [The State of Deep Coral Ecosystems of the United States](#). NOAA Technical Memorandum CRCP-3. Silver Spring, MD
- Sánchez JA, Dueñas LF, Rowley SJ, Gonzalez-Zapata FL, Vergara DC, Montaña-Salazar SM, Calixto-Botía I, Gómez CE, Abeytia R, Colin PL, Cordeiro RTS, Pérez CD (2019) Gorgonian Corals. In: Loya Y, Puglise KA, Bridge TC (eds) *Mesophotic Coral Ecosystems, Book Coral Reefs of the World 12*. Springer, Switzerland
- Schubert N, Brown D, Rossi S (2016) Symbiotic versus non-symbiotic octocorals: Physiological and ecological implications. In: Rossi S, Bramanti L, Orejas C (eds) *Marine Animal Forests: The Ecology of Benthic Biodiversity Hotspots*. Springer

References

1. National Museum of Natural History (NMNH) (2020) Invertebrate Zoology Collections - Online Collection Database; Accessed 07/20/2020. US National Museum of Natural History, Smithsonian Institution, Washington D.C.
2. Opresko DM (2009) Antipatharia (Cnidaria) of the Gulf of Mexico. In: Felder DL, Camp DK (eds) *Gulf of Mexico—Origins, Waters, and Biota Biodiversity*. Texas A&M University Press, College Station, Texas
3. Opresko DM, Sanchez JA (2005) Caribbean Shallow-water Black Corals (Cnidaria: Anthozoa: Antipatharia). *Caribbean Journal of Science* 41:492-507
4. Harter S, Reed JK, Farrington S, David A (2015) South Atlantic MPAs and Deepwater Coral HAPCs: Characterization of Fish Communities, Benthic Habitat, and Benthic Macrobiota; Final Report for 2012-2014 NOAA Cruises.
5. Reed JK Harbor Branch Oceanographic Museum Collection (unpublished).
6. Bo M, Opresko DM (2015) Redescription of *Stichopathes pourtalesi* Brook, 1889 (Cnidaria: Anthozoa: Antipatharia: Antipathidae). *Breviora* 540:1-18

7. Reed JK, Weaver DC, Pomponi SA (2006) Habitat and fauna of deep-water *Lophelia pertusa* coral reefs off the Southeastern USA: Blake Plateau, Straits of Florida, and Gulf of Mexico. *Bulletin of Marine Science* 78:343-375
8. Opresko DM, de Laia Loiola L (2008) Two New Species of *Chrysopathes* (Cnidaria: Anthozoa: Antipatharia) from the Western Atlantic. *Zootaxa* 1707:49-59
9. Opresko DM (2003) Revision of the Antipatharia (Cnidaria: Anthozoa). Part III. Cladopathidae. *Zool Med Leiden* 77:495-536
10. Williams B, Risk M, Ross SW, Sulak KJ (2007) Stable isotope records from deep-water antipatharians: 400-year records from the south-eastern coast of the United States of America. *Bulletin of Marine Science* 81:437-447
11. Opresko DM (1974) A study of the classification of the Antipatharia (Coelenterata:Anthozoa) with redescrptions of eleven species. Ph.D., University of Miami, Miami, FL
12. Reed JK, Pomponi SA, Weaver D, Paull CK, Wright AE (2005) Deep-water sinkholes and bioherms of South Florida and Pourtales Terrace- habitat and fauna. *Bulletin of Marine Science* 77:267-296
13. Opresko DM (2006) Revision of the Antipatharia (Cnidaria: Anthozoa). Part V. Establishment of a new family, Stylopathidae. *Zool Med Leiden* 80-4:109-138
14. Cairns SD, Chapman RE Biogeographic affinities of the North Atlantic deep-water Scleractinia. In: Willison JHM, Hall J, Gass SE, Kenchington ELR, Butler M, Doherty P (eds). *Proc Proceedings of the First International Symposium on Deep-Sea Corals*. Ecology Action Centre and Nova Scotia Museum
15. Cairns SD (1979) The deepwater Scleractinia of the Caribbean Sea and adjacent waters. *Studies on the fauna of Curaçao and other Caribbean Islands* 57:1-341
16. Ross SW, Nizinski MS (2007) State of Deep Coral Ecosystems in the U.S. Southeast Region: Cape Hatteras to Southeastern Florida. In: Lumsden SE, Hourigan TF, Bruckner AW, Dorr G (eds) *The State of Deep Coral Ecosystems of the United States*; NOAA Technical Memorandum CRCP-3. Dept. of Commerce, Silver Spring MD
17. Cairns SD (2000) A revision of the shallow-water azooxanthellate Scleractinia of the western Atlantic. *Studies on the Fauna of Curacao and other Caribbean Islands* 75:1-240
18. Cairns SD, Hoeksema BW, van der Land J (1999) Appendix: List of extant stony corals. *Atoll Research Bulletin* 459:13-46
19. Correa ML (2012) Environmental Controls of Cold-Water Coral Mound Distribution. Ph.D. Dissertation, University of Miami, Coral Gables, FL
20. Reed JK (1980) Distribution and structure of deep-water *Oculina varicosa* coral reeds off central eastern Florida. *Bulletin of Marine Science* 30:667-677
21. Cairns SD (1988) *Cryptotrochus*, new genus and two new species of deep-water corals (Scleractinia: Turbinoliidae). *Proc Biol Soc Wash* 101:709-716
22. Watling L, France SC, Pante E, Simpson A (2011) Biology of deep-water octocorals. *Adv Mar Biol* 60:41-122
23. Opresko DM (1980) Taxonomic description of some deep-sea octocorals of the Mid and North Atlantic. In: Hecker B, Blechschmidt G, Gibson P (eds) Appendix B in Canyon Assessment Study No BLM-AA551-CT8-49 US Department of Interior Bureau of Land Management, Washington, DC, 418 pp
24. Ross S, et al. Unpublished.
25. DeVictor ST, Morton SL (2010) Identification guide to the shallow-water (0-200 m) octocorals of the South Atlantic Bight. *Zootaxa* 2599:1-62
26. Moore KM, Alderslade P, Miller KJ (2017) A taxonomic revision of *Anthothela* (Octocorallia: Scleraxonia: Anthothelidae) and related genera, with the addition of new taxa, using morphological and molecular data. *Zootaxa* 4304:1-212
27. Bayer FM (1961) The shallow-water Octocorallia of the West Indian Region. *Studies on the Fauna of Curaçao and other Caribbean Islands* 12:1-373
28. Cairns SD (2001) Studies on western Atlantic Octocorallia (Coelenterata: Anthozoa). Part 1: The genus *Chrysogorgia* Duchassaing & Michelotti, 1864. *Proceedings of the Biological Society of Washington* 114:746-787

29. Bayer FM (1979) *Distichogorgia sconsa*, a new genus and species of chrysogorgiid octocoral (Coelenterata: Anthozoa) from the Blake Plateau off northern Florida, USA. *Proceedings of The Biological Society of Washington* 92:876-882
30. Cairns SD, Cordeiro RTS (2017) A new genus and species of golden coral (Anthozoa, Octocorallia, Chrysogorgiidae) from the Northwest Atlantic. *ZooKeys* 668:1-10
31. YPN (2015) Yale Peabody Museum of Natural History - Invertebrate Zoology Collections.
32. Watling L (2007) A review of the genus *Iridogorgia* (Octocorallia: Chrysogorgiidae) and its relatives, chiefly from the North Atlantic Ocean. *Journal of the Marine Biological Association of the UK* 87:393
33. NOAA (2019) Deep-Sea Coral and Sponge Database. NOAA EX1806 Video Observation
34. Salgado EJ, Etnoyer PJ (2020) Photographic catalog of deep-sea corals collected from the US West Atlantic margin by NOAA Ship *Okeanos Explorer* in years 2017- 2019. NOAA Technical Memorandum, NOS NCCOS 273 132 pp
35. Tu TH, Altuna A, Jeng MS (2015) Coralliidae (Anthozoa: Octocorallia) from the INDEMARES 2010 expedition to north and northwest Spain (northeast Atlantic), with delimitation of a new species using both morphological and molecular approaches. *Zootaxa* 3926:301-328
36. Cairns SD, Bayer FM (2009) Octocorallia (Cnidaria) of the Gulf of Mexico. In: Felder DL, Camp DK (eds) *Gulf of Mexico—Origins, Waters, and Biota Volume 1 Biodiversity*. Texas A&M Press, College Station, Texas
37. Deichmann E (1936) The Alcyonaria of the western part of the Atlantic Ocean. *Memoirs of the Museum of Comparative Zoölogy at Harvard College* 53:1–317, 337 pls.
38. Thoma JN (2013) Molecular and Morphological Diversity of Sea Fans with Emphasis on Deep-sea Octocorals of the Order Alcyonacea Lamouroux, 1812. Ph.D. Doctoral Dissertation, University of Louisiana at Lafayette,
39. Bayer FM (1952) New western Atlantic records of octocorals (Coelenterata: Anthozoa), with descriptions of three new species. *Journal of the Washington Academy of Sciences* 42:183-189
40. Museum of Comparative Zoology HUM (2016) Invertebrate Zoology Collections - Online Collection Database; Accessed 11/15/2016.
41. Cairns SD, Bayer FM (2004) Studies on Western Atlantic Octocorallia (Coelenterata: Anthozoa). Part 5: The Genera *Plumarella* Gray, 1870; *Acanthoprismoa*, n. gen.; and *Candidella* Bayer, 1954. *Proceedings of the Biological Society of Washington* 117:447-487
42. Cairns SD, Bayer FM (2002) Studies on western Atlantic Octocorallia (Coelenterata, Anthozoa): Part 2: The genus *Callogorgia* Gray, 1858. *Proceedings of the Biological Society of Washington* 115:840-867
43. Bayer FM, Cairns SD, Cordeiro RTS, Pérez CD (2015) New records of the genus *Callogorgia* (Anthozoa: Octocorallia) in the western Atlantic, including the description of a new species. *Journal of the Marine Biological Association of the United Kingdom* 95:905-911
44. Bayer FM (2001) New species of *Calyptraphora* (Coelenterata: Octocorallia: Primnoidae) from the western part of the Atlantic Ocean. *Proceedings of the Biological Society of Washington* 114:367-380
45. Cairns SD, Bayer FM (2003) Studies on western Atlantic Octocorallia (Coelenterata: Anthozoa). Part 3. The genus *Narella* Gray, 1870. *Proceedings of the Biological Society of Washington* 116:617-648
46. Cairns SD (2006) Studies on western Atlantic Octocorallia (Coelenterata, Anthozoa): Part 6: The genera *Primnoella* Gray, 1858; *Thouarella* Gray, 1870; *Dasystenella* Versluys, 1906. *Proceedings of the Biological Society of Washington* 119:161-194
47. Cairns SD, Wirshing HH (2015) Phylogenetic reconstruction of scleraxonian octocorals supports the resurrection of the family Spongiodermidae (Cnidaria, Alcyonacea). *Invertebrate Systematics* 29:345–368
48. Williams GC (2011) The Global Diversity of Sea Pens (Cnidaria: Octocorallia: Pennatulacea). *PLoS-One* 6(7): e22747
49. Bayer FM (1958) Additional records of western Atlantic octocorals. *Journal of the Washington Academy of Sciences* 47:379-390
50. Cairns SD (1986) A Revision of the Northwest Atlantic Stylasteridae (Coelenterata: Hydrozoa). *Smithsonian Contributions to Zoology* 418:1-131
51. Lindner A, Cairns SD, Cunningham CW (2008) From offshore to onshore: multiple origins of shallow-water corals

from deep-sea ancestors. Plos One 3:e2429

52. Calder DR, Cairns SD (2009) Hydroids (Cnidaria: Hydrozoa) of the Gulf of Mexico. In: Felder DL, Camp DK (eds) Gulf of Mexico: Origin, waters, and biota Volume 1, Biodiversity, Book 1. Texas A&M University, College Station, TX