



## Research article

[urn:lsid:zoobank.org:pub:A5B10105-A73F-4AB1-BF59-2095D95638F3](http://urn.lsid:zoobank.org:pub:A5B10105-A73F-4AB1-BF59-2095D95638F3)

# Redescription of *Psolus tessellatus* Koehler, 1896 (Echinodermata, Holothuroidea) with neotype designation

Claude MASSIN

Royal Belgian Institute of Natural Sciences, Department of Invertebrate Biology,  
29 rue Vautier, 1000 Brussels (Belgium), E-mail: [claude.massin@naturalsciences.be](mailto:claude.massin@naturalsciences.be)  
[urn:lsid:zoobank.org:author:896580A2-73E2-4562-AFCD-6AA69E5125B4](http://urn.lsid:zoobank.org:author:896580A2-73E2-4562-AFCD-6AA69E5125B4)

**Abstract.** In the bycatch of a scientific campaign in the Bay of Biscay (VITAL 2002) a third specimen of *Psolus tessellatus* Koehler, 1896 was collected very close to the type locality. This specimen is here fully illustrated and designated as the neotype.

**Key words.** Psolidae, Bay of Biscay, redescription, first illustration, neotype.

Massin C. 2013. Redescription of *Psolus tessellatus* Koehler, 1896 (Echinodermata, Holothuroidea) with neotype designation. *European Journal of Taxonomy* 38: 1-5. <http://dx.doi.org/10.5852/ejt.2013.38>

## Introduction

In 1895, the University of Lyon organized a campaign on board of the “Caudan” in the Bay of Biscay. During that expedition, material was collected between 110 and 1.710 m depth (Koehler 1896). Four new holothurian species were described and housed in the collections of the University of Lyon: *Holothuria roulei* Koehler, 1896 (= *Mesothuria roulei* (Koehler, 1896)), *Paroriza pallens* Koehler, 1896 (= *Stichopus pallens* (Koehler, 1896)), *Benthogone rosea* Koehler, 1896 and *Psolus tessellatus* Koehler, 1896. In 1985, Jangoux revisited the types in the collections of Lyon and only found the type material of *Holothuria roulei* and *Benthogone rosea*. The type material of *Paroriza pallens* and *Psolus tessellatus* was missing and must be considered as lost. Moreover, *P. tessellatus* was not illustrated by Koehler (1896).

A second catch of *P. tessellatus* was mentioned between the Azores and Madeira at 1.100 m depth by Fiege & Janssen (1998), who first illustrated the species.

In 2002, IFREMER Nantes organized a campaign (VITAL) in the Bay of Biscay for the evaluation of fish populations. In the bycatch of trawled material one specimen of *Psolus tessellatus* was collected. It is the object of the present note.

## Material and methods

The specimen of *Psolus tessellatus* was collected at Station 14 of the VITAL campaign (47°36' N – 8°25' W, 1399-1442 m, collecting date 23 Aug. 2002). It was fixed on a small stone (40 x 44 x 50 mm) and housed in the collection of the RBINS under the catalogue number IG31016-HOL 1304.

### Abbreviations

BIOGAS = BIOlogie golf GAScogne  
IFREMER = Institut Français de Recherche pour l'Exploitation de la Mer.  
RBINS = Royal Belgian Institute of Natural Sciences  
VITAL = Victor rove TALus continental

### Results

Phylum Echinodermata Brugière, 1791  
Order Denrochirotida Grube, 1840  
Family Psolidae Forbes, 1841  
Genus *Psolus* Oken, 1815

*Psolus tessellatus* Koehler, 1896  
Figs 1A-D, 2A-G

*Psolus tessellatus* Koehler, 1896: 119.

*Psolus tessellatus* – Mortensen 1927: 414. — Fiege & Janssen 1998: 341, figs 2-3.

*Psolus tessellatus* – Costelloe *et al.* 2001: 348.





Fig. 1. *Psolus tessellatus* Koehler, 1896. A. Dorsal view. B. Buccal cone. C. Anal cone. D. Ventral view.

**Redescription**

Small oval specimen, 17.4 mm long, 10.6 mm broad and 2.4 mm thick (Fig. 1A). Dorsal body wall covered by large scales (from 850 x 550 to 1350 x 1100  $\mu\text{m}$ ) (Figs 1A-C, 2A). The scale consists of one layer at the edge but rapidly becomes multilayered when going to the centre (Fig. 2B). No tube



**Fig. 2.** *Psolus tessellatus* Koehler, 1896. **A.** Dorsal scale. **B.** Detail of dorsal scale. **C.** Plates of ventral sole. **D.** Rods of large tube feet. **E.** End plate of large tube feet. **F.** End plate of small tube feet. **G.** Perforated plates at the base of the large tube feet.

feet dorsally. Ten scales between mouth and anus. Anus surrounded by irregular small plates (Fig. 1C). Mouth surrounded by 10 scales: five large triangular scales alternating with 5 small, elongated scales (Fig. 1A, B). Tentacles fully retracted, not observed.

Ventral sole very thin, translucent, calcareous ring and digestive tract visible by transparency (Fig. 1D). Ossicles as perforated plates, 100-160  $\mu\text{m}$  long with 1-8 holes (Fig. 2C). Ventral sole with 2 rows of tube feet at the edge (Fig. 1D). Inner row of tube feet large, with perforated rods, 125-280  $\mu\text{m}$  long (Fig. 2D) and large end plate in one piece (more or less 300  $\mu\text{m}$  across) (Fig. 2F). External row with very small tube feet with small end plates made of one piece (more or less 100  $\mu\text{m}$  across) (Fig. 2E). At the base of the large tube feet, perforated plates, 100-170  $\mu\text{m}$  long, similar to those of the ventral sole, sometimes knobbed (Fig. 2G). No mid ventral tube feet except 1-2 at the rear (Fig. 1D).

## Discussion

The specimen at hand belongs without any doubt to the genus *Psolus*: thick dorsal surface covered by scales, thin ventral sole, tube feet restricted to the ventral sole. Only 7 *Psolus* species are known from the Northeast Atlantic Ocean, viz *P. fabricii* (Duben & Koren, 1846), *P. hypsinosus* Heding, 1942, *P. nummularis* R. Perrier, 1899, *P. phantapus* (Strussenfeld, 1767), *P. pourtalesi* Théel, 1886, *P. squamatus* (Müller, 1776), and *P. tessellatus* Koehler, 1896.

*P. tessellatus* is readily separated from the other mentioned species by the valves surrounding the mouth: 5 large triangular ones alternating with 5 narrow ones. It is also the only *Psolus* species mentioned here with 10 scales present between mouth and anus. For *P. hypsinosus* and *P. fabricii* (see Heding 1942) there are only 5-7 scales between mouth and anus, whereas for *P. pourtalesi*, *P. squamatus* and *P. nummularis* (see Théel 1886; Deichmann 1930; Madsen & Hansen 1994; Perrier 1902) there are over 15. *P. tessellatus* is also the only *Psolus* species present in the Bay of Biscay.

The species seems to have a very restricted distribution and bathymetry in the Bay of Biscay: 45°47' N – 6°15' W and 1.700 m for the holotype, and 47°36' N – 8°25' W and 1.400 m for the present specimen. In deeper water (1900-4800 m) holothurians are abundant (34 species, Sibuet 1977) but *P. tessellatus* was not mentioned in spite of intensive sampling by IFREMER Brest (BIOGAS, Sibuet 1977; Sibuet & Segonzac 1985). IFREMER Nantes has intensively sampled at depths of 0-500 m with a few samples at 1000 m (J. Martin, personal communication; Sea Cruises Inventory 2012), but no *P. tessellatus* were collected.

The rarity of *P. tessellatus* could be an artefact linked to the low intensity of sampling between 1.000 and 1.700 m, a zone obviously favourable to the species (Fiege & Janssen 1998).

Since the holotype is lost and Koehler (1896) did not give any illustrations of the holotype (general aspect and ossicles) and since the specimen studied here was collected very close to the type locality, it is here designated as the neotype of *P. tessellatus* (see International Code of Zoological Nomenclature, articles 75, 75.3.4, 75.3.6.).

## Acknowledgements

It is a pleasure to thank Dr. Y. Samyn (RBINS) for comments on the first draft of this manuscript, Mme J. Martin (IFREMER Nantes) for allowing me to study material from the VITAL Campaign and Mr. A. Van Haelen for the photographs on Plate 1.

## References

- Costelloe M., Emblow C. & White R. 2001. *European Register of Marine Species: a check-list of the marine species in Europe and a bibliography of guides to their identification*. Patrimoines Naturels 50, Muséum National d'Histoire Naturelle, Paris.
- Deichmann E. 1930. The holothurians of the western part of the Atlantic Ocean. *Bulletin of the Museum of Comparative Zoology at Harvard College* 71 (3): 41-226.
- Fiege D. & Janssen R. 1998. Ein Fall von Parasitismus bei *Psolus tessellatus* einer ungewöhnlichen Holothurie. *Natur und Museum* 128 (10): 341-344
- Heding S.G. 1942. Holothuroidea II. *The Danish Ingolf-Expedition IV* (13): 1-39.
- Jangoux M. 1985. Catalogue commenté des types d'Echinodermes actuels conservés dans les collections lyonnaises. *Nouvelles Archives du Muséum d'Histoire naturelle de Lyon* 23: 3-11.
- Koehler R. 1896. Echinodermes. *Résultats Scientifiques de la campagne du «Caudan» dans le golfe de Gascogne* 1: 33-127.
- Madsen F.J. & Hansen B. 1994. *Echinodermata Holothuroidea*. Marine Invertebrates of Scandinavia 9, Scandinavian University Press, Norway.
- Mortensen T.H. 1927. *Handbook of the Echinoderms of the British Isles*. Oxford University Press, London.
- Perrier R. 1902. Holothuries. In: Perrier R. (ed.) *Expéditions Scientifiques du «Travailleur» et du «Talisman»*: 273-554. Masson, Paris.
- Sea Cruises Inventory. 2012. [http://www.ifremer.fr/sismer/UK/catal/campagne/indexorga2\\_armat.htql?corg=IFREM](http://www.ifremer.fr/sismer/UK/catal/campagne/indexorga2_armat.htql?corg=IFREM)
- Sibuet M. 1977. Répartition et diversité des Echinodermes (Holothurides-Astérides) en zone profonde dans le Golfe de Gascogne. *Deep-Sea Research* 24: 549-563.
- Sibuet M. & Segonzac M. 1985. Abondance et répartition de l'épifaune mégabenthique. In: Laubier L. & Moniot C. (eds) *Peuplements profonds du Golfe de Gascogne, campagne Biogas*: 143-156. IFREMER, Brest.
- Théel H. 1886. Reports on the Results of Dredging in the Gulf of Mexico (1877-78), in the Caribbean Sea (1879-80), and along the Eastern Coast of the United States during the Summer of 1880 by the U. S. Coast Survey Steamer "Blake". *Bulletin of the Museum of Comparative Zoology at Harvard College* 13 (1): 1-21.

*Manuscript received: 5 December 2012*

*Manuscript accepted: 28 January 2013*

*Published on: 7 February 2013*

*Topic editor: Rudy Jocqué*

*Desk editor: Kristiaan Hoedemakers*

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the *EJT* consortium: Muséum National d'Histoire Naturelle, Paris, France; National Botanic Garden of Belgium, Meise, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Natural History Museum, London, United Kingdom; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark.